

## **Response to the Editor's comments**

### **Comments to the Author:**

The language and grammar of the Main text should be improved and the Conclusions section is too much a repetition of the Abstract; it should also be improved. Both these comments can be dealt with after the ACPD discussion phase.

**Response:** Thanks for the handling of our paper. Following the Editor's recommendation, we have corrected the grammatical mistakes and significantly improved the English of the revised MS. We also revised the conclusion section carefully.

## **Responses to the Anonymous Referee's comments**

### **Referee #1 Evaluations**

The work by Haque et al presents a range of organic compounds in atmospheric aerosols from a year-long campaign in South Korea. The authors used well established analytical techniques for their analysis. The experimental procedure is well designed and in general the data is well presented. My detailed comments are listed below.

**Response:** Thanks for the careful reading and valuable suggestions to improve the scientific content of the manuscript. Following the reviewer's comment, we have carefully made corrections in the revised MS. Our responses to all comments made by the reviewer are given below. Please refer to the revised MS, in which changes are highlighted in yellow.

**Comment:** First of all, although the work is of relatively good quality (as can be judged from the draft); I felt it is largely of a routine nature (at least the way it is presented). I believe that such type of publications can be useful but if considered for publication at ACP they need to go under the Measurement report's category.

**Response:** We respectfully do not comply with the reviewer's argument but would like to highlight the two key features that make this study suitable for publication in ACP under the Research article category.

1. Here, we present a new finding on the contribution of coal combustion sources in East Asia in controlling the atmospheric levels of levoglucosan (*Lev*) apart from the traditional biomass/biofuel burning emissions. This is based on the prevailing linear relationship between the radiocarbon based nonfossil-EC and *Lev* in the year-round TSP samples collected from the Korean Climate Observatory at Gosan (KCOG) site in Jeju Island. The Gosan supersite is the best location to understand how the chemical composition of source-emissions from East Asia affects the outflow regions in winter and spring. Recent studies have highlighted the potential contribution of *Lev* from residential coal combustion in China (Yan et al., 2018), with estimated annual emission of ~2.2 Gg of *Lev* from domestic coal combustion (Wu et al., 2021). Given this background information, the prevailing significant linear relationship between *Lev* and nonfossil-EC (P-value < 0.05) over the KCOG clearly emphasizes the need

for reconsideration of the previous assessments on the impact of biomass burning in East Asian outflow to the western North Pacific.

2. Another important highlight is the detailed study on the atmospheric abundances of biogenic SOA tracers along with those of other primary biological particles from the KCOG for the year 2013. This dataset is further compared with the molecular distributions and relative abundances of these organic tracers in the TSP samples collected over Gosan during 2001, a decade ago (Fu et al., 2012). This comparison allows us to better understand the regional changes in the emission sources (e.g., fugitive dust, biomass burning, and fossil-fuel combustion) on a decadal basis.

These points have been added in the revised MS. Please see lines 682-697.

Major comment:

**Comment:** I would like to know if the authors considered atmospheric stability of the measured organic compounds especially in the light of a long-range transport or/and correlation and statistical analysis of the reported species. For example, it has been demonstrated that levoglucosan can decay in particles exposed to OH. *“The extent of decay ranged from 20 to 90% and was strongly correlated to the integrated OH exposure. Increased relative humidity did not enhance or impede reaction. Relative kinetics indicate that levoglucosan has an atmospheric lifetime of 0.7–2.2 days when biomass burning particles are exposed to  $1 \times 10^6$  molecules  $\text{cm}^{-3}$  of OH (typical average summertime conditions). This implies that levoglucosan reacts with OH on a timescale similar to that of transport and deposition, which has important implications for the use of levoglucosan as a tracer for biomass burning emissions in source apportionment studies”* Hennigan et al. 2010.

I trust the above also applies to other measured organic compounds for some of which the information on reactivity is not available. My question to the authors - how does this affect the presented results? The authors present correlation of different organic compounds and discuss their potential similar source based on these correlations and statistical analysis. Does the stability of considered organic compounds can play a role in the observed absence of correlation? The same applies to the discussion on long range transport of the observed organics. I doubt that stability of considered sugars in Figure 4 is the same for all compounds.

**Response:** As the reviewer rightly pointed out that levoglucosan (*Lev*) undergoes a photooxidation with OH radicals during atmospheric transport (half-life: <2.2 days) and, this would cause lower abundances of this anhydrosugar (Hennigan et al., 2010). *Lev* concentration measured over the KCOG includes contributions from both fossil and nonfossil sources in the East Asian outflow. In such a case, we found the moderately significant correlations of *Lev* with nonfossil EC (i.e., biomass burning) and fossil-EC (e.g., Miocene lignite coals) over the KCOG. It is very likely that this weakened relationship is because of the photo-degradation of *Lev* during atmospheric transport. This result would be an otherwise high atmospheric abundance of *Lev* and its pronounced linear relationships with the nonfossil- and fossil-EC, which implies a much stronger impact of both these source emissions in the East Asian continental outflow in winter and spring. This is also applicable for low concentrations and poor correlations of other organics such as primary saccharides and BSOA tracers. Considering the feasibility of photooxidation, it is likely that secondary OA undergoes much faster cycling than the primary sugar compounds. This would mean a slight underestimation of BSOA over the KCOG in the East Asian outflow and their atmospheric abundances over Gosan reflect a lower limit.

These explanations have been added in the revised MS. Please see lines 370-374, 452-453, 458-460, and 676-681.

**Comment:** Moreover, besides anhydrosugars, the authors state that “*Cis-pinonic acid showed poor correlation with pinic acid ( $r = 0.35$ ,  $p = 0.12$ ) (Figure 6f), suggesting that they formed from different monoterpenes such as  $\alpha$ -pinene,  $\beta$ -pinene, or  $\delta$ -limonene.*” Such assumptions can be made only if the measurements are taken very close to their emission sources **and/or** assuming that both compounds have similar atmospheric stability or reactivity. Is it the case for all considered compounds in this study? I believe this needs to be considered when discussing correlations or long range transport.

**Response:** We comply with the reviewer's view. Likely, the extent of loss of BSOA by the photooxidation during transport would also cause such poor correlations, influencing their abundances over KCOG. We have added a few sentences in the revised MS. Please see lines 476-478 and 499-501.

Minor comment:

**Comment:** It is hard to read Figure 2 (especially Figure 2a). Please either split the concentration axis so that all minor compounds can be clearly seen in the plot or move it into the SI and increase the size of the figure.

**Response:** Following the reviewer's suggestion, we have added an additional figure in the supplement for the visualization of minor compounds in the temporal plot. Please see Figure S2.

Reference:

Hennigan, C.J., Sullivan, A. P., Collett, J. L., Robinson, A. L.: Levoglucosan stability in biomass burning particles exposed to hydroxyl radicals, *Geophys. Res. Lett.*, 37, <https://doi.org/10.1029/2010GL043088>, 2010.

**Response:** Added in the reference list of the main text.

## **Reviewer #2 Evaluations**

The authors present the results of the chemical composition of organic aerosol in South Korea with a focus on primary organic compounds and biogenic secondary organic aerosol tracers. Seasonal variations and sources of studied compounds were analysed. The paper is suitable for publication in the journal Atmospheric Chemistry and Physics, however, several comments reported below should be addressed before acceptance for publication. Minor revisions of the paper are requested.

**Response:** We thank the reviewer for the positive assessment of this work. We have carefully revised the manuscript following the reviewer's comments and suggestions. Our responses to all comments made by the reviewer are given below. Please refer to the revised MS, in which changes are highlighted in yellow.

### **Comments:**

**Comment:** Lines 277-281: Glucose and other saccharides may be partly formed also during biomass burning, which explains their contribution during the winter period.

**Response:** Following the reviewer's comment, we have added the following sentence in the revised MS as: "Since glucose, fructose, and sucrose showed moderately significant correlations ( $R^2 = 0.44-0.48$ ,  $p < 0.01$ ) with levoglucosan in winter, it is somewhat possible that BB source emission could also influence the concentrations of these saccharides in this season (Haque et al., 2019; Fu et al., 2008)." Please see lines 279-283.

It can be noted that levoglucosan showed a poor/negative correlation with other saccharides except xylose (lines 404-405) and inositol in winter (discussed in the main text). Please see lines 302-305 in the revised MS.

**Comment:** Line 283-294: Sources of inositol and other polyols are only perfunctorily characterized focusing on prevailing sources. Sources of inositol are entirely missing.

**Response:** Following the reviewer's suggestion, more discussion on sources of polyols along with inositol has been added in the revised MS as: "The major sources of arabitol and mannitol are airborne fungal spores (Bauer et al., 2008), accompanying detritus from mature leaves (Pashynska et al., 2002). Heald and Spracklen (2009) reported that mannitol and arabitol are considerably associated with terrestrial biosphere activity. Inositol is largely derived from the developing leaves in summer (Pashynska et al., 2002) and BB in winter (Fu

et al., 2010b). Zhu et al. (2015b) found similar seasonal behavior of inositol with those of other sugar alcohols with the predominance in summer, associated with microbial activities in local forests from Okinawa. Inositol showed a moderately significant correlation with levoglucosan ( $R^2 = 0.33$ ,  $p < 0.01$ ) in winter; however, there were no positive linear relationships between levoglucosan and other sugar alcohols, implying a partial emission of inositol from the BB during winter in Gosan aerosols.” Please see lines 296-305. More information regarding the sources of polyols are already interpreted in section 3.4 (lines 409-419).

**Comment:** Line 306-307: Unlike other studies, the concentration of mannosan in this study is surprisingly lower than those of galactosan, do you have an explanation of this fact?

**Response:** Following the reviewer’s comment, we have added the following sentences in the revised MS as: “Galactosan is more abundant in crop-residue burning emissions than mannosan (Engling et al., 2009; Sheesley et al., 2003). It is very much likely that the impact of crop-residue burning emissions in East Asia over Gosan is more prominent in winter/spring. Such high abundances of galactosan over mannosan were found in the North China Plain (Fu et al., 2008) and in the Indo-Gangetic Plain outflow sampled over the Bay of Bengal (Bikkina et al., 2019).” Please see lines 320-325.

**Comment:** Line 400-404: Ratios levoglucosan/mannosan and levoglucosan/(mannosan + galactosan) also allow to distinguish biomass burning and lignite combustion as the source of these anhydrosugars. In addition, the empirical equation using levoglucosan and mannosan data allows you to calculate the contribution of softwood and hardwood to the total amount of combusted wood.

**Response:** Based on the reviewer’s comment, we have added the results of these diagnostic ratios in the revised MS as “Ratios of *Lev/Man* and *Lev/(Man + Gal)* can be useful to distinguish BB and coal combustion contributions. The average ratios of *Lev/Man* ( $15.1 \pm 6.76$ ) and *Lev/(Man + Gal)* ( $4.27 \pm 1.23$ ) in Gosan aerosols are much closer to those from wood burning and coal combustion sources emissions, respectively (Yan et al., 2018). It reveals that *Lev* could originate from both biomass and coal burning source emissions, which is consistent with the linear relationship between *Lev*-C and fossil-/nonfossil carbon fraction

(section 3.7). Please see lines 422-427 in the main text. Unfortunately, we are not familiar with the empirical equation to calculate the contribution of softwood and hardwood.

**Comment:** Line 426: Add references to levoglucosan degradation.

**Response:** Added. Please see line 453 in the revised MS.

**Comment:** Line 536: Add missing information into Table 3, such as analysed PM fraction and studied season.

**Response:** Added. Please see Table 3 in the revised MS.

**Comment:** Line 562: Another key factor is a higher concentration of ozone and other oxidation agents in summer.

**Response:** Added. Please see lines 591-592 in the revised MS.



**Regional heterogeneities in the emission of airborne primary sugar compounds and biogenic secondary organic aerosols in the East Asian outflow: Evidence for coal combustion as a source of levoglucosan**

**M. Mozammel Haque<sup>1,2,3</sup>, Yanlin Zhang<sup>1,2\*</sup> Srinivas Bikkina<sup>4</sup>, Meehye Lee<sup>5</sup>, and Kimitaka Kawamura<sup>3,4\*</sup>**

<sup>1</sup>*Yale-NUIST Center on Atmospheric Environment, International Joint Laboratory on Climate and Environment Change (ILCEC), Nanjing University of Information Science & Technology, Nanjing, 210044, China*

<sup>2</sup>*School of Applied Meteorology, Nanjing University of Information Science & Technology, Nanjing 210044, China*

<sup>3</sup>*Institute of Low Temperature Science, Hokkaido University, Sapporo 060-0819, Japan*

<sup>4</sup>*Chubu Institute for Advanced Studies, Chubu University, Kasugai 487-8501, Japan*

<sup>5</sup>*Department of Earth and Environmental Sciences, Korea University, Anam-dong, Sungbuk-gu, Seoul 136-701, South Korea*

*\*Corresponding author*

E-mail: dryanlinzhang@outlook.com (Yan-Lin Zhang)

E-mail: kkawamura@isc.chubu.ac.jp (Kimitaka Kawamura)

## ABSTRACT

Biomass burning (BB) significantly influences the chemical composition of organic aerosols (OA) in the East Asian outflow. Source apportionment of BB-derived OA is an influential factor for understanding their regional emissions, which is crucial for reducing uncertainties in their projected climate and health-effects. We analyzed here three different classes of atmospheric sugar compounds (anhydrosugars, primary sugars, and sugar alcohols) and two types of biogenic secondary organic aerosol (BSOA) tracers (isoprene- and monoterpene derived SOA products) in a year-long collected total suspended particulate matter (TSP) from an island-based receptor site in South Korea, the Gosan. We investigate seasonal variations in the source-emissions of BB-derived OA using mass concentrations of anhydrosugars and radiocarbon ( $^{14}\text{C}$ -) isotopic composition of organic carbon (OC) and elemental carbon (EC) in ambient aerosols. Levoglucosan (*Lev*) is the most abundant anhydrosugar, followed by galactosan (*Gal*) and mannosan (*Man*). Strong correlations of *Lev* with *Gal* and *Man*, along with their ratios (*Lev/Gal*:  $6.65 \pm 2.26$ ; *Lev/Man*:  $15.1 \pm 6.76$ ) indicate the contribution from hardwood burning emissions. The seasonal trends revealed that the BB impact is more pronounced in winter and fall, as evidenced by the high concentrations of anhydrosugars. Likewise, significant correlations among three primary sugars (i.e., glucose, fructose, and sucrose) emphasized the contribution of airborne pollen. The primary sugars showed higher concentrations in spring/summer than winter/fall. The fungal spore tracer compounds (i.e., arabinol, mannitol, and erythritol) correlated well with trehalose (i.e., a proxy for soil organic carbon), suggesting the origin from airborne fungal spores and soil microbes in the East Asian outflow. These sugar alcohols peaked in summer, followed by spring/fall and winter. Monoterpene-derived SOA tracers were most abundant compared to isoprene-SOA tracers. Both BSOA tracers were dominant in summer, followed by fall, spring, and winter. The source apportionment based on multiple linear regressions, diagnostic mass ratios, and positive matrix factorization analysis altogether revealed that biomass burning (41.9%) and biogenic SOA (21.1%) mostly dictate the OA loading in the ambient aerosols from East Asian outflow. We also found significant positive linear relationships of  $^{14}\text{C}$ -based nonfossil- and fossil-derived organic carbon fractions with *Lev-C* along with the comparable regression slopes, suggesting the importance of BB and coal combustion sources in the East Asian outflow.

**Keywords:** Biomass burning tracers, primary biological aerosol particles, biogenic SOA tracers, radiocarbon-based source apportionment, organic aerosols, East Asian outflow

## 1. Introduction

Organic aerosols (OA), which account for a major fraction of up to 50% of airborne total suspended particulate matter, have considerable effects on regional and global climate by absorbing or scattering sunlight (Kanakidou et al., 2005). However, the climate effects of OA are involved with large uncertainties due to our limited understanding of the contributing sources. OA can be derived from both primary emissions and secondarily formed species. Sugars are an important group of water-soluble, primarily organic compounds whose concentrations are significant in atmospheric aerosols over the continent (Jia and Fraser, 2011; Fu et al., 2008; Yttri et al., 2007; Graham et al., 2003). Anhydrosugars such as levoglucosan, galactosan, and mannosan are the key tracers of biomass burning (BB) emissions (Simoneit, 2002). Sugar alcohols, along with glucose, trehalose and sucrose are mostly originated from primary biological particles such as fungal spores, pollen, bacteria, and viruses, and vegetative debris (Graham et al., 2003; Simoneit et al., 2004a; Bauer et al., 2008; Deguillaume et al., 2008). Primary sugars and sugar alcohols are predominantly present in the coarse mode aerosols, accounting for 0.5-10% of atmospheric aerosol carbon matter (Yttri et al., 2007; Pio et al., 2008).

Secondary organic aerosol (SOA) is a large fraction of OA, while there were only limited studies about the key factors controlling SOA formation. The SOA formation significantly increased with the enhancement of the ambient aerosol mass (Liu et al., 2018). SOA is formed by both homogenous and heterogeneous reactions of volatile organic compounds (VOCs) in the atmosphere (Surratt et al., 2010; Robinson et al., 2007; Claeys et al., 2004). On a global estimation, biogenic VOCs (BVOCs) such as isoprene, monoterpenes (e.g.,  $\alpha/\beta$ -pinene), and sesquiterpenes (e.g.,  $\beta$ -caryophyllene) are one order of magnitude higher than those of anthropogenic VOCs (e.g., toluene) (Guenther et al., 2006). The global emissions of annual BVOCs were estimated to be 1150 TgC yr<sup>-1</sup>, accounting for 44% isoprene and 11% monoterpenes (Guenther et al., 1995). Isoprene is highly reactive and promptly reacts with oxidants such as O<sub>3</sub>, OH, and NO<sub>x</sub> in the atmosphere to form SOA (Kroll et al., 2005, 2006; Ng et al., 2008; Surratt et al., 2010; Bikkina et al., 2021), estimated to be 19.2 TgC yr<sup>-1</sup>, consisting of ~70% of the total SOA budget (Heald et al., 2008). Monoterpenes are important sources of biogenic secondary organic aerosol (BSOA), accounting for ~35% of the global BVOCs emissions (Griffin et al., 1999).

Anthropogenic activities such as coal and biofuel combustion over East Asia, including China, are responsible for the vast emission of OA (Huebert et al., 2003; Zhang et al., 2016). Understanding the ambient levels OA in the East Asian outflow is crucial for

assessing their regional climatic effects. As part of this effort, Korean Climate Observatory at Gosan (KCOG), a super site located in South Korea, is an ideal location for investigating the atmospheric outflow characteristics from East Asia (Fu et al., 2010a; Kundu et al., 2010; Ramanathan et al., 2007; Kawamura et al., 2004; Arimoto et al., 1996). For instance, primary OA associated with soil/desert dust in East Asia, along with forest fires in Siberia/northeastern China, are transported over Gosan in spring (Wang et al., 2009a). BSOA during long-range transport from the continent and Open Ocean, as well as local vegetation, can significantly contribute to Gosan aerosols. Although these investigations were carried out almost a decade ago, no such observations are available in contemporary times from Gosan. Here, we attempt to understand the current states of East Asian OA using both the molecular marker approach and radiocarbon data of carbonaceous components.

The KCOG, located on the western side of Jeju Island adjacent to the Yellow Sea and the East China Sea, is facing the Asian continent but is isolated from public areas of the island (Kawamura et al., 2004). Simoneit et al. (2004b) have documented during the ACE-Asia campaign that OA from the BB and fossil fuel combustion sources are transported along with desert dust to the KCOG during continental outflow. An intensive campaign was organized at the KCOG during spring 2005 to observe the physical properties of East Asian aerosols while two dust events were detected (Nakajima et al., 2007). Here, we focus on the characterization of airborne anhydrosugars, primary sugars, sugar alcohols, and BSOA tracers from the KCOG. Gosan is influenced by the continental outflow from East Asia during winter, spring and fall, whereas the site is influenced by the maritime air masses from the Pacific Ocean and other marginal seas. This makes the KCOG ideal for characterizing the regional heterogeneities in the emissions of organic compounds in the East Asian outflow based on the TSP samples collected during April 2013-April 2014.


## 2. Methods

### 2.1. Aerosol sampling and prevailing meteorology

Total suspended particles (TSP) were collected on pre-combusted (450°C for 6 h) quartz fiber filters (20 cm × 25 cm, Pallflex) at the KCOG (33.17 °N, 126.10 °E, see Figure 1), South Korea. To get enough signal for the radiocarbon measurements, each TSP sample was collected for 10–14 days from April 2013 to April 2014. Twenty-one samples were collected using a high-volume air sampler (Kimoto AS-810, ~65 m<sup>3</sup> h<sup>-1</sup>) installed on the rooftop of a trailer house (~3 m above the ground). After the collection, aerosol filters were transferred to a pre-combusted (450°C for 6 h) glass jar (150 mL) equipped with a Teflon-lined screw cap and transported to the laboratory in Sapporo. These TSP samples were stored in a dark

freezer room at -20°C until the analysis. Three field blank filters were also collected during the campaign.

The ambient temperatures at the Gosan site were on average 6.9°C in winter, 14.1°C in spring, 27.0°C in summer, and 17.1°C in fall. Likewise, the average relative humidity was found to be highest in summer (71.3%), followed by spring (64.9%), fall (63.5%), and winter (54.7%). Gosan is influenced by the pollution sources in East Asia during winter as well as other transition periods (spring and fall) due to the prevailing westerlies. In contrast, winds in summer blew mostly from the western North Pacific (WNP) by the easterly winds. The spring season is, in particular, important for the transport of mineral dust mixed with polluted OA to Gosan (Kundu et al., 2010).



**Figure 1.** Cluster analysis of backward air mass trajectories over Gosan (indicated by a star symbol) for the TSP collected during (a) spring, (b) summer, (c) fall, and (d) winter seasons.

## 2.2. Extraction and analysis of organic compounds

Approximately 3.14 cm<sup>2</sup> filter cuts were extracted with dichloromethane/methanol (2:1; v/v). The extracts were concentrated using a rotary evaporator under vacuum and then blown down to near dryness with pure nitrogen gas. The dried residues were subsequently reacted with N, O-bis(trimethylsilyl)trifluoroacetamide containing 1% trimethylchlorosilane (BSTFA+1%

TMCS, SUPELCO<sup>®</sup>, Sigmaaldrich<sup>®</sup>) and pyridine at 70 °C for 3 h to derive OH and COOH groups of polar organic compounds to trimethylsilyl ethers and esters, respectively. After the derivatization followed by the addition of a known amount of internal standard solution (Tridecane; 1.43 ng L<sup>-1</sup> in n-hexane), the derivatized extracts were injected onto a gas chromatograph (Hewlett-Packard model 6890 GC) coupled to a mass spectrometer (Hewlett-Packard model 5973, MSD) (GC-MS). More details on the quantification of polar organic compounds using GC-MS are described in Haque et al. (2019).

The target compounds (anhydrosugars, primary sugars, sugar alcohols, and BSOA tracers) were separated on a DB-5MS fused silica capillary column (30 m x 0.25 mm i.d., 0.5 µm film thickness) using helium as a carrier gas at a flow rate of 1.0 ml min<sup>-1</sup>. The GC oven temperature was programmed from 50°C for 2 min and then increased from 50 to 120°C at 30°C min<sup>-1</sup> and to 300°C at 6°C min<sup>-1</sup> with a final isotherm hold at 300°C for 16 min. The sample was injected in a splitless mode with the injector temperature at 280°C. The MS was operated at 70 eV and scanned from 50 to 650 Da on an electron impact (EI) mode. Mass spectral data were acquired and processed using the Chemstation software. The organic compounds were identified individually by comparison with retention times and mass spectra of authentic standards and NIST library and literature data of mass fragmentation patterns (Medeiros and Simoneit, 2007). For assessing the recoveries, ~100-200 ng of the standard solution was spiked on the blank filter and analyzed as a real sample. Overall, the average recoveries were found to be 80-104% for target compounds. The field and laboratory blank filters (n = 3) were also analyzed by the same procedures as a real sample. Target compounds were not found in the field blanks. The analytical errors based on concentrations by replicate sample analyses (n = 3) were less than 15%.

### 2.3. Radiocarbon isotopic composition of TC and EC

The concentrations of total carbon (TC) in TSP samples were analyzed using an elemental analyzer. For the radiocarbon isotopic composition ( $\Delta^{14}\text{C}$ ), the aerosol filter punches were exposed for ~12 h to HCl fumes in a vacuum desiccator. Subsequently, these filters were analyzed for  $\Delta^{14}\text{C}$  on a modified elemental analyzer coupled via a gas interface to Accelerator Mass Spectrometer Mini Carbon Dating System (MICADAS) at the University of Bern, Switzerland (Salazar et al., 2015). The evolved CO<sub>2</sub> of TC from the elemental analyzer was passed through a moisture trap (Sicapent, Merck) and isolated it from other residual gasses using a temperature-controlled zeolite trap. The purified CO<sub>2</sub> was introduced through a gas interface system to MICADAS, where <sup>14</sup>C/<sup>12</sup>C ratios are measured according to the analytical



procedures detailed in Zhang et al. (2016). Likewise, the evolved CO<sub>2</sub> of elemental carbon from the Sunset Lab OC/EC analyzer using Swiss 4S protocol (Zhang et al., 2012), was directed to the MICDAS and measured for the <sup>14</sup>C/<sup>12</sup>C ratio relative to standard calibration gas. These results were expressed as fractions of modern carbon ( $f_M$ ) by normalizing with a  $\delta^{13}C$  value of the reference standard in the year 1950 (−25‰) according to Stuiver and Polach (1997) for the fractionation effects. The  $f_M(OC)$  can be estimated by using the  $f_M(TC)$  and  $f_M(EC)$  in an isotope mass balance equation (Zhang et al., 2015). Additionally, we estimated the relative contributions of OC and EC from the nonfossil and fossil sources ( $f_{nonfossil}$  and  $f_{fossil}$ , respectively) using the following equations.

$$f_{nonfossil-OC} = f_M(OC-sample)/f_M(OC-ref); f_M(OC-ref) \approx 1.07 \pm 0.04 \quad (1)$$

$$f_{nonfossil-EC} = f_M(EC-sample)/f_M(EC-ref); f_M(EC-ref) \approx 1.10 \pm 0.05 \quad (2)$$

$$f_{fossil-OC} = 1 - f_{nonfossil-OC} \quad (3)$$

$$f_{fossil-EC} = 1 - f_{nonfossil-EC} \quad (4)$$

The reference values of OC and EC were obtained from Mohn et al. (2008). Using the fractions of  $f_{fossil-OC}$  and  $f_{nonfossil-OC}$ , we can, therefore, estimate the mass concentration of ambient organic carbon (OC-ambient) from fossil and nonfossil sources (OC<sub>fossil</sub> and OC<sub>nonfossil</sub>, respectively).

$$OC_{nonfossil} = f_{nonfossil-OC} \times [OC]_{ambient} \quad (5)$$

$$OC_{fossil} = f_{fossil-OC} \times [OC]_{ambient} \quad (6)$$

More details on the radiocarbon isotopic composition data over Gosan were reported elsewhere (Zhang et al., 2016).

### 3. Results and discussion

#### 3.1. Trajectory and cluster analysis


Backward air mass trajectories are useful for assessing the impact of local versus regional source emissions over Gosan. Seven-day isentropic backward air mass trajectories were computed using the hybrid single-particle Lagrangian integrated trajectory model (HYSPLIT, version 4: [Stein et al., 2015](#)) over the KCOG for the sampling period using the meteorological datasets of the Global Data Assimilation System (GDAS) network. The trajectory endpoint files from the HYSPLIT model were further used for the cluster analysis using the Trajstat package (Wang et al., 2009b) for all four seasons (Figure 1). Although cluster analysis revealed the predominance of continental transport in spring, fall, and winter seasons, the air masses over the KCOG in summer mostly originated from the WNP. Since

spring is a transition of winds switching from westerlies to easterlies, Gosan is likely influenced by the long-range transport of dust, pollution, and sea-salt aerosols.

The vertical mixing of pollutants within the boundary layer height also plays an important role in controlling the strength of continental outflow alongside regional meteorology. For instance, the mixing height of air parcels from the HYSPLIT model is mostly confined to 1000 m in winter but somewhat increased towards spring and fall seasons (Figure S1). This vertical enhancement in the boundary layer height facilitates the transport of mineral dust particles from the arid and semiarid regions in East Asia along with urban pollutants to Gosan in spring and fall compared to winter. However, the strength of continental outflow somewhat depends on several factors, including source emissions, meteorology, and mixing height of air parcels.

Gosan is influenced by three types of air masses in spring (Figure 1a), from the Mongolian Desert (cluster 1: 15%), North China (cluster 2: 45%), and from the Yellow Sea (cluster 3: 40%). In contrast, the easterlies from the WNP in summer mostly influenced the composition of TSP over the KCOG. This inference is based on the cluster analysis for summer samples (Figure 1b), which showed four regimes, including transport from the Sea of Japan (cluster 1: 13%), WNP (cluster 2: 22%), South China Sea (cluster 3: 46%), and East China Sea (cluster 4: 19%). In contrast, cluster analysis revealed three major transport regimes from East Asia in fall and winter (Figure 1c-d). However, there are subtle differences that exist between winter and fall in terms of influence from nearby versus distant pollution sources. For instance, long-range transport of air masses from west Mongolia (cluster 1: 21%) and the Russian Far East (cluster 3: 14%) exerted a weak influence on the TSP sampled over Gosan in winter. Besides, we observed a somewhat larger impact of air masses from the North China Plain over Gosan (cluster 2: 64%) in winter. In contrast, Gosan is less influenced by air masses originating from the North China Plain, contributing ~28% (cluster 1) than those from Mongolia (cluster 2: 39%) and Russian Far East (cluster 3: 33%) in fall. Therefore, the impact of East Asian outflow is stronger in winter than in spring and fall.





**Figure 2.** (a) Cumulative concentration levels of anhydrosugars, primary sugars, and sugar alcohols (i.e., represented by bars), and depicting the contributions of nonfossil (green color) and fossil (black color) organic carbon (i.e., pie charts), (b) Cumulative concentration levels of isoprene- and monoterpene SOA tracers in each TSP sample collected over Gosan.

### 3.2. Temporal and seasonal variability of sugars


The temporal/seasonal trends of sugar compounds over the KCOG provide useful information on the emission strengths of various sources in the East Asian outflow. All three anhydrosugars showed similar temporal and seasonal trends with higher concentrations in winter and fall than spring and summer (Figures 2a and S2). As levoglucosan and other two anhydrosugars (mannosan and galactosan) are the pyrolysis products of cellulose/hemicellulose, their higher concentrations along with an increase in nonfossil fraction of OC (Figure 2a; pie charts) in TSP from winter and fall revealed the impact of BB emissions. The MODIS satellite-based fire counts (Figure S1) together with cluster analysis in winter and fall (Figure 1) have revealed an influence of active BB emissions in the North China Plain, Mongolia, and Russian Far East. The temporal trends of glucose, fructose, and sucrose exhibited less variability throughout the sampling period; however, we observed a

slight increase in their concentration towards spring/summer (Figure 2a). Glucose and fructose have origins from leaf fragments and pollen species (Fu et al., 2012a). Sucrose is a potential tracer for airborne pollen (Fu et al., 2012a) and late spring/early summer is often regarded as a season of “pollen-allergies”. Therefore, similar temporal trends of glucose and fructose with sucrose indicate their common source, pollens (Figure 2a). Since glucose, fructose, and sucrose showed moderately significant correlations ( $R^2 = 0.44-0.48$ ,  $p < 0.01$ ) with levoglucosan in winter, it is somewhat possible that BB source emission could also influence the concentrations of these saccharides in this season (Haque et al., 2019; Fu et al., 2008).

BB also contributes to xylose and, hence, the temporal variability of xylose is mimicking that of anhydrosugars. Trehalose is a primary sugar and a useful tracer for organic carbon associated with soil dust particles (Fu et al., 2012a). The temporal variability of trehalose closely resembled that of fungal spore tracers (arabitol, mannitol, and erythritol), showing high concentrations in spring, summer, and fall seasons (Figures 2a and S2) (Zhu et al., 2015a; Fu et al., 2012a). The KCOG is under the influence of a large-scale advection of mineral dust from East Asia to the WNP during these three seasons (Tyagi et al., 2017; Huebert et al., 2003). The mineral dust transport from East Asia to the WNP can be traced by the high concentrations of non-sea-salt  $\text{Ca}^{2+}$  in the TSP samples from Gosan (Arimoto et al., 1996). Similar temporal trends of trehalose and nss- $\text{Ca}^{2+}$ , particularly in spring samples (Figure S3), suggest that the abundance of OA specific to fungal spores over Gosan is likely associated with the Kosa (Asian dust) events.

The major sources of arabitol and mannitol are airborne fungal spores (Bauer et al., 2008), accompanying detritus from mature leaves (Pashynska et al., 2002). Heald and Spracklen (2009) reported that mannitol and arabitol are considerably associated with terrestrial biosphere activity. Inositol is largely derived from the developing leaves in summer (Pashynska et al., 2002) and BB in winter (Fu et al., 2010b). Zhu et al. (2015b) found similar seasonal behavior of inositol with those of other sugar alcohols with the predominance in summer, associated with microbial activities in local forests from Okinawa. Inositol showed a moderately significant correlation with levoglucosan ( $R^2 = 0.33$ ,  $p < 0.01$ ) in winter; however, there were no positive linear relationships between levoglucosan and other sugar alcohols, implying a partial emission of inositol from the BB during winter in Gosan aerosols. Therefore, the temporal variability of inositol differs from other sugar alcohols (Figure S2). The sources of sugar compounds are further discussed in section 3.4.

The seasonally averaged concentrations of all the anhydrosugars and xylose are higher in winter/fall than spring/summer (Figure 3a-d), possibly due to a greater influence of long-range transport from East Asia. In contrast, glucose, fructose, and sucrose peaked in spring but decreased in other seasons (Figure 3e-g), mainly because of the contribution of airborne pollen. Trehalose showed higher concentrations in fall and summer, followed by spring and winter (Figure 3h). Arabitol, mannitol, and erythritol showed higher concentrations in summer/fall than in winter and spring (Figure 3i-k). This seasonal trend is consistent with those of soil-derived fungal spores. This feature is consistent with earlier observations from a remote oceanic island in the WNP (Okinawa) during the impact of East Asian outflow (Zhu et al., 2015a). The seasonally averaged mass concentrations of inositol are highest in spring, followed by summer, fall, and winter (Figure 3l). Overall, the molecular compositions of anhydrosugars showed the predominance of levoglucosan followed by galactosan and mannosan (Figure S4). Galactosan is more abundant in crop-residue burning emissions than mannosan (Engling et al., 2009; Sheesley et al., 2003). It is very much likely that the impact of crop-residue burning emissions in East Asia over Gosan is more prominent in winter/spring. Such high abundances of galactosan over mannosan were found in the North China Plain (Fu et al., 2008) and in the Indo-Gangetic Plain outflow sampled over the Bay of Bengal (Bikkina et al., 2019). Although the temporal variability of primary sugars in the TSP samples from Gosan showed a characteristic peak of glucose and sucrose (Figure 2a), the seasonally averaged distributions are different (Figure S4). The molecular distributions of sugar alcohols are characterized by high loadings of arabitol and mannitol, followed by erythritol and inositol (Figure S4).



**Figure 3.** Seasonal variability of atmospheric levels of sugar compounds and BSOA tracers in TSP samples from Gosan during April 2013-April 2014.

### 3.3. Temporal and seasonal variability of BSOA tracers


We identified six isoprene-SOA tracers such as 2-methylglyceric acid (2-MGA), three C<sub>5</sub>-alkene triols, and two 2-methyltetrols (2-methylthreitol and 2-methylerythritol) (2-MTs) in Gosan aerosol samples (Table 1). The sum of isoprene-SOA tracers ranged from 0.35 to 15.1 ng m<sup>-3</sup> (avg. 3.69 ng m<sup>-3</sup>) with the predominance of 2-MTs (avg. 2.09 ng m<sup>-3</sup>). 2-MGA is the second most abundant isoprene-SOA tracer (avg. 0.99 ng m<sup>-3</sup>), a high-generation product probably formed by further photooxidation of methacrolein and methacrylic acid. A similar molecular composition was observed over the North Pacific and California Coast (Fu et al., 2011). All the isoprene SOA tracers exhibited similar temporal variations with higher concentrations in the summer/spring months compared to autumn and winter (Figure 2b). Conversely, four monoterpene-SOA tracers, i.e., *cis*-pinonic acid, pinic acid, 3-hydroxyglutaric acid (3-HGA), and 3-methyl-1,2,3-butanetricarboxylic acid (MBTCA), were detected in Gosan samples (Table 1). Total concentrations of monoterpene-SOA tracers were found to be 1.65 to 35.5 ng m<sup>-3</sup> (avg. 9.24 ng m<sup>-3</sup>) with a high concentration of MBTCA (avg. 5.11 ng m<sup>-3</sup>). All the monoterpene-SOA tracers showed similar temporal trends with high values in summer/spring periods than autumn/winter (Figure 2b). Nevertheless, *cis*-

pinonic acid was ascribed somewhat different temporal variability with other monoterpene-SOA tracers. It is likely that *cis*-pinonic acid might be further photo-oxidized to form MBTCA (Szmigielski et al., 2007).

The seasonal average of isoprene-SOA tracers showed high concentrations in summer, followed by spring/fall and winter (Figure 3m-o). One key feature of the data presented here is that two fall samples (KOS984; 2-12 September and KOS986; 27 September to 07 October) exhibited high concentrations for 2-MTs over Gosan (Figure 2b). We presumed that local vegetation might contribute significantly to the formation of 2-MTs as it is a first-generation product. Moreover, 2-MTs can be derived from the open ocean under low NO<sub>x</sub> conditions (Hu et al., 2013). 3-HGA and pinic acid showed somewhat higher concentrations in summer/spring than fall/winter due to the growing vegetation (Figure 3q-r). *Cis*-pinonic acid was more abundant in spring compared to summer (Figure 3p) because of its photo-degradation, as discussed above. In contrast, MBTCA was dominant in summer/fall than spring/winter (Figure 3s). Here, the formation of MBTCA could be enhanced in fall during atmospheric transport from East Asia. The molecular distributions of isoprene-SOA tracers were characterized by a high loading of 2-MTs, followed by 2-MGA and C<sub>5</sub>-alkene triols in all seasons (Figure S4). The Molecular composition of monoterpene-SOA tracers was dominated by MBTCA, followed by 3-HGA, pinic acid, and *cis*-pinonic acid in all seasons (Figure S4). Overall, BSOA tracers were found to be most abundant in summer, followed by fall, spring, and winter (Table 1). Interestingly, it is likely that secondary OA undergoes much faster cycling than the primary sugar compounds, considering the feasibility of photooxidation. This would mean a slight underestimation of BSOA over the KCOG in the East Asian outflow and, hence, their atmospheric abundances over Gosan reflect a lower limit.

Kang et al. (2018b) reported that monoterpene-SOA tracers were more abundant than isoprene-SOA tracers in spring-summer over the East China Sea, which is consistent with this study. Although the mass budget calculations showed that isoprene and monoterpenes are largely emitted by terrestrial plants; however, the open ocean can also contribute to isoprene and monoterpenes significantly (Conte et al., 2020; Shaw et al., 2010; Broadgate et al., 1997). Air mass back trajectory and cluster analysis (Figure 1) implied that air masses mostly originated from the ocean during summer. This means the open ocean significantly contributed to isoprene-SOA production. However, terrestrial sources from the continent also substantially enhanced the formation of BSOA. For example, one sample (KOS: 979; 1-11

June 2013) during summer showed the highest loading of BSOA tracers (Figure 2b) when air masses are transported from the continent (Zhang et al., 2016).



**Figure 4.** Multiple linear regression (MLR) analysis of airborne sugar compounds in TSP collected over Gosan.

### 3.4. Source apportionment- regression analysis and diagnostic ratios

Anhydrosugars strongly correlated with xylose (Figure 4), suggesting their common source as BB emission in East Asia. Fu et al. (2012a) analyzed pollen from different tree species (*e.g.*, White birch, Chinese willow, Peking willow, Sugi, Hinoki), which are enriched in sucrose (182-37,300  $\mu\text{g g}^{-1}$ ), glucose (378-3601  $\mu\text{g g}^{-1}$ ) and fructose (162-1813  $\mu\text{g g}^{-1}$ ). In our samples, sucrose strongly correlated with glucose and fructose (Figure 4), suggesting their origin from plant-derived airborne pollen. Likewise, a strong correlation was found between arabitol and mannitol, indicating a mutual origin from a similar type of fungal spores (Zhu et al., 2015a; Fu et al., 2012a; Bauer et al., 2008; Yttri et al., 2007). Bauer et al. (2008) ascribed


weak correlations between arabitol and mannitol to the diverse nature of ambient fungal spores. Furthermore, both sugar alcohols correlated well with trehalose, a tracer for soil organic carbon (Fu et al., 2012a). This observation suggests their common origin from soil organic matter associated with fungal spores. Erythritol also originates from fungal spores; however, its abundance is affected by the atmospheric aging process. Kessler et al. (2010) reported that erythritol is degraded during long-range transport in 12.7 days. Consequently, arabitol and mannitol were moderately correlated with erythritol in the Gosan samples due to the degradation of latter sugar alcohol in the East Asian outflow.

The linear relationship of levoglucosan (*Lev*) with mannosan (*Man*), galactosan (*Gal*), and  $\text{nss-K}^+$  provide useful information on the type of BB emissions (hardwood, softwood, or crop-residue). Ratios of *Lev/Man* and *Lev/(Man + Gal)* can be useful to distinguish BB and coal combustion contributions. The average ratios of *Lev/Man* ( $15.1 \pm 6.76$ ) and *Lev/(Man + Gal)* ( $4.27 \pm 1.23$ ) in Gosan aerosols are much closer to those from wood burning and coal combustion sources emissions, respectively (Yan et al., 2018). It reveals that *Lev* could originate from both biomass and coal burning source emissions, which is consistent with the linear relationship between *Lev-C* and fossil-/nonfossil carbon fraction (section 3.7). Furthermore, different types of biomass are characterized by distinct *Lev/Man* ratios. For instance, *Lev/Man* ratios from the softwood burning emissions (3-10) differ from those of hardwood (15-25) and crop-residues (>40) (Singh et al., 2017; Schmidl et al., 2008a, b; Fu et al., 2008; Engling et al., 2006, 2009; Fine et al., 2001, 2004). We found the *Lev/Man* ratios (Table 2) over the KCOG overlap between seasons and somewhat close to that of hardwood burning emissions in northern China, Mongolia, and the Russian Far East, as corroborated by the backward air mass trajectories and MODIS fire counts (Figures 1 and S1). Besides, *Lev/K<sup>+</sup>* and *Man/Gal* ratios in summer differ from those of other seasons (Table 2). Cheng et al. (2013) have apportioned qualitatively the source contributions of anhydrosugars over a receptor site based on the comparison of *Lev/K<sup>+</sup>* and *Lev/Man* ratios in aerosols to those from various sources profiles compiled from the literature. This approach of using the mass ratios of *Lev/Man* and *Lev/K<sup>+</sup>* has been proven useful for deciphering the difference in BB-derived OA (Bikkina et al., 2019).

Here, we adopted this methodology to ascertain the likely contributing sources of anhydrosugars, which are BB tracers from different seasons (Figure 5). This source apportionment relies on the fact that *Lev/K<sup>+</sup>* from the softwood burning (10-1000) is higher than hardwood (1-100) (Fine et al., 2004). In contrast, *Lev/Man* ratios for softwood are lower than those of hardwood burning (10-100) (Fine et al., 2004). Likewise, the *Lev/Man* and




$Lev/K^+$  ratios from grasses and crop-residues are 10-100 and 0.01-1.0, respectively (Bikkina et al., 2019). On a similar note, the  $Lev/K^+$  ratios from the burning of pine needles (0.1-1.0) somewhat overlap with hardwood burning emissions but are characterized by distinct  $Lev/Man$  ratios (Bikkina et al., 2019). The burning of dead leaves (duff) showed higher  $Lev/K^+$  than pine needles and grasses, but their  $Lev/Man$  is on the lower side than the former biomass type and the softwood burning emissions. However,  $Lev$  is more susceptible to degradation by photooxidation with OH radicals during atmospheric transport (half-life: <2.2 days) (Hennigan et al., 2010) and, this would cause lower abundances of this anhydrosugar. Hence, the hardwood  $Lev/K^+$  ratios could slightly shift downwards. Therefore, caution is required while interpreting the ambient data from a receptor site (Bikkina et al., 2019). Overlapping the seasonal data on this scatter plot of  $Lev/K^+$  versus  $Lev/Man$  (Figure 5) clearly revealed a mixed contribution of burning of hardwood and crop-residue in the East Asian outflow. It should be noted that the photooxidation process during atmospheric transport is also applicable for low concentrations and poor correlations of other primary saccharides.



**Figure 5.** Scatter plot of levoglucosan/ $K^+$  versus levoglucosan/mannosan ratios in TSP collected over Gosan during April 2013-April 2014. The data for Siberian forest fires and burning leaves were adopted from Sullivan et al. (2008).



The correlation coefficients and diagnostic ratios of BSOA tracers specify their source origin or formation pathway. Nevertheless, atmospheric stability or reactivity of BSOA tracers through the photooxidation during long-range transport may bias the correlation coefficients in Gosan aerosols. 2-MGA showed significant correlation with 2-MTs ( $r = 0.79$ ,  $p < 0.01$ ) and C<sub>5</sub>-alkene triols ( $r = 0.50$ ,  $p < 0.05$ ) (Figure 6a-b), suggesting their similar formation pathway or common sources of isoprene SOA tracers. However, a poor correlation coefficient between 2-MTs and C<sub>5</sub>-alkene triols ( $r = 0.33$ ,  $p = 0.13$ ) (Figure 6c) indicates their different formation process over the Gosan atmosphere. Wang et al. (2005) documented that polyols are formed from isoprene through diepoxy derivatives, which further convert into 2-MTs by acid-catalyzed hydrolysis. On the other hand, C<sub>5</sub>-alkene triols are produced from the precursor of hydroxyp peroxy radicals that are initially derived from isoprene through rearrangement reactions (Surratt et al., 2006). It can be noted that the formation mechanisms of 2-MGA and 2-MTs are different while depending on the NO<sub>x</sub> concentrations (Surratt et al., 2010). Thus, the ratio of 2-MGA/2-MTs attributes to the influence of NO<sub>x</sub> on isoprene SOA formation. We found a low ratio of 2-MGA/2-MTs (0.67) (Figure S5a, Table 2) in summer, implying enhancement of 2-MTs formation over the open ocean due to the low-NO<sub>x</sub> environment in the ocean atmosphere. On the contrary, 2-MGA/2-MTs ratios for other seasons were  $>1.0$  (Figure S5a, Table 2), indicating an elevated formation of 2-MGA through continental high NO<sub>x</sub> condition, which is consistent with air masses back trajectory.



**Figure 6.** Pearson linear correlation coefficient analysis of BSOA tracers in Gosan TSP aerosols during April 2013-April 2014.


*Cis*-pinonic acid showed a weak correlation with pinic acid ( $r = 0.35$ ,  $p = 0.12$ ) (Figure 6f), suggesting that different atmospheric reactivity of *cis*-pinonic/pinic acids during transport would cause such poor correlation. In contrast, pinic acid exhibited a strong positive linear correlation with 3-HGA ( $r = 0.90$ ,  $p < 0.01$ ) and MBTCA ( $r = 0.90$ ,  $p < 0.01$ ) (Figure 6d-e), implying their similar sources. It should be noted that the formation processes of pinic acid, 3-HGA, and MBTCA are different because pinic acid is a first-generation product, and 3-HGA and MBTCA are high-generation products (Claeys et al., 2007; Müller et al., 2012; Szmigielski et al., 2007). The ratio of *cis*-pinonic acid + pinic acid to MBTCA (P/M) is used to evaluate the aging of monoterpene SOA.

A low P/M ratio suggests the transformation of *cis*-pinonic and pinic acids to MBTCA and thus relatively aged monoterpene-SOA, whereas a high ratio reflects relatively fresh monoterpene-SOA (Gómez-González et al., 2012; Ding et al., 2014). Gómez-González et al. (2012) reported aged monoterpene-SOA ( $P/M = 0.84$ ) from a Belgian forest site while fresh chamber-produced  $\alpha$ -pinene-SOA tracers showed P/M ratios of 1.51 to 3.21 (Offenberg

et al., 2007). The average ratio of P/M in this study showed 0.62 with the low value in summer (Figure S5b, Table 2), which is lower than those of Guangzhou (fresh monoterpene-SOA; 28.9) while the air masses originated from southern China (Ding et al., 2014). This observation indicates that monoterpenes SOA have undergone substantial aging during transport to Gosan, particularly in summer when extensive photochemical oxidation occurred due to the high temperature and intense solar radiation.

### 3.5. Source apportionment - Positive Matrix Factorization (PMF) analysis

We used the positive matrix factorization (PMF) analysis of airborne sugar compounds and BSOA tracers to assess their sources over Gosan. Five-factor profiles were chosen that could best explain the data (Figure 7). In factor 1, we found higher loadings of sucrose (88%), fructose (57%), and glucose (40%). Sucrose is a well-known tracer for airborne pollen as these types of biogenic sources also contain significant levels of glucose and fructose; we attribute this factor to contribution from "pollen". Likewise, factor 2 has higher loadings of C<sub>5</sub>-alkene triols (54%), 2-MGA (38%), *cis*-pinonic acid (34%), pinic acid (33%), and 3-HGA (33%). These SOA tracers are formed by the photochemical oxidation of BVOCs (e.g., isoprene and monoterpene), which are emitted from terrestrial vegetation and oceanic phytoplankton. Therefore, we attribute factor 2 to biogenic secondary oxidation products. Factor 3 was abundantly loaded by non-sea-salt fraction of calcium (nss-Ca<sup>2+</sup>), implying that the component is associated with soil dust as nss-Ca<sup>2+</sup> is a specific tracer of Earth surface soil (Athanasopoulou et al., 2010; Brahney et al., 2013). It should be noted that significant loadings were also observed for trehalose (39%), glucose (35%), and fructose (35%) in factor 3, indicating their association with soil microbes.




**Figure 7.** Positive Matrix Factorization (PMF)- based Source profiles resolved for the measured primary saccharides in TSP collected over Gosan during April 2013-April 2014.

We found high loadings of levoglucosan (83%), nss-K<sup>+</sup> (80%), and xylose (73%) in factor 4. Such high contributions of levoglucosan and K<sup>+</sup> (hemicellulose/cellulose pyrolysis tracers) and xylose (wood sugar) are typical of wood burning emissions. We, therefore, ascribe this factor to the influence of BB emissions in East Asia over Gosan. Factor 5 is dominated by 2-MTs (73%), MBTCA (53%), mannitol (55%), erythritol (54%), and arabitol (52%). 2-MTs and MBTCA are BSOA tracers, which are produced by the photooxidation of isoprene and monoterpene, respectively. In contrast, arabitol, mannitol, and erythritol are typical tracers of airborne fungal spores (Lewis and Smith, 1967; Bauer et al., 2008). Thus,

factor 5 is associated with mixed sources of ‘fungal spores’ and ‘biogenic secondary oxidation products’. PMF results were further utilized to calculate the relative contributions from various sources to OC over Gosan (Figure S6). BB was the dominant source of OC over Gosan (41.9%), followed by BSOA (21.1%), pollen (17.1%), and soil dust (11.4%).

### 3.6. Relative abundances in WSOC and OC


Levoglucosan is the most abundant anhydrosugar, contributing 0.05-1.54% of WSOC and 0.03-1.02% of OC. Likewise, sucrose, glucose, and fructose were more abundant primary sugars, contributing 0.01-2.83%, 0.03-1.22%, and 0.02-0.74% of WSOC, respectively. Contributions of the three primary sugars varied from 0.05% to 3.41% of OC. Arabitol and mannitol are the most abundant sugar alcohols, whose contribution to WSOC and OC ranged from 0.02% to 0.93% and 0.01% to 0.85%, respectively. Figure 8 depicts the contribution of sugar compounds to WSOC and OC in TSP collected over Gosan during the study period. We also compared the atmospheric abundances of sugar compounds from Gosan with the literature data (Table 3). This comparison revealed the less influence of BB tracer compounds (i.e., anhydrosugar levels) over Gosan, a factor of 5-10 times lower than those reported for the BB-influenced source regions in China and East Asia (Fu et al., 2012b; Kang et al., 2018a; Wang and Kawamura, 2005; Wang et al., 2012). However, the levels of anhydrosugar over Gosan are higher than those observed over the remote Canadian High Arctic site (Fu et al., 2009a). In contrast, Gosan is characterized by high concentrations of primary sugars compared to other remote sampling sites in Table 3. This is because of the overwhelming contribution of primary sugars associated with soil dust particles over Gosan during the East Asian outflow. Such high loadings of primary sugars were observed from other remote island receptor sites in the WNP (Okinawa) during the spring season (Zhu et al., 2015a). Likewise, the concentrations of sugar alcohols from Gosan are similar to those from other receptor sites influenced by the East Asian outflow (Verma et al., 2018; Zhu et al., 2015a).



**Figure 8.** Contribution of primary saccharide-C in (a) WSOC(%) and (b) OC(%) in TSP collected over Gosan during April 2013-April 2014.

Contributions of isoprene-SOA tracers to ambient OC (0.01-0.40%, avg. 0.09%) and WSOC (0.02-0.57%, 0.13%) were lower than monoterpene-derived SOA (0.04-0.63%, 0.23% for OC and 0.06-0.82%, 0.33% for WSOC). The contributions of isoprene oxidation products to OC and WSOC were found to be highest in summer (0.23% and 0.32%, respectively), followed by fall (0.09% and 0.13%), spring (0.04% and 0.06%), and winter (0.02% and 0.03%). Likewise, the contributions of monoterpene-SOA products to aerosol OC and WSOC exhibited highest value in summer (0.40% and 0.55%, respectively), followed by fall (0.24% and 0.35%), spring (0.18% and 0.27%), and winter (0.10% and 0.14%). We found that the contribution of BSOA products to carbonaceous components was twice in summer (Figure 9, Table 2). This means SOA formation was occurred in summer to a greater extent due to the intensive BVOCs emission with key factors of meteorological parameters

(higher temperature and radiation), i.e., higher concentrations of ozone, and other oxidizing agents (NO<sub>x</sub>, OH, etc.).



**Figure 9.** Contribution of (a) isoprene- and monoterpene SOA tracers-C in ambient OC(%) and (b) isoprene- and monoterpene SOA tracers-SOC in ambient OC(%) in Gosan aerosol samples during April 2013-April 2014.

We estimated secondary organic carbon (SOC) derived from isoprene and monoterpenes, using the measured values of BSOA tracers and following the SOA tracer-based method first proposed by Kleindienst et al. (2007). A summary of the estimated SOC is provided in Table 2. The contribution of isoprene to SOC was calculated 2.26 to 97.4 ngC m<sup>-3</sup> (avg. 23.7 ngC m<sup>-3</sup>), accounting for 1.45% of OC and 35.5% of total BSOC with the predominance in summer (3.56% and 46.6% for OC and SOC, respectively). The estimation of monoterpene SOA to SOC (avg. 40.1 ngC m<sup>-3</sup>) was observed around two times higher than isoprene (Table 2). Interestingly, the contribution of monoterpene-derived SOC to ambient


OC (3.65%) was dominant in summer, but monoterpene SOC to total SOC (72.1%) was most abundant in spring (Figure 9b). The seasonal distributions of biogenic SOC (Table 2) implied that a substantial amount of SOC was formed from monoterpenes in spring. The estimated biogenic SOC at the KCOG is almost one order of magnitude lower than those from other continental sites in China urban areas (e.g., Pearl River Delta: 446 ngC m<sup>-3</sup>) (Ding et al., 2012). However, the estimated biogenic SOC load from the KCOG is much higher than that reported from a remote site in the Canadian High Arctic (Alert: 9.4 ngC m<sup>-3</sup>) (Fu et al., 2009b) and comparable with over the East China Sea (Kang et al., 2018b).

It should be noted that concentrations of SOA tracers cannot always provide the actual contribution of the biogenic source to ambient organic aerosol mass. For example, loadings of monoterpene-SOA tracers were lower in sample KOS999 (28 March - 9 April 2014) (3.02 ng m<sup>-3</sup>) compared to sample KOS1000 (9 - 18 April 2014) (14.4 ng m<sup>-3</sup>), whereas the estimated contribution of SOC to ambient OC showed an opposite trend (KOS999: 2.08%; KOS1000: 1.56%). This result demonstrates that the estimation of SOC is an important factor in evaluating the contribution of BSOA to organic aerosol mass. We calculated biogenic OC using radiocarbon (<sup>14</sup>C) data following the method proposed by Szidat et al. (2006). Biogenic OC showed poor correlation with biogenic SOC ( $r = 0.36$ ,  $p = 0.09$ ) but significant linear relationship with primary sugars (i.e., glucose, fructose and sucrose) ( $r = 0.54$ ,  $p < 0.5$ ), suggesting that primary bioaerosols from plant-derived airborne pollen dictate on biogenic OC over Gosan.

### 3.7. Significance of fossil fuel as a source for levoglucosan

The ambient levoglucosan (*Lev*) levels showed a significant linear relationship with the OC<sub>fossil</sub>, suggesting the fossil source contribution of this molecular marker (Figure 10a). However, such a significant correlation was not evident between OC<sub>fossil</sub> and other major sugar compounds. Until recently, *Lev* has been thought to originate primarily from the hemicellulose/cellulose pyrolysis of vegetation and, hence, can be employed as a powerful tracer for biomass smoke particles (Fraser and Lakshmanan, 2000; Simoneit et al., 1999). Nevertheless, residential coals (e.g., lignite and bituminous coal) have been shown to contain high concentrations of '*Lev*' but also emit traces of *Man* and *Gal* (Kourtchev et al., 2011; Fabbri et al., 2008). Recently, Yan et al. (2018) found a significant linear relationship between the <sup>14</sup>C-based fossil fraction of WSOC and *Lev-C* in the aerosols generated from coal combustion and the ambient aerosol samples. Therefore, the prevailing linear relationship between OC<sub>fossil</sub> and *Lev-C* in Gosan samples (Figure 10a) is likely due to a common source contribution from coal combustion in East Asia.





641

642 **Figure 10.** Linear regression analysis between levoglucosan in terms of their carbon content  
 643 (*Lev-C*) and <sup>14</sup>C-based mass concentrations of (a) organic carbon and (b) elemental carbon of  
 644 fossil origin (OC<sub>fossil</sub> and EC<sub>fossil</sub>, respectively), (c) nss-SO<sub>4</sub><sup>2-</sup>, (d) nonfossil derived organic  
 645 carbon (OC<sub>nonfossil</sub>), (e) nonfossil derived elemental carbon (EC<sub>nonfossil</sub>), and (f) biomass  
 646 burning derived primary OC (POC<sub>BB</sub>) in TSP collected over Gosan during April 2013-April  
 647 2014. In panel (b), the squares represent three outliers (i.e., samples with rather high and low  
 648 *Lev/EC* ratios; please see text for more details).

The slope of the linear regression between *Lev*-C and OC<sub>fossil</sub> (0.0094; Figure 10a) is higher than those documented for the coal combustion source in China ( $\sim 0.004 \pm 0.007$ ) (Yan et al., 2018). Moreover, *Lev*-C moderately correlated with the EC<sub>fossil</sub> with the regression slope ( $\sim 0.01$ ) in Gosan samples (Figure 10b), being comparable to that observed for the coal combustion in China ( $0.044 \pm 0.076$ ) (Yan et al., 2018). It should be noted that excluding the three outliers as shown in Figure 10b (black square), *Lev*-C showed a stronger correlation with EC<sub>fossil</sub> ( $R^2 = 0.74$ ,  $p < 0.05$ ). Of these, two outliers in winter (KOS995: 22-29 January 2014; KOS996: 21 February - 3 March 2014) have higher *Lev*-C levels over that of EC<sub>fossil</sub>, when air mass trajectories showed the impact of BB emissions in the North China Plain. In contrast, the third outlier in summer (KOS979: 1-11 Jun'2013) has a lower *Lev*-C/EC<sub>fossil</sub>, while air parcels transported from nearby cities in China, Korea, and Japan, thus, have more contribution from vehicular emissions. Overall, both the regression slopes are, thus, the representative nature of *Lev*-C/OC<sub>fossil</sub> and *Lev*-C/EC<sub>fossil</sub> in the East Asian outflow. *Lev*-C and nss-SO<sub>4</sub><sup>2-</sup> exhibited a poor correlation (Figure 10c), although both were transported from East Asia.

*Lev*-C exhibited a rather weak correlation with OC<sub>nonfossil</sub> ( $R^2 = 0.19$ ) than that with EC<sub>nonfossil</sub> ( $R^2 = 0.42$ ) over Gosan during the study period (Figure 10d-e). This could be likely because OC<sub>nonfossil</sub> has contributions from the BB and the secondary formation process or the primary biogenic sources. The contribution of primary OC generated from BB (POC<sub>BB</sub>) to OC<sub>nonfossil</sub> was taken from Zhang et al. (2016). In their study, the <sup>14</sup>C-based EC<sub>nonfossil</sub> levels were scaled by a factor to constrain the POC<sub>BB</sub> (Zhang et al., 2016). Here the conversion factor is '4.5' (range: 3-10), which is a median value representing the primary OC/EC ratio from BB emissions ((POC/EC)<sub>BB</sub>).

$$\text{POC}_{\text{BB}} = \text{EC}_{\text{nonfossil}} \times (\text{POC}/\text{EC})_{\text{BB}} \quad (7)$$

*Lev*-C showed a somewhat improved linear correlation with POC<sub>BB</sub> than with the OC<sub>nonfossil</sub> (Figure 10f). It is apparent from Figure 10 that the regression slopes are comparable, indicating their contribution to *Lev* from both coal combustion and BB emissions over Gosan.

The prevailing weak linear relationship (moderate correlation) of *Lev* with nonfossil and fossil carbon fractions is likely being the result of photo-degradation of *Lev* during atmospheric transport. This result would mean that the higher atmospheric abundance of *Lev* and its pronounced linear relationships with the nonfossil and fossil carbon fractions implies a much stronger impact of both source emissions in East Asia during the continental outflow in winter and spring.

Overall, we present a new finding on the contribution of coal combustion sources in East Asia in controlling the atmospheric levels of *Lev* apart from the traditional biomass/biofuel burning emissions. This is based on the prevailing linear relationship between the radiocarbon based nonfossil-EC and *Lev* in the year-round TSP samples collected from the KCOG site in Jeju Island. The Gosan supersite is the best location to understand how the chemical composition of source-emissions from East Asia affects the outflow regions in winter and spring. Recent studies have highlighted the potential contribution of *Lev* from residential coal combustion in China (Yan et al., 2018), with estimated annual emission of ~2.2 Gg of *Lev* from domestic coal combustion (Wu et al., 2021). Given this background information, the prevailing significant linear relationship between *Lev* and nonfossil-EC ( $P$ -value  $< 0.05$ ) over the KCOG clearly emphasizes the need for reconsideration of the previous assessments on the impact of BB in East Asian outflow to the WNP. Additionally, this dataset is further compared with the molecular distributions and relative abundances of organic tracers in the TSP samples collected over Gosan during 2001, a decade ago (Fu et al., 2012). This comparison allows us to better understand the regional changes in the emission sources (*e.g.*, fugitive dust, BB, and fossil-fuel combustion) on a decadal basis.

#### 4. Conclusions

We investigated seasonal variations of primary organic components such as anhydrosugars, primary sugars, and sugar alcohols and biogenic secondary organic aerosol (BSOA) tracers (isoprene- and monoterpene-derived SOA products) in ambient aerosols from Gosan, Jeju Island. Among the detected sugar compounds, levoglucosan was dominant in winter/fall, whereas glucose and sucrose were more abundant in spring/summer. The seasonal trends documented that BB impact is more significant in winter/fall and the primary bioaerosol particles are important in spring/summer. Diagnostic ratios of levoglucosan, galactosan, and mannosan reflect that emissions from BB are mostly dominated by hardwood. The significant linear relationship of sucrose with glucose and fructose suggests their origin from airborne pollen. On a similar note, trehalose showed a significant positive correlation with arabinol, mannitol, and erythritol, implying their contribution from airborne fungal spores and soil microbes over the KCOG.

Distributions of biogenic SOA tracers were characterized by a predominance of monoterpene- than isoprene-derived oxidation products in Gosan aerosols. The BSOA tracers were formed in summer to a greater extent, followed by fall/spring and then winter. The low ratio of *cis*-pinonic acid + pinic acid to MBTCA (P/M) demonstrated that monoterpene SOA

was relatively aged over Gosan aerosols. The estimated secondary organic carbon (SOC) with the predominance in summer **shows** that substantial SOA formation occurred in summer due to favorable meteorological conditions. The backward air mass trajectories and source apportionment studies entirely demonstrated that BB and biogenic SOA contribution **significantly dominate the ambient OA mass over KCOG. Interestingly, levoglucosan-C exhibited a significant positive correlation with nonfossil and fossil organic carbon fractions, along with the comparable regression slopes.** This result reveals that BB and coal (lignite) combustion both are **the prominent source** for levoglucosan in the East Asian outflow.

Although there is some evidence that levoglucosan could originate from the combustion of brown coals (*e.g.*, lignite) in China; however, our observations are from the KCOG (receptor site) also hinted **at** the fossil source contribution of this molecular marker in the regional influx of the East Asian outflow. Therefore, attribution of ambient levoglucosan levels over the WNP to the impact of BB emission may cause large uncertainty.

#### **Data availability**

The data used in this paper are available upon request from the corresponding author.

#### **Author contributions**

KK and YLZ designed the research. ML collected the aerosol samples. MMH and SB performed the analysis of aerosol samples. MMH wrote the paper under the guidance of YLZ and KK. All authors were actively involved in the discussion of the paper.

#### **Competing interests**

The authors declare that they have no conflict of interest.

#### **Acknowledgments**

We acknowledge the financial supports of the Japan Society for the Promotion of Science (JSPS) through Grant-in-Aid No. 24221001 and the National Natural Science Foundation of China (Grant No 41977305).

#### **References**

- Arimoto, R., Duce, R. A., Savoie, D. L., Prospero, J. M., Talbot, R., Cullen, J. D., Tomza, U., Lewis, N. F., and Ray, B. J.: Relationships among aerosol constituents from Asia and the North Pacific during PEM-West A, *J. Geophys. Res. Atmos.*, 101, 2011-2023, <https://doi.org/10.1029/95JD01071>, 1996.
- Athanasopoulou, E., Tombrou, M., Russell, A. G., Karanasiou, A., Eleftheriadis, K., and Dandou, A.: Implementation of road and soil dust emission parameterizations in the aerosol model CAMx: Applications over the greater Athens urban area affected by natural sources, *J. Geophys. Res.*, 115, D17301, <https://doi.org/10.1029/2009JD013207>, 2010.

- Bauer, H., Claeys, M., Vermeylen, R., Schueller, E., Weinke, G., Berger, A., and Puxbaum, H.: Arabitol and mannitol as tracers for the quantification of airborne fungal spores, *Atmos. Environ.*, 42, 588–593, 2008.
- Bikkina, S., Haque, M. M., Sarin, M., and Kawamura, K.: Tracing the relative significance of primary versus secondary organic aerosols from biomass burning plumes over coastal ocean using sugar compounds and stable carbon isotopes, *ACS Earth Space Chem.*, 3, 1471–1484, <https://doi.org/10.1021/acsearthspacechem.9b00140>, 2019.
- Bikkina, S., Kawamura, K., Sakamoto, Y., and Hirokawa, J.: Low molecular weight dicarboxylic acids, oxocarboxylic acids and  $\alpha$ -dicarbonyls as ozonolysis products of isoprene: Implication for the gaseous-phase formation of secondary organic aerosols, *Sci. Total Environ.*, 769, 144472, <https://doi.org/10.1016/j.scitotenv.2020.144472>, 2021.
- Brahney, J., Ballantyne, A. P., Sievers, C., and Neff, J. C.: Increasing  $\text{Ca}^{2+}$  deposition in the western US: The role of mineral aerosols, *Aeolian Res.*, 10, 77–87, 2013.
- Broadgate, W. J., Liss, P. S., and Penkett, S. A.: Seasonal emissions of isoprene and other reactive hydrocarbon gases from the ocean, *Geophys. Res. Lett.*, 24, 2675–2678, <https://doi.org/10.1029/97GL02736>, 1997.
- Cheng, Y., Engling, G., He, K.-B., Duan, F.-K., Ma, Y.-L., Du, Z.-Y., Liu, J.-M., Zheng, M., and Weber, R. J.: Biomass burning contribution to Beijing aerosol, *Atmos. Chem. Phys.*, 13, 7765–7781, 2013.
- Claeys, M., Graham, B., Vas, G., Wang, W., Vermeylen, R., Pashynska, V., Cafmeyer, J., Guyon, P., Andreae, M. O., Artaxo, P., and Maenhaut, W.: Formation of secondary organic aerosols through photooxidation of isoprene, *Science*, 303(5661), 1173–1176, 2004.
- Claeys, M., Szmigielski, R., Kourtchev, I., van der Veken, P., Vermeylen, R., Maenhaut, W., Jaoui, M., Kleindienst, T. E., Lewandowski, M., Offenberg, J., and Edney, E. O.: Hydroxydicarboxylic acids: Markers for secondary organic aerosol from the photooxidation of  $\alpha$ -pinene, *Environ. Sci. Technol.*, 41(5), 1628–1634, 2007.
- Conte, L., Szopa, S., Aumont, O., Gros, V., and Bopp, L.: Sources and sinks of isoprene in the global open ocean: Simulated patterns and emissions to the atmosphere, *J. Geophys. Res. Ocean.*, 9, e2019JC015946, <https://doi.org/10.1029/2019JC015946>, 2020.
- Deguillaume, L., Leriche, M., Amato, P., Ariya, P. A., Delort, A.-M., Pöschl, U., Chaumerliac, N., Bauer, H., Flossmann, A. I., and Morris, C. E.: Microbiology and atmospheric processes: chemical interactions of primary biological aerosols, *Biogeosciences*, 5, 1073–1084, 2008.
- Ding, X., Wang, X. M., Gao, B., Fu, X. X., He, Q. F., Zhao, X. Y., Yu, J. Z., and Zheng, M.: Tracer-based estimation of secondary organic carbon in the Pearl River Delta, south China, *J. Geophys. Res. Atmos.*, 117, D05313, <https://doi.org/10.1029/2011JD016596>, 2012.
- Ding, X., He, Q. F., Shen, R. Q., Yu, Q. Q., and Wang, X. M.: Spatial distributions of secondary organic aerosols from isoprene, monoterpenes,  $\beta$ -caryophyllene, and aromatics over China during summer, *J. Geophys. Res.*, 119, 11877–11891, <https://doi.org/10.1002/2014JD021748>, 2014.
- Engling, G., Carrico, C. M., Kreidenweis, S. M., Collett, J. L., Day, D. E., Malm, W. C., Lincoln, E., Hao, W. M., Iinuma, Y., and Herrmann, H.: Determination of levoglucosan

- in biomass combustion aerosol by high-performance anion-exchange chromatography with pulsed amperometric detection, *Atmos. Environ.*, 40, S299–S311, 2006.
- Engling, G., Lee, J. J., Tsai, Y. W., Lung, S. C. C., Chou, C. C. K., and Chan, C. Y.: Size-resolved anhydrosugar composition in smoke aerosol from controlled field burning of rice straw, *Aerosol Sci. Technol.*, 43(7), 662–672, <https://doi.org/10.1080/02786820902825113>, 2009.
- Fabbri, D., Marynowski, L., Fabianska, M. J., Zaton, M., and Simoneit, B. R. T.: Levoglucosan and other cellulose markers in pyrolysates of miocene lignites: Geochemical and environmental implications, *Environ. Sci. Technol.*, 42(8), 2957–2963, <https://doi.org/10.1021/Es7021472>, 2008.
- Fine, P. M., Cass, G. R., and Simoneit, B. R. T.: Chemical characterization of fine particle emissions from fireplace combustion of woods grown in the northeastern United States, *Environ. Sci. Technol.*, 35(13), 2665–2675, 2001.
- Fine, P. M., Cass, G. R., and Simoneit, B. R. T.: Chemical characterization of fine particle emissions from the fireplace combustion of wood types grown in the Midwestern and Western United States, *Environ. Eng. Sci.*, 21(3), 387–409, 2004.
- Fraser, M. P. and Lakshmanan, K.: Using levoglucosan as a molecular marker for the long-range transport of biomass combustion aerosols, *Environ. Sci. Technol.*, 34(21), 4560–4564, <https://doi.org/10.1021/es991229l>, 2000.
- Fu, P., Kawamura, K., Kanaya, Y., and Wang, Z.: Contributions of biogenic volatile organic compounds to the formation of secondary organic aerosols over Mt. Tai, Central East China, *Atmos. Environ.*, 44, 4817–4826, <https://doi.org/10.1016/j.atmosenv.2010.08.040>, 2010a.
- Fu, P., Kawamura, K., Kobayashi, M., and Simoneit, B. R. T.: Seasonal variations of sugars in atmospheric particulate matter from Gosan, Jeju Island: Significant contributions of airborne pollen and Asian dust in spring, *Atmos. Environ.*, 55, 234–239, <https://doi.org/10.1016/j.atmosenv.2012.02.061>, 2012a.
- Fu, P. Q., Kawamura, K., Okuzawa, K., Aggarwal, S. G., Wang, G., Kanaya, Y., and Wang, Z.: Organic molecular compositions and temporal variations of summertime mountain aerosols over Mt. Tai, North China Plain, *J. Geophys. Res. Atmos.*, 113, D1910, <https://doi.org/10.1029/2008JD009900>, 2008.
- Fu, P. Q., Kawamura, K., and Barrie, L. A.: Photochemical and other sources of organic compounds in the Canadian high Arctic aerosol pollution during winter-spring, *Environ. Sci. Technol.*, 43(2), 286–292, 2009a.
- Fu, P. Q., Kawamura, K., Chen, J., and Barrie, L. A.: Isoprene, monoterpene, and sesquiterpene oxidation products in the high Arctic aerosols during late winter to early summer, *Environ. Sci. Technol.*, 43(11), 4022–4028, 2009b.
- Fu, P. Q., Kawamura, K., Pavuluri, C. M., Swaminathan, T., and Chen, J.: Molecular characterization of urban organic aerosol in tropical India: contributions of primary emissions and secondary photooxidation, *Atmos. Chem. Phys.*, 10(6), 2663–2689, 2010b.
- Fu, P. Q., Kawamura, K., and Miura, K.: Molecular characterization of marine organic aerosols collected during a round-the-world cruise, *J. Geophys. Res.*, 116, D13302, <https://doi.org/10.1029/2011jd015604>, 2011.

- Fu, P. Q., Kawamura, K., Chen, J., Li, J., Sun, Y. L., Liu, Y., Tachibana, E., Aggarwal, S. G., Okuzawa, K., Tanimoto, H., Kanaya, Y., and Wang, Z. F.: Diurnal variations of organic molecular tracers and stable carbon isotopic composition in atmospheric aerosols over Mt. Tai in the North China Plain: an influence of biomass burning, *Atmos. Chem. Phys.*, 12(18), 8359–8375, <https://doi.org/10.5194/Acp-12-8359-2012>, 2012b.
- Gómez-González, Y., Wang, W., Vermeylen, R., Chi, X., Neirynck, J., Janssens, I. A., Maenhaut, W., and Claeys, M.: Chemical characterisation of atmospheric aerosols during a 2007 summer field campaign at Brasschaat, Belgium: Sources and source processes of biogenic secondary organic aerosol, *Atmos. Chem. Phys.*, 12, 125–138, <https://doi.org/10.5194/acp-12-125-2012>, 2012.
- Graham, B., Guyon, P., Taylor, P. E., Artaxo, P., Maenhaut, W., Glovsky, M. M., Flagan, R. C., and Andreae, M. O.: Organic compounds present in the natural Amazonian aerosol: Characterization by gas chromatography-mass spectrometry, *J. Geophys. Res. Atmos.*, 108, D24, 4766, <https://doi.org/10.1029/2003JD003990>, 2003.
- Griffin, R. J., Cocker, D. R., Seinfeld, J. H., and Dabdub, D.: Estimate of global atmospheric organic aerosol from oxidation of biogenic hydrocarbons, *Geophys. Res. Lett.*, 26, 2721–2724, 1999.
- Guenther, A., Hewitt, C. N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdau, M., McKay, W. A., Pierce, T., Scholes, B. R. S., Tallamraju, R., Taylor, J., and Zimmerman, P.: A global model of natural volatile organic compound emissions, *J. Geophys. Res.*, 100(D5), 8873–8892, 1995.
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210, 2006.
- Haque, M. M., Kawamura, K., Deshmukh, D. K., Fang, C., Song, W., Mengying, B., and Zhang, Y. L.: Characterization of organic aerosols from a Chinese megacity during winter: Predominance of fossil fuel combustion, *Atmos. Chem. Phys.*, 19, 5147–5164, <https://doi.org/10.5194/acp-19-5147-2019>, 2019.
- Heald, C. L., and Spracklen, D. V.: Atmospheric budget of primary biological aerosol particles from fungal spores, *Geophys. Res. Lett.*, 36, L09806, <https://doi.org/10.1029/2009GL037493>, 2009.
- Heald, C. L., Henze, D. K., Horowitz, L. W., Feddema, J., Lamarque, J. F., Guenther, A., Hess, P. G., Vitt, F., Seinfeld, J. H., Goldstein, A. H., and Fung, I.: Predicted change in global secondary organic aerosol concentrations in response to future climate, emissions, and land use change, *J. Geophys. Res.*, 113(D5), <https://doi.org/10.1029/2007jd009092>, 2008.
- Hennigan, C. J., Sullivan, A. P., Collett, J. L., and Robinson, A. L.: Levoglucosan stability in biomass burning particles exposed to hydroxyl radicals, *Geophys. Res. Lett.*, 37, <https://doi.org/10.1029/2010GL043088>, 2010.
- Hu, Q. H., Xie, Z. Q., Wang, X. M., Kang, H., He, Q. F., and Zhang, P.: Secondary organic aerosols over oceans via oxidation of isoprene and monoterpenes from Arctic to Antarctic, *Sci. Rep.*, 3, 3119, <https://doi.org/10.1038/srep02280>, 2013.
- Huebert, B. J., Bates, T., Russell, P. B., Shi, G. Y., Kim, Y. J., Kawamura, K., Carmichael, G., and Nakajima, T.: An overview of ACE-Asia: Strategies for quantifying the



- relationships between Asian aerosols and their climatic impacts, *J. Geophys. Res. Atmos.*, 108(D23), 8633, <https://doi.org/10.1029/2003JD003550>, 2003.
- Jia, Y. L. and Fraser, M.: Characterization of saccharides in size-fractionated ambient particulate matter and aerosol sources: The contribution of primary biological aerosol particles (PBAPs) and soil to ambient particulate matter, *Environ. Sci. Technol.*, 45(3), 930–936, <https://doi.org/10.1021/es103104e>, 2011.
- Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y., Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati, E., Stephanou, E. G., and Wilson, J.: Organic aerosol and global climate modelling: a review, *Atmos. Chem. Phys.*, 5, 1053–1123, 2005.
- Kang, M., Ren, L., Ren, H., Zhao, Y., Kawamura, K., Zhang, H., Wei, L., Sun, Y., Wang, Z., and Fu, P.: Primary biogenic and anthropogenic sources of organic aerosols in Beijing, China: Insights from saccharides and n-alkanes, *Environ. Pollut.*, 243, 1579–1587, <https://doi.org/10.1016/j.envpol.2018.09.118>, 2018a.
- Kang, M., Fu, P., Kawamura, K., Yang, F., Zhang, H., Zang, Z., Ren, H., Ren, L., Zhao, Y., Sun, Y., and Wang, Z.: Characterization of biogenic primary and secondary organic aerosols in the marine atmosphere over the East China Sea, *Atmos. Chem. Phys.*, 19, 13947–13967, <https://doi.org/10.5194/acp-18-13947-2018>, 2018b.
- Kawamura, K., Kobayashi, M., Tsubonuma, N., Mochida, M., Watanabe, T., and Lee, M.: Organic and inorganic compositions of marine aerosols from East Asia: Seasonal variations of water-soluble dicarboxylic acids, major ions, total carbon and nitrogen, and stable C and N isotopic composition, in *geochemical investigations in earth and space science: A Tribute to Isaac R. Kaplan*, edited by R. J. Hill, J. Leventhal, Z. Aizenshtat, M. J. Baedeker, G. Claypool, R. Eganhouse, M. Goldhaber, and K. Peters, pp. 243–265, The Geochemical Society, 2004.
- Kessler, S. H., Smith, J. D., Che, D. L., Worsnop, D. R., Wilson, K. R., and Kroll, J. H.: Chemical sinks of organic aerosol: Kinetics and products of the heterogeneous oxidation of erythritol and levoglucosan, *Environ. Sci. Technol.*, 44(18), 7005–7010, <https://doi.org/10.1021/Es101465m>, 2010.
- Kourtchev, I., Hellebust, S., Bell, J. M., O'Connor, I. P., Healy, R. M., Allanic, A., Healy, D., Wenger, J. C., and Sodeau, J. R.: The use of polar organic compounds to estimate the contribution of domestic solid fuel combustion and biogenic sources to ambient levels of organic carbon and PM<sub>2.5</sub> in Cork Harbour, Ireland, *Sci. Total Environ.*, 11, 2143–2155, <https://doi.org/10.1016/j.scitotenv.2011.02.027>, 2011.
- Kroll, J. H., Ng, N. L., Murphy, S. M., Flagan, R. C., and Seinfeld, J. H.: Secondary organic aerosol formation from isoprene photooxidation under high-NO<sub>x</sub> conditions, *Geophys. Res. Lett.*, 32(18), L18808, <https://doi.org/10.1029/2005gl023637>, 2005.
- Kroll, J. H., Ng, N. L., Murphy, S. M., Flagan, R. C., and Seinfeld, J. H.: Secondary organic aerosol formation from isoprene photooxidation, *Environ. Sci. Technol.*, 40, 1869–1877, 2006.
- Kundu, S., Kawamura, K., and Lee, M.: Seasonal variations of diacids, ketoacids, and  $\alpha$ -dicarbonyls in aerosols at Gosan, Jeju Island, South Korea: Implications for sources,



- formation, and degradation during long-range transport, *J. Geophys. Res.*, 115, D19307, <https://doi.org/10.1029/2010jd013973>, 2010.
- Lewis, D. H. and Smith, D. C.: Sugar alcohols (polyols) in fungi and green plants: 1. Distributions, physiology and metabolism, *New Phytol.*, 66, 143–184, 1967.
- Liu, J., Chu, B., Chen, T., Liu, C., Wang, L., Bao, X., and He, H.: Secondary organic aerosol formation from ambient air at an urban site in Beijing: Effects of OH exposure and precursor concentrations, *Environ. Sci. Technol.*, 52, 6834–6841, <https://doi.org/10.1021/acs.est.7b05701>, 2018.
- Medeiros, P. M. and Simoneit, B. R. T.: Analysis of sugars in environmental samples by gas chromatography-mass spectrometry, *J. Chromatogr. A*, 1141(2), 271–278, <https://doi.org/10.1016/j.chroma.2006.12.017>, 2007.
- Medeiros, P. M., Conte, M. H., Weber, J. C., and Simoneit, B. R. T.: Sugars as source indicators of biogenic organic carbon in aerosols collected above the Howland Experimental Forest, Maine, *Atmos. Environ.*, 40(9), 1694–1705, 2006.
- Mohn, J., Szidat, S., Fellner, J., Rechberger, H., Quartier, R., Buchmann, B., and Emmenegger, L.: Determination of biogenic and fossil CO<sub>2</sub> emitted by waste incineration based on <sup>14</sup>CO<sub>2</sub> and mass balances, *Bioresour. Technol.*, 99, 6471–6479, <https://doi.org/10.1016/j.biortech.2007.11.042>, 2008.
- Müller, L., Reinnig, M. C., Naumann, K. H., Saathoff, H., Mentel, T. F., Donahue, N. M., and Hoffmann, T.: Formation of 3-methyl-1,2,3-butanetricarboxylic acid via gas phase oxidation of pinonic acid - A mass spectrometric study of SOA aging, *Atmos. Chem. Phys.*, 12, 1483–1496, <https://doi.org/10.5194/acp-12-1483-2012>, 2012.
- Nakajima, T., Yoon, S. C., Ramanathan, V., Shi, G. Y., Takemura, T., Higurashi, A., Takamura, T., Aoki, K., Sohn, B. J., Kim, S. W., Tsuruta, H., Sugimoto, N., Shimizu, A., Tanimoto, H., Sawa, Y., Lin, N. H., Lee, C. T., Goto, D., and Schutgens, N.: Overview of the atmospheric brown cloud east Asian regional experiment 2005 and a study of the aerosol direct radiative forcing in east Asia, *J. Geophys. Res. Atmos.*, 112, D24S91, <https://doi.org/10.1029/2007JD009009>, 2007.
- Ng, N. L., Kwan, A. J., Surratt, J. D., Chan, A. W. H., Chhabra, P. S., Sorooshian, A., Pye, H. O. T., Crounse, J. D., Wennberg, P. O., Flagan, R. C., and Seinfeld, J. H.: Secondary organic aerosol (SOA) formation from reaction of isoprene with nitrate radicals (NO<sub>3</sub>), *Atmos. Chem. Phys.*, 8(14), 4117–4140, 2008.
- Offenberg, J., Lewis, C., Lewandowski, M., Jaoui, M., Kleindienst, T. E., and Edney, E. O.: Contributions of toluene and  $\alpha$ -pinene to SOA formed in an irradiated toluene/ $\alpha$ -pinene/NO<sub>x</sub>/air mixture: Comparison of results using <sup>14</sup>C content and SOA organic tracer methods, *Environ. Sci. Tec.*, 41, 3972–3976, 2007.
- Pashynska, V., Vermeylen, R., Vas, G., Maenhaut, W., and Claeys, M.: Development of a gas chromatographic/ion trap mass spectrometric method for the determination of levoglucosan and saccharidic compounds in atmospheric aerosols. Application to urban aerosols, *J. Mass Spectrom.*, 37(12), 1249–1257, 2002.
- Pio, C., Legrand, M., Alves, C. A., Oliveira, T., Afonso, J., Caseiro, A., Puxbaum, H., Sánchez-Ochoa, A., and Gelencsér, A.: Chemical composition of atmospheric aerosols during the 2003 summer intense forest fire period, *Atmos. Environ.*, 42, 7530–7543, 2008.

- Ramanathan, V., Li, F., Ramana, M. V., Praveen, P. S., Kim, D., Corrigan, C. E., Nguyen, H., Stone, E. A., Schauer, J. J., Carmichael, G. R., Adhikary, B., and Yoon, S. C.: Atmospheric brown clouds: Hemispherical and regional variations in long-range transport, absorption, and radiative forcing, *J. Geophys. Res. Atmos.*, 112, D22S21, <https://doi.org/10.1029/2006JD008124>, 2007.
- Robinson, A. L., Donahue, N. M., Shrivastava, M. K., Weitkamp, E. A., Sage, A. M., Grieshop, A. P., Lane, T. E., Pierce, J. R., and Pandis, S. N.: Rethinking organic aerosols: Semivolatile emissions and photochemical aging, *Science*, 315, 1259–1262, 2007.
- Salazar, G., Zhang, Y. L., Agrios, K., and Szidat, S.: Development of a method for fast and automatic radiocarbon measurement of aerosol samples by online coupling of an elemental analyzer with a MICADAS AMS, *Nucl. Instruments Methods Phys. Res. Sect. B Beam Interact. with Mater. Atoms*, 361, 163–167, <https://doi.org/10.1016/j.nimb.2015.03.051>, 2015.
- Schmidl, C., Marr, I. L., Caseiro, A., Kotianova, P., Berner, A., Bauer, H., Kasper-Giebl, A., and Puxbaum, H.: Chemical characterisation of fine particle emissions from wood stove combustion of common woods growing in mid-European Alpine regions, *Atmos. Environ.*, 42, 126–141, 2008a.
- Schmidl, C., Bauer, H., Dattler, A., Hitzenberger, R., Weissenboeck, G., Marr, I. L., and Puxbaum, H.: Chemical characterisation of particle emissions from burning leaves, *Atmos. Environ.*, 42(40), 9070–9079, <https://doi.org/10.1016/J.Atmosenv.2008.09.010>, 2008b.
- Shaw, S. L., Gantt, B., and Meskhidze, N.: Production and emissions of marine isoprene and monoterpenes: A Review, *Adv. Meteorol.*, 2010, 408696, <https://doi.org/10.1155/2010/408696>, 2010.
- Simoneit, B. R. T.: Biomass burning-a review of organic tracers for smoke from incomplete combustion, *Appl. Geochem.*, 17, 129–162, 2002.
- Simoneit, B. R. T., Elias, V. O., Kobayashi, M., Kawamura, K., Rushdi, A. I., Medeiros, P. M., Rogge, W. F., and Didyk, B. M.: Sugars-dominant water-soluble organic compounds in soils and characterization as tracers in atmospheric particulate matter, *Environ. Sci. Technol.*, 38(22), 5939–5949, 2004a.
- Simoneit, B. R. T., Kobayashi, M., Mochida, M., Kawamura, K., Lee, M., Lim, H-J., Turpin, B. J., and Komazaki, Y.: Composition and major sources of organic compounds of aerosol particulate matter sampled during the ACE-Asia campaign, *J. Geophys. Res.*, 109(D19), D19S10, <https://doi.org/10.1029/2004jd004598>, 2004b.
- Simoneit, B. R. T., Schauer, J. J., Nolte, C. G., Oros, D. R., Elias, V. O., Fraser, M. P., Rogge, W. F., and Cass, G. R.: Levoglucosan, a tracer for cellulose in biomass burning and atmospheric particles, *Atmos. Environ.*, 33(2), 173–182, [https://doi.org/10.1016/S1352-2310\(98\)00145-9](https://doi.org/10.1016/S1352-2310(98)00145-9), 1999.
- Singh, N., Mhawish, A., Deboudt, K., Singh, R. S., and Banerjee, T.: Organic aerosols over Indo-Gangetic Plain: Sources, distributions and climatic implications, *Atmos. Environ.*, 157, 59–74, <https://doi.org/10.1016/j.atmosenv.2017.03.008>, 2017.
- Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA's hysplit atmospheric transport and dispersion modeling system, *Bull. Am. Meteorol. Soc.*, 96, 2059–2077, <https://doi.org/10.1175/BAMS-D-14-00110.1>, 2015.

1013 Stuiver, M. and Polach, H. A.: Discussion: Reporting of  $^{14}\text{C}$  data, *Radiocarbon*, 19(3),  
 1014 355–363, 1997.

1015 Sullivan, A. P., Holden, A. S., Patterson, L. A., McMeeking, G. R., Kreidenweis, S. M.,  
 1016 Malm, W. C., Hao, W. M., Wold, C. E., and Collett, Jr., J. L.: A method for smoke  
 1017 marker measurements and its potential application for determining the contribution of  
 1018 biomass burning from wildfires and prescribed fires to ambient  $\text{PM}_{2.5}$  organic carbon, *J.*  
 1019 *Geophys. Res.*, 113, D22302, <https://doi.org/10.1029/2008JD010216>, 2008.

1020 Surratt, J. D., Murphy, S. M., Kroll, J. H., Ng, N. L., Hildebrandt, L., Sorooshian, A.,  
 1021 Szmigielski, R., Vermeylen, R., Maenhaut, W., Claeys, M., Flagan, R. C., and Seinfeld, J.  
 1022 H.: Chemical composition of secondary organic aerosol formed from the photooxidation  
 1023 of isoprene, *J. Phys. Chem. A*, 110(31), 9665–9690, 2006.

1024 Surratt, J. D., Chan, A. W. H., Eddingsaas, N. C., Chan, M. N., Loza, C. L., Kwan, A. J.,  
 1025 Hersey, S. P., Flagan, R. C., Wennberg, P. O., and Seinfeld, J. H.: Reactive intermediates  
 1026 revealed in secondary organic aerosol formation from isoprene, *Proc. Natl. Acad. Sci.*  
 1027 *USA*, 107(15), 6640–6645, 2010.

1028 Szidat, S., Jenk, T. M., Synal, H. A., Kalberer, M., Wacker, L., Hajdas, I., Kasper-Giebl, A.,  
 1029 and Baltensperger, U.: Contributions of fossil fuel, biomass-burning, and biogenic  
 1030 emissions to carbonaceous aerosols in Zurich as traced by  $^{14}\text{C}$ , *J. Geophys. Res. Atmos.*,  
 1031 111, D07206, <https://doi.org/10.1029/2005JD006590>, 2006.

1032 Szmigielski, R., Surratt, J. D., Gómez-González, G., Van der Veken, P., Kourtchev, I.,  
 1033 Vermeylen, R., Blockhuys, F., Jaoui, M., Kleindienst, T. E., Lewandowski, M.,  
 1034 Offenberg, J. H., Edney, E. O., Seinfeld, J. H., Maenhaut, W., and Claeys, M.: 3-Methyl-  
 1035 1,2,3-butanetricarboxylic acid: An atmospheric tracer for terpene secondary organic  
 1036 aerosol, *Geophys. Res. Lett.*, 34, L24811, <https://doi.org/10.1029/2007GL031338>, 2007.

1037 Theodosi, C., Panagiotopoulos, C., Nouara, A., Zampas, P., Nicolaou, P., Violaki, K.,  
 1038 Kanakidou, M., Sempéré, R., and Mihalopoulos, N.: Sugars in atmospheric aerosols over  
 1039 the Eastern Mediterranean, *Prog. Oceanogr.*, 163, Special Issue: SI, 70-81,  
 1040 <https://doi.org/10.1016/j.pocean.2017.09.001>, 2018.

1041 Tyagi, P., Kawamura, K., Kariya, T., Bikkina, S., Fu, P., and Lee, M.: Tracing atmospheric  
 1042 transport of soil microorganisms and higher plant waxes in the East Asian outflow to the  
 1043 North Pacific Rim by using hydroxy fatty acids: Year-round observations at Gosan, Jeju  
 1044 Island, *J. Geophys. Res.*, 122, 4112–4131, <https://doi.org/10.1002/2016JD025496>, 2017.

1045 Verma, S. K., Kawamura, K., Chen, J., Fu, P., and Zhu, C.: Thirteen years of observations on  
 1046 biomass burning organic tracers over Chichijima Island in the western North Pacific: An  
 1047 outflow region of Asian aerosols, *J. Geophys. Res.*, 120, 4155–4168,  
 1048 <https://doi.org/10.1002/2014JD022224>, 2015.

1049 Verma, S. K., Kawamura, K., Chen, J., and Fu, P.: Thirteen years of observations on primary  
 1050 sugars and sugar alcohols over remote Chichijima Island in the western North Pacific,  
 1051 *Atmos. Chem. Phys.*, 18, 81–101, <https://doi.org/10.5194/acp-18-81-2018>, 2018.

1052 Wang, G. and Kawamura, K.: Molecular characteristics of urban organic aerosols from  
 1053 Nanjing: A case study of a mega-city in China, *Environ. Sci. Technol.*, 39(19), 7430–  
 1054 7438, 2005.

- Wang, G. H., Kawamura, K., and Lee, M.: Comparison of organic compositions in dust storm and normal aerosol samples collected at Gosan, Jeju Island, during spring 2005, *Atmos. Environ.*, 43(2), 219–227, <https://doi.org/10.1016/J.Atmosenv.2008.09.046>, 2009a.
- Wang, G. H., Li, J. J., Cheng, C. L., Zhou, B. H., Xie, M. J., Hu, S. Y., Meng, J. J., Sun, T., Ren, Y. Q., Cao, J. J., Liu, S. X., Zhang, T., and Zhao, Z. Z.:** Observation of atmospheric aerosols at Mt. Hua and Mt. Tai in central and east China during spring 2009-Part 2: Impact of dust storm on organic aerosol composition and size distribution, *Atmos. Chem. Phys.*, 12(9), 4065–4080, 2012.
- Wang, W., Kourtchev, I., Graham, B., Cafmeyer, J., Maenhaut, W., and Claeys, M.: Characterization of oxygenated derivatives of isoprene related to 2-methyltetrols in Amazonian aerosols using trimethylsilylation and gas chromatography/ion trap mass spectrometry, *Rapid Commun. Mass Spectrom.*, 19, 1343–1351, 2005.
- Wang, Y. Q., Zhang, X. Y., and Draxler, R. R.: TrajStat: GIS-based software that uses various trajectory statistical analysis methods to identify potential sources from long-term air pollution measurement data, *Environ. Model. Softw.*, 24, 938–939, <https://doi.org/10.1016/j.envsoft.2009.01.004>, 2009b.
- Wu, J., Kong, S., Zeng, X., Cheng, Y., Yan, Q., Zheng, H., Yan, Y., Zheng, S., Liu, D., Zhang, X., Fu, P., Wang, S., and Qi, S.:** First high-resolution emission inventory of levoglucosan for biomass burning and non-biomass burning sources in China, *Environ. Sci. Technol.*, 55, 3, 1497–1507, 2021.
- Yan, C., Zheng, M., Sullivan, A. P., Shen, G., Chen, Y., Wang, S., Zhao, B., Cai, S., Desyaterik, Y., Li, X., Zhou, T., Gustafsson, Ö., and Collett, J. L.: Residential coal combustion as a source of levoglucosan in China, *Environ. Sci. Technol.*, 52, 3, 1665–1674, <https://doi.org/10.1021/acs.est.7b05858>, 2018.
- Yttri, K. E., Dye, C., and Kiss, G.: Ambient aerosol concentrations of sugars and sugar-alcohols at four different sites in Norway, *Atmos. Chem. Phys.*, 7, 4267–4279, 2007.
- Zangrando, R., Barbaro, E., Kirchgeorg, T., Vecchiato, M., Scalabrin, E., Radaelli, M., Dorđević, D., Barbante, C., and Gambaro, A.: Five primary sources of organic aerosols in the urban atmosphere of Belgrade (Serbia), *Sci. Total Environ.*, 571, 1441–1453, <https://doi.org/10.1016/j.scitotenv.2016.06.188>, 2016.
- Zhang, Y. L., Perron, N., Ciobanu, V. G., Zotter, P., Minguillón, M. C., Wacker, L., Prévôt, A. S. H., Baltensperger, U., and Szidat, S.: On the isolation of OC and EC and the optimal strategy of radiocarbon-based source apportionment of carbonaceous aerosols, *Atmos. Chem. Phys.*, 12, 10841–10856, <https://doi.org/10.5194/acp-12-10841-2012>, 2012.
- Zhang, Y. L., Huang, R. J., El Haddad, I., Ho, K. F., Cao, J. J., Han, Y., Zotter, P., Bozzetti, C., Daellenbach, K. R., Canonaco, F., Slowik, J. G., Salazar, G., Schwikowski, M., Schnelle-Kreis, J., Abbaszade, G., Zimmermann, R., Baltensperger, U., Prévôt, A. S. H., and Szidat, S.: Fossil vs. non-fossil sources of fine carbonaceous aerosols in four Chinese cities during the extreme winter haze episode of 2013, *Atmos. Chem. Phys.*, 15(3), 1299–1312, <https://doi.org/10.5194/acp-15-1299-2015>, 2015.
- Zhang, Y. L., Kawamura, K., Agrios, K., Lee, M., Salazar, G., and Szidat, S.: Fossil and nonfossil sources of organic and elemental carbon aerosols in the outflow from Northeast China, *Environ. Sci. Technol.*, 50, 6284–6292, <https://doi.org/10.1021/acs.est.6b00351>, 2016.

1099 Zhu, C., Kawamura, K., and Kunwar, B.: Effect of biomass burning over the western North  
1100 Pacific Rim: wintertime maxima of anhydrosugars in ambient aerosols from Okinawa.,  
1101 Atmos. Chem. Phys., 15, 1959–1973, <https://doi.org/10.5194/acp-15-1959-2015>, 2015a.

1102 Zhu, C., Kawamura, K., and Kunwar, B.: Organic tracers of primary biological aerosol  
1103 particles at subtropical Okinawa island in the western North pacific rim, J. Geophys. Res.,  
1104 120, 5504-5523, <https://doi.org/10.1002/2015JD023611>, 2015b.

1105

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119

1120

1121

1122

1123

1124

1125

1126 **Table 1.** Concentrations of identified sugar compounds and BSOA tracers (ng m<sup>-3</sup>) in the atmospheric  
1127 aerosol samples from Gosan.

Species	Annual	Summer	Fall	Winter	Spring
	Avg. <sup>a</sup> ± S.D. <sup>b</sup> Min. <sup>c</sup> , Max. <sup>d</sup>	Avg. ± S.D. Min., Max.	Avg. ± S.D. Min., Max.	Avg. ± S.D. Min., Max.	Avg. ± S.D. Min., Max.
<b>Anhydrosugars</b>					
Levoglucozan (Lev)	17.6 ± 16.8 0.60, 45.9	2.92 ± 3.89 0.60, 9.81	21.7 ± 19.0 2.30, 43.9	39.2 ± 6.60 32.0, 45.9	12.7 ± 11.6 1.45, 33.4
Mannosan (Man)	1.57 ± 1.82 0.05, 6.74	0.18 ± 0.24 0.05, 0.61	1.69 ± 1.49 0.13, 3.66	3.63 ± 2.28 1.47, 6.74	1.31 ± 1.57 0.08, 4.08
Galactosan (Gal)	2.28 ± 2.10 0.14, 6.78	0.64 ± 0.68 0.14, 1.82	2.45 ± 2.13 0.35, 4.92	5.21 ± 1.64 3.40, 6.78	1.65 ± 1.26 0.50, 3.88
<b>Primary Sugars</b>					
Glucose	18.8 ± 27.1 2.45, 122	13.4 ± 18.2 2.45, 45.6	16.5 ± 15.7 2.68, 33.6	4.74 ± 3.14 2.88, 9.44	32.4 ± 41.1 4.87, 122
Fructose	10.3 ± 15.9 0.97, 74.0	4.90 ± 5.15 0.97, 13.7	7.48 ± 6.99 1.71, 16.2	3.82 ± 4.45 1.56, 10.5	19.8 ± 25.0 2.69, 74.0
Sucrose	16.1 ± 32.2 0.26, 140	1.46 ± 1.05 0.68, 3.28	9.74 ± 12.1 0.76, 27.2	8.87 ± 16.3 0.42, 33.3	35.1 ± 50.5 0.26, 140
Trehalose	2.42 ± 1.97 0.65, 7.03	2.72 ± 2.53 0.97, 6.98	3.71 ± 2.29 1.18, 7.03	1.21 ± 0.42 0.72, 1.63	1.98 ± 1.55 0.65, 5.33
Xylose	0.81 ± 0.65 0.04, 2.03	0.23 ± 0.23 0.04, 0.63	0.86 ± 0.68 0.14, 1.70	1.59 ± 0.37 1.21, 2.03	0.74 ± 0.55 0.16, 1.68
<b>Sugar alcohols</b>					
Arabitol	3.96 ± 4.24 0.47, 18.7	5.64 ± 7.46 1.20, 18.7	5.27 ± 3.51 2.27, 10.9	1.06 ± 0.60 0.47, 1.91	3.47 ± 2.19 1.18, 6.30
Mannitol	4.61 ± 5.54 0.25, 22.0	7.39 ± 8.65 1.71, 22.0	6.64 ± 6.03 1.69, 16.7	0.99 ± 0.45 0.55, 1.60	3.24 ± 2.67 0.25, 7.03
Erythritol	0.62 ± 0.43 0.12, 1.52	0.92 ± 0.53 0.42, 1.52	0.93 ± 0.33 0.55, 1.27	0.42 ± 0.27 0.16, 0.80	0.30 ± 0.12 0.12, 0.48
Inositol	0.34 ± 0.39 0.04, 1.53	0.29 ± 0.41 0.08, 1.03	0.56 ± 0.60 0.10, 1.53	0.13 ± 0.07 0.08, 0.23	0.35 ± 0.28 0.04, 0.73
<b>Isoprene SOA tracers</b>					
2-MGA	0.99 ± 0.70 0.17, 2.79	1.61 ± 1.17 0.17, 2.79	0.95 ± 0.43 0.51, 1.61	0.67 ± 0.14 0.49, 0.81	0.76 ± 0.35 0.20, 1.32
Σ2-MLTs	1.04 ± 1.40 0.05, 5.81	2.48 ± 1.98 0.51, 5.81	1.34 ± 1.16 0.33, 2.74	0.20 ± 0.05 0.15, 0.26	0.29 ± 0.25 0.05, 0.82
ΣC5-alkene triols	0.20 ± 0.25 0.02, 1.17	0.46 ± 0.44 0.02, 1.17	0.13 ± 0.05 0.05, 0.18	0.09 ± 0.03 0.05, 0.13	0.14 ± 0.11 0.02, 0.30
<b>Monoterpene SOA tracers</b>					
<i>cis</i> -pinonic acid	0.15 ± 0.14 0.02, 0.52	0.12 ± 0.13 0.03, 0.36	0.08 ± 0.06 0.02, 0.16	0.07 ± 0.03 0.02, 0.10	0.26 ± 0.17 0.08, 0.52
pinic acid	1.67 ± 1.01 0.72, 4.81	2.40 ± 1.67 0.99, 4.81	1.53 ± 0.57 0.75, 2.11	1.07 ± 0.25 0.84, 1.42	1.61 ± 0.78 0.72, 2.94
3-HGA	2.30 ± 2.38 0.19, 10.6	3.46 ± 4.39 0.19, 10.6	1.52 ± 0.77 0.76, 2.79	1.40 ± 0.39 0.94, 1.88	2.54 ± 1.83 0.38, 4.57
MBTCA	5.11 ± 4.54 0.29, 19.6	8.44 ± 6.88 3.00, 19.6	6.24 ± 3.64 1.17, 9.49	1.90 ± 0.85 0.97, 2.83	3.77 ± 2.96 0.29, 8.08

<sup>a</sup>Average, <sup>b</sup>Standard deviation, <sup>c</sup>Minimum, <sup>d</sup>Maximum. 2-MGA: 2-methylglyceric acid, 2-MLTs: 2-methyltetrols, 3-HGA: 3-hydroxyglutaric acid, MBTCA: 3-methyl-1,2,3-butanetricarboxylic acid.

**Table 2.** Statistical summary of diagnostic ratios and carbonaceous components contribution in Gosan aerosols.

Species	Annual	Summer	Fall	Winter	Spring
	Avg. <sup>a</sup> ± S.D. <sup>b</sup> Min. <sup>c</sup> , Max. <sup>d</sup>	Avg. ± S.D. Min., Max.	Avg. ± S.D. Min., Max.	Avg. ± S.D. Min., Max.	Avg. ± S.D. Min., Max.
<b>Diagnostic ratios</b>					
Lev/Man	15.1 ± 6.76 6.81, 31.3	17.1 ± 8.21 8.58, 28.4	13.7 ± 2.51 11.1, 17.3	13.5 ± 6.24 6.81, 21.7	15.4 ± 8.76 6.98, 31.3
Man/Gal	0.55 ± 0.32 0.10, 1.07	0.28 ± 0.13 0.10, 0.42	0.64 ± 0.20 0.37, 0.90	0.66 ± 0.27 0.43, 1.05	0.61 ± 0.43 0.16, 1.07
Lev/(Man + Gal)	4.27 ± 1.23 2.18, 6.56	3.12 ± 0.78 2.18, 4.03	5.08 ± 0.32 4.71, 5.51	4.85 ± 1.35 3.49, 6.56	4.17 ± 1.33 2.50, 6.45
Lev/K <sup>+</sup> × 10 <sup>-2</sup>	5.73 ± 5.65 0.65, 23.2	1.00 ± 0.52 0.65, 1.91	3.94 ± 2.51 1.26, 6.30	12.3 ± 7.87 6.42, 23.2	6.65 ± 4.48 1.77, 15.3
Lev/OC × 10 <sup>-2</sup>	0.73 ± 0.66 0.08, 2.29	0.14 ± 0.10 0.08, 0.31	0.84 ± 0.61 0.12, 1.56	1.60 ± 0.67 0.99, 2.29	0.56 ± 0.40 0.11, 1.05
Lev/WSOC × 10 <sup>-2</sup>	1.09 ± 0.97 0.10, 3.46	0.20 ± 0.13 0.10, 0.42	1.25 ± 0.94 0.23, 2.21	2.33 ± 0.90 1.48, 3.46	0.92 ± 0.65 0.13, 1.69
MGA/MLTs	2.18 ± 1.59 0.33, 5.40	0.67 ± 0.43 0.33, 1.33	1.04 ± 0.49 0.41, 1.55	3.60 ± 1.43 1.90, 5.40	3.27 ± 1.17 1.60, 4.75
<sup>e</sup> P/MBTCA	0.62 ± 0.61 0.17, 5.90	3.22 ± 0.62 2.47, 3.89	3.62 ± 1.67 1.52, 5.90	1.68 ± 0.64 0.83, 2.38	1.73 ± 1.07 0.34, 3.32
<b>Carbonaceous components</b>					
Isoprene derived SOC (µgC m <sup>-3</sup> )	23.7 ± 23.9 2.26, 97.4	51.0 ± 32.4 8.08, 97.4	25.9 ± 17.9 8.54, 45.7	8.52 ± 1.07 7.46, 9.87	11.3 ± 6.80 2.26, 24.9
Isoprene SOC to OC (%)	1.45 ± 1.74 0.21, 6.40	3.56 ± 2.18 1.03, 6.40	1.46 ± 1.38 0.35, 3.48	0.35 ± 0.17 0.21, 0.57	0.57 ± 0.38 0.24, 1.35
Isoprene SOC to total SOC (%)	35.5 ± 15.5 13.5, 71.1	46.6 ± 20.3 26.0, 71.1	38.1 ± 10.4 21.1, 47.3	31.6 ± 7.05 24.0, 39.7	27.9 ± 15.5 13.5, 54.0
Monoterpene SOC (µgC m <sup>-3</sup> )	40.1 ± 33.3 7.18, 154	62.7 ± 56.5 19.2, 154	40.7 ± 20.0 11.8, 57.7	19.3 ± 5.61 14.0, 25.2	35.5 ± 23.5 7.18, 62.7
Monoterpene SOC to OC (%)	2.0 ± 1.47 0.37, 5.74	3.65 ± 1.58 2.44, 5.74	2.15 ± 1.45 0.48, 3.88	0.87 ± 0.58 0.37, 1.48	1.58 ± 0.84 0.54, 3.13
Monoterpene SOC to total SOC (%)	64.5 ± 15.5 28.9, 86.5	53.4 ± 20.3 28.9, 74.0	61.9 ± 10.4 52.7, 78.9	68.4 ± 7.05 60.3, 76.1	72.1 ± 15.5 46.0, 86.5

<sup>e</sup>P: *cis*-pinonic acid + pinic acid



**Table 3.** Comparisons of the **mean** concentration (ng m<sup>-3</sup>) of anhydrosugars, sugar, and sugar alcohols in Gosan aerosols with those from different sites around the world.

Sampling sites	Sampling type	Sampling time	Anhydro-sugars	Primary sugars	Sugar alcohols	References
Gosan, South Korea	TSP	Summer Fall Winter Spring	<b>3.74</b> <b>25.9</b> <b>48.0</b> <b>15.7</b>	<b>22.8</b> <b>38.3</b> <b>20.2</b> <b>90.0</b>	<b>14.3</b> <b>13.4</b> <b>2.60</b> <b>7.36</b>	<b>This study</b>
Chennai, India	PM10	Summer Winter	127 134	15.5 11.4	7.44 4.81	Fu et al., 2010b
Mt. Tai, China	TSP	Summer (June)	224	61.1	125	Fu et al., 2012b
Alert, Canada	TSP	Winter Spring	0.32 0.02	1.14 0.18	0.25 0.36	Fu et al., 2009a
Okinawa, western North Pacific	TSP	Summer Autumn Winter Spring	0.93 2.58 6.04 3.44	73.5 56.0 34.4 101	62.9 30.7 6.03 31.2	Zhu et al., 2015a, b
Chichijima, western North Pacific	TSP	Summer Autumn Winter Spring	0.32 0.85 2.40 0.94	32.8 22.0 14.2 24.2	38.6 35.1 3.93 15.5	Verma et al., 2015, 2018
Mt. Hua, China (Non-dust storm)	PM10	April	57.8	92.5	22.4	Wang et al., 2012
Mt. Hua, China (Dust storm)	PM10	April	44.5	162	25.7	Wang et al., 2012
Nanjing, China	PM2.5	Summer Winter	151 (Lev) 268 (Lev)	59.3 42.3	11.8 13.4	Wang and Kawamura, 2005
Beijing, China	PM2.5	Summer Autumn Winter Spring	14.5 129 254 81.4	6.63 17.2 41.5 33.9	3.31 13.7 17.8 12.3	Kang et al., 2018a
Belgrade, Serbia	TSP	Autumn	425 (Lev)	116	98.4	Zangrando et al., 2016
Maine, USA	PM1	May-October	13.9	28	8.31	Medeiros et al., 2006
Crete, Greece	PM10	Year-round	14.4	32.3	6.53	Theodosi et al., 2018

1132

1133

1134

1135

Supporting information for

**Regional heterogeneities in the emission of airborne primary sugar compounds and biogenic secondary organic aerosols in the East Asian outflow: Evidence for coal combustion as a source of levoglucosan**

**Md. Mozammel Haque<sup>1,2,3</sup>, Yanlin Zhang<sup>1,2\*</sup> Srinivas Bikkina<sup>4</sup>, Meehye Lee<sup>5</sup>, and Kimitaka Kawamura<sup>3,4\*</sup>**


<sup>1</sup>*Yale-NUIST Center on Atmospheric Environment, International Joint Laboratory on Climate and Environment Change (ILCEC), Nanjing University of Information Science & Technology, Nanjing, 210044, China*

<sup>2</sup>*School of Applied Meteorology, Nanjing University of Information Science & Technology, Nanjing 210044, China*


<sup>3</sup>*Institute of Low Temperature Science, Hokkaido University, Sapporo 060-0819, Japan*

<sup>4</sup>*Chubu Institute for Advanced Studies, Chubu University, Kasugai 487-8501, Japan*


<sup>5</sup>*Department of Earth and Environmental Sciences, Korea University, Anam-dong, Sungbuk-gu, Seoul 136-701, South Korea*




**Figure S1.** HYSPLIT-based backward air mass trajectories as a function of arrival height (see color bar in each panel) and MODIS-firecounts for the TSP collected over Gosan according the four seasons, (a) winter, (b) spring, (c) summer, and (d) autumn.




**Figure S2.** Time series of the concentrations of anhydrosugars, primary sugars, and sugar alcohols during the period of study in Gosan.



**Figure S3.** Temporal variability of mass concentrations of trehalose and non-sea-salt Ca<sup>2+</sup> in TSP collected over Gosan during April 2013-April 2014.



**Figure S4.** Seasonal variability of molecular distributions of anhydrosugars, primary sugars, sugar alcohols, isoprene SOA tracers and monoterpene SOA tracers in their total mass concentration of respective group-types in TSP samples collected over Gosan during April 2013-April 2014.




**Figure S5.** Diagnostic ratios of 2-methylglyceric acid/2-methyltetrols (2-MGA/2-MTs) and *cis*-pinonic acid+pinic acid/3-methyl-1,2,3-butanetricarboxylic acid (P+P/MBTCA) in Gosan TSP samples collected during April 2013-April 2014.



1224

1225

1226



1227

1228 **Figure S6.** PMF- resolved source contributions to the organic carbon in Gosan aerosols.

1229