# **1** Comparison of the influence of two types of cold surge

# 2 on haze dispersion in Eastern China

3 Shiyue Zhang<sup>1</sup>, Gang Zeng<sup>1</sup>, Xiaoye Yang<sup>1</sup>, Ruixi Wu<sup>2</sup>, Zhicong Yin<sup>1, 3</sup>

4 <sup>1</sup> Key Laboratory of Meteorological Disaster of Ministry of Education (KLME),

5 Collaborative Innovation Center on Forecast and Evaluation of Meteorological Disasters (CIC-FEMD),

- 6 Nanjing University of Information Science and Technology, Nanjing, 210044, China
- 7 <sup>2</sup> Meteorological Bureau of Jiading District, Shanghai 201815, China

8 <sup>3</sup> Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai), Zhuhai, 519080, China

9 Correspondence to: Gang Zeng (zenggang@nuist.edu.cn)

10 Abstract. Cold surge (CS) is considered as a favorable weather process to improve air quality and is

11 widely recognized. However, there is no detailed study on the differences in the dispersion ability of

12 different types of CSs to haze days in eastern China (HD<sub>EC</sub>). This paper uses the hierarchical clustering

13 algorithm to classify the cool season (November to February of the following year) CSs across eastern

- 14 China into blocking CSs and wave-train CSs and compares their influences on the number of HD<sub>EC</sub> from
- 15 1980 to 2017. Results show that the wave-train CSs can significantly improve the visibility in eastern
- 16 China and generally improve air quality for about 2 days longer than the blocking CSs, which indicates
- 17 that the blocking CSs have a weaker ability to dissipate HD<sub>EC</sub> compared with the wave-train CSs. The
- $18 \qquad CSs \ affect \ the \ HD_{EC} \ by \ changing \ meteorological \ elements \ like \ thermal \ inversion \ potential, \ horizontal$
- 19 surface wind, sea level pressure (SLP), and surface air temperature (SAT). 4 days after the CSs outbreak,
- 20 the variations of thermal inversion potential and horizontal surface wind of two types of CSs tend to be
- 21 consistent. However, the negative SAT anomalies and the positive SLP anomalies caused by the blocking
- 22 CSs lasted shorter than those caused by the wave-train CSs, which forms favorable conditions for the
- 23 rapid growth of HD<sub>EC</sub>. Furthermore, results show that in recent years, especially after the 1990s, the
- 24 frequency of wave-train CSs has decreased significantly, while the frequency of blocking CSs has slightly

1

25 increased, indicating that the overall ability of CSs to dissipate HD<sub>EC</sub> has weakened in general. This work

26 may provide reference for the future formulation of haze control policies in East Asia.

27

批注[张1]: Reviewer#1 Technical corrections (1)

批注[张2]: Reviewer#1 Technical corrections (2)

批注[张3]: Reviewer#1 Specific comments (1)

## 28 1. Introduction

29 Haze in eastern China is generally referred to as the polluted particulate aerosols suspended in the 30 air (Yin et al., 2019a), and can reduce visibility and affect traffic and ecological sustainability (Xu et al., 31 2013; Xie et al., 2014; Wang et al., 2016). Studies have shown that the haze in China is mainly 32 concentrated in the eastern region of China (EC), and its peak is noticeable in winter and spring (Wang 33 et al., 2015, 2016). During haze days, the concentration of aerosol particles increases and results in a 34 wide range of visibility decline (Luo et al., 2001; Xu, 2001; Wu et al., 2012; Fu et al., 2013; Wu et al., 35 2014). For example, in the winter of 2015, severe haze in the Beijing-Tianjin-Hebei region affected more 36 than 500,000 square kilometers, causing heavy pollution in 37 cities (Chang et al., 2016; Zhang et al., 37 2016). After this event, researchers and policymakers paid more attention to the studies related to haze 38 events. Besides, strict control measures of air pollution and energy emissions have also been put in place. 39 Many studies indicated that the long-term trends of haze are closely related to fossil-fuel emissions 40 (Shi et al., 2008; Wei et al., 2017). On the other hand, meteorological conditions also play an important 41 role in determining regional air quality. In addition to the influence of human activities, the formation of 42 haze is closely related to static and calm weather conditions, such as strong thermal inversion potential 43 (TIP), negative sea level pressure (SLP) anomaly, and weak wind speed (Niu et al., 2010; Cai et al., 44 2017). In recent years, due to the decreased relative humidity, it is difficult for haze particles to transform 45 into fog drops, making the number of haze days present a rising trend (Ding and Liu, 2014). In addition, 46 the anomalies of atmospheric circulation caused by global warming may also enhance the stability of the 47 lower atmosphere, which leads to more severe and frequent haze pollution (Cai et al., 2017). All these 48 emphasize that the threat of haze to human society could be more serious in the near future. 49 Global warming leads to the decrease of cold days and cold surges (CSs) by raising the surface air 50 temperature (SAT), which also provides favorable conditions for the increase of haze days (Lin et al., 51 2009). CS is a typical extreme weather process in East Asia, which significantly impacts the atmospheric 52 circulation to improve the local air quality (Hu et al., 2000; Qu et al., 2015; Wang et al., 2016). With the 53 outbreak of CSs, a series of abrupt variations of meteorological elements such as the positive SLP 54 anomaly, the decrease of SAT, and the enhancement of north wind component will occur in the areas 55 where the CSs pass (Compo et al., 1999). When a CS occurs, the arrival of fresh and dry cold air can

56 dissipate and reduce local air pollutants (Lin et al., 2008). Wang et al. (2016) proposed that the " early in

批注 [张4]: Reviewer#1 Specific comments (2)

批注[张5]: Reviewer#1 Technical corrections (4) the north and late in the south " feature of air quality improvement in mainland China results from the cold air masses moving southward from high latitudes to low latitudes after the outbreak of CSs. Although some studies have shown that the weakening of East Asian Winter Monsoon and global warming leads to the decrease of CSs (Qu et al., 2015; Wang et al., 2006), extreme low-temperature events are still frequent (Park et al., 2011a), which makes the assessment of haze dispersion capacity of cold air activities still full of uncertainty.

63 Previous studies have shown that the outbreak of CSs has an obvious effect on haze dispersion (Lin 64 et al., 2009; Hien et al., 2011; Ashfold et al., 2017). However, most of them analyzed the haze variation 65 during the CSs based on case analyses or considering the interannual influence of CS frequency on haze. Furthermore, studies have shown that there are large differences between individual cases of CSs in terms 66 67 of circulation anomalies, influence path and range (Park et al., 2014; Cai et al., 2019). Therefore, it is 68 necessary to consider the influence of classified CSs on haze. Based on this limitation, the following two 69 questions are proposed in this paper: Are there different effects of CSs' types on the haze days in EC 70 (HD<sub>EC</sub>)? If so, what is the physical mechanism that makes the difference? The solution to these issues 71 will help us understand the mechanism of CSs in dissipating the haze and improving its predictability in 72 the future. The variation of HD<sub>EC</sub> and its relationship with different types of CSs are described in section 73 3.1. 74 The rest of this paper is organized as follows: Section 2 introduces the data and methods, while

The rest of this paper is organized as follows: Section 2 introduces the data and methods, while section 3 presents the study findings. The variation of  $HD_{EC}$  and its relationship with two types of CSs are shown in section 3.1. Section 3.2 explains the reason why different types of CSs have different abilities to dissipate  $HD_{EC}$ . Finally, the main conclusions and discussion are presented in section 4.

#### 78 2. Data and Methods

## 79 2.1 Data

The datasets employed in this study were: (1) daily ERA-Interim atmospheric fields including SLP, air temperature at different levels, SAT, horizontal wind, and geopotential height (GPH) provided by the European Center for Medium-Range Weather Forecasts (ECMWF) (Dee et al., 2011). They have a horizontal resolution of 1.0°×1.0°. (2) daily observational datasets for 756 meteorological stations from 1980 to 2017 collected by the National Meteorological Information Center of China Meteorological 批注[张6]: Reviewer#1 Specific comments (3)

85	Administration, including relative humidity (Rhum), visibility, and weather phenomena (It refers to the
86	physical phenomena of precipitation, surface condensation, visual range obstacle, atmospheric optics,
87	lightning, and wind in the atmosphere and near-surface). These datasets were observed four times per
88	day (02:00, 08:00, 14:00, and 20:00LT). Stations with more than 5% missing data were eliminated, while
89	sporadic missing data (less than 3 days) were filled by cubic spline interpolation. Successive (3 days and
90	more) missing data were discarded.
91	In addition, we also used PM2.5 concentration data (acquired from the China National
92	Environmental Monitoring Centre and were widely used in the research of PM2.5 in China, refer to Yin
93	et al. (2021) and Wang et al. (2021)) together with NCEP/NCAR Reanalysis datasets (Kalnay et al., 1996)
94	to verify the response of PM2.5 to the two types of CSs from 2014 to 2019. Such a scheme can effectively
95	avoid the dependence of conclusions on datasets and ensure that our results are based on many cold surge
96	samples.
97	2.2 Definition of HD <sub>EC</sub>
98	The visibility and relative humidity are routinely used in meteorology to distinguish the haze (Yin
99	et al., 2017). After filtering the other weather parameters affecting visibility (i.e., dust, precipitation,
100	sandstorm), we defined a haze day as a day with visibility lower than 10 km and the Rhum less than 90 $\%$
101	occurring at any of the four times (02:00, 08:00, 14:00, and 20:00LT) (Yin et al., 2019a) from 1980-2013.
102	However, the visibility observation in China was switched from manual observation to high temporal
103	resolution automated observation after 2013 (Yin et al., 2017). Therefore, because of systematic biases
104	between manual and automated observation, the 7.5 km automated observed visibility (Zhang et al., 2021)
105	and Rhum less than 90 % are suggested as the occurrence of haze. Figure S1 shows the climatology of
106	haze days in China from 1980 to 2017. The haze days are mainly concentrated in the EC (22°N-37°N,
107	106°E-121°E), which is selected as the target area in the present study. The number of the monthly
108	average of $HD_{EC}$ by regional average indicated that the $HD_{EC}$ mainly peaks (Figure S1b) in the cool
109	season (November to February of the following year (NDJF)).
110	
111	2.3 Definition of CS

The CS is a cooling process superimposed on a cold day (Park et al., 2011a). The outbreak of the

4

112

批注[张7]: Reviewer#1 Technical corrections (7)

批注 [张8]: Reviewer#1 Technical corrections (5)

批注[张9]: Reviewer#1 Technical corrections (6)

批注 [张10]: Reviewer#1 Specific comments (10)

批注 [张11]: Reviewer#2 Comments (1)

批注 [张12]: Reviewer#1 Specific comments (5)

- 113 CSs in East Asia is closely related to the Siberian high, known as the Siberian high surge (Compo et al.,
- 114 1999). In this study, we first divided EC into  $5^{\circ} \times 5^{\circ}$  grid boxes as shown in Figure 1 and then calculated
- 115 the average SAT for each box to avoid the extreme SAT anomaly in a single grid. To explore the impact
- 116 of CSs on HD<sub>EC</sub>, the selection of CS in this paper fulfills the following three criteria (Park et al., 2008,
- 117 2011b, 2015): (1) the maximum pressure center in the domain of the Siberian high (Figure 1) should
- exceed 1,035 hPa on the day of the CS outbreak. (2) the daily temperature drops  $(SAT_t SAT_{t-1})$  and the
- 119 SAT anomalies should exceed -1.5 standard deviation (i.e., the standard deviation of the SAT from 1980
- 120 to 2017) at least one box (setting the threshold to 1.5 times the standard deviation can not only obtain the
- 121 CS time with sufficient intensity relative to the local climate but also ensure enough CS samples). (3) the
- 122 haze day appeared in the box where the CS occurs from -2 days to 0 day related to the occurred CS. A
- total of 187 CSs that might affect a haze day in EC were identified.



批注 [张13]: Reviewer#2 Comments (2)

批注[张14]: Reviewer#1 Specific comments (6)



- 125 Figure 1. The domain of Siberian high (thick solid box; 40°N–60°N, 80°E–100°E) and EC (dotted box; 22°N-37°N,
- 126 106°E-121°E) divided into 5°× 5° grid boxes. Shadings indicate the cool season mean SLP.

## 127 2.4 Classification of CS

Referring to the research of Park et al. (2008) and Yang et al. (2020b), this paper uses the hierarchical clustering algorithm (HCA) to classify the CS types. The HCA (Rokach et al., 2005) creates a hierarchical nested clustering tree by calculating the similarity between different categories of data samples. In the clustering tree, the original data samples of different types are at the lowest level of the tree, and the top level of the tree is the root point of a cluster. This paper uses Euclidean distance to calculate the distance (similarity) between different samples. Here, we introduce the silhouette coefficient 134 to determine the best classification number (Rousseeuw, 1987). For any sample *i*, the silhouette

136 
$$s(i) = \frac{b(i) - a(i)}{max\{a(i), b(i)\}}$$

137 a(i) means the average distance from sample *i* to all other samples in the cluster it belongs to, and b(i)138 means the lowest average distance from sample *i* to all samples in any other cluster. The silhouette 139 coefficient of the clustering result is the average of the silhouette coefficients of all samples. The range 140 of silhouette coefficient is - 1 to 1. The closer to 1, the better the classification results. In this paper, the 141 GPH anomalies in the region of 30°E-170°E, 0°-80°N at 300 hPa on the outbreak day of CS was used to 142 perform HCA.

143

#### 144 3. Results and Discussion

## 145 **3.1** The influence of two types of CSs on HD<sub>EC</sub>

The circulation evolution with different types of CSs is quite different (Park et al., 2013), which 146 147 leads to the different distribution of surface meteorological conditions and haze. Here we display the 148 evolutions of two typical CS events (Figure 2). These two events were selected because they belong to 149 different types of CSs, referring to Park et al. (2008 and 2014), and have a large different effect on HD<sub>EC</sub>. Figures 2a, 2e, 2i, 2m show a CS that occurred on December 31, 2000, with positive and negative GPH 150 151 anomalies over the sub-arctic and East Asian coast, respectively, which meet the definition of the 152 blocking CS (Park et al., 2015). The blocking structure has a relatively stable lifecycle, so the HD<sub>EC</sub> only 153 has a certain dispersion on the day of the CS outbreak, and heavy HD<sub>EC</sub> begins to emerge 2 days after 154 the CS outbreaks (Figures 2j and 2n). Figures 2c, 2g, 2k, 2o indicate a CS that occurred on January 7, 155 1983, which meets the definition of the wave-train CS (Chai et al., 2002; Park et al., 2015). The CS is associated with the wave-train structure of "-+ - +" at the upper troposphere. The cold air moves from 156 west to east and invades EC along with this zonal wave-train (Yang et al., 2020a). Therefore, the wave-157 158 train CS has a better ability to disperse HD<sub>EC</sub>, and no HD<sub>EC</sub> appears for about a week after the wave-train 159 CS erupts (Figures 21 and 2p).

批注[张16]: Reviewer#1 Specific comments (19)

批注 [张15]: Reviewer#1 Specific comments (7)

(1)

批注[张17]: Reviewer#1 Technical corrections (8)



161 Figure 2. Composite of GPH anomalies (shading; gpm) at 300 hPa from day -2 to day 4 for case1 CS outbreak on
162 December 31, 2000 (a, e, i, m), and case2 CS outbreak on January 7, 1983 (c, g, k, o), and the related spatial
163 distribution of HD<sub>EC</sub> (shading) (b, f, j, n and d, h, l, p).
164

165 The case analysis results indicate that the ability of different types of CSs to haze dispersion is different. According to the principle of maximum distance between clusters, the CSs from 1980-2017 166 167 can be classified into two categories (Figure 3a). The silhouette coefficient of the clustering model shows 168 that when all CSs are divided into two types, the difference between them is the largest. Figures 3c and 169 3d show the composite GPH anomalies at 300 hPa that depict the blocking CSs and wave-train CSs. 170 Such classification results are consistent with previous studies (Park et al., 2014, 2015), which are mainly manifested in the location of the center of circulation anomalies. The cold air of the blocking CSs mainly 171 172 moves in a north-south direction that invades from Siberia to EC, and the cold air of the wave-train CSs

173 originating from the Ural Mountains converged near Lake Baikal then invaded EC.

批注 [张18]: Reviewer#1 Specific comments (7)



174

179

Figure 3. Hierarchical clustering tree (a) and silhouette coefficient (b) of cool season CSs in the EC. Composite of
GPH anomalies at 300 hPa (shading; gpm. Dotted areas are statistically significant at the 95% confidence level.) and
horizontal wind anomalies at 850 hPa (vectors; m s<sup>-1</sup>. Only shows the areas which are statistically significant at the
95% confidence level) relative to blocking CSs (c) and wave-train CSs (d).

180 Figure 4 presents the circulation anomalies from day -2 to day 6 of the CSs events in the two types 181 and the related evolution of HD<sub>EC</sub>. For the blocking CSs, largely positive and negative GPH anomalies 182 at 300 hPa are found over the Arctic and EC. The HD<sub>EC</sub> tends to dissipate first and then increase rapidly 183 after the CSs erupt. By 6 days after the CSs erupt, the haze reaches a relatively large value (Figure 4i). For the wave-train CSs, a zonal wave-train structure of GPH anomalies can be seen in the midlatitude of 184 185 the Eurasian landmass. From day -2 to day 6, the zonal wave-train appears to move toward EC. With the 186 movement of the wave-train, the haze dissipates rapidly, and EC can maintain high air quality weather 187 for a longer time. Sporadic HD<sub>EC</sub> does not appear until 6 days after, which is different from the existence 188 of HD<sub>EC</sub> when the blocking CSs occur. It shows that blocking CSs have a weak ability to dissipate haze 189 compared with wave-train CSs. This conclusion is also consistent with the individual cases mentioned 190 above (Figure 2). In addition, we verify the response of PM2.5 to the two types of CSs from 2014 to 191 2019, and similar results were obtained (Figure S2). It shows that the selection of data sets does not affect 192 the main conclusions of this paper. 8

批注 [张19]: Reviewer#1 Specific comments (8)

批注 [张20]: Reviewer#1 Technical corrections (10)

Specific comments (10)

批注 [张21]: Reviewer#1							
	Technical corrections (11)						
	批注 [张22]: Reviewer#1						
	Technical corrections (12)						

批注 [张23]: Reviewer#1
Technical corrections (13)
批注 [张24]: Reviewer#1



# 批注 [张25]: Reviewer#1

Specific comments (9)

Reviewer#2

Comments (1)

193 194

Figure 4. Composite of GPH anomalies at 300 hPa (contour; in intervals of 20 gpm) from day -2 to day 6 relative

195 to the outbreak of CSs and the corresponding spatial distribution of HD<sub>EC</sub> (shading, only shows the areas which are

- 196 statistically significant at the 95% confidence level by t-test.) for blocking CSs (a, c, e, g, i) and wave-train CSs (b,
- 197 d, f, h, j). The number in the lower right corner of each figure represents the ratio of the grids of HDEc to that of EC.

批注 [张26]: Reviewer#1 Specific comments (9) and (11)

#### 198 3.2 Why are two types of CSs different in dispersing HD<sub>EC</sub>?

199 According to the definition of HD<sub>EC</sub>, which combines visibility and Rhum in this study, we 200 composite the daily visibility anomalies and Rhum anomalies for 9 days before and after the outbreak of 201 the blocking CSs and wave-train CSs, respectively (Figure 5). This helps to understand why the two 202 types of CSs have different abilities to disperse HD<sub>EC</sub>. According to our definition, haze is determined 203 by visibility and Rhum. Considering two types of CSs, it was found that there is no significant difference in Rhum between the two kinds of CSs, which were reflected in the trend and difference between the 204 205 lower limit of blocking CS and the upper limit of wave-train CS after the outbreak of CS. However, the 206 blocking CSs are generally less effective in improving visibility than the wave-train CSs. When the 207 blocking CSs outbreak, the visibility shows an increasing trend; however, it begins to deteriorate 208 continuously 3 days later. Though the visibility in EC has a noticeable downward trend 5 days before the 209 outbreak of the wave-train CSs, it improves significantly on the day of the wave-train CSs outbreak and 210 rapidly deteriorates again about 3 days after the wave-train CSs occur. To further understand the 211 differences between the two types of CS, a closer investigation of additional meteorological parameters 212 was performed.



批注 [张28]: Reviewer#1 Specific comments (13)

批注 [张27]: Reviewer#1 Specific comments (12)

批注 [张29]: Reviewer#2 Comments (3)

214

215 before and after the outbreak of the blocking CSs (blue lines) and wave-train CSs (red lines), respectively. Shading

216 represents plus/minus one standard deviation among the CSs.

217

218 Previous studies show that haze is influenced by surface meteorological conditions (Wang et al.,

219 2015a; Yin et al., 2019a), which have significant variations after the outbreak of CSs (Park et al., 2014).

220 Figure 6 reveals the thermal inversion potential anomalies (TIP, defined as the air temperature at 850 hPa 10

221	minus SAT referring to Yin et al. (2019b)), surface horizontal wind speed (UV_sfc) anomalies, SAT
222	anomalies, and SLP anomalies for 4 days before and after the outbreak of the blocking CSs and wave-
223	train CSs. The results show that the high variations of meteorological elements reached the strongest on
224	the day of the CSs outbreak, and their anomalies weakened in the next 4 days. The variation of
225	meteorological elements during wave-train CSs is larger than during blocking CSs. 4 days after the
226	outbreak of the two types of CSs, the difference of TIP and UV_sfc between the two types of CSs tended
227	to be the same. However, the negative SAT anomalies and the positive SLP anomalies caused by the
228	wave-train CSs lasted longer than those caused by the blocking CSs. This is in line with the difference
229	in HD <sub>EC</sub> dispersion ability between the two types of CSs. Namely, the negative anomaly of temperature
230	and the positive anomaly of pressure change are not conducive to the maintenance of haze (Yin et al.,
231	2019a).
232	It should be noted that the SAT and SLP anomalies caused by the two types of CSs in this paper are
233	different from those of Park et al. (2014), who identified CSs in a different region, which included the
234	northern part of Northeast Asia. This is because the invasion of cold air is generally from north to south,
235	so their research covers more CSs in Northeast Asia, while the present study only focuses on CSs in

236 eastern China with heavy haze. Nevertheless, if we choose the same region, similar results can be

237 obtained (Figure S3).

批注 [张30]: Reviewer#1 Technical corrections (17)

批注[张31]: Reviewer#1 Technical corrections (18)



(b)UV sfo

(a)TIP

238 239

240

241

242 243

244

245

246

247 248 批注 [张32]: Reviewer#1 Technical corrections (14) and (19)

批注[张34]: Reviewer#1 Specific comments (14)

批注 [张33]: Reviewer#2 Comments (3)

批注 [张35]: Reviewer#1 Technical corrections (21)

blocking CSs maintained for a longer time and a larger control region in EC, which may cause the weak
 dispersion ability to HD<sub>EC</sub>. From the perspective of UV\_sfc, the cold air was limited to the north, and the

251 warm and humid conditions in EC were maintained before the outbreak of the CSs. Therefore, the CSs

252 cause the airflow with the northern wind component to invade EC, rapidly dispersing the haze and

- 253  $\,$  causing the visibility to rise. However, after the outbreak of the CSs, the anomalies of UV\_sfc in EC  $\,$
- 254 decrease, providing conducive conditions to the generation and maintenance of haze. The anomalies of
- $255 \qquad UV\_sfc \ in \ EC \ after \ the \ outbreak \ of \ blocking \ CSs \ are \ weaker \ and \ have \ a \ shorter \ duration \ than \ wave-train$
- 256 CSs.



**批注 [张36]:** Reviewer#1 Technical corrections (23)

257

Figure 7. Composite anomalies of TIP (shading; K; dotted areas are statistically significant at the 95% confidence
level) and UV\_sfc (vectors; m s<sup>-1</sup>) from day -2 to day 6 relative to the outbreak of blocking CSs (a, c, e, g, i) and the
wave-train CSs (b, d, f, h, j).

261

262 The increase of Siberian high accompanies the outbreak of CSs, and the splitting and southward movement of the Siberian high leads cold air into EC. Comparatively speaking, the distribution of SLP 263 264 anomalies in Eurasia before the blocking CSs form a pattern similar to the negative phase of the Arctic 265 oscillation. Figure 8 shows that when the blocking CSs occur, the positive SLP anomalies and the 266 negative SAT anomalies in the high-latitudes move southward. At the same time, the positive SLP 267 anomalies control EC. 2 days after the outbreak of the blocking CSs, the positive SLP anomalies, and the 268 negative SAT anomalies in the EC decline rapidly, providing favorable conditions for the accumulation 269 of pollutants. On the other hand, the occurrence of wave-train CSs is accompanied by the eastward 270 movement of significant positive SLP anomalies and negative SAT anomalies. 2 days after, the positive 271 SLP anomalies affect EC continuously, resulting in a longer period of high visibility in EC. In addition, 272 Rhum also has significant effects on the hygroscopic growth of particles, which will change the mass 273 concentration of aerosols and in turn the visibility (Wang et al., 2019). After the occurrence of the two 274 types of CSs, most regions of EC present negative Rhum anomalies. The negative Rhum anomalies 275 caused by wave-train CSs have a longer duration, stronger intensity, and wider influence range (Figure

276 S4). This shows that the wave-train CS has a stronger ability to dissipate haze.

批注 [张37]: Reviewer#2 Comments (4)



批注 [张38]: Reviewer#1 Specific comments (15) and technical corrections (23)





279 level) and SAT (contour; K) from day -2 to day 6 relative to the outbreaks of blocking CSs (a, c, e, g, i) and the

 $280 \qquad \text{wave-train CSs (b, d, f, h, j). The thick black isoline represents the 0 value of SAT anomalies.}$ 

281	The results discussed earlier indicate that the blocking CSs have a weak ability to dissipate the
282	HD <sub>EC</sub> , while the outbreak of wave-train CSs can make EC maintain high air quality for a longer time.
283	Thus, the frequency variations of the two types of CSs may also affect the trend of HD <sub>EC</sub> in recent years.
284	Figure 9a displays the time series of the frequency of blocking CSs and wave-train CSs. The results show
285	that the wave-train CSs have an obvious downward trend, and the blocking CSs have a slight upward
286	trend. It means that in recent years, the ability of CSs to dissipate $HD_{EC}$ has decreased in general. We
287	further calculated the partial correlation coefficients between the frequency of the two types of CS and
288	HD <sub>EC</sub> to exclude the influence of the other type of CSs. It is found that there is a significant positive
289	correlation between blocking CSs and HD <sub>EC</sub> (Figure 9b). It should be noted that this does not mean that
290	more blocking CSs cause more haze but reflects the weak dispersion ability of blocking CSs to $HD_{EC}$ ,
291	resulting in relatively more $HD_{EC}$ . The negative correlation between wave-train CSs and $HD_{EC}$ is
292	significant (Figure 9c), which is consistent with the result above.
293	In addition, we also evaluated the relationship between the trend of the total number of CSs (of both
294	types) and HD <sub>EC</sub> (Figure 9d). The results show that the correlation is weaker than that between a single
295	type of CSs and $HD_{EC}$ , which is the interference caused by the difference in the ability of the two types
296	of CSs to dissipate haze. In fact, the relatively more $HD_{EC}$ in the central EC caused by variations of total
297	CSs could be supported in a previous study (Yang et al., 2020a). The pattern of total CSs changes from
298	wave-train type to blocking type, especially after the mid-1990s.
299	Furthermore, previous studies also have shown that with the appearance of a warm Arctic-cold
300	Eurasian pattern more blocking high is expected to be maintained in the winter (Cohen et al., 2014; Luo
301	et al., 2016), causing the pattern of CS changed from wave-train type to blocking type (Yang et al., 2020b).
302	Therefore, it can be considered that the ability of CSs to dissipate haze in East Asia weakened in the
303	future is mainly due to the significant reduction of wave-train CS, and policymakers are required to
304	consider the problem of air pollutions.

批注 [张39]: Reviewer#1 Specific comments (17)

批注 [张40]: Reviewer#1 Technical corrections (24)

批注 [张41]: Reviewer#1 Specific comments (16)

批注 [张42]: Reviewer#1 Specific comments (1) and (18)



## 批注 [张43]: Reviewer#1 Technical corrections (27)

305

Figure 9. (a) Time series of the blocking CSs and the wave-train CSs (The solid line is the linear regression of the time series, and the text in the upper right corner indicates the trend of the solid lines). The partial correlation coefficient between HD<sub>EC</sub> and the frequency of blocking CSs (b) and the wave-train CSs (c), and the correlation coefficient between HD<sub>EC</sub> and the frequency of all CSs (d). Dotted areas are statistically significant at the 95% confidence level.

## 311 4. Conclusions

312 This paper investigates the connection between the CSs and the cool season haze over the EC based 313 on the observational and reanalysis datasets from 1980 to 2017. The 187 CSs over EC are classified into 314 two types by HCA, blocking CSs and wave-train CSs. Usually, the blocking CSs are accompanied by a 315 meridional dipole in the upper-tropospheric GPH anomalies, which consists of a stable blocking structure. 316 The blocking structure tends to control the EC for a long time and forms a relatively stable meteorological 317 condition, which has the disadvantage of dissipating the HD<sub>EC</sub>. Correspondingly, the local meteorological 318 conditions, especially TIP and the quiescent wind band, rapidly appear after the blocking CSs outbreak, 319 provide a haze-prone background. In addition, the positive SLP anomalies induced by the outbreak of

批注 [张44]: Reviewer#1 Specific comments (19) and (21)

320	the blocking	CSs can	rapidly	restore t	o normal.	and	the SAT	warm u	unde unde	r the	influence	of the	he
0-0			/										

- 321 weakening of the north wind component. Therefore, the ability of blocking CSs to dissipate HD<sub>EC</sub> is
- 322 limited. On the contrary, high air quality in EC can last longer due to the shorter duration of TIP and
- 323 longer duration of positive SLP anomalies after the wave-train CS. HD<sub>EC</sub> can generally remain at a low
- 324 level for a shorter (longer) time after the outbreak of blocking (wave-train) CSs. It is confirmed that
- 325 blocking CSs has been increased over the past few years (Park et al., 2011a; Luo et al., 2018).
- 326 Furthermore, the decreasing trend of wave-train CSs is likely to continue in the future, while the
- 327 frequency of blocking CSs is expected to remain stable, which may weaken the dispersion of haze and
- 328 worsen the HD<sub>EC</sub>.

## 329 Data availability

The ground observations are from the website: <u>http://data.cma.cn</u>. Daily mean meteorological data
 are obtained from the ERA-Interim reanalysis data archive: <u>http://www.ecmwf.int/en/research/climate-</u>
 <u>reanalysis/era-interim</u>.

## 333 Author contributions

- 334 SZ and GZ put forward the idea and design research, RW provided observational data including
- 335 relative humidity, visibility, and weather phenomena. SZ and XY performed research, and ZY provided
- 336 valuable suggestions. SZ wrote the manuscript with contributions from all co-authors.

#### 337 Competing interests

338 The authors declare that they have no conflict of interest.

## 339 Acknowledgements

- 340 This research is supported by the National Key Research and Development Program of China
- 341 (2017YFA0603804) and the National Natural Science Foundations of China (42175035; 42175034;
- 342 41575085).

批注 [张45]: Reviewer#1 Technical corrections (29)

批注[张46]: Reviewer#1 Specific comments (20)

## 343 References

- cold surges on air quality in Eastern Asia, Atmospheric Environment, 166, 498-509,
  https://doi.org/10.1016/j.atmosenv.2017.07.047, 2017.
- 347 Cai, B., Zeng, G., Zhang, G. and Li, Z.: Autumn cold surge paths over North China and the associated
- atmospheric circulation, Atmosphere, 10, 134, <u>https://doi.org/10.3390/atmos10030134</u>, 2019.
- 349 Cai, W. J., Li, K., Liao, H., Wang, H. J., and Wu, L. X.: Weather Conditions Conducive to Beijing Severe
- Haze More Frequent under Climate Change, Nat. Clim. Change, 7, 257–262,
  https://doi.org/10.1038/nclimate3249, 2017.
- 352 Compo, G. P., Kiladis, G. N., Webster, P. J.: The horizontal and vertical structure of East Asia winter
- monsoon pressure surges, QJR Meteorol. Soc, 125, 29–54, <u>https://doi.org/10.1002/qi.49712555304</u>,
  1999.
- Chai, D. H., Wu, M. H., Li, J. X., Zhao, Y. Q.: Analysis of greenhouse effect of cold wave low temperature
  on different structures in 2000 (in Chinese), Meteorological Science, 2002(03), 367-371, 2002.
- Chang, L. Y., Xu, J. M., Tie, X. X., and Wu, J. B.: Impact of the 2015 El Nino event on winter air quality
  in China, Sci. Rep., 6, 34275, <u>https://doi.org/10.1038/srep34275</u>, 2016.
- 359 Chen, H. P., Wang, H.J.: Haze Days in North China and the associated atmospheric circulations based on
- daily visibility data from 1960 to 2012, Journal of Geophysical Research: Atmospheres, 120(12), 5895-
- 361 5909, <u>https://doi.org/10.1002/2015JD023225</u>, 2015.
- Cohen, J., Screen, J., Furtado, J., et al.: Recent Arctic amplification and extreme mid-latitude weather,
  Nature Geoscience, 7, 627–637. <u>https://doi.org/10.1038/ngeo2234</u>, 2014.
- 364 Compo, G. P., Kiladis, G. N., and Webster, P. J.: The horizontal and vertical structure of east Asian winter
- 365 monsoon pressure surges, Q. J. R. Meteorol. Soc., 125, 29–54,
   366 <u>https://doi.org/10.1002/qj.49712555304</u>, 1999.
- 367 Dee, D., and Coauthors.: The ERA-interim reanalysis: configuration and performance of the data
- 368 assimilation system, Quarterly Journal of the Royal Meteorological Society, 137, 553-597,
- 369 <u>https://doi.org/10.1002/qj.828</u>, 2011.
- 370 Ding, Y. H., and Liu, Y. J.: Analysis of long-term variations of fog and haze in China in recent 50 years
- and their relations with atmospheric humidity, Sci. China Ser. D: Earth Sci., 57, 36-46,

## 372 https://doi.org/10.1007/s11430-013-4792-1, 2014.

- 373 Fu, C. B., Wu, J., Gao, Y. C., Zhao, D. M., Han, Z.W.: ConECutive extreme visibility events in China
- during1960–2009, Atmos. Environ., 68, 1–7, <u>https://doi.org/10.1016/j.atmosenv.2012.11.035</u>, 2013.
- 375 Han, S. Q., Bian, H., Tie, X., Xie, Y., Sun, M., Liu, A.: Impact measurements of nocturnal planetary
- boundary layer on urban air pollutants: From a 250-m tower over Tianjin, China. J. Hazard. Mater., 162,
  264–269,2009.
- $\label{eq:states} 378 \qquad \text{Hien, P.D., Loc, P.D., Dao, N.V.: Air pollution episodes associated with East Asian winter monsoons, Sci.}$
- 379 Total. Environ, 409, 5063–5068, <u>https://doi.org/10.1016/j.scitotenv.2011.08.049</u>, 2011.
- 380 Hu, Z.Z., Bengtsson, L., Arpe, K.: Impact of global warming on the Asian winter monsoon in a coupled
- 381 GCM, J. Geophys. Res., 05, 4607–4624, <u>https://doi.org/10.1029/1999JD901031</u>, 2000.
- 382 Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White,
- 383 G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, R., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak,
- 384 J., Mo, K. C., Ropelewski, C., Wang, J., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-year reanalysis
- 385 project, B.Am. Meteorol. Soc., 77, 437–471, <u>https://doi.org/10.1175/1520-</u>
   386 <u>0477(1996)077<0437:TNYRP>2.0.CO;2, 1996.</u>
- Lin, C. Y., Lung, S. C.C., Guo, H. R. et al.: Climate variability of cold surge and its impact on the air
  quality of Taiwan, Climatic Change 94, 457–471, <u>https://doi.org/10.1007/s10584-008-9495-9</u>, 2009.
- 389 Luo, D. H., Chen, X. D., Dai, A. G et al.: Changes in Atmospheric Blocking Circulations Linked with
- Winter Arctic Warming: A New Perspective, Journal of Climate, 31 (18), 7661-7678,
   https://doi.org/10.1175/JCLI-D-18-0040.1, 2018.
- 392 Luo, D. H., Xiao, Y. Q., Yao, Y., Dai, A. G. et al.: Impact of Ural Blocking on Winter Warm Arctic-Cold
- 393 Eurasian Anomalies. Part I: Blocking-Induced Amplification, Journal of Climate, 29 (11), 3925-3947,
- 394 <u>https://doi.org/10.1175/JCLI-D-15-0611.1</u>, 2016.
- 395 Luo, Y. F., Lu, D., Zhou, X. J.; Li, W. L.: Characteristics of the spatial distribution and yearly variation
- 396 of aerosol optical depth over China in last 30 years, J. Geophys. Res., 106(D13), 14,501–14,513,
   397 https://doi.org/10.1029/2001JD900030, 2001.
- 398 Niu, F., Li, Z. Q., Li, C., Lee, K.-H., Wang, M. Y.: Increase of wintertime fog in China: Potential impacts
- 399 of weakening of the eastern Asian monsoon circulation and increasing aerosol loading, J. Geophys. Res.,
- 400 115, D00K20, https://doi.org/10.1029/2009JD013484, 2010.
- 401 Park, T. W., Ho, C. H., Jeong, J. H., et al.: Different characteristics of cold day and cold surge frequency 21

402 over East Asia in a global warming situation, Journal of Geophysical Research: Atmospheres,

403 116(D12), D12118, <u>https://doi.org/10.1029/2010JD015369</u>, 2011a.

- 404 Park, T. W., Ho, C. H., Yang, S.: Relationship between the Arctic Oscillation and Cold Surges over East
  405 Asia, J Clim, 24(1), 68–83, https://doi.org/10.1175/2010jcli3529.1, 2011b.
- 406 Park, T. W., Jeong, J.H., Ho, C.H., et al.: Characteristics of atmospheric circulation associated with cold
- 407 surge occurrences in East Asia: A case study during 2005/06 winter, Adv. Atmos. Sci, 25, 791–804,
   408 <u>https://doi.org/10.1007/s00376-008-0791-0</u>, 2008.
- 409 Park, T. W, Ho, C. H, Jeong, J. H, et al.: A new dynamical index for classification of cold surge types

410 over East Asia, Climate dynamics, 45(9): 2469-2484, <u>https://doi.org/10.1007/s00382-015-2483-7</u>,

- 411 2015.
- 412 Park, T. W., Ho, C. H., Deng, Y.: A synoptic and dynamical characterization of wave-train and blocking
- 413 cold surge over East Asia, Climate dynamics 43, 753–770, <u>https://doi.org/10.1007/s00382-013-1817-</u>
  414 <u>6</u>, 2014.
- 415 Qian, Y., Leung, L. R., Ghan, S. J. Giorgi, F.: Regional climate effects of aerosols over China: Modeling
- 416 and observation, Tellus, 55B, 914–934, <u>https://doi.org/10.3402/tellusb.v55i4.16379</u>, 2003.
- 417 Qu, W., Wang, J., Zhang, X., Yang, Z., Gao, S.: Effect of cold wave on winter visibility over eastern
- 418 China, J. Geophys. Res, 120, 2394–2406, <u>https://doi.org/10.1002/2014JD021958</u>, 2015.
- Rokach, L., Maimon, O.: Clustering methods. Data mining and knowledge discovery handbook, Springer
  US, 321-352, 2005.
- 421 Rousseeuw, P.: Silhouettes: A graphical aid to the interpretation and validation of cluster analysis. Journal
- 422 of Computational and Applied Mathematics, 20, 5365, <u>https://doi.org/10.1016/0377-0427(87)90125-</u>
  423 7, 1987.
- 424 Shi, C., Zhang, H., Roth, M., Li, Z.: Impacts of urbanization on long-term fog variation in Anhui Province,
- 425 China. Atmos. Environ, 42, 8484–8492, <u>https://doi.org/10.1016/j.atmosenv.2008.08.002</u>, 2008.
- 426 Wang, H. J., and Chen, H. P.: Understanding the recent trend of haze pollution in eastern China: roles of
- 427 climate change, Atmos. Chem. Phys., 16, 4205–4211, <u>https://doi.org/10.5194/acp-16-4205-2016</u>, 2016.
- 428 Wang, H. J., Chen, H. P., Liu, J. P.: Arctic Sea Ice Decline Intensified Haze Pollution in Eastern China,
- 429 Atmospheric and Oceanic Science Letters, 8:1, 1-9, <u>https://doi.org/10.3878/AOSL20140081</u>, 2015.
- 430 Wang, X, Zhang, R, Tan, Y, et al.: Dominant synoptic patterns associated with the decay process of PM
- 431 2.5 pollution episodes around Beijing, Atmospheric Chemistry and Physics, 21(4): 2491-2508,

22

## 432 https://doi.org/10.5194/acp-21-2491-2021, 2021.

- 433 Wang, X, Zhang, R, Yu, W.: The Effects of PM 2.5 Concentrations and Relative Humidity on
- 434 Atmospheric Visibility in Beijing, Journal of Geophysical Research Atmospheres, 2019.
- 435 Wang, Z. S., Liu, X. D., Xie, X. N.: Effects of Strong East Asian Cold Surges on Improving the Air
- 436 Quality over Mainland China, Atmosphere, 7, 38, https://doi.org/10.3390/atmos7030038, 2016.
- 437 Wang, Z., and Ding, Y.: Climate change of the cold wave frequency of China in the last 53 years and the
- 438 possible reasons (in Chinese). Chin. J. Atmos. Sci., 30, 1068–1076, 2006.
- 439 Wei, Y., Li, J., Wang, Z., Chem, H., Wu, Q., Li, J., Wang, Y., and Wang, W.: Trends of surface PM2:5
- 440 over Beijing-Tianjin-Hebei in 2013-2015 and their causes: emission controls vs. meteorological
- 441 conditions, Atmos. Oceanic Sci. Lett., 10, 276–283, <u>https://doi.org/10.1080/16742834.2017.1315631</u>,
- 442 2017.
- 443 Wu, J., Fu, C. B., Zhang, L. Y., Tang, J. P.: Trends of visibility on sunny days in China in the recent 50
- 444 years, Atmos. Environ., 55, 339–342, <u>https://doi.org/10.1016/j.atmosenv.2012.03.037</u>, 2012.
- 445 Wu, J., Luo, J. G., Zhang, L. Y., Xia, L., Zhao, D. M., Tang, J. P.: Improvement of aerosol optical depth
- retrieval using visibility data in China during the past 50 years, J. Geophys. Res. Atmos., 119, 13,370–
  13,387, <u>https://doi.org/10.1002/2014JD021550</u>, 2014.
- 448 Xie, Y. B., Chen, J., and Li, W.: An assessment of PM2:5 related health risks and impaired values of
- 449 Beijing residents in a consecutive high-level exposure during heavy haze days, Environ. Sci., 35, 1–8,
- 450 2014.
- 451 Xu, P., Chen, Y. F., and Ye, X. J.: Haze, air pollution, and health in China, Lancet, 382, 2067,
  452 https://doi.org/10.1016/S0140-6736(13)62693-8, 2013.
- 453 Xu, Q.: Abrupt change of themid-summer climate in central east China by the influence of atmospheric
- 454 pollution, Atmos. Environ., 35, 5029–5040, https://doi.org/10.1016/S1352-2310(01)00315-6, 2001.
- 455 Yang, X. Y., Zeng, G., Zhang, G. W., Li, Z.X.: Interdecadal Variation of Winter Cold Surge Path in East
- 456 Asia and Its Relationship with Arctic Sea Ice, Journal of Climate, 33(11), 4907-4925,
- 457 <u>https://doi.org/10.1175/JCLI-D-19-0751.1</u>, 2020a.
- 458 Yang, X. Y., Zeng, G., Zhang, G. W., Vedaste, I., Xu, Y.: Future projections of winter cold surge paths
- 459 over East Asia from CMIP6 models, International Journal of Climatology, 1-16,
  460 https://doi.org/10.1002/joc.6797, 2020b.
- 461 Yin, Z, Zhang, Y, Wang, H, et al.: Evident PM 2.5 drops in the east of China due to the COVID-19 23

463	https://doi.org/10.5194/acp-21-1581-2021, 2021.	
464	Yin, Z. C., Wang, H. J.: Possible Relationship between the Chukchi Sea Ice in the Early Winter and the	
465	February Haze Pollution in the North China Plain., 32(16), 5179-5190, https://doi.org/10.1175/JCLI-	
466	<u>D-18-0634.1</u> , 2019b.	
467	Yin, Z. C., Wang, H. J.: Role of atmospheric circulations in haze pollution in December 2016., 17, 11673-	
468	11681, https://doi.org/10.5194/acp-17-11673-2017, 2017.	
469	Yin, Z.C., Li, Y. Y., Wang, H. J.: Response of early winter haze in the North China Plain to autumn	
470	Beaufort sea ice, Atmos. Chem. Phys., 19, 1439-1453, https://doi.org/10.5194/acp-19-1439-2019,	
471	2019a.	
472	Yin, Z.C., Wang, H.J., and Chen, H. P.: Understanding severe winter haze events in the North China	
473	Plain in 2014: Roles of climate anomalies, Atmos. Chem. Phys., 17, 1641-1651,	
474	https://doi.org/10.5194/acp-17-1641-2017, 2017.	
475	Zhang X. Y., Yin, Z. C., Wang, H. J., and Duan M. K.: Monthly Variations of Atmospheric Circulations	
476	Associated with Haze Pollution in the Yangtze River Delta and North China, Adv. Atmos. Sci., 38(4),	
477	569-580, https://doi.org/10.1007/s00376-020-0227-z, 2021.	
478	Zhang, L. Y., Lu, X. B., Yang, L. L., Ding, F., Zhu, Z. F., and Rui, D. M.: Study on the Effects of the	
479	Strong El Nino Event on Air Quality of Eastern China in Winter, The Administration and Technique of	

quarantine measures in February, Atmospheric Chemistry and Physics, 21(3): 1581-1592,

480 Environmental Monitoring, 28, 23–27, 2016.

462

- 481 Zhong, W. G., Yin, Z. C., Wang, H. J.: The relationship between anticyclonic anomalies in northeastern
- 482 Asia and severe haze in the Beijing-Tianjin-Hebei region, Atmos. Chem. Phys., 19, 5941-5957,
- 483 <u>https://doi.org/10.5194/acp-19-5941-2019</u>, 2019.

批注 [张47]: Cited references

批注 [张48]: Cited references