

Response to Referee #1

We would like to thank the reviewer #1 for taking the time to review this manuscript and provide valuable and constructive feedback that have improved the manuscript.

In this author comment all the points one-by-one raised by the reviewer are copied here and shown in bold text, along with the corresponding reply from the authors in plain text. We will go through our manuscript text and figures and try to shorten the paper where appropriate and we will submit a revised and restructured version of the manuscript when the second referee comment becomes available (for including all suggestions).

- 1. The Introduction takes a very long time to get to the point, and at the end we still don't have a clear statement of the problem. Is the paper to quantify emissions from Madrid, or from three landfills, or is this the same thing? Why should we care about Madrid? What is actually new here?**

Sorry for this vaguely-referred information. We will make this clearer in the introduction. We estimate emissions from waste disposal sites as stated in the title of the paper. These three landfills are so close to the Madrid metropolitan area that they can be considered as city sources, and they are the most significant (also the unique significant) CH₄ sources in Madrid area. Meanwhile, these together disposal sites are rather strong source as compared with the inventory in Madrid. The metropolitan cities are continuously growing due to population movements, industries, etc., and thus, more and more cities incorporate landfills (and other CH₄ potential sources) into their limits/influential areas. This study might be also interesting for different big/medium cities and the method can be applied there.

- 2. Section 2.1.2: give reference for the specific TROPOMI product that you are using. TROPOMI has a very low success rate (3% globally), is that an issue here? Or is Madrid sunny enough?**

(1) The specific TROPOMI products are from Lorente et al. (2021)*. We will add this to Section 2.1.2. according to the referee's comment.

(2) Yes, you are right that TROPOMI globally has a low success rate. This is more problematic at high latitudes and in winter. Southern Europe, like Madrid is pretty sunny and the weather during the field campaign was sunny as well.

The TROPOMI data with high quality (qa=1.0) used in this study are further collocated with the IASI data, covering the time period from November 2017 to October 2020. There are nearly 29,000 measurements. Although these observations are from many different days they can be used consistently for the local emission estimates, because we remove the daily varying background signal. This background removal is very important and allows for using data from many different days for achieving good coverage. The multi-year data set provides a large and consistent observational dataset (horizontal: for the whole area around Madrid; temporal: for different wind regimes) for studying the local emissions.

*Lorente, A., Borsdorff, T., Butz, A., Hasekamp, O., aan de Brugh, J., Schneider, A., Wu, L., Hase, F., Kivi, R., Wunch, D., 740 Pollard, D. F., Shiomi, K., Deutscher, N. M., Velasco, V. A., Roehl, C. M., Wennberg, P. O., Warneke, T., and Landgraf, J.: Methane retrieved from TROPOMI: improvement of the data product and validation of the first 2 years of measurements, *Atmos. Meas. Tech.*, 14, 665–684, <https://doi.org/10.5194/amt-14-665-2021>, 2021.


3. Section 2.1.3: this combined IASI+TROPOMI TXCH₄ product is probably totally dominated by TROPOMI information in the PBL, which is what matters here. So how is it independent from TROPOMI?

TROPOMI measures column integrated methane (XCH₄). XCH₄ is affected by tropospheric CH₄ concentrations but also strongly by the altitude of the tropopause (a high tropopause causes low stratospheric contributions to XCH₄ and thus high XCH₄ values, a low tropopause vice versa). One possibility to avoid this contribution that affects the study of lower tropospheric CH₄ emissions is to remove the background and work only with anomalies. If one assumes that the background captures all the tropospheric background and in addition the stratospheric contribution, the anomaly contains the interesting signal. However, if the background calculation misses some of the stratospheric contribution signals, the interpretation using the anomaly for investigating local emission can lead to large errors. For instance, it might be that winds from the north-east and south-west are somehow correlated with tropopause altitudes (because both wind directions and tropopause altitudes have a seasonal cycle) and if this seasonal cycle is not well resolved in the background, the wind-assigned anomaly can have an artificial signal that comes from the stratosphere.

IASI has good information about the stratospheric contribution and combining IASI with TROPOMI allows for the generation of a product (TXCH₄) that is largely unaffected by the stratospheric contribution. Consequently, the risk for the TXCH₄ anomalies to be affected by the stratospheric contributions is much lower than for XCH₄. We calculate the emission for XCH₄ and for TXCH₄ and get similar emission rates for both data sets. This means our background correction is working correctly already for XCH₄. Furthermore, we repeat the calculations for the upper tropospheric and stratospheric CH₄ (UTSXCH₄, information from the IASI data) and found that there is no emission signal. All these prove that the emission signals we observe are not by accident, instead they come from local surface emissions and that our method for background calculation together with the wind-assigned anomaly method is able to detect these signals correctly in the XCH₄ as well as in the TXCH₄ data.

4. Figure 1 is very difficult to read.

Thanks for the comment. The figure is redone as below:



5. Section 2.3: this section is very confusing because it is not clear what the authors are trying to optimize. Are the ‘daily plumes’ for the individual landfills? Are they summed over the three landfills? Are the three landfills treated as a single plume? The cone model is surely wrong for instantaneous plumes but is reasonable for time-averaged plumes, which is what is fitted but it takes the paper a while to explain this.

Each individual landfill is considered as an individual point source. The daily plumes from the individual landfills are super-positioned to have a total daily plume. The final results we provide are emission rates averaged for the whole three years.

6. Line 206: I don’t get the point about seeking an analogy with NO₂. The landfills don’t emit NO_x, NO_x is an area source, and the decay of the NO₂ plume is by oxidation rather than dilution into background.

Yes, you are right that no NO_x is emitted from the landfills (while the Madrid metropolitan area is a strong source of NO₂). The usage of NO₂ in this study is to check if our method is reliable. The NO₂ is exploited as a tracer for atmospheric transport, offering sufficient chemical lifetime for forming a nice plume structure, which is a simple target gas for TROPOMI. NO₂ is a suitable (approximately stable) tracer for qualitatively demonstrating the method developed for the wind-assigned anomaly. We do not correctly consider the photochemical loss of NO₂, and the demonstration is not intended to provide a high-quality quantitative analysis of the NO₂ source strength (a refined model including NO₂ decay would generate slightly reduced outer plume lobes. The lifetime of NO_x should be in the order between 6.3h during night and 29h during daytime in winter (Kenagy et al, 2018) and 5.9h in summer (Shah et al., 2020)). Our intention is simply to check the implementation of our approach and the data we feed into the simulation: if our wind data or plume dispersion modelling would be incorrect, we would not be able to reasonably reproduce the plume properly in our model runs.


* Kenagy, H. S., Sparks, T. L., Ebben, C. J., Wooldridge, P. J., Lopez-Hilfiker, F. D., Lee, B. H., Thornton, J. A., McDuffie, E. E., Fibiger, D. L., Brown, S. S., Montzka, D. D., Weinheimer, A. J., Schroder, J. C., Campuzano-

Jost, P., Day, D. A., Jimenez, J. L., Dibb, J. E., Campos, T., Shah, V., Jaeglé, L. and Cohen, R. C.: NO_x Lifetime and NO_y Partitioning During WINTER, *J. Geophys. Res. Atmos.*, 123(17), 9813–9827, doi:<https://doi.org/10.1029/2018JD028736>, 2018.

Shah, V., Jacob, D. J., Li, K., Silvern, R. F., Zhai, S., Liu, M., Lin, J., and Zhang, Q.: Effect of changing NO_x lifetime on the seasonality and long-term trends of satellite-observed tropospheric NO₂ columns over China, *Atmos. Chem. Phys.*, 20, 1483–1495, <https://doi.org/10.5194/acp-20-1483-2020>, 2020.

7. Wind speed is denoted v in the text, w in Figure 2.

Many thanks to point out this mistake. We corrected it.



8. Equation (9): not clear how you get y_{BG}

For clarification, Eq. 9 y_{BG} should be replaced by y (the original satellite data $y = y_{BG} + y_{plume}$, see Eq. 7). We use y to estimate the coefficients that describe the background. Information from y is only used when the observations are not affected by the plume. In that case \mathbf{K}_{BG}^* and thus \mathbf{G}_{BG} are zero. On the other hand, \mathbf{K}_{BG}^* (and thus \mathbf{G}_{BG}) is set to zero whenever $y_{plume} = 0$. Then $y_{BG} = y$ and Eq. 9 is correct as it. But in order to make it clearer we will write in Eq. 9 y instead of y_{BG} .

9. Section 2.3: there are many uncertainties in the procedure for inferring emissions. How can it be validated? An obvious way would be to use the independent COCCON observations to evaluate the posterior concentrations resulting from the TROPOMI inversion.

Estimated emission rates can include large uncertainties. In this study, the uncertainty of the estimated emission rates derived from TROPOMI XCH₄ ($7.4 \times 10^{25} \pm 6.4 \times 10^{24}$ molec s⁻¹) based on our method is about 9% ($6.4 \times 10^{24} / 7.4 \times 10^{25}$). Please note that this is only the uncertainty due to the background uncertainty. Figure 10 shows sensitivity analysis due to wind, emission source, and opening angle, and reveals that there are other important uncertainty sources.

The referee gives a very good suggestion that COCCON is an independent data to evaluate the results. We applied this strategy and tried to estimate the emission rate from COCCON measurements on October 4 (Section 3.1), when a significant enhancement was observed by the downwind-side COCCON SN69. The estimated rate is about 3.7×10^{25} molec s⁻¹. This value is about half of that derived from TROPOMI XCH₄ or combined TXCH₄, which is a plausible match, because the satellite covers the complete area, whereas the COCCON plume observation primarily detects the emission from a single nearby landfill.

10. Figure 3: the agreement between TROPOMI and COCCON in that Figure strikes me as very poor, despite the authors' claim to the contrary. I'm not surprised by this in view of the known TROPOMI biases, but it undermines confidence in the results of the TROPOMI inversion. The paper goes on about the problems on Sept 25 and Oct 4 but that seems anecdotal and those two days don't seem any worse than the rest of the population in Figure 3.

We do not recognize an apparent bias between COCCON and TROPOMI (in Fig 3 we see some scatter, but the whole ensemble follows the 1:1 line quite well). Larger variations of XCH₄ in a metropolitan area containing localized sources as Madrid are to be expected (which will induce some scatter between the datasets, because the spatial resolution of the space-based sensor is much lower). Studies under background conditions have revealed very good agreement and low bias between COCCON and TROPOMI (Tu et al., 2020*). Moreover, it needs to be emphasized that our conclusions on emission strengths depend on averaged values of observed *gradients* in each dataset, so a general bias due to a calibration mismatch between satellite and ground-based would remain largely without effect.

Tu, Q., Hase, F., Blumenstock, T., Kivi, R., Heikkinen, P., Sha, M. K., Raffalski, U., Landgraf, J., Lorente, A., Borsdorff, T., Chen, H., Dietrich, F., and Chen, J.: Intercomparison of atmospheric CO₂ and CH₄ abundances on regional scales in boreal areas using Copernicus Atmosphere Monitoring Service (CAMS) analysis, Collaborative Carbon Column Observing Network (COCCON) spectrometers, and Sentinel-5 Precursor satellite observations, Atmos. Meas. Tech., 13, 4751–4771, <https://doi.org/10.5194/amt-13-4751-2020>, 2020.

11. Lines 329-330: how do we know that the ‘COCCON instruments show a very good ability to detect the source’? No specific results or data from COCCON are shown.

The COCCON SN69 was located in the northwest of the landfill Valdemingómez with a quite close distance (4.5 km). When the wind came from southeast, the COCCON SN69 was located downwind of the landfill and detected a significant plume (nearly up to 100 ppb on October 4, 2018, Figure 4), whereas the other COCCON sites did not observe any enhancements. Another example is October 1, 2018 (Figure A-2): the wind direction was north to northeast and the COCCON stations were not on the downwind side of the landfills, which resulted in no enhancement at any of the COCCON stations. The observations on different days were largely depended on the wind situation. The obvious downwind enhancement observed by the COCCON instruments demonstrates that they have the ability to detect the emissions of the source, which has been demonstrated in Kille et al. (2019*) as well.

* Kille, N., Chiu, R., Frey, M., Hase, F., Sha, M. K., Blumenstock, T., Hannigan, J. W., Orphal, J., Bon, D. and Volkamer, R.: Separation of Methane Emissions From Agricultural and Natural Gas Sources in the Colorado Front Range, Geophys. Res. Lett., 46(7), 3990–3998, doi:<https://doi.org/10.1029/2019GL082132>, 2019.

12. Figure 8 is cryptic. What domain is shown? What are we learning from it?

We have tried to visualize as good as possible the different steps of the data treatment. Figure 7 shows the time series of the different satellite data (y from Eq. 7), their estimated background (y_{BG} from Eq. 12), and the anomaly signal due to the local emissions (y_{plume} from Eq. 13). Then Figure 8 shows the anomalies (i.e. y_{plume}) horizontally averaged for different wind directions. This data is used for the Δ -calculations. Eq.15 captures both: the horizontal averaging according to the wind directions as well as the Δ calculations. The results of these calculations are then shown in Fig. 9. These ΔCH_4 data are finally used to estimate the emission rates according to Eq. 18.

Furthermore, Figure 8 is very useful here, because it demonstrates that the CH_4 hotspots are south-east of Madrid and not in the center of the city. South-east of Madrid is where the waste disposal sites are located.

13. Table 5: I don't see the relevance of this Table to the paper.

This Table shows some results from other studies as a reference for our results. It helps to demonstrate that our results are reliable and lie in the reasonable range. The inventory only lists the active landfill cells and does not include the closed ones, which probably still emit for many years (Sánchez et al., 2019*). This is an additional argument for the relevance of the kind of work we are presenting here.

*Sánchez, C., de la Fuente, M. del M., Narros, A., del Peso, I. and Rodríguez, E.: Comparison of modeling with empirical calculation of diffuse and fugitive methane emissions in a Spanish landfill, J. Air Waste Manage. Assoc., 69(3), 362–372, doi:10.1080/10962247.2018.1541029, 2019.

Response to Referee #2

We would like to thank the reviewer #2 for taking the time to review this manuscript and provide valuable and constructive feedback that have improved the manuscript.

In this author comment all the points raised by the reviewer are copied here one-by-one and shown in bold text, along with the corresponding reply from the authors in plain text.

- 1. The Introduction gives extensively credit on former works and is therefore also an important part, which should not be missing even it might not help the reader in understanding of the new technics.**

Therefore, I would ask for a kind of road map at the beginning of second section “method”, before 2.1 which gives an idea, why and how the measurements and models are combined, or maybe it could be included directly at the end of introduction where the different sections are already mentioned.

Here, we add two more sentences to describe our methodology in the last paragraph of the introduction:

“Section 2 describes our methodology. We calculate the difference of the satellite data maps for two opposite wind regimes (we refer to the resulting signals as wind-assigned anomalies). A simple plume model is applied to predict the wind-assigned anomalies for a chosen position and strength of a source. The results of our study are presented and discussed in Section 3.....”

2. TROPOMI+IASI

- 2.1 You report two estimations in the abstract, which are slightly contra dictionary. The error in the estimation is higher than the error estimation from the combined retrieval, I imagine that this is more or less the effect that partial columns are constructed with more errors (Error of TROPOMI + error of IASI), but there should be another systematic error, which is reduced if the tropospheric partial column anomaly is sub or over estimated. It would be easier for us readers, if the authors conclude on a single best estimation, and report the other option using just TROPOMI just in the results and discussion of the article.**

As stated correctly by the referee, the stratospheric contribution causes larger uncertainties when using the XCH₄ data. However, it is very difficult to estimate this error. In the paper, we do not estimate the uncertainty of the stratosphere contribution. Instead, we use combined data (IASI+TROPOMI, where the stratospheric contribution can be neglected) to show that our background removal method works well and removes the stratospheric effects, also in the XCH₄ data.

2.2 The combination of IASI and TROPOMI assumes measurements of the same air mass, and therefore at the same time, but IASI comes around 10:00 and TROPOMI around 13:00, does this matter here, o IASI characterize more the background CH₄ and is therefore less critical. Maybe it would be interested in a discussion how the time difference between IASI and TROPOMI of some hours, might affect the combined retrieval.

Please also describe a bit more how much the estimation improves due the combination with IASI, I could imagine, that this advantage will increase if you use long term averages, so that the random error loses its importance in relation to improved sensitivity in the lower atmosphere.

In Schneider et al. (2021b) we found that a temporal mismatch of up to four hours is no problem. The noise error is larger than the error introduced by combining two measurements that have a temporal mismatch of four hours. The mismatch error just increases slightly the statistical error budget.

Yes, the noise error is larger in the combined product (see Schneider et al., 2021b). And this error will get smaller, when more data are used for averaging. So, if the emission rate keeps constant over a long time, the estimation of these emission rate will become the better, the longer the time series is (the more data are used for averaging). However, in practice the emission rates might not be fully constant over time.

3. Plume model

3.1 The authors realize their proper idea and do not have to cite other works, which might be partly similar ideas, but it would help some reader to understand it easier, which is similar to other approaches (up, downwind Rotating method, Gaussian's Law).


The comment is very helpful to make reader better understand of our simple model. We try to explain our model more:

Our plume model is a simplified version of the Gaussian plume model (https://link.springer.com/chapter/10.1007/978-1-4757-4465-1_7, Figure 7-1). Our simple model treats a two-dimensional concentration field (without height being involved) generated by a point source (or an array of point sources) and the gas molecules in the downwind side are evenly distributed into a sector (not a Gaussian distribution) centered along the wind direction. The opening angle of the sector is the only free model parameter, which is adjusted in order to reasonably reproduce the observe NO₂ plume.

3.2 Do I understand correctly the point, that the plume model is retrieved using the City centre of Madrid as central source, but applied to the source located at a different location (landfills). So orography should be similar or not be important. I would have been interested in the required conditions on the regional topography for this approach to be transfer also to other cities or waste deposal sites. And I also would like to know the typical mixing layer height in the area around Madrid.


For NO₂ we use the Madrid city center as a central source for the plume model. However, we use three appropriately located point sources in case of CH₄. The a priori information about the CH₄ emissions from the landfills (Table 3) is taken from the Spanish Register of Emissions and Pollutant Sources. Each individual landfill is considered as a point source. The contributions from the individual landfills are super-positioned to generate a total daily plume.

The regional topography from Google map and the altitude derived from TROPOMI are presented as below. The mountain ranges locate along the NE-SW direction, which forces the wind flowing along this direction. We, however, do expect similar wind conditions in the city center and the landfills located nearby, as the topography is rather flat in this sub-region.



Our Spanish colleagues kindly compiled information on the Planetary Boundary Layer (PBL) height: the height of the PBL in Madrid, and its seasonal evolution, is the typical from continental areas. Figure below shows the averaged seasonal cycle of the height of PBL using meteorological radiosondes launched twice a day from the Madrid-Barajas station (WMO #08221) in the period 1981-2015. It is calculated using the Heffter method (Heffter JL. 1980. “Transport Layer Depth Calculations.” Second Joint Conference on Applications of Air Pollution Meteorology, New Orleans, Louisiana). Note that the radiosonde launch station is the same used in the MEGEI-MAD campaign.

So, the noon PBL reaches typically altitudes of about 1300 m in wintertime and up to 2500 m in summertime. We discuss possible corrections due to increasing wind speed with altitude on our emission estimates further below.



3.3 The test using NO₂ from TROPOMI results in a lower emission rate compared to estimation from the literature. The authors identify the lifetime as the reason and maybe also the value in the literature might have errors, or might not be comparable, different timeHowever a validation of the wind speed estimation from a 10 m altitude would be nice.

For CH₄ and its long lifetime the anomaly in the column should be definitively given by 1/d, as described by the equation 3.

The forward model contains the velocity and actually the quotient emission/velocity determine the concentration, or vice versa in the inversion. That is why the wind speed estimation might be so important. The sensitivity study using the wind at 10 m is very helpful and important, never the less I did not get completely get the point how you decided that 10 m is the best. I think the wind velocity you look for is the velocity you can multiply with the column anomaly to get the total CH₄ flux. Ad hoc I would take the average windspeed in the atmospheric boundary layer.

The COCCON sites at “Jose Echegaray” and “Barajas” have a distance of more or less around 8km, on 25 September 2018 (Figure 4) we see that anomalies in CH₄ arrives around 1-2 hours later in Barajas than it appears in “Jose Echegaray”.

So the velocity of propagation of this plume might be 8 km/2 h to 8 km/h or 1.11m/s to 2.2 m/s respectively, this fits actually very well to the assumed and modelled wind velocity in the plot below. Surely that is the intention showing the plot, but maybe it should be also explicitly be mentioned, that you can also use the 5 FTIR sites to validate the wind estimation strategy you have chosen, as here the interest in the effective flux of CH₄ and it might be the best validate the velocity using the FTIR EM27 measurements of columnar CH₄, so that you are independent of the vertical distribution of CH₄.

EM27 measurements take typically a little bit less than a 1 minute, maybe you could do a cross correlation to retrieve the delay after interpolate data to a common 1 minute grid, and then include a point in the figure showing the wind speed or projection, but it is sufficient just to mention it.

We thank the referee for the careful consideration concerning the wind. We fully agree that the limited quality of the available wind data is a significant source of uncertainty.

According to WMO (WMO, 2018), the measurement representative of the surface winds is the wind records at 10 m a.g.l. to avoid the roughness of surrounding terrain. In this sense, the winds at 10 m are usually taken as a proxy for surface emission estimations (e.g. Viatte et al., 2017), such as in the Madrid case. In addition, we chose ERA5 wind at 10 m, because it can be directly compared to the in situ wind observations at 10m at the Cuatro Vientos Airport, which helps to estimate the uncertainty introduced by the wind data.

The wind at ~500 m a.g.l (900 hPa) would be a more appropriate choice for transport modelling if we allow for vertical mixing of the plume within the PBL. As shown in the Table 1 and Figure 1 below, there is a significant increase of wind speed with altitude. The ERA5 wind data at 10 m and ~500 m do not differ significantly concerning the wind direction, but the wind velocity increases with altitude. The wind speed increases by 60%, i.e. using ERA5 wind information at ~500 m instead of that at 10 m would yield 60% increase in the emission rate.

Table 1: ERA5 wind at 10 m and 100 m in TROPOMI overpass days.

Wind direction range	10 m		~500 m/900 hPa	
	Number of days in total (%)	Averaged wind speed \pm standard deviation (m s^{-1})	Number of days in total (%)	Averaged wind speed \pm standard deviation (m s^{-1})
NE / $>315^\circ$ or $<135^\circ$	28.4	2.3 ± 1.2	30.9	3.8 ± 2.1
SW / $135^\circ - 315^\circ$	61.8	2.3 ± 1.4	56.7	3.6 ± 2.0




Figure 1: Wind roses for daytime (08:00 UTC – 19:00 UTC) for the ERA5 model wind at 10 m (left) and at ~500 m (900 hPa) (right).

It is really a good hint to mention that we should use the array of COCCON stations itself to validate the assumptions on the wind field. We will add this statement to our manuscript:

“These five COCCON stations can serve as an independent source of information for constraining the wind speed. For example, the distance between the Jose Echegaray and Barajas is about 10 km. The highest anomalies of XCH₄ arrived around 1.5 hours later at Barajas station than it appeared at the Jose Echegaray station on 25 September 2018, which indicates an averaged wind speed of 1.8 m/s. This value fits well to the ERA5 model wind velocity.”

WMO, Guide to Instruments and Methods of Observation Volume I –Measurement of Meteorological Variables, Report No. 8, ISBN 978-92-63-10008-5, Geneva, Switzerland, 2018.

Viatte, C., Lauvaux, T., Hedelius, J. K., Parker, H., Chen, J., Jones, T., Franklin, J. E., Deng, A. J., Gaudet, B., Verhulst, K., Duren, R., Wunch, D., Roehl, C., Dubey, M. K., Wofsy, S., and Wennberg, P. O.: Methane emissions from dairies in the Los Angeles Basin, Atmos. Chem. Phys., 17, 7509–7528, <https://doi.org/10.5194/acp-17-7509-2017>, 2017.

4. Page 2, Line 37:

4.1 “The wind-assigned plume method is also applied to the tropospheric and upper tropospheric/stratospheric column averaged CH₄ mixing ratio products (in the following referred to as TXCH₄ and UTSXCH₄) derived from a-posteriori merged Infrared Atmospheric Sounding Interferometer (IASI) profile and TROPOMI total column data.” Maybe you could split the sentence into 2 and find somehow a different description, which is easy to understand:

For the CH₄ emission estimation, the wind-assigned plume method is applied to the lower tropospheric methane /dry air column ratio (TXCH) of the combined TROPOMI Infrared Atmospheric Sounding Interferometer (IASI) Product. TXCH₄ and the upper tropospheric/stratospheric column averaged CH₄ mixing ratio (UTSXCH₄) are derived from a-posteriori merged Infrared Atmospheric Sounding Interferometer (IASI) profile and the TROPOMI total column data.

We modify the sentence according to referee’s comments:

“For the CH₄ emission estimation, the wind-assigned plume method is applied to the lower tropospheric CH₄/dry air column ratio (TXCH₄) of the combined TROPOMI + IASI (Infrared Atmospheric Sounding Interferometer) product. TXCH₄ and the upper tropospheric/stratospheric column averaged CH₄ mixing ratio (UTSXCH₄) are derived from a-posteriori merged IASI profile and the TROPOMI total column data.”

4.2 Does it make sense to apply the method to the UTSXCH₄ Product? As mentioned earlier I would just use TXCH₄ in the abstract and conclusion.

The very low emission rate derived from the UTSXCH₄ demonstrates that all is working well: from TXCH₄ we get the same as for XCH₄ (if background is carefully removed). And in UTSXCH₄ we see no emission signal (i.e. the background removal and the calculation of wind-assigned anomalies introduce no artificial signals).

5. **Line 40: The first sentence is a bit redundant, maybe you could drop here the sentence “Based on the NE and SW wind fields, we developed a simple plume model locating the source at three waste disposal sites east of Madrid for CH₄.” or move it to line 37.**

We remove this sentence according to the referee’s comment.

6. **Line 44: “day and. All em...**

Maybe you could rewrite the sentence which is also not clear.” The COCCON observations indicate a weaker CH₄ emission strength of around 3.7×10^{25} molec s⁻¹ from local source (near to the Valdemingómez waste plant) in accordance with observations in a single day and. “Please write very clear if COCCON is used to estimate an independent “local source”, if this source is part and included in the Tropomi estimation given quantitatively in the line above. Please state if TROPOMI and COCCON based estimations are contradiction, complementary and/or consistent. If you want, you might move the last sentence to line 25, as it is finally the most important finding.

The comment helps us to explain the results clearer. We change this sentence to:

“COCCON observations are investigated to estimate the local source as an independent method. The COCCON observations indicate a weaker CH₄ emission strength of around 3.7×10^{25} molec s⁻¹ from local source (near to the Valdemingómez waste plant) in accordance with observations in a single day. That this figure is lower than the one derived from the satellite observations is a plausible result, because the analysis of the satellite data refers to a larger area, covering further emission sources in the study region, whereas the signal observed by COCCON is generated by a nearby local source.”

7. **Figure 8,9:**

Please use a fixed color scale in Fig 9 and maybe also in Fig 8. The modelled XCH₄ for total column, lower and upper troposphere seems to be exactly the same in fig 9 b,e,h. Maybe when you talk about emission rates line 474 and line 477 you cloud use CH₄ and not the dry air mol fraction.

XCH₄-> CH₄

To remove unwanted signals due to validations of ground pressure and atmospheric humidity we estimated the emission rates from total and partial column averaged dry air mixing ratios and not directly from total column amounts. The color scales in Figure 8 and 9 are chosen according to the typical dry air columns. The dry air total column is typically by a factor of 1.9 larger than the dry air tropospheric partial column. And it is typically by a factor of 2.1 larger than the dry air upper tropospheric/stratospheric partial column. These factors explain the different color scales.