



Global Evidence of Aerosol-induced Invigoration in Marine Cumulus Cloud

Alyson Douglas¹ and Tristan L'Ecuyer²

¹Atmospheric, Oceanic, and Planetary Physics Department, University of Oxford, Sherrington Rd, Oxford OX1 3PU

²Atmospheric and Oceanic Sciences Department, University of Wisconsin-Madison 1225 W Dayton St, Madison, WI 53706

Correspondence: Alyson Douglas (alyson.douglas@physics.ox.ac.uk)

Abstract. Aerosol-cloud-precipitation interactions can lead to a myriad of responses within shallow cumulus clouds including an invigoration response, whereby aerosol loading results in a higher rain rate, more turbulence, and deepening of the cloud layer. However few global studies have found direct evidence that invigoration occurs. The few satellite based studies that report evidence for such effects generally focus on only the deepening response. Here, we show evidence of invigoration beyond a deepening response. Using latent heating and vertical motion profiles derived from CloudSat radar observations, we show precipitating cumulus clouds in unstable, polluted environments exhibit a marked increase in precipitation formation rates and cloud top entrainment rates. However, invigoration is only discernible when the stability of the boundary layer is explicitly accounted for in the analysis. Without this environmental constraint, the mean polluted and pristine cloud responses are indistinguishable from each other due to offsetting cloud responses in stable and unstable environments. Invigoration, or suppression depending on the environment, may induce possible feedbacks in both stable and unstable conditions that could subdue or enhance these effects, respectively. The strength of the invigoration response is found to additionally depend on cloud organization defined here by the size of the warm rain system. These results suggest that warm cloud parameterizations must account for not only the possibility of aerosol-induced cloud invigoration, but also the dependence of this invigorated state on the environment and the organization of the rain system.

1 Introduction

Aerosol-cloud interactions remain one of the largest sources of uncertainty in future climate projections (Boucher et al., 2013). Further, their role in climate feedbacks, particularly how they affect low clouds, controls the magnitude of the climate sensitivity (Zelinka et al.). However, despite the importance of tropical low clouds to the global climate, understanding their response to anthropogenic activity, including aerosol loading, remains a challenge (Bony and Dufresne, 2005). In particular, invigoration, or the enhanced size, depth, precipitation rate, or turbulence, of low clouds was hypothesized as a potential outcome of aerosol-cloud interactions decades ago but remains relatively unconfirmed from observations (Pincus and Baker, 1994; Rosenfeld et al., 2008). If invigoration of warm cloud precipitation occurs, it not only affects where and how much clouds precipitate, but the entire hydrological cycle (Li et al., 2011). Invigoration of warm cloud structure also has the potential to alter deep convection, making eventual storms more intense and turbulent (Chen et al., 2017).



25 Previous studies focused on detecting warm cloud deepening as a signal of invigoration, as it theoretically implies increased
turbulence and precipitation within the cloud (Altaratz et al., 2014). L'Ecuyer et al. (2009) showed warm, polluted precipitating
clouds grow deeper than those in more pristine environments using space-borne radar observations from CloudSat. Christensen
and Stephens (2011) similarly found as ships passed below marine stratocumulus, the locally affected clouds, as identified using
a combination of radar and passive satellites, deepened. Yuan et al. (2011) found evidence of cloud deepening in trade cumulus
30 when interacting with nearby volcanic emissions while Kubar et al. (2009) found evidence of cloud deepening in highly
polluted environments in all warm cloud types. On the other hand, Dey et al. (2012) found no evidence of cloud deepening
in the smallest clouds studied, only an increase in extent. Never-the-less, while numerous observational studies have been
able to discern a cloud depth response using both passive and active sensors, few have controlled for the environment in their
estimates in order to certify that this response is due to aerosol-forced invigoration and not a confounding environmental signal.
35 Additionally, a deepening cloud does not conclusively establish the physical processes associated with invigoration, such as
increased turbulence and precipitation formation rate, only that another cloud adjustment process occurs.

Modeling efforts have proven more promising in, at least hypothetically, demonstrating invigoration of warm clouds is
possible by aerosol. Heiblum et al. (2019) used a LES model to show that clouds formed in higher aerosol environments
release more latent heat and promote a larger rain cell size. Jiang et al. (2009) similarly used a LES and found clouds in
40 polluted environments produced more evaporation at the cloud edge in simulated trade cumuli, producing more vertical motion.
(Spill et al., 2019) found shallow convective clouds in high aerosol loading environments are more likely to deepen with a
variable response of the rain rate. Clouds formed in polluted environments may experience an increase in droplet mobility,
which delays collision coalescence and changes the organization of liquid water within the cloud to a more invigorated state
reaffirming Albrecht's original theory of a second aerosol indirect effect (Koren et al., 2015; Albrecht, 1989; Berg et al., 2008).
45 (Seifert et al., 2015) saw a decrease in cloud lifetime with increasing lifetime due to desiccation of warm cloud cover from
precipitation induced environmental feedbacks. Depending on the environmental conditions, the liquid water path of the cloud
may decrease, signaling a curtailment, not invigoration, response (Jiang et al., 2006).

The environment plays a strong role in modulating warm rain processes and therefore must be considered when using
observations to imply aerosol-forced invigoration of warm clouds (Stevens and Feingold, 2009). Prior work has shown that
50 the environment controls the amount of suppression of precipitation within the cloud, which may modulate the amount of
invigoration (L'Ecuyer et al., 2009). The strength of the marine boundary layer inversion controls cloud top height in many
warm clouds (Wood, 2012) and has been shown to heavily influence rain formation rates in warm clouds (Nelson and L'Ecuyer,
2018). The magnitude and sign of warm cloud aerosol-cloud interactions is likewise heavily modulated by both the inversion
strength and free atmospheric relative humidity (Douglas and L'Ecuyer, 2019). The humidity of the free atmosphere affects
55 how aerosol impacts the distribution of liquid water throughout the cloud layer due to entrainment processes (Ackerman et al.,
2004). Both are considered within this study in order to constrain these confounding factors.

To a first order, the liquid water path controls the probability of a cloud raining (L'Ecuyer et al., 2009; Berg et al., 2006).
Aerosols, in turn, impacts the liquid water path as part of a cloud adjustment process, which then further alters the probability of
precipitation. The relationship between aerosol-cloud interactions and cloud liquid water are neither universal nor well known.



60 In order to reduce the uncertainty interpreting our results, we limit our observations to clouds with liquid water paths in a narrow range between 150 to 200 g/m^{-2} . In doing so, we focus only on how aerosol alters the organization of rain formation and evaporation within the cloud layer, not its influence on cloud liquid water. Invigoration in this context includes how aerosol enhances rain formation within the cloud, alters evaporation in the entrainment zones, and induces more turbulence.

Using latent heating and vertical motion profiles from the Wisconsin Algorithm for Latent heating and Rainfall Using Satel-
65 lites (WALRUS), we show that there is a discernible signal of invigoration in warm clouds due to aerosol. Observations are limited to cumulus clouds discerned using CloudSat and Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation (CALIPSO) observations. The Moderate Resolution Imaging Spectroradiometer (MODIS) aerosol index (product of aerosol optical depth and Angstrom exponent) is used as a proxy for how aerosol concentrations affect the number of cloud condensation nuclei. A series of constraints are implemented in order to control for the role of stability in modulating (or confounding
70 signals of) invigoration.

2 Data and Methods

2.1 Data

All observations are from instruments aboard NASA A-Train satellites from 2007 to 2010 and from 60° south to 60° north. Aerosol index (AI) from MODIS serves as our aerosol concentration proxy while the AMSR-E provides the mean cloud liquid
75 water path of the scene. CloudSat's cloud profiling radar (CPR) is used to define cloud extent and we employ WALRUS to infer changes in latent heating and vertical motion within cloud profile.

Aerosol index is the product of the Angstrom exponent and the aerosol optical depth measured at 550 nm and is better correlated with cloud droplet concentrations than AOD (Ångström, 1964; Hasekamp et al., 2019). MODIS AI is available in clear sky scenes over the ocean, meaning cloudy AI must be interpolated from nearby cloud-free scenes (Levy et al., 2010). We
80 remove AI within 2 km of the clouds in order to reduce the influence of aerosol swelling in high humidity scenes (Christensen et al., 2017). We define pristine conditions as those with an AI less than .042 and polluted as those with an AI higher than .09. These roughly correspond to the lower and upper 20 percentiles of our dataset. Avoiding intermediate AIs reduces the possibility our analysis captures possible transition states as clouds move out of the aerosol limited regime (Koren et al., 2014).

Clouds are limited to LWPs between 150 to 200 gm^{-2} using AMSR-E (Wentz and Meissner, 2007). Although AMSR-E
85 LWP is derived using a larger field-of-view, a rough constraint on cloud liquid water content provided the spatial extent and depth of the cloud are limited using CPR observations. Cloud extents are defined using CloudSat's 2B-CLDCLASS-LIDAR product by sorting clouds by the number of contiguous cloudy pixels and limiting the analysis to clouds with at most 15 contiguous, cloudy pixels, approximately the size of an AMSR-E footprints (Sassen et al., 2008). This cloud based partitioning is analogous to the cloud object based partitioning used by Igel et al. (2014) except while Igel et al. (2014) focused on deep
90 convective systems, our clouds are constrained to shallow convective types. We focus on cumulus warm clouds, rather than stratus or stratocumulus, in order isolate the effects of aerosol on shallow convection.



Environmental information is provided by MERRA-2 reanalysis. We define the stability of the atmosphere using the estimated inversion strength (EIS) (Wood and Bretherton, 2006). Stability of the boundary layer controls the depth of the cloud making it imperative that this relationship is constrained in order to separate aerosol effects from environmental forcings (Zuidema et al., 2009). Unstable environments are defined as having an EIS below 1 K while stable environments are defined as having an EIS above 3 K. This partitions environments into two main regimes: trade cumuli (unstable) and cumuli from stratocumulus to cumulus transitions (stable). A dry free atmosphere alters the distribution of liquid throughout the cloud layer, thereby directly impacting precipitation formation processes as well. In order to control for these interactions, clouds are further subset into a dry regime whereby the RH_{700} is below 30% to analyze how dry air entrainment may impact invigoration processes.

2.2 Latent heating profiles

The Wisconsin Algorithm for Latent heating and Rainfall Using Satellites (WALRUS) provides information on the latent heating and vertical motion profiles in the atmosphere. The algorithm combines CloudSat's CPR observations with a database of warm rain states derived from the Regional Atmospheric Modeling System (RAMS) simulations to emulate realistic latent heating rates and related vertical motion (Nelson et al., 2016). WALRUS limits our analysis to maritime clouds with heights less than the freezing level and only those that exhibit reflectivity greater than 0 dBZ somewhere in the column, consistent with the Rain Certain flag in CloudSat's 2C-PRECIP-COLUMN product (Haynes et al., 2009). Our results do not include the effects of drizzle on possible invigoration processes. This should also focus our results on only the growing and mature stages of shallow convection. Signals of invigoration are derived based on changes in the latent heating within the cloud. Precipitation formation rates correspond to the latent heat release within the cloud, while evaporation due to entrainment at the cloud top or below the cloud are indicated by cooling from WALRUS. Enhanced turbulence, or the change in vertical velocity, is determined by the difference in vertical velocity between polluted and pristine environments.

WALRUS employs a Bayesian Monte Carlo method in order to derive probabilistic latent heating profile. While precipitation amounts alone can be used to infer total latent heating in the column, vertically-resolved reflectivity profiles allow the inference of the distribution of latent heating throughout the profile, below, within, and above the cloud. The Bayesian Monte Carlo method relies on an a priori distribution of possible characteristics to connect to the CloudSat observations. The a priori database is created using the RAMS model with simulations based on the Atlantic Trade Wind Experiment field campaign. The model is run at a 250 m horizontal and 100 m vertical resolution for a set of sea surface temperatures (293 K, 298 K, and 303 K). Quick Beam produces radar reflectivity profiles and attenuation signals from the RAMS simulation, which are sampled every 40 minutes for the database. Overall, WALRUS had 1.4 million possible a priori warm rain structures against which observed CloudSat reflectivities are compared to retrieve the most physically realistic latent heating and associated vertical motion rates. For more information please refer to Nelson et al. (2016).




Figure 1. An example of the latent heating profiles for clouds in unstable (top) and stable (bottom) environments with a 7 km rain system size.

2.3 Partitioning clouds

Cloud profiles are partitioned according to the individual cloud base and cloud top heights determined for each profile using
125 CloudSat's 2B-CLDCASS-LIDAR CloudLayerBase and CloudLayerTop products. These heights are used to distinguish the
in-cloud region from the environment below or above it. The maximum above cloud cooling due to evaporation is found by
taking the maximum of all the cooling rates (which include cooling by evaporation) starting at the cloud top to the top of the
profile. The cloud top is obvious in the latent heating profiles (Figure 1); the abrupt shift from heating to cooling indicates
the entrainment zone of the cloud near the cloud top. The mean below cloud cooling rate is similarly found using the cloud
130 base height from 2B-CLDCASS-LIDAR and taking the sum of all evaporative cooling rates at the cloud base to the bottom
of the profiles (approximately ground level). The geometrical center of the cloud is the midpoint of the cloud (e.g. for a 7 km
cloud as seen in Figure 1, the midpoint is 3.5 km) therefore the profiles on either side are used to determine the behavior of the
geometrical center of the cloud.

3 Results and Discussion

135 3.1 Aerosol effects on warm rain formation rates

Theoretical arguments for warm rain invigoration predict that in a more polluted environment, the rate of collision coalescence
and therefore precipitation production increases. Our analysis suggests that, on average, clouds in polluted environments do
not show an increased rates of precipitation formation relative to those in pristine environments (black solid line, Figure 2). The




Figure 2. The mean amount of latent heating released due to precipitation formation rate in the geometrical center of the rain system as a function of rain system size for all (top) and dry (bottom) warm clouds with an extent of 15 km. Black is for all stabilities, blue is for unstable environments, red is for stable environments; dashed represents pristine and solid represents polluted surroundings.



140 difference between polluted (solid) and pristine (dashed) conditions is minimal when clouds in environments are considered together. However, when separated according to the environmental stability, it is evident that the reason for this is not that the warm rate intensity is unaffected by aerosol loading, it is that clouds react differently under stable and unstable conditions. In unstable environments, polluted conditions lead to a marked increase in precipitation rate relative to unstable, pristine conditions (blue, dotted line) for all rain systems smaller than ~ 6 km. Conversely, stable, polluted conditions (red, solid line) lead to a decrease in precipitation rate relative to stable, pristine conditions (red, solid line). The opposite reactions in stable vs. 145 unstable conditions offset each other, giving the impression that warm rain is unaffected by aerosol loading when in actuality its sensitivity is environmentally dependent. Invigoration is only identifiable when stability is accounted for and this suggests that aerosol-induced invigoration of shallow convection may exhibit marked spatial patterns globally.

Polluted clouds exhibit a non-linear relationship between the size of the rain system and the maximum rain formation rate, pristine clouds show a steady, linear increase in the rain formation rate as the size of the rain system increases (Figure 2). Rain 150 formation in polluted clouds, on the other hand, exhibits an inflection point first increasing gradually with size up to 5 km and then after that size they show much greater rates of rain formation. This inflection point depends on both the stability of the boundary layer and the humidity of the free atmosphere. Pristine conditions do not show this acceleration.

The core of a warm convective system should theoretically exhibit the greatest invigoration of precipitation. Our results indicate this conceptual model is correct: as invigoration of the warm rain formation rate due to aerosol is most pronounced 155 in the geometrical center of the rain system (Figure 3). Mean precipitation rates increases in the center of unstable, polluted clouds relative to both cleaner and more stable conditions. This effect is exacerbated in dry conditions (Figure 3, bottom) until the rain system seems to hit a size inflection point around 7 km. While instability in polluted clouds leads to greater formation rates in the center, clouds in stable but equally polluted environments show a decrease in rain production relative to pristine conditions. This suppressive behavior is observed regardless of the overlying free atmosphere, as clouds in dry environments 160 (Figure 3 bottom) show the same behavior as all clouds (Figure 3 top).

3.2 Aerosol effects on evaporative processes

However, that is not to say that the free atmosphere does not play a role in altering the thermodynamics or possible invigorate state of warm rain systems. Evaporative processes link entrainment, below cloud evaporation, precipitation formation, and the energy budget of a cloud. When focusing on how aerosol may affect entrainment, the moisture content of the free atmosphere 165 becomes a controlling factor. A drier atmosphere fosters greater evaporation rates above the cloud in more polluted environments (Figure 4). While increased mixing with the free atmosphere may lead to cloud deepening, it may also lead to an early onset of cloud breakup processes through evaporation-entrainment (Small et al., 2009). In some cases, increased entrainment and evaporation at the cloud top could lead to reduced cloud top heights, opposite of an invigoration effect (Xue and Feingold, 2006). Whether the growth of a particular cloud is enhanced or inhibited may depend on the distribution of liquid water near 170 the cloud top and the ability of the cloud to penetrate the free atmosphere.

A drier atmosphere enhances cloud top evaporation in only unstable conditions; clouds in stable conditions are unaffected by a drier free atmosphere. This is likely due to the stronger capping inversion in stable conditions which limits mixing with




Figure 3. The maximum rate latent heat release due to precipitation within the cloud as a function of rain system size for all (top) and dry (bottom) warm clouds with an extent of 15 km. Black is for all stabilities, blue is for unstable environments, red is for stable environments; dashed represents pristine and solid represents polluted surroundings.




Figure 4. The maximum rate latent heating released due to evaporation above the cloud as a function of rain system size for all (top) and dry (bottom) warm clouds with an extent of 15 km. Black is for all stabilities, blue is for unstable environments, red is for stable environments; dashed represents pristine and solid represents polluted surroundings.



the dry free atmosphere, limiting its effects on the cloud layer and, by extension, the invigoration process (Christensen and Stephens, 2011). While clouds in stable environments have similar responses in precipitation formation rates, the inversion's
175 role in limiting effects is more pronounced in evaporation due to entrainment mixing. By limiting the amount of mixing with the free atmosphere, the inversion damps the ability of stable, polluted clouds to experience a cloud deepening effect compared to unstable, polluted clouds. While these clouds do not show signs of invigoration, stable, polluted conditions may prolong cloud lifetime by lessening cloud thinning processes (Van der Dussen et al., 2014).

That the cores of dry, polluted, unstable systems experience significantly greater rain formation rates than all other environ-
180 ments may suggest these clouds undergo some aggregation process focusing the majority of precipitation formation within the core of the cloud. This results agrees with a theoretical model posed by Morrison (2017), where entrainment of dry air leads to narrowing effect on the cumuli and enhancement of the core. Aerosol may act to invigorate this specific response by increasing the entrainment-evaporation at the cloud top, promoting turbulence within the cloud layer.

Below cloud evaporation and its associated cooling destabilizes the boundary layer, which could then further invigorate
185 the cloud layer through amplified turbulence (Xue and Feingold, 2006). Figure 5 demonstrates that larger warm rain systems exhibit considerably more below cloud evaporation than smaller systems. Evaporation, even when weighted by the size of the rain system, scales with the total amount of rain forming, as more rain means more possible below cloud evaporation. Polluted clouds exhibit less below cloud evaporation regardless of the stability and size of the rain system. This may imply that pristine conditions stabilize the boundary through below cloud cooling and increasing the boundary layer temperature inversion.

190 There are two possible mechanisms that may lead to polluted environments having a relatively lower rate of evaporation below cloud compared to their pristine counterparts. The first mechanism relies on the change in droplet size due to the differences in where rain is being produced in the cloud under pristine and polluted conditions. In polluted conditions, rain may form higher within the cloud; as precipitation forms and drops fall, the drop grows larger as its path increases, decreasing the amount of evaporation below cloud base (Dagan et al., 2016). So although aerosol loading decreases the mean cloud drop
195 size, rain droplet size experiences an inverse effect, the magnitude of which is determined by the height within the cloud where precipitation forms.

The second possibility is that in environments with drier free atmospheres, clouds in unstable environments (both pristine and polluted) have much greater rates of evaporation below the cloud (Figure 5 top). The increased evaporation below cloud may be driven by the turbulence created by increased evaporation in the entraining layer of the cloud (Figure 4), leading to
200 more mixing throughout the cloud layer. This in turn leads to more activation of available CCN, decreasing the mean drop size and increasing the rate of evaporation below the cloud in both polluted and pristine cases. It is possible that in polluted, unstable environments, smaller droplets evaporate efficiently, quickly increasing the humidity of the lower boundary layer, resulting in an overall decrease in the rate of evaporation (relative to pristine conditions) as the cloud continues to precipitate (Pincus and Baker, 1994; Jiang et al., 2009). Or it may be that stable, polluted boundary layers show the lowest rates of below
205 cloud evaporation because they also have lower rates of precipitation formation. Establishing the specific processes responsible for the observed invigoration signatures is not possible from current satellite observations.




Figure 5. The mean rate of latent heating due to evaporation below the cloud as a function of rain system size for all (top) and dry (bottom) warm clouds with an extent of 15 km. Black is for all stabilities, blue is for unstable environments, red is for stable environments; dashed represents pristine and solid represents polluted surroundings.



The implied effects on cloud lifetime in unstable, polluted conditions agree with Albrecht's original hypothesis, whereby aerosol works to increase cloud lifetime. The overall effect of aerosol loading on not just the thermodynamics, but lifetime of the cloud depend heavily on the environment surrounding the cloud (Albrecht, 1993). In unstable, polluted environments, below cloud evaporation cools the lower boundary layer, while latent heating due to rain formation warms cloudy portion of the boundary layer, stabilizing the boundary layer. This stabilizing effect would decrease in time, however, as the magnitudes of both the cooling below cloud and warming in cloud depend on the instability (Dagan et al., 2017). The same stabilizing effect may be seen in pristine scenes as well, as clean clouds in unstable conditions also showed greater rates of below cloud cooling due to evaporation. So while aerosol may work to help prolong lifetime through this stabilizing mechanism, the environment through a stabilizing feedback works to lengthen the cloud lifetime regardless of the aerosol conditions.

3.3 Aerosol Effects on Vertical Motion

Results are consistent with the hypothesis that invigoration will increase turbulence, indicated by changes in vertical motion, within the cloud layer due to greater amounts of latent heat release (Rosenfeld et al., 2008). Figure 6 shows that clouds in polluted environments display higher updraft speeds within the cloud layer than those in pristine environments. This reaffirms an ongoing hypothesis that cloud deepening is driven by enhanced updrafts (Christensen and Stephens, 2011). Aerosol may act to redistribute water throughout the cloud resulting in changes to the distribution of latent heating (Dagan et al., 2018). Modifying where latent heat is released, especially increasing the difference between the center where rain formation is occurring and edge evaporation due to cloud edge entrainment, alters turbulence and flow within the cloud layer. As seen in Figure 1, edge and core behavior and latent heating signatures are markedly different. While it remains unclear how aerosol may affect the absolute amount of water within a cloud, it is clear aerosol affects how water is distributed within the cloud (Toll et al., 2019; Rosenfeld et al., 2019).

When separated into stable and unstable environments (Figure 6) it becomes obvious how *strongly* stable environments damp invigoration. While unstable environments intensify the turbulence within the cloud layer, stable environments show only a faint increase in turbulence at the middle of the cloud. This may explain why stable, polluted environments also manifest the smallest rates of evaporation due to cloud top entrainment, as these clouds have less overturning motion near the cloud tops. Though stable, polluted environments displayed a reduced core precipitation formation rate compared to their pristine counterparts, the reduced size of the polluted droplets may allow greater mobility and therefore rates of vertical motion (Koren et al., 2015). Unstable, large rain systems (~ 6 km) may have downdrafts within the cloud due to enhanced evaporation entrainment, leading to more in-cloud turbulence, overturning motion, and mixing.

It should be noted that increased rates of rain formation in unstable conditions could induce a positive feedback: latent heating increases from faster rain formation which leads to more vertical motion and turbulence, greater updraft speeds, more entrainment and therefore buoyancy fluctuations within the cloud, which in turn leads to more collision-coalescence and latent heating. This could explain both the greatly enhanced rain formation rates in the center (Figure 2) and intensified turbulence (Figure 6). However, this feedback may be particularly sensitive to the size of the rain system, as larger systems may decrease the chance of the feedback occurring. The scattered increases in turbulence seen in rain systems of size 6 km and the sloping




Figure 6. The difference (polluted - pristine) in vertical motion in unstable (top) and stable (bottom) environments for clouds of rain size 4 km (top), 5 km (middle), and 6 km (bottom).

off of rain formation rates at the same size imply a sensitivity to the size and organization of the system (Fan et al., 2016). Additionally, any possible invigoration feedback may be moderated by a stabilizing effect induced in unstable conditions, whereby increased cooling due to evaporation below cloud and warming due to rain formation within the cloudy layer stabilize the boundary layer.

245 4 Conclusions

Cumulus clouds in polluted, unstable environments display greater rates of peak and core precipitation formation along with greater amounts of vertical motion and therefore turbulence within the cloud. Dry environments act to increase this response, along with inducing further invigoration effects by increasing the amount of entrainment mixing. Stable environments act to dampen invigoration by capping entrainment effects and reducing precipitation formation rates. In polluted environments, a stable boundary layer and strong inversion acts to inhibit rain production relative to pristine environments. This reverse response may be driven by reduced amounts of vertical motion in polluted cloud tops and bases, hindering precipitation formation throughout the cloud.

Invigoration is an "elusive" effect in the aerosol-cloud interaction community perhaps because observing it depends on the definition used. Based on our definition, whereby aerosol loading in warm clouds increases the precipitation formation rate and in-cloud vertical motion (a proxy for turbulence), there is evidence invigoration may occur. However from the results shown, two important aspects of invigoration emerge:



260 1. The magnitude of invigoration in marine cumuli depends strongly on the size of the cloud. As the rain system grows, all possible signs of invigoration, from more rain formation to increased turbulence, are a function of the size of the rain system. This implies the organization of the cloud plays a role in defining how aerosol loading may impact aspects of invigoration. The dependence of these processes on the size of the system may explain why many components of invigoration (LWP response, cloud deepening, etc.) are non-linear when regressed against aerosol alone.

2. Stability can reduce and/or reverse all aspects of invigoration within the cloud layer. The mean warm cloud signal of invigoration is completely buffered by the environment unless stability is accounted for.

265 The mean formation rate in polluted environments closely tracked the mean pristine rate (Figures 2, 3), only by accounting for both the stability and humidity effects do distinct signals appear. Even with stability and the environment accounted for through sets of constraints, we had to impose limits on liquid water path in order to isolate the invigoration. Without these constraints, on the environment and cloud state, invigoration would be indiscernible using only the mean response. Models must account for these factors when parameterizing aerosol impacts on precipitation.

270 Our analysis is heavily predicated on latent heating and vertical motion from WALRUS. As such, some uncertainty is inherent in the results due to the indirect nature of a satellite-based latent heating and vertical motion estimates. However, the vertical structure of reflectivity and integral constraint provided by the path-integrated attenuation (PIA) provide strong constraints on hydrometeor distributions and integrated water path in warm clouds (Lebsock and L'Ecuyer, 2011; Nelson et al., 2016). The results presented here imply that aerosols induce systematic changes in observed reflectivity profiles and attenuation
275 from raindrops that are indicative of different precipitation formation and above/below cloud evaporation rates, though the precise magnitudes may be uncertain. Further support for the plausibility of the WALRUS products is provided by Nelson and L'Ecuyer (2018) who document systematic regional variations in latent heating and inferred vertical motion in global warm rain systems. This study is limited to only a "snapshot" view of clouds, unable to account for the individual lifetimes of individual cumuli. Tracking clouds throughout their lifetime, similar to the tracking employed by (Christensen et al., 2020)
280 to follow clouds through the stratocumulus to cumulus transition, would offer insights into regime specific processes at the root of the signatures seen here. However, geostationary satellite observations provide much more limited insights into cloud structure. The lack of vertical structure precludes the retrieval of in-cloud latent heating and vertical motion. Future analysis should attempt to blend vertical structure information with observations of the state of the cloud throughout its lifetime in order to understand how the trajectory of a cloud affects its response to aerosol loading.

285 *Author contributions.* Alyson Douglas completed all analysis and writing of the paper. Tristan L'Ecuyer provided guidance on the analysis and critical feedback on the story of the paper.

Competing interests. The authors have no competing interests to declare.



Acknowledgements. Thank you to Guy Dagan for feedback on the manuscript. Thank you to Philip Stier and the EU Horizon Grant for funding this work.



290 References

- Ackerman, A. S., Kirkpatrick, M. P., Stevens, D. E., and Toon, O. B.: The impact of humidity above stratiform clouds on indirect aerosol climate forcing, *Nature*, 432, 1014–1017, 2004.
- Albrecht, B. A.: Aerosols, cloud microphysics, and fractional cloudiness, *Science*, 245, 1227–1230, 1989.
- Albrecht, B. A.: Effects of precipitation on the thermodynamic structure of the trade wind boundary layer, *Journal of Geophysical Research: Atmospheres*, 98, 7327–7337, 1993.
- 295 Altaratz, O., Koren, I., Remer, L., and Hirsch, E.: Cloud invigoration by aerosols—Coupling between microphysics and dynamics, *Atmospheric Research*, 140, 38–60, 2014.
- Ångström, A.: The parameters of atmospheric turbidity, *Tellus*, 16, 64–75, 1964.
- Berg, W., L'Ecuyer, T., and Kummerow, C.: Rainfall climate regimes: The relationship of regional TRMM rainfall biases to the environment, *Journal of Applied Meteorology and Climatology*, 45, 434–454, 2006.
- 300 Berg, W., L'Ecuyer, T., and van den Heever, S.: Evidence for the impact of aerosols on the onset and microphysical properties of rainfall from a combination of satellite observations and cloud-resolving model simulations, *Journal of Geophysical Research: Atmospheres*, 113, 2008.
- Bony, S. and Dufresne, J.-L.: Marine boundary layer clouds at the heart of tropical cloud feedback uncertainties in climate models, *Geophysical Research Letters*, 32, 2005.
- 305 Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., et al.: Clouds and aerosols, in: *Climate change 2013: the physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, pp. 571–657, Cambridge University Press, 2013.
- Chen, Q., Koren, I., Altaratz, O., Heiblum, R. H., Dagan, G., and Pinto, L.: How do changes in warm-phase microphysics affect deep convective clouds?, *Atmospheric Chemistry and Physics*, 17, 2017.
- 310 Christensen, M. W. and Stephens, G. L.: Microphysical and macrophysical responses of marine stratocumulus polluted by underlying ships: Evidence of cloud deepening, *Journal of Geophysical Research: Atmospheres*, 116, 2011.
- Christensen, M. W., Neubauer, D., Poulsen, C. A., Thomas, G. E., McGarragh, G. R., Povey, A. C., Proud, S. R., and Grainger, R. G.: Unveiling aerosol–cloud interactions—Part 1: Cloud contamination in satellite products enhances the aerosol indirect forcing estimate, *Atmospheric Chemistry and Physics*, 17, 2017.
- 315 Christensen, M. W., Jones, W. K., and Stier, P.: Aerosols enhance cloud lifetime and brightness along the stratus-to-cumulus transition, *Proceedings of the National Academy of Sciences*, 117, 17 591–17 598, 2020.
- Dagan, G., Koren, I., Altaratz, O., and Heiblum, R. H.: Aerosol effect on the evolution of the thermodynamic properties of warm convective cloud fields, *Scientific reports*, 6, 1–8, 2016.
- 320 Dagan, G., Koren, I., Altaratz, O., and Heiblum, R. H.: Time-dependent, non-monotonic response of warm convective cloud fields to changes in aerosol loading, *Atmospheric Chemistry and Physics*, 17, 7435–7444, 2017.
- Dagan, G., Koren, I., and Altaratz, O.: Quantifying the effect of aerosol on vertical velocity and effective terminal velocity in warm convective clouds, *Atmospheric Chemistry and Physics*, 18, 2018.
- Dey, S., Sengupta, K., Basil, G., Das, S., Nidhi, N., Dash, S., Sarkar, A., Srivastava, P., Singh, A., and Agarwal, P.: Satellite-based 3D structure of cloud and aerosols over the Indian Monsoon region: implications for aerosol–cloud interaction, in: *Remote sensing and modeling of the atmosphere, oceans, and interactions iv*, vol. 8529, p. 852907, International Society for Optics and Photonics, 2012.
- 325



- Douglas, A. and L'Ecuyer, T.: Quantifying variations in shortwave aerosol-cloud-radiation interactions using local meteorology and cloud state constraints, *Atmospheric Chemistry & Physics*, 19, 6251–6268, 2019.
- Fan, J., Wang, Y., Rosenfeld, D., and Liu, X.: Review of aerosol–cloud interactions: Mechanisms, significance, and challenges, *Journal of the Atmospheric Sciences*, 73, 4221–4252, 2016.
- 330 Hasekamp, O. P., Gryspeerdt, E., and Quaas, J.: Analysis of polarimetric satellite measurements suggests stronger cooling due to aerosol-cloud interactions, *Nature communications*, 10, 1–7, 2019.
- Haynes, J. M., L'Ecuyer, T. S., Stephens, G. L., Miller, S. D., Mitrescu, C., Wood, N. B., and Tanelli, S.: Rainfall retrieval over the ocean with spaceborne W-band radar, *Journal of Geophysical Research: Atmospheres*, 114, 2009.
- 335 Heiblum, R. H., Pinto, L., Altaratz, O., Dagan, G., and Koren, I.: Core and margin in warm convective clouds–Part 2: Aerosol effects on core properties, *Atmospheric Chemistry and Physics*, 19, 2019.
- Igel, M. R., Drager, A. J., and Van Den Heever, S. C.: A CloudSat cloud object partitioning technique and assessment and integration of deep convective anvil sensitivities to sea surface temperature, *Journal of Geophysical Research: Atmospheres*, 119, 10 515–10 535, 2014.
- Jiang, H., Xue, H., Teller, A., Feingold, G., and Levin, Z.: Aerosol effects on the lifetime of shallow cumulus, *Geophysical Research Letters*, 33, 2006.
- 340 Jiang, H., Feingold, G., and Koren, I.: Effect of aerosol on trade cumulus cloud morphology, *Journal of Geophysical Research: Atmospheres*, 114, 2009.
- Koren, I., Dagan, G., and Altaratz, O.: From aerosol-limited to invigoration of warm convective clouds, *science*, 344, 1143–1146, 2014.
- Koren, I., Altaratz, O., and Dagan, G.: Aerosol effect on the mobility of cloud droplets, *Environmental Research Letters*, 10, 104 011, 2015.
- 345 Kubar, T. L., Hartmann, D. L., and Wood, R.: Understanding the importance of microphysics and macrophysics for warm rain in marine low clouds. Part I: Satellite observations, *Journal of the atmospheric sciences*, 66, 2953–2972, 2009.
- Lebsock, M. D. and L'Ecuyer, T. S.: The retrieval of warm rain from CloudSat, *Journal of Geophysical Research: Atmospheres*, 116, 2011.
- L'Ecuyer, T. S., Berg, W., Haynes, J., Lebsock, M., and Takemura, T.: Global observations of aerosol impacts on precipitation occurrence in warm maritime clouds, *Journal of Geophysical Research: Atmospheres*, 114, 2009.
- 350 Levy, R., Remer, L., Kleidman, R., Mattoo, S., Ichoku, C., Kahn, R., and Eck, T.: Global evaluation of the Collection 5 MODIS dark-target aerosol products over land, *Atmospheric Chemistry and Physics*, 10, 10 399, 2010.
- Li, Z., Niu, F., Fan, J., Liu, Y., Rosenfeld, D., and Ding, Y.: Long-term impacts of aerosols on the vertical development of clouds and precipitation, *Nature Geoscience*, 4, 888–894, 2011.
- Morrison, H.: An analytic description of the structure and evolution of growing deep cumulus updrafts, *Journal of the Atmospheric Sciences*, 74, 809–834, 2017.
- 355 Nelson, E. L. and L'Ecuyer, T. S.: Global character of latent heat release in oceanic warm rain systems, *Journal of Geophysical Research: Atmospheres*, 123, 4797–4817, 2018.
- Nelson, E. L., L'Ecuyer, T. S., Saleeby, S. M., Berg, W., Herbener, S. R., and Van Den Heever, S. C.: Toward an algorithm for estimating latent heat release in warm rain systems, *Journal of Atmospheric and Oceanic Technology*, 33, 1309–1329, 2016.
- 360 Pincus, R. and Baker, M.: Precipitation, solar absorption, and albedo susceptibility in marine boundary layer clouds, *Nature*, 372, 250–252, 1994.
- Rosenfeld, D., Lohmann, U., Raga, G. B., O'Dowd, C. D., Kulmala, M., Fuzzi, S., Reissell, A., and Andreae, M. O.: Flood or drought: how do aerosols affect precipitation?, *science*, 321, 1309–1313, 2008.



- Rosenfeld, D., Zhu, Y., Wang, M., Zheng, Y., Goren, T., and Yu, S.: Aerosol-driven droplet concentrations dominate coverage and water of
365 oceanic low-level clouds, *Science*, 363, eaav0566, 2019.
- Sassen, K., Wang, Z., and Liu, D.: Global distribution of cirrus clouds from CloudSat/Cloud-Aerosol lidar and infrared pathfinder satellite
observations (CALIPSO) measurements, *Journal of Geophysical Research: Atmospheres*, 113, 2008.
- Seifert, A., Heus, T., Pincus, R., and Stevens, B.: Large-eddy simulation of the transient and near-equilibrium behavior of precipitating
shallow convection, *Journal of Advances in Modeling Earth Systems*, 7, 1918–1937, 2015.
- 370 Small, J. D., Chuang, P. Y., Feingold, G., and Jiang, H.: Can aerosol decrease cloud lifetime?, *Geophysical Research Letters*, 36, 2009.
- Spill, G., Stier, P., Field, P. R., and Dagan, G.: Effects of aerosol in simulations of realistic shallow cumulus cloud fields in a large domain,
Atmospheric Chemistry and Physics, 19, 13 507–13 517, 2019.
- Stevens, B. and Feingold, G.: Untangling aerosol effects on clouds and precipitation in a buffered system, *Nature*, 461, 607–613, 2009.
- Toll, V., Christensen, M., Quaas, J., and Bellouin, N.: Weak average liquid-cloud-water response to anthropogenic aerosols, *Nature*, 572,
375 51–55, 2019.
- Van der Dussen, J., De Roode, S., and Siebesma, A.: Factors controlling rapid stratocumulus cloud thinning, *Journal of the Atmospheric
Sciences*, 71, 655–664, 2014.
- Wentz, F. J. and Meissner, T.: Supplement 1 algorithm theoretical basis document for AMSR-E ocean algorithms, NASA: Santa Rosa, CA,
USA, 2007.
- 380 Wood, R.: Stratocumulus clouds, *Monthly Weather Review*, 140, 2373–2423, 2012.
- Wood, R. and Bretherton, C. S.: On the relationship between stratiform low cloud cover and lower-tropospheric stability, *Journal of climate*,
19, 6425–6432, 2006.
- Xue, H. and Feingold, G.: Large-eddy simulations of trade wind cumuli: Investigation of aerosol indirect effects, *Journal of the atmospheric
sciences*, 63, 1605–1622, 2006.
- 385 Yuan, T., Remer, L., and Yu, H.: Microphysical, macrophysical and radiative signatures of volcanic aerosols in trade wind cumulus observed
by the A-Train, 2011.
- Zelinka, M. D., Myers, T. A., McCoy, D. T., Po-Chedley, S., Caldwell, P. M., Ceppi, P., Klein, S. A., and Taylor, K. E.: Causes of higher
climate sensitivity in CMIP6 models, *Geophysical Research Letters*.
- Zuidema, P., Painemal, D., De Szoeko, S., and Fairall, C.: Stratocumulus cloud-top height estimates and their climatic implications, *Journal
390 of Climate*, 22, 4652–4666, 2009.