

# Controls on surface aerosol particle number concentrations and aerosol-limited cloud regimes over the central Greenland Ice Sheet.

Heather Guy<sup>1,2</sup>, Ian M. Brooks<sup>2</sup>, Ken S. Carslaw<sup>2</sup>, Benjamin J. Murray<sup>2</sup>, Von P. Walden<sup>3</sup>, Matthew D. Shupe<sup>4,5</sup>, Claire Pettersen<sup>6</sup>, David D. Turner<sup>8</sup>, Christopher J. Cox<sup>5</sup>, William D. Neff<sup>4,5</sup>, Ralf Bennartz<sup>6,7</sup>, and Ryan R. Neely III<sup>1,2</sup>

<sup>1</sup>National Centre for Atmospheric Science, Leeds, U.K.

<sup>2</sup>School of Earth and Environment, University of Leeds, Leeds, U.K.

<sup>3</sup>Department of Civil and Environmental Engineering, Laboratory for Atmospheric Research, Washington State University, Pullman, WA, USA

<sup>4</sup>Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO, USA

<sup>5</sup>Physical Sciences Laboratory, National Oceanic and Atmospheric Administration, Boulder, CO, USA

<sup>6</sup>Space Science and Engineering Center, UW-Madison, Madison, WI, USA

<sup>7</sup>Earth and Environmental Sciences, Vanderbilt University, Nashville, TN, USA

<sup>8</sup>Global Systems Laboratory, National Oceanic and Atmospheric Administration, Boulder, CO, USA

**Correspondence:** Heather Guy (heather.guy@ncas.ac.uk)

**Abstract.** This study presents the first full annual cycle (2019-2020) of ambient surface aerosol particle number concentration measurements (condensation nuclei  $> 20$  nm,  $N_{20}$ ) collected at Summit Station, in the centre of the Greenland Ice Sheet ( $72.58^\circ$  N,  $-38.45^\circ$  E, 3250 m asl). The mean surface concentration in 2019 was  $129 \text{ cm}^{-3}$ , with the 6 h mean ranging between  $1 \text{ cm}^{-3}$  and  $1441 \text{ cm}^{-3}$ . The highest monthly mean concentrations occurred during the late spring and summer, with the minimum concentrations occurring in February (mean:  $18 \text{ cm}^{-3}$ ). High  $N_{20}$  events are linked to anomalous anticyclonic circulation over Greenland and the descent of free tropospheric aerosol down to the surface, whereas low  $N_{20}$  events are linked to anomalous cyclonic circulation over south-east Greenland that drives upslope flow and enhances precipitation en-route to Summit. Fog strongly effects particle number concentrations, on average reducing  $N_{20}$  by 20% during the first three hours of fog formation. Extremely low  $N_{20}$  events ( $< 10 \text{ cm}^{-3}$ ) occur in all seasons, and we suggest that fog, and potentially cloud formation, can be limited by low aerosol particle concentrations over central Greenland.

## 1 Introduction

The Greenland Ice Sheet (GrIS) has been losing mass at an unprecedented and accelerating rate since the early 21st century (Rignot et al., 2008, 2011; Van Den Broeke et al., 2016; Fettweis et al., 2017; Trusel et al., 2018; Shepherd et al., 2020), and as a result, has become the largest single contributor to global sea level rise (Van Den Broeke et al., 2016; Bamber et al., 2018; Slater et al., 2020). The majority of this mass loss is due to changes in the ice sheet surface mass balance (Slater et al., 2020), and in particular, increased surface melt and run off (Enderlin et al., 2014; Van Den Broeke et al., 2016; Shepherd et al., 2020). Clouds play a critical role in the ice sheet surface mass balance, both by providing mass input in the form of precipitation, and by modulating the net radiation at the surface, thus influencing surface melt and run-off (Bennartz et al., 2013; Van Tricht

et al., 2016; Hofer et al., 2017; Miller et al., 2017). To make accurate projections of the future contribution of the GrIS to  
20 sea level rise, models must correctly represent the properties of clouds and their interaction with the surface energy budget.  
Although circulation anomalies drive a larger proportion of surface melt, discrepancies in cloud microphysical properties  
between different models currently result in larger uncertainties in future GrIS melt projections than the difference between  
low and high greenhouse emission scenarios (Hofer et al., 2019). Amongst the largest uncertainties in cloud microphysical  
modelling are the type, concentration and sources of aerosol particles (e.g. Seinfeld et al., 2016). Improving our understanding  
25 of aerosols and their relationship with cloud properties over the GrIS is therefore key to reducing the uncertainty in future  
projections of GrIS melt and global sea level rise.

Cloud properties are sensitive to the type and concentration of tropospheric aerosol particles (e.g. Twomey, 1977; Curry  
et al., 1996; Storelvmo, 2017). Mixed-phase clouds in particular, which contribute significantly to surface warming over the  
GrIS (Miller et al., 2015; Van Tricht et al., 2016), are sensitive to the number concentration of cloud condensation nuclei and  
30 ice-nucleating particles (e.g. Norgren et al., 2018; Solomon et al., 2018); where cloud condensation nuclei (CCN) are a subset  
of aerosol particles on which liquid droplets can form, and ice-nucleating particles (INP) are a subset of aerosols that can  
catalyze the formation of ice crystals.


In ice-covered polar regions, CCN concentrations can be very low; surface CCN concentrations at 0.2% supersaturation  
are usually less than  $100 \text{ cm}^{-3}$  and can regularly fall below  $10 \text{ cm}^{-3}$  in the high Arctic (e.g. Mauritsen et al., 2011; Leck  
35 and Svensson, 2015), compared to typical values of over  $1000 \text{ cm}^{-3}$  at rural mid-latitude sites (e.g. Schmale et al., 2018). In  
cases where CCN are extremely low ( $< 10 \text{ cm}^{-3}$ ), the small number of sites for droplet activation limits cloud droplet number  
concentration, and high supersaturations cause all available CCN to activate and grow to relatively large sizes, facilitating  
further growth by collision and coalescence and resulting in precipitation as drizzle (Mauritsen et al., 2011). This generates  
a positive feedback where the lack of CCN can result in total dissipation of the cloud (Mauritsen et al., 2011; Stevens et al.,  
40 2018). Thus, within this CCN-limited regime, the availability of CCN becomes a dominant control on cloud formation and  
longevity such that a small increase in concentration can lead to a decrease in droplet size that serves to reduce precipitation  
efficiency, leading to a relative increase in cloud liquid water path (LWP) (Mauritsen et al., 2011). The change in LWP in turn  
modulates the cloud longwave radiative effect (Mauritsen et al., 2011; Miller et al., 2015). Alternatively, the addition of CCN  
when a cloud is not in the CCN-limited regime can have a cooling effect at the surface in the summer due to the associated  
45 increase in cloud reflectivity of incoming solar radiation (Twomey, 1977; Intrieri et al., 2002). For optically thin clouds ( $< 40 \text{ g}$   
 $\text{m}^{-2}$ ), that are common at Summit (Shupe et al., 2013b; Miller et al., 2015), the smaller droplet size associated with increased  
CCN results in higher cloud emissivity, increasing the downwelling longwave radiative flux and having a relative warming  
effect at the surface (Lubin and Vogelmann, 2006; Garrett and Zhao, 2006). Understanding when and where each of these  
processes dominate is extremely important for understanding cloud radiative forcing and the surface energy budget (Schmale  
50 et al., 2021).

The concentration of ice-nucleating particles (INP) is also an important control on the longevity and radiative impact of  
clouds. INP are required to form primary ice in supercooled liquid clouds that are warmer than the homogeneous freezing  
temperature (approximately  $-38 \text{ }^\circ\text{C}$ ) (e.g. Kanji et al., 2017). Because the low-level clouds that have the largest radiative effect

at the Arctic surface usually have temperatures between  $-38\text{ }^{\circ}\text{C}$  and  $0\text{ }^{\circ}\text{C}$  (Shupe and Intrieri, 2004; Shupe et al., 2013b; Miller et al., 2015), INP concentrations are an important control on the ice and liquid water contents of these clouds. Clouds containing ice crystals are optically thinner than those containing only supercooled water droplets, and therefore emit less longwave radiation towards the surface, having a relative cooling effect (e.g. Prenni et al., 2007). Even more importantly, once ice crystals are present in a supercooled cloud, the lower saturation vapor pressure of ice versus liquid water results in the preferential growth of ice crystals at the expense of liquid droplets when the environment is subsaturated with respect to water but supersaturated with respect to ice. This is known as the Wegener-Bergeron-Findeisen (WBF) process, the result of which is a decrease in LWP as droplets evaporate and an increase in precipitation due to the growth of relatively large ice crystals, ultimately leading to cloud dissipation (e.g. Lohmann and Feichter, 2005). INP concentrations are typically orders of magnitude lower than CCN concentrations, and are particularly low in the Arctic based on limited existing measurements ( $\sim 10^{-7}$  to  $10^{-5}\text{ cm}^{-3}$ , Wex et al., 2019). The lack of INP in the Arctic may contribute to the unusual persistence of low-level mixed-phase stratocumulus clouds (Morrison et al., 2012), that are highly important for radiative forcing at the surface, and played a role in the anomalous GrIS surface melt event in 2012 (Bennartz et al., 2013).

Both CCN and INP concentrations are also important for precipitation accumulation. In liquid clouds, the increase in cloud droplet number concentration and associated decrease in cloud droplet size under high CCN concentrations reduces the opportunities for droplet collision and coalescence and hence reduces precipitation relative to equivalent situations with lower droplet concentrations (e.g. Lohmann and Feichter, 2005). In mixed-phase clouds this process is more complex, since changes in the cloud droplet size distribution can have both positive and negative effects on the efficiency of ice production (Cheng et al., 2010; Lance et al., 2011; Possner et al., 2017). Cloud phase partitioning is also important since ice phase clouds have markedly different precipitation characteristics to those containing super-cooled liquid water (Pettersen et al., 2018; McIlhattan et al., 2020). Model simulations generally overestimate precipitation accumulation over the GrIS (McIlhattan et al., 2017; Kay et al., 2018; Lenaerts et al., 2020), and in particular the contribution from mixed-phase clouds. McIlhattan et al. (2017) find that the CESM model overestimates snow frequency from mixed-phase clouds by 52% and underestimates the occurrence frequency of liquid-bearing clouds by 21% over the central GrIS. This is consistent with an overly active WBF process in the model; a process that is strongly controlled by INP concentrations.

To date, all observations of the CCN-limited regime (Mauritsen et al., 2011; Leaitch et al., 2016), and INP concentrations (Wex et al., 2019), in the Arctic are located at marine or coastal sites. However, the central GrIS is a distinct Arctic environment due to its high elevation (3250 m asl at its highest point, Fig. 1) and persistent ice cover ( $1.7 \times 10^6\text{ km}^2$ ) that results in a year-round high surface albedo. There are no substantial local sources of aerosol from the surface for over 400 km in any direction from the center of the ice sheet. The year-round high surface albedo of the central GrIS (Box et al., 2012) results in unique seasonality in cloud radiative forcing. Most parts of the Arctic have less snow and ice cover in the summer and hence a lower albedo, during this time clouds can have a net cooling effect at the surface due their relatively high albedo (e.g. Shupe and Intrieri, 2004). In contrast, over the central GrIS the seasonal change in surface albedo is negligible and clouds have a net warming effect at the surface year-round (Miller et al., 2015; Van Tricht et al., 2016).



**Figure 1.** Location of Summit Station at the highest point on the Greenland Ice Sheet. Ice elevation contours are from the Greenland Ice Mapping Project (Howat et al., 2017).

The high elevation and extreme radiative cooling from the center of the GrIS drives low-level katabatic winds that radiate towards the ice sheet edge and, combined with synoptic and large scale circulation patterns, support the formation of a persistent high pressure system over Greenland (Heinemann and Klein, 2002; Hanna et al., 2016). For this reason, Greenland has been referred to as the ‘northern wind pole’, where upper level air currents driven by the Hadley circulation descend and return to lower latitudes (Hobbs, 1945; Heinemann and Klein, 2002). The descent of upper tropospheric air to the surface of the central GrIS results in a larger contribution of well-mixed free tropospheric aerosol (Stohl, 2006). Hence the transport processes and source regions controlling the concentrations of aerosol particles over the central GrIS are distinct from other Arctic sites (Hirdman et al., 2009; Backman et al., 2021).

The presumed insignificance of local aerosol sources at the surface of the GrIS suggests that both low CCN concentrations with the potential to limit cloud formation, and low INP concentrations that can control cloud phase, could certainly occur. The difference in aerosol transport pathways to the GrIS when compared to coastal or marine Arctic sites implies that the processes controlling aerosol-limited cloud regimes, and their frequency of occurrence, might differ substantially from other Arctic locations. Hence, a thorough analysis of the role of the aerosol-limited conditions over the GrIS is warranted, especially given the unique sensitivity of the GrIS to longwave cloud forcing.

Despite the potential for aerosol-limited clouds to effect the surface mass balance of the GrIS, and the large uncertainties in modelled cloud microphysical properties over Greenland (Hofer et al., 2019; Schmale et al., 2021), there are very few observations of aerosol particle number concentration over the central GrIS, and those that do exist are mostly limited to the

105 summer season (Ziemba et al., 2010; Flyger et al., 1976; Hogan et al., 1984; Davidson et al., 1993; Bergin et al., 1994, 1995).  
This study presents the first full year of surface aerosol particle number concentration measurements from Summit Station,  
in the central GrIS, which can be used as a baseline for future modelling studies investigating the effect of cloud-aerosol  
interactions on the GrIS surface energy budget and mass balance. We assess local and synoptic controls on surface aerosol  
particle concentrations at Summit and present three case studies where extremely low total particle number concentrations (<  
110  $10 \text{ cm}^{-3}$ ) coincide with cloud dissipation, indicating that CCN-limited clouds occur over the central GrIS and could be an  
important contributor to the surface energy budget.

## 2 Measurements and methods

All observations in this study were made at Summit Station (Summit), a scientific research base funded by the U.S. National  
Science Foundation. Summit is located at the highest point on the GrIS (3250 m asl) and is over 400 km from the coast in the  
115 east and west directions, and over 1,000 km from the south west and south east coasts (Fig. 1). Aerosol, cloud and atmospheric  
profile measurements were collected as part of the ICECAPS-ACE project: ICECAPS (Integrated Characterisation of Energy,  
Clouds and Precipitation at Summit) has been operating at Summit since 2010 and consists of a suite of ground based remote  
sensing instrumentation and twice-daily radiosonde launches (Shupe et al., 2013b). The ACE (Aerosol Cloud Experiment)  
addition to ICECAPS began collecting data in February 2019 and includes measurements of surface aerosol particle number  
120 concentration and size distribution in addition to turbulent and radiative fluxes used to characterize the surface energy budget.  
This study uses a subset of ICECAPS-ACE data listed in Table 1, as well as meteorological measurements from the NOAA  
Global Monitoring Laboratory (GMLMET, 2021). The references in Table 1 provide additional information on the instruments  
and methodologies for the derived parameters. Section 2.1 provides the details of the aerosol particle number concentration  
sampling and quality control.

125 To investigate the effect of near-surface local processes that have the potential to modify surface aerosol particle concentra-  
tions, we look at four event types: fog, precipitation, blowing snow (BLSN), and strong surface-based temperature inversions  
(SBIs). For each type, we examine the change in aerosol particle concentrations across multiple events. To qualify, events of  
each type must last at least 60 minutes and separate events of the same type must be at least 5 hours apart. Subsections 2.2 to  
2.4 provide specific details about how each event type is defined.

130 To assess the synoptic controls on surface aerosol particle concentrations, we use ERA5 reanalysis data (Hersbach et al.,  
2020) made available by the European Centre for Medium-Range Weather Forecasts (ECMWF). ERA5 is the highest resolution  
global reanalysis product to date, with  $\sim 15$  km horizontal resolution over Greenland, 137 pressure levels up to 80 km, and 1 h  
temporal resolution. We also use ERA5 reanalysis to drive the FLEXPART Lagrangian particle dispersion model (Pisso et al.,  
2019) to simulate aerosol transport pathways and surface emission sensitivities. Section 2.5 provides further details about the  
135 FLEXPART experimental design.

## 2.1 Surface aerosol particle number concentrations


A condensation particle counter (GRIMM CPC 5.400) measured the ambient number concentration of condensation nuclei at 1 Hz frequency. The omni-directional conical inlet head was located  $\sim 3$  m above the surface (this varied slightly throughout the observation period with snow drifting and accumulation) and air was sampled with a flow rate of  $0.3 \text{ L min}^{-1}$ . The inlet  
140 was connected to the CPC via a 6 m length of conductive silicone tubing with an 8 mm inner diameter. Although the CPC is calibrated to measure condensation nuclei  $> 5$  nm diameter, the addition of the long inlet results in a loss of particles inside the tubing. Figure 2 shows an estimation of the loss of aerosol particles inside the inlet generated by the Particle Loss Calculator (Von Der Weiden et al., 2009). Smaller particles are increasingly lost due to diffusion to the walls of the inlet, and larger particles due to sedimentation and deposition. The Particle Loss Calculator does not account for the temperature  
145 gradient within the tubing, however, because the cold air in the inlet stream transitions into a warmer inlet (inside the heated building), this will act to reduce the loss of particles (Von Der Weiden et al., 2009). Also, because particle concentrations are small ( $\ll 100,000 \text{ cm}^{-3}$ ), loss due to coagulation is negligible (Von Der Weiden et al., 2009). Based on these modelled inlet losses, the CPC measured condensation nuclei with diameters between 20 nm and  $2.3 \mu\text{m}$  with over 50% efficiency (Fig 2). For this reason we henceforth refer to the CPC concentration measurements as  $N_{20}$ , indicating number concentrations of  
150 particles with diameter  $> 20$  nm. Modelled inlet losses are  $< 15 \%$  for particles with diameters between 0.08 and  $1 \mu\text{m}$ , which is representative of the typical size range of CCN in clean Arctic environments (Hudson and Da, 1996; Leaitch et al., 2016).

Ziembra et al. (2010) made measurements of surface aerosol particle size distribution between 5.5 and 195 nm at Summit in May and June 2007. Their observations suggest that high concentrations of nucleation mode particles ( $< 30$  nm diameter) occur periodically during the Summer at Summit. The reduced collection efficiency of our CPC between 20 and 40 nm would  
155 have resulted in an under-count of the total  $N_{20}$  by up to 27% during the 2007 measurement period reported by Ziembra et al. (2010), but only 8% in the accumulation mode (100 to 200 nm). The concentration of ultrafine particles ( $< 100$  nm diameter) at Summit likely varies seasonally and on shorter timescales. In the absence of year-round measurements of particle size distribution at Summit, it is not possible to fully quantify the uncertainties in  $N_{20}$  reported here.

To filter out data that may have been impacted by local station pollution, we omitted measurements collected when wind  
160 speeds are  $< 1 \text{ m s}^{-1}$  and when the wind direction is such that contaminated air from station operations may have advected across the inlet (between  $270^\circ$  and  $360^\circ$  from true north). A comparison between two OPC-N3 optical particle counters (described further in section 2.2), located at the opposite sides of camp, confirmed that these criteria are sufficient to account for the impact of local station pollution (not shown). The removal of data associated with particular surface wind conditions may bias the dataset, however during the measurement period considered in this study wind speeds  $< 1 \text{ m s}^{-1}$  only occur 3.4 % of  
165 the time and polluting wind directions only occur 9.1 % of the time.

## 2.2 Detection of fog

Supercooled liquid fog is common at Summit and occurs in all seasons, with a minimum occurrence in April and maximum in September (Cox et al., 2019). Fog droplets form on CCN and grow by condensation to typical diameters of 15 to  $25 \mu\text{m}$




**Figure 2.** Modelled particle loss as a function of particle diameter in the CPC inlet, as estimated by the Particle Loss Calculator (Von Der Weiden et al., 2009)

(Cox et al., 2019). Particles larger than  $\sim 3 \mu\text{m}$  cannot pass through the CPC inlet (Fig. 2), hence during fog events, the CPC  
 170 measures the interstitial aerosol particle concentration. In this way, fog can result in extremely low surface aerosol particle  
 concentration measurements that are not representative of the aerosol population outside of the fog (Bergin et al., 1995). In  
 the absence of an instrument designed specifically to detect fog at Summit, we use data from an Alphasense Optical Particle  
 Counter (OPC-N3, Crilley et al., 2018) located next to the CPC inlet to identify fog periods.

The OPC-N3 resolves particle size distribution in 24 bins between  $0.35 \mu\text{m}$  and  $40 \mu\text{m}$  diameter. Natural aerosol particles  
 175 with diameters greater than  $10 \mu\text{m}$  are highly unlikely to be present in central Greenland due to the large distance from the  
 source of any coarse mode aerosol particles and the large dry deposition velocity of such particles (Giorgi, 1986; Jaenicke,  
 1990). Under this assumption, particles detected by the OPC-N3 with diameters over  $10 \mu\text{m}$  must be fog droplets or ice  
 crystals. Real-time data monitoring at Summit and comparison with visual observations for 6 months confirm that the OPC-N3  
 180 detects particles within this size range during both fog and blowing snow. At Summit, 80% of cases of drifting or blowing  
 snow reported by onsite observers in 2019 occurred when the 3h mean 10 m wind speed was  $> 6 \text{ m s}^{-1}$ ; we remove all cases  
 with wind speeds above this threshold to separate fog events from possible blowing snow events. We classify fog events as  
 when the total concentration of particles with diameters  $> 10 \mu\text{m}$  is greater than  $0.1 \text{ cm}^{-3}$ . Figure 3 provides an example of  
 the detection of fog using this methodology and the associated reduction in  $N_{20}$  measured by the CPC.

Comparing this OPC-N3 fog classification to manual onsite observations reported at 00:00, 12:00 and 18:00 UTC daily,  
 185 the OPC-N3 does not detect fog when fog is reported by the observer (false negatives) in 35 out of 152 cases (23%). Six of



**Figure 3.** (a) Particle size distribution from the OPC-N3 from 31 July 2019 16:00 UTC to 01 August 2019 16:00 UTC (1 min averages). (b)  $N_{20}$  particle number concentration from the CPC during the same period (1 min averages). The duration of the fog event identified by the methodology described in Section 2.2 is shaded in green.

these cases can be attributed to inconsistent observer log entries or to logged issues with the OPC-N3, some others may result from discrepancies between the actual and reported observation time. However, false positive detection is rare, occurring in only 6 cases (1%). Therefore, although some fog events might be missed by the OPC-N3 fog classification, it is an accurate indicator of fog presence. The OPC-N3 was in operation between June and December 2019, and during this time the data are 190 96% complete.

### 2.3 Detection of precipitation and blowing snow

Below-cloud wet scavenging during snowfall can also reduce surface aerosol particle concentrations (e.g. Martin et al., 1980; Paramonov et al., 2011). A Precipitation Occurrence Sensor System (POSS) (Sheppard and Joe, 2008) located about 2 m above ground level at Summit measures the Doppler velocity spectrum of hydrometeors within a 1 m<sup>3</sup> sampling volume. 195 Surface snowfall rate retrieved from the POSS agrees well with retrievals from the lowest reliable range gate of the Millimeter Cloud Radar (MMCR) at Summit, with a root mean squared error of 0.08 mm h<sup>-1</sup> (Castellani et al., 2015). The ‘POSS power unit’ (the zeroth moment of the Doppler spectrum) can be used as a binary indicator of precipitation, and in this study we use a threshold of 2 POSS power units to identify precipitation events and exclude blowing snow, as per Pettersen et al. (2018). POSS data are 95% complete between June and December 2019.

200 The wind speed threshold for blowing snow (BLSN) varies depending on temperature and the properties of surface snow (Schmidt, 1982; Mann et al., 2000). In section 2.2 we used a 6 m s<sup>-1</sup> threshold as a minimum to avoid cases of possible blowing snow. However, to positively identify BLSN events we use a 10 m wind speed threshold of  $\geq 9$  m s<sup>-1</sup>. During 2019, on-site observers reported blowing or drifting snow 99% of the time when the 3 h mean was above this threshold.




## 2.4 Detection of surface-based temperature inversions

205 Surface-based temperature inversions (SBIs) occur at Summit in all seasons due to strong and persistent radiative cooling of the surface. SBIs are most common in the winter (Oct-Mar) where they occur over 70% of the time with a typical magnitude of  $\sim 5$  °C between 10 m and 2 m above the surface (Miller et al., 2013). In the summer (JJA), the amplitude of SBIs is weaker, and they only occur  $\sim 30\%$  of the time (Miller et al., 2013). SBIs limit the turbulent mixing of air (and aerosols) down to the surface, and as a result, aerosol particle concentrations measured at the surface may not be representative of concentrations at  
210 cloud level (Igel et al., 2017). To explore the effect of SBIs on surface aerosol particle concentrations in this study we classify SBI events where the 15 m minus the 2 m (above ground level) temperature difference must be greater than 3 °C ( $> 0.23$  °C  $m^{-1}$ ). Detection of SBI events is limited to June through October 2019 due to outages in the 15 m temperature sensor, but during this time data were 97% complete.

## 2.5 Aerosol source regions and transport pathways

215 The FLEXPART Lagrangian particle dispersion model (Pisso et al., 2019) is used to simulate aerosol transport pathways and surface emission sensitivities throughout 2019. FLEXPART simulations are run every 6 hours and driven by reanalysis data from ERA5, at the same horizontal and vertical resolution as the input data. In each simulation, 40,000 ‘particles’ are released at 100 m above the surface at Summit, and FLEXPART traces each particle back in time for 20 days. Particles follow the mean 3D wind field from ERA5 combined with a stochastic 3D turbulence field and parameterized convection (Forster et al., 2007).  
220 FLEXPART also simulates wet and dry deposition as linear decay constants based on a user input particle mean diameter, density, water and ice nucleation efficiency. In both cases deposition acts to reduce the total ‘mass’ of each particle, and a particle’s back trajectory stops when its mass reaches zero. Due to limited prior information about aerosols at Summit, we used the default aerosol tracer species; which assumes a particle mean diameter of  $0.25$   $\mu m$ , density of  $1400$   $kg\ m^{-3}$ , and water and ice nucleation efficiencies of 0.9 and 0.1 respectively. Particles of  $0.25$   $\mu m$  diameter are efficiently measured by the  
225 CPC at Summit (Fig. 2), fall within the typical size range of Arctic CCN (e.g. Jung et al., 2018), and have the relatively long atmosphere lifetimes necessary for advection over the GrIS ( $> 10$  days in the middle-upper troposphere; Jaenicke, 1990). FLEXPART outputs gridded emission sensitivity and supplementary back trajectory data that include the mean (centroid) back trajectory of all particles for each simulation, and the percentage of particles within the planetary boundary layer (PBL) at each time step. The surface emission sensitivity is proportional to the total amount of time that all particle back trajectories  
230 have spent near the surface (0-2,000 m) during the simulation period, representing the probability that aerosol particles emitted from each grid cell would have been detected at Summit at the simulation start time. We plot surface emission sensitivity as a percentage of the maximum value, to facilitate comparisons between figures.



**Figure 4.** Surface  $N_{20}$  from the CPC at Summit from February 2019 until May 2020.

### 3 Results

#### 3.1 Surface aerosol particle number concentrations at Summit, 2019-2020


235 The mean surface  $N_{20}$  in 2019 was  $129 \text{ cm}^{-3}$ , with the 6 h mean ranging between  $1 \text{ cm}^{-3}$  and  $1441 \text{ cm}^{-3}$  (Fig. 4). The minimum  $N_{20}$  in 2019 at Summit occurs in late February and early March, followed by a sharp increase of 2 orders of magnitude throughout March and April (Fig. 4). Between May and October, concentrations are fairly consistent and on the order of  $100 \text{ cm}^{-3}$  before decreasing again between October and December. Although data in early 2020 is limited, a similar increase in concentrations between February and May is apparent (Fig. 4).

#### 240 3.2 The effect of local surface processes on aerosol particle concentrations

The OPC-N3 identified 48 distinct fog events whilst it was operational between June and December 2019. The longest cumulative fog duration was in August (Fig. 5a) when fog was present for  $\sim 23\%$  of the month, consistent with previous multi year observations of supercooled liquid fogs at Summit (Cox et al., 2019). The mean duration of fog events was 3.3 hours and the longest event lasted 9.8 hours.

245 SBI events were also present in all months, and increased in total duration from summer to winter (Fig. 5b), again consistent with previous observations (Miller et al., 2013). The average duration of SBI events was 8.4 hours and the longest individual event lasted 5.8 days. SBI and fog events are not independent since fog condensate often forms due to surface cooling associated with the establishment of SBIs (e.g. Cox et al., 2019). Just under half all detected SBI events also contained fog (Fig. 5b), although because fog events are typically shorter, this only accounted for 17% of the total SBI duration.


250 Precipitation frequency and duration was highest in the summer and lowest in November and December (Fig. 5c). The average duration of precipitation events was 2.9 hours and the longest event lasted 14.1 hours. In contrast, BLSN events occurred most frequently in November and December, with an average duration of 6.9 hours (Fig. 5d). The seasonal distribution and duration of precipitation and BLSN events are also consistent with previous multi-year observations (Castellani et al., 2015;



**Figure 5.** Frequency and duration of (a) fog events, (b) surface-based temperature inversion events, (c) precipitation events and (d) blowing snow events, detected between June and December 2019 using the methodology described in Sections 2.2-2.4. Blue bars include all events and green bars show the change in distribution for SBI and precipitation events after the removal of events containing fog. (e) The distribution of  $N_{20}$  for the same months, excluding outliers.

Pettersen et al., 2018; Bennartz et al., 2019; Cox et al., 2019). Fog was detected during 23 of the 54 precipitation events (Fig. 255 5c). Because the OPC-N3 does not distinguish between fog and BLSN, it is not possible to determine how often fog might have been present during BLSN events. However, because of the high concentrations of ice crystals during blowing snow events, any supercooled water droplets are likely to be removed either through riming or the WBF process.


Figure 6 shows the median change in  $N_{20}$  during the first 3 hours of each event type. Only during fog events is there a consistent change: After three hours, the majority of fog events show a reduction in  $N_{20}$  by up to 35% (Fig. 6a). For SBI 260 events, there is very little discernible change in  $N_{20}$  during the first two hours (Fig. 6b). After  $\sim 140$  min there is a small median reduction in  $N_{20}$  that is not present when events that contain fog are omitted. During both precipitation and BLSN events, the median change in  $N_{20}$  remains close to zero (Fig. 6c, d).



**Figure 6.** The change in surface aerosol particle concentration (%) over time during the first 3 h of each event for (a) fog, (b) surface-based temperature inversions, (c) precipitation and (d) blowing snow events. The thick blue line and blue shading are the median and interquartile range of all events, the thick green line and green shading are the median and interquartile range for SBI and precipitation events that do not contain fog. The pink line indicates the total number of events at each time step for all events (solid) and excluding fog events (dashed).

### 3.3 Synoptic controls on surface aerosol particle concentrations

Here we explore the general relationship between  $N_{20}$  and synoptic conditions during 2019. Because both  $N_{20}$  and variables that change on synoptic timescales (i.e. surface pressure, geopotential height) vary seasonally, this seasonal dependence is removed prior to analysis. To calculate  $N_{20}$  anomalies we subtract the monthly median value for 2019. For all other variables (from GML-MET, 2020, and ERA5) anomalies are calculated by subtracting the 10 year (2009-2019) monthly mean climatology. Generally throughout 2019 anomalous changes in the 3 day mean surface pressure are in phase with anomalous 3 day median  $N_{20}$ , with some exceptions (Fig. 7a). To look at typical synoptic conditions associated with anomalous  $N_{20}$  at Summit, we look at ‘high’ and ‘low’  $N_{20}$  events, where the 3 day median  $N_{20}$  anomaly is greater than the 75th percentile or less than the 25th percentile respectively. To avoid oversampling, any events separated by less than four days are combined into a single event. The resulting high and low  $N_{20}$  events are highlighted in Fig. 7a, and are spread evenly throughout the annual cycle (15 high events and 14 low events).




**Figure 7.** (a) Surface pressure anomaly (3 day mean, solid black line) and  $N_{20}$  anomaly (3 day median, dashed) during 2019. Red and blue shading highlight high and low aerosol concentration events respectively (high/ low events are where the  $N_{20}$  anomaly is above the 75th/ below the 25th percentile). (b) The mean and interquartile range in surface pressure anomaly across all high (red) and low (blue)  $N_{20}$  events, for the 72 hours before and after the maximum (*minimum*)  $N_{20}$  for each high (*low*) event.

On average, an increase in surface pressure anomaly precedes anomalously high  $N_{20}$  events, with the maximum  $N_{20}$  coinciding with surface pressure anomalies leveling off (Fig. 7b). In contrast, a decrease in surface pressure anomaly precedes the majority of low  $N_{20}$  events, with the minimum  $N_{20}$  coinciding with the minimum surface pressure anomaly on average (Fig. 7b). Averaged over all high  $N_{20}$  events, 500 hPa geopotential heights are anomalously high (by over 75 m in central Greenland) and there is an anomalous anti-cyclonic circulation over the GrIS (Fig. 8a). In contrast, when averaged over the low  $N_{20}$  events, there is a region of anomalously low geopotential heights and anomalous cyclonic circulation centered on south-east Greenland (Fig. 8b).

FLEXPART simulations of surface emission sensitivity during the high  $N_{20}$  events show that sensitivity to surface emissions in the 20 days prior to detection at Summit outside of the ice sheet itself is rare (Fig. 9a), although there is some sensitivity to emissions from North America and Europe. Because there are no significant aerosol sources over the ice sheet itself, this implies that most of the simulated particles arriving at Summit during these events have been high in the atmosphere (>2,000 m agl) for over 20 days prior to detection at Summit. This is supported by the low percentages of simulated particles in the planetary boundary layer, and relatively high mean altitude of all simulated particles during the high  $N_{20}$  events (Fig. 9a). In contrast, the surface emission sensitivity during the low  $N_{20}$  events covers a broader area, encompassing coastal Greenland, Iceland, the Canadian Arctic and the intervening north Atlantic Ocean (Fig. 9b). There is a much higher percentage of simulated particles in the boundary layer in the week preceding detection at Summit during the low  $N_{20}$  events than during the high  $N_{20}$  events, and during the low  $N_{20}$  events the simulated particles are transported up to the highest point of the ice sheet from lower elevations (Fig. 9b).

### 3.4 Case studies of potential aerosol-limited cloud regimes at Summit


Fig. 4 demonstrates that  $N_{20}$  falls below  $10 \text{ cm}^{-3}$  in all seasons at Summit, suggesting that surface CCN concentrations fall below this threshold even more frequently. Given the existing evidence that aerosol particle concentrations this low can limit



**Figure 8.** ERA5 mean 500 hPa geopotential height anomaly (shaded) and 500 hPa horizontal wind anomalies (barbed) for all high  $N_{20}$  events (a) and low  $N_{20}$  events (b).

295 cloud formation elsewhere in the Arctic (Mauritsen et al., 2011; Stevens et al., 2018), we hypothesize that fog formation can be limited by low CCN concentrations over central Greenland, and if there are occasions where the surface aerosol particle concentration is representative of concentration at cloud height, that cloud formation can be limited by low CCN concentrations too. In this section we look in detail at three events where extremely low aerosol particle concentrations ( $N_{20} < 10 \text{ cm}^{-3}$  for  $> 3$  hours) coincided with cloud dissipation in the absence of fog, to look for further evidence of CCN-limited cloud regimes  
 300 at Summit. All times throughout the discussion of these case studies are given in UTC.

For each of the three cases considered (3 July 2019, 10 August 2019, and 21 November 2019), air is advected to the top of the ice sheet from different directions: On 3 July 2019, the primary aerosol source region is northern Siberia (Fig. 10a), on 10 August 2019, air approaches Summit from the north via the Canadian Arctic Archipelago (Fig. 10b), and on 21 November 2019, air approaches Summit from the south east, and is sensitive to emissions from northern Quebec (Fig. 10c). Two of the  
 305 three case studies (3 Jul 2019 and 21 November 2019) occur in the presence of anomalously low 500 hPa geopotential heights over south-east Greenland, with a stronger than usual south easterly wind component drawing air up the ice sheet from the south east coast (Fig. 10a and 10c). On both of these occasions  $> 50 \%$  of particles are within the PBL 4-6 days prior to arrival at Summit. On the 10 August 2019 case, there is an anomalous region of high 500 hPa geopotential heights over north-west Greenland and a stronger than usual northerly wind component over Summit (Fig. 10b). Although the FLEXPART simulated  
 310 particles remain closer to the ground for a longer period of time, the percentage of particles within the PBL in the 10 days prior




**Figure 9.** Results from FLEXPART back trajectory simulations averaged over the high  $N_{20}$  events (a) and the low  $N_{20}$  events (b). Upper: Surface emission sensitivity aggregated over 20 days prior to detection at Summit (as a percentage of the maximum value). Lower: Mean altitude and fraction of particles within the planetary boundary layer (PBL) for all simulated particles over the 10 days prior to detection.

to detection at Summit is much lower on 10 August 2019 than in the other two cases (Fig. 10b). On all three occasions, air is advected up to the ice sheet to Summit from lower elevations and spends  $> 1$  day prior to detection at Summit within the lowest 800 m above the surface of Greenland. The local conditions associated with each case are outlined below.

### 3.4.1 03 July 2019

315 On 02 July 2019,  $N_{20}$  dropped rapidly from  $\sim 200 \text{ cm}^{-3}$  to  $< 10 \text{ cm}^{-3}$  over a period of  $\sim 9$  h (Fig. 11a). The 12:00 equivalent potential temperature profile on 02 July 2019 shows that the lowest layer of broken stratocumulus cloud existed within a well mixed boundary layer (Fig. 11e). Shortly after 18:00,  $N_{20}$  dropped below  $10 \text{ cm}^{-3}$ , and there was a reduction in cloud cover (Fig. 11a-c). The on-site observer log recorded a transition from broken altocumulus at 18:00 to few clouds and unlimited visibility at 00:00 on 03 July 2019, despite the fact that the lowest 200 m above the surface remained saturated with respect  
 320 to water (Fig. 11f). On this occasion, the 00:00 radiosonde was launched from the surface at 23:15, and typically the weather observation is recorded at the time of launch. Photographs from a webcam viewing the aerosol inlet (taken every 15 minutes and orientated towards the eastern horizon) confirm that skies were clear and visibility was good at 23:15, but by 23:30 there was a clearly visible fog bow, indicating liquid fog droplets (Fig. 12). Notably on this occasion,  $N_{20}$  fell to  $< 10 \text{ cm}^{-3}$  in the absence of fog. At 00:15 visibility was obscured and the OPC-N3 detected fog at the surface (Fig. 12). The fact that the  
 325 OPC-N3 did not detect fog droplets until 00:15, despite an increase in LWP at 23:30 (Fig. 11d), could be explained by either a) the droplets forming the fog bow at 23:30 were too large to be detected by the OPC-N3 ( $> 40 \mu\text{m}$  diameter) or b) the fog




**Figure 10.** Synoptic anomaly plots and aerosol transport pathways during the three low  $N_{20}$  cases studies. Upper row: 500 hPa geopotential height and horizontal wind anomalies from ERA-5. Middle row: FLEXPART surface emission sensitivity (as a percentage of the maximum value) over the 10 days prior to aerosol detection at Summit. Lower row: FLEXPART mean aerosol transport height (back bold line) and percentage of particles within the planetary boundary layer (orange line) over the 10 days prior to detection at Summit. The shaded green area represents the mean height of topography beneath all particles.

was in the process of descending to the surface. In either case, both a) and b) support the hypothesis of Mauritsen et al. (2011), that in the absence of sufficient CCN, any existing CCN activate and grow to relatively large sizes, falling to the surface as ‘drizzle’.

330 The rapid transition from clear skies (despite a saturated surface layer) at 23:15 to fog at 00:15 coincided with  $N_{20}$  beginning to increase again (Fig. 11a). As  $N_{20}$  continued to increase, a thin low-level mixed-phase cloud returned and gradually lifted and thickened. By 12:00 on 03 July 2019,  $N_{20}$  had returned to  $\sim 200 \text{ cm}^{-3}$  and the lowest cloud layer had developed into a typical Arctic mixed-phase cloud (Shupe et al., 2006; Morrison et al., 2012; Shupe et al., 2013b) with a cloud top close to 500 m (Fig 11c), capping a well-mixed boundary layer (Fig. 11e).






**Figure 11.** Conditions during the 03 July 2019 low aerosol case study. (a) Surface  $N_{20}$  (CPC), with occurrences of fog (OPC-N3), precipitation (POSS), and SBI  $> 3^{\circ}\text{C}$  events indicated. (b) Radar reflectivity (MMCR). (c) Lidar depolarisation ratio (MPL), blue colours represent liquid droplets and reds are ice crystals. (d) Column integrated liquid water path and precipitable water vapour (MWR). (e) Temperature (dashed) and equivalent potential temperature (solid) radiosonde profiles. (f) Relative humidity with respect to water (solid) and ice (dashed) from radiosonde profiles. The coloured vertical lines on the left-hand plots correspond to the time of each vertical radiosonde profile in the right-hand plots.

### 335 3.4.2 10 August 2019

At 12:00 on 09 August 2019 there was a 2.5 km deep cloud over Summit (Fig. 13b) and surface  $N_{20}$  was  $\sim 100 \text{ cm}^{-3}$  (Fig. 13a). Between 14:00 and 18:00, there was a sharp decrease in liquid water path (Fig. 13d) as the cloud thinned until there was nothing detected by the radar at 19:00 (Fig. 13b). At this time, the surface-based temperature inversion strengthened to



**Figure 12.** Photographs from a webcam oriented towards the eastern horizon on 02-03 July 2019. The aerosol inlet is visible mounted on the mast in the SE corner.

$> 0.23 \text{ } ^\circ\text{C m}^{-1}$  (Fig. 13a), likely due to the increase in longwave cooling at the surface after the reduction in cloud cover.

340 In this case, it was only after the strengthening of the surface temperature inversion that  $N_{20}$  began to decrease. At 21:40 the OPC-N3 detected fog, and  $N_{20}$  decreased more rapidly, falling below  $10 \text{ cm}^{-3}$  at 23:35 and reaching a minimum of  $0.5 \text{ cm}^{-3}$  at 01:00 on 10 August 2019 after which the fog thinned and cleared (Fig. 13a).  $N_{20}$  began to rise again from 02:15, and when the particle concentration increased above  $10 \text{ cm}^{-3}$  at 04:00, there was a sudden sharp increase in liquid water path (Fig. 13d), and a thin low mixed-phase cloud developed (Fig. 13c). The cloud thickened as  $N_{20}$  continued to increase back to  $\sim 100 \text{ cm}^{-3}$


345 at 07:00. Fig. 14 shows the transition from cloudy to clear skies, then to thin fog and back to overcast again throughout this event.

### 3.4.3 21 November 2019

Surface  $N_{20}$  decreased from  $50 \text{ cm}^{-3}$  at 06:00 on 20 November 2019 to a minimum of  $0.5 \text{ cm}^{-3}$  at 06:30 on 21 November 2019, and remained below  $10 \text{ cm}^{-3}$  for a total of 24 hours (Fig. 15a). As  $N_{20}$  decreased, a low-level mixed-phase cloud thinned

350 and liquid water path fell to  $0 \text{ g m}^{-2}$  by 09:00 (Fig. 15c,d). The 20 November 12:00 radiosonde shows that the boundary layer was neutrally stratified up to about 300 m, above a very shallow stable surface layer (where the air temperature increased  $7 \text{ } ^\circ\text{C}$  in the 4 m immediately above the surface) (Fig. 15e). At 00:00 on 21 November 2019 the temperature inversion in the lowest 4 m of the atmosphere strengthened to  $12 \text{ } ^\circ\text{C}$ , the sky above Summit was clear, and  $N_{20}$  continued to fall until 06:00. At 12:00 on 21 November 2019 a 3 km deep ice cloud moved across Summit (Fig. 15b,c) and  $N_{20}$  began to increase again

355 (Fig. 15a). Liquid water path initially remained close to zero but increased sharply when  $N_{20}$  rose above  $10 \text{ cm}^{-3}$  at 00:00 on




**Figure 13.** Same as Fig. 11 but during the 10 August 2019 low aerosol case study.

22 November 2019. Between the 20 November 2019 12:00 and the 22 November 2019 00:00 radiosonde profile, the 3 km agl potential temperature decreased by  $> 5$  °C (Fig. 15e), possibly indicating an air mass transition during this period.

## 4 Discussion

### 4.1 The seasonal cycle of surface aerosol particle concentrations at Summit

360 Despite differences in the measured size ranges, the  $N_{20}$  reported in the present study are of the same order of magnitude as previous summertime measurements of condensation nuclei at Summit (100-500  $\text{cm}^{-3}$  in the first week of July 1992, Bergin et al., 1994), and from DYE III on the south east GrIS ( $\sim 6$ -1000  $\text{cm}^{-3}$  in July and August 1982, Hogan et al., 1984). These




**Figure 14.** Photographs from a webcam oriented towards the eastern horizon on 09-10 August 2019. The aerosol inlet is visible mounted on the mast in the SE corner.

results are also comparable in magnitude to  $N_{10}$  concentrations measured at other Arctic stations ( $\sim 1\text{-}2000\text{ cm}^{-3}$  at Utqiagvik, Alaska, and  $\sim 5\text{-}3000\text{ cm}^{-3}$  at Pallas, Finland; Asmi et al., 2013), however the seasonal cycle is notably different. The seasonal cycle in surface aerosol particle concentration at many sea level Arctic sites is dominated by the cycle of ‘Arctic haze’ (e.g. Shaw, 1995); where anthropogenic pollutants build up in the winter resulting in maximum aerosol particle concentrations in early spring, followed by a sharp reduction of particles in the summer. In contrast, at Summit we see minimum surface aerosol particle concentrations in the winter and higher concentrations in the late spring and Summer. The seasonal cycle at Summit in 2019 is consistent with multi year seasonal cycles of mineral particles in snowpit samples at Summit (Drab et al., 2002), and with measurements of bulk aerosol light scattering and absorption coefficients that are related to aerosol particle concentrations (Schmeisser et al., 2018), suggesting that it is a persistent annual feature.

Hirdman et al. (2009) used FLEXPART back trajectory simulations to show that surface aerosol particle concentrations at Summit are an order of magnitude less sensitive to surface emissions from within the Arctic compared to lower altitude Arctic sites, which is a possible explanation for why Summit does not experience Arctic haze build up during the winter. In contrast, the GrIS is more sensitive to aerosol sources above the boundary layer, often originating further south and descending to the GrIS via subsidence driven by radiative cooling (Stohl, 2006; Hirdman et al., 2009).

At sea level Arctic sites (both marine and coastal), the extremely low aerosol particle concentrations observed in the summer are largely attributed to increases in wet deposition (Garrett et al., 2010; Browse et al., 2012). An important distinction between Summit (where the  $0\text{ }^{\circ}\text{C}$  isotherm is always below the surface except in extreme situations; Shupe et al., 2013b; Bennartz et al., 2013) and sea level Arctic sites, is that Summit does not currently experience rain during the summer. The fact that mean




**Figure 15.** Same as Fig. 11 but during the 21 November 2019 low aerosol case study. Note that there are no SBI ‘events’ recorded during this period due to missing data, but that the radiosonde profiles indicate a constant shallow surface-based temperature inversion throughout.

monthly  $N_{20}$  is relatively high in the summer at Summit could be related to the fact that wet deposition is much less efficient in ice-bearing clouds (Henning et al., 2004). In this case, future increases in the height of the  $0^\circ\text{C}$  isotherm over the GrIS could result in lower summertime aerosol particle concentrations.

## 4.2 Controls on surface aerosol particle concentrations at Summit

385 The processes controlling surface aerosol particle concentrations over the central GrIS form a complex system, integrating local meteorological conditions, air mass history during aerosol transport, source regions and transport pathways. Fig. 16 illustrates some of the key components of this system, distinguishing between those processes that are supported by evidence in this study, and those for which uncertainties still remain. We have made the assumption that there are no local sources of aerosol



**Figure 16.** Conceptual model illustrating the key components controlling surface aerosol particle concentrations over the central GrIS, highlighting factors that are supported by evidence in this study and important areas for future research (see legend inset).

at the surface. There is a possibility that particle growth via condensation of precursor gases, possibly released from organic material in the snowpack, could occasionally contribute to near surface CCN concentrations (Ziemba et al., 2010). We do not consider this process in the present study, but the contribution of ultra-fine particle growth to CCN concentrations over the GrIS remains unclear and warrants further investigation.

Out of the four surface processes considered in this study (fog, SBIs, precipitation and BLSN), only fog events have a strong and consistent effect on measured  $N_{20}$ . The effect of fog on surface aerosol particle concentrations is consistent with previous studies that were limited to the summer months (Bergin et al., 1994, 1995). Future studies should ensure that fog is accounted for before generalising sampled particle concentrations over wider regions or altitude ranges. Importantly, the observed decrease in aerosol particle number concentration during fog events reflects the number of particles that are incorporated into droplets too large to pass through the CPC inlet, either through CCN activation and growth or scavenging by fog droplets, and these particles are not necessarily deposited at the surface.

Despite the potential for SBIs to act as a barrier for turbulent mixing and hence reduce the rate that aerosol particles are transported down to the surface (Dibb et al., 1992; Li et al., 2019; Thomas et al., 2019), we found no consistent change in  $N_{20}$  during the first 3 hours of SBI events, and no relationship between the change in  $N_{20}$  and the mean intensity of the SBI, which ranges between  $0.23 \text{ }^{\circ}\text{C m}^{-1}$  and  $0.92 \text{ }^{\circ}\text{C m}^{-1}$  (not shown). SBIs may have a more important role on surface aerosol particle concentrations over longer timescales, especially because the loss of aerosol particles to the surface by dry deposition is slow (Garrett et al., 2010), but because fog regularly forms during SBI events, it is difficult to isolate the influence of the SBI from the influence of fog scavenging on aerosol particle concentrations during longer events. SBIs may also contribute to observed reduction in  $N_{20}$  during fog events by restricting turbulent mixing. This study does not consider changes in mechanically

induced turbulence over time or elevated temperature inversions, further studies are required to understand the role of changes in turbulent mixing on controlling surface aerosol particle concentrations.

410  $N_{20}$  also does not respond consistently to the precipitation or BLSN events considered in this study (Fig. 6c,d). This is in agreement with Bergin et al. (1995), who did not observe a significant effect of precipitation on surface aerosol particle concentrations at Summit during the summer. below-cloud scavenging rates are sensitive to a wide range of parameters that we do not consider here, including snow crystal size and habit, degree of riming, relative humidity and Reynolds number (Feng, 2009; Browse et al., 2012). We also do not distinguish between below-cloud precipitation and clear sky precipitation (diamond  
415 dust). However, although the rate of wet deposition might vary between events, below-cloud scavenging should reduce  $N_{20}$ , and the fact that we do not consistently observe this suggests that other processes are acting to maintain surface aerosol particle concentrations during precipitation. For example, in both the 03 July 2019 and 21 November 2019 case studies (Section 3.4), aerosol particle concentrations increase during precipitation. One explanation for this could be the release of particles near the surface via below-cloud evaporation of hydrometeors. Low-level mixed-phase clouds in particular can act to facilitate the  
420 transport of particles from the free troposphere into the boundary layer through entrainment and activation at cloud top and release through evaporation at cloud base (Igel et al., 2017). At Summit, the majority of precipitation occurs in the presence of these low-level mixed-phase clouds (Pettersen et al., 2018). Clouds can also act to increase the efficiency of turbulent mixing down to the surface because below-cloud turbulent mixing driven by the sinking of radiatively cooled air near the cloud top can extend down into the mechanically driven surface mixed layer (Brooks et al., 2017; Shupe et al., 2013a). Precipitation  
425 itself might also contribute to this increase in turbulent mixing via drag. Given that most of the aerosol particles arriving at Summit descend from the free troposphere (Hirdman et al., 2009), the role of clouds in the transport of aerosol particles into the boundary layer of the GrIS is an important area for future research.

Synoptic conditions play an important role in controlling  $N_{20}$  at Summit, with most anomalously high  $N_{20}$  events coinciding with anomalously high surface pressure during 2019 (Fig. 7). The difference in aerosol emission sensitivity and transport  
430 pathway simulations between anomalously high and low  $N_{20}$  events (Fig. 9), combined with the difference in mean synoptic anomalies (Fig. 8), imply that high surface aerosol particle concentrations occur at Summit when air is transported down to the surface from high in the free troposphere, associated with subsidence related to anomalous strong high pressure systems over Greenland (Fig. 16b). This free tropospheric air is sensitive to emissions from mid and low latitudes that can release particles high into the atmosphere where they remain suspended for long periods of time (i.e. > 20 days) (Stohl, 2006; Hirdman et al.,  
435 2009; Roiger et al., 2011). This result is consistent with previous studies investigating the transport pathways of aerosol particles that arrive at Summit. Both Hirdman et al. (2009) and Schmeisser et al. (2018) conclude that because on average the majority of air arriving at Summit has only been in contact with the surface over the ice sheet itself, particles measured at Summit must have descended from the free troposphere after transportation at high altitudes over time scales > 20 days. Persistent anomalously high geopotential heights over central Greenland are also associated with the occurrence of precipitating low-  
440 level mixed-phase stratocumulous clouds (McIlhattan et al., 2020) that can encourage the transport of aerosol particles from the free tropospheric into the boundary layer (Igel et al., 2017).

Anomalously low aerosol particle concentrations typically occur in the presence of anomalous cyclonic circulation and low geopotential heights off the south east coast of Greenland (Fig. 8b) that drive air up to the top of the GrIS from the coast and surrounding ocean (Fig. 9b, 16b). During such events, adiabatic cooling due to orographic lifting as the air is advected up the GrIS results in increased condensation and precipitation (Schuenemann et al., 2009). These conditions are associated with deep glaciated clouds advecting over Summit from south east Greenland (Pettersen et al., 2018). The associated increase in wet deposition en-route to Summit could contribute to the relatively low  $N_{20}$ . These events are more common in the winter season, when the north Atlantic storm track is more active (Schuenemann et al., 2009). Hogan et al. (1984) also reached a similar conclusion based on surface aerosol particle measurements at DYE III during the summer; they observed that low particle concentrations followed moist upslope flow and precipitation driven by a low pressure system to the south of Greenland, and that concentrations increased after the establishment of a high pressure system and downslope flow.

This study does not consider changes in emission or removal rates along the aerosol transport pathway. Emission rates vary seasonally within the Arctic due to changes in ice cover and biomass burning (Willis et al., 2018), and isolated events such as volcanic eruptions can have large impacts on background aerosol particle concentrations (e.g. Friberg et al., 2015). Removal rates vary along a particular transport pathway with changes in precipitation amount and phase (Garrett et al., 2010; Browse et al., 2012). Given this, it is quite remarkable that the relationship between anomalous aerosol particle concentrations at Summit and anomalous synoptic conditions is so evident. The strength of this relationship implies that future changes in Arctic large scale circulation could effect aerosol particle concentrations and aerosol-cloud-radiation interactions over the GrIS. In particular, changes in the frequency of storms moving up the southeast coast of Greenland (Ulbrich et al., 2008) or the position of the Icelandic low (Berdahl et al., 2018) might effect the frequency of extremely low aerosol particle concentration events over the central GrIS.

### 4.3 Potential for cloud formation to be limited by low CCN concentrations and discussion of case studies

$N_{20}$  fell below  $10 \text{ cm}^{-3}$  on multiple occasions year-round at Summit in 2019 (Fig. 4). Because CCN are a subset of total condensation nuclei concentration, it is likely that CCN concentrations fall low enough to limit cloud and fog formation, based on approximate threshold estimates determined from past observational and modelling studies over the Arctic Ocean (Mauritsen et al., 2011; Stevens et al., 2018). The ratio of  $\text{CCN}/N_{10}$  at a supersaturation of 0.55% measured over Summit during a research flight in 2008 was 0.52 (Latham et al., 2013), which is similar to the mean  $\text{CCN}/N_{10}$  ratio observed at the Zeppelin Observatory in Svalbard outside of the Arctic haze season (Jung et al., 2018). However this ratio is a function of supersaturation, and at very high supersaturations (that can occur under extremely low CCN concentrations) small particles that do not typically act as CCN can activate (Leaith et al., 2016; Baccarini et al., 2020). If we make the assumption that all CCN are activated after the first 3 hours of fog formation during the events in Fig. 6a, the fact that we see a median 20% reduction in total  $N_{20}$  during these events implies a  $\text{CCN}/N_{20}$  ratio of 0.2, and for the individual event example in Fig. 3, the  $\text{CCN}/N_{20}$  would have been 0.46. Using the more conservative ratio estimation of 0.46, surface CCN concentrations will have fallen below  $10 \text{ cm}^{-3}$  for 46 days or 15% of the measurement period during 2019. Because supercooled liquid fog can have a large effect on surface radiative fluxes at Summit (liquid fog at Summit has an average total (SW+LW) cloud radiative forcing



of  $26.1 \text{ W m}^{-2}$  compared to clear skies, Cox et al., 2019), if fog formation is limited by low CCN concentrations, this could have an important effect on the ice sheet surface energy budget, especially over individual events which can play a role in pre-conditioning the snow surface in advance of melt (Miller et al., 2017). The same could be true for clouds where surface concentrations are representative of CCN concentrations at cloud level. For example, the exceptional July 2012 Greenland melt event was enhanced by the presence of low-level mixed-phase clouds with a LWP of  $\sim 30 \text{ g m}^{-2}$  (Bennartz et al., 2013), in this case, if small changes in CCN concentrations acted to either increase or decrease the cloud LWP, they could have controlled the presence versus absence of surface melt.

For all three of the case studies in section 3.4, decreasing aerosol particle concentrations were associated with a reduction in cloud cover, and the reverse was also true. However, differences in timing and boundary layer structure imply that different processes were involved in each case. This demonstrates that it is not sufficient to use simple correlations between cloud properties and aerosol particle concentrations to investigate cloud-aerosol interactions, since there are many additional confounding variables. Although we cannot delineate the individual drivers of the changes in  $N_{20}$  during these case studies based purely on observations, the near-zero liquid water path is convincing evidence that low CCN concentrations are limiting the formation of liquid water droplets at the surface despite supersaturation when  $N_{20}$  is  $< 10 \text{ cm}^{-3}$  in all three case studies (i.e. fog formation). Note the only other events where we observed  $N_{20} < 10 \text{ cm}^{-3}$  and  $\text{LWP} > 10 \text{ g m}^{-2}$  occurred in February and March 2019 and were associated with clouds with base heights between 250 and 1,000 m above the surface. The static stability of the surface layer in these cases means it is not possible to know whether the surface  $N_{20}$  was representative of aerosol particle concentrations in the cloud layer.

Finally, for all three case studies, back trajectory simulations indicate that aerosol particles were transported upslope to Summit from lower elevations (Fig. 10), and two of the cases (July and November) occurred in the presence of cyclonic circulation off the southeast coast of Greenland - the typical synoptic condition associated with anomalously low aerosol particle concentrations at Summit (Fig. 8). Although the simulated aerosol source regions are all from high latitudes ( $> 50^\circ\text{N}$ ; Fig. 10), they originate from very different directions (Siberia on 3 July 2019, the Canadian Archipelago on 10 August 2019, and south west of Greenland on 21 November 2019). This suggests that the upslope transport pathway to Summit, which is strongly linked to precipitation over the GrIS (Schuenemann et al., 2009) and notably from glaciated as opposed to mixed-phase clouds (Pettersen et al., 2018), has a stronger influence on  $N_{20}$  than source region. These upslope flow enhanced precipitation events are also coupled to anomalously warm temperatures over the GrIS, which likely results in a higher percentage of rain (and hence increased wet deposition) en-route to Summit (Pettersen et al., in review). These results imply that increased wet deposition during transport may play a large role in driving CCN concentrations below the threshold where they can sustain cloud formation. The role of wet deposition in controlling aerosol particle concentrations over the central GrIS is therefore an important area for future research.

## 5 Summary and conclusions

This study presents the first full year of surface aerosol particle number concentration measurements from the central Greenland Ice Sheet and assesses the local and synoptic controls on surface  $N_{20}$ . In 2019, the minimum aerosol particle concentrations occur in February (which has a monthly average concentration of just  $18 \text{ cm}^{-3}$ , and a standard deviation,  $\sigma_N = 16 \text{ cm}^{-3}$ ), and the maximum concentrations occur in April and May (monthly average concentrations of  $247 \text{ cm}^{-3}$ ,  $\sigma_N = 130 \text{ cm}^{-3}$  and  $206 \text{ cm}^{-3}$ ,  $\sigma_N = 165 \text{ cm}^{-3}$ , respectively). Between May and October, concentrations remain on the order of  $100 \text{ cm}^{-3}$  before they decrease again between October and December. This seasonal cycle is distinct from that of many sea level Arctic sites which experience minimum surface aerosol concentrations in the summer (Freud et al., 2017; Schmeisser et al., 2018).

Changes in synoptic conditions strongly control  $N_{20}$ , with almost all anomalously high  $N_{20}$  events associated with anomalously high surface pressure over Summit. High  $N_{20}$  occurs under anomalously high geopotential heights and strong anticyclonic circulation over Greenland, which act to enhance the descent of free tropospheric air to the ice sheet surface. Low  $N_{20}$  occurs in the presence of anomalous cyclonic circulation over south east Greenland, when low pressure systems drive up slope flow that is associated with increased precipitation (Schuenemann et al., 2009; Pettersen et al., 2018). Below average aerosol particle concentrations occur more often in the winter, when the frequency of low pressure systems driven by the North Atlantic storm track increases (Schuenemann et al., 2009). The distinction between upslope flow and descent from higher altitudes appears to be a stronger control on  $N_{20}$  than aerosol source region, suggesting an important role for wet deposition along aerosol transport pathways.

We find that fog strongly effects surface aerosol particle concentration measurements, in agreement with previous studies that look at isolated events during the summer (Bergin et al., 1994, 1995). On average, there is a 20% reduction in  $N_{20}$  after the first 3 h of a fog event. Because fog significantly modifies local surface aerosol particle concentrations, future studies should ensure that fog is accounted for before generalising sampled aerosol particle concentrations over wider regions or altitude ranges. In contrast, precipitation, blowing snow, and strong surface-based temperature inversions ( $> 0.23 \text{ }^\circ\text{C m}^{-1}$ ) do not have a consistent effect on  $N_{20}$  during the first 3 h of the event. Competing influences of advection, or either cloud or mechanically induced changes in the turbulent structure of the boundary layer, might play roles in modulating aerosol particle concentrations during these events and are not considered in this study.

This study uses a conservative estimate to determine that surface aerosol particle concentrations low enough to limit cloud and or fog formation (based on observations and model simulations over the Arctic ocean; Mauritsen et al., 2011; Stevens et al., 2018) do occur in both winter and summer over the central GrIS. However, long term vertical profiles of CCN concentrations are necessary to determine how often this is relevant at cloud height. Although practically difficult, continuous vertical profiles of aerosol particle concentrations above the GrIS are essential for understanding the interaction between clouds, aerosols, and the ice sheet surface energy budget, and should be a priority for future campaigns. Vertical aerosol profiles are particularly important over the central GrIS where most of the aerosol particles arriving at the surface descend from higher elevations in the free troposphere (Hirdman et al., 2009; Schmeisser et al., 2018, this study). The unique transport pathway and resulting seasonal cycle of aerosol particles over the central GrIS demonstrate that observations of aerosol properties at sea level Arctic

sites cannot be generalised over the GrIS, in agreement with previous studies (e.g. Hirdman et al., 2009; Schmeisser et al., 2018; Schmale et al., 2021).

*Data availability.* All data are publicly available. ICECAPS-ACE aerosol measurements and multi level temperature sensor data can be accessed through the CEDA archive at <http://catalogue.ceda.ac.uk/uuid/f06c6aa727404ca788ee3dd0515ea61a>. NOAA GML meteorological data are available at <ftp://aftp.cmdl.noaa.gov/data/645met/sum/> (last access: March 2021). All additional ICECAPS data are available from the Arctic Data Center: MMCR (doi:10.18739/A2Q52FD4V), MPL (doi:10.18739/A2862BC30), POSS (doi:10.18739/A2GQ6R30G), HATPRO and MWRHF (doi:10.18739/A2TX3568P) and radiosonde profiles (doi:10.18739/A20P0WR53). ERA5 reanalysis data are made available by the European Centre for Medium-Range Weather Forecasts (ECMWF) and can be accessed at the Copernicus Climate Data Store, DOI: 10.24381/cds.bd0915c6.

*Author contributions.* The original ICECAPS project proposal was conceived by MDS, VPW, DDT and RB. HG led data collection and curation of ICECAPS-ACE aerosol data supervised by RRN and IMB. HG led the analysis with contributions from RRN, IMB, BJM, CP, MDS and CJC. HG prepared the manuscript with contributions from all co-authors.

*Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* The efforts of technicians at Summit Station and science support provided by Polar Field Services were crucial to maintaining data quality and continuity at Summit. We acknowledge Bethany Wyld for assistance during the ICECAPS-ACE field installation, and Richard Rigby for his support setting up and running the FLEXPART model. We also gratefully acknowledge support from the National Centre of Atmospheric Science (NCAS), the NCAS Atmospheric Measurement and Observation Facility, and notably Dr. Barbara Brooks for support in instrument troubleshooting and data quality control. ICECAPS is a long-term research program with a large number of collaborators and we are grateful for all their efforts in developing and maintaining the various instruments and data products used in this study. Financial support for ICECAPS-ACE was provided by NSFGEO-NERC grant 1801477, and HG was funded by the NERC SPHERES DTP grant number NE/L002574/1. MDS and WDN were supported by the National Science Foundation grant OPP1801477. We are grateful for the thoughtful suggestions of two anonymous reviewers that have improved the quality of this manuscript.

## References

- Asmi, A., Collaud Coen, M., Ogren, J., Andrews, E., Sheridan, P., Jefferson, A., Weingartner, E., Baltensperger, U., Bukowiecki, N., Li-  
565 havainen, H., et al.: Aerosol decadal trends–Part 2: In-situ aerosol particle number concentrations at GAW and ACTRIS stations, *Atmospheric Chemistry and Physics*, 13, 895–916, 2013.
- Baccarini, A., Karlsson, L., Dommen, J., Duplessis, P., Vüllers, J., Brooks, I. M., Saiz-Lopez, A., Salter, M., Tjernström, M., Baltensperger, U., et al.: Frequent new particle formation over the high Arctic pack ice by enhanced iodine emissions, *Nature communications*, 11, 1–11, 2020.
- 570 Backman, J., Schmeisser, L., and Asmi, E.: Asian emissions explain much of the Arctic Black Carbon events, *Geophysical Research Letters*, 2021.
- Bamber, J. L., Westaway, R. M., Marzeion, B., and Wouters, B.: The land ice contribution to sea level during the satellite era, *Environmental Research Letters*, 13, 2018.
- Bennartz, R., Shupe, M. D., Turner, D. D., Walden, V. P., Steffen, K., Cox, C. J., Kulie, M. S., Miller, N. B., and Pettersen, C.: July 2012  
575 Greenland melt extent enhanced by low-level liquid clouds, *Nature*, 496, 83, 2013.
- Berdahl, M., Rennermalm, A., Hammann, A., Mioduszewski, J., Hameed, S., Tedesco, M., Stroeve, J., Mote, T., Koyama, T., and McConnell, J. R.: Southeast Greenland winter precipitation strongly linked to the Icelandic Low position, *Journal of Climate*, 31, 4483–4500, 2018.
- Bergin, M. H., Jaffrezo, J. L., Davidson, C. I., Caldow, R., and Dibb, J.: Fluxes of chemical species to the Greenland ice sheet at Summit by fog and dry deposition, *Geochimica et Cosmochimica Acta*, 58, 3207–3215, 1994.
- 580 Bergin, M. H., Jaffrezo, J.-L., Davidson, C. I., Dibb, J. E., Pandis, S. N., Hillamo, R., Maenhaut, W., Kuhns, H. D., and Makela, T.: The contributions of snow, fog, and dry deposition to the summer flux of anions and cations at Summit, Greenland, *Journal of Geophysical Research*, 100, 16275, 1995.
- Box, J. E., Fettweis, X., Stroeve, J. C., Tedesco, M., Hall, D. K., and Steffen, K.: Greenland ice sheet albedo feedback: thermodynamics and atmospheric drivers, *The Cryosphere*, 6, 821–839, 2012.
- 585 Brooks, I. M., Tjernström, M., Persson, P. O. G., Shupe, M. D., Atkinson, R. A., Canut, G., Birch, C. E., Mauritsen, T., Sedlar, J., and Brooks, B. J.: The Turbulent Structure of the Arctic Summer Boundary Layer During The Arctic Summer Cloud-Ocean Study, *Journal of Geophysical Research: Atmospheres*, 122, 9685–9704, 2017.
- Browse, J., Carslaw, K. S., Arnold, S. R., Pringle, K., and Boucher, O.: The scavenging processes controlling the seasonal cycle in Arctic sulphate and black carbon aerosol, *Atmos. Chem. Phys*, 12, 6775–6798, 2012.
- 590 Castellani, B. B., Shupe, M. D., Hudak, D. R., and Sheppard, B. E.: The annual cycle of snowfall at Summit, Greenland, *Journal of Geophysical Research: Atmospheres*, 120, 6654–6668, 2015.
- Cheng, C.-T., Wang, W.-C., and Chen, J.-P.: Simulation of the effects of increasing cloud condensation nuclei on mixed-phase clouds and precipitation of a front system, *Atmospheric Research*, 96, 461–476, 2010.
- Cox, C. J., Noone, D. C., Berkelhammer, M., Shupe, M. D., Neff, W. D., Miller, N. B., Walden, V. P., and Steffen, K.: Supercooled liquid  
595 fogs over the central Greenland Ice Sheet, *Atmospheric Chemistry and Physics*, 19, 7467–7485, 2019.
- Crilley, L. R., Shaw, M., Pound, R., Kramer, L. J., Price, R., Young, S., Lewis, A. C., and Pope, F. D.: Evaluation of a low-cost optical particle counter (Alphasense OPC-N2) for ambient air monitoring, *Atmospheric Measurement Techniques*, 11, 709–720, 2018.
- Curry, J. A., Rossow, W. B., Randall, D., Schramm, J. L., Rossow, W. B., Randall, D., and Schramm, J. L.: Overview of Arctic cloud and radiation characteristics, *Journal of Climate*, 9, 1731–1764, 1996.

- 600 Davidson, C., Jaffrezo, J.-L., Mosher, B., Dibb, J. E., Borys, R., Bodhaine, B., Rasmussen, R., Boutron, C., Gurlach, U., Cachier, H., et al.:  
Chemical constituents in the air and snow at Dye 3, Greenland—I. Seasonal variations, *Atmospheric Environment. Part A. General Topics*,  
27, 2709–2722, 1993.
- Dibb, J. E., Jaffrezo, J. L., and Legrand, M.: Initial findings of recent investigations of air-snow relationships in the summit region of the  
Greenland ice sheet, *Journal of Atmospheric Chemistry*, 14, 167–180, 1992.
- 605 Drab, E., Gaudichet, A., Jaffrezo, J. L., and Colin, J. L.: Mineral particles content in recent snow at Summit (Greenland), *Atmospheric  
Environment*, 36, 5365–5376, 2002.
- Enderlin, E. M., Howat, I. M., Jeong, S., Noh, M.-J., VanAngelen, J. H., and Van den Broeke, M. R.: An improved mass budget for the  
Greenland ice sheet, *Geophysical Research Letters*, pp. 866–872, 2014.
- Feng, J.: A size-resolved model for below-cloud scavenging of aerosols by snowfall, *Journal of Geophysical Research Atmospheres*, 114,  
610 1–8, 2009.
- Fettweis, X., Box, J. E., Agosta, C., Amory, C., Kittel, C., Lang, C., Van As, D., Machguth, H., and Gallée, H.: Greenland ice sheet surface  
mass balance using the regional climate MAR model, *The Cryosphere*, 11, 20–22, 2017.
- Flyger, H., Heidam, N., Hansen, K., Megaw, W., Walther, E., and Hogan, A.: The background level of the summer tropospheric aerosol,  
sulphur dioxide and ozone over Greenland and the North Atlantic Ocean, *Journal of Aerosol Science*, 7, 103–140, 1976.
- 615 Flynn, C. J., Mendoza, A., Zheng, Y., and Mathurb, S.: Novel polarization-sensitive micropulse lidar measurement technique, *Optics  
express*, 15, 2785–2790, 2007.
- Forster, C., Stohl, A., and Seibert, P.: Parameterization of Convective Transport in a Lagrangian Particle Dispersion Model and Its Evaluation,  
*Journal of Applied Meteorology and Climatology*, 46, 403–422, 2007.
- Freud, E., Krejci, R., Tunved, P., Leaitch, R., Nguyen, Q. T., Massling, A., Skov, H., and Barrie, L.: Pan-Arctic aerosol number size distribu-  
620 tions: seasonality and transport patterns, *Atmospheric Chemistry and Physics*, 17, 8101–8128, 2017.
- Friberg, J., Martinsson, B. G., Sporre, M. K., Andersson, S. M., Brenninkmeijer, C. A. M., Hermann, M., Velthoven, P. F. J., and Zahn, A.:  
Influence of volcanic eruptions on midlatitude upper tropospheric aerosol and consequences for cirrus clouds, *Earth and Space Science*,  
2, 285–300, 2015.
- Garrett, T. J. and Zhao, C.: Increased Arctic cloud longwave emissivity associated with pollution from mid-latitudes, *Nature*, 440, 787–789,  
625 2006.
- Garrett, T. J., Zhao, C., and Novelli, P. C.: Assessing the relative contributions of transport efficiency and scavenging to seasonal variability  
in Arctic aerosol, *Tellus, Series B: Chemical and Physical Meteorology*, 62, 190–196, 2010.
- Giorgi, F.: A particle dry-deposition parameterization scheme for use in tracer transport models, *Journal of Geophysical Research*, 91, 9794,  
1986.
- 630 GMLMET: Meteorology Measurements from the NOAA/ESRL/GMD Baseline Observatory Summit Station, [ftp://ftp.cmdl.noaa.gov/data/  
met/sum/](ftp://ftp.cmdl.noaa.gov/data/met/sum/), 2021.
- Guy, H., Neely III, R. R., and Brooks, I.: ICECAPS-ACE: Integrated Characterization of Energy, Clouds, Atmospheric state, and Precipitation  
at Summit, Greenland - Aerosol Cloud Experiment measurements., Centre for Environmental Data Analysis, [http://catalogue.ceda.ac.uk/  
uuid/f06c6aa727404ca788ee3dd0515ea61a](http://catalogue.ceda.ac.uk/uuid/f06c6aa727404ca788ee3dd0515ea61a), 2020.
- 635 Hanna, E., Cropper, T. E., Hall, R. J., and Cappelen, J.: Greenland Blocking Index 1851-2015: a regional climate change signal, *International  
Journal of Climatology*, 36, 4847–4861, 2016.

- Heinemann, G. and Klein, T.: Modelling and observations of the katabatic flow dynamics over Greenland, *Tellus A: Dynamic Meteorology and Oceanography*, 54, 542–554, 2002.
- Henning, S., Bojinski, S., Diehl, K., Ghan, S., Nyeki, S., Weingartner, E., Wurzler, S., and Baltensperger, U.: Aerosol partitioning in natural  
640 mixed-phase clouds, *Geophysical Research Letters*, 31, 2004.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut,  
645 J. N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, 2020.
- Hirdman, D., Sodemann, H., Eckhardt, S., Burkhardt, J. F., Jefferson, A., Mefford, T., Quinn, P. K., Sharma, S., Ström, J., and Stohl, A.: Source identification of short-lived air pollutants in the Arctic using statistical analysis of measurement data and particle dispersion model output, *Atmospheric Chemistry and Physics Discussions*, 9, 19 879–19 937, 2009.
- Hobbs, W. H.: The Greenland Glacial Anticyclone, *Journal of Meteorology*, 2, 143–153, 1945.
- 650 Hofer, S., Tedstone, A. J., Fettweis, X., and Bamber, J. L.: Decreasing cloud cover drives the recent mass loss on the Greenland Ice Sheet, *Science Advances*, 3, 2017.
- Hofer, S., Tedstone, A. J., Fettweis, X., and Bamber, J. L.: Cloud microphysics and circulation anomalies control differences in future Greenland melt, *Nature Climate Change*, 9, 523–528, 2019.
- Hogan, A. W., Barnard, S. C., Kerschull, K., Townsend, R., and Samson, J. A.: Aerosol variation in the western hemisphere Arctic, *Journal*  
655 *of Aerosol Science*, 15, 13–33, 1984.
- Howat, I., Negrete, A., and Smith, B.: The Greenland Ice Mapping Project (GIMP) Land Ice and Ocean Classification Mask, Version 1., NASA National Snow and Ice Data Center Distributed Active Archive Center., <https://doi.org/https://doi.org/10.5067/B8X58MQBFUPA>, 2017.
- Hudson, J. G. and Da, X.: Volatility and size of cloud condensation nuclei, *Journal of Geophysical Research Atmospheres*, 101, 4435–4442,  
660 1996.
- Igel, A. L., Ekman, A. M. L., Leck, C., Tjernström, M., Savre, J., and Sedlar, J.: The free troposphere as a potential source of arctic boundary layer aerosol particles, *Geophysical Research Letters*, 44, 7053–7060, 2017.
- Intrieri, J. M., Fairall, C. W., Shupe, M. D., Persson, P. O. G., Andreas, E. L., Guest, P. S., and Moritz, R. E.: An annual cycle of Arctic surface cloud forcing at SHEBA, *Journal of Geophysical Research: Oceans*, 107, 2002.
- 665 Jaenicke, R.: Problems of the Distribution of the Global Aerosol, *Russian Chemical Reviews*, 59, 959, 1990.
- Jung, C. H., Yoon, Y. J., Kang, H. J., Gim, Y., Lee, B. Y., Ström, J., Krejci, R., and Tunved, P.: The seasonal characteristics of cloud condensation nuclei (CCN) in the arctic lower troposphere, *Tellus B: Chemical and Physical Meteorology*, 70, 1–13, 2018.
- Kanji, Z. A., Ladino, L. A., Wex, H., Boose, Y., Burkert-Kohn, M., Cziczo, D. J., and Krämer, M.: Overview of ice nucleating particles, *Meteorological Monographs*, 58, 1–1, 2017.
- 670 Kay, J. E., L’Ecuyer, T., Pendergrass, A., Chepfer, H., Guzman, R., and Yettella, V.: Scale-aware and definition-aware evaluation of modeled near-surface precipitation frequency using CloudSat observations, *Journal of Geophysical Research: Atmospheres*, 123, 4294–4309, 2018.
- Lance, S., Shupe, M., Feingold, G., Brock, C., Cozic, J., Holloway, J., Moore, R., Nenes, A., Schwarz, J., Spackman, J. R., et al.: Cloud condensation nuclei as a modulator of ice processes in Arctic mixed-phase clouds, *Atmospheric Chemistry and Physics*, 11, 8003–8015, 2011.

- 675 Latham, T., Beyersdorf, A., Thornhill, K., Winstead, E., Cubison, M., Hecobian, A., Jimenez, J., Weber, R., Anderson, B., and Nenes, A.: Analysis of CCN activity of Arctic aerosol and Canadian biomass burning during summer 2008, *Atmospheric Chemistry and Physics*, 13, 2735–2756, 2013.
- Leaitch, W. R., Korolev, A., Aliabadi, A. A., Burkart, J., Willis, M. D., Abbatt, J. P. D., Bozem, H., Hoor, P., Köllner, F., Schneider, J., Herber, A., Konrad, C., and Brauner, R.: Effects of 20-100 nm particles on liquid clouds in the clean summertime Arctic, *Atmos. Chem. Phys.*, 16, 11 107–11 124, 2016.
- 680 Leck, C. and Svensson, E.: Importance of aerosol composition and mixing state for cloud droplet activation over the Arctic pack ice in summer, *Atmospheric Chemistry and Physics*, 15, 2545–2568, 2015.
- Lenaerts, J., Camron, M. D., Wyburn-Powell, C. R., and Kay, J. E.: Present-day and future Greenland Ice Sheet precipitation frequency from CloudSat observations and the Community Earth System Model, *The Cryosphere*, 14, 2253–2265, 2020.
- 685 Li, J., Chen, H., Li, Z., Wang, P., Fan, X., He, W., and Zhang, J.: Analysis of Low-level Temperature Inversions and Their Effects on Aerosols in the Lower Atmosphere, *Advances in Atmospheric Sciences*, 36, 1235–1250, 2019.
- Lohmann, U. and Feichter, J.: Global indirect aerosol effects: a review, *Atmospheric Chemistry and Physics*, 5, 715–737, 2005.
- Lubin, D. and Vogelmann, A. M.: A climatologically significant aerosol longwave indirect effect in the Arctic, *Nature*, 439, 453–456, 2006.
- Mann, G. W., Anderson, P. S., and Mobbs, S. D.: Profile measurements of blowing snow at Halley, Antarctica, *Journal of Geophysical Research*, 105, 491–508, 2000.
- 690 Martin, S. J., Wang, P. K., and Pruppacher, H. R.: A Theoretical Determination of the Efficiency with which Aerosol Particles are Collected by Simple Ice Crystal Plates, *Journal of the Atmospheric Sciences*, 37, 1628–1638, 1980.
- Mauritsen, T., Sedlar, J., Tjernström, M., Leck, C., Martin, M., Shupe, M., Sjogren, S., Sierau, B., Persson, P. O. G., Brooks, I. M., Others, Tjernström, M., Leck, C., Martin, M., Shupe, M., Sjogren, S., Sierau, B., Persson, P. O. G., Brooks, I. M., and Swietlicki, E.: An Arctic CCN-limited cloud-aerosol regime, *Atmospheric Chemistry and Physics*, 11, 165–173, 2011.
- 695 McIlhatten, E. A., L’Ecuyer, T. S., and Miller, N. B.: Observational evidence linking Arctic supercooled liquid cloud biases in CESM to snowfall processes, *Journal of Climate*, 30, 4477–4495, 2017.
- McIlhatten, E. A., Pettersen, C., Wood, N. B., and L’Ecuyer, T. S.: Satellite observations of snowfall regimes over the Greenland Ice Sheet, *The Cryosphere*, 14, 4379–4404, 2020.
- 700 Miller, N. B., Turner, D. D., Bennartz, R., Shupe, M. D., Kulie, M. S., Cadeddu, M. P., and Walden, V. P.: Surface-based inversions above central Greenland, *Journal of Geophysical Research Atmospheres*, 118, 495–506, 2013.
- Miller, N. B., Shupe, M. D., Cox, C. J., Walden, V. P., Turner, D. D., and Steffen, K.: Cloud radiative forcing at Summit, Greenland, *Journal of Climate*, 28, 6267–6280, 2015.
- 705 Miller, N. B., Shupe, M. D., Cox, C. J., Noone, D., Persson, P. O. G., and Steffen, K.: Surface energy budget responses to radiative forcing at Summit, Greenland, *The Cryosphere*, 11, 497–516, 2017.
- Moran, K. P., Mariner, B. E., Post, M. J., Kropfli, R. A., Welsh, D. C., and Widener, K. B.: An Unattended Cloud-Profiling Radar for Use in Climate Research, *Bulletin of the American Meteorological Society*, 79, 443–455, 1998.
- Morrison, H., De Boer, G., Feingold, G., Harrington, J., Shupe, M. D., and Sulia, K.: Resilience of persistent Arctic mixed-phase clouds, *Nature Geoscience*, 5, 11–17, 2012.
- 710 Norgren, M. S., Boer, G. d., and Shupe, M. D.: Observed aerosol suppression of cloud ice in low-level Arctic mixed-phase clouds, *Atmospheric Chemistry and Physics*, 18, 13 345–13 361, 2018.

- Paramonov, M., Grönholm, T., and Virkkula, A.: Below-cloud scavenging of aerosol particles by snow at an urban site in Finland, *Boreal Environment Research*, 16, 304–320, 2011.
- Pettersen, C., Bennartz, R., Merrelli, A. J., Shupe, M. D., Turner, D. D., and Walden, V. P.: Precipitation regimes over central Greenland  
715 inferred from 5 years of ICECAPS observations., *Atmospheric Chemistry & Physics*, 18, 2018.
- Pettersen, C., Henderson, S., Mattingly, K., Bennartz, R., and Breeden, M.: The Critical Role of Euro-Atlantic Blocking in Promoting  
Snowfall in Central Greenland, *JGR – Atmospheres*, in review.
- Pisso, I., Sollum, E., Grythe, H., Kristiansen, N. I., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. L., Groot Zwaafink,  
C. D., Evangeliou, N., Sodemann, H., Haimberger, L., Henne, S., Brunner, D., Burkhardt, J. F., Fouilloux, A., Brioude, J., Philipp, A.,  
720 Seibert, P., and Stohl, A.: The Lagrangian particle dispersion model FLEXPART version 10.4, *Geosci. Model Dev*, 12, 4955–4997, 2019.
- Possner, A., Ekman, A. M., and Lohmann, U.: Cloud response and feedback processes in stratiform mixed-phase clouds perturbed by ship  
exhaust, *Geophysical Research Letters*, 44, 1964–1972, 2017.
- Prenni, A. J., Harrington, J. Y., Tjernström, M., DeMott, P. J., Avramov, A., Long, C. N., Kreidenweis, S. M., Olsson, P. Q., and Verlinde, J.:  
Can ice-nucleating aerosols affect Arctic seasonal climate?, *Bulletin of the American Meteorological Society*, 88, 541–550, 2007.
- 725 Rignot, E., Box, J. E., Burgess, E., and Hanna, E.: Mass balance of the Greenland ice sheet from 1958 to 2007, *Geophysical Research Letters*,  
35, 1–5, 2008.
- Rignot, E., Velicogna, I., van den Broeke, M. R., Monaghan, A., and Lenaerts, J. T.: Acceleration of the contribution of the Greenland and  
Antarctic ice sheets to sea level rise, *Geophysical Research Letters*, 38, 2011.
- Roiger, A., Schlager, H., Schäfler, A., Huntrieser, H., Scheibe, M., Aufmhoff, H., Cooper, O. R., Sodemann, H., Stohl, A., Burkhardt, J.,  
730 et al.: In-situ observation of Asian pollution transported into the Arctic lowermost stratosphere, *Atmospheric Chemistry and Physics*, 11,  
10975–10994, 2011.
- Schmale, J., Henning, S., Decesari, S., Henzing, B., Keskinen, H., Sellegri, K., Ovadnevaite, J., Pöhlker, M. L., Brito, J., Bougiatioti, A., et al.:  
Long-term cloud condensation nuclei number concentration, particle number size distribution and chemical composition measurements at  
regionally representative observatories, *Atmospheric Chemistry and Physics*, 18, 2853–2881, 2018.
- 735 Schmale, J., Zieger, P., and Ekman, A. M. L.: Aerosols in current and future Arctic climate, *Nature Climate Change*, 2021.
- Schmeisser, L., Backman, J., Ogren, J. A., Andrews, E., Asmi, E., Starkweather, S., Uttal, T., Fiebig, M., Sharma, S., Eleftheriadis, K., Others,  
Vratolis, S., Bergin, M., Tunved, P., and Jefferson, A.: Seasonality of aerosol optical properties in the Arctic, *Atmospheric Chemistry and  
Physics Discussions*, pp. 1–41, 2018.
- Schmidt, R. A.: Properties of blowing snow, *Reviews of Geophysics*, 20, 39–44, 1982.
- 740 Schuenemann, K. C., Cassano, J. J., and Finniss, J.: Synoptic forcing of precipitation over Greenland: Climatology for 1961–99, *Journal of  
Hydrometeorology*, 10, 60–78, 2009.
- Seinfeld, J. H., Bretherton, C., Carslaw, K. S., Coe, H., DeMott, P. J., Dunlea, E. J., Feingold, G., Ghan, S., Guenther, A. B., Kahn, R., et al.:  
Improving our fundamental understanding of the role of aerosol–cloud interactions in the climate system, *Proceedings of the National  
Academy of Sciences*, 113, 5781–5790, 2016.
- 745 Shaw, G. E.: The Arctic haze phenomenon, *Bulletin of the American Meteorological Society*, 76, 2403–2414, 1995.
- Shepherd, A., Ivins, E., Rignot, E., Smith, B., van Den Broeke, M., Velicogna, I., Whitehouse, P., Briggs, K., Joughin, I., Krinner, G., et al.:  
Mass balance of the Greenland Ice Sheet from 1992 to 2018, *Nature*, 579, 233–239, 2020.
- Sheppard, B. E. and Joe, P. I.: Performance of the precipitation occurrence sensor system as a precipitation gauge, *Journal of Atmospheric  
and Oceanic Technology*, 25, 196–212, 2008.



- 750 Shupe, M., Persson, P., Brooks, I., Tjernström, M., Sedlar, J., Mauritsen, T., Sjogren, S., and Leck, C.: Cloud and boundary layer interactions over the Arctic sea ice in late summer, *Atmospheric Chemistry and Physics*, 13, 9379–9399, 2013a.
- Shupe, M. D. and Intrieri, J. M.: Cloud radiative forcing of the Arctic surface: The influence of cloud properties, surface albedo, and solar zenith angle, *Journal of Climate*, 17, 616–628, 2004.
- Shupe, M. D., Matrosov, S. Y., and Uttal, T.: Arctic mixed-phase cloud properties derived from surface-based sensors at SHEBA, *Journal of the atmospheric sciences*, 63, 697–711, 2006.
- 755 Shupe, M. D., Turner, D. D., Walden, V. P., Bennartz, R., Cadetdu, M. P., Castellani, B. B., Cox, C. J., Hudak, D. R., Kulie, M. S., Miller, N. B., Others, Neely, R. R., Neff, W. D., and Rowe, P. M.: High and dry: New observations of tropospheric and cloud properties above the Greenland Ice Sheet, *Bulletin of the American Meteorological Society*, 94, 169–186, 2013b.
- Slater, T., Hogg, A. E., and Mottram, R.: Ice-sheet losses track high-end sea-level rise projections, 2020.
- 760 Solomon, A., Boer, G. d., Creamean, J. M., McComiskey, A., Shupe, M. D., Maahn, M., and Cox, C.: The relative impact of cloud condensation nuclei and ice nucleating particle concentrations on phase partitioning in Arctic mixed-phase stratocumulus clouds, *Atmospheric Chemistry and Physics*, 18, 17 047–17 059, 2018.
- Stevens, R. G., Loewe, K., Dearden, C., Dimitrellos, A., Possner, A., Eirund, G. K., Raatikainen, T., Hill, A. A., Shipway, B. J., Wilkinson, J., Romakkaniemi, S., Tonttila, J., Laaksonen, A., Korhonen, H., Connolly, P., Lohmann, U., Hoose, C., Ekman, A. M., Carslaw, K. S., and 765 Field, P. R.: A model intercomparison of CCN-limited tenuous clouds in the high Arctic, *Atmospheric Chemistry and Physics*, 18, 2018.
- Stohl, A.: Characteristics of atmospheric transport into the Arctic troposphere, *Journal of Geophysical Research Atmospheres*, 111, 1–17, 2006.
- Storelvmo, T.: Aerosol effects on climate via mixed-phase and ice clouds, *Annual Review of Earth and Planetary Sciences*, 45, 199–222, 2017.
- 770 Thomas, M. A., Devasthale, A., Tjernström, M., and Ekman, A. M. L.: The Relation Between Aerosol Vertical Distribution and Temperature Inversions in the Arctic in Winter and Spring, *Geophysical Research Letters*, 46, 2836–2845, 2019.
- Trusel, L. D., Das, S. B., Osman, M. B., Evans, M. J., Smith, B. E., Fettweis, X., McConnell, J. R., Noël, B. P. Y., and van den Broeke, M. R.: Nonlinear rise in Greenland runoff in response to post-industrial Arctic warming, *Nature*, 564, 104–108, 2018.
- Turner, D. D., Clough, S. A., Liljegren, J. C., Clothiaux, E. E., Cady-Pereira, K. E., and Gaustad, K. L.: Retrieving Liquid Water Path and Precipitable Water Vapor From the Atmospheric Radiation Measurement (ARM) Microwave Radiometers, *IEEE Transactions on 775 Geoscience and Remote Sensing*, 45, 3680–3690, 2007.
- Twomey, S.: The influence of pollution on the shortwave albedo of clouds, *Journal of the atmospheric sciences*, 34, 1149–1152, 1977.
- Ulbrich, U., Pinto, J. G., Kupfer, H., Leckebusch, G. C., Spanghel, T., and Reyers, M.: Changing Northern Hemisphere storm tracks in an ensemble of IPCC climate change simulations, *Journal of Climate*, 21, 1669–1679, 2008.
- 780 Van Den Broeke, M. R., Enderlin, E. M., Howat, I. M., Kuipers Munneke, P., Noël, B. P., Jan Van De Berg, W., Van Meijgaard, E., and Wouters, B.: On the recent contribution of the Greenland ice sheet to sea level change, *Cryosphere*, 10, 1933–1946, 2016.
- Van Tricht, K., Lhermitte, S., Lenaerts, J., Gorodetskaya, I., L'Ecuyer, S., Noël, B., Van Den Broeke, M., Turner, D., and Van Lipzig, N.: Clouds enhance Greenland ice sheet meltwater runoff, *Nature Communications*, 7, 2016.
- Von Der Weiden, S.-L., Drewnick, F., and Borrmann, S.: Atmospheric Measurement Techniques Particle Loss Calculator-a new software tool 785 for the assessment of the performance of aerosol inlet systems, *Tech. rep.*, 2009.

Wex, H., Huang, L., Zhang, W., Hung, H., Traversi, R., Becagli, S., Sheesley, R. J., Moffett, C. E., Barrett, T. E., Bossi, R., et al.: Annual variability of ice-nucleating particle concentrations at different Arctic locations, *Atmospheric Chemistry and Physics*, 19, 5293–5311, 2019.

790 Willis, M. D., Richard, L. W., and P.D., A. J.: Processes Controlling the Composition and Abundance of Arctic Aerosol, *Reviews of Geophysics*, 56, 621–671, 2018.

Ziemba, L. D., Dibb, J. E., Griffin, R. J., Huey, L. G., and Beckman, P.: Observations of particle growth at a remote, Arctic site, *Atmospheric Environment*, 44, 1649–1657, 2010.

**Table 1.** Measurements used in this study, references provide additional instrument information and methodologies for derived parameters.

Instrument	Measured / <i>derived</i> parameters (used in this study)	Data availability (used in this study)	Reference
Condensation Particle Counter (CPC)	Ambient condensation nuclei number concentration > 5 nm	Feb '19-May '20 (exc. Jan '20)	Guy et al. (2020)
Alphasense Optical Particle Counter (OPC-N3)	Aerosol particle size distribution 0.35 to 40 $\mu\text{m}$	Jun-Dec '19	Crilley et al. (2018)
NOAA Met. suite	10 m wind speed and direction, Surface pressure	Feb '19-May 20'	GMLMET (2021)
Vaisala HMP155 T/RH probe	2 m air temperature, 15 m air temperature	Jun-Oct '19	Guy et al. (2020)
Precipitation Occurrence Sensor System (POSS), X-band (10.5 GHz)	<i>Precipitation occurrence</i> ( <i>POSS power unit</i> )	Mar-Dec '19	Sheppard and Joe (2008)
Millimeter cloud radar (MMCR) Ka band (35 GHz)	Radar reflectivity	.	Moran et al. (1998)
Radiosondes (00:00 and 12:00 UTC)	Vertical temperature and humidity profiles	Case studies only	Shupe et al. (2013b)
HATPro and MWRHF microwave radiometers (23, 21, 90 and 150 GHz)	<i>Liquid water path</i> , <i>precipitable water vapour</i>	Case studies only	Turner et al. (2007)
Micropulse lidar (MPL)	<i>Lidar depolarisation ratio</i>	Case studies only	Shupe et al. (2013b) Flynn et al. (2007)