

What caused a record high PM₁₀ episode in northern Europe in October 2020?

Christine D. Groot Zwaaftink, Wenche Aas, Sabine Eckhardt, Nikolaos Evangeliou, Paul Hamer, Mona Johnsrud, Arve Kylling, Stephen M. Platt, Kerstin Stebel, Hilde Uggerud, Karl Espen Yttri

5 NILU - Norwegian Institute for Air Research, P.O. Box 100, N-2027 Kjeller, Norway

Correspondence to: Christine Groot Zwaaftink (cgz@nilu.no)

Abstract

Early October 2020, northern Europe experienced an episode with poor air quality due to high concentrations of particulate matter (PM). At several sites in Norway, recorded weekly values exceeded historical maximum PM₁₀ concentrations from the
10 past 4 to 10 years. Daily mean PM₁₀ values at Norwegian sites were up to 97 $\mu\text{g m}^{-3}$ and had a median value of 59 $\mu\text{g m}^{-3}$. We analysed this severe pollution episode caused by long-range atmospheric transport based on surface and remote sensing observations and transport model simulations to understand its causes. Samples from three sites in mainland Norway and the Arctic remote station Zeppelin (Svalbard) showed strong contributions from mineral dust to PM₁₀ (23-36% as a minimum and 31-45% as a maximum) and biomass burning (8-16% – 19-21%). Atmospheric transport simulations indicate that Central Asia
15 was the main source region for mineral dust observed in this episode. The biomass burning fraction can be attributed to forest fires in Ukraine and southern Russia, but we cannot exclude other sources contributing, like fires elsewhere, because the model underestimates observed concentrations. The combined use of remote sensing, surface measurements and transport modelling proved effective in describing the episode and distinguishing its causes.

1 Introduction

20 Early October 2020, atmospheric concentrations of PM₁₀ and PM_{2.5}, i.e. particulate matter (PM) with an equivalent aerodynamic diameter of $\leq 10 \mu\text{m}$ and $\leq 2.5 \mu\text{m}$, were elevated at many stations in northern Europe (Fig. 1), exceeding international and national air quality guidelines (AQG) (WHO, 2006). On 2 and/or 3 October, daily mean EU-limit values for PM₁₀ of 50 $\mu\text{g m}^{-3}$ were exceeded at 45 of the European Environment Agency's (EEA) air quality urban and regional background sites (Fig. 1). In Norway, daily averaged PM₁₀ values exceeded previous maximum values recorded at several
25 sites. Visibility was strongly reduced, and questions concerning the source of these pollutants and possible influence on health arose (e.g. Aftenposten, 2020).

PM levels are usually increased in winter and spring compared to summer in urban areas in northern Europe, largely explained by increased emissions from residential wood burning and resuspension of road dust generated by use of studded tires, contributing to the fine and the coarse fraction of PM₁₀, respectively (Kukkonen et al., 2020; Laupsa et al., 2009; Yttri et al.,


30 2009; Yttri et al., 2005). However, in this case the widespread exceedance of PM₁₀ values, not only in Norway, pointed to long-range transport (LRT) of PM. Violation of limit values caused by LRT of PM in Norway rarely occur. Historically, high levels of LRT of PM have been associated with secondary inorganic aerosol species, SO₄²⁻, NO₃⁻ and NH₄⁺, formed by atmospheric oxidation of European SO₂, NH₃ and NO_x emissions (e.g. Tarrason et al., 2019; Yttri et al., 2021). A fast-track analysis showed that the source of the increased concentrations could have been biomass burning in Eastern Europe. However, 35 the fact that concentrations of coarse fraction PM₁₀ were also enhanced suggested that sources such as dust, ash and sea salt aerosol could also be important. Out of these, mineral dust (from agricultural or natural sources) is the most likely aerosol based on the region with enhanced PM levels (Fig. 1).

Events of LRT of mineral dust to Scandinavia are rare. Mineral dust outbreaks from the Sahara can reach the surface in Scandinavia (e.g. Ansmann et al., 2003), although the dust plumes likely reside at higher altitudes. An aerosol transport event 40 with combined contributions from biomass burning from the Iberian Peninsula and dust emissions from North Africa, affecting several regions in Europe, was described by Akritidis et al. (2020). Also, transport events of mineral dust from Kazakhstan have been shown to affect Scandinavia (e.g. Hongisto and Sofiev, 2004). Besides affecting surface concentrations of PM and air quality, airborne dust can strongly influence the radiation budget of the atmosphere, both directly and indirectly (e.g. Kylling et al., 2018; Myhre et al., 2013).

45 LRT events of aerosol particles from biomass burning have been frequently observed in Scandinavia (e.g., Yttri et al., 2007b; Saarikoski et al., 2007) including the European Arctic (Stohl et al., 2007), and can be related to wildfires and agricultural waste burning in the Baltic countries, western Russia, Belarus, and Ukraine. Fires release various particulate and gaseous substances such as organic and black carbon (OC and BC), CO₂, CH₄, and polycyclic aromatic hydrocarbons (PAHs) (Hao et al., 2016; Hao and Ward, 1993; Shi et al., 2015), some of which have an adverse effect on human health, such as benzo(a)pyrene, a 50 Group 1 carcinogen (IARC, 2021).

To quantify biomass burning contributions to PM we analyse filter samples for carbonaceous aerosol as well as levoglucosan (a biomass burning tracer). LRT modelling of the biomass burning aerosol in this study is based on BC, which is formed by the incomplete combustion of e.g., fossil fuels, biofuels, and biomass (Bond et al., 2013). BC not only affects human health (Lelieveld et al., 2015), but also climate (Myhre et al., 2013).

55 Here, we show how LRT of mineral dust from Central Asia and biomass burning aerosol, mainly from Ukraine, caused elevated PM levels in Norway. Our results are based on a combination of in-situ observations, satellite images and model simulations of long-range atmospheric transport.



60 **Figure 1** Mean PM_{2.5} (a,c) and PM₁₀ (b,d) values on 2 (a,b) and 3 (c,d) October 2020 (24-hour mean) at all EEA air quality background sites (includes urban as well as rural, but sites close to roads are excluded). Triangles indicate three Norwegian rural background sites with extensive PM speciation (section 2.1.1). For comparison: The WHO air quality guidelines for maximum daily averaged PM₁₀ and PM_{2.5} are 50 µg m⁻³ and 25 µg m⁻³, respectively, and the EU limit value for PM₁₀ is 50 µg m⁻³.

2. Data and methods

65 2.1 Surface observations

2.1.1 Sites and instrumentation

There are 48 (urban categorized) sites in Norwegian cities and towns measuring both PM₁₀ and PM_{2.5} mass concentration as part of their air quality programme Hak (2015). Rural background stations include the Birkenes Observatory (58°23'N, 8°15'E, 219 m above sea level, asl), Hurdal (60°22'N, 11°4'E, 300 m asl) and Kårvatn (62°47'N, 8°53'E, 210 m asl). These
70 are all EMEP (European Monitoring and Evaluation Programme) sites located in southern Norway (Fig. 1). The Birkenes Observatory is situated close to the Skagerrak coast (~20 km), whereas Hurdal and Kårvatn are further inland. Forests dominate the land use at all three sites, the remainder being meadows, low intensity agricultural areas and freshwater lakes. All three sites are thus considered rural background sites for air quality measurements. The Zeppelin Observatory (78°54' N 11°52' E, 472 m asl) is located on the Zeppelin Mountain 2 km south of the Ny-Ålesund settlement at
75 the west coast of Svalbard (Norway) and is considered an Arctic remote/ global background site. PM mass concentration (PM₁₀ and PM_{2.5}), organic and elemental carbon (OC/EC) (in PM₁₀ and PM_{2.5}), organic tracers (levoglucosan, mannosan, galactosan, arabitol, mannitol, glucose, trehalose, 2-methylerythritol, and 2-methylthreitol) (in PM₁₀), and crustal elements (Al, Fe, Mn, and Ti) (in PM₁₀) were obtained from low-volume samplers (flow rate of 38 L min⁻¹) with a sampling time of one week at the rural background sites. Prefired (850 °C; 3 h) quartz fiber filters (Whatman QM-A; 47 mm in diameter) were
80 conditioned (20±1 °C; 50±5% RH (relative humidity); 48 h) before and after exposure. Filters were stored at 4 °C after weighing and before OC/EC analysis and at -18 °C prior to organic tracer analysis. Please find further details on the analysis methods in appendix A1.

At the Zeppelin Observatory, OC/EC, organic tracers and crustal elements were obtained from a high-volume sampler with a PM₁₀ inlet (flow rate of 667 L min⁻¹) and with a sampling time of one week. Quartz fiber filters (PALLFLEX Tissuequartz
85 2500QAT-UP; 150 mm in diameter) were prefired (850 °C; 3 h) but not conditioned, and stored at -18 °C.

Major inorganic anions (SO₄²⁻, NO₃⁻, Cl⁻) and cations (Ca²⁺, Mg²⁺, K⁺, Na⁺, NH₄⁺) were collected using a NILU stacked filter unit (SFU) collecting aerosol particles on a Teflon filter (2 µm pore, 47 mm Zefluor Teflon, Gelman sciences) with a sampling time of 24 h. The SFU does not have a pre-impactor but has a downward facing inlet that effectively reduces the sampling efficiency for aerosol particles with an equivalent aerodynamic diameter larger than 10 µm.

90 PM₁₀ mass concentration was measured using a TEOM 1400a (Tapered Element Oscillating Monitor) (Patashnick and Rupprecht, 1991) instrument operating at a one-minute time resolution at the Birkenes Observatory. The aggregated weekly mean PM₁₀ concentration obtained by the TEOM instrument is proven equivalent to the weekly time resolved, filter based PM₁₀ measurements at Birkenes, which is determined according to the reference method EN 12341 (CEN, 2014).

95 Absorption coefficients (B_{Abs}) were measured using a seven-wavelength ($\lambda=370; 470; 520; 590; 660; 880; 950$ nm) Aethalometer (AE33, Magee Scientific) operating downstream of a PM_{10} inlet and calculated according to Drinovec et al. (2015) at the Birkenes and Zeppelin observatories. B_{Abs} was converted to eBC using co-located EC measurements.

2.1.2 Mineral dust mass estimation

100 We assumed observed levels of Al, Fe, Mn and Ti to be associated with mineral dust exclusively, and that Al was present as Al_2O_3 , Fe as Fe_2O_3 , Mn as MnO and Ti as TiO_2 (Alastuey et al., 2016). Not all mineral dust elements were measured such as Si, nor is the mineralogy of the mineral dust collected on the filter samples known, thus, SiO_2 was estimated based on an empirical factor (eq.1) (Alastuey et al., 2016). We also made assumptions regarding the speciation of CO_3^{2-} , assuming 70% was present as $CaCO_3$, 20% as $MgCO_3$ and 10% as K_2CO_3 (ref.). The exact CO_3^{2-} speciation is however of minor importance, as the CO_3^{2-} - fraction increases the lower mineral dust estimate (eq.2) by only 6.3% (assuming all is $MgCO_3$) to 10% (assuming all is K_2CO_3), using data for Zeppelin as an example. Ca^{2+} and Mg^{2+} assumed to be associated with CO_3^{2-} were not considered part of the water-soluble fraction determined by ion chromatography (See Appendix 1 for a description of the ion chromatography method). Excess Ca^{2+} , Mg^{2+} , and K^+ (i.e., not part of sea salt aerosol or K_2CO_3) were assumed to be present as oxides (CaO , MgO , and K_2O), which together with the carbonates and metal oxides provided the upper mineral dust estimate (eq.3). We consider this an approximation, as chemical reactions likely take place during atmospheric transport. Both NO_3^- and SO_4^{2-} were enhanced during the episode and could indicate the presence of e.g. $Mg(NO_3)_2$ and $MgSO_4$ formed from reactions between $MgCO_3$ and acids such as HNO_3 or H_2SO_4 (Laskin et al., 2004). Accounting for such reactions would increase the upper mineral dust estimate (eq.3) by 12 – 21%. However, it is not apparent that nitrates and sulphates formed this way should be apportioned to the mineral dust fraction. Finally, ion equivalent calculations suggest that all SO_4^{2-} and NO_3^- were associated with NH_4^+ , thus we do not include these potential reactions in the upper estimate of mineral dust.

CO_3^{2-} can also be part of wildfire emissions and is a dominant species of ash produced at around 500 °C with $CaCO_3$ being the most abundant species followed by $MgCO_3$ and K_2CO_3 (Bodí et al., 2014). Hence, it is questionable whether CO_3^{2-} should be apportioned as mineral dust or biomass burning when such emissions are mixed in the atmosphere. At temperatures >580 °C, carbonates dissociate to oxides (Bodí et al., 2014), which are partially soluble in water, thus, as for CO_3^{2-} , it is not clear if CaO , MgO , and K_2O should be part of mineral dust or biomass burning particles. Consequently, any attempt to reconstruct the mineral dust mass concentration should be considered semi-quantitative. We thus calculate a lower (eq.2) and an upper (eq.3) estimate of the mineral dust mass concentration including no or all CO_3^{2-} and oxides, respectively. With CO_3^{2-} and oxides apportioned to mineral dust (eq.3) an upper estimate of BB (eq.8) would not be possible and vice versa.

125

$$[SiO_2] = 2.5 \times [Al_2O_3] \text{ (eq.1)}$$

$$[Mineral\ dust_{mass}]_{lower} = [SiO_2] + [Al_2O_3] + [Fe_2O_3] + [MnO] + [TiO_2] \text{ (eq.2)}$$

$$130 \quad [Mineral\ dust_{mass}]_{upper} = [SiO_2] + [Al_2O_3] + [Fe_2O_3] + [MnO] + [TiO_2] + [(n)CO_3^{2-}] + [(n)O] \text{ (eq.3)}$$

$$(n = Ca^{2+}, Mg^{2+}, 2Na^+, 2K^+)$$

2.1.3 Biomass burning mass estimation

135 Levoglucosan is formed from the thermal degradation of cellulose and is a proven tracer of biomass burning (BB) emissions (Locker, 1988; Simoneit et al., 1999) as demonstrated in numerous papers (e.g., Zdrahal et al., 2002; Puxbaum et al., 2007; Szidat et al., 2009; Yttri et al., 2019). Emission ratios of levoglucosan from wildfires are likely to vary widely reflecting combustion conditions and vegetation, and source region. Here we apply emission ratios for total carbon and organic carbon (Yttri et al., 2014) to calculate OC_{BB} and EC_{BB} , which are derived from ambient
140 sampling of wildfires emissions in Eastern Europe, including Ukraine (Saarikoski et al., 2007). Calculated concentrations of OC_{BB} and EC_{BB} should be considered semiquantitative given the uncertainty of the emission ratios and the potential atmospheric degradation of levoglucosan.

Although BB aerosol from wildfires is dominated by carbonaceous aerosol, carbonates and oxides can result from wildfires, as well as being a part of mineral dust. Hence, we calculated a lower estimate of the biomass burning aerosol accounting for
145 the carbonaceous fraction ($TC_{BB}=OC_{BB}+EC_{BB}$) (eq.4-7) and an upper estimate accounting for carbonates and oxides, in addition to the carbonaceous fraction (eq.8). For eq.4, eq.5 and eq.10, notations in parentheses are emissions ratios.

$$[TC_{BB}] = [Levoglucosan] \times (TC/levoglucosan)_{BB} \quad \text{(eq.4)}$$

$$[OC_{BB}] = [TC_{BB}] \times (OC/TC)_{BB} \quad \text{(eq.5)}$$

$$150 \quad [EC_{BB}] = [TC_{BB}] - [OC_{BB}] \quad \text{(eq.6)}$$

$$[BB_{mass}]_{Lower} = [OC_{BB} \times 2.2] + [EC_{BB} \times 1.1] \quad \text{(eq.7)}$$

$$[BB_{mass}]_{Upper} = [OC_{BB} \times 2.2] + [EC_{BB} \times 1.1] + [(n)CO_3^{2-}] + [(n)O] \quad (n = Ca^{2+}, Mg^{2+}, 2Na^+, 2K^+) \text{ (eq. 8)}$$

155 where BB_{mass} is the mass concentration ($\mu\text{g m}^{-3}$) of the biomass burning aerosol.

For aerosol particles dominated by biomass burning, a factor of 2.2-2.6 to convert OC_{BB} ($\mu\text{g C m}^{-3}$) to OM_{BB} (organic matter; $\mu\text{g m}^{-3}$) is recommended, whereas 1.9-2.2 is suggested for aged aerosol particles (Turpin and Lim, 2001). Here we used a factor of 2.2 both for OC_{BB} and OC , as OC at rural background and remote sites largely are long-range transported and thus aged. Similarly, a factor of 1.1 was used for both EC_{BB} and EC (Kiss et al., 2002).

160 We estimated levels of primary biological aerosol particles (PBAP), biogenic secondary organic aerosol (BSOA), and fossil
fuel sources (FF), as these are complementary to BB and yield a better constraint on the biomass burning source. For PBAP,
we used observed levels of arabinol, mannitol, glucose, and trehalose ($\Sigma_{\text{PBAP-Tracers}}$) (Table 1), an OC to $\Sigma_{\text{PBAP-Tracers}}$ ratio of
14.6 \pm 2.1 derived from Yttri et al. (2021), and an OM:OC conversion factor of 1.75 (Yttri et al., 2011a) to calculate OM_{PBAP}
(unit: $\mu\text{g m}^{-3}$). OM_{BSOA} was estimated based on observed levels of 2-methyltetrols ($\Sigma_{\text{BSOA-tracers}}$ =2-methylerythritol and 2-
165 methylthreitol) (Tab. 2), an OC to $\Sigma_{\text{BSOA-tracers}}$ ratio of 165 \pm 18 derived from Yttri et al. (2021) and an OM:OC conversion factor
of 2.2 (Turpin and Lim, 2001). Yttri et al. (2021) found that 2-methyltetrols traced local BSOA, which was 30% of total
identified BSOA constituents in PM₁₀, hence total OM_{BSOA} was adjusted accordingly. EC from fossil fuel sources (EC_{FF}) was
calculated according to eq.9:

$$170 \quad [\text{EC}_{\text{FF}}] = [\text{EC}] - [\text{EC}_{\text{BB}}] \quad (\text{eq.9})$$

where EC_{BB} is obtained from equation 6.

OC from fossil fuel sources (OC_{FF}) was calculated according to (eq.10):

175

$$[\text{OC}_{\text{FF}}] = [\text{EC}_{\text{FF}}] \times (\text{OC/EC})_{\text{FF}} \quad (\text{eq. 10})$$

where the (OC/EC)_{FF} ratio (2.0 \pm 0.25) derived from Yttri et al. (2021) include both primary and secondary OC. An OM:OC
conversion factor of 1.2 was applied for OC_{FF} and 1.1 for EC_{FF}. As for BB_{mass} we used the index “mass” in PBAP_{mass}, BSOA_{mass}
180 and FF_{mass} to impress that the unit is $\mu\text{g m}^{-3}$.

2.1.4 Source apportionment of the absorption coefficient into eBC_{BB} and eBC_{FF}

We used positive matrix factorisation with a multilinear engine (PMF-ME2) (Canonaco et al., 2013) and SoFI Pro software
(Canonaco et al., 2021) to apportion eBC_{BB} and eBC_{FF} based on observations of the absorption coefficient. PMF yields factor
185 profiles (here the wavelength dependant absorption, see Yttri et al., 2021) and time series of the emission sources. Two-factor
solutions from repeat bootstrapped PMF runs were mapped via Ångström exponents (AAE, calculated from the factor profiles)
using factor 1= lowest AAE, factor 2=highest AAE. An averaged 2 factor solution for each site was then determined, with
AAE factor 1; factor 2 of 0.94; 2.01 at Birkenes and 0.8; 1.7 at Zeppelin. Noting that traffic emissions may be partly biofuel
derived, and that residential coal combustion may be partly responsible for the high AAE factor, we identify factor 1 as
190 fossil/liquid fuel eBC and factor 2 as biomass burning/solid fuel eBC consistent with literature AAEs for these sources
(Sandradewi et al., 2008, Zotter et al., 2017).

2.2 Satellite observations

The Ocean and Land Colour Instrument (OLCI) on board the Sentinel-3 satellites (Donlon et al., 2012), measures the solar radiation reflected by the Earth's surface and atmosphere in 21 spectral bands from the visible to the near infrared. Over land and within 300 km of charted land, the ground spatial resolution is 300 m. The primary objective of OLCI products is to provide accurate measurements of ocean colour and land surface reflectance. We use OLCI measurements to visualize the aerosol when passing over central Norway and Sweden and to investigate the effect of the aerosol on the measured OLCI radiances by comparison to radiative transfer simulations.

Furthermore, we use observations from the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) on board of the Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO) platform. CALIPSO was launched April 2006 (Winker et al., 2009). CALIOP is a two-wavelength (1064 and 532 nm), polarization-sensitive (at 532 nm) elastic backscatter lidar, which provides globally day- and nighttime profiles of aerosol backscatter, extinction (with an extinction-to-backscatter a priori), and linear particle depolarization with altitude resolution between 30 m and 300 m, below 8.3 km and between 30.1 and 40.0 km, respectively. CALIOP has a small horizontal footprint of 335 m and a revisit time of ~16 days. Here, we utilize the level 2 data products (version 4.21) of the aerosol extinction at 532 nm to evaluate the representation of the dust and BC (from BB) plume in our atmospheric transport simulations. It is given at a spatial resolution of 60 m vertically and 5 km horizontally. The V4 level 2 cloud-aerosol discrimination (CAD) algorithm distinguishes between following tropospheric aerosol subtypes: clean marine, polluted continental/smoke, clean continental, polluted dust, elevated smoke, and dusty marine (Kim et al., 2018). The data were downloaded from the ICARE Data and Services Center (<http://www.icare.univ-lille1.fr/>, last access: 19 January 2021).

2.3 Radiative transfer model

To understand the processes influencing the OLCI radiances, radiative transfer simulations were made using the DISORT model (Stamnes et al., 1988; Buras et al., 2011) within the libRadtran framework (Emde et al., 2016). The Subarctic summer atmosphere (Anderson et al., 1986) was adopted as the ambient atmosphere. Visible Infrared Imaging Radiometer Suite (VIIRS) measurements indicate cloud top height between about 2-4 km and cloud optical depth between 50-100 (worldview.earthdata.nasa.gov). Hence, for the cloudy simulations, a cloud layer of optical depth 70 was included between 2 and 3 km. For simulations over water the Cox and Munk (1954a, b) ocean bidirectional reflectance distribution function (BRDF) was adopted. Aerosols were included using a profile based on FLEXPART simulation results over Norway (section 3.1). The aerosol optical properties were prepared for input to libRadtran with the MOPSMAP tool (Gasteiger and Wiegner, 2018).

¹ <https://sentinel.esa.int/web/sentinel/user-guides/sentinel-3-olci/overview>

2.4 Atmospheric transport models and emission datasets

With the Lagrangian particle dispersion model FLEXPART version 10.4 (Pisso et al., 2019) we model mineral dust in forward and BC in both forward and backward mode. FLEXPART calculates trajectories of particles to describe transport and diffusion of tracers in the atmosphere. Particles are assumed to be spherical and influenced by gravitational settling, dry deposition, and in-cloud and below-cloud scavenging (Grythe et al., 2017). The model is widely applied for LRT modelling of fire emissions (Evangelou et al., 2016; 2019) and dust sources (e.g. Sodemann et al., 2015).

Emissions from BB were adopted from Copernicus Atmosphere Monitoring Services (CAMS) Global Fire Assimilation System (GFAS). CAMS GFAS assimilates fire radiative power (FRP) observations from satellite-based sensors converting the energy released during fire combustion into gases and aerosol daily fluxes (Di Giuseppe et al., 2016; Kaiser et al., 2012). Data are available globally on a regular grid with horizontal resolution of 0.1 degrees from 2003 to present. FRP observations assimilated in GFAS are the NASA Terra MODIS and Aqua MODIS active fire products (<http://modis-fire.umd.edu/>, (Kaufman et al., 2003)). FRP measures the heat power emitted by fires, as a result of the combustion process and is directly related to the total biomass combusted (Wooster et al., 2005). Using land-use dependent conversion factors, GFAS converts FRP into emission estimates of 44 smoke constituents, such as CO, CO₂, CH₄, and black-carbon and organic matter components of the aerosol (Kaiser et al., 2012). Here, we used emissions of BC that were subsequently ingested into FLEXPART, which simulated it forward to track atmospheric LRT. The simulations were driven by operational meteorological analysis data from European Centre for Medium-range Weather Forecast (ECMWF) of 1-degree spatial resolution and 3-hourly temporal resolution. The spatial resolution of the output was set to 0.5 degrees and the temporal resolution to daily. The simulations accounted for wet and dry deposition, assuming a particle density of 1500 kg m⁻³ and a logarithmic size distribution with an aerodynamic mean diameter of 0.25 μm and a standard deviation of 0.3 (Hu et al., 2018; Long et al., 2013).

Besides forward simulations of BC, we also performed backward simulations based on 3-hourly releases from the rural background and remote stations to obtain the emission sensitivity and distinguish sources contributing to BC concentrations at these locations. We thereby assumed the same properties for BC as in the forward simulation. Emission sensitivity in the bottom 500 m was linked to fields of BC emissions from biomass burning based on GFAS and BC emissions from fossil fuels retrieved from the Evaluating the Climate and Air Quality Impacts of Short-Lived Pollutants (ECLIPSE) emission data set (Stohl et al., 2015; Klimont et al., 2017) on 0.5-degrees resolution.

Emissions of mineral dust are calculated with the FLEXDUST module (Groot Zwaafink et al., 2016). This module describes dust mobilization and emission as a function of (threshold) friction velocity following the approach introduced by Marticorena and Bergametti (1995). Modelled threshold friction velocity is influenced by soil moisture (Fécan et al., 1999) and sediment regions were identified based on large-scale topography (Ginoux et al., 2001). Emissions are calculated at 0.25 degrees resolution and 3-hourly interval. The forward simulations include ten size bins for dust smaller than 20 micrometres (diameter). For comparison to measurements based on PM₁₀ we will here only consider the size bins with dust up to 10 micrometres. Based on FLEXDUST emissions, we run two forward simulations of atmospheric transport of dust with FLEXPART. The

255 main simulation included global dust emissions, while the additional simulation included only dust from the Central Asian
desert region, here defined as a square region extending from 42°E to 82°E and 35°N to 50°N. We chose this setup rather than
backwards simulations (like we did for BC) because backwards simulations for mineral dust would have required separate
simulations for each size bin and station and this approach was thus more efficient. FLEXPART and FLEXDUST simulations
of mineral dust were driven with the same meteorological forcing data as for biomass burning FLEXPART simulations.
260 For comparison, we also include estimates from an operational air quality forecast system. The CAMS regional ensemble
forecast (Marécal et al., 2015) is composed of nine air quality models run over a European domain (30°W to 75°E, and 25°N
to 75°N). Forecasts are produced daily and run for 96 hours. The ensemble is taken as the median of the nine models, which
has higher skill than any of individual nine models. The dust product is the median of the prognostic simulations of mineral
dust from each model. Mineral dust is represented in each model by differing size bins and physics, so the dust concentration
265 represents PM of all sizes associated with mineral dust, which may include size bins up to PM₂₀. While each model uses its
own schemes to represent the dust emissions, they all use the dust from the CAMS global model to represent initial and
boundary conditions (Collin, 2021). The CAMS global model is run using IFS (Rémy et al., 2019) which simulates dust
emissions, uses three size bins (0.03-0.55 μm, 0.55-0.9 μm, and 0.9-20 μm), and performs assimilation of satellite AOD to
update aerosol concentrations. The CAMS model data were downloaded from the Copernicus Atmospheric Data Store
270 (<https://ads.atmosphere.copernicus.eu/>, last access: 18 March, 2021).


3. Results and discussions

Although the episode was initially mostly detected based on strong impacts on air quality at the surface, the aerosols were not
only present near the surface. We will first use satellite observations to assess the vertical extent and origin of the plumes.
Figure 2 presents an RGB (Red, Green, Blue) composite of OCLI observations for 2 October 2020. First, the image clearly
275 shows the presence of aerosols above the cloud layer over Norway/Sweden. Aerosols were thus present, in considerable
amounts, at an altitude of at least 2-3 km. We will further use the OLCI measurements in combination with radiative transfer
simulations to determine the absorption and scattering properties of the aerosol. Four points are marked in Figure 2, indicating
cloud only (Cloud), aerosol over cloud (Aerosol A), aerosol over water (Aerosol B) and water without visible cloud and aerosol
(Water). For these four cases OLCI top of the atmosphere radiances spectra were averaged over 9×9km². The spatially
280 averaged spectra and their standard deviations are shown as solid lines in Figure 3.

The aerosol reduces the radiance by about a factor of 2 when above the cloud layer due to increased absorption by
the aerosol compared to the non-absorbing cloud (compare Aerosol A and Cloud in Fig. 3), while above water the aerosol
increases the radiance due to increased backscattering (compare Aerosol B and Water in Fig. 3).


To elucidate the aerosol type(s) that may reproduce the OLCI radiances, radiative transfer simulations were made. These
285 simulations required aerosol optical depth, single scattering albedo (SSA) and phase function as input. Aerosol optical depth
is available from several sensors for the episode (see for example worldview.earthdata.nasa.gov). However, information about
aerosol type and absorption and scattering properties is generally not available. Thus, simulations were made with a
combination of highly scattering (resembling mineral dust) and highly absorbing aerosols (soot). The microphysical properties
of the aerosol were taken from Hess et al. (2018) and the optical properties calculated by the MOPSMAP tool (Gasteiger and

290 Wiegner, 2018). The scattering aerosols have a SSA of 0.98 at 400 nm and it decreases to about 0.93 at 1000 nm. The absorbing aerosols have a SSA of 0.28 at 400 nm and it decreases to about 0.1 at 1000 nm. No single aerosol type was able to reproduce the measurements, rather various combinations of the highly scattering and highly absorbing aerosols were needed to match the measurements. The amounts of scattering and absorbing aerosols were determined by scaling the scattering and absorption aerosol optical depths to get a best match between the OLCI radiances and the simulations, solid and dotted lines in Figure 3
295 respectively. For the water case (blue lines in Fig. 3), a highly scattering optical depth (OD) of 0.45 and highly absorbing OD of 0.2, both at 550 nm, were used. For Aerosol B (clearly visible aerosol over water, red lines in Fig. 3), highly scattering OD=2.5 and highly absorbing OD=0.7, while for Aerosol A (aerosol over cloud, black lines in Fig. 3) highly scattering OD=2.1 and highly absorbing OD=0.7. Thus, the reproduction of the satellite measurements indicates that the aerosol had an absorbing component.



300

Figure 2 RGB from OLCI bands 3, 6 and 8 centred at 442.5, 560 and 665 nm. Data from 2 Oct 2020, 09:33 UTC.




305 **Figure 3** The OLCI radiances (solid lines) at the four locations marked in Fig. 2. The error bars are standard deviation of the OLCI radiances within the square markers in Fig. 2. Note that for Aerosol B and Water the error bars are small and thus not shown. The dotted lines are radiative transfer model simulations of the OLCI measurements. See section 2.3 for further details.

The episode was clearly visible from a satellite perspective and the impact on the near-surface air quality was considerable. 24-hour mean PM_{10} concentrations on 2 and 3 October ranged from 8 to $97 \mu g m^{-3}$ at Norwegian urban sites, with a median value of $59 \mu g m^{-3}$. The episode thereby caused violation of the 24-hour PM_{10} EU-limit of $50 \mu g m^{-3}$ at 39 out of 48 Norwegian sites on 2 and/or 3 October, corresponding to 70 exceedance days in total. In comparison to years 2018 and 2019, for a selection of 30 sites with measurements in those years, this means that the number of exceedance days during the episode corresponded to 18% of the average total exceedance days for 2018 and 2019. The number of exceedance days is used to determine whether measures to improve air quality need to be taken, and it does not distinguish whether the cause of poor air quality is due to LRT or local pollutions.

315 During this episode, it appears that local sources were of minor importance, since 24-hour mean concentrations observed at the rural background site Birkenes, of $66 \mu g m^{-3}$ and $61 \mu g m^{-3}$, were in the same range as the urban sites. Weekly mean concentrations at the three rural background sites ranged from $18.6-22.2 \mu g m^{-3}$ for PM_{10} and from $4.1-16.1 \mu g m^{-3}$ for $PM_{2.5}$, being new maximum values for September–November 2010/11–2019 for all but $PM_{2.5}$ at Birkenes (Fig. 4; Table 1). Reconstructed PM_{10} mass concentration at Zeppelin amounted to $6.3 \mu g m^{-3}$. A prevailing coarse fraction ($PM_{10-2.5}$) of PM_{10} was seen at Birkenes (78% of PM_{10}) and Hurdal (61%), whereas it was minor at K arvatn (23%). The spatial variability is not easily explained but was seen at Norwegian urban sites during the episode as well (27–78% in $PM_{10-2.5}$, Table 1). A pronounced and even dominating coarse fraction can occur when sources such as sea salt aerosol and mineral dust are prominent.

325 We thus see indications of influences from mineral dust and BB aerosols, which can be confirmed by chemical speciation of PM, available at the three rural background sites and the Zeppelin Observatory (Fig. 4; Table 1). For a selection of species, we also show long-term measurements for comparison. With the chemical analysis performed (Table 1), and with assumptions made regarding mineral dust composition (lower estimate) (Sect. 2.1.2), and OC and EC conversion factors (Sect. 2.1.3), we were able to explain 92% (Birkenes), 74% (Hurdal), and 75% (K arvatn) of the PM_{10} mass at the rural sites. Note that secondary inorganic aerosol constituents and inorganic anions and cations were not available for the K arvatn site. The

range of mass closure obtained for the sites is comparable to previous studies (e.g. Putaud et al., 2010; Yttri et al., 2021; Aas et al., 2021) but should be considered a conservative estimate, using the lower estimate of the mineral dust fraction.



330

335

Figure 4 Panels show concentrations observed in the October 2020 episode (red diamonds: Birkenes, orange circles: Hurdal, purple triangles: Kårvatn, blue diamonds: Zeppelin) along with box plots (5, 25, 50, 75, 95 percentiles and outliers) at the rural background sites Birkenes, Hurdal and Kårvatn, and remote site Zeppelin of weekly data in the period Sept.-Nov. for a) PM₁₀ mass concentration (2010-2019); b) PM_{2.5} mass concentration (2010-2019); c) Organic Carbon (OC) in PM₁₀ (2010-2019 for Birkenes, 2011-2019 for Hurdal and Kårvatn 2017-2019 for Zeppelin); d) OC in PM_{2.5} (similar as for c except Zeppelin with no measurements); e) Elemental Carbon (EC) in PM₁₀ (similar as for c); f) EC in PM_{2.5} (similar as for d); g) Crustal elements in PM₁₀ (2014-2019); h) Levoglucosan (2010-2019), mannitol, and 2-methylerythritol in PM₁₀ (2016-2018); i)

Crustal elements in PM₁₀ at Zeppelin (2018-2019); j) Levoglucosan, mannitol, and 2-methylerythritol in PM₁₀ at Zeppelin (2017-2019). In panel g) and h) the box plots are for Birkenes only.

340 3.1 Mineral dust

Mineral dust elements (Al, Fe, Mn, and Ti) were all highly elevated during the episode. At Birkenes, the Al and Fe levels were over 30 times higher than the long-term seasonal mean (2014-2019), whereas the corresponding factors for Ti and Mn were respectively 27 and 16 (Table 1; Fig. 4). Maximum ever concentrations were observed for all mineral dust elements at Birkenes by a fair margin, suggesting that an event of this magnitude is rare. Note that long term time series of mineral dust are available only for Birkenes. K arvatn experienced the highest concentration for all the mineral dust elements during the episode, followed by Birkenes and Hurdal, although with a minor difference. The relative composition of the mineral dust elements was indistinguishable between the three rural background sites and the Zeppelin Observatory. Al was most abundant at all sites and concentrations of other elements relative to Al are similar at all sites: Al:Fe:Ti:Mn = 1:(1.5-1.6):(26-38):(56-65). This similar relative composition points at a common origin of the observed mineral dust. The Al:Fe ratio points to Eurasian mineral dust sources (Crocchianti et al., 2021), which is supported by our FLEXPART/FLEXDUST model simulations and CAMS regional mineral dust product, discussed below. Observed elements at Zeppelin were of similar composition yet observed masses were 2.4 – 3.7 times lower than the mean at the rural background sites, which is not surprising given the larger distance to the source region.

355 **Table 1: Mean concentrations of PM and associated species observed during the episode compared to the long-term weekly mean (\pm SD) for Sept.-Nov. (2010/11-2019) for Birkenes, Hurdal, K arvatn and Zeppelin, except crustal elements at Birkenes (2014-2019) and Zeppelin (2018-2019), EC/OC and organic tracers at Zeppelin (2017-2019).**

	Birkenes		Hurdal		K�arvatn		Zeppelin	
	Observatory						Observatory	
<i>Mass concentration</i>	<i>Episode</i>	<i>Mean (\pmSD)</i>	<i>Episode</i>	<i>Mean</i>	<i>Episode</i>	<i>Mean</i>	<i>Episode</i>	<i>Mean (\pmSD)</i>
<i>(μg m⁻³)</i>				<i>(\pmSD)</i>		<i>(\pmSD)</i>		
PM ₁₀	18.7	4.9 \pm 3.4	22.2	4.3 \pm 2.6	21.0	2.4 \pm 2.1	6.3 ¹⁾	N/A
PM _{2.5}	4.1	2.5 \pm 2.2	8.6	2.4 \pm 1.6	16.1	1.4 \pm 1.3	N/A	N/A
PM _{10-2.5}	14.6	2.6 \pm 2.0	13.6	2.0 \pm 1.4	4.9	1.1 \pm 1.0	N/A	N/A
<i>Carbonaceous</i>								
<i>aerosol (μg C m⁻³)</i>								
<i>PM₁₀</i>								
OC	2.0	0.72 \pm 0.49	2.9	1.2 \pm 0.77	3.3	0.72 \pm 0.61	0.81	0.051 \pm 0.079

	Birkenes		Hurdal		Kårvatn		Zeppelin	
	Observatory						Observatory	
EC	0.41	0.09±0.07	0.38	0.13±0.06	0.37	0.05±0.04	0.07	0.006±0.010
CO ₃ ²⁻	0.05	N/A	0.08	N/A	0.11	N/A	0.02	N/A
<i>PM_{2.5}</i>								
OC	0.67	0.48±0.33	1.5	0.59±0.30	2.4	0.42±0.29	N/A	N/A
EC	0.20	0.09±0.07	0.30	0.12±0.06	0.32	0.05±0.04	N/A	N/A
CO ₃ ²⁻	0.03	N/A	0.06	N/A	0.09	N/A	N/A	N/A
<i>PM_{10-2.5}</i>								
OC	1.3	0.25±0.24	1.3	0.67±0.68	0.93	0.26±0.28	N/A	N/A
EC	0.20		0.08		0.06		N/A	N/A
CO ₃ ²⁻	0.02	N/A	0.02	N/A	0.02	N/A	N/A	N/A
Secondary inorganic								
<i>aerosol (µg m⁻³)</i>								
SO ₄ ²⁻	1.6	0.80±0.77	1.5	0.50±0.46	N/A	0.27±0.37	0.44	0.25±0.26
NO ₃ ⁻	1.7	0.90±0.98	0.71	0.54±0.73	N/A	0.24±0.43	0.23	0.20±0.38
NH ₄ ⁺	0.60	0.26±0.35	0.39	0.19±0.25	N/A	0.10±0.21	0.037	0.039±0.075
Inorganic anions								
<i>and cations</i>								
<i>(ng m⁻³)</i>								
Ca ²⁺	500	40.8±29.2	540	28.8±21.6	N/A	23.7±20.1	201	43.7±59.2
Mg ²⁺	122	61.8±44.5	58.5	19.1±13.5	N/A	15.1±11.1	35	36.2±26.6
Na ⁺	583	501±371	85.9	136±113	N/A	102±98.7	80	239±178
K ⁺	170	57.4±36.7	147	49.3±28.6	N/A	27.4±20.2	160	22.8±23.6
Cl ⁻	666	627±538	81.2	115±119	N/A	123±149	114	339±291
Crustal elements								
<i>(ng m⁻³)</i>								
Al	819	24.7±16.2	734	N/A	1000	N/A	254	53.8±36.6

	Birkenes		Hurdal		Kårvatn		Zeppelin	
	Observatory						Observatory	
Fe	540	16.5±16.6	485	N/A	650	N/A	162	26.1±17.5
Ti	21.5	0.80±0.67	19.6	N/A	28.1	N/A	9.7	1.65±0.1.11
Mn	14.3	0.89±0.83	13.1	N/A	16.6	N/A	3.9	0.45±0.29
Organic tracers								
<i>(ng m⁻³)</i>								
<i>Biomass burning</i>								
Levoglucosan	28.5	10.6±11.3	27.8	N/A	33.9	N/A	5.0	0.53±0.65
Mannosan	5.92	2.04±2.63	6.66	N/A	5.48	N/A	0.98	0.082±0.104
Galactosan	1.03	0.52±0.66	1.79	N/A	1.60	N/A	0.27	0.024±0.028
<i>Biogenic Secondary</i>								
<i>Organic Aerosol</i>								
2-methylerythritol	0.667	0.20±0.34	0.551	N/A	0.349	N/A	0.105	0.085±0.199
2-methylthreitol	0.242	0.081±0.128	0.267	N/A	0.122	N/A	0.055	0.042±0.086
<i>Primary Biological</i>								
<i>Aerosol Particles</i>								
Arabitol	16.0	6.57±6.30	17.9	N/A	34.6	N/A	0.66	0.14±0.30
Mannitol	16.0	7.08±6.60	18.1	N/A	34.7	N/A	0.82	0.16±0.34
Glucose	11.7	5.32±4.47	9.00	N/A	13.6	N/A	1.61	0.37±0.47
Trehalose	9.15	4.58±4.94	8.94	N/A	22.2	N/A	1.27	0.21±0.41

1) Based on reconstructed mass


360 We find a lower estimate of the mineral dust concentration at the rural background sites ranging from 5.6-7.6 $\mu\text{g m}^{-3}$ (eq.2)
and an upper estimate ranging from 7.2-8.6 $\mu\text{g m}^{-3}$ (eq.3), whereas 1.9-2.6 $\mu\text{g m}^{-3}$ was attributed to mineral dust at the remote
site. Only CO_3^{2-} was included in the upper estimate for Kårvatn, as data needed to calculate the oxides were missing. The
oxides increased the upper estimate at Birkenes and Hurdal by 14%. If we assume a similar increase at Kårvatn, the upper
estimate increases from 8.6 $\mu\text{g m}^{-3}$ to 9.8 $\mu\text{g m}^{-3}$. The lower estimate provides a 25-36% mass contribution to PM_{10} at the rural
365 background sites whereas the upper estimate ranges from 32-47%. Similarly, at Zeppelin, we found that 31-41% of PM_{10} was
mineral dust. Note though, that we do not have measurements of PM_{10} concentrations at Zeppelin. We therefore assumed that

our PM reconstruction from different constituents (sections 2.1.2 and 2.1.3) explains as much of the total PM₁₀ as it did for the background sites Birkenes and Hurdal, which is on average 83%.

370 Simulations with FLEXPART, based on dust emissions from FLEXDUST, help us to further demonstrate and understand the LRT of mineral dust during this episode. Figure 5 shows the modelled surface concentrations on 27 September 2020, during the storm in Central Asia that caused the large mineral dust emissions, and on 2 October 2020. These time steps were chosen based on the availability of CALIOP overpasses that captured the dust (or BC) plume, which will be discussed later. In addition, we provide an animation of modelled dust surface concentrations between 25 September 2020 and 8 October 2020 as a supplement. FLEXPART results show a distinct dust plume transporting dust from the regions east of the Caspian
375 Sea towards north-west Europe. The dust plume partly overlaps with a wildfire plume starting in Ukraine, as is illustrated by the black lines in the right hand side figures. The contour lines of 500hPa geopotential height illustrate how transport of mineral dust originating from Central Asia is forced on a front between a high-pressure region over Russia and low pressure in Europe. We see enhanced mineral dust concentrations over North Africa and a plume of dust transported over eastern Europe. Dust originating from North Africa is partly mixed into the dust plume from Central Asia yet contributes little to the increased
380 surface concentrations of mineral dust observed in Norway during this episode, as will be further discussed below. Natural dust sources in Central Asia include, e.g., the Karakum and Aralkum deserts in Turkmenistan and Kazakhstan. These deserts are part of what is sometimes referred to as the dust belt, extending from the west coast of Africa to China (Prospero et al., 2002). There is a variety of deserts in Central Asia with different characteristics. Most of these dust sources are active between March and October (Shen et al., 2016; Indoitu et al., 2012). According to our FLEXDUST simulations, a dust storm occurred
385 in Central Asia in the end of September. From 25 September 2020 total dust emissions in this region started increasing, reaching maximum values on 27 September 2020, and slowly decreasing again until the 3 October 2020. Dust emissions were up to 2 g m⁻² h⁻¹ and total emissions from this region amounted to 5.1 Tg in 8 days.

Figure 6 shows the curtain of aerosol extinction profiles at 532 nm (colour coded, the overpass is plotted in Fig 5.) with FLEXPART mineral dust and BC concentration contour lines overlayed. The top panel shows CALIOP aerosol extinction
390 from 26 Sept. 2020, time of first record: 23:48:09, between 36° and 45°E, with dominating aerosol subtype (not shown here) dust and polluted dust, at the surface around 40.5° and 43°E, elsewhere and at elevated levels. The FLEXPART simulation captured both the elevated concentrations near the surface and the vertical spread of the dust plume at the time of emissions and near the source region of the dust (white contours). The dust plumes reached Norway on 2 October 2020. Figure 6 shows CALIOP aerosol extinction from that day, at around midnight (middle panel, 13.5-29°E) and around noon (lower panel, 0-
395 9°E). The modelled dust plumes partly coincide with the satellite images. The night-time CALIOP curtain shows an area with enhanced aerosol loading extending to above 5 km of polluted continental/dust west of 16°E, and a lower region with dust/polluted dust east of 20°E. Also note that the lack of absorbing aerosols around 20°E in the satellite images is locally due to obstructions by clouds. Twelve hours later the CALIOP overpass the lowermost tip of Norway, showing mineral dust reaches over 4 km height (around 5°E). An animation of dust RGB (Red, Green, Blue) composite images, which are based on infrared
400 channel data from the Spinning Enhanced Visible and Infrared Imager (SEVIRI) is provided as a supplement. Regions with


bright pink colours² visualises the dust transport over several days (30 September 2020 – 3 October 2020). Images were downloaded from EUMETSAT (<https://eumetview.eumetsat.int/mapviewer/?product=EO:EUM:DAT:MSG:DUST>).




405 **Figure 5** Modelled surface concentrations of BC from biomass burning (a, c, e) and mineral dust (b, d, f) on 27 September 2020 and 2 October 2020 (midnight and noon). Blue contours: ECMWF 500 hPa geopotential height (m). In the right column black contour lines indicate the regions of the modelled wildfire plume ($BC > 0.03 \mu\text{g m}^{-3}$) as shown in the left column. BC simulations started only after 27 September 2020 because no relevant emissions were observed for the episode in Norway. The red lines indicate the location of CALIOP overpasses shown in Fig. 6.

² See https://www-cdn.eumetsat.int/files/2020-04/pdf_rgb_quick_guide_dust.pdf


a) 27.09.2020 03:00



b) 02.10.2020 00:00



c) 02.10.2020 12:00




Aerosol extinction at 532 nm [units: km^{-1}]

410 **Figure 6 Aerosol extinction at 532 nm [units: km⁻¹] during three overpasses (see Fig. 5 for location) is shown as colour plot (a. CALIOP overpass from 26 Sept. 2020, time of first record 23:48, b: 2. October 2020 01:18, and c. 2 October 2020 11:55). The titles of panels a-c indicate the FLEXPART 3-hourly time step corresponding to the overpass. White contours show FLEXPART mineral dust concentrations (thick and thin lines, top: 40 and 6 $\mu\text{g m}^{-3}$, middle: 50 and 10 $\mu\text{g m}^{-3}$, bottom: 15 and 3 $\mu\text{g m}^{-3}$). Magenta contours show FLEXPART BC concentrations (thick and thin lines, top: none, middle: 50 and 5 ng m^{-3} , bottom: 40 and 10 ng m^{-3}).**
415

In Figure 7 we compare time series of the FLEXPART modelled surface concentrations of mineral dust to the observed PM₁₀ concentrations at Birkenes. (Time series of PM₁₀ are not available at the other stations included in the model results.) We further show surface concentrations from the CAMS regional model product over Europe, which includes the three stations in Norway but not the remote station in the Arctic. Both FLEXPART and CAMS regional show increased surface concentrations of mineral dust at all three stations between 28 and 30 September 2020 and 1 and 4 October 2020. While the first peak is not seen clearly in the observation the second peak is visible in both models for all stations. The timing of the second peak value appears similar, yet continuation of the event is different. Moreover, the estimated values of surface concentrations are larger in FLEXPART than in CAMS regional. This is even more obvious for the weekly averages shown in the bottom panel of Figure 7, where FLEXPART modelled concentration are a factor of 1.7 to 3.4 higher than CAMS. The lower estimates of mineral dust by CAMS are probably a result of various factors. One issue was apparent when we examined the dust concentrations from each of the nine ensemble members. At Birkenes all models and at K arvatn all except for one of the models underestimated the dust concentration. We also found that the EURAD and MOCAGE models tended to greatly underestimate the dust concentrations (values ranging from 0.1 – 3.1 $\mu\text{g m}^{-3}$), which will contribute to an overall lowering of the median. For the MOCAGE model, this has been linked to the deposition of desert dust being too high and an incorrect treatment of the boundary conditions from the CAMS global model in a previous MOCAGE model cycle and both issues have been subsequently corrected (Mathieu Joly, personal comm.). Furthermore, likely causes of differences between the models are for instance included dust sources, emission and scavenging parameterizations. The dust plume from Central Asia did not reach Zeppelin until 4 October and peak surface concentrations are lower than for the rural stations, complying with the observations. Comparison to average dust concentrations based on 8-10 days sample measurements (Figure 7, bottom) indicates a nice agreement between the model and observations, although modelled dust concentrations are overestimated at Zeppelin. From an additional simulation that only included mineral dust emissions from Central Asia, we find that desert regions in Central Asia are major contributors to the dust episode, accounting for 88% of surface dust concentrations at the three rural background sites. The modelled relative contribution of mineral dust from Central Asia is similar at all sites, pointing at common sources. This agrees with the observed relative composition of the mineral dust elements between the three sites being indistinguishable. At Zeppelin, the model finds a relatively larger contribution from other, most likely local, sources. No indication of this was seen in the observations, suggesting that local mineral dust emissions were overestimated by the model.

420
425
430
435
440



445 **Figure 7** PM₁₀ concentrations observed at Birkenes (a). Mineral dust surface concentrations as simulated with FLEXPART (b) and CAMS-regional (c) at three stations in Norway and the remote station (outside CAMS domain). d); mean concentrations from simulations compared to observations from weekly samples for Birkenes (30 September – 7 October 2020), Hurdal and Kårvatn (28 September – 5 October 2020) and Zeppelin (2 – 12 October 2020). For FLEXPART estimates in d) light green colours indicate dust from global sources and dark green colours indicate the contribution of dust from Central Asia only.

3.2 Biomass burning

450 We calculated that carbonaceous aerosol, which generally is the major fraction of biomass burning aerosol, made a 26-36% contribution to PM_{10} , 34-44% to $PM_{2.5}$, and 21-43% to $PM_{10-2.5}$ (Table 1) when converting OC and EC to account for other elements than just Carbon. Adding CO_3^{2-} , increased the contribution only to a minor extent (1-4%). The carbonaceous aerosol made a similar contribution to reconstructed PM_{10} at Zeppelin (29%) as for the rural background sites, as did CO_3^{2-} (1%).

455 BB aerosol mainly consist of EC and OC. Observed EC levels at the rural background sites were either a record high or top six, considering both the annual (not shown) and the Sept.-Nov. 2010/11-2019 time series (Table 1). OC (in PM_{10}) levels were also highly elevated, i.e., within the 95-99th percentile. The OC level was noticeably higher at Kårvatn and Hurdal compared to Birkenes, whereas EC (in PM_{10}) was highest at Birkenes, although by a short margin (Fig. 2). The OC and EC levels observed at Zeppelin were the highest reported since regular measurements started in 2017, but still 3 times lower compared to the major wildfire episode influencing the site in April/May 2006 (Stohl et al., 2007).

460 The split between the fine and coarse fraction of PM_{10} varied substantially between EC and OC at the rural background sites, and between sites. EC results from incomplete combustion of biomass and fossil fuel and is almost exclusively associated with fine aerosol particles. Hence, one would expect rather high EC values for the fine fraction of PM_{10} . The 50:50 percent split between the fine and coarse fraction of PM_{10} seen at Birkenes therefore is a rare finding, whereas the 79:21 percent and 84:16 percent splits for the two other rural background sites are closer to the long-term mean (97:3). OC was even more skewed towards the coarse fraction than EC, with a fine/coarse split ranging from 34:66 percent at Birkenes to 72:28 percent at Kårvatn. 465 The reason for such an atypically high coarse fraction could be due to condensation, agglomeration, and heterogenic chemical reactions influencing the size distribution of carbonaceous aerosol, and, for OC, contribution of PBAP. Another possibility is that pyro convection entrains large, partly combusted particles from the ground to such high altitudes that they can be subject to LRT. Further, Dusek et al. (2017) and Yttri et al. (2021) pointed to charring of coarse fraction PBAP during thermal-optical analysis (TOA), forming pyrolytic carbon that is erroneously interpreted as coarse EC. In fact, the episode coincided with the time of the year when PBAP peak (early fall) (Yttri et al., 2007a; Yttri et al., 2021), and the PBAP tracer levels were highly elevated but still comparable to levels previously seen at this time of the year (Fig.4, Table 1). 470

A lower estimate of 12-16% (Eq. 7) was calculated for BB_{mass} to PM_{10} at the rural background sites, whereas the upper estimate ranged from 17-21%. The lower estimate apportioned only $7.6 \pm 1.8\%$ of PM_{10} to BB_{mass} at Zeppelin, which likely experienced 475 a more pronounced degradation of levoglucosan due to its remote location. The upper estimate of 17% is in line with that observed at the rural background sites.

Estimates of the major carbonaceous aerosol (OC and EC) sources (BB, BSOA, FF, and PBAP) can be derived from the source specific organic tracers listed in Table 1 (see section 2.1.3 for details). Here we estimated the $PBAP_{mass}$, $BSOA_{mass}$ and FF_{mass} contribution to PM_{10} for a better constraint and understanding of the BB source, finding that that their joint contribution made 480 an equally large contribution to PM_{10} as the lower estimate of BB at both the rural background sites (13-17%) and at the remote

site (6.4%), underlining the importance of the BB source. Combined, the contribution of BB_{mass} (lower estimate), $PBAP_{\text{mass}}$, $BSOA_{\text{mass}}$ and FF_{mass} ought to match the observed level of the carbonaceous aerosol (here: Sum of OM and $EC \times 1.1$). For the rural background sites this matched quite well ($99 \pm 21\%$), whereas it only amounted to 48% at the remote Arctic site. The most likely explanation of this discrepancy is failure to account for degradation of organic tracers, and particularly levoglucosan.

485 There are indications that 2-methyltetrols have short atmospheric lifetimes as well (Yttri et al., 2021), whereas the low levels of PBAP tracers at Zeppelin likely reflect the scarce vegetation of the Arctic biome. Further, the ER ratios used for PBAP and BSOA (sect. 2.1.4) are derived from measurements in the Boreo-Nemoral biome, thus their suitability in the Arctic biome is a matter of discussion. Consequently, the apportionment of BB_{mass} , $PBAP_{\text{mass}}$, $BSOA_{\text{mass}}$ and FF_{mass} is associated with greater uncertainty than for the rural background sites.

490 High time resolution measurements of eBC at Birkenes was attributed to a biomass burning fraction (eBC_{BB}) and a fossil fuel combustion fraction (eBC_{FF}), apportioning 43% to eBC_{BB} and 57% to eBC_{FF} for the episode in question (midnight 1-2 October to midnight 3-4 October 2020). Extending the period to match that of the weekly sample (30 September–7 October 2020) reduced the eBC_{BB} fraction to 35%, whereas the eBC_{FF} fraction increased to 65%. The $eBC_{\text{BB}}/eBC_{\text{FF}}$ split is thus comparable to the levoglucosan approach, which apportioned equally large shares to EC_{BB} and EC_{FF} for 30 September-7 October but note

495 that the range ($50 \pm 20\%$) of the levoglucosan approach is very wide. eBC_{BB} ($R^2=0.82$) correlated higher with the high time resolution measurements of PM_{10} at Birkenes than eBC_{FF} ($R^2=0.67$), suggesting that biomass burning emissions were more important for the evolution of PM than fossil fuel sources. eBC_{FF} explained 60% of eBC at Zeppelin and eBC_{BB} 40%, considering both the episode (2-7 October 2020) and the longer period covered by the filter sample (2-12 October 2020). As for Birkenes, this corresponds with the levoglucosan estimate for EC, which apportions an equally large fraction to EC_{FF} and

500 EC_{BB} .

Time series of eBC at Birkenes and Zeppelin are shown in Figures 8 and 9, indicate that eBC values from both biomass burning and fossil fuel were of similar magnitude and both reached peak values on 3 October 2020 at Birkenes and on 5 October 2020 at Zeppelin. Also, modelled black carbon from biomass burning peaks on 3 October 2020 at Birkenes, yet simulated values are underestimated by roughly a factor 10 compared to observations. BC concentrations at the other background stations are

505 of similar magnitude, while there is a delay in peak concentration at Kårvatn compared to the other stations. The high correlation ($R^2=0.92$) between eBC and the major OM fraction at Zeppelin (Fig. 9), obtained from the collocated Aerosol Chemical Speciation Monitor Time-of-Flight (ACSM-ToF) instrument, points to combustion sources as the origin of OM. The correlation was more pronounced for eBC_{FF} ($R^2=0.83$) than for eBC_{BB} ($R^2=0.66$). The episode of enhanced BC from fossil fuel combustion is not reproduced in our model results for the rural background stations and is strongly underestimated at Zeppelin.

510 Explanations can be found in the use of monthly mean fossil fuel BC emissions that could average out effects of short-term emissions.




Figure 8 eBC_{BB} (a) and eBC_{FF} (b) concentrations retrieved with PMF from observations at Birkenes (black) and simulated with FLEXPART at Birkenes, Hurdal and Kårvatn.

515




Figure 9 eBC_{BB} (a) and eBC_{FF} (b) concentrations retrieved with PMF from observations (black) and simulated with FLEXPART (blue) and observed OM concentrations (green) at Zeppelin.


520 We further investigate the LRT of BC from biomass burning on maps and cross-sections coinciding with CALIOP
 observations, like for mineral dust (Figure 5 and Figure 6). In addition, we provide an animation of modelled BC from biomass
 burning surface concentrations between 27 September 2020 and 8 October 2020 as a supplement. Although BC from biomass
 burning is strongly underestimated at Birkenes, the model results do show elevated BC concentrations over southern Norway
 on 2 October 2020. In the CALIOP profiles, modelled BC plumes (magenta contour lines) coincide with regions of very strong
 525 aerosol extinction, though being mainly categorized as dust/polluted (22-28°E). While not explicitly shown here, we would

like to note, that also the TROPOspheric Monitoring Instrument (TROPOMI) on board of the Copernicus Sentinel-5 Precursor satellite, detects enhanced aerosol index values and increased CO total column over the Baltic countries on October 1-2, which confirms the presence of UV absorbing aerosols, including biomass burning aerosols, in this region. In the region at approximately 20°E on 2 October 2020 00:00 a clear BC signal is modelled, yet absent in the CALIOP profile due to clouds.

530 We use backward modelling to characterize the source regions for the observed air pollution. The source -receptor relationship (SRR) for BC at Birkenes during this episode is shown in Figure 10 (top panel). The SRR indicates how sensitive the concentrations at the receptor are to emissions in different source regions. As for the dust plume, LRT is strongly influenced by a low- and high-pressure system (Fig. 5). The strong winds from the east confine the SRR mostly to a region extending between Birkenes and Central Asia. Wildfires contributing to the BC concentrations at Birkenes are thus mostly restricted to

535 this region, and largest contributions are seen in Ukraine and southern Russia, although some contributions from fires in North America and northern Russia are seen as well. Due to the combination of a strong underestimation of modelled BC from biomass burning and good representation of dust concentrations at our stations, we expect that the BC emissions are a larger cause of error than the actual atmospheric transport modelling. It is thus likely that we are missing some sources contributing to the plume (Figure 10, top panel) or that locations are correct, yet total emissions are underestimated. Fossil fuel emissions

540 are more widespread and mostly emissions in southern Sweden and from the Baltic States down to Ukraine contribute to the BC concentrations at Birkenes (Figure 10, bottom) in this event. Similar results were seen at the stations Kårvatn and Hurdal (not shown). At Zeppelin, there was an enhanced influence of fossil fuel emissions in northern Europe on BC levels.



545 **Figure 10 BC source-receptor relationship at Birkenes (2 October 2020 00:00 – 4 October 2020 00:00) modelled with FLEXPART in backwards mode (a) and BC fossil fuel emissions (b). Markers in the top panel indicate locations with BC emissions from biomass burning that contribute $0.1 > \text{ng m}^{-3}$ to BC concentrations at Birkenes. Only emissions that contribute to the modelled concentrations at Birkenes are included in the lower panel. The source-receptor relationship includes air masses up to 100 m above ground level.**

550 **4. Conclusions**

An exceptional episode of elevated PM_{10} concentrations at several measurement sites, with levels exceeding those on any other occasion in the last 10 years, was observed on 2 and 3 October 2020 in Norway, and elsewhere in northern Europe. We have analysed this episode based on surface observations, satellite observations, and atmospheric transport modelling.

LRT of PM was recorded with satellite observations, showing aerosols above the cloud layer. Radiative transport simulations with concentration profiles of different aerosol types were performed to gain an understanding of the processes influencing radiances observed by the OLCI (Sentinel-3). These indicated that there was a contribution from both absorbing and scattering particles for this episode. The rather larger contribution of coarse fraction relative to the fine fraction of PM_{10} , however, pointed

more in the direction of mineral dust. Chemical analysis of surface samples was thus necessary to determine the sources. In our surface observations, a clear influence of biomass burning was revealed (12-21%), as well as mineral dust (25-47%).

560 Surface concentrations of crustal elements Al, Fe, Ti and Mn all strongly exceeded previous maximum recordings. At Birkenes, crustal element levels were 16 to 30 times higher than long-term mean values. Contributions of mineral dust were estimated to be 25-36% of PM₁₀ as a minimum and 31-47% as a maximum. The ratios between crustal element levels were similar at all stations and pointed to a common dust source. Based on atmospheric transport simulations, we concluded that Central Asia, including the Karakum desert, was the main source of mineral dust observed in Norway during this episode, contributing
565 roughly 88% to surface dust concentrations.

Coincidentally, biomass burning emissions in the same transport pathway as the mineral dust plume, caused simultaneous peaks in PM₁₀. Contributions of biomass burning to PM₁₀ were estimated to be 12-16% up to 17-21% at the rural background stations and 8-17% at the remote site. Fires in Ukraine were a source of LRT to the background stations, as shown with backwards transport modelling. The model results however, underestimated the retrieved BC levels at Birkenes and Zeppelin
570 based on surface eBC observations. It could thus be that emissions in this region were underestimated, or that additional sources were relevant. A qualitative comparison between CALIOP observations and FLEXPART model output, although limited due to cloud coverage, suggested that the model does capture the location of the BC plume.

Our analysis reveals how unrelated emission sources can combine during long range transport to cause extreme adverse air quality events in Norway. Considering the 24-hour PM₁₀ EU-limit of 50 µg m⁻³, a total of 70 exceedance days at 39 stations
575 were observed in this single event. It thereby corresponds to 18% of the annual mean number of exceedance days in Norway. This shows the large impact LRTepisodes may have on air quality regulations. The combined use of remote sensing, high quality measurements and transport modelling proved effective in describing the episode and distinguishing its causes.

Further research is needed to assess whether there are reasons to assume that this kind of episodes may occur more frequently in future. Emissions from wildfires and mineral dust sources are sensitive to changes in climate, land use and human activities
580 and different temporal and spatial scales should be considered. Also, changes in atmospheric transport patterns could affect the occurrence of LRT-episodes in Europe. Future studies should thus include earth system model simulations to give us a better understanding in the occurrence of such episodes in future, although especially the complexity due to human component will make it difficult to draw conclusions.

Data availability

585 All in situ data are reported to the EMEP monitoring programme (Tørseth et al., 2012) and are available from the database infrastructure EBAS (<http://ebas.nilu.no/>) hosted at NILU. Measurement data of PM from Norwegian cities and municipalities are available at: api.nilu.no. FLEXPART simulation results are available from the authors upon request.

Author contributions

KEY, WA and MJ provided analysis and interpretation of surface observations. SMP provided eBC PMF analysis. HU
590 provided (historical) elemental data. AK prepared analysis of the OLCI data and radiative transfer modelling. KS and SE
prepared visualisation and interpretation of CALIOP data. PH contributed to retrieval and interpretation of CAMS regional
dust product. NE, SE and CGZ performed FLEXDUST and FLEPXART simulations and prepared analysis and visualisation
of model results. CGZ and KEY wrote the original draft of the paper. All authors contributed with critical review, commentary,
and editing to the manuscript.

595 Acknowledgements

We thank NASA and CNES engineers and scientists for making CALIOP data available. The lidar data were downloaded from
the ICARE Data and Service Center. Furthermore, we would like to acknowledge EUMETSAT for making Dust RGBs
available for download. The in-situ data used are part of the Norwegian national monitoring program (Aas et al., 2020) funded
by the Norwegian Environment Agency, except the tracer analysis at Zeppelin which is funded by the Ministry of Climate and
600 Environment, while tracer analysis at Birkenes is funded by NILU. The research leading to these results has benefited from
Aerosols, Clouds, and Trace gases Research InfraStructure (ACTRIS), funding from the European Union Seventh Framework
Programme (FP7/2007–2013) under ACTRIS-2 and the grant agreement no. 262254, and the COST Action CA16109,
Chemical On-Line cOmpoSition and Source Apportionment of fine aerosol (COLOSSAL).

The manuscript contains modified Copernicus Atmosphere Monitoring Service Information 2020, neither the European
605 Commission nor ECMWF is responsible for any use that may be made of the information it contains.

References

- Aas, W., Eckhardt, S., Fiebig, M., Platt, S.M., Solberg, S., Yttri, K. E., and Zwaafink Groot, C.: Monitoring of long-range
transported air pollutants in Norway, annual report 2020, Miljødirektoratet rapport, NILU, Kjeller, Norway, M-2072/2021
610 NILU OR 13/2021, 2021.
- Aftenposten: Mystisk røyk i Midt-Norge kan komme fra Øst-Europa, in: Aftenposten,
<https://www.aftenposten.no/norge/i/JJ1PyP/mystisk-roeyk-i-midt-norge-kan-komme-fra-oest-europa>, 2020.
- Akritidis, D., Katragkou, E., Georgoulas, A. K., Zanis, P., Kartsios, S., Flemming, J., Inness, A., Douros, J., and Eskes, H.: A
complex aerosol transport event over Europe during the 2017 Storm Ophelia in CAMS forecast systems: analysis and
615 evaluation, *Atmos. Chem. Phys.*, 20, 13557-13578, 10.5194/acp-20-13557-2020, 2020.
- Alastuey, A., Querol, X., Aas, W., Lucarelli, F., Pérez, N., Moreno, T., Cavalli, F., Areskoug, H., Balan, V., Catrambone, M.,
Ceburnis, D., Cerro, J. C., Conil, S., Gevorgyan, L., Hueglin, C., Imre, K., Jaffrezo, J. L., Leeson, S. R., Mihalopoulos, N.,
Mitosinkova, M., O'Dowd, C. D., Pey, J., Putaud, J. P., Riffault, V., Ripoll, A., Sciare, J., Sullegrì, K., Spindler, G., and

- Yttri, K. E.: Geochemistry of PM₁₀ over Europe during the EMEP intensive measurement periods in summer 2012 and
620 winter 2013, *Atmos. Chem. Phys.*, 16, 6107-6129, 10.5194/acp-16-6107-2016, 2016.
- Anderson, G. P., Clough, S. A., Kneizys, F., Chetwynd, J. H., and Shettle, E. P.: AFGL atmospheric constituent profiles (0.120
km), AIR FORCE GEOPHYSICS LAB HANSCOM AFB MA, 1986.
- Ansmann, A., Bösenberg, J., Chaikovsky, A., Comerón, A., Eckhardt, S., Eixmann, R., Freudenthaler, V., Ginoux, P.,
Komguem, L., Linné, H., Márquez, M. Á. L., Matthias, V., Mattis, I., Mitev, V., Müller, D., Music, S., Nickovic, S., Pelon,
625 J., Sauvage, L., Sobolewsky, P., Srivastava, M. K., Stohl, A., Torres, O., Vaughan, G., Wandinger, U., and Wiegner, M.:
Long-range transport of Saharan dust to northern Europe: The 11–16 October 2001 outbreak observed with EARLINET,
J. Geophys. Res., 108, <https://doi.org/10.1029/2003JD003757>, 2003.
- Bodí, M. B., Martín, D. A., Balfour, V. N., Santín, C., Doerr, S. H., Pereira, P., Cerdà, A., and Mataix-Solera, J.: Wildland fire
ash: Production, composition and eco-hydro-geomorphic effects, *Earth-Science Reviews*, 130, 103-127,
630 <https://doi.org/10.1016/j.earscirev.2013.12.007>, 2014.
- Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B.,
Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H.,
Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U.,
Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the
635 climate system: A scientific assessment, *J. Geophys. Res.*, 118, 5380-5552, 10.1002/jgrd.50171, 2013.
- Buras, R., Dowling, T., and Emde, C.: New secondary-scattering correction in DISORT with increased efficiency for forward
scattering, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 112, 2028-2034, 2011.
- Canonaco, F., Crippa, M., Slowik, J. G., Baltensperger, U., and Prévôt, A. S.: SoFi, an IGOR-based interface for the efficient
use of the generalized multilinear engine (ME-2) for the source apportionment: ME-2 application to aerosol mass
640 spectrometer data, *Atmospheric Measurement Techniques*, 6, 3649-3661, 2013.
- Canonaco, F., Tobler, A., Chen, G., Sosedova, Y., Slowik, J. G., Bozzetti, C., Daellenbach, K. R., El Haddad, I., Crippa, M.,
and Huang, R.-J.: A new method for long-term source apportionment with time-dependent factor profiles and uncertainty
assessment using SoFi Pro: application to 1 year of organic aerosol data, *Atmospheric Measurement Techniques*, 14, 923-
943, 2021.
- 645 Cavalli, F., Viana, M., Yttri, K. E., Genberg, J. and Putaud, J.-P.: Toward a standardised thermal-optical protocol for
measuring atmospheric organic and elemental carbon: the EUSAAR protocol, *Atmos. Meas. Tech.*, 3(1), 79–89,
[doi:10.5194/amt-3-79-2010](https://doi.org/10.5194/amt-3-79-2010), 2010.
- CEN: Ambient air - Standard gravimetric measurement method for the determination of the PM₁₀ or PM_{2.5} mass
650 concentration of suspended particulate matter European Committee for Standardization, Brussels, EN 12341:12014, 2014.
- Collin, G.: Regional Production, Updated documentation covering all Regional operational systems and the ENSEMBLE,
METEO-FRANCE, CAMS50_2018SC2012_D2012.2010.2012-U2013_Models_documentation_202101_v202101, 2021.

- Cox, C., and Munk, W.: Statistics of the sea surface derived from sun glitter, *J. Mar. Res.*, 13, 198-227, 1954a.
- Cox, C., and Munk, W.: Measurement of the roughness of the sea surface from photographs of the sun's glitter, *Josa*, 44, 838-850, 1954b.
- Di Giuseppe, F., Remy, S., Pappenberger, F., and Wetterhall, F.: Improving CAMS biomass burning estimations by means of the Global ECMWF Fire Forecast system (GEFF), *ECMWF Tech. Memo.* 790, 18 pp., [https://www.ecmwf.int/sites/default/files ...](https://www.ecmwf.int/sites/default/files...), 2016.
- Donlon, C., Berruti, B., Buongiorno, A., Ferreira, M. H., Féménias, P., Frerick, J., Goryl, P., Klein, U., Laur, H., Mavrocordatos, C., Nieke, J., Rebhan, H., Seitz, B., Stroede, J., and Sciarra, R.: The Global Monitoring for Environment and Security (GMES) Sentinel-3 mission, *Remote Sensing of Environment*, 120, 37-57, <https://doi.org/10.1016/j.rse.2011.07.024>, 2012.
- Dusek, U., Hitzenberger, R., Kasper-Giebl, A., Kistler, M., Meijer, H. A. J., Szidat, S., Wacker, L., Holzinger, R., and Röckmann, T.: Sources and formation mechanisms of carbonaceous aerosol at a regional background site in the Netherlands: insights from a year-long radiocarbon study, *Atmos. Chem. Phys.*, 17, 3233-3251, 10.5194/acp-17-3233-2017, 2017.
- Emde, C., Buras-Schnell, R., Kylling, A., Mayer, B., Gasteiger, J., Hamann, U., Kylling, J., Richter, B., Pause, C., and Dowling, T.: The libRadtran software package for radiative transfer calculations (version 2.0. 1), *Geosci. Mod. Dev.*, 9, 1647-1672, 2016.
- Evangelou, N., Balkanski, Y., Hao, W. M., Petkov, A., Silverstein, R. P., Corley, R., Nordgren, B. L., Urbanski, S. P., Eckhardt, S., and Stohl, A.: Wildfires in northern Eurasia affect the budget of black carbon in the Arctic—a 12-year retrospective synopsis (2002–2013), *Atmos. Chem. & Phys.*, 16, 7587-7604, 2016.
- Evangelou, N., Kylling, A., Eckhardt, S., Myroniuk, V., Stebel, K., Paugam, R., Zibitsev, S., and Stohl, A.: Open fires in Greenland in summer 2017: transport, deposition and radiative effects of BC, OC and BrC emissions, *Atmos. Chem. & Phys.*, 19, 1393-1411, 2019.
- Fécan, F., Marticorena, B., and Bergametti, G.: Parametrization of the increase of the aeolian erosion threshold wind friction velocity due to soil moisture for arid and semi-arid areas, *Annales Geophysicae*, 17, 149-157, 10.1007/s00585-999-0149-7, 1999.
- Gasteiger, J., and Wiegner, M.: MOPSMAP v1.0: a versatile tool for the modeling of aerosol optical properties, *Geosci. Model Dev.*, 11, 2739-2762, 10.5194/gmd-11-2739-2018, 2018.
- Ginoux, P., Chin, M., Tegen, I., Prospero, J. M., Holben, B., Dubovik, O., and Lin, S. J.: Sources and distributions of dust aerosols simulated with the GOCART model, *Journal of Geophysical Research-Atmospheres*, 106, 20255-20273, 10.1029/2000jd000053, 2001.
- Groot Zwaafink, C. D., Grythe, H., Skov, H., and Stohl, A.: Substantial contribution of northern high-latitude sources to mineral dust in the Arctic, *J. Geophys. Res.*, 121, 13,678-613,697, 10.1002/2016JD025482, 2016.

- Grythe, H., Kristiansen, N. I., Groot Zwaafink, C. D., Eckhardt, S., Ström, J., Tunved, P., Krejci, R., and Stohl, A.: A new aerosol wet removal scheme for the Lagrangian particle model FLEXPART v10, *Geosci. Model Dev.*, 10, 1447-1466, 10.5194/gmd-10-1447-2017, 2017.
- 690 Hak, C.: Norway's air quality monitoring network. Assessment of station siting according to regulations in EU's air quality directive, NILU, Norwegian Environment Agency Report M-358/2015, 2015.
- Hao, W. M., and Ward, D. E.: Methane production from global biomass burning, *J. Geophys. Res.*, 98, 20657-20661, 1993.
- Hao, W. M., Petkov, A., Nordgren, B. L., Corley, R. E., Silverstein, R. P., Urbanski, S. P., Evangeliou, N., Balkanski, Y., and Kinder, B. L.: Daily black carbon emissions from fires in northern Eurasia for 2002-2015, *Geoscientific Model Development*. 9: 4461-4474., 9, 4461-4474, 2016.
- 695 Hess, M., Koepke, P., and Schult, I.: Optical Properties of Aerosols and Clouds: The Software Package OPAC, *B. Am. Meteorol. Soc.*, 79, 831-844, 1998.
- Hongisto, M., and Sofiev, M.: Long-range transport of dust to the Baltic Sea region, *Int. J. Environ. Pollut.*, 22, 72-86, 2004.
- Hu, Y., Fernandez-Anez, N., Smith, T., and Rein, G.: Review of emissions from smouldering peat fires and their contribution to regional haze episodes, *Int. J. Wildland Fire*, 27, 293-312, 2018.
- 700 IARC: List of Classifications. Agents classified by the IARC Monographs, in: *IARC Monographs on the Evaluation of Risk to Humans*, 2021.
- Indoitu, R., Orlovsky, L., and Orlovsky, N.: Dust storms in Central Asia: spatial and temporal variations, *Journal of Arid Environments*, 85, 62-70, 2012.
- Jankowski, N., Schmidl, C., Marr, I. L., Bauer, H. and Puxbaum, H.: Comparison of methods for the quantification of carbonate carbon in atmospheric PM10 aerosol samples, *Atmos. Environ.*, 42(34), 8055-8064, doi:10.1016/j.atmosenv.2008.06.012, 2008
- 705 Kaiser, J., Heil, A., Andreae, M., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J.-J., Razinger, M., Schultz, M., and Suttie, M.: Biomass burning emissions estimated with a global fire assimilation system based on observed fire radiative power, *Biogeosc.*, 9, 527, 2012.
- 710 Kaufman, Y., Ichoku, C., Giglio, L., Korontzi, S., Chu, D., Hao, W., Li, R.-R., and Justice, C.: Fire and smoke observed from the Earth Observing System MODIS instrument--products, validation, and operational use, *Int. J. Remote Sens.*, 24, 1765-1781, 2003.
- Klimont, Z., Kupiainen, K., Heyes, C., Purohit, P., Cofala, J., Rafaj, P., Borken-Kleefeld, J., and Schöpp, W.: Global anthropogenic emissions of particulate matter including black carbon, *Atmos. Chem. & Phys.*, 17, 8681-8723, 2017.
- 715 Kukkonen, J., López-Aparicio, S., Segersson, D., Geels, C., Kangas, L., Kauhaniemi, M., Maragkidou, A., Jensen, A., Assmuth, T., Karppinen, A., Sofiev, M., Hellén, H., Riikonen, K., Nikmo, J., Kousa, A., Niemi, J. V., Karvosenoja, N., Santos, G. S., Sundvor, I., Im, U., Christensen, J. H., Nielsen, O. K., Plejdrup, M. S., Nøjgaard, J. K., Omstedt, G., Andersson, C., Forsberg, B., and Brandt, J.: The influence of residential wood combustion on the concentrations of PM2.5 in four Nordic cities, *Atmos. Chem. Phys.*, 20, 4333-4365, 10.5194/acp-20-4333-2020, 2020.

- 720 Kylling, A., Groot Zwaaftink, C. D., and Stohl, A.: Mineral Dust Instantaneous Radiative Forcing in the Arctic, *Geophys. Res. Lett.*, 45, 4290-4298, doi:10.1029/2018GL077346, 2018.
- Laskin, A., Wietsma, T. W., Krueger, B. J., and Grassian, V. H.: Heterogeneous chemistry of individual mineral dust particles with nitric acid: A combined CCSEM/EDX, ESEM, and ICP-MS study, *J. Geophys. Res.*, 110, <https://doi.org/10.1029/2004JD005206>, 2005.
- 725 Laupsa, H., Denby, B., Larssen, S., and Schaug, J.: Source apportionment of particulate matter (PM_{2.5}) in an urban area using dispersion, receptor and inverse modelling, *Atmospheric Environment*, 43, 4733-4744, <https://doi.org/10.1016/j.atmosenv.2008.07.010>, 2009.
- Lelieveld, J., Evans, J. S., Fnais, M., Giannadaki, D., and Pozzer, A.: The contribution of outdoor air pollution sources to premature mortality on a global scale, *Nature*, 525, 367-371, 2015.
- 730 Locker, H. B.: The use of levoglucosan to assess the environmental impact of residential wood-burning on air quality, Dartmouth Coll., Hanover, NH (USA), 1988.
- Long, C. M., Nascarella, M. A., and Valberg, P. A.: Carbon black vs. black carbon and other airborne materials containing elemental carbon: Physical and chemical distinctions, *Environ. Pollut.*, 181, 271-286, <https://doi.org/10.1016/j.envpol.2013.06.009>, 2013.
- 735 Marécal, V., Peuch, V. H., Andersson, C., Andersson, S., Arteta, J., Beekmann, M., Benedictow, A., Bergström, R., Bessagnet, B., Cansado, A., Chéroux, F., Colette, A., Coman, A., Curier, R. L., Denier van der Gon, H. A. C., Drouin, A., Elbern, H., Emili, E., Engelen, R. J., Eskes, H. J., Foret, G., Friese, E., Gauss, M., Giannaros, C., Guth, J., Joly, M., Jaumouillé, E., Josse, B., Kadyrov, N., Kaiser, J. W., Krajsek, K., Kuenen, J., Kumar, U., Liora, N., Lopez, E., Malherbe, L., Martinez, I., Melas, D., Meleux, F., Menut, L., Moinat, P., Morales, T., Parmentier, J., Piacentini, A., Plu, M., Poupkou, A.,
- 740 Queguiner, S., Robertson, L., Rouïl, L., Schaap, M., Segers, A., Sofiev, M., Tarasson, L., Thomas, M., Timmermans, R., Valdebenito, Á., van Velthoven, P., van Versendaal, R., Vira, J., and Ung, A.: A regional air quality forecasting system over Europe: the MACC-II daily ensemble production, *Geosci. Model Dev.*, 8, 2777-2813, 10.5194/gmd-8-2777-2015, 2015.
- Marticorena, B., and Bergametti, G.: Modeling the atmospheric dust cycle: 1. Design of a soil-derived dust emission scheme, *J. Geophys. Res.*, 100, 16415-16430, 10.1029/95JD00690, 1995.
- 745 Myhre, G., D. Shindell, F.-M. Bréon, W. Collins, J. Fuglestedt, J. Huang, D. Koch, J.-F. Lamarque, D. Lee, B. Mendoza, T. Nakajima, Robock, A., Stephens, G., Takemura, T., and Zhang, H.: Anthropogenic and Natural Radiative Forcing, Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, in: *Climate Change 2013: The Physical Science Basis.*, edited by: Stocker, T. F., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA., 2013.
- 750 WHO: WHO Air quality guidelines for particulate matter, ozone, nitrogen dioxide and sulfur dioxide: global update 2005: summary of risk assessment, World Health Organization, 2006.

- Patashnick, H., and Rupprecht, E. G.: Continuous PM-10 Measurements Using the Tapered Element Oscillating Microbalance, *J. Air Waste Manage. Assoc.*, 41, 1079-1083, 10.1080/10473289.1991.10466903, 1991.
- 755 Pisso, I., Sollum, E., Grythe, H., Kristiansen, N. I., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. L., Groot Zwaafink, C. D., Evangeliou, N., Sodemann, H., Haimberger, L., Henne, S., Brunner, D., Burkhart, J. F., Fouilloux, A., Brioude, J., Philipp, A., Seibert, P., and Stohl, A.: The Lagrangian particle dispersion model FLEXPART version 10.4, *Geosci. Model Dev.*, 12, 4955-4997, 10.5194/gmd-12-4955-2019, 2019.
- Prospero, J. M., Ginoux, P., Torres, O., Nicholson, S. E., and Gill, T. E.: ENVIRONMENTAL CHARACTERIZATION OF
760 GLOBAL SOURCES OF ATMOSPHERIC SOIL DUST IDENTIFIED WITH THE NIMBUS 7 TOTAL OZONE MAPPING SPECTROMETER (TOMS) ABSORBING AEROSOL PRODUCT, *Rev. Geophys.*, 40, 2-1-2-31, doi:10.1029/2000RG000095, 2002.
- Putaud, J.-P., Van Dingenen, R., Alastuey, A., Bauer, H., Birmili, W., Cyrys, J., Flentje, H., Fuzzi, S., Gehrig, R., Hansson, H. C., Harrison, R. M., Herrmann, H., Hitzenberger, R., Hüglin, C., Jones, A. M., Kasper-Giebl, A., Kiss, G., Kousa, A.,
765 Kuhlbusch, T. A. J., Löschau, G., Maenhaut, W., Molnar, A., Moreno, T., Pekkanen, J., Perrino, C., Pitz, M., Puxbaum, H., Querol, X., Rodriguez, S., Salma, I., Schwarz, J., Smolik, J., Schneider, J., Spindler, G., ten Brink, H., Tursic, J., Viana, M., Wiedensohler, A., and Raes, F.: A European aerosol phenomenology – 3: Physical and chemical characteristics of particulate matter from 60 rural, urban, and kerbside sites across Europe, *Atmos. Environ.*, 44, 1308–1320, 2010.
- 770 Puxbaum, H., Caseiro, A., Sánchez-Ochoa, A., Kasper-Giebl, A., Claeys, M., Gelencsér, A., Legrand, M., Preunkert, S., and Pio, C.: Levoglucosan levels at background sites in Europe for assessing the impact of biomass combustion on the European aerosol background, *J. Geophys. Res.*, 112, 2007.
- Rémy, S., Kipling, Z., Flemming, J., Boucher, O., Nabat, P., Michou, M., Bozzo, A., Ades, M., Huijnen, V., Benedetti, A., Engelen, R., Peuch, V. H., and Morcrette, J. J.: Description and evaluation of the tropospheric aerosol scheme in the
775 European Centre for Medium-Range Weather Forecasts (ECMWF) Integrated Forecasting System (IFS-AER, cycle 45R1), *Geosci. Model Dev.*, 12, 4627-4659, 10.5194/gmd-12-4627-2019, 2019.
- Saarikoski, S., Sillanpää, M., Sofiev, M., Timonen, H., Saarnio, K., Teinilä, K., Karppinen, A., Kukkonen, J., and Hillamo, R.: Chemical composition of aerosols during a major biomass burning episode over northern Europe in spring 2006: Experimental and modelling assessments, *Atmospheric Environment*, 41, 3577-3589,
780 <https://doi.org/10.1016/j.atmosenv.2006.12.053>, 2007.
- Sandradewi, J., Prévôt, A. S., Szidat, S., Perron, N., Alfarra, M. R., Lanz, V. A., Weingartner, E., and Baltensperger, U.: Using aerosol light absorption measurements for the quantitative determination of wood burning and traffic emission contributions to particulate matter, *Environmental science & technology*, 42, 3316-3323, 2008.
- Shen, H., Abuduwaili, J., Samat, A., and Ma, L.: A review on the research of modern aeolian dust in Central Asia, *Arabian
785 Journal of Geosciences*, 9, 625, 2016.
- Shi, Y., Matsunaga, T., Saito, M., Yamaguchi, Y., and Chen, X.: Comparison of global inventories of CO₂ emissions from biomass burning during 2002–2011 derived from multiple satellite products, *Environ. Pollut.*, 206, 479-487, 2015.

- 790 Simoneit, B. R. T., Schauer, J. J., Nolte, C. G., Oros, D. R., Elias, V. O., Fraser, M. P., Rogge, W. F., and Cass, G. R.:
Levoglucosan, a tracer for cellulose in biomass burning and atmospheric particles, *Atmospheric Environment*, 33, 173-
182, [https://doi.org/10.1016/S1352-2310\(98\)00145-9](https://doi.org/10.1016/S1352-2310(98)00145-9), 1999.
- Sodemann, H., Lai, T. M., Marengo, F., Ryder, C. L., Flamant, C., Knippertz, P., Rosenberg, P., Bart, M., and McQuaid, J. B.:
Lagrangian dust model simulations for a case of moist convective dust emission and transport in the western Sahara region
during Fennec/LADUNEX, *J. Geophys. Res.*, 120, 6117-6144, 10.1002/2015JD023283, 2015.
- 795 Stamnes, K., Tsay, S.-C., Wiscombe, W., and Jayaweera, K.: Numerically stable algorithm for discrete-ordinate-method
radiative transfer in multiple scattering and emitting layered media, *Appl. Opt.*, 27, 2502-2509, 1988.
- Stohl, A., Berg, T., Burkhardt, J., Fjæraa, A., Forster, C., Herber, A., Hov, Ø., Lunder, C., McMillan, W., and Oltmans, S.:
Arctic smoke—record high air pollution levels in the European Arctic due to agricultural fires in Eastern Europe in spring
2006, *Atmos. Chem. & Phys.*, 7, 511-534, 2007.
- 800 Stohl, A., Aamaas, B., Amann, M., Baker, L. H., Bellouin, N., Berntsen, T. K., Boucher, O., Cherian, R., Collins, W., and
Daskalakis, N.: Evaluating the climate and air quality impacts of short-lived pollutants, *Atmos. Chem. & Phys.*, 15, 10529-
10566, 2015.
- Szidat, S., Ruff, M., Perron, N., Wacker, L., Synal, H.-A., Hallquist, M., Shannigrahi, A. S., Yttri, K. E., Dye, C., and Simpson,
D.: Fossil and non-fossil sources of organic carbon (OC) and elemental carbon (EC) in Göteborg, Sweden, *Atmos. Chem.
& Phys.*, 9, 1521-1535, 2009.
- 805 Tarrasón, L. H., Paul David; Guerreiro, Cristina de Brito Beirao; Meleux, Frederik; Rouil, Laurence Interim Annual
Assessment Report for 2018. European air quality in 2018, 2019.
- Turpin, B. J., and Lim, H.-J.: Species contributions to PM_{2.5} mass concentrations: Revisiting common assumptions for
estimating organic mass, *Aerosol Science & Technology*, 35, 602-610, 2001.
- 810 Tørseth, K., Aas, W., Breivik, K., Fjæraa, A. M., Fiebig, M., Hjellbrekke, A.-G., Lund Myhre, C., Solberg, S., and Yttri, K.
E.: Introduction to the European Monitoring and Evaluation Programme (EMEP) and observed atmospheric composition
change during 1972–2009, *Atmos. Chem. & Phys.*, 12, 5447-5481, 2012.
- Wooster, M. J., Roberts, G., Perry, G. L. W., and Kaufman, Y. J.: Retrieval of biomass combustion rates and totals from fire
radiative power observations: FRP derivation and calibration relationships between biomass consumption and fire radiative
energy release, *J. Geophys. Res.*, 110, <https://doi.org/10.1029/2005JD006318>, 2005.
- 815 Yttri, K. E., Canonaco, F., Eckhardt, S., Evangeliou, N., Fiebig, M., Gundersen, H., Hjellbrekke, A.-G., Lund Myhre, C., Platt,
S. M., Prévôt, A. S. H., Simpson, D., Solberg, S., Surratt, J., Tørseth, K., Uggerud, H., Vadset, M., Wan, X., and Aas, W.:
Trends, composition, and sources of carbonaceous aerosol at the Birkenes Observatory, northern Europe, 2001–2018,
Atmos. Chem. Phys., 21, 7149–7170, <https://doi.org/10.5194/acp-21-7149-2021>, 2021
- 820 Yttri, K., Dye, C., and Kiss, G.: Ambient aerosol concentrations of sugars and sugar-alcohols at four different sites in Norway,
Atmos. Chem. & Phys., 7, 4267-4279, 2007a.

- Yttri, K. E., Dye, C., Slørdal, L. H., and Braathen, O.-A.: Quantification of monosaccharide anhydrides by liquid chromatography combined with mass spectrometry: application to aerosol samples from an urban and a suburban site influenced by small-scale wood burning, *J. Air Waste Manage. Assoc.*, 55, 1169-1177, 2005.
- 825 Yttri, K. E., Aas, W., Bjerke, A., Cape, J. N., Cavalli, F., Ceburnis, D., Dye, C., Emblico, L., Facchini, M. C., Forster, C., Hanssen, J. E., Hansson, H. C., Jennings, S. G., Maenhaut, W., Putaud, J. P., and Tørseth, K.: Elemental and organic carbon in PM₁₀: a one year measurement campaign within the European Monitoring and Evaluation Programme EMEP, *Atmos. Chem. Phys.*, 7, 5711-5725, 10.5194/acp-7-5711-2007, 2007b.
- Yttri, K. E., Dye, C., Braathen, O.-A., Simpson, D., and Steinnes, E.: Carbonaceous aerosols in Norwegian urban areas, *Atmos. Chem. & Phys.*, 9, 2007-2020, 2009.
- 830 Yttri, K. E., Lund Myhre, C., Eckhardt, S., Fiebig, M., Dye, C., Hirdman, D., Ström, J., Klimont, Z., and Stohl, A.: Quantifying black carbon from biomass burning by means of levoglucosan—a one-year time series at the Arctic observatory Zeppelin, *Atmos. Chem. & Phys.*, 14, 6427-6442, 2014.
- Yttri, K. E., Simpson, D., Bergström, R., Kiss, G., Szidat, S., Ceburnis, D., Eckhardt, S., Hueglin, C., Nøjgaard, J. K., and Perrino, C.: The EMEP Intensive Measurement Period campaign, 2008–2009: characterizing carbonaceous aerosol at nine
835 rural sites in Europe, *Atmos. Chem. & Phys.*, 19, 4211-4233, 2019.
- Zdráhal, Z., Oliveira, J., Vermeylen, R., Claeys, M., and Maenhaut, W.: Improved method for quantifying levoglucosan and related monosaccharide anhydrides in atmospheric aerosols and application to samples from urban and tropical locations, *Environmental Science & Technology*, 36, 747-753, 2002.
- Zotter, P., Herich, H., Gysel, M., El-Haddad, I., Zhang, Y., Močnik, G., Hüglin, C., Baltensperger, U., Szidat, S., and Prévôt,
840 A. S.: Evaluation of the absorption Ångström exponents for traffic and wood burning in the Aethalometer-based source apportionment using radiocarbon measurements of ambient aerosol, *Atmos. Chem. & Phys.*, 17, 4229-4249, 2017.

Appendix A1: Sample analysis

OC/EC was obtained by Thermal-optical analysis (TOA), using the Lab OC-EC Aerosol Analyser (Sunset Laboratory Inc) and
845 according to the EUSAAR-2 temperature program (Cavalli et al., 2010). The samples content of CO_3^{2-} - carbon was
determined as for OC and EC but after subjecting a punch of the filter sample to thermal-oxidative pre-treatment (Jankowski
et al., 2008; Evangelidou et al., 2016). The samples OC and EC content were corrected with respect to the CO_3^{2-} - carbon.

Organic tracers (levoglucosan, mannosan, galactosan, arabitol, mannitol, glucose, trehalose, 2-methylerythritol and 2-
methylthreitol) were analyzed using Waters Acquity ultra-performance liquid chromatography (UPLC) in combination with
850 Waters Premier XE high-resolution time-of-flight mass spectrometry (HR-TOF-MS) operated in the negative electrospray
ionization (ESI) mode. The analytical methodology is based on that of Dye and Yttri (2005) but deviates by choice of column
(two 2.1_150mm HSS T3, 1.8 μm , Waters Inc.). All species were identified by retention time and mass spectra of authentic
standards and isotope-labelled standards of levoglucosan, galactosan, mannitol, arabitol, trehalose and glucose were used as
recovery standard (see Yttri et al., 2021).

855 Al, Fe, Mn and Ti were analyzed by ICP-MS (Inductively Coupled Plasma/Mass Spectrometry) (Agilent 7700x). Prior to
analysis, aerosol filter samples were extracted (diluted supra pure HNO_3), digested (75 min; max temp of 250 $^\circ\text{C}$ for 15
min) using an UltraClave microwave system (Milestone, Italy), and diluted to 10 ml (ion exchanged H_2O). External calibration
was applied, and calibration standards made of HNO_3 (supra pure) (10% v/v) to adapt to the sample matrix. Indium was used
as recovery standard and applied to all samples, standards, blank filters, and reference materials.

860 Prior to ion chromatography analysis, filter samples were soaked in Milli-Q water (10 ml) and subjected to ultrasonic agitation
(30 min). Extracts were analyzed with respect to Ca^{2+} , K^+ , Mg^{2+} , Na^+ , and NH_4^+ on a Dionex Integriion ion chromatograph,
using a Dionex cation exchange CS16 column (3 mm x 250 mm), and a conductivity detector. Samples were eluted
using methane sulphonic acid (34 mM) at a flow rate of 0.5 ml min^{-1} . Cl^- , NO_3^- , and SO_4^{2-} were analyzed on
a Dionex Integrion ion chromatograph, using a Dionex anion exchange AS9-SC column (4 mm x 250 mm), and a conductivity
865 detector. Samples were eluted using carbonate (K_2CO_3 , 2.0 mM; HCO_3^- , 0.75mM) at a flow rate of 2 ml min^{-1} .