



Atmospheric Measurements at the Foot and the Summit of Mt. Tai - Part I: HONO Formation and Its Role in the Oxidizing Capacity of the Upper Boundary Layer

5

Chaoyang Xue^{1, 2*}, Can Ye^{1, 9}, Jörg Kleffmann³, Chenglong Zhang^{1, 4}, Valéry Catoire², Fengxia Bao⁵, Abdelwahid Mellouki^{6, 7}, Likun Xue⁷, Jianmin Chen⁸, Keding Lu⁹, Yong Zhao¹⁰, Hengde Liu¹⁰, Zhaoxin Guo¹⁰, Yujing Mu^{1, 4*}

¹ Research Centre for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085, China

10 ² Laboratoire de Physique et Chimie de l'Environnement et de l'Espace (LPC2E), CNRS–Université Orléans–CNES, Cedex 2, Orléans 45071, France

³ Physical and Theoretical Chemistry, University of Wuppertal, Gaußstrasse 20, Wuppertal 42119, Germany

⁴ Centre for Excellence in Regional Atmospheric Environment, Institute of Urban Environment, Chinese Academy of Sciences, Xiamen 361021, China

15 ⁵ Multiphase Chemistry Department, Max Planck Institute for Chemistry, Mainz 55128, Germany

⁶ Institut de Combustion Aérothermique, Réactivité et Environnement, Centre National de la Recherche Scientifique (ICARE-CNRS), Cedex 2, Orléans 45071, France

⁷ Environmental Research Institute, Shandong University, Qingdao, Shandong 266237, China

20 ⁸ Shanghai Key Laboratory of Atmospheric Particle Pollution and Prevention, Department of Environmental Science and Engineering, Institute of Atmospheric Sciences, Fudan University, Shanghai 200438, China

⁹ State Key Joint Laboratory of Environment Simulation and Pollution Control, College of Environmental Sciences and Engineering, Peking University, Beijing, 100871, China

¹⁰ Taishan National Reference Climatological Station, Tai'an, Shandong, 271000, China

Correspondence to:

25 Chaoyang Xue (chaoyang.xue@cirs-orleans.fr; 86chaoyang.xue@gmail.com)

Yujing Mu (yjmu@rcees.ac.cn)



Abstract

A comprehensive field campaign, with measurements of HONO and related parameters, was conducted in summer 2018 at the
30 foot (150 m a.s.l.) and the summit (1534 m a.s.l.) of Mt. Tai (Shandong province, China). At the summit station, high HONO
mixing ratios were observed during this campaign (mean $\pm 1\sigma$: 133 ± 106 pptv, maximum: 880 pptv), with a diurnal noontime
peak (mean $\pm 1\sigma$: 133 ± 72 pptv at 12:30 local time). Constraints on the kinetics of aerosol-derived HONO sources (NO_2
uptake on the aerosol surface and particulate nitrate photolysis) were performed and discussed, which enables a better
understanding of the interaction of HONO and aerosols, especially in the polluted North China Plain. Various shreds of
35 evidence of air mass transport from the ground to the summit levels were provided. Furthermore, daytime HONO formation
from different paths and its role in radical production were quantified and discussed.

We found that the homogeneous reaction $\text{NO} + \text{OH}$ could only explain 8.0% of the daytime HONO formation, resulting in
strong unknown sources (P_{un}). Campaign-averaged P_{un} was about 290 ± 280 pptv h^{-1} with a maximum of about 1800 pptv
 h^{-1} . Aerosol-derived HONO formation mechanisms were not the major sources of P_{un} . Their contributions to daytime HONO
40 formation varied from negligible to moderate (similar to $\text{NO} + \text{OH}$), depending on the used chemical kinetics. Coupled with
sensitivity tests on the used kinetics, the NO_2 uptake on the aerosol surface and particulate nitrate photolysis contributed 1.5 –
19% and 0.6 – 9.6% of the observed P_{un} , respectively. Based on synchronous measurements at the foot and the summit stations,
a bunch of field evidence was proposed to support that the remaining majority (70 – 98%) of P_{un} was dominated by the rapid
vertical transport from the ground to the summit levels and heterogeneous formation on the ground surfaces during the transport.
45 HONO photolysis at the summit level initialized daytime photochemistry and represented an essential HO_x ($\text{OH} + \text{HO}_2$) source
in the daytime, with a contribution of 26%, more than one-third of that of O_3 . We provided evidence that ground-derived
HONO played a significant role in the oxidizing capacity of the upper boundary layer through the enhanced vertical air mass
exchange driven by mountain winds. The follow-up impacts should be considered in the regional chemistry-transport models.

1 Introduction

50 In the past two decades, atmospheric nitrous acid (HONO) has attracted numerous laboratory experiments and field campaigns
because of its significant contribution to the atmospheric concentration of hydroxyl radicals (OH) and the incomplete
understanding of its sources (Kleffmann, 2007). Besides the homogeneous reaction of NO with OH, various HONO formation
pathways were proposed, including: a) emission from combustion processes, e.g., vehicle exhaust, domestic combustion and
biomass burning (Klosterkötter et al., 2021; Kramer et al., 2020; Kurtenbach et al., 2001; Liu et al., 2017; Peng et al., 2020;
55 Theys et al., 2020); b) heterogeneous dark and photosensitized reactions of NO_2 on surfaces such as soot (Ammann et al., 1998;
Monge et al., 2010), organic compounds (George et al., 2005; Han et al., 2017; Stemmler et al., 2006, 2007), acids (Kleffmann
et al., 1998), urban grime (Liu et al., 2019a), MgO (Ma et al., 2017), mineral dust (Ndour et al., 2008), vegetation leaves
(Marion et al., 2021), etc.; c) photolytic reactions of total nitrate (particulate nitrate and adsorbed nitric acid) (Bao et al., 2018;
Laufs and Kleffmann, 2016; Ye et al., 2016; Zhou et al., 2003, 2011) and ortho-nitrophenols (Bejan et al., 2006); d) emissions



60 from soil (Donaldson et al., 2014; Oswald et al., 2013; Su et al., 2011), etc. Even though many potential HONO sources have been identified in the past, there is still a significant gap between model results and observations (Fu et al., 2019; Liu et al., 2017; Xue et al., 2020; Zhang et al., 2019a, 2019b). One of the critical puzzles is the quantity of HONO formation from the aerosol-derived sources, particularly NO₂ uptake on aerosol surfaces and aerosol nitrate photolysis in high-aerosol regions such as the North China Plain (NCP).

65 The NO₂ uptake on aerosol surfaces was proposed to be much less important than that on ground surfaces in previous studies because of the low S/V (surface to volume ratio) of particles compared to ground surfaces and the similar reaction kinetics on the same types of surfaces (Nie et al., 2015; Stemmler et al., 2007). However, the contribution of NO₂ uptake on aerosol surfaces to HONO formation in the extremely polluted region is not well constrained. For example, previous studies using box models or regional transport chemistry models found the NO₂ uptake on aerosol surfaces lead to a negligible impact on daytime
70 HONO formation in the polluted NCP (Liu et al., 2019b; Xue et al., 2020; Zhang et al., 2019a, 2019b). Nevertheless, a recent chamber study (Ge et al., 2019) found a high dark NO₂ uptake coefficient (2.0×10^{-5} to 1.7×10^{-4}) on NaCl particles under high RH (90%), NH₃ (50-2000 ppbv), and SO₂ (600 ppbv) conditions. First, such severe pollution rarely occurred. Second, if such a high NO₂ coefficient on the aerosol surface was applied in night-time HONO budget analysis, the dominant role of NO₂ uptake on the ground surface in night-time HONO formation, which was already generally accepted, might be challenged
75 (Kleffmann, 2007; Kurtenbach et al., 2001; Stutz et al., 2002; Xue et al., 2020). Besides, recent nocturnal vertical measurements of HONO in Beijing found both ground-based and aerosol-derived sources may play important roles in HONO formation during the clean period and haze period, respectively (Meng et al., 2020). Therefore, the contribution of NO₂ uptake on the aerosol surface to HONO formation still needs more field constraints.

The photolysis of particulate nitrate (pNO₃) was found to be an important HONO source in low NO_x areas such as forest
80 canopy and marine boundary layer. High enhancement factors ($EF = J(\text{pNO}_3)/J(\text{HNO}_3)$), within the range of tens to thousands, were proposed in forest area, marine boundary layer, and polluted areas like the NCP (Bao et al., 2020; Ye et al., 2016, 2017; Zhou et al., 2007, 2011). However, model studies with field constraints (Romer et al., 2018; Xue et al., 2020) found that the EF was moderate (7-30) rather than tens to thousands obtained in laboratory studies (Bao et al., 2020; Ye et al., 2016, 2017; Zhou et al., 2007). Moreover, a recent laboratory flow tube study (Wang et al., 2021) revealed that the EF was lower than 1 in
85 the aqueous phase. Another flow tube study (Laufs and Kleffmann, 2016) also reported a slow HONO formation from secondary heterogeneous reactions of NO₂ produced during HNO₃ photolysis. Besides, a very recent chamber study (Shi et al., 2021) found that the EF values of airborne nitrate were lower than 10 (generally around 1), which also indicates an insignificant contribution of nitrate photolysis to HONO formation. Furthermore, when considering the large variation of EF values (from digits to thousands) in the model, model performance on HONO simulations could be improved but accompanied by large
90 uncertainties (Fu et al., 2019; Liu et al., 2019b). Therefore, HONO formation from nitrate photolysis still needs more field constraints.

In addition, the role of HONO photolysis in the oxidizing capacity of the upper boundary layer remains unclear. As there exists a significant gradient in HONO distribution from the ground level to the upper troposphere, HONO photolysis was accounted



to be much less important compared to O_3 photolysis (Ye et al., 2018; Zhang et al., 2009). However, in mountainous regions,
95 mountain winds, including mountain breeze (downslope) and valley breeze (upslope) can accelerate the air mass exchange
between the mountain top and the ground levels, which may affect HONO levels and the atmospheric oxidizing capacity at
the summit level (Jiang et al., 2020; Schmid et al., 2020; Ye et al., 1987).


Herein, atmospheric measurements at the foot (~150 m a.s.l.) and the summit (~1534 m a.s.l.) of Mt. Tai allow us to understand
more about 1) the transport of ground-formed HONO and its role in the upper boundary layer; 2) HONO formation from the
100 aerosol-derived sources as the ground sources might be less effective; 3) the oxidizing capacity of the upper boundary layer
and its contributors.

2 Experimental

2.1 Site Description

HONO was alternately measured at two locations: the foot and the summit of Mt. Tai (Figure 1 and S1). The foot station was
105 inside Shandong College of Electric Power, a typical urban site (36.18°N, 117.11°E). HONO, VOCs, OVOCs, CO, O_3 , SO_2 ,
 NO_x , $PM_{2.5}$, PM_{10} , $J(NO_2)$, and meteorological parameters were continuously measured at this station. Details about the foot
station and the used instruments can be found in the companion study to be submitted to the same journal (entitled
“Atmospheric Measurements at the Foot and the Summit of Mt. Tai (1534 m a.s.l.) Part II: HONO Budget and Radical (RO_x
+ NO_3) Chemistry in the Lower Boundary Layer”). The summit station (36.23°N, 117.11°E) is located inside a meteorological
110 observatory at the eastern part of the summit of Mt. Tai, with an altitude of about 1534 m a.s.l. It is in the north part of Tai’an
city (altitude: ~150 m, population: ~5.6 million), and about 60 km south of Jinan city (the capital city of Shandong province,
altitude: ~20 m, population: ~8.7 million).

Since Mt. Tai is a famous tourist place, most of the tourist activities on the summit happen around the Southern Heavenly Gate,
the Bixia Temple, and the Jade Emperor Peak. The most crowded period is around sunrise when visitors come for the view of
115 sunrise. The Southern Heavenly Gate is about 1 km west of and about 100 m lower than our station. There are several small
restaurants nearby, but they don’t cause significant emissions as they only use electricity for the energy supply. The Bixia
Temple is about 200 m west to and about 50 m lower than our station, and small anthropogenic emissions may be produced
here because of the incense burning, but the impact on our measurements is expected to be negligible as a result of the fast
dilution process at the summit level. The Jade Emperor Peak is about 200 m northwest of and has a similar altitude to our
120 station. Visitors generally stay there for a short time and don’t have activities that may produce significant emissions. A detailed
discussion about the influence of anthropogenic emissions at the summit level on our measurement is presented in Section
3.2.1.



125 **Figure 1: Locations of the Mt. Tai, the summit station, and the ground station. (A): Locations of Mt. Tai and nearby cities (Tai'an and Jinan) colored by altitude; (B): Contour map of Mt. Tai; The black and the yellow stars represent the locations of the summit and the ground stations, respectively. (C): A view of the station from the southwest. Map data were taken from the National Catalogue Service for Geographic Information (<https://www.webmap.cn>). Photo copyright: Chaoyang Xue.**

2.2 Instrumentation

During the campaign, HONO was continuously measured by the LOPAP technique (LONG Path Absorption Photometer, Model-03, QUMA GmbH, Germany) with a detection limit of 1.5 pptv for 5 min average (Heland et al., 2001; Kleffmann et al., 2006). The performance of LOPAP was well assessed and recorded in different environmental conditions (Heland et al., 2001), including low-NO_x and high-altitude sites (Kleffmann and Wiesen, 2008). The LOPAP instrument was installed at the foot station from 29th May to 8th July 2017, and then transported to the summit station with successful measurements from 9th to 31st July 2017. At the summit station, NO₂ was measured by a Model-T500U-CAPS-NO₂-analyzer (Teledyne API, USA) that utilizes a patented Cavity Attenuated Phase Shift (CAPS) technique to measure NO₂ in the air directly. NO and NO_y were measured by API-T200U-NO_y-analyzer (Teledyne API, USA) based on the chemiluminescence principle coupled with a remote NO_y converter via umbilical to allow measurements with a lower detectable limit of 50 pptv. PM_{2.5} was measured by a SHARP 5030 monitor (Thermo Scientific, USA). CO and SO₂ were measured by a T300U-CO monitor (Teledyne API, USA) and a Model 43C SO₂ monitor (Thermo Scientific, USA), respectively. J(NO₂) was measured by a 4-π J(NO₂) filter radiometer (Metcon GmbH, Germany). Other J-values used in this study, including J(HONO), J(O(¹D))), and J(HNO₃), are calculated by the trigonometric SZA function (MCM default photolysis frequency calculation, see the companion paper and Jenkin et al. (1997)) and scaled by the measured J(NO₂). For instance, $J(\text{HONO}) = J(\text{HONO})_{\text{model}} * J(\text{NO}_2)_{\text{measured}} / J(\text{NO}_2)_{\text{model}}$.

Water-soluble ions, including particulate nitrate (pNO₃) of PM_{2.5}, were collected by filter method and analyzed by an ion chromatograph (Liu et al., 2020) every 2 hours late June and early July, but it suffered a sampling problem after 12th July. Aerosol size (13.6 – 763.5 nm) distribution was measured by a Scanning Mobility Particle Sizer (SMPS, Model 3938, TSI Inc., USA) equipped with a Differential Mobility Analyzer (DMA, Model 3082, TSI Inc., USA) and a Condensation Particle



Counter (CPC, Model 3775, TSI Inc., USA). Meteorological parameters (temperature, relative humidity, wind speed, wind direction) were measured by instruments from the Shandong Taishan Meteorological Station simultaneously, and details can be found in previous studies at this station (Jiang et al., 2020). In this study, 10-min averaged data were used for the following analysis. Details about the instrumentation at the foot station could be found in the companion ACP paper. Measurements at the foot station ended on 16th July. To compare pollutants between the foot and the summit levels during the same period (Section 3.2.2), measurements (only hourly CO, NO₂, PM_{2.5}, PM₁₀, O₃, and SO₂ were available) from the monitoring station (~200 m east to the foot station) were used.

3 Results and Discussion

3.1 Overview of the Observations

Figure 2 shows the meteorological parameters measured at the summit of Mt. Tai during the campaign. The air temperature (T in °C) was slightly lower (~17 °C) in the first two days compared to the period after 10th July (~20 °C). As clouds were frequently formed at the summit (Li et al., 2020), the observed relative humidity (RH) commonly reached 100%, with a mean of 96%. Based on the wind measurements, air mass at the summit mainly came from the south (direction of Tai'an city), with a mean wind speed (WS) of 5.1 m s⁻¹. In particular, during the period of 23rd to 26th July, high wind speed (1-min max: 19.5 m s⁻¹, 10-min max: 18.5 m s⁻¹) was observed, accompanied by a relatively low temperature, low pressure (p), low radiation (J(NO₂)), and high RH.




Figure 2: Meteorological parameters measured at the summit of Mt. Tai during the campaign.

Figure 3 illustrates the time series of HONO and related pollutants measured during the campaign. Several pollution events were observed. For example, the measured daytime PM_{2.5} was generally larger than 20 μg m⁻³, and high SO₂ mixing ratios (>1



ppbv) were observed during daytime on some days (i.e., from 14th to 26th July). NO mixing ratios were generally lower than 0.5 ppbv due to significant suppression by high O₃ levels of usually higher than 50 ppbv. NO₂ was generally lower than 2 ppbv with several events, during which NO₂ was relatively higher. Besides, the measured HONO mixing ratio varies from 1.1 pptv (close to the detection limit) to 880 pptv, with a mean of 133 pptv and a median of 101 pptv, respectively (Table 1). For the same sampling site at the summit of Mt. Tai, as listed in

Table 2, the observed mean HONO mixing ratios in summer is similar to those observed at the same site in winter (150 pptv, December 2017) and spring (130 pptv, March – April 2018) reported by Jiang et al. (2020), but the variation of HONO mixing ratios in summer was within a much narrower range (1 – 880 pptv) than in winter (0 – 1140 pptv) and spring (0.5 – 3230 pptv).

With an exception for relatively lower HONO levels at altitudes higher than 2000 m or in the free troposphere (Ye et al., 2018), HONO mixing ratios are significantly higher at the summit of Mt. Tai than at other mountain sites (

Table 2). For example, mean HONO mixing ratios observed at Mt. Whiteface in the USA (Zhou et al., 2007) and Mt. Hohenpeissenberg in Germany (Acker et al., 2006) were 46 and 100 (daytime)/30 (night-time) pptv, respectively. This phenomenon could be explained by fewer human activities around these mountains, while Mt. Tai locates in the middle of the NCP with a relatively high pollution level.

Note that high HONO mixing ratios were observed during the periods from 14th to 26th July, with the co-occurrence of high SO₂ (a primary pollutant generally emitted at the ground level with a relatively short lifetime). To better understand HONO formation and its role at different pollution levels, data were classified into two periods: high HONO period (HP, 14th to 26th) and low HONO period (LP) that covers all the other days. Statistics of observations during the two periods are summarized in Table 1. Average HONO, NO_y, SO₂, and PM_{2.5} during LP are 76 pptv, 4.7 ppbv, 0.3 ppbv, and 12 μg m⁻³, respectively, slightly lower than those during HP (194 pptv, 7.0 ppbv, 0.8 ppbv, and 17 μg m⁻³, respectively).




Figure 3: HONO and related species measured at the summit of Mt. Tai during the campaign.



190 **Table 1: Statistics of observations from 9th to 31st July 2018 at the summit of Mt. Tai. High HONO period (HP): 14th – 26th July; Low HONO period (LP): 9th – 13th and 27th – 31st July.**

Parameters	Whole campaign				LP/HP			
	Min	Max	Mean	Median	Min	Max	Mean	Median
HONO (pptv)	1.1*	880	133	101	1.1*/14	806/880	76/194	71/172
NO (ppbv)	0.01*	1.89	0.27	0.25	0.01*/0.07	1.9/1.1	0.27/0.27	0.24/0.26
NO ₂ (ppbv)	0.1*	8.3	1.9	1.8	0.3/0.1*	7.1/8.3	1.4/2.2	1.2/2.0
NO _y (ppbv)	0.2	21.0	6.1	6.0	0.2/0.4	13.8/21.0	4.7/7.0	4.7/6.9
O ₃ (ppbv)	20	142	74	76	22/20	102/142	66/79	66/82
CO (ppbv)	167	1101	345	343	180/167	989/1101	348/344	335/347
SO ₂ (ppbv)	0.1*	7.1	0.6	0.4	0.1*/0.1*	3.4/7.1	0.3/0.8	0.2/0.6
PM _{2.5} (µg m ⁻³)	1*	65	15	10	1*/1*	65/61	12/17	8/13
S _a (m ⁻¹)	2.3E-6	1.2E-3	3.0E-4	2.5E-4	2.3/3.5E-6	12/8.3E-4	2.1/3.6E-4	1.7/3.4E-4
S _a ×f(RH) (m ⁻¹)	7.0E-6	3.5E-3	8.3E-4	7.1E-4	7.0/11E-6	3.5/2.5E-3	6.1/9.7E-4	4.9/9.0E-4
p (kPa)	83.5	84.7	84.4	84.5	84.3/83.5	84.7/84.7	84.5/84.4	84.5/84.5
T (°C)	14.8	24.4	19.7	19.7	14.8/17.4	22.0/24.4	18.6/20.4	18.9/20.3
RH (%)	46	100	96	100	77/46	100/100	98/95	100/100
WD (°)	5	356	170	186	10/5	356/349	160/179	167/195
WS (m s ⁻¹)	0*	18.5	5.1	4.4	0*/0.3	10.6/18.5	4.2/5.7	4.1/4.8
J(NO ₂) (10 ⁻³ s ⁻¹)	-/-	7.9	1.2	2.2	-/-	7.6/7.9	1.0/1.3	0.18/0.24

*: near or below the detection limit of the used instrument

Table 2: Summary of ground-based or aircraft-based HONO measurements at background/remote sites (including mountain or pole sites) and cities near Mt. Tai.

Location	Altitude (m)	Period	Technique	Mean (pptv)	Range (pptv)	HONO/NO _x (%)	Reference
Background sites							
Zugspitze, Germany	2650	9-16 June 2001	LOPAP	12.6*	2-35	2.5*	(Kleffmann et al., 2002)
Cimone, Italy	2165	8-17 May 2004	Coil-HPLC		0-40		(Beine et al., 2005)
Hohenpeissenberg, Germany	980	3-12 Jul 2002	Denuder-IC	100 ^a /30 ^b	<10-200	6.3 ^c	(Acker et al., 2006)
Whiteface, USA	1483	14 Jun-20 Jul 1999	Coil-HPLC	46	<5-400	23	(Zhou et al., 2007)
Jungfrauoch, Switzerland	3580	2-7 Nov 2005	LOPAP	7.5	<0.5-50	4.6	(Kleffmann and Wiesen, 2008)
Barrow, USA	~3	13 Mar-14 Apr 2009	LOPAP	27	<0.4-500	6.0	(Villena et al., 2011)
Concordia station, Antarctic plateau (CSAP)	3233	22 Dec 2010-18 Jan 2011	LOPAP	28	5-59		(Kerbrat et al., 2012)
CSAP	3233	9-23 Feb 2011	LOPAP	3	0-14		(Kerbrat et al., 2012)
CSAP	3233	4 Dec 2011-13 Jan 2012	LOPAP	35 ^d /30 ^e			(Legrand et al., 2014)
Southeastern US	PBL	1 June-15 July 2013	LPAP	11.2	3-34		(Ye et al., 2018)
Southeastern US	FT	1 June-15 July 2013	LPAP	5.6	1-15		(Ye et al., 2018)
Mt. Tai	1534	1-31 Dec 2017	LOPAP	150	0-1140	3.2	(Jiang et al., 2020)
Mt. Tai	1534	5 Mar-8 Apr 2018	LOPAP	130	0.5-3230	6.0	(Jiang et al., 2020)
Mt. Tai	1534	9-31 Jul 2018	LOPAP	133	1-880	6.4	This study
Nearby cities							
Jinan	~150	26 Nov 2013-5 Jan 2016	MARGA	350	<3340		(Wang et al., 2015)
Jinan	~150	1 Sep 2015-31 Aug 2016	LOPAP	1150	17-8360	7.9 ^a /5.6 ^b	(Li et al., 2018)
Tai'an	~150	29 May-8 Jul 2018	LOPAP	620	50-2970	4.2	This study

195 *: data published in Kleffmann and Wiesen (2008) and some unpublished data from the study of Kleffmann et al. (2002).



^a: noontime, ^b: night-time, ^c: HONO/NO₂, ^d, and ^e: mean values in December and January, respectively.

PBL and FT: the planetary boundary layer and the free troposphere.

3.2 Impact of Anthropogenic Emissions on the Measured HONO


3.2.1 Impact of Emissions at the Summit Level (1534 m a.s.l.)

200 High values of NO_x/NO_y were expected in a very fresh plume with significant local emissions. Throughout the campaign, the average NO_x/NO_y ratio was 0.43 ± 0.28 , which was much lower than fresh plumes observed in the nearest city of Tai'an with an average of 0.93 ± 0.05 (from the measurement at the foot station), indicating an aged air mass and a general small impact of nearby anthropogenic emissions at the summit level.

205 However, regular local emissions caused rapid increases of some pollutants. As an example, the most rapid increase of HONO and other pollutants, which was observed between 5:20 and 6:20 on 29th July 2018 is shown in Figure 4. During this event, HONO rapidly increased from 18 to 700 pptv, in concert with rises in NO, CO, PM_{2.5}, NO_x, and NO_y but a decrease in O₃ (Table 3). The synchronous increase in NO (a primary pollutant of combustion) and the decrease in O₃ indicates a relatively fresh plume due to the fast titration reaction, as shown in R-1:



210 As there was no significant anthropogenic emission at the summit level, the polluted plume was expected to originate from the level below rather than long transport. During this period, the $\Delta\text{HONO}/\Delta\text{NO}_x$ was 8%, much larger than that inferred from direct emissions (typically inferred as less than 1%). The ratio could be enhanced by: 1) night-time NO₂-to-HONO conversion at the ground level and 2) in-plume NO₂-to-HONO conversion along the mountain slope (rock and vegetation surfaces, etc.) and 3) particle surfaces as both the boundary layer height (BLH) elevation and the valley breeze are initialized after sunrise.



215

Figure 4: HONO and related species measured on the morning of 29th July 2018.




Table 3: Concentrations of HONO and related species measured at 05:20 and 06:20 on 29th July 2018

Species	05:20	06:20	Δ
HONO/ppbv	18	700	682
CO/ppbv	214	659	445
NO/ppbv	0.1	1.9	1.8
NO ₂ /ppbv	0.4	7.1	6.7
NO _x /ppbv	0.5	9.0	8.5
NO _y /ppbv	0.4	13.3	12.9
PM _{2.5} /μg m ⁻³	2.1	12.4	10.3
O ₃ /ppbv	59	51	-8

3.2.2 Impacts from the Level Below the Summit

3.2.2.1 Insight on the Morning Peaks of the Diurnal Profiles



220

Figure 5: Diurnal variations of HONO and related parameters observed at the summit (orange) and the foot (blue) stations. Data for the summit station have a time resolution of 10 min. Hourly PM_{2.5}, NO₂, and CO were available but NO and NO_y were not available at the foot station during this period. All the data were in the same measurement period from 9th to 31st July, except for HONO, HONO/NO_x, and J(NO₂) for the foot station measured from 29th May to 8th July.

225

In Figure 5 the campaign averaged diurnal data is shown, in which most observed species, including HONO, NO, NO₂, NO_y, CO, and PM_{2.5}, showed small peaks during 6:00 – 6:30. This suggests a regular process responsible for this phenomenon rather than an accidental event. Note that the sun started to rise and to heat the ground surface, as well as the mountain surface, one hour before those peaks, leading to an increasing BLH (Anisimov et al., 2017). On the other hand, sunrise would initiate the



daytime upslope valley breeze wind (Kalthoff et al., 2000; Schmid et al., 2020; Ye et al., 1987), which could also be supported
230 by the increasing pressure and temperature (1 hour after sunrise) observed at the summit (Figure S2). Hence, it can be inferred
that the morning peaks resulted from the rising air parcel, within which pollutants accumulated during night-time. Interestingly,
similar morning peaks were also observed in winter and spring (Figure S3A), indicating the persistent impact of this process.

3.2.2.2 Insight on the Seasonal HONO Variations

In addition to the morning peaks analysis, seasonal HONO variations at the summit were also summarized (Figure 6 and S3),
235 including measurements in winter, spring, and summer. Distinctly higher $PM_{2.5}$ and NO_2 were observed in winter (Figure 6B
and 6C) than in summer. However, HONO levels in winter/spring/summer were similar (Figure 6A), indicating that the
aerosol-derived sources did not dominate HONO formation at the summit level. In general, HONO levels observed at the
ground level of the NCP were significantly higher in winter than in summer (Li et al., 2018; Nie et al., 2015; Xue et al., 2020).
A similar HONO level observed in summer was possible because of a more rapid vertical exchange between the ground level
240 and the summit level (see Section 3.2.3).




Figure 6: Statistic summaries of (A): HONO, (B): $PM_{2.5}$, and (C): NO_2 in the three seasons. Error bars represent the standard deviation. The top and the bottom of each diamond represent the 25% and 75% percentages, respectively. The star and the line inside each diamond denote the average and the median, respectively. Data for winter and spring was taken from Jiang et al. (2020).


3.2.2.3 Insight on the Comparison of Pollutants at the Foot and Summit Level

Comparison of daytime (5:00 – 18:00) average $PM_{2.5}$, CO, O_3 , and SO_2 observed at the foot and the summit stations are shown
in Figure 7. It is apparent that all the average daytime levels of primary pollutants (CO and SO_2), partially primary pollutant
($PM_{2.5}$), and secondary pollutant (O_3) show very similar variation trends at both monitoring station, revealing 1) a significant
or even dominant impact of pollutants at the foot level on that at the summit level, and 2) the presence of a pathway that
250 enables the vertical air mass exchange between the summit and the foot levels. This was also consistent with the higher daytime
HONO (Figure S3A) observed at the summit station in winter than in summer because the regional pollution was generally
much severer in winter than in summer.

Besides, during night-time, the summit (~1500 m altitude) is above the boundary layer (in the residual layer), and similar
variation trends of pollutants were also found at the foot and the summit stations (Figure S4), indicating still the presence of



255 vertical air mass exchange at night. This could also be inferred from the higher night-time HONO (Figure S3A) in summer than in winter because 1) more south winds (the direction to Tai'an city) were observed in summer (Figure S5) and 2) the nocturnal boundary layer height was generally much lower in winter than that in summer.



260 **Figure 7: Comparison of daytime (5:00 – 18:00) average (A): SO₂, (B): O₃, (C): CO, and (D): PM_{2.5} observed at the foot (left axis in blue) and summit (right axis in orange) stations during the same period from 9th to 31st July.**

3.2.3 Impact from Tai'an City (150 m a.s.l.)

Besides the discussion in Section 3.2.1, five arguments point to the potential impact from pollution in the nearest city (Tai'an city, ~150 m a.s.l.) on the summit HONO level:

- 265 a) the “∩” shape of HONO variation in the daytime was different from that of NO₂ (a constant level during the daytime), NO_y (which increased in the early morning and then remained stable at noontime, followed by a continuous increase in the late afternoon) and PM_{2.5} (which also showed a “∩” shape variation but its peak was 3 hours later than the HONO peak). These observations indicate that the observed HONO at the summit was not dominated by in-situ aerosol-derived formation (Figure 5) but an external HONO source such as transport;
- 270 b) high-level HONO was frequently observed at the ground level (150 m a.s.l.) in Tai'an city (Table 2), and almost the same variation trends of HONO/NO_x were observed at both the summit and foot stations (Figure 5G);
- c) HONO peaks at the summit occurred at noontime when the BLH was high, and valley breeze wind was strong;



- d) high-level HONO (>200 pptv) observed at the summit mainly appeared when the air mass came from south or southwest (the direction to Tai'an city, see Figure 8);
- e) HONO peaks occurred synchronously with the peaks of SO₂, which is mainly emitted at the ground level, and NO₂, which is an important HONO precursor (Figure 3).

The impact could be achieved through: 1) the air mass ascending by valley breeze upslope wind (daytime) and by the north wind (daytime and night-time, Section 3.2.2.3), and 2) HONO formation during the air mass ascending process, i.e., HONO formation through the NO₂ heterogeneous uptake on the mountain slope surfaces (George et al., 2005; Marion et al., 2021; Stemmler et al., 2006). The HONO production from the above processes was defined as P(HONO)_{transport} and will be discussed in Section 3.6.




Figure 8: Pollution rose plot of HONO against the wind direction. The frequency contribution of counts by wind direction is also shown on the left axis.

The length of the hypotenuse from the foot to the summit is about 4.2 km, with an average elevation angle of about 20°. In the daytime, the valley breeze could occur with a upslope wind speed of 2 – 5 m s⁻¹ reported in previous measurements (Kalthoff et al., 2000; Schmid et al., 2020; Ye et al., 1987; also see https://glossary.ametsoc.org/wiki/Upvalley_wind), it takes about 14 – 35 min ($t_{\text{transport}}$) for the air mass to be transported from the foot to the summit. The south wind could enhance the upslope valley breeze wind; for example, the mean south winds measured at the ground and summit stations are >2 and >5 m s⁻¹, respectively. If the integrated wind speed along the mountain slope is 4 – 10 m s⁻¹, the calculated $t_{\text{transport}}$ will be reduced to 7 – 17.5 min.

The key question is the quantity of HONO that still exists after transport from the foot to the summit levels regarding its photolysis in the daytime. Assuming first-order decay of HONO by photolysis during the transport, the remaining HONO and ratio at the summit can be calculated:

$$c_t = c_0 e^{-J(\text{HONO}) \cdot t_{\text{transport}}}, \quad \text{Eq-1}$$

$$\alpha = \frac{c_t}{c_0} = e^{-J(\text{HONO}) \cdot t_{\text{transport}}}, \quad \text{Eq-2}$$



where c_t , c_0 , $J(\text{HONO})$, and α represent the remaining HONO after a transport period ($t_{\text{transport}}$), the initial HONO concentration at the foot, the HONO photolysis frequency, and the remaining proportion of HONO.

Figure 9 shows the calculated α with $t_{\text{transport}} = 7$ or 17.5 min during the daytime. It is apparent that α was larger than 40% with $t_{\text{transport}} = 17.5$ min and larger than 70% with $t_{\text{transport}} = 7$ min, providing a theoretical basis for the potential role of vertical HONO transport from the ground to the summit levels. This calculation only included the daytime HONO sink through photolysis, but the sources, such as $\text{NO} + \text{OH}$ and heterogeneous NO_2 reactions, were not considered, and hence, the calculated α represents a lower limit. Thus, the impact of transport was expected to be larger when 1) taking other HONO formation paths (e.g., NO_2 heterogeneous reactions on the mountain surfaces and the vegetation surfaces) into account, and 2) vegetation shadows on the mountain surface slow down HONO photolysis during the transport. Therefore, ground level (~150 m a.s.l.) HONO as well as its formation during transport may affect the HONO measurement at the summit significantly. The quantification of the contribution will be discussed in Section 3.6.






Figure 9: Diurnal profiles of the remaining proportion of HONO (α) after a period of transport ($t_{\text{transport}}$) from the ground to the summit levels and $J(\text{HONO})$ during the daytime.

3.3 Daytime Unknown HONO Source Strength

The photo-stationary state (PSS), presented by the following equations, is valid to calculate the unknown HONO source strength (P_{un}) when local emission was negligible (Crilley et al., 2016; Kleffmann et al., 2005; Michoud et al., 2012). The predicted HONO concentration by PSS ($[\text{HONO}]_{\text{PSS}}$) and P_{un} could be calculated by Eq-3 and Eq-4, respectively.



$$[\text{HONO}]_{\text{PSS}} = \frac{k_1 \times [\text{NO}] \times [\text{OH}]}{J(\text{HONO}) + k_2 \times [\text{OH}]}, \quad \text{Eq-3}$$

$$P_{\text{un}} = ([\text{HONO}] - [\text{HONO}]_{\text{PSS}}) \times (J(\text{HONO}) + k_2 \times [\text{OH}]), \quad \text{Eq-4}$$



OH measurements were not available during this campaign but were available for the summit station in June 2006 (Kanaya et al., 2009). The significant correlation of OH and $J(\text{O}^1\text{D})$ observed during the former campaign was used here to estimate OH concentrations. Since $\text{NO} + \text{OH}$ was not the dominant HONO source (8.0%) or $\text{HONO} + \text{OH}$ not the dominant sink (1.9%), the uncertainty caused by the estimated OH should be not significant.

The diurnal variation of the calculated noontime (10:00 – 16:00) P_{un} is shown in Figure 10. Campaign-averaged P_{un} was about 290 ± 280 pptv h^{-1} with a maximum of about 1800 pptv h^{-1} . The maximum P_{un} value appeared at midday (13:00), indicating a photo-enhanced HONO source. Similarly, high correlations ($r = 0.79, 0.83, \text{ or } 0.83$) were found between P_{un} and $\text{NO}_2^*J(\text{NO}_2)$, $\text{pNO}_3^*J(\text{HNO}_3)$, or $\text{NO}_y^*J(\text{HNO}_3)$ (Table 4), suggesting the potential HONO formation from photosensitized NO_2 reactions or photolysis of NO_z ($\text{NO}_z = \text{NO}_y - \text{NO} - \text{NO}_2$) species such as particulate nitrate (pNO_3). Moreover, the relatively poor correlations ($r = 0.17$ or 0.64) between P_{un} and $\text{NO}_2^*S_a$ or $\text{NO}_2^*S_a^*J(\text{NO}_2)$ (Table 4) suggested a minor role of NO_2 uptake on the aerosol surface in the HONO formation. Besides, a high correlation between P_{un} and HONO ($r = 0.76$) was obtained. A possible reason could be that HONO and other pollutants were not dominated by in situ formation but by transport, as discussed in Section 3.2.2. As correlation analysis is only a preliminary indicator and it might not be instructive for HONO budget analysis when the vertical air mass exchange occurs, further investigation of NO_2 uptake on aerosol surface and photolysis of pNO_3 is presented in Section 3.4 and Section 3.5, respectively.

Table 4: The correlation coefficients (r) between HONO or P_{un} and other parameters (first column).

Correlations	HONO	P_{un}
CO	0.20	-0.10
NO	0.06	0.09
NO_2	0.53	0.05
NO_x	0.52	0.06
NO_y	0.49	0.45
NO_z	0.38	0.49
$\text{PM}_{2.5}$	0.07	0.21
HONO	--	0.76
$\text{HONO}^*J(\text{HONO})$	--	0.988
$J(\text{NO}_2)$	-0.04	0.77
$\text{NO}_2^*\text{PM}_{2.5}$	0.28	0.25
$\text{NO}_2^*J(\text{NO}_2)$	0.03	0.79
$\text{pNO}_3^*J(\text{HNO}_3)$	0.05	0.83
$\text{NO}_y^*J(\text{HNO}_3)$	0.06	0.83
$\text{NO}_2^*S_a$	0.47	0.17
$\text{NO}_2^*S_a^*J(\text{NO}_2)$	0.10	0.64



335 3.4 Constraint on HONO Formation from NO₂ Uptake on the Aerosol Surface

During the daytime, the HONO production rate from the NO₂ uptake on the aerosol surface (P(HONO)_a) with the photo-enhanced effect was parameterized by the following equation.

$$P(\text{HONO})_a = \frac{v(\text{NO}_2) \times S_a \times [\text{NO}_2]}{4} \times [\gamma_a \times \frac{J(\text{NO}_2)_{\text{measured}}}{0.005 \text{ s}^{-1}}], \quad \text{Eq-5}$$


where $v(\text{NO}_2)$, S_a , γ_a , $J(\text{NO}_2)_{\text{measured}}/0.005$ are the molecular speed of NO₂ (m s⁻¹), aerosol surface density (m⁻¹), the photo-enhanced uptake coefficient of NO₂ on the aerosol surfaces, and the photo-enhancement factor normalized to a $J(\text{NO}_2) = 0.005$ s⁻¹. In Eq-5, an upper limit HONO yield for the NO₂ conversion of 100% was assumed. Additionally, RH was proposed to significantly influence aerosol surface density, especially at our site, with frequently high RH up to 100%. Then besides calculating the aerosol surface density based on the measured aerosol size distribution ($S_{a_measured}$), we estimated the effective aerosol surface density in m⁻¹ with an RH enhancement factor $f(\text{RH})$ ($S_a = S_{a_measured} * f(\text{RH})$) using the following equation:

$$345 \quad f(\text{RH}) = 1 + a \times (\text{RH}/100)^b, \quad \text{Eq-6}$$

where a and b are empirical values of 2.06 and 3.60, respectively (Liu et al., 2008). The average S_a without and with RH enhancement is 3.0×10^{-4} and 8.3×10^{-4} m⁻¹, respectively. Note that the uncertainty of S_a is not expected to cause a significant uncertainty on HONO budget analysis as P(HONO)_a was not the dominant source (Section 3.6).

As P_{un} includes all the sources except NO + OH, then P(HONO)_a << P_{un} can always be obtained. Hence, the real γ_a value should be much lower than the inferred ones ($\gamma_{a_inferred}$) from $P_{\text{un}} = P(\text{HONO})_a$. In total, 606 γ_a values were inferred based on the measurements, varying from 1.3×10^{-4} to 8.5×10^{-3} , with a mean of $(8.3 \pm 7.5) \times 10^{-4}$. However, the minimum ($\gamma_{a_inferred_mini}$) of 1.3×10^{-4} is still very high, compared to the results of most lab studies, in which values of γ_a of typically at few times 10⁻⁵ or even less were observed (Han et al., 2016; Liu et al., 2019a; Ndour et al., 2008; Sosedova et al., 2011; Stemmler et al., 2007). Hence a popularly used value of $\gamma_a = 2 \times 10^{-5}$ was used to calculate P(HONO)_a and $\gamma_{a_inferred_mini}$ of 1.3×10^{-4} was also used for uncertainty analysis as the upper limit.

The calculated P(HONO)_a with these γ_a values are shown in Figure 10. It is obvious that P(HONO)_a is significantly lower than P_{un} with either lab-based $\gamma_a = 2 \times 10^{-5}$ or even $\gamma_{a_inferred_mini} = 1.3 \times 10^{-4}$, pointing out the minor role of NO₂ uptake on the aerosol surface in daytime HONO formation. With the lab-based $\gamma_a = 2 \times 10^{-5}$, P(HONO)_a could only explain 3% of P_{un} , which is similar to previous model studies (Liu et al., 2019b; Xue et al., 2020; Zhang et al., 2016). The contribution of P(HONO)_a to P_{un} increased when using $\gamma_{a_inferred_mini}$, but resulting from an overestimated γ_a as discussed before. Nevertheless, analysis in this study still could be an important effort in the field constraints on the NO₂-to-HONO conversion on the aerosol surface.



365 **Figure 10: Unknown HONO source strength (P_{un}), and HONO production rates from $NO + OH$ ($P(HONO)_{NO+OH}$), NO_2 uptake on the aerosol surface ($P(HONO)_a$) with the inferred $\gamma_{a_inferred_mini} = 1.3 \times 10^{-4}$ and the popular used $\gamma_a = 2 \times 10^{-5}$, and nitrate photolysis ($P(HONO)_{nitrate}$) with EF values of 1, 7 and 15.6.**

3.5 Constraint on HONO Formation from the Photolysis of Particulate Nitrate

As one of the important inorganic components of aerosols, particulate nitrate (pNO_3) could undergo photolysis, with the production of HONO. This process needs more field constraints as discussed in the Introduction section. During the present campaign, pNO_3 at the summit was measured by a filter method every 2 hours (Liu et al., 2020), but it suffered a sampling
 370 problem after 12th July. Because NO_z ($NO_z = NO_y - NO - NO_2$) mainly contains pNO_3 and its precursors, e.g., HNO_3 and N_2O_5 , similar variations were expected between NO_z and pNO_3 . As shown in Figure S6, NO_z and pNO_3 exhibited a very high correlation ($R^2 = 0.895$), for which pNO_3 makes 44% of the NO_z and this fraction was used to estimate pNO_3 in the period when it was not measured. The uncertainty of the estimated pNO_3 should have no significant impact on daytime HONO formation concerning its small contribution to daytime HONO formation (see Section 3.6).

375 A high correlation between P_{un} and $pNO_3 * J(HNO_3)$ was found (Table 4), suggesting a possible impact of pNO_3 on HONO formation. But one should bear in mind that the high correlation might also be caused by the remarkable impact on pNO_3 formation from HONO-related reactions (e.g., $HONO \xrightarrow{hv} OH \xrightarrow{NO_2} HNO_3 \rightarrow pNO_3$) (Xue et al., 2020) or other photolytic processes. For parameterization, an enhancement factor (EF) was defined as the ratio of photolysis frequencies of pNO_3 to gas-phase HNO_3 . Then HONO production from pNO_3 photolysis ($P(HONO)_{nitrate}$) could be quantified by Eq-7:

380
$$P(HONO)_{nitrate} = pNO_3 * J(HNO_3) * EF, \quad \text{Eq-7}$$

Similar to NO_2 uptake on the aerosol surface, one can always find $P(HONO)_{nitrate} \ll P_{un}$. Hence, it is expected that the real EF should be much lower than the inferred ones ($EF_{inferred}$) from $P(HONO)_{nitrate} = P_{un}$. Therefore, 606 EF values were inferred, in the range of 15.6 to 1072, with a mean of 173 ± 98 , which is much higher than those (around 1) determined in recent flow tube



or smog chamber studies (Shi et al., 2021; Wang et al., 2021). The minimum ($EF_{\text{inferred_mini}} = 15.6$) is at a similar level to field
385 studies of Romer et al. (2018) and Zhou et al. (2003), and the lower values in the laboratory studies (Bao et al., 2018; Ye et
al., 2016, 2017; Zhou et al., 2011). To quantify the HONO production from $p\text{NO}_3$ photolysis, the EF value of 7 from a recent
field study (Romer et al. 2018) was used for $P(\text{HONO})_{\text{nitrate}}$ calculation, and $EF_{\text{inferred_mini}}$ (15.6) from this study and EF values
of ~ 1 from recent laboratory studies (Shi et al., 2021; Wang et al., 2021) were also used for the uncertainty analysis and
comparison.

390 The calculated $P(\text{HONO})_{\text{nitrate}}$ with these EF values is shown in Figure 10. With the $EF = 7$, $P(\text{HONO})_{\text{nitrate}}$ was at a level of
half of $\text{NO} + \text{OH}$ but much lower than P_{un} , which was also observed at the summit of Mt. Whiteface (Zhou et al., 2007).
 $P(\text{HONO})_{\text{nitrate}}$ could explain 4.3% of the observed P_{un} . While its contribution varied from 0.6% to 9.6%, depending on EF
values varying from 1 to 15.6 in the sensitivity tests. Therefore, with a $P(\text{HONO})_{\text{a}}$ ($\gamma_{\text{a}} = 2 \times 10^{-5}$) contribution of 3% (Section
3.4) and a $P(\text{HONO})_{\text{nitrate}}$ ($EF = 7$) contribution of 4.3%, the other sources (defined as $P(\text{HONO})_{\text{other}} = P_{\text{un}} - P(\text{HONO})_{\text{a}} -$
395 $P(\text{HONO})_{\text{nitrate}}$, mainly transported from the ground level as discussed in Section 3.2.2), made a dominated contribution of
92.7% to the observed P_{un} .

Moreover, significant differences between EF values obtained from field studies and laboratory studies indicate a complex
process of $p\text{NO}_3$ photolysis that may be influenced by various environmental parameters, e.g., the aerosol $p\text{NO}_3$ loading and
the aerosol composition (Bao et al., 2018, 2020; Ye et al., 2016, 2017), and experimental laboratory conditions, e.g., collected
400 particles on the filter or generated airborne particles (Shi et al., 2021; Wang et al., 2021). We, therefore, suggest that this
process still needs further field or laboratory constraints. To our knowledge, this study provided the first field constraint on the
aerosol-derived HONO sources in the NCP region, where the abundance of aerosol was frequently observed, and its role in
HONO formation is still highly controversial.

The landscape (e.g., mountains) enhances the vertical air mass exchange, leading to a weak vertical HONO distribution within
405 the boundary layer, which is not yet considered in previous studies. This will underestimate the role of ground-derived sources
in HONO formation in the upper boundary layer over mountain regions. Regarding the short enough lifetime of OH ($\ll 1$ s),
it is expected that OH cannot be transported. However, part of HONO was formed at the ground level through $\text{NO} + \text{OH}$, and
the same amount of OH consumed at the ground level would be released at the summit level through HONO photolysis,
constituting a potential pathway of OH transport through the reservoir of HONO. Moreover, the enhanced vertical air mass
410 exchange could also lead to fast transport of other pollutants ($\text{PM}_{2.5}$, O_3 , CO, SO_2 , etc.) from the ground to the summit levels,
which is expected to significantly impact the atmospheric composition as well as its chemistry in the upper boundary layer or
the residual layer. The discussion and implications in this study will be instructive for further laboratory or model studies.

3.6 Role of HONO in the Oxidizing Capacity of the Upper Boundary Layer

O_3 was typically the major HO_x ($\text{HO}_x = \text{OH} + \text{HO}_2$) source at high altitude regions, including the upper boundary layer. Then
415 we compared the HO_x production rates from O_3 and HONO photolysis to investigate whether HONO could play a significant
role in the oxidizing capacity of the atmosphere at this high-altitude site. Photolysis of HONO and O_3 with their integrated



HO_x production is shown in R-2 and R-5 to R-7, respectively. OH loss through $\text{HONO} + \text{OH}$ and $\text{NO} + \text{OH}$ was not subtracted from $P(\text{HO}_x)_{\text{HONO}}$ because of the dominated role of vertical transport in the HONO formation at the summit level (net HO_x production at the summit level and comparison with the foot level is calculated and discussed in the companion ACP paper).



$P(\text{HO}_x)_{\text{HONO}} = [\text{HONO}] * J(\text{HONO}), \quad \text{Eq-8}$

$P(\text{HO}_x)_{\text{O}_3} = [\text{O}_3] * J(\text{O}(^1\text{D})) * \phi, \quad \text{Eq-9}$

425 where the reaction constants were taken from the IUPAC kinetic database (<https://iupac-aeris.ipsl.fr>). The atmospheric RH and temperature largely influenced the branching ratio of R-6 to R-7. The average OH yield (ϕ) during the campaign of 20% was used for calculating OH production from O_3 photolysis.

Figure 11 displays the diurnal profiles of HO_x production rates from HONO and O_3 during the clean and polluted periods. It is apparent that O_3 photolysis contributed mostly to HO_x production, owing to that $P(\text{HO}_x)_{\text{O}_3}$ was generally more than two
430 times of $P(\text{HO}_x)_{\text{HONO}}$ during daytime except in the early morning and later afternoon. Variations of $P(\text{HO}_x)_{\text{HONO}}$ in the clean period started to increase after sunrise, followed by a decline after the peak at 11:00. The rapid decline of $P(\text{HO}_x)_{\text{HONO}}$ and $P(\text{HO}_x)_{\text{O}_3}$ around noon was caused by low solar intensity as a result of frequently (near-) saturated humidity and resulting cloud formation (Figure 2 and S1F). The largest $P(\text{HO}_x)_{\text{HONO}}$ of the diurnal profile in the clean period was 1.9×10^6 molecules $\text{cm}^{-3} \text{s}^{-1}$, and contributed 21% of total HO_x production through the photolysis of HONO and O_3 ($P(\text{HO}_x)_{\text{sum}}$), which was close to the
435 average contribution (24%) in the clean period. During the polluted period, however, the curve of diurnal $P(\text{HO}_x)_{\text{HONO}}$ was broader than that in the clean period, with an unexpectedly sustainable high level ($>2.0 \times 10^6$ molecules $\text{cm}^{-3} \text{s}^{-1}$) from 10:20 to 16:00, increasing the contribution of $P(\text{HO}_x)_{\text{HONO}}$ to $P(\text{HO}_x)_{\text{sum}}$ to 27.5%. At 15:00, the maximum average $P(\text{HO}_x)_{\text{HONO}}$ of 3.3×10^6 molecules $\text{cm}^{-3} \text{s}^{-1}$ was observed in the polluted period, contributing about 34% of $P(\text{HO}_x)_{\text{sum}}$.

In particular, in the early morning at the ground level, HONO photolysis is found to initialize daytime photochemistry after
440 night-time accumulation (Alicke et al., 2002; Kleffmann, 2007; Platt et al., 1980). If the air mass exchange between the ground level and the summit level is fast enough, HONO photolysis may also initialize daytime photochemistry at the summit level. Then we compared the relative contribution of HO_x production rates from HONO and O_3 at different hours. As shown in Figure 12, $P(\text{HO}_x)_{\text{HONO}}$ arose earlier than $P(\text{HO}_x)_{\text{O}_3}$ and dominated HO_x production ($>50\%$) in the early morning (5:00 – 7:20), demonstrating that daytime atmospheric photochemistry at the summit level is also initialized by HONO photolysis.

445 Taken together, the average contribution of $P(\text{HO}_x)_{\text{HONO}}$ to $P(\text{HO}_x)_{\text{sum}}$ was about 26%, more than one-third of $P(\text{HO}_x)_{\text{O}_3}$ (the contribution is 18% if OH loss through $\text{HONO} + \text{OH}$ and $\text{NO} + \text{OH}$ is subtracted, see the companion paper). As discussed before, the transport from the ground level to the summit level contributed to the majority of HONO observed at the summit level, pointing to a new insight that ground-derived HONO played an important role in the oxidizing capacity, not only at the



ground level but also in the upper boundary layer (~1500 m) in mountainous regions. Yet this vertical exchange might be only
450 valid in the mountainous areas, and the follow-up regional impact still needs to be quantified by further model studies.





Figure 11: Diurnal profiles of HO_x production rates from HONO and O₃ during (A): the clean and (B) the polluted periods. The two pie charts in each parcel represent the relative contribution of HO_x production rates from HONO and O₃ throughout the whole day (left) and when the maximum P(HO_x)_{HONO} occurred (right), respectively.



455

Figure 12: The relative contribution of integrated HO_x production rates from HONO and O₃ during the daytime.

4 Summary

Observations of HONO and related parameters at the summit of Mt. Tai (1534 m a.s.l.) in July 2018 were presented. The average HONO mixing ratio is 133 ± 106 pptv, with a maximum of 880 pptv, significantly higher than observations at other
460 mountain summits worldwide. Along with observations at the ground level (the nearest city, Tai'an city), HONO formation from different paths and its role in the atmospheric oxidizing capacity of the upper boundary layer were explored and discussed. The main conclusions are listed as follows:

1. Constraints on the kinetics of NO₂ uptake coefficient on the aerosol surface and photolysis of pNO₃ were obtained based on the assumption that P_{un} could be solely explained by NO₂ uptake on the aerosol surface, P(HONO)_a or
465 particulate nitrate photolysis, P(HONO)_{nitrate}. The inferred λ_a and EF values were much higher than most values



obtained from recent laboratory studies, indicating that the aerosol-derived HONO could not explain the observed P_{un} . In the NCP region, the abundance of aerosol was frequently observed, but its role in HONO formation is still highly controversial as a result of uncertain kinetics. This study provided the first field constraints on aerosol-derived HONO sources in this region and will be instructive for further laboratory or model studies.

- 470 2. With a λ_a value of 2×10^{-5} and an EF value of 7, $P(\text{HONO})_a$ and $P(\text{HONO})_{\text{nitrate}}$ showed small contributions (3% and 4.3%, respectively) to daytime HONO formation at the summit station. Both $P(\text{HONO})_a$ and $P(\text{HONO})_{\text{nitrate}}$ varied from negligible to moderate levels (similar to $\text{NO} + \text{OH}$), depending on γ_a and EF values, suggesting the necessity to further study the related kinetics. Additionally, although high values of γ_a (1.3×10^{-4}) and EF (15.6) compared with recent studies were tested here, both sources were still much lower than the observed P_{un} . The remaining majority
- 475 (92.7%) of P_{un} was dominated by the rapid vertical transport from the ground to the summit levels including heterogeneous HONO formation on surfaces of the mountain slope, which was inferred from comprehensive evidence presented in this study.
3. Photolysis of HONO initialized daytime photochemistry in the early morning. In addition, for the daytime average, it contributed 26% of $P(\text{HO}_x)_{\text{sum}}$, more than one-third of $P(\text{HO}_x)_{\text{O}_3}$, indicating the important role of HONO in the oxidizing capacity of the atmosphere in mountainous areas. HONO formation at the ground level could significantly influence the HONO mixing ratios and the atmospheric oxidizing capacity at the summit level through the vertical air mass exchange. Moreover, the enhanced vertical air mass exchange could also lead to a fast exchange of other pollutants between the ground and the summit levels and significantly impact the atmospheric composition as well as the chemistry in the upper boundary layer or the residential layer. However, those follow-up impacts, by far, are not
- 480 quantified by the current model studies.
- 485

Acknowledgment

We are grateful to Xiaowei He, Pengfei Liu, Chao Zhu, Jiarong Li, Hui Chen, Xianmang Xu, Hongyong Li, Pengcheng Zhang, and Jinhe Wang for their help on the measurement at the summit of Mt. Tai. We thank all researchers involved in this campaign from the Research Centre for Eco-Environmental Sciences-Chinese Academy of Sciences, Fudan University, Shandong

490 Jianzhu University, Shandong University, and the Municipal Environmental Protection Bureau of Tai'an. We thank Yunqiao Zhou for his help with the map plotting.

Funding: This work was supported by the National Natural Science Foundation of China (Nos. 91544211, 41727805, 41931287, and 41975164), and the PIVOTS project provided by the Region Centre – Val de Loire (ARD 2020 program and CPER 2015 – 2020).

495 **Author Contribution:** C.X., C.Y., C.Z., Y.Z., H.L., and Z.G. performed the field measurements. C.X. analyzed the observation data and wrote the paper with inputs from all co-authors. C.Y. and J.K. helped with the data analysis and manuscript writing. J.K., C.Y., V.C., A.M., L.X., J.C., K.L., F.B., and Y.M. revised the manuscript.



Competing Interests: The authors declare no competing financial interest.

Data Availability: All the summertime data used in this study is available upon request from the corresponding authors.

500 References

- Acker, K., Möller, D., Wieprecht, W., Meixner, F. X., Bohn, B., Gilge, S., Plass-Dülmer, C. and Berresheim, H.: Strong daytime production of OH from HNO₂ at a rural mountain site, *Geophys. Res. Lett.*, 33(2), L02809, doi:10.1029/2005GL024643, 2006.
- Alicke, B., Platt, U. and Stutz, J.: Impact of nitrous acid photolysis on the total hydroxyl radical budget during the Limitation of Oxidant Production/Pianura Padana Produzione di Ozono study in Milan, *J. Geophys. Res. Atmos.*, 107(D22), 8196, doi:10.1029/2000JD000075, 2002.
- 505 Ammann, M., Kalberer, M., Jost, D. T., Tobler, L., Rössler, E., Piguet, D., Gäggeler, H. W. and Baltensperger, U.: Heterogeneous production of nitrous acid on soot in polluted air masses, *Nature*, 395(6698), 157–160, doi:10.1038/25965, 1998.
- 510 Anisimov, S. V., Galichenko, S. V. and Mareev, E. A.: Electrodynamic properties and height of atmospheric convective boundary layer, *Atmos. Res.*, 194, 119–129, doi:10.1016/j.atmosres.2017.04.012, 2017.
- Bao, F., Li, M., Zhang, Y., Chen, C. and Zhao, J.: Photochemical Aging of Beijing Urban PM_{2.5}: HONO Production, *Environ. Sci. Technol.*, 52(11), 6309–6316, doi:10.1021/acs.est.8b00538, 2018.
- Bao, F., Jiang, H., Zhang, Y., Li, M., Ye, C., Wang, W., Ge, M., Chen, C. and Zhao, J.: The Key Role of Sulfate in the
515 Photochemical Renoxification on Real PM_{2.5}, *Environ. Sci. Technol.*, 54(6), 3121–3128, doi:10.1021/acs.est.9b06764, 2020.
- Beine, H. J., Amoroso, A., Esposito, G., Sparapani, R., Ianniello, A., Georgiadis, T., Nardino, M., Bonasoni, P., Cristofanelli, P. and Dominé, F.: Deposition of atmospheric nitrous acid on alkaline snow surfaces, *Geophys. Res. Lett.*, 32(10), L10808, doi:10.1029/2005GL022589, 2005.
- Bejan, I., Abd El Aal, Y., Barnes, I., Benter, T., Bohn, B., Wiesen, P. and Kleffmann, J.: The photolysis of ortho-nitrophenols:
520 A new gas phase source of HONO, *Phys. Chem. Chem. Phys.*, 8(17), 2028–2035, doi:10.1039/b516590c, 2006.
- Crilley, L. R., Kramer, L., Pope, F. D., Whalley, L. K., Cryer, D. R., Heard, D. E., Lee, J. D., Reed, C. and Bloss, W. J.: On the interpretation of: In situ HONO observations via photochemical steady state, *Faraday Discuss.*, 189, 191–212, doi:10.1039/c5fd00224a, 2016.
- Donaldson, M. A., Bish, D. L. and Raff, J. D.: Soil surface acidity plays a determining role in the atmospheric-terrestrial exchange of nitrous acid, *Proc. Natl. Acad. Sci. U.S.A.*, 111(52), 18472–18477, doi:10.1073/pnas.1418545112, 2014.
- Fu, X., Wang, T., Zhang, L., Li, Q., Wang, Z., Xia, M., Yun, H., Wang, W., Yu, C., Yue, D., Zhou, Y., Zheng, J. and Han, R.: The significant contribution of HONO to secondary pollutants during a severe winter pollution event in southern China, *Atmos. Chem. Phys.*, 19(1), 1–14, doi:10.5194/acp-19-1-2019, 2019.
- Ge, S., Wang, G., Zhang, S., Li, D., Xie, Y., Wu, C., Yuan, Q., Chen, J. and Zhang, H.: Abundant NH₃ in China Enhances
530 Atmospheric HONO Production by Promoting the Heterogeneous Reaction of SO₂ with NO₂, *Environ. Sci. Technol.*, 53(24), 14339–14347, doi:10.1021/acs.est.9b04196, 2019.
- George, C., Strekowski, R. S., Kleffmann, J., Stemmler, K. and Ammann, M.: Photoenhanced uptake of gaseous NO₂ on solid organic compounds: A photochemical source of HONO?, *Faraday Discuss.*, 130, 195–210, doi:10.1039/b417888m, 2005.
- Han, C., Yang, W., Wu, Q., Yang, H. and Xue, X.: Heterogeneous Photochemical Conversion of NO₂ to HONO on the Humic
535 Acid Surface under Simulated Sunlight, *Environ. Sci. Technol.*, 50(10), 5017–5023, doi:10.1021/acs.est.5b05101, 2016.
- Han, C., Yang, W., Yang, H. and Xue, X.: Enhanced photochemical conversion of NO₂ to HONO on humic acids in the



- presence of benzophenone, *Environ. Pollut.*, 231(1), 979–986, doi:10.1016/j.envpol.2017.08.107, 2017.
- Heland, J., Kleffmann, J., Kurtenbach, R. and Wiesen, P.: A New Instrument To Measure Gaseous Nitrous Acid (HONO) in the Atmosphere, *Environ. Sci. Technol.*, 35(15), 3207–3212, doi:10.1021/es000303t, 2001.
- 540 Jenkin, M. E., Saunders, S. M. and Pilling, M. J.: The tropospheric degradation of volatile organic compounds: a protocol for mechanism development, *Atmos. Environ.*, 31(1), 81–104, doi:10.1016/S1352-2310(96)00105-7, 1997.
- Jiang, Y., Xue, L., Gu, R., Jia, M., Zhang, Y., Wen, L., Zheng, P., Chen, T., Li, H., Shan, Y., Zhao, Y., Guo, Z., Bi, Y., Liu, H., Ding, A., Zhang, Q. and Wang, W.: Sources of nitrous acid (HONO) in the upper boundary layer and lower free troposphere of the North China Plain: insights from the Mount Tai Observatory, *Atmos. Chem. Phys.*, 20(20), 12115–12131, doi:10.5194/acp-20-12115-2020, 2020.
- 545 Kalthoff, N., Horlacher, V., Corsmeier, U., Volz-Thomas, A., Kolahgar, B., Geiß, H., Möllmann-Coers, M. and Knaps, A.: Influence of valley winds on transport and dispersion of airborne pollutants in the Freiburg-Schauinsland area, *J. Geophys. Res. Atmos.*, 105(D1), 1585–1597, doi:10.1029/1999JD900999, 2000.
- Kanaya, Y., Pochanart, P., Liu, Y., Li, J., Tanimoto, H., Kato, S., Suthawaree, J., Inomata, S., Taketani, F., Okuzawa, K., 550 Kawamura, K., Akimoto, H. and Wang, Z. F.: Rates and regimes of photochemical ozone production over Central East China in June 2006: A box model analysis using comprehensive measurements of ozone precursors, *Atmos. Chem. Phys.*, 9(20), 7711–7723, doi:10.5194/acp-9-7711-2009, 2009.
- Kerbrat, M., Legrand, M., Preunkert, S., Gallée, H. and Kleffmann, J.: Nitrous acid at Concordia (inland site) and Dumont d’Urville (coastal site), East Antarctica, *J. Geophys. Res. Atmos.*, 117(8), D08303, doi:10.1029/2011JD017149, 2012.
- 555 Kleffmann, J.: Daytime Sources of Nitrous Acid (HONO) in the Atmospheric Boundary Layer, *ChemPhysChem*, 8(8), 1137–1144, doi:10.1002/cphc.200700016, 2007.
- Kleffmann, J. and Wiesen, P.: Technical Note: Quantification of interferences of wet chemical HONO LOPAP measurements under simulated polar conditions, *Atmos. Chem. Phys.*, 8(22), 6813–6822, doi:10.5194/acp-8-6813-2008, 2008.
- 560 Kleffmann, J., Becker, K. H. and Wiesen, P.: Heterogeneous NO₂ conversion processes on acid surfaces: Possible atmospheric implications, *Atmos. Environ.*, 32(16), 2721–2729, doi:10.1016/S1352-2310(98)00065-X, 1998.
- Kleffmann, J., Heland, J., Kurtenbach, R., Lörzer, J. and Wiesen, P.: A New Instrument (LOPAP) for the Detection of Nitrous Acid (HONO), *Environ. Sci. Pollut. Res.*, 9(special issue 4), 48–54, 2002.
- Kleffmann, J., Gavriloaiei, T., Hofzumahaus, A., Holland, F., Koppmann, R., Rupp, L., Schlosser, E., Siese, M. and Wahner, A.: Daytime formation of nitrous acid: A major source of OH radicals in a forest, *Geophys. Res. Lett.*, 32(5), L05818, 565 doi:10.1029/2005GL022524, 2005.
- Kleffmann, J., Lörzer, J. C., Wiesen, P., Kern, C., Trick, S., Volkamer, R., Rodenas, M. and Wirtz, K.: Intercomparison of the DOAS and LOPAP techniques for the detection of nitrous acid (HONO), *Atmos. Environ.*, 40(20), 3640–3652, doi:10.1016/j.atmosenv.2006.03.027, 2006.
- 570 Klosterkötter, A., Kurtenbach, R., Wiesen, P. and Kleffmann, J.: Determination of the emission indices for NO, NO₂, HONO, HCHO, CO, and particles emitted from candles, *Indoor Air*, 31(1), 116–127, doi:10.1111/ina.12714, 2021.
- Kramer, L. J., Crilley, L. R., Adams, T. J., Ball, S. M., Pope, F. D. and Bloss, W. J.: Nitrous acid (HONO) emissions under real-world driving conditions from vehicles in a UK road tunnel, *Atmos. Chem. Phys.*, 20(9), 5231–5248, doi:10.5194/acp-20-5231-2020, 2020.
- 575 Kurtenbach, R., Becker, K. H., Gomes, J. A. G., Kleffmann, J., Lörzer, J. C., Spittler, M., Wiesen, P., Ackermann, R., Geyer, A. and Platt, U.: Investigations of emissions and heterogeneous formation of HONO in a road traffic tunnel, *Atmos. Environ.*, 35(20), 3385–3394, doi:10.1016/S1352-2310(01)00138-8, 2001.
- Laufs, S. and Kleffmann, J.: Investigations on HONO formation from photolysis of adsorbed HNO₃ on quartz glass surfaces, *Phys. Chem. Chem. Phys.*, 18(14), 9616–9625, doi:10.1039/c6cp00436a, 2016.



- 580 Legrand, M., Preunkert, S., Frey, M., Bartels-Rausch, T., Kukui, A., King, M. D., Savarino, J., Kerbrat, M. and Jourdain, B.: Large mixing ratios of atmospheric nitrous acid (HONO) at Concordia (East Antarctic Plateau) in summer: A strong source from surface snow?, *Atmos. Chem. Phys.*, 14(18), 9963–9976, doi:10.5194/acp-14-9963-2014, 2014.
- Li, D., Xue, L., Wen, L., Wang, X., Chen, T., Mellouki, A., Chen, J. and Wang, W.: Characteristics and sources of nitrous acid in an urban atmosphere of northern China: Results from 1-yr continuous observations, *Atmos. Environ.*, 182, 296–306, doi:10.1016/j.atmosenv.2018.03.033, 2018.
- 585 Li, J., Zhu, C., Chen, H., Zhao, D., Xue, L., Wang, X., Li, H., Liu, P., Liu, J., Zhang, C., Mu, Y., Zhang, W., Zhang, L., Herrmann, H., Li, K., Liu, M. and Chen, J.: The evolution of cloud and aerosol microphysics at the summit of Mt. Tai, China, *Atmos. Chem. Phys.*, 20(22), 13735–13751, doi:10.5194/acp-20-13735-2020, 2020.
- Liu, J., Li, S., Mekic, M., Jiang, H., Zhou, W., Loisel, G., Song, W., Wang, X. and Gligorovski, S.: Photoenhanced Uptake of NO₂ and HONO Formation on Real Urban Grime, *Environ. Sci. Technol. Lett.*, 6(7), 413–417, doi:10.1021/acs.estlett.9b00308, 590 2019a.
- Liu, P., Ye, C., Xue, C., Zhang, C., Mu, Y. and Sun, X.: Formation mechanisms of atmospheric nitrate and sulfate during the winter haze pollution periods in Beijing: gas-phase, heterogeneous and aqueous-phase chemistry, *Atmos. Chem. Phys.*, 20(7), 4153–4165, doi:10.5194/acp-20-4153-2020, 2020.
- Liu, X., Cheng, Y., Zhang, Y., Jung, J., Sugimoto, N., Chang, S.-Y., Kim, Y. J., Fan, S. and Zeng, L.: Influences of relative humidity and particle chemical composition on aerosol scattering properties during the 2006 PRD campaign, *Atmos. Environ.*, 595 42(7), 1525–1536, doi:10.1016/j.atmosenv.2007.10.077, 2008.
- Liu, Y., Lu, K., Ma, Y., Yang, X., Zhang, W., Wu, Y., Peng, J., Shuai, S., Hu, M. and Zhang, Y.: Direct emission of nitrous acid (HONO) from gasoline cars in China determined by vehicle chassis dynamometer experiments, *Atmos. Environ.*, 169, 89–96, doi:10.1016/j.atmosenv.2017.07.019, 2017.
- 600 Liu, Y., Lu, K., Li, X., Dong, H., Tan, Z., Wang, H., Zou, Q., Wu, Y., Zeng, L., Hu, M., Min, K.-E., Kecorius, S., Wiedensohler, A. and Zhang, Y.: A Comprehensive Model Test of the HONO Sources Constrained to Field Measurements at Rural North China Plain, *Environ. Sci. Technol.*, 53(7), 3517–3525, doi:10.1021/acs.est.8b06367, 2019b.
- Ma, Q., Wang, T., Liu, C., He, H., Wang, Z., Wang, W. and Liang, Y.: SO₂ Initiates the Efficient Conversion of NO₂ to HONO on MgO Surface, *Environ. Sci. Technol.*, 51(7), 3767–3775, doi:10.1021/acs.est.6b05724, 2017.
- 605 Marion, A., Morin, J., Gandolfo, A., Ormeño, E., D’Anna, B. and Wortham, H.: Nitrous acid formation on Zea mays leaves by heterogeneous reaction of nitrogen dioxide in the laboratory, *Environ. Res.*, 193, 110543, doi:10.1016/j.envres.2020.110543, 2021.
- Meng, F., Qin, M., Tang, K., Duan, J., Fang, W., Liang, S., Ye, K., Xie, P., Sun, Y., Xie, C., Ye, C., Fu, P., Liu, J. and Liu, W.: High-resolution vertical distribution and sources of HONO and NO₂ in the nocturnal boundary layer in urban Beijing, 610 China, *Atmos. Chem. Phys.*, 20(8), 5071–5092, doi:10.5194/acp-20-5071-2020, 2020.
- Michoud, V., Kukui, A., Camredon, M., Colomb, A., Borbon, A., Miet, K., Aumont, B., Beekmann, M., Durand-Jolibois, R., Perrier, S., Zapf, P., Siour, G., Ait-Helal, W., Locoge, N., Sauvage, S., Afif, C., Gros, V., Furger, M., Ancellet, G. and Doussin, J. F.: Radical budget analysis in a suburban European site during the MEGAPOLI summer field campaign, *Atmos. Chem. Phys.*, 12(24), 11951–11974, doi:10.5194/acp-12-11951-2012, 2012.
- 615 Monge, M. E., D’Anna, B., Mazri, L., Giroir-Fendler, A., Ammann, M., Donaldson, D. J. and George, C.: Light changes the atmospheric reactivity of soot, *Proc. Natl. Acad. Sci. U.S.A.*, 107(15), 6605–6609, doi:10.1073/pnas.0908341107, 2010.
- Ndour, M., D’Anna, B., George, C., Ka, O., Balkanski, Y., Kleffmann, J., Stemmler, K. and Ammann, M.: Photoenhanced uptake of NO₂ on mineral dust: Laboratory experiments and model simulations, *Geophys. Res. Lett.*, 35(5), L05812, doi:10.1029/2007GL032006, 2008.
- 620 Nie, W., Ding, A. J., Xie, Y. N., Xu, Z., Mao, H., Kerminen, V.-M., Zheng, L. F., Qi, X. M., Huang, X., Yang, X.-Q., Sun, J. N., Herrmann, E., Petäjä, T., Kulmala, M. and Fu, C. B.: Influence of biomass burning plumes on HONO chemistry in eastern



- China, *Atmos. Chem. Phys.*, 15(3), 1147–1159, doi:10.5194/acp-15-1147-2015, 2015.
- Oswald, R., Behrendt, T., Ermel, M., Wu, D., Su, H., Cheng, Y., Breuninger, C., Moravek, A., Mougín, E., Delon, C., Loubet, B., Pommerening-Röser, A., Sörgel, M., Pöschl, U., Hoffmann, T., Andreae, M. O., Meixner, F. X. and Trebs, I.: HONO emissions from soil bacteria as a major source of atmospheric reactive nitrogen, *Science*, 341(6151), 1233–1235, doi:10.1126/science.1242266, 2013.
- Peng, Q., Palm, B. B., Melander, K. E., Lee, B. H., Hall, S. R., Ullmann, K., Campos, T., Weinheimer, A. J., Apel, E. C., Hornbrook, R. S., Hills, A. J., Montzka, D. D., Flocke, F., Hu, L., Permar, W., Wielgasz, C., Lindaas, J., Pollack, I. B., Fischer, E. V., Bertram, T. H. and Thornton, J. A.: HONO Emissions from Western U.S. Wildfires Provide Dominant Radical Source in Fresh Wildfire Smoke, *Environ. Sci. Technol.*, 54(10), 5954–5963, doi:10.1021/acs.est.0c00126, 2020.
- Platt, U., Perner, D., Harris, G. W., Winer, A. M. and Pitts Jr, J. N.: Observations of nitrous acid in an urban atmosphere by differential optical absorption, *Nature*, 285(5763), 312–314, doi:10.1038/285312a0, 1980.
- Romer, P. S., Wooldridge, P. J., Crouse, J. D., Kim, M. J., Wennberg, P. O., Dibb, J. E., Scheuer, E., Blake, D. R., Meinardi, S., Brosius, A. L., Thames, A. B., Miller, D. O., Brune, W. H., Hall, S. R., Ryerson, T. B. and Cohen, R. C.: Constraints on Aerosol Nitrate Photolysis as a Potential Source of HONO and NO_x, *Environ. Sci. Technol.*, 52(23), 13738–13746, doi:10.1021/acs.est.8b03861, 2018.
- Schmid, F., Schmidli, J., Hervo, M. and Haeefe, A.: Diurnal Valley Winds in a Deep Alpine Valley: Observations, *Atmosphere*, 11(1), 54, doi:10.3390/atmos11010054, 2020.
- Shi, Q., Tao, Y., Krechmer, J. E., Heald, C. L., Murphy, J. G., Kroll, J. H. and Ye, Q.: Laboratory Investigation of Renoxification from the Photolysis of Inorganic Particulate Nitrate, *Environ. Sci. Technol.*, 55(2), 854–861, doi:10.1021/acs.est.0c06049, 2021.
- Sosedova, Y., Rouvière, A., Bartels-Rausch, T. and Ammann, M.: UVA/Vis-induced nitrous acid formation on polyphenolic films exposed to gaseous NO₂, *Photochem. Photobiol. Sci.*, 10(10), 1680–1690, doi:10.1039/c1pp05113j, 2011.
- Stemmler, K., Ammann, M., Donders, C., Kleffmann, J. and George, C.: Photosensitized reduction of nitrogen dioxide on humic acid as a source of nitrous acid, *Nature*, 440(7081), 195–198, doi:10.1038/nature04603, 2006.
- Stemmler, K., Ndour, M., Elshorbany, Y., Kleffmann, J., D’Anna, B., George, C., Bohn, B. and Ammann, M.: Light induced conversion of nitrogen dioxide into nitrous acid on submicron humic acid aerosol, *Atmos. Chem. Phys.*, 7(16), 4237–4248, doi:10.5194/acp-7-4237-2007, 2007.
- Stutz, J., Alicke, B. and Neftel, A.: Nitrous acid formation in the urban atmosphere: Gradient measurements of NO₂ and HONO over grass in Milan, Italy, *J. Geophys. Res. Atmos.*, 107(22), 8192, doi:10.1029/2001JD000390, 2002.
- Su, H., Cheng, Y., Oswald, R., Behrendt, T., Trebs, I., Meixner, F. X., Andreae, M. O., Cheng, P., Zhang, Y. and Pöschl, U.: Soil Nitrite as a Source of Atmospheric HONO and OH Radicals, *Science*, 333(6049), 1616–1618, doi:10.1126/science.1207687, 2011.
- Theys, N., Volkamer, R., Müller, J.-F., Zarzana, K. J., Kille, N., Clarisse, L., De Smedt, I., Lerot, C., Finkenzeller, H., Hendrick, F., Koenig, T. K., Lee, C. F., Knote, C., Yu, H. and Van Roozendaal, M.: Global nitrous acid emissions and levels of regional oxidants enhanced by wildfires, *Nat. Geosci.*, 13(10), 681–686, doi:10.1038/s41561-020-0637-7, 2020.
- Villena, G., Wiesen, P., Cantrell, C. A., Flocke, F., Fried, A., Hall, S. R., Hornbrook, R. S., Knapp, D., Kosciuch, E., Mauldin, R. L., McGrath, J. A., Montzka, D., Richter, D., Ullmann, K., Walega, J., Weibring, P., Weinheimer, A., Staebler, R. M., Liao, J., Huey, L. G. and Kleffmann, J.: Nitrous acid (HONO) during polar spring in Barrow, Alaska: A net source of OH radicals?, *J. Geophys. Res. Atmos.*, 116(24), D00R07, doi:10.1029/2011JD016643, 2011.
- Wang, L., Wen, L., Xu, C., Chen, J., Wang, X., Yang, L., Wang, W., Yang, X., Sui, X., Yao, L. and Zhang, Q.: HONO and its potential source particulate nitrite at an urban site in North China during the cold season, *Sci. Total Environ.*, 538, 93–101, doi:10.1016/j.scitotenv.2015.08.032, 2015.
- Wang, X., Dalton, E. Z., Payne, Z. C., Perrier, S., Riva, M., Raff, J. D. and George, C.: Superoxide and Nitrous Acid Production



- 665 from Nitrate Photolysis Is Enhanced by Dissolved Aliphatic Organic Matter, *Environ. Sci. Technol. Lett.*, 8(1), 53–58, doi:10.1021/acs.estlett.0c00806, 2021.
- Xue, C., Zhang, C., Ye, C., Liu, P., Catoire, V., Krysztofiak, G., Chen, H., Ren, Y., Zhao, X., Wang, J., Zhang, F., Zhang, C., Zhang, J., An, J., Wang, T., Chen, J., Kleffmann, J., Mellouki, A. and Mu, Y.: HONO Budget and Its Role in Nitrate Formation in the Rural North China Plain, *Environ. Sci. Technol.*, 54(18), 11048–11057, doi:10.1021/acs.est.0c01832, 2020.
- 670 Ye, C., Zhou, X., Pu, D., Stutz, J., Festa, J., Spolaor, M., Tsai, C., Cantrell, C., Mauldin, R. L., Campos, T., Weinheimer, A., Hornbrook, R. S., Apel, E. C., Guenther, A., Kaser, L., Yuan, B., Karl, T., Haggerty, J., Hall, S., Ullmann, K., Smith, J. N., Ortega, J. and Knote, C.: Rapid cycling of reactive nitrogen in the marine boundary layer, *Nature*, 532(7600), 489–491, doi:10.1038/nature17195, 2016.
- Ye, C., Zhang, N., Gao, H. and Zhou, X.: Photolysis of Particulate Nitrate as a Source of HONO and NO_x, *Environ. Sci. Technol.*, 51(12), 6849–6856, doi:10.1021/acs.est.7b00387, 2017.
- Ye, C., Zhou, X., Pu, D., Stutz, J., Festa, J., Spolaor, M., Tsai, C., Cantrell, C., Mauldin III, R. L., Weinheimer, A., Hornbrook, R. S., Apel, E. C., Guenther, A., Kaser, L., Yuan, B., Karl, T., Haggerty, J., Hall, S., Ullmann, K., Smith, J. and Ortega, J.: Tropospheric HONO distribution and chemistry in the southeastern US, *Atmos. Chem. Phys.*, 18(12), 9107–9120, doi:10.5194/acp-18-9107-2018, 2018.
- 680 Ye, Z. J., Segal, M. and Pielke, R. A.: Effects of Atmospheric Thermal Stability and Slope Steepness on the Development of Daytime Thermally Induced Upslope Flow, *J. Atmos. Sci.*, 44(22), 3341–3354, doi:10.1175/1520-0469(1987)044<3341:EOATSA>2.0.CO;2, 1987.
- Zhang, J., An, J., Qu, Y., Liu, X. and Chen, Y.: Impacts of potential HONO sources on the concentrations of oxidants and secondary organic aerosols in the Beijing-Tianjin-Hebei region of China, *Sci. Total Environ.*, 647, 836–852, doi:10.1016/j.scitotenv.2018.08.030, 2019a.
- 685 Zhang, J., Chen, J., Xue, C., Chen, H., Zhang, Q., Liu, X., Mu, Y., Guo, Y., Wang, D., Chen, Y., Li, J., Qu, Y. and An, J.: Impacts of six potential HONO sources on HO_x budgets and SOA formation during a wintertime heavy haze period in the North China Plain, *Sci. Total Environ.*, 681, 110–123, doi:10.1016/j.scitotenv.2019.05.100, 2019b.
- Zhang, L., Wang, T., Zhang, Q., Zheng, J., Xu, Z. and Lv, M.: Potential sources of nitrous acid (HONO) and their impacts on ozone: A WRF-Chem study in a polluted subtropical region, *J. Geophys. Res. Atmos.*, 121(7), 3645–3662, doi:10.1002/2015JD024468, 2016.
- 690 Zhang, N., Zhou, X., Shepson, P. B., Gao, H., Alaghmand, M. and Stirm, B.: Aircraft measurement of HONO vertical profiles over a forested region, *Geophys. Res. Lett.*, 36(15), L15820, doi:10.1029/2009GL038999, 2009.
- Zhou, X., Gao, H., He, Y., Huang, G., Bertman, S. B., Civerolo, K. and Schwab, J.: Nitric acid photolysis on surfaces in low-NO_x environments: Significant atmospheric implications, *Geophys. Res. Lett.*, 30(23), 2217, doi:10.1029/2003GL018620, 2003.
- Zhou, X., Huang, G., Civerolo, K., Roychowdhury, U. and Demerjian, K. L.: Summertime observations of HONO, HCHO, and O₃ at the summit of Whiteface Mountain, New York, *J. Geophys. Res. Atmos.*, 112(8), 1–13, doi:10.1029/2006JD007256, 2007.
- 700 Zhou, X., Zhang, N., TeraVest, M., Tang, D., Hou, J., Bertman, S., Alaghmand, M., Shepson, P. B., Carroll, M. A., Griffith, S., Dusanter, S. and Stevens, P. S.: Nitric acid photolysis on forest canopy surface as a source for tropospheric nitrous acid, *Nat. Geosci.*, 4(7), 440–443, doi:10.1038/ngeo1164, 2011.