

Quantifying urban, industrial, and background changes in NO₂ during the COVID-19 lockdown period based on TROPOMI satellite observations

5 Vitali Fioletov¹, Chris A. McLinden¹, Debora Griffin¹, Nickolay Krotkov², Fei Liu², Henk Eskes³

¹Air Quality Research Division, Environment and Climate Change Canada, Toronto, Canada

²Atmospheric Chemistry and Dynamics Laboratory, NASA Goddard Space Flight Center, Greenbelt, Maryland, USA

³Royal Netherlands Meteorological Institute, De Bilt, the Netherlands

10

Correspondence to: Vitali Fioletov (Vitali.Fioletov@outlook.com or Vitali.Fioletov@ec.gc.ca)

Abstract. The COVID-19 lockdown had a large impact on anthropogenic emissions of air pollutants and particularly on nitrogen dioxide (NO₂). While the overall NO₂ decline over some large cities is well-established, ~~understanding the details is a challenge~~ ~~since multiple source categories contribute because of a variety of sources of NO₂~~. In this study, a new method of isolation of three components: background NO₂, NO₂ from urban sources, and from industrial point sources is applied to estimate the ~~COVID-19 lockdown~~ ~~impact of the COVID-19 lockdown~~ on each of them. The approach is based on fitting satellite data by a statistical model with empirical plume dispersion functions ~~driven by a meteorological reanalysis~~ ~~driven by the observed winds~~. Population density and surface elevation data as well as coordinates of industrial sources were used in the analysis. The ~~tropospheric~~ NO₂ vertical column density (VCD) values measured by ~~the~~ Tropospheric Monitoring Instrument (TROPOMI) on board Sentinel-5 Precursor over ~~262261 (3° by 4°)~~ urban areas for the period from March 16 to June 15, 2020, were compared with the average VCD values for the same period in 2018 and 2019. While background NO₂ component remained almost unchanged, the urban NO₂ component declined by -18% ~~to -28%~~ over most regions. India, South America, and a part of Europe (particularly, Italy, France, and Spain) demonstrated a ~~-40% to -50%~~ urban emissions decline. In contrast, ~~the~~ decline over urban areas in China, where the lockdown was over during the analyzed period, was ~~on average, only -4.43% ± 8% except for Wuhan, where more than 60% decline was observed~~. Emissions from large industrial sources in the analyzed urban areas varied ~~largely~~ from region to region from ~~+54.8% ± 6%~~ for China to ~~-40% ± 10%~~ for India. Changes in urban emissions are correlated with changes in Google mobility data (the correlation coefficient is 0.6662) confirming that changes in traffic was one of the key elements in decline of urban NO₂ emissions. No correlation was found between changes in background NO₂ and Google mobility data. On the global scale, the background and urban component were remarkably stable in 2018, 2019, and 2021, with averages of all analysed areas being all within ±2.5% and suggesting that there were no substantial drifts or shifts in TROPOMI data. The 2020 data are clearly an outlier. The mean background component was -6.0%±1.2% and the mean urban component was -26.7%±2.6% or 20-σ below the baseline level from the other years.

15

20

25

30

1 Introduction

Nitrogen oxides ($\text{NO}_x = \text{NO}_2 + \text{NO}$) are air pollutants that originate from various anthropogenic (fuel combustion) and natural (e.g., biomass burning, lightning) sources and ~~NO_x -whose~~ emissions are regulated in many countries. ~~Satellite measurements of o~~

5 One component of NO_x , NO_2 ~~has-hav~~e a long history ~~of satellite measurements~~. In the stratosphere, the SAGE (Stratospheric Aerosol and Gas Experiment) instrument provided NO_2 profile information through the stratosphere beginning in the mid-1980s (Cunnold et al., 1991). Satellite observations of tropospheric NO_2 columns are more recent and began with the nadir-viewing GOME (Global Ozone Monitoring Experiment) in 1996 (Martin et al., 2002) with several successors, chief among these OMI (Ozone Monitoring Instrument) (Duncan et al., 2015; Krotkov et al., 2016; Lamsal et al., 2015, 2021; Levelt et al., 2018) and, most recently, TROPOMI (Tropospheric Monitoring Instrument) (~~Van-van~~ Geffen et al., 2020; Veeffkind et al., 2012). Collectively these instruments have been used to better understand NO_2 sources, sinks, distributions, and trends (Beirle et al., 2011, 2019; Liu et al., 2016; Lorente et al., 2019; Lu et al., 2015; Martin et al., 2002; McLinden et al., 2012; Stavrakou et al., 2020; Virghileanu et al., 2020) .

15 One primary NO_2 characteristic provided by satellites is tropospheric vertical column density (VCD), a geophysical quantity representing the total number of molecules or total mass per unit of area in the troposphere. The main features of the tropospheric NO_2 VCD distribution are well established. Due to its relatively short lifetime, a few hours within a plume ~~at~~ during the daytime, NO_2 is elevated near sources such as urban areas (Beirle et al., 2019; Lorente et al., 2019; Lu et al., 2015) and industrial ~~sources-locations~~ such as power plants and oil refineries (Liu et al., 2016; McLinden et al., 2012). Over high mountains, NO_2 VCDs are relatively small as the troposphere there is “thinner” with fewer emission sources. Ship tracks and major highways also create elevated NO_2 values on satellite maps (Beirle et al., 2004; Georgoulias et al., 2020; Liu et al., 2020; 20 Richter et al., 2004).

~~Satellite data are widely used to estimate emissions and lifetimes from large NO_2 point sources (Streets et al., 2013). Methods such as inverse modelling (Konovalov et al., 2006; Mijling and Van Der A, 2012) and, more recently, flux divergence (Beirle et al., 2019) are used for such purpose. One common approach is based on a rotation of satellite NO_2 pixels around the source to align the wind data to a common direction, integrating the data across the wind direction and then fitting the results with an exponentially modified Gaussian (EMG) function (Lange et al., 2021; Pommier et al., 2013). The two unknown parameters, the emission strength and lifetime are estimated directly from the fit in one dimensional space. The method works well for isolated stationary point sources and with steady winds. Another approach employs a two dimensional plume function of the wind speed (Dammers et al., 2019; Fioletov et al., 2015). The plume function depends on three parameters: the plume width, lifetime and emission strengths. While all three parameters can be estimated from the fit, the algorithm works better if the plume width and lifetime are estimated in advance and then their prescribed values are used to estimate the emission. This algorithm can be further improved to account for multiple sources (Fioletov et al., 2017).~~ The COVID-19 lockdown had an impact on the tropospheric NO_2 emissions levels, first in China and then worldwide (Bao and Zhang, 2020; Bauwens et al., 2020; Ding et al., 2020; Gkatzelis et al., 2021; Kanniah et al., 2020; Keller et al., 2021; Koukouli et al., 2021; Liu et al., 2020;

Vadrevu et al., 2020; Virghileanu et al., 2020; Zhang et al., 2021). It was demonstrated that NO₂ surface concentrations and VCDs have significantly declined in the U.S. and Canada after mid-March 2020 (Bauwens et al., 2020; Goldberg et al., 2020; Griffin et al., 2020). A decline of about 20% to 25% was observed in the U.S. megacities, as well as over some rural areas. A decline was also reported over Europe (e.g., Bar et al., 2021; Barré et al., 2021), India (Mirsa et al., 2021; Hassan et al., 2021), Pakistan (Ghaffar et al., 2021; Mehmood et al., 2021), Brazil (Dantas et al., 2020; Siciliano et al., 2020), and other parts of the world (Ass et al., 2020; Aydin et al., 2020; Fu et al., 2020) as also discussed in overview papers (Gkatzelis et al., 2021; Levelt et al., 2021). The results of satellite data analysis, however, may depend on the selected area around the city. Changes in industrial point source emissions could be different from urban emissions and there is also free tropospheric NO₂ (Silvern et al., 2019) that could mask the lockdown-related changes in anthropogenic emissions.

The impact of the lockdown on tropospheric NO₂ VCD from satellite data was often estimated by comparing mean or median values over a certain area for the periods before and after the lockdown (e.g., Qu, et al., 2021; Barré et al., 2021; Mehmood et al., 2021; Hassan et al., 2021; see also online tool https://so2.gsfc.nasa.gov/no2/no2_index.html accessed Dec. 18, 2021) or as the values weighted according to the population density (Sannigrahi et al., 2021). This makes the results dependent on the area analyzed and sensitive to the wind speed (Goldberg et al., 2021). There is also free tropospheric NO₂ that could mask the lockdown-related changes in anthropogenic emissions (Silvern et al., 2019). Moreover, meteorological variability was also a contributing factor to the differences (e.g., Barré et al., 2021; Griffin et al., 2020) although some studies found that its impact may not be very large (Bar et al., 2021).

An alternative approach is based on estimation of NO₂ emissions using satellite data and then comparing the emissions estimates before and after the lockdown started (Lange et al., 2021). There are several methods to estimate the emissions (Streets et al., 2013). Methods such as inverse modelling (Konovalov et al., 2006; Mijling and van Der A, 2012) and, more recently, flux divergence (Beirle et al., 2019; 2021) are used for such purpose. One common technique is based on a rotation of satellite NO₂ pixels around the source so the NO₂ data would appear if the wind is from one common direction, allowing many overpasses to be combined. These rotated data are then integrated in the cross-wind direction and then fitting the results with an exponentially modified Gaussian (EMG) function (Lange et al., 2021; Pommier et al., 2013). The two unknown parameters, the emission strength and lifetime are estimated directly from the fit in one-dimensional space. The method works well for isolated stationary point sources and with steady winds (Beirle et al., 2021) but may not work in the areas where emissions from closely located multiple sources are mixed with urban emissions. Another approach employs a two-dimensional EMG plume function of the wind speed (Dammers et al., 2019; Fioletov et al., 2015, McLinden et al., 2020). The plume function depends on three parameters: the plume width, lifetime, and emission strengths. While all three parameters can be estimated from the fit, the algorithm works better if the plume width and lifetime are estimated in advance and then prescribed in the fit to estimate the emission strength. This algorithm was further improved to account for multiple sources or area (Fioletov et al., 2017).

A different approach was used in this study. A statistical model was used to describe the TROPOMI NO₂ data over 3° by 4° areas (roughly, 330 km by 330 km at 42°N) around major cities and isolate three components related to (1) plumes

from urban sources, (2) plumes from industrial point sources, and (3) background NO₂. The parameters of the statistical model link the satellite NO₂ values to proxies related to elevation and population density as well as to locations of large industrial point sources. Then the three components in 2020 were compared to their values in 2018-2019 to study the COVID-19 lockdown impact. The parameters have simple physical interpretation such as point source or area emission rates and the background NO₂ distribution. As the model has only a few parameters and their estimates are based on several hundreds of TROPOMI pixels in each area, statistical uncertainties of the parameter estimates are very small. The variability of NO₂ itself due to meteorological or observational conditions was studied by comparing the parameter estimates for 2018, 2019, and 2021 that were not affected by lockdowns.

In this study, a new method of isolation of three main components: background NO₂, NO₂ from urban sources, and from industrial point sources is applied to estimate the COVID-19 lockdown impact on each of them. The algorithm is based on a multisource plume dispersion function fitting approach developed for sulfur dioxide (SO₂) point and area sources (Fioletov et al., 2017; McLinden et al., 2020). It assumes that each source produces a plume that depends on unknown emission strength and these emission strengths are derived from the best fit to the satellite data. The algorithm was adapted for NO₂ where emissions from urban areas, which tend to be dominated by residential and mobile emission sources, were often a major source sector one of the major factors. Population density data were used as a proxy for such emissions (Lamsal et al., 2013). As the elevation of the landscape plays an important role in the NO₂ spatial distribution, it too was taken into account by the algorithm. Since the approach is based on statistical methods with a large number of parameters to estimate, it was necessary to have a sufficiently long data set to reduce the impact of natural factors such as meteorology that can cause NO₂ VCD differences of ~15% over monthly timescales (Goldberg et al., 2020; Levelt et al., 2021). We use three-month periods, with the averages for estimates for three-month long the period from March 16 to June 15, 2020, are compared to similar estimates those for the same period in 2018 and 2019 for the 3° by 4° areas around 263-261 major cities worldwide. As the study is focused on relative NO₂ changes due to the lockdown, possible systematic errors related to the TROPOMI retrievals (Verhoelst et al., 2021) and the algorithm fitting parameters (Fioletov et al., 2016) play a much smaller role than in the case of absolute emission estimates.

This paper is organized as follows: Section 2 describes various data sets used in the study; the analysis algorithm is discussed in Section 3. In Section 4, the COVID-19 lockdown impact is studied. USA and Canada are analyzed in detail to illustrate the method, then statistics for Europe is are provided and finally results for the entire world are presented. Discussion and conclusions are given in Section 5. The algorithm is described in Appendix. Additional technical information and statistics for individual regions are are given in the Supplement.

2 Data Sets

2.1 TROPOMI NO₂ VCD data

TROPOMI, onboard of the European Space Agency (ESA) and EU Copernicus Sentinel 5 Precursor (S5p) satellite, was launched on 13 October 2017 (van Geffen et al., 2020; Veefkind et al., 2012). The satellite follows a Sun-synchronous, low-

Earth (825 km) orbit with a daily equator crossing time of approximately 13:30 local solar time (van Geffen et al., 2019). At nadir, TROPOMI pixel sizes were $3.5 \times 7 \text{ km}^2$ at the beginning of operation and were reduced to $3.5 \times 5.6 \text{ km}^2$ on 6 August 2019 and the swath width is 2,600 km. TROPOMI NO₂ VCD values represent the total number of molecules or total mass per unit area below tropopause and are often given in molecules or moles (one mole is equal to 6.022×10^{23} molecules) per square metre or centimetre as well as in Dobson Units (DU, 1 DU = $2.69 \times 10^{16} \text{ molec} \cdot \text{cm}^{-2}$). In this study, level 2 TROPOMI data available from the Copernicus open data access hub (<https://s5phub.copernicus.eu>) were used. The reprocessed (RPRO) data version V1.2.2 was used for 2018 and offline mode (OFFL) data of version V1.2.2 to version V1.3.2 were used for 2019-2020. The 2021 data (V1.4.0) were used only to estimate the interannual variability. The standard TROPOMI product, tropospheric vertical columns, based on air mass factors (AMFs) calculated using the vertical profile of NO₂ from the TM5-MP model at $1^\circ \times 1^\circ$ resolution (Williams et al., 2017) was used. In the analysis, we use only data for which the quality assurance value is higher than 0.75 (Van van Geffen et al., 2018). Also, satellite pixels with snow on the ground, a solar zenith angle greater than 80-75 degrees and with cloud radiance fraction above 0.3 were excluded from the analysis.

The specified random uncertainty of a single TROPOMI tropospheric NO₂ VCD measurement is $7 \times 10^{14} \text{ molec cm}^{-2}$ (or 0.026 DU) (ESA EOP-GMO, 2017). Tack et al., (2021) estimated this uncertainty and found it to be $5.6 \pm 0.4 \times 10^{14} \text{ molec cm}^{-2}$. There is some evidence that TROPOMI NO₂ is biased low by 14%-40% over polluted areas due to a limited spatial resolution of the model used to calculate the AMFs (Judd et al., 2020; Verhoelst et al., 2021; Zhao et al., 2020). This bias can be reduced by recalculation of AMFs with higher spatial resolution (Zhao et al., 2020; Tack et al., 2021). The biases caused by limited spatial resolution of AMFs, however, may be lower if spatially averaged data are considered (Ialongo et al., 2020). Moreover, the differences between 2018-2019 and 2020 data, that are studied here, should not be affected by such AMF-related biases if the vertical profiles shapes have not changed (Levelt et al., 2021).

The TROPOMI NO₂ distribution over the US and southern Canada is shown in Fig.1. The data are stratified by the wind speed to highlight some of the features of the NO₂ VCD distribution. NO₂ values are elevated over large cities/highly populated areas, as is particularly evident from the maps for low wind speed where the NO₂ remains close to the source before chemical or physical removal. Fig.1 also illustrates the fact that TROPOMI NO₂ values over megacities are higher under calm winds and lower over high winds (e.g., Goldberg et al., 2020). Elevated NO₂ values are also observed over power plants and mining operations (Goldberg et al., 2021). The NO₂ VCD distribution also depends on local topography (Kim et al., 2021). For example, smaller values over elevated areas such as the Rocky Mountains and Appalachians and higher values over valleys such the California Central Valley are evident from the map. There is also some background NO₂ that can be seen even over remote areas with no major anthropogenic sources: NO₂ VCDs are not negligible (about $5 \cdot 10^{14} \text{ cm}^{-2}$) over vast remote areas such as National Forests in Montana or Algonquin Provincial Park in Ontario as well as over the oceans.

As TROPOMI has only one daily overpass at most locations, diurnal NO₂ variations may affect emission estimates. Measurements from the ground demonstrate, that unlike surface concentrations, the diurnal variations of NO₂ VCDs are relatively small, particularly in spring (Herman et al., 2009; Chong et al., 2018). However, since night time NO₂ information is not available from satellite, we should say that all the results presented here are limited to daytime emissions only.

2.2 Wind data

As in several previous studies (Fioletov et al., 2015; McLinden et al., 2020; Zoogman et al., 2016) the plume dispersion function (discussed below in Sect. 3) is based on the wind speed and direction obtained from the meteorological reanalysis. For each satellite pixel, wind speed and direction were ~~taken from~~ calculated based on European Centre for Medium-Range Weather Forecasts (ECMWF) ERA5 reanalysis data (C3S, 2017; Dee et al., 2011), which were merged with TROPOMI measurements. The wind profile data have one hour temporal resolution and are available on a 0.25° horizontal grid. U- and V- (west-east and south-north, respectively) wind-speed components were then linearly interpolated to the location of the centre of each TROPOMI pixel and to overpass time. The ERA5 wind components at 1000, 950, and 900 hPa were averaged (that approximately corresponds to the mean winds between 0 and 1 km). This interval was comparable to the wind data used
5 in other similar studies: Beirle et al. (2019) used data at 450 m, while Lange et al., (2021), used data from 100 m. The results are not very sensitive to the wind ~~height-profile~~ within this range as was previously investigated by (Beirle et al., (2011) because the boundary layer wind is relatively constant, except close to the surface.

2.3 ~~Elevation, emission, and population~~ Population density data

The Gridded Population of the World (GPW) dataset (SEDAC, 2017) was used as a proxy for the urban component. GPW
15 data are on 0.042 degree (2.5 arc-minute) grid and consists of estimates of human population density (number of persons per square kilometre) based on counts consistent with national censuses and population registers. When lower resolution data were required, they were obtained by averaging the original data within the new grid cells. Information about large city location and population, that was used to select cities for the analysis, was obtained from the World Cities Database available from <https://simplemaps.com/data/world-cities> (accessed on May 10, 2021). ~~To verify the obtained results, another proxy, night lights, was used. The night lights data are from Visible Infrared Imaging Radiometer Suite (VIIRS) onboard Suomi National Polar orbiting Partnership (NPP) available from <https://earthobservatory.nasa.gov/features/NightLights/page3.php> (accessed on May 10, 2021) (Miller et al., 2012).~~
20

2.4 Industrial point source locations

The algorithm of this study requires coordinates of industrial point sources as an input. In addition, emissions data from the
25 U.S. and Canada are used to verify the emissions estimated from TROPOMI data. Information about emission sources was used in two main ways. First, coordinates of selected sources were used to establish locations of point sources in the fitting algorithm. Second, location and emission strengths were used to calculate NO₂ VCD spatial distribution used for comparisons with satellite based estimates. The analysis was done in three stages. The algorithms were originally developed and tested for the U.S. and Canada, then it was applied for Europe. For these regions, detailed information about emission sources was
30 available. Finally, it was applied worldwide, where information about the emission sources was often limited.

For the U.S., 2018-2020 ~~and Canada, monthly~~ point source NO_x emissions from the U.S. Environmental Protection Agency (EPA) National Emissions Inventory (NEI) (EPA, 2020) ~~and annual emissions from the Canadian National Pollutant Release Inventory (NPRI, 2020) were used. Note that, unlike most of the other sources of emission information, the U.S. EPA emissions data are~~ based on a continuous emissions monitoring system (CEMS) ~~are used. , i.e., Note that CEMS database is based~~ on real emission measurements ~~reported with 1-hour resolution that were then averaged over the analysed period. This database includes most of the sources including all large power plants. At the time of this study, the U.S. monthly emissions data were available up to October 2020~~ For sources that are not available from NEI-CEMS, emissions from eGRID database (<https://www.epa.gov/egrid/download-data>, accessed on August 5, 2021) for 2018 and 2019 were used. ~~They are reported as annual emission estimates and we assume that the emission rates are the same throughout the year. This database includes emissions from oil refineries and cement factories that are often not available from NEI-CEMS. Finally, U.S. airport emissions are obtained from 2017 NEI version released in January 2021 (<https://www.epa.gov/air-emissions-inventories/2017-national-emissions-inventory-nei-data>). For Canada, annual emissions from the Canadian National Pollutant Release Inventory (NPRI, 2020) are used. For Canadian sites, only annual emissions were available up to 2018 and so monthly values were calculated by dividing 2018 annual emissions by 12~~ Only sources with annual emissions ~~greater than 1~~ greater than 1 0.5 kt of NO_x per year were selected and used in this study.

~~Information about annual~~ Coordinates of the European ~~emissions industrial point sources were obtained~~ is available from European Pollutant Release and Transfer Register (<https://prtr.eea.europa.eu/>) for 2007-2017 (accessed on March 2, 2021). ~~And those that emitted more than 0.5 kt yr⁻¹ of NO_x are included in the analysis. The emissions database includes two files: one with data for 2007-2017 and the most recent database for 2017-2019. However, the second database does not include data from many countries and the inventories are incomplete. For this reason, we used the data from the former database to establish locations of European point emission sources.~~ The world powerplant database (<https://globalenergymonitor.org/projects/global-coal-plant-tracker/>) was used to find locations of power plants for the global analysis. ~~Note that for the analysis, only source locations, not emissions themselves, are required. Other~~ Missing sources ~~that were detected~~ added from satellite data based on the analysis of the NO₂ residuals maps (see Section 3) ~~and were then identified confirmed~~ using satellite imagery as was previously done in other studies (e.g., McLinden et al., 2016; Fioletov et al., 2016, Dammers et al., 2019; Beirle et al., 2021). Satellite images from such as Google (<https://www.google.com/maps>), Microsoft Bing (<https://www.bing.com/maps>), and Sentinel 2 (<https://apps.sentinel-hub.com/eo-browser/>) maps. Multiple sources were used since some of the images from Google maps are not always up to date and may not show recently-build factories.

2.5 Elevation data

Elevation data used in this study are from 0.03 degree (two-arc minute) gridded global relief ETOPO2v2 database (NOAA, 2006). When lower resolution data were required, they were obtained by averaging the original data within the new

grid cells. Google Each Community Mobility Report data were obtained from <https://www.google.com/covid19/mobility/> (accessed on March 2, 2021).

2.6 Google mobility data and analyzed period

The lockdown periods due to the COVID restrictions varied from country to country, but in most countries, they started in the second half of March 2020. In the analysed Canadian cities, the lockdown started between March 12 and 17. In the U.S., it started between March 18-19 (Atlanta, Los Angeles) and April 2 (Houston). In Europe, the lockdown started as early as on March 8 (Milan), but for most of the cities the lockdown was introduced in after March 14. The second half of March is also the time when the lockdown measures started in many other cities around the world including Auckland, Baghdad, Buenos Aires, Johannesburg, Lagos, Manila, New Delhi, Sydney, and many others (Levelt et al., 2021, their Appendix B).

It is more difficult to determine the time of return to normal activities because the restrictions were often lifted in phases. For example, in the U.S., a “stay at home” order was lifted between April 30 (Texas) and June 11 (New Hampshire). Moreover, there was no formal lockdown in some countries (Belarus, Japan, South Korea, Sweden, Taiwan), but a decline in public activities can be seen even in these countries as well. For example, there was a up to 40% decline in road transport emissions in Sweden even in absence of any formal lockdowns (Guevara et al., 2021). For this reason, we use mobility data as a proxy instead on the lockdown dates to select the analysed period.

The Google Community Mobility Report data (available from <https://www.google.com/covid19/mobility/>, accessed on March 1, 2021) were used to determine a common time period for our analysis. These data represent the changes in the number of people at locations of various types compared to a baseline level. A baseline day represents a normal value for that day of the week. The baseline day is the median value from the 5-week period Jan 3 – Feb 6, 2020. These mobility data can be used as a proxy for the urban traffic (e.g., Guevara, 2021) and are known to be correlated with urban NO₂ emissions (Venter et al., 2020; Bar et al., 2021; Misra et al., 2021). In this study, they were compared to urban and background NO₂ levels in different countries. The mobility data are available for several categories. Results for mobility for “retail and recreation” are presented as this category demonstrated the highest correlation with estimated urban emissions. The “retail and recreation” category covers visits to restaurants, cafes, shopping centres, theme parks, museums, libraries, movie theatres, and similar locations. We will refer to this category as to the “Google mobility data” for brevity.

Fig. 2 shows changes in Google mobility data relative to the baseline period for the regions analysed in this study (see Section 4). During the three month-long period from March 16 to June 15, 2020, the mobility data were below the baseline level in all analysed regions. Note that for the thirteen region, China, Google mobility data are not available, and the lockdown there occurred earlier (in February) except for Wuhan, where the lockdown was lifted on April 8. Note that there was no formal lockdown in the Japan, South Korea, and Taiwan region, although we still see some decline in mobility data there.

3 The Emissions fitting algorithms

This section provides only a general description of the method. The calculation formulas are given in the Appendix. The approach used in this study is based on a linear regression model. All satellite measurements over a certain area during a certain period are linked to locations of industrial point sources as well as to population density and elevation-related proxies by a few parameters that characterise these links. Thus, information from thousands satellite measurements is “compressed” into a handful of parameters and therefore their estimates can have very low statistical uncertainties. Then, satellite measurements can be “reconstructed” using the regression model and contribution of three terms of the model (industrial, urban and background) can be studied. Such model may not be very accurate in “predicting” values of individual satellite pixels, but we will show that it performs well when it is used to describe a three-month mean NO₂ VCD distribution over the analysed areas.

~~This section provides only a general description of the method. The calculation formulas are given in the Appendix.~~ The method is adapted from the previously designed algorithm for multi-sources SO₂ emission estimates (Fioletov et al., 2017) where the emissions are determined from the best fit of satellite observation by a set of plume functions (one per source) scaled by parameters of estimation representing the emission strength. Unlike SO₂, where emissions are mostly generated by point sources, NO₂ emissions also originate from area sources such as large cities. As shown in Fig.1, landscape also has a major impact on the NO₂ distribution. To accommodate these features, the statistical model from Fioletov et al (2017) was modified to:

$$TROPOMI\ NO_2 = \alpha_0 + (\beta_0 + \beta_1(\theta - \theta_0) + \beta_2(\varphi - \varphi_0)) \cdot \exp(-H/H_0) + \alpha_p \Omega_p + \sum \alpha_i \Omega_i + (\beta_0 + \beta_1(\theta - \theta_0) + \beta_2(\varphi - \varphi_0)) \cdot \exp(-H/H_0) + \varepsilon \quad (1)$$

where α_0 , α_e , α_p , α_i , β_0 , β_1 , and β_2 are the unknown regression parameters representing population density-related proxies and emissions from individual point sources and a background with contribution from the elevation; Ω_p is the source plume function for the population density-related distributed source (or area source); Ω_i are the source plume functions for industrial point sources; H is the elevation above sea level and the empirical scaling factor $H_0=1.0$ km was introduced to make the exponential argument dimensionless and to account for altitudinal dependence better; and ε is the residual noise.

Eq.1 is a linear regression statistical model with unknown coefficient sets α and β . There are three main components in the model: the background term, $\alpha_0 + (\beta_0 + \beta_1(\theta - \theta_0) + \beta_2(\varphi - \varphi_0)) \cdot \exp(-H/H_0)$, related to background and elevation (four fitted coefficients), the urban component term, $\alpha_p \Omega_p$, related to the population density (one coefficient), and the industrial term, $\sum \alpha_i \Omega_i$, that represents the contribution from industrial point sources (variable number of coefficients from zero to few dozens). We will refer to them as to background, urban and industrial components.

The fitting was done for 3° by 4° areas around large cities by minimization of the squares of the residuals (ε). The fitting and parameter estimation was done using all individual level 2 pixels for the period from March 16 to June 15 three times: for 2018, 2019, and 2020. So, three sets of coefficients were obtained and then used to estimate the background levels and emissions. Then, the average results for 2018 and 2019 were compared with these for 2020. The fitting was done for all satellite pixels centered within 3° by 4° areas around large cities and collected during a three-month period by minimization

of the squares of the residuals (ϵ). The size of the area is based on the following considerations: The larger the area the less accurate assumptions about a linear gradient of background NO₂ and constant emissions per capita are. The algorithm is based on fitting plumes. For typical plume characteristics (discussed below), the size of fitting area should be long, in order of 100 km, to have enough data for the fit. Finally, the area should be large enough to avoid correlation between the elevation and population density proxies.

As in Fioletov et al., (2017), the plume from an industrial point source i is described by a plume function $\Omega(\theta, \varphi, \omega, s, \theta_i, \varphi_i)$ where θ and φ are the satellite pixel coordinates; ω and s are the wind direction and speed for that pixel; and θ_i and φ_i are the source coordinates. An unknown parameter (α_i) represents the total NO₂ mass emitted from the source i . The emission rate for source i can be expressed as $E_i = \alpha_i/\tau$, where τ is a prescribed NO₂ lifetime (or, more accurately, decay time, but we use the term “lifetime” because it is more common). Note that τ is different from the chemical lifetime (de Foy et al., 2015). Once the emission rate is established, it can be used to reconstruct how distribution of NO₂ emitted by that source would be seen by a satellite, i.e., estimate the industrial component in satellite data.

The plume functions Ω are EMG functions that are commonly used to approximate plumes of VCDs of trace gases such as NO₂, SO₂, and ammonia (Beirle et al., 2011, 2014; Dammers et al., 2019; Fioletov et al., 2017, 2015; de Foy et al., 2015; Liu et al., 2016; McLinden et al., 2020). Similar in concept to a Gaussian plume function, they also take into account the finite physical size of the source and the spatial resolution of the satellite instrument being utilized. The lifetime τ reflects the rate at which NO₂ is removed from the plume due to chemical conversion or physical removal such as deposition; it depends on several factors such as season and NO₂ concentration. It is about 2-6 hours in summer and longer in winter (de Foy et al., 2014; Liu et al., 2016). Moreover, for some sources, the lifetime may be changing over time (Laughner and Cohen, 2019) as NO₂ concentration declines, although other studies suggest that such changes are minor (Stavrakou et al., 2020). Recent TROPOMI-based estimates show that a typical lifetime in urban areas is between 2 and 5 hours in spring and autumn with shorter lifetimes at low latitudes (Lange et al., 2021). While the lifetime has a large impact on the emission estimates, relative changes are less sensitive to it. In addition to τ , the shape of the EMG function depends on the prescribed plume width (w), that depends on the size of the source and the size of satellite pixel. The value of $w=8$ km for plume width was used in this study for TROPOMI along with a constant value of $\tau=3.3$ hours. The switch from 7 to 5.6 km along-track resolution in 2019 might have some impact on the optimal plume width, but as the sensitivity analysis shows that small changes in w have only a minor impact on the results. We estimated that, for the urban component, on average, a 1-hour deviation from the used τ value (3.33 hour) or a 2 km variation in w changes the differences between 2020 and 2018-2019 values only by about 1%.

Unlike many previous studies (Beirle et al., 2011; Fioletov et al., 2016; Lange et al., 2021) where the background offset was presumed to be constant and estimated from, for example, upwind NO₂ data, we included a special term that is responsible for it. In equation (1), the $\alpha_0 + (\beta_0 + \beta_1(\theta - \theta_0) + \beta_2(\varphi - \varphi_0)) \cdot \exp(-H/H_0)$ term represents the “background” NO₂ that is assumed to be declining exponentially with elevation, i.e., within the analyzed 3° by 4° area, the higher is the elevation the lower is the background tropospheric column NO₂ VCD. It was also assumed that this contribution from elevation depends on geographical coordinates only and not on the winds. Even in absence of any sources, there could be some gradient in

5 tropospheric NO₂ over the analyzed area, as for example, over some regions in northern Canada or along the east coast of the U.S. (Fig. 1). To account for such gradients, the linear term $\beta_1(\theta-\theta_0)+\beta_2(\varphi-\varphi_0)$, where θ_0 and φ_0 are the coordinates of the centre of the analyzed area, was added. In other words, it was assumed that there is a linear gradient of background NO₂ within the analyzed area and NO₂ VCD declines exponentially with height over elevated regions. Finally, α_0 was added to the model to account for remaining free-tropospheric NO₂ at high elevations where $\exp(-H/H_0)$ is very close to 0. Its presence gives a better agreement of the fitting results with the satellite data for areas with a high range of elevations. Since this term is a part of the statistical model, all parameters α_0 , β_0 , β_1 , and β_2 are estimated from the fitting. Once they are estimated, the term can be calculated for each location within the analysed 3°×4° area that gives a “background” value for that location that depends on the coordinates and elevation only. For simplicity, we will refer to the term discussed in this paragraph as to the “background” component.

10 Finally, the $\alpha_p \Omega_p$ term represents the emissions contribution from factors, related to human-urban activity. Such emissions can be estimated by establishing a regular grid and then estimating emission for each grid point as was previously done for SO₂ (Fioletov et al., 2017, McLinden et al., 2020). If, for example, we use a 0.2° by 0.2° grid (i.e., 336 (16 x 21) grid cells), this would mean that we need to add 336 unknown coefficients to Eq.1. It would make the coefficient estimates less
15 robust and prevent us from estimating emissions form individual industrial point sources because their plume functions would be highly correlated with the plume functions of the neighboring grid cells. Instead, we assumed that emissions from each grid cell are proportional to the cell population and the coefficient of proportionality is the same for the entire analysed 3° by 4° area. Thus, we just need to estimate one coefficient (α_p) that is proportional annual emissions per capita. This makes the statistical uncertainty of such coefficient very small. In fact, for most analyzed areas, the uncertainty were at least 10 times
20 less that the coefficient itself. The composite plume function Ω_p is, ~~therefore,~~ a sum of plume functions of all individual ~~0.2° by 0.2° grid~~ cell centres multiplied by the grid cell population. Thus, Ω_p depends on geographical coordinates, population density and local winds. The original population density data were converted to a 0.2° by 0.2° by averaging population density data within each grid cell. Smaller grids such as 0.1° by 0.1° were also considered, but it was found that the reduction of the grid size does not change the results, while increase the computation time.

25 The downside of this approach is that the estimates would produce mean emission per capita for a rather large area. This may not be very representative if there are cities with different economical conditions within the analyzed area as, for example, at the border of North and South Korea. Such cases are easily identifiable from the maps of the fitting residuals: such cities would appear as areas of large positive and negative anomalies. We did find several such cases and manually adjusted the area to include only one highly populated area.

30 ~~,where the population density is used as a proxy. ___grid and then each of 336 (16 x 21) grid-cells were considered as point sources located at the grid centre with emissions proportional to the population. However, instead of estimating emission for each such “source” separately, the emissions were parameterized by a single unknown parameter α_p . This means that emissions per capita are assumed to be the same everywhere within the analysed 3° by 4° area Eq.1 is a linear regression statistical model with unknown coefficient sets α and β . The proxy plume functions used in the model preferably should be uncorrelated, because~~

otherwise the coefficients have correlated errors making their interpretation difficult. ~~There are three main components in the model: related to background and elevation (four fitted coefficients), to the population density (one coefficient) and to industrial sources (variable number of coefficients from zero to few dozens). We will refer to them as to background, urban and industrial components.~~ For a typical urban area, these components should be independent: high population density zones typically occupy a small part of the area and industrial sources are typically located away from such highly populated zones. Note that the NO_2 lifetime is relatively short and the median wind speed in, for example, the eastern U.S. is about 10 km per hour, so sources located 30-40 km apart typically have uncorrelated plume functions.

High correlation between the population and landscape-related proxies is possible if a city is in a valley surrounded by high mountains. The correlation could be reduced by increasing the size of the analysed area, but if the area is too large, the assumption that the background level has a linear gradient in the area may not be valid. Therefore, we limited the area to 3° by 4° . The correlation coefficients between the site elevation and population density for 3° by 4° areas are typically small. For example, in the U.S., correlations are positive over Florida (about 0.2) with the population density is higher in the inland area ~~above sea level~~, and negative in the Portland-Seattle-Vancouver area (about -0.35), where it is higher near the ocean and lower in the mountains. As the plume functions of individual industrial sources are very local (~50 km footprint), they do not correlate with the elevation. With such low correlation coefficients, elevation does not affect estimates of other parameters of the regression model.

When industrial point sources are located in close proximity, their plume functions in the statistical model (Eq.1) are highly correlated. In practice, it often appears as if, for example, estimated emissions from one source are unrealistically high, while emissions from the other near-by source are low or even negative. In such cases, emissions from individual industrial sources often cannot be estimated. However, the sources can be grouped into independent clusters and total emission from such clusters can be estimated. Such grouping could be done manually on a case-by-case basis, but it would be subjective and very time consuming. Instead, we applied an algorithm based on factor analysis. We would like to emphasize, that the factor analysis, described in the next two paragraphs, was used to improve emission estimation for individual sources or clusters of sources. It is not required if only total emissions from all point sources in the area are estimated in order to separate them from urban emissions or if all industrial sources are isolated remote sources.

To group industrial sources into clusters, an orthogonalization process was applied to the plume functions of individual industrial sources. First, the correlation matrix for the plume functions of individual point sources (Ω_i) was calculated and eigenvalues and eigenvectors (or “factors”) of the correlation matrix were determined. The correlation matrix was calculated just once using March 16 – June 15 data from all three years. An isolated remote source would appear as an eigenvector with an eigenvalue of 1. Two (or more) closely located, but isolated from others, sources, would have one corresponding eigenvector and an eigenvalue of 2 (or more). Eigenvalues lower than 1 mean that the corresponding sources are already partially included in other eigenvectors. To reduce the number of “factors”, only “factors” with eigenvalues > 0.6 were kept.

The approach based on eigenvalues of the correlation matrix creates proxies that are not correlated and reduces the number of the fitting coefficients. While they correctly describe the total contribution of all industrial sources in the area in the total NO₂ variability (or total emissions), individual ~~eigenvalues~~eigenvectors, i.e., linear combinations of the original plume functions, may not have clear interpretation. For example, they may include the original plume functions with negative coefficients. In order to avoid that and obtain proxies that have a meaningful interpretation, the ~~eigenvalues~~eigenvectors were linearly transformed, so they became as close to the original plume functions as possible, while the correlation coefficients between them remained low. ——— This was done using the varimax factor analysis method that is implemented in modern statistical software packages such as R and SAS (Belhekar, 2013). It orthogonally rotates the established “factors” to maximize the sum of squared correlations between the original variables and factors. Then, the algorithm uses a linear combination of the original variables that have the highest correlations with the rotated “factors”. I.e., the condition of orthogonality is removed in order to find the simplest linear combination of the original variables. In practice, the algorithm produces a set of “clusters”, i.e., linear combinations of the original plume functions, that have low correlation coefficients (typically less than 0.2) between them, and each cluster has high correlation coefficient (typically more than 0.95) with one orthogonal “factor”. To simplify this further, if a linear combination has a weight for an original variable under 0.2, its weight was set to 0. As a result, all non-isolated point sources were grouped into small clusters and emissions estimates were done for such clusters instead of individual sources, while each isolated remote source forms a single-source cluster that corresponds to only that source. It is possible that a single source contributes to more than one cluster that makes interpretation of emissions for such clusters more difficult, but such cases are rare.

As in any regression analysis-based study, correlation between the proxies is one of the main obstacles in the result interpretation. The “orthogonalization” of plume functions from industrial sources largely reduces cross-correlations between the proxies, but high correlations between industrial and population density-related plume functions are still possible if industrial sources are located in highly populated areas. In such cases, it may be difficult to separate its signal from the contribution of the population density-related proxy. For example, in one case (Edmonton, Canada) this correlation coefficient was as high as 0.94 and it was not possible to separate urban and industrial emissions there. ~~For seven cities in the U.S. and Canada it was between 0.81 and 0.85 (for the remaining sources, it is less than 0.77).~~ Note that for large cities and small industrial sources, high correlation means that the ~~industrial~~-emissions from such industrial sources cannot be reliably estimated, although the impact on estimation of the population density-related signal is small. For this reason, industrial point sources located in the 0.2° by 0.2° cells where population is greater than 600,000 people were excluded. This is an empirically estimated limit, and, in a few cases of very large cities (New York, Moscow), it was manually adjusted. We monitored the correlation coefficients between industrial and population density related plume functions and, in some cases, excluded certain sources or even certain urban areas from the analysis.

The fitting and parameter estimation was done using all individual TROPOMI level 2 pixels for the period from March 16 to June 15 four times: for 2018, 2019, 2020 and 2021. So, four sets of coefficients (one set per year) were obtained and then used to estimate the background levels and emissions. Then, the results for 2018 and 2019 were compared with these

for 2020. We also performed the same analysis for 2021, but these results were only used to analyze interannual variability because COVID-19 lockdowns may still have some impact on NO₂ in 2021.

As the regression model has three main terms ~~urban or~~ (background, urban and industrial), the NO₂ VCD for each TROPOMI pixel is represented in Eq. 1 as a sum of three values (components) plus a residual error. Then the values of individual components and residuals can be analyzed the same way as the original TROPOMI measurements, e.g., mean values over a certain period (in our case, March 16– June 15) can be calculated as a function of latitude and longitude.

This is illustrated in Fig. 32, where individual terms of Eq. 1 are shown for an area ~~centered~~ centered on Montreal. The area includes two large cities, Montreal (4.2 million) and Ottawa (1.4 million, including the sister city of Gatineau). The terrain elevations in the analysed area are in the range from just a few metres above sea level along the Saint Lawrence River to more than 500 meters 100 km north of Montreal. For this plot (as well as for Fig. 1 and other figures), we used a non-linear scale that is more sensitive to small quantities in order to make small deviations more pronounced. The top row of Fig. 32 shows the mean TROPOMI NO₂ data (Fig. 32a), the fitting results (Fig. 32b), and the difference between them or the residuals (Fig. 32c). Individual fitting components such as ~~T~~ the background-term ($\alpha_0 + (\beta_0 + \beta_1(\theta - \theta_0) + \beta_2(\varphi - \varphi_0)) \exp(-H/H_0)$), urban ($\alpha_p \Omega_p$), and industrial components are shown in Fig 32 panels d, e, and f respectively.

The contribution of industrial point sources ($\sum \alpha_i \Omega_i$) is illustrated by Fig. 32 g-i. In the case of Montreal, total emissions from industrial sources are relatively small, less than 1.8 kt per year. Note that unlike the previous algorithm (Fioletov et al., 2017), where Ω_i represented plume functions from individual sources, this new Ω_i represents plume functions of clusters of closely located individual sources determined by factor analysis. The estimated parameter α_i represents total NO₂ mass of the entire cluster, while Ω_i is a weighted sum of plume functions of individual sources in the cluster. The weighting coefficients are determined by the varimax technique, described above. In the case of Fig. 32, the first cluster is comprised of two sources and the second and third clusters are each just single point sources. The estimated parameter α_i represents emission from the entire cluster, required that $\alpha_i \geq 0$.

The background and urban component maps have a simple interpretation. Figures 3 j and k shows maps of the elevation and population density respectively. Not surprisingly, the background component, that is dominated by scaled elevation, looks similar to the elevation map itself. The urban component is the population density map convoluted with EMG functions and therefore it looks like a smoothed population density map.

The suggested algorithm essentially finds the emission levels that give the best agreement with the TROPOMI data NO₂ VCD and then uses these estimates to “reconstruct” the spatial NO₂ distribution as well as contribution from each source. As explained by Fioletov et al., (2017), the technique of satellite VCD reconstruction from fitted coefficients α_i using Eq.1 to isolate different components, can be applied to the reported emissions E_i , by using $\alpha_i = E_i \tau$. This produces a map of VCD that would be seen by satellites if these reported emissions are the only sources of NO₂. The same approach was employed here using U.S. emission inventories. For such estimates, the ratio between NO_x and NO₂ is required. Beirle et al., (2021, their Fig. 2), has recently estimated the NO_x to NO₂ ratio for different parts of the world and found that the ratio is about 1.4 over the U.S. and typically between 1.2 and 1.6. elsewhere. The value of 1.4 was used in this study.

The quality of a regression model can be described in terms of the correlation coefficient between the original and predicted values. In case of Montreal, the correlation coefficient of about 0.55. I.e., a set of about half of million original TROPOMI observations over the 3° by 4° area during a three-month period can be described by just 8 parameters (a_p , a_0 , β_0 , β_{1_s} and β_2 plus 3 coefficients α_i for industrial sources). The model can be further improved by adding parameters responsible for workday- weekend differences, seasonal changes, and meteorological proxies (Goldberg et al., 2021; Kim et al., 2021). However, we focused on the mean NO₂ changes over a 3-months period, and they can be successfully estimated without such additional parameters. Fig. 3 shows that the fitting results are able to reproduce such mean data accurately: in the case of Montreal, the coefficient of determination (R^2), i.e., the ratio of the variance of the residuals (Fig. 3 c) to the variance of the averaged TROPOMI data (Fig.3 a) is between 0.9 (in 2019) to 0.93 (in 2020) meaning that fitting results “explain” from 90% to 93% of the observed variance. The Pearson correlation coefficient I between the mean TROPOMI data and the fitting results is about 0.96 (Fig. 3 d).

The necessity of both linear gradient- and elevation-related components in the background term in Eq.1 is illustrated by Fig. 4. If the surface is nearly flat in the analysed area (as, for example, in the case of Minneapolis, Fig. 4 h), the background component is dominated by the linear gradient. However, the elevation affects the NO₂ distribution near mountain areas as, for example, in case of Seattle, where mountains as high as 2000 m are located east of the city (Fig. 4 i). It is interesting to note that the background components are practically identical for both periods that gives a high confidence in the obtained results. The influence of the landscape on the NO₂ distribution also explains why the distribution near Seattle does not look like a “hotspot” NO₂ distribution near a typical large urban area.

Since the fitting results are based on just a handful parameters, the approach of this study is to investigate changes in these parameters or the three regression terms themselves between 2020 and 2018-2019.

4 NO₂ VCDs over some urban areas

To test the method, the described technique was applied to the 22 largest urban areas in the U.S. and 5 in Canada (Fig. 5). Four examples with detailed analysis of the components of the NO₂ distribution are discussed below with results shown in Fig. 6. Eight types of maps are shown. They include mean values for the analysed period for the actual TROPOMI data (column a), the fitting results (b), the residuals (c), i.e., (a) minus (b), as well as individual components of the fitting: the background (d) and urban (e) components, and the industrial source clusters (f).

As mentioned in Section 3, reported emissions can be used to “reconstruct” VCD distribution for NO₂ emitted from these sources using Eq.1. The suggested algorithm essentially finds the emission levels that give the best agreement with the TROPOMI data and then uses these estimates to “reconstruct” the spatial NO₂ distribution as well as contribution from each source. As explained by (Fioletov et al., 2017), the technique of VCD reconstruction from fitted coefficients α_i using Eq.1 used to isolate different components, can be applied to the reported emissions E_i by using $\alpha_i = E_i \times \tau$. This produces a map of VCD that would be seen by satellites if these reported emissions are the only sources of NO₂. The same approach was employed

here. In the U.S., directly measured industrial NO_x emissions from point sources are available. It was assumed that the NO_2 to NO_x ratio is constant 0.7. The maps of NO_2 VCDs from the reported bottom-up emissions is shown in Fig. 64 (column *g*). We would like to emphasise that such reconstruction is based on industrial emissions data only, without any satellite NO_2 observations (although τ and w in the plume functions were the same as in the satellite-based estimates). Finally, the maps of the difference between TROPOMI industrial sources-related component (Fig. 6, column *f*) and NO_2 VCD from reported “bottom-emissionup” emission-based reconstruction (Fig. 6, column *g*) is also shown in Fig. 6 (column *h*).

4.1 Case studies

Four examples that represent different cases of NO_2 distributions around large urban areas are discussed below. In the case of Boston, there is a single urban source with no large industrial sources nearby and with relatively small impact from the terrain. The Atlanta area represents the case where the urban component is well-separated from industrial sources and the area also contains the world largest airport. In the Pittsburgh area, industrial and urban sources have comparable contributions, and the TROPOMI-base industrial emissions estimates can be validated by EPA NEI CEMS measured emissions. Multiple industrial sources in the Houston area are missing from the used EPA NEI CEMS emission database and in this example emissions from the EPA eGRID database can be compared with TROPOMI-based estimates. this example illustrates why it was necessary to add locations of additional sources to explain the observed NO_2 distribution in the area.

Boston is a major urban area with a population of more than 8 million (for the Combined statistical area of Greater Boston). On the TROPOMI NO_2 map (Fig. 64, column *a*), it appears as a large “hotspot” that can be successfully reproduced by the statistical model (Eq.1) using the population density as a proxy. From our estimates, there is a -25.24%±2% decline (the error bars correspond to 2- σ for random uncertainty, see Section 4.3) in the urban emissions in 2020 compared to the 2018-2019 average. Our estimates of urban emissions changes in 2020 are similar to the -22.8% drop in TROPOMI NO_2 values estimated by Goldberg et al., (2020) for the period from March 15 to April 30; and -18.3% drop estimated by Bar et al., (2021) for the period from March 22 to May 30. Boston also shows one of the largest in the U.S. declines in the background component (about +5.20%±0.5%). Although the background component is not linked to particular plumes, it is likely that very high emissions from the largest in the U.S. NO_2 hotspot over the New York-Philadelphia contributed to the background NO_2 over Boston and decline in emissions there caused changes in the background NO_2 over Boston.

Our estimate of the urban emission decline for Atlanta is about -35%±2%. It is more than a -20% decline estimated by Goldberg et al., (2020). However, changes in the background component were only about -13%±0.4%, while the background component over Atlanta is comparable to the urban component, so decline of the sum of the two components over Atlanta should be about -23%. The urban component estimates are based on fitting of the plume from the city itself where NO_2 is dominated by on-road vehicle emissions. Kondragunta et al., (2021) estimated that the decline in on-road emissions is about -28%, that is closer to our estimate. The Atlanta area also hosts the Hartsfield–Jackson Atlanta International Airport (labelled as “1” in Fig 6, column *h*), the world’s busiest airport with more than 100 million passengers per year in 2018-2019 (<https://aci.aero/data-centre/annual-traffic-data/passengers/2017-passenger-summary-annual-traffic-data/>). — The Atlanta

airport NO₂ signal can be easily isolated since the airport is located far away from industrial sources (the correlation coefficients between the plume functions are less than 0.2) and ~~on-at~~ a distance from Atlanta city's most populated area (the correlation coefficient is 0.54). VCDs estimated for the industrial source clusters (column *f* in Fig. 64) are in line with those based on reported emissions (column *g* in Fig. 64). ~~When examining their difference (column *h*), the airport appears as a hotspot as its emissions are not included in the reported emissions inventory.~~ Our estimated annual emissions for the airport are 5.1 ± 0.2 , 6.4 ± 0.2 , 2.9 ± 0.2 kt y⁻¹ in 2018, 2019, and 2020 respectively, while the EPA emissions inventory value is 3.7 kt y⁻¹ for 2017 (the last available year). Thus, our estimates show a 55% decline in airport emissions between 2019 and 2020. The decline in aircraft operations for the analysed period was about 75% for passenger flights and 25% for cargo operations (according to Department of Aviation, Hartsfield-Jackson Atlanta International Airport, <https://www.atl.com/business-information/statistics/>, accessed on Nov. 15, 2021). All industrial sources (power plants) are also far away from the populated area and signals from these sources can be clearly separated from the population density related signal. For illustration purpose only, for Hartsfield-Jackson Atlanta International Airport, the 2017 EPA emissions inventory value was used to calculate NO₂ VCD in column *g* of Fig. 6 for 2018 and 2019 and ~~The area is flat, and the background level does not change very much within the area except for mountains to the north.~~ a half of that value for 2020.

The Pittsburgh area ~~includes the cities Pittsburgh (population of ~2.4 million) and Cleveland (~3.6 million) and~~ has one of the highest emissions from industrial sources among the analysed areas. Several coal-burning power plants are located east, west, and south of the city. Their emissions ~~from them~~ are comparable or even larger than from ~~Pittsburgh the city~~ itself. The NO₂ distribution around major industrial sources “reconstructed” from the reported emissions (Fig. 6, column *g*) is similar to the NO₂ distribution from industrial sources based on satellite estimates (Fig. 6, column *f*). The differences (column *f* minus column *g*) are small although NO₂ from ~~the~~ reported emissions are slightly larger for the cluster of power plants east of the city. The total reported emissions from all industrial sources in the Pittsburgh area are 43, 37, 26 kt y⁻¹ for 2018, 2019, 2020 respectively, while our estimates are 36, 34, 24 kt y⁻¹ (with 2-σ uncertainty of about 1.5 kt y⁻¹), i.e., the 2020 decline from our estimates is 35%, while the decline in reported emissions is 31%. The urban emissions declined from ~~about 4972 ± 1.1 ± 2.3 kt y⁻¹ in 2018- and 2019 to 2536 ± 1.2 kt y⁻¹ in 2020, i.e., by -54.50%.~~

In the case of Houston, the EPA NEI CEMS emissions inventory contains emissions from the power plants in the area, but not from large oil refineries that are responsible for hotspots seen on the TROPOMI mean NO₂ plot. Their coordinates and emissions estimates were obtained from the eGRID inventory. The reported industrial emissions values for the analyzed Houston area in 2018-2019 are 17 kt y⁻¹, while our estimates are 36 and 31 kt y⁻¹ for 2018 and 2019 respectively and the estimated value for 2020 is 33 kt y⁻¹. It appears that TROPOMI-based emissions estimates agree with emissions from the power plants from CEMS but noticeably larger than emissions from oil refineries available from the eGRID inventory (Fig. 6, column *h*). Our estimated changes in background and urban components for the Houston area are $-2.3\% \pm 0.4\%$ and $-18\% \pm 1.6\%$ respectively, i.e., we see a decline in the urban component and practically no changes in the two other components. Goldberg et al., (2020) estimated the decline over Houston in -15.6% , although the spread between three used methods of estimation is large, from -26.3% to -1.9% . Note that the lockdown period in Houston was relatively short, from April 2 to April 30. ~~In order~~

to reduce the residuals, we added the coordinates of 10 known refineries as industrial sources in Eq. 1. As a result, the residuals became small, although some elevated NO₂ values can still be seen in the industrial areas suggesting the list of refineries included in this analysis may not be complete. This example illustrates that it is necessary to add some sources to the list of sources based on the used EPA inventory, derived from measured emissions. A total of 56 sources across the US and Canada were added and the list of sources in the Supplement (file Additional_Sources_Canada_US.xls)

4.2 Relative contribution of different components

NO₂ VCD represents total number of molecules, and equivalently mass, per area unit. When ~~individual background, urban, and industrial~~ components affecting of the NO₂ distribution are ~~known~~ estimated as described in Section 3, it is possible to calculate the total NO₂ mass, of each of the components and estimate their relative contribution to the total NO₂ mass-based on Eq.1. The diagram in Fig. 76 shows such contribution of individual components for the Montreal area (Fig. 3 d, e, f). Most NO₂ mass is associated with the term related to the background component. For the Montreal area, the contribution of industrial sources is four times less than the contribution of the urban component and the two of these components are responsible for less than a quarter of the total NO₂ mass in the area. Relative contribution of the three components in the other areas for the 2018-2019 period are shown in Fig. S17. Most of NO₂ mass belongs to the background component that is not directly linked to particular urban or industrial sources. Plumes from urban and industrial sources, These plumes, or to be precise, $\alpha_p Q_p$ and $\Sigma \alpha_i Q_i$ terms in Eq.1, are responsible for ~~less than~~ about a third of a third of total satellite estimated NO₂ mass in New York and Los Angeles and far less in the other all of the analysed 3° by 4° urban areas in the U.S and Canada. This result depends on a particular size of the area, but the fraction of the background component is larger for larger areas as all major urban areas are already included in the analysis. ~~Most of NO₂ mass belongs to the background component that is not directly linked to particular urban or industrial sources.~~ Fig. S17 also shows that NO₂ mass emitted from cities are larger than emissions from the industrial sources for most of analysed areas in the U.S. and Canada.

————— Note that characteristics such as the mean background value and annual emissions per capita are much less dependent on the area size and the rest of the study is focused on them. ~~The total industrial emissions obviously depend on the sources located in the analysed area.~~

The mean NO₂ distribution near major emission sources has sharp gradients ~~and estimates of the NO₂ lifetime from satellite data that~~ suggest that this the NO₂ lifetime time is relatively short, on the order of a few hours that is also confirmed by direct estimates (Beirle et al., 2011; de Foy et al., 2015). However, large background component may suggest that the lifetime should be relatively long since NO₂ distribution follows the terrain over large areas. This difference in the lifetime could be reconciled if we assume that a fraction of NO₂ emitted from cities and industrial sources gets into free troposphere and have a longer lifetime there than near the ground. Also, levels of the OH radical, the main chemical NO_x sink, within a plume can be much larger than under “clean” conditions and NO₂ lifetime could be longer under such condition than in the plume (Juncosa Calahorrano et al., 2021). Other sources, e.g., lightning or soil emissions may contribute to background component NO₂ directly. The background term can also include components of stratospheric NO₂ that was imperfectly

removed as part of the retrieval algorithm (von Geffen et al., 2020). Finally, estimates of NO₂ lifetime from TROPOMI data (e.g., de Foy et al., 2015; Liu et al., 2016; Lange et al., 2021) are based on daytime observations only. However, the lifetime at night could be different (Kenagy et al., 2018) and nighttime emissions and NO_x evolution during the nights are not reflected in our estimates.

5 **4.3. Variability and uncertainty estimates**

Two characteristics of uncertainties of the estimated NO₂ components are calculated, and the results are presented in Table 1. Uncertainties related to the random measurement errors can be estimated assuming that the residuals ϵ in Eq. 1 are uncorrelated and have the same variance. They represent how precise are the estimates of individual components. Since the total number of satellite pixels in the statistical model is very large, several hundred thousand, and the number of parameters is small, such uncertainties are typically low. These uncertainties are calculated for the three components in each analysed year and the average value for each area (in percent) are given in Table 1 as “random error”. On average, these random errors are about 0.25%, 1% and 3% for the baseline, urban, and industrial components respectively. The random uncertainty represents how precise the component value is calculated and provides the lowest limit of the total uncertainty.

Interannual variability is another characteristic that reflect uncertainties related to the contributions from meteorology, possible instrument or algorithm-related issues, differences in sampling due to variations in cloud cover, and perhaps, other factors. It is also affected by the changes in emissions themselves. Internal variability can be estimated by comparing the components, estimated for different years. The 2020 data are not used this estimate since they were largely affected the lockdowns. Instead, we added estimates for 2021 and calculated the standard deviations from the three values (Table 1). The uncertainties estimated from year to year variability. Although estimates from just three data points are not very reliable, they show similar results for most of the analyzed areas and their average can be used as a characteristic of the interannual variability. The average standard deviation of the interannual variability for the background component is only 7.5%. The interannual variabilities for the urban and industrial components are 10% and 18% respectively. The interannual variability represents the upper limit of the total uncertainty.

The uncertainty of the percentage change between 2018-2019 and 2020 values is a combination of the uncertainty of the baseline, estimated from just two years and the uncertainty of the 2020 value. This gives the following values for 2- σ confidence limits for the percent changes: 18%, 24%, and 44% for the baseline, urban, and industrial components respectively.

4.4 The COVID-19 lockdown impact: the U.S. and Canada

The ability of the method to isolate individual components of the satellite measured total NO₂ mass makes it possible to estimate the impact of the COVID-19-related lockdown on these components separately. As mentioned, we compared the averages for the period from March 16 to June 15 in 2018 and 2019 to the same period averages for 2020.

To illustrate the changes in the background component, Fig. 8 (top) shows the mean VCD values of that component shown in Fig 46, column d (or, in other words, the mean value of $\alpha_0 + (\beta_0 + \beta_1(\theta - \theta_0) + \beta_2(\varphi - \varphi_0)) \cdot \exp(-H/H_0)$) for the

analyzed areas for the two time intervals (left column) as well as the percentage change in 2020 vs. 2018-2019 values (right column). The mean value of decline for the background components among all urban areas is $-46.59\% \pm 3.03\%$. As mentioned, the largest decline of the background component was observed at Boston. The decline was also large, about -20%, over two areas (Edmonton and Calgary) in Canadian province of Alberta. It is unlikely that this decline is related to the lockdown: the restrictions in Alberta were not as tight as in many other areas: only some non-essential services were closed on March 27 and the restriction started to be eased in May (<https://edmonton.citynews.ca/2020/12/24/2020-look-back-albertas-pandemic-response/> accessed on Nov. 20, 2021). In 2020, Alberta had a “historically low” level of forest fires: by June of 2020, fires had burned just about 450 hectares of forest, compared to 650,000 hectares by June of 2019. (<https://globalnews.ca/news/7396849/alberta-2020-slow-wildfire-season/> accessed on Nov. 10, 2021) and therefore, lower than normal natural NO₂ emissions and lower background levels.

The changes in the urban component are shown in Fig. 8 (middle row) expressed as annual NO₂ emissions per capita. Recall, that emission rate is the mass divided by the constant lifetime (that was, therefore, the percent changes in emissions per capita and the changes in total mass are identical. The relative changes for the urban component (Fig. 8 right column) are typically larger than those for the background component. The average total emissions per capita declined by -28% in 2020 compared to 2018-2019 average (from 5.6 to 4.2 kg per capita per year). The median value of decline among all urban areas is -3226% and the mean value of percentage decline is $-2627\% \pm 5.66.2\%$. The changes in emissions per capita are rather uniform except Vancouver where 2020 emissions are 15% larger than the average 2018-2019 emissions. The reason for that is not immediately clear but may be related to unusual meteorology and persistent cloud cover there in 2020. Information about statistical significance of such “outlier” ~~can be obtain from estimates of interannual variability based on 2018 and 2019 data. The variance of the difference between 2018 and 2019 mean values is equal to the variance of the mean NO₂-value multiplied by two. While two years from one area is not enough to estimate the variance, the variance can be reliably estimated if we assume that the interannual variability is the same for all analysed areas (See Supplement A). This gives us the error bars for the Vancouver 2020 value: $15\% \pm 19.5\%$, i.e., this value is within the natural variability range discussed in section 4.4.~~ Edmonton is excluded from this panel ~~excluded from this analysis~~ because ~~the plume functions from~~ two industrial sources are ~~highly correlated to the population density related plume function (see section 3) located in the city itself~~ and, therefore, it is hard to separate their emissions from ~~them~~ the urban emissions. Without such separation, industrial emissions are counted as population density-related that makes Edmonton annual per capita emissions nearly twice larger than emissions for other cities (see Supplement B).

The number of sources and emission strength from large industrial sources varies from area to area. Some areas, e.g., Las Vegas, do not have such emissions sources at all. The total emission from all large industrial emission sources and percentage change in emissions are shown in Fig. 8 (bottom row). The mean value of percentage decline in the areas with industrial sources is $-2822\% \pm 611\%$. Unlike background and urban components, changes in emissions from industrial point sources demonstrate rather large scattering from one area to another. It is not a surprise since in addition to the difference in

~~the strength and length of the lockdown between the areas, there is a difference in the lockdown impact on various industrial sectors. It is not a surprise since different industrial sectors were affected by COVID-19 differently.~~

Overall, the 2020 values for the baseline and industrial components at individual sites are within $2\text{-}\sigma$ limits of the interannual variability (the grey dashed lines in the right panel of Fig. 8) with just few exceptions, while 16 of 27 urban component values are outside these limits.~~For this reason, we region with very different meteorological conditions (assuming that deviations for individual areas are not correlated), i.e., well outside the limits of natural variability. The approach that is focused on regional statistics rather than on individual areas is used for all other regions in this study.~~

Since industrial point source emission estimates are obtained as part of our TROPOMI NO₂ VCD data analysis, such estimated emissions can be compared to the reported ones. In general, there is an agreement between estimated and reported emissions, as was already demonstrated in Fig. 46 a. The scatter plot of estimated vs. reported emissions in 2018-2019 is shown in Fig. 9 for the U.S. urban areas. ~~Each dot on the plot corresponds to one urban area emissions in either 2018 or 2019 with the total of 40 data points. For this analysis, in the fitting we used only point sources where reported emissions data were available.~~ The correlation coefficients between the two data sets from Fig. 9 is 0.984. ~~The slope of the regression line is about 0.7 suggesting that, on average, our estimates are 30% higher than reported emissions. The standard deviation of the residuals is about 5 kt y⁻¹, and that gives an approximate uncertainty for point source of NO₂ emissions that includes all possible sources of errors. Note that 2020 US EPA NEI reported emission were incomplete at the time of this study.~~

~~As noted in Section 4.3, the interannual variability is rather large and therefore, the decline for individual areas is often not significant. For this reason, we analysed 27 individual areas covering a vast region with very different meteorological conditions, so the average of individual area estimates can be calculated with high confidence. Indeed, the ratios of changes in urban and industrial components to their standard deviations are 8.5 and 5.9 respectively (assuming that deviations for individual areas are not correlated), i.e., well outside the limits of interannual variability. The approach that is focused on regional statistics rather than on individual areas is used for all other regions in this study.~~

4.4.5 The COVID-19 lockdown impact: Europe

The described technique was applied to the European Union countries (plus non-members from former Yugoslavia) where detailed information about the industrial emissions sources is available. The analysis was also done for 3° by 4° areas around 36 largest European cities with population greater than 1 million plus some national capitals with population more than 500,000. ~~At the time of this study, the 2020 emissions data were not available, and the 2018-2019 emissions inventory was incomplete with data from many countries missing. Unlike the U.S., the available data were only annual, not monthly, and emissions were typically estimates and not direct measurements. We used the 2017 inventory to obtain the coordinates of the largest sources and then use them in the fitting algorithm.~~ Note, that to avoid double-counting, if more than one city located within an area, we used that area just once (e.g., Manchester and Birmingham are in one area).

The absolute and relative changes between 2018-2019 and 2020 for the three components are shown in Fig. 10. The NO₂ decline of the urban component was particularly large, more than 50%, for the countries in the most western part of the

continent where the strictest lockdown measures were taken: France, Spain, and UK (Fig. 10 a). In contrast, the decline in the German, Czech and some other East European cities was only 20-25% (Fig. 10 b). For this reason, two sub-regions were formed for the analysis: Europe-1 (Italy, France, Spain, Portugal, Belgium, Ireland, and UK) and Europe-2 with all other countries. In general, the mean background values and estimated NO₂ emissions rates per capita in Europe are similar to those in the U.S. and Canada. However, relative changes are somewhat different.

In 2018-2019, the estimated annual emissions per capita for both European regions were very similar to those for the U.S. and Canada. In 2020, the urban component declined in almost every analysed area. The average declines for Europe-1 and Europe-2 regions were -52.54%±4% and -14.13% ±8% respectively. This is in general agreement with total NO_x emission reduction for these two European sub-regions: -50% for Europe-1 countries (Italy, Spain, France) and -15% to -25% for Europe-2 countries (Germany, Sweden) with -85% of the total reduction attributable to on-road transport (Guevara et al., 2021). The decline in Europe-1 was rather uniform with all but one area demonstrating a decline of more than -40%. In contrast, only two areas demonstrated a -40% decline in Europe-2, while most of the areas had a decline under -20%. Two areas in Europe-2 (Budapest and Belgrade) demonstrated an increase in NO₂. They are located 320 km apart and it is possible that relatively high NO₂ values there were caused by some specific meteorological conditions in spring of 2020: the NASA GEOS Composition Forecasting (GEOS-CF) simulations show a positive NO₂ anomaly over Hungary in April-May 2020.

As in the case of the U.S. and Canada, the mean background component in Europe shows a smaller decline than the urban component. On average, it was -5.9%±2% and -11.5±3% lower in 2020 than in 2018-2019 for Europe-1 and Europe-2 regions respectively, but it was pretty consistent as almost all individual areas demonstrated a decline. Large decline in population-related emissions and relatively small decline in the background component for Europe-1 and the opposite for Europe-2 may create an impression that here is anticorrelation between the background level and population-related component, but it is not true. The large decline in average background for Europe-2 was caused by large negative background values for the Scandinavian countries in 2020 that also had large negative changes in the urban components. As discussed later in Section 4.6.5, there is no correlation between the changes in the background levels and urban component.

The emissions from industrial sources also demonstrated a decline, although the scattering of the values is large as the changes varied from country to country and from sector to sector. Guevara et al., (2021) estimated that the emission decline in the energy industry was up to -30% Italy, but under -5% in Sweden. The emission decline from the manufacturing industry was smaller, from about -15% in Italy, Spain, France, UK to -5% in Germany and near zero in Sweden. Emissions from aviation were reduced by -90% in all European countries (Guevara et al., 2021). We estimated changes in emissions only for industrial sources that are located in the analysed areas around major cities and therefore do not represent the entire industrial emissions, but our estimates also show a difference between Europe-1 and Europe-2 regions: the average decline values -34%±10 and -13%±16%, respectively.

The uncertainty estimates are also in general similar to those for the U.S. and Canada: the random uncertainty is about 0.25% for the background component and 1% for the urban component. The natural-interannual variability estimates are also similar for the background component (6.8% and 7.5% for Europe-1 and Europe-2 respectively). The natural-interannual

variability for the urban component for Europe-1 (10%) is also the same as that for the Canada-U.S. region, but higher (15%) for the Europe-2 region (Supplement, Tables S1 and S2).

For illustration purpose, four areas are examined in greater detail below in Fig. 11. The Manchester-~~Birmingham plot~~ map (Fig. 11 top row) illustrates a large area of high population density in Central England with several power plants to the East. Recall, that the urban component is essentially the population density convoluted with EMG functions and the two large hotspots in the urban component corresponds to Manchester and Liverpool area to the north and Birmingham area at the south. The Our TROPOMI data analysis shows a $-40\% \pm 1.4\%$ decline in the ~~population density related urban~~ component and about $-25.18\% \pm 5.6$ decline from total emission from the power plants. These numbers are close the reported by the UK Office of National Statistics decline in road traffic (-35% and -50% for April and May, respectively) and industrial activity (-20%) (Potts et al., 2021). The estimated decline in the urban component is very close to -42% decline in the surface NO₂ concentrations reported by Lee et al., (2020). In contrast, the background component shows almost no change in 2020 compared to 2018-2019. For this reason, the total decline in TROPOMI NO₂ VCD over Manchester (-27% according Barré et al., 2021, and -32% according to Potts et al., 2021) is smaller than our estimates for the urban component alone.

Paris is an example of a city that appears as a large, isolated urban source. The change between the two periods in the background component is about $-10\% \pm 0.5\%$, while ~~t~~The decline in the ~~population related signal~~ urban component~~t~~ is about $-55.57\% \pm 1.5\%$ and can be clearly seen on the plot. ~~The only relatively large point source in the Paris area is Charles de Gaulle Airport that is evident on the 2018-2019 plot and practically disappeared on the 2020 plot. The terrain does not play a major role in background component of the Paris area. There is some north-south gradient in the background component with higher values at the north-eastern corner of the area.~~ As in the other cases, this value of decline is larger than the decline from TROPOMI NO₂ VCD data without separation of the two component and closer to the changes in NO₂ surface concentrations. Estimated decline in TROPOMI NO₂ VCDs over Paris was about -30% (Bauwens et al., 2020; Barré et al., 2021), while the estimated decline in NO₂ concentrations was -40-50% (Keller et al., 2021; Barré et al., 2021). The terrain does not play a major role in background component for the Paris area. ~~There is some north-south gradient in the background component with higher values at the north-eastern corner of the area. The only relatively large industrial point source in the Paris area is Charles de Gaulle Airport that~~ which is evident on the 2018-2019 plot and practically disappeared on the 2020 plot. Our estimates show about 90% decline in NO₂ emissions from 2018-2019 averages (from about 6.1 ± 2.2 kt y⁻¹ to 0.5 ± 0.2 kt y⁻¹) that is in line with more than -95% decline in the passenger traffic of Charles de Gaulle Airport in April and May 2020 (<https://www.parisaeroport.fr/en/group/finance/investor-relations/traffic> accessed on Nov. 10, 2021).

Milan was one of the first European cities where some lockdown measures were imposed in late February and reduction in NO₂ was reported (Collivignarelli et al., 2020). Complex terrain affects the NO₂ distribution creating large differences between VCD values over the mountains in and valleys and also makes it more difficult to fit the observations with the plume functions based on assumption of straight-line plumes resulting in relatively high residuals. ~~The urban component demonstrates a more than 50% decline in 2020, while the~~ The background component shows practically no difference between the two periods. ~~Complex terrain also makes it more difficult to fit the observations with the plume functions based on~~

~~assumption of straight line plumes resulting in relatively high residuals.~~ The contribution from industrial point sources for that area is ~~very~~ small. The urban component demonstrates a $-53\pm 1.5\%$ decline in 2020 that is similar to other TROPOMI-based estimates for Milan: -38 ± 10 (Bauwens et al., 2020) and about -50 (Barré et al., 2021), while the estimated decline in surface NO_2 concentrations was ~~only just~~ slightly larger, -41% (Keller et al., 2021) and about -52% (Barré et al., 2021). Such small difference in the decline between VCDs and surface concentrations may be due to a relatively small contribution of the background component to the total VCD: it is just about one third of the urban component over Milan

The ~~plots maps~~ for the Prague-Dresden area illustrate how changes in NO_2 from industrial sources reflect differences in COVID-19 lockdown policies in Germany and Czech Republic. The decline in the urban component was only about ~~2016~~ $\pm 3\%$. In addition to Prague, that component also includes cities in East Germany (Dresden, Leipzig), but the changes over these cities and Prague are similar and close to -20% (Barré et al., 2021). Otherwise, the difference would appear in the residuals (Fig. 119, column c). The main industrial sources in the Prague area are coal mines and coal-burning power plants in Czech Republic west of Prague near the German border and in Germany north of Prague, near the Polish border. In Czech Republic, the NO_2 values of the industrial component remain unchanged, while the values over German industrial sources declined by a factor of 2. This is likely the result of different approaches to coal power industry in two countries. In the Czech case, power plants remained fully functioning and certain steps were taken to assure smooth operation and protect the workers: employees of power plants stayed on their job for longer periods, to avoid the risk of infection at home (EC, 2020). In Germany, ~~with its focus on renewable energy sources, a different approach was taken.~~ The power generation from coal-burning plants was reduced by 60% (from 13.4 TWh to 5.6 TWh per month) in April-May 2020 compared to 2019 (<https://www.energy-charts.info/charts/energy/chart.htm?l=en&c=DE&year=2020> accessed on March 1, 2021); ~~while production from renewable energy sources was increased.~~ As a result, we see a large difference in NO_2 ~~signals-VCDs~~ from the power plants in two countries.

4.5-6 The global COVID-19 lockdown impact

To evaluate the COVID-19 lockdown impact worldwide, the analysis described earlier in Section 4 was performed for 261 ~~263~~ urban areas around the world. ~~It should be noted that the NO_2 “footprints” of cities with the same population vary greatly from region to region with the highest values in Northern Eurasia and Australia, and the smallest in Africa and India. To illustrate these large differences, Fig. 12 shows the examples of NO_2 distribution near cities with population of about 5-6 million with very large (Saint Petersburg, Russia) and very small (Dar es Salaam, Tanzania) per capita emissions. The total mass of NO_2 per capita related to the urban component for Saint Petersburg was 40 times larger than for Dar es Salaam (all for 2018-2019).~~ While all All cities with population greater than 1 million were considered ~~in this study.~~ However, some of them, particularly in Africa, do not ~~even~~ produce significant NO_2 emissions that can be measured by TROPOMI over the three-month period selected for this study. Another obstacle is in Western Africa, where biomass burning made it difficult to estimate “background” levels as they were very different from year to year. Biomass burning areas appears as large anomalies on the maps of the residuals making the standard deviations of the residuals much higher than at the other African cities. For this

reason, several areas with population over a million in Western Africa were not included in the analysis. In case of China, there are too many cities with population over one million. We raised the limit for China and considered only cities with population greater than 6 million to keep the number of analysed areas similar to other regions. 2

5 It should be noted that the NO₂ “footprints” of cities with the same population vary greatly from region to region with the highest values in Northern Eurasia and Australia, and the smallest in Africa and India. To illustrate these large differences, Fig. 12 shows the examples of NO₂ distribution near cities with population of about 5-6 million with very large (Saint Petersburg, Russia) and very small (Dar es Salaam, Tanzania) per capita emissions. The total mass of NO₂ per capita related to the urban component for Saint Petersburg was 40 times larger than for Dar es Salaam (all for 2018-2019).

10 The analysis algorithm requires coordinates of individual industrial sources in order to separate them from the urban component. The world power plant database (see section 2.3) was used to locate most of the power plants, while other sources were identified from hotspots on the NO₂ residual maps. ~~However, detailed information about industrial source locations is often not available in many regions. Note that the residuals map can be used for detecting of missing in the inventory emission sources. “Hotspots” on the residual map as typically may correspond to emission sources that are not included in the original fitting.~~ Coordinates of such sources are determined from high-resolution satellite imagery and added to the point source list and then the fitting process is repeated. A total of 357 such additional sources were identified. Most of them were cement and steel factories, and oil refineries. In addition, the world busiest airports were included as “industrial” emission sources. However, other sources, ~~in particular, e.g.~~, ship tracks or major highways may still be missing that may affect estimates for some areas. Some of these sources are identifiable in the residual maps and could be added to the statistical model in the future.

20 The map of mean background and urban components values for all 262261 sites in 2018-2019 is shown in Fig. 123 (top). The map of mean background component values for all 262 sites is shown in Fig. 13 (top). The analysed period from mid-March to mid-June is close to spring in the Northern Hemisphere and autumn in the Southern Hemisphere, i.e. the seasons with very similar values of lifetime (Lange et al., 2021). Therefore, seasonal differences between the two hemispheres should be minimal, and maps of the main estimated components should well represent their global distribution. The highest background values are seen over East China and the northern part of Central Europe, while the lowest are mostly over South America and East Africa. In general, it is similar to the global NO₂ distribution (Krotkov et al., 2016), although the background levels are much lower than those seen over NO₂ hotspots on the mean satellite NO₂ map.

25 ~~The values of the urban component, expressed as annual emission rate per capita, are shown in Fig. 13 (bottom).~~ The urban component demonstrates the highest values are over Siberian region of Russia. They ~~and~~ are likely related to additional NO_x emissions due to heating there since the climatological temperatures there are in a relatively cold period relatively low in March-April compared to other regions. Another hotspot is Edmonton, but as mentioned, its high value is due to poor separation of urban emissions from emissions from two and industrial sources there. Annual emissions per capita are also high over Middle East. However, in this region we found that the population density data in some areas including, for example,

Riyadh, may not be reliable and emissions per capita may be overestimated. The population density maps there do not match Google map satellite images and other proxies such as night light data. This requires further investigation.

The lowest annual emissions per capita are in South America, Africa, and India. Although emissions per capita were calculated for each area independently and the population and industrial sources vary greatly from area to area, the per capita values are uniform: for example, almost all areas in India marked by green dots, most of European areas are orange, etc. This further gives a confidence in the obtained estimates.

Note that the urban NO₂ “footprints” of cities with the same population vary greatly from region to region with the highest values in Northern Eurasia and Australia, and the smallest in Africa and India. To illustrate these large differences, Fig. S2 shows the examples of NO₂ distribution near cities with population of about 5-6 million with very large (Saint Petersburg, Russia) and very small (Dar es Salaam, Tanzania) per capita emissions. The total mass of NO₂ per capita related to the urban component for Saint Petersburg was 40 times larger than for Dar es Salaam.

Fig. 13 shows the maps of percent changes for individual areas for the background and urban components. Relative changes of the background component are typically within $\pm 10\%$ and much smaller than in the urban component. One of the exceptions is the Scandinavian countries and Germany that caused an overall 13% change in the background value for the Europe 2 region of the regions with large negative changes in the background component is Middle East. As mentioned, the population density data are not always accurate in that region and the background component may not be perfectly separated from the urban and industrial components. In contrast, the urban component demonstrates a much larger decline, particularly over Europe-1 and India. Note that the changes in the urban and background components are fairly independent: a analysis of all 262261 areas revealed that there is no correlation between changes in urban NO₂ emissions and changes in background values (the correlation coefficient between them is -0.0079).

The estimates for individual areas were then grouped into 13 large regions with 10-20 areas in each: the U.S. and Canada, Europe-1 and -2, China, India, South-East Asia (also includes also Pakistan and Bangladesh), Japan with Taiwan and South Korea, Northern Eurasia (former USSR countries and Mongolia), Middle East, Africa, Australia and New Zealand, Central America, and South America. The regions are based on geographical location with similarities in economic development and reactions to the COVID-19 pandemic were also taken into account. Then, the average characteristics of the background, urban and industrial components were calculated for each region. Relative contributions of the three components (similar to Fig.7) and the bar charts of the mean values of these components as well as the decline between 2020 and 2018-2019 for all regions are available from the Supplement. Three areas, Johannesburg (South Africa), Nicosia (Cyprus), and Pyongyang (North Korea) were not included in any particular region because their NO₂ emissions were very different from those from neighbouring countries and therefore may bias regional statistics.

_____ $\bar{}$
_____ The summary results for the regions are shown in Fig. 144, while the results for individual areas within each region as well as the obtained emissions values are available from the Supplement. The regions in Fig. 144 are sorted by relative

decline in the urban component (from smallest to largest). The regional changes were calculated as the average of percent changes for individual areas for that region ~~and areas with components below some threshold levels were excluded.~~ - ~~The uncertainty values in Fig. 14 are based on variation of the values for individual areas within the region. The background component has the smallest variability among the three components typically between 5% and 9%. The urban component variability is between 7% and 17% and the decline observed in the urban component for South America, Europe-1 and India is outside 3- σ limits even for individual areas in these regions. The industrial component was added to separate emission from large industrial sources in the urban areas from urban emissions themselves. Emissions from such industrial sources are typically similar or smaller than urban emissions and the variability of the industrial component (10%-30%) is similar or larger than that for the urban component.~~

~~Fig. 15 shows the maps of changes for individual areas for the background and urban components.~~

~~Fig. 15 shows the maps of changes for individual areas for the background and urban components. China shows the smallest and not significant decline in the urban component, while the majority of the regions demonstrated statistically significant urban emissions decline within the range -18 -28%. The decline was the largest, -36 -52%, in three regions: Europe 1, South America and India. The map of the urban emission changes (Fig.15, bottom) shows that the first two regions indeed contain countries with large decline of urban emissions. In case of India, a similar decline can be seen in neighbouring Pakistan and Bangladesh. In Africa, a decline is seen at the south and the north of the continent, while countries in West Africa mostly show no decline and even some increase probably due to a contribution from forest fires. China shows the smallest and not significant decline in the urban component over the analysed period as the main COVID-19 lockdown in China occurred earlier (in February). Most of the regions demonstrated statistically significant urban emissions decline within the range -18 to -28%. The decline was the largest, -36 to -52%, in three regions: Europe-1, South America and India. The map of the urban emission changes (Fig.13, bottom) shows that the first two regions indeed contain countries with large decline of urban emissions. In case of India, a similar decline can be seen in neighbouring Pakistan and Bangladesh. In Africa, a decline is seen at the south and the north of the continent, while countries in West Africa mostly show no decline and even some increase probably due to a contribution from forest fires.~~

~~Changes in the background component are shown in Fig. 14 and Fig. 15 (top). Relative changes of the background component are typically within $\pm 10\%$ and much smaller than in the urban component. One of the exceptions is the Scandinavian countries and Germany that caused an overall -13% change in the background value for the Europe 2 region. The urban and background components are fairly independent. Analysis of all 262 areas revealed that there is no correlation between changes in urban NO_2 emissions and changes in background values (the correlation coefficient is -0.009).~~

As mentioned in section 4.3, the industrial NO_2 component varies from area to area and from one type of NO_2 source to another, although there are some clear regional differences. Chinese cities demonstrated small changes in both urban and industrial components (-2.8% and +5% respectively) with one exception. Emissions from Wuhan, the city where the pandemic began, declined by more than -60%. Industrial emissions there also declined, but only by -30%. The background component

shows no change there. A very strict Wuhan lockdown ended on April 8, 2020, but during that lockdown, NO₂ emissions in Wuhan declined -82% relative to the 2019 level (Ghahremanloo et al., 2021). That strict lockdown period lasted for less than one third of the analysed period, but apparently, it took some time for NO₂ emissions to return to the pre-lockdown levels.

It is more difficult to interpret changes in industrial source emissions because they are changing over time for various reasons that may require an investigation on a case-by-case basis. For example, a large uncertainties in the industrial emission changes for Central America in Fig. 14 are caused by doubling of emissions from power plants near Havana. This increase is likely caused by emissions from three powerships (power plants on ships) with total capacity of 184 MW, that started their operation in Port de Mariel near Havana in the second half of 2019 (<https://karpowership.com/en/project-cuba>, accessed Nov. 4, 2021). The largest regional industrial emissions decline was observed over Europe-1 and India, i.e., where largest urban emissions decline was observed. It is likely the severe restrictions during the COVID-19 lockdown period there affected the industrial activity. However, on a larger scale, this link is not that obvious. Although the lockdown had impact on industrial sources, the correlation coefficient between changes in urban and industrial emissions among all analysed areas is only 0.180.01.

As mentioned, statistical errors related to the fitting procedure are relatively small due to a very large number of satellite pixels used in the fit. For the urban component, they are between 1% and 10% for the cities analysed in this study. However, the year-to-year variability could be high. Table 2 summarizes the uncertainty for the 13 regions analysed in this study. It is similar to Table 1; however individual rows contain the averages of uncertainty estimates for all individual areas in the region. The uncertainties for the background component are between 4.9 and 9%. The urban component demonstrates the interannual variability between 9% and 22% with the largest value over the Middle East were, as mentioned, there could be a problem with the population density data quality. It is far more difficult to interpret the estimated interannual variability for the industrial sources because it depends on multiple factors from the meteorological conditions to the emission strength itself. The main conclusion here is that it is typically -10...% to -20% for emissions from large (about 5 kt y⁻¹ or more) sources estimated based on 3 months of data.

The uncertainties values in Fig. 14 for the regions are based on variation of the values for individual areas within the region. The uncertainties for individual area emissions can be estimated from the differences between 2018 and 2019 values as discussed in Supplement A (see also Table S1). The background component has the smallest variability among the three components typically between 5% and 9%. The urban component variability is between 7% and 17% and the decline observed in the urban component for South America, Europe 1 and India is outside 3-σ limits even for individual areas in these regions. The industrial component was largely added to separate emission from large industrial sources in the urban areas from urban emissions themselves. Emission from such industrial sources are typically similar or smaller than urban emissions and the variability of the industrial component (10%–30%) is similar or larger than that for the urban component.

To demonstrate that the observed NO₂ changes in urban emissions are indeed linked to the restricting measures taken by different countries, the estimated percent NO₂ changes in annual emissions per capita were compared to the Google Each Community Mobility Report data (available from <https://www.google.com/covid19/mobility/>, accessed on March 1, 2021).

The mobility data represent the changes in the number of people at locations of various type and can be used as a proxy for the urban traffic. The changes in the background and urban components ~~were calculated for every country were-and~~ compared ~~to~~ changes in mobility data. ~~The mean values of the three components of the NO₂ distribution (background, urban, and industrial) were calculated for every country by averaging the corresponding values for individual cities.~~ Only countries with two or more cities were used in the comparison. Note that the mobility data were averages of all regions for the entire country, while the NO₂ changes were estimated for areas around large cities only. Mobility data for China, Korea and some other countries were not available.

The scatter plot of the mobility and the NO₂ VCD changes (Fig. 156) demonstrates a very different relationship between the urban and background components. Changes in mobility and urban components are correlated (Fig. 156 left). As expected, the relative changes in the urban component are smaller than the mobility changes as the urban component includes more than just mobility-related traffic. The highest correlation is observed when changes in the NO₂ urban component are compared with mobility for “retail and recreation”, covering visits to restaurants, cafes, shopping centers, theme parks, museums, libraries, movie theatres, and similar locations. The correlation coefficient between the percent changes in per capita emissions and “retail and recreation” mobility is 0.66-62 (the probability that there is no correlation is less than 0.00040003). There is no statistically significant correlation (the correlation coefficient is -0.083) between the background NO₂ and mobility data (Fig. 156 right). ~~The industrial component shows some correlation (the correlation coefficient is 0.5) with the mobility data, although not as high as for the urban component.~~

For individual areas, the uncertainties due to the interannual variability rather large, so the observed 2020 decline of the urban component in many areas is within that uncertainty. Regional averages are more accurate and declines in urban emissions are statistically significant for all regions except China. Finally, mean 2020 declines of all areas (except China) are -6.0%±1.2% and -26.7%±2.6% for the background and urban components respectively that corresponds to 10-σ and 20-σ levels (Supplement, Fig. S3). In 2018, 2019, and 2021, such global averages are remarkable stable, as all mean values are within ±2.5%, and within ±5% if the means are taken with the two-sigma uncertainties. This suggests that there are no substantial drifts or shifts on the global scale in TROPOM data and the 2020 data is clearly an outlier.

~~To compare our results with other similar estimates of NO_x emission, we looked at a recent study by Lange et al., 2021. In their study, TROPOMI NO₂ data were used to estimate emissions from 45 sources worldwide and compared them with the available emissions inventories and some other satellite based emission estimates. As our study also provides total emissions estimates (the sum of urban and industrial emissions) for 33 of them, we compare our results with those from Lange et al., 2021. While the emission estimation algorithms and approaches are different in the two studies, the results show a good agreement and the correlation coefficient between the two sets of emission estimates is 0.78. As expected, the emission estimated in this study are higher than from Lange et al., 2021. This is because we used larger areas and, typically, there was more than one emission source in the analysed 3° by 4° areas of this study. The details of this comparison are available from Supplement C.~~

~~————— In this study, the population density was used as a proxy for urban emissions. We found that this proxy works well unless the population density was very high and the link between population density and emissions might not be linear anymore. Such exceptions were rare (e.g., some regions of New York City). To test the impact of such nonlinearity, we used a different proxy, the night light data set and all calculations were repeated with them. Unlike the population density, the used night light data reached saturation at highly populated areas and did not grow any further. It was found that the results were similar and the major findings about different reaction of the background and urban components to the COVID-19 lockdown were still valid, although with some differences in absolute values (see section Supplement E). The correlation coefficient between mobility and urban emissions estimated from night light is lower (0.57) than for those from population density suggesting that the population density based analysis captures the NO₂ emissions reduction due to lower traffic better.~~

10 5 Discussion and conclusion

Statistical regression analysis was used to separate contribution from industrial sources, urban areas, and background levels to the satellite observed tropospheric NO₂ columns (VCDs) and to study the impact of the COVID-19 lockdown on each component separately. The analysis was done for 262261 major urban areas around the world grouped into 13 large regions. The algorithm also estimates urban and industrial emissions assuming a constant NO₂ lifetime (or, more accurately, decay time). A constant value of 3.3 hours was used as the lifetime.

To verify the obtained emissions estimates, we compared our result with the estimates from a similar study by Lange et al., (2021). In that study, TROPOMI NO₂ data were used to estimate emissions from 45 sources worldwide and then the results were compared with the available emissions inventories and some other satellite-based emission estimates. There were 33 sources common to both works. In order to compare them we first calculated our total emissions, the of the two studies for them i.e., the sum of urban and industrial emissions, and then converted them to the same lifetimes as in Lange et al., (2021), and then multiplied them by 1.4 to calculate NO_x emission. As expected, our emission estimates were higher than from Lange et al., (2021) because there is typically more than one emission source in the analysed 3° by 4° areas of this study. Nevertheless, there is 0.78 correlation coefficient between the two estimates.

Unlike other similar studies that simply removed the background offset (e.g., ~~(~~Beirle et al., (2011); Lange et al., (2021)), this study included the background component in the analysis. On a scale of several hundred km (as we analyzed 3° by 4° areas), most of the NO₂ mass is typically related to the background component. Even in the areas such as New York City, the background component accounts for 2/3 of the total mass. This explains why the estimated impact of the COVID-19 lockdown in urban areas depends on the size of the analysed area: the bigger larger the size-area the more background NO₂ it includes and, therefore, the smaller ~~is~~ the NO₂ difference between the COVID-19 lockdown and reference periods.

In most of the analysed areas, changes in the background components between the COVID-19 lockdown period analysed here (from March 16 to June 15, 2020) were typically within 10% from the 2018-2019 levels. In contrast, the urban component, based on population density, demonstrated a substantial and rather uniform decline of about -18% ~~...to~~ -28% in

most of the regions. ~~Three-Two~~ regions (most western part of Europe ~~and~~; India ~~and South America~~) demonstrated a larger decline, about -40% ~~to~~ -50%. China showed a much smaller decline ($-4.43\% \pm 98\%$) because the lockdown there occurred prior to the analysed period. As for industrial point sources, emissions from them varied from region to region and from sector to sector. ~~On average,~~ They demonstrate a ~~decline of about a about -20% or less decline~~ except for India and Europe-1 regions.

5 Abrupt changes ~~and in~~ urban and industrial emissions due to COVID-19 lockdown did not immediately result in a similar decline in the background component. This may explain why large changes in NO₂ emissions in urban areas produced a relatively small, about 9% decline in global NO₂ (Bray et al., 2021). The importance of background NO₂ VCD was previously noted by Qu et al., (2021) and Silvern et al., (2019) when they found that the observed satellite tropospheric NO₂ VCD satellite trends in remote areas do not match the expected changes. The origins of background NO₂ are still largely related to urban and
10 industrial sources as it is clearly higher in the northern hemisphere, particularly over China, Central Europe and Eastern U.S., than in the southern hemisphere and tropics. However, the analysed three-month period may simply not be ~~not~~ long enough for the lockdown to cause large changes in the background levels. There are also other NO_x sources such as soil emissions (Hudman et al., 2012; Sha et al., 2021). They as well as sources aloft, such as lightning, and to a lesser extent, aircraft NO_x directly contribute to the background component. It is estimated that lightning is responsible for roughly 16% of global
15 production and most of this NO_x is found in the free troposphere (Bucselá et al., 2019). Furthermore, Zhang et al., (2012) estimated that sources such as lightning, soils, and wildfires that account for about 20% of emissions annually and up to 39% in summer. Satellite measurements are also more sensitive to NO₂ in the free troposphere than in the boundary layer and a relatively small amounts of NO₂ there produce a larger signal in satellite data. Another possible explanation is that at low NO₂ concentration in the boundary layer and free troposphere may have longer lifetimes than in the plumes. The fact that NO₂
20 fluctuations remain persistent over longer time in clean conditions than over polluted areas (Vinnikov et al., 2017) indirectly confirms that.

Barré et al., (2021), noticed a different lockdown-related decline between NO₂ VCDs and surface concentrations (-23% and -43% over Europe respectively). Moreover, Qu et al., (2021) reported that VCDs and surface concentrations had a similar decline between 2019 and 2020 at only 5% of the most pollutes sites. At the other sites, TROPOMI NO₂ VCDs data
25 demonstrated, on average, a smaller decline than surface concentrations. Different changes in background and urban components in TROPOMI NO₂ could explain this inconsistency between the surface and satellite VCD-based results. The urban component is directly linked to city plumes and therefore is a better proxy for surface concentrations in polluted areas, while the background component includes contribution from other sources that were not affected by the lockdown.

The urban and industrial components are based on plume dispersion functions ~~and that~~ correspond to NO₂ near the
30 ground, ~~probably-almost always~~ in the boundary layer. The urban component is based on the population density and the assumption that annual emissions per capita are uniform ~~everywhere~~ in the analysed 3° by 4° area. There are very large differences, up to factor of 40, in estimated emissions per capita ~~between-among~~ the different areas. The estimates were done for 3-month periods. For such a short time interval, most of the cities with population more than 1 million produce a statistically significant signal, that can be readily detected in TROPOMI NO₂ data. As estimated emissions per capita are rather uniform,

they can be used to account for urban component outside large cities. Thus, it should be possible to estimate background, urban, and industrial components on the global scale and analyse the residuals in search of other factors contributing to the NO₂ budget.

5 The approach described in this study can be used to estimate emissions from cities and industrial point sources. For the latter, only source coordinates are required. A comparison of reported and ~~TROPOMI-derived from TROPOMI data~~ NO_{x2} emissions for the U.S. demonstrated a good correlation (~~0.9~~) between them. As source coordinates can be also detected from satellite data alone (Beirle et al., 2019; Ding et al., 2020; McLinden et al., 2016), it may be possible to develop an independent “top-down” NO_{x2} emission inventory from satellite measurements to complement and ~~emissions~~ improve available “bottom-up” inventories as it was done for SO₂ (Liu et al., 2018). This could be important for regions, where no other emission
10 information is available.

Data availability

The TROPOMI NO₂ product is publicly available on the Copernicus Sentinel-5P data hub (<https://s5phub.copernicus.eu>). ~~The reprocessed (RPRO) and offline mode (OFFL) data of version V1.2.2 to version V1.3.2 were used. The Google Each Community Mobility Report data are available from https://www.google.com/covid19/mobility/. The Gridded Population of the World (GPW) dataset is available from NASA Socioeconomic Data and Applications Center at https://sedac.ciesin.columbia.edu/data/collection/gpw-v4. The European Centre for Medium-Range Weather Forecasts (ECMWF) ERA5 reanalysis data are available from https://www.ecmwf.int/en/forecasts/datasets/reanalysis-datasets/era5. Elevation data are from gridded global relief ETOPO2v2 database (https://www.ngdc.noaa.gov/mgg/global/etopo2.html).~~
15

20 Author contributions

VF analysed the data and prepared the paper with input from CM and critical feedback from all the co-authors. CM and DG generated the TROPOMI data subsets for the analysis. NK and FL contributed to the interpretation of the results. HE provided the TROPOMI NO₂ data product and related information. All authors read and agreed on the published version of the paper.

Competing interests

25 The authors declare that they have no conflict of interest.

Acknowledgments. Sentinel 5 Precursor TROPOMI Level 2 product is developed with funding from the Netherlands Space Office (NSO) and processed with funding from the European Space Agency (ESA).

Appendix

This appendix contains additional details of the used fitting algorithm that is largely based on the algorithm for multiple point source emission estimates (Fioletov et al., 2017). TROPOMI NO₂ VCD can be expressed as a sum of contributions $\alpha_i \cdot \Omega_i$ from all individual industrial sources (i), a population density-related term $\alpha_p \Omega_p$, an elevation-related background, and noise (ε):

$$5 \quad TROPOMI \text{ NO}_2(\theta, \varphi) = \alpha_0 + \alpha_p \Omega_p(\theta, \varphi) + \sum \alpha_i \Omega_i(\theta, \varphi) + (\beta_0 + \beta_1(\theta - \theta_0) + \beta_2(\varphi - \varphi_0)) \cdot \exp(-H(\theta, \varphi)/H_0) + \varepsilon(\theta, \varphi) \quad (A1)$$

All Ω function are normalized (i.e., their total integral equals to 1) plume functions: the value of that function for a particular pixel with latitude θ and longitude φ , is proportional to the value of the plume parameterization from the source i located at the latitude θ_i and longitude (φ_i) (all in radian). The parameterization assumes that the plume is moving downwind along a straight line has a Gaussian shape spread across that line. To describe the plume, we can rotate satellite pixels for a particular
 10 day around the source, so the plume would always be moving from north to south, apply the plume parameterization, and then rotate the pixels back. If (x_i, y_i) and (x'_i, y'_i) are the pixel's Cartesian coordinates (km) in the system with the origin at the source i before and after the rotation respectively, then they can be calculated from the pixel and source latitudes and longitudes as:

$$15 \quad \begin{aligned} x_i &= r \cdot (\varphi - \varphi_i) \cdot \cos(\theta_i); \\ y_i &= r \cdot (\theta - \theta_i); \\ x'_i &= x_i \cdot \cos(-\omega) + y_i \cdot \sin(-\omega); \\ y'_i &= -x_i \cdot \sin(-\omega) + y_i \cdot \cos(-\omega); \end{aligned}$$

where $r=111.3 \text{ km} \cdot 180 / \pi$ (or $r=6371 \text{ km} \cdot \pi / 180$ for latitude and longitude in degrees); ω is the pixel wind direction (0 for
 20 north); φ_i and θ_i are the source i longitude and latitude (all in radian). Note that there was a typo in this original formula for r in Fioletov et al., (2017).

Following Fioletov et al., (2017), the contribution $\alpha_i \cdot \Omega_i = \alpha_i \Omega(\theta, \varphi, \omega, s, \theta_i, \varphi_i)$ from the source i can be expressed as $\alpha_i \cdot \Omega_i = \alpha_i \cdot f(x'_i, y'_i) \cdot g(y'_i, s)$, where:

$$25 \quad \begin{aligned} f(x'_i, y'_i) &= \frac{1}{\sigma_1 \sqrt{2\pi}} \exp\left(-\frac{x_i'^2}{2\sigma_1^2}\right); \\ g(y'_i, s) &= \frac{\lambda_1}{2} \exp\left(\frac{\lambda_1(\lambda_1 \sigma^2 + 2y'_i)}{2}\right) \cdot \operatorname{erfc}\left(\frac{\lambda_1 \sigma^2 + y'_i}{\sqrt{2}\sigma}\right); \\ \sigma_1 &= \begin{cases} \sqrt{\sigma^2 - 1.5y'_i}, & y'_i < 0; \\ \sigma, & y'_i \geq 0; \end{cases} \\ \lambda_1 &= \lambda/s; \end{aligned} \quad (A2)$$

It is assumed that NO₂ emitted from a point source decline exponentially (i.e., as $\exp(-\lambda t)$) with time (t) with a constant “lifetime” (or decay rate) $\tau=1/\lambda$. The second parameter is the plume width (σ).

Note that $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) \cdot g(y, s) dx dy = \int_{-\infty}^{\infty} (\int_{-\infty}^{\infty} f(x, y) dx) \cdot g(y, s) dy = \int_{-\infty}^{\infty} g(y, s) dy = 1$, therefore the parameter α_i represents the total observed number of NO₂ molecules (or the NO₂ mass) near the source i . If *TROPOMI NO₂* is in DU, and σ is in km, then a is in $2.69 \cdot 10^{26}$ molec. or 0.021 T(NO₂). Furthermore, the emission strength (E) can be calculated as $E = a/\tau$ assuming a simple mass balance.

5 As mentioned in section 3, some of the sources used in the analysis are not point sources but clusters. In that case, $\Omega_i = \sum_j w_j \Omega_j(\theta, \phi, \omega, s, \theta_j, \phi_j)$, where Ω_j is the plume function for source j and w_j is the weighting coefficient established by the factor analysis.

Similarly, $\alpha_p \Omega_p$ represent the contribution from the population density-related component, where Ω_p is the plume function from an area-distributed source. Ω_p is a weighted sum of plume functions from a grid with the weighting coefficients proportional to the population of at the grid points $\Omega_p = \sum_{ij} \rho_{ij} \Omega(\theta, \phi, \omega, s, \theta_{ij}, \phi_{ij})$, where θ_{ij}, ϕ_{ij} are the grid points
 10 coordinates and ρ_{ij} is the population associated with that grid points. Thus, α_p is the coefficient that represent the total NO₂ mass the corresponds to one person. In our calculations we used 3° by 4° area with a 0.2° by 0.2° grid with 336 (16 x 21) grid cell.

Finally, the elevation-related background term $\alpha_0 + (\beta_0 + \beta_1(\theta - \theta_0) + \beta_2(\varphi - \varphi_0)) \cdot \exp(-H/H_0)$, where θ_0 and
 15 φ_0 are the coordinates of the centre of the analyzed area and E is the elevation in km and $H_0=1$ km, is determined by three parameters.

Equation (A1) represents a linear regression model where the unknown parameters α_p, α_i can be estimated from the measured variable (*TROPOMI NO₂*) at many pixels and known regressors. The fitting was done three times using all data for the analysed period (March 16 – June 15) in 2018, 2019, and 2020.

References

- 5 [Ass, KE, Eddaif, A, Radey, O, Aitzaouit, O, Yakoubi, ME, Chelhaoui, Y.: Effect of restricted emissions during Covid-19 lockdown on air quality in Ra bat—Morocco. Global NEST Journal 22.DOI: <http://dx.doi.org/10.30955/gnj.003431>, 2020.](https://doi.org/10.30955/gnj.003431)
- [Aydın, S, Nakiyingi, BA, Esmen, C, Guñeysu, S, Ejjada, M.: Environmental impact of coronavirus \(COVID-19\) from Turkish perceptive. Environment, Development and Sustainability. DOI: <http://dx.doi.org/10.1007/s10668-020-00933-5>, 2020.](https://doi.org/10.1007/s10668-020-00933-5)
- 10 Bao, R. and Zhang, A.: Does lockdown reduce air pollution? Evidence from 44 cities in northern China, *Sci. Total Environ.*, 731, doi:10.1016/j.scitotenv.2020.139052, 2020.
- [Bar, S., Parida, B.R., Mandal, S.P., Pandey, A.C., Kumar, N., Mishra, B.: Impacts of partial to complete COVID-19 lockdown on NO₂ and PM_{2.5} levels in major urban cities of Europe and USA, *Cities*, Volume 117, 103308, ISSN 0264-2751, <https://doi.org/10.1016/j.cities.2021.103308>, 2021.](https://doi.org/10.1016/j.cities.2021.103308)
- 15 [Barré, J., Petetin, H., Colette, A., Guevara, M., Peuch, V.-H., Rouil, L., Engelen, R., Inness, A., Flemming, J., Pérez García-Pando, C., Bowdalo, D., Meleux, F., Geels, C., Christensen, J. H., Gauss, M., Benedictow, A., Tsyro, S., Friese, E., Struzewska, J., Kaminski, J. W., Douros, J., Timmermans, R., Robertson, L., Adani, M., Jorba, O., Joly, M., and Kouznetsov, R.: Estimating lockdown-induced European NO₂ changes using satellite and surface observations and air quality models. *Atmos. Chem. Phys.*, 21, 7373–7394, <https://doi.org/10.5194/acp-21-7373-2021>, 2021.](https://doi.org/10.5194/acp-21-7373-2021)
- 20 Bauwens, M., Compornolle, S., Stavrakou, T., Müller, J. F., van Gent, J., Eskes, H., Levelt, P. F., van der A, R., Veeffkind, J. P., Vlietinck, J., Yu, H. and Zehner, C.: Impact of Coronavirus Outbreak on NO₂ Pollution Assessed Using TROPOMI and OMI Observations, *Geophys. Res. Lett.*, 47(11), 1–9, doi:10.1029/2020GL087978, 2020.
- Beirle, S., Platt, U., von Glasow, R., Wenig, M. and Wagner, T.: Estimate of nitrogen oxide emissions from shipping by satellite remote sensing, *Geophys. Res. Lett.*, 31(18), L18102, doi:10.1029/2004GL020312, 2004.
- 25 Beirle, S., Boersma, K. F., Platt, U., Lawrence, M. G. and Wagner, T.: Megacity emissions and lifetimes of nitrogen oxides probed from space, *Science* (80-.), 333(6050), 1737–1739, doi:10.1126/science.1207824, 2011.
- Beirle, S., Hörmann, C., Penning de Vries, M., Dörner, S., Kern, C. and Wagner, T.: Estimating the volcanic emission rate and atmospheric lifetime of SO₂ from space: a case study for Kīlauea volcano, Hawai`i, *Atmos. Chem. Phys.*, 14(16), 8309–8322, doi:10.5194/acp-14-8309-2014, 2014.
- 30 Beirle, S., Borger, C., Dörner, S., Li, A., Hu, Z., Liu, F., Wang, Y. and Wagner, T.: Pinpointing nitrogen oxide emissions from space, *Sci. Adv.*, 5(11), doi:10.1126/sciadv.aax9800, 2019.
- [Beirle, S., Borger, C., Dörner, S., Eskes, H., Kumar, V., de Laat, A., and Wagner, T.: Catalog of NO_x emissions from point sources as derived from the divergence of the NO₂ flux for TROPOMI, *Earth Syst. Sci. Data*, 13, 2995–3012, <https://doi.org/10.5194/essd-13-2995-2021>, 2021.](https://doi.org/10.5194/essd-13-2995-2021)

- Belhekar, V. M.: Factor Analysis and Structural Equation Modeling, in A Step-by-Step Approach to Using SAS® for Factor Analysis and Structural Equation Modeling, Second Edition, edited by N. O'Rourke and L. Hatcher, pp. 314–361, SAS Institute., 2013.
- Bray, C. D., Nahas, A., Battye, W. H. and Aneja, V. P.: Impact of lockdown during the COVID-19 outbreak on multi-scale air quality, *Atmos. Environ.*, 254(August 2020), 118386, doi:10.1016/j.atmosenv.2021.118386, 2021.
- 5 Bucsela, E. J., Pickering, K. E., Allen, D. J., Holzworth, R. H. and Krotkov, N. A.: Midlatitude Lightning NO_x Production Efficiency Inferred From OMI and WLLN Data, *J. Geophys. Res. Atmos.*, 124(23), 13475–13497, doi:10.1029/2019JD030561, 2019.
- C3S: Copernicus Climate Change Service (C3S) : ERA5: Fifth generation of ECMWF atmospheric reanalyses of the global climate. Copernicus Climate Change Service Climate Data Store (CDS), [online] Available from: <https://cds.climate.copernicus.eu/cdsapp#!/home> (Accessed 20 June 2020), 2017.
- 10 Chong, H., Lee, H., Koo, J.H., Kim, J., Jeong, U., Kim, W., Kim, S.W., Herman, J.R., Abuhassan, N.K., Ahn, J.Y., Park, J.H., Kim, S.K., Moon, K.J., Choi, W.J. and Park, S.S.: Regional Characteristics of NO₂ Column Densities from Pandora Observations during the MAPS-Seoul Campaign. *Aerosol Air Qual. Res.* 18: 2207-2219. <https://doi.org/10.4209/aaqr.2017.09.0341>, 2018.
- 15 Collivignarelli, M. C., Abbà, A., Bertanza, G., Pedraz-zani, R., Ricciardi, P., and Carnevale Miino, M.: Lockdown for CoViD-2019 in Milan: What are the effects on air quality?. *Sci Total Environ.* 732, 139280–139280. <https://doi.org/10.1016/j.scitotenv.2020.139280>, 2020.
- Cunnold, D. M., Zawodny, J. M., Chu, W. P., Pommereau, J. P., Goutail, F., Lenoble, J., McCormick, M. P., Veiga, R. E., 20 Murcray, D., Iwagami, N., Shibasaki, K., Simon, P. C. and Peetermans, W.: Validation of SAGE II NO₂ measurements, *J. Geophys. Res.*, 96(D7), 12913, doi:10.1029/91JD01344, 1991.
- Dammers, E., McLinden, C. A. , Griffin, D., Shephard, M. W., ~~v~~Van Der Graaf, S., Lutsch, E., Schaap, M., Gainairu-Matz, Y., Fioletov, V., ~~v~~Van Damme, M., Whitburn, S., Clarisse, L., Cady-Pereira, K., Clerbaux, C., Francois Coheur, P. and Erismann, J.-W. J. W.: NH₃ emissions from large point sources derived from CrIS and IASI satellite observations, 25 *Atmos. Chem. Phys.*, 19(19), 12261–12293, doi:10.5194/acp-19-12261-2019, 2019.
- Dantas, G, Siciliano, B, França, BB, da Silva, CM, Ar-billa, G. The impact of COVID-19 partial lock- down on the air quality of the city of Rio de Janeiro, Brazil. *Science of the Total Environment* 729. DOI: <http://dx.doi.org/10.1016/j.scitotenv.2020.139085>. 2020.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., 30 Bauer, P., Bechtold, P., Beljaars, A. C. M., Berg, L. van de, Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., Rosnay, P. de, Tavolato, C., Thépaut, J.-N. and Vitart, F.: The ERA-Interim reanalysis: Configuration and performance of the data assimilation system, *Q. J. R. Meteorol. Soc.*, 137, 553–597, doi:DOI: 10.1002/qj.828, 2011.

- Ding, J., van der A, R. J., Eskes, H. J., Mijling, B., Stavrou, T., van Geffen, J. H. G. M. and Veefkind, J. P.: NO_x Emissions Reduction and Rebound in China Due to the COVID-19 Crisis, *Geophys. Res. Lett.*, 47(19), e2020GL089912, doi:10.1029/2020GL089912, 2020.
- 5 Duncan, B. N., Lamsal, L. N., Thompson, A. M., Yoshida, Y., Lu, Z., Streets, D. G., Hurwitz, M. M. and Pickering, K. E.: A space-based, high-resolution view of notable changes in urban NO_x pollution around the world (2005–2014), *J. Geophys. Res. Atmos.*, 121(X), 976–996, doi:10.1002/2015JD024121. Received, 2015.
- EC: European Commission: COVID-19 in European coal regions. [online] Available from: https://ec.europa.eu/energy/sites/ener/files/documents/covid-19_in_european_coal_regions.pdf (Accessed 1 March 2021), 2020.
- 10 EPA: U.S. Environmental Protection Agency (EPA) National Emissions Inventory, [online] Available from: <https://www.epa.gov/air-emissions-inventories> (Accessed 17 July 2020), 2020.
- [ESA EOP-GMQ: Sentinel-5 Precursor Calibration and Validation Plan for the Operational Phase, available at:https://sentinel.esa.int/documents/247904/2474724/ Sentinel-5P-Calibration-and-Validation-Plan.pdf, \(last access: 29 August 2021\), 2017.](https://sentinel.esa.int/documents/247904/2474724/Sentinel-5P-Calibration-and-Validation-Plan.pdf)
- 15 Fioletov, V., McLinden, C. A., Kharol, S. K., Krotkov, N. A., Li, C., Joiner, J., Moran, M. D., Vet, R., Visschedijk, A. J. H. and Denier Van Der Gon, H. A. C.: Multi-source SO₂ emission retrievals and consistency of satellite and surface measurements with reported emissions, *Atmos. Chem. Phys.*, 17(20), doi:10.5194/acp-17-12597-2017, 2017.
- Fioletov, V. E., McLinden, C. A., Krotkov, N. and Li, C.: Lifetimes and emissions of SO₂ from point sources estimated from OMI, *Geophys. Res. Lett.*, 42(6), doi:10.1002/2015GL063148, 2015.
- 20 Fioletov, V. E., McLinden, C. A., Krotkov, N. A., Li, C., Joiner, J., Theys, N., Carn, S. and Moran, M. D.: A global catalogue of large SO₂ sources and emissions derived from the Ozone Monitoring Instrument, *Atmos. Chem. Phys.*, 16(18), 11497–11519, doi:doi:10.5194/acp-16-11497-2016, 2016.
- de Foy, B., Wilkins, J. L., Lu, Z., Streets, D. G. and Duncan, B. N.: Model evaluation of methods for estimating surface emissions and chemical lifetimes from satellite data, *Atmos. Environ.*, 98, 66–77, doi:10.1016/j.atmosenv.2014.08.051, 2014.
- 25 de Foy, B., Lu, Z., Streets, D. G., Lamsal, L. N. and Duncan, B. N.: Estimates of power plant NO_x emissions and lifetimes from OMI NO₂ satellite retrievals, *Atmos. Environ.*, 116, 1–11, doi:10.1016/j.atmosenv.2015.05.056, 2015.
- van Geffen, J. H. G. M., Eskes, H. J., Boersma, K. F., Maasakkers, J. D. and Veefkind, J. P.: TROPOMI ATBD of the total and tropospheric NO₂ data products., [online] Available from: http://www.tropomi.eu/sites/default/files/files/publicS5P-KNMI-L2-0005-RP-ATBD_NO2_data_products-20190206_v140.pdf (Accessed 17 June 2020), 2019.
- 30 ~~v~~an Geffen, J., Eskes, H., Boersma, K., Maasakkers, J. and Veefkind, J.: TROPOMI ATBD of the total and tropospheric NO₂ data products(issue 1.2. 0), De Bilt, the Netherlands, s5P-KNMI-L2-0005-RP., 2018.
- ~~v~~an Geffen, J., Folkert Boersma, K., Eskes, H., Sneep, M., Ter Linden, M., Zara, M. and Pepijn Veefkind, J.: S5P TROPOMI

NO₂ slant column retrieval: Method, stability, uncertainties and comparisons with OMI, *Atmos. Meas. Tech.*, 13(3), 1315–1335, doi:10.5194/amt-13-1315-2020, 2020.

Gkatzelis, G. I., Gilman, J. B., Brown, S. S., Eskes, H., Gomes, A. R., Lange, A. C., McDonald, B. C., Peischl, J., Petzold, A., Thompson, C. R., and Kiendler-Scharr, A.: The Global Impacts of COVID-19 Lockdowns on Urban Air Quality: A Critical Review and Recommendations, *Elem. Sci. Anth.*, 9, doi:10.1525/elementa.2021.00176, 2021.

Fu, F., Purvis-Roberts, K.L., Williams, B. Impact of the COVID-19 pandemic lockdown on air pollution in 20 major cities around the world. *Atmosphere* 11(11). DOI: <http://dx.doi.org/10.3390/atmos11111189>, 2020.

Georgoulias, A. K., Boersma, K. F., Vliet, J. van, Zhang, X., A1, R. van der, Zanis, P. and Laatz, J. de: Detection of NO₂ pollution plumes from individual ships with the TROPOMI/S5P satellite sensor, *Environ. Res. Lett.*, 10(2), 0–6, 2020.

10 Ghahremanloo, M., Lops, Y., Choi, Y. and Mousavinezhad, S.: Impact of the COVID-19 outbreak on air pollution levels in East Asia, *Sci. Total Environ.*, 754, doi:10.1016/j.scitotenv.2020.142226, 2021.

Gkatzelis, G. I., Gilman, J. B., Brown, S. S., Eskes, H., Gomes, A. R., Lange, A. C., McDonald, B. C., Peischl, J., Petzold, A., Thompson, C. R. and Kiendler-Scharr, A.: The global impacts of COVID-19 lockdowns on urban air pollution: A critical review and recommendations, *Elementa*, 9(1), doi:10.1525/elementa.2021.00176, 2021.

15 Goldberg, D. L., Anenberg, S. C., Griffin, D., McLinden, C. A., Lu, Z. and Streets, D. G.: Disentangling the impact of the COVID-19 lockdowns on urban NO₂ from natural variability, *Geophys. Res. Lett.*, doi:10.1029/2020GL089269, 2020.

Griffin, D., McLinden, C. A., Racine, J., Moran, M. D., Fioletov, V., Pavlovic, R., Mashayekhi, R., Zhao, X. and Eskes, H.: Assessing the impact of corona-virus-19 on nitrogen dioxide levels over southern Ontario, Canada, *Remote Sens.*, 12(24), 1–13, doi:10.3390/rs12244112, 2020.

20 Guevara, M., Jorba, O., Soret, A., Petetin, H., Bowdalo, D., Serradell, K., Tena, C., Van Der Gon, H. D., Kuenen, J., Peuch, V. H. and Pérez García-Pando, C.: Time-resolved emission reductions for atmospheric chemistry modelling in Europe during the COVID-19 lockdowns, *Atmos. Chem. Phys.*, 21(2), 773–797, doi:10.5194/acp-21-773-2021, 2021.

Herman, J., Cede, A., Spinei, E., Mount, G., Tzortziou, M., and Abuhassan, N.: NO₂ column amounts from ground-based Pandora and MFDOAS spectrometers using the direct-sun DOAS technique: Intercomparisons and application to OMI validation, *J. Geophys. Res.*, 114, D13307, doi:10.1029/2009JD011848, 2009.

25 Hudman, R. C., Moore, N. E., Mebust, A. K., Martin, R. V., Russell, A. R., Valin, L. C. and Cohen, R. C.: Steps towards a mechanistic model of global soil nitric oxide emissions: Implementation and space based-constraints, *Atmos. Chem. Phys.*, 12(16), 7779–7795, doi:10.5194/acp-12-7779-2012, 2012.

30 Judd, L. M., Al-Saadi, J. A., Szykman, J. J., Valin, L. C., Janz, S. J., Kowalewski, M. G., et al.: valuating Sentinel-5P TROPOMI tropospheric NO₂ column densities with airborne and Pandora spectrometers near New York City and Long Island Sound. *Atmospheric Measurement Techniques*, 13(11), 6113–6140. <https://doi.org/10.5194/amt-13-6113-2020>, 2020.

- Ialongo, I., Virta, H., Eskes, H., Hovila, J., and Douros, J.: Comparison of TROPOMI/Sentinel-5 Precursor NO₂ observations with ground-based measurements in Helsinki, *Atmos. Meas. Tech.*, **13**, 205–218, <https://doi.org/10.5194/amt-13-205-2020>, 2020.
- 5 Juncosa Calahorrano, J. F., Lindaas, J., O'Dell, K., Palm, B. B., Peng, Q., Flocke, F., Pollack, I. B., Garofalo, L. A., Farmer, D. K., Pierce, J. R., Collett, J. L., Weinheimer, A., Campos, T., Hornbrook, R. S., Hall, S. R., Ullmann, K., Pothier, M. A., Apel, E. C., Permar, W., Hu, L., Hills, A. J., Montzka, D., Tyndall, G., Thornton, J. A. and Fischer, E. V.: Daytime Oxidized Reactive Nitrogen Partitioning in Western U.S. Wildfire Smoke Plumes, *J. Geophys. Res. Atmos.*, **126**(4), e2020JD033484, doi:10.1029/2020JD033484, 2021.
- 10 Kenagy, H. S., Sparks, T. L., Ebben, C. J., Wooldrige, P. J., Lopez-Hilfiker, F. D., Lee, B. H., et al.: NO_x lifetime and NO_y partitioning during WINTER. *Journal of Geophysical Research: Atmospheres*, **123**, 9813–9827, <https://doi.org/10.1029/2018JD028736>, 2018
- Kanniah, K. D., Kamarul Zaman, N. A. F., Kaskaoutis, D. G. and Latif, M. T.: COVID-19's impact on the atmospheric environment in the Southeast Asia region, *Sci. Total Environ.*, **736**, 139658, doi:10.1016/j.scitotenv.2020.139658, 2020.
- 15 Keller, C. A., Evans, M. J., Emma Knowland, K., Hasenkopf, C. A., Modekurty, S., Lucchesi, R. A., Oda, T., Franca, B. B., Mandarino, F. C., Valeria Díaz Suárez, M., Ryan, R. G., Fakes, L. H. and Pawson, S.: Global impact of COVID-19 restrictions on the surface concentrations of nitrogen dioxide and ozone, *Atmos. Chem. Phys.*, **21**(5), 3555–3592, doi:10.5194/acp-21-3555-2021, 2021.
- 20 Kim, Minsu, Gerrit, Kuhlmann, Dominik, Brunner, Dataset for: Importance of satellite observations for high-resolution mapping of near-surface NO₂ by machine learning. *Remote Sensing of Environment*, **264**, art. no. 112573, <https://doi.org/10.1016/j.rse.2021.112573>, 2021.
- Kondragunta, S., Wei, Z., McDonald, B. C., Goldberg, D. L., & Tong, D. Q. COVID-19 induced fingerprints of a new normal urban air quality in the United States. *Journal of Geophysical Research: Atmospheres*, **126**, e2021JD034797, <https://doi.org/10.1029/2021JD034797>, 2021.
- 25 Konovalov, I. B., Beekmann, M., Richter, A. and Burrows, J. P.: Inverse modelling of the spatial distribution of NO_x emissions on a continental scale using satellite data, *Atmos. Chem. Phys.*, **6**(7), 1747–1770, doi:10.5194/acp-6-1747-2006, 2006.
- Koukouli, M. E., Skoulidou, I., Karavias, A., Parcharidis, I., Balis, D., Manders, A., Segers, A., Eskes, H. and Van Geffen, J.: Sudden changes in nitrogen dioxide emissions over Greece due to lockdown after the outbreak of COVID-19, *Atmos. Chem. Phys.*, **21**(3), 1759–1774, doi:10.5194/acp-21-1759-2021, 2021.
- 30 Krotkov, N. A., McLinden, C. A., Li, C., Lamsal, L. N., Celarier, E. A., Marchenko, S. V., Swartz, W. H., Bucsela, E. J., Joiner, J., Duncan, B. N., Boersma, K. F., Veefkind, J. P., Levelt, P. F., Fioletov, V. E., Dickerson, R. R., He, H., Lu, Z. and Streets, D. G.: Aura OMI observations of regional SO₂ and NO₂ pollution changes from 2005 to 2015, *Atmos. Chem. Phys.*, **16**(7), 4605–4629, doi:10.5194/acp-16-4605-2016, 2016.

- Lamsal, L. N., Martin, R. V., Parrish, D. D. and Krotkov, N. A.: Scaling relationship for NO₂ pollution and urban population size: A satellite perspective, *Environ. Sci. Technol.*, 47(14), 7855–7861, doi:10.1021/es400744g, 2013.
- Lamsal, L. N., Duncan, B. N., Yoshida, Y., Krotkov, N. a., Pickering, K. E., Streets, D. G. and Lu, Z.: U.S. NO₂ trends (2005–2013): EPA Air Quality System (AQS) data versus improved observations from the Ozone Monitoring Instrument (OMI), *Atmos. Environ.*, 110(2), 130–143, doi:10.1016/j.atmosenv.2015.03.055, 2015.
- 5 [Lamsal, L. N., Krotkov, N. A., Vasilkov, A., Marchenko, S., Qin, W., Yang, E.-S., Fasnacht, Z., Joiner, J., Choi, S., Haffner, D., Swartz, W. H., Fisher, B., and Bucsela, E.: Ozone Monitoring Instrument \(OMI\) Aura nitrogen dioxide standard product version 4.0 with improved surface and cloud treatments, *Atmos. Meas. Tech.*, 14, 455–479, <https://doi.org/10.5194/amt-14-455-2021>, 2021.](#)
- 10 Lange, K., Richter, A. and Burrows, J. P.: Variability of nitrogen oxide lifetimes and emission fluxes estimated by Sentinel-5P observations, *Atmos. Chem. Phys. Discuss.*, doi:<https://doi.org/10.5194/acp-2021-273>, 2021.
- Laughner, J. and Cohen, R. C.: Direct observation of changing NO_x lifetime in North American cities, *Science* ~~(80-)~~, 366 (6466), 723–727 [online] Available from: <https://science.sciencemag.org/content/366/6466/723.full>, 2019.
- Lee, J. D., Drysdale, W. S., Finch, D. P., Wilde, S. E., and Palmer, P. I.: UK surface NO₂ levels dropped by 42 % during the
- 15 COVID-19 lockdown: impact on surface O₃, *Atmos. Chem. Phys.*, 20, 15743–15759, <https://doi.org/10.5194/acp-20-15743-2020>, 2020.
- Levelt, P. F., Joiner, J., Tamminen, J., Veefkind, J. P., Bhartia, P. K., Fioletov, V., Carn, S., Laatz, J. De, Deland, M., Marchenko, S. and Mcpeters, R.: The Ozone Monitoring Instrument : overview of 14 years in space, *Atmos. Chem. Phys.*, 18, 5699–5745, doi:<https://doi.org/10.5194/acp-18-5699-2018>, 2018.
- 20 [Levelt, P. F., Stein Zweers, D. C., Aben, I., Bauwens, M., Borsdorff, T., De Smedt, I., Eskes, H. J., Lerot, C., Loyola, D. G., Romahn, F., Stavrou, T., Theys, N., Van Roozendaal, M., Veefkind, J. P., and Verhoelst, T.: Air quality impacts of COVID-19 lockdown measures detected from space using high spatial resolution observations of multiple trace gases from Sentinel-5P/TROPOMI, *Atmos. Chem. Phys. Discuss.* \[preprint\], <https://doi.org/10.5194/acp-2021-534>, in review, 2021.](#)
- 25 Liu, F., Beirle, S., Zhang, Q., Dörner, S., He, K. and Wagner, T.: NO_x lifetimes and emissions of cities and power plants in polluted background estimated by satellite observations, *Atmos. Chem. Phys.*, 16(8), 5283–5298, doi:10.5194/acp-16-5283-2016, 2016.
- Liu, F., Choi, S., Li, C., Fioletov, V. E., Mclinden, C. A. and Joiner, J.: A new global anthropogenic SO₂ emission inventory for the last decade : a mosaic of satellite-derived and bottom-up emissions, *Atmos. Chem. Phys.*, 18, 16571–16586,
- 30 2018.
- Liu, F., Page, A., Strode, S. A., Yoshida, Y., Choi, S., Zheng, B., Lamsal, L. N., Li, C., Krotkov, N. A., Eskes, H., Ronald Vander, A., Veefkind, P., Levelt, P., Joiner, J. and Hauser, O. P.: Abrupt declines in tropospheric nitrogen dioxide over China after the outbreak of COVID-19, *arXiv*, (2), 2–7, 2020.
- Lorente, A., Boersma, K. F., Eskes, H. J., Veefkind, J. P., van Geffen, J. H. G. M., de Zeeuw, M. B., Denier van der Gon, H.

- A. C., Beirle, S. and Krol, M. C.: Quantification of nitrogen oxides emissions from build-up of pollution over Paris with TROPOMI, *Sci. Rep.*, 9(1), 1–10, doi:10.1038/s41598-019-56428-5, 2019.
- Lu, Z., Streets, D. G., De Foy, B., Lamsal, L. N., Duncan, B. N. and Xing, J.: Emissions of nitrogen oxides from US urban areas: Estimation from Ozone Monitoring Instrument retrievals for 2005-2014, *Atmos. Chem. Phys.*, 15(18), 10367–10383, doi:10.5194/acp-15-10367-2015, 2015.
- Martin, R. V., Chance, K., Jacob, D. J., Kurosu, T. P., Spurr, R. J. D., Bucsel, E., Gleason, J. F., Palmer, P. I., Bey, I., Fiore, A. M., Li, Q., Yantosca, R. M. and Koelemeijer, R. B. A.: An improved retrieval of tropospheric nitrogen dioxide from GOME, *J. Geophys. Res. D Atmos.*, 107(20), doi:10.1029/2001JD001027, 2002.
- McLinden, C. A., Fioletov, V., Boersma, K. F., Krotkov, N., Sioris, C. E., Veeckind, J. P. and Yang, K.: Air quality over the Canadian oil sands: A first assessment using satellite observations, *Geophys. Res. Lett.*, 39(4), 1–8, doi:10.1029/2011GL050273, 2012.
- McLinden, C. A., Fioletov, V., Shephard, M. W., Krotkov, N., Li, C., Martin, R. V., Moran, M. D. and Joiner, J.: Space-based detection of missing sulfur dioxide sources of global air pollution, *Nat. Geosci.*, 9(7), 496–500, doi:10.1038/ngeo2724, 2016.
- McLinden, C. A., Adams, C. L. F., Fioletov, V., Griffin, D., Makar, P. A., Zhao, X., Kovachik, A., Dickson, N. M., Brown, C., Krotkov, N., Li, C., Theys, N., Hedelt, P. and Loyola, D. G.: Inconsistencies in sulphur dioxide emissions from the Canadian oil sands and potential implications, *Environ. Res. Lett.*, 16, doi:10.1088/1748-9326/abcbbb, 2020.
- [Mehmood K, Bao Y, Petropoulos GP, Abbas R, Abrar MM, Saifullah, Mustafa A, Soban A, Saud S, Ahmad M, Hussain I, Fahad S. Investigating connections between COVID-19 pandemic, air pollution and community interventions for Pakistan employing geoinformation technologies. Chemosphere. 2021 Jun;272:129809. doi: 10.1016/j.chemosphere.2021.129809. Epub 2021 Jan 29. PMID: 33582510; PMCID: PMC7846247.](#)
- Mijling, B. and Van Der A, R. J.: Using daily satellite observations to estimate emissions of short-lived air pollutants on a mesoscopic scale, *J. Geophys. Res. Atmos.*, 117(17), 17302, doi:10.1029/2012JD017817, 2012.
- Miller, S. D., Mills, S. P., Elvidge, C. D., Lindsey, D. T., Lee, T. F. and Hawkins, J. D.: Suomi satellite brings to light a unique frontier of nighttime environmental sensing capabilities, *Proc. Natl. Acad. Sci. U. S. A.*, 109(39), 15706–15711, doi:10.1073/pnas.1207034109, 2012.
- [Misra, P., Takigawa, M., Khatri, P., Dhaka, S.K., Dimri, A.P., Yamaji, K., Kajino, M., Takeuchi, W., Imasu, R., Nitta, K., Patra, P.K., Hayashida, S.; Nitrogen oxides concentration and emission change detection during COVID-19 restrictions in North India \(2021\) Scientific Reports, 11 \(1\), art. no. 9800](#)
- NOAA: 2-Minute Gridded Global Relief Data (ETOPO2v2), [online] Available from: <https://www.ngdc.noaa.gov/mgg/global/etopo2.html> (Accessed 17 June 2020), 2006.
- NPRI: National Pollutant Release Inventory (NPRI). Sulfur oxide emissions for Canada, [online] Available from: www.canada.ca/en/services/environment/pollution-waste-management/national-pollutant-release-inventory.html (Accessed 17 July 2020), 2020.

- Pommier, M., McLinden, C. A. and Deeter, M.: Relative changes in CO emissions over megacities based on observations from space, *Geophys. Res. Lett.*, 40(14), 3766–3771, doi:10.1002/grl.50704, 2013.
- 5 [Potts, D.A., Marais, E.A., Boesch, H., Pope, R.J., Lee, J., Drysdale, W., Chipperfield, M.P., Kerridge, B., Siddans, R., Moore, D.P., Remedios, J. Diagnosing air quality changes in the UK during the COVID-19 lockdown using TROPOMI and GEOS-Chem \(2021\) Environmental Research Letters, 16 \(5\), art. no. 054031, 2021.](#)
- Qu, Z., Jacob, D. J., Silvern, R. F., Shah, V., Campbell, P. C., Valin, L. C. and Murray, L. T.: US COVID-19 shutdown demonstrates importance of background NO₂ in inferring NO_x emissions from satellite NO₂ observations, *Geophys. Res. Lett.*, 48(10), doi:10.1029/2021GL092783, 2021.
- 10 Richter, A., Eyring, V., Burrows, J. P., Bovensmann, H., Lauer, A., Sierk, B. and Crutzen, P. J.: Satellite measurements of NO₂ from international shipping emissions, *Geophys. Res. Lett.*, 31(23), 1–4, doi:10.1029/2004GL020822, 2004.
- 15 [Sannigrahi S., P. Kumar, A. Molter, Q. Zhang, B. Basu, A. S. Basu, F. Pilla, Examining the status of improved air quality in world cities due to COVID-19 led temporary reduction in anthropogenic emissions, Environmental Research, 196., 110927, https://doi.org/10.1016/j.envres.2021.110927, 2021](#)
- SEDAC: The Gridded Population of the World (GPW), [online] Available from:
20 <https://sedac.ciesin.columbia.edu/data/collection/gpw-v4>, 2017.
- Sha, T., Ma, X., Zhang, H., Janecek, N., Wang, Y., Wang, Y., Castro Garcíá, L., Jenerette, G. D. and Wang, J.: Impacts of Soil NO_xEmission on O₃Air Quality in Rural California, *Environ. Sci. Technol.*, doi:10.1021/acs.est.0c06834, 2021.
- 25 [Siciliano, B.; Carvalho, G.; Da Silva, C.M.; Arbilla, G. The Impact of COVID-19 Partial Lockdown on Primary Pollutant Concentrations in the Atmosphere of Rio de Janeiro and São Paulo Megacities \(Brazil\). Bull. Environ. Contam. Toxicol., 105, 2–8, 2020.](#)
- Silvern, R. F., Jacob, D. J., Mickley, L. J., Sulprizio, M. P., Travis, K. R., Marais, E. A., Cohen, R. C., Laughner, J. L., Choi, S., Joiner, J. and Lamsal, L. N.: Using satellite observations of tropospheric NO₂ columns to infer long-term trends in US NO_x emissions: The importance of accounting for the free tropospheric NO₂ background, *Atmos. Chem. Phys.*, 19(13), 8863–8878, doi:10.5194/acp-19-8863-2019, 2019.
- 30 Stavrakou, T., Müller, J. F., Bauwens, M., Boersma, K. F. and van Geffen, J.: Satellite evidence for changes in the NO₂ weekly cycle over large cities, *Sci. Rep.*, 10(1), 1–9, doi:10.1038/s41598-020-66891-0, 2020.
- Streets, D. G., Canty, T., Carmichael, G. R., De Foy, B., Dickerson, R. R., Duncan, B. N., Edwards, D. P., Haynes, J. A., Henze, D. K., Houyoux, M. R., Jacob, D. J., Krotkov, N. A., Lamsal, L. N., Liu, Y., Lu, Z., Martin, R. V., Pfister, G. G., Pinder, R. W., Salawitch, R. J. and Wecht, K. J.: Emissions estimation from satellite retrievals: A review of current capability, *Atmos. Environ.*, 77, 1011–1042, doi:10.1016/j.atmosenv.2013.05.051, 2013.
- 35 [Tack, F., Merlaud, A., Iordache, M.-D., Pinardi, G., Dimitropoulou, E., Eskes, H., Bomans, B., Veeffkind, P., and Van Roozendael, M.: Assessment of the TROPOMI tropospheric NO₂ product based on airborne APEX observations, Atmos. Meas. Tech., 14, 615–646, https://doi.org/10.5194/amt-14-615-2021, 2021.](#)
- Vadrevu, K. P., Eaturu, A., Biswas, S., Lasko, K., Sahu, S., Garg, J. K. and Justice, C.: Spatial and temporal variations of air

pollution over 41 cities of India during the COVID-19 lockdown period, *Sci. Rep.*, 10(1), 1–15, doi:10.1038/s41598-020-72271-5, 2020.

- 5 Veefkind, J. P. P., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., Claas, J., Eskes, H. J. J., de Haan, J. F. F., Kleipool, Q., van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser, H. and Levelt, P. F. F.: TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications, *Remote Sens. Environ.*, 120(2012), 70–83, doi:10.1016/j.rse.2011.09.027, 2012.
- [Venter, Z. S., Aunan, K., Chowdhury, S. Jos Lelieveld, J.: COVID-19 lockdowns cause global air pollution declines, PNAS 117 \(32\) 18984-18990.; https://doi.org/10.1073/pnas.2006853117, 2020.](https://doi.org/10.1073/pnas.2006853117)
- 10 Verhoelst, T., Compernolle, S., Pinardi, G., Lambert, J. C., Eskes, H. J., Eichmann, K. U., Fjæraa, A. M., Granville, J., Niemeijer, S., Cede, A., Tiefengraber, M., Hendrick, F., Pazmiño, A., Bais, A., Bazureau, A., Folkert Boersma, K., Bogner, K., Dehn, A., Donner, S., Elokhov, A., Gebetsberger, M., Goutail, F., Grutter De La Mora, M., Gruzdev, A., Gratsea, M., Hansen, G. H., Irie, H., Jepsen, N., Kanaya, Y., Karagkiozidis, D., Kivi, R., Kreher, K., Levelt, P. F., Liu, C., Müller, M., Navarro Comas, M., PETERS, A. J. M., Pommereau, J. P., Portafaix, T., Prados-Roman, C., 15 Puertedura, O., Querel, R., Remmers, J., Richter, A., Rimmer, J., Cárdenas, C. R., De Miguel, L. S., Sinyakov, V. P., Stremme, W., Strong, K., Van Roozendaal, M., Pepijn Veefkind, J., Wagner, T., Wittrock, F., Yela González, M. and Zehner, C.: Ground-based validation of the Copernicus Sentinel-5P TROPOMI NO₂ measurements with the NDACC ZSL-DOAS, MAX-DOAS and Pandonia global networks, *Atmos. Meas. Tech.*, 14(1), 481–510, doi:10.5194/amt-14-481-2021, 2021.
- 20 Vinnikov, K. Y., Dickerson, R. R., Krotkov, N. A., Edgerton, E. S. and Schwab, J. J.: The net decay time of anomalies in concentrations of atmospheric pollutants, *Atmos. Environ.*, 160, 19–26, doi:10.1016/j.atmosenv.2017.04.006, 2017.
- Virghileanu, M., Săvulescu, I., Mihai, B. A., Nistor, C. and Dobre, R.: Nitrogen dioxide (No₂) pollution monitoring with sentinel-5p satellite imagery over europe during the coronavirus pandemic outbreak, *Remote Sens.*, 12(21), 1–29, doi:10.3390/rs12213575, 2020.
- 25 Williams, J. E., Folkert Boersma, K., Le Sager, P. and Verstraeten, W. W.: The high-resolution version of TM5-MP for optimized satellite retrievals: Description and validation, *Geosci. Model Dev.*, 10(2), 721–750, doi:10.5194/gmd-10-721-2017, 2017.
- [Zhao, X., Griffin, D., Fioletov, V., McLinden, C., Cede, A., Tiefengraber, M., Müller, M., Bogner, K., Strong, K., Boersma, F., Eskes, H., Davies, J., Ogyu, A., and Lee, S. C.: Assessment of the quality of TROPOMI high-spatial-resolution NO₂ data products in the Greater Toronto Area, Atmos. Meas. Tech., 13, 2131–2159, https://doi.org/10.5194/amt-13-2131-2020, 2020.](https://doi.org/10.5194/amt-13-2131-2020)
- 30 Zhang, H., Lin, Y., Wei, S., Loo, B. P. Y., Lai, P. C., Lam, Y. F., Wan, L. and Li, Y.: Global association between satellite-derived nitrogen dioxide (NO₂) and lockdown policies under the COVID-19 pandemic, *Sci. Total Environ.*, 761, 144148, doi:10.1016/j.scitotenv.2020.144148, 2021.

Zhang, L., Jacob, D. J., Knipping, E. M., Kumar, N., Munger, J. W., Carouge, C. C., van Donkelaar, A., Wang, Y. X., and Chen, D.: Nitrogen deposition to the United States: distribution, sources, and processes, Atmos. Chem. Phys., 12, 4539–4554, <https://doi.org/10.5194/acp-12-4539-2012>, 2012.

5 Zoogman, P., Liu, X., Suleiman, R. M., Pennington, W. F., Flittner, D. E., Al-Saadi, J. A., Hilton, B. B., Nicks, D. K.,
Newchurch, M. J., Carr, J. L., Janz, S. J., Andraschko, M. R., Arola, A., Baker, B. D., Canova, B. P., Chan Miller,
C., Cohen, R. C., Davis, J. E., Dussault, M. E., Edwards, D. P., Fishman, J., Ghulam, A., González Abad, G., Grutter,
M., Herman, J. R., Houck, J., Jacob, D. J., Joiner, J., Kerridge, B. J., Kim, J., Krotkov, N. A., Lamsal, L., Li, C.,
Lindfors, A., Martin, R. V., McElroy, C. T., McLinden, C., Natraj, V., Neil, D. O., Nowlan, C. R., O’Sullivan, E. J.,
10 Palmer, P. I., Pierce, R. B., Pippin, M. R., Saiz-Lopez, A., Spurr, R. J. D., Szykman, J. J., Torres, O., Veeffkind, J. P.,
Veihelmann, B., Wang, H., Wang, J. and Chance, K.: Tropospheric emissions: Monitoring of pollution (TEMPO), J.
Quant. Spectrosc. Radiat. Transf., doi:10.1016/j.jqsrt.2016.05.008, 2016.

Table 1. The standard deviations of the random errors and interannual variability for background, urban and industrial components for the U.S. and Canada in percent. The random errors are calculated as the averages of estimates for individual years. The interannual variability estimates are the standard deviations calculated from three years (2018, 2019, and 2021). Interannual variability of the industrial component is calculated for regions with estimated total emissions greater than 1 kt yr⁻¹.

5

Area	Random Error (%)			Interannual Variability (%)		
	Background	Urban	Industrial	Background	Urban	Industrial
Atlanta	0.20	1.04	1.47	5.9	9.7	11.3
Boston	0.29	0.93	5.68	4.0	10.8	39.3
Calgary	0.27	1.62	2.40	7.5	3.5	23.8
Charlotte	0.25	1.80	2.64	5.1	7.0	19.2
Chicago	0.27	0.73	1.74	17.5	9.3	11.6
Dallas	0.20	0.82	2.34	5.8	9.2	6.2
Denver	0.27	0.89	2.29	4.0	7.8	17.5
Detroit	0.27	0.99	2.17	13.3	5.5	16.8
Edmonton	0.34	0.80	1.91	11.0	15.6	8.1
Houston	0.22	0.78	1.25	6.7	13.0	10.5
Las Vegas	0.12	0.48		7.6	16.7	
Los Angeles	0.22	0.22		1.4	9.2	
Miami	0.15	0.88	2.66	1.9	5.8	40.7
Minneapolis	0.20	1.42	3.28	15.1	10.1	29.5
Montreal	0.29	1.21	3.24	11.8	7.3	19.7
New York	0.36	0.41	5.17	8.2	5.8	8.0
Orlando	0.17	1.17	2.16	4.0	10.6	16.5
Phoenix	0.17	0.75	2.84	4.9	17.0	23.5
Pittsburgh	0.33	1.62	2.20	6.5	6.3	7.0
Portland	0.29	0.87	4.32	5.8	21.2	23.2
San Antonio	0.18	1.50	1.20	8.1	5.9	42.0
San Francisco	0.17	0.64	4.64	1.1	12.8	20.9
Seattle	0.31	1.14	2.71	9.0	11.0	18.9
St. Louis	0.20	1.48	1.87	16.3	4.5	7.8
Toronto	0.29	0.78	1.97	10.8	12.9	13.7
Vancouver	0.49	0.74	5.74	2.1	9.6	14.7
Washington	0.24	1.03	2.76	6.8	13.2	8.0
Average	0.25	0.99	2.82	7.5	10.0	18.3
<u>Standard deviation</u>	<u>0.03</u>	<u>0.08</u>	<u>0.39</u>	<u>1.3</u>	<u>4.4</u>	<u>4.3</u>

Table 2. The standard deviations of the random errors and interannual variability for background, urban and industrial components for 13 regions in percent. The random errors and interannual variability are calculated from three years (2018, 2019, and 2021) for each area and then averaged for all areas in the region. Interannual variability of the industrial component is calculated for regions with estimated total emissions greater than 1 kt yr⁻¹.

Area	Random Error (%)			Interannual Variability (%)		
	Background	Urban	Industrial	Background	Urban	Industrial
Africa	0.21	1.56	2.42	7.3	19.8	21.3
Australia and New Zealand	0.34	0.80	1.22	8.6	9.2	12.2
Canada and US	0.25	0.99	2.82	7.5	10.0	18.3
Central America	0.21	0.95	1.72	5.4	11.0	20.3
China	0.30	1.06	1.90	7.3	19.6	16.4
Europe-1	0.23	0.96	4.06	5.6	12.3	27.6
Europe-2	0.25	1.30	3.13	6.8	15.4	17.1
India	0.23	1.78	0.77	9.0	16.6	12.4
Japan Korea, Taiwan	0.31	0.86	1.29	4.9	12.9	13.2
Middle East	0.29	1.43	0.86	7.5	22.2	19.9
Northern Eurasia	0.29	1.42	2.78	8.8	13.5	25.4
South America	0.39	2.97	6.72	7.8	18.6	23.6
South-East Asia	0.29	1.03	1.68	8.0	14.1	16.1
Average	0.28	1.32	2.41	7.3	15.0	18.8
Standard deviation	0.05	0.58	1.61	1.3	4.1	4.9




Figure 1. Mean TROPOMI NO₂ VCDs over the US and southern Canada for March 16–June 15, in (top) 2018-2019 and (bottom) 2020. The main features of the NO₂ distribution such as elevated NO₂ values over large cities, industrial sources and in the valleys such the California Central Valley (1)-and lower values over the mountains such as Appalachian (2) are evident from the plot. Also Note, that NO₂ VCDs are not negligible (about $5 \cdot 10^{14} \text{ cm}^{-2}$) even over remote vast remote areas such as National Forests in Montana (3) or Algonquin Provincial Park in Ontario (4) as well as over the oceans areas suggesting some background levels. The maps are based on Level 2 data gridded on 0.1° by 0.1° grid grouped by the wind speed: (left) less than 10 km per hour and (right) more than 10 km per hour.






Figure 2. Changes in weekly Google mobility data (for “retail and recreation” category) relative to the baseline period (Jan 3 – Feb 6, 2020) for twelve regions analysed in this study. The black vertical lines represent the beginning and the end of the period analyzed in this study (March 16 – June 15).








Figure 32. (a) Mean TROPOMI NO₂ for March 16– June 15, 2018-2019, over the Montreal area, (b) the fitting results and (c) the residuals (i.e., the difference between (a) and (b)). Tropospheric NO₂ VCDs have a large “background” level that is reflected by (d) the elevation-related component. (e) The population density-related and (f) industrial sources-related components. Panel (b) is the sum of panels (d), (e) and (f). Emission point sources are shown by the black dots and the airport by the slightly larger gray dot. The industrial sources-related component is comprised of three clusters: one (g) with two sources and two (h, i) with one source each. The data are smoothed by the oversampling technique with the averaging radius R=10 km. Panel (b) is the sum of panels (d), (e) and (f). Proxies used by the statistical model (Eq. 1): (j) Elevation map on the colour scale that is similar to that for the elevation-related (“background”) component; (k) population density map. (l) Mean

TROPOMI values (panel *a*) vs. the fitting results (panel *b*). Each dot represents the mean value for a cell on 0.2 by 0.2 grid for 2018 or 2019. The number of data points (N), the correlation coefficient (R) and the slope are also shown.




Figure 4. Mean TROPOMI NO₂ for March 16 – June 15 over a flat area around Minneapolis and mountain area around Seattle as indicated. The columns represent: mean TROPOMI NO₂ values, (column *a, d*), the fitting results (*b, e*), and the elevation-related background component (*c, f*). Elevation map on the colour scale that is similar to the background component (*g, i*) and the population density maps (*h, j*). The “hotspots” on the population density maps correspond to Minneapolis (*h*) and Seattle (*j*).





Figure 35. The map of locations of the analysed 27 most populated urban cities in (red) the US and (blue) Canada (22 and 5 areas respectively). The analysis was done for 3° (latitude) by 4° (longitude) areas around the sites. The mean NO₂ values for eight areas for the period from March 16 to June 15 in 2018-2019 and 2020 are also shown.






Figure 46. Mean TROPOMI NO₂ for March 16–June 15 over the four areas as indicated. The columns represent: mean TROPOMI NO₂ VCD values, (column *a*), the fitting results (*b*), the residuals (*c*) ~~and as~~ well as individual components of the fitting: the elevation-related (*d*), the population density-related (*e*), and the industrial sources-related (*f*). VCDs estimated from reported emissions are in column (*g*) and the difference between columns *f* and *g* is in column *h*. 1- Hartsfield–Jackson Atlanta International Airport, 2- oil refineries near Houston.




Figure 5. Mean TROPOMI NO₂ for March 16–June 15 over a flat area around Minneapolis and mountain area around Seattle as indicated. The columns represent: mean TROPOMI NO₂ values, (column *a, d*), the fitting results (*b, e*), and the elevation-related background component (*c, f*).

5




Figure 67. The contribution of the three components to the total NO₂ mass in the Montreal area for March 16 – June 15 (average for 2018-2019). The total mass can be represented as a sum of three components shown in Figure 2.

5

2018 -2019
Region=Canada and U.S.










Figure 7. Relative contribution of the three components (background, urban, and industrial) components for all 27 Canadian and U.S. areas for March 16 – June 15 (average for 2018-2019). The contribution from industrial sources and cities are responsible for less than a third of total NO₂ mass of the analysed 3° by 4° urban areas.








Figure 8. (left) The background (top row), urban (middle row), and industrial (bottom row) components for all 27 analysed areas in the U.S. and Canada in 2018-2019 (blue) and 2020 (orange). (right) The decline 2020 mean-values in percent from the mean-2018-2019 values-values. The background component is expressed as the mean value of that component for the analysed area. The urban component is expressed as annual emissions per capita, and the industrial component is expressed as total emissions from the point sources for the period from March 16 to June 15, in 2018-2019 (blue) and 2020 (orange) values for the three components for the (top) background, (middle) urban, and (bottom) industrial components for all 27 analysed area The grey dashed lines on the right panels indicate the 2- σ level for the interannual variability. (right) The decline 2020 mean values in percent from the mean 2018-2019 values.







Estimated vs. reported industrial source emission rates in 2018-2019 in kt/year



5 **Figure 9.** Estimated and reported annual NO₂ emissions rates for U.S. sources for 2018-2019. Each dot represents the sum of all emissions in one urban area in 2018 or 2019 and there are 40 dots on the plot. The emissions are expressed as annual rates.

~~TROPOMI emissions were calculated only for the sites that reported their emissions.~~ The correlation coefficient between the two data sets is ~~0.849~~ and the slope is 0.71 ± 0.15 . ~~The standard deviation of the residuals is about 5 kt y^{-1} .~~ The plot also shows the predicted regression line (blue), 95% confidence intervals for the regression mean (the shaded area), and 95% prediction intervals (dashed lines).

- ~~5 Areas that have no reported emissions data for 2019 are not included in the plot.~~












Figure 10a. The same as Fig. 8, but for Europe-1 sub-region (Italy, France, Spain, Portugal, Belgium, Ireland, and UK).






Figure 10b. The same as Fig. 8, but for Europe-2 sub-region (other EU countries and non-members from former Yugoslavia).




Figure 11. Similar to Figure 4-6 columns a-f, for areas around four European cities: Manchester, Paris, Milan, and Prague.
1- Charles de Gaulle Airport, 2- power plants in Germany, 3- power plants in Czech Republic.






Figure 12. Similar to Figure 4 columns a-e, for Saint Petersburg, and Dar es Salaam, cities with similar population of about 6 million people, but very different NO₂ “footprint”. There are no large industrial sources in these two areas.















Figure 123. (top) The map of the background component NO_2 in 2018-2019 for the period March 16 – June 15 estimated from TROPOMI. (bottom) The map of emissions-annual per capita urban NO_2 emissions for the same period. The analysis was done using estimates for large-cities with population greater than cities (> 1 million)6 million in China and 1 million for the rest of the world-.



5 **Figure 13.** The map of NO₂ percent changes between 2018-2019 and 2020 for the period March 16 – June 15 estimated from TROPOMI data for (top) background component and (bottom) urban annual emissions per capita. The analysis was done using estimates for cities with population greater than 6 million in China and greater than 1 million for the rest of the world.






Figure 144. (left) The mean values for March 16–June 15 in 2018-2019 (blue) and 2020 (orange) values for ~~the three~~ **the three** components for the (top) **the** background-, (middle) urban-, and (bottom) industrial components for the 13 regions. (right) The decline of 2020 mean values in percent from the mean 2018-2019 values. The data are sorted according to the changes between 2020 and 2018-2019 in the urban component (middle right panel). ~~The error bars represent 2σ intervals.~~ Mean values for each region were calculated as a mean of **the values from** all areas for that region. The uncertainty (σ) was calculated as a standard error of the mean. The error bars represent 2σ intervals.




Figure 156. (left) A scatter plot of Google mobility statistic (for “retail and recreation” sites) changes vs. TROPOMI NO₂ VCD changes for (left) the urban (population density related) and (right) background components during the period from March 16 to June 15, 2020 compared to the baseline period. The Google mobility statistic changes show the difference with the pre-lockdown period (Jan 3 – Feb 6, 2020) in percent. For TROPOMI, the difference is between the 2020 values and the average for the 2018-2019 average. Each dot symbol represents one country/country; the dot colour demonstrates the region as shown in the legend. Only countries with at least two cities used in this study are included in this plot. The correlation coefficient between the two data sets is 0.6662. The dashed Y=X line is shown for reference. The error bars represent the standard errors.