

Modeling secondary organic aerosol formation from volatile chemical products

Elyse A. Pennington^{1,2}, Karl M. Seltzer¹, Benjamin N. Murphy³, Momei Qin^{1,4}, John H. Seinfeld²,

5 Havala O.T. Pye³

¹Oak Ridge Institute for Science and Education Fellow in the Office of Research and Development, US Environmental Protection Agency, Research Triangle Park, NC 27711

²Department of Chemical Engineering, California Institute of Technology, Pasadena, CA 91125

³Office of Research and Development, US Environmental Protection Agency, Research Triangle Park, NC 27711

10 ⁴Jiangsu Key Laboratory of Atmospheric Environment Monitoring and Pollution Control, Collaborative Innovation Center of Atmospheric Environment and Equipment Technology, School of Environmental Science and Engineering, Nanjing University of Information Science & Technology, Nanjing, China

Correspondence to: Havala O.T. Pye (Pye.Havala@epa.gov)

Abstract. Volatile chemical products (VCPs) are commonly-used consumer and industrial items that are an important source of anthropogenic emissions. Organic compounds from VCPs evaporate on atmospherically relevant time scales and include many species that are secondary organic aerosol (SOA) precursors. However, the chemistry leading to SOA, particularly that of intermediate volatility organic compounds (IVOCs), has not been fully represented in regional-scale models such as the Community Multiscale Air Quality (CMAQ) model, which tend to underpredict SOA concentrations in urban areas. Here we develop a model to represent SOA formation from VCP emissions. The model incorporates a new VCP emissions inventory and employs three new classes of emissions: siloxanes, oxygenated IVOCs, and nonoxygenated IVOCs. VCPs are estimated to produce $1.67 \mu\text{g m}^{-3}$ of noontime SOA, doubling the current model predictions and reducing the SOA mass concentration bias from -75% to -58% when compared to observations in Los Angeles in 2010. While oxygenated and nonoxygenated intermediate volatility VCP species are emitted in similar quantities, SOA formation is dominated by the nonoxygenated IVOCs. Formaldehyde and SOA show similar relationships to temperature and bias signatures indicating common sources and/or chemistry. This work suggests that VCPs contribute up to half of anthropogenic SOA in Los Angeles and models must better represent SOA precursors from VCPs to predict the urban enhancement of SOA.

1 Introduction

Organic aerosol (OA) is a major component of fine particulate matter ($\text{PM}_{2.5}$) in urban areas throughout the world (Zhang et al., 2007). $\text{PM}_{2.5}$ influences human health (Lim et al., 2012), climate (Intergovernmental Panel on Climate Change, 2014), and visibility (Hyslop, 2009), so understanding OA composition is an important step in mitigating the adverse effects of $\text{PM}_{2.5}$. Secondary organic aerosol (SOA) is often the dominant component of OA (Jimenez et al., 2009) and is formed when gas-phase volatile organic compounds (VOCs) react with atmospheric oxidants to form products that condense into the aerosol phase, where they can undergo further reaction. SOA is formed via thousands of atmospheric reactions (Goldstein & Galbally, 2007), so understanding its sources remains a challenge.

Volatile chemical products (VCPs) are an important source of organic emissions that lead to SOA formation (McDonald et al., 2018; Qin et al., 2021). As vehicle exhaust becomes cleaner and mobile source emissions decline, the relative importance of VCP emissions increases (Khare & Gentner, 2018). Previous work suggests that during the 2010 California Nexus of Air Quality and Climate Change (CalNex) campaign in Southern California (Ryerson et al., 2013), VCPs contributed approximately $1.1 \mu\text{g m}^{-3}$, or 41%, of observed SOA above background levels in the Los Angeles Basin (Qin et al., 2021).

Modeling the formation of SOA in three-dimensional (3D) chemical transport models (CTMs) is challenging due to the complexity of VOC chemistry and computational constraints of regional-scale modeling. Models have tended to underpredict SOA mass in urban locations for a variety of reasons. For one, the SOA formation potential of intermediate

volatility organic compounds (IVOCs) and semivolatile organic compounds (SVOCs) – or S/IVOCs – is not well constrained. Observations made during the CalNex campaign demonstrate that S/IVOCs are important sources of SOA, making up 10% of total gas-phase organic compound concentrations (Zhao et al., 2014) while contributing up to 80% of above-background SOA mass (Hayes et al., 2015). Although it is often impossible to identify all individual species contributing to ambient S/IVOCs, these compounds may be classified based on their properties (e.g. volatility). Volatility basis set (VBS) models (Donahue et al., 2011) are often used to represent S/IVOC chemistry and partitioning, and have improved model estimates of SOA (Woody et al., 2016; Hayes et al., 2015; Robinson et al., 2007). Murphy et al. (2017) integrated a VBS model into the Community Multiscale Air Quality (CMAQ) model version 5.2 to represent the multigenerational aging of semivolatile primary organic aerosol (POA) leading to the production of SOA. Other studies have parameterized VBS models to represent S/IVOCs from mobile emissions (Lu et al., 2020; Jathar et al., 2017), but none have parameterized SOA formation from VCP S/IVOCs emissions. Additionally, the emissions of S/IVOCs are not well constrained and are often not included in detailed emissions inventories (Zhao et al., 2015). However even when S/IVOCs are included in emissions inventories, they are often assigned to nonreactive or nonvolatile model surrogates that do not participate in model chemistry (T. Shah et al., 2020). Improving the representation of SOA chemistry in CMAQ will allow for more accurate exposure estimates in health studies and source apportionment for air quality management decisions.

Deleted: only

Another source of error in CTMs is the lack of representation of oxygenated SOA precursors. Historically, mechanism development has focused on the oxidation chemistry of species emitted primarily from vehicles (e.g. BTEX: benzene, toluene, ethylbenzene, and xylene) or biogenic sources (e.g. isoprene, monoterpenes). While VCPs do emit some of these species, they also emit many oxygenated compounds (Seltzer et al., 2021; McDonald et al., 2018). The implications of a few important oxygenated precursors on air quality have recently been quantified (e.g. Janecek et al., 2017; Charan et al., 2020; L. Li & Cocker, 2018; W. Li et al., 2018), but many oxygenated precursors have not been studied in a laboratory setting. For the few oxygenated VCPs that have been studied in laboratory chambers, SOA yields were reported under unrealistic atmospheric conditions, e.g. high OH and aerosol seed concentrations (Charan et al., 2021). So, the SOA yields of these compounds have primarily been estimated using models such as the Statistical Oxidation Model (SOM; Cappa & Wilson, 2012) or VBS (McDonald et al., 2018; R. U. Shah et al., 2020). These oxygenated species are not included as SOA precursors in most models, and their chemistry is needed to improve predictions of SOA mass.

In this work, we introduce a chemical mechanism to represent SOA formation from VCPs. Specifically, the potential of both oxygenated and nonoxygenated IVOCs to form SOA is developed and evaluated. We utilize a new VCP emissions inventory known as VCPy (Seltzer et al., 2021) to represent organic emissions from VCPs and to parameterize model species behavior in the chemical mechanism. The chemistry and emissions inventory are implemented in the CMAQ model version 5.3.2 to simulate air quality during the CalNex campaign in California in 2010. The model predictions are compared to measurements made in Pasadena during CalNex and the speciation of predicted SOA is examined.

Deleted: Community Multiscale Air Quality (

Deleted:)

Deleted: ¶

2 Methods

2.1 VCPy emissions inventory implementation

90 VCPy is a modeling framework that estimates reactive organic carbon emissions from VCPs (Seltzer et al., 2021). Within
this framework, the complete VCP sector is disaggregated into several product use categories (PUCs; e.g. cleaning products,
personal care products, adhesives and sealants, paints and coatings). U.S. nationwide usage of each PUC is estimated, and
survey data are then used to quantify the mass fraction of organic, inorganic, and water proportions, as well as speciate the
95 organic fraction. Physiochemical properties of each organic component are used to estimate the characteristic evaporation
timescale, which is then compared to an assigned use timescale to determine whether a compound is retained or evaporated
from each PUC. In the initial implementation of VCPy (version 1.0), which is representative of 2016 conditions, the
predicted nationwide and Los Angeles County VCP emission rates were 9.5 kg person⁻¹ year⁻¹ and 8.2 kg person⁻¹ year⁻¹,
respectively. These emission rates are consistent with the low end of values seen in a previous study that used a top-down
100 approach to estimate VCP emissions (Qin et al., 2021). In our work, product use is based on data from 2010 with
composition specified using data from the early 2000s to overlap with the CalNex campaign.

Since the speciation of organic emissions from VCPy is explicit, the underlying chemical and physical properties of
emissions are output from the framework. These properties, many of which are relevant to atmospheric oxidation and
subsequent SOA formation, include the oxidation rate with the hydroxyl radical (k_{OH}), molecular weight (MW), ~~effective~~
105 saturation concentration (C*), and oxygen-to-carbon ratio (O:C). SOA mass yields, which are defined as the mass of SOA
formed per mass of ROC precursor reacted, were assigned based on compound-specific structure and volatility (Seltzer et al.,
2021).

110 A key step in implementing this inventory into CMAQ is ensuring that all compounds predicted to be emitted by VCPy are
mapped to either an existing or a new model surrogate. Emissions of low-volatility organic vapors (C* < 10^{6.5} µg m⁻³) from
all sources are prime SOA precursors but traditionally discarded from the gas-phase chemical mechanism used in many
CTMs (e.g. represented as nonvolatile (NVOL), nonreactive (NROG), or unspecified IVOC species that are not used in the
chemical mechanism of CMAQ). As a result, these species do not participate in atmospheric chemistry and thus do not
115 impact radical concentrations or SOA mass. In addition, oxygenated compounds are not currently included as SOA
precursors in many mechanisms because of the historic focus on SOA formation from nonoxygenated vehicle exhaust and
traditional VOCs like single-ring aromatics and biogenic hydrocarbons. The work of Qin et al. (2021) specifically identifies
this loss of emitted reactive carbon mass as a reason for underestimated SOA from the personal care sector in the CMAQ
model. To account for the SOA potential of this previously neglected organic mass, all compounds currently mapped to

Deleted: volatility

Formatted: Normal, No bullets or numbering

Deleted: ; Carter, 2010).

NROG, NVOL, and IVOC are reviewed, with most of this mass routed to one of three newly added categories of model surrogates: siloxanes (SILOX), oxygenated IVOCs (SOAOXY), ~~or~~ nonoxygenated IVOCs (IVOCP3, IVOCP4, IVOCP5, IVOCP6, IVOCP5ARO, and IVOCP6ARO). The updated mechanism (with SOA pathways described in Section 2.2) with the newly implemented speciation mapping is henceforth described as SAPRC07TIC_AE7I_VCP and the complete list of assignment rules is provided in the SI Methods.

Deleted: and

County-level VCPy emissions (Seltzer et al., 2021) were gridded at 4-km scale to fit the CalNex domain (Baker et al., 2015) using a variety of spatial surrogates. The spatial surrogates used depend on the category of VCP emissions being described: agricultural land is used as a proxy for all agricultural pesticide emissions, the density of oil and gas wells for the oil and gas solvent emissions, and population for all remaining VCP sources. While some categories of VCP emissions could feature more refined spatial surrogate proxies, the uncertainty associated with spatial allocation of sources may be lower than uncertainty in individual source strength. More specifically, if an entire VCP category could be matched to a single surrogate, allocation methods would still assume there is no variation in the strength of individuals within the population of that surrogate (Y. Li et al., 2021).

Deleted: In summary:

All silicon-containing species (including siloxanes and silanes) are assigned to SILOX. Species assigned to low-volatility, inactive species (i.e. NVOL, NROG, or IVOC) whose ozone and SOA formation are not otherwise captured in existing gas-phase surrogates are assigned to the new IVOC surrogates (SOAOXY, IVOCP3/4/5/6/5ARO/6ARO) or an existing long alkane surrogate (SOAALK) according to their structure, O:C, C*, and SOA yield. All species assigned to SOAOXY, IVOCP3/4/5/6/5ARO/6ARO, and SOAALK are also mapped to existing alkane surrogates based on k_{OH} to participate in ozone chemistry. Siloxanes are not mapped to alkane surrogates because of their low k_{OH} . Since these alkane species do not participate in the SOA module and the VCP surrogates do not affect ozone chemistry, this double-mapping does not double-count chemistry due to the lack of overlap between these parts of the gas- and aerosol-phase mechanisms. Some compounds were manually reassigned depending on their SOA yield. All other compounds retained their assignments to existing model surrogates.

All VCP emissions feature a sinusoidal diurnal profile with a peak at noon, with no application of day-of-week or seasonal profiles. Since the simulation period used in this study is a single month, no seasonal changes would be observable over this time frame and previous work suggests little seasonal variability in VCP emissions (Gkatzelis et al., 2021). Other emission sectors (e.g. mobile sources, agriculture) are adjusted for seasonal impacts based on meteorological conditions and known activity data.

Deleted: The seasonal effects applied to other sectors of emissions, e.g. mobile source emissions, are a function of the meteorological conditions. ...

2.2 Parameterizing SOA formation from VCPs

To better represent the atmospheric chemistry of VCPs, SOA formation is added for the three new categories of emissions (siloxanes, oxygenated IVOCs, and nonoxygenated IVOCs) in the SAPRC07TIC_AE7I_VCP chemical mechanism within CMAQ (Table 1).

Formatted: Font: 10 pt, Not Bold

Deleted: Table 1

Cyclic volatile methylsiloxanes (cVMS), or siloxanes for short, are present in many personal care products, adhesives, and sealants. Collectively, siloxanes represent a large fraction of VCP emissions (Seltzer et al., 2021). Decamethylcyclopentasiloxane (D₅-siloxane) is the most prevalent siloxane in urban atmospheres (D.-G. Wang et al., 2013) and laboratory studies have found D₅-siloxane SOA yields ranging from 0% (Charan et al., 2021) to 50% (Janecek et al., 2019). The explicit oxidation mechanism is unknown and the SOA yields of other siloxanes are not well understood (Coggon et al., 2018). Here, siloxanes are treated separately from other oxygenated VCP species due to their anomalously low OH oxidation rate (Table 1). The mechanism of SOA formation used here utilizes an existing two-product model from

Formatted: Font: 10 pt, Not Bold

Deleted: Table 1

Janecek et al. (2019) that was parameterized using oxidation flow reactor (OFR) experiments and photooxidation chamber data from Wu & Johnston (2017). In this implementation, the OH oxidation rate constant for D₅-siloxane matches the rate reported in Janecek et al. (2017) and the hydroxyl radical is replenished after reaction.

Few laboratory chamber studies have investigated the oxidation processes of other oxygenated gas-phase species (e.g. Charan et al., 2020; L. Li & Cocker, 2018), so little experimental data exist about the SOA yields or oxidation products of oxygenated SOA precursors. Additionally, many models that predict the products of oxidation reactions (e.g. SOM and VBS) have not been parameterized or evaluated using oxygenated precursors. Without these models and laboratory studies, little is known about the oxidation products of these precursors, which limits our ability to develop a detailed model of their SOA formation. Therefore, all non-siloxane oxygenated IVOC emissions are represented by a single surrogate (SOAOXY) that undergoes a one-step gas-phase reaction with the hydroxyl radical to form a nonvolatile aerosol surrogate (AOIVOC). This simple mechanism reduces the reliance on many parameters that are not well-constrained. The MW, k_{OH} , C*, and SOA yield of this surrogate are calculated as a mass-weighted average of the oxygenated IVOC emissions from VCPs in Los Angeles County which are generally consistent with what would be calculated using nationwide information.

Nonoxygenated IVOC emissions are represented using the model described by Lu et al. (2020), which uses a VBS model and multigenerational aging scheme to represent the SOA from gasoline, diesel, and aircraft sources. Six surrogates are differentiated by structure (alkane vs. aromatic) and effective saturation concentration, and each is assigned a four-product yield distribution, generating SVOCs after one oxidation step. Many of the nonoxygenated IVOC species from mobile and VCP emission sources have similar structures (i.e. long and branched alkanes and aromatics), volatilities, and SOA yields (see Fig. S1), making the Lu et al. (2020) model a good representation of oxidation and SOA formation from nonoxygenated VCP IVOCs.

Deleted: volatility

Table 1. Properties of the VCP surrogates added to CMAQ version 5.3.2.

	MW (g mol ⁻¹)	$k_{OH} \times 10^{11}$ (cm ³ molec ⁻¹ sec ⁻¹)	α_i	C* (μ g m ⁻³)	SOA mass yield (at 10 μ g m ⁻³)	H _{vap} (kJ mol ⁻¹)	κ_{org}	H (M atm ⁻¹)	OM/ OC
SILOX	368.66 ^a	0.155 ^b	-	-	-	-	-	3.87×10^{2f}	-
SV SILOX1/ AS SILOX1/	416.66 ^a	-	0.14 ^c	0.95 ^c	0.13	131 ^d	0.09 ^e	2.97×10^{6f}	3.49 ^e
SV SILOX2/	384.66 ^a	-	0.82 ^c	484 ^c	0.017	101 ^d	0.05 ^e	7.99×10^{4f}	3.22 ^e

ASILOX2J									
SOAOXY ^g	170.95	2.54	-	-	-	-	-	2.85 x 10 ³ ^f	-
AOIVOCJ ^g	186.95	-	-	-	0.0628	-	0.09 ^e	-	1.73 ^e
IVOCP3 ^h	296.6	2.65	^h	10 ³	0.43	52	-	2 x 10 ⁸	-
IVOCP4 ^h	254.9	2.25	^h	10 ⁴	0.43	41	-	2 x 10 ⁸	-
IVOCP5 ^h	219.4	1.89	^h	10 ⁵	0.35	30	-	2 x 10 ⁸	-
IVOCP6 ^h	184.4	1.55	^h	10 ⁶	0.15	19	-	2 x 10 ⁸	-
IVOCP5ARO ^h	197.3	7.56	^h	10 ⁵	0.36	30	-	2 x 10 ⁸	-
IVOCP6ARO ^h	162.3	3.05	^h	10 ⁶	0.25	19	-	2 x 10 ⁸	-

210 ^aThe gas-phase siloxane (SILOX) MW is the average of the MW of all VCPy siloxane and silane species weighted by Los Angeles County emission rates. The MW of the higher-volatility siloxane products (SVSILOX2/ASILOX2J) is approximated as the sum of the MW of SILOX and one oxygen. The MW of the lower-volatility products (SVSILOX1/ASILOX1J) has an additional two oxygens to represent its significant decrease in volatility.

^bThe gas-phase siloxane (SILOX) _{KOH} is given in Janecek et al. (2017).

215 ^cThe stoichiometric product yields (α_i) and C* of the siloxanes are given in Janecek et al. (2019).

^dEnthalpy of vaporization (H_{vap}) values for the siloxanes are estimated according to the method in Epstein et al. (2010).

^eAll OM/OC ratios and hygroscopicity parameters (κ_{org}) are estimated using equations 5 and 12, respectively, in Pye et al. (2017).

^fHenry's Law constants (H) at 298.15 K are estimated using the surrogate-based method in Hodzic et al. (2014).

220 ^gThe MW, _{KOH}, C*, and SOA yield of SOAOXY (gas) and AOIVOCJ (aerosol) are calculated as a mass-weighted average of the oxygenated IVOC emissions from VCPs in Los Angeles County. Because AOIVOC is formed via a single reaction with a constant SOA yield, it is treated as nonvolatile and therefore is not assigned a C* or H_{vap} .

^hAll nonoxygenated IVOC surrogate properties – including four stoichiometric product yields (α_i) for each surrogate used in the multigenerational scheme – are described in Lu et al. (2020).

225

2.3 CMAQ model implementation

2.3.1 CalNex model configuration

230 The updated chemical mechanism and VCPy-derived emissions were implemented in CMAQ version 5.3.2 (US EPA Office of Research and Development, 2020). CMAQ version 5.3 and the subsequent minor releases are documented in Appel et al. (2021). The model was used to simulate air quality during the CalNex campaign from May 15 to June 15, 2010, with an additional 14-day spin-up period. Outside the VCP updates, the model configuration matches the implementation used in Qin et al. (2021) and Lu et al. (2020). The model domain has 4-km x 4-km horizontal resolution (325 x 225 grid cells) covering
235 California and Nevada with 36 vertical levels reaching 50 mbar. Meteorological inputs are derived from the Weather Research and Forecasting (WRF) Advanced Research WRF core Model version 3.8.1 (Skamarock et al., 2008). Gas-phase chemistry is represented using SAPRC07TIC (Pye et al., 2013; Xie et al., 2013) with the addition of the VCP chemical mechanism summarized in [Table 1](#). Aerosol-phase chemistry is simulated using an extended version of the AERO7 mechanism, depicted in [Figure 1](#), including all AERO7 reactions plus those of the new VCP mechanism (boxed in red) and mobile IVOCs (boxed in red in the lower left) that participate in the multigenerational aging shown in the orange boxes (Lu et al., 2020). This diagram also includes a representation of the aqueous-phase cloud chemistry and removal used in the Asymmetric Convection Model (ACM) version 2 module (Binkowski & Roselle, 2003), which has been updated to include wet deposition properties for the new aerosol surrogates ([Table 1](#)).
240
245 All non-VCP anthropogenic emissions are based on the 2011 National Emissions Inventory (NEI) version 2 (US EPA, 2015). VCP emissions in the NEI are removed and replaced with VCPy predicted emissions using the Detailed Emissions Scaling, Isolation, and Diagnostic (DESID) module (Murphy et al., 2021). Mobile NO_x emissions were reduced by 25% in all simulations to better match observational data from the CalNex campaign (Qin et al., 2021). Mobile IVOC emissions and the semivolatile treatment of mobile POA were treated according to the methods described in Lu et al. (2020). The IVOCs
250 are assigned to the appropriate IVOC_{P3/4/5/6/5ARO/6ARO} surrogates that are also used to treat nonoxygenated IVOCs from VCPs. Wind-blown dust emissions are neglected in this study. Biogenic emissions are calculated online using the Biogenic Emission Inventory System (BEIS) version 3.6.1 (Bash et al., 2016) as are sea spray aerosol emissions.

Deleted: Table 1

Formatted: Font: 10 pt, Not Bold

Deleted: Fig. 1

Formatted: Font: 10 pt, Not Bold

Deleted: Table 1





Figure 1. Treatment of OA chemistry in the CMAQv5.3.2+VCP model. The thick black box surrounds all aerosol-phase species. All smaller black boxes depict species undergoing gas-phase oxidation from VOCs to semivolatile or nonvolatile SOA species. Orange font depicts the VBS model for S/IVOCs. Red font depicts particle-phase accretion reactions while purple font depicts particle-phase hydrolysis reactions. Green font represents heterogeneous processes. Blue font shows cloud-processed aerosol and yellow font shows aerosol water associated with the organic phase. Gray boxes are nonvolatile primary organic aerosol (POA) species. Double-sided arrows represent reversible processes and one-sided arrows represent irreversible processes. Dashed lines represent processes that are dependent on relative humidity. The diagram includes the AERO7 mechanism plus the three VCP-forming pathways specific to this work (thick boxes in red). See AE7I Species Table (2016/2021) for species descriptions.



Deleted:

Deleted: d

2.3.2 Simulation cases

Three simulations were evaluated against the observations collected during the CalNex campaign. A “zero VCP” case removes all VCP emissions. The “CMAQv5.3.2” case is a standard CMAQ simulation with base emissions (i.e. VCP emissions from the NEI) and base chemistry (i.e. no new VCP chemistry). Finally, the “CMAQv5.3.2+VCP” case adds both the VCP chemistry described above (i.e. SAPRC07TIC_AE7I_VCP) and replaces all NEI VCP emissions with VCPy-derived VCP emissions. Comparisons between the “zero VCP” case and the “CMAQv5.3.2+VCP” case illustrate the complete impact of VCPy emissions on modeled SOA. In contrast, comparisons between the “CMAQv5.3.2” case and the

“CMAQv5.3.2+VCP” case illustrate the impact of the new representation of VCP emissions and chemistry against the current status of VCPs in CMAQ. Results from the CMAQv5.3.2 case are presented primarily in the SI.

2.3.3 Comparison with observations

280

Observational data are provided by a suite of instruments deployed during the 2010 CalNex campaign in Pasadena. There were two data collection sites in the CalNex campaign – Pasadena and Bakersfield – and model predictions are compared to measurements made at the Pasadena site, which is located in the Los Angeles Basin approximately 18 km downwind of the urban core (Ryerson et al., 2013). PM₁ (fine particulate matter with diameter < 1 μm) OA data were obtained with an aerosol mass spectrometer (AMS) and have been analyzed via positive matrix factorization (PMF) to determine its composition (Hayes et al., 2013). Formaldehyde (HCHO) data are provided in Warneke et al. (2011) and carbon monoxide (CO) data are available from Santoni et al. (2014). Ozone data throughout California were obtained from the EPA AQS monitoring network for 178 sites operating during the simulation period (US EPA, 2013). Hourly ozone concentrations were used to calculate daily maximum 8-hour average (MDA8) ozone concentrations.

285

290

3 Results & Discussion

3.1 VCP emissions and implications for SOA

295

VCP emissions were split almost equally between species that do and do not form SOA. The SAPRC07TIC_AE7I_VCP speciation mapping (Figure 2) indicates 56.4% (4.8×10^7 kg year⁻¹) of Los Angeles County VCP emitted mass does not form SOA. This mass includes small species commonly used as solvents, such as ethanol, acetone, and small alkanes. The remaining 43.6% (3.7×10^7 kg year⁻¹) of Los Angeles County emissions are assigned to model surrogates that form SOA. 3.5% of the total emissions are assigned to siloxanes, 7.8% to oxygenated IVOCs, 11.8% to nonoxygenated IVOCs, and 20.4% to traditional SOA precursors, such as VOC alkanes, toluene, and other aromatics. The volatility and SOA yields of species in each category are summarized in Fig. S1.

300

305

Figure 2 indicates that in traditional model processing, precursors to SOA are systematically discarded from chemistry calculations. As described in Section 2.1, low-volatility emissions (i.e. NROG, NVOL, and IVOC) do not participate in SOA or radical chemistry in traditional SAPRC07TIC_AE7I which is a key issue in representing SOA mass. The inner ring of Figure 2 depicts the fraction of each category that was originally assigned to inactive species (NROG, NVOL, and IVOC; hatched) versus other existing surrogates (solid). 2.6×10^7 kg year⁻¹ (30.7%) of the total VCP emissions were originally assigned to these surrogates and did not participate in any atmospheric chemistry processes. Using the new speciation and mechanism, 1.8×10^7 kg year⁻¹ (21.2% of total VCP emissions) were reassigned to surrogates that form SOA in the model

Deleted: Fig. 2

Deleted: long


Deleted: Figure 2

Deleted: there are

Deleted: that

Deleted: Fig. 2

(hatched inner ring: red, blue, orange, and purple). The remaining $8.0 \times 10^6 \text{ kg year}^{-1}$ (9.4% of total VCP emissions; inner ring hatched green) is comprised of species with SOA yields of zero and were not reassigned to SOA-forming surrogates.



320 **Figure 2. Percentage of the VCP emissions assigned to each category of CMAQ surrogates using the SAPRC07TIC_AE7L_VCP**
speciation profiles. The total rate of VCP emissions in Los Angeles County is $8.3 \times 10^7 \text{ kg yr}^{-1}$. The outer ring depicts the
percentage of total VCPy-derived emissions assigned to each of the three new VCP categories (siloxanes in red, oxygenated IVOCs
in blue, and nonoxygenated IVOCs in orange), the traditional SOA precursors described by existing model surrogates (purple),
and existing surrogates that do not form SOA (green). The inner ring gives an indication of the original assignments of each of the
325 **outer ring categories. Hatching indicates emissions originally assigned to model surrogates that do not participate in model**
chemistry: IVOC, NVOL, and NROG. Solid colors represent other surrogate assignments.

Averaged over the duration of the CalNex campaign, VCPs are predicted to be a larger source of IVOCs than mobile sources, as shown by the increase in gas-phase IVOC mass in the CMAQv5.3.2+VCP case compared to the zero VCP case (Fig. S2). Across mobile and VCP sources during CalNex, CMAQ predicts $6.4 \mu\text{g m}^{-3}$ of the gas-phase IVOC mass is nonoxygenated and $2.6 \mu\text{g m}^{-3}$ of the IVOC mass is oxygenated (Fig. S2). The observed campaign-average total IVOC concentration was $10.5 \mu\text{g m}^{-3}$ (Zhao et al., 2014), with $6.3 \mu\text{g m}^{-3}$ attributed to hydrocarbon-like IVOCs and $4.2 \mu\text{g m}^{-3}$ attributed to oxygenated IVOCs. However, this observed estimate of oxygenated IVOCs is conservative (lower bound) based on the experimental method employed by Zhao et al. (2014). Thus, the predicted nonoxygenated IVOC mass, which includes contributions from both mobile and VCP sources, reproduces observations with high fidelity. CMAQ, which only considers
335 IVOCs from VCP and mobile sectors, underpredicts the mass of oxygenated IVOCs by 38%, suggesting additional missing products of oxidation or emissions.

The new SOA systems combined with traditional SOA precursors in CMAQ resulted in an effective SOA yield for the VCP sector – defined as the emission-weighted average of the individual species’ mass-based SOA yields – of 5.6% for Los Angeles County. This Los Angeles County yield is in good agreement with the work of Qin et al. (2021), that found a 5% yield led to SOA predictions consistent with ambient observational constraints. The U.S. effective VCP SOA yield (5.3%) is
340

only slightly lower than the yield expected for Los Angeles, due to differences stemming from the variability in the composition of VCP emissions nationwide versus in Los Angeles.

3.2 CMAQ results: SOA, ozone, and formaldehyde

Modeled PM_{10} SOA increased considerably in response to the newly implemented VCP emissions and chemistry, bringing model predictions into closer agreement with observations. Daily maximum PM_{10} SOA concentrations increased from $1.4 \mu g m^{-3}$ (-79% mean bias) in the zero VCP case to $2.8 \mu g m^{-3}$ (-58% mean bias) in the CMAQv5.3.2+VCP case, compared to the observed peak value of $6.6 \mu g m^{-3}$ (Figure 3a). The diurnal distributions resulted from photochemistry and the sinusoidal distribution of VCP emissions that peak at 12:00 local time. Modeled $PM_{2.5}$ SOA concentrations improved for all mass loadings and all hours of the day, with the slope of modeled versus observed concentrations increasing from 0.23 in the zero VCP case to 0.43 in the CMAQv5.3.2+VCP case (Figure 4a). Results for the CMAQv5.3.2 case are given in Figs. S3 and S4. Modeled $PM_{2.5}$ SOA displayed similar behavior as PM_{10} SOA; i.e. the organic fraction and secondary organic fraction of $PM_{2.5}$ was only marginally smaller than the corresponding fractions of PM_{10} and followed the same diurnal pattern.




Figure 3. a) Average hourly concentrations of background-corrected PM_{10} SOA observed and simulated by the zero VCP and CMAQv5.3.2+VCP modeling cases May 15-June 15. Box and whiskers show all hourly concentrations observed by AMS at the CalNex site. A constant background value was removed from all observed concentrations according to the method in Hayes et al. (2015). The background value of each simulation was determined by averaging the lower 50% of hourly concentrations from 00:00 LT to 04:00 LT and subtracting that from each curve. b) Average hourly concentration of total (not size-resolved) SOA for the two simulation cases and their difference (CMAQv5.3.2+VCP – zero VCP). c) Difference in hourly concentrations of total SOA by category.

The difference between hourly averaged total (i.e. not size-resolved) SOA concentrations in the zero VCP and CMAQv5.3.2+VCP case are shown in Figure 3b and the contributions to that difference from categories of SOA surrogates are shown in Figure 3c. Of the three new categories of VCP emissions, nonoxygenated IVOC precursors formed the most

Deleted: ¶
3.2.1 SOA ¶

Deleted: Fig. 3

Formatted: Subscript

Formatted: Font: Not Bold, Check spelling and grammar

Deleted: Fig. 4

Deleted: Fig. 3

Deleted: Fig. 3

375 SOA in CMAQ. The increased SOA from the nonoxygenated IVOC VCP precursors reached a peak concentration of 1.14 $\mu\text{g m}^{-3}$, equal to 69% of the total noontime difference. This can be explained by the high SOA yields of the individual species (Fig. S1) and the model surrogates.

380 SOA from oxygenated IVOC VCP precursors reached a peak concentration of 0.11 $\mu\text{g m}^{-3}$ (6.7% of the SOA difference). While oxygenated IVOC emissions were similar in abundance to nonoxygenated IVOC emissions (Figure 2), these species lead to less SOA formation due to their lower SOA yields (Fig. S1): higher degrees of oxygenation tend to promote fragmentation upon reaction with OH (Jimenez et al., 2009), producing smaller molecules with higher volatilities and lower potential to form SOA. It is possible that the net yield of modeled SOA from oxygenated IVOC precursors will increase as the results from more laboratory studies become available, or if a more detailed model is used. For example, particle-phase oligomerization reactions from oxygenated IVOC precursors would produce nonvolatile aerosol products, but this chemistry has not yet been investigated in an atmospheric chamber.

385 Siloxanes formed very little SOA, reaching a maximum of 21 ng m^{-3} (1.3% of the SOA difference) at noon. Despite having nonnegligible SOA yields (Fig. S1) and emission rates (Fig. 2), siloxanes react with OH on long time scales (Table 1). As such, this results in low localized SOA mass, which is consistent with other modeling and laboratory studies that have predicted siloxanes to form SOA on the order of ng m^{-3} or less (Charan et al., 2021; Milani et al., 2021; Janecek et al., 2017). The low resultant SOA mass demonstrates that while gas-phase siloxanes serve as a useful tracer for personal care product and adhesive emissions from VCPs (Gkatzelis et al., 2021), particle-phase products from siloxane oxidation may not form quickly enough to serve as a reliable tracer for these emissions.

395 While traditional species accounted for the greatest fraction of VCP SOA precursor emissions that lead to SOA formation (Figure 2), they contributed only 23% (0.39 $\mu\text{g m}^{-3}$) of the increased noontime SOA in the CMAQv5.3.2+VCP case. These traditional SOA precursors form SOA less efficiently than the IVOC surrogates (Fig. S1), so they result in less SOA formation than IVOCs despite higher emissions.

400 While this work indicates oxygenated IVOCs form much less SOA than nonoxygenated IVOCs, more work is needed to determine if this result is robust across all emission sectors and in future conditions. Oxygenated IVOCs represent a class of emissions that has traditionally been discarded from regional models, but have become an important research focus with the rising importance of VCP emissions (Khare & Gentner, 2018). The contribution of oxygenated IVOCs and siloxanes to ambient conditions may be spatially variable and continue to evolve as product formulations shift towards exempt VOCs which tend to be oxygenated. Oxygenated IVOCs from other emissions sources, such as meat cooking or wood burning, could be abundant but were not considered here. Additionally, we do not know if SOA from these precursors has a health impact higher or lower than average $\text{PM}_{2.5}$.

Deleted: Fig. 2

Deleted: .H

Deleted: similarly

Formatted: Font: 10 pt, Not Bold

Deleted: Table 1

Deleted: Despite

Deleted: ing

Deleted: Fig. 2

415

The SOA from VCP IVOCs reached a daily maximum of $1.25 \mu\text{g m}^{-3}$ on average at noon (Figure 3c). IVOCs from mobile sources contributed an additional $1.1 \mu\text{g m}^{-3}$ at noon (Lu et al., 2020). Therefore this updated CMAQ model predicted a total IVOC-derived SOA concentration of $2.35 \mu\text{g m}^{-3}$, equivalent to 35% of the total observed above-background PM_{10} SOA concentration ($6.6 \mu\text{g m}^{-3}$). Previous work stated that 40-85% of above-background SOA concentrations in Pasadena are attributable to S/IVOCs (Hayes et al., 2015), suggesting that additional processes are still needed in the model. This will be discussed further in Section 3.3.

420




Figure 4. Modeled concentrations predicted by CMAQ zero VCP case (green) and CMAQv5.3.2+VCP case (blue) vs. observations from the CalNex Pasadena ground site. The line with a slope of 1 is indicated with a gray dashed line. a) Hourly PM_{10} SOA. b) Hourly formaldehyde (HCHO). c) MDA8 O_3 . Background values were not removed from any panels.

425

Formaldehyde is one of the most abundant VOCs in the atmosphere and observations of this compound can serve many purposes. Biomass burning, vehicles, and other urban sources emit formaldehyde, and because of its short lifetime (~hours), it can serve as a proxy for local organic emissions. It is also formed in the atmosphere when VOCs undergo radical reactions, oxidize, and fragment, so it serves as an indicator for SOA chemistry since it is formed by many of the same reactions that also lead to SOA formation (Seinfeld & Pandis, 2016). In addition, it is depleted by photolysis and is an important source of radical initiation reactions (Griffith et al., 2016). Formaldehyde can be retrieved directly by satellites (Levelt et al., 2018), which can be used to validate ground data, evaluate model predictions, and predict OA concentrations remotely (Liao et al., 2019). For all of these reasons, formaldehyde is a useful indicator of VOC chemistry in a model.

435

Predicted formaldehyde concentrations improved in response to the new VCP emissions and chemistry, indicating that model updates improve the representation of VOC chemistry beyond SOA in the model. Similar to predicted SOA, formaldehyde concentrations increased at all times, with the ratio of modeled to observed values increasing from 0.58 in the zero VCP case to 0.75 in the CMAQv5.3.2+VCP case (Figure 4b). The diurnal profile of hourly averaged formaldehyde

Moved (insertion) [1]

Deleted: Fig. 3

Deleted:

Formatted: Caption, Centered, Line spacing: 1.5 lines

Deleted: 4

Deleted: ¶

Moved up [1]: The SOA from VCP IVOCs reached a daily maximum of $1.25 \mu\text{g m}^{-3}$ on average at noon (Fig. 3c). IVOCs from mobile sources contributed an additional $1.1 \mu\text{g m}^{-3}$ at noon (Lu et al., 2020). Therefore this updated CMAQ model predicted a total IVOC-derived SOA concentration of $2.35 \mu\text{g m}^{-3}$, equivalent to 35% of the total observed above-background PM_{10} SOA concentration ($6.6 \mu\text{g m}^{-3}$). Previous work stated that 40-85% of above-background SOA concentrations in Pasadena are attributable to S/IVOCs (Hayes et al., 2015), suggesting that additional processes are still needed in the model. This will be discussed further in Section 3.3.¶

Deleted: ¶
3.2.2 Formaldehyde¶

Formatted: Font: Not Bold, Check spelling and grammar

Deleted: Fig. 4

concentrations is given in Fig. S3. This work focused primarily on improving the representation of SOA from VCPs, so radical chemistry for the new SOA precursors was treated using existing alkane-like behavior (surrogates ALK1/2/3/4/5).

460 With a more detailed representation of VCP radical chemistry, predicted formaldehyde concentrations may improve further.

The bias in predicted ozone concentrations was also reduced by including VCP chemistry. The ratio of modeled to observed concentrations increased from 0.72 in the zero VCP case to 0.95 in the CMAQv5.3.2+VCP case (Figure 4c). Improved ozone is also seen for all operational AQS sites in the California modeling domain, with the modeled to observed ratio increasing from 0.63 in the zero VCP case to 0.70 in the CMAQv5.3.2+VCP case (Fig. S5). The diurnal profile of hourly averaged ozone concentrations is given in Fig. S3. This study focused on VCP behavior in relation to SOA formation and used existing model species to capture ozone formation. Future work focusing on the ozone chemistry of VCPs could change the magnitude and diurnal profile of predicted ozone.

470 SOA can be facilitated by increases in oxidant abundance and chemical pathways from precursors to semivolatile or low-volatility products. Average noontime total SOA mass increased from 1.96 $\mu\text{g m}^{-3}$ in the zero VCP case to 3.62 $\mu\text{g m}^{-3}$ in the CMAQv5.3.2+VCP case (Figure 3b), an increase of 84.7%. Ozone concentration can be used as an indicator of oxidant burden and oxidation rates due to its high responsiveness, while OH concentrations may be less responsive (Qin et al., 2021).

The average noontime ozone concentration increased from 43.0 ppb in the zero VCP case to 49.2 ppb in the CMAQv5.3.2+VCP case (Fig. S3c), an increase of 14.4%. Assuming ozone can serve as a proxy for oxidation rates, the improved ozone concentration suggests that ~14.4% of increased model SOA concentrations are due to an increase in the oxidant burden and oxidation rates. The SOA mass increased by a larger percentage (84.7%), indicating emissions and chemistry updates combined were approximately 5 times $[(84.7\% - 14.4\%) / 14.4\%]$ more effective than enhanced oxidant levels alone in increasing SOA. This is consistent with the work of Qin et al. (2021), which found that the lack of key

480 emitted precursors in models – rather than their associated radical chemistry – had the largest impact on PM_{2.5} formation. Additionally, we note that the default CMAQ model (CMAQv5.3.2) with baseline chemistry and VCP emissions predicted about the same amount of SOA as the zero VCP case (Fig. S3a). In contrast, ozone increased in the default CMAQv5.3.2 model with VCPs (Fig. S3c). Since the oxidant burden increased noticeably in the CMAQv5.3.2 case but did not equate to a large increase in PM₁ SOA, results suggest the oxidant level alone does not have a large influence on enhancing SOA if the relevant precursor pathways are not also implemented.

490 The response of formaldehyde can similarly be compared to the change in oxidant burden due to VCPs. At noontime, average formaldehyde increased from 2.41 ppb in the zero VCP case to 2.80 ppb in the CMAQv5.3.2+VCP case, an increase of 16.2%. As above, we attribute ~14.4% of the increase in pollutant concentration to the increase in oxidation rates. While formaldehyde does contribute to the oxidant burden via photolysis and radical initiation, the contribution of formaldehyde to the RO_x radical budget is likely small and on the order of 10% (e.g. Griffith et al., 2016; Kaiser et al., 2015; Luecken et al.,

Deleted: The increase in formaldehyde was predominantly due to the increased magnitude of VCP emissions. This work focused

Deleted: it is expected that

Deleted: will

Deleted: 3.2.3 Ozone

Formatted: Font: Not Bold, Check spelling and grammar

Deleted: Fig. 4

Deleted: As stated in Section 3.2.2,

Deleted: t

Deleted: Assuming ozone can serve as a proxy for oxidation rates, the increase in ozone concentration relative to the increase in SOA concentration provides information about the proportional impact of oxidant level versus combined emissions and chemistry on total SOA mass. Average noontime total SOA mass increased from 1.96

Deleted: Fig. 3

Deleted: This suggests that the

Deleted: from VCPs

Deleted: PM₁ SOA increased most in the CMAQv5.3.2+VCP case, with little increase in the CMAQv5.3.2 case compared to the zero VCP case (Fig. S3a). ...

Deleted: linearly between the three simulation cases

Deleted: T

Deleted: supporting the idea that

2018). Thus, the increase in formaldehyde concentrations between simulation cases is likely due primarily to the increase in oxidation rate. The increase in formaldehyde between simulation cases, therefore, cannot be largely attributed to the addition of S/IVOC emissions and their ability to form formaldehyde as a byproduct of oxidation. This is consistent with the work of Coggon et al. (2021), which showed that vehicle VOCs perturb formaldehyde to a larger degree than VCP VOCs do, suggesting that VCP emissions and fragmentation chemistry may not be directly responsible for formaldehyde, but rather modulate formaldehyde formation via changes in oxidant abundance.

3.3 Features of remaining model bias

The residual PM₁ SOA bias in Pasadena is well-correlated with ambient temperature (Figure 5a). PM₁ SOA bias is defined as modeled hourly concentrations minus observed hourly concentrations. At cooler temperatures in the overnight hours, bias is low and fluctuates around zero. However, as temperature increases towards midday and SOA concentrations increase, the bias becomes more negative, indicating greater model underprediction.

SOA concentrations can be a function of temperature based on precursor emissions and chemistry throughout the day.

Previous work demonstrated that observed OA in Los Angeles is positively correlated with temperature, and declining OA concentrations have been due largely to reductions of temperature-independent OA. Because this corresponds to a decline in anthropogenic emissions, they suggest that anthropogenically-derived OA is largely temperature-independent while biogenically-derived OA is largely temperature-dependent (Nussbaumer & Cohen, 2021). Modeled OA is positively correlated with temperature, consistent with the observed Los Angeles OA, and is driven by the larger, secondary portion of OA, rather than POA (Fig. S7). However, the improvement to predicted SOA between simulation cases was seen unequally at different temperatures, as indicated by the larger reduction in absolute model bias at higher temperatures (Figure 5a). This suggests that the SOA derived from VCP species have a temperature-dependent response, in addition to the biogenic emissions cited in Nussbaumer & Cohen (2021). In particular, because nonoxygenated IVOCs were the dominant source of increased SOA predicted by the CMAQv5.3.2+VCP simulation, this work suggests that S/IVOCs are an important source of temperature-dependent SOA in Los Angeles.

Because S/IVOCs have been shown to be a major constituent of modeled SOA and contribute to the correlation between SOA bias and temperature, other sources of S/IVOCs emissions may account for some of the remaining residual SOA bias in the model. For example, asphalt emissions are proposed to contribute 8-30% of total S/IVOC emissions in the South Coast Air Basin in Southern California and have SOA mass yields exceeding 10% (Khare et al., 2020). Their potential to form SOA is very large, and because asphalt emissions are highly temperature-dependent, the SOA increase would be seen largely during midday resulting in an improvement of high-temperature SOA bias. In addition, the underprediction of oxygenated gas-phase IVOCs (Section 3.1) suggests that additional sources of oxygenated IVOC precursors may be missing from the

Deleted: Fig. 5

Deleted: The correlation between SOA bias and temperature is likely caused in part by the inclusion and representation of S/IVOCs emissions and their chemistry in the model. ...

Deleted: Our results indicate that

Deleted: concentrations

Deleted: are

Deleted: . The correlation

Deleted: Fig. 5

complete inventory. One possible explanation of the temperature-dependence of the SOA bias is that modeled SOA volatility is too high. But, oxygenated SOA is nonvolatile and nonoxygenated IVOC SOA is continually processed to lower volatility through gas-phase OH oxidation.

560




Figure 5. Bias (modeled - observed) of hourly concentrations vs. modeled temperature for the zero VCP case (green) and CMAQv5.3.2+VCP case (blue). Hourly concentrations are binned into five temperature ranges of 5°C each and the data in each bin is represented by a box-and-whisker plot. The horizontal midline depicts the median of the data, the edges of the box extend from the lower to upper quartile of the data, and the whiskers extend from the minimum to the maximum of the data. a) PM₁ SOA bias (μg m⁻³). b) PM₁ POA bias (μg m⁻³). c) Formaldehyde (HCHO) bias (ppb). d) CO bias (ppb).

565

Formaldehyde, CO, and POA are often used to understand the atmospheric evolution of SOA because they are products of the same anthropogenic activity and/or VOC oxidation chemistry that forms SOA. As such, they can be used to better understand the remaining sources of error in the model. POA is formed via combustion from vehicles, industrial processes, cooking, and biomass burning (Jathar et al., 2014; Huffman et al., 2009). CO and formaldehyde are emitted from many processes and formed as products of atmospheric VOC oxidation (Seinfeld & Pandis, 2016). These species are often used to understand the effect of dilution on SOA (Hayes et al., 2013). Dilution is caused both by atmospheric transport away from

570

emission sources, as well as the change in planetary boundary layer (PBL) height over the diurnal cycle. VCPs do not emit
575 POA, CO, or formaldehyde, so any changes observed in their simulated concentrations were caused by chemical and
physical processing in the existing model.

The POA bias did not express the same temperature dependence as SOA, and thus POA is not affected in the same way in
the model by the processes causing the temperature-dependence of SOA bias. Since VCPs do not emit POA and all other
580 emission sources were consistent between simulation cases, the slight increase in POA concentrations between the zero VCP
and CMAQv5.3.2+VCP cases (Figs. 5 and S7) is due to increased partitioning of semivolatile POA into the particle-phase
resulting from higher total OA mass loadings (the treatment of semivolatile POA in CMAQ is described in Murphy et al.
(2017)). The POA bias can be exclusively attributed to errors in combustion source emissions inventories and meteorological
585 effects. The combustion source inventories also include emissions of gaseous SOA precursors, which may be incorrectly
modeled even if the POA emissions are accurate, especially for cooking and biomass burning sources. While the POA bias
does decrease with increasing temperature, it is positive at all temperatures and does not have larger underpredictions at
higher temperatures (Figure 5b). Due to the inconsistency between POA and SOA behavior, errors influencing the emission
and transport of POA can likely not be used to describe the temperature dependence of SOA bias. The POA bias also does
not provide information about the error in vapor emissions from combustion sources – including S/IVOCs – and their
590 temperature-dependence, and improving combustion emissions inventories may help to close the model-observation gap for
SOA.

CO is often used to account for the effects of dilution by scaling SOA to CO enhancement ($\Delta\text{CO} = \text{CO} - \text{CO}_{\text{background}}$).
Negligible changes in the CO concentration were found between simulation cases considered here (Fig. S3) and the model
595 CO bias is uncorrelated with temperature (Figure 5d). The consistency of predicted CO concentration between cases implies
that CO is not affected by the emissions changes to the VCP sector and thus cannot separate SOA formation efficiency from
lack of emitted precursors. CO enhancement serves as an effective indicator and correction factor for mobile source
emissions in urban areas (e.g. Hayes et al., 2013; Ensberg et al., 2014; Woody et al., 2016), but this work indicates that CO is
not an effective tracer for distinguishing VCPs from other sources. The lack of correlation between CO and temperature also
600 implies that errors in the modeled PBL height at different times of day (and potential impact on dilution of pollutant
concentrations) is not an important driver of the SOA bias temperature-dependence.

In contrast to POA and CO, the formaldehyde bias demonstrated the same trend with temperature as SOA (Figure 5c). This
suggests that formaldehyde is affected by emissions, chemistry, and dilution changes similarly to SOA. This is supported by
605 the stronger correlation seen between SOA and formaldehyde compared to the correlation between SOA and POA or CO
(Fig. S8). Therefore, formaldehyde may provide more information about the errors in modeling VOC chemistry and possibly
SOA formation. It is possible that remaining formaldehyde bias is due to missing formaldehyde emissions. The VCP

Deleted: Fig. 5

Deleted: Thus POA is not affected in the same way in the model by the processes causing the temperature-dependence of SOA bias.

Deleted: Fig. 5

Deleted: ¶

Another potential factor that could influence the temperature-dependence of SOA bias is modeled PBL height. Previous studies that used the same modeled meteorological data as this study (Lu et al., 2020) showed that the modeled PBL height in Pasadena tends to be overestimated during midday and underestimated overnight compared to ceilometer data, which could result in a dilution effect matching the temperature dependence seen in Fig. 5a. However, the predicted CO bias does not depend on temperature, which implies that modeled PBL height is not an important driver of the SOA bias

Deleted: Fig. 5

Moved down [2]: As stated above, the behavior of POA and CO bias suggest that errors in combustion emissions and PBL height cannot fully describe the temperature-dependence of SOA bias, and POA and CO are better indicators of mobile and industrial sources.

Deleted: Formaldehyde may instead serve as a better indicator of SOA production in urban areas where VCPs are important atmospheric constituents. This may be particularly true for SOA formed from S/IVOCs, which we have shown to be a dominant source of SOA in Los Angeles. Therefore, formaldehyde may

Deleted: modeling

635 inventory includes near-zero emissions of formaldehyde, but formaldehyde is emitted from wooden furniture and emission
rates increase with temperature (Y. Wang et al., 2021). This may account for some of the temperature-dependence of
formaldehyde bias, but likely not the entirety since the VCP emissions inventory has been evaluated with select ambient
VOC measurements with low error (Seltzer et al., 2021). One possible explanation of the temperature-dependence of both
the SOA and formaldehyde biases is missing sources of emissions and resulting chemistry. Previous work has shown that
formaldehyde formation is particularly sensitive to the emissions/chemistry of alkenes (e.g. isoprene) and, to a lesser extent,
640 alkanes and aromatics (Luecken et al., 2018), so these precursors likely indicate missing emissions as a source of error in
our model. While the radical chemistry of these hydrocarbon precursors are included in the model, additional missing
chemistry may be causing some of the error. Chemical processes that have not been included in the mechanism include
autooxidation (Crouse et al., 2013) – which forms low-volatility SOA – and formaldehyde potentially formed from the
fragmentation of S/IVOC precursors to SOA. The inclusion of these missing emissions and/or chemistry would further
645 impact oxidant levels, which we have shown to be an important source of modeled SOA and formaldehyde. As stated above,
the behavior of POA and CO bias suggest that errors in combustion emissions and PBL height cannot fully describe the
temperature-dependence of SOA bias, and POA and CO are better indicators of mobile and industrial sources. Formaldehyde
may instead serve as a better indicator of SOA production in urban areas where VCPs are important atmospheric
constituents. While many factors may contribute to the temperature-dependence of SOA and formaldehyde bias, future work
650 must investigate the importance of these factors and tracking the response of formaldehyde to these changes alongside SOA
could provide insight.

Deleted: Because formaldehyde is a major product of VOC oxidation, the temperature-dependence of both SOA and formaldehyde may suggest that additional missing chemistry may be

Deleted: , detailed chemistry of oxygenated precursors,

Moved (insertion) [2]

Deleted: F

Deleted: As mobile-source emissions decline and VCP emissions become the dominant source of aerosol in urban areas, formaldehyde should be considered an important indicator for SOA production from VCPs.

4 Conclusions and future work

655 We have shown that VCPs are a major source of SOA in urban atmospheres by introducing updated emissions and VCP-
relevant chemistry into CMAQ that better represents SOA precursors emitted from these sources. This includes three new
categories of emissions: siloxanes, oxygenated IVOCs, and nonoxygenated IVOCs. VCP emissions from the VCPy
framework (Seltzer et al., 2021) were used to parameterize the new chemistry, and the mapping of VCP emitted species to
model surrogates was reviewed and updated based on species structure, volatility, and estimated SOA yield.

660 The new model chemistry and emissions inventory doubles the predicted SOA concentrations above background levels,
increasing the average daily maximum PM₁ SOA concentration by 1.4 µg m⁻³, equating to a 21% decrease in the absolute
mean bias. Most of the increased SOA mass was formed from nonoxygenated IVOC VCP precursors, followed by SOA
formed from traditional VOC precursors and oxygenated IVOC precursors, with little SOA formed from siloxanes.
665 Improvements were additionally seen in simulated formaldehyde and ozone concentrations.

Future work should consider how VCP emissions have evolved over time. VCPy version 1.0 requires information about VCP product composition and usage patterns from broad sources, including product surveys, economic statistics, and population distributions. These metrics change over time and will affect both the speciation and emission rates of organic compounds from VCPs. Diurnal and seasonal patterns of VCP emissions should also be updated to reflect more recent observations (Gkatzelis et al., 2021).

The remaining error in VCP-derived SOA predictions may reflect our lack of understanding about the oxidation pathways of low-volatility and/or oxygenated species. More information is needed about the structure, volatility, and reactivity of the products of atmospheric oxidation reactions, plus the impacts of wall loss and NO_x concentrations on SOA yields from experiments, so that models and parameterizations like the VBS can be developed. As this data become available, models can be improved to represent SOA formation from oxygenated precursors and S/IVOCs emitted from VCPs. In addition, the correlation between SOA concentration bias and temperature suggests residual model error is associated with missing sources of S/IVOC emissions, including emissions from asphalt (Khare et al., 2020), combustion sources, ~~or~~ other S/IVOCs that have large potential to form SOA. The formaldehyde bias demonstrates a similar relationship to temperature as the SOA bias, implying that investigations of formaldehyde could provide insight into VOC chemistry leading to the formation of SOA from VCPs. Including S/IVOC emissions and their atmospheric chemistry will be important for future air quality models.

Deleted: and

Deleted: can serve as a representative tracer of

Deleted: to investigate

Data Availability

CalNex observations are publicly available at <https://csl.noaa.gov/groups/csl7/measurements/2010calnex/>. The full VCPy dataset is available by downloading VCPy1.0 at <https://doi.org/10.23719/1520157>. The SAPRC07TIC_AE7I_VCP speciation profile, CMAQ chemical mechanism source code, and CMAQ output are posted at _____

Deleted: will be posted on data.gov upon final publication.

Supplement

The supplement related to this article is available online at: _____

Author Contributions

HOTP, KMS, and EAP designed the research. EAP and KMS implemented the mechanism in CMAQv5.3.2 and ran the simulations. All authors participated in data analysis and discussions. EAP drafted the paper with input from all coauthors.

Acknowledgements

The authors thank Christopher Cappa, Wyatt Appel, and Ben Schulze for modeling support and helpful discussions.

Financial Support

715 Karl Seltzer and Elyse Pennington were supported by the Oak Ridge Institute for Science and Education (ORISE) Research
Participation Program for the U.S. Environmental Protection Agency (EPA). Elyse Pennington was also supported by a
Global Research Outreach (GRO) award from the Samsung Advanced Institute of Technology (SAIT).

Competing Interests

The authors declare that they have no competing interests.

720

Disclaimer

Although this work was contributed by research staff in the Environmental Protection Agency and has been reviewed and
approved for publication, it does not reflect official policy of the EPA. The views expressed in this document are solely those
of authors and do not necessarily reflect those of the Agency. EPA does not endorse any products or commercial services
725 mentioned in this publication.

References

AE7I Species Table. (2021). U.S. Environmental Protection Agency.

[https://github.com/USEPA/CMAQ/blob/72ec4e5681e1ed0b4917792a9a240b0302a303a7/CCTM/src/MECHS/mec
hanism_information/saprc07tic_ae7i_aq/AE7I_species_table.md](https://github.com/USEPA/CMAQ/blob/72ec4e5681e1ed0b4917792a9a240b0302a303a7/CCTM/src/MECHS/mec
hanism_information/saprc07tic_ae7i_aq/AE7I_species_table.md) (Original work published 2016)

730

Appel, K. W., Bash, J. O., Fahey, K. M., Foley, K. M., Gilliam, R. C., Hogrefe, C., Hutzell, W. T., Kang, D., Mathur, R.,
Murphy, B. N., Napelenok, S. L., Nolte, C. G., Pleim, J. E., Pouliot, G. A., Pye, H. O. T., Ran, L., Roselle, S. J.,
Sarwar, G., Schwede, D. B., ... Wong, D. C. (2021). The Community Multiscale Air Quality (CMAQ) model
versions 5.3 and 5.3.1: System updates and evaluation. *Geoscientific Model Development*, *14*(5), 2867–2897.

735

<https://doi.org/10.5194/gmd-14-2867-2021>

Baker, K. R., Carlton, A. G., Kleindienst, T. E., Offenberg, J. H., Beaver, M. R., Gentner, D. R., Goldstein, A. H., Hayes, P.
L., Jimenez, J. L., Gilman, J. B., Gouw, J. A. de, Woody, M. C., Pye, H. O. T., Kelly, J. T., Lewandowski, M.,
Jaoui, M., Stevens, P. S., Brune, W. H., Lin, Y.-H., ... Surratt, J. D. (2015). Gas and aerosol carbon in California:
Comparison of measurements and model predictions in Pasadena and Bakersfield. *Atmospheric Chemistry and
Physics*, *15*(9), 5243–5258. <https://doi.org/10.5194/acp-15-5243-2015>

740

Bash, J. O., Baker, K. R., & Beaver, M. R. (2016). Evaluation of improved land use and canopy representation in BEIS
v3.61 with biogenic VOC measurements in California. *Geoscientific Model Development*, *9*(6), 2191–2207.
<https://doi.org/10.5194/gmd-9-2191-2016>

Binkowski, F. S., & Roselle, S. J. (2003). Models-3 Community Multiscale Air Quality (CMAQ) model aerosol component
1. Model description. *Journal of Geophysical Research: Atmospheres*, *108*(D6).
<https://doi.org/10.1029/2001JD001409>

745

Deleted: ¶

Deleted: ¶

Formatted: Font: 10 pt

Formatted: Line spacing: 1.5 lines

- 750 Cappa, C. D., & Wilson, K. R. (2012). Multi-generation gas-phase oxidation, equilibrium partitioning, and the formation and evolution of secondary organic aerosol. *Atmospheric Chemistry and Physics*, *12*(20), 9505–9528. <https://doi.org/10.5194/acp-12-9505-2012>
- Charan, S. M., Buenconsejo, R. S., & Seinfeld, J. H. (2020). Secondary organic aerosol yields from the oxidation of benzyl alcohol. *Atmospheric Chemistry and Physics*, *20*(21), 13167–13190. <https://doi.org/10.5194/acp-20-13167-2020>
- 755 Charan, S. M., Huang, Y., Buenconsejo, R. S., Li, Q., Cocker III, D. R., & Seinfeld, J. H. (2021). Secondary Organic Aerosol Formation from the Oxidation of Decamethylcyclopentasiloxane at Atmospherically Relevant OH Concentrations. *Atmospheric Chemistry and Physics Discussions*, 1–17. <https://doi.org/10.5194/acp-2021-353>
- Coggon, M. M., Gkatzelis, G. I., McDonald, B. C., Gilman, J. B., Schwantes, R. H., Abuhassan, N., Aikin, K. C., Arend, M. F., Berkoff, T. A., Brown, S. S., Campos, T. L., Dickerson, R. R., Gronoff, G., Hurley, J. F., Isaacman-VanWertz, G., Koss, A. R., Li, M., McKeen, S. A., Moshary, F., ... Warneke, C. (2021). Volatile chemical product emissions enhance ozone and modulate urban chemistry. *Proceedings of the National Academy of Sciences*, *118*(32). <https://doi.org/10.1073/pnas.2026653118>
- 760 Coggon, M. M., McDonald, B. C., Vlasenko, A., Veres, P. R., Bernard, F., Koss, A. R., Yuan, B., Gilman, J. B., Peischl, J., Aikin, K. C., DuRant, J., Warneke, C., Li, S.-M., & de Gouw, J. A. (2018). Diurnal Variability and Emission Pattern of Decamethylcyclopentasiloxane (D5) from the Application of Personal Care Products in Two North American Cities. *Environmental Science & Technology*, *52*(10), 5610–5618. <https://doi.org/10.1021/acs.est.8b00506>
- 765 Crounse, J. D., Nielsen, L. B., Jørgensen, S., Kjaergaard, H. G., & Wennberg, P. O. (2013). Autoxidation of Organic Compounds in the Atmosphere. *The Journal of Physical Chemistry Letters*, *4*(20), 3513–3520. <https://doi.org/10.1021/jz4019207>
- 770 Donahue, N. M., Epstein, S. A., Pandis, S. N., & Robinson, A. L. (2011). A two-dimensional volatility basis set: 1. organic-aerosol mixing thermodynamics. *Atmospheric Chemistry and Physics*, *11*(7), 3303–3318. <https://doi.org/10.5194/acp-11-3303-2011>
- 775 Ensberg, J. J., Hayes, P. L., Jimenez, J. L., Gilman, J. B., Kuster, W. C., de Gouw, J. A., Holloway, J. S., Gordon, T. D., Jathar, S., Robinson, A. L., & Seinfeld, J. H. (2014). Emission factor ratios, SOA mass yields, and the impact of vehicular emissions on SOA formation. *Atmospheric Chemistry and Physics*, *14*(5), 2383–2397. <https://doi.org/10.5194/acp-14-2383-2014>
- Epstein, S. A., Riipinen, I., & Donahue, N. M. (2010). A Semiempirical Correlation between Enthalpy of Vaporization and Saturation Concentration for Organic Aerosol. *Environmental Science & Technology*, *44*(2), 743–748. <https://doi.org/10.1021/es902497z>
- 780 Gkatzelis, G. I., Coggon, M. M., McDonald, B. C., Peischl, J., Aikin, K. C., Gilman, J. B., Trainer, M., & Warneke, C. (2021). Identifying Volatile Chemical Product Tracer Compounds in U.S. Cities. *Environmental Science & Technology*, *55*(1), 188–199. <https://doi.org/10.1021/acs.est.0c05467>

Deleted: Carter, W. P. L. (2010). Development of the SAPRC-07 chemical mechanism. *Atmospheric Environment*, *44*(40), 5324–5335. <https://doi.org/10.1016/j.atmosenv.2010.01.026>

- Goldstein, A. H., & Galbally, I. E. (2007). Known and Unexplored Organic Constituents in the Earth's Atmosphere. *Environmental Science & Technology*, 41(5), 1514–1521. <https://doi.org/10.1021/es072476p>
- Griffith, S. M., Hansen, R. F., Dusanter, S., Michoud, V., Gilman, J. B., Kuster, W. C., Veres, P. R., Graus, M., Gouw, J. A. de, Roberts, J., Young, C., Washenfelder, R., Brown, S. S., Thalman, R., Waxman, E., Volkamer, R., Tsai, C., Stutz, J., Flynn, J. H., ... Stevens, P. S. (2016). Measurements of hydroxyl and hydroperoxy radicals during CalNex-LA: Model comparisons and radical budgets. *Journal of Geophysical Research: Atmospheres*, 121(8), 4211–4232. <https://doi.org/10.1002/2015JD024358>
- Hayes, P. L., Carlton, A. G., Baker, K. R., Ahmadov, R., Washenfelder, R. A., Alvarez, S., Rappenglück, B., Gilman, J. B., Kuster, W. C., de Gouw, J. A., Zotter, P., Prévôt, A. S. H., Szidat, S., Kleindienst, T. E., Offenberg, J. H., Ma, P. K., & Jimenez, J. L. (2015). Modeling the formation and aging of secondary organic aerosols in Los Angeles during CalNex 2010. *Atmospheric Chemistry and Physics*, 15(10), 5773–5801. <https://doi.org/10.5194/acp-15-5773-2015>
- Hayes, P. L., Ortega, A. M., Cubison, M. J., Froyd, K. D., Zhao, Y., Cliff, S. S., Hu, W. W., Toohey, D. W., Flynn, J. H., Lefer, B. L., Grossberg, N., Alvarez, S., Rappenglück, B., Taylor, J. W., Allan, J. D., Holloway, J. S., Gilman, J. B., Kuster, W. C., Gouw, J. A. de, ... Jimenez, J. L. (2013). Organic aerosol composition and sources in Pasadena, California, during the 2010 CalNex campaign. *Journal of Geophysical Research: Atmospheres*, 118(16), 9233–9257. <https://doi.org/10.1002/jgrd.50530>
- Hodzic, A., Aumont, B., Knote, C., Lee-Taylor, J., Madronich, S., & Tyndall, G. (2014). Volatility dependence of Henry's law constants of condensable organics: Application to estimate depositional loss of secondary organic aerosols. *Geophysical Research Letters*, 41(13), 4795–4804. <https://doi.org/10.1002/2014GL060649>
- Huffman, J. A., Docherty, K. S., Mohr, C., Cubison, M. J., Ulbrich, I. M., Ziemann, P. J., Onasch, T. B., & Jimenez, J. L. (2009). Chemically-Resolved Volatility Measurements of Organic Aerosol from Different Sources. *Environmental Science & Technology*, 43(14), 5351–5357. <https://doi.org/10.1021/es803539d>
- Hyslop, N. P. (2009). Impaired visibility: The air pollution people see. *Atmospheric Environment*, 43(1), 182–195. <https://doi.org/10.1016/j.atmosenv.2008.09.067>
- Intergovernmental Panel on Climate Change (Ed.). (2014). Anthropogenic and Natural Radiative Forcing. In *Climate Change 2013 – The Physical Science Basis: Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (pp. 659–740). Cambridge University Press. <https://doi.org/10.1017/CBO9781107415324.018>
- Janecek, N. J., Hansen, K. M., & Stanier, C. O. (2017). Comprehensive atmospheric modeling of reactive cyclic siloxanes and their oxidation products. *Atmospheric Chemistry and Physics*, 17(13), 8357–8370. <https://doi.org/10.5194/acp-17-8357-2017>
- Janecek, N. J., Marek, R. F., Bryngelson, N., Singh, A., Bullard, R. L., Brune, W. H., & Stanier, C. O. (2019). Physical properties of secondary photochemical aerosol from OH oxidation of a cyclic siloxane. *Atmospheric Chemistry and Physics*, 19(3), 1649–1664. <https://doi.org/10.5194/acp-19-1649-2019>

- 820 Jathar, S. H., Gordon, T. D., Hennigan, C. J., Pye, H. O. T., Pouliot, G., Adams, P. J., Donahue, N. M., & Robinson, A. L. (2014). Unspeciated organic emissions from combustion sources and their influence on the secondary organic aerosol budget in the United States. *Proceedings of the National Academy of Sciences*, *111*(29), 10473–10478. <https://doi.org/10.1073/pnas.1323740111>
- Jathar, S. H., Woody, M., Pye, H. O. T., Baker, K. R., & Robinson, A. L. (2017). Chemical transport model simulations of organic aerosol in southern California: Model evaluation and gasoline and diesel source contributions. *Atmospheric Chemistry and Physics*, *17*(6), 4305–4318. <https://doi.org/10.5194/acp-17-4305-2017>
- 825 Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., ... Worsnop, D. R. (2009). Evolution of Organic Aerosols in the Atmosphere. *Science*, *326*(5959), 1525–1529. <https://doi.org/10.1126/science.1180353>
- 830 [Kaiser, J., Wolfe, G. M., Bohn, B., Broch, S., Fuchs, H., Ganzeveld, L. N., Gomm, S., Häsel, R., Hofzumahaus, A., Holland, F., Jäger, J., Li, X., Lohse, I., Lu, K., Prévôt, A. S. H., Rohrer, F., Wegener, R., Wolf, R., Mentel, T. F., ... Keutsch, F. N. \(2015\). Evidence for an unidentified non-photochemical ground-level source of formaldehyde in the Po Valley with potential implications for ozone production. *Atmospheric Chemistry and Physics*, *15*\(3\), 1289–1298. <https://doi.org/10.5194/acp-15-1289-2015>](https://doi.org/10.5194/acp-15-1289-2015)
- 835 Khare, P., & Gentner, D. R. (2018). Considering the future of anthropogenic gas-phase organic compound emissions and the increasing influence of non-combustion sources on urban air quality. *Atmospheric Chemistry and Physics*, *18*(8), 5391–5413. <https://doi.org/10.5194/acp-18-5391-2018>
- Khare, P., Machesky, J., Soto, R., He, M., Presto, A. A., & Gentner, D. R. (2020). Asphalt-related emissions are a major missing nontraditional source of secondary organic aerosol precursors. *Science Advances*, *6*(36), eabb9785. <https://doi.org/10.1126/sciadv.abb9785>
- 840 Levelt, P. F., Joiner, J., Tamminen, J., Veefkind, J. P., Bhartia, P. K., Stein Zweers, D. C., Duncan, B. N., Streets, D. G., Eskes, H., van der A, R., McLinden, C., Fioletov, V., Carn, S., de Laat, J., DeLand, M., Marchenko, S., McPeters, R., Ziemke, J., Fu, D., ... Wargan, K. (2018). The Ozone Monitoring Instrument: Overview of 14 years in space. *Atmospheric Chemistry and Physics*, *18*(8), 5699–5745. <https://doi.org/10.5194/acp-18-5699-2018>
- 845 Li, L., & Cocker, D. R. (2018). Molecular structure impacts on secondary organic aerosol formation from glycol ethers. *Atmospheric Environment*, *180*, 206–215. <https://doi.org/10.1016/j.atmosenv.2017.12.025>
- Li, W., Li, L., Chen, C., Kacarab, M., Peng, W., Price, D., Xu, J., & Cocker, D. R. (2018). Potential of select intermediate-volatility organic compounds and consumer products for secondary organic aerosol and ozone formation under relevant urban conditions. *Atmospheric Environment*, *178*, 109–117. <https://doi.org/10.1016/j.atmosenv.2017.12.019>
- 850 Li, Y., Rodier, C., Lea, J. D., Harvey, J., & Kleeman, M. J. (2021). Improving spatial surrogates for area source emissions inventories in California. *Atmospheric Environment*, *247*, 117665. <https://doi.org/10.1016/j.atmosenv.2020.117665>

- 855 Liao, J., Hanisco, T. F., Wolfe, G. M., St. Clair, J., Jimenez, J. L., Campuzano-Jost, P., Nault, B. A., Fried, A., Marais, E. A.,
Gonzalez Abad, G., Chance, K., Jethva, H. T., Ryerson, T. B., Warneke, C., & Wisthaler, A. (2019). Towards a
satellite formaldehyde – in situ hybrid estimate for organic aerosol abundance. *Atmospheric Chemistry and Physics*,
19(5), 2765–2785. <https://doi.org/10.5194/acp-19-2765-2019>
- Lim, S. S., Vos, T., Flaxman, A. D., Danaei, G., Shibuya, K., Adair-Rohani, H., AlMazroa, M. A., Amann, M., Anderson, H.
860 R., Andrews, K. G., Aryee, M., Atkinson, C., Bacchus, L. J., Bahalim, A. N., Balakrishnan, K., Balmes, J., Barker-
Collo, S., Baxter, A., Bell, M. L., ... Ezzati, M. (2012). A comparative risk assessment of burden of disease and
injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: A systematic analysis for the
Global Burden of Disease Study 2010. *The Lancet*, 380(9859), 2224–2260. [https://doi.org/10.1016/S0140-6736\(12\)61766-8](https://doi.org/10.1016/S0140-6736(12)61766-8)
- Lu, Q., Murphy, B. N., Qin, M., Adams, P. J., Zhao, Y., Pye, H. O. T., Efstathiou, C., Allen, C., & Robinson, A. L. (2020).
865 Simulation of organic aerosol formation during the CalNex study: Updated mobile emissions and secondary organic
aerosol parameterization for intermediate-volatility organic compounds. *Atmospheric Chemistry and Physics*, 20(7),
4313–4332. <https://doi.org/10.5194/acp-20-4313-2020>
- Luecken, D. J., Napelenok, S. L., Strum, M., Scheffe, R., & Phillips, S. (2018). Sensitivity of Ambient Atmospheric
870 Formaldehyde and Ozone to Precursor Species and Source Types Across the United States. *Environmental Science
& Technology*, 52(8), 4668–4675. <https://doi.org/10.1021/acs.est.7b05509>
- McDonald, B. C., Gouw, J. A. de, Gilman, J. B., Jathar, S. H., Akherati, A., Cappa, C. D., Jimenez, J. L., Lee-Taylor, J.,
Hayes, P. L., McKeen, S. A., Cui, Y. Y., Kim, S.-W., Gentner, D. R., Isaacman-VanWertz, G., Goldstein, A. H.,
Harley, R. A., Frost, G. J., Roberts, J. M., Ryerson, T. B., & Trainer, M. (2018). Volatile chemical products
emerging as largest petrochemical source of urban organic emissions. *Science*, 359(6377), 760–764.
875 <https://doi.org/10.1126/science.aaq0524>
- Milani, A., Al-Naiema, I. M., & Stone, E. A. (2021). Detection of a secondary organic aerosol tracer derived from personal
care products. *Atmospheric Environment*, 246, 118078. <https://doi.org/10.1016/j.atmosenv.2020.118078>
- Murphy, B. N., Nolte, C. G., Sidi, F., Bash, J. O., Appel, K. W., Jang, C., Kang, D., Kelly, J., Mathur, R., Napelenok, S.,
880 Pouliot, G., & Pye, H. O. T. (2021). The Detailed Emissions Scaling, Isolation, and Diagnostic (DESID) module in
the Community Multiscale Air Quality (CMAQ) modeling system version 5.3.2. *Geoscientific Model Development*,
14(6), 3407–3420. <https://doi.org/10.5194/gmd-14-3407-2021>
- Murphy, B. N., Woody, M. C., Jimenez, J. L., Carlton, A. M. G., Hayes, P. L., Liu, S., Ng, N. L., Russell, L. M., Setyan, A.,
Xu, L., Young, J., Zaveri, R. A., Zhang, Q., & Pye, H. O. T. (2017). Semivolatile POA and parameterized total
combustion SOA in CMAQv5.2: Impacts on source strength and partitioning. *Atmospheric Chemistry and Physics*,
885 17(18), 11107–11133. <https://doi.org/10.5194/acp-17-11107-2017>
- Nussbaumer, C. M., & Cohen, R. C. (2021). Impact of OA on the Temperature Dependence of PM 2.5 in the Los Angeles
Basin. *Environmental Science & Technology*, 55(6), 3549–3558. <https://doi.org/10.1021/acs.est.0c07144>

- Pye, H. O. T., Murphy, B. N., Xu, L., Ng, N. L., Carlton, A. G., Guo, H., Weber, R., Vasilakos, P., Appel, K. W., Budisulistiorini, S. H., Surratt, J. D., Nenes, A., Hu, W., Jimenez, J. L., Isaacman-VanWertz, G., Misztal, P. K., & Goldstein, A. H. (2017). On the implications of aerosol liquid water and phase separation for organic aerosol mass. *Atmospheric Chemistry and Physics*, *17*(1), 343–369. <https://doi.org/10.5194/acp-17-343-2017>
- Pye, H. O. T., Pinder, R. W., Piletic, I. R., Xie, Y., Capps, S. L., Lin, Y.-H., Surratt, J. D., Zhang, Z., Gold, A., Luecken, D. J., Hutzell, W. T., Jaoui, M., Offenberg, J. H., Kleindienst, T. E., Lewandowski, M., & Edney, E. O. (2013). Epoxide Pathways Improve Model Predictions of Isoprene Markers and Reveal Key Role of Acidity in Aerosol Formation. *Environmental Science & Technology*, *47*(19), 11056–11064. <https://doi.org/10.1021/es402106h>
- Qin, M., Murphy, B. N., Isaacs, K. K., McDonald, B. C., Lu, Q., McKeen, S. A., Koval, L., Robinson, A. L., Efstathiou, C., Allen, C., & Pye, H. O. T. (2021). Criteria pollutant impacts of volatile chemical products informed by near-field modelling. *Nature Sustainability*, *4*(2), 129–137. <https://doi.org/10.1038/s41893-020-00614-1>
- Robinson, A. L., Donahue, N. M., Shrivastava, M. K., Weitkamp, E. A., Sage, A. M., Grieshop, A. P., Lane, T. E., Pierce, J. R., & Pandis, S. N. (2007). Rethinking Organic Aerosols: Semivolatile Emissions and Photochemical Aging. *Science*, *315*(5816), 1259–1262. <https://doi.org/10.1126/science.1133061>
- Ryerson, T. B., Andrews, A. E., Angevine, W. M., Bates, T. S., Brock, C. A., Cairns, B., Cohen, R. C., Cooper, O. R., Gouw, J. A. de, Fehsenfeld, F. C., Ferrare, R. A., Fischer, M. L., Flagan, R. C., Goldstein, A. H., Hair, J. W., Hardesty, R. M., Hostetler, C. A., Jimenez, J. L., Langford, A. O., ... Wofsy, S. C. (2013). The 2010 California Research at the Nexus of Air Quality and Climate Change (CalNex) field study. *Journal of Geophysical Research: Atmospheres*, *118*(11), 5830–5866. <https://doi.org/10.1002/jgrd.50331>
- Santoni, G. W., Daube, B. C., Kort, E. A., Jiménez, R., Park, S., Pittman, J. V., Gottlieb, E., Xiang, B., Zahniser, M. S., Nelson, D. D., McManus, J. B., Peischl, J., Ryerson, T. B., Holloway, J. S., Andrews, A. E., Sweeney, C., Hall, B., Hints, E. J., Moore, F. L., ... Wofsy, S. C. (2014). Evaluation of the airborne quantum cascade laser spectrometer (QCLS) measurements of the carbon and greenhouse gas suite – CO₂, CH₄, N₂O, and CO – during the CalNex and HIPPO campaigns. *Atmospheric Measurement Techniques*, *7*(6), 1509–1526. <https://doi.org/10.5194/amt-7-1509-2014>
- Seinfeld, J. H., & Pandis, S. N. (2016). *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change* (3rd ed.). John Wiley & Sons, Inc.
- Seltzer, K. M., Pennington, E., Rao, V., Murphy, B. N., Strum, M., Isaacs, K. K., & Pye, H. O. T. (2021). Reactive organic carbon emissions from volatile chemical products. *Atmospheric Chemistry and Physics*, *21*(6), 5079–5100. <https://doi.org/10.5194/acp-21-5079-2021>
- Shah, R. U., Coggon, M. M., Gkatzelis, G. I., McDonald, B. C., Tasoglou, A., Huber, H., Gilman, J., Warneke, C., Robinson, A. L., & Presto, A. A. (2020). Urban Oxidation Flow Reactor Measurements Reveal Significant Secondary Organic Aerosol Contributions from Volatile Emissions of Emerging Importance. *Environmental Science & Technology*, *54*(2), 714–725. <https://doi.org/10.1021/acs.est.9b06531>

- Shah, T., Shi, Y., Beardsley, R., & Yarwood, G. (2020). *Speciation Tool User's Guide Version 5.0*. Ramboll US Corporation.
- Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., & Barker, D. (2008). *A Description of the Advanced Research WRF Version 3* (NCAR/TN-475+STR). University Corporation for Atmospheric Research.
925 <http://dx.doi.org/10.5065/D68S4MVH>
- US EPA, O. (2013, August 1). *Air Quality System (AQS)* [Data and Tools]. US EPA. <https://www.epa.gov/aqs>
- US EPA, O. (2015, June 2). *National Emissions Inventory (NEI)* [Other Policies and Guidance]. US EPA.
<https://www.epa.gov/air-emissions-inventories/national-emissions-inventory-nei>
- 930 US EPA Office of Research and Development. (2020). *CMAQ*. Zenodo. <https://doi.org/10.5281/zenodo.4081737>
- Wang, D.-G., Norwood, W., Alaei, M., Byer, J. D., & Brimble, S. (2013). Review of recent advances in research on the toxicity, detection, occurrence and fate of cyclic volatile methyl siloxanes in the environment. *Chemosphere*, *93*(5), 711–725. <https://doi.org/10.1016/j.chemosphere.2012.10.041>
- Wang, Y., Wang, H., Tan, Y., Liu, J., Wang, K., Ji, W., Sun, L., Yu, X., Zhao, J., Xu, B., & Xiong, J. (2021). Measurement of the key parameters of VOC emissions from wooden furniture, and the impact of temperature. *Atmospheric Environment*, 118510. <https://doi.org/10.1016/j.atmosenv.2021.118510>
- 935 Warneke, C., Veres, P., Holloway, J. S., Stutz, J., Tsai, C., Alvarez, S., Rappenglueck, B., Fehsenfeld, F. C., Graus, M., Gilman, J. B., & de Gouw, J. A. (2011). Airborne formaldehyde measurements using PTR-MS: Calibration, humidity dependence, inter-comparison and initial results. *Atmospheric Measurement Techniques*, *4*(10), 2345–2358. <https://doi.org/10.5194/amt-4-2345-2011>
- 940 Woody, M. C., Baker, K. R., Hayes, P. L., Jimenez, J. L., Koo, B., & Pye, H. O. T. (2016). Understanding sources of organic aerosol during CalNex-2010 using the CMAQ-VBS. *Atmospheric Chemistry and Physics*, *16*(6), 4081–4100. <https://doi.org/10.5194/acp-16-4081-2016>
- Wu, Y., & Johnston, M. V. (2017). Aerosol Formation from OH Oxidation of the Volatile Cyclic Methyl Siloxane (cVMS) Decamethylcyclopentasiloxane. *Environmental Science & Technology*, *51*(8), 4445–4451. <https://doi.org/10.1021/acs.est.7b00655>
- 945 Xie, Y., Paulot, F., Carter, W. P. L., Nolte, C. G., Luecken, D. J., Hutzell, W. T., Wennberg, P. O., Cohen, R. C., & Pinder, R. W. (2013). Understanding the impact of recent advances in isoprene photooxidation on simulations of regional air quality. *Atmospheric Chemistry and Physics*, *13*(16), 8439–8455. <https://doi.org/10.5194/acp-13-8439-2013>
- 950 Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami, A., Middlebrook, A. M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P. F., Salcedo, D., Onasch, T., Jayne, J. T., Miyoshi, T., Shimojo, A., ... Worsnop, D. R. (2007). Ubiquity and dominance of oxygenated species in organic aerosols in anthropogenically-influenced Northern Hemisphere midlatitudes. *Geophysical Research Letters*, *34*(13). <https://doi.org/10.1029/2007GL029979>

- 955 Zhao, Y., Hennigan, C. J., May, A. A., Tkacik, D. S., de Gouw, J. A., Gilman, J. B., Kuster, W. C., Borbon, A., & Robinson, A. L. (2014). Intermediate-Volatility Organic Compounds: A Large Source of Secondary Organic Aerosol. *Environmental Science & Technology*, *48*(23), 13743–13750. <https://doi.org/10.1021/es5035188>
- Zhao, Y., Nguyen, N. T., Presto, A. A., Hennigan, C. J., May, A. A., & Robinson, A. L. (2015). Intermediate Volatility Organic Compound Emissions from On-Road Diesel Vehicles: Chemical Composition, Emission Factors, and Estimated Secondary Organic Aerosol Production. *Environmental Science & Technology*, *49*(19), 11516–11526. <https://doi.org/10.1021/acs.est.5b02841>
- 960

Formatted: Line spacing: 1.5 lines