



# Impact of non-ideality on reconstructing spatial and temporal variations of aerosol acidity with multiphase buffer theory

Guangjie Zheng<sup>1</sup>, Hang Su<sup>2</sup>, Siwen Wang<sup>2</sup>, Andrea Pozzer<sup>3</sup>, Yafang Cheng<sup>1\*</sup>

- 5 <sup>1</sup> Minerva Research Group, Max Planck Institute for Chemistry, Mainz 55128, Germany.  
<sup>2</sup> Multiphase Chemistry Department, Max Planck Institute for Chemistry, Mainz 55128, Germany.  
<sup>3</sup> Air Chemistry Department, Max Planck Institute for Chemistry, Mainz 55128, Germany.

Correspondence to: Y. Cheng ([yafang.cheng@mpic.de](mailto:yafang.cheng@mpic.de))

10 **Abstract.** Aerosol acidity is a key parameter in atmospheric aqueous chemistry and strongly influence the interactions of air pollutants and ecosystem. The recently proposed multiphase buffer theory provides a framework to reconstruct long-term trends and spatial variations of aerosol pH based on the effective acid dissociation constant of ammonia ( $K_{a,NH_3}^*$ ). However, non-ideality in aerosol droplets is a major challenge limiting its broad applications. Here, we introduced a non-ideality correction factor ( $c_{ni}$ ) and investigated its governing factors. We found that besides relative humidity (RH) and temperature,  $c_{ni}$  is mainly determined by the molar fraction of  $NO_3^-$  in aqueous-phase anions, due to different  $NH_4^+$  activity coefficients between  $(NH_4)_2SO_4$ - and  $NH_4NO_3$ -dominated aerosols. A parameterization method is thus proposed to estimate  $c_{ni}$  at given RH, temperature and  $NO_3^-$  fraction, and is validated against long-term observations and global simulations. In the ammonia-buffered regime, with  $c_{ni}$  correction the buffer theory can well reproduce the  $K_{a,NH_3}^*$  predicted by comprehensive thermodynamic models, with root-mean-square deviation  $\sim 0.1$  and correlation coefficient  $\sim 1$ . Note that, while  $c_{ni}$  is needed to predict  $K_{a,NH_3}^*$  levels, it is usually not the dominant contributor to its variations, as  $\sim 90\%$  of the temporal or spatial variations in  $K_{a,NH_3}^*$  is due to variations in aerosol water and temperature.

15  
20



## 1. Introduction

Aerosol acidity strongly influences the thermodynamics and chemical kinetics of atmospheric aerosols and is therefore one essential parameter in evaluating their environmental, health and climate effects (Pye et al., 2020; Zheng et al., 2020). However, direct measurements of aerosol pH in the real atmosphere are not available so far (Pye et al., 2020; Li et al., 2020). The fast equilibrium with ambient air, tiny volume and high ionic strength and nucleation potential are the main challenges for measurements, especially online or in-situ measurements. Several groups are developing new techniques for this purpose (Wei et al., 2018; Craig et al., 2018; Li et al., 2020; Ault, 2020). For example, Wei et al. (2018) developed an in-situ Raman microscopy method for pH measurements in microdroplets (diameter  $\sim 20 \mu\text{m}$ ), with an uncertainty of  $\sim 0.5$  pH units. Craig et al. (2018) and Li et al. (2020) developed colorimetric analyses on pH-indicator papers for aerosol pH measurement, which exhibit uncertainties around 0.4-0.5 pH units. These currently available techniques, however, still need to be developed further for real atmospheric applications.

Due to the lack of direct measurements, modelling tools have been intensively used to calculate the aerosol pH (Fountoukis and Nenes, 2007; Fountoukis et al., 2009; Clegg et al., 2001; Zuend et al., 2008). Results of thermodynamic models are subject to uncertainties in the input parameters (Fountoukis et al., 2009; Pye et al., 2020; Weber et al., 2016; Guo et al., 2016; Guo et al., 2017; Pye et al., 2018; Tao and Murphy, 2019; Hennigan et al., 2015; Guo et al., 2015; Peng et al., 2019). For example, Hennigan et al. (2015) revealed the importance of including gas-phase species in the input, in addition to the full aerosol composition measurements (Fountoukis et al., 2009). Guo et al. (2015) suggests overall uncertainties of  $\sim 0.2$ -0.5 pH units related to aerosol composition. Pye et al. (2020) reviewed major thermodynamic models and show that the estimated acidity among different models were on average 0.3 pH units, but sometimes as much as 1 pH unit.

The recently proposed multiphase buffer theory shows that globally most of the populated urban areas are within the multiphase ammonia-buffered regime (Zheng et al., 2020). In the buffered regions/periods,  $\text{p}K_{\text{a},\text{NH}_3}^*$  can serve as a proxy of aerosol pH, where  $K_{\text{a},\text{NH}_3}^*$  is the effective acid dissociation constant of  $\text{NH}_3$  in multiphase systems (section 2). Ideally,  $\text{p}K_{\text{a},\text{NH}_3}^*$  is fully determined by aerosol water content (AWC) and temperature. However, the non-ideality in aerosols may introduce deviations from the ideal conditions. Here we investigated such deviation and derived a non-ideality correction factor for using  $\text{p}K_{\text{a},\text{NH}_3}^*$  as a proxy of aerosol pH. Governing factors of the non-ideality correction factor in aerosol droplet are further explored and discussed, based on which a parameterization method to estimate the non-ideality correction factors are proposed. We also estimated that a constant correction factor of  $\text{p}K_{\text{a},\text{NH}_3}^*$  is often good enough to predict pH over a period at a given site, or to explain the global pH variations. We thereby provided a way for pH retrieval when chemical measurements are unavailable for the ammonia-buffered regions and periods.




## 2. Methods

### 2.1 Effective acid dissociation constant as a proxy of aerosol pH

#### Acid dissociation constant of NH<sub>3</sub> in bulk solutions, $K_a$

55 The definition of acids and bases have been evolving over time (Zheng et al., 2020). The pioneering Arrhenius theory defined base as a substance that dissociates in water to form hydroxide (OH<sup>-</sup>) ions (Pfenning, 2015). Therefore, an Arrhenius base can be expressed as B<sub>A</sub>OH, which dissociation in water as:




with the corresponding base dissociation constant  $K_b$  being:

60 
$$K_b = [B_A^+] [OH^-] / [B_AOH] \quad (1b)$$

In combination with the water dissociation of:



The corresponding acid dissociation constant, defined as  $K_a = K_w / K_b$ , is thus (reaction 2 – reaction 1):



65 The later Bronsted-Lowry theory defined base as proton acceptor (Pfenning, 2015), and is expressed as B<sub>BL</sub> here. In this sense, an Arrhenius base B<sub>A</sub>OH is not considered as a Brønsted base, but rather salts. The dissociation reaction for a Brønsted base is expressed as:



and the corresponding  $K_a$  is thus (reaction 2 – reaction 4):

70 
$$B_{BL}H^+ \rightleftharpoons B_{BL} + H^+, \quad K_a = [B_{BL}] [H^+] / [B_{BL}H^+] \quad (5)$$

As NH<sub>3</sub>(aq) is actually the water adduct of NH<sub>3</sub>, it is often be expressed equivalently as NH<sub>3</sub>(aq) = NH<sub>3</sub>·H<sub>2</sub>O(aq) = NH<sub>4</sub>OH(aq). In this sense, it can fit in the category of both definitions. In the Arrhenius definition, the base B<sub>A</sub>OH = NH<sub>4</sub>OH, namely B<sub>A</sub> = NH<sub>4</sub><sup>+</sup>. Therefore,  $K_{a,NH_3}$  is (Eq. 3):

$$K_{a,NH_3} = [NH_4OH(aq)] [H^+] / [NH_4^+] = [NH_3(aq)] [H^+] / [NH_4^+] \quad (6)$$

75 While with the Bronsted-Lowry definition, the base is B<sub>BL</sub> = NH<sub>3</sub>(aq), and  $K_{a,NH_3}$  is (Eq. 5):

$$K_{a,NH_3} = [NH_3(aq)] [H^+] / [NH_4^+] \quad (7)$$

which is the same as Eq. 6. Therefore, different definition of bases for the ammonia family (B<sub>A</sub> = NH<sub>4</sub><sup>+</sup> or B<sub>BL</sub> = NH<sub>3</sub>(aq)) will led to the same expression of  $K_{a,NH_3}$ , as defined in Zheng et al. (2020). The same applies for other volatile weak bases.



### Ideal multiphase acid dissociation constant of NH<sub>3</sub>

- 80 The multiphase effective acid dissociation constant of NH<sub>3</sub> under ideal conditions,  $K_{a,NH_3}^{*,i}$ , depends only on AWC and temperature as (Zheng et al., 2020):

$$K_{a,NH_3}^{*,i} = \frac{[H^+(aq)][NH_3(aq)] + [NH_3(g)]}{[NH_4^+(aq)]} = K_{a,NH_3} \left( 1 + \frac{\rho_w}{H_{NH_3} R T AWC} \right)$$

- where AWC is mainly determined by air particulate matter concentrations and RH. The [NH<sub>3</sub>(g)] represents equivalent molality of gaseous NH<sub>3</sub> in solution (see details in Zheng et al. (2020)). The  $H_{NH_3}$  is Henry's law constant of NH<sub>3</sub> in mol L<sup>-1</sup> atm<sup>-1</sup>,  $R$  is the gas constant of 0.08205 atm L mol<sup>-1</sup> K<sup>-1</sup>,  $T$  is temperature in K, AWC is in μg m<sup>-3</sup>, and  $\rho_w$  is water density in μg m<sup>-3</sup>.

For typical ambient conditions when AWC varies between 1 to 1000 μg m<sup>-3</sup>, the [NH<sub>3</sub>(g)] is usually 10<sup>5</sup> to 10<sup>8</sup> times larger than [NH<sub>3</sub>(aq)], and the above equation can be simplified into:

$$K_{a,NH_3}^{*,i} = \frac{[H^+(aq)][NH_3(g)]}{[NH_4^+(aq)]} = K_{a,NH_3} \frac{\rho_w}{H_{NH_3} R T AWC} \quad (8a)$$

- 90 And taking negative lognormal on both sides, we have pH is related to  $pK_{a,NH_3}^{*,i}$  (i.e.,  $-\log K_{a,NH_3}^{*,i}$ ) as (Zheng et al., 2020):

$$pH = pK_{a,NH_3}^{*,i} + \log \frac{[NH_3(g)]}{[NH_4^+(aq)]} \quad (8b)$$

### 2.2 Influences of non-ideality on aerosol pH

For ambient aerosols, the ionic strength ( $I$ ) is high, and the non-ideality must be considered. Under such non-ideal conditions, the multiphase equilibrium of NH<sub>3</sub> can be expressed as (Zheng et al., 2020):

$$95 \quad K_{a,NH_3}^{*,i} = \frac{(\gamma_{NH_3(g)}[NH_3(g)] + \gamma_{NH_3(aq)}[NH_3(aq)])(\gamma_{H^+}[H^+(aq)])}{\gamma_{NH_4^+}[NH_4^+(aq)]} \quad (9)$$

where  $\gamma_X$  is the activity coefficient for species X.

Activity coefficients for gases, like  $\gamma_{NH_3(g)}$ , are usually treated as unity. Again, for typical ambient conditions [NH<sub>3</sub>(g)] is much larger than [NH<sub>3</sub>(aq)], and  $\gamma_{NH_3(aq)}[NH_3(aq)]$  can be omitted. Eq. 9 can thus be simplified into:

$$K_{a,NH_3}^{*,i} = \frac{\gamma_{H^+}}{\gamma_{NH_4^+}} \frac{[H^+(aq)][NH_3(g)]}{[NH_4^+(aq)]} \quad (10)$$

- 100 Under non-ideal conditions, pH is usually defined by the proton activity, i.e.:

$$pH_a = -\log(\gamma_{H^+}[H^+]) \quad (11)$$

However, in thermodynamic models that are most commonly applied in current global models (ISORROPIA II, MOSAIC, etc.), the pH is usually defined as free-H<sup>+</sup> molality (Pye et al., 2020), i.e.:

$$pH = pH_F = -\log([H^+]) \quad (12)$$



105 The difference of activity- and molality-defined pH (i.e.,  $\text{pH}_a$  and  $\text{pH}_F$ ) is discussed in a previous study (Pye et al., 2020), which show that deviations of  $\text{pH}_F$  from  $\text{pH}_a$  is larger at lower RH, and is usually within 1 unit when  $\text{RH} > 60\%$  (Pye et al., 2020). To be comparable with results in previous studies, the pH we discussed hereinafter follow the free- $\text{H}^+$  molality definition. Discussion based on activity-defined pH is detailed in Appendices A and B.

With the free- $\text{H}^+$  molality pH definition, multiphase buffer theory can be rewritten as (Eqs. 10 and 12):

$$110 \quad K_{a,\text{NH}_3}^{*,ni} = \frac{[\text{NH}_3(\text{g})][\text{H}^+(\text{aq})]}{[\text{NH}_4^+(\text{aq})]} = \frac{\gamma_{\text{NH}_4^+}}{\gamma_{\text{H}^+}} K_{a,\text{NH}_3}^{*,i} = \frac{\gamma_{\text{NH}_4^+}}{\gamma_{\text{H}^+}} K_{a,\text{NH}_3} \frac{\rho_w}{H_{\text{NH}_3} R T \text{ AWC}} \quad (13a)$$

$$\text{pH} = -\log([\text{H}^+(\text{aq})]) = \text{p}K_{a,\text{NH}_3}^{*,ni} + \log \frac{[\text{NH}_3(\text{g})]}{[\text{NH}_4^+(\text{aq})]} \quad (13b)$$

where  $K_a^{*,ni}$  is the multiphase effective acid dissociation constant under non-ideal conditions. The difference of pH caused by non-ideality is therefore (Eqs. 8 and 13):

$$c_{ni} = \text{p}K_a^{*,ni} - \text{p}K_a^{*,i} = -\log \frac{\gamma_{\text{NH}_4^+}}{\gamma_{\text{H}^+}} \quad (14a)$$

115 where  $c_{ni}$  is hereinafter denoted as the non-ideality correction factor.

Another way to calculate  $c_{ni}$  is by definition, i.e.:

$$c_{ni} = \text{p}K_a^{*,ni} - \text{p}K_a^{*,i} = -\log \left( \frac{[\text{NH}_3(\text{g})][\text{H}^+(\text{aq})]}{[\text{NH}_4^+(\text{aq})]} \right) + \log \left( K_{a,\text{NH}_3} \frac{\rho_w}{H_{\text{NH}_3} R T \text{ AWC}} \right) \quad (14b)$$

For some thermodynamics models that predict both the activity coefficients of ions and the gas-particle partitioning of species like the E-AIM model (section 2.3),  $c_{ni}$  can be derived either from Eq. 14a (activity-based) or Eq. 14b (gas-particle partitioning based). However, current atmospheric chemical transport models usually adopted the more computation-efficient thermodynamic models (ISORROPIA II, MOSAIC, etc.), in which only the mean activity coefficient of an electrolyte species  $ij$  in water,  $\gamma_{ij}$ , are derived, where  $i$  is a cation while  $j$  is an anion (Pye et al., 2020; Fountoukis and Nenes, 2007; Zaveri et al., 2005). For these models, we cannot directly derive  $\gamma_{\text{NH}_4^+}$  or  $\gamma_{\text{H}^+}$ , and  $c_{ni}$  are derived through Eq. 14b (i.e., from the predicted  $[\text{NH}_3]$ ,  $[\text{NH}_4^+]$ ,  $[\text{H}^+]$ , and AWC).

## 125 2.3 Model simulations

**Thermodynamic models.** Here we used E-AIM model (model IV; <http://www.aim.env.uea.ac.uk/aim/aim.php>) (Clegg et al., 1992b; Wexler and Clegg, 2002; Friese and Ebel, 2010) to predict both the activity coefficients for individual ions and the gas-particle partitioning. The E-AIM model adopted the Pitzer-Simonson-Clegg model (Clegg et al., 1992a; Clegg et al., 1998) to calculate single-ion activity coefficients, which included most comprehensive conditions and have been used as a benchmark (Clegg et al., 1992b; Hennigan et al., 2015; Pye et al., 2020). Therefore, both the activity-based pH ( $\text{pH}_a$ , Eq. 11) and the free- $\text{H}^+$  molality pH ( $\text{pH}_F$ , Eq. 12) can be derived (Appendix B). In addition, we also adopted the ISORROPIA v2.3 model



(Fountoukis and Nenes, 2007) for comparison, which is computational effective and has been commonly adopted in global and regional models. To reduce the computational cost, the ISORROPIA model calculated only the binary activity coefficients  $\gamma_{ij}$  using the Kusik-Meissner relationship and the Bromley's formula (Fountoukis and Nenes, 2007). Therefore, only the free-  
135  $H^+$  molality pH ( $pH_F$ , Eq. 12) can be derived in ISORROPIA (Appendix D). For example, for a HCl droplet, both the  $\gamma_{H+(aq)}$  and  $\gamma_{Cl-(aq)}$  are calculated in E-AIM, while only the mean binary activity coefficient of  $\gamma_{H-Cl} = \sqrt{\gamma_{H+(aq)}\gamma_{Cl-(aq)}}$  is estimated in ISORROPIA.

**Global models.** Spatial variation of  $c_{ni}$  was studied based on the two global models. The global GEOS-Chem model simulations (v11-01) were conducted at a resolution of  $2.5^\circ$  longitude  $\times$   $2^\circ$  latitude with 47 vertical layers for 2016. Detailed  
140 model settings are provided elsewhere (Zheng et al., 2020). The global EMAC (ECHAM5/MESSy2 for Atmospheric Chemistry) model were conducted at a resolution of T63 (i.e.,  $\sim 1.8^\circ \times 1.8^\circ$  at the equator) with 31 vertical levels for 2016. Detailed EMAC model settings are provided in Appendix C.

### 3. Results and discussion

#### 3.1 Influencing factors of the non-ideality coefficient

145 All activity coefficients first depends on RH and temperature. In addition, for ammonium-buffered ambient aerosols, major anions in pair with  $NH_4^+$  or  $H^+$  is  $NO_3^-$  and  $SO_4^{2-}$ . The ratio of mean activity coefficients is therefore expected to differ when they're mainly combined with  $SO_4^{2-}$  (i.e.,  $\gamma_{NH_4HSO_4}/\gamma_{H-HSO_4}$ ) or  $NO_3^-$  (i.e.,  $\gamma_{NH_4NO_3}/\gamma_{HNO_3}$ ).

Figure 1 shows the dependence of  $c_{ni}$  under different systems (Appendix A), as predicted by the gas-particle partitioning (Eq. 14b) with E-AIM (Fig. 1 a, c, e) and ISORROPIA II (Fig. 1b, d, f), respectively. Based on both models,  $c_{ni}$  differs much  
150 between  $NH_3$ - $H_2SO_4$  system (Fig. 1a, b) and  $NH_3$ - $HNO_3$ - $H_2SO_4$  system (Fig. 1c,d), even at the same RH and temperature. The difference is still large when compared at the same ionic strength and temperature (Fig. A1), illustrating that the difference is mainly due to the ion-pair specific binary activity coefficients,  $\gamma_{ij}^0$  (Zaveri et al., 2005; Fountoukis and Nenes, 2007; Clegg et al., 1992b) (Appendix B; Fig. B1).

Due to the large difference in  $c_{ni}$  between  $NH_4NO_3$  and  $(NH_4)_2SO_4$  dominated aerosols, the  $c_{ni}$  at given RH and temperature  
155 conditions is therefore sensitive to the anion profiles, as characterized by the fraction of  $NO_3^-$  in anions(aq),  $f_{NO_3^-}$ , of:

$$f_{NO_3^-} (\mu eq/\mu eq) = [NO_3^-(aq)]/[Anions(aq)] \quad (15a)$$

$$[Anions(aq)] = 2 [SO_4^{2-}(aq)] + [NO_3^-(aq)] + [Cl^-(aq)] \quad (15b)$$

The  $f_{NO_3^-}$  is proportional to  $NO_3^-/SO_4^{2-}$  molar ratios when  $Cl^-$  is negligible. In comparison, the cation profiles, or the relative abundances of non-volatile cations (NVCs; total cations from  $Na^+$ ,  $Ca^{2+}$ ,  $K^+$ , and  $Mg^{2+}$ ), play a minor role as their influence  
160 is more indirect (Fig. 1e, f).



### 3.2 Comparison of $c_{ni}$ estimated by E-AIM and ISORROPIA

As discussed in section 2.2, for E-AIM  $c_{ni}$  can be estimated either by activity coefficients (Eq. 14a) or gas-particle partitioning (Eq. 14b), and the results agreed perfectly (black lines in Fig. 2). Therefore, the  $c_{ni}$  estimation with E-AIM is calculated by the gas-particle partitioning (Eq. 14b) hereinafter, the same as ISORROPIA.

165 Although showing the same influencing factors,  $c_{ni}$  estimated by E-AIM and ISORROPIA are not identical (Fig. 1). Especially for the  $\text{NH}_3\text{-H}_2\text{SO}_4\text{-H}_2\text{O}$  system (i.e.,  $(\text{NH}_4)_2\text{SO}_4$  dominated aerosols), E-AIM (Fig. 1a) and ISORROPIA (Fig. 1b) even predicted reversed trends in  $c_{ni}$  dependence on RH and temperature. This is more clearly shown in Fig. 2 (blue dots), where  $c_{ni}$  by E-AIM and ISORROPIA at the same conditions (i.e., same RH, temperature and chemical profiles) are compared. As shown in Fig. 2a, while  $c_{ni}$  predicted by E-AIM ranged -0.3 to 0.5 for  $(\text{NH}_4)_2\text{SO}_4$  dominated aerosols, that by ISORROPIA is always  
170 larger than 0.1.

For the  $\text{NH}_3\text{-H}_2\text{SO}_4\text{-H}_2\text{O}$  system, we found that these two models generate similar prediction of AWC (and therefore similar ideal constant,  $K_{a,\text{NH}_3}^{*i}$ ) (Fig. D1a). The different  $c_{ni}$  is mainly due to disagreement in the predicted molar ratios of  $\text{NH}_3(\text{g})/\text{NH}_4^+$  (Fig. D1b). This is caused by the difference of calculated activity coefficients between ISORROPIA (Song et al., 2018; Fountoukis and Nenes, 2007) and E-AIM (see details in Appendix D and Fig. D2). Despite the difference in estimated  $c_{ni}$ , the  
175 difference in pH predictions by E-AIM and ISORROPIA is relatively small, as pH was mainly controlled by  $\text{p}K_{a,\text{NH}_3}^{*i}$  (Fig. D1c).

Unlike the  $\text{NH}_3\text{-H}_2\text{SO}_4\text{-H}_2\text{O}$  system,  $c_{ni}$  estimated by ISORROPIA generally agrees well with (while tends to be somewhat higher than) E-AIM when  $\text{HNO}_3$  is present in the system (Fig. 2b, c). This indicates that constraint from  $\text{NH}_3\text{-HNO}_3$  equilibriums are quite important in estimating  $c_{ni}$  with ISORROPIA (see details in Appendix D). Under ambient conditions,  
180 there's barely places with negligible  $\text{HNO}_3$ , thus the ISORROPIA predicted  $c_{ni}$  generally agreed with E-AIM (section 3.4).

With the known governing factors, here we propose a parameterization method to estimate  $c_{ni}$  at given RH, temperature and  $f_{\text{NO}_3}$ , with lookup tables generated by comprehensive thermodynamic models, E-AIM and ISORROPIA (“AIM\_molality” database and “ISORROPIA\_molality” database as in Data S1). In addition, the parameterized  $c_{ni}$  for activity-based pH (Eq. 11; Appendix B; Fig. B1) is also available (“AIM\_activity” database in Data S1). A Matlab code to get  $c_{ni}$  is also provided  
185 (Data S1). Example slices of this  $c_{ni}$  parameterization based on “AIM\_molality” estimations are shown in Fig. 3. Note that this parameterization method aimed at  $\text{NH}_3\text{-HNO}_3\text{-H}_2\text{SO}_4\text{-H}_2\text{O}$  system, assuming no NVCs. We will show that this assumption is acceptable under most cases in the following sections.

### 3.3 Validation and applications with long-term observations

To validate the  $c_{ni}$  parameterization method under actual ambient conditions, we here show an example application based on  
190 the long-term measurements in Toronto (Tao and Murphy, 2019) (Fig. 4). From 2007 to 2016, Toronto resides in the ammonia-buffered regime for ~80% of the times, and the model-predicted pH based on the measured chemical compositions follows nicely with the variation of actual  $\text{p}K_{a}^{*,ni}$  estimated by thermodynamic models (Eq. 14), for both E-AIM (Fig. 4a) and



ISORROPIA (Fig. 4c). Parameterized  $c_{\text{ni}}$  agreed quite well with the actual ones for both models (Fig. 4b, d, black circles), with  $R^2$  both being 0.99, and the corresponding root-mean-square deviation (RMSD) both being  $\sim 0.1$ .

195 Figure 4 also suggest that most of the variation of actual  $\text{p}K_{\text{a}}^{*,\text{ni}}$  comes from the variation of ideal constants ( $\text{p}K_{\text{a}}^{*,\text{i}}$ ), not the non-ideality. For example, assuming the full aerosol and gas measurements were conducted only in a calibration year of 2012, based on which the annual mean and monthly mean  $c_{\text{ni}}$  can be derived (Fig. E1). Annual mean  $c_{\text{ni}}$  is 0.4 for E-AIM and 0.8 for ISORROPIA estimations. When we use the annual mean  $c_{\text{ni}}$  as a constant correction (i.e., estimated  $\text{p}K_{\text{a}}^{*,\text{ni}} = \text{p}K_{\text{a}}^{*,\text{i}} + \text{annual mean}$ ), fluctuation in the estimated  $\text{p}K_{\text{a}}^{*,\text{ni}}$  would actually all come from  $\text{p}K_{\text{a}}^{*,\text{i}}$ . However, this estimated  $\text{p}K_{\text{a}}^{*,\text{ni}}$  can already

200 explain  $\sim 90\%$  of the variations in actual  $\text{p}K_{\text{a}}^{*,\text{ni}}$  (red dots in Fig. 4b, d), illustrating the dominance of  $\text{p}K_{\text{a}}^{*,\text{i}}$  (i.e., AWC and temperature fluctuations) over non-ideality. In comparison, applying the month-dependent  $c_{\text{ni}}$  values (blue dots in Fig. 4b, d) makes little difference with the annual constant estimations ( $R^2$  differed only by 1%).

Figure 4 and Fig. E1 illustrate that a constant  $c_{\text{ni}}$  is often good enough at a given site. Full aerosol species measurements for a whole year, or under periods representative of annual-average conditions (like spring or fall seasons for Toronto; Fig. E1) is

205 recommended in determining the localized  $c_{\text{ni}}$ , which, together with AWC and temperature measurements, could already provide a good approximation of the aerosol pH. This is especially useful in retrieving the acidity variations when full chemical measurements are not available in the long run.

### 3.4 Validation and application against global model simulations

We further investigated the influence of non-ideality in explaining the spatial variations of aerosol acidity based on global

210 model simulations. On the global scale, fraction of  $\text{NO}_3^-$  in aqueous phase anions depends on two factors: the total nitrate (gas + particle phase) to sulfate ratios, and the partitioning of total nitrates. When total nitrate  $\ll$  sulfates, the aerosols would be dominated by  $(\text{NH}_4)_2\text{SO}_4$  even if all the nitrates are partitioned into the particle phase. In this case, non-ideality correction factor can be estimated from Fig. 1a, b at known RH and temperature. However, both GEOS-Chem and the EMAC results show that this criterion is barely met for the ammonia-buffered regions. Besides, for all the reported observation results we

215 know of, only summertime south-eastern U.S. (Weber et al., 2016) has a total nitrate that is  $< 5\%$  of the sulfate (charge ratios). Therefore, under most conditions,  $c_{\text{ni}}$  largely depends on the partitioning of total nitrates, and an estimation of  $f_{\text{NO}_3}$  is needed to derive the correction factor.

Figure 5 shows the estimated  $\text{p}K_{\text{a}}^{*,\text{ni}}$  against actual  $\text{p}K_{\text{a}}^{*,\text{ni}}$  based on GEOS-Chem simulations, and that based on EMAC simulations are shown in Fig. C1. Three scenarios are assumed to examine the sensitivity of  $\text{p}K_{\text{a}}^{*,\text{ni}}$  prediction with  $c_{\text{ni}}$  values:

220 (a) constant temperature ( $T$ ) of 288 K and RH of 73%, (b) constant RH of 73%, but with annual-average temperatures for each site; and (c) annual-average  $T$  and RH for each site. For all scenarios, annual mean chemical compositions for the ammonia-buffered surface regions (Zheng et al., 2020) are used, and  $c_{\text{ni}}$  is estimated by both E-AIM and ISORROPIA II models. Similar with Fig. 4, in the “parameterized” series  $c_{\text{ni}}$  is estimated by the parameterization method proposed in this study with RH,  $T$  and  $f_{\text{NO}_3}$  at certain model grid, while in the “global mean” series,  $c_{\text{ni}}$  is assumed to be constant as the average of actual  $c_{\text{ni}}$





225 estimated by the thermodynamic models under each scenario, which is  $\sim 0.6$  for E-AIM model and  $\sim 0.8$  for ISORROPIA model.

Based on GEOS-Chem simulations, the parameterized  $c_{ni}$  (black dots in Fig. 5) work nicely in reproducing actual  $pK_a^{*,ni}$ , with  $R^2$  near 1 under all scenarios, and the RMSD of  $< 0.03$  for ISORROPIA model and  $\sim 0.1$  for AIM model. Again, we found that variations of  $c_{ni}$  is much smaller than the variation of  $pK_a^{*,i}$  caused by particulate matter concentrations and temperatures. With a constant global-mean  $c_{ni}$  correction (i.e., assuming a global average  $f_{NO_3}$ ) (blue dots in Fig. 5), the estimated  $pK_a^{*,ni}$  can already explain over 93 % of the variations in actual  $pK_a^{*,ni}$ , with/without considering the influence of meteorology on non-ideality alike. Correspondingly, it can already explain  $\sim 70\%$  of the aerosol pH variations (Zheng et al., 2020), where the pH is further subject to variations in  $NH_3(g)$  and NVCs (Eq. 8; Zheng et al. (2020)).

The EMAC simulations show similar patterns with GEOS-Chem results. Estimated  $pK_a^{*,ni}$  with the parameterized  $c_{ni}$  corrections agreed well with actual  $pK_a^{*,ni}$ , with  $R^2$  over 0.94 for E-AIM model and over 0.91 for ISORROPIA model (Fig. C1). This is somewhat lower than the Toronto site (Fig. 4) or the GEOS-Chem result (Fig. 5), which is due to the larger variations in the simulated chemical profiles (e.g., importance of NVCs and  $Cl^-$ , etc.). The constant  $c_{ni}$  assumption (blue dots in Fig. C1) works similarly with the parameterized ones when influence of meteorology is excluded (Fig. C1 a, d) or when spatial variations of temperatures are considered (Fig. C1 b,e). When spatial variations of both temperature and RH (Fig. C1c,f) are considered, the constant  $c_{ni}$  assumption works worse than the parameterized ones, but is still acceptable ( $R^2$  being 0.75 for E-AIM and 0.69 for ISORROPIA).

Note that under all conditions, the “global mean” method tend to overestimate  $c_{ni}$  when actual  $pK_a^{*,ni}$  of  $NH_3$  is smaller than 2 (Fig. 5, Fig. C1). That is caused by  $f_{NO_3}$ . The low  $pK_a^{*,ni}$  indicates low AWC levels (Zheng et al., 2020) and relatively low pH levels (Eq. 13). Under such conditions,  $HNO_3$  tends to stay in the gas phase (Nenes et al., 2020), corresponding to a low  $f_{NO_3}$  of  $\sim 0$ . In comparison, the global-mean  $c_{ni}$  correspond to the global-mean simulated  $f_{NO_3}$  of  $\sim 0.4$ . As  $c_{ni}$  increase with increasing  $f_{NO_3}$  (Fig. 1e,f), the global-mean  $c_{ni}$  would tend to overestimate actual low  $pK_a^{*,ni}$  conditions (i.e.,  $< 2$ ).

#### 4. Conclusions

Overall, we found that the non-ideality correction is needed for using  $pK_a^{*,ni}$  of  $NH_3$  as a proxy of aerosol pH in ammonia-buffered regimes. This correction factor,  $c_{ni}$ , generally ranging 0.3  $\sim$  1.1, and mainly depends on RH, temperature and the fraction of nitration in aqueous-phase anions. E-AIM generally predicted a lower  $c_{ni}$  than the ISORROPIA model. We proposed a parameterization method to estimate the  $c_{ni}$ , which works quite well, as validated against both long-term observations and global simulations. Although the correction is needed in estimating the ammonia  $pK_a^{*,ni}$  levels, the variations in  $pK_a^{*,ni}$  is often much less sensitive to the non-ideality than to aerosol water content and temperature. Therefore, a constant correction factor of  $pK_{a,NH_3}^*$  is often good enough to predict pH over a period at a given site, or to explain the global pH variations. We thereby provided a way for pH retrieval when chemical measurements are unavailable for the ammonia-buffered regions.



*Data availability.* All data used in this study are described in the manuscript and supporting information.

## 260 Appendix A. Scenario settings for different systems

In Figs. 1, 2, A1, B1, D1 and D2, we assumed three systems, with the settings detailed below.

**NH<sub>3</sub>-H<sub>2</sub>SO<sub>4</sub>-H<sub>2</sub>O system.** For this system, we assumed a constant input with 0.5 μmol m<sup>-3</sup> of total sulfate (i.e., 1 μeq m<sup>-3</sup> of anions) and 2 μmol m<sup>-3</sup> of total ammonia. This ratio is to ensure that the system pH is around the maximum buffering capacity of ammonia. However, we found that for ISORROPIA model, the solver with only ammonia and sulfates inputs is not stable, with predicted pH often larger than 7 (SI S4). We thereby introduced 0.015 μmol m<sup>-3</sup> of Na<sup>+</sup> (3% of the total sulfate molar concentrations, or 1.5% of the anions), which exerted little influence on the ionic environments (difference in E-AIM results less than 3%) but will change the ISORROPIA subroutine solver called. The RH and temperature are then varied at different values to check the influence.

**NH<sub>3</sub>-HNO<sub>3</sub>-H<sub>2</sub>O system.** For this system, we assumed a constant input with 1 μmol m<sup>-3</sup> of total nitrate (also 1 μeq m<sup>-3</sup> of anions) and 2 μmol m<sup>-3</sup> of total ammonia, and then varied the RH and temperatures to derive non-ideality correction factors.

**Na<sup>+</sup>-NH<sub>3</sub>-HNO<sub>3</sub>-H<sub>2</sub>SO<sub>4</sub>-H<sub>2</sub>O system.** For this system, we fixed the RH at 73% and temperature at 288.15 K, 2 μmol m<sup>-3</sup> of total ammonia and a fixed concentration of total anions as 1 μeq m<sup>-3</sup>. The nitrate/sulfate ratios are then varied (but keeping their total charges the same) to get different nitrate fractions. For example, when the input sulfate is 0.25 μmol m<sup>-3</sup> equalling 0.5 μeq m<sup>-3</sup> of anions, the input total nitrate is then set to 0.5 μmol m<sup>-3</sup>, corresponding to a total anion of the system as 1 μeq m<sup>-3</sup>. Meanwhile, the ratio of NVCs (here assumed to be Na<sup>+</sup> only) to anions is also varied, and combined with the different nitrate/sulfate ratios to generate different simulation conditions.

## Appendix B: Non-ideality correction factor for activity-based pH definitions

With activity-based pH definition (i.e., pH = -log(γ<sub>H<sup>+</sup></sub> [H<sup>+</sup>])), multiphase buffer theory can be rewritten as:

$$K_{a,\text{NH}_3}^{*,nia} = \frac{[\text{NH}_3(\text{g})]}{[\text{NH}_4^+(\text{aq})]} (\gamma_{\text{H}^+} [\text{H}^+(\text{aq})]) = \gamma_{\text{NH}_4^+} K_{a,\text{NH}_3}^{*,i} = \gamma_{\text{NH}_4^+} K_{a,\text{NH}_3} \left(1 + \frac{\rho_w}{H_{\text{NH}_3} R T \text{AWC}}\right) \quad (\text{A1a})$$

$$\text{pH} = -\log(\gamma_{\text{H}^+} [\text{H}^+(\text{aq})]) = \text{p}K_{a,\text{NH}_3}^{*,nia} + \log \frac{[\text{NH}_3(\text{g})]}{[\text{NH}_4^+(\text{aq})]} \quad (\text{A1b})$$

where  $K_{a}^{*,nia}$  is the multiphase effective acid dissociation constant under non-ideal conditions. The difference of pH caused by non-ideality,  $c_{nia}$ , is therefore:



$$c_{nia} = pK_a^{*,nia} - pK_a^{*,i} = -\log \gamma_{NH_4^+} \quad (A2)$$

That is, the non-ideality correction factor for activity-based pH is actually the  $\gamma_{NH_4^+}$ , which can be calculated with the more  
285 comprehensive models like E-AIM. The E-AIM calculated mole-fraction-based activity coefficient ( $f_i$ ) that can be converted  
to the molality-based activity coefficient ( $\gamma_i$ ) by (Pye et al., 2020; Peng et al., 2019):

$$\gamma_i = f_i x_w = f_i x_i / (m_i M_w) \quad (A3)$$

where  $x_i$  and  $m_i$  are respectively the mole fraction and molality of species  $i$ , and  $x_w$  and  $M_w$  are respectively the mole fraction  
and molecular weight of water. All these variables are given in E-AIM outputs. Major influencing factors of  $c_{nia}$  is also RH,  
290 temperature and fraction of  $NO_3^-$  in anions in the aqueous phase (aq), as shown in Fig. B1.

### Appendix C. EMAC model settings

In this section, we will only focus on the model settings for EMAC simulations, while for the GEOS-chem model settings,  
please refer to Zheng et al. (2020). We used the global ECHAM/MESy Atmospheric chemistry – Climate (EMAC) model,  
which is a numerical chemistry and climate simulation system that includes sub-models describing tropospheric and middle  
295 atmosphere processes and their interaction with oceans, land and human influences (Jöckel et al., 2010). The core atmospheric  
model is the 5th generation European Centre Hamburg general circulation model (ECHAM5) (Roeckner et al., 2006), which  
has been modularized, and to which improved submodels and updates of boundary layer, radiation, cloud and convection  
routines have been introduced. The EMAC model development is coordinated within an international consortium: see  
<https://www.mesyy-interface.org>. For the present study we applied EMAC (ECHAM5 version 5.3.02, MESy version 2.54.0)  
300 in the T63L31-resolution, i.e., with a spherical truncation of T63 (corresponding to a quadratic Gaussian grid of approx. 1.8  
by 1.8 degrees in latitude and longitude) with 31 vertical hybrid terrain-following pressure levels up to 10 hPa in the lower  
stratosphere. Meteorological conditions as in ERA-interim data from European Centre for Medium- range Weather Forecasts  
(ECMWF) were simulated by the model by applying a “nudging” technique (Jöckel et al., 2006). EMAC simulates gas-phase  
and heterogeneous chemistry through the MECCA submodel, which accounts for the photochemical oxidation of natural and  
305 anthropogenic emissions, including a comprehensive account of volatile organic carbon compounds (Sander et al., 2019).  
Aerosol microphysical processes and gas/particle partitioning are simulated with the GMXe submodel (Pringle et al., 2010;  
Pozzer et al., 2012), which describes the aerosol size distribution by seven interacting lognormal modes (four hydrophilic and  
three hydrophobic). The aerosol composition can vary between these modes (externally mixed) and is uniform within each  
mode (internally mixed). The inorganic aerosol composition is computed with the ISORROPIA-II thermodynamic equilibrium  
310 submodel (Fountoukis and Nenes, 2007). It calculates the gas/liquid/solid equilibrium partitioning of inorganic compounds  
and water. The composition and atmospheric evolution of organic aerosol compounds are simulated with the ORACLE  
submodel, which represents volatility classes of organics through their effective saturation concentrations (Tsimpidi et al.,  
2018). For this work the anthropogenic emissions EDGAR (Emissions Database for Global Atmospheric Research v4.3.2)



(Crippa et al., 2018) were applied, as well as the GFAS (Global Fire Assimilation System, v1.0) (Kaiser et al., 2012) for biomass burning emissions. The EMAC results are shown in Fig. C1.

#### Appendix D. Potential reasons for discrepancies in predicting aerosol pH by ISORROPIA and E-AIM for the $\text{NH}_3\text{-H}_2\text{SO}_4\text{-H}_2\text{O}$ system

In this study, we applied ISORROPIA version 2.1 (Fountoukis and Nenes, 2007) and E-AIM (model IV; <http://www.aim.env.uea.ac.uk/aim/aim.php>) (Clegg et al., 1992b; Wexler and Clegg, 2002; Friese and Ebel, 2010), and the following description and discussion refer to these versions of the models. For  $\text{NH}_3\text{-H}_2\text{SO}_4\text{-H}_2\text{O}$  system, we found that by assuming the same input of  $0.5 \mu\text{mol m}^{-3}$  of total sulfate (i.e.,  $1 \mu\text{eq m}^{-3}$  of anions) and  $2 \mu\text{mol m}^{-3}$  of total ammonia and varies the RH (60% - 90%) and temperatures (265 K - 300 K), ISORROPIA predicted a very high aerosol pH of about 13 (12.6-13.2) while the E-AIM predicted pH ranged 2~5, which is obviously more realistic. However, by introducing only a small amount of  $\text{Na}^+$  ( $0.015 \mu\text{mol m}^{-3}$ , or 3% of the total sulfate), the ISORROPIA predicted pH dropped dramatically to 2~5 (Fig. D1), while the E-AIM predicted pH changed little than the no- $\text{Na}^+$  predictions (pH increased systematically by 0.03 with both  $R^2$  and slope being 1). Besides, the predicted pH assuming only  $\text{HNO}_3$  and  $\text{NH}_3$  inputs ( $\text{NH}_3\text{-HNO}_3\text{-H}_2\text{O}$  system) agreed well between ISORROPIA and E-AIM (Fig. D1).

We found that the dramatic changes in ISORROPIA predicted pH levels with or without small amount of  $\text{Na}^+$  and  $\text{NO}_3^-$  additions are related to the different calculation procedures among subcases. Here we focused on subcases under the metastable and sulphate-poor (i.e., total potential cations, including total ammonia ( $[\text{NH}_3]_t$ ) and NVCs, exceed twice the molar ratios of total sulfate ( $[\text{H}_2\text{SO}_4]_t$ )) conditions.

In ISORROPIA, when there's only  $\text{NH}_3$  and  $\text{H}_2\text{SO}_4$  (i.e., "pure"  $\text{NH}_3\text{-H}_2\text{SO}_4\text{-H}_2\text{O}$  system), the corresponding subcase is "A2". As detailed below, for this subcase, activity coefficients included in the final calculations are  $\gamma_{\text{H-HSO}_4}$ ,  $\gamma_{\text{2H-SO}_4}$ , and  $\gamma_{\text{NH}_4\text{-HSO}_4}$ . As shown in Fig. D2a-c, for all these three values, there's large difference between E-AIM and ISORROPIA estimations (note that log scales are used for  $\gamma_{\text{H-HSO}_4}$  and  $\gamma_{\text{2H-SO}_4}$  plots). Therefore, it's not surprising that there is large discrepancy between the predicted pH from subcase A2 of ISORROPIA and E-AIM.

In comparison, the subcase would change to "D3" when  $\text{HNO}_3$  is introduced to the system. As detailed below, for this subcase, only  $\gamma_{\text{NH}_4\text{-NO}_3}$  is involved in the calculations. As shown in Fig. D2d, although the ISORROPIA still shows a different trend than the E-AIM, it is, however, at least on the same order of magnitude as the one predicted by E-AIM.

By introducing a small amount of  $\text{Na}^+$  into the  $\text{NH}_3\text{-H}_2\text{SO}_4\text{-H}_2\text{O}$  system, the calculation procedure of ISORROPIA would change from A2 to G5 (a  $\text{Na}^+\text{-NH}_3\text{-H}_2\text{SO}_4\text{-HNO}_3\text{-HCl-H}_2\text{O}$  aerosol system). For G5 subcase, we noticed two issues: (1) although the total  $\text{HNO}_3$  is zero, the model still tried to predict  $\gamma_{\text{H-NO}_3}$  and  $\gamma_{\text{NH}_4\text{-NO}_3}$ ; (2) as it was using  $\text{Cl}^-$  as the  $x$  variable at the final solutions, a small amount of  $\text{Cl}^-$  is always present, which is introduced by the model so the calculation procedures could go on. The relevant values are shown in Fig. D2e. In comparison, the E-AIM predicted no  $\text{NO}_3^-$  or  $\text{Cl}^-$ , and the activity coefficients of other relevant species change little with the no  $\text{Na}^+$  case. Therefore, we could not perform a comparison between



ISORROPIA and E-AIM for this case (as there's no  $\gamma_{\text{NO}_3}$  or  $\gamma_{\text{Cl}}$  in E-AIM). Based on the pH and non-ideality comparisons (Fig. D1), however, we could see that the  $\text{NH}_3$  partitioning estimated this way is far more realistic than the A2 subcase.

**Calculation principles for subcase A2 (an  $\text{NH}_3\text{-H}_2\text{SO}_4\text{-H}_2\text{O}$  aerosol system).** For the subcase A2, the major constraining equations include the  $[\text{SO}_4^{2-}]/[\text{HSO}_4^-]$  equilibriums, gas-particle partitioning of ammonia, and charge balance:

$$\text{HSO}_4^- \rightleftharpoons \text{H}^+ + \text{SO}_4^{2-}, \quad K_{\text{a,HSO}_4} = \frac{[\text{H}^+(\text{aq})][\text{SO}_4^{2-}(\text{aq})] \gamma_{\text{H}^+(\text{aq})} \gamma_{\text{SO}_4^{2-}(\text{aq})}}{[\text{HSO}_4^-(\text{aq})] \gamma_{\text{HSO}_4^-(\text{aq})}} = \frac{[\text{H}^+(\text{aq})][\text{SO}_4^{2-}(\text{aq})] \gamma_{2\text{H-SO}_4(\text{aq})}^3}{[\text{HSO}_4^-(\text{aq})] \gamma_{\text{H-HSO}_4(\text{aq})}^2} \quad (\text{D1a})$$

$$\text{NH}_4^+ \rightleftharpoons \text{NH}_3(\text{aq}) + \text{H}^+, \quad K_{\text{ag,NH}_3} = K_w / K_{\text{bg,NH}_3} = \frac{[\text{H}^+(\text{aq})][\text{NH}_3(\text{g})] \gamma_{\text{H}^+(\text{aq})}}{[\text{NH}_4^+(\text{aq})] \gamma_{\text{NH}_4^+(\text{aq})}} = \frac{[\text{H}^+(\text{aq})][\text{NH}_3(\text{g})] \gamma_{\text{H-HSO}_4(\text{aq})}^2}{[\text{NH}_4^+(\text{aq})] \gamma_{\text{NH}_4\text{-HSO}_4(\text{aq})}^2} \quad (\text{D1b})$$

$$([\text{NH}_4^+] + [\text{H}^+]) / (2[\text{SO}_4^{2-}] - [\text{HSO}_4^-]) - 1 = 0 \quad (\text{D1c})$$

With these three equations and known total ammonia ( $[\text{NH}_3]_t$ ) and total sulfate ( $[\text{H}_2\text{SO}_4]_t$ ), we have:

$$[\text{NH}_3]_t C_{2S} [\text{H}^+] / (1 + C_{2S} [\text{H}^+]) + [\text{H}^+] - [\text{H}_2\text{SO}_4]_t (2 C_1 / [\text{H}^+] + 1) / (1 + C_1 / [\text{H}^+]) = 0 \quad (\text{D2})$$

where  $C_1 = K_{\text{a,HSO}_4} \frac{\gamma_{\text{H-HSO}_4(\text{aq})}^2}{\gamma_{2\text{H-SO}_4(\text{aq})}^3}$  while  $C_{2S} = \frac{\gamma_{\text{H-HSO}_4(\text{aq})}^2}{K_{\text{ag,NH}_3} \gamma_{\text{NH}_4\text{-HSO}_4(\text{aq})}^2}$ . The only unknown is thus  $[\text{H}^+]$ , which can thus be solved by

bisection solution processes. As shown in the equation, activity coefficients that matters in solving this system include

$$\frac{\gamma_{\text{H-HSO}_4(\text{aq})}^2}{\gamma_{2\text{H-SO}_4(\text{aq})}^3} \text{ in } C_1 \text{ and } \frac{\gamma_{\text{H-HSO}_4(\text{aq})}^2}{\gamma_{\text{NH}_4\text{-HSO}_4(\text{aq})}^2} \text{ in } C_{2S}.$$

**Calculation principles for subcase D3 (an  $\text{NH}_3\text{-H}_2\text{SO}_4\text{-HNO}_3\text{-H}_2\text{O}$  aerosol system).** For the subcase D3, the major equilibriums considered is the gas-particle partitioning of ammonia and nitrates of:

$$\text{NH}_4^+ \rightleftharpoons \text{NH}_3(\text{aq}) + \text{H}^+, \quad K_{\text{ag,NH}_3} = K_w / K_{\text{bg,NH}_3} = \frac{[\text{H}^+(\text{aq})][\text{NH}_3(\text{g})] \gamma_{\text{H}^+(\text{aq})}}{[\text{NH}_4^+(\text{aq})] \gamma_{\text{NH}_4^+(\text{aq})}} = \frac{[\text{H}^+(\text{aq})][\text{NH}_3(\text{g})] \gamma_{\text{H-NO}_3(\text{aq})}^2}{[\text{NH}_4^+(\text{aq})] \gamma_{\text{NH}_4\text{-NO}_3(\text{aq})}^2} \quad (\text{D3})$$

$$\text{HNO}_3(\text{g}) \rightleftharpoons \text{H}^+ + \text{NO}_3^-, \quad K_{\text{ag,HNO}_3} = \frac{[\text{H}^+(\text{aq})][\text{NO}_3^-(\text{aq})] \gamma_{\text{H}^+(\text{aq})} \gamma_{\text{NO}_3^-(\text{aq})}}{[\text{HNO}_3(\text{g})] \gamma_{\text{HNO}_3(\text{g})}} \quad (\text{D4})$$

Note that in subcase D3 the  $\gamma_{\text{H}^+}/\gamma_{\text{NH}_4^+}$  is estimated by  $(\gamma_{\text{H-NO}_3}/\gamma_{\text{NH}_4\text{-NO}_3})^2$ , not the  $(\gamma_{\text{H-HSO}_4}/\gamma_{\text{NH}_4\text{-HSO}_4})^2$  as in subcase A2.

These two equilibriums are further combined to be:

$$C_3 = \frac{[\text{NO}_3^-(\text{aq})][\text{NH}_4^+(\text{aq})]}{[\text{HNO}_3(\text{g})][\text{NH}_3(\text{g})]} = \frac{K_{\text{ag,HNO}_3}}{K_{\text{ag,NH}_3} \gamma_{\text{NH}_4\text{-NO}_3}^2} \quad (\text{D5a})$$

As to the charge balance, here only major species are considered as:

$$[\text{NH}_4^+(\text{aq})] = 2[\text{SO}_4^{2-}(\text{aq})] + [\text{NO}_3^-(\text{aq})] = 2[\text{H}_2\text{SO}_4]_t + [\text{NO}_3^-(\text{aq})] \quad (\text{D5b})$$



Combining Eqs. D5a and D5b, at given total nitrate ( $[\text{HNO}_3]_t$ , namely  $[\text{NO}_3^-(\text{aq})] + [\text{HNO}_3(\text{g})]$ ) and  $[\text{NH}_3]_t (= [\text{NH}_4^+(\text{aq})] + [\text{NH}_3(\text{g})])$  levels, the solution function can be expressed as:

$$370 \quad \frac{[\text{NH}_4^+(\text{aq})]}{[\text{NH}_3(\text{g})]} \frac{[\text{NO}_3^-(\text{aq})]}{[\text{HNO}_3(\text{g})]} - C_3 = 0$$

$$\frac{[\text{NH}_4^+(\text{aq})]}{([\text{NH}_3]_t - [\text{NH}_4^+(\text{aq})])} \frac{([\text{NH}_4^+(\text{aq})] - 2[\text{H}_2\text{SO}_4]_t)}{([\text{HNO}_3]_t + 2[\text{H}_2\text{SO}_4]_t - [\text{NH}_4^+(\text{aq})])} - C_3 = 0 \quad (\text{D6})$$

Where the only unknown is  $[\text{NH}_4^+(\text{aq})]$  and can be solved through bisection method. As shown in the equation, the only activity coefficients that matters in solving this system is  $(\gamma_{\text{NH}_4\text{-NO}_3})^2$  in  $C_3$ .

**Calculation principles for subcase G5 (a  $\text{Na}^+\text{-NH}_3\text{-H}_2\text{SO}_4\text{-HNO}_3\text{-HCl-H}_2\text{O}$  aerosol system).** For the subcase G5, the major  
 375 equilibriums considered is the gas-particle partitioning of  $\text{NH}_3$ ,  $\text{HNO}_3$  and  $\text{HCl}$ , while sulfate is considered to exist mainly as  $[\text{SO}_4^{2-}(\text{aq})]$ . General derivation processes are similar with D3 and is also detailed in a previous study (Song et al., 2018). Briefly, the key equilibriums include that of  $\text{HNO}_3$  (Eq. D4) and  $\text{HCl}$  of:

$$\text{HCl}(\text{g}) \rightleftharpoons \text{H}^+(\text{aq}) + \text{Cl}^-(\text{aq}), \quad K_{\text{ag,HCl}} = \frac{[\text{H}^+(\text{aq})][\text{Cl}^-(\text{aq})]\gamma_{\text{H}^+(\text{aq})}\gamma_{\text{Cl}^-(\text{aq})}}{[\text{HCl}(\text{g})]\gamma_{\text{HCl}(\text{g})}} = \frac{[\text{H}^+(\text{aq})][\text{Cl}^-(\text{aq})]\gamma_{\text{H-Cl}}^2}{[\text{HCl}(\text{g})]} \quad (\text{D7})$$

Which can be combined into:

$$380 \quad C_4 = \frac{\xi_{\text{HNO}_3}}{\xi_{\text{HCl}}} = \frac{[\text{NO}_3^-(\text{aq})][\text{HCl}(\text{g})]}{[\text{Cl}^-(\text{aq})][\text{HNO}_3(\text{g})]} = \frac{K_{\text{ag,HNO}_3}}{K_{\text{ag,HCl}}} \frac{\gamma_{\text{H-Cl}}^2}{\gamma_{\text{H-NO}_3}^2} \quad (\text{D8})$$

Therefore  $[\text{NO}_3^-(\text{aq})]$  and  $[\text{HNO}_3(\text{g})] (= [\text{HNO}_3]_t - [\text{NO}_3^-(\text{aq})])$  can be solved at known assumed  $[\text{Cl}^-(\text{aq})]$ .

And the  $[\text{NH}_4^+(\text{aq})]$  associated with  $\text{Cl}^-(\text{aq})$  and  $\text{NO}_3^-(\text{aq})$ ,  $[\text{NH}_4^+(\text{aq})]_{\text{NC}}$ , is solved by:

$$([\text{NH}_4^+(\text{aq})]_{\text{NC}})^2 - B [\text{NH}_4^+(\text{aq})]_{\text{NC}} + C = 0 \quad (\text{D9})$$

where

$$385 \quad B = [\text{NH}_3]_t + [\text{Na}^+] - 2[\text{H}_2\text{SO}_4]_t + [\text{Cl}^-(\text{aq})] + [\text{NO}_3^-(\text{aq})] + C_{2N}^{-1}$$

$$C = ([\text{NH}_3]_t + [\text{Na}^+] - 2[\text{H}_2\text{SO}_4]_t) ([\text{Cl}^-(\text{aq})] + [\text{NO}_3^-(\text{aq})]) - C_{2N}^{-1} (2[\text{H}_2\text{SO}_4]_t - [\text{Na}^+])$$

where  $C_{2N} = \frac{\gamma_{\text{H-NO}_3(\text{aq})}^2}{K_{\text{ag,NH}_3}\gamma_{\text{NH}_4\text{-NO}_3(\text{aq})}^2}$ . And with  $[\text{NH}_4^+(\text{aq})]_{\text{NC}}$ , we have:

$$[\text{NH}_4^+(\text{aq})] = [\text{NH}_4^+(\text{aq})]_{\text{NC}} + 2[\text{H}_2\text{SO}_4]_t - [\text{Na}^+]$$

$$[\text{NH}_4^+(\text{g})] = [\text{NH}_3]_t - [\text{NH}_4^+(\text{aq})]$$

390 The system then solves the equation sets through bisection method by assuming a series of  $[\text{Cl}^-(\text{aq})]$  levels.

As shown in the equations above, activity coefficients that matters in solving this system (Eqs. D8~D9) include  $\gamma_{\text{H-NO}_3}$ ,  $\gamma_{\text{H-Cl}}$ , and  $\gamma_{\text{NH}_4\text{-NO}_3}$ .



## Appendix E. Information for the assumed calibration year of 2012 in Toronto site (Fig. E1).

395

*Author contributions.* Y.C., H.S. and G.Z. designed the study. G.Z. performed the study. S.W. provided GEOS-Chem model results. A.P. provided the global numerical EMAC model results. G.Z., Y.C, and H.S. wrote the manuscript with input from all coauthors.

400 *Competing interests.* The authors declare no competing interests.

*Acknowledgments.* The research was supported by the Max Planck Society (MPG). Y.C. acknowledges the Minerva Program of MPG. We would like to thank Ulrich Pöschl for the helpful discussion and support.

## 405 References

- Ault, A. P.: Aerosol Acidity: Novel Measurements and Implications for Atmospheric Chemistry, *Accounts of Chemical Research*, 53, 1703-1714, 10.1021/acs.accounts.0c00303, 2020.
- Clegg, S. L., Pitzer, K. S., and Brimblecombe, P.: Thermodynamics of multicomponent, miscible, ionic-solutions. 2. Mixtures including unsymmetrical electrolytes, *J. Phys. Chem.*, 96, 9470-9479, 10.1021/j100202a074, 1992a.
- 410 Clegg, S. L., Pitzer, K. S., and Brimblecombe, P.: Thermodynamics of multicomponent, miscible, ionic solutions. Mixtures including unsymmetrical electrolytes, *The Journal of Physical Chemistry*, 96, 9470-9479, 10.1021/j100202a074, 1992b.
- Clegg, S. L., Brimblecombe, P., and Wexler, A. S.: Thermodynamic model of the system  $\text{H}^+\text{-NH}_4^+\text{-Na}^+\text{-SO}_4^{2-}\text{-NO}_3^-\text{-Cl}^-\text{-H}_2\text{O}$  at 298.15 K, *J. Phys. Chem. A*, 102, 2155-2171, 10.1021/jp973043j, 1998.
- Clegg, S. L., Seinfeld, J. H., and Brimblecombe, P.: Thermodynamic modelling of aqueous aerosols containing electrolytes and dissolved organic compounds, *Journal of Aerosol Science*, 32, 713-738, [https://doi.org/10.1016/S0021-8502\(00\)00105-1](https://doi.org/10.1016/S0021-8502(00)00105-1), 2001.
- 415 Craig, R. L., Peterson, P. K., Nandy, L., Lei, Z., Hossain, M. A., Camarena, S., Dodson, R. A., Cook, R. D., Dutcher, C. S., and Ault, A. P.: Direct Determination of Aerosol pH: Size-Resolved Measurements of Submicrometer and Supermicrometer Aqueous Particles, *Analytical Chemistry*, 90, 11232-11239, 10.1021/acs.analchem.8b00586, 2018.
- 420 Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Dentener, F., van Aardenne, J. A., Monni, S., Doering, U., Olivier, J. G. J., Pagliari, V., and Janssens-Maenhout, G.: Gridded emissions of air pollutants for the period 1970–2012 within EDGAR v4.3.2, *Earth Syst. Sci. Data*, 10, 1987-2013, 10.5194/essd-10-1987-2018, 2018.
- Fountoukis, C., and Nenes, A.: ISORROPIA II: a computationally efficient thermodynamic equilibrium model for  $\text{K}^+\text{-Ca}^{2+}\text{-Mg}^{2+}\text{-NH}_4^+\text{-Na}^+\text{-SO}_4^{2-}\text{-NO}_3^-\text{-Cl}^-\text{-H}_2\text{O}$  aerosols, *Atmos. Chem. Phys.*, 7, 4639-4659, 10.5194/acp-7-4639-2007, 2007.



- 425 Fountoukis, C., Nenes, A., Sullivan, A., Weber, R., Van Reken, T., Fischer, M., Matías, E., Moya, M., Farmer, D., and Cohen, R. C.: Thermodynamic characterization of Mexico City aerosol during MILAGRO 2006, *Atmos. Chem. Phys.*, 9, 2141-2156, 10.5194/acp-9-2141-2009, 2009.
- Friese, E., and Ebel, A.: Temperature Dependent Thermodynamic Model of the System  $H^+-NH_4^+-Na^+-SO_4^{2-}-NO_3^- -Cl^- -H_2O$ , *The Journal of Physical Chemistry A*, 114, 11595-11631, 10.1021/jp101041j, 430 2010.
- Guo, H., Xu, L., Bougiatioti, A., Cerully, K. M., Capps, S. L., Hite Jr, J. R., Carlton, A. G., Lee, S. H., Bergin, M. H., Ng, N. L., Nenes, A., and Weber, R. J.: Fine-particle water and pH in the southeastern United States, *Atmos. Chem. Phys.*, 15, 5211-5228, 10.5194/acp-15-5211-2015, 2015.
- Guo, H., Sullivan, A. P., Campuzano-Jost, P., Schroder, J. C., Lopez-Hilfiker, F. D., Dibb, J. E., Jimenez, J. L., Thornton, J. 435 A., Brown, S. S., Nenes, A., and Weber, R. J.: Fine particle pH and the partitioning of nitric acid during winter in the northeastern United States, *Journal of Geophysical Research: Atmospheres*, 121, 10,355-310,376, 10.1002/2016JD025311, 2016.
- Guo, H., Nenes, A., and Weber, R. J.: The underappreciated role of nonvolatile cations on aerosol ammonium-sulfate molar ratios, *Atmos. Chem. Phys. Discuss.*, 2017, 1-19, 10.5194/acp-2017-737, 2017.
- 440 Hennigan, C. J., Izumi, J., Sullivan, A. P., Weber, R. J., and Nenes, A.: A critical evaluation of proxy methods used to estimate the acidity of atmospheric particles, *Atmos. Chem. Phys.*, 15, 2775-2790, 10.5194/acp-15-2775-2015, 2015.
- Jöckel, P., Tost, H., Pozzer, A., Brühl, C., Buchholz, J., Ganzeveld, L., Hoor, P., Kerkweg, A., Lawrence, M. G., Sander, R., Steil, B., Stiller, G., Tanarhte, M., Taraborrelli, D., van Aardenne, J., and Lelieveld, J.: The atmospheric chemistry general circulation model ECHAM5/MESSy1: consistent simulation of ozone from the surface to the mesosphere, *Atmos. Chem. 445 Phys.*, 6, 5067-5104, 10.5194/acp-6-5067-2006, 2006.
- Jöckel, P., Kerkweg, A., Pozzer, A., Sander, R., Tost, H., Riede, H., Baumgaertner, A., Gromov, S., and Kern, B.: Development cycle 2 of the Modular Earth Submodel System (MESSy2), *Geosci. Model Dev.*, 3, 717-752, 10.5194/gmd-3-717-2010, 2010.
- Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J. J., Razingerg, M., Schultz, M. G., Suttie, M., and van der Werf, G. R.: Biomass burning emissions estimated with a global fire assimilation system based on 450 observed fire radiative power, *Biogeosciences*, 9, 527-554, 10.5194/bg-9-527-2012, 2012.
- Li, G., Su, H., Ma, N., Zheng, G., Kuhn, U., Li, M., Klimach, T., Pöschl, U., and Cheng, Y.: Multifactor colorimetric analysis on pH-indicator papers: an optimized approach for direct determination of ambient aerosol pH, *Atmos. Meas. Tech.*, 13, 6053-6065, 10.5194/amt-13-6053-2020, 2020.
- Nenes, A., Pandis, S. N., Weber, R. J., and Russell, A.: Aerosol pH and liquid water content determine when particulate matter 455 is sensitive to ammonia and nitrate availability, *Atmos. Chem. Phys.*, 20, 3249-3258, 10.5194/acp-20-3249-2020, 2020.
- Peng, X., Vasilakos, P., Nenes, A., Shi, G., Qian, Y., Shi, X., Xiao, Z., Chen, K., Feng, Y., and Russell, A. G.: Detailed Analysis of Estimated pH, Activity Coefficients, and Ion Concentrations between the Three Aerosol Thermodynamic Models, *Environmental Science & Technology*, 53, 8903-8913, 10.1021/acs.est.9b00181, 2019.






- Pfennig, B. W.: Principles of inorganic chemistry, John Wiley & Sons, 2015.
- 460 Pozzer, A., de Meij, A., Pringle, K. J., Tost, H., Doering, U. M., van Aardenne, J., and Lelieveld, J.: Distributions and regional budgets of aerosols and their precursors simulated with the EMAC chemistry-climate model, *Atmos. Chem. Phys.*, 12, 961-987, 10.5194/acp-12-961-2012, 2012.
- Pringle, K. J., Tost, H., Message, S., Steil, B., Giannadaki, D., Nenes, A., Fountoukis, C., Stier, P., Vignati, E., and Lelieveld, J.: Description and evaluation of GMXe: a new aerosol submodel for global simulations (v1), *Geosci. Model Dev.*, 3, 391-465 412, 10.5194/gmd-3-391-2010, 2010.
- Pye, H. O. T., Zuend, A., Fry, J. L., Isaacman-VanWertz, G., Capps, S. L., Appel, K. W., Foroutan, H., Xu, L., Ng, N. L., and Goldstein, A. H.: Coupling of organic and inorganic aerosol systems and the effect on gas-particle partitioning in the southeastern US, *Atmos. Chem. Phys.*, 18, 357-370, 10.5194/acp-18-357-2018, 2018.
- Pye, H. O. T., Nenes, A., Alexander, B., Ault, A. P., Barth, M. C., Clegg, S. L., Collett Jr, J. L., Fahey, K. M., Hennigan, C. 470 J., Herrmann, H., Kanakidou, M., Kelly, J. T., Ku, I. T., McNeill, V. F., Riemer, N., Schaefer, T., Shi, G., Tilgner, A., Walker, J. T., Wang, T., Weber, R., Xing, J., Zaveri, R. A., and Zuend, A.: The acidity of atmospheric particles and clouds, *Atmos. Chem. Phys.*, 20, 4809-4888, 10.5194/acp-20-4809-2020, 2020.
- Roeckner, E., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kornblueh, L., Manzini, E., Schlese, U., and Schulzweida, U.: Sensitivity of Simulated Climate to Horizontal and Vertical Resolution in the ECHAM5 Atmosphere Model, *Journal of* 475 *Climate*, 19, 3771-3791, 10.1175/jcli3824.1, 2006.
- Sander, R., Baumgaertner, A., Cabrera-Perez, D., Frank, F., Gromov, S., Grooß, J. U., Harder, H., Huijnen, V., Jöckel, P., Karydis, V. A., Niemeyer, K. E., Pozzer, A., Riede, H., Schultz, M. G., Taraborrelli, D., and Tauer, S.: The community atmospheric chemistry box model CAABA/MECCA-4.0, *Geosci. Model Dev.*, 12, 1365-1385, 10.5194/gmd-12-1365-2019, 2019.
- 480 Song, S., Gao, M., Xu, W., Shao, J., Shi, G., Wang, S., Wang, Y., Sun, Y., and McElroy, M. B.: Fine-particle pH for Beijing winter haze as inferred from different thermodynamic equilibrium models, *Atmos. Chem. Phys.*, 18, 7423-7438, 10.5194/acp-18-7423-2018, 2018.
- Tao, Y., and Murphy, J. G.: The sensitivity of PM<sub>2.5</sub> acidity to meteorological parameters and chemical composition changes: 10-year records from six Canadian monitoring sites, *Atmos. Chem. Phys.*, 19, 9309-9320, 10.5194/acp-19-9309-2019, 2019.
- 485 Tsimpidi, A. P., Karydis, V. A., Pozzer, A., Pandis, S. N., and Lelieveld, J.: ORACLE 2-D (v2.0): an efficient module to compute the volatility and oxygen content of organic aerosol with a global chemistry-climate model, *Geosci. Model Dev.*, 11, 3369-3389, 10.5194/gmd-11-3369-2018, 2018.
- Weber, R. J., Guo, H., Russell, A. G., and Nenes, A.: High aerosol acidity despite declining atmospheric sulfate concentrations over the past 15 years, *Nature Geosci.*, 9, 282-285, 10.1038/ngeo2665
- 490 <http://www.nature.com/ngeo/journal/v9/n4/abs/ngeo2665.html#supplementary-information>, 2016.



- Wei, H., Vejerano, E. P., Leng, W., Huang, Q., Willner, M. R., Marr, L. C., and Vikesland, P. J.: Aerosol microdroplets exhibit a stable pH gradient, *Proceedings of the National Academy of Sciences of the United States of America*, 115, 7272-7277, 10.1073/pnas.1720488115, 2018.
- Wexler, A. S., and Clegg, S. L.: Atmospheric aerosol models for systems including the ions  $H^+$ ,  $NH_4^+$ ,  $Na^+$ ,  $SO_4^{2-}$ ,  $NO_3^-$ ,  $Cl^-$ ,  $Br^-$ , and  $H_2O$ , *Journal of Geophysical Research: Atmospheres*, 107, ACH 14-11-ACH 14-14, 10.1029/2001jd000451, 2002.
- Zaveri, R. A., Easter, R. C., and Wexler, A. S.: A new method for multicomponent activity coefficients of electrolytes in aqueous atmospheric aerosols, *Journal of Geophysical Research: Atmospheres*, 110, 10.1029/2004jd004681, 2005.
- Zheng, G., Su, H., Wang, S., Andreae, M. O., Pöschl, U., and Cheng, Y.: Multiphase buffer theory explains contrasts in atmospheric aerosol acidity, *Science*, 369, 1374-1377, 10.1126/science.aba3719, 2020.
- Zuend, A., Marcolli, C., Luo, B. P., and Peter, T.: A thermodynamic model of mixed organic-inorganic aerosols to predict activity coefficients, *Atmos. Chem. Phys.*, 8, 4559-4593, 10.5194/acp-8-4559-2008, 2008.




505 Figures




**Figure 1. The non-ideality correction factor,  $c_{ni}$ , estimated by E-AIM (a, c, e) and ISORROPIA (b, d, f) for different aerosol systems. (a, b)  $\text{NH}_3\text{-H}_2\text{SO}_4\text{-H}_2\text{O}$  system with aerosols dominated by  $(\text{NH}_4)_2\text{SO}_4$  at varying RH and temperature conditions; (c, d)  $\text{NH}_3\text{-HNO}_3\text{-H}_2\text{O}$  system with aerosols dominated by  $\text{NH}_4\text{NO}_3$  at varying RH and temperature conditions, and (e, f)  $\text{Na}^+\text{-NH}_3\text{-HNO}_3\text{-H}_2\text{SO}_4\text{-H}_2\text{O}$  system with varying chemical profiles at 288.15 K and RH of 73%. The chemical profiles in (e, f) are characterized by the fraction of  $\text{NO}_3^-$  in anions(aq) and NVCs/anions(aq), where the non-volatile cations (NVCs) are assumed to be  $\text{Na}^+$  only here. The assumed RH and T conditions in (e, f) are marked as blacked stars in (a-d), while the chemical profiles for (a-d) and (e, f) are marked by the corresponding letter in (e) and (f), respectively.**

510




515 **Figure 2. Comparison of different  $c_{\text{ni}}$  estimation methods for three representative aerosol systems.** The  $c_{\text{ni}}$  are compared at the same conditions (i.e., same RH, temperature and chemical profiles). The  $x$  values are  $c_{\text{ni}}$  estimated by activity coefficients (Eq. 13a) with E-AIM model, and the  $y$  values include  $c_{\text{ni}}$  estimated by gas-particle ratios (Eq. 13b) with E-AIM (black lines) and ISORROPIA (blue dots) models. The systems are the same as Fig. 1.




520

Figure 3. Example slices of the  $c_{ni}$  parameterization based on “AIM\_molality” estimations as given in Data S1.



525 **Figure 4.** Comparison of pH, actual and estimated  $pK_{a,ni}^*$  based on the ten-year observations in Toronto. Data were taken from Canada's National Air Pollution Surveillance (NAPS) Program, as detailed in ref. <sup>14</sup>. Predictions are based on (a-b) E-AIM model and (c-d) ISORROPIA model. The "parameterized" series in (b, d) are predicted by the parameterization method proposed with input of the observed RH, temperature and model predicted fraction of nitrates in anions. The annual mean and monthly mean are based on mean  $c_{ni}$  of an arbitrary example year of 2012.




530

**Figure 5.** Comparison of actual and estimated  $pK_{a,ni}^*$  based on the GEOS-Chem global simulations in 2016. Predictions are based on (a-b) E-AIM model and (c-d) ISORROPIA model. The “parameterized” series are based on the parameterization method proposed in this study, while the global means are based on mean  $c_{ni}$  calculated from thermodynamic models under each scenario.




535 Figures for Appendices



**Figure A1.** Ionic strength ( $I$ ) and the non-ideality correction factor,  $c_{\text{ni}}$ , as calculated by E-AIM (a, c, e) and ISORROPIA (b, d, f) under different aerosol systems. The systems are the same as Fig. 1, while the RH in Fig. 1(a-d) and  $c_{\text{ni}}$  in Fig. 1(e-f) are replaced into  $I$ .





540

**Figure B1.** Dependence of the non-ideality correction factor for activity-based pH definitions,  $c_{nia}$  (i.e.,  $-\log(\gamma_{NH_4})$ ), as estimated by E-AIM. (a) NH<sub>3</sub>-HNO<sub>3</sub>-H<sub>2</sub>O system with aerosols dominated by NH<sub>4</sub>NO<sub>3</sub> at varying RH and temperature conditions, and (b) Na<sup>+</sup>-NH<sub>3</sub>-HNO<sub>3</sub>-H<sub>2</sub>SO<sub>4</sub>-H<sub>2</sub>O system with varying chemical profiles at 288.15 K and RH of 73%. Note that the  $c_{nia}$  for NH<sub>3</sub>-H<sub>2</sub>SO<sub>4</sub>-H<sub>2</sub>O system (i.e., (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>-dominated aerosols) is not shown, as it varied little (ranging 0.44~0.47) over the whole RH and temperature ranges explored.

545







Figure C1. Same as Fig. 5, but based on EMAC results.



550 **Figure D1.** Drivers of the difference in  $c_{ni}$  estimated by ISORROPIA and E-AIM models for the  $\text{NH}_3\text{-H}_2\text{SO}_4\text{-H}_2\text{O}$  system. The  $\zeta_{g/p}$  of  $\text{NH}_3$  indicates the molar ratios of  $\text{NH}_3(\text{g})$  to particle-phase  $\text{NH}_4^+$ .



555 **Figure D2. Comparison of activity coefficients for different species.** (a)-(c) Comparison of activity coefficients involved in ISORROPIA A2 subcase calculations, as predicted by ISORROPIA and E-AIM. (d) Comparison of activity coefficients involved in ISORROPIA D3 subcase calculations, as predicted by ISORROPIA and E-AIM. (e) Mean activity coefficients predicted by ISORROPIA that are involved in ISORROPIA G5 subcase calculations.



560

**Figure E1. Monthly variation of (a, c)  $\text{NO}_3^-$  fraction in anions(aq), and (b, d) the corresponding non-ideality correction factors for Toronto site in 2012.** The data are estimated by (a-b) E-AIM model and (c-d) ISORROPIA model. The black dash lines represent the annual mean levels. The box and whiskers represent the 10<sup>th</sup>, 25<sup>th</sup>, 50<sup>th</sup>, 75<sup>th</sup> and 90<sup>th</sup> percentiles, respectively, while the red markers represent the monthly means.