

Response to comments #1

RC1 comments:

This study evaluated the temporal and spatial distributions of yield and economic losses due to long-term ozone exposures for major crops in China between 2010 and 2017. The results of this study are interesting, however, the novelty is not clearly stated, and the writing is very poor which requires almost a total revision. Please carefully address the comments below.

Response: We really appreciate the reviewer's efforts in providing constructive comments for our paper. We have made several modifications and implemented the suggestions as needed. We describe a few major changes, followed by our response to individual comments. We have made great efforts to work on the Results and Discussion sections. We also re-emphasized the novelty of our study in the Introduction related to the reviewer's specific question (below).

Major comments:

About the Introduction:

There are only two paragraphs in the introduction, with the first paragraph does not mention the impact of ozone on crops at all. Moreover, the scientific questions did not raise clearly in the second paragraph. Overall, I feel the introduction is not well structured, and the science question targeted in this study needs to be clearly stated.

Response: We thank the reviewer's suggestion. We now rewrite the sentences from L64-74: "To date, very few studies have investigated the long-term trends and spatial patterns of ozone impacts on crop production in China. Previous studies have been mainly focus on a specific region of China, such as NCP (Zhang et al., 2017; Hu et al., 2020; Feng et al., 2020), or Yangtze River Delta (Wang et al., 2012). In this study, we focus on the long-term ozone-exposure impact analysis from 2010 to 2017 in China to assess the yield losses of four major crops (wheat, maize, rice, and soybean) and evaluate their associated economic losses. The specific period of 2010-2017 was chosen to cover the emission changes before and after the APPCAP established in 2013. Previous studies have been reporting the crop yield losses in one year (e.g., Lin et al., 2018; Yi et al., 2018; Feng et al., 2019a,b), or several years after the APPCAP (Zhao et al., 2019; Wang et al., 2022), and our study aims to present a comprehensive analysis of ozone-induce crop yield losses and economic impacts in the agriculture sector before and after the China APPCAP. Such an analysis is expected to provide scientific support to policymakers for their decision making."

Reference:

Feng, Z., Kobayashi, K., Li, P., Xu, Y., Tang, H., Guo, A., Paoletti, E. and Calatayud, V.: Impacts of current ozone pollution on wheat yield in China as estimated with observed ozone, meteorology and day of flowering, *Atmos. Environ.*, 217(March), doi:10.1016/j.atmosenv.2019.116945, 2019a.

Feng, Z., De Marco, A., Anav, A., Gualtieri, M., Sicard, P., Tian, H., Fornasier, F., Tao, F., Guo, A. and Paoletti, E.: Economic losses due to ozone impacts on human health, forest productivity and crop yield across China, *Environ. Int.*, 131(June), doi:10.1016/j.envint.2019.104966, 2019b.

Feng, Z., Hu, T., Tai, A. P. K. and Calatayud, V.: Yield and economic losses in maize caused by ambient ozone in the North China Plain (2014–2017), *Sci. Total Environ.*, 722, 137958, doi:10.1016/j.scitotenv.2020.137958, 2020.

Hu, T., Liu, S., Xu, Y., Feng, Z. and Calatayud, V.: Assessment of O₃-induced yield and economic losses for wheat in the North China Plain from 2014 to 2017, China, *Environ. Pollut.*, 258, 113828, doi:10.1016/j.envpol.2019.113828, 2020.

Lin, Y., Jiang, F., Zhao, J., Zhu, G., He, X., Ma, X., Li, S., Sabel, C. E. and Wang, H.: Impacts of O₃ on premature mortality and crop yield loss across China, *Atmos. Environ.*, 194(July), 41–47, doi:10.1016/j.atmosenv.2018.09.024, 2018.

Wang, X., Zhang, Q., Zheng, F., Zheng, Q., Yao, F., Chen, Z., Zhang, W., Hou, P., Feng, Z., Song, W., Feng, Z. and Lu, F.: Effects of elevated O₃ concentration on winter wheat and rice yields in the Yangtze River Delta, China, *Environ. Pollut.*, 171, 118–125, doi:10.1016/j.envpol.2012.07.028, 2012.

Wang, Y., Wild, O., Ashworth, K., Chen, X., Wu, Q., Qi, Y. and Wang, Z.: Reductions in crop yields across China from elevated ozone, *Environ. Pollut.*, 292(118218), doi:https://doi.org/10.1016/j.envpol.2021.118218, 2022.

Yi, F., McCarl, B. A., Zhou, X. and Jiang, F.: Damages of surface ozone: Evidence from agricultural sector in China, *Environ. Res. Lett.*, 13(3), doi:10.1088/1748-9326/aaa6d9, 2018.

Zhang, W., Feng, Z., Wang, X., Liu, X. and Hu, E.: Quantification of ozone exposure- and stomatal uptake-yield response relationships for soybean in Northeast China, *Sci. Total Environ.*, 599–600, 710–720, doi:10.1016/j.scitotenv.2017.04.231, 2017.

Zhao, H., Zheng, Y., Zhang, Y. and Li, T.: Evaluating the effects of surface O₃ on three main food crops across China during 2015–2018, *Environ. Pollut.*, 258, 113794, doi:10.1016/j.envpol.2019.113794, 2020.

About the section 2.2

Line 85-90: The authors mentioned that there are many different crop-ozone matrixes available, and they adopted AOT40 in their study. The authors should at least lay out a few crop-ozone matrixes, and talk about the possible advantage and disadvantage among different matrixes. At the end, give a reason why AOT40 is selected.

Response: AOT40 metric is the European standard for the protection of vegetation, and widely used in both America and Asia (Tang et al., 2013; Lefohn et al., 2018; Lin et al., 2018). The AOT40 metric is also considered as more accurate at high levels of ozone concentration (Tuovinen, 2000; Hollaway et al., 2012), which is the case for ozone pollution in China (Lu et al., 2018, 2020). To clarify this, we modify the sentence in line 97:

“In this study, we adopted the ozone metric of AOT40 which is the European standard for the protection of vegetation, and also the commonly used and reliable indicator in both America and Asia for crop yield assessment (UNECE, 2017; Tang et al., 2013; Lefohn et al., 2018; Lin et al., 2018; Feng et al., 2019a,b). The AOT40 metric is also considered as more accurate at high levels of ozone concentration (Tuovinen, 2000; Hollaway et al., 2012), which is the case for China (Lu et al., 2018, 2020).”

Reference:

Hollaway, M. J., Arnold, S. R., Challinor, A. J., and Emberson, L. D.: Intercontinental trans-boundary contributions to ozone-induced crop yield losses in the Northern Hemisphere, *Biogeosciences*, 9, 271–292, https://doi.org/10.5194/bg-9-271-2012, 2012.

Lefohn, A. S., Malley, C. S., Smith, L., Wells, B., Hazucha, M., Simon, H., Naik, V., Mills, G., Schultz, M. G., Paoletti, E., De Marco, A., Xu, X., Zhang, L., Wang, T., Neufeld, H. S., Musselman, R. C., Tarasick, D., Brauer, M., Feng, Z., Tang, H., Kobayashi, K., Sicard, P., Solberg, S. and Gerosa, G.: Tropospheric ozone assessment report: Global ozone metrics for climate change, human health, and crop/ecosystem research, *Elementa*, 6, doi:10.1525/elementa.279, 2018.

Lu, X., Zhang, L., Chen, Y., Zhou, M., Zheng, B., Li, K., Liu, Y., Lin, J., Fu, T.-M., and Zhang, Q.: Exploring 2016–2017 surface ozone pollution over China: source contributions and meteorological influences, *Atmos. Chem. Phys.*, 19, 8339–8361, <https://doi.org/10.5194/acp-19-8339-2019>, 2019.

Lu, X., Zhang, L., Wang, X., Gao, M., Li, K., Zhang, Y., Yue, X. and Zhang, Y.: Rapid Increases in Warm-Season Surface Ozone and Resulting Health Impact in China Since 2013, *Environ. Sci. Technol. Lett.*, doi:10.1021/acs.estlett.0c00171, 2020.

Tang, H., Takigawa, M., Liu, G., Zhu, J., & Kobayashi, K. (2013). A projection of ozone-induced wheat production loss in China and India for the years 2000 and 2020 with exposure-based and flux-based approaches. *Global Change Biology*, 19(9), 2739-2752.

Tuovinen, J. P.: Assessing vegetation exposure to ozone: Properties of the AOT40 index and modifications by deposition modelling, *Environ. Pollut.*, 109(3), 361–372, doi:10.1016/S0269-7491(00)00040-3, 2000.

UNECE, 2017. Chapter 3: mapping critical levels for vegetation. International cooperative programme on effects of air pollution on natural vegetation and crops. Bangor, UK. <http://icpvegetation.ceh.ac.uk/>, last accessed 24 November 2021.

Line 94-95: The authors should specifically define or point out the crops corresponding to what growing season, as multiple growing seasons were mentioned in the manuscript.

Response: We thank the reviewer pointing this out. We now add the following sentence to point the readers to the growing season for different crops:

“Growing seasons for major crops in China were indicated in Table 1, and acquired from Major World Crop Areas and Climate Profiles (MWCACP), and the Food and Agriculture Organization of the United Nation (FAO) (Lin et al., 2018; Zhao et al., 2020).”

Line 96-97: Are there any differences in the definition of the growing season between NWCACP and FAO? The authors only mentioned NWCACP and FAO with no details.

Response: The definitions of the growing season between NWCACP and FAO are the same.

About the section 3.3

There are redundant descriptions. For instance, the authors made some comparison to some literature, i.e., Lines 180, 181, 199, 205 and 206. However, they repeat similar discussions later (i.e., Line 270-273). In addition, the comparison does not specifically mention what year or period, making the comparison invalid.

Response: We thank the reviewer pointing this out. We now delete the comparisons between the crop yields losses in China with the annual crop productions in other countries in section 3.3.

Instead, we only kept them at the discussions line 249-line 254:

“Combing the annual crop production from the Statistical Yearbook of China, we estimated that the surface ozone in China could cause an average of 26.42 million metric tons losses (Mt) of

wheat production from 2010 to 2017. These losses are even comparable to the annual average wheat production during the same period in Paris, which is the fifth largest wheat production in the world (<http://www.fao.org/faostat/en/#data/QC>, accessed December 12, 2021). We also estimated that the surface ozone exposure could cause 18.58 Mt losses of rice production in China, comparable to the annual rice production in Philippines, the world's 8th largest rice production.”

Line 191-193

The authors highlight the differences in the calculations of the growing season lead to different years for the lowest values. In my opinion, this is nothing new worth highlighting.

Response: We agree with the reviewer that we should not highlight the calculations of different growing seasons. Instead, we revised the sentence in line 192, and highlighted the seasonality of O₃ concentration:

“The CPL for double early and late rice both peak in 2014, but with different years for the lowest values (Tables S9 and S10), highlighting the seasonal variations of O₃ concentration between different growing seasons (Table 1).”

Line 212-213:

It seems to be contradictory that the authors stated the lowest CPL in northeast China, and then emphasize Heilongjiang is the highest. Later on, I realize the statement of highest yield in Heilongjiang is probably in another year. The authors need to carefully check out the entire manuscript to make the statement clear and readable.

Response: We thank the reviewer pointing this out. We now rewrite this sentence to avoid confusion in line 216-217:

“We estimated that the ozone-induced CPL for soybean ranges from 1.09 Mt in 2017 to 1.84 Mt in 2010, with 8-year annual average of 1.52 Mt (Fig. 4; Table S13). Heilongjiang, Anhui, and Henan are the three provinces with the highest CPL, with 0.69, 0.17, 0.16 Mt loss on average individually (Table S13).”

The Discussions section needs a total revision.

Response: We appreciate the reviewer's comment about the discussion. We now rewrite this part. In the new discussion section, we talked about the decreasing trend of ozone-induced crop yields losses in China after 2013, the future climate and population changes on crops, and also the uncertainties for our study originating from the model, the emission inventories and the concentration-response function we used. We also rewrote the Results and Summary section to show the results only.

“4 Discussions

Surface ozone emerged as an important environmental issue in China, and were shown increasing trend in major megacities for the past few years using both modelling and observation data (Lu et al., 2018, 2019; 2020; Li et als., 2020; Liu and Wang, 2020a,b; Ni et al., 2018; Wang et al., 2020), though strict clean air regulations have been implemented after 2013. Exposure to high concentrations of surface ozone not only poses threat to human health, but also cause damages to crop. Our study presented a comprehensive analysis on the impact of surface ozone exposure on four major crop production loss in China, including wheat, rice (double early and late rice, single rice), maize (north maize and south maize), and soybean. Unlike the surface ozone trend, we showed that the national crop yields for major crops in China usually peaks in

2014 or 2015, shortly after the strict clean air regulations after 2013. The decreasing trend of crop yield losses associated with surface ozone exposure was mainly explained by the fact that the surface ozone in China were increasing in urban areas, while decreasing in the rural areas (Li et al., 2022), where the major crops are planted. Nonetheless, the relatively higher ozone, especially compared with developed countries, such as United States and Japan (Lu et al., 2018), are still posing great threats to crop productions in China. Combing the annual crop production from the Statistical Yearbook of China, we estimated that the surface ozone in China could cause an average of 26.42 million metric tons losses (Mt) of wheat production from 2010 to 2017. These losses are even comparable to the annual average wheat production during the same period in Paris, which is the fifth largest wheat production in the world (<http://www.fao.org/faostat/en/#data/QC>, accessed December 12, 2021). We also estimated that the surface ozone exposure could cause 18.58 Mt losses of rice production in China, comparable to the annual rice production in Philippines, the world's 8th largest rice production. Transferring to economic values, we estimated the surface ozone exposure could cost more than 20 billion \$ losses, representing more than 0.20% of annual average Gross Domestic Product (GDP) in China from 2010 to 2017. The latest edition of the State of Food Security and Nutrition in the World estimated that between 720 and 811 million people in the world faced hunger in 2020, with 161 million increasing compared with 2019, and nearly 2.37 billion people did not have access to adequate food, with no regions spared (FAO, 2021). Therefore, reducing surface ozone pollution could not only bring the benefits of reducing ozone-related premature deaths, but also bring the benefits of control the global hunger and malnutrition issues, thus helping to reach the Sustainable Development Goal 2 of "Zero Hunger". Meanwhile, Chinese population are projected to continue to increase and peak around 2025 under all the shared socioeconomic pathways (SSPs, Chen et al., 2020), making it more urgent to improve the crop productions by all means.

Uncertainties exist in the design of our study, including the coarse resolution of the global transport model we used, the regional emission inventories, as well as the concentration-response functions. From the model evaluation, we learnt that our model tends to overestimate the annual MDA8 O₃ concentration in China. However, through sensitivity experiences, Wang et al. (2022) showed that model biases in ozone were likely to have a relatively small impact on estimated production losses. The uncertainties from the changes in growing seasons, and the concentration-response functions tend to have larger effects. We propose that further studies, using high-resolution bias-corrected ozone concentration data and region-specific response functions, need to be carried out to quantify the negative effects of surface ozone on crops. In our study, we also did not consider the possible climate changes on the crop productions. However, previous studies have demonstrated that temperature increases could significantly reduce the crop productions as well (Asseng et al., 2015; Wiebe et al., 2015; Liu et al., 2016; Zhao et al., 2016, 2017). Despite these limitations and uncertainties, our study strives to estimate the long-term negative effects from surface ozone exposure in China before and after the clean air action in China. These estimations could provide the government and policy-makers useful references to be taken into account of the detrimental effects of ozone exposure on crop productions in China when making regional-specific ozone control policies."

There are many sentences duplicated in several places. For instance, Line 81-82, "In general, the model simulated AOT40 values were lower than the observation data, with normalized mean bias ranging from -5% in 2015 to -28% in 2017". Line 244-246, and 278-279

Response: We thank the reviewer pointing this out. We now remove the repeated sentences in 244-246, 278-279.

Lines 238-246: very wordy, should be trimmed substantially and make clear what major message the authors want to convey.

Response: We thank the reviewer's comment. We now shorten this sentence, and put them into the beginning of the section "5 Conclusions and Summary", line 277:

"In this study, we applied chemical transport model simulation with updated annual anthropogenic emission inventory to study the long-term trend of O₃-induced crop production losses from 2010 to 2017 in China."

The second paragraph of Discussions only lay out many results without any depth.

Response: Please see our response above. We have rewritten our Discussion and Conclusion sections. All the detailed results were put into the new "Conclusions and Summary" section, and we discussed the uncertainties in the design of our study, the possible shortage of this study, and the policy implications as indicated from this study. We believe our discussion is more clear now.

The authors mentioned many times of the year 2014 (i.e., Lines 185, 192, and 219), but what special with the year has never been mentioned.

Response: We thank the reviewer pointing this out. In our study design, the specific period of 2010-2017 was chosen to cover the emission changes before and after the China's Air Pollution Prevention and Control Action Plan (APPCAP) which was established in 2013. Studies have shown that the anthropogenic emissions for major air pollutants are seen significant decline (Zheng et al., 2018; Zhang et al., 2019) after 2013. However, the summertime ozone in China urban regions have been reported to continue to increase after 2013 (Lu et al., 2018, 2020; Li K. et al., 2019, 2020; Li X. et al., 2022). In our study, we showed that crop yield losses associated with ozone exposure generally peak before 2014, and then decrease thereafter, demonstrating the fact that the surface ozone in rural China have a decreasing trend, consistent with the long-term observations (Li X. et al., 2022).

Reference:

Li, K., Jacob, D. J., Zhang, Q., Liao, H., Bates, K. H. and Shen, L.: Anthropogenic drivers of 2013–2017 trends in summer surface ozone in China, *Proc. Natl. Acad. Sci.*, 116(2), 422–427, doi:10.1073/pnas.1812168116, 2019a.

Li, K., Jacob, D. J., Shen, L., Lu, X., De Smedt, I., and Liao, H.: Increases in surface ozone pollution in China from 2013 to 2019: anthropogenic and meteorological influences, *Atmos. Chem. Phys.*, 20, 11423–11433, <https://doi.org/10.5194/acp-20-11423-2020>, 2020.

Li, X., Yuan, B., Parrish, D. D., Chen, D., Song, Y., Yang, S., Liu, Z. and Shao, M.: Long-term trend of ozone in southern China reveals future mitigation strategy for air pollution, *Atmos. Environ.*, 269(118869), doi:<https://doi.org/10.1016/j.atmosenv.2021.118869>, 2022.

Lu, X., Hong, J., Zhang, L., Cooper, O. R., Schultz, M. G., Xu, X., Wang, T., Gao, M., Zhao, Y. and Zhang, Y.: Severe Surface Ozone Pollution in China: A Global Perspective, *Environ. Sci. Technol. Lett.*, 5(8), 487–494, doi:10.1021/acs.estlett.8b00366, 2018.

Lu, X., Zhang, L., Wang, X., Gao, M., Li, K., Zhang, Y., Yue, X. and Zhang, Y.: Rapid Increases in Warm-Season Surface Ozone and Resulting Health Impact in China Since 2013, *Environ. Sci. Technol. Lett.*, doi:10.1021/acs.estlett.0c00171, 2020.

Zhang, Q., Zheng, Y., Tong, D., Shao, M., Wang, S., Zhang, Y., Xu, X., Wang, J., He, H., Liu, W., Ding, Y., Lei, Y., Li, J., Wang, Z., Zhang, X., Wang, Y., Cheng, J., Liu, Y., Shi, Q., Yan, L., Geng, G., Hong, C., Li, M., Liu, F., Zheng, B., Cao, J., Ding, A., Gao, J., Fu, Q., Huo, J., Liu, B., Liu, Z., Yang, F., He, K. and Hao, J.: Drivers of improved PM_{2.5} air quality in China from 2013 to 2017, *Proc. Natl. Acad. Sci. U. S. A.*, 116(49), 24463–24469, doi:10.1073/pnas.1907956116, 2019.

Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., Qi, J., Yan, L., Zhang, Y., Zhao, H., Zheng, Y., He, K. and Zhang, Q.: Trends in China's anthropogenic emissions since 2010 as the consequence of clean air actions, *Atmos. Chem. Phys.*, 18(19), 14095–14111, doi:10.5194/acp-18-14095-2018, 2018.

The authors mentioned spatial heterogeneity across regions and provinces, but with no details.
Response: We now delete this sentence in our new Section 5 “Conclusions and Summary”.

The last few sentences talked about these ozone pollution control over different regions. However, the major role of ozone on CPL over these regions have not been well discussed at all.
Response: We appreciate the reviewer's question. We now remove this sentence since we made great efforts to reconstruct our results and discussions and felt that these last few sentences are not necessary anymore.

Minor comments:

Line 59-60: This message of the sentence is not clearly stated. The sentence writes that previous studies have focused on crop production loss from ozone at the global scale. Have any of the studies focused on China?

Response: There are published studies focusing on the China crop yields loss. To avoid confusion, we now rewrite the sentences about the novelty about our study:
“To date, very few studies have investigated the long-term trends and spatial patterns of ozone impacts on crop production in China. Previous studies have been mainly focus on a specific region of China, such as NCP (Zhang et al., 2017; Hu et al., 2020; Feng et al., 2020), or Yangtze River Delta (Wang et al., 2012). In this study, we focus on the long-term ozone-exposure impact analysis from 2010 to 2017 in China to assess the yield losses of four major crops (wheat, maize, rice, and soybean) and evaluate their associated economic losses. The specific period of 2010-2017 was chosen to cover the emission changes before and after the APPCAP established in 2013. Previous studies have been reporting the crop yield losses in one year (e.g., Lin et al., 2018; Yi et al., 2018; Feng et al., 2019a,b), or several years after the APPCAP (Zhao et al., 2019; Wang et al., 2022), and our study aims to present a comprehensive analysis of ozone-induced crop yield losses and economic impacts in the agriculture sector before and after the China APPCAP. Such an analysis is expected to provide scientific support to policymakers for their decision making.”

Line 85: Matrixes: should be metrics?

Response: We changed to “metrics”

Line 129: The price for each crop during 2010-2017 is given based on the min/max, however, the readers do not know the specific price corresponding to each year. A table might be useful to lay out the prices.

Response: We appreciate the question. We now add a new Table in the supporting:

Table S1. The crop market prices for the major crops in China, acquired from the FAOSTAT (unit of \$ per ton; <http://www.fao.org/faostat/>, last accessed 26th, March, 2020).

Year	2010	2011	2012	2013	2014	2015	2016	2017 ¹
Wheat	279.5	321.5	323.3	355.1	377	391.4	385.1	384.5
Soybean	738.5	803.4	841.5	677.9	869.7	808.2	581.5	753.1
Maize	273.3	321.8	383.5	489.1	441.9	432.4	264.3	252.2
Rice	296.6	403.9	456.4	492.3	457.9	508.9	559.9	508.9

Line 162: RYLs for specific crops should be clearly written.

Response: We now revise this sentence:

“The RYLs for the double rice, range from 10.71% in Anhui to 7.11% in Yunan for the 8-year average (Table S4).”

Line 178: The citation of the Statistical Yearbook of China should be added.

Response: Thanks for pointing out. We now add the citation for the Statistical Yearbook of China.

“From the Statistical Yearbook of China, the national wheat production increased from 115.19 million Mt in 2010 to 134.34 million Mt in 2017, which are mainly planted in the NCP (<http://www.stats.gov.cn/tjsj/ndsj/2019/indexeh.htm>, last accessed December 9th, 2021).”

Line 183: “CPL” should be replaced by “wheat CPL”

Response: we made the change following the reviewer’s suggestion.

“Fig. 5 shows the wheat CPL for each province in China from 2010 to 2017”

Line 223: studies changed to study.

Response: We changed to “study”.

Table S13: The production loss of China in 2017 was miscalculated and “374” should be replaced by “74”. Please carefully check all the calculations in the tables.

Response: We thank the reviewer finding this out. We now change to the right number.

Line 235: References should be added.

Response: We now add the following references here:

“Exposure to high concentrations of surface ozone not only poses threat to human health, but also cause damages to crops (Krupa et al., 1998; EPA, 1996; EEA 1999; Mauzerall & Wang, 2001).”

Reference

EEA, 1999. Environmental assessment Report No. 2. Environment in the European Union at the turn of the century. European Environmental Agency, Copenhagen, 446pp.

EPA, 1996. Air quality criteria for ozone and related photochemical oxidants. United States Environmental Protection Agency (EPA), pp. 1–1 to 1–33.

Krupa, S. V., Nosal, M. and Legge, A. H.: Short communication A numerical analysis of the combined open-top chamber data from the USA and Europe on ambient ozone and negative crop responses, *101*, 157–160, 1998.

Mauzerall, D. L. and Wang, X.: Protecting agricultural crops from the effects of tropospheric ozone exposure: reconciling Science and Standard Setting in the United States, Europe, and Asia, *Annu. Rev. Energy Environ.*, *26*(1), 237–268, doi:10.1146/annurev.energy.26.1.237, 2001.

Line 237: The authors said the previous studies only focused on small regions. However, in the introduction, the authors mentioned there are studies with a focus of the globe. This seems to be contradictory.

Response: We thank the reviewer pointing this out. Here we meant small regions in China. There are significant number of studies focusing on global. We now rewrite this sentence to be more precise:

“Previous studies have been using modelling results or observation data to study the crop production losses in China for a single year (Lin et al., 2018; Feng et al., 2019b), or several years at specific regions, such as North China Plain (Zhang et al., 2017; Hu et al., 2020; Feng et al., 2020), or Yangtze River Delta (Wang et al., 2012). Some studies also estimated crop yield losses for three or four years in China after 2013 (Zhao et al., 2020; Wang et al., 2022), when the Chinese government implemented the stringent Air Pollution Prevention and Control Action Plan (APPCAP).”

Reference:

Feng, Z., De Marco, A., Anav, A., Gualtieri, M., Sicard, P., Tian, H., Fornasier, F., Tao, F., Guo, A. and Paoletti, E.: Economic losses due to ozone impacts on human health, forest productivity and crop yield across China, *Environ. Int.*, *131*(June), doi:10.1016/j.envint.2019.104966, 2019b.

Feng, Z., Hu, T., Tai, A. P. K. and Calatayud, V.: Yield and economic losses in maize caused by ambient ozone in the North China Plain (2014–2017), *Sci. Total Environ.*, *722*, 137958, doi:10.1016/j.scitotenv.2020.137958, 2020.

Hu, T., Liu, S., Xu, Y., Feng, Z. and Calatayud, V.: Assessment of O₃-induced yield and economic losses for wheat in the North China Plain from 2014 to 2017, *China, Environ. Pollut.*, *258*, 113828, doi:10.1016/j.envpol.2019.113828, 2020.

Lin, Y., Jiang, F., Zhao, J., Zhu, G., He, X., Ma, X., Li, S., Sabel, C. E. and Wang, H.: Impacts of O₃ on premature mortality and crop yield loss across China, *Atmos. Environ.*, *194*(July), 41–47, doi:10.1016/j.atmosenv.2018.09.024, 2018.

Wang, X., Zhang, Q., Zheng, F., Zheng, Q., Yao, F., Chen, Z., Zhang, W., Hou, P., Feng, Z., Song, W., Feng, Z. and Lu, F.: Effects of elevated O₃ concentration on winter wheat and rice yields in the Yangtze River Delta, *China, Environ. Pollut.*, *171*, 118–125, doi:10.1016/j.envpol.2012.07.028, 2012.

Wang, Y., Wild, O., Ashworth, K., Chen, X., Wu, Q., Qi, Y. and Wang, Z.: Reductions in crop yields across China from elevated ozone, *Environ. Pollut.*, 292(118218), doi:<https://doi.org/10.1016/j.envpol.2021.118218>, 2022.

Yi, F., McCarl, B. A., Zhou, X. and Jiang, F.: Damages of surface ozone: Evidence from agricultural sector in China, *Environ. Res. Lett.*, 13(3), doi:[10.1088/1748-9326/aaa6d9](https://doi.org/10.1088/1748-9326/aaa6d9), 2018.

Zhang, W., Feng, Z., Wang, X., Liu, X. and Hu, E.: Quantification of ozone exposure- and stomatal uptake-yield response relationships for soybean in Northeast China, *Sci. Total Environ.*, 599–600, 710–720, doi:[10.1016/j.scitotenv.2017.04.231](https://doi.org/10.1016/j.scitotenv.2017.04.231), 2017.

Zhao, H., Zheng, Y., Zhang, Y. and Li, T.: Evaluating the effects of surface O₃ on three main food crops across China during 2015–2018, *Environ. Pollut.*, 258, 113794, doi:[10.1016/j.envpol.2019.113794](https://doi.org/10.1016/j.envpol.2019.113794), 2020.

Line 255: It should be written clearly whether the CPL and EL for a particular crop or the total CPL and EL for all four crops.

Response: We appreciate the reviewer's question. We now remove this sentence since we made great efforts to reconstruct our results and discussions and felt that this sentence is not necessary anymore.