



Atmospheric photo-oxidation of myrcene: OH reaction rate constant, gas phase oxidation products and radical budgets

Zhaofeng Tan¹, Luisa Hantschke¹, Martin Kaminski^{1,a}, Ismail-Hakki Acir^{1,b}, Birger Bohn¹, Changmin Cho¹, Hans-Peter Dorn¹, Xin Li^{1,c}, Anna Novelli¹, Sascha Nehr^{1,d}, Franz Rohrer¹, Ralf Tillmann¹, Robert Wegener¹, Andreas Hofzumahaus¹, Astrid Kiendler-Scharr¹, Andreas Wahner¹, and Hendrik Fuchs¹

¹Institute of Energy and Climate Research, IEK-8: Troposphere, Forschungszentrum Jülich GmbH, Jülich, Germany

^anow at: Federal Office of Consumer Protection and Food Safety, Department 5: Method Standardisation, Reference Laboratories, Resistance to Antibiotics, Berlin, Germany

^bnow at: Institute of Nutrition and Food Sciences, Food Science, University of Bonn, Bonn, Germany

^cnow at: State Key Joint Laboratory of Environmental Simulation and Pollution Control, College of Environmental Sciences and Engineering, Peking University, Beijing, China

^dnow at: European University of Applied Sciences, Brühl, Germany

Correspondence: Hendrik Fuchs (h.fuchs@fz-juelich.de)

Abstract. The photo-oxidation of myrcene, a monoterpene species emitted by plants, was investigated at atmospheric conditions in the outdoor simulation chamber SAPHIR. The chemical structure of myrcene consists of one moiety that is a conjugated π -system (similar to isoprene) and another moiety that is a triple-substituted olefinic unit (similar to 2-methyl-2-butene). Hydrogen shift reactions of organic peroxy radicals (RO_2) formed in the reaction of isoprene with atmospheric OH radicals are known to be of importance for the regeneration of OH. Structure-activity relationships (SAR) suggest that similar hydrogen shift reactions like in isoprene may apply to the isoprenyl part of RO_2 radicals formed during the OH oxidation of myrcene. In addition, SAR predicts further isomerization reactions that would be competitive with bi-molecular RO_2 reactions for chemical conditions that are typical for forested environments with low concentrations of nitric oxide. Assuming that OH peroxy radicals can rapidly interconvert by addition and elimination of O_2 like in isoprene, bulk isomerization rate constants of 0.21 s^{-1} and 0.097 s^{-1} ($T = 298 \text{ K}$) for the 3 isomers resulting from the 3'-OH and 1-OH addition, respectively, can be derived from SAR. Measurements of radicals and trace gases in the experiments allowed to calculate radical production and destruction rates, which are expected to be balanced. Largest discrepancies between production and destruction rates were found for RO_2 . Additional loss of organic peroxy radicals due to isomerization reactions could explain the observed discrepancies. The uncertainty of the total radical ($\text{RO}_x = \text{OH} + \text{HO}_2 + \text{RO}_2$) production rates were high due to the uncertainty in the yield of radicals from myrcene ozonolysis. However, results indicate that radical production can only be balanced, if the reaction rate constant of the reaction between hydroperoxy (HO_2) and RO_2 radicals derived from myrcene is lower (0.9 to $1.6 \times 10^{-11} \text{ cm}^3 \text{ s}^{-1}$) than predicted by SAR. Another explanation of the discrepancies would be that a significant fraction of products (yield: 0.3 to 0.6) from these reactions include OH and HO_2 radicals instead of radical terminating organic peroxides. Experiments also allowed to determine the yields of organic oxidation products acetone (yield: 0.45 ± 0.08) and formaldehyde (yield: 0.35 ± 0.08). Acetone and formaldehyde are produced from different oxidation pathways, so that yields of these compounds reflect the branching ratios of the initial OH addition to myrcene. Yields determined in the experiments are consistent with branching ratios expected



from SAR. The yield of organic nitrate was determined from the gas-phase budget analysis of reactive oxidized nitrogen in the chamber giving a value of 0.13 ± 0.03 . In addition, the reaction rate constant for myrcene + OH was determined from the measured myrcene concentration yielding a value of $(2.3 \pm 0.3) \times 10^{-10} \text{ cm}^3 \text{ s}^{-1}$.

25 1 Introduction

Monoterpenes are emitted from vegetation accounting for approximately 160 Tg of the 1000 Tg of biogenic volatile organic compounds (VOCs) that are released into the atmosphere per year (Guenther et al., 2012; Sindelarova et al., 2014). Monoterpenes are highly reactive to the major oxidants in the atmosphere, hydroxyl radicals (OH), ozone (O_3) and nitrate radicals (NO_3) (Atkinson and Arey, 2003) and thus play an important role in ozone and secondary organic aerosol formation (Xu et al.,
30 2015; Zhang et al., 2018; Schwantes et al., 2020).

Myrcene emissions contribute to the total biogenic monoterpene emissions in the range of 2 to 10 % (Guenther et al., 2012). Emission rates highly depend on the type of tree and season of the year (Helmig et al., 2013). In addition, there are hints for anthropogenic sources from the analysis of the composition of indoor air (Kostiainen, 1995). Few studies have been conducted to investigate the oxidation of myrcene (Deng et al., 2018; Atkinson, 1997; Reissell et al., 2002; Kim et al., 2011). In these
35 studies, acetone, formaldehyde and 4-vinyl-pentenal have been identified as major oxidation products from the reaction of myrcene with OH, but yields determined in these studies vary. Lee et al. (2006) reported an organic nitrate yield of 10 % from the oxidation of myrcene by direct measurements using mass spectrometry. The reaction rate constant of myrcene + OH was determined to be $(2.1 \pm 0.2) \times 10^{-10}$ (Atkinson et al., 1986) and $(3.4_{-1.0}^{+1.5}) \times 10^{-10}$ (Hites and Turner, 2009) at room temperature with a discrepancy of up to 60 %. These results demonstrate the photo-oxidation of myrcene requires further investigation.

40 There is no detailed chemical mechanism of myrcene degradation by OH in literature. The acyclic structure of myrcene consists of two parts, an isoprenyl part ($\text{CH}_2=\text{CH}-\text{C}(=\text{CH}_2)\text{CH}_2-$ moiety) and another part that is structurally similar to 2-methyl-2-butene ($(\text{CH}_3)_2\text{C}=\text{CH}-\text{CH}_2-$ moiety) (Fig. 1). Recent investigations of the oxidation of isoprene (Fuchs et al., 2013; Peeters et al., 2014; Wennberg et al., 2018; Novelli et al., 2020) revealed that organic peroxy radicals (RO_2) formed after the attack of OH to isoprene can rapidly interconvert, so that fast H-atom migration reactions of specific RO_2 isomers with
45 initially low yield can significantly gain in importance. These reactions impact atmospheric chemistry most, if they become competitive with bi-molecular loss reactions such as reactions with nitric oxide (NO) and hydroperoxy radicals (HO_2). In the case of isoprene, these isomerization reactions can eventually regenerate OH radicals, so that a high OH regeneration efficiency of 0.5 can be sustained in the atmosphere also in regions, where radical termination reactions have been thought to dominate the fate of radicals (Novelli et al., 2020). Therefore, radical regeneration from isomerization reactions helped partly
50 to explain observations of unexpectedly high OH concentrations in field experiments (Lelieveld et al., 2008; Hofzumahaus et al., 2009; Whalley et al., 2011). Radical regeneration from RO_2 isomerization reactions have also been shown to play a role in the oxidation of methacrolein (Fuchs et al., 2014; Crounse et al., 2012), 3-pentanone (Crounse et al., 2013), glyoxal (Lockhart et al., 2013), n-hexane, 2-hexanol (Praske et al., 2018), hydroxymethyl hydroperoxides (Allen et al., 2018), and 2-hydroperoxy-2-methylpentane (Praske et al., 2019).



55 In this study, the oxidation of myrcene by OH was investigated in the atmospheric simulation chamber SAPHIR (Simulation of Atmospheric Photochemistry In a Large Reaction Chamber) at Forschungszentrum Jülich. Experiments were performed under controlled conditions with concentrations of trace gases and radicals typical for atmospheric conditions. Experiments aim for adding to the determination of a degradation mechanism for myrcene.

2 Oxidation mechanism of myrcene

60 Myrcene is an acyclic hydrocarbon with three unsaturated carbon double bonds to which OH preferentially adds. One end of the myrcene molecule is structurally similar to isoprene and the other end is similar to 2-methyl-2-butene (Fig. 1). The myrcene oxidation by OH forms ten isomers of peroxy radicals (MyO_2) (Fig. 2). Structure-activity relationships (SAR) by Peeters et al. (2007) suggest that OH preferably adds to the double bond of the isolated $-\text{CH}=\text{C}(\text{CH}_3)_2$ moiety producing two peroxy radical species with yields of 17 % (7-OH-6-OO radical) and 31 % (6-OH-7-OO radical) (Fig. 2). When OH adds
65 to the conjugated carbon double bonds, six isomeric hydroxy peroxy radicals are formed like in the OH or NO_3 oxidation of isoprene (Peeters et al., 2009, 2014; Vereecken et al., 2021a). Addition to the $\text{C}_{3'}$ position yields two allyl-like hydroxy radical isomers producing E-3'-OH-1-OO, 3'-OH-3-OO and Z-3'-OH-1-OO peroxy radicals after the subsequent addition of O_2 . A total yield of 31 % is estimated by SAR (Peeters et al., 2007) for these three isomers. Similarly, OH addition to the C_1 position followed by addition of O_2 to the allyl radical structure leads to the formation of E-1-OH-3'-OO, 1-OH-2-OO
70 and Z-1-OH-3'-OO radicals with a total yield of 17 %. In the isoprene mechanism developed by Peeters et al. (2009, 2014); Novelli et al. (2020); Vereecken et al. (2021a), a central element is the fast interconversion reactions between the OH-adducts and corresponding OH peroxy radicals that proceed by addition and elimination of O_2 . Similar reactions are expected for the isoprene structure in myrcene (Fig. 2) establishing a coupled equilibrium between MyO_2 isomers.

For OH attack on the C_2 or C_3 position of myrcene, the resulting OH adducts are not resonance stabilized and are hence
75 not as favorable as the allylic-type radicals produced from C_1 or $\text{C}_{3'}$ addition. Yields of these MyO_2 radicals are expected to be less than 2 % and therefore the chemistry of these two minor isomers is not further discussed in this work.

All MyO_2 isomers can undergo bi-molecular reactions with NO, HO_2 and other RO_2 species. Specific reaction rate constants have not been measured, but values are expected to be within the range of typical rate constants for organic peroxy radicals. For example, SAR by Jenkin et al. (2019) suggests a value of $2.2 \times 10^{-11} \text{ cm}^3 \text{ s}^{-1}$ ($T = 298 \text{ K}$) of the reaction rate
80 constants for the reaction of monoterpene derived RO_2 (including MyO_2) with HO_2 . Reaction products of the reaction of RO_2 with HO_2 are expected to be organic peroxides.

The reaction of the 6-OH-7-OO and 7-OH-6-OO radicals with NO produce mainly 4-vinyl-4-pentenal, acetone and HO_2 (Fig. 3) as shown in experimental studies (Orlando et al., 2000; Reissell et al., 2002; Lee et al., 2006) as well as predicted by SAR (Vereecken and Peters, 2009, 2010). Reissell et al. (2002) and Lee et al. (2006) suggested additional pathways for
85 the alkoxy radical that is formed in the reaction with NO of these two MyO_2 species. These pathways would include rearrangement reactions followed by reaction with NO and fragmentation. This was suggested to explain the observation of organic species with various masses ($m/z=71$, $m/z=113$, $m/z=115$) detected by the PTR-MS instrument in Lee et al. (2006).



Similarly, Böge et al. (2013) suggested a reaction pathway of 6-OH-7-OO and 7-OH-6-OO radicals to explain observed terpenylic acid in their experiments. The reaction of the other two most abundant radicals, 1-OH-2-OO and 3'-OH-3-OO, with
90 NO would lead to the formation of HO₂ and formaldehyde together with 2-methyldiene-6-methyl-5-heptenal and 1-vinyl-5-methyl-4-hexanone, respectively (Fig. 3).

In addition to bi-molecular reactions, uni-molecular can be of importance for specific MyO₂. Due to the similarity of the conjugated double bond structure in myrcene and isoprene, it can be expected that H-shift reactions found to be important in isoprene (Peeters et al., 2009, 2014; Fuchs et al., 2013; Wennberg et al., 2018; Novelli et al., 2020) apply for corresponding
95 MyO₂ radicals. Therefore, Z-3'-OH-1-OO and Z-1-OH-3'-OO radicals are expected to undergo α-OH 1,6-H migration followed by O₂ addition leading to hydroperoxy peroxy radicals (Fig. 4). As a first approximation, reaction rate constants can be assumed to be in the order of 1 s⁻¹ at 298 K like for corresponding reactions in isoprene (Peeters et al., 2014). Similar products from further isomerization and decomposition reactions could be expected as predicted for isoprene (Peeters et al., 2014). This would also lead to the regeneration of HO_x radicals.

100 According to SAR by Vereecken and Nozière (2020), E-3'-OH-1-OO undergo an allylic 1,7H-shift (Fig. 4). The reaction rate constant is estimated by this SAR to be high with a value of 8 s⁻¹ (T = 298 K). In addition, E-1-OH-3'-OO can undergo an allylic 1,6H-shift isomerization reaction with a fast isomerization reaction rate constant of 2 s⁻¹ (T = 298 K). Products likely undergo fast ring-closure reactions on the dimethyl double bond with rates on the order of 1 s⁻¹ (pers. comm. Vereecken, 2021b).

105 As a consequence of the equilibration between different MyO₂ isomers that originate from the 3'-OH or 1-OH addition, a significant fraction of the MyO₂ can be removed through the H-shift reaction channels, if rates of competing bi-molecular reactions of all RO₂ isomers are low enough. In order to estimate effective bulk MyO₂ isomerization reaction rates, the distribution of MyO₂ isomers is estimated by using reaction rate constants for the oxygen addition and elimination reactions recommended for isoprene (Novelli et al., 2020). This results in a total bulk MyO₂ loss rate of approximately 0.21 s⁻¹ and
110 0.097 s⁻¹ (T = 298 K) for the 3 isomers resulting from the 3'-OH and 1-OH addition, respectively. This means that isomerization reactions are becoming competitive for nearly half of the total MyO₂ (Fig. 1) isomers for chemical conditions with NO mixing ratios lower than 1 ppbv. It is worth noting that isomerization reactions rate constants have a strong temperature dependence, so that their impact can be significantly enhanced at higher temperatures. However, all rate constants by SAR predictions have also a high uncertainty of at least a factor of two and the uncertainty might be as high as a factor of 10 (Vereecken
115 and Nozière, 2020).

3 Methods

3.1 Experiments in the SAPHIR chamber

The experiments were conducted in the outdoor atmospheric simulation chamber SAPHIR. SAPHIR has a cylindrical shape with double walls made of Teflon (FEP) film (length: 18 m diameter: 5 m, volume: 270 m³). The space between the double
120 walls is permanently purged with clean air to avoid diffusion of impurities from ambient air into the inner chamber. The walls



are transmissive for the entire solar UV and visible spectrum. The chamber is operated with synthetic air that is produced from evaporated liquid nitrogen and oxygen of highest purity (Linde, purity > 99.99990 %). It is kept at a slight overpressure of 35 Pa that is maintained by a replenishment flow, which compensates for leakages and the sampling flow of analytical instruments. As a consequence, trace gases are diluted with a rate constant that is equivalent to a lifetime of approximately
125 17 hours. The air in the chamber can be exposed to sunlight by opening a shutter system. When the chamber film is exposed to solar radiation, nitrous acid (HONO), formaldehyde and acetone are released. The source strengths range between 100 to 200 pptv/h. The photolysis of HONO leads to a continuous increase of nitrogen oxide concentrations in the chamber and is a significant source for hydroxyl radicals (OH). More details of the SAPHIR can be found in previous publications (Bohn et al., 2005; Rohrer et al., 2005, e.g.).

130 In total 4 experiments investigating the oxidation of myrcene by OH were performed (Table 1), two of which were done at medium levels of nitric oxide (NO) (18 August 2012: 0.18 to 0.43 ppbv NO (Fig. 5) and 22 August 2012: 0.15 to 0.30 ppbv of NO (Fig. S1 Supplement)) while lower NO mixing ratios below 0.11 ppbv were achieved in the other two experiments (17 July 2013, Fig. S2 Supplement, and 18 July 2013, Fig. S3 Supplement).

The experimental procedure was similar in all experiments. The chamber was cleaned in the night before the experiment by
135 flushing the chamber with a high flow of synthetic air to remove all trace gases from previous experiments. Figure 5 shows, as an example, time series of trace gases for the experiment conducted on 22 August 2012. Experiments started in the morning with humidification of the chamber air in the dark. This was achieved by flushing evaporated Milli-Q water into the chamber together with a high flow of synthetic air. The chamber air was exposed to sunlight approximately for 1 hour without the presence of any additional reactant to determine the source strengths of chamber sources for formaldehyde for the specific
140 experiment. In the experiments in 2013, approximately 50 ppbv ozone produced by a silent discharge ozonizer (O3onia) was injected to suppress NO in the reaction with O₃.

Air mixtures of myrcene (Sigma-Aldrich, purity 99 %) were premixed in a canister (stainless steel with Silconert coating). The mixture was injected into the chamber air by calibrated mass flow-controllers to reach myrcene mixing ratios of several ppbv. Injections were done 2 times (3 ppbv each) in experiments with medium NO and four injections with smaller concen-
145 trations (approximately 2 ppbv) were done in the other experiments.

Additional reference experiments were performed using the same procedure like for the experiments with myrcene, but either with one injection of 150 ppmv of methane (29 May 2020 (Fig. S4 Supplement) and 10 July 2013) or three injections of 5 ppbv of α -pinene (03 September 2019 (Fig. S5 Supplement)). These experiments were used to evaluate the accuracies of the procedures that were used to analyse the organic nitrate formation and radical budgets in the experiments with myrcene.

150 3.2 Measurement of trace gas concentrations

Table 2 summarizes properties of instruments used in this work. The set of instruments was similar like used in previous experiments investigating the photochemistry of organic compounds (Kaminski et al., 2017; Fuchs et al., 2018; Novelli et al., 2018; Rolletter et al., 2019). Therefore, only a brief description is given here. Ozone was detected by a UV photometer (Ansyco). NO concentrations were measured by a chemiluminescence instrument (Eco Physics) that also detected nitrogen



155 dioxide (NO_2) after conversion to NO in a photolytic converter. Methane, water vapour and carbon monoxide concentrations were measured by a cavity ring-down instrument (Picarro). Nitrous acid (HONO) was detected by a custom-built long-path absorption photometry (LOPAP) (Li et al., 2014). Photolysis frequencies were derived from solar actinic flux measurements by a spectroradiometer outside the chamber. Calculations take into account the reduction of radiation by the chamber construction elements and the Teflon film (Bohn and Zilken, 2005).

160 Myrcene was detected by gas chromatography coupled with a flame ionization detector (GC-FID) (Wegener et al., 2007) and by a proton-transfer-reaction mass-spectrometer (PTR-MS, Ionicon). The PTR-MS instrument was not calibrated for myrcene. Therefore, the signal was scaled to match the GC-FID measurements and used because of its higher time resolution (40 s) compared to that of GC-FID (30 min). Acetone and acetaldehyde were also measured by GC-FID. Formaldehyde (HCHO) measurements were performed by a Hantzsch instrument (AeroLaser) or by differential optical absorption spectrometry (DOAS).

165 OH reactivity (k_{OH}), which is the pseudo first-order rate constant of the OH radical loss, was measured by a laser flash photolysis - laser induced fluorescence instrument (Lou et al., 2010; Fuchs et al., 2017).

3.3 Measurement of radical concentrations

Measurements of OH radicals were performed by differential optical absorption spectrometry (DOAS) (Dorn et al., 1995) and laser-induced fluorescence (LIF) (Tan et al. (2017) and references therein). The DOAS instrument was not available in the

170 experiment on 16 August 2012. In the experiment 6 days later (22 August 2012), OH concentrations measured by the LIF instrument were consistently 20 % higher than those measured by DOAS that is considered to be an absolute standard. The difference is larger than the combined 1σ uncertainty (14.5 %) of the measurements and is probably caused by a calibration error of the LIF instrument. This assumption is supported by the observed decay of myrcene which is caused by the reaction with OH. The decay can be accurately described with LIF data if these are reduced by 20 % in both experiments (Fig. S6

175 Supplement, Section 6.1). Therefore, OH measurements by LIF were scaled by a factor of 0.8 for both experiments. For the experiments in 2013, OH measurements by LIF and DOAS well agreed within 5 %. For the analysis of this work, OH measurements by the DOAS instrument are used if available.

In addition to OH, peroxy radicals (HO_2 and RO_2) can be detected by the LIF instrument after chemical conversion to OH. The conversion of HO_2 radicals is accomplished by adding excess NO in a second low-pressure fluorescence cell, in which the

180 sum of HO_2 and OH concentrations is measured (Fuchs et al., 2011). The measurement of RO_2 is accomplished in a 2-stage system (ROxLIF) in which all atmospheric RO_x radicals are first converted to HO_2 by added NO and CO in a flow reactor (Fuchs et al., 2008). This is followed by HO_2 to OH conversion with additionally added NO in a fluorescence cell. In this case, RO_2 is determined as the difference of RO_x and OH + HO_2 . The operational parameters of the reactor are optimized to maximize the detection sensitivity for methyl peroxy radicals (CH_3O_2). However, it was found in previous studies that

185 specific RO_2 species from other VOCs may be converted less efficiently in the reactor, if their conversion to HO_2 needs more reaction steps and therefore more reaction time than in case of CH_3O_2 (Fuchs et al., 2008). Alternative explanation for the lower sensitivity could be caused by the reversibility of the MyO_2 formation at low O_2 in the reactor, which slow down the conversion to HO_2 .



For the evaluation of data in this work, the ROxLIF instrument was calibrated for CH₃O₂ and myrcene RO₂ radicals (MyO₂) that were produced by reaction of OH with the corresponding VOC in the radical calibration source as described by Fuchs et al. (2008). The detection sensitivity was only half as high for MyO₂ compared to CH₃O₂. The lower sensitivity could hint that some specific MyO₂ isomers are not efficiently converted to HO₂ for conditions inside the conversion reactor. The difference in the detection sensitivity for MyO₂ and other RO₂ species like CH₃O₂ adds to the uncertainty of measurements, because a mixture of different peroxy radicals is present during the experiments and the exact distribution of RO₂ species is not known. For the analysis in this work, the fraction of MyO₂ is estimated as described in the next subsection, in order to account for the lower sensitivity.

3.4 VOC reactivity and RO₂ speciation in myrcene experiments

The total OH reactivity (k_{OH}) that was measured in the experiments consists of reactivity from organic and inorganic compounds. The reaction of OH with most of the organic compounds leads to the formation of RO₂ radicals. Therefore, it is useful to distinguish between OH reactivity from those compounds unmeasured (k_{OHVOC}) and OH reactivity calculated from measured concentrations of inorganic compounds (CO, NO, NO₂) and formaldehyde:

$$k_{\text{OHVOC}} = k_{\text{OH}} - (k_8[\text{HCHO}] + k_9[\text{CO}] + k_{12}[\text{NO}_2] + k_{13}[\text{NO}]) \quad (1)$$

k_i are bimolecular rate constants of reactions R_{*i*} listed in Table 3. For the reference experiment with methane, only methane contributed to the VOC reactivity, because its oxidation product is formaldehyde, which does not produce RO₂ in the reaction with OH. Therefore, methyl peroxy radicals are the only RO₂ radical expected in that experiment. For the experiments with myrcene, the VOC reactivity includes the reactivity from myrcene and from partly unmeasured oxygenated organic compounds (OVOC) that are products of the myrcene oxidation. The OH reactivity from myrcene ($k_{\text{OHmyrcene}}$) can be calculated from measured myrcene concentrations and the rate constant of its reaction with OH (k_7 , Table 3):

$$k_{\text{OHmyrcene}} = k_7[\text{myrcene}] \quad (2)$$

Assuming that each OH reaction with an organic compound except formaldehyde in the experiments in this work leads to the formation of one RO₂ radical and all RO₂ radicals have similar chemical lifetimes, the distribution of RO₂ species from different VOCs is similar to the distribution of the OH reactivity from the VOCs. Therefore, the concentration of RO₂ derived from myrcene (MyO₂) can be approximated by scaling the total measured organic peroxy radical concentration ($[\text{RO}_2]_m$) that is determined assuming the same instrument sensitivity for all RO₂ with the ratio of OH reactivity from myrcene ($k_{\text{OHmyrcene}}$) to the total OH reactivity from VOCs (k_{OHVOC}). Taking also into account that the sensitivity of the instrument for MyO₂ (S_{MyO_2}) is reduced compared to the sensitivity for CH₃O₂ ($S_{\text{CH}_3\text{O}_2}$), the concentration of MyO₂ can be calculated from observed quantities:



$$[\text{MyO}_2] = \frac{1}{\frac{S_{\text{MyO}_2}}{S_{\text{CH}_3\text{O}_2}} + \frac{k_{\text{OHVOC}}}{k_{\text{OHmyrcene}}} - 1} [\text{RO}_2]_m \quad (3)$$

RO₂ species originating from other hydrocarbons than myrcene (RO_{2,other}) can be calculated using the remaining fraction
220 of the VOC reactivity:

$$[\text{RO}_{2,\text{other}}] = [\text{MyO}_2] \left(\frac{k_{\text{OHVOC}}}{k_{\text{OHmyrcene}}} - 1 \right) \quad (4)$$

Equations 3 and 4 allow calculating a more realistic total RO₂ concentration by taking the sum of MyO₂ and RO_{2,other}
instead of using the measured RO₂ concentration, if the same instrument sensitivity for all RO₂ is assumed. An example of the
result of this calculation is shown in Fig. 5 demonstrating that MyO₂ was the dominant RO₂ species right after each myrcene
225 injection. However, these values have a high uncertainty that cannot easily be quantified, because, for example, the detection
sensitivity for RO₂ produced in the reaction of OH with oxidation products may also differ from that of methyl peroxy radicals.
The uncertainty of calculated RO₂ concentrations is lowest after each injection of myrcene, because the total OH reactivity
and therefore RO₂ production is dominated by myrcene. Therefore, the analysis of radical production and destruction rates in
this work (Section 6.4) focuses on the times right after each myrcene injection.

230 4 Experimental determination of organic nitrate yields

The determination of organic nitrate yields from photochemical VOC oxidation makes use of a budget analysis of the sum of
NO_x and HONO in the chamber air. In the following, the sum of NO, NO₂ and HONO concentrations is defined as NO_y^{*},
which can be calculated from measurements of these species (Table 2).

In the sunlit SAPHIR chamber, the only source of reactive nitrogen is the emission of HONO from the chamber film
235 (Section 3.1). The source strength $Q(\text{HONO})$ is variable and depends on solar ultraviolet radiation, temperature and relative
humidity (Rohrer et al., 2005). HONO is photolyzed to OH and NO (R1), which is further oxidized to NO₂ by reactions with
O₃, HO₂ (Reaction R5), and RO₂ (Reaction R11). The sum of NO_y^{*} is chemically lost in the chamber by reactions forming
nitric acid (HNO₃) (Reaction R12) and organic nitrates (RONO₂) (R14). In addition, the NO_y^{*} species are removed ($L(\text{NO}_y^*)$)
by transport due to the replenishment flow that compensates for chamber leakage and gas sampling of analytical instruments
240 with a first-order rate constant of $k_d \approx 1.6 \times 10^{-5} \text{ s}^{-1}$ (Section 3.1):

$$L(\text{NO}_y^*) = ([\text{NO}] + [\text{NO}_2] + [\text{HONO}]) k_d \quad (5)$$

The dilution rate is monitored by the input flow rate of synthetic air which yield an accuracy of k_d better than 1 %. Hence,
the uncertainty due to the dilution is neglected due to its small contribution. The concentration of NO_y^{*} at a given time is then
determined by the difference between the time-integrated production and loss terms.



$$245 \quad [\text{NO}_y^*] = \int_t (Q(\text{HONO}) - k_{12}[\text{OH}][\text{NO}_2] - k_{14}[\text{RO}_2][\text{NO}] - L(\text{NO}_y^*)) dt \quad (6)$$

It is assumed that the formation of HNO_3 and RONO_2 are effective sinks for NO_x and that reformation of NO_x by their reactions with OH or photolysis does not play a role. This assumption is justified because the low reaction rate constant with OH and small absorption cross sections of HNO_3 and RONO_2 (Burkholder et al., 2020; Browne et al., 2014) lead to their lifetimes in the range of several days, much slower than the time scale of the experiments. The loss of NO_2 due to reaction with
 250 ozone which forms nitrate radicals (NO_3) is neglected in Equation 6, because the NO_3 radical is efficiently converted back to NO_2 by photolysis and reaction with NO in the photo-oxidation experiments in this work. The formation of other oxidized nitrogen species such as peroxy nitric acid (HNO_4) and acetyl peroxy nitrate (PAN) is also assumed to be negligible because their mixing ratios are expected to be only a few tens pptv for conditions of the experiments.

The formation rate of HONO that is within the range of a few hundred pptv/h in the chamber experiments can be determined
 255 from measurements of OH, NO, HONO and $j(\text{HONO})$ by assuming a photo-stationary state for the HONO concentration:

$$\begin{aligned} \frac{d[\text{HONO}]}{dt} &= Q(\text{HONO}) - j_{\text{HONO}}[\text{HONO}] + k_{\text{OH}+\text{NO}}[\text{OH}][\text{NO}] = 0 \\ Q(\text{HONO}) &= j_{\text{HONO}}[\text{HONO}] - k_{13}[\text{OH}][\text{NO}] \end{aligned} \quad (7)$$

For the experimental conditions ($j(\text{HONO}) = 8 \times 10^{-4} \text{ s}^{-1}$) photo-stationary state is reached within approximately 20 minutes. On this time scale, the dilution of HONO by the chamber replenishment flow is only 2 % and is therefore neglected.
 260 The uncertainty of the HONO formation rate (Equation 7 is dominated by the uncertainty in the HONO measurement (22 %, Table 3).

Using measured concentrations of NO, NO_2 and HONO, the concentration of NO that is converted to organic nitrates during the experiment (ΔRONO_2) is determined by the balance of nitrogen oxide concentrations:

$$\begin{aligned} \Delta[\text{RONO}_2] &= \int_t k_{14}[\text{RO}_2][\text{NO}] dt \\ 265 \quad &= \Phi_{\text{RONO}_2} \int_t (k_{11} + k_{14})[\text{RO}_2][\text{NO}] dt \\ &= \int_t (Q(\text{HONO}) - k_{12}[\text{OH}][\text{NO}_2] - L(\text{NO}_y^*)) dt - ([\text{NO}] + [\text{NO}_2] + [\text{HONO}]) \end{aligned} \quad (8)$$

The reaction yield of organic nitrates (Φ_{RONO_2}) can be then derived as slope of a linear fit (Equation 8). The accuracy of the yield is mainly determined by the accuracy the kinetic reaction rate constants and measurements that vary between the specific experiments.



270 4.1 Test experiments with methane and α -pinene

In order to test the method described above, organic nitrate yields from the photo-oxidation of methane and α -pinene were experimentally determined and compared with literature values. NO mixing ratios were between 100 to 300 pptv in both experiments, so that at least 60 % of the RO₂ radicals reacted with NO.

Figure 6 shows time series of NO_y species in α -pinene and methane experiments. In the methane experiment, the total mixing ratio of NO_y species calculated from the HONO chamber source (Equation 7) is around 1.3 ppbv and is mainly explained by measured NO₂ and NO concentrations. In the experiment with α -pinene, the calculated NO_y mixing ratio increased from 1 to 1.3 ppbv within two hours. Approximately 1 ppbv was present as NO₂ and NO and the remaining fraction can be attributed to the formation of organic nitrates from α -pinene. The alkyl nitrate yield is derived from Equation 8 by a linear fit as described above.

280 For the experiment with methane, the alkyl nitrate yield is smaller than error of the calculation (0.00 ± 0.04 , Fig. 6). This small value well agrees with literature values of < 0.003 (Scholtens et al., 1995) and 0.0039 ± 0.0011 (Butkovskaya et al., 2012) at room temperature.

The alkyl nitrate yield determined from the OH oxidation experiment with α -pinene is 0.32 ± 0.06 . The accuracy is mainly due to the uncertainty of RO₂ measurements. Reported literature values are 0.01 (Aschmann et al., 2002), 0.18 ± 0.09 (Nozière et al., 1999) and 0.26 ± 0.07 (Rindelaub et al., 2015). All experiments were conducted at measurement conditions where the reaction of RO₂ with NO was the dominant RO₂ reaction pathway, so that the obtained organic nitrate yields can be attributed to the yield from reaction of α -pinene derived RO₂. The value determined in this work agrees well with values determined from direct measurements using FT-IR spectroscopy in experiments by Nozière et al. (1999) and Rindelaub et al. (2015) within the specified errors. In contrast, Aschmann et al. (2002) approximated the nitrate yield from mass spectrometer measurements with a high uncertainty that likely explains the much lower yield in their experiments compared to the other two studies and results in this work.

290 The good agreement between nitrate yields determined in the experiments with methane and α -pinene with reported values in literature demonstrates that the applied method for the determination of alkyl nitrate formation in the chamber gives reliable results.

295 5 Experimental analysis of the chemical budgets of OH, HO₂, RO₂ and RO_x

The chamber experiments are also used to study the chemical budgets of OH, HO₂, RO₂, and RO_x during the photochemical oxidation of myrcene. Radical production and destruction rates are determined from reaction kinetic data and measured trace gas concentrations for reactions that are known to produce or consume radicals in the experiments. Because the chemical lifetime of radicals is short (seconds to minutes), the radical concentrations are expected to be in steady state. Therefore, the total loss rates for each radical species is balanced by its total production rate. Analysing radical production and destruction rates using experimental data gives an indication if reactions taken into account can describe observations or if there are contributions from further reactions. A similar method has been used before by Tan et al. (2019, 2020) to analyse radical

300



budgets in atmospheric air in China. Here, the method is tested for the photochemical degradation of methane and then applied to the more complex degradation mechanism of myrcene.

305 5.1 Radical production, destruction and regeneration reactions

In the chamber experiments, OH, HO₂ and RO₂ radicals are primarily formed from photolysis of HONO (Reaction R1), O₃ (Reaction R2), formaldehyde (Reaction R3), and from the ozonolysis of myrcene (Reaction R4). Radical termination processes include the formation of nitrates (HNO₃, HONO and RONO₂, Reactions R12, R13 and 14) and peroxides (ROOH and H₂O₂, Reaction R15 and R16). The contribution from RO₂ self-combination reactions is negligible and not considered
310 here. The photolysis of peroxides that can lead to the production of radicals is neglected due to the slow photolysis frequency (typical value for example for CH₃OOH: 2 × 10⁻⁶ s⁻¹ at noontime). The rates for the primary production of RO_x radicals ($P(\text{RO}_x)$) and for the termination loss rate ($L(\text{RO}_x)$) can be calculated as:

$$P(\text{RO}_x) = j_{\text{HONO}}[\text{HONO}] + \Phi_{\text{OH},2}j_{\text{O}(^1\text{D})}[\text{O}_3] + 2j_{\text{HCHO}}[\text{HCHO}] + (\Phi_{\text{OH},4} + \Phi_{\text{RO}_2,4})k_4[\text{VOC}][\text{O}_3] \quad (9)$$

$$L(\text{RO}_x) = (k_{12}[\text{NO}_2] + k_{13}[\text{NO}])[\text{OH}] + (k_{14}[\text{NO}] + 2k_{15}[\text{HO}_2])[\text{RO}_2] + 2k_{16}[\text{HO}_2]^2 \quad (10)$$

315 For the calculation of $P(\text{RO}_x)$ and $L(\text{RO}_x)$, measured radical and trace gas concentrations are used. For the analysis of the myrcene experiments, total RO₂ concentrations include corrections for the reduced detection sensitivity of MyO₂ as explained in Section 3.4. The applied rate constants are listed in Table 3. The lumped rate constant for the reactions between RO₂ and HO₂ (R15) has a rather large uncertainty, because rate constants of different RO₂ species can be different by up to a factor of 4 (Jenkin et al., 2019). The uncertainty of other reaction rate constants is typically around 10%. In addition, the accuracies
320 of measurements (Table 2) add to the total uncertainty in the calculation of loss and production rates. In these experiments, there is a small background OH reactivity (< 1 s⁻¹) which can be a permanent loss of radicals, but could also regenerate HO₂ and/or RO₂. However, this background reactivity is small compared to the OH reactivity from methane and myrcene during the experiments (> 15 s⁻¹), so that it does not affect the analysis.

If production and destruction rates of single RO_x species (OH, HO₂ and RO₂) are calculated, radical conversion reactions
325 need to be additionally taken into account. The total OH loss rate ($L(\text{OH})$) can be quantified by the product of measured OH concentrations and total OH reactivity:

$$L(\text{OH}) = [\text{OH}]k_{\text{OH}} \quad (11)$$

The OH production rate ($P(\text{OH})$) can be calculated from the sum of production from HONO (Reaction R1) and O₃ photolysis followed by water reaction (Reaction R2), ozonolysis of VOCs (Reaction R4) and radical regeneration from HO₂ reacting
330 with NO (Reaction R5) or O₃ (Reaction R6):

$$P(\text{OH}) = j_{\text{HONO}}[\text{HONO}] + \Phi_{\text{OH},2}j_{\text{O}(^1\text{D})}[\text{O}_3] + \Phi_{\text{OH},4}k_4[\text{VOC}][\text{O}_3] + (k_5[\text{NO}] + k_6[\text{O}_3])[\text{HO}_2] \quad (12)$$



The loss of HO₂ ($L(\text{HO}_2)$) and RO₂ ($L(\text{RO}_2)$) radicals are dominated by their reactions with NO for conditions of the experiments here (Reactions R5, R11, R14) in addition to recombination of peroxy radicals including reactions of HO₂ with RO₂ (Reaction R15) and HO₂ self-reactions (Reaction R16):

$$L(\text{HO}_2) = (k_5[\text{NO}] + k_6[\text{O}_3] + k_{15}[\text{RO}_2] + 2k_{16}[\text{HO}_2])[\text{HO}_2] \quad (13)$$

$$L(\text{RO}_2) = ((k_{11} + k_{14})[\text{NO}] + k_{15}[\text{HO}_2])[\text{RO}_2] \quad (14)$$

The HO₂ production ($P(\text{HO}_2)$) rate can be calculated from the photolysis of aldehydes, of which HCHO (Reaction R3) were measured in these experiments, and reactions that convert OH or RO₂ to HO₂. OH to HO₂ conversion occurs in the reaction of OH with HCHO (Reaction R6), CO (Reaction R7) and O₃ (Reaction R8). The reaction of RO₂ with NO (Reaction R9) produces either HO₂ or organic nitrates (Reaction R14). The total HO₂ production rate is then calculated as:

$$P(\text{HO}_2) = 2j_{\text{HCHO}}[\text{HCHO}] + (k_6[\text{HCHO}] + k_7[\text{CO}] + k_8[\text{O}_3])[\text{OH}] + k_{11}[\text{NO}][\text{RO}_2] \quad (15)$$

RO₂ primary production consists of the ozonolysis of VOCs (Reaction R4). In addition RO₂ is produced from radical propagation reactions via the reaction of OH with VOCs. The total production rate ($P(\text{RO}_2)$) can be calculated from the VOC reactivity, k_{OHVOC} , (Equation 1) assuming that each reaction of a VOC with OH produces one RO₂ radical:

$$P(\text{RO}_2) = \Phi_{\text{RO}_2} k_4 [\text{VOC}][\text{O}_3] + k_{\text{OHVOC}}[\text{OH}] \quad (16)$$

5.2 Radical production and destruction in a test experiment with methane

The chemical radical budget analysis in SAPHIR experiments was tested in a photo-oxidation experiment with methane. The chemical oxidation mechanism of methane is much simpler than that of monoterpenes. (1) Ozonolysis reactions do not play a role. (2) Organic peroxy radicals that are formed in the chemical mechanism are methyl peroxy radicals (CH₃O₂), which can be accurately measured by the ROxLIF system. (3) There are recommendations for rate constants for reactions involving methyl peroxy radicals (IUPAC, 2020).

Reaction rates for the methane oxidation experiment (29 May 2020) are shown in Fig. 7. Since no DOAS OH measurements were available for this experiment, OH measurements by the LIF instrument were used (Fig. S4). Total turnover rates are similar for all single radical species (OH, HO₂, RO₂) with values between 8 and 12 ppbv/h and do not vary much over the course of the experiment. The loss of peroxy radicals is dominated by the radical regeneration reaction with NO whereas radical recombination reactions contribute less than 10 % to the entire loss of peroxy radicals. OH is nearly only lost by its reaction with methane and formaldehyde.

The production and destruction rates of total RO_x are significantly smaller than those of single radical species with values that rise from 2 to 4 ppbv/h over the course of the experiment as radical regeneration reactions cancel out. The increase is due



360 to radicals from the photolysis of formaldehyde that is continuously produced from the chamber wall (Section 3.1) and in the reaction of OH + methane. The increase in radical production is balanced by increasing rates of peroxy radical recombination reactions and of the reaction of OH with NO₂. The latter is due to the increase of nitrogen oxide concentrations from the chamber source of nitrous acid.

Radical production and destruction rates of each radical species and of total RO_x are roughly balanced within the uncertainty of the calculation (Fig. 7). Maximum deviations are less than 0.5 ppbv/h for the production and destruction rates of RO_x and HO₂. Differences are much smaller than the accuracy of the calculation (1.5 ppbv/h for RO_x and 2 ppbv/h for HO₂). Higher deviations with values of up to 4 ppbv/h are seen for OH and RO₂ but opposite behaviour. The accuracy of OH and RO₂ radical production and destruction rates are in the range of 2 to 3 ppbv/h, which cannot explain the discrepancies. As the RO_x and HO₂ budget are closed using measured OH concentrations, the imbalance between OH and RO₂ production and destruction indicates an unknown systematic error in the conversion rate from OH to RO₂. One possible explanation for the observed imbalances would be that the calculated reaction rate of OH + CH₄ is too large. This could either be caused by the measured OH or CH₄ concentrations being too high, or by the applied reaction rate constant $k_{\text{OH}+\text{CH}_4}$ being too large.

The methane concentrations in the chamber were above the measurement range specified by the manufacturer (Picarro, 0-20 ppmv). However, an instrumental test of the instrument done prior to the chamber experiment showed that the stated measurement accuracy holds at the concentration (150 ppmv) used in the presented experiment. A small background reactivity of 2 s⁻¹ was found before the injection of methane. The OH reactivity calculated from measured methane concentration and the reaction rate constant is about 26 s⁻¹, which is consistent with the direct OH reactivity measurement considering the contribution of background reactivity (Fig. S4). This gives the confidence on the accuracy of the methane measurement.

Another reason for the imbalances could be LIF calibration errors that are larger than given in Table 2. The specification for OH in Table 2 was generally confirmed in previous intercomparisons with the OH-DOAS instrument. However, in some instances the LIF measurements were found to deviate more than expected. An example are the experiments on 16 Aug and 22 Aug 2012 (Fig. 5 and Fig. S1), where the LIF measurements are higher than the DOAS data by a factor 1.2 probably due to an incorrect calibration.

In the following Section, radical budgets for the chemical degradation of myrcene are investigated. In these cases, also OH-DOAS data were available which have a generally better accuracy than LIF. Therefore, the DOAS data were used for the budget analysis (Fig. 10 - 12). Note that LIF and DOAS showed good agreement in these experiments (Fig. S2 and S3).

The rate constant for OH + CH₄ has an uncertainty factor of $f=1.1$ according to NASA/JPL 2020 and $f=1.15$ according to IUPAC web (4 June 2013). Thus, 10-15% difference in the radical budgets would be explainable by a too large rate constant.

This partly unresolved discrepancy shows the limitation of the analysis of radical production and destruction rates and needs to be further investigated. The analysis of the experiment with the well-defined chemistry of methane indicates that the maximum accuracy of calculated differences between production and destruction rates for single radical species is 20 % in the chamber experiments.



6 Results and discussion of the experiments with myrcene

6.1 Reaction rate constant of the OH reaction with myrcene

395 The rate constant of the reaction of myrcene with OH is determined from the measured temporal decay of myrcene and the measured OH concentration. The time series of measured myrcene concentrations are compared to calculations using a chemical box model that only includes chemical loss reactions with OH and O₃ and dilution. The model is constrained to measured values of temperature, pressure, the dilution rate constant, OH and O₃ concentrations. The initial myrcene concentration is set to the injected amount of myrcene. The reaction rate constant for myrcene with O₃ is taken from the work by ? who measured a value of $2.21 \times 10^{-15} \exp(-(520 \pm 109)K/T) \text{ cm}^3\text{s}^{-1}$ using relative rate technique. Atkinson et al. (1986) suggested a nearly 20 % faster reaction rate constant of $4.7 \times 10^{-16} \text{ cm}^3\text{s}^{-1}$ ($T = 298 \text{ K}$) that is still consistent with the value by ? within the experimental uncertainties. The chemical lifetime of myrcene with respect to the ozonolysis reaction was approximately 44 min in the low NO_x experiments with O₃ concentrations of 40 ppbv but was up to 3 hours in the experiments with medium NO_x and O₃ concentrations of less than 10 ppbv. In comparison, the chemical lifetime of myrcene with respect to the reaction with OH was within the range of 15 min at OH concentrations of approximately $5 \times 10^6 \text{ cm}^{-3}$ observed in all experiments. Hence, the contribution of ozonolysis to the total chemical loss of myrcene in the experiments was 10 % to 26 %.

The rate constant of the reaction of myrcene with OH is optimized, such that the difference between measured and modelled myrcene concentration time series is minimized. For this optimization the OH concentration observed from DOAS is used. This procedure is applied to all three experiments resulting in a rate constant of $(2.3 \pm 0.3) \times 10^{-10} \text{ cm}^3\text{s}^{-1}$ ($T = 298 \text{ K}$, Table 4, Fig. S6 Supplement). The error results mainly from the variability of values determined in the different experiments.

The rate constant determined in this study is in good agreement with those reported in the literature (Table 4). Relative rate technique was used in experiments by Atkinson et al. (1986) and Grimsrud et al. (1975). The value reported by Hites and Turner (2009) ($3.4 \times 10^{-10} \text{ cm}^3\text{s}^{-1}$) is higher than those of this work and reported by Atkinson et al. (1986), but has a high uncertainty of $\pm 35 \%$ that is explained by experimental difficulties in the handling of myrcene. The rate constant calculated from structure-activity relationship (SAR) by Peeters et al. (2007) is nearly 20 % lower than the experimentally derived reaction rate constants, but this difference is within the accuracy of SAR predictions.

6.2 Product yields of the reactions of myrcene hydroxy peroxy radicals with NO

When OH reacts with myrcene, about half (48 %) of the OH adds to the $-\text{CH}=\text{C}(\text{CH}_3)_2$ moiety and the other half (48 %) to the isoprenyl part (Peeters et al., 2007). In the first case, 4-vinyl-4-pentenal and acetone are formed from the reaction of MyO₂ with NO (Section 2). In the second case, 2-methylidene-6-methyl-5-heptenal or 1-vinyl-5-methyl-4-hexenone are produced together with formaldehyde. Thus, the yields of acetone and formaldehyde are indicators for the yields of the OH addition to myrcene.

Acetone and formaldehyde yields are calculated from measured time series of product species and myrcene in the experiments on 16 and 22 August 2012, when NO mixing ratios were 200 pptv, so that $>90 \%$ of RO₂ reacted with NO (Fig. 5). Only measurements after the first myrcene injection are used here to avoid that secondary chemistry impacts the yield deter-



mination. The two other experiments are not considered for three reasons. The large amount of ozone (up to 50 ppbv) would require considerable correction for myrcene ozonolysis. Second, due to the lower NO concentration, reactions of RO₂ with HO₂ would be competitive with the reaction of RO₂ with NO. Third, the smaller amount of injected myrcene produced less oxidation products.

430 Corrections are applied to measured myrcene, formaldehyde and acetone time series, in order to relate myrcene that reacted with OH and product species that are chemically formed from this reaction following the procedure described in previous work (Galloway et al., 2011; Kaminski et al., 2017). Product concentrations are corrected for losses due to dilution, photolysis and reaction with OH. In addition, formation of formaldehyde and acetone from chamber sources need to be subtracted from the measured concentrations. Myrcene concentrations are also corrected for dilution and the fraction of myrcene that reacted
435 with ozone. After corrections have been applied, the relationships between consumed myrcene and product concentrations are linear, if species are mainly formed as first-generation products in the reaction of myrcene and OH. The slopes give acetone and formaldehyde yields of 0.45 ± 0.08 and 0.35 ± 0.08 , respectively (Fig. 8). The error is caused by the uncertainty in the source strengths of chamber sources for acetone and formaldehyde and the accuracy of measurements.

The acetone yield is in excellent agreement with SAR predictions that 48 % of the OH adds to the double bond of the
440 $-\text{CH}=\text{C}(\text{CH}_3)_2$ moiety (Peeters et al., 2007). The formaldehyde yield of 0.35 ± 0.08 is lower than the SAR prediction for the OH attack to the isoprenyl part (0.48). One possible reason for the smaller than expected formaldehyde yield could be isomerization reactions of the myrcene peroxy radicals shown (Fig. 3), which may not lead to the production of HCHO. The estimated bulk isomerization rates of 0.097 s^{-1} and 0.21 s^{-1} (Section 2) are sufficiently fast to compete with the loss rate of 0.04 s^{-1} for MyO₂ with 200 pptv of NO, so that the reduced formaldehyde yield could be fully explained by isomerization
445 reaction pathways. A different explanation for the lower formaldehyde yield compared to acetone is that the nitrate yield is larger for the MyO₂ that would otherwise end up as formaldehyde. In fact, the total product yields is closed to unity (acetone (0.45 ± 0.08) + formaldehyde (0.35 ± 0.08) + nitrate (0.13 ± 0.03) = 0.93 ± 0.12).

Acetone and formaldehyde yields agree with published results by Reissell et al. (2002) and Orlando et al. (2000) (Table 5). Lee et al. (2006) reported a smaller acetone yield of 0.22 ± 0.02 and a significantly higher HCHO yield of 0.74 ± 0.08 .
450 However, their HCHO yield carries a large uncertainty because concentrations were outside the range for which the instrument was calibrated.

The oxygenated organic compound 4-vinyl-4-pentenal has been detected in previous studies from the reaction of myrcene with OH (Reissell et al., 2002; Lee et al., 2006). Fragmentation in the PTR-MS and further oxidation of 4-vinyl-4-pentenal complicated the unambiguous yield determination. Therefore, Lee et al. (2006) reported a high yield of 0.4, but also state a
455 lower limit of 0.09. Acetone is the co-product of 4-vinyl-4-pentenal (Fig. 3). The high limit yield of 0.4 for 4-vinyl-4-pentenal by Lee et al. (2006) is therefore consistent with the acetone yield determined in this work. The yield of 4-vinyl-4-pentenal was also measured by Reissell et al. (2002), but a lower yield of 0.19 ± 0.04 was found. This low yield is apparently inconsistent with the yield of 0.45 ± 0.06 for the co-product acetone determined in the same experiments (Table 5). The authors suggest that there could be re-arrangement of the hydroxyalkoxy radical that compete with the decomposition to 4-vinyl-4-pentenal,
460 but may still lead to acetone production, so that the yield for acetone could become higher than that of 4-vinyl-4-pentenal.



The yield of organic nitrate from reactions of MyO₂ with NO is determined from the analysis of reactive nitrogen oxides in the chamber as described in Section 4. This results in an organic nitrate yield of 0.13 ± 0.03 (Fig. 9). This value is consistent with the yield of 0.10 ± 0.03 reported by Lee et al. (2006), who directly measured organic nitrates by mass spectrometry. Values are lower than the yield expected from SAR described in Jenkin et al. (2019), which predicts a yield of 0.19 that would also
465 apply for other monoterpenes, but may still be within the uncertainty of the SAR predictions. Like for the formaldehyde yield, the smaller than predicted organic nitrate yield may also be partly due to competing MyO₂ isomerization reactions.

6.3 Primary radical production and termination

The primary radical production is due to photolysis (Reactions R1 to R3) and ozonolysis reactions (Reaction R4). Production rates by photolysis of O₃, HONO and HCHO are calculated using measured trace gas concentrations and photolysis frequen-
470 cies. The calculation of the production rate from myrcene ozonolysis requires the knowledge of both the reaction rate constant (Section 6.1) and the yield of OH and RO₂ radicals. The uncertainty of yields are high. Values range from 0.71 determined from high-level quantum chemistry calculations and kinetic calculations (Deng et al., 2018) to experimental values of 1.15 with an uncertainty of a factor of 1.5 (Atkinson et al., 1992). It is worth noting that additional radicals could be produced from ozonolysis reactions of oxidation products at later times of the experiment because first-generation organic products from the
475 reaction of myrcene with OH still contain C–C double bonds that can react with ozone.

The radical termination reactions include reactions with nitrogen oxides (Reactions R12–R14) and radical self-reactions (Reactions R15, R16). Loss of radicals in the reaction with NO_x and the HO₂ self-reaction contribute less than 1 ppbv/h to the total loss rate each. Contributions from RO₂ self-reactions are expected to be negligible (< 2 %) because their reaction rate constants are typically much smaller than those of RO₂+HO₂ reactions (Table 3). The reaction rate constant of the MyO₂+HO₂
480 reaction is estimated from SAR by Jenkin et al. (2019) and therefore, the value could also have a high uncertainty.

In the experiment with medium NO concentrations (22 August 2012), the total radical production is low with values of less than 1.5 ppbv/h and radical production and destruction are balanced (Fig. 10). No ozone is added, so that radical production from ozonolysis plays a minor role, and radical loss by radical recombination is suppressed due to the competition of peroxy radical reactions with NO. Therefore, the uncertainties in the radical yield of myrcene ozonolysis and the reaction rate constant
485 of the MyO₂+HO₂ reaction do not impact the results.

In contrast, radical production and destruction rates in experiments performed at low NO concentrations (17 July 2013, 18 July 2013) that is achieved by suppressing NO in the reaction with O₃ are dominated by radical production from myrcene ozonolysis and radical destruction by MyO₂+HO₂ reactions right after the injection of myrcene. Taking the upper limit of the radical yield from the ozonolysis reaction of 1.42 (0.71 for OH and 0.71 for MyO₂), the production is 4 ppbv/h. The radical
490 primary production could be smaller, if the unspecified radical yield of 1.15 suggested by Atkinson et al. (1992) is applied. Applying the reaction rate constant of the MyO₂+HO₂ derived by SAR (Section 2) results in a maximum radical loss rate of 13.5 ppbv/h also right after the injection of myrcene when MyO₂ concentrations are highest. Because this high destruction rate cannot be balanced by radical production, either the reaction rate constant must be lower than SAR predictions or there are reaction pathways that do not lead to radical termination products (organic peroxide), but regenerate radicals. Due to the



495 high uncertainty of radical production from ozonolysis in these experiments, no firm conclusion can be drawn. The reaction rate constant would need to be reduced between a factor of 0.4 and 0.7 (0.9 to $1.6 \times 10^{-11} \text{ cm}^3\text{s}^{-1}$ ($T = 298 \text{ K}$, Table 3) or the yield of of radical products would need to be in the range of 0.3 and 0.6 to match the range of radical production.

A reduced $\text{MyO}_2 + \text{HO}_2$ reaction rate constant is significantly different from the SAR predictions by Jenkin et al. (2019). The SAR value is similar for all monoterpenes and well agrees with direct measurements for the reaction rate constant of HO_2 with
500 RO_2 derived from α -pinene ($2.1 \times 10^{-11} \text{ cm}^3\text{s}^{-1}$), γ -terpinene ($2.0 \times 10^{-11} \text{ cm}^3\text{s}^{-1}$) and limonene ($2.1 \times 10^{-11} \text{ cm}^3\text{s}^{-1}$) reported by Boyd et al. (2003). A high yield of radical products from the $\text{MyO}_2 + \text{HO}_2$ reaction may not be expected from analogies of isoprene and 2-methyl-2-butene (Liu et al., 2012; Paulot et al., 2009). For these reasons there is no clear conclusion, how exactly rate constants and yields need to be adjusted to balance the RO_x production and destruction rates.

6.4 RO_x radical chain propagation reactions

505 The total production and loss rates of OH, HO_2 and RO_2 radicals do not exhibit much variability over the course of the experiments with medium NO concentrations (Fig. 11, Fig. S7 Supplement). In the experiments with low NO, the turnover rates of radical production and destruction reactions for OH, HO_2 , and RO_2 exhibit peak values when myrcene is injected (Fig. 12 and Fig. S8 Supplement). This distinct feature is related to the elevated radical production of up to 2 ppbv/h from the ozonolysis of myrcene and peak concentrations of peroxy radical concentrations.

510 Production and destruction rates of OH radicals are balanced within the accuracies of the calculation in all experiments. Rates are similar in all experiments with values around 4 ppbv/h. Maximum RO_2 turnover rates are lower (2 to 3 ppbv/h) compared to those of OH and HO_2 , because up to 25 % of the OH radicals are directly converted to HO_2 , for example in the reaction of OH with CO or HCHO. Because myrcene is consumed within a few hours, the turnover rate of RO_2 is decreasing over time after each myrcene injection. In experiments with medium NO concentrations, HO_2 radical production and destruction rates
515 are not balanced by 0.5 of 1.5 ppbv/h in contrast to those of OH. Discrepancies, however, are only slightly larger than the $1-\sigma$ uncertainty of the calculated values. The total uncertainty is dominated by the accuracy of RO_2 measurements because of the low detection sensitivity of the instrument for MyO_2 and the uncertainty in the sensitivity of other RO_2 formed from oxidation products that are dominant, once myrcene reacted away.

Even higher discrepancies of up to 2 ppbv/h between HO_2 and RO_2 production and destruction rates are observed in the
520 experiments with low NO mixing ratios. Values are highest right after each myrcene injection with the HO_2 production rate being lower than the destruction rate and the RO_2 destruction rate being higher than the production rate. Because differences are highest when the chemical system is dominated by the oxidation of myrcene and therefore the presence of MyO_2 , reaction pathways of MyO_2 other than reactions with NO and HO_2 could be responsible for the imbalances.

As described in Section 2, MyO_2 isomerization reactions can become competitive at low NO mixing ratios like in these
525 experiments (< 0.1 ppbv). Isomerization rates derived from SAR are applied (Section 2). Products of the 1,6 H-shift reactions in myrcene that correspond to similar reactions in isoprene could produce radicals like in the case of isoprene (Fig. 3). Products from the two additional isomerization reactions likely undergo fast ring-closure reactions on the di-methyl double bond and it is uncertain, if OH or HO_2 are directly produced. As an estimate for the potential impact of MyO_2 isomerization reactions on



radical regeneration, production of one HO_x radical from each isomerization reaction is assumed in the following. The results
530 of this sensitivity analysis are shown in Fig. 11 and 12.

The bulk MyO₂ loss due to isomerization for the isomers that can rapidly interconvert (Fig. 2) are competitive with the
reaction of MyO₂ with NO at mixing ratios of 0.5 to 1 ppbv making these reaction very competitive for conditions of the
experiments. As shown in Fig. 12, the discrepancy between RO₂ radical production and destruction rates would be even over-
compensated, if isomerization rates determined by SAR are applied. Sensitivity model runs suggest that a bulk isomerization
535 reaction rate of 0.05 s⁻¹ would be sufficient to balance the RO₂ production rate. This value is a factor 2 to 4 lower than bulk
reaction rates calculated by SAR, but are still within the uncertainty of calculations.

Although this analysis shows the potential for high contributions of MyO₂ isomerization to the total loss rate of RO₂ for
conditions that are typical for forested areas, the uncertainties are high concerning the isomerization rate constants and the
yield of HO_x radicals. In addition, discrepancies of radical production and destruction rates within the range of 20 % of the
540 total rate for single radical species can be observed even for well-known chemical systems as methane as shown in Section 5.2.
This is likely due to uncertainties in the radical concentration measurements. Unaccounted systematic errors in the analysis
may therefore also explain a significant fraction of the differences in radical production and destruction rates in the experiments
with myrcene.

7 Summary and conclusions

545 The photooxidation of the monoterpene myrcene was investigated at atmospheric conditions in the simulation chamber SAPHIR
at two different levels of NO. The chemical mechanism of the oxidation of myrcene by OH has not been investigated in detail
so far. Based on the structural similarity with isoprene (CH₂=CH-C(=CH₂)CH₂- moiety) and with 2-methyl-2-butene
(-CH₂-CH=C(CH₃)₂ moiety) and structure-activity relationship (Peeters et al., 2007; Jenkin et al., 2019; Vereecken and
Nozière, 2020) a chemical mechanism for the first oxidation step is proposed that includes rapid RO₂ interconversion by re-
versible oxygen addition and H-shift reactions of RO₂ like in isoprene. In addition to RO₂ isomerization reactions similar to
550 that of isoprene derived RO₂, additional H-shift reactions are suggested by SAR, so that the overall impact of isomerization
reactions in myrcene can be high, although only approximately half of the attack of OH to myrcene is on the isoprenyl part.
Assuming that the interconversion of RO₂ is similar as in isoprene, bulk isomerization rate constants of 0.21 s⁻¹ and 0.097 s⁻¹
(*T* = 298 K) for the 3'-OH and 1-OH addition, respectively, are calculated using rate constants
555 derived by SAR.

Experiments in the chamber allowed to determine the reaction rate constant of the reaction of myrcene with OH resulting in
a value of $(2.3 \pm 0.3) \times 10^{-10}$ cm³s⁻¹, which is in good agreement with values reported in the literature within the uncertainties
(Table 4). Product yields of acetone and formaldehyde calculated from measured time series of these species were found to
be 0.45 ± 0.08 and 0.35 ± 0.08, respectively. Although these values well agree with studies by Reissell et al. (2002); Orlando
560 et al. (2000) within the uncertainties, the formaldehyde yield is lower than expected for conditions of the experiment with NO
mixing ratios of 200 pptv, if there are no RO₂ isomerization reactions. The lower yield would be consistent with competing



RO₂ isomerization reactions that may not produce formaldehyde. From the analysis of nitrogen oxides in the chamber, the yield of organic nitrates in the myrcene oxidation could be determined to be 0.13 ± 0.03 in agreement with measurements by Lee et al. (2006) (0.10 ± 0.03). Both values are lower than typical values predicted by SAR (0.19, Jenkin et al. (2019) and
565 found for monoterpene species which often range between 0.15 and 0.25. This may impact the formation of secondary organic aerosol from the OH oxidation of myrcene compared to other monoterpene species. The applicability of the procedure for the determination of the organic nitrate yield was tested in photo-oxidation experiments with methane and α -pinene, both of which resulted in organic nitrate yields that well agree with expected values. This demonstrates that the method gives reliable results and can be applied in chamber experiments.

570 Radical production and destruction rates can be calculated using measured radical and trace gas concentrations. Total radical (RO_x) production and destruction rates are balanced in the experiments in this work, if the rate of the radical loss terminating the radical reaction chain due to recombination reactions of HO₂ and RO₂ was only 40 % of the loss expected from the reaction rate constant predicted by SAR or if this reaction would regenerate radicals instead of producing radical termination products (hydroperoxides). However, this conclusion has a high uncertainty of approximately a factor of 2 due to the uncertainty of
575 radical production from myrcene ozonolysis.

Imbalances between radical production and destruction rates are observed for RO₂ radicals in the experiments with low NO concentrations, when MyO₂ isomerization reaction can become competitive with bi-molecular reactions. Bulk reaction rates constants around 0.05 s^{-1} for MyO₂ isomers which can quickly interconvert and which partly can isomerize, are a factor of 2 to 4 lower than reaction rates constants calculated by SAR, but values are consistent regarding the high uncertainty of the
580 determination from experiments and the uncertainty of SAR calculations.

The analysis of radical concentrations measurements in field campaigns, where monoterpene emissions dominated the mix of organic compounds such as in Finland (Hens et al., 2014) and in the Rocky Mountains (Kim et al., 2013) showed that current chemical models cannot explain measured OH and HO₂ concentrations. The analysis of chamber experiments revealed that the OH oxidation of the most abundant monoterpene species α -pinene and β -pinene are not well understood (Kaminski
585 et al., 2017; Rolletter et al., 2019). Experiments here show that also the reaction of OH with myrcene is complex due to the rapid RO₂ inter-conversion and H-shift RO₂ reactions. Therefore, short-comings in the description of the photo-oxidation of myrcene could contribute to explaining model-measurement discrepancies found in field campaigns.

Data availability. Data are available from the Eurochamp database (16 August 2012: <https://doi.org/10.25326/5XRG-Y765>, Fuchs et al. (2021a), 22 August 2012: <https://doi.org/10.25326/9RAV-5450>, Fuchs et al. (2021b), 17 July 2013: <https://doi.org/10.25326/GP2K-R926>,
590 Fuchs et al. (2021c), 18 July 2013: <https://doi.org/10.25326/JJ16-1S54>, Fuchs et al. (2021d), 03 September 2019: <https://doi.org/10.25326/PBBV-WF18>, Fuchs et al. (2021e), 29 May 2020: <https://doi.org/10.25326/MG6T-TW58>, Fuchs et al. (2021f))



Author contributions. Z.T., L.H. and H.F. wrote the manuscript. M.K., R.W. and H.F. designed and led the experiments in the chamber. I.A. (organic compounds), B.B. (radiation), H.-P. D. (radicals), X.L. (HONO), S.N. (OH reactivity), F.R. (nitrogen oxides, ozone), R.T. (organic compounds), A.N. (radicals), C.C. (radicals) were responsible for measurements used in this work. All co-authors commented and discussed the manuscript and contributed thereby to the writing of the manuscript.

595

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This work was supported by the EU FP-7 program EUROCHAMP-2 (grant agreement no. 228335). This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement no. 681529). S. Nehr and B. Bohn thank the Deutsche Forschungsgemeinschaft for funding (grant no. 1580/3-1). The authors thank Luc Vereecken for the discussion of the chemical mechanism.

600



References

- Allen, H. M., Crounse, J. D., Bates, K. H., Teng, A. P., Krawiec-Thayer, M. P., Rivera-Rios, J. C., Keutsch, F. N., St. Clair, J. M., Hanisco, T. F., Møller, K. H., Kjaergaard, H. G., and Wennberg, P. O.: Kinetics and product yields of the OH initiated oxidation of hydroxymethyl hydroperoxide, *J. Phys. Chem. A*, 122, 6292–6302, <https://doi.org/10.1021/acs.jpca.8b04577>, 2018.
- 605 Aschmann, S. M., Atkinson, R., and Arey, J.: Products of reaction of OH radicals with α -pinene, *J. Geophys. Res.*, 107, ACH 6–1–ACH 6–7, <https://doi.org/10.1029/2001JD001098>, 2002.
- Atkinson, R.: Gase-phase tropospheric chemistry of volatile organic compounds: 1. alkanes and alkenes, *J. Phys. Chem. Ref. Data*, 26, 217–290, 1997.
- Atkinson, R. and Arey, J.: Atmospheric degradation of volatile organic compounds, *Chem. Rev.*, 103, 4605–4638, <https://doi.org/10.1021/cr0206420>, 2003.
- 610 Atkinson, R., Aschmann, S. M., and Pitts, J. N.: Rate constants for the gas-phase reactions of the OH radical with a series of monoterpenes at 294 ± 1 K, *Int. J. Chem. Kin.*, 18, 287–299, <https://doi.org/10.1002/kin.550180303>, 1986.
- Atkinson, R., Aschmann, S. M., Arey, J., and Shorees, B.: Formation of OH radicals in the gas phase reactions of O₃ with a series of terpenes, *J. Geophys. Res.*, 97, 6065–6073, <https://doi.org/10.1029/92jd00062>, 1992.
- 615 Böge, O., Mutzel, A., Iinuma, Y., Yli-Pirilä, P., Kahnt, A., Joutsensaari, J., and Herrmann, H.: Gas-phase products and secondary organic aerosol formation from the ozonolysis and photooxidation of myrcene, *Atmos. Environ.*, 79, 553–560, <https://doi.org/10.1016/j.atmosenv.2013.07.034>, 2013.
- Bohn, B. and Zilken, H.: Model-aided radiometric determination of photolysis frequencies in a sunlit atmosphere simulation chamber, *Atmos. Chem. Phys.*, 5, 191–206, <https://doi.org/10.5194/acp-5-191-2005>, 2005.
- 620 Bohn, B., Rohrer, F., Brauers, T., and Wahner, A.: Actinometric measurements of NO₂ photolysis frequencies in the atmosphere simulation chamber SAPHIR, *Atmos. Chem. Phys.*, 5, 493–503, <https://doi.org/10.5194/acp-5-493-2005>, 2005.
- Boyd, A. A., Flaud, P.-M., Daugey, N., and Lesclaux, R.: Rate constants for RO₂ + HO₂ reactions measured under a large excess of HO₂, *J. Phys. Chem. A*, 107, 818–821, <https://doi.org/10.1021/jp026581r>, 2003.
- Browne, E. C., Wooldridge, P. J., Min, K. E., and Cohen, R. C.: On the role of monoterpene chemistry in the remote continental boundary layer, *Atmos. Chem. Phys.*, 14, 1225–1238, <https://doi.org/10.5194/acp-14-1225-2014>, 2014.
- 625 Burkholder, J. B., Sander, S. P., Abbatt, J. P. D., Barker, J. R., Huie, R. E., Kolb, C. E., Kurylo, M. J., Orkin, V. L., Wilmouth, D. M., and Wine, P. H.: Chemical kinetics and photochemical data for use in atmospheric studies—evaluation number 19 ([https://jpldataeval.jpl.nasa.gov/pdf/NASA-JPLpanel for data evaluation technical report, 19-5, 2020](https://jpldataeval.jpl.nasa.gov/pdf/NASA-JPLpanel%20for%20data%20evaluation%20technical%20report,%2019-5,%202020)).
- Butkovskaya, N., Kukui, A., and Le Bras, G.: Pressure and temperature dependence of methyl nitrate formation in the CH₃O₂ + NO reaction, *J. Phys. Chem. A*, <https://doi.org/10.1021/jp210710d>, 2012.
- 630 Crounse, J. D., Knap, H. C., Oronso, K. B., Jorgensen, S., Paulot, F., Kjaergaard, H. G., and Wennberg, P. O.: On the atmospheric fate of methacrolein: 1. Peroxy radical isomerization following addition of OH and O₂, *J. Phys. Chem. A*, 116, 5756–5762, <https://doi.org/10.1021/jp211560u>, 2012.
- Crounse, J. D., Nielsen, L. B., Jørgensen, S., Kjaergaard, H. G., and Wennberg, P. O.: Autoxidation of organic compounds in the atmosphere, *J. Phys. Chem. Lett.*, 4, 3513–3520, <https://doi.org/10.1021/jz4019207>, 2013.
- 635 Deng, P., Wang, L., and Wang, L.: Mechanism of gas-phase ozonolysis of β -myrcene in the atmosphere, *J. Phys. Chem. A*, 122, 3013–3020, <https://doi.org/10.1021/acs.jpca.8b00983>, 2018.



- Dorn, H.-P., Brandenburger, U., Brauers, T., and Hausmann, M.: A new in-situ laser long-path absorption instrument for the measurement of tropospheric OH radicals, *J. Atmos. Sci.*, 52, 3373–3380, 1995.
- 640 Fuchs, H., Hofzumahaus, A., and Holland, F.: Measurement of tropospheric RO₂ and HO₂ radicals by a laser-induced fluorescence instrument, *Rev. Sci. Instrum.*, 79, 084 104, <https://doi.org/10.1063/1.2968712>, 2008.
- Fuchs, H., Bohn, B., Hofzumahaus, A., Holland, F., Lu, K. D., Nehr, S., Rohrer, F., and Wahner, A.: Detection of HO₂ by laser-induced fluorescence: calibration and interferences from RO₂ radicals, *Atmos. Meas. Tech.*, 4, 1209–1255, <https://doi.org/10.5194/amt-4-1209-2011>, 2011.
- 645 Fuchs, H., Hofzumahaus, A., Rohrer, F., Bohn, B., Brauers, T., Dorn, H.-P., Häsel, R., Holland, F., Kaminski, M., Li, X., Lu, K., Nehr, S., Tillmann, R., Wegener, R., and Wahner, A.: Experimental evidence for efficient hydroxyl radical regeneration in isoprene oxidation, *Nature Geosci.*, 6, 1023–1026, <https://doi.org/10.1038/NGEO1964>, 2013.
- Fuchs, H., Acir, I. H., Bohn, B., Brauers, T., Dorn, H. P., Häsel, R., Hofzumahaus, A., Holland, F., Kaminski, M., Li, X., Lu, K., Lutz, A., Nehr, S., Rohrer, F., Tillmann, R., Wegener, R., and Wahner, A.: OH regeneration from methacrolein oxidation investigated in the atmosphere simulation chamber SAPHIR, *Atmos. Chem. Phys.*, 14, 7895–7908, <https://doi.org/10.5194/acp-14-7895-2014>, 2014.
- 650 Fuchs, H., Tan, Z., Lu, K., Bohn, B., Broch, S., Brown, S. S., Dong, H., Gomm, S., Häsel, R., He, L., Hofzumahaus, A., Holland, F., Li, X., Liu, Y., Lu, S., Min, K. E., Rohrer, F., Shao, M., Wang, B., Wang, M., Wu, Y., Zeng, L., Zhang, Y., Wahner, A., and Zhang, Y.: OH reactivity at a rural site (Wangdu) in the North China Plain: contributions from OH reactants and experimental OH budget, *Atmos. Chem. Phys.*, 17, 645–661, <https://doi.org/10.5194/acp-17-645-2017>, 2017.
- 655 Fuchs, H., Albrecht, S., Acir, I. H., Bohn, B., Breitenlechner, M., Dorn, H. P., Gkatzelis, G. I., Hofzumahaus, A., Holland, F., Kaminski, M., Keutsch, F. N., Novelli, A., Reimer, D., Rohrer, F., Tillmann, R., Vereecken, L., Wegener, R., Zaytsev, A., Kiendler-Scharr, A., and Wahner, A.: Investigation of the oxidation of methyl vinyl ketone (MVK) by OH radicals in the atmospheric simulation chamber SAPHIR, *Atmos. Chem. Phys.*, 2018, 8001–8016, <https://doi.org/10.5194/acp-18-8001-2018>, 2018.
- Fuchs, H., Kaminski, M., Acir, I.-H., Bohn, B., Dorn, H.-P., Nehr, S., Rohrer, F., Tillmann, R., Wegener, R., and Li, X.: Atmospheric simulation chamber study: myrcene + OH - Gas-phase oxidation - product study [Data set]. AERIS. <https://doi.org/10.25326/5XRG-Y765>, 2021a.
- 660 Fuchs, H., Kaminski, M., Acir, I.-H., Bohn, B., Dorn, H.-P., Nehr, S., Rohrer, F., Tillmann, R., Wegener, R., and Li, X.: Atmospheric simulation chamber study: myrcene + OH - Gas-phase oxidation - product study [Data set]. AERIS. <https://doi.org/10.25326/9RAV-54505>, 2021b.
- 665 Fuchs, H., Kaminski, M., Acir, I.-H., Bohn, B., Dorn, H.-P., Nehr, S., Rohrer, F., Tillmann, R., Wegener, R., and Li, X.: Atmospheric simulation chamber study: myrcene + OH - Gas-phase oxidation - product study [Data set]. AERIS. <https://doi.org/10.25326/GP2K-R926>, 2021c.
- Fuchs, H., Kaminski, M., Acir, I.-H., Bohn, B., Dorn, H.-P., Nehr, S., Rohrer, F., Tillmann, R., Wegener, R., and Li, X.: Atmospheric simulation chamber study: myrcene + OH - Gas-phase oxidation - product study [Data set]. AERIS. <https://doi.org/10.25326/JJ16-1S54>, 2021d.
- 670 Fuchs, H., Bohn, B., Rohrer, F., Tillmann, R., Cho, C., and Novelli, A.: Atmospheric simulation chamber study: alpha-pinene + OH - Gas-phase oxidation - product study [Data set]. AERIS. <https://doi.org/10.25326/JJ16-1S54>, 2021e.
- Fuchs, H., Rohrer, F., Bohn, B., Cho, C., and Novelli, A.: Atmospheric simulation chamber study: myrcene + OH - Reference experiment [Data set]. AERIS. <https://doi.org/10.25326/MG6T-TW58>, 2021f.



- 675 Galloway, M. M., Huisman, A. J., Yee, L. D., Chan, A. W. H., Loza, C. L., Seinfeld, J. H., and Keutsch, F. N.: Yields of oxidized volatile organic compounds during the OH radical initiated oxidation of isoprene, methyl vinyl ketone, and methacrolein under high-NO_x conditions, *Atmos. Chem. Phys.*, 11, 10 779–10 790, <https://doi.org/10.5194/acp-11-10779-2011>, 2011.
- Grimsrud, E. P., Westberg, H. H., and Rasmussen, R. A.: Atmospheric reactivity of monoterpene hydrocarbons, NO, photooxidation and ozonolysis, *Int. J. Chem. Kin.*, 7:Suppl. 1, 183–195, 1975.
- 680 Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The model of emissions of gases and aerosols from nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geosci. Model Dev.*, 5, 1471–1492, <https://doi.org/10.5194/gmd-5-1471-2012>, 2012.
- Helmig, D., Daly, R. W., Milford, J., and Guenther, A.: Seasonal trends of biogenic terpene emissions, *Chemosphere*, 93, 35–46, <https://doi.org/10.1016/j.chemosphere.2013.04.058>, 2013.
- 685 Hens, K., Novelli, A., Martinez, M., Auld, J., Axinte, R., Bohn, B., Fischer, H., Keronen, P., Kubistin, D., Nölscher, A. C., Oswald, R., Paasonen, P., Petäjä, T., Regelin, E., Sander, R., Sinha, V., Sipilä, M., Taraborrelli, D., Tatum Ernest, C., Williams, J., Lelieveld, J., and Harder, H.: Observation and modelling of HO_x radicals in a boreal forest, *Atmos. Chem. Phys.*, 14, 8723–8747, <https://doi.org/10.5194/acp-14-8723-2014>, 2014.
- Hites, R. A. and Turner, A. M.: Rate constants for the gas-phase β -myrcene + OH and isoprene + OH reactions as a function of temperature, *Int. J. Chem. Kin.*, 41, 407–413, <https://doi.org/10.1002/kin.20413>, 2009.
- Hofzumahaus, A., Rohrer, F., Lu, K., Bohn, B., Brauers, T., Chang, C.-C., Fuchs, H., Holland, F., Kita, K., Kondo, Y., Li, X., Lou, S., Shao, M., Zeng, L., Wahner, A., and Zhang, Y.: Amplified trace gas removal in the troposphere, *Science*, 324, 1702–1704, <https://doi.org/10.1126/science.1164566>, 2009.
- IUPAC: IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation, s, p. last access 23 November 2020, 2020.
- 695 Jenkin, M. E., Valorso, R., Aumont, B., and Rickard, A. R.: Estimation of rate coefficients and branching ratios for reactions of organic peroxy radicals for use in automated mechanism construction, *Atmos. Chem. Phys.*, 19, 7691–7717, <https://doi.org/10.5194/acp-19-7691-2019>, 2019.
- Kaminski, M., Fuchs, H., Acir, I. H., Bohn, B., Brauers, T., Dorn, H. P., Häsel, R., Hofzumahaus, A., Li, X., Lutz, A., Nehr, S., Rohrer, F., Tillmann, R., Vereecken, L., Wegener, R., and Wahner, A.: Investigation of the β -pinene photooxidation by OH in the atmosphere simulation chamber SAPHIR, *Atmos. Chem. Phys.*, 17, 6631–6650, <https://doi.org/10.5194/acp-17-6631-2017>, 2017.
- 700 Kim, D., Stevens, P. S., and Hites, R. A.: Rate constants for the gas-phase reactions of OH and O₃ with β -ocimene, β -myrcene, and α - and β -farnesene as a function of temperature, *J. Chem. Phys. A*, 115, 500–506, <https://doi.org/10.1021/jp111173s>, 2011.
- Kim, S., Wolfe, G. M., Mauldin, L., Cantrell, C., Guenther, A., Karl, T., Turnipseed, A., Greenberg, J., Hall, S. R., Ullmann, K., Apel, E., Hornbrook, R., Kajii, Y., Nakashima, Y., Keutsch, F. N., DiGangi, J. P., Henry, S. B., Kaser, L., Schnitzhofer, R., Graus, M., Hansel, A., Zheng, W., and Flocke, F. F.: Evaluation of HO_x sources and cycling using measurement-constrained model calculations in a 2-methyl-3-butene-2-ol (MBO) and monoterpene (MT) dominated ecosystem, *Atmos. Chem. Phys.*, 13, 2031–2044, <https://doi.org/10.5194/acp-13-2031-2013>, 2013.
- 705 Kostianinen, R.: Volatile organic compounds in the indoor air of normal and sick houses, *Atmos. Environ.*, 29, 693–702, [https://doi.org/10.1016/1352-2310\(94\)00309-9](https://doi.org/10.1016/1352-2310(94)00309-9), 1995.
- Lee, A., Goldstein, A. H., Kroll, J. H., Ng, N. L., Varutbangkul, V., Flagan, R. C., and Seinfeld, J. H.: Gas-phase products and secondary aerosol yields from the photooxidation of 16 different terpenes, *J. Geophys. Res.*, 111, n/a–n/a, <https://doi.org/10.1029/2006JD007050>, 2006.



- Lelieveld, J., Butler, T. M., Crowley, J. N., Dillon, T. J., Fischer, H., Ganzeveld, L., Harder, H., Lawrence, M. G., Martinez, M., Taraborrelli, D., and Williams, J.: Atmospheric oxidation capacity sustained by a tropical forest, *Nature*, 452, 737–740, <https://doi.org/10.1038/nature06870>, 2008.
- Li, X., Rohrer, F., Hofzumahaus, A., Brauers, T., Häsel, R., Bohn, B., Broch, S., Fuchs, H., Gomm, S., Holland, F., Jäger, J., Kaiser, J., Keutsch, F. N., Lohse, I., Lu, K., Tillmann, R., Wegener, R., Wolfe, G. M., Mentel, T. F., Kiendler-Scharr, A., and Wahner, A.: Missing gas-phase source of HONO inferred from Zeppelin measurements in the troposphere, *Science*, 344, 292–296, <https://doi.org/10.1126/science.1248999>, 2014.
- Liu, Y. J., Herdinger-Blatt, I., McKinney, K. A., and Martin, S. T.: Production of methyl vinyl ketone and methacrolein via the hydroperoxyl pathway of isoprene oxidation, *Atmos. Chem. Phys. Discuss.*, 12, 33 323–33 358, <https://doi.org/10.5194/acpd-12-33323-2012>, 2012.
- Lockhart, J., Blitz, M., Heard, D., Seakins, P., and Shannon, R.: Kinetic study of the OH + glyoxal reaction: Experimental evidence and quantification of direct OH recycling, *J. Phys. Chem. A*, 117, 11 027–11 037, <https://doi.org/10.1021/jp4076806>, 2013.
- Lou, S., Holland, F., Rohrer, F., Lu, K., Bohn, B., Brauers, T., Chang, C. C., Fuchs, H., Häsel, R., Kita, K., Kondo, Y., Li, X., Shao, M., Zeng, L., Wahner, A., Zhang, Y., Wang, W., and Hofzumahaus, A.: Atmospheric OH reactivities in the Pearl River Delta - China in summer 2006: measurement and model results, *Atmos. Chem. Phys.*, 10, 11 243–11 260, <https://doi.org/10.5194/acp-10-11243-2010>, 2010.
- Novelli, A., Kaminski, M., Rolletter, M., Acir, I. H., Bohn, B., Dorn, H. P., Li, X., Lutz, A., Nehr, S., Rohrer, F., Tillmann, R., Wegener, R., Holland, F., Hofzumahaus, A., Kiendler-Scharr, A., Wahner, A., and Fuchs, H.: Evaluation of OH and HO₂ concentrations and their budgets during photooxidation of 2-methyl-3-butene-2-ol (MBO) in the atmospheric simulation chamber SAPHIR, *Atmos. Chem. Phys.*, 18, 11 409–11 422, <https://doi.org/10.5194/acp-18-11409-2018>, 2018.
- Novelli, A., Vereecken, L., Bohn, B., Dorn, H. P., Gkatzelis, G. I., Hofzumahaus, A., Holland, F., Reimer, D., Rohrer, F., Rosanka, S., Taraborrelli, D., Tillmann, R., Wegener, R., Yu, Z., Kiendler-Scharr, A., Wahner, A., and Fuchs, H.: Importance of isomerization reactions for OH radical regeneration from the photo-oxidation of isoprene investigated in the atmospheric simulation chamber SAPHIR, *Atmos. Chem. Phys.*, 20, 3333–3355, <https://doi.org/10.5194/acp-20-3333-2020>, 2020.
- Nozière, B., Barnes, I., and Becker, K.-H.: Product study and mechanisms of the reactions of α -pinene and of pinonaldehyde with OH radicals, *J. Geophys. Res.*, 104, 23 645–23 656, <https://doi.org/10.1029/1999JD900778>, 1999.
- Orlando, J. J., Nozière, B., Tyndall, G. S., Orzechowska, G. E., Grazyńska, E., Paulson, S. E., and Rudich, Y.: Product studies of the OH- and ozone-initiated oxidation of some monoterpenes, *J. Geophys. Res.*, 105, 11561–11572, <https://doi.org/10.1029/2000JD900005>, 2000.
- Paulot, F., Crouse, J. D., Kjaergaard, H. G., Kroll, J. H., Seinfeld, J. H., and Wennberg, P. O.: Isoprene photooxidation: new insights into the production of acids and organic nitrates, *Atmos. Chem. Phys.*, 9, 1479–1501, <https://doi.org/10.5194/acp-9-1479-2009>, 2009.
- Peeters, J., Boullart, W., Pultau, V., Vandenberk, S., and Vereecken, L.: Structure–activity relationship for the addition of OH to (poly)alkenes: site-specific and total rate constants, *J. Phys. Chem. A*, 111, 1618–1631, <https://doi.org/10.1021/jp066973o>, 2007.
- Peeters, J., Nguyen, T. L., and Vereecken, L.: HO_X radical regeneration in the oxidation of isoprene, *Phys. Chem. Chem. Phys.*, 11, 5935–5939, <https://doi.org/10.1039/b908511d>, 2009.
- Peeters, J., Müller, J.-F., Stavrou, T., and Nguyen, V. S.: Hydroxyl radical recycling in isoprene oxidation driven by hydrogen bonding and hydrogen tunneling: The upgraded LIM1 mechanism, *J. Phys. Chem. A*, 118, 8625–8643, <https://doi.org/10.1021/jp5033146>, 2014.
- Praske, E., Otkjær, R. V., Crouse, J. D., Hethcox, J. C., Stoltz, B. M., Kjaergaard, H. G., and Wennberg, P. O.: Atmospheric autoxidation is increasingly important in urban and suburban North America, *Proc. Nat. Acad. Sci.*, 115, 64–69, <https://doi.org/10.1073/pnas.1715540115>, 2018.



- 750 Praske, E., Otkjaer, R. V., Crounse, J. D., Hethcox, J. C., Stoltz, B. M., Kjaergaard, H. G., and Wennberg, P. O.: Intramolecular hydrogen shift chemistry of hydroperoxy-substituted peroxy radicals, *J. Phys. Chem. A*, 123, 590–600, <https://doi.org/10.1021/acs.jpca.8b09745>, 2019.
- Reissell, A., Aschmann, S. M., Atkinson, R., and Arey, J.: Products of the OH radical- and O₃-initiated reactions of myrcene and ocimene, *J. Geophys. Res.*, 107, ACH 3–1–ACH 3–6, <https://doi.org/10.1029/2001JD001234>, 2002.
- 755 Rindelaub, J. D., McAvey, K. M., and Shepson, P. B.: The photochemical production of organic nitrates from α -pinene and loss via acid-dependent particle phase hydrolysis, *Atmos. Environ.*, 100, 193–201, <https://doi.org/10.1016/j.atmosenv.2014.11.010>, 2015.
- Rohrer, F., Bohn, B., Brauers, T., Brüning, D., Johnen, F.-J., Wahner, A., and Kleffmann, J.: Characterisation of the photolytic HONO-source in the atmosphere simulation chamber SAPHIR, *Atmos. Chem. Phys.*, 5, 2189–2201, <https://doi.org/10.5194/acp-5-2189-2005>, 2005.
- Rolletter, M., Kaminski, M., Acir, I. H., Bohn, B., Dorn, H. P., Li, X., Lutz, A., Nehr, S., Rohrer, F., Tillmann, R., Wegener, R., Hofzumahaus, A., Kiendler-Scharr, A., Wahner, A., and Fuchs, H.: Investigation of the α -pinene photooxidation by OH in the atmospheric simulation chamber SAPHIR, *Atmos. Chem. Phys.*, 19, 11 635–11 649, <https://doi.org/10.5194/acp-19-11635-2019>, 2019.
- 760 Rolletter, M., Blocquet, M., Kaminski, M., Bohn, B., Dorn, H. P., Hofzumahaus, A., Holland, F., Li, X., Rohrer, F., Tillmann, R., Wegener, R., Kiendler-Scharr, A., Wahner, A., and Fuchs, H.: Photooxidation of pinonaldehyde at ambient conditions investigated in the atmospheric simulation chamber SAPHIR, *Atmos. Chem. Phys.*, 20, 13 701–13 719, <https://doi.org/10.5194/acp-20-13701-2020>, 2020.
- 765 Schlosser, E., Brauers, T., Dorn, H.-P., Fuchs, H., Häsel, R., Hofzumahaus, A., Holland, F., Wahner, A., Kanaya, Y., Kajii, Y., Miyamoto, K., Nishida, S., Watanabe, K., Yoshino, A., Kubistin, D., Martinez, M., Rudolf, M., Harder, H., Berresheim, H., Elste, T., Plass-Dülmer, C., Stange, G., and Schurath, U.: Technical Note: Formal blind intercomparison of OH measurements: results from the international campaign HOxComp, *Atmos. Chem. Phys.*, 9, 7923–7948, <https://doi.org/10.5194/acp-9-7923-2009>, 2009.
- Scholtens, K. W., Messer, B. M., Cappa, C. D., and Elrod, M. J.: Kinetics of the CH₃O₂+NO reaction: Temperature dependence of the overall rate constant and improved upper limit for the CH₃ONO₂ branching channel, *J. Geophys. Res.*, 100, 18 811–18 816, <https://doi.org/10.1021/jp990469k>, 1995.
- 770 Schwantes, R. H., Emmons, L. K., Orlando, J. J., Barth, M. C., Tyndall, G. S., Hall, S. R., Ullmann, K., St. Clair, J. M., Blake, D. R., Wisthaler, A., and Bui, T. P. V.: Comprehensive isoprene and terpene gas-phase chemistry improves simulated surface ozone in the southeastern US, *Atmos. Chem. Phys.*, 20, 3739–3776, <https://doi.org/10.5194/acp-20-3739-2020>, 2020.
- 775 Sindelarova, K., Granier, C., Bouarar, I., Guenther, A., Tilmes, S., Stavrou, T., Müller, J. F., Kuhn, U., Stefani, P., and Knorr, W.: Global data set of biogenic VOC emissions calculated by the MEGAN model over the last 30 years, *Atmos. Chem. Phys.*, 14, 9317–9341, <https://doi.org/10.5194/acp-14-9317-2014>, 2014.
- Tan, Z., Fuchs, H., Lu, K. D., Hofzumahaus, A., Bohn, B., Broch, S., Dong, H. B., Gomm, S., Haseler, R., He, L. Y., Holland, F., Li, X., Liu, Y., Lu, S. H., Rohrer, F., Shao, M., Wang, B. L., Wang, M., Wu, Y. S., Zeng, L. M., Zhang, Y. S., Wahner, A., and Zhang, Y. H.: Radical chemistry at a rural site (Wangdu) in the North China Plain: observation and model calculations of OH, HO₂, and RO₂ radicals, *Atmos. Chem. Phys.*, 17, 663–690, <https://doi.org/10.5194/acp-17-663-2017>, 2017.
- 780 Tan, Z., Lu, K., Hofzumahaus, A., Fuchs, H., Bohn, B., Holland, F., Liu, Y., Rohrer, F., Shao, M., Sun, K., Wu, Y., Zeng, L., Zhang, Y., Zou, Q., Kiendler-Scharr, A., Wahner, A., and Zhang, Y.: Experimental budgets of OH, HO₂, and RO₂ radicals and implications for ozone formation in the Pearl River Delta in China 2014, *Atmos. Chem. Phys.*, 19, 7129–7150, <https://doi.org/10.5194/acp-19-7129-2019>, 2019.
- 785 Tan, Z., Hofzumahaus, A., Lu, K., Brown, S. S., Holland, F., Huey, L. G., Kiendler-Scharr, A., Li, X., Liu, X., Ma, N., Min, K.-E., Rohrer, F., Shao, M., Wahner, A., Wang, Y., Wiedensohler, A., Wu, Y., Wu, Z., Zeng, L., Zhang, Y., and Fuchs, H.: No evidence for a significant impact



- of heterogeneous chemistry on radical concentrations in the North China Plain in summer 2014, *Environ. Sci. Technol.*, **54**, 5973–5979, <https://doi.org/10.1021/acs.est.0c00525>, 2020.
- 790 Teng, A. P., Crouse, J. D., and Wennberg, P. O.: Isoprene peroxy radical dynamics, *J. Am. Chem. Soc.*, **139**, 5367–5377, <https://doi.org/10.1021/jacs.6b12838>, 2017.
- Vereecken, L. and Peters, J.: Decomposition of substituted alkoxy radicals - Part I: a generalized structure-activity relationship for reaction barrier heights, *Phys. Chem. Chem. Phys.*, **11**, 9062–9074, <https://doi.org/10.1039/B909712K>, 2009.
- Vereecken, L. and Peters, J.: A structure-activity relationship for the rate coefficients of H-migration in substituted alkoxy radicals, *Phys. Chem. Chem. Phys.*, **12**, 120608–12620, <https://doi.org/10.1039/C0CP00387E>, 2010.
- 795 Vereecken, L. and Nozière, B.: H migration in peroxy radicals under atmospheric conditions, *Atmos. Chem. Phys.*, **20**, 7429–7458, <https://doi.org/10.5194/acp-20-7429-2020>, 2020.
- Vereecken, L., Carlsson, P. T. M., Novelli, A., Bernard, F., Brown, S. S., Cho, C., Crowley, J. N., Fuchs, H., Mellouki, W., Reimer, D., Shenolikar, J., Tillmann, R., Zhou, L., Kiendler-Scharr, A. and Wahner, A.: Theoretical and experimental study of peroxy and alkoxy radicals in the NO₃-initiated oxidation of isoprene, *Phys. Chem. Chem. Phys.*, **23**, 5496–5515, <https://doi.org/10.1039/D0CP06267G>, 800 2021a.
- Vereecken, L.: personal communication, 2021b.
- Wegener, R., Brauers, T., Koppmann, R., Bares, S. R., Rohrer, F., Tillmann, R., Wahner, A., Hansel, A., and Wisthaler, A.: Simulation chamber investigation of the reactions of ozone with short-chained alkenes, *J. Geophys. Res.*, **112**, D13 301, <https://doi.org/10.1029/2006JD007531>, 2007.
- 805 Wennberg, P. O., Bates, K. H., Crouse, J. D., Dodson, L. G., McVay, R. C., Mertens, L. A., Nguyen, T. B., Praske, E., Schwantes, R. H., Smarte, M. D., St Clair, J. M., Teng, A. P., Zhang, X., and Seinfeld, J. H.: Gas-Phase Reactions of Isoprene and Its Major Oxidation Products, *Chem. Rev.*, **118**, 3337–3390, <https://doi.org/10.1021/acs.chemrev.7b00439>, 2018.
- Whalley, L. K., Edwards, P. M., Furneaux, K. L., Goddard, A., Ingham, T., Evans, M. J., Stone, D., Hopkins, J. R., Jones, C. E., Karunaharan, A., Lee, J. D., Lewis, A. C., Monks, P. S., Moller, S. J., and Heard, D. E.: Quantifying the magnitude of a missing hydroxyl radical source 810 in a tropical rainforest, *Atmos. Chem. Phys.*, **11**, 7223–7233, <https://doi.org/10.5194/acp-11-7223-2011>, 2011.
- Xu, L., Guo, H., Boyd, C. M., Klein, M., Bougiatioti, A., Cerully, K. M., Hite, J. R., Isaacman-VanWertz, G., Kreisberg, N. M., Knote, C., Olson, K., Koss, A., Goldstein, A. H., Hering, S. V., de Gouw, J., Baumann, K., Lee, S.-H., Nenes, A., Weber, R. J., and Ng, N. L.: Effects of anthropogenic emissions on aerosol formation from isoprene and monoterpenes in the southeastern United States, *Proc. Nat. Acad. Sci.*, **112**, 37–42, <https://doi.org/10.1073/pnas.1417609112>, 2015.
- 815 Zhang, H. F., Yee, L. D., Lee, B. H., Curtis, M. P., Worton, D. R., Isaacman-VanWertz, G., Offenberg, J. H., Lewandowski, M., Kleindienst, T. E., Beaver, M. R., Holder, A. L., Lonneman, W. A., Docherty, K. S., Jaoui, M., Pye, H. O. T., Hu, W. W., Day, D. A., Campuzano-Jost, P., Jimenez, J. L., Guo, H. Y., Weber, R. J., de Gouw, J., Koss, A. R., Edgerton, E. S., Brune, W., Mohr, C., Lopez-Hilfiker, F. D., Lutz, A., Kreisberg, N. M., Spielman, S. R., Hering, S. V., Wilson, K. R., Thornton, J. A., and Goldstein, A. H.: Monoterpenes are the largest source of summertime organic aerosol in the southeastern United States, *Proc. Nat. Acad. Sci.*, **115**, 2038–2043, 820 <https://doi.org/10.1073/pnas.1717513115>, 2018.



Table 1. Experimental conditions for the oxidation experiments. Concentrations are given for the conditions in the SAPHIR chamber at the time of the first VOC injection.

date	type of VOC	VOC / ppbv	NO / ppbv	O ₃ / ppbv	T / K
16 August 2012	myrcene	2.3	0.2	< 20	305
22 August 2012	myrcene	2.3	0.2	< 15	310
17 July 2013	myrcene	0.8	0.06	40	305
18 July 2013	myrcene	0.8	0.09	40	308
29 May 2020	methane	140000	0.23	60 - 120	303
03 September 2019	α -pinene	8.5	0.1	8	296



Table 2. Instrumentation for radical and trace gas detection in the experiments.

measured quantity	measurement technique	time	precision	accuracy
		resolution	(1 σ)	(1 σ)
OH	differential optical absorption spectroscopy	205 s	$0.6 \times 10^6 \text{ cm}^{-3}$	6.5 %
OH	laser-induced fluorescence (LIF)	47 s	$0.4 \times 10^6 \text{ cm}^{-3}$	13 %
HO ₂ , RO ₂	laser-induced fluorescence (LIF)	47 s	$1.5 \times 10^7 \text{ cm}^{-3}$	16 %
MyO ₂				50 %
k_{OH}	laser photolysis + LIF	180 s	0.3 s^{-1}	10 %, $\pm 0.7 \text{ s}^{-1}$
myrcene, α -pinene	proton-transfer-reaction mass-spectrometry	40 s	15 pptv	7 %
myrcene, acetone	gas-chromatography	30 min	100 pptv	6 %
acetaldehyd				
CH ₄	cavity ring-down spectroscopy	60 s	1 ppbv	3 ppbv
CO	cavity ring-down spectroscopy	60 s	25 ppbv	1 ppbv
NO	chemiluminescence	180 s	4 pptv	5 %
NO ₂	photolytic converter + chemiluminescence	180 s	2 pptv	5 %
HONO	long-path absorption photometry	300 s	7 pptv	20 %
O ₃	UV-absorption	10 s	1 ppbv	5 %
HCHO	differential optical absorption spectroscopy	100 s	20 %	10 %
HCHO	Hantzsch method (Aerolaser)	90 s	100 pptv	10 %
photolysis freq.	actinic flux spectroradiometry	60 s	10 %	10 %



Table 3. Chemical reactions considered in the radical budget analysis of OH, HO₂ and RO₂. Some reactions are only applicable in the experiment with either methane, α -pinene or myrcene, in which specific RO₂ radicals, CH₃O₂, MyO₂ or APO₂ are formed. RO₂+RO₂ reactions are not listed, because they do not significantly contribute to the loss of radicals for conditions of experiments here.

Reaction	$k(T=298\text{ K}, P=1\text{ atm})$	uncertainty ^a	reference
radical initiation reactions			
R1 HONO+h ν (< 400 nm) \rightarrow OH+NO	j_{HONO}	22 %	measured
R2 O ₃ +h ν (< 340 nm) \rightarrow O(¹ D)+O ₂	$j_{\text{O}(\text{1D})}$	11 %	measured
O(¹ D)+H ₂ O \rightarrow OH+OH	$2.1 \times 10^{-10}\text{ cm}^3\text{ s}^{-1}$		IUPAC (2020)
O(¹ D)+M \rightarrow O(³ P)+M	$3.3 \times 10^{-11}\text{ cm}^3\text{ s}^{-1}$		IUPAC (2020)
R3 HCHO+h ν (< 335 nm)+2O ₂ \rightarrow 2HO ₂ +CO	j_{HCHO}	11 %	measured
R4 O ₃ +myrc \rightarrow 0.71 OH+0.71 RO ₂	$3.9 \times 10^{-16}\text{ cm}^3\text{ s}^{-1}$	22 %	Deng et al. (2018); ?
radical propagation reactions			
R5 HO ₂ +NO \rightarrow OH+ NO ₂	$8.5 \times 10^{-12}\text{ cm}^3\text{ s}^{-1}$	27 %	IUPAC (2020)
R6 HO ₂ +O ₃ \rightarrow OH+ 2 O ₂	$2.0 \times 10^{-15}\text{ cm}^3\text{ s}^{-1}$	20 %	IUPAC (2020)
R7 OH+VOC+O ₂ \rightarrow RO ₂			
RO ₂ = CH ₃ O ₂	$6.4 \times 10^{-15}\text{ cm}^3\text{ s}^{-1}$	17 %	IUPAC (2020)
RO ₂ = MyO ₂	$2.3 \times 10^{-10}\text{ cm}^3\text{ s}^{-1}$	25 %	this work
R8 OH+HCHO+O ₂ \rightarrow HO ₂ +CO +H ₂ O	$8.4 \times 10^{-12}\text{ cm}^3\text{ s}^{-1}$	25 %	IUPAC (2020)
R9 OH+CO+O ₂ \rightarrow HO ₂ + CO ₂	$2.3 \times 10^{-13}\text{ cm}^3\text{ s}^{-1}$	17 %	IUPAC (2020)
R10 OH+O ₃ \rightarrow HO ₂ +O ₂	$7.3 \times 10^{-14}\text{ cm}^3\text{ s}^{-1}$	17 %	IUPAC (2020)
R11 RO ₂ +NO \rightarrow HO ₂ +NO ₂ +R'CHO ^c			
RO ₂ = CH ₃ O ₂	$7.7 \times 10^{-12}\text{ cm}^3\text{ s}^{-1}$	27 %	IUPAC (2020)
RO ₂ = MyO ₂	$0.87 \times 9.1 \times 10^{-12}\text{ cm}^3\text{ s}^{-1}$	55 %	IUPAC (2020) ^b
radical termination reactions			
R12 OH+NO ₂ \rightarrow HNO ₃	$1.2 \times 10^{-11}\text{ cm}^3\text{ s}^{-1}$	28 %	IUPAC (2020)
R13 OH+NO \rightarrow HONO	$9.8 \times 10^{-12}\text{ cm}^3\text{ s}^{-1}$	28 %	IUPAC (2020)
R14 RO ₂ +NO \rightarrow RONO ₂			
RO ₂ = CH ₃ O ₂	≈ 0	-	IUPAC (2020)
RO ₂ = APO ₂	$0.18 \times 9.1 \times 10^{-12}\text{ cm}^3\text{ s}^{-1}$	27 %	IUPAC (2020) ^b
RO ₂ = MyO ₂	$0.13 \times 9.1 \times 10^{-12}\text{ cm}^3\text{ s}^{-1}$	55 %	IUPAC (2020) ^b
R15 RO _{2,myrc} +HO ₂ \rightarrow ROOH +O ₂	$9.1 \times 10^{-12}\text{ cm}^3\text{ s}^{-1}$	25 %	this work
R16 HO ₂ +HO ₂ + H ₂ O \rightarrow H ₂ O ₂ +O ₂ + H ₂ O	$4.0 \times 10^{-13}\text{ cm}^3\text{ s}^{-1}$ ^d	25 %	IUPAC (2020)

^a Uncertainty measurements (Table 2) and kinetic rate constants (10 % for reactions of inorganic and methyl peroxy radicals and 15 % for other)

^b Reaction rate constant: IUPAC (2020); the branching ratio of 0.87 is taken from this work

^c The dominant reaction of alkoxy radical is to form HO₂ which is not rate limiting for RO₂ to HO₂ conversion

^d Effective reaction rate constant for 1 % H₂O mixing ratio



Table 4. Rate constants of the myrcene + OH reaction at 298 K.

$k \text{ cm}^3 \text{ s}^{-1}$	method	reference
$(2.1 \pm 0.2) \times 10^{-10}$	relative rate technique	Atkinson et al. (1986)
$(3.4^{+1.5}_{-1.0}) \times 10^{-10}$	relative rate technique	Hites and Turner (2009)
1.8×10^{-10}	structure-activity relationship	Peeters et al. (2007)
$(2.3 \pm 0.3) \times 10^{-10}$	direct measurement	this work



Table 5. Products yields from the reaction of myrcene with OH.

product	yield	temperature / K	reference
formaldehyde	0.30 ± 0.06	n.a.	Orlando et al. (2000)
	0.74 ± 0.08	294	Lee et al. (2006)
	0.35 ± 0.08	293-330	this work
acetone	0.36 ± 0.05	n.a.	Orlando et al. (2000)
	0.45 ± 0.06	296±2	Reissell et al. (2002)
	0.22 ± 0.02	294	Lee et al. (2006)
	0.45 ± 0.08	293-330	this work
organic nitrate	0.10 ± 0.03	294	Lee et al. (2006)
	0.13 ± 0.03	293-330	this work




Figure 1. Chemical structures of myrcene, isoprene and 2-methyl 2-butene.




Figure 2. Peroxy radicals formed in the addition of OH to the double bonds in myrcene. MyO₂ formed from the OH addition to the isoprenyl part of myrcene leads to the formation of MyO₂ species that are expected to rapidly interconvert by oxygen abstraction and elimination like in isoprene (Peeters et al., 2014). Assuming reaction rate constants for oxygen abstraction and elimination like in isoprene, 3'-OH-3-OO and 1-OH-2-OO are expected to have the highest yields among the MyO₂ species that can interconvert. MyO₂ yields are predicted using SAR by Peeters et al. (2007).




Figure 3. Reactions of the four most abundant myrcene hydroxy peroxy radicals with NO forming HO₂ and carbonyl compounds.




Figure 4. Fast H-shift reactions of myrcene peroxy radicals. Rate constants apply to 298 K. They are adopted for the Z-3'-OH-1-OO and Z-1-OH-3'-OO isomers from the corresponding 1,6-H-shift reactions of isoprene peroxy radicals (Peeters et al., 2014). Similar subsequent chemistry like for isoprene leads to the formation of peroxy radicals, some of which rapidly decompose. The isomerization reaction rate constants for the E-3'-OH-1-OO and E-1-OH-3'-OO peroxy radicals are calculated using SAR by Vereecken and Nozière (2020). Subsequent chemistry of products from these reactions likely undergo fast ring-closure reactions on the dimethyl double bond (not shown here).




Figure 5. Trace gas and radical concentrations in the chamber experiment investigating the OH oxidation of myrcene (myrc) at NO mixing ratios between 200 and 300 pptv on 22 August 2012. Peroxy radicals from the reaction of OH with myrcene (MyO₂) and total organic peroxy radical (RO₂=MyO₂+RO_{2,other}) concentrations are calculated from observed radical concentrations as explained in Section 3.4. In the lowest right panel, the loss rates of total RO₂ with respect to their reaction with NO and radical recombination reactions are shown.




Figure 6. Time series of nitrogen oxide species in the experiments with methane (10 July 2013, (a)) and α -pinene (03 September 2019, (c)). The red lines in panels (a) and (c) are time integrated HONO emissions calculated as described in the text. NO_x and HONO concentrations were measured, the time integrated NO_2 loss by HNO_3 formation from the reaction of NO_2 with OH and the time integrated NO_2 loss by RONO_2 formation from the reaction of RO_2 with NO were calculated as described in the text. The alkyl nitrate yield (Φ_{RONO_2}) is determined from the regression analysis (red line, panel (d)) using Equation 8. Errors are derived from error propagation of measurements.




Figure 7. Calculated production and destruction rates of reactions involving radicals for the reference experiment with methane (29 May 2020). Production rates for OH, HO₂, RO₂ and total RO_x are shown in panels (a), (b), (c) and (d), respectively and loss rates for OH, HO₂, RO₂ and total RO_x are shown in panels (f), (g), (h) and (i). Lower panels (j) – (m) show the differences between loss and production rates ($L - P$) with the associated accuracy (grey areas).




Figure 8. Corrected product concentrations versus the myrcene that reacted away with OH for two experiments with medium NO (light colors: 16 August 2012, dark colors: 22 August 2012). Corrections are applied to account for formation of product species not connected to the oxidation of myrcene (chamber sources) and loss processes (reaction with OH, photolysis). The slopes of linear fits (solid lines) give the product yields of HCHO and acetone from the myrcene reaction with OH. The uncertainties of the yield calculations are indicated by dashed lines.




Figure 9. Determination of the organic nitrate yield from the production of nitrogen oxide species ($Q(\text{HONO})$) in the chamber experiment. Upper panel: Example of cumulated reactive nitrogen oxide mixing ratios over the course of the experiment on 18 July 2013. The blue line denotes the total production of nitrogen oxides including organic nitrate applying a nitrate yield (Φ_{RONO_2}) of 0.13. Dashed lines show the error of the total nitrogen oxide calculation. Lower panel: Scatter plot of integrated turnover rate of the reaction of RO_2 with NO versus the unaccounted nitrogen oxide mixing ratios (ΔNO_y , Equation 8). Results from all four experiments are included. The organic nitrate yield is determined by the slope of regression analysis (red line, $R^2 = 0.92$). Error bars denotes experimental uncertainty derived from the accuracy of the measurements (Table 2) and dashed red lines the resulting uncertainty in the slope of the regression analysis.




Figure 10. RO_x primary production, P , (upper panels), termination L (middle panels) rates and their difference (lower panels). Grey areas in the lower panels give the uncertainty of the difference between radical destruction and production ($L - P$). The rate constant of the reaction of MyO_2 with HO_2 was adjusted, to minimize the difference between radical production and destruction (see text for more explanation).




Figure 11. Rates of radical conversion reactions and imbalances between production and destruction rates ($L - P$) for the experiment with medium NO mixing ratios (0.15 to 0.30 ppbv) on 22 August 2012. Grey areas in the lower panels give the uncertainty of the calculation ($L - P$). In the lowest right panel, the black and cyan lines denote RO₂ loss rates without and with considering MyO₂ isomerization reactions.




Figure 12. Rates of radical production and destruction reactions for OH, HO₂ and RO₂ for experiments at low NO mixing ratio (<0.11 ppbv) on 18 July 2013. Grey areas in the lower panels give the uncertainty of $L - P$. In the lowest right panel, the black and cyan lines denote RO₂ budget without and with considering MyO₂ isomerization reactions.