	1	Aerosol optical properties calculated from size distributions, filter samples and absorption photometer data	
	2	at Dome C, Antarctica and their relationships between seasonal cycles of sources	
	3		
	4	Aki Virkkula ^{1,3} , Henrik Grythe ² , John Backman ¹ , Tuukka Petäjä ³ , Maurizio Busetto ⁴ , Christian Lanconelli ^{4,56} ,	
	5	Angelo Lupi ⁴⁵ , Silvia Becagli ^{67,8} , Rita Traversi ^{67,8} , Mirko Severi ^{67,8} , Vito Vitale ⁷⁵ ,	Formatted: Superscript
	6	Patrick Sheridan ⁸⁹ , Elisabeth Andrews ^{8,9,<u>10</u>}	
•	7		
	8	¹ Finnish Meteorological Institute (FMI), Erik Palmenin aukio 1, P.O.Box 503, FI-00101 Helsinki	
	9	² Norwegian Institute for Air Research (NILU), P.O.Box 100, 2027 KJELLER	
	10	³ University of Helsinki, Institute for Atmospheric and Earth System Research (INAR) / Physics, Faculty of	
	11	Science, Helsinki, Finland	
1	12	⁴ <u>National Research Council of Italy,</u> Institute of Atmospheric Sciences and Climate, National Research Council,	
	13	(<u>CNR-</u> ISAC -CNR), <u>Via P. Gobetti 101, 40129</u> Bologna, Italy	
	14	⁵ National Research Council of Italy, Institute of Polar Sciences (CNR-ISP), Via P. Gobetti 101, 40129 Bologna,	
	15	Italy	
	16	⁵⁶ European Commission, Joint Research Centre, Ispra (VA), Italy	
	17	⁶² University of Florence, Dept. of Chemistry, Sesto Fiorentino (Florence) – Italy	
	18	⁷⁸ National Research Council of Italy, Institute of Polar Sciences, (CNR-ISP)-CNR, University of Venice, V. Torino	
•	19	155, 30172 Venice-Mestre, Italy.	
	20	⁹⁸ NOAA Global Monitoring Laboratory, Boulder (CO), USA	
	21	⁹¹⁰ Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder (CO), USA	
	22		
	23	Correspondence to: A. Virkkula (aki.virkkula@fmi.fi)	
	24		
	25	Abstract	
	26	Optical properties of surface aerosols at Dome C, Antarctica in 2007-2013 and their potential source areas are	
	27	presented. Scattering coefficients (σ_{sp}) were calculated from measured particle number size distributions with a	
	28	Mie code and from filter samples using mass scattering efficiencies. Absorption coefficients ($\sigma_{\mbox{\scriptsize ap}}$) were	
	29	determined with a 3-wavelength Particle Soot Absorption Photometer (PSAP) and corrected for scattering by	
	30	using two different algorithms. The scattering coefficients were also compared with σ_{sp} measured with a	
	31	nephelometer at the South Pole Station (SPO). The minimum σ_{ap} was observed in the austral autumn and the	
	32	maximum in the austral spring, similar to other Antarctic sites. The darkest aerosol, i.e., the lowest single	

1 scattering albedo $\omega_o \approx 0.91$ was observed in September and October and the highest $\omega_o \approx 0.99$ in February and 2 March. The uncertainty of the absorption Ångström exponent α_{ap} is high. The lowest α_{ap} monthly medians 3 were observed in March and the highest in August - October. The equivalent black carbon (eBC) mass 4 concentrations were compared with eBC measured at three other Antarctic sites: the SPO and two coastal 5 sites, Neumayer and Syowa. The maximum monthly median eBC concentrations are almost the same (~3 ± 1 ng 6 m³-) at all these sites in October-November. This suggests that there is no significant difference in eBC 7 concentrations between the coastal and plateau sites. The seasonal cycle of the eBC mass fraction exhibits a 8 minimum f(eBC) \approx 0.1% in February-March and a maximum ~4-5% in August-October. Source areas were 9 calculated using 50-day FLEXPART footprints. The highest eBC concentrations and the lowest ω_o were 10 associated with air masses coming from South America, Australia and Africa. Vertical simulations that take BC 11 particle removal processes into account show that there would be essentially no BC particles arriving at Dome 12 C from north of latitude 10°S at altitudes < 1600 m. The main biomass-burning regions Africa, Australia and 13 Brazil are more to the south and their smoke plumes have been observed at higher altitudes than that so they 14 can get transported to Antarctica. The seasonal cycle of BC emissions from wildfires and agricultural burning 15 and other fires in South America, Africa and Australia were calculated from data downloaded from the Global 16 Fire Emissions Database (GFED). The maximum total emissions were in August-September but the peak of 17 monthly average eBC concentrations is observed 2 - 3 months later in November not only at Dome C but also 18 at SPO and the coastal stations. The air mass residence-time-weighted BC emissions from South America are 19 approximately an order of magnitude larger than from Africa and Oceania suggesting that South American BC 20 emissions are the largest contributors to eBC at Dome C. At Dome C the maximum and minimum scattering 21 coefficients were observed in austral summer and winter, respectively. At SPO σ_{sp} was similar to that observed 22 at Dome C in the austral summer but there was a large difference in winter, suggesting that in winter SPO is 23 more influenced by sea spray emissions than Dome C. The seasonal cycles of σ_{sp} at Dome C and at the SPO 24 were compared with the seasonal cycles of secondary and primary marine aerosol emissions. The σ_{sp} measured 25 at SPO correlated much better with the sea-spray aerosol emission fluxes in the Southern Ocean than σ_{sp} at 26 Dome C. The seasonal cycles of biogenic secondary aerosols were estimated from monthly average 27 phytoplankton biomass concentrations obtained from the CALIOP satellite sensor data. The analysis suggests 28 that a large fraction of the biogenic scattering aerosol observed at Dome C has been formed in the polar zone 29 but it may take a month for the aerosol to be formed, grown and get transported from the sea level to Dome C.

30

32

1 1. Introduction

2 The Antarctic interior region has scarce observations of atmospheric constituents and many aspects of the 3 atmospheric properties are underdetermined. The Antarctic dome or the polar vortex, which is much stronger 4 than its northern counterpart and present throughout the year (Karpetchko et al., 2005), at most times 5 efficiently prevents transport into the Antarctic troposphere from lower latitudes. However, wildfires and 6 agricultural burning emissions from Africa, South America and Australia do affect vast regions of the southern 7 hemisphere, including Antarctica. For instance, Hara et al. (2010) found that haze episodes at Syowa Station, 8 during which visibility can drop to 10 km for periods of ~30 h, were caused by biomass burning aerosol from 9 South America transported to the Antarctic coast via the eastward approach of cyclones. At the Neumayer 10 station large-scale meridional transport of biomass-burning derived black carbon, preferentially from South 11 America, seems to determine the BC burden and causes a distinct and consistent spring / early summer 12 concentration maximum (Weller et al., 2013).

14 Concordia station lies on Dome C (75°06'S, 123°23'E), at 3233 m above sea level (.a.s.l.) on the East Antarctic 15 plateau, about 1100 km from the nearest coastline, the Ross Sea. The base is French / Italian operated, with 16 research fields within astronomy and glaciology as well as atmospheric sciences. The atmospheric 17 instrumentation is located in a small cabin southwest of the main base (at the site described by Udisti et al., 18 2012) where it is upwind of the base at the prevailing wind directions. Concordia is one of only three 19 permanent year-round stations operated on the Antarctic Plateau, the others being the American Amundsen-20 Scott observatory (South Pole (SPO), 2835 m.a.s.l., about 1300 km from the nearest open sea, 1600 km away 21 from Dome C) and the Russian Vostok station (78°28'S, 106°51'E, 3488 m.a.s.l., 600km away). Thus, there are 22 large spatial distances between the continuous atmospheric observation. However, properties of the Antarctic 23 atmosphere tend to extend both over longer temporal and spatial scales than elsewhere (Fiebig et al., 2014) 24 suggesting that the scarce observations that exist can be assumed to be representative of larger areas than 25 typical in other climate regions. This would imply that Dome C is an important indicator for the entire Antarctic 26 inland. Though measurement conditions are harsh the continuous long-term monitoring provided here can be 27 a baseline for the aerosol optical properties of the Antarctic inland and may provide indications of changes in 28 atmospheric constituents and aerosol levels.

29

13

30 There are several studies on the aerosol chemical composition at Dome C (e.g., Jourdain et al., 20098; Becagli

31 et al., 2012, 2021; Udisti et al., 2004, 2012; Legrand et al., 2016, 2017a, 2017b), and also the aerosol optical

32 depth (AOD) has been measured there (Tomasi et al., 2007). However, in situ surface aerosol scattering and

Formatted: English (United States)

Formatted: Highlight

1 absorption coefficients at Dome C have not been presented. The light absorption coefficient and particle 2 number size distributions (PNSD) have been measured continuously with a 3-wavelength Particle Soot 3 Absorption Photometer (PSAP) and a differential mobility particle sizer (DMPS) since 2007. The PNSD data have 4 already been used in several papers. Järvinen et al. (2013) analyzed the seasonal cycle and modal structure of 5 PNSD measured with the DMPS, Chen et al. (2017) analyzed number size distribution of air ions measured with 6 an Air Ion Spectrometer (AIS) and the PNSD measured with the DMPS and Lachlan-Cope et al. (2020) used the 7 Dome C DMPS data for comparing with the PNSD measured at the coastal site Halley. The PSAP data, however, 8 have not been presented in detail. Caiazzo et al. (2021) used some of the PSAP data mainly for evaluating 9 elemental carbon (EC) sample contamination. Grythe (2017) used the data from 2007-2013 as part of his PhD 10 thesis but in the present paper we will analyze that period in more detail. Here we will describe the methods 11 for measuring absorption and calculating scattering from the size distributions and filter samples.

12

The goals of the paper are to present descriptive statistics of extensive and intensive aerosol optical properties at Dome C in 2007 – 2013, their seasonal cycles and the relationships between the seasonal cycles of major sources of absorbing and scattering aerosols. The AOPs will be compared with other observations from other Antarctic sites, in most detail the scattering coefficients measured at the South Pole.

17

18 2. Methods

19 2.1 Sampling site

20 Concordia station is a permanently operated French / Italian Antarctic research base on East Antarctic plateau. 21 The observations are performed in isolated sites around the main base. The Dome C sampling site is the same 22 as used by Udisti et al. (2012), Becagli et al. (2012), and Järvinen et al. (2013). It is located about 1 km 23 southwest of the station main buildings, upwind in the direction of the prevailing wind. The northeastern 24 direction (10°-90°) has been declared as the contaminated sector. Below the validity of the contaminated 25 sector will be analyzed by using the absorption photometer data. For in situ aerosol instrumentation the 26 sample air was taken at the flowrate of 5 Liters Per Minute (LPM) from the roof of the cabin with a straight 2-m 27 long 25-mm diameter stainless steel tube inlet. It was covered with a protective cap to protect against snow fall 28 and ice buildup.

- 29
- 30
- 31
- 32

1 2.2 Instruments

2 2.2.1 Aerosol measurements

3 Light absorption by particles was measured with a Radiance Research 3 λ PSAP at three wavelengths, λ = 467 4 nm, 530 nm, and 660 nm. There was no nephelometer measuring scattering coefficient so it was calculated 5 from particle size distributions and filter sample data as described below. Particle number size distributions 6 were measured at 10-minute time resolution in the size range 10 - 620 nm with a custom-built differential 7 mobility particle sizer (DMPS) as described by Järvinen et al. (2013) and in the size range 0.3 – 20 μ m with a 8 Grimm model 1.108 optical particle counter (OPC) in 2007 – 2009. RH was not measured in the Dome C sample 9 air but it can be safely claimed that it was dry. The absolute humidity in the air on the upper plateau is very low 10 and temperature varies from colder than about -20°C in the austral summer down to about 80°C in the austral 11 winter. When air is sampled to the instruments in the measurement containers where temperature is >+10 $^{\circ}{
m C}$ 12 RH decreases to very low values. In addition to the in-situ instruments, PM₁ and PM₁₀ filter samples were 13 collected for chemical analyses by ion chromatography. The length of the sampling period of the PM_1 and PM_{10} 14 samples was 3 or 4 days and 1 day, respectively.

15

The data coverage for the PSAP, the DMPS, the OPC and the PM₁ and PM₁₀ filter sample data are presented in Fig. 1. The number of hours of accepted data and the number of samples are shown in parentheses for the continuous instruments and the filter samplers, respectively. The filtering criteria will be presented below (section 2.4).

20

21 2.2.2 Meteorological measurements

Ambient air temperature (t), relative humidity (RH), wind speed (WS) and wind direction (WD) data were from the routine meteorological observation at Station Concordia as part of the IPEV/PNRA Project - a collaborative project between "Programma Nazionale di Ricerche in Antartide" (PNRA) and Institut Polaire Français Paul-Emile Victor (IPEV) (<u>www.climantartide.it</u>).

26

27 2.3 Data processing

28 2.3.1 Mass concentrations from size distributions

29 60-minute average size distributions n(D_p) were first calculated from the original 10-minute data and corrected

30 for STP (p = 1013 hPa, T = 273.15 K). The DMPS $n(D_p)$ data were corrected for diffusion losses during the

- 31 inversion (Järvinen et al., 2013; Chen et al., 2017), Mass concentration were calculated from the number size
- 32 distributions measured with the DMPS from

Formatted: Highlight

Formatted: Highlight

$$m(DMPS) = \rho_{p} V(DMPS) = \rho_{p} \int_{10nm}^{620nm} \frac{\pi}{6} D_{p}^{3} n(D_{p}) dD_{p}$$
(1)

where the density $\rho_p = 1.7 \text{ g cm}^3$ was used. For particle density of 1.7 g cm⁻³ the particle diameter 620 nm corresponds to the aerodynamic diameter $D_a = \sqrt{\rho_p / \rho_0} D_p = \sqrt{1.7} \times 620 \text{nm} \approx 808 \text{ nm}$, where $\rho_0 = 1 \text{ g cm}^{-3}$. To be consistent with the definitions of filter-sample size ranges that typically show the upper aerodynamic diameter of a sampler inlet the mass concentration calculated from Eq. (1) will be referred to as m(DMPS,PM_{0.8}) and the volume concentration as V(DMPS,PM_{0.8}).

7

1

8 In December 2007 – July 2009 there was also the Grimm 1.108 OPC that measures number concentrations of

9 particles in the D_p range of $0.3 - 20 \ \mu$ m. The particle number concentrations in the size range Dp > 1 μ m were 10 first corrected for wind-speed dependent and particle diameter dependent inlet and sampling tube losses as 11 described in the supplement. The three largest channels of the OPC measure the number concentrations in D_p 12 range of 7.5 - 20 μ m. For an assumed density $\rho_p = 1.7 \ \text{g cm}^{-3}$ the diameter D_p = 7.5 μ m corresponds to the 13 aerodynamic diameter D_a = 9.8 μ m. Assuming that ρ_p is constant over the whole size range the mass 14 concentration of particles smaller than D_a = 10 μ m is calculated from the number size distributions by excluding 15 the three largest particle OPC channels as

16
$$m(n(D_p), PM_{10}) = \rho_p V(n(D_p), PM_{10}) = \rho_p (V(DMPS, PM_{0.8}) + V(OPC, PM_{0.8-10}))$$
 (2)

17 The fraction of volume concentration measured by the DMPS equals

18
$$fV(DMPS) = \frac{V(DMPS, PM_{0.8})}{V(n(D_p), PM_{10})}$$

19 This fraction was calculated from data collected during the simultaneous operation of the DMPS and the OPC.

20 The monthly average fV(DMPS) values presented in Table 1 were used for the period 2008 – 2013 to calculate 21 mass concentrations in the size range $D_a < 10 \ \mu m$ from

22
$$m(DMPS, PM_{10}) = \frac{\rho_p V(DMPS, PM_{0.8})}{fV(DMPS)} = \frac{m(DMPS, PM_{0.8})}{fV(DMPS)}$$
(4)

In other words, the variable names m(DMPS,PM_{0.8}) and m(DMPS,PM₁₀) will be used below to emphasize that
 these mass concentrations were calculated from DMPS data. The mass concentrations m(DMPS,PM_{0.8}) and
 m(DMPS,PM₁₀) can be considered to be the lower and upper estimates of m.

26

27 2.3.2 Scattering coefficients from the size distributions

28 Scattering coefficients were calculated using the 60-minute average size distributions from

Formatted: Highlight Formatted: Highlight Formatted: Highlight Formatted: Highlight

(3)

$$\sigma_{\rm sp}({\rm m},\lambda) = \left| Q_{\rm s}({\rm D}_{\rm p},{\rm m},\lambda)\frac{\pi}{4}{\rm D}_{\rm p}^2{\rm n}({\rm D}_{\rm p}){\rm d}{\rm D}_{\rm p} \right|$$

where Q_s is the scattering efficiency calculated using the Mie code by Barber and Hill (1990), m is the refractive index, λ is the wavelength and $n(D_p)$ is the particle number size distribution. Analogous to the mass concentrations the scattering coefficients were determined from the simultaneous DMPS and OPC measurements in December 2007 – July 2009 from

$$\sigma_{sp}(n(D_p), PM_{10}) = \sigma_{sp}(DMPS, PM_{0.8}) + \sigma_{sp}(OPC, PM_{0.8-10})$$

7 where $\sigma_{sp}(OPC,PM_{0.8})$ and $\sigma_{sp}(OPC,PM_{0.8-10})$ are the scattering coefficient calculated from the particle number 8 size distributions in the size ranges measured by the DMPS and the OPC, respectively. As explained above, the 9 <u>number size distributions, for D_A > 1 μ m were corrected for the inlet and sampling tube losses.</u> For 10 $\sigma_{sp}(\frac{OPEDMPS}{2}, PM_{0.8})$ the refractive index of sulfuric acid (SA, H₂SO₄, m_r = 1.426 + 0i, Seinfeld and Pandis, 1998) 11 was used. This refractive index is slightly lower than that estimated for submicron aerosols at two low-altitude 12 Antarctic stations Aboa, and Neumayer in Queen Maud Land. Virkkula et al. (2006) -measured particle number 13 size distributions in the size range D_p < 800 nm with a DMPS and light scattering of submicron particles with a 14 nephelometer at the Finnish a site about 130 km inland from the open Weddell Sea in January 2000. With an 15 iteration procedure matching nephelometer-measured and size-distribution-derived scattering coefficients the 16 real refractive indices were 1.43 \pm 0.07 and 1.45 \pm 0.04 at λ = 550 nm for all data and excluding new particle 17 formation, respectively. Jurányi and Weller (2019) measured size distributions with an SMPS and a laser 18 aerosol spectrometer (LAS) for a full year at the coastal site Neumayer and by fitting data of the two 19 instruments in the overlapping range of 120 - 340 nm obtained $m_r = 1.44 \pm 0.08$. Considering that both Aboa 20 and Neumayer are closer to sources of ammonia that neutralizes aerosol and increases the refractive index 21 above that of pure sulfuric acid (1.426) it was assumed here that the use of 1.426 for the calculation of σ_{sp} from 22 the size range measured with the DMPS is reasonable. For the larger particle size range, σ_{sp} (OPC,PM_{0.8-10}) the 23 refractive index of NaCl (mr = 1.544, Seinfeld and Pandis, 1998) was used. This value is in line with the average 24 refractive index of 1.54 with a range from 1.50 to 1.58 in the particle size range 0.3 - 12 µm in impactor 25 samples taken at the South Pole (Hogan et al., 1979) and with the supermicron particle refractive index of 1.53 26 ± 0.02 calculated from the chemical composition of 12-stage impactor samples taken at the coastal site Aboa 27 (Virkkula et al., 2006).

28

1

6

 $29 \qquad \text{The fraction of scattering coefficient measured by the DMPS was calculated from} \\$

30
$$f\sigma_{sp}(DMPS, \lambda) = \frac{\sigma_{sp}(DMPS, PM_{0.8}, \lambda)}{\sigma_{sp}(n(D_p), PM_{10}, \lambda)}$$

Formatted: Highlight Formatted: Highlight Formatted: Subscript, Highlight
Formatted: Highlight Formatted: Subscript, Highlight
Formatted: Subscript, Highlight
1 · 5 5
Formatted: Highlight



(6)

(7)

1 The wavelengths of λ = 467 nm, 530 nm, and 660 nm were used to match the PSAP data. Similar to fV(DMPS), 2 fo_{sp}(DMPS, λ) was calculated from data collected during the simultaneous operation of the DMPS and the OPC, 3 the seasonal monthly statistics were calculated (Table 1) and the respective monthly averages were applied to 4 the period 2008 – 2013 to calculate o_{sp} in the size range D_a < 10 µm from

5
$$\sigma_{\rm sp}(\rm DMPS, \rm PM_{10}, \lambda) = \frac{\sigma_{\rm sp}(\rm DMPS, \rm PM_{0.8}, \lambda)}{f\sigma_{\rm sp}(\rm DMPS, \lambda)}$$
(8)

6 The wavelength symbol λ will be used below only when necessary. The variable names $\sigma_{sp}(DMPS,PM_{0.8})$ and 7 $\sigma_{sp}(DMPS,PM_{10})$ will be used to emphasize that these scattering coefficients were calculated from DMPS data in 8 the aerodynamic particle size ranges $D_a < 0.8 \ \mu m$ and $D_a < 10 \ \mu m$. The scattering coefficients $\sigma_{sp}(DMPS,PM_{0.8})$ 9 and $\sigma_{sp}(DMPS,PM_{10})$ can also be considered to be the lower and upper estimates of σ_{sp} at the given 10 wavelength.

11

12 Figure 2 shows the average particle number, volume and scattering size distributions at λ = 530 nm in the size 13 range 10 nm - 10 μ m and the respective normalized cumulative size distributions in the size range of 10 nm -14 7.5 µm during the period from 14 December 2007 to 14 July 2009 in summer and in winter. Fig. 2a and 2b show that for the number concentrations the OPC size range plays an insignificant role whereas the larger particles 15 16 contribute significantly to both total particle volume concentration (Figs. 2c and 2d) and scattering coefficients 17 (Figs. 2e and 2f) and that this contribution varies seasonally. The contributions of fV(DMPS) and $f\sigma_{sp}(DMPS,\lambda)$ 18 were calculated for hourly-averaged size distributions from Eqs. (3) and (7), the monthly seasonal statistics 19 were calculated and presented in Table 1 and as a boxplot in Fig. 3. Both the table and the boxplot show that 20 both fV(DMPS) and $f\sigma_{sp}(DMPS,\lambda)$ have maxima in summer and minima in winter. They also show that the 21 ranges are large. Consequently the use of the monthly averages presented in Table 1 for calculating 22 m(DMPS,PM₁₀) and σ_{so} (DMPS,PM₁₀), Eqs. (4) and (8), creates an additional uncertainty to the results. Another 23 important result is that the wavelength dependency of $f\sigma_{sp}(DMPS,\lambda)$ is clear and it also has a seasonal cycle.

24

25 The wavelength dependency of the scattering coefficient can be described by the scattering Ångström 26 exponent

27
$$\alpha_{sp} = -\frac{\ln(\sigma_{sp}(\lambda_1)/\sigma_{sp}(\lambda_2))}{\ln(\lambda_1/\lambda_2)}$$
(9)

that was calculated by using the wavelength pair 467/660 nm. The variable names $\alpha_{sp}(DMPS,PM_{0.8})$ and $\alpha_{sp}(DMPS,PM_{10})$ will be used below for α_{sp} calculated from $\sigma_{sp}(DMPS,PM_{0.8})$ and $\sigma_{sp}(DMPS,PM_{10})$, respectively.

1 2.3.3 Absorption coefficients and equivalent black carbon concentrations

The PSAP data were first corrected for flow and spot size. The flow was calibrated 37 times during 2007 - 2013
with a TSI flow meter. The slopes and offsets of the calibrations were interpolated for each hour and the PSAP
flows were corrected accordingly. All absorption coefficients were corrected to STP (1013.25 hPa and 273.15
K).

6

7 The PSAP measures signal and reference detector counts and the respective sums, $\sum SIG$ and $\sum REF$ are used 8 for calculating non-scattering-corrected absorption coefficient, here $\sigma_{ap,nsc}$, from

9
$$\sigma_{ap,nuc} = f(Tr) \frac{A}{Q\Delta t} \ln \left(\frac{\left(\sum_{\Delta t} SIG / \sum_{\Delta t} REF \right)_{t,\Delta t}}{\left(\sum_{\Delta t} SIG / \sum_{\Delta t} REF \right)_{t}} \right) = f(Tr)\sigma_{0}$$
(10)

where A is the filter spot area, Q the flow rate, $Tr = (\Sigma SIG/\Sigma REF)_t/(\Sigma SIG/\Sigma REF)_{t=0}$ is the transmittance, f(Tr) the loading correction function and Δt the count integration time. The PSAP reports $\sigma_{ap,nsc}$ with a 0.1 Mm⁻¹ resolution at a 1-second time resolution. Averaging the 1-sec data is not good enough since at Dome C absorption coefficients are most of the time clearly lower than 0.1 Mm⁻¹. Therefore the signal and reference counts ΣSIG and ΣREF were used in (10) with $\Delta t = 60$ min. Manufacturer-cut spots of the standard filter material Pallflex E70-2075W were used in the PSAP. The spot diameter was measured to be 4.9 ± 0.1 mm, so the spot area A was 18.9 ± 0.6 mm². The uncertainty of A is ~3%.

17

Transmittance is reduced mainly by light absorption but also also due to scattering aerosol which results in the so-called apparent absorption and has to be taken into account in the data processing. There are different algorithms for processing PSAP data, e.g. by Bond et al. (1999), Virkkula et al. (2005), Müller et al. (2014), and Li et al. (2020). Here we will use both the algorithm presented by Bond et al. (1999) (here B1999) with the adjustment presented by Ogren (2010):

23
$$\sigma_{ap} = \frac{1}{1.22} \left(\frac{0.97 \cdot 0.873}{1.0796 \cdot Tr + 0.71} \sigma_0 - 0.02 \cdot \sigma_{sp} \right) = \frac{1}{1.5557 \cdot Tr + 1.0227} \sigma_0 - 0.0164 \cdot \sigma_{sp} = \sigma_{ap,nsc} - 0.0164 \cdot \sigma_{sp}$$
(11)

and the algorithm presented by Virkkula et al. (2005) with the constants updated by Virkkula (2010) (hereV2010):

26
$$\sigma_{ap} = \left(k_0 + k_1(h_0 + h_1\omega_0)\ln(Tr)\right)\sigma_0 - s \cdot \sigma_{sp}$$
(12)

27 where

$$28 \qquad \qquad \omega_0 = \sigma_{sp} / \left(\sigma_{sp} + \sigma_{ap} \right) \tag{13}$$

Formatted: English (United States)

1 is the single-scattering albedo and k_0 , k_1 , h_0 , h_1 , and s are wavelength-dependent constants. In the rest of the 2 paper the symbol $\sigma_{ap,nsc}$ will be used to present the non-scattering-corrected absorption coefficient, corrected 3 with the constants and formula in Eq. (11) excluding the subtraction of σ_{sp} .

4

5 Since there are the above-explained size-dependent uncertainties of scattering coefficient, additional 6 absorption coefficient estimates were calculated by using both algorithms. The upper estimates of absorption 7 coefficients $\sigma_{ap}(\sigma_{sp}(DMPS,PM_{0.8}))$ were calculated by using the lower estimate of scattering coefficient σ_{sp} = 8 $\sigma_{sp}(DMPS,PM_{0.8})$ in the scattering corrections in Eqs. (11) and (12) and the lower estimates of absorption 9 coeffcient $\sigma_{ap}(\sigma_{sp}(DMPS,PM_{10}))$ were calculated by using the upper estimate of scattering coefficient σ_{sp} = 10 σ_{so} (DMPS,PM₁₀) in the scattering corrections. Consequently the lower and upper estimates of ω_o are denoted 11 as $\omega_o(\sigma_{sp}(DMPS,PM_{0.8}))$ and $\omega_o(\sigma_{sp}(DMPS,PM_{10}))$, respectively. They were calculated by using both Eqs. (11) and 12 (12) for calculating σ_{ap} .

13

14 Considering that the period with the simultaneous measurements with the DMPS and the OPC showed that the 15 DMPS size range always leads to an underestimation of both aerosol mass and scattering coefficient, it is likely 16 that σ_{ap} corrected for scattering with σ_{sp} (DMPS, PM₁₀) is closer to the true σ_{ap} than that corrected with 17 σ_{sp} (DMPS, PM_{0.8}). In the results both $\sigma_{ap,nsc}$, σ_{ap} (σ_{sp} (DMPS,PM_{0.8})) and σ_{ap} (σ_{sp} (DMPS,PM₁₀)) will be presented to 18 evaluate the effect of using only the size range mesured with the DMPS for the scattering correction.

19

22

Similar to σ_{sp} , the wavelength dependency of light absorption by particles can roughly be described by the absorption Ångström exponent:

$$\alpha_{ap} = -\frac{\ln\left(\sigma_{ap}(\lambda_1)/\sigma_{ap}(\lambda_2)\right)}{\ln\left(\lambda_1/\lambda_2\right)} \tag{14}$$

that was calculated by using λ = 467 nm and 660 nm for $\sigma_{ap,nsc}$, $\sigma_{ap}(\sigma_{sp}(DMPS,PM_{10}))$ and both Eqs. (11) and (12). The variable names $\alpha_{ap}(\sigma_{ap,nsc})$, $\alpha_{ap}(\sigma_{sp}(DMPS,PM_{10}),B1999)$ and $\alpha_{ap}(\sigma_{sp}(DMPS,PM_{10}),V2010)$, respectively, will be used to denote the α_{ap} calculated in different ways. These calculations were conducted to study the uncertainty of α_{ap} due to scattering corrections.

27

28 The absorption coefficient was used to estimate the concentration of equivalent black carbon, eBC (Petzold et 29 al. 2013) from:

$$30 \qquad eBC = \frac{\sigma_{ap}}{MAC} \tag{15}$$

where MAC is the mass absorption coefficient. For freshly-emitted BC the MAC value is approximately 7.5 m² g⁻ 1 2 $^{-1}$ at λ = 550 nm (Bond et al., 2013). By assuming a wavelength-dependency of λ^{-1} this corresponds to MAC \approx 3 7.8 m² g⁻¹ at λ = 530 nm. This can be considered to yield an upper estimate for eBC concentrations since for 4 coated BC particles MAC is larger (Bond et al., 2013). eBC was calculated by using $\sigma_{ap,nsc}$ and 5 $\sigma_{ap}(\sigma_{sp}(DMPS,PM_{10}))$ calculated with both algorithms, Eq. (11) and (12). The corresponding variable names 6 $eBC(\sigma_{ap,nsc})$ and $eBC(\sigma_{sp}(DMPS,PM_{10}))$ will be used below for them. The scattering-corrected 7 $eBC(\sigma_{so}(DMPS,PM_{10}))$ can be considered to be closer to the true eBC concentration. The reason for also 8 presenting eBC($\sigma_{ap,nsc}$) is that often an estimate of BC concentrations is needed even if it is known that it is an 9 upper estimate (Caiazzo et al., 2021). It is also comparable with the eBC often presented from Aethalometer 10 measurements. Presenting both yields a quantitative estimate of the bias due to not correcting the data for 11 scattering.

12

13 The eBC mass fractions in the two size ranges $D_a < 0.8 \ \mu m$ and $D_a < 10 \ \mu m$ were calculated from

14
$$feBC(m(DMPS, PM_{0.8})) = 100\% \frac{eBC}{m(DMPS, PM_{0.8})}$$
 (16)

15
$$feBC(m(DMPS, PM_{10})) = 100\% \frac{eBC}{m(DMPS, PM_{10})}$$
 (17)

where the mass concentrations m(DMPS, PM_{0.8}) and m(DMPS, PM₁₀) were defined in Eq. (4) and eBC calculated from Eq. (15). Mass fractions were calculated for eBC($\sigma_{ap,nsc}$) and eBC(σ_{sp} (DMPS, PM₁₀)).

18 19 20

21 2.3.4 Noise of scattering and absorption coefficients and eBC

The uncertainty of scattering coefficients should in principle be calculated from the error propagation formula 22 23 $(\delta\sigma_{sp})^2 = \sum (\partial\sigma_{sp}/\partial x_i)^2 (\delta x_i)^2$, where ∂x_i is the uncertainty of variable x_i in calculating σ_{sp} (e.g., Sherman et 24 al., 2015). That would require taking into account all uncertainties of the size distribution measurements and 25 Mie modeling. However, a simplified approach was used here. The σ_{sp} calculated from the size distribution data 26 and the uncertainty of the size distribution range were used for calculating lower and upper estimates of σ_{sp} as 27 explained above. In addition to that the noise of σ_{sp} was estimated from the average of the absolute 28 differences of all two consecutive hourly-averaged scattering coefficients $\delta \sigma_{sp}$ (average, 1h) = 29 average($|\Delta \sigma_{sp}(\mathbf{1}h)|$) = average($|\sigma_{sp}(t_{i+1}) - \sigma_{sp}(t_i)|$). The average noise of 24-h averages was calculated from $\delta\sigma_{sp}(24h) = \delta\sigma_{sp}(average, 1h)/\sqrt{24}$. The noises were calculated for both $\sigma_{sp}(DMPS, PM_{0.8})$) and 30

1 $\sigma_{sp}(DMPS,PM_{10})$. The noises are presented in Table 2. Note that the difference $|\sigma_{sp}(t_{i+1}) - \sigma_{sp}(t_i)|$ is not only due 2 to random noise so higher $|\Delta\sigma_{sp}|$ values are observed when σ_{sp} is in reality increasing or decreasing so the true 3 random noise is slightly lower. When σ_{sp} is used in calculating the scattering correction of σ_{ap} in B1999 (Eq. 4 (11)) σ_{sp} is multiplied by 0.0164. Consequently, the σ_{sp} noise for the 24-h averages results in a 0.0164 σ_{sp} noise 5 for σ_{ap} . These noises are also presented in Table 2.

6

7 The uncertainty of the absorption coefficient should also be calculated from the error propagation formula, 8 similar to Sherman et al. (2015). However, here only the uncertainty of the spot size (~3%) and the statistical 9 noise are taken into account. The noise of the non-scattering-corrected hourly $\sigma_{ap,nsc}$ was estimated from the 10 average of the absolute differences of all two consecutive absorption measurements $\delta \sigma_{ap,nsc}$ (average) = 11 $average(|\Delta \sigma_{ap,nsc}|) = average(|\sigma_{ap,nsc}(t_i) - \sigma_{ap,nsc}(t_i)|)$ similar to the noise estimate of σ_{sp} . The noise of 24-hour averages was estimated from $\delta\sigma_{ap,nsc}(24h) = \delta\sigma_{ap,nsc}(average, 1h)/\sqrt{24}$. The noise in the scattering-12 13 corrected absorption cofficients were calculated from $\delta \sigma_{ap} = \delta \sigma_{ap,nsc} + 0.0164 \delta \sigma_{sp}$ for both $\sigma_{ap}(\sigma_{sp}(DMPS,PM_{0.8}))$ 14 and $\sigma_{ap}(\sigma_{sp}(DMPS, PM_{10}))$ and for 1h and 24h averages (Table 2). The noise determined this way is formally 15 correct only for σ_{ap} calculated with the B1999 formula, Eq. (11), not for V2010. However, calculated directly 16 fom the absolute differences, the average $|\Delta\sigma_{ap}(B1999)|\approx average |\Delta\sigma_{ap}(V2010)|$, but the contribution of 17 scattering to the noise was only determined for B1999 as explained above. For V2010, Eq. (12) a formal error 18 propagation calculation is more complicated due to the iterative form of the procedure and it is out of the 19 scope of the present paper. The noise of eBC was calculated from $\delta(eBC(\sigma_{ap})) = \delta\sigma_{ap}/MAC$ for both non-20 scattering-corrected and scattering-corrected eBC. The detection limits were defined as $2\times\delta(eBC(\sigma_{ap}))$. The 21 results are presented in Table 3.

22 23

The largest uncertainty factor for σ_{ap} , ω_{o} , α_{ap} , and eBC is not related to noise. It is due to the uncertainty of the refractive index and size distributions used for calculating σ_{sp} and the algorithm. This was evaluated by calculating σ_{ap} by using the lower and upper estimates of σ_{sp} in both scattering correction algorithms. These four values were used then for calculating ω_{o} , α_{ap} , and eBC and they are presented below in relevant tables and figures.

29

30 2.4 Filtering and preprocessing the in situ data

Both PSAP absorption and DMPS-derived scattering coefficient data were filtered manually by removing rapidly changing values since they can be assumed to result from contamination from the station or from some technical problem. The PSAP transmittance data were used to filter out data measured at Tr < 0.7 following
 recommendations in WMO/GAW Report No. 227 (2016) and the PSAP handbook (Springston, 2018). During
 most of 2010 the PSAP flow was extremely unstable so practically the whole year was removed.

5 All major sources of light absorbing aerosol other than the Dome C base are so far away that rapid variations in 6 $\sigma_{ap,nsc}$ are due to either instrument malfunction or influence from the base, for instance emissions from 7 vehicles. Further filtering of the data was done by removing data in which 10-minute averages of $\sigma_{ap,nsc}$ were 8 more than 10 times larger than the hourly $\sigma_{ap,nsc}$. This was done to remove short events that are local but do 9 not appear to come directly from the base, based on wind direction. In all roughly 13% of the data were 10 deemed contaminated.

12 Additionally, wind data were used to remove clear contamination from the station. The sampling site is located 13 upwind of the base itself by the prevailing wind directions. The base has a year-round diesel generator and 14 vehicles operated within the base-area move around the base from November to February. Fig. 4 shows the 15 distribution of $\sigma_{ap,nsc}$ in 5° wind direction (WD) sectors at wind speed WS > 2 m s⁻¹. The generator at the base is 16 clearly observed as a pronounced peak of in $\sigma_{ap,nsc}$ at WD 60°. If the 75th percentile of the $\sigma_{ap,nsc}$ cumulative 17 distribution is used as the criterion for the contaminated sector data when sector data when the wind direction 18 was between 30°<WD<90° would be filtered which is 6 % of the of data. If the 99th percentile of $\sigma_{ap,nsc}$ is used 19 the contamination sector is wider, 20°<WD<110°, and 10% of the data would be filtered. Here the latter, i.e., 20 the stricter criterion was used. The distribution of $\sigma_{\text{ap,nsc}}$ in the same WD sectors at several wind speed intervals 21 are shown in the supplement, Fig. S1. It is obvious that at low wind speeds contaminated air can come from all 22 directions. Therefore, when WS < 2 m s⁻¹ all data were filtered out, regardless of WD.

Since the size distribution and absorption measurements are done in the same cabin, the DMPS and OPC data were also removed when the PSAP observations indicated contamination. Fig. 1 shows the instruments' operational time in hours. The DMPS measurements had more gaps than the PSAP. The three instruments required for a valid measurement were not always operational at the same time. After filtering, altogether 15815 hours of data remained for the statistical analyses. No filtering was applied to the PM₁ and PM₁₀ filter samples. The contamination is mainly BC so it was be assumed that the effect on ion concentrations was not significant.

31

23

4

11

1 The calculations were done using hourly-averaged data. These data were filtered to remove contaminated data 2 as explained above. The filtered data were then averaged over 24 hours to reduce noise and improve detection 3 limits. In the discussions below, the running 24-hour averages were used, centered at each hour, i.e. $\sigma_{ap}(t,24H)$ 4 = average($\sigma_{ap}(t-12,1H),...,\sigma_{ap}(t+11,1H)$) which means, for instance, that at noon $\sigma_{ap}(t=12,24H)$ = 5 average($\sigma_{ap}(t=0,1H), ..., \sigma_{ap}(t=23,1H)$) so the noon average represents all absorption coefficients measured 6 during that day. If, during any period to be averaged, there were less than 12 hours of non-contaminated data 7 then that 24-hour average was excluded from further analysis.

8

9 2.5 Filter sample analyses and data processing

There were two samplers in the immediate vicinity of the cabin where the other in situ measurements were made. There was a PM_{10} sampling head operating following the CSN EN 12341 European Standard. The PM_1 samples were collected on the backup filter of a Dekati PM_{10} impactor. In both of these particles were sampled on Teflon filters (Pall-Gelman, 47-mm diameter, 2- μ m nominal porosity). PM_{10} and PM_1 load is obtained by summing the mass of the ions determined on Teflon filters. Note that this can be considered to be the lower estimate since there could be unidentified compounds, such as organic carbon on the filters.

16

17 Just before the analysis, half of each filter was extracted with 10 mL of ultrapure water (18 M Ω Milli-Q) in 18 ultrasonic bath for 20 min. Every filter manipulation was carried out under a class-100 laminar-flow hood, to 19 minimize contamination risks. Inorganic anions and cations, as well as selected organic anions, were 20 simultaneously measured by using a three Thermo Scientific Dionex ion-chromatography system, equipped 21 with electrochemical-suppressed conductivity detectors. The sample handling during the IC injection was 22 minimized by using a specifically-designed Flow-Injection Analysis (IC-FIA) device (Morganti et al., 2007). 23 Cations (Na⁺, NH₄⁺, K⁺, Mg²⁺ and Ca²⁺) were determined by using a Thermo Scientific Dionex CS12A-4 mm 24 analytical column with 20 mM H_2SO_4 eluent. Inorganic anions (Cl⁻, NO₃⁻, SO₄²⁻ and C₂O₄²⁻) were measured by a 25 Thermo Scientific Dionex AS4A-4 mm analytical column with a 1.8 mM Na₂CO₃/1.7 mM NaHCO₃ eluent. F⁻ and 26 some organic anions (acetate, glycolate, formate and methanesulfonate) were determined by a Thermo 27 Scientific Dionex AS11 separation column by a gradient elution (0.075-2.5 mM Na2B4O7 eluent). Further details 28 on the ion chromatographic measurements are reported in Udisti et al. -(2004)-and and Becagli et al. (2011, 29 2021). All concentrations were corrected to STP (1013.25 hPa and 273.15 K). The ion data used in the present 30 work is a subset of the data from 2005 to 2013 that Becagli et al. (2021) K)

Formatted: Highlight

Formatted: Highlight

31 <u>used for an analysis of the relationships between non-sea-salt sulfate, MSA, biogenic sources and</u>
 32 <u>environmental constraints.</u>

2 3 In addition to calculating scattering coefficients from the DMPS data, PM1 and PM10 mass concentrations were 4 also used tofor calculatinge the scattering coefficients. The scattering coefficients were calculated by 5 multiplying the mass concentrations with mass scattering efficiencies (MSE) presented by Hand and Malm 6 (2007). The PM₁₀ mass concentrations were multiplied by the mass scattering efficiency of 1.9 m² g⁻¹ and the 7 PM1 concentrations were multiplied by 3.6 m² g¹. These are the MSE for "total mixed" aerosol and "fine mixed" 8 aerosol in the Table 5 in Hand and Malm (2007), respectively. It has to be kept in mind that the MSE values in 9 the above-mentioned paper were derived from measurements in the continental USA so they most likely have 10 a high uncertainty when applied to the Dome C aerosol. The MSE values presented by Quinn et al. (2002) were 11 used for calculating scattering coefficient of nss sulfate in PM₁ filters.

14 **2.6 Scattering data from the South Pole**

At the South Pole Station (SPO) light scattering coefficient has been measured for more than 40 years. An integrating nephelometer was installed in 1979 and used to measure σ_{sp} , at four wavelengths (450, 550, 700, 850 nm). This nephelometer (Meteorology Research Inc. (MRI), Altadena, CA) was used until its failure in 2002, and a TSI Model 3563 3-wavelength nephelometer (λ = 450 nm, 550 nm, and 700 nm) replaced it in November 2002. (Sheridan et al., 2016). Running 24-hour averages of σ_{sp} (550 nm) were calculated for the years 2007 – 2013 the same way as was done for the Dome C data. The data were used for comparisons with σ_{sp} calculated from the Dome C data.

23 2.7 Source area analyses

The airmass history and transport of aerosols to Dome C were calculated with the Lagrangian dispersion model FLEXPART (Stohl et al., 2005; Pisso et al, 202019). ECMWF reanalysis meteorology was used to run 60000 trajectories every 6 hour 50 days backwards from Dome C to make a statistical sampling of the air measured there. The FLEXPART trajectories follow the mean flow of the atmosphere plus random perturbations to account for turbulence.

29

22

1

12 13

In backward mode, the FLEXPART output is emission sensitivity <u>S</u> or that is proportional to residence time within a grid <u>cell</u> (Stohl et al., 2005; Hirdman et al., 2010; Pisso et al, 2019S). <u>Depending on the settings the output unit of FLEXPART in the backward runs, can be s, s m³ kg⁻¹ or s kg m⁻³. In the present work the unit of S</u>

Formatted: Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Formatted: Highlight

Formatted: Highlight
Formatted: Highlight

1 is seconds (s). When coupled with emissions, FLEXPART emission sensitivity creates a concentration at the 2 release point that is equivalent to forward simulations from emissions, except for some small numerical 3 differences (Seibert and Frank, 2004). One advantage of using a backward simulation in a case like this is that 4 the emission sensitivity fields can be used not only to simulate concentrations but also directly to quantitatively 5 describe exactly where the air that reaches Dome C originates, and, thus, potential emissions influences. 6 Emission sensitivity close to the surface – here at levels < 1000 m a.g.l – is often called the footprint (e.g., 7 Hirdman et al. 2010). If a footprint were multiplied with emission mass flux in kg m³ s¹ at some grid cell the 8 result would be a concentration due to that emission at the receptor site (Stohl et al., 2005). In the present 9 work this step was not done<u>.</u>

Formatted: Highli	ght
Formatted: Super	script
Formatted: Super	script
Formatted: Highli	ght

11 To investigate the role of removal processes during transport, for all model runs, two different tracers were 12 used, one atmospheric tracer with no removal and simulated BC particles with a lognormal size distribution 13 (geometric mean diameter = 150 nm, geometric standard deviation 1.5) experiencing both dry and wet 14 deposition. All tracers were run backwards for 50 days, in most cases sufficient for the aerosol tracer to have 15 less than 1e-12 of the emission sensitivity of the inert air tracer, meaning any emission prior to this would have 16 been removed by the time of arrival at Dome C. The wet removal differentiates removal within and below 17 clouds, also considering the water phase of the clouds and the precipitation type. The FLEXPART removal 18 parameters are the efficiency of aerosols to serve as cloud condensation nuclei (CCN_{eff}) and ice nuclei (IN_{eff}). 19 The values used for them were CCN_{eff} = 0.9 and IN_{eff} = 0.1 as in Table 4 of Grythe et al. (2017). The FLEXPART 20 below-cloud scavenging is a scheme based on Laakso et al. (2003) and Kyrö et al. (2009), both described in 21 Grythe et al. (2017). The model includes a realistic distribution of clouds by incorporating three-dimensional 22 cloud information from ECMWF. For a detailed description see Grythe et al. (2017).

23

10

24 2.7.1 Footprint difference calculations

A statistical analysis was applied to differentiate types of air pathways using a method derived from Hirdman et al. (2010). With the main aim to investigate the different pathways to Dome C, each 6hr interval was given a rank in regards to eBC concentration and single-scattering albedo. The emission sensitivity of the 50-day transport for an aerosol tracer was sorted according to its relative type. The emission sensitivities of the highest (S_H) and the lowest (S_L) 10% of eBC concentration and ω_o were calculated by averaging their emission sensitivities for a given grid cell $\frac{1}{2}$ j, and by:

31
$$S_* = \frac{1}{M} \sum_{m=1}^{M} S(i, j, m)$$

(18)

- 1 Where M is the number of measurements, and S_{*} can be any of the sorting criteria. The relative difference
- 2 between two emission sensitivities S_1 and S_2 in % is then calculated as:

$$RD_{1,2}(i,j) = 100\% \frac{S_1(i,j) - S_2(i,j)}{S_1(i,j) + S_2(i,j)}$$

4 where S₁ and S₂ are the two footprints. In the calculation the emission sensitivities close to the surface, at
5 1000 m a.g.l. were used and so Eq. (19) can be called the relative difference of footprints. This analysis of the
6 footprints can be used to differentiate between different influencing factors on the airmass. This can be either
7 the influence of transport, or removal or combination of these (transport efficiency) or the emission strength.

9 2.7.2 Emissions used for interpreting the footprint statistics and observed seasonal cycles

10 The Global Fire Emissions Database (GFED) is a satellite information-based fire activity map. Monthly gridded 11 burned area and emissions from fires are included in the product (http://www.globalfiredata.org). Emitted BC 12 is calculated based on emission factors, which depend on the type of vegetation that is burning. Satellites give 13 snapshots collected to give pseudo global coverage and not continuous coverage. GFED v3.1 is based on the 14 area burned, which is derived by coupling Moderate resolution Imaging Spectroradiometer (MODIS) fire pixel 15 counts with surface reflectance images (Giglio et al., 2006, 2009, 2010). This widely used emission inventory 16 has uncertainties that arrive both from the emission factors and also from the amount of burnt material. A 17 comparison of this bottom-up inventory with top-down inventories found large regional differences, and top-18 down estimates were about 30% higher (Bond et al., 2013).

19

3

8

20 For the scattering aerosol two sources were considered. An off-line tool (FLEX-SSA) developed by Grythe et al. 21 (2014) and Grythe (2017) to simulate sea spray aerosol (SSA) with FLEXPART was used. It uses inputs from the 22 ECMWF model. These inputs are the wind speed at 10 m above the surface (U10) and the sea surface 23 temperature (SST). The tool takes into account the sea ice fraction which is important to the Southern Ocean 24 SSA emissions. The other major marine scattering aerosols discussed below are biogenic secondary aerosols. 25 Behrenfeld et al. (20176) _estimated monthly average phytoplankton biomass (Cphyto) concentrations in 2007 – 26 2015 from the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) satellite sensor data in three zones: 27 zone $1 = 45^{\circ} - 55^{\circ}$ S, zone $2 = 55^{\circ} - 65^{\circ}$ S and zone $3 = 65^{\circ} - 75^{\circ}$ S. The data provided by Behrenfeld (2021, personal 28 communication) were used for calculating seasonal monthly C_{phyto} averages in the three zones in 2008 – 2013. 29 C_{phyto} can be used as a proxy of biological activity and emissions of dimethyl sulfide (DMS), a precursor of 30 secondary biogenic aerosols.

31

Formatted: Highlight

Formatted: Highlight

(19)

17

2

1 3. Results and discusion

2 3.1 Overview of the data

The time series of σ_{sp} calculated from the size distributions and from the PM₁ and PM₁₀ concentrations and the $\sigma_{ap,nsc}$ at Dome C and σ_{sp} measured with the nephelometer at the SPO are presented in Fig. 5. For the DMPSderived σ_{sp} only the upper estimate, σ_{sp} (DMPS,PM₁₀), Eq. (8) is shown. The descriptive statistics of aerosol optical properties and mass concentrations in the whole period are presented in Tables 4 and 5.

8 Several observations can be made from the time series in Fig. 5. First, the scattering coefficients calculated 9 from the size distributions and the filter samples follow each other relatively well. There is a clear seasonal 10 cycle of both σ_{sp} and $\sigma_{ap,nsc}$. It is clearly seen that $\sigma_{ap,nsc}$ follows the temporal variations of σ_{sp} (DMPS), the high 11 and low values occur mainly simultaneously which is good, considering that these two AOPs were measured 12 with independent instruments. Since the PSAP and other filter-based absorption photometers are sensitive not 13 only to absorbing but also to scattering aerosol and since Dome C is far from BC sources it is possible that the 14 good correlation is due to the apparent absorption only. Below this will be studied simply by using Eqs. (11) and 15 (12) to account for the scattering artifact on the absorption measurement.

16

7

17 The σ_{sp} at Dome C and SPO agree better in austral summer than in winter. However, many high-concentration 18 episodes are observed also in winter almost simultaneous at Dome C and SPO. As an example, a four-month 19 period in May-August 2011 is presented in more detail in Fig. 6. The figure shows 24-hour running averages of 20 σ_{sp} , σ_{ap} and ω_o at λ = 530 nm at Dome C and σ_{sp} at λ = 550 nm at SPO. Fig. 6a shows the upper estimate σ_{sp} = 21 $\sigma_{sp}(DMPS,PM_{10})$, the corresponding lower estimate of $\sigma_{ap} = \sigma_{ap}(\sigma_{sp}(DMPS,PM_{10}))$ (corrected according to 22 B1999, Eq. 11) and the upper estimate of $\omega_{0.}$ Fig. 6b presents the lower estimate of $\sigma_{sp} = \sigma_{sp}$ (DMPS, PM_{0.8}), the 23 corresponding upper estimate of $\sigma_{ap} = \sigma_{ap}(\sigma_{sp}(DMPS, PM_{0.8}))$ and the lower estimate of ω_o . In both Fig. 6a and 24 6b also the non-scattering-corrected absorption coefficient $\sigma_{ab,nsc}$ is shown.

25

In Fig. 6a, the period denoted by (1) shows an episode in which ω_o decreases significantly for several days, being an example of long-range-transported eBC. Episodes (2) and (4) are examples of periods when σ_{sp} is approximately an order of magnitude higher at SPO than at Dome C. There are also events such as episode (3) when σ_{sp} is approximately the same at both sites. The peaks often seem to appear slightly earlier at SPO than at Dome C, suggesting transport from SPO to Dome C rather than the other way around. An example of this is shown in the footprint (Fig. 7) calculated for the episode denoted by (3) in Fig. 6. The footprint shows that the air masses came from the direction of the Antarctic peninsula via SPO to Dome C. Air flow from the direction of 1 the Weddell Sea to SPO and then to Dome C is consistent with a very long known winter-time circulation 2 pattern (Alt et al., 1959) as reviewed by Shaw (1979). During the event denoted by episode (3) $\sigma_{ap,nsc}$ was also 3 high. However, when the scattering correction (Eq. 11) was applied the resulting σ_{ap} was not especially high 4 and ω_o was in the range of 0.98 – 1.00 for both the upper and lower estimates of σ_{sp} , which indicates that non-5 scattering-corrected absorption coefficients may be considerably overestimated when σ_{sp} is high.

6

7 The scattering coefficients calculated from the size distributions, averaged over the filter sampling periods

8 correlate positively with the scattering coefficients calculated from the PM₁ and PM₁₀ filters (Fig. 8). Ordinary 9 least squares regression was used here. The main purpose of the regression was to study whether there is a 10 statistically significant correlation between the scattering coefficients calculated from the size distributions and 11 the filter samples. According to the slopes 0.78 ± 0.02 and 0.76 ± 0.04 of the regression lines in Figd. 8a and 8c 12 σ_{sp} (DMPS, PM_{0.8}) seems to be the lower estimate of σ_{sp} also when it is compared with the filter-sample-13 derived σ_{sp} . According to the slope of 1.29 ± 0.04 in Fig. 8b σ_{sp} (DMPS, PM₁₀) is an upper estimate of σ_{sp} 14 compared with $\sigma_{sp}(PM_{10})$ but when $\sigma_{sp}(DMPS, PM_{10})$ it is compared with $\sigma_{sp}(PM_1)$ the slope is 1.01 ± 0.06 which 15 appears to be somewhat controversial. There are also other peculiarities in the scatter plots. The scatter plot of 16 σ_{sp} (DMPS, PM_{0.8}) vs. σ_{sp} (PM₁₀) (Fig. 8a) have data points where σ_{sp} (DMPS) is low, in the range of ~0.02 – 0.03 17 Mm⁻¹ but $\sigma_{sp}(PM_{10})$ varies in a much larger range from ~0.02 to ~0.9 Mm⁻¹. This also occurs when $\sigma_{sp}(DMPS)$, 18 PM_{10}) is compared with $\sigma_{sp}(PM_{10})$ (Fig. 8b). The pattern could be explained by too low values of of both 19 $\sigma_{sp}(DMPS, PM_{0.8})$ and $\sigma_{sp}(DMPS, PM_{10})$ or by too high values of $\sigma_{sp}(PM_{10})$. Similar suspicious pattern is not 20 observed in the comparison with the PM₁ filters (Fig. 8c and 8d) suggesting the problem may be with $\sigma_{sp}(PM_{10})$. 21 It is clear that this is not a calibration of either the size-distribution-derived or the filter-sample-derived σ_{sp} but 22 the main message of the regressions is that the values are in the same order of magnitude and that there is a 23 statistically significant positive correlation between them which increases confidence in the results. When the 24 regressions are compared with each other it has to be kept in mind that the sampling periods and the number 25 of samples of the PM₁ and PM₁₀ data were not the same. 26

Other reasons for the wide scatter of the data points are the mass scattering efficiencies (MSE) used for calculating scattering coefficients from the filter samples (see section 2.5), uncertainties in ion analyses from the filters and uncertainties in calculating scattering coefficient from the size distributions, especially the estimation of $\sigma_{sp}(DMPS,PM_{10})$ from size distributions measured with the DMPS only. In spite of all these uncertainties the statistical values (averages and percentiles of the cumulative distributions) of the scattering coefficients are reasonably similar. For instance, the medians of $\sigma_{sp}(PM_{10}, \lambda=550 \text{ nm}), \sigma_{sp}(PM_{1}, \lambda=550 \text{ nm})$,

1 $\sigma_{sp}(\text{DMPS, PM}_{0.8}, \lambda\text{=}530 \text{ nm})$ and $\sigma_{sp}(\text{DMPS, PM}_{10}, \lambda\text{=}530 \text{ nm}),$ were 0.24 Mm $^{-1}$, 0.24 Mm $^{-1}$, 0.15 Mm $^{-1}$, and 2 0.2²³ Mm⁻¹, respectively (Table 4). The fact that the medians of $\sigma_{sp}(PM_{10})$ and $\sigma_{sp}(PM_1)$ are the same is 3 somewhat suspicious, it would be expected that $\sigma_{s0}(PM_1) < \sigma_{s0}(PM_{10})$. At this point it is worth paying attention 4 to the statistics of the mass concentrations calculated from the size distributions and from the sum of ions in 5 the filter samples (Table 5). The median mass concentrations of the PM1 and PM10 filters were 66 ng m⁻³ and 6 126 ng m⁻³, respectively, in the expected order. These mass concentrations are also in reasonably good 7 agreement with median m(DMPS, PM_{0.8}) of 70 ng m⁻³ and median m(DMPS, PM₁₀) of 110^{08} ng m⁻³ (Table 5). 8 This suggests that the MSE values used for calculating scattering coefficients from the filter masses were not 9 correct. As it was written in section 2.5 the MSE values were taken from Hand and Malm (2007) who derived 10 them from measurements conducted mainly in US national parks. Considering this, the agreement of the filter-11 sample-derived with the size-distribution-derived σ_{sp} is reasonable.

13 **3.2 Seasonal cycles of AOPs**

12

28

14 **3.2.1** Seasonal cycles of scattering and absorption coefficients

15 The seasonal cycles of scattering and absorption coefficients are presented in Fig. 9. The SPO scattering 16 coefficients presented in Fig. 9a-d were measured using the TSI nephelometer and the Dome C scattering 17 coefficients were calculated using the PM1 (Fig. 9a) and PM10 (Fig. 9b) filter sample data as explained in section 18 2.5 and and from the number size distributions (Figs. 9c-d). The maximum and minimum monthly average and 19 median scattering coefficients were observed in austral summer and winter, respectively. At SPO the 20 scattering coefficient was similar to that at Dome C in austral summer but there was a large difference in 21 austral winter. At SPO the maximum monthy average scattering coefficients were observed in austral winter 22 but at Dome C in austral summer. This suggests that in austral winter SPO is more influenced by sea spray 23 emissions than Dome C. However, even though the averages and medians are lower at Dome C high scattering 24 coefficients are also occasionally observed there in austral winter, as is shown by the 95th percentiles in Fig. 9c 25 and 9d and above in the time series of winter 2011 (Fig. 6). The data does not explain the reasons of the 26 difference between Dome C and SPO in austral winter. It may either be due to different geographical locations, 27 different size ranges measured by the instruments or both.

	29	A hypothetical explanation for the difference between the scattering coefficients at SPO and Dome C could be
--	----	--

- 30 that in the very dry conditions the particles are not spherical. It is true that the shape of particles affects light
- 31 scattering. However, it mainly affects the polarization of scattered light; spherical particles do not change the
- 32 state of the polarization of scattered light but nonspherical particle do. This is used for example in polarization

1	Formatted: Highlight
1	Formatted: Highlight

1	lidars to discriminate ice crystals, dust particles and droplets. However, integral photometric characteristics,
2	such as extinction, scattering and absorption cross sections and single-scattering albedo do not depend
3	significantly on particle shape as is shown in chapter, 10 of the textbook by Mischenko et al. (2002). Therefore
4	nonsphericity is not a likely explanation for the difference.
5	

6 The minimum monthly means and medians of σ_{ap} at Dome C were observed in austral autumn (MAM) and the 7 maximum monthly means and medians in austral spring (SON), which is different than the seasonal cycle of σ_{sp} . 8 (Fig 9e and 9f, Tables S2 and S4). As a result, the seasonal cycle of the single-scattering albedo ω_o is such that 9 the darkest aerosol, i.e., the lowest ω_o is observed in September and October and the highest ω_o in February 10 and March (Fig. 9g and 9h, Table S5). When the lower estimate for σ_{sp} (i.e., σ_{sp} (DMPS, PM_{0.8})) is used for the 11 scattering correction (Eqs. (11) and (12)) the October monthly medians of ω_{0} are 0.862 and 0.868 when using 12 the B1999 and V2010 algorithms, respectively, and when the upper estimate σ_{sp} (DMPS, PM₁₀) is used for the scattering corrections the October monthly medians of ω_o are 0.91109 and 0.9146 when using the B1999 and 13 14 V2010 algorithms, respectively (Table S5). The highest monthly median single-scattering albedos are ~0.98 and 15 > 0.99 with both algorithms when using the σ_{sp} lower and upper estimates for the scattering corrections, 16 respectively. These results show that when σ_{sp} is not measured but calculated from the size distributions the 17 σ_{ap} and ω_{o} are clearly less sensitive to the selection of the algorithm (B1999 or V2010) than to the scattering 18 coefficient used for the scattering correction. But as was noted in section 2.3.4, it is likely that 19 $\sigma_{ap}(\sigma_{sp}(DMPS,PM_{10}))$ is closer to the true absorption coefficient than $\sigma_{ap}(\sigma_{sp}(DMPS,PM_{0.8}))$ so we can also 20 consider the seasonal cycles presented in Figs. 9d, 9f and 9h to be the closest to the true ones.

21

22 **3.2.2** Seasonal cycles of scattering and absorption Ångström exponents

23 The wavelength dependency of both scattering and absorption have clear seasonal cycles. The average 24 scattering Ångström exponent of particles in the DMPS size range, α_{sp} (DMPS,PM_{0.8}) varies from ~2.6 in austral 25 summer (DJF) to ~2.1 in austral winter (JJA) indicating that in austral summer the size distributions are 26 dominated by smaller particles than in winter (Fig. 10a, Table S3). This cycle is much clearer, when α_{sp} is 27 calculated from the upper estimate of scattering: average α_{sp} (DMPS, PM₁₀) varies from ~1.9 in austral summer 28 to ~0.8 in winter. The seasonal cycle of α_{sp} (DMPS,PM₁₀) is actually strikingly similar to the seasonal cycle of α_{sp} 29 of σ_{sp} measured at SPO. This supports the use of the wavelength-dependent formula (Eq. 8) for calculating 30 σ_{sp} (DMPS,PM₁₀, λ) from σ_{sp} (DMPS,PM_{0.8}, λ). The range of α_{sp} is much larger at SPO than at Dome C, however. The 31 main reason is probably that when $\sigma_{sp}(DMPS, PM_{10}, \lambda)$ was calculated with Eq. (8) only the monthly averages of

22

Formatted: Highlight

1 $f\sigma_{sp}(DMPS,\lambda)$ (Eq. 7) were used but the $f\sigma_{sp}(DMPS,\lambda)$ range is actually quite large (Fig. 3). The SPO values were 2 calculated from direct PM₁₀ scattering measurements from a nephelometer.

3

The absorption Ångström exponent α_{ap} was calculated for the non-scattering corrected absorption coefficient $\sigma_{ap,nsc}$ and for the scattering-corrected $\sigma_{ap}(\sigma_{sp}(DMPS,PM_{10}))$ with the two algorithms. Close to the σ_{ap} detection limit the ratios of σ_{ap} at two wavelengths are very noisy so Fig. 10b, Table 4 and Table S6 present α_{ap} statistics of absorption coefficients for $\sigma_{ap} > 3 \times \delta \sigma_{ap}$ where $\delta \sigma_{ap}$ is the wavelength-dependent 24-h average noise at $\lambda =$ 467 nm and $\lambda = 660$ nm (Table2). Note that the number of accepted data points is lower for the scatteringcorrected than for the non-scattering-corrected α_{ap} (Table 4). The reason is that the scattering correction often decreases σ_{ap} below $3 \times \delta \sigma_{ap}$.

11

17

12 The first observation that can be made from looking at the statistics (Fig. 10b, Table 4 and Table S6) is that 13 $\alpha_{ap}(\sigma_{sp}(DMPS,PM_{10}),V2010)$ is always larger than $\alpha_{ap}(\sigma_{ap,nsc})$ and $\alpha_{ap}(\sigma_{sp}(DMPS,PM_{10}),B1999)$. The main 14 explanation of this is that the constants in the V2010 algorithm (Eq. 12) depend on wavelength but the B1999 15 algorithm (Eq. 11) uses the same constants for all wavelengths. The differences between the α_{ap} obtained from 16 different algorithms were also discussed by Backman et al. (2014) and Luoma et al. (20<u>21</u>19).

The seasonal cycles of $\alpha_{ap}(\sigma_{ap,nsc})$ and $\alpha_{ap}(\sigma_{sp}(DMPS,PM_{10}),B1999)$ are qualitatively similar: the lowest medians are observed in March and the maxima in August – October. This cycle is approximately anticorrelated with the ω_{o} seasonal cycle: in March the median ω_{o} is the highest and the lowest in August – October. In March the median $\alpha_{ap}(\sigma_{ap,nsc})$ and $\alpha_{ap}(\sigma_{sp}(DMPS,PM_{10}),B1999)$ were ~0.6 and 0.37 and in August-September 0.96 and ~0.92-0.965, respectively (Table S6), essentially the value generally used for pure BC. The seasonal cycle of $\alpha_{ap}(\sigma_{sp}(DMPS,PM_{10}),V2010)$ is a little bit different: the minimum median of ~1.2 is in February and the maximum of ~1.7 occurs in June (Table S6).

25

The interpretation of α_{ap} is complicated. The α_{ap} is related to the dominant absorbing aerosol type but physical properties of the particles also affect it. For externally mixed BC particles it is generally assumed to be around 1 (Hegg et al., 2002; Bond and Bergstrom, 2006; Bond et al., 2013a) and higher for some organic aerosol from biomass smoke and mineral dust (Kirchstetter et al., 2004; Russell et al., 2010; Devi et al, 2016). However, α_{ap} also depends on the size of BC cores and coating thickness. It is easy to show with Mie models that for single non-coated BC particles with $D_p < ~20$ nm α_{ap} is indeed close to 1, but when $D_p \approx 100$ nm $\alpha_{ap} \approx 1.3$ depending on the wavelength pair used for the calculation and < 1 when $D_p > ~150$ nm. For BC particle size distributions

1 the width and the dominant particle size affect α_{ap} . Coating of BC cores affects α_{ap} even more: when BC 2 particles are coated either with a light-absorbing shell or even with a light-scattering shell α_{ap} can be clearly 3 larger than 1 (e.g., Gyawali et al., 2009; Lack and Cappa, 2010; Virkkula, 2021). Core-shell simulations of size 4 distributions of BC particles coated with a light-scattering shell show that $\alpha_{ap} > -1.4$ for the wavelength pair of 5 470/950 nm could be obtained for BC particle size distributions when the shell volume fraction is > ~89 - 90% 6 and the geometric mean diameter of the BC particles is in the range of ~70 - 100 nm (Virkkula, 2021). Higher 7 α_{ap} would be obtained also by coating with a light-absorbing shell such as brown carbon. In the present work 8 such α_{ap} values were obtained for $\alpha_{ap}(\sigma_{sp}(DMPS,PM_{10}),V2010)$ for the wavelength pair 467/660 nm. So, if these 9 values are closer to the truth it seems that the BC particles that are observed at Dome C are thickly coated and 10 their dominant particle size is < ~100 nm. On the other hand, if the average $\alpha_{ap} \approx 0.8$ obtained for $\alpha_{ap}(\sigma_{ap,nsc})$ 11 and $\alpha_{ap}(\sigma_{sp}(DMPS, PM_{10}), B1999)$ is closer to the truth, the core-shell simulation of Virkkula (2021) suggests that 12 BC particle size distributions would dominated by thinly-coated particles in the size range > 100 nm.

13

14 3.3 Seasonal cycles of mass concentrations, eBC mass concentrations and mass fractions

15 The seasonal cycles of the mass concentrations m(DMPS,PM_{0.8}) and m(DMPS,PM₁₀), see section 2.3.2, the 16 mass concentrations of the PM1 and PM10 filter samples, the mass fraction of the sum of secondary sulfur ions, 17 the eBC mass concentrations and the eBC mass fractions feBC(m(DMPS,PM0.8)) and feBC(m(DMPS,PM10)) (Eqs. 18 16 and 17) presented in Fig. 11 and in Tables S1, S7, S8, and S9. Some corresponding published Antarctic data 19 are also plotted in Fig. 11 for comparison. The m(DMPS,PM_{0.8}) and the m(PM₁) are consistent with each other 20 in that the minimum median mass concentrations are observed in May and June and maximum medians in 21 February. This cycle is very similar to that observed at the Norwegian station Troll in 2007-2011 (Fiebig et al., 22 2014). The monthly average volume concentrations of particles in the size range 33 – 830 nm in Fig. 9 of Fiebig 23 et al. (2014) were digitized and multiplied by the same particle density $\rho = 1.7$ g cm⁻³ that was used for the 24 Dome C data and plotted in Fig 11a. The average (± standard deviation) of the ratio m(DMPS,PM0.8,Dome 25 C)/m(DMPS,Troll) of the monthly averages is \sim 0.6 ± 0.2, i.e., about 40% lower at Dome C. Fiebig et al. (2014) 26 reasoned that the seasonal cycle of particles in the size range measured by the DMPS, i.e., m(DMPS) is 27 controlled by photo-oxidation-limited aerosol formation. This is obviously true for Dome C also. In February, 28 when the maximum monthly average PM_1 and PM_{10} concentrations were observed also the contribution of the 29 sum of secondary sulfur ions (nss SO_4^{2-} + MSA⁻) was the highest (Fig. 11b): the average (± standard deviation) 30 contributions to sum of ions in the PM₁ and PM₁₀ filters was then \sim 81 ± 12% and \sim 61 ± 23%, respectively. The 31 concentrations and the contributions of nss SO₄²⁻ + MSA⁻ were the lowest in July, ~9 ±5 % and ~ 5 ± 5% for PM₁ 32 and PM₁₀, respectively.

2 The seasonal cycle of larger particles ($m(PM_{10})$ and $m(DMPS,PM_{10})$) is much weaker (Fig. 11a) than the $m(PM_1)$ 3 and m(DMPS,PM_{0.8}) cycle. The explanation is that the contribution of sea salt to aerosol mass is the highest in 4 winter (Fig. 11b) and that a large fraction of sea-salt particles is in the supermicron size range, in line with 5 other studies of aerosols at Dome C (e.g., Jourdain et al., 20098; Udisti et al., 2012; Legrand et al., 2017a, 6 2017b). Note that the seasonal cycle of the mass fraction of secondary sulfur ions is qualitatively similar to the 7 seasonal cycle of the scattering Ångström exponent α_{sp} (Fig. 10a): both have the highest values in the austral 8 summer and the lowest values in the austral winter. This is especially clear for the PM₁₀ filters. The small insert 9 in Fig. 11b shows the scatter plot of the monthly average α_{sp} (DMPS,PM₁₀) vs. (nss SO₄²⁻ + MSA⁻)/PM₁₀. The 10 relationship is essentially linear and the correlation coefficient is high, $r^2 = 0.93$. Since the usual interpretation 11 of the size dependence of α_{sp} is that it is inversely proportional to dominating particle size it indicates that 12 when the mass fraction of secondary aerosol is the highest the dominating particle size is the smallest. As such 13 this is not a surprising observation but it is an additional piece of information that links the chemical 14 composition and aerosol optical properties.

The estimated m(PM₁₀) values are consistent with the concentrations measured gravimetrically by Annibaldi et al. (2011) in December 2005 – January 2006. The average PM₁₀ mass concentration they obtained was 134 ± 12 ng m⁻³ at p = 1013 hPa and T = 298 K which equals 146 ± 12 ng m⁻³ at p = 1013 hPa and T = 273 K used in the present paper. The average (and median) PM₁₀ mass concentrations in the present work were 167 ng m⁻³ (140 ng m⁻³) and 167 ng m⁻³ (143 ng m⁻³) in December and January, respectively (Table S1), in a good agreement with the gravimetric measurement of Annibaldi et al. (2011) even though their measurements were not conducted in the same period as ours.

23

15

1

24 In the austral summer the mass concentration calculated from the size distributions (m(DMPS,PM0.8) and 25 m(DMPS,PM₁₀)) were ~100 ng m⁻³ higher than the sums of ions in the PM₁ and PM₁₀ filters (Table S1). Part of 26 the explanation could in principle be that the density 1.7 g cm⁻³ used for calculating mass concentrations from 27 the size distibutions was too high but it cannot explain all of it. Another possible explanation is that there were 28 organic compounds not observed with ion chromatography. Caiazzo et al. (2021) took filter samples at Dome C 29 in a different period, December 2016 - January 2018, analyzed them for organic and elemental carbon with an 30 OC/EC analyzer. The average OC concentration was 86 ± 29 ng m⁻³, approximately the concentration difference 31 between the size distribution-derived and the sums of ions in the filter samples.

32

The eBC concentrations eBC($\sigma_{ap,nsc}$) and eBC(σ_{sp} (DMPS,PM₁₀)) were calculated from Eq. (15). For the scattering-1 2 corrected σ_{ap} the two algorithms, Eq. (11) ad (12) yielded essentially the same absorption coefficients at $\lambda =$ 3 530 nm. Therefore only one of them is shown in the seasonal cycle plot in Fig 11c but both are presented in the 4 supplement Table S7. On the other hand, eBC(Gap,nsc) is also plotted to show how much the scattering 5 correction affects the calculated eBC concentrations in different seasons. For comparison, published monthly 6 median eBC seasonal cycles at three other Antarctic sites are plotted in Fig 11c: at Neumayer, a coastal site in 7 Queen Maud Land, using two two methods, an Aethalometer and a MAAP (Weller et al., 2013), at Syowa, 8 another Queen Maud Land coastal site using an Aethalometer (Hara et al., 2019), and at SPO using an 9 Aethalometer (Sheridan et al., 2016). The maximum median eBC concentrations are observed in October-10 November at all sites. The maximum eBC in October-November is $\sim 3 \pm 1$ ng m⁻³, quite similar at all sites. For 11 eBC +it appears that there is no significant difference between the coastal and plateau sites. The highest 12 monthly median eBC concentrations are those measured with the MAAP at Neumayer in October but, for the 13 same month, the median Aethalometer-derived eBC at Neumayer is the lowest. The lowest monthly median 14 eBC concentrations are observed in April-May at Neumayer, SPO and Dome C and three months earlier in 15 February at Syowa. The lowest monthly medians, ~0.2 ng m⁻³ and ~0.3 ng m⁻³ were observed at Dome C and 16 SPO in May, respectively. The minima were higher at the coastal sites. Note, however, that the eBC 17 concentrations measured with the Aethalometer in Fig. 11bc were not corrected for scattering. This correction 18 was done only for the PSAP data from Dome C and automatically for the MAAP data from Neumayer. After the 19 corrections the Dome C monthly median eBC(σ_{sp} (DMPS,PM₁₀)) ranged from ~0.2 in May to ~3 ng m⁻³ in 20 October-November, i.e., approximately by an order of magnitude and approximately the same as at SPO. The 21 range is smaller at the coastal sites. This might be due to not correcting for the scattering artifact even though 22 the range of MAAP-derived eBC concentrations at Neumayer is also smaller than on the plateau sites. 23

24 The seasonal cycle of eBC is somewhat different from that of the mass concentration. Consequently, the 25 minimum eBC mass fractions in both size ranges (feBC(m(DMPS,PM_{0.8}) and feBC(m(DMPS,PM₁₀)), Eqs. (16) and 26 (17), were in February-March and the maxima in August-October (Fig. 11d, Tables S8 and S9). The eBC mass 27 fractions during this peak were actually quite high. Especially, if it is assumed that all eBC is in the size range 28 measured with the DMPS even for the scattering-corrected eBC monthly medians and averages of feBC varied 29 around 4-5% and the 75th percentiles around 6-7% by using both algorithms (Table S8). These are BC mass fractions typically observed in urban locations (e.g., Liu et al., 2014; Shen et al., 2018), in airborne 30 31 measurements over Europe (McMeeking et al., 2010) and in biomass burning plumes (Pratt et al., 2011), 32 suggesting that in these periods a large fraction of aerosol was long-range transported aerosol from other

1 continents or highly processed air with larger more scattering aerosol preferentially removed. The highest eBC 2 monthly average and median mass concentrations were observed in November but then feBC was lower than 3 its maximum. This can be explained by the increase of the amount of new, non-absorbing natural secondary 4 particles and condensational growth of BC cores by compounds originating from the sea austral during spring 5 and summer. Järvinen et al. (2013) classified new particle formation (NPF) events observed at Dome C and the 6 highest fraction of new particle formation events was in November while in austral spring the particle growth 7 rate was also the highest. The minimum feBC monthly averages were < ~0.5% and medians < ~0.3% in 8 February-March (Tables S8 and S9). This minimum occurs simultaneously also with the minimum eBC 9 concentrations. This suggests that during this time of the year the amount of long-range transported aerosol 10 from other continents is at minimum at the same time when the biogenic aerosol production from the oceans 11 is still high.

12

13 The seasonal cycles of single-scattering albedo (Fig 9) and eBC mass fraction (Fig. 11d) are anticorrelated with 14 each other. It is logical: the lower the feBC is, the higher is the fraction of scattering aerosol and ω_{0} . Their 15 relationships can be used for assessing whether their observed seasonal cycles could be explained by internal 16 mixing of BC particles and scattering components. Linear regressions of monthly average and median ω_0 vs. 17 feBC yield high correlation coefficients but the regression lines would yield negative values at feBC = 100%. So 18 an exponential function of the form of ω_0 (feBC) = ω_0 (feBC=0)exp(-k·feBC) was fitted with the data (Fig. 12). The 19 correlation coefficients were slightly worse, ~0.89 \pm 0.01 for ω_o vs. feBC(m(DMPS,PM_{0.8}) (Fig. 12a) than the 20 ~0.98 ± 0.01 for ω_o vs. feBC(m(DMPS,PM₁₀) (Fig. 12b). If the fitted exponential functions were valid up to feBC = 21 100% the ω_0 (feBC(m(DMPS,PM_{0.8}) would predict that the average $\omega_o \approx 0.2$ and the ω_0 (feBC(m(DMPS,PM_{10})) 22 would predict that ω_o (feBC=100%) \approx 0.06. These are reasonable values for pure BC: it has been measured that 23 for fresh pure BC ω_0 is approximately 0.2 ± 0.1 (e.g., Bond and Bergstrom, 2006; Mikhailov et al, 2006; Bond et 24 al., 2013).

25

To evaluate whether these relationships could be explained by coated spherical BC particles, lognormal size distributions $n(D_p, GMD, GSD)$ were generated where GMD is the geometric mean diameter and GSD the geometric standard deviation and D_p range is 3 nm – 10 μ m. In the simulations all particles had a BC core and a scattering shell and the core volume fraction was the same for all particles in the size distribution. The coreshell model N-Mie (Voshchinnikov and Mathis, 1999) that is based on a recursive algorithm of Wu and Wang (1991) was used for calculating scattering and absorption efficiencies Q_s and Q_a and scattering and absorption coefficients. See Virkkula (2021) for a detailed description of the simulations.

2 The BC core volume fraction varied from 0.25 to 4% to cover the range of feBC(m(DMPS,PM10)) shown in Fig. 3 12b. The simulated ω_0 vs. feBC(PM₁₀) agrees best with the observation-derived values when GSD = 1.8 and 4 GMD = 200 nm. Smaller GMD and GSD yield lower ω_o vs. feBC(PM₁₀). A rigorous error-minimizing procedure 5 was not done since the main goal of this simulation was to show that core-shell geometry can explain the 6 observed ω_o vs. feBC(PM₁₀) relationship. However, a deeper analysis of the modal structure of the size 7 distributions was considered to be out of the scope of the present paper. Note that it was not assumed above 8 that all particles observed at Dome C had a BC core. The simulations did not use any of the measured number 9 concentrations or absorption coefficients. They were pure simulations with generated size distributions and 10 variable BC core and scattering shell volume fractions. They were conducted to study whether the observed ω_{lpha} 11 vs. feBC(PM10) relationship could be explained with size distributions of coated BC particles. However, 12 considering that the BC particles observed at Dome C most probably have been transported from the surrounding continents for thousands of kilometers it is very likely that at most of the observed BC particles 13 14 have been coated during aging processes.

Formatted: Highlight
Formatted: Highlight

Formatted: Highlight
Formatted: Highlight

17 **3.4** Aerosol sources and transport and their seasonal cycles

Source areas were modeled by using FLEXPART as described in section 2.7. The 60000 50-day backtrajectories were used for estimating vertical transport routes and the source areas of the observed aerosols. Sources of individual plumes are not analyzed, the main goal is to find explanations of the observed seasonal cycles of aerosol optical properties.

22

15 16

1

23 3.4.1 Vertical and horizontal transport

24 The average residence time of the trajectories in each grid cell in the altitude range 50 - 14400 m a.g.l. south of 25 the Equator is shown for the inert tracer and for the simulated BC aerosol tracer (Fig. 13). The average altitude 26 of both the inert tracer and the BC tracer decreases sharply as the latitude approaches 75°S which shows that 27 on the average air masses descend over the high plateau. This is in line with several studies that show that 28 stratospheric air is brought down into the lower troposphere by descending air masses above the Antarctic 29 continent (Ito, 1989; James, 1989; Parish and Bromwich, 1991; King and Turner, 1997; Krinner and Genthon, 30 2003; Stohl and Sodemann, 2010). Even though the average altitude of the trajectories shows descending air 31 masses above to Dome C for the inert tracer the low-level residence times are high also at low levels (<1000 m a.g.l.) at latitudes south of about 40°S indicating that it is not uncommon that air masses flow near the surface
 up to Dome C.

3

14

4 When transport was modeled with FLEXPART using in-cloud and below-cloud scavenging of aged BC particles as 5 described by Grythe et al. (2017), for the BC tracer the residence times are lower than for the inert tracer 6 except near Dome C. Additionally, the average trajectory altitude is clearly different than for the inert tracer. 7 Far from Antarctica the BC tracer residence times are clearly smaller than for the inert tracer. This is due to the 8 removal of BC particles by in-cloud and below-cloud scavenging and dry deposition. As a result there would be 9 essentially no BC particles arriving at Dome C from north of latitude 10°S at altitudes < 1600 m which is 10 indicated by the respective area in Fig. 13b just white. Consequently, those BC particles that do get lifted above 11 the clouds increase the average altitude of the BC tracer backtrajectories compared with the inert tracer 12 average altitude as is shown by the respective black and white lines in Fig 13b. At latitudes ~20°S - 50-°S the BC 13 tracer average altitude is approximately 1000 m higher than that of the inert tracer.

15 Between Figs. 13a and 13b there are grey text boxes that show the approximate latitude range of some major 16 BC emitting regions. Indonesia is north of 10°S so BC particles should rise to about 2 km altitude in order to get 17 transported to Dome C. The Indonesian smoke plumes are generally lower than that (Tosca et al., 2011) so its 18 contribution to eBC observed at Dome C is probably negligible. The other three main biomass-burning regions 19 Africa, Australia and Brazil are more to the south and the smoke plume heights have been observed at clearly 20 higher altitudes (Pereira et al., 2016; Rémy Remy et al., 2017; Gonzalez-Alonso et al., 2019). Smoke plumes 21 were recently observed at a height of 4 km over Brazil and even higher, at 6 km over Australia (Shikwambana 22 and Kganyago, 2021). Comparison with Fig. 13b shows that from these altitudes it is likely that BC particles 23 reach also Dome C. All this is in agreement with Fiebig et al. (2009) who established a source-receptor 24 relationship between biomass burning events in Central Brazil and the aerosol observed at the Norwegian 25 station Troll.

26

Near Dome C and over all continental Antarctica the BC tracer average altitude is lower than that of the inert tracer. When BC has a lower average altitude than tracer it means that the air at that point has undergone more removal above than below the average altitude. This suggests that near Dome C most removal takes place in the clouds, higher up than the release point. This implies that there is more in-cloud removal than precipitation scavenging or dry deposition over the plateau.

32

1 3.4.2 Source areas of eBC and high and low ω_o using footprint differences

Source areas were next estimated by using FLEXPART and calculating the footprint differences as described in section 2.7.1. The relative difference (RD) (Eq. 19) of the emission sensitivities in the two lowest layers (> 1000 m a.g.l.) was calculated for the highest and the lowest 10% of eBC concentration and ω_0 . As it was shown in section 3.3 the highest and lowest ω_0 percentiles correspond to the lowest and the highest eBC mass fraction percentiles.

8 The highest eBC concentrations have an increased signal from the surrounding continents Australia, Africa and 9 South America (Fig. 14a). However, high RD values are also associated with large areas over the Pacific Ocean, 10 the Indian Ocean and the Atlantic Ocean. These high RD values on the oceans may in principle be due to 11 shipping emissions but it is more likely that they are due to BC emissions from the continents where the 12 emissions are considerably higher than from shipping. Smoke from South America is predominantly 13 transported southeast into the South Atlantic (Freitas et al., 2005; Reid et al., 2009; Ulke et al., 2011) and 14 further to the east. African smoke gets first transported westward to the Atlantic Ocean by easterly winds (Reid 15 et al., 2009). Most of this smoke wraps around the the South Atlantic Subtropical High, gets mixed with the 16 South American smoke plume and flows east in the subtropical jet region towards Australia and even further to 17 the remote South Pacific (Singh et al., 2000; Staudt et al., 2002; Chatfield et al., 2002). During the transport to higher latitudes the smoke typically rises to about 5 – 7 km (Singh et al., 2000; Staudt et al., 2002; Reid et al., 18 19 2009). 20 21 In addition to the eastward transport in the subtropical jet smoke outflow associated with the trade winds, i.e., 22 the dominating easterlies from South America towards the west to the Pacific Ocean has been observed 23 between Equator and 20°S (Freitas et al., 2005, Ulke et al., 2011; Bourgeois et al., 2015). Bourgeois et al. (2015) 24 studied aerosol transport over the Andes from the Amazon Basin to the remote Pacific Ocean by using six years 25 (2007–2012) of data from the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) satellite instrument. 26 They found that mainly biomass burning particles emitted during the dry season in the Amazon Basin, are lifted 27 in significant amounts over the Andes. Aerosols reaching the top of the Andes, at altitudes typically between 4 28 and 5 km, are entrained into the free troposphere (FT) over the southeast Pacific Ocean and they can be traced 29 on average over 4000 km away from the continent indicating an aerosol residence time of 8–9 days in the FT 30 over the Pacific Ocean.

31

7

Formatted: Highlight

Formatted: Highlight

1	Koch et al. (2009) evaluated BC model predictions of several global models from the AeroCom model
2	intercomparison project. Among several results they presented global maps of modeled BC surface
3	concentrations and annual mean column BC loads predicted by the different models. Fig. 14a displays
4	similarities with both the measurement studies and the global model intercomparison results of Koch et al.
5	(2009). Especially the global maps of annual mean column BC load for GISS sensitivity simulations (Fig. 8 of
6	Koch et al., 2009) have several patterns that are similar to the relative difference RD calculated using Eq. (19)
7	for eBC concentrations measured at Dome C. Here the features of these two figures are compared qualitatively
8	by moving eastwards starting fom Africa. The RD in Fig 14a is high above Africa and on the Atlantic Ocean west
9	of Africa, in agreement with the column BC load predicted by all the GISS sensitivity simulations. To the east,
10	between Africa and Australia there is a belt of high RD values, similar to most of the GISS sensitivity
11	simulations. Interesting is that there is even a local minimum over the Indian Ocean in both the RD values and
12	the GISS simulations. The values over Australia are also high in both figures. There is a clear difference east of
13	Australia: the RD values are high even at the meridian of about 160°W – 180°W although the BC load in most
14	of the GISS simulations decreases clearly immediately to the east of Australia. The RD values then decrease to
15	another local minimum at the meridian of about 150°W and then increase towards South America, in
16	agreement with the GISS sensitivity simulations. Over South America the RD values are positive but decrease to
17	negative values on the South Atlantic immediately to the east of the continent. Such a decrease is not in the
18	GISS sensitivity simulations. So the RD values over the western half of the South Atlantic are the most
19	suspicious ones in Fig. 14a. They suggest that the lowest eBC concentrations observed at Dome C were
20	associated with air masses from there. This is not reasonable since the studies cited above show that South
21	American smoke plumes generally flow to the South Atlantic. This discrepancy remains unexplained.
22	
23	The highest ω_o are obviously related to sea-spray aerosol (SSA) and biogenic secondary aerosol emissions from
24	the Southern Ocean (Fig. 14b). The differences between the oceanic sectors are not big but there some: in the
25	Atlantic and Pacific sectors the RD of ω_p is clearly positive south of latitude 30°S but in the Indian Ocean sector
26	RD is positive south of latitude ~40°S. This can be compared with the Chlorophyll-a (Chl-a) concentrations in
27	five oceanic sectors presented by Becagli et al. (2021). They determined seasonal cycle of Chl-a concentrations
28	in five sectors: Weddell Sea (WS, 60° W–20° E), Indian Ocean (IO, 20–90° E), Western Pacific Ocean (WP, 90–
29	160° E), Ross Sea (RS, 160° E–130° W), and the combined Bellingshausen and Amundsen Seas (BA, 130–60°W).
30	They found that the highest Chl-a concentrations were in the WS, RS and BA sectors and the lowest in the IO
31	and WP sectors. This is in line with the above-described geographical distribution of RD of ω_{o} in Fig. 14b
32	suggesting that the differences in the RD distributions are due to biogenic aerosols.

-	Formatted: Highlight
-1	Formatted: Highlight
-	Formatted: Highlight
1	Formatted: Highlight
Ϊ	Formatted: Highlight
-	Formatted: Highlight

1 2 3 3.4.3 Seasonal cycles of air mass transport from different regions 4 Seasonal cycles of meteorological processes and conditions also affect properties of aerosols observed at Dome 5 C. The most evident is solar radiation that affects new particle formation (e.g., Ito, 1989; Fiebig et al., 2014; 6 Järvinen et al., 2013) but important are also the seasonal cycles of transport from the surrounding continents 7 and from the marine boundary layer of the Southern Ocean (Stohl and Sodemann, 2010). The seasonal 8 contributions of air masses from different regions were calculated as the fraction of time the FLEXPART 9 trajectories spent over Antarctica, the surrounding oceans and continents (Fig. 15). 10 11 In the austral summer (DJF) the contribution of continental air other than Antarctica is low and its minimum is 12 in April. This is true also for marine air masses other than the Southern Ocean. In summer the contribution of 13 Antarctic air is at the highest. Then in May the air mass transport from other continents increases clearly and it 14 reaches a maximum in July, a smaller fraction in August and a second maximum in September. The seasonal 15 cycle of transport from all the surrounding continental areas is qualitatively similar. It can be explained by the 16 seasonal cycle of the tropospheric polar vortex. Waugh et al. (2017) described the structures, seasonality, and 17 dynamics of the stratospheric and tropospheric polar vortices and their connections to extreme events at 18 Earth's surface. Estimating from their Fig. 2, the approximate edge of the tropospheric polar vortex of the 19 Southern Hemisphere is at the latitudes of 50°S in January and 30°S in July. This means that in the austral 20 winter the southern parts of South America, Africa and Australia are within the tropospheric polar vortex but 21 not in the austral summer. This is in line with Fig. 15. The next question is how well do the seasonal cycles of 22 the aerosol optical properties observed at Dome C follow the emissions of major absorbing and scattering 23 aerosols in different regions. 24 25 3.4.4 Relationships of seasonal cycles of BC emissions in the surrounding continents and eBC at Dome C 26 Monthly BC emissions from wildfires and agricultural burning and other fires in South America, Africa and 27 Oceania (Australia, New Zealand, Melanesia, Micronesia and Polynesia) in 2006-2012 were downloaded from 28 the Global Fire Emissions Database (GFED, V3.1). The seasonal cycles are presented in Fig. 16a together with

the monthly average eBC concentrations – corrected for scattering – observed at Dome C. It is obvious in Fig 16a that the maximum of the sum of the emissions occurs three months earlier than the maximum eBC concentration at Dome C. Monthly BC emissions from wildfires and agricultural burning and other fires in South America, Africa and Oceania (Australia, New Zealand, Melanesia, Micronesia and Polynesia) in 2006-2012 were Formatted: Highlight Formatted: Highlight Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

32

1	downloaded from the Global Fire Emissions Database (GFED, V3.1). The seasonal cycles are presented in Fig.
2	16a together with the monthly average eBC concentrations – corrected for scattering – observed at Dome C. It
3	is obvious in Fig 16a that the maximum of the sum of the emissions occurs three months earlier than the
4	maximum eBC concentration at Dome C.

6 The seasonal cycles of air mass residence time over each of the major source areas plays a crucial role in 7 explaining the relationship between eBC at DomeC and BC emissions. The BC emissions were multiplied with 8 the fraction of time (=f(t)) the FLEXPART trajectories spent over each of the three major source areas, shown in 9 Fig. 15. These quantities (f(t)×BC emissions) will be called residence-time-weighted BC emissions, RTW BC 10 emissions. They are plotted in Fig. 16b together with the normalized eBC concentrations. Linear regressions 11 between the eBC concentration and RTW BC emissions were calculated but only the respective squared 12 correlation coefficients R^2 are shown. The correlation between eBC concentrations and RTW BC emissions is 13 weak, $R^2 < 0.3$ (Fig. 16b). But the main reason is that the residence times shown in Fig. 15 present the fraction 14 of time that trajectories arriving at Dome C at any given month have spent over each of the regions before 15 arrival at Dome C. So next it was assumed that the transport time from each of the source areas to Dome C is 16 one month so f(t) of each month was multiplied with the BC emission one month earlier. This shifts the RTW BC 17 emission peaks of all the source areas so that all R² increase somewhat (Fig. 16c). When it is assumed that the 18 transport time is 2 months R² increase clearly, 0.688 for Oceania and 0.665 for South America, for Africa 19 correlation still remains low (Fig. 16d). In October-November, when the eBC concentrations at Dome C are the 20 largest the 2-month-shifted RTW BC emissions from South America are approximately an order of magnitude 21 larger than from Africa and Oceania suggesting that South American BC emissions are the largest contributor to 22 eBC at Dome C.

23

5

	24	The highest correlation between the eBC concentrations and RTW BC emissions with the 2-month time shift
	25	suggests further that transport time is considerably longer than the 30 days Stohl and Sodemann (2010)
I	26	obtained in a 5.5-year climatology of atmospheric transport into the Antarctic troposphere. The two-month
	27	transport time naturally implies that BC aerosol residence time is also so long. Williams et al. (2002) estimated
	28	by modeling that in the upper atmosphere aerosols of 0.065 μm in size have residence times of approximately
	29	1 month and can be transported on a hemispheric scale. Papastefanou (2006) determined residence times of
	30	tropospheric aerosols with different methods and obtained a much shorter mean value of 8 days. Kristiansen et
	31	al. (2012) estimated accumulation-mode aerosol removal times from the atmosphere using a global
	32	measurement data set collected over several months after the accident at the Fukushima Dai-ichi nuclear
_		

1	power plant in March 2011. After correction for redisactive decay, the ¹³⁷ Cr. ⁽¹³³ Ve ratios can be accurated to		
2	power plant in March 2011. After correction for radioactive decay, the CSY "Xe ratios can be assumed to		
2	removal times for 137 Cs of 10.0–12.9 days. The simulations of Croft et al. (2014) with instantaneous injections		
1	inte layers at 5 km and 7 km violded mean lifetimes of 14.7 days and 21.1 days respectively. At higher altitudes		
4	into layers at 5 km and 7 km yielded mean incumes of 14.7 days and 21.1 days, respectively. At higher attitudes		
5	removal mechanisms are weaker so lifetimes are longer. These are altitudes where blomass-burning smoke		
6	plumes have been observed, as discussed above. Note that these lifetimes are the e-folding lifetimes t from		
/	$C(t) = C(t_0)exp(-t/\tau)$. That means, for example, that if biomass burning smoke where BC concentration is 1000		
8	ng m ⁻³ – similar to concentrations observed in a Brazilian biomass burning measurement campaign (Brito et al.,		Formatted: Highlight
9	2014) – rises to 5 km and gets transported for 60 days the BC concentration decreases to 16.9 ng m ⁻³ . With τ =		
10	10 d the concentration would decrease to 2.5 ng m ⁻³ . At Dome C the 25 th to 75 th percentile range of eBC		
11	concentration in November is $\sim 2 - 6$ ng m ⁻³ (Table S7). As the simple estimation shows this range is consistent		
12	with BC emissions of about 1 µg m ⁻³ and transport for two months.		
13			
14	The above discussion does not mean that pollution episodes from the surrounding continents could not be		Formatted: Highlight
15	transported faster than in two months. For instance, Fiebig et al. (2009) calculated the source areas and		
16	transport routes of an elevated aerosol concentration event observed at the Norwegian Troll station in Queen		
17	Maud Land. The analysis showed that the origin of the aerosol was biomass burning in Brazil 11 – 12 days		Formatted: Highlight
18	before the arrival at Troll. Hara et al. (2010) analyzed several haze events observed at Syowa, <u>the Japanese</u>		Formatted: Highlight
19	Antarctic coastal station. They found that the traveling time from Southern Africa to Syowa varied in the range		Formatted: Highlight
20	of 4 to 8 days, even faster than to Troll in the episode analyzed by Fiebig et al. (2009). The explanation for the	_	Formatted: Highlight
21	considerably longer transport time to Dome C, 2 months, is that it was obtained from the correlation of		
22	seasonal cycles of BC emissions and observed concentrations so it is more like a statistical estimate of seasonal	_	Formatted: Highlight
23	transport times. That suggests that on the average the BC emitted in the biomass burning in the surrounding		
24	continents circulates the Southern Hemisphere and migrates slowly towards the Antarctic upper plateau.		
25	۸		Formatted: English (United States)
26	Fig. 16 also shows that the observed concentration varies by an order of magnitude only although the		
27	emissions vary by two orders of magnitude. This suggests that at the observed seasonal minimum eBC in		
28	March-April the sources are something else than wildfires and agricultural burning BC emissions in the GFED		
29	data. Possible sources can be other anthropogenic emissions such as traffic including shipping, industrial		
30	emissions and heating. <u>Stohl and Sodemann (2010) estimated that</u>	_	Formatted: Highlight
31	ship emissions south of 60°S account for half of the total BC concentrations in the lowest 1000 m of the		
32	atmosphere south of 70°S in December. However, due to the particle removal processes as described in section		Formatted: Highlight
I			

1	3.4.1 the contribution of ship emissions to BC at Dome C much lower. No quantitative estimations were	
2	calculated. Xu et al. (2021) recently presented a new global BC emission inventory for all countries in the world	
3	for the years 1960 – 2017. In an Excel file supplement Xu et al. (2021) presents the emissions classified into	
4	energy sector, industrial, residential, on-road motor vehicles, other anthropogenic and wildfires. The sums of	
5	South American, Southern hemispheric African and Oceanian emissions in the period 2008 – 2013 and the	
6	respective contributions of wildfire emissions were calculated. They were ~40 \pm 8 %, ~49 \pm 2 % and ~59 \pm 9 %	
7	for South American, Southern hemispheric African and Oceanian emissions, respectively. However, only	
8	wildfire-emitted BC-particles have the potential to rise high enough to avoid in-cloud or below-cloud	
9	scavenging during the transport, as was discussed above.	
10		
11	3.4.5 Relationships of seasonal cycles of scattering aerosol emissions and σ_{sp} at Dome C and SPO	
12	Ŧ	
13	The seasonal cycles of scattering coefficient at Dome C and at the SPO were compared with the seasonal cycles	
14	of secondary and primary marine aerosol emissions. The sea-spray aerosol (SSA) emissions calculated with the	
15	FLEX-SSA offline tool (section 2.7.2) show that SSA emissions from the Southern Ocean peak in the austral	
16	winter (Fig. 17a). Sea ice and the glacier surfaces especially in the coastal regions are also important sources of	-(
17	and rate approaches (and a final a 2020). Here at al. 2020). Hereaver, the size was to find a qualitative	
1/	sea-sait aerosois (e.g., Frey et al., 2020; Hara et al., 2020). However, the aim was to find a qualitative	
17	explanation of the high scattering coefficients observed in winter especially at SPO. The FLEX-SSA offline tool	
17 18 19	explanation of the high scattering coefficients observed in winter especially at SPO. The FLEX-SSA offline tool was used for that and it yields a seasonal cycle that at at least partially explains the observations. There was no	
17 18 19 20	sea-sait derosols (e.g., Frey et al., 2020; Hara et al., 2020). However, the aim was to find a qualitative explanation of the high scattering coefficients observed in winter especially at SPO. The FLEX-SSA offline tool was used for that and it yields a seasonal cycle that at at least partially explains the observations. There was no tool for estimating seasonal cycles of sea-salt emissions from sea ice.	(
17 18 19 20 21	explanation of the high scattering coefficients observed in winter especially at SPO. The FLEX-SSA offline tool was used for that and it yields a seasonal cycle that at at least partially explains the observations. There was no tool for estimating seasonal cycles of sea-salt emissions from sea ice. -Secondary marine aerosols are formed from the oxidation of emission products of phytoplankton. Becagli et	
17 18 19 20 21 22	explanation of the high scattering coefficients observed in winter especially at SPO. The FLEX-SSA offline tool was used for that and it yields a seasonal cycle that at at least partially explains the observations. There was no tool for estimating seasonal cycles of sea-salt emissions from sea ice. -Secondary marine aerosols are formed from the oxidation of emission products of phytoplankton. <u>Becagli et</u> al. (2021) discussed the relationships between biogenic aerosols (nss SO4 ² , and MSA;), Chlorophyll-a	
17 18 19 20 21 22 23	explanation of the high scattering coefficients observed in winter especially at SPO. The FLEX-SSA offline tool was used for that and it yields a seasonal cycle that at at least partially explains the observations. There was no tool for estimating seasonal cycles of sea-salt emissions from sea ice. -Secondary marine aerosols are formed from the oxidation of emission products of phytoplankton. Becagli et al. (2021) discussed the relationships between biogenic aerosols (nss SO4 ² , and MSA;), Chlorophyll-a concentration which is a proxy for phytoplankton, sea ice extent and area, source areas and the transport	
17 18 19 20 21 22 23 24	 sea-sait derosols (e.g., Frey et al., 2020; Hara et al., 2020). However, the aim was to find a qualitative explanation of the high scattering coefficients observed in winter especially at SPO. The FLEX-SSA offline tool was used for that and it yields a seasonal cycle that at at least partially explains the observations. There was no tool for estimating seasonal cycles of sea-salt emissions from sea ice. Secondary marine aerosols are formed from the oxidation of emission products of phytoplankton. <u>Becagli et al.</u> (2021) discussed the relationships between biogenic aerosols (nss SO4², and MSA'), Chlorophyll-a concentration which is a proxy for phytoplankton, sea ice extent and area, source areas and the transport routes to Dome C and the Southern Annular Mode (SAM). They showed that these relationships are different in 	
17 18 19 20 21 22 23 24 25	explanation of the high scattering coefficients observed in winter especially at SPO. The FLEX-SSA offline tool was used for that and it yields a seasonal cycle that at at least partially explains the observations. There was no tool for estimating seasonal cycles of sea-salt emissions from sea ice. -Secondary marine aerosols are formed from the oxidation of emission products of phytoplankton. Becagli et al. (2021) discussed the relationships between biogenic aerosols (nss SO ₄ ²⁺ , and MSA ²), Chlorophyll-a concentration which is a proxy for phytoplankton, sea ice extent and area, source areas and the transport routes to Dome C and the Southern Annular Mode (SAM). They showed that these relationships are different in early, middle, and late austral summer. Here a subset of the data is analyzed in a method that is	
17 18 19 20 21 22 23 24 25 26	 sea-sait derosols (e.g., Frey et al., 2020; Hara et al., 2020). However, the aim was to hind a qualitative explanation of the high scattering coefficients observed in winter especially at SPO. The FLEX-SSA offline tool was used for that and it yields a seasonal cycle that at at least partially explains the observations. There was no tool for estimating seasonal cycles of sea-salt emissions from sea ice. Secondary marine aerosols are formed from the oxidation of emission products of phytoplankton. Becagli et al. (2021) discussed the relationships between biogenic aerosols (nss SO₄², and MSA;). Chlorophyll-a concentration which is a proxy for phytoplankton, sea ice extent and area, source areas and the transport routes to Dome C and the Southern Annular Mode (SAM). They showed that these relationships are different in early, middle, and late austral summer. Here a subset of the data is analyzed in a method that is complementary to that of Becagli et al. (2021). The phytoplankton concentration maximum is in summer and 	
17 18 19 20 21 22 23 24 25 26 27	 sea-sait derosols (e.g., Frey et al., 2020; Hara et al., 2020). However, the aim was to find a qualitative explanation of the high scattering coefficients observed in winter especially at SPO. The FLEX-SSA offline tool was used for that and it yields a seasonal cycle that at at least partially explains the observations. There was no tool for estimating seasonal cycles of sea-salt emissions from sea ice. Secondary marine aerosols are formed from the oxidation of emission products of phytoplankton. Becagli et al. (2021) discussed the relationships between biogenic aerosols (nss SO₄²⁺, and MSA²), Chlorophyll-a concentration which is a proxy for phytoplankton, sea ice extent and area, source areas and the transport routes to Dome C and the Southern Annular Mode (SAM). They showed that these relationships are different in early, middle, and late austral summer. Here a subset of the data is analyzed in a method that is complementary to that of Becagli et al. (2021). The phytoplankton concentration maximum is in summer and minimum in winter in the satellite data analysis of Behrenfeld et al. (2016) (Fig. 17a). However, phytoplankton 	
 17 18 19 20 21 22 23 24 25 26 27 28 	sea-sait derosols (e.g., Frey et al., 2020; Hara et al., 2020). However, the aim was to find a qualitative explanation of the high scattering coefficients observed in winter especially at SPO. The FLEX-SSA offline tool was used for that and it yields a seasonal cycle that at at least partially explains the observations. There was no tool for estimating seasonal cycles of sea-salt emissions from sea ice. -Secondary marine aerosols are formed from the oxidation of emission products of phytoplankton. Becagli et al. (2021) discussed the relationships between biogenic aerosols (nss SO_4^2 , and MSA). Chlorophyll-a concentration which is a proxy for phytoplankton, sea ice extent and area, source areas and the transport routes to Dome C and the Southern Annular Mode (SAM). They showed that these relationships are different in early, middle, and late austral summer. Here a subset of the data is analyzed in a method that is complementary to that of Becagli et al. (2021). The phytoplankton concentration maximum is in summer and minimum in winter in the satellite data analysis of Behrenfeld et al. (2016) (Fig. 17a). However, phytoplankton concentration alone does not explain the seasonal cycle of secondary aerosol scattering coefficient. The DMS	
17 18 19 20 21 22 23 24 25 26 27 28 29	 sea-sait derosols (e.g., Frey et al., 2020; Hara et al., 2020). However, the aim was to find a qualitative explanation of the high scattering coefficients observed in winter especially at SPO. The FLEX-SSA offline tool was used for that and it yields a seasonal cycle that at at least partially explains the observations. There was no tool for estimating seasonal cycles of gea-salt emissions from sea ice. Secondary marine aerosols are formed from the oxidation of emission products of phytoplankton. Becagli et al. (2021) discussed the relationships between biogenic aerosols (nss SO4², and MSA), Chlorophyll-a concentration which is a proxy for phytoplankton, sea ice extent and area, source areas and the transport routes to Dome C and the Southern Annular Mode (SAM). They showed that these relationships are different in early, middle, and late austral summer. Here a subset of the data is analyzed in a method that is complementary to that of Becagli et al. (2021). The phytoplankton concentration maximum is in summer and minimum in winter in the satellite data analysis of Behrenfeld et al. (2016) (Fig. 17a). However, phytoplankton concentration alone does not explain the seasonal cycle of secondary aerosol scattering coefficient. The DMS oxidation is a photochemical process so a simplified method was used for estimating solar radiation intensity. 	
 17 18 19 20 21 22 23 24 25 26 27 28 29 30 	 sea-sait derosols (e.g., Frey et al., 2020; Hara et al., 2020). However, the aim was to find a qualitative explanation of the high scattering coefficients observed in winter especially at SPO. The FLEX-SSA offline tool was used for that and it yields a seasonal cycle that at at least partially explains the observations. There was no tool for estimating seasonal cycles of sea-salt emissions from sea ice. -Secondary marine aerosols are formed from the oxidation of emission products of phytoplankton. Becagli et al. (2021) discussed the relationships between biogenic aerosols (nss SO₄², and MSA²). Chlorophyll-a concentration which is a proxy for phytoplankton, sea ice extent and area, source areas and the transport routes to Dome C and the Southern Annular Mode (SAM). They showed that these relationships are different in early, middle, and late austral summer. Here a subset of the data is analyzed in a method that is complementary to that of Becagli et al. (2021). The phytoplankton concentration maximum is in summer and minimum in winter in the satellite data analysis of Behrenfeld et al. (2016) (Fig. 17a). However, phytoplankton concentration is a photochemical process so a simplified method was used for estimating solar radiation intensity. Global radiation intensity Ir (W m⁻²) was calculated by using a clear sky model for direct and diffuse insolation 	
 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 	 Sea-salt aerosols (e.g., Frey et al., 2020; Hara et al., 2020). However, the aim was to find a dualitative explanation of the high scattering coefficients observed in winter especially at SPO. The FLEX-SSA offline tool was used for that and it yields a seasonal cycle that at at least partially explains the observations. There was no tool for estimating seasonal cycles of sea-salt emissions from sea ice. Secondary marine aerosols are formed from the oxidation of emission products of phytoplankton. Becagli et al. (2021) discussed the relationships between biogenic aerosols (nss SO₄²) and MSA'). Chlorophyll-a concentration which is a proxy for phytoplankton, sea ice extent and area, source areas and the transport routes to Dome C and the Southern Annular Mode (SAM). They showed that these relationships are different in early, middle, and late austral summer. Here a subset of the data is analyzed in a method that is complementary to that of Becagli et al. (2021). The phytoplankton concentration maximum is in summer and minimum in winter in the satellite data analysis of Behrenfeld et al. (2016) (Fig. 17a). However, phytoplankton concentration alone does not explain the seasonal cycle of secondary aerosol scattering coefficient. The DMS oxidation is a photochemical process so a simplified method was used for estimating solar radiation intensity. Global radiation intensity I_T (W m⁻²) was calculated by using a clear sky model for direct and diffuse insolation (Bird and Hulstrom, 1981). I_T was calculated at the surface (p =1000 mbar) for each hour of the year at the 	

Formatted: Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

-	Formatted: Highlight
1	Formatted: Highlight
1	Formatted: Font: +Body (Calibri), 11 pt, Highlight
-	Formatted: Highlight
	Formatted: Font: (Default) +Body (Calibri), 11 pt, Highlight
Ù	Formatted: Highlight
$\langle \rangle$	Formatted: Font: +Body (Calibri), 11 pt, Highlight
Ì	Formatted: Font: +Body (Calibri), 11 pt, Highlight
Υ	Formatted: Font: +Body (Calibri), 11 pt. Highlight

1 (45°-55°S), zone 2 (55°-65°S), and the polar zone, zone 3 (65°-75°S). The monthly averages of I_T at 10:00 – 2 14:00 in each day of a month were calculated. The reasoning for using this time is that new particle formation 3 is typically, although not exclusively, a daytime phenomenon at numerous locations around the world, also in 4 Antarctica (e.g., Weller et al., 2015; Kerminen et al., 2018; Kim et al., 2019; Brean et al, 2021). The monthly 5 product C_{phytolT}, was then used as a proxy of biogenic secondary aerosol emissions. Behrenfeld et al. (2016) also 6 presented the photosynthetically active radiation (PAR) in the polar zone obtained from the MODIS Aqua 7 ocean colour sensor acquired from http://oceancolor.gsfc.nasa.gov/cms. In addition to the product C_{phytol}T, also 8 the monthly product C_{phyto}PAR was calculated for the zone 3 and used as a proxy for biogenic secondary aerosol 9 emissions. In Zone 3 the two emission proxies C_{phyto}I_T and C_{phyto}PAR agree well (Fig. 17a).

10

27

11 The normalized seasonal cycles of the SSA flux and the secondary aerosol emission proxy C_{phyto}PAR in the polar 12 zone are plotted together with the normalized seasonal cycles of the scattering coefficients at Dome C (Fig. 13 17b) and at SPO (Fig. 17c). At Dome C both σ_{sp} (DMPS,PM_{0.8}) and σ_{sp} (DMPS,PM₁₀) approximately follow the 14 seasonal cycle of the secondary aerosol emissions in October - April. In the austral winter the effect of SSA is 15 visible especially in the high monthly σ_{sp} averages. The medians are cleary lower, they follow the secondary 16 aerosol emission proxy even in May. Also the normalized seasonal cycle of $\sigma_{so}(nss-SO_4^{22},PM_1)$, the scattering 17 coefficient calculated from non-seasalt sulfate concentrations in PM₁ filter samples is presented in Fig. 17b. It 18 also follows approximately the seasonal cycle of C_{phyto} PAR for the whole year. However, there appears to be a 19 time lag. $C_{phyto}PAR$ grows fast in October but $\sigma_{sp}(nss-SO_4^{-2};PM_1)$ in November. When $C_{phyto}PAR$ is shifted by one 20 month they agree clearly better. The linear regression of of $\sigma_{sp}(nssSO_4^{2-}, PM_1)$ vs. C_{phyto} PAR yields the correlation 21 coefficient $R^2 = 0.65$ when there is no C_{phyto} PAR time shift and $R^2 = 0.92$ when C_{phyto} PAR is shifted by one month 22 (Fig. 18). This suggests that on the average it takes approximately a month for the secondary aerosol to be 23 formed, grown, get mixed in the upper atmospheric layers and be transported to the upper plateau. As it was 24 discussed above, Williams et al. (2002) estimated by modeling that in the upper atmosphere aerosol residence 25 time can be \sim 1 month and Croft et al. (2014) obtained e-folding lifetimes would be \sim 15 days and \sim 21 days at 5 26 km and 7 km altitudes, respectively, so it is reasonable.

Formatted: Highlight

Formatted: Highlight
Formatted: Highlight

²⁸ -There is one thing that the time shift cannot explain. The maximum σ_{sp} – and mass concentrations as discussed ²⁹ in Section 3.3 – was in February. This cannot be explained by the seasonal cycle of the biogenic secondary ³⁰ aerosol emission proxy: the maximum C_{phyto}PAR was in December so even a one month transport time does not ³¹ make February the maximum. Above it was assumed that the proxy is simply the product C_{phyto}PAR. However,

32 the underlying hypothesis of this proxy is that the emissions of the precursor gases of new particle formation,
1	mainly DMS, are linearly related to the phytoplankton mass concentrations. This is probably not the case, there
2	are other factors. Another explanation could be meteorological. Becagli et al. (2021) showed that the MSA and
3	nss-SO ₄ ²⁻ maxima in February could possibly be explained by faster transport from the surrounding oceans with
4	air masses traveling at lower elevation than in December-JanuaryFurther analyses of that relationship are out
5	of the scope of the present paper, however. For more details, see Becagli et al. (2021). Note that $\sigma_{sp}(nssSO_4^{2^2},$
6	PM_1) correlates well also with $C_{phyto}I_T$ of zones 1 and 2 shown in Fig. 17a which means the secondary marine
7	aerosol may have been formed and grown also further to the north than in the polar zone.

At SPO σ_{sp} is more strongly affected by the SSA emissions than at Dome C, especially in the austral winter and spring (Fig. 17c). Note that at SPO the seasonal cycle of the normalized median σ_{sp} follows roughly the normalized SSA emission flux in July – December. The seasonal cycle of the normalized average σ_{sp} is different because averages are affected more by individual high transport events and medians represent better the prevailing background aerosol of the season. As explained above, the difference between Dome C and SPO may either be due to geographical locations, different size ranges measured by the instruments or both.

15

16 4. Summary and conclusions

17 Aerosol optical properties have been measured at several Antarctic sites but scattering and absorption data 18 measured at Dome C have not been not examined in detail earlier. This work fills that gap using light 19 absorption from a 3 λ PSAP and light scattering coefficients (σ_{sp}) calculated from particle number size 20 distributions measured with a Differential Mobility Particle Sizer (DMPS) and an optical particle counter (OPC). 21 Additionally, single scattering albedo (ω_0), absorption Ångström exponent (α_{av}), scattering Ångström exponent 22 (α_{sp}) and equivalent black carbon (eBC) concentrations were calculated. The sources of the aerosol were 23 estimated by calculating footprints with FLEXPART and by calculating seasonal cycles of transport of both 24 scattering and absorbing particles from different source areas.

25

Aerosol light scattering coefficient was calculated from the DMPS size distributions using two different refractive indices and the contribution of scattering due to particles larger than those measured with the DMPS to provide a lower and upper estimate of σ_{sp} . Light scattering was also estimated using mass scattering efficiencies in conjunction with mass concentrations obtained from PM₁ and PM₁₀ filter samples. The two most frequently used algorithms to calculate aerosol absorption coefficients (σ_{ap}) from PSAP measurements require scattering coefficients. Both algorithms were applied and both the upper and lower estimate of σ_{sp} were used in order to provide understanding of the differences between the algorithms and the impact of the scattering

Formatted: Highlight

Formatted: Highlight

1 adjustment. The absorption coefficient calculated using the σ_{sp} upper estimate was considered to be the best 2 σ_{ap} estimate.

3

4 There were clear seasonal cycles of σ_{sp} and σ_{ap} at Dome C. The maximum and minimum of σ_{sp} were observed in 5 austral summer and winter, respectively. The Dome C scattering coefficients were also compared with σ_{so} 6 measured with a nephelometer at the South Pole (SPO). At SPO the scattering coefficient was similar to that 7 measured at Dome C in austral summer but there was a large difference in the austral winter. At SPO the 8 maximum monthly averages were observed in austral winter. This suggests that, in winter, SPO is more 9 influenced by sea spray emissions than Dome C. At Dome C the σ_{ab} exhibited a different seasonal cycle of than 10 σ_{sp--} the minimum σ_{ap} was observed in the austral autumn and the maximum in spring. As a result the 11 seasonal cycle of the single-scattering albedo ω_o is such that the darkest aerosol, i.e., the lowest $\omega_o \approx 0.91$ is 12 observed October and the highest ω_{o} > 0.99 in February and March.

13

14 The scattering Ångström exponent α_{sp} calculated from the σ_{ssp} lower estimate varied from ~2.6 in austral 15 summer to ~2.1 in austral winter indicating that in austral summer the size distributions are dominated by 16 smaller particles than in winter. For the σ_{sp} upper estimate α_{sp} varied from ~1.9 in austral summer to ~0.8 in 17 winter. This seasonal cycle is quite similar to the seasonal cycle of α_{sp} of σ_{sp} measured at SPO. The uncertainty 18 of the absorption Ångström exponent α_{ap} is high, particularly in the clean conditions existing in Antarctica. 19 However, despite the high uncertainties the seasonal cycles of α_{ap} , with and without the scattering correction, 20 are qualitatively similar: the lowest monthly medians are observed in March while the maxima occur in August 21 – October. This cycle – even that of the non-scattering corrected α_{ap} – is anticorrelated with the ω_o seasonal 22 cycle.

23

24 The eBC mass concentrations were compared with eBC measured at three other sites: the South Pole and two 25 coastal sites Neumayer and Syowa. The maximum monthly median eBC concentrations are almost the same 26 (~3 ± 1 ng m⁻³) at all these sites in October-November. This suggests that, as far as eBC is concerned, there is no 27 significant difference between the coastal and plateau sites. The seasonal cycle of eBC is slightly different from 28 the mass concentration calculated from the number size distributions measured with the DMPS. Consequently, 29 the seasonal cycle of the eBC mass fraction (f(eBC)) is such that f(eBC) minimum is in February-March and the 30 maximum is in August-October, anticorrelating with ω_o . The eBC mass fractions in this peak are actually quite 31 high and vary around 4-5% with the 75th percentiles > 6-7%, only slightly depending on PSAP correction Formatted: Highlight

algorithm. These levels of eBC mass fractions are typically observed in polluted air, suggesting that in these
 periods a large fraction of aerosol is long-range transported aerosol from other continents.

3

Source areas were calculated with 50-day FLEXPART footprints. The relative differences of the footprints calculated for the highest and lowest 10% of eBC concentrations and ω_o showed that the highest eBC concentrations and the lowest ω_o were associated with air masses coming from South America, Australia and Africa. Vertical simulations that take BC particle removal processes into account show that there would be essentially no BC particles arriving at Dome C from north of latitude 10°S at altitudes < 1600 m. The main biomass-burning regions Africa, Australia and Brazil are more to the south and their smoke plumes have been observed at higher altitudes than that so they can get transported to Antarctica.

11

12 The seasonal cycle of BC emissions from wildfires and agricultural burning and other fires in South America, 13 Africa and Australia were calculated from data downloaded from the Global Fire Emissions Database (GFED). 14 The maximum total emissions were in August-September but the peak of monthly average eBC concentrations 15 is observed November, 2 - 3 months later not only at Dome C but also at the South Pole and the coastal 16 stations. This is considerably longer than the 30 days presented in an earlier study. If this peak eBC 17 concentration is really due to the peak emissions from the above-mentioned fires in the surrounding 18 continents it means that the aerosol from these fires remains in air for several months and gets mixed 19 essentially over the entire Southern Hemisphere. This does not mean that pollution episodes from the 20 surrounding continents could not be transported faster than in two months. Earlier studies have clearly shown 21 that biomass-burning smoke can get transported frm South America or Africa to Antarctica in 1-2 weeks. The 22 explanation for the considerably longer transport time to Dome C, 2 months, is that it was obtained from the 23 correlation of seasonal cycles of BC emissions and observed concentrations so it is more like a statistical 24 estimate of seasonal transport times. That suggests that on the average the BC emitted in the biomass burning 25 in the surrounding continents circulates the Southern Hemisphere and migrates slowly towards the Antarctic 26 upper plateau.

27

The seasonal contributions of air masses from different regions were calculated as the fraction of time the FLEXPART trajectories spent over Antarctica, the surrounding oceans and continents. The BC emissions were multiplied with the fraction of time the trajectories spent over each of the three major source areas. In October-November, when the eBC concentrations at Dome C are the largest the 2-month-shifted residencetime-weighted BC emissions from South America are approximately an order of magnitude larger than from
 Africa and Oceania suggesting that South American BC emissions are the largest contributor to eBC at Dome C.

3

4 The seasonal cycles of scattering coefficient at Dome C and at the SPO were compared with the seasonal cycles 5 of secondary and primary marine aerosol emissions. The seasonal cycles of sea spray aerosol (SSA) emissions 6 were simulated with the FLEX-SSA offline tool. The seasonal cycles of biogenic secondary aerosols were 7 estimated from monthly average phytoplankton biomass concentrations obtained from the CALIOP satellite 8 sensor data. The correlation coefficients between scattering coefficients measured at Dome C and 9 phytoplankton biomass concentrations and a biogenic secondary aerosol emission proxy are high. It may take a 10 month for the biogenic aerosol to be formed and get transported from the sea level to Dome C. The scattering 11 coefficients measured at SPO correlated much better with the SSA emission fluxes in the Southern Ocean than 12 the scattering coefficients measured at Dome C. The difference between the scattering coefficients at these 13 sites may either be due to geographical locations, different size ranges measured by the instruments or both.

14

15 Data Availability

The Dome C aerosol physical measurement data will be made openly available for the final publication at the EBAS database (http://ebas.nilu.no). The SPO data are there already. The Dome C aerosol chemical Data Availability The Dome C aerosol physical measurement data will be made openly available for the final publication at the

20 EBAS database (<u>http://ebas.nilu.no</u>). The SPO data are there already. The Dome C aerosol chemical 21 composition data will be available upon request by writing to RT, BS or MS.

22 23

24 Author contributions

25 AV contributed to designing the measurements of aerosol physical properties at Dome C, processing and 26 analyzing data and writing most of the manuscript with contributions from all coauthors; HG did all the 27 FLEXPART modeling and processed GFED data and wrote the associated text; JB wrote the codes for data processing; TP reviewed and commented the manuscript; MB, CL and AL were responsible for the installation 28 29 and long-term operation of the aerosol instruments at Dome C; SB, RT, and MS were responsible for the 30 sampling and analyses of the PM1 and PM10 filter samples; VV contributed to designing and administration of 31 the measurements at Dome C; PS and EA were responsible for the SPO aerosol measurements. All authors 32 reviewed and commented the manuscript.

3 Competing interests

4 Tuukka Petäjä is editor of ACP.

5

6 Financial support

AV was funded by the Academy of Finland project Antarctic Climate Forcing Aerosol (ACFA) (decision nr. 335845) and by the Business Finland project Black Carbon Footprint (BCF) (grant nr. 528/31/2019). HG was supported by NILU's internal financing for publications (G-106057). Aerosol sampling at Dome C was carried out in the framework of the Air-Glacs1181 'Station Concordia' project, which was funded by the MIUR through the Italian 'Programma Nazionale di Ricerche in Antartide' (PNRA) and in the framework of 'DECA-POL' (2010/A3.05) PNRA project. EA was funded by the NOAA Climate Program Office and PS by NOAA base funds.

13

14 Acknowledgments

The field operations benefitted from the support of the French-Italian Concordia Station. Meteorological data and information were obtained from IPEV/PNRA Project "Routine Meteorological Observation at Station Concordia, <u>www.climantartide.it</u>. The technicians at SPO are acknowledged for operating the instruments and Derek Hageman for data processing. Prof. Michael J. Behrenfeld of Oregon State University is acknowledged for providing the phytoplankton biomass concentrations from the CALIOP satellite sensor and the photosynthetically active radiation data from the MODIS Aqua ocean colour sensor.

1		
2	References	
3	Alt, J., P. Astapenko, and N.J. Ropar, Jr.: Some aspects of the antarctic atmospheric circulation in 1958, IGY Rep.	
4	4, 113 pp., Nat. Acad. of Sci., Washington, D.C., 1959.	
5		
6	Annibaldi, A., Truzzi, C., Illuminati, S., Scarponi, G.: Direct gravimetric determination of aerosol mass	
7	concentration in central Antarctica. Anal. Chem., 83, 143-151, doi:10.1021/AC202408Z, 2011.	
8		
9	Backman, J., Virkkula, A., Vakkari, V., Beukes, J. P., Van Zyl, P. G., Josipovic, M., Piketh, S., Tiitta, P., Chiloane, K.,	
10	Petäjä, T., Kulmala, M., and Laakso, L.: Differences in aerosol absorption Ångström exponents between	
11	correction algorithms for a particle soot absorption photometer measured on the South African Highveld,	
12	Atmos. Meas. Tech., 7, 4285–4298, https://doi.org/10.5194/amt-7-4285-2014, 2014.	
13		
14	Barber, P. W. and Hill, S. C.: Light scattering by particles: Computational methods, World Scientific Publishing,	
15	Singapore, 1990.	
16		
17	Becagli, S., Ghedini, C., Peeters, S., Rottiers, A., Traversi, R., Udisti, R., Chiari, M., Jalba, A., Despiau, S., Dayan,	
18	U., Temara, A.: MBAS (Methylene Blue Active Substances) and LAS (Linear Alkylbenzene Sulphonates) in	
19	Mediterranean coastal aerosols: sources and transport processes. Atmos. Environ., 45, 6788–6801,	
20	doi:10.1016/j.atmosenv.2011.04.041, 2011.	
21		
22	Becagli, S., Scarchilli, C., Traversi, R., Dayan, U., Severi, M., Frosini, D., Vitale, V., Mazzola, M., Lupi, A., Nava, S.,	
23	and Udisti, R.: Study of present-day sources and transport processes affecting oxidised sulphur compounds in	
24	atmospheric aerosols at Dome C (Antarctica) from year-round sampling campaigns, Atmos. Environ., 52, 98-	
25	108, 2012.	
26		
27	Becagli, S., Marchese, C., Caiazzo, L., Ciardini, V., Lazzara, L., Mori, G., Nuccio, C., Scarchilli, C., Severi, M., and	Formatted: Highlight
28	Traversi, R.: Biogenic aerosol in central East Antarctic Plateau as a proxy for the ocean-atmosphere interaction	
29	in the Southern Ocean. Science of The Total Environment, 151285,	
30	https://doi.org/10.1016/j.scitotenv.2021.151285, 2021.	Formatted: Highlight

Formatted: Highlight

1	Behrenfeld, M. J., Hu, Y., O'Malley, R. T., Boss, E. S., Hostetler, C. A., Siegel, D. A., Sarmiento, J. L., Schulien, J.,	
2	Hair, J. W., Lu, X., Rodier, S., and Scarino, A. J.: Annual boom-bust cycles of polar phytoplankton biomass	
3	revealed by space-based lidar, Nat. Geosci., 10, 118–122, https://doi.org/10.1038/ngeo2861, 2017.	
4		
5	Bird, R. E. and Hulstrom, R. L., Simplified clear sky model for direct and diffuse insolation on horizontal	
6	surfaces, SERI/TR-642-761, 1981.	
7		
8	Bond, T. C., Anderson, T. L., and Campbell, D.: Calibration and Intercomparison of Filter-Based Measurements	
9	of Visible Light Absorption by Aerosols, Aerosol Sci. Tech., 30, 582–600,	
10	https://doi.org/10.1080/027868299304435, 1999.	
11		
12	Bond, T. C. and Bergstrom, R. W.: Light Absorption by Carbonaceous Particles: An Investigative Review, Aerosol	
13	Sci. Technol., 40, 27–67, 2006.	
14		
15	Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S.,	
16	Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman,	
17	C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z.,	
18	Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of	
19	black carbon in the climate system: A scientific assessment, J. Geophys. Res., 118, 5380–5552,	
20	doi:10.1002/jgrd.50171, 2013.	
21		
22	Bourgeois, Q., Ekman, A. M. L., and Krejci, R.: Aerosol transport over the Andes from the Amazon Basin to the	Formatted: Highlight
23	remote Pacific Ocean: A multiyear CALIOP assessment, J. Geophys. Res., 120, 8411–8425,	
24	https://doi.org/10.1002/2015JD023254, 2015.	
25		
26		
27	Brean, J., Dall'Osto, M., Simó, R., Shi, Z., Beddows, D.C.S, and Harrison, R.: Open ocean and coastal new particle	
28	formation from sulfuric acid and amines around the Antarctic Peninsula. Nat. Geosci.	
29	<u>https://doi.org/10.1038/s41561-021-00751-γ</u> . 2021.	
30		
31	Brito, J., Rizzo, L. V., Morgan, W. T., Coe, H., Johnson, B., Haywood, J., Longo, K., Freitas, S., Andreae, M. O., and	Formatted: Highlight
32	Artaxo, P.: Ground-based aerosol characterization during the South American Biomass Burning Analysis	

1	<u>(</u> SAMBBA) field experiment, Atmos. Chem. Phys., 14, 12069–12083, https://doi.org/10.5194/acp-14-12069-	
2	<u>2014, 2014</u> .	
3		
4	Caiazzo, L., Calzolai, G., Becagli, S., Severi, M., Amore, A., Nardin, R., Chiari, M., Giardi, F., Nava, S., Lucarelli, F.,	
5	Pazzi, G., Cristofanelli, P., Virkkula, A., Gambaro, A., Barbaro, E., Traversi, R.: Carbonaceous Aerosol in Polar	
6	Areas: First Results and Improvements of the Sampling Strategies. Atmosphere 2021, 12, 320.	
7	https://doi.org/10.3390/atmos12030320, 2021.	
8		
9	Chatfield, R., Guo, Z., Sachse, G., Blake, D., and Blake, N.: The subtropical global plume in the Pacific Exploratory	
10	Mission Tropics A (PEM-Tropics A), PEM-Tropics B, and the Global Atmospheric Sampling Program (GASP): How	
11	tropical emissions affect the remote Pacific, J. Geophys. Res., 107(D16), doi:10.1029/2001JD000497, 2002.	
12		
13	Chen, X., Virkkula, A., Kerminen, VM., Manninen, H. E., Busetto, M., Lanconelli, C., Lupi, A., Vitale, V., Del	
14	Guasta, M., Grigioni, P., Väänänen, R., Duplissy, EM., Petäjä, T., and Kulmala, M.: Features in air ions	
15	measured by an air ion spectrometer (AIS) at Dome C, Atmos. Chem. Phys., 17, 13783–13800,	
16	https://doi.org/10.5194/acp-17-13783-2017, 2017.	
17		
18	Croft, B., Pierce, J. R., and Martin, R. V.: Interpreting aerosol lifetimes using the GEOS-Chem model and	Formatted: Highligh
19	<u>constraints from radionuclide measurements, Atmos. Chem. Phys., 14, 4313–4325,</u>	
20	https://doi.org/10.5194/acp-14-4313-2014, 2014.	
21		
22		
23	CSN EN 12341 European Standard: Ambient air Standard gravimetric measurement method for the	
24	determination of the PM10 or PM2,5 mass concentration of suspended particulate matter, https://www.en-	
25	standard.eu/csn-en-12341-ambient-air-standard-gravimetric-measurement-method-for-the-determination-of-	
26	the-pm10-or-pm2-5-mass-concentration-of-suspended-particulate-matter/	
27		
28	Devi, J. J., Bergin, M. H., Mckenzie, M., Schauer, J. J., and Weber, R. J.: Contribution of particulate brown carbon	
29	to light absorption in the rural and urban Southeast US. Atmospheric Environment, 136, 95-104, 2016.	
30		
31	Fiebig, M., Lunder, C. R., and Stohl, A.: Tracing biomass burning aerosol from South America to Troll Research	
32	Station, Antarctica, Geophys. Res. Lett., 36, L14815, doi:10.1029/2009GL038531, 2009.	

1		
2	Fiebig, M., Hirdman, D., Lunder, C. R., Ogren, J. A., Solberg, S., Stohl, A., and Thompson, R. L.: Annual cycle of	
3	Antarctic baseline aerosol: controlled by photooxidation-limited aerosol formation, Atmos. Chem. Phys., 14,	
4	3083–3093, https://doi.org/10.5194/acp-14-3083-2014, 2014.	
5		
6	Freitas, S. R., Longo, K. M., Silva Dias, M., Silva Dias, P., Chatfield, R., Prins, E., Artaxo, P., Grell, G., and Recuero,	 Formatted: Highlight
7	F.: Monitoring the transport of biomass burning emissions in South America, Environmental Fluid Mechanics,	
8	doi:10.1007/s10652-005-0243-7, 5(1–2), 135–167, 2005.	
9		
10	Frey, M. M., Norris, S. J., Brooks, I. M., Anderson, P. S., Nishimura, K., Yang, X., Jones, A. E., Nerentorp	 Formatted: Highlight
11	Mastromonaco, M. G., Jones, D. H., and Wolff, E. W.: First direct observation of sea salt aerosol production	
12	from blowing snow above sea ice, Atmos. Chem. Phys., 20, 2549–2578, https://doi.org/10.5194/acp-20-2549-	 Formatted: Highlight
13	<u>2020, 2020.</u>	 Formatted: Highlight
14		
15	Giglio, L., I. Csiszar, and C. O. Justice : Global distribution and seasonality of active fires as observed with the	
16	Terra and Aqua Moderate Resolution Imaging Spectroradiometer (MODIS) sensors, J. Geophys. Res., 111,	
17	G02016,doi:10.1029/2005JG000142, 2006.	
18		
19	Giglio, L., Loboda, T., Roy, D. P., Quayle, B., and Justice, C. O.: An active-fire based burned area mapping	
20	algorithm for the MODIS sensor, Remote. Sens. Environ., 113, 408-420, doi:10.1016/j.rse.2008.10.006, 2009.	
21		
22	Giglio, L., Randerson, J. T., van der Werf, G. R., Kasibhatla, P. S., Collatz, G. J., Morton, D. C., and DeFries, R. S.:	
23	Assessing variability and long-term trends in burned area by merging multiple satellite fire products,	
24	Biogeosciences, 7, 1171–1186, doi:10.5194/bg-7-1171-2010, 2010.	
25		
26	Gonzalez-Alonso, L., Val Martin, M., and Kahn, R. A.: Biomass-burning smoke heights over the Amazon	
27	observed from space, Atmos. Chem. Phys., 19, 1685–1702, https://doi.org/10.5194/acp-19-1685-2019, 2019.	
28		
29	Grythe, H., Ström, J., Krejci, R., Quinn, P., and Stohl, A.: A review of sea-spray aerosol source functions using a	
30	large global set of sea salt aerosol concentration measurements, Atmos. Chem. Phys., 14, 1277-1297,	
31	https://doi.org/10.5194/acp-14-1277-2014, 2014.	
32		

1	Grythe, H., Kristiansen, N. I., Groot Zwaaftink, C. D., Eckhardt, S., Ström, J., Tunved, P., Krejci, R., and Stohl, A.: A			
2	new aerosol wet removal scheme for the Lagrangian particle model FLEXPART v10, Geosci. Model Dev., 10,			
3	1447–1466, https://doi.org/10.5194/gmd-10-1447-2017, 2017.			
4				
5	Grythe, H.: Quantification of sources and removal mechanisms of atmospheric aerosol particles, PhD thesis,			
6	5 Stockholm University, 2017 <u>.</u>			
7				
8	Gyawali, M., Arnott, W.P., Lewis, K., Moosmüller, H.: In situ aerosol optics in Reno, NV, USA during and after			
9	the summer 2008 California wildfires and the influence of absorbing and non-absorbing organic coatings on			
10	spectral light absorption. Atmos. Chem. Phys. 9, 8007-8015, 2009.			
11				
12	Hand, J. L., and Malm, W. C.: Review of aerosol mass scattering efficiencies from ground-based measurements			
13	since 1990, J. Geophys. Res., 112, D16203, doi:10.1029/2007JD008484, 2007.			
14				
15	Hara, K., K. Osada, M. Yabuki, G. Hashida, T. Yamanouchi, M. Hayashi, M. Shiobara, C.Nishita, and M.Wada:			
16	Haze episodes at Syowa Station, coastal Antarctica: Where did they come from?, J. Geophys. Res., 115,			
17	D14205, doi:10.1029/2009JD012582, 2010.			
18				
19	Hara, K., Sudo, K., Ohnishi, T., Osada, K., Yabuki, M., Shiobara, M., and Yamanouchi, T.: Seasonal features and			
20	origins of carbonaceous aerosols at Syowa Station, coastal Antarctica, Atmos. Chem. Phys., 19, 7817-7837,			
21	https://doi.org/10.5194/acp-19-7817-2019, 2019.			
22				
23	Hara, K., Osada, K., Yabuki, M., Matoba, S., Hirabayashi, M., Fujita, S., Nakazawa, F., and Yamanouchi, T.:		Formatted: Highlight	_
24	Atmospheric seasalt and halogen cycles in the Antarctic, Environ. Sci.: Process Impacts, 22, 2003–2022,			
25	https://doi.org/10.1039/d0em00092b, 2020.	\triangleleft	Formatted: Highlight	
26	i de la constante d		Formatted: Highlight	
27	Hegg, D., Covert, D., Crahan, K., and Jonssen, H.: The dependence of aerosol light-scattering on RH over the			
28	Pacific Ocean, Geophys. Res. Lett., 29, 2002.			
29				
30	Hirdman, D., Sodemann, H., Eckhardt, S., Burkhart, J. F., Jefferson, A., Mefford, T., Quinn, P. K., Sharma, S.,		Formatted: Highlight	
31	Ström, J., and Stohl, A.: Source identification of short-lived air pollutants in the Arctic using statistical analysis			

1	of measurement data and particle dispersion model output, Atmos. Chem. Phys., 10, 669–693,	
2	https://doi.org/10.5194/acp-10-669-2010, 2010.	
3		
4	Hogan, A.W., S. Barnard, and J. Bortniak: Physical properties of the aerosol at the South Pole, Geophys. Res.	
5	<u>Lett., 6, 845–848, 1979.</u>	
6		
7	Ito, T.: Antarctic submicron aerosols and long-range transport of pollutants, Ambio,18,34–41, 1989.	
8		
9		
10	Ito, A., A. Ito, and H. Akimoto: Seasonal and interannual variations in CO and BC emissions from open biomass	
11	burning in Southern Africa during 1998 – 2005, Global Biogeochem. Cycles,21, GB2011,	
12	doi:10.1029/2006GB002848, 2007.	
13		
14	James, I. N.: The Antarctic drainage flow: Implications for hemispheric flow on the Southern Hemisphere,	
15	Antarct. Sci.,1,279–290, doi:10.1017/S0954102089000404., 1989.	
16		
17	Jourdain, B., S. Preunkert, O. Cerri, H. Castebrunet, R. Udisti, and M. Legrand: Year-round record of size	
18	segregated aerosol composition in central Antarctica (Concordia station): Implications for the degree of	
19	fractionation of sea-salt particles, J. Geophys. Res., 113, D14308, doi:10.1029/2007JD009584, 2008	
20		
21	Jurányi, Z. and Weller, R.: One year of aerosol refractive index measurement from a coastal Antarctic site,	
22	Atmos. Chem. Phys., 19, 14417–14430, https://doi.org/10.5194/acp-19-14417-2019, 2019.	
23		
24	Järvinen, E., Virkkula, A., Nieminen, T., Aalto, P. P., Asmi, E., Lanconelli, C., Busetto, M., Lupi, A., Schioppo, R.,	
25	Vitale, V., Mazzola, M., Petäjä, T., Kerminen, VM., and Kulmala, M.: Seasonal cycle and modal structure of	
26	particle number size distribution at Dome C, Antarctica, Atmos. Chem. Phys., 13, 7473-7487, doi:10.5194/acp-	
27	13-7473-2013, 2013.	
28		
29	Karpetchko, A., Kyro, E., and Knudsen, B.: Arctic and Antarctic polar vortices 1957-2002 as seen from the ERA-	
30	40 reanalyses, J. Geophys. Res., 110, D21109, doi:10.1029/2005JD006113, 2005.	
31		

1	Kerminen, V. M., Chen, X., Vakkari, V., Petäjä, T., Kulmala, M.,and Bianchi, F.: Atmospheric new particle	
2	formation and growth:review of field observations, Environ. Res. Lett., 13,	
3	103003,https://doi.org/10.1088/1748-9326/aadf3c, 2018.	
4		
5	Kim, J., Yoon, Y. J., Gim, Y., Choi, J. H., Kang, H. J., Park, KT., Park, J., and Lee, B. Y.: New particle formation	
6	events observed at King Sejong Station, Antarctic Peninsula - Part 1: Physical characteristics and contribution	
7	to cloud condensation nuclei, Atmos. Chem. Phys., 19, 7583-7594, https://doi.org/10.5194/acp-19-7583-2019,	
8	2019.	
9		
10	King, J. C., and Turner, J.: Antarctic Meteorology and Climatology, 409 pp., Cambridge Univ. Press, New York,	
11	doi:10.1017/CBO9780511524967, 1997.	
12		
13	Koch, D., Schulz, M., Kinne, S., McNaughton, C., Spackman, J. R., Balkanski, Y., Bauer, S., Berntsen, T., Bond, T.	Formatted: Highlight
14	C., Boucher, O., Chin, M., Clarke, A., De Luca, N., Dentener, F., Diehl, T., Dubovik, O., Easter, R., Fahey, D. W.,	
15	Feichter, J., Fillmore, D., Freitag, S., Ghan, S., Ginoux, P., Gong, S., Horowitz, L., Iversen, T., Kirkevåg, A., Klimont,	
16	Z., Kondo, Y., Krol, M., Liu, X., Miller, R., Montanaro, V., Moteki, N., Myhre, G., Penner, J. E., Perlwitz, J., Pitari,	
17	G., Reddy, S., Sahu, L., Sakamoto, H., Schuster, G., Schwarz, J. P., Seland, Ø., Stier, P., Takegawa, N., Takemura,	
18	T., Textor, C., van Aardenne, J. A., and Zhao, Y.: Evaluation of black carbon estimations in global aerosol models,	
19	<u> Atmos. Chem. Phys., 9, 9001–9026, https://doi.org/10.5194/acp-9-9001-2009, 2009.</u>	
20		
21	Krinner G. and Genthon C.: Tropospheric transport of continental tracers towards Antarctica under varying	
22	climatic conditions, Tellus B, 55:1, 54-70, DOI: 10.3402/tellusb.v55i1.16354, 2003.	
23		
24	Kirchstetter, T. W., Novakov, T., and Hobbs, P. V.: Evidence that the spectral dependence of light absorption by	
25	aerosols is affected by organic carbon, J. Geophys. Res., 109, D21208, https://doi.org/10.1029/2004jd004999,	
26	2004. Kirchstetter, T., Novakov, T., and Hobbs, P.: Evidence that the spectral dependence of light absorption	
27	by aerosols is affected by organic carbon, Journal of Geophysical Research, 109, 2004	
28		
29		
30	Kristiansen, N. I., Stohl, A., and Wotawa, G.: Atmospheric removal times of the aerosol-bound radionuclides	Formatted: Highlight
31	¹³⁷ Cs and ¹³¹ I measured after the Fukushima Dai-ichi nuclear accident – a constraint for air quality and climate	
32	models, Atmos. Chem. Phys., 12, 10759–10769, https://doi.org/10.5194/acp-12-10759-2012, 2012.	

2 Kyrö, E.-M., Grönholm, T., Vuollekoski, H., Virkkula, A., Kulmala, M., and Laakso, L.: Snow scavenging of 3 ultrafine particles: field measurements and parameterization, Boreal Environ. Res., 14,527–538, 2009. 4 5 Laakso, L., Grönholm, Rannika, Ü., Kosmalea, M., Fiedlera, V., Vehkamäki, H., and Kulmalaa, M.: Ultrafine 6 particle scavenging coefficients calculated from 6 years field measurements, Atmos. Environ., 37, 3605–3613, 7 doi:10.1016/S1352-2310(03)00326-1, 2003. 8 9 Lachlan-Cope, T., Beddows, D. C. S., Brough, N., Jones, A. E., Harrison, R. M., Lupi, A., Yoon, Y. J., Virkkula, A., 10 and Dall'Osto, M.: On the annual variability of Antarctic aerosol size distributions at Halley Research Station, 11 Atmos. Chem. Phys., 20, 4461–4476, https://doi.org/10.5194/acp-20-4461-2020, 2020. 12 13 Lack, D. A. and Cappa, C. D.: Impact of brown and clear carbon on light absorption enhancement, single scatter 14 albedo and absorption wavelength dependence of black carbon, Atmos. Chem. Phys., 10, 4207-4220, 15 doi:10.5194/acp-10-4207-2010, 2010. 16 17 Legrand, M., Yang, X., Preunkert, S., and Theys, N.: Year-round records of sea salt, gaseous, and particulate 18 inorganic bromine in the atmospheric boundary layer at coastal (Dumont d'Urville) and central (Concordia) East 19 Antarctic sites, J. Geophys. Res. Atmos., 121, 997–1023, https://doi.org/10.1002/2015JD024066, 2016. 20 21 Legrand, M., Preunkert, S., Wolff, E., Weller, R., Jourdain, B., and Wagenbach, D.: Year-round records of bulk 22 and size segregated aerosol composition in central Antarctica (Concordia site) - Part 1: Fractionation of sea-salt 23 particles, Atmos. Chem. Phys., 17, 14039-14054, https://doi.org/10.5194/acp-17-14039-2017, 2017a. 24 25 Legrand, M., Preunkert, S., Weller, R., Zipf, L., Elsässer, C., Merchel, S., Rugel, G., and Wagenbach, D.: Year-26 round record of bulk and size-segregated aerosol composition in central Antarctica (Concordia site) - Part 2: 27 Biogenic sulfur (sulfate and methanesulfonate) aerosol, Atmos. Chem. Phys., 17, 14055-14073, 28 https://doi.org/10.5194/acp-17-14055-2017, 2017b. 29 30 Li, H., McMeeking, G. R., and May, A. A.: Development of a new correction algorithm applicable to any filter-31 based absorption photometer, Atmos. Meas. Tech., 13, 2865-2886, https://doi.org/10.5194/amt-13-2865-32 2020, 2020.

1		
2	Liu, D., Allan, J. D., Young, D. E., Coe, H., Beddows, D., Fleming, Z. L., Flynn, M. J., Gallagher, M. W., Harrison, R.	
3	M., Lee, J., Prevot, A. S. H., Taylor, J. W., Yin, J., Williams, P. I., and Zotter, P.: Size distribution, mixing state and	
4	source apportionment of black carbon aerosol in London during wintertime, Atmos. Chem. Phys., 14, 10061-	
5	10084, https://doi.org/10.5194/acp-14-10061-2014, 2014.	
6		
7	Luoma, K., Virkkula, A., Aalto, P., Lehtipalo, K., Petäjä, T., and Kulmala, M.: Effects of different correction	
8	algorithms on absorption coefficient – a comparison of three optical absorption photometers at a boreal forest	
9	site, Atmos. Meas. Tech., 14, 6419–6441, https://doi.org/10.5194/amt-14-6419-2021, 2021.Luoma, K.,	
10	Virkkula, A., Aalto, P., Lehtipalo, K., Petäjä, T., and Kulmala, M.: A comparison of three optical absorption	
11	photometers at a boreal forest site - effects of different correction algorithms, Atmos. Meas. Tech. Discuss.,	
12	https://doi.org/10.5194/amt-2020-325, in review, 2020.	
13		
14		
15	McMeeking, G. R., Hamburger, T., Liu, D., Flynn, M., Morgan, W. T., Northway, M., Highwood, E. J., Krejci, R.,	
16	Allan, J. D., Minikin, A., and Coe, H.: Black carbon measurements in the boundary layer over western and	
17	northern Europe, Atmos. Chem. Phys., 10, 9393–9414, https://doi.org/10.5194/acp-10-9393-2010, 2010.	
18		
19	Mikhailov, E., Vlasenko, S., Podgorny, I., Ramanathan, V., and Corrigan, C.: Optical properties of soot-water	
20	drop agglomerates: An experimental study, J. Geophys. Res., 111, 1–16, 2006.	
21	Mishchenko, M. I., Travis, L., and Lacis, A.: Scattering, Absorption, and Emission of Light by Small Particles,	Formatted: Highlight
22	Cambridge University Press, 2002.	
23		
24	MorgantiA, Becagli_ S, Castellano_ E, Severi_ M, Traversi_ R, Udisti R.: An improved flow analysis-ion	
25	chromatography method for determination of cationic and anionic species at trace levels in Antarctic ice cores.	
26	Anal_ Chim_ Acta_ 603;_190–198, 2007.	
27		
28	Müller, T., Virkkula, A., and Ogren, J. A.: Constrained two-stream algorithm for calculating aerosol light	
29	absorption coefficient from the Particle Soot Absorption Photometer, Atmos. Meas. Tech., 7, 4049-4070,	
30	https://doi.org/10.5194/amt-7-4049-2014, 2014.	
31		Formatted: Tab stops: 12.25 cm, Let
I		

 Absorption by Aerosols," Aerosol Sci. Technol., 44(8), 589–591, doi:10.1080/02786826.2010.482111, 2010. Parastefanou, C. Residence time of tropospheric aerosols in association with radioactive nuclides, Appl. Radiat Isotones, 64, 93–100, 2006. Parish, T.R. and Bromwich, D.H.: Continental-scale simulation of the Antarctic katabatic wind regime, J. Clim., 4, 135–146, https://doi.org/10.1175/1520-0442(1991)004<0135:cssota>2.0.co;2, 1991. Pereira, G., Siqueira, R., Rosário, N. E., Longo, K. L., Freitas, S. R., Cardozo, F. S., Kaiser, J. W., and Wooster, M. J.: Assessment of fire emission inventories during the South American Biomass Burning Analysis (SAMBBA) experiment, Atmos. Chem. Phys., 16, 6961–6975, https://doi.org/10.5194/acp-16-6961-2016, 2016. Petcold, A., Ogren, J. A., Flebig, M., Laj, P., Li, SM., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto, N., Wehrli, C., Wiedensohler, A., and Zhang, XY.: Recommendations for reporting "black carbon" measurements, Atmos. Chem. Phys., 13, 8365–8379, https://doi.org/10.5194/acp-13-8365-2013, 2013. Piso, L., Sollum, E., Grythe, H., Kristansen, N. L. Castiani, M., Letchardt, S., Arnold, D., Morton, D., Thompson, R. J.: Groot Zwaaflink, C. D. Evengeliou, N., Sodemann, H., Haimberzer, L., Henne, S., Brunner, O., Burkhart, J. F. Foullioux, A., Brioude, J., Philipo, A., Selbert, P., and Stohl, A.: The Lagrangian particle dispersion model HLXPART version 10.4, Geosci. Model Dev, 12, 4955–4997, https://doi.org/10.5194/acp-13-935-2019, 2019. Pratt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, A. J., Heymsfield, A. J., Seinfield, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11-12549-20	1	Ogren, J. A.: Comment on "Calibration and Intercomparison of Filter-Based Measurements of Visible Light	
 Peasatefanou, C.: Residence time of tropospheric aerosols in association with radioactive nuclides, April Formatted: Highlight Rodint Rotrose, 64, 92–100, 2006 Parish, T.R. and Bromwich, D.H.: Continental-scale simulation of the Antarctic katabatic wind regime, J. Clim., 4, 135–146, https://doi.org/10.1175/1520-0442(1991)004<0135:cssota>2.0.co;2, 1991. Pereira, G., Siqueira, R., Rosário, N. E., Longo, K. L., Freitas, S. R., Cardozo, F. S., Kaiser, J. W., and Wooster, M. J.: Assessment of fire emission inventories during the South American Biomass Burning Analysis (SAMBBA) experiment, Atmos. Chem. Phys., 16, 6961–6975, https://doi.org/10.5194/acp-16-6961-2016, 2016. Petzold, A., Ogren, J. A., Fiebig, M., Laj, P., Li, S-M., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto, N., Wehrli, C., Wiedensohler, A., and Zhang, XY.: Recommendations for reporting "black carbon" measurements, Atmos. Chem. Phys., 13, 8365–8379, https://doi.org/10.5194/acp-13-8365-2013, 2013. Piszo, I. Sollum, E., Grythe, H., Kristinsen, N. I., Cassiani, M., Erkhardt, S., Arnold, D., Morton, D., Thomoson, R. J.: Groot Zwastink, C. D., Evangeliou, N., Sodemann, H., Haimberger, L., Henne, S., Brunner, D., Barkhart, J. F., Fouilloux, A., Brioude, J., Philipp, A., Seibert, P., and Stohl, A. The Lagrangian particle dispersion model Pitzt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, A. J., Heymsfield, A. J., Seinfeld, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11-12549-2011, 2011. 	2	Absorption by Aerosols," Aerosol Sci. Technol., 44(8), 589–591, doi:10.1080/02786826.2010.482111, 2010.	
 Papastefanou, C. Residence time of tropospheric aerosols in association with radioactive nuclides, Appl. Formatted: Highlight Fadiat. Isotones, 64, 93-100, 2006. Parish, T.R. and Bromwich, D.H.: Continental-scale simulation of the Antarctic katabatic wind regime, J. Clim., 4, 135–146, https://doi.org/10.1175/1520-0442(1991)004-0135:cssota>2.0.co;2, 1991. Pereira, G., Siqueira, R., Rosário, N. E., Longo, K. L., Freitas, S. R., Cardozo, F. S., Kaiser, J. W., and Wooster, M. J.: Assessment of fire emission inventories during the South American Biomass Burning Analysis (SAMBBA) experiment, Atmos. Chem. Phys., 16, 6961–6975, https://doi.org/10.5194/acp-16-6961-2016, 2016. Petzold, A., Ogren, J. A., Flebig, M., Laj, P., Li, SM., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto, N., Wehrli, C., Wiedensohler, A., and Zhang, XY.: Recommendations for reporting "black carbon" measurements, Atmos. Chem. Phys., 13, 8365–8379, https://doi.org/10.5194/acp-13-8365-2013, 2013. Petzold, A., Ogren, J. A., Flebig, M., Laj, P., Li, SM., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto, N., Wehrli, C., Wiedensohler, A., and Zhang, XY.: Recommendations for reporting "black carbon" measurements, Atmos. Chem. Phys., 13, 8365–8379, https://doi.org/10.5194/acp-13-8365-2013, 2013. Pisso, L. Sollum, E., Grythe, H., Kristiansen, N. J., Casslani, M., Etkhardt, S., Annold, D., Morton, D., Thompson, R. Formatted: Highlight L. Groot Zwaaftink, C. O., Evangeliou, N., Sodemann, H., Haimberger, L., Henne, S., Brunner, D., Burkhart, J. F. Poulloux, A., Brioude, J., Philipp, A., Selbert, P. and Stohl, A., The Lagrangian particle dispersion model PitzNART version 10.4, Geosci. Model Dev, 12, 4955–4997, https://doi.org/10.5194/gmd:12.4955-2019, 2010 Pratt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L.,	3		
 Radiat. Isotopes, 64, 93–100, 2005 Parish, T.R. and Bromwich, D.H.: Continental-scale simulation of the Antarctic katabatic wind regime, J. Clim., 4, 135–146, https://doi.org/10.1175/1520-0442(1991)004<0135:cssota>2.0.co;2, 1991. Pereira, G., Siqueira, R., Rosário, N. E., Longo, K. L., Freitas, S. R., Cardozo, F. S., Kaiser, J. W., and Wooster, M. J.: Assessment of fire emission inventories during the South American Biomass Burning Analysis (SAMBBA) experiment, Atmos. Chem. Phys., 16, 6961–6975, https://doi.org/10.5194/acp-16-6961-2016, 2016. Petzold, A., Ogren, J. A., Fiebig, M., Laj, P., Li, SM., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto, N., Wehrli, C., Wiedensohler, A., and Zhang, XY.: Recommendations for reporting "black carbon" measurements, Atmos. Chem. Phys., 13, 8365–8379, https://doi.org/10.5194/acp-13-8365-2013, 2013. Petzold, Zaoftink, C. D., Evangeliou, N., Sodemann, H., Haimberger, L., Henne, S., Brunner, D., Burkhart, J. F., Pouilloux, A., Brioude, J., Philipo, A., Seibert, P., and Stohl, A.: The Lagrangian particle dispersion model REXPART version 10.4, Geosai: Model Dev., 12, 4955–4997, https://doi.org/10.5194/gacf-12-4955-2019, 2019 Pratt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, A. J., Heymsfield, A. J., Seinfeld, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11-12549-2011, 2011. 	4	Papastefanou, C.: Residence time of tropospheric aerosols in association with radioactive nuclides, Appl.	Formatted: Highlight
 Parish, T.R. and Bromwich, D.H.: Continental-scale simulation of the Antarctic katabatic wind regime, J. Clim., 4, 135–146, https://doi.org/10.1175/1520-0442(1991)004-0135:cssota>2.0.co;2, 1991. Pereira, G., Siqueira, R., Rosário, N. E., Longo, K. L., Freitas, S. R., Cardozo, F. S., Kaiser, J. W., and Wooster, M. J.: Assessment of fire emission inventories during the South American Biomass Burning Analysis (SAMBBA) experiment, Atmos. Chem. Phys., 16, 6961–6975, https://doi.org/10.5194/acp-16-6961-2016, 2016. Petzold, A., Ogren, J. A., Fiebig, M., Laj, P., Li, SM., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto, N., Wehrli, C., Wiedensohler, A., and Zhang, XY.: Recommendations for reporting "black carbon" measurements, Atmos. Chem. Phys., 13, 8365–8379, https://doi.org/10.5194/acp-13-8365-2013, 2013. Pisso, I., Sollum, E., Grythe, H., Kristiansen, N. I., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. L. Groot Zwaaflink. C. D., Evangeliou, N., Sodemann, H., Haimberger, L. Henne, S., Brunner, D., Burkhart, J. F. Fouilloux, A., Brioude, J., Philipo, A., Seibert, P., and Stohl, A. The Lagrangian particle dispersion model BLEXPART version 10.4, Geosci. Model Dev., 12, 4955–4997, https://doi.org/10.5194/gmd-12.4955-2019, 2019 Pratt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, A. J., Heymsfield, A. J., Seinfeld, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11-12549-2011, 2011. 	5	Radiat. Isotopes, 64, 93–100, 2006.	
 Parish, T.R. and Bromwich, D.H.: Continental-scale simulation of the Antarctic katabatic wind regime, J. Clim, 4, 135–146, https://doi.org/10.1175/1520-0442(1991)004<0135:cssota>2.0.co;2, 1991. Pereira, G., Siqueira, R., Rosário, N. E., Longo, K. L., Freitas, S. R., Cardozo, F. S., Kaiser, J. W., and Wooster, M. J.: Assessment of fire emission inventories during the South American Biomass Burning Analysis (SAMBBA) experiment, Atmos. Chem. Phys., 16, 6961–6975, https://doi.org/10.5194/acp-16-6961-2016, 2016. Petzold, A., Ogren, J. A., Fiebig, M., Laj, P., Li, SM., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto, N., Wehrli, C., Wiedensohler, A., and Zhang, XY:: Recommendations for reporting "black carbon" measurements, Atmos. Chem. Phys., 13, 8365–8379, https://doi.org/10.5194/acp-13-8365-2013, 2013. Pisso, I., Solum, E., Grythe, H., Kristiansen, N. L. Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. Groot Zwaafink, C. D., Evangeliou, N., Sodemann, H., Haimberger, L., Henne, S., Brunner, D., Burkhardt, J. F. Fouilloux, A., Broude, J., Philipo, A., Seibert, P., and Stohl, A.: The Lagrangian particle dispersion model FLXPART version 10.4, Geosci. Model Dev., 12, 4955–4997, https://doi.org/10.5194/gmd-12-4955-2019, 2019 Pratt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, A. J., Heymsfield, A. J., Seinfeld, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11-12549-2011, 2011. 	6		
 4, 135–146, https://doi.org/10.1175/1520-0442(1991)004<0135:cssota>2.0.co;2, 1991. Pereira, G., Siqueira, R., Rosário, N. E., Longo, K. L., Freitas, S. R., Cardozo, F. S., Kaiser, J. W., and Wooster, M. J.: Assessment of fire emission inventories during the South American Biomass Burning Analysis (SAMBBA) experiment, Atmos. Chem. Phys., 16, 6961–6975, https://doi.org/10.5194/acp-16-6961-2016, 2016. Petzold, A., Ogren, J. A., Fiebig, M., Laj, P., Li, SM., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto, N., Wehrli, C., Wiedensohler, A., and Zhang, XY.: Recommendations for reporting "black carbon" measurements, Atmos. Chem. Phys., 13, 8365–8379, https://doi.org/10.5194/acp-13-8365-2013, 2013. Pisso, I., Sollum, E., Grythe, H., Kristansen, N. I., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. L., Groot Zwaaffink, C. D., Evangeliou, N., Sødemann, H., Halmberger, L., Henne, S., Brunner, D., Burkhart, J. F., Foullioux, A., Brioude, J., Philipp, A., Seibert, P., and Stohl, A: The Lagrangian particle dispersion model ELEXPART version 10.4, Geosci. Model Dev, 12, 4955–4997, https://doi.org/10.5194/gmd-12-4955-2019, 2019 Pratt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, A. J., Heymsfield, A. J., Seinfeld, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11-12549-2011, 2011. 	7	Parish, T.R. and Bromwich, D.H.: Continental-scale simulation of the Antarctic katabatic wind regime, J. Clim.,	
 Pereira, G., Siqueira, R., Rosário, N. E., Longo, K. L., Freitas, S. R., Cardozo, F. S., Kaiser, J. W., and Wooster, M. J.: Assessment of fire emission inventories during the South American Biomass Burning Analysis (SAMBBA) experiment, Atmos. Chem. Phys., 16, 6961–6975, https://doi.org/10.5194/acp-16-6961-2016, 2016. Petzold, A., Ogren, J. A., Fiebig, M., Laj, P., Li, SM., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto, N., Wehrli, C., Wiedensohler, A., and Zhang, XY.: Recommendations for reporting "black carbon" measurements, Atmos. Chem. Phys., 13, 8365–8379, https://doi.org/10.5194/acp-13-8365-2013, 2013. Pisso, I., Sollum, E., Grythe, H., Kristiansen, N. I., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. Formatted: Highlight L., Groot Zwaaflink, C. D., Evangeliou, N., Sodemann, H., Haimberger, L., Henne, S., Brunner, D., Burkhart, J. F., fouilloux, A., Brioude, J., Philipp, A., Selbert, P., and Stohl, A.: The Lagrangian particle dispersion model FLEXPART version 10.4, Geosci. Model Dev., 12, 4955–4997, https://doi.org/10.5194/gmd-12-4955-2019, 2019 Pratt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, A. J., Heymsfield, A. J., Seinfeld, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11-12549-2011, 2011. 	8	4, 135–146, https://doi.org/10.1175/1520-0442(1991)004<0135:cssota>2.0.co;2, 1991.	
 Pereira, G., Siqueira, R., Rosário, N. E., Longo, K. L., Freitas, S. R., Cardozo, F. S., Kaiser, J. W., and Wooster, M. J.: Assessment of fire emission inventories during the South American Biomass Burning Analysis (SAMBBA) experiment, Atmos. Chem. Phys., 16, 6961–6975, https://doi.org/10.5194/acp-16-6961-2016, 2016. Petzold, A., Ogren, J. A., Fiebig, M., Laj, P., Li, SM., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto, N., Wehrli, C., Wiedensohler, A., and Zhang, XY.: Recommendations for reporting "black carbon" measurements, Atmos. Chem. Phys., 13, 8365–8379, https://doi.org/10.5194/acp-13-8365-2013, 2013. Pisso, I., Sollum, E., Grythe, H., Kristiansen, N. I., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. Formatted: Highlight L., Groot Zwaaftink, C. D., Evangeliou, N., Sodemann, H., Haimberger, L., Henne, S., Brunner, D., Burkhart, J. F., fouilloux, A., Brioude, J., Philipo, A., Seibert, P., and Stohl, A.: The Lagrangian particle dispersion model FLEXPART version 10.4, Geosci. Model Dev., 12, 4955–4997, https://doi.org/10.5194/gmd-12-4955-2019, 2019, Pratt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, A. J., Heymsfield, A. J., Seinfeld, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11-12549-2011, 2011. 	9		
 J.: Assessment of fire emission inventories during the South American Biomass Burning Analysis (SAMBBA) experiment, Atmos. Chem. Phys., 16, 6961–6975, https://doi.org/10.5194/acp-16-6961-2016, 2016. Petzold, A., Ogren, J. A., Fiebig, M., Laj, P., Li, SM., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto, N., Wehrli, C., Wiedensohler, A., and Zhang, XY.: Recommendations for reporting "black carbon" measurements, Atmos. Chem. Phys., 13, 8365–8379, https://doi.org/10.5194/acp-13-8365-2013, 2013. Pisso, L., Sollum, E., Grythe, H., Kristiansen, N. I., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. g. Groot Zwaaftink, C. D., Evangeliou, N., Sodemann, H., Haimberger, L., Henne, S., Brunner, D., Burkhart, J. E., fouilloux, A., Brioude, J., Philipp, A., Seibert, P., and Stohl, A.: The Lagrangian particle dispersion model <u>flEXPART version 10.4, Geosci. Model Dev., 12, 4955–4997, https://doi.org/10.5194/gmd-12-4955-2019, 2019.</u> Pratt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, A. J., Heymsfield, A. J., Seinfeld, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11-12549-2011, 2011. 	10	Pereira, G., Siqueira, R., Rosário, N. E., Longo, K. L., Freitas, S. R., Cardozo, F. S., Kaiser, J. W., and Wooster, M.	
 experiment, Atmos. Chem. Phys., 16, 6961–6975, https://doi.org/10.5194/acp-16-6961-2016, 2016. Petzold, A., Ogren, J. A., Fiebig, M., Laj, P., Li, SM., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto, N., Wehrli, C., Wiedensohler, A., and Zhang, XY.: Recommendations for reporting "black carbon" measurements, Atmos. Chem. Phys., 13, 8365–8379, https://doi.org/10.5194/acp-13-8365-2013, 2013. Pisso, I., Sollum, E., Grythe, H., Kristiansen, N. I., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. Pisso, I., Sollum, E., Grythe, H., Kristiansen, N. I., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. Pisso, I., Sollum, E., Grythe, H., Kristiansen, N. I., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. Pisso, I., Sollum, E., Grythe, H., Kristiansen, N. I., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. Pisso, I., Sollum, E., Grythe, H., Kristiansen, N. I., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. Pisso, I., Sollum, E., Grythe, H., Kristiansen, N. I., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. Pisso, I., Sollum, E., Grythe, H., Kristiansen, N. I., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. Pisso, I., Sollum, E., Grythe, H., Kristiansen, N. I., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. Pisso, I., Soluta, A., Brioude, J., Philipp, A., Selbert, P., and Stohl, A.: The Lagrangian particle dispersion model Pitzt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, A. J., Heymsfield, A. J., Seinfeld, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11.12549-2011, 2011. 	11	J.: Assessment of fire emission inventories during the South American Biomass Burning Analysis (SAMBBA)	
 Petzold, A., Ogren, J. A., Fiebig, M., Laj, P., Li, SM., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto, N., Wehrli, C., Wiedensohler, A., and Zhang, XY.: Recommendations for reporting "black carbon" measurements, Atmos. Chem. Phys., 13, 8365–8379, https://doi.org/10.5194/acp-13-8365-2013, 2013. Pisso, I., Sollum, E., Grythe, H., Kristiansen, N. I., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. <u>Pisso, I., Sollum, E., Grythe, H., Kristiansen, N. I., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R.</u> <u>Fouilloux, A., Brioude, J., Philipp, A., Seibert, P., and Stohl, A.: The Lagrangian particle dispersion model</u> <u>FLEXPART version 10.4, Geosci. Model Dev., 12, 4955–4997, https://doi.org/10.5194/gmd-12-4955-2019, 2019</u> Pratt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, A. J., Heymsfield, A. J., Seinfeld, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11-12549-2011, 2011. 	12	experiment, Atmos. Chem. Phys., 16, 6961–6975, https://doi.org/10.5194/acp-16-6961-2016, 2016.	
 Petzold, A., Ogren, J. A., Fiebig, M., Laj, P., Li, SM., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto, N., Wehrli, C., Wiedensohler, A., and Zhang, XY.: Recommendations for reporting "black carbon" measurements, Atmos. Chem. Phys., 13, 8365–8379, https://doi.org/10.5194/acp-13-8365-2013, 2013. Pisso, I., Sollum, E., Grythe, H., Kristiansen, N. L., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. L., Groot Zwaaftink, C. D., Evangeliou, N., Sodemann, H., Haimberger, L., Henne, S., Brunner, D., Burkhart, J. F., Fouilloux, A., Brioude, J., Philipp, A., Seibert, P., and Stohl, A.: The Lagrangian particle dispersion model FLEXPART version 10.4, Geosci. Model Dev., 12, 4955–4997, https://doi.org/10.5194/gmd-12-4955-2019, 2019 Pratt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, A. J., Heymsfield, A. J., Seinfeld, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11-12549-2011, 2011. 	13		
 G., Sugimoto, N., Wehrli, C., Wiedensohler, A., and Zhang, XY.: Recommendations for reporting "black carbon" measurements, Atmos. Chem. Phys., 13, 8365–8379, https://doi.org/10.5194/acp-13-8365-2013, 2013. Pisso, I., Sollum, E., Grythe, H., Kristiansen, N. I., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. L., Groot Zwaaftink, C. D., Evangeliou, N., Sodemann, H., Haimberger, L., Henne, S., Brunner, D., Burkhart, J. F., Fouilloux, A., Brioude, J., Philipp, A., Seibert, P., and Stohl, A.: The Lagrangian particle dispersion model FLEXPART version 10.4, Geosci. Model Dev., 12, 4955–4997, https://doi.org/10.5194/gmd-12-4955-2019, 2019. Pratt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, A. J., Heymsfield, A. J., Seinfeld, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11-12549-2011, 2011. 	14	Petzold, A., Ogren, J. A., Fiebig, M., Laj, P., Li, SM., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo,	
 measurements, Atmos. Chem. Phys., 13, 8365–8379, https://doi.org/10.5194/acp-13-8365-2013, 2013. Pisso, I., Sollum, E., Grythe, H., Kristiansen, N. L. Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. L., Groot Zwaaftink, C. D., Evangeliou, N., Sodemann, H., Haimberger, L., Henne, S., Brunner, D., Burkhart, J. F., Fouilloux, A., Brioude, J., Philipp, A., Seibert, P., and Stohl, A.: The Lagrangian particle dispersion model FLEXPART version 10.4, Geosci. Model Dev., 12, 4955–4997, https://doi.org/10.5194/gmd-12-4955-2019, 2019. Pratt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, A. J., Heymsfield, A. J., Seinfeld, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11-12549-2011, 2011. 	15	G., Sugimoto, N., Wehrli, C., Wiedensohler, A., and Zhang, XY.: Recommendations for reporting "black carbon"	
 Pisso, I., Sollum, E., Grythe, H., Kristiansen, N. L., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. L., Groot Zwaaftink, C. D., Evangeliou, N., Sodemann, H., Haimberger, L., Henne, S., Brunner, D., Burkhart, J. F., Fouilloux, A., Brioude, J., Philipp, A., Seibert, P., and Stohl, A.: The Lagrangian particle dispersion model FLEXPART version 10.4, Geosci. Model Dev., 12, 4955–4997, https://doi.org/10.5194/gmd-12-4955-2019, 2019. Pratt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, A. J., Heymsfield, A. J., Seinfeld, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11-12549-2011, 2011. 	16	measurements, Atmos. Chem. Phys., 13, 8365–8379, https://doi.org/10.5194/acp-13-8365-2013, 2013.	
 Pisso, I., Sollum, E., Grythe, H., Kristiansen, N. I., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R. L., Groot Zwaaftink, C. D., Evangeliou, N., Sodemann, H., Haimberger, L., Henne, S., Brunner, D., Burkhart, J. F., Fouilloux, A., Brioude, J., Philipp, A., Seibert, P., and Stohl, A.: The Lagrangian particle dispersion model FLEXPART version 10.4, Geosci. Model Dev., 12, 4955–4997, https://doi.org/10.5194/gmd-12-4955-2019, 2019. Pratt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, A. J., Heymsfield, A. J., Seinfeld, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11-12549-2011, 2011. 	17		
 L., Groot Zwaaftink, C. D., Evangeliou, N., Sodemann, H., Haimberger, L., Henne, S., Brunner, D., Burkhart, J. F., Fouilloux, A., Brioude, J., Philipp, A., Seibert, P., and Stohl, A.: The Lagrangian particle dispersion model FLEXPART version 10.4, Geosci. Model Dev., 12, 4955–4997, https://doi.org/10.5194/gmd-12-4955-2019, 2019. Pratt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, A. J., Heymsfield, A. J., Seinfeld, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11-12549-2011, 2011. 	18	Pisso, I., Sollum, E., Grythe, H., Kristiansen, N. I., Cassiani, M., Eckhardt, S., Arnold, D., Morton, D., Thompson, R.	 Formatted: Highlight
 Fouilloux, A., Brioude, J., Philipp, A., Seibert, P., and Stohl, A.: The Lagrangian particle dispersion model FLEXPART version 10.4, Geosci. Model Dev., 12, 4955–4997, https://doi.org/10.5194/gmd-12-4955-2019, 2019. Pratt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, A. J., Heymsfield, A. J., Seinfeld, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11-12549-2011, 2011. 	19	L., Groot Zwaaftink, C. D., Evangeliou, N., Sodemann, H., Haimberger, L., Henne, S., Brunner, D., Burkhart, J. F.,	
 FLEXPART version 10.4, Geosci. Model Dev., 12, 4955–4997, https://doi.org/10.5194/gmd-12-4955-2019, 2019. Pratt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, A. J., Heymsfield, A. J., Seinfeld, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11-12549-2011, 2011. 	20	Fouilloux, A., Brioude, J., Philipp, A., Seibert, P., and Stohl, A.: The Lagrangian particle dispersion model	
 Pratt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, A. J., Heymsfield, A. J., Seinfeld, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11-12549-2011, 2011. 	21	FLEXPART version 10.4, Geosci. Model Dev., 12, 4955–4997, https://doi.org/10.5194/gmd-12-4955-2019, 2019.	
 Pratt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, A. J., Heymsfield, A. J., Seinfeld, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11-12549-2011, 2011. 	22		
 Heymsfield, A. J., Seinfeld, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11-12549-2011, 2011. Ouine D. K. Müller, T. J., Dates, T. S., Orens, J. A. Anderen, F. and Shara, C. S. A. Dansen model of 	23	Pratt, K. A., Murphy, S. M., Subramanian, R., DeMott, P. J., Kok, G. L., Campos, T., Rogers, D. C., Prenni, A. J.,	
 aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549–12565, https://doi.org/10.5194/acp-11-12549-2011, 2011. Outron D. K. Atilka, J. J. & Dates J. S. Oraza J. A. Andrean, E. and Shan, G. S. A. Danara and af 	24	Heymsfield, A. J., Seinfeld, J. H., and Prather, K. A.: Flight-based chemical characterization of biomass burning	
26 https://doi.org/10.5194/acp-11-12549-2011, 2011. 27 28 Quine, Duty, Million, Tudy, Datas, Tudy, Constant, Million, Constant, Structure, Structur	25	aerosols within two prescribed burn smoke plumes, Atmos. Chem. Phys., 11, 12549-12565,	
27 28 - Outre D. K. Miller T. L. Dates T. S. Orres J. A. Andress F. and Share G. F. A. Dares month of	26	https://doi.org/10.5194/acp-11-12549-2011, 2011.	
29. Output D. K. Miller, T. L. Dates, T. C. Orace, J. A. Andrews, F. and Shaw, C. F. A. Davies moved of	27		
28 Quinn, P. K., Miller, T. L., Bates, T. S., Ogren, J. A., Andrews, E., and Snaw, G. E.: A 3-year record of	28	Quinn, P. K., Miller, T. L., Bates, T. S., Ogren, J. A., Andrews, E., and Shaw, G. E.: A 3-year record of	
29 simultaneously measured aerosol chemical and optical properties at Barrow, Alaska, J. Geophys. ResAtmos.,	29	simultaneously measured aerosol chemical and optical properties at Barrow, Alaska, J. Geophys. ResAtmos.,	
30 107, 4130, doi:10.1029/2001JD001248, 2002.	30	107, 4130, doi:10.1029/2001JD001248, 2002.	

1	Reid, J. S., Hyer, E. J., Prins, E. M., Westphal, D. L., Zhang, J., Wang, J., Christopher, S. A., Curtis, C. A., Schmidt,
2	C. C., Eleuterio, D. P., Richardson, K. A., and Hoffman, J. P.: Global monitoring and forecasting of biomass-
3	burning smoke: Description and lessons from the Fire Locating and Modeling of Burning Emissions (FLAMBE)
4	program, J. Sel. Topics Appl. Earth Obs. Rem. Sens., 2, 144–162, 2009.
5	
6	Rémy, S., Veira, A., Paugam, R., Sofiev, M., Kaiser, J. W., Marenco, F., Burton, S. P., Benedetti, A., Engelen, R. J.,
7	Ferrare, R., and Hair, J. W.: Two global data sets of daily fire emission injection heights since 2003, Atmos.
8	Chem. Phys., 17, 2921–2942, https://doi.org/10.5194/acp-17-2921-2017, 2017.
9	
10	Russell, P., Bergstrom, R., Shinozuka, Y., Clarke, A., DeCarlo, P., Jimenez, J., Livingston, J., Redemann, J.,
11	Dubovik, O., Strawa, A.: Absorption Angstrom Exponent in AERONET and related data as an indicator of aerosol
12	composition. Atmos. Chem. Phys. 10, 1155-1169, 2010.
13	
14	Seibert, P. and Frank, A.: Source-receptor matrix calculation with a Lagrangian particle dispersion model in
15	backward mode, Atmos. Chem. Phys., 4, 51–63, 2004, http://www.atmos-chem-phys.net/4/51/2004/.
16	
17	Seinfeld, J. H., and S. N. Pandis: Atmospheric Chemistry and Physics: From Air Pollution to Climate Change, John
18	Wiley, Hoboken, N. J., 1998.
19	
20	Shaw, G.: Considerations on the Origin and Properties of the Antarctic Aerosol, Rev. Geophys. 8, 1983 – 1998,
21	1979.
22	
23	Shen, Y., Virkkula, A., Ding, A., Wang, J., Chi, X., Nie, W., Qi, X., Huang, X., Liu, Q., Zheng, L., Xu, Z., Petäjä, T.,
24	Aalto, P. P., Fu, C., and Kulmala, M.: Aerosol optical properties at SORPES in Nanjing, east China, Atmos. Chem.
25	Phys., 18, 5265–5292, https://doi.org/10.5194/acp-18-5265-2018, 2018.
26	
27	Sheridan, P., Andrews, E., Schmeisser, L., Vasel, B., and Ogren, J.: Aerosol measurements at South Pole:
28	climatology and impact of local contamination. Aerosol Air Qual. Res. 16, 855–872.
29	https://doi.org/10.4209/aagr.2015.05.0358, 2016.

1	Sherman, J. P., Sheridan, P. J., Ogren, J. A., Andrews, E., Hageman, D., Schmeisser, L., Jefferson, A., and Sharma,		
2	S.: A multi-year study of lower tropospheric aerosol variability and systematic relationships from four North		
3	American regions, Atmos. Chem. Phys., 15, 12487–12517, https://doi.org/10.5194/acp-15-12487-2015, 2015.		
4			
5	Shikwambana, L. and Kganyago, M.: Observations of Emissions and the Influence of Meteorological Conditions		
6	during Wildfires: A Case Study in the USA, Brazil, and Australia during the 2018/19 Period. Atmosphere, 12, 11.		
7	https://dx.doi.org/10.3390/atmos12010011, 2021.		
8			
9	Singh, H. B., Viezee, W., Chen, Y., Bradshaw, J., Sandholm, S., Blake, D., Blake, N., Heikes, B., Snow, J., Talbot, R.,	 Formatted: Highlight	
10	Browell, E., Gregory, G., Sachse, G., and Vay, S.: Biomass burning influences on the composition of the remote		
11	South Pacific troposphere: analysis based on observations from PEM-Tropics-A, Atmos. Environ., 34(4), 635–		
12	<u>644, 2000.</u>		
13			
14			
15	Springston, S.R.: Particle Soot Absorption Photometer (PSAP) Instrument Handbook; DOE ARM Climate		
16	Research		
17	Facility: Washington, DC, USA, 2018.		
18			
19	Staudt, A. C., Jacob, D. J., Logan, J. A., Bachiochi, D., Krishnamurti, T. N., and Poisson, N.: Global chemical model		
20	analysis of biomass burning and lightning influences over the South Pacific in austral spring, J. Geophys. Res.,		
21	107, 4200, doi:10.1029/2000JD000296, 2002.		
22			
23	Stohl, A. and Sodemann, H.: Characteristics of atmospheric transport into the Antarctic troposphere, J.	 Formatted: Highlight	
24	Geophys. Res., 115, D02305, doi:10.1029/2009JD012536, 2010.		
25			
26	Stohl, A., Forster, C., Frank, A., Seibert, P., and Wotawa, G.: Technical note: The Lagrangian particle dispersion	 Formatted: Highlight	
27	model FLEXPART version 6.2, Atmos. Chem. Phys., 5, 2461–2474, https://doi.org/10.5194/acp-5-2461-2005,		
28	<u>2005.</u>		
29			
30	Tomasi, C., Vitale, V., Lupi, A., Di Carmine, C., Campanelli, M., Herber, A., Treffeisen, R., Stone, R-S., Andrews,		
31	E., Sharma, S., Radionov, V., von Hoyningen-Huene, W., Stebel, K., Hansen, G.H., Myhre, C.L., Wehrli, C.,		

Aaltonen, V., Lihavainen, H., Virkkula, A., Hillamo, R., Ström, J., Toledano, C., Cachorro, V.E., Ortiz, P., de Frutos,

 polar regions: A historical overview on the basis of optical depth and in situ observations, J. Geophys. Res., 112, D16205, doi:10.1029/2007/D0084322007. Tosca, M. G., Randerson, J. T., Zender, C. S., Nelson, D. L., Diner, D. J., and Logan, J. A.: Dynamics of fire plumes and smoke clouds associated with peat and deforestation fires in Indonesia, J. Geophys. Res., 116, D08207, doi:10.1029/2010/D015148, 2011. Udisti, R., Becagli, S., Benassai, S., Castellano, E., Fattori, I., Innocenti, M., Migliori, A., Traversi, R.: Atmosphere-is snow interaction by a comparison between aerosol and uppermost snow layers composition at Dome C (East Antarctica). Ann. Glaciol, 39, 53–61, 2004. Udisti, R., Dayan, U., Becagli, S., Busetto, M., Frosini, D., Legrand, M., Lucarelli, F., Preunkert, S., Severi, M., Traversi, R., and Vitale, V.: Sea spray aerosol in central Antarctica. Present atmospheric behaviour and implications for paleoclimatic reconstructions, Atmos. Environ., 52, 109–120, 2012. Ulke, A. G., Longo, K. M., and de Freitas, S. R.: Biomass Burning in South America: Transport Patterns and Immacts, in: Biomass – Detection, Production and Usage, edited by M. D. Matovic, InTech, Rijeka, Croatia, https://doi.org/10.5772/19264.2011. Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP), Aerosol Sci. Tech., 44, 706–712, https://doi.org/10.1080/02786826.2010.482110, 2010. Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, https://doi.org/10.1080/02786826.2010.482110, 2010. Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, https://doi.org/10.5194/amt.14-3707-2021, 2021. Virkkula, A.: Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a fi	1	A.M., Blindheim, S., Frioud, M., Gausa, M., Zielinski, T., Petelski, T., and Yamanouchi, T. (2007): Aerosols in					
 112, D16205, doi:10.1029/2007JD008432.,2007. Tosca, M. G., Randerson, J. T., Zender, C. S., Nelson, D. L., Diner, D. J., and Logan, J. A.: Dynamics of fire plumes and smoke clouds associated with peat and deforestation fires in Indonesia, J. Geophys. Res., 116, D08207, doi:10.1029/2010JD015148, 2011. Udisti, R., Becagli, S., Benassai, S., Castellano, E., Fattori, I., Innocenti, M., Migliori, A., Traversi, R.: Atmosphere- snow interaction by a comparison between aerosol and uppermost snow layers composition at Dome C (East Antarctica). Ann. Giaciol, 39, 53–61, 2004. Udisti, R., Dayan, U., Becagli, S., Busetto, M., Frosini, D., Legrand, M., Lucarelli, F., Preunkert, S., Severi, M., Traversi, R., and Vitale, V.: Sea spray aerosol in central Antarctica. Present atmospheric behaviour and implications for paleoclimatic reconstructions, Atmos. Environ., 52, 109–120, 2012. Ulke, A. G., Longo, K. M., and de Freitas, S. R.; Biomass Burning, in South America: Transport Patterns and Impacts, In: Biomass – Detection, Production and Usage, edited by M. D. Matovic, InTech, Rijeka, Croatia, https://doi.org/10.5772/19264, 2011. Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP), Aerosol Sci. Tech., 44, 706–712, https://doi.org/10.1080/02786826.2010.482110, 2010. Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, https://doi.org/10.5194/amt.14-3707-2021, 2021. Virkkula, A.: Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	2	polar regions: A historical overview on the basis of optical depth and in situ observations, J. Geophys. Res.,					
 Tosca, M. G., Randerson, J. T., Zender, C. S., Nelson, D. L., Diner, D. J., and Logan, J. A.: Dynamics of fire plumes and smoke clouds associated with peet and deforestation fires in Indonesia, J. Geophys. Res., 116, D08207, doi:10.1029/2010/D015148, 2011. Udisti, R., Becagli, S., Benassai, S., Castellano, E., Fattori, I., Innocenti, M., Migliori, A., Traversi, R.: Atmosphere-snow interaction by a comparison between aerosol and uppermost snow layers composition at Dome C (East Antarctica). Ann. Glaciol, 39, 53–61, 2004. Udisti, R., Dayan, U., Becagli, S., Busetto, M., Frosini, D., Legrand, M., Lucarelli, F., Preunkert, S., Severi, M., Traversi, R., and Vitale, V.: Sea spray aerosol in central Antarctica. Present atmospheric behaviour and implications for paleoclimatic reconstructions, Atmos. Environ., 52, 109–120, 2012. Ulke, A. G., Longo, K. M., and de Freitas, S. R.: Biomass Burning in South America: Transport Patterns and Impacts, in: Biomass – Detection, Production and Usage, edited by M. D. Matovic, InTech, Riieka, Croatia, https://doi.org/10.5772/19264, 2011. Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP), Aerosol Sci. Tech., 44, 706–712, https://doi.org/10.1080/02786826.2010.482110, 2010. Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, https://doi.org/10.5194/amt.14-3707-2021, 2021. Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	3	112, D16205, doi:10.1029/2007JD008432- <u>, 2007.</u>					
 Tosca, M. G., Randerson, J. T., Zender, C. S., Nelson, D. L., Diner, D. J., and Logan, J. A.: Dynamics of fire plumes and smoke clouds associated with peat and deforestation fires in Indonesia, J. Geophys. Res., 116, D08207, doi:10.1029/2010JD015148, 2011. Udisti, R., Becagli, S., Benassai, S., Castellano, E., Fattori, I., Innocenti, M., Migliori, A., Traversi, R.: Atmosphere- snow interaction by a comparison between aerosol and uppermost snow layers composition at Dome C (East Antarctica). Ann. Glaciol, 39, 53–61, 2004. Udisti, R., Dayan, U., Becagli, S., Busetto, M., Frosini, D., Legrand, M., Lucarelli, F., Preunkert, S., Severi, M., Traversi, R., and Vitale, V.: Sea spray aerosol in central Antarctica. Present atmospheric behaviour and implications for paleoclimatic reconstructions, Atmos. Environ., 52, 109–120, 2012. Ulke, A. G., Longo, K. M., and de Freitas, S. R.: Biomass Burning in South America: Transport Patterns and Impacts, in: Biomass – Detection, Production and Usage, edited by M. D. Matovic, InTech, Rieka, Croatia, https://doi.org/10.5772/19264, 2011. Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP), Aerosol Sci. Tech., 44, 706–712, <u>https://doi.org/10.1080/02786826.2010.482110</u>, 2010. Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, <u>https://doi.org/10.5194/amt-14-3707-2021</u>, 2021. Virkkula, A.: Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	4						
 and smoke clouds associated with peat and deforestation fires in Indonesia, J. Geophys. Res., 116, D08207, doi:10.1029/2010JD015148, 2011. Udisti, R., Becagli, S., Benassai, S., Castellano, E., Fattori, I., Innocenti, M., Migliori, A., Traversi, R.: Atmosphere-snow interaction by a comparison between aerosol and uppermost snow layers composition at Dome C (East Antarctica). Ann. Glaciol, 39, 53–61, 2004. Udisti, R., Dayan, U., Becagli, S., Busetto, M., Frosini, D., Legrand, M., Lucarelli, F., Preunkert, S., Severi, M., Traversi, R., and Vitale, V.: Sea spray aerosol in central Antarctica. Present atmospheric behaviour and implications for paleoclimatic reconstructions, Atmos. Environ., 52, 109–120, 2012. Ulke, A. G., Longo, K. M., and de freitas, S. R.: Biomass Burning in South America: Transport Patterns and Impacts, in: Biomass – Detection, Production and Usage, edited by M. D. Matovic, InTech, Rijeka, Croatia, https://doi.org/10.5124/2011. Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP), Aerosol Sci. Tech., 44, 706–712, https://doi.org/10.5194/amt-14-3707-2021, 2010. Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, https://doi.org/10.5194/amt-14-3707-2021, 2021. Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	5	Tosca, M. G., Randerson, J. T., Zender, C. S., Nelson, D. L., Diner, D. J., and Logan, J. A.: Dynamics of fire plumes					
 doi:10.1029/2010/D015148, 2011. Udisti, R., Becagli, S., Benassai, S., Castellano, E., Fattori, I., Innocenti, M., Migliori, A., Traversi, R.: Atmosphere- snow interaction by a comparison between aerosol and uppermost snow layers composition at Dome C (East Antarctica). Ann. Glaciol, 39, 53–61, 2004. Udisti, R., Dayan, U., Becagli, S., Busetto, M., Frosini, D., Legrand, M., Lucarelli, F., Preunkert, S., Severi, M., Traversi, R., and Vitale, V.: Sea spray aerosol in central Antarctica. Present atmospheric behaviour and implications for paleoclimatic reconstructions, Atmos. Environ., 52, 109–120, 2012. Uike, A. G., Longo, K. M., and de Freitas, S. R.: Biomass Burning in South America: Transport Patterns and Impacts, in: Biomass – Detection, Production and Usage, edited by M. D. Matovic, InTech, Rijeka, Croatia, https://doi.org/10.5772/19264, 2011. Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP), Aerosol Sci. Tech., 44, 706–712, https://doi.org/10.1080/02786826.2010.482110, 2010. Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, https://doi.org/10.5194/amt.14-3707-2021, 2021. Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	6	and smoke clouds associated with peat and deforestation fires in Indonesia, J. Geophys. Res., 116, D08207,					
 8 9 10 Udisti, R., Becagli, S., Benassai, S., Castellano, E., Fattori, I., Innocenti, M., Migliori, A., Traversi, R.: Atmosphere- snow interaction by a comparison between aerosol and uppermost snow layers composition at Dome C (East Antarctica). Ann. Glaciol, 39, 53–61, 2004. 13 14 Udisti, R., Dayan, U., Becagli, S., Busetto, M., Frosini, D., Legrand, M., Lucarelli, F., Preunkert, S., Severi, M., Traversi, R., and Vitale, V.: Sea spray aerosol in central Antarctica. Present atmospheric behaviour and implications for paleoclimatic reconstructions, Atmos. Environ., 52, 109–120, 2012. 17 18 Ulke, A. G., Longo, K. M., and de Freitas, S. R.: Biomass Burning in South America: Transport Patterns and Impacts. in: Biomass – Detection, Production and Usage, edited by M. D. Matovic, InTech, Rijeka, Croatia, https://doi.org/10.5772/19264.2011. 23 Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP), Aerosol Sci. Tech., 44, 706–712, https://doi.org/10.1080/02786826.2010.482110, 2010. 25 Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, https://doi.org/10.5194/amt-14-3707-2021, 2021. 28 29 Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	7	doi:10.1029/2010JD015148, 2011.					
 Udisti, R., Becagli, S., Benassai, S., Castellano, E., Fattori, I., Innocenti, M., Migliori, A., Traversi, R.: Atmosphere- snow interaction by a comparison between aerosol and uppermost snow layers composition at Dome C (East Antarctica). Ann. Glaciol, 39, 53–61, 2004. Udisti, R., Dayan, U., Becagli, S., Busetto, M., Frosini, D., Legrand, M., Lucarelli, F., Preunkert, S., Severi, M., Traversi, R., and Vitale, V.: Sea spray aerosol in central Antarctica. Present atmospheric behaviour and implications for paleoclimatic reconstructions, Atmos. Environ., 52, 109–120, 2012. Ulke, A. G., Longo, K. M., and de Freitas, S. R.: Biomass Burning in South America: Transport Patterns and Impacts, in: Biomass – Detection, Production and Usage, edited by M. D. Matovic, InTech, Rijeka, Croatia, https://doi.org/10.5772/19264.2011. Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP), Aerosol Sci. Tech., 44, 706–712, <u>https://doi.org/10.1080/02786826.2010.482110</u>, 2010. Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, <u>https://doi.org/10.5194/amt-14-3707-2021</u>, 2021. Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	8						
 Udisti, R., Becagli, S., Benassai, S., Castellano, E., Fattori, I., Innocenti, M., Migliori, A., Traversi, R.: Atmosphere- snow interaction by a comparison between aerosol and uppermost snow layers composition at Dome C (East Antarctica). Ann. Glaciol, 39, 53–61, 2004. Udisti, R., Dayan, U., Becagli, S., Busetto, M., Frosini, D., Legrand, M., Lucarelli, F., Preunkert, S., Severi, M., Traversi, R., and Vitale, V.: Sea spray aerosol in central Antarctica. Present atmospheric behaviour and implications for paleoclimatic reconstructions, Atmos. Environ., 52, 109–120, 2012. Ulke, A. G., Longo, K. M., and de Freitas, S. R.: Biomass Burning in South America: Transport Patterns and Impacts, in: Biomass – Detection, Production and Usage, edited by M. D. Matovic, InTech, Rijeka, Croatia, https://doi.org/10.5772/19264, 2011. Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP), Aerosol Sci. Tech., 44, 706–712, https://doi.org/10.572/19264, 2011. Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, https://doi.org/10.5194/amt-14-3707-2021, 2021. Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	9						
 snow interaction by a comparison between aerosol and uppermost snow layers composition at Dome C (East Antarctica). Ann. Glaciol, 39, 53–61, 2004. Udisti, R., Dayan, U., Becagli, S., Busetto, M., Frosini, D., Legrand, M., Lucarelli, F., Preunkert, S., Severi, M., Traversi, R., and Vitale, V.: Sea spray aerosol in central Antarctica. Present atmospheric behaviour and implications for paleoclimatic reconstructions, Atmos. Environ., 52, 109–120, 2012. Ulke, A. G., Longo, K. M., and de Freitas, S. R.: Biomass Burning in South America: Transport Patterns and Impacts, in: Biomass – Detection, Production and Usage, edited by M. D. Matovic, InTech, Rijeka, Croatia, https://doi.org/10.5772/19264, 2011. Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP), Aerosol Sci. Tech., 44, 706–712, https://doi.org/10.1080/02786826.2010.482110, 2010. Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, https://doi.org/10.5194/amt-14-3707-2021, 2021. Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	10	Udisti, R., Becagli, S., Benassai, S., Castellano, E., Fattori, I., Innocenti, M., Migliori, A., Traversi, R.: Atmosphere-					
 Antarctica). Ann. Glaciol, 39, 53–61, 2004. Udisti, R., Dayan, U., Becagli, S., Busetto, M., Frosini, D., Legrand, M., Lucarelli, F., Preunkert, S., Severi, M., Traversi, R., and Vitale, V.: Sea spray aerosol in central Antarctica. Present atmospheric behaviour and implications for paleoclimatic reconstructions, Atmos. Environ., 52, 109–120, 2012. Ulke, A. G., Longo, K. M., and de Freitas, S. R.: Biomass Burning in South America: Transport Patterns and Impacts, in: Biomass – Detection, Production and Usage, edited by M. D. Matovic, InTech, Rijeka, Croatia, https://doi.org/10.5772/19264, 2011. Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP), Aerosol Sci. Tech., 44, 706–712, https://doi.org/10.1080/02786826.2010.482110, 2010. Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, https://doi.org/10.5194/amt-14-3707-2021, 2021. Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	11	snow interaction by a comparison between aerosol and uppermost snow layers composition at Dome C (East					
 Udisti, R., Dayan, U., Becagli, S., Busetto, M., Frosini, D., Legrand, M., Lucarelli, F., Preunkert, S., Severi, M., Traversi, R., and Vitale, V.: Sea spray aerosol in central Antarctica. Present atmospheric behaviour and implications for paleoclimatic reconstructions, Atmos. Environ., 52, 109–120, 2012. Ulke, A. G., Longo, K. M., and de Freitas, S. R.: Biomass Burning in South America: Transport Patterns and Impacts, in: Biomass – Detection, Production and Usage, edited by M. D. Matovic, InTech, Rijeka, Croatia, https://doi.org/10.5772/19264, 2011. Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP), Aerosol Sci. Tech., 44, 706–712, https://doi.org/10.1080/02786826.2010.482110, 2010. Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, https://doi.org/10.5194/amt-14-3707-2021, 2021. Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	12	Antarctica). Ann. Glaciol, 39, 53–61, 2004.					
 Udisti, R., Dayan, U., Becagli, S., Busetto, M., Frosini, D., Legrand, M., Lucarelli, F., Preunkert, S., Severi, M., Traversi, R., and Vitale, V.: Sea spray aerosol in central Antarctica. Present atmospheric behaviour and implications for paleoclimatic reconstructions, Atmos. Environ., 52, 109–120, 2012. Ulke, A. G., Longo, K. M., and de Freitas, S. R.: Biomass Burning in South America: Transport Patterns and Impacts, in: Biomass – Detection, Production and Usage, edited by M. D. Matovic, InTech, Rijeka, Croatia, https://doi.org/10.5772/19264, 2011. Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP), Aerosol Sci. Tech., 44, 706–712, https://doi.org/10.1080/027868265.2010.482110, 2010. Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, https://doi.org/10.5194/amt-14-3707-2021, 2021. Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	13						
 15 Traversi, R., and Vitale, V.: Sea spray aerosol in central Antarctica. Present atmospheric behaviour and implications for paleoclimatic reconstructions, Atmos. Environ., 52, 109–120, 2012. 17 18 Ulke, A. G., Longo, K. M., and de Freitas, S. R.: Biomass Burning in South America: Transport Patterns and 19 Impacts, in: Biomass – Detection, Production and Usage, edited by M. D. Matovic, InTech, Rijeka, Croatia, 20 https://doi.org/10.5772/19264, 2011. 21 23 Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP), 24 Aerosol Sci. Tech., 44, 706–712, https://doi.org/10.1080/02786826.2010.482110, 2010. 25 26 Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, 27 Atmos. Meas. Tech., 14, 3707–3719, https://doi.org/10.5194/amt-14-3707-2021, 2021. 28 29 Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: 30 Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	14	Udisti, R., Dayan, U., Becagli, S., Busetto, M., Frosini, D., Legrand, M., Lucarelli, F., Preunkert, S., Severi, M.,					
 implications for paleoclimatic reconstructions, Atmos. Environ., 52, 109–120, 2012. Ulke, A. G., Longo, K. M., and de Freitas, S. R.: Biomass Burning in South America: Transport Patterns and Impacts, in: Biomass – Detection, Production and Usage, edited by M. D. Matovic, InTech, Rijeka, Croatia, https://doi.org/10.5772/19264, 2011. Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP), Aerosol Sci. Tech., 44, 706–712, https://doi.org/10.1080/02786826.2010.482110, 2010. Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, https://doi.org/10.5194/amt-14-3707-2021, 2021. Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	15	Traversi, R., and Vitale, V.: Sea spray aerosol in central Antarctica. Present atmospheric behaviour and					
 Ulke, A. G., Longo, K. M., and de Freitas, S. R.: Biomass Burning in South America: Transport Patterns and Impacts, in: Biomass – Detection, Production and Usage, edited by M. D. Matovic, InTech, Rijeka, Croatia, https://doi.org/10.5772/19264, 2011. 	16	implications for paleoclimatic reconstructions, Atmos. Environ., 52, 109–120, 2012.					
 18 Ulke, A. G., Longo, K. M., and de Freitas, S. R.: Biomass Burning in South America: Transport Patterns and Impacts, in: Biomass – Detection, Production and Usage, edited by M. D. Matovic, InTech, Rijeka, Croatia, https://doi.org/10.5772/19264, 2011. 21	17						
 Impacts, in: Biomass – Detection, Production and Usage, edited by M. D. Matovic, InTech, Rijeka, Croatia, https://doi.org/10.5772/19264, 2011. Kormatted: English Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP), Aerosol Sci. Tech., 44, 706–712, https://doi.org/10.1080/02786826.2010.482110, 2010. Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, https://doi.org/10.5194/amt-14-3707-2021, 2021. Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	18	Ulke, A. G., Longo, K. M., and de Freitas, S. R.: Biomass Burning in South America: Transport Patterns and					
 https://doi.org/10.5772/19264, 2011. Formatted: English Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP), Aerosol Sci. Tech., 44, 706–712, https://doi.org/10.1080/02786826.2010.482110, 2010. Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, https://doi.org/10.5194/amt-14-3707-2021, 2021. Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	19	Impacts, in: Biomass – Detection, Production and Usage, edited by M. D. Matovic, InTech, Rijeka, Croatia,					
 Formatted: English Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP), Aerosol Sci. Tech., 44, 706–712, https://doi.org/10.1080/02786826.2010.482110, 2010. Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, https://doi.org/10.5194/amt-14-3707-2021, 2021. Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	20	https://doi.org/10.5772/19264, 2011.					
 Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP), Aerosol Sci. Tech., 44, 706–712, https://doi.org/10.1080/02786826.2010.482110, 2010. Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, https://doi.org/10.5194/amt-14-3707-2021, 2021. Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	21	A	 For	matteo	l: Englisł	n (United	States)
 Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP), Aerosol Sci. Tech., 44, 706–712, <u>https://doi.org/10.1080/02786826.2010.482110</u>, 2010. Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, <u>https://doi.org/10.5194/amt-14-3707-2021</u>, 2021. Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	22						
 Aerosol Sci. Tech., 44, 706–712, https://doi.org/10.1080/02786826.2010.482110, 2010. Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, https://doi.org/10.5194/amt-14-3707-2021, 2021. Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	23	Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP),					
 Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, <u>https://doi.org/10.5194/amt-14-3707-2021</u>, 2021. Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	24	Aerosol Sci. Tech., 44, 706–712, <u>https://doi.org/10.1080/02786826.2010.482110</u> , 2010.					
 Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, Atmos. Meas. Tech., 14, 3707–3719, https://doi.org/10.5194/amt-14-3707-2021, 2021. Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	25						
 Atmos. Meas. Tech., 14, 3707–3719, <u>https://doi.org/10.5194/amt-14-3707-2021</u>, 2021. Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	26	Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell,					
 Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	27	Atmos. Meas. Tech., 14, 3707–3719, <u>https://doi.org/10.5194/amt-14-3707-2021</u> , 2021.					
 Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.: Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	28						
 Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 	29	Virkkula, A., Ahlquist, N. C., Covert, D. S., Arnott, W. P., Sheridan, P. J., Quinn, P. K., and Coffman, D. J.:					
31 Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005. 32	30	Modification, calibration and a field test of an instrument for measuring light absorption by particles, Aerosol					
32	31	Sci. Tech., 39, 68–83, https://doi.org/10.1080/027868290901963, 2005.					
	32						

1	Virkkula, A., Koponen, I. K., Teinilä, K., Hillamo, R., Kerminen, VM., and Kulmala, M.: Effective real refractive	
2	index of dry aerosols in the Antarctic boundary layer, Geophys. Res. Lett., 33, L06805,	
3	https://doi.org/10.1029/2005GL024602, 2006.	
4		
5	Voshchinnikov, N.V. and Mathis, J.S.: Calculating Cross Sections of Composite Interstellar Grains, Astrophys. J.,	
6	526, 257–264, <u>https://doi.org/10.1086/307997</u> , 1999.	
7		
8	Waugh, D. W., Sobel, A. H., and Polvani, L. M.: What is the polar vortex and how does it influence weather, B.	 Formatted: Highlight
9	Am. Meteorol. Soc., 98, 37–44, https://doi.org/10.1175/BAMS-D-15-00212.1, 2017.	
10		
11	Weller, R., Minikin, A., Petzold, A., Wagenbach, D., and König-Langlo, G.: Characterization of long-term and	
12	seasonal variations of black carbon (BC) concentrations at Neumayer, Antarctica, Atmos. Chem. Phys., 13,	
13	1579–1590, https://doi.org/10.5194/acp-13-1579-2013, 2013.	
14		
15	Weller, R., Schmidt, K., Teinilä, K., and Hillamo, R.: Natural new particle formation at the coastal Antarctic	
16	site Neumayer, Atmos. Chem. Phys., 15, 11399–11410, doi:10.5194/acp-15-11399-2015, 2015.	
17		
18	Williams, J., de Reus, M., Krejci, R., Fischer, H., and Ström, J.: Application of the variability-size relationship to	 Formatted: Highlight
19	atmospheric aerosol studies: estimating aerosol lifetimes and ages, Atmos. Chem. Phys., 2, 133–145, 2002,	
20	http://www.atmos-chem-phys.net/2/133/2002/.	
21		
22	WMO/GAW Report No. 227: Aerosol Measurement Procedures, Guidelines and Recommendations, 2nd	
23	Edn., 103 pp., August2016 (WMO-No. 1177), Geneva, Switzerland, 2016.	
24		
25	Wu Z.P., Wang Y.P.: Electromagnetic scattering for multilayered spheres: recursive algorithms, Radio Science,	
26	26, 1393-1401, 1991 <u>.</u>	
27		
28	Xu, H., Ren, Y., Zhang, W., Meng, W., Yun, X., Yu, X., Li, J., Zhang, Y., Shen, G., Ma, J., Li, B., Cheng, H., Wang, X.,	Formatted: Highlight
29	Wan, Y, and Tao, S.: Updated Global Black Carbon Emissions from 1960 to 2017: Improvements, Trends, and	
30	Drivers, Environmental Science and Technology, 55 (12), 7869-7879, DOI: 10.1021/acs.est.1c0311, 2021	
31		

1 Tables

2 Table 1. Seasonal variation of the fractions of volume concentration and scattering coefficients in the size range

3 measured by the DMPS of the respective values calculated from the combined size distributions measured with

4 the DMPS and the OPC at Dome C in December 2007 – July 2009.

	fV(DMPS), %		%	fσ _{sp} (DMPS,46	7), %	fσ _{sp} (DMPS,53	0), %	fσ _{sp} (DMPS,660), %		
	month	average ± std	median	average ± std	median	average ± std	median	average ± std	median	
	JAN	86.8 ± 10.7	91.6	85.2 ± 9.9	88.7	78.5 ± 12.6	82.5	66.1 ± 15.8	69.6	
	FEB	85.7 ± 14.0	90.6	82.8 ± 14.3	87.2	76.1 ± 15.9	81.1	64.0 ± 17.5	68.6	
	MAR	79.0 ± 20.9	88.3	77.0 ± 20.2	85.0	70.2 ± 21.6	78.2	58.6 ± 22.2	65.2	
	APR	72.8 ± 22.7	83.2	73.6 ± 20.4	80.5	66.4 ± 22.0	72.9	55.0 ± 23.0	59.4	
	MAY	55.4 ± 24.3	53.1	62.5 ± 20.9	63.9	54.5 ± 22.0	54.6	42.9 ± 22.1	40.7	
	JUN	49.8 ± 18.3	51.2	61.7 ± 14.6	64.5	52.7 ± 14.6	54.9	39.8 ± 13.2	41.1	
	JUL	49.5 ± 17.9	50.2	62.2 ± 15.1	65.3	53.4 ± 15.2	55.7	40.8 ± 14.2	41.9	
	AUG	54.4 ± 15.5	56.0	68.3 ± 12.2	69.9	60.0 ± 13.0	60.7	47.4 ± 13.5	46.7	
	SEP	62.6 ± 14.6	64.4	73.8 ± 9.8	74.6	66.1 ± 11.4	66.4	54.0 ± 13.3	53.4	
	OCT	64.6 ± 14.5	66.3	74.1 ± 10.5	74.4	66.4 ± 12.2	66.4	54.2 ± 14.0	53.5	
	NOV	74.7 ± 13.7	79.1	77.4 ± 10.5	80.2	69.4 ± 12.0	72.0	56.0 ± 13.2	57.7	
	DEC	80.4 ± 14.4	84.1	80.1 ± 10.9	83.4	72.1 ± 12.1	75.4	57.7 ± 12.4	60.5	
5	year	70.5 ± 21.8	76.7	74.3 ± 17.0	78.4	66.7 ± 18.4	70.3	54.3 ± 19.1	56.2	
		fV(DMPS), %		fosp(DMPS,46	7), %	fosp(DMPS,53	0), %	fosp(DMPS,66	60), %	
	month	average ± std	median	average ± std	median	average ± std	median	average ± std	median	
	JAN	85.7 ± 11.4	91.2	84.2 ± 10.4	88.3	77.3 ± 13.1	81.8	64.6 ± 16.3	68.7	
	FEB	85.1 ± 14.3	90.5	82.4 ± 14.6	87.1	75.6 ± 16.2	81.0	63.4 ± 17.8	68.4	
	MAR	78.9 ± 21.2	88.7	76.8 ± 20.7	85.6	70.0 ± 22.1	78.9	58.7 ± 22.7	66.3	
	APR	72.3 ± 23.5	83.8	72.7 ± 21.7	80.9	65.6 ± 23.2	73.1	54.5 ± 24.0	59.6	
	MAY	53.8 ± 24.8	48.9	60.5 ± 22.0	61.8	52.6 ± 23.0	51.9	41.4 ± 23.0	37.9	
	JUN	48.2 ± 18.3	48.7	59.5 ± 15.8	62.7	50.6 ± 15.5	53.0	38.0 ± 13.8	39.4	
	JUL	48.4 ± 18.3	47.9	60.6 ± 16.0	63.5	52.0 ± 15.9	54.2	39.6 ± 14.7	40.5	
	AUG	53.5 ± 15.5	55.3	66.3 ± 14.7	69.5	58.1 ± 14.6	60.4	45.7 ± 14.0	46.4	
	SEP	61.9 ± 14.3	64.1	72.2 ± 13.3	73.8	64.5 ± 13.6	65.9	52.4 ± 14.0	52.8	
	OCT	63.7 ± 14.2	65.5	72.5 ± 13.2	73.8	64.9 ± 13.7	65.8	52.6 ± 14.2	52.7	
	NOV	74.0 ± 13.2	77.0	76.9 ± 10.1	79.0	68.8 ± 11.5	70.8	55.2 ± 12.6	56.4	
	DEC	80.3 ± 13.6	84.0	79.9 ± 10.3	82.9	71.7 ± 11.4	74.9	57.2 ± 11.9	59.8	

7 8

9

Table 2. Noise of scattering and absorption coefficients calculated from the particle number size distributions

and the PSAP data. $\sigma_{sp}(DMPS,PM_{0.8})$ and $\sigma_{sp}(DMPS,PM_{10})$. Noise was estimated as explained in section 2.5.

	Average n								
Scattering	$\delta\sigma_s$	_{ep} (DMPS,PN	Л _{о.8})	δσ	_p (DMPS,PI	M ₁₀)			
∆t \ λ(nm)	467	530	660	467	530	660			
1 h	5.2×10 ⁻²	4.2×10 ⁻²	2.7×10 ⁻²	7.2×10 ⁻²	6.5×10 ⁻²	5.3×10 ⁻²			
24 h	1.1×10 ⁻²	0.85×10 ⁻²	0.55×10 ⁻²	1.5×10 ⁻²	1.3×10 ⁻²	1.1×10 ⁻²			
	Average n	oise of PSA	P data scatt	ering correction	on, Mm ⁻¹				
	0.0164	×δσ _{sp} (DMP\$	S,PM _{0.8})	0.0164	0.0164×δσ _{sp} (DMPS,PM ₁₀)				
1 h	0.85×10 ⁻³	0.68×10 ⁻³	0.44×10 ⁻³	1.2×10 ⁻³	1.1×10 ⁻³	0.87×10 ⁻³			
24 h	0.17×10 ⁻³	0.14×10 ⁻³	0.091×10 ⁻³	0.24×10 ⁻³	0.22×10 ⁻³	0.18×10 ⁻³			
Absorption $\delta \sigma_{ap,nsc}$			δα	$\delta\sigma_{ap}(\sigma_{sp}(PM_{0.8}))$			δα	$\delta\sigma_{ap}(\sigma_{sp}(PM_{1}$	
Δt \ λ(nm)	467	530	660	467	530	660		467	467 530
1 h	3.2×10 ⁻³	2.5×10 ⁻³	2.7×10 ⁻³	4.0×10 ⁻³	3.2×10 ⁻³	3.2×10 ⁻³		4.3×10 ⁻³	4.3×10 ⁻³ 3.6×10 ⁻³
24 h	0.65×10 ⁻³	0.52×10 ⁻³	0.56×10 ⁻³	0.82×10 ⁻³	0.66×10 ⁻³	0.65×10 ⁻³		0.89×10 ⁻³	0.89×10 ⁻³ 0.74×10 ⁻³

Table 3. Noise and detection limits of eBC concentration calculated from the noise of the absorption

3 coefficients presented in Table 2.

		eBC noise, ng m ⁻³	
	$\delta(eBC(\sigma_{ap,nsc}))$	$\delta(eBC(\sigma_{ap}(\sigma_{sp}(DMPS,PM_{0.8}))))$	$\delta(eBC(\sigma_{ap}(\sigma_{sp}(DMPS,PM_{10}))))$
1 h	0.327	0.415	0.464
24 h	0.067	0.085	0.100
		eBC detection limits, ng m ⁻³	
	$2\delta(eBC(\sigma_{ap,nsc}))$	$2\delta(eBC(\sigma_{ap}(\sigma_{sp}(DMPS,PM_{0.8}))))$	$2\delta(eBC(\sigma_{ap}(\sigma_{sp}(DMPS,PM_{10}))))$
1 h	0.65	0.83	0.93
24 h	0.13	0.17	0.20

1 Table 4. Descriptive statistics of aerosol optical properties at Dome C in 2008 – 2013.

Percentiles								
	λ (nm)	Ν	Ave ± std	5	25	50	75	95
		E:	tensive AOPs					
	Scattering co	pefficient, σ_s	_p , Mm⁻¹					
Filter samples								
$\sigma_{sp}(PM_{10})$	550	1765	0.31 ± 0.31	0.058	0.139	0.24	0.38	0.75
$\sigma_{sp}(PM_1)$	550	468	0.30 ± 0.23	0.046	0.118	0.24	0.41	0.75
DMPS-data								
σ _{sp} (DMPS, PM _{0.8}))	467	21987	0.35 ± 0.64	0.034	0.095	0.19	0.40	1.06
	530	21987	0.27 ± 0.51	0.026	0.073	0.15	0.30	0.80
	660	21987	0.16 ± 0.32	0.016	0.044	0.09	0.18	0.46
$\sigma_{sp}(\text{DMPS}, \text{PM}_{10}))$	467	21987	0.48 ± 0.96	0.052	0.138	0.26	0.52	1.34
	530	21987	0.41 ± 0.88	0.046	0.120	0.22	0.44	1.12
	660	21987	0.31 ± 0.73	0.034	0.089	0.16	0.31	0.78
	Absorption c	oefficient, σ	_{ap} , Mm ⁻¹					
	σ_{ap} , calcul	ated using I	 31999					
$\sigma_{ap}(\sigma_{sp}(DMPS,PM_{10}))$	467	15778	0.015 ± 0.018	< 0.001	0.0029	0.0096	0.021	0.051
	530	15815	0.013 ± 0.015	< 0.001	0.0029	0.0085	0.019	0.044
	660	15657	0.012 ± 0.012	< 0.001	0.0033	0.0077	0.016	0.036
	σ_{ap} , calcul	ated using V	/2010					
$\sigma_{ap}(\sigma_{sp}(DMPS,PM_{10}))$	467	15778	0.016 ± 0.018	< 0.001	0.0033	0.0099	0.022	0.054
	530	15815	0.012 ± 0.015	< 0.001	0.0023	0.0076	0.017	0.041
	660	15657	0.009 ± 0.011	< 0.001	0.0016	0.0057	0.013	0.031
		In	ensive AOPs					
	Single-scatte	ering albedo	$\omega_{o} = \sigma_{sp}(DMPS,PM)$	10)/(σ _{sp} (DMPS	S,PM ₁₀) + c	_{ap})		
	σ	ap calculated	d using B1999					
$\omega_{o}(\sigma_{ap,nsc})$	530	15815	0.93 ± 0.05	0.84	0.91	0.94	0.97	0.99
$\omega_o(\sigma_{sp}(DMPS,PM_{10}))$	530	15815	0.95 ± 0.05	0.85	0.92	0.96	0.99	1.00
	σ	ap calculated	d using V2010					
$\omega_{o}(\sigma_{sp}(\text{DMPS},\text{PM}_{10}))$	530	15815	0.95 ± 0.05	0.86	0.93	0.96	0.99	1.00
	Scattering År	ngström exp	ponent α_{so} (λ =467/6	60 nm)				
$\alpha_{sp}(\sigma_{sp}(DMPS,PM_{0.8}))$	467/660	21987	2.31 ± 0.28	1.90	2.10	2.25	2.54	2.79
$\alpha_{\text{sp}}(\sigma_{\text{sp}}(\text{DMPS},\text{PM}_{10}))$	467/660	21987	1.35 ± 0.40	0.77	1.05	1.30	1.66	2.04
	Absorption Å	ngström ex	oonent (λ=467/660	nm) α_{abs} for	σ _{ap} > 3 δσ _a	D		
	σ	ap calculated	d using B1999			•		
$\alpha_{ap}(\sigma_{ap,nsc})$	467/660	15607	0.86 ± 0.34	0.29	0.67	0.87	1.02	1.37
$\alpha_{ap}(\sigma_{sp}(DMPS,PM_{10}))$	467/660	11475	0.76 ± 0.47	< 0	0.56	0.82	0.99	1.40
	σ	ap calculated	d using V2010					
$\alpha_{ap}(\sigma_{sp}(DMPS,PM_{10}))$	467/660	10270	1.37 ± 0.38	0.72	1.17	1.37	1.59	2.03

Percentiles								
	λ (nm)	N	Ave ± std	5	25	50	75	95
		E	xtensive AOPs					
	Scattering co	pefficient, σ	_{sp} ,Mm⁻¹					
Filter samples								
$\sigma_{sp}(PM_{10})$	550	1765	0.31 ± 0.31	0.058	0.139	0.24	0.38	0.75
σ _{sp} (PM ₁)	550	468	0.30 ± 0.23	0.046	0.118	0.24	0.41	0.75
DMPS-data	407	04007	0.05 0.04	0.004	0.005	0.40	0.40	4.00
$\sigma_{sp}(DMPS, PM_{0.8}))$	467	21987	0.35 ± 0.64	0.034	0.095	0.19	0.40	1.06
	530 660	21987	0.27 ± 0.51 0.16 ± 0.32	0.026	0.073	0.15	0.30	0.80
	000	21307	0.10 ± 0.52	0.010	0.044	0.05	0.10	0.40
σ₅ (DMPS, PM₁0))	467	21987	0.49 ± 0.98	0.054	0.141	0.27	0.52	1.36
φ(, , , , , , , , , , , , , , , , , , ,	530	21987	0.42 ± 0.90	0.047	0.123	0.23	0.44	1.13
	660	21987	0.31 ± 0.75	0.035	0.091	0.16	0.32	0.80
			1					
	Absorption c	oefficient, σ	ap, Mm ⁻ '					
	σ_{ap} , calcul	ated using	B1999					
$\sigma_{ap}(\sigma_{sp}(DMPS,PM_{10}))$	467	15778	0.015 ± 0.017	< 0.001	0.0028	0.0095	0.021	0.051
	530	15815	0.013 ± 0.015	< 0.001	0.0028	0.0085	0.019	0.044
	000	10001	0.011 ± 0.012	< 0.001	0.0032	0.0076	0.016	0.036
((51,50,514,))	σ _{ap} , calcul	ated using	V2010					
$\sigma_{ap}(\sigma_{sp}(DMPS,PM_{10}))$	467	15778	0.016 ± 0.018	< 0.001	0.0032	0.0099	0.022	0.053
	530 660	15815 15657	0.012 ± 0.015 0.009 ± 0.011	< 0.001	0.0022	0.0076	0.017	0.041
	000	15057	0.000 ± 0.011	< 0.001	0.0010	0.0000	0.015	0.000
		In	tensive AOPs					
	Single-scatte	ering albedo	$\omega_{o} = \sigma_{sp}(\text{DMPS},\text{PM})$	10)/(σ _{sp} (DMPS	S,PM ₁₀) + c	s _{ap})		
	σ	ap calculate	d using B1999					
$\omega_{o}(\sigma_{ap,nsc})$	530	15815	0.933 ± 0.048	0.845	0.909	0.944	0.972	0.985
$ω_o(σ_{so}(DMPS,PM_{10}))$	530	15815	0.948 ± 0.049	0.857	0.923	0.959	0.988	1.000
·	σ	an calculate	d using V2010					
$ω_0(\sigma_{en}(DMPS,PM_{10}))$	530	15815	0.952 ± 0.047	0.864	0.928	0.963	0.990	1.000
0 (apr) 10//								
	Scattering År	ngström exp	ponent α_{sp} (λ =467/6	60 nm)				
$\alpha_{so}(\sigma_{so}(DMPS,PM_{0.8}))$	467/660	21987	2.31 ± 0.28	1.90	2.10	2.25	2.54	2.79
$\alpha_{so}(\sigma_{so}(DMPS,PM_{10}))$	467/660	21987	1.34 ± 0.40	0.75	1.03	1.28	1.65	2.03
φ(φ())))								
	Absorption Å	ngström ex	ponent (λ=467/660	nm) α_{abs} for	$σ_{ap} > 3 \delta σ_a$	р		
	σ	ap calculate	d using B1999					
$\alpha_{an}(\sigma_{an}, \infty)$	467/660	15607	0.86 ± 0.34	0.29	0.67	0.87	1.02	1.37
$\alpha_{ab}(\sigma_{ab}(DMPS,PM_{ab}))$	467/660	11475	0.76 ± 0.47	< 0	0.56	0.82	0.99	1.41
······································		calculate	d using V2010		5.00	5.02	2.00	
	467/660	10270	1 38 + 0 30	0.73	1 1 2	1 38	1.60	2.03
ap(0sp(Divir 0, riving))	+07/000	10270	1.50 ± 0.58	0.73	1.10	1.00	1.00	2.03

Table 5. Statistical summary of mass concentrations estimated from particle number size distributions, sums of 1 2 ion concentrations of PM₁ and PM₁₀ filter samples and the PSAP data at Dome C in 2008 – 2013. The statistical 3 values of the PM_1 and PM_{10} are those of all individual filters, the statistical values calculated from the DMPS 4 and PSAP data are those of running 24h-averaged data, see details in the text. m(DMPS,PM0.8): mass 5 concentration calculated from the particle number size distributions measured with the DMPS assuming 6 particle density of 1.7 g cm⁻³; eBC: equivalent Black Carbon concentration calculated from the absorption 7 coefficients at λ =530 nm calculated by using the B1999 algorithm without any scattering corrections and with 8 B1999 and V2010 algorithms using $\sigma_{sp} = \sigma_{sp}(DMPS, PM_{10})$ for the scattering corrections and assuming MAC = 9 7.78 m² g⁻¹. fPM₁₀: scattering-corrected eBC mass fraction calculated from eBC/m(DMPS,PM₁₀)×100%; fPM_{0.8}: 10 scattering-corrected eBC mass fraction calculated from eBC/m(DMPS,PM_{0.8})×100%.

				Pe	ercentiles		
	Ν	Ave ± std	5	25	50	75	95
Mass concentrations							
PM ₁₀ filters, ng m ⁻³	1765	162 ± 161	30	73	126	201	394
PM ₁ filters, ng m ⁻³	468	82 ± 64	13	33	66	114	208
m(DMPS, PM _{0.8}), ng m ⁻³	21987	123 ± 161	12	31	70	150	382
m(DMPS, PM_{10}), ng m ⁻³		171 ± 255	23	54	108	199	483
eBC from $\sigma_{ap}(\lambda = 530$) nm) calcula	ated using B1999					
eBC(no σ_{sp} correction), ng m ⁻³	15815	2.6 ± 2.6	0.36	0.94	1.9	3.2	7.6
eBC($\sigma_{sp}(DMPS,PM_{10})))$, ng m ⁻³	15815	1.7 ± 2.0	< 0.2	0.37	1.10	2.4	5.7
feBC(m(DMPS,PM _{0.8})), %	15815	2.6 ± 2.7	< 0.1	0.53	1.62	4.0	7.8
feBC(m(DMPS,PM ₁₀)), %	15815	1.6 ± 1.7	< 0.1	0.35	1.16	2.5	4.7
eBC from $\sigma_{ap}(\lambda = 530)$) nm) calcula	ated using V2010					
$eBC(\sigma_{sp}(DMPS,PM_{10})))$, ng m ⁻³	15815	1.6 ± 1.9	< 0.2	0.29	0.98	2.2	5.3
feBC(m(DMPS,PM _{0.8})), %	15815	2.4 ± 2.6	< 0.1	0.41	1.48	3.7	7.3
feBC(m(DMPS,PM ₁₀)), %	15815	1.5 ± 1.6	< 0.1	0.28	1.06	2.3	4.5
		-		Pe	ercentiles		
	Ν	Ave ± std	5	25	50	75	95
Mass concentrations	1705	100 101		70	100		
PM ₁₀ filters, ng m ⁻³	1765	162 ± 161	30	73	126	201	394
PM ₁ filters, ng m [°]	468	82 ± 64	13	33	66	114	208
m(DMPS, PM _{0.8}), ng m ⁻³	21987	123 ± 161	12	31	70	150	382
m(DMPS, PM ₁₀), ng m ^{-s}	21987	173 ± 260	23	55	110	200	488
eBC from $\sigma_{ap}(\lambda = 530$) nm) calcula	ated using B1999					
eBC(no σ_{sp} correction), ng m ⁻³	15815	2.6 ± 2.6	0.36	0.94	1.86	3.2	7.6
$eBC(\sigma_{sp}(DMPS,PM_{10})))$, ng m ⁻³	15815	1.7 ± 2.0	< 0.2	0.36	1.09	2.4	5.7
feBC(m(DMPS,PM _{0.8})), %	15815	2.6 ± 2.7	< 0.1	0.51	1.60	4.0	7.7
feBC(m(DMPS,PM ₁₀)), %	15815	1.6 ± 1.7	< 0.1	0.34	1.14	2.4	4.7
eBC from $\sigma_{ap}(\lambda = 530)$) nm) calcula	ated using V2010					
$eBC(\sigma_{sp}(DMPS,PM_{10})))$, ng m ⁻³	15815	1.6 ± 1.9	< 0.2	0.28	0.97	2.2	5.3
feBC(m(DMPS,PM _{0.8})), %	15815	2.4 ± 2.6	< 0.1	0.39	1.47	3.7	7.3
feBC(m(DMPS PM)) %	15815	1.5 ± 1.6	< 0.1	0.26	1.04	2.2	4.4

1 Figures

2008 2009	2010	2011	2012	2013	

2

3 Figure 1. The periods of the PSAP, the DMPS, the Grimm OPC and the PM_1 and PM_{10} filter sample data. The

4 number of hours of accepted data and the number of samples are shown in parentheses for the continuous

6

⁵ instruments and the filter samplers, respectively.



2

3 Figure 2. Average particle size distributions in summer (DJF) and in winter (JJA) in December 2007 – July 2009 4 when both the DMPS and the Grimm OPC were operational. Left: average and median (a) number, (c) volume, 5 and (e) scattering size distributions at λ =530 nm; right (b, d, and f): cumulative fractions of the respective



1 parameters in the size range $D_p < 7.5 \ \mu$ m. which corresponds to the aerodynamic particle size range $D_a < 9.8$

Figure 3. Seasonal cycle of the contribution of the size range measured by the DMPS to a) volume
concentration and b) scattering coefficient at the PSAP wavelengths in December 2007 – July 2009 when both
the DMPS and the Grimm OPC were operational. The circle shows the average, the horizontal line the median,
the box the 25th to 75th percentile range, and the whiskers the 5th to 95th percentile range in each month.

9

2

μm.



1

Figure 4. Wind and absorption coefficient. a) Hourly-averaged non-scattering-corrected absorption coefficients ($\sigma_{ap,nsc}$, Eq. (1)) observed at wind speed WS > 2 m/s in 5° wind direction (WD) sectors. The lines present the percentiles of the cumulative $\sigma_{ap,nsc}$ distribution in each WD sector. f(WD sector): fraction of wind data from each sector. CS 75th perc.: Contamination sector determined from the 75th percentiles of the cumulative $\sigma_{ap,nsc}$ distribution. CS 99th perc.: Contamination sector determined from the 99th percentiles of the cumulative $\sigma_{ap,nsc}$ distribution. b) Distribution of WS and WD as a wind rose.



Figure 5. Time series of scattering coefficients calculated from the DMPS ($\sigma_{sp}(DMPS,PM_{10})$ at $\lambda = 530$), PM₁ and PM₁₀ ($\lambda = 550$) filter data measured at Dome C and measured with the nephelometer at the South Pole Station (SPO) ($\lambda = 550$) and $\sigma_{ap,nsc}(\lambda = 530)$ measured with the PSAP at Dome C. The σ_{sp} from the DMPS and the nephelometer and $\sigma_{ap,nsc}$ are running 24-hour averages at each hour (± 12 hours), and the σ_{sp} from the PM1 and PM10 filters are those calculated for each filter. The red box within the 2011 time series shows the period presented in more detail in Fig. 6 and the red asterisk symbol (*) for which the footprint in Fig. 7 was calculated.



Figure 6. Four-month time series (May-August 2011) of σ_{sp} , σ_{ap} and ω_o at $\lambda = 530$ nm at Dome C and σ_{sp} at at $\lambda = 550$ nm at the South Pole Station (SPO): a) upper estimate of σ_{sp} (σ_{sp} (DMPS, PM₁₀)), lower estimate of σ_{ap} (= $\sigma_{ap}(\sigma_{sp}(DMPS, PM_{10}))$) and upper estimate of ω_o . b) lower estimate of σ_{sp} (= $\sigma_{sp}(DMPS, PM_{0.8})$)), upper estimate of σ_{ap} (= $\sigma_{ap}(\sigma_{sp}(DMPS, PM_{0.8}))$) and lower estimate of ω_o . In both a) and b) also the non-scattering-corrected absorption coefficient is shown. All values are running 24-hour averages at each hour (± 12 hours). The numbers 1 – 4 are discussed in the text.





4 Figure 7. FLEXPART footprint of the overall highest day of scattering in winter 2011, on Jul 28, 2011, indicated

5 by the number (3) in Fig. 6.



1 2 3

3 Figure 8. Comparison of scattering coefficients calculated from the DMPS vs. scattering coefficients calculated 4 from the PM₁ and PM₁₀ filter sample data at Dome C, all at λ = 550 nm. The scattering coefficients calculated 5 from the DMPS data were averaged for the sampling times of the PM₁ and PM₁₀ samples and interpolated to λ 6 = 550 nm. a) lower estimate of σ_{sp} (= σ_{sp} (DMPS, PM_{0.8})) vs. σ_{sp} (PM₁₀), b) upper estimate of σ_{sp} (= 7 σ_{sp} (DMPS,PM₁₀)) vs. σ_{sp} (PM₁₀), c) σ_{sp} (DMPS, PM_{0.8})) vs. σ_{sp} (PM₁), d) σ_{sp} (DMPS,PM₁₀)) vs. σ_{sp} (PM₁). N: number of 8 data points. The red line shows the linear regression line that is forced through zero. The regression equations 9 show the slope ± standard error of the slope, the squared correlation coefficient and the p value of the slope.



2 Figure 9. Seasonal cycles of scattering and absorption coefficients and single-scattering albedo. a) Scattering 3 coefficient (σ_{sp}) calculated from the sums of analyzed ion concentrations in PM₁ filters at λ = 550 nm, b) σ_{sp} 4 calculated from the sums of analyzed ion concentrations in PM_{10} filters, c) the lower estimate of σ_{sp} = 5 $\sigma_{sp}(DMPS,PM_{0.8})$, d) the upper estimate of $\sigma_{sp} = \sigma_{sp}(DMPS,PM_{10})$, e) absorption coefficient σ_{ap} calculated with 6 the algorithms of B1999 and V2010 (Eqs. (17) and (18)) by using the σ_{sp} lower estimate for scattering 7 correction, f) σ_{ap} calculated with the two algorithms by using the σ_{sp} upper estimate for scattering correction, g) 8 single-scattering albedo ω_o calculated by using the σ_{sp} lower estimate for both σ_{sp} and σ_{ap} and h) ω_o calculated 9 by using the σ_{sp} upper estimate for both σ_{sp} and σ_{ap} .



1

2 Figure 10. Seasonal cycles of the wavelength dependency of a) scattering and b) absorption. In a) the Ångström 3 exponent α_{sp} was calculated from the size distributions measured at Dome C ($\sigma_{sp}(DMPS,PM_{0.8})$ and 4 $\sigma_{sp}(DMPS,PM_{10})$ for the wavelength range 467 – 660 and measured at the South Pole Station with a 5 nephelometer. The SPO α_{sp} was calculated for the wavelength range 550 – 700 nm. In b) the absorption 6 Ångström exponent α_{ap} was calculated for the σ_{ap} without scattering correction, and by using the B1999 and 7 V2010 algorithms with scattering corrected using $\sigma_{sp} = \sigma_{sp}(DMPS,PM_{10})$, In all of them the data with $\sigma_{ap} > 3\delta\sigma_{ap}$ 8 were used.

9


2 Figure 11. Seasonal cycles of a) aerosol mass concentration calculated from the particle number size 3 distributions m(DMPS,PM₁₀) and m(DMPS,PM_{0.8}), the sum of ions analyzed from PM₁ and PM₁₀ filters, b) mass 4 fraction of the sum of $nssSO_4^{2-}$ and MSA in PM₁ and PM₁₀ filters, c) equivalent black carbon (eBC) concentration 5 calculated from the non-scattering-corrected absorption coefficients, and from σ_{ap} corrected with the σ_{sp} upper 6 estimate (σ_{sp} (DMPS,PM₁₀)), d) mass fraction eBC calculated as the ratio of eBC corrected with the σ_{sp} upper 7 estimate to m(DMPS,PM0.8) and m(DMPS,PM10). Comparison values: a) monthly average mass concentration 8 calculated from particle volume concentrations at Troll (F2014: Fiebig et al., 2014), average gravimetric PM₁₀ 9 mass concentration at Dome C (A2011: Annibaldi et al., 2011), c) monthly median eBC concentrations 10 measured at Neumayer (NM) with a MAAP and an Aethalometer (AE) (W2013: Weller et al., 2013), at SPO with 11 an Aethalometer (S2016: Sheridan et al., 2016), and at Syowa with an Aethalometer (H2019: Hara et al., 2019). 12 The small insert in b) shows the scatter plot and linear regression of monthly average α_{sp} (DMPS,PM10) vs. 13 monthly average mass fraction of the sum of $nssSO_4^{2-}$ and MSA in PM₁₀ filters.



Figure 12. Monthly average and median ω_0 vs. eBC mass fraction calculated as the ratio of eBC to a) m(DMPS,PM_{0.8}) and b) m(DMPS,PM₁₀). The dashed lines represent fittings of ω_0 (feBC) = ω_0 (0)exp(-k·feBC) with the data. The continuous lines in b) represent simulations with a core-shell (CS) model for lognormal number size distributions with geometric standard deviation GSD = 1.8 and geometric mean diameter GMD shown in the legend.



2 Figure 13. The residence time of 60000 50-day FLEXPART backtrajectories arriving at Dome C in 2008 3 - 2013 as a function of latitude and altitude above ground level. The residence time is colorcoded 4 from blue to yellow to show increasing time spent in each grid cell. a) The residence times of the inert 5 tracer backtrajectories. White line: the average altitude of trajectories at each latitude. b) Same as the in (a) but for the BC tracer. Black line: the average altitude of the BC tracer, the white line is for 6 7 comparison average altitude of the inert tracer. The black triangle shows the latitude of Dome C 8 (75°06'S). The grey text boxes between a) and b) show the approximate latitude range of some major 9 BC emitting regions.



3 Figure 14. The relative difference RD, Eq. (19), of transport for the two lowest layers (>1000 magl) between the

4 highest and the lowest 10% of a) eBC concentration and b) single-scattering albedo. Positive values indicate a

- 5 relative increase in transport, and negative values a decreased transport from a given area by the highest 10%
- 6 of measurements.
- 7



 $\,$ Figure 15. Seasonal air mass transport by region calculated as the fraction of time the FLEXPART 50-day $\,$

3 trajectories spent over Antarctica, the surrounding oceans and continents.



Figure 16. Seasonal cycles of fire-related BC emissions from the surrounding continents and eBC concentration
at Dome C. a) BC emissions and eBC concentrations, b) RTW BC emissions, c) f(t) × BC emissions 1 month
earlier, d) f(t) × BC emissions 2 months earlier. The numbers show the squared correlation coefficient of linear
regression between eBC concentrations and f(t) × BC emissions with the shown time shifts.



2 Figure 17. Normalized seasonal cycles of a) major natural sources of light-scattering aerosols from the oceans, 3 b) scattering coefficients at Dome C and c) scattering coefficients at SPO. $\sigma_{sp}(DMPS,PM_{0.8})$ and $\sigma_{sp}(DMPS,PM_{10})$: 4 scattering coefficient calculated from particle number size distributions; $\sigma_{sp}(nss-SO_4^{2-},PM_1)$: scattering 5 coefficient calculated from non-seasalt sulfate concentrations in PM_1 filter samples; σ_{sp} (nephelometer, SPO): 6 scattering coefficient measured with the nephelometer at SPO; Cphyto: phytoplankton biomass concentration in 7 2008 - 2013 calculated as monthly averages from the time series presented by Behrenfeld et al. (2016) using 8 CALIOP satellite data in Zone 1 (45° – 55°S), Zone 2 (55° – 65°S), and Zone 3 (65° – 75°S); I_T (50°S), I_T (60°S) and 9 I₁(70°S): average global radiation at latitudes 50°S, 60°S and 70°S at longitude 0° at 10 – 14 UTC; PAR(Zone 3): 10 photosynthetically active radiation from Behrenfeld et al. (2016); SSA: Sea Spray Aerosol flux modeled 11 according to Grythe et al. (2014).

12







- $3 \quad \text{ the } PM_1 \text{ filter samples } \sigma_{sp}(\text{nssSO}_4{}^2,\text{PM}_1) \text{ at } \text{ Dome } C \text{ vs. normalized monthly averages of } C_{phyto}\text{PAR in polar}$
- 4 latitude zone with and without a time shift.
- 5