



Acidity and the multiphase chemistry of atmospheric aqueous particles and clouds

Andreas Tilgner¹, Thomas Schaefer¹, Becky Alexander², Mary Barth³, Jeffrey L. Collett, Jr.⁴,
Kathleen M. Fahey⁵, Athanasios Nenes^{6,7}, Havalala O. T. Pye⁵, Hartmut Herrmann^{1*}, and V. Faye
5 McNeill^{8*}

¹Leibniz Institute for Tropospheric Research (TROPOS), Atmospheric Chemistry Department (ACD), Leipzig, 04318, Germany

²Department of Atmospheric Science, University of Washington, Seattle, WA, 98195, USA

³National Center for Atmospheric Research, Boulder, CO, 80307, USA

10 ⁴Department of Atmospheric Science, Colorado State University, Fort Collins, CO, 80523, USA

⁵Office of Research and Development, US Environmental Protection Agency, Research Triangle Park, NC, 27711, USA

⁶School of Architecture, Civil and Environmental Engineering, Ecole Polytechnique Fédérale de Lausanne, Lausanne, CH-1015, Switzerland

⁷Institute for Chemical Engineering Sciences, Foundation for Research and Technology Hellas, Patras, GR-26504, Greece

15 ⁸Departments of Chemical Engineering and Earth and Environmental Sciences, Columbia University, New York, NY, 10027, USA

Correspondence to: V. Faye McNeill (vfm2103@columbia.edu), H. Herrmann (herrmann@tropos.de)

Abstract.

The acidity of aqueous atmospheric solutions is a key parameter driving both the partitioning of semi-volatile acidic and basic
20 trace gases and their aqueous-phase chemistry. In addition, the acidity of atmospheric aqueous phases, e.g. deliquesced aerosol particles, cloud and fog droplets, is also dictated by aqueous-phase chemistry. These feedbacks between acidity and chemistry have crucial implications for the tropospheric lifetime of air pollutants, atmospheric composition, deposition to terrestrial and oceanic ecosystems, visibility, climate, and human health. Atmospheric research has made substantial progress in understanding feedbacks between acidity and multiphase chemistry during recent decades. This paper reviews the current state
25 of knowledge on these feedbacks with a focus on aerosol and cloud systems, involving both inorganic and organic aqueous-phase chemistry. Here, we describe the impacts of acidity on the phase partitioning of acidic and basic gases and buffering phenomena. Next, we review feedbacks of different acidity regimes on key chemical reaction mechanisms and kinetics, as well as uncertainties and chemical subsystems with incomplete information.

30 Finally, we discuss atmospheric implications and highlight needs for future investigations, particularly with respect to reducing emissions of key acid precursors in a changing world, and needs for advancements of field and laboratory measurements and model tools.



1 Introduction

The acidity of the atmospheric aqueous phase (i.e., deliquesced aerosol particles, cloud and fog droplets) impacts human health, climate, and terrestrial/oceanic ecosystems (e.g., Pye et al. (2020) and references therein). Changes in acidity in these aqueous media can arise due to uptake of acidic or basic gases, coalescence, or chemical reactions in the aqueous phase. In turn, acidity of aerosols influences the phase partitioning of semi-volatile species, particulate matter (e.g., Nenes et al. (2020b)), their deposition rates (e.g., Nenes et al. (2020a)) and the rates and types of their chemical transformations. As a result of this two-way coupling between acidity and chemistry, acidity in atmospheric aqueous-aerosol matrices is controlled not only by thermodynamic equilibrium, but also by mass transfer and chemical reaction kinetics, and emissions. Multiphase oxidation and reduction processes in atmospheric waters are strongly linked to the acidity-dependent uptake of acidic or basic compounds, which in turn affects the phase partitioning and the composition of aerosol particles. Moreover, the acidity level directly impacts chemical transformations, but the acidity itself is also influenced as a consequence of such processes. Figure 1 illustrates important tropospheric chemical processes in aqueous atmospheric matrices that are influenced by acidity and affecting acidity.

The most important source of acidity in aqueous aerosols in the troposphere is the uptake and in-situ formation of strong acids, including sulfuric acid, a classic and important compound connected to anthropogenic pollution. Acid formation in aqueous atmospheric phases is itself influenced by acidity, but, more importantly, it also substantially increases the acidity of those media. Important acidity-influenced chemical processes, such as the conversion of sulfur(IV) to sulfur(VI) (Calvert et al., 1985; Faloon, 2009; Harris et al., 2013; Turnock et al., 2019), as well as acid-driven and acid-catalyzed reactions of organic compounds (McNeill et al., 2012; Herrmann et al., 2015), contribute significantly to both secondary inorganic (SIA) and organic (SOA) aerosol formation. These constituents are often responsible for a large fraction of fine particulate matter (Jimenez et al., 2009). Due to their importance, they are strongly associated with aerosol effects on climate (Charlson et al., 1992; Boucher et al., 2013; Seinfeld et al., 2016; McNeill, 2017), air quality (Fuzzi et al., 2015), visibility (Hyslop, 2009), ecosystems (Keene and Galloway, 1984; Adriano and Johnson, 1989; Baker et al., 2020) and human health (Pöschl, 2005a; Pope and Dockery, 2012; Lelieveld et al., 2015). Therefore, changes in acidity can significantly affect the global impacts of aerosols (Turnock et al., 2019).

Acidity-dependent chemical reactions also modify the tropospheric multiphase oxidant budget. For instance, the activation of halogen radicals is promoted by acidity (see Fig. 1 and Sect. 4.2) and can substantially affect the tropospheric oxidative capacity (Vogt et al., 1996; von Glasow et al., 2002a; Pechtl and von Glasow, 2007; Sherwen et al., 2016; Sherwen et al., 2017; Hoffmann et al., 2019b). Acidity can indirectly affect aerosol and cloud composition by promoting the solubilization of transition metals and other bioavailable nutrients such as phosphorous (Meskhidze et al., 2005; Nenes et al., 2011; Shi et al., 2011; Stockdale et al., 2016). Soluble transition metal ions (TMIs) can initiate enhanced HO_x chemistry in aqueous aerosol particles and clouds or catalyze S(IV) oxidation. Moreover, these solubilized metals, phosphorous and semi-volatile inorganic reactive nitrogen molecules (NH₃, HNO₃) can deposit to the ocean surface, contribute to the bioavailable nutrient budget, and

thus impact biological activity and the carbon cycle. TMI solubilization also influences the impacts of atmospheric aerosols on human health (Fang et al., 2017). On the other hand, chemical interactions of marine and crustal primary aerosol constituents (e.g., carbonates, phosphates, halogens), dissolved weak organic acids, and other weak acids (e.g., HNO_3 , HCl , HONO) and bases (e.g., ammonia and amines) can lead to a buffering of the acidity of aqueous solutions (see Fig. 1; Weber et al. (2016); Song et al. (2019a); and Sect. 2.2 for details).




Figure 1. Schematic of chemical processes influenced by and effects on acidity in tropospheric aerosols.

75 In comparison to other aqueous environments, such as sea water and continental surface waters, which are characterized by rather small acidity variations, atmospheric aqueous environments show much higher diversity (see Pye et al. (2020) for details). This is in part because of the huge concentration range of dissolved species in atmospheric waters, but also due to the decoupled exchange of acidic and basic species between the gas and condensed phases. Due to the technical challenges of



sampling and/or characterizing the pH of aerosols, fogs, and cloud water, there is also comparatively limited data on the acidity
80 of these phases in time and space. A companion article, Pye et al. (2020), provides a more complete overview of literature data
on the acidity of atmospheric waters, which we briefly summarize here: Typical pH values for cloud and fog droplets lie
between 2-7, while pH values for continental and marine aerosol particles have a larger range, -1-5 and 0-8, respectively
(Herrmann et al. (2015); Pye et al. (2020) and references therein). Because of the importance of aerosol and cloud acidity for
atmospheric processes and the environment, acidity has been a key subject of research for three decades. The majority of those
85 studies were focused on clouds, motivated by acid rain as well as SIA formation. A detailed review on observations,
thermodynamic processes and implications of atmospheric acidity is given in the companion paper (Pye et al., 2020).
Here, we review in detail the impact of acidity on the chemical transformations of atmospheric aerosols, clouds, and fog water,
with a focus on aqueous-phase chemical reaction kinetics and mechanisms. We also highlight how chemical reactions control
acidity in atmospheric aqueous media. We first discuss the uptake of acidic and basic gases and buffering phenomena, then
90 describe feedbacks between particle and droplet acidity and inorganic chemical reactions (SO₂ oxidation and halogen
chemistry) and aqueous-phase organic chemistry. Finally, a summary addresses atmospheric implications and needs for future
investigations, for example, in the context of reduced fossil fuel combustion emissions of key acid precursors in a changing
world.

2 Fundamental physical and chemical processes of importance for acidity

95 2.1 Aqueous-phase partitioning of acidic and basic gases

The partitioning of acidic or basic gases to atmospheric aerosols or cloud/fog droplets can have a major influence on
condensed-phase acidity. Likewise, the acidity of the aqueous phase itself influences the partitioning of dissociating species
from the gas phase. Condensed-phase acidity also governs back transfer or evaporation of dissociating compounds into the gas
phase (see Sect. 2.2).

100

The phase partitioning of acids and bases

The partitioning of a compound, between the gas phase, aqueous-phase, and its ionic forms, is usually achieved in < 1 hour
for fine-mode aqueous aerosols and small cloud droplets (Dassios and Pandis, 1998; Ervens et al., 2003; Ip et al., 2009; Koop
et al., 2011). Therefore, equilibrium conditions are often assumed in order to estimate the aqueous-phase concentrations.

105

Exceptions include large droplets with higher pH-values, droplets or particles with surface coatings, viscous aerosol particles,
or highly reactive dissolving compounds where mass transfer limitations in the gas or aqueous phase can prevent the attainment
of equilibrium partitioning on relevant timescales. The assumption of a thermodynamic equilibrium in such a case may result
in model biases (Ervens et al., 2003).



Assuming an ideal aqueous solution at equilibrium, i.e., neglecting, for example, mass transport limitations, chemical
110 production and degradation processes and non-ideal solution effects (i.e., considering the activity of ions in solution equal to
their aqueous concentration), the aqueous-phase concentration of a soluble compound ($[A]_{aq}$) is proportional to the partial
pressure of the compound in the gas-phase ($p_{A(\text{air})}$) and its Henry's law constant H_A . The Henry's law constant (in $\text{mol L}^{-1} \text{atm}^{-1}$)
is defined as:

$$H_A = \frac{[A]_{aq}}{p_{A(\text{air})}} \quad (1)$$

115 Once an acid is taken up into an aqueous solution, it can dissociate into a hydrogen ion (H^+) and anions (A^{z-}), the degree of
which depends on its tendency for dissociation, characterized by an equilibrium dissociation constant K_a , and the acidity of
the aqueous environment. Consequently, an effective Henry's Law constant, H_A^* , e.g. for a diacid, is defined by Eq. 2a. For a
monoprotic acid, the third term in the parenthesis is omitted ($K_{a2} = 0$). For typical atmospheric monoprotic bases, such as NH_3
or dimethylamine, the corresponding effective Henry's Law constant, H_A^* , is defined by Eq. 2b. In Eq. 2b, K_a is the equilibrium
120 dissociation constant K_a of the base cation.

$$H_{A(\text{acid})}^* = H_A \left(1 + \frac{K_{a1}}{[\text{H}^+]} + \frac{K_{a1}K_{a2}}{[\text{H}^+]^2} \right) \quad (2a)$$

$$H_{A(\text{base})}^* = H_A \left(1 + \frac{[\text{H}^+]}{K_a} \right) \quad (2b)$$

Together with the liquid water content (LWC), the acidity of an aqueous solution can substantially affect the partitioning of
dissociating compounds to the aqueous aerosol or cloud phase. Increasing acidity leads to a decrease of the effective
125 partitioning of acids and to an increase in the effective partitioning of bases, and vice versa. For example, the partitioning of
nitrate to the particle phase varies dramatically across the typical range of aerosol pH, with nearly 100% of nitrate existing as
 HNO_3 in the gas phase at pH 1, and near-complete particle-phase partitioning at pH 4. As a result, even small biases in predicted
particle pH in air quality models can result in over- or under-predictions of fine particle mass (Vasilakos et al., 2018). Since
atmospheric waters are typically acidic, bases are predominantly present in their protonated form and their partitioning is not
130 greatly altered by typical variations in pH. Hence, this section mainly focuses on the impact of acidity on the partitioning of
weak acids into aqueous aerosols and cloud/fog droplets.

From Eq. 1 and the ideal gas law, the concentration of the dissociating compound in the gas ($C_{A_{\text{air}}}$) and aqueous ($C_{A_{\text{aq}}}$) phase
with respect to the volume of air can be determined. Moreover, the aqueous-phase fraction of A ($X_{A_{\text{aq}}}$), i.e. the ratio of the
aqueous-phase concentration of compound A and the overall multiphase concentration (sum of A in gas and aqueous phase
135 (including undissociated and dissociated forms of A)) can accordingly be calculated by Eq. 3 (see Seinfeld and Pandis (2006)
for details).

$$X_{A_{\text{aq}}} = \frac{C_{A_{\text{aq}}}}{(C_{A_{\text{aq}}} + C_{A_{\text{air}}})} = \frac{H_A^* \cdot R^* \cdot T \cdot \text{LWC} \cdot 10^{-6}}{1 + H_A^* \cdot R^* \cdot T \cdot \text{LWC} \cdot 10^{-6}} \quad (3)$$

Here, $C_{A_{\text{air}}}$ is the concentration of A in air [$\text{mol L}^{-1}_{\text{air}}$], $C_{A_{\text{aq}}}$ is the aqueous-phase concentration of A in the volume of air
[$\text{mol L}^{-1}_{\text{air}}$], R^* is the universal gas constant [$0.082058 \text{ atm L}_{\text{air}} \text{ mol}^{-1} \text{ K}^{-1}$]; T [K] is the temperature, H_A^* is the effective Henry's




140 Law constant [$\text{mol L}^{-1} \text{water atm}^{-1}$] and LWC is the liquid water content [$\text{g m}^{-3} \text{air}$]. Considering activities instead of concentrations, Eq. 3 modifies to Eq. 3a and 3b for monoprotic acids and bases (see Nenes et al. (2020b) and Guo et al. (2017) for details):

$$X_{aq,acid} = \frac{H_A \cdot K_{a1} \cdot R^* \cdot T \cdot LWC \cdot 10^{-6}}{\gamma_{H^+} \cdot \gamma_{A^-} \cdot [H^+] + H_A \cdot K_{a1} \cdot R^* \cdot T \cdot LWC \cdot 10^{-6}} \quad (3a)$$

$$X_{aq,base} = \frac{H_A \cdot K_{a1} \cdot R^* \cdot T \cdot LWC \cdot 10^{-6}}{1 + \frac{\gamma_{H^+} \cdot H_A \cdot [H^+]}{\gamma_{B^+} \cdot K_{a1}} \cdot R^* \cdot T \cdot LWC \cdot 10^{-6}} \quad (3b)$$

where γ_{H^+} , γ_{A^-} , γ_{B^+} are the single-ion activity coefficients for H^+ , the acid anion (A^-) and the base cation (B^+), respectively, which can be calculated for a known ion composition using thermodynamic models (e.g., ISORROPIA-II (Fountoukis and Nenes, 2007), E-AIM (Clegg and Seinfeld, 2006), AIOMFAC (Zuend et al., 2008)).

Figure 2 displays the aqueous fraction, $X_{A_{aq}}$, of 8 weak atmospheric acids (sulfurous acid, nitrous acid, formic acid, acetic acid, glycolic acid, lactic acid, benzoic acid, phthalic acid, 2-nitrophenol, 2,4-dinitrophenol) and two important atmospheric bases (ammonia and dimethylamine) as a function of the LWC and acidity, calculated by Eq. 3. For the plots, an acidity range ($[H^+] = 10^{-1}$ - $10^{-7} \text{ mol L}^{-1}$) and a liquid water content range (10^{-6} - 1 g m^{-3}) have been considered that represent typical values for tropospheric aqueous aerosols, cloud/fog droplets and haze (see Herrmann et al. (2015)). A temperature of 298 K was assumed. It should be noted that temperature plays an important role on the effective solubility of trace gases. In general, as temperature decreases, the trace gas effective solubility increases. Thus, clouds at the top of the mixing layer height ($\sim 285 \text{ K}$ typically) have higher aqueous fractions than aerosol water near the surface on a hot summer day. Similarly, winter hazes should also have higher aqueous fractions than summertime haze events. Therefore, the aqueous fractions shown in Fig. 2 should be used carefully. Note, the H_A and pK_a values applied for the idealized calculation of LWC and acidity dependent aqueous fraction $X_{A_{aq}}$ are listed in Table S1 in the Supporting Information.



160 **Figure 2.** Calculated aqueous-phase fraction $X_{A_{aq}}$ of 8 selected weak acids (a: SO_2 , b: HONO, c: Formic acid, d: Acetic acid, e: Glycolic acid, f: Phthalic acid, g: 2-nitrophenol, h: 2,4-dinitrophenol) and bases (i: Ammonia, j: Dimethylamine) as a function of the LWC and acidity. The black lines are the isolines of the aqueous fractions of 10^{-i} ($i = 1, \dots, 6$). The dashed white lines indicate pK_a values of the corresponding acids (except for the two bases and for 2-nitrophenol due to the very high pK_a of 7.2 (see Table S1)).



165 Examples in Fig. 2 illustrate that acidity, along with the LWC, strongly influences the phase partitioning of weak acids and
bases into the aqueous phase. The partitioning into the aqueous phase is efficient for pH values well above the individual pK_a
values of each acidic compound as long as the LWC does not limit the uptake. High LWCs ($0.1\text{--}1\text{ g m}^{-3}$) typically associated
with cloud/fog conditions and, accordingly, less acidic media ($\text{pH} > 4$), favor phase partitioning towards the aqueous phase for
most of the weak acids as well as for ammonia. Less water-soluble acids (i.e., with lower H values), such as dissolved SO_2
and HONO, display fractions above 0.1 only under less acidic conditions for typical cloud LWC values. Thus, even at colder
170 cloud temperatures than the 298 K used in Fig. 2, where H_A^* is larger, SO_2 and HONO largely remain in the gas phase under
typical cloud acidity conditions. Hence, note that $X_{A_{aq}}$ values of SO_2 are typically in the range of 0.005 to 0.5 depending on
both the cloud acidity and temperature. Under typical aerosol conditions, the LWC restricts uptake and only very small
fractions of the less water-soluble acids can partition in the aqueous particle phase. Moreover, very weak acids, with pK_a values
larger than 7 (e.g., 2-nitrophenol) show almost no acidity dependency in the plotted acidic range. On the other hand, for
175 stronger acids the LWC and acidity impact is even lower due to their lower and/or multiple pK_a values. For example, phthalic
acid partitions in substantial amounts into the aqueous phase for a large range of acidity and LWC conditions. The implication
is that only very water-soluble and strong acids are expected to remain in acidic aerosol solutions. However, it is worth
mentioning again that this treatment neglects several other factors/processes affecting the partitioning of acids in the aqueous
phase, particularly under concentrated aqueous-aerosol conditions. Specifically, volatile acids (e.g., formic and acetic) often
180 show substantial deviations from this theory (see Nah et al. (2018)) for instance because of the formation of organic salts which
can increase their particle partitioning by two orders of magnitude (Meng et al., 2007). In practice, weak acid anions are often
measured in non-negligible fractions in the particle phase (Tanner and Law, 2003; Limbeck et al., 2005; van Pinxteren and
Herrmann, 2007; Bao et al., 2012; Nah et al., 2018; Teich et al., 2019).

185 *Non-ideal solutions*

At less than 100% relative humidity, aqueous aerosol solutions exist as a highly concentrated, complex mixture of electrolytes.
Interionic and ion-molecule interactions are critically important under those conditions, leading to thermodynamically non-
ideal behavior (Pitzer, 1991; Zaveri, 2005; Cappa et al., 2008; Zuend et al., 2008; Herrmann et al., 2015; Rusumdar et al.,
2016; Rusumdar et al., 2020). Therefore, parameters that have been developed for dilute aqueous solutions do not strictly apply
190 to aerosol-phase chemistry.

Nevertheless, several such principles, such as Henry's Law, have been shown experimentally to hold for the aqueous aerosol
phase (Kroll et al., 2005; Sumner et al., 2014), although it may be necessary to account for phenomena such as salting effects
(Kampf et al., 2013; Waxman et al., 2015). Factors such as ionic strength, the different chemical composition of the
concentrated solution, other favored chemical pathways, shifted chemical equilibria (salting-in/out, hydration, metal
195 complexes, dimer/polymer etc.) and more can significantly affect overall phase partitioning and reaction rates. The inclusion
of these factors into the calculation of the effective Henry's Law constant can explain increased or decreased aqueous-phase



partitioning of chemical compounds such as atmospheric carbonyl compounds (Kampf et al., 2013; Waxman et al., 2015) and organic monocarboxylic acids (Limbeck et al., 2005; Meng et al., 2007) compared to what may be expected based on aqueous solubility alone. Ionic strength effects are also believed to be critically important for acidity-producing in-particle chemical reactions such as S(IV) oxidation (Martin and Hill, 1987a; Lagrange et al., 1993; Lagrange et al., 1994; Maaß et al., 1999; Ali et al., 2014; Cheng et al., 2016), although experimental data at the extremely high ionic strengths typical of atmospheric aerosols are limited. The first models treating both non-ideal solution effects as well as their feedbacks on occurring chemical processes in detail, have been developed in the last couple of years enabling advanced investigations, e.g. on the phase partitioning issues (Rusumdar et al., 2016).

2.2 Acidity buffering

The response of pH in the atmospheric aqueous phases to a perturbation in acidity can be strongly affected by the presence and ability of weak acids or bases to buffer against that change. A buffer is a mixture of a weak acid and its conjugate base (e.g., formic acid and formate) or a mix of a weak base and its conjugate acid (e.g., ammonia and ammonium). The buffering effect, a resistance to pH change, comes from changes in the equilibrium between concentrations, for example, of a weak acid and a conjugate base. The Henderson-Hasselbach equation (Eq. 4)

$$pH = pK_a + \log \left(\frac{[A^-]}{[HA]} \right) \quad (4)$$

is used to calculate the pH of a buffer solution based on the acid dissociation constant (K_a) and the concentrations of the acid [HA] and its conjugate base $[A^-]$. Ion speciation curves for a wide range of atmospherically relevant weak acids are shown in Fig. 3. The magnitude of the buffering effect is greatest when the solution pH is equal to the pK_a of the weak acid buffer (intersection points of the speciation curves as shown in Fig. 3). Consider, for example, the case of formic acid ($pK_a = 3.75$ at 298 K) and formate (see Fig. 3c). If protons are added (e.g., through addition of a strong acid such as sulfuric acid) to a solution containing formate or formic acid, and the solution pH is far above or below the 3.75 pK_a of formic acid, each added proton will directly increase the H^+ concentration in solution. When the solution pH, however, is close to the formic acid pK_a (where the concentrations of formic acid and formate are equal), many of the added protons will be consumed in converting formate to formic acid, thereby slowing the pH decline of the solution. For diprotic acids, buffering occurs at each of the two acid dissociation steps. Carbonate buffering is a relevant example for atmospheric cloud and fog droplets. The pK_a values for carbonic acid and bicarbonate are 6.35 and 10.33 at 298 K. A titration by acid addition beginning at pH 12, therefore, would show strong buffering at pH 10.3 and again at pH 6.4, the latter being much more relevant for atmospheric water. Moreover, in mineral dust and volcanic particles that can bear phosphate minerals such as apatite, dissolved phosphate can act as a buffer. But, unlike carbonates, the phosphate buffer cannot be not lost owing to volatilization. Nevertheless, the buffering by phosphate in other kind of atmospheric aerosol particles is negligible because of the typically extremely low phosphate concentration.





Figure 3. Ion speciation of dissolved (a) SO_2 , (b) HONO, (c) formic acid, (d) acetic acid, (e) phthalic acid, and (f) 2-nitrophenol as a function of pH.

230 The buffering capacity (β), a measure to quantitatively express the resistance of an aqueous solution towards acidity changes, is defined for a monoprotic acid by Eq. 5 (see Urbansky and Schock (2000) for details). The buffering capacity β expresses the amount of an acid or base concentration addition ($d[C_{a/b}]$) needed to cause a certain change in pH ($d(\text{pH})$).

$$\beta = \frac{d[C_{a/b}]}{d(\text{pH})} = \ln 10 \cdot \left(\frac{K_W}{[H]^+} + [H^+] + \sum_i \frac{[C]_i \cdot K_{a,i} [H^+]}{(K_{a,i} + [H^+])^2} \right) \quad (5)$$



Eq. 5 and the plotted examples in Fig. 4 reveal that very high and very low acidity conditions show significantly increased buffering capacities. Moreover, added buffers in the solution lead to local maxima of β between very acidic and very alkaline conditions. In the case of one monoprotic acid present in an aqueous solution, the maximum of the buffering capacity occurs close to the pK_a value of the acid, as mentioned above (within approximately ± 1 pH unit). Furthermore, Eq. 5 shows that buffering capacity, i.e., the amplitude of the local maxima, depends on the concentration of the buffer compound. This agrees with findings in the field, e.g. in fog samples analyzed by Collett et al. (1999) (see discussion below). Furthermore, this dependency implies a rather high buffering capacity in regions with high multiphase concentrations of weak inorganic and organic acids and bases or high amounts of particulate buffers such as carbonate components. However, the latter are most important in buffering the acidity of supermicron particles or fog/cloud droplets that activate on them.



245 **Figure 4.** Buffering capacity β of water, ammonia/ammonium and carbonate/bicarbonate/carbonic acid (top: from left to right) as well as formic and acetic acid (bottom: from left to right) as a function of pH. The atmospherically relevant range of cloud and aerosol pH is marked in yellow.



250 Titrations of actual cloud and fog samples have exhibited buffering across a wide pH range, suggesting the importance of pH buffering by a variety of compounds with different pK_a values. For example, Collett et al. (1999) report titrations of fog samples collected at urban and rural locations in California's San Joaquin Valley. Observed buffering in rural fogs in the study could be nearly accounted for based on ammonia and bicarbonate concentrations present in the fog samples. By contrast, significant additional buffering (β up to 10^{-4} mol L⁻¹) was observed in urban fogs over a broad pH range from 4 to 7. The amount of additional buffering was strongly correlated with concentrations of organic compounds in fogs from these environments, with relevant organic buffering agents likely including carboxylic and dicarboxylic acids and phenols.

255 The buffering phenomenon described above is often referred to as "internal buffering," since it derives from shifts in equilibrium concentrations of compounds present in solution. The exchange of material with the gas phase can also lead to "external buffering." Perhaps the most important form of external buffering is the uptake of additional ammonia from the gas phase in response to a drop in solution pH, as outlined by Liljestr and (1985) and Jacob et al. (1986). Corresponding buffering in atmospheric aerosols from semi-volatile partitioning also occurs, as shown by Meng et al. (2007), Weber et al. (2016) and
260 Song et al. (2019a) as well as recently by Zheng et al. (2020). However, it should be noted that the effect of aerosol pH buffering from semi-volatile gases on relevant chemical processes has not been studied comprehensively and still represents an issue for future research.

One important consequence of pH buffering in fog and cloud drops is an effect on rates of pH-sensitive aqueous reactions. The presence of (internal and/or external) acid buffering in cloud and fog droplets can slow droplet acidification and maintain
265 greater rates of reaction for strongly pH-dependent aqueous chemical pathways (e.g., the oxidation of S(IV) by ozone) which are favored by high pH.

3 Sources of acidity and alkalinity

Acidic and alkaline components of tropospheric aerosols result from primary gas and aerosol particle emissions as well as secondary gas-phase and aqueous-phase formation processes (P oschl, 2005b; Seinfeld and Pandis, 2006; Seinfeld, 2015; Zhang
270 et al., 2015). The most important acidic chemical components of aerosols and cloud/fog droplets are sulfuric acid, nitric acid, nitrous acid, and hydrochloric acid, as well as organic mono- and dicarboxylic acids (e.g., formic acid, acetic acid, oxalic acid etc.) (Vet et al., 2014; Zhang et al., 2015). Then, the most important basic components of aerosols and cloud/fog droplets are ammonium, amines and alkali/alkaline earth metals (U.S.EPA, 2000; Vet et al., 2014; Zhang et al., 2015). The global contribution of different acid and base ions to precipitation has been assessed by Vet et al. (2014). As precipitation samples
275 provide both compositional and acidity information for the vertical column, these data represent a useful means to point out the spatial sources and sinks of gas and aqueous-phase acidity and alkalinity components.

Gaseous acids can be directly emitted into the troposphere from primary sources such as biomass combustion, traffic (fuel combustion), domestic heating, industrial burning, agriculture, soil, and vegetation (Chebbi and Carlier, 1996; Paulot et al., 2011; Spataro and Ianniello, 2014; Kawamura and Bikkina, 2016). Moreover, gaseous acids can be formed secondarily by



280 gas-phase oxidations of emitted acid precursor compounds such as SO₂, NO_x, and VOCs (Chebbi and Carlier, 1996; Paulot et al., 2011; Spataro and Ianniello, 2014; von Schneidemesser et al., 2015; Zhang et al., 2015; Kawamura and Bikkina, 2016). The gas-phase OH oxidation of SO₂ is an important source of gaseous sulfuric acid and, after condensation, of particulate sulfate (von Schneidemesser et al., 2015; Zhang et al., 2015). SO₂ is emitted from anthropogenic activities such as the combustion of sulfur-containing fuels and various natural sources such as volcanos (Smith et al., 2001; Smith et al., 2011; Seinfeld, 2015). Moreover, it is formed from the oxidation of natural precursors such as dimethyl sulfide (DMS, CH₃SCH₃) emitted by oceanic phytoplankton (Seinfeld and Pandis, 2006). The gaseous oxidation pathway of SO₂ contributes also to newly formed particles (nucleation, (Zhang et al., 2015)) which are expected to be quite acidic. Furthermore, the gaseous oxidation of NO_x and VOCs can lead to the formation of nitric/nitrous acid and organic acids (e.g., formic, acetic and oxalic acid) (Chebbi and Carlier, 1996; Paulot et al., 2011; Spataro and Ianniello, 2014; Zhang et al., 2015; Kawamura and Bikkina, 2016). By contrast, gaseous bases such as ammonia and amines are almost exclusively emitted into the troposphere, mainly from agriculture due to intensive stock farming and the use of NH₃-based fertilizer applications. Moreover, bases are released from biomass burning, vehicles, industrial processes, and as a consequence of volatilization from soils and oceans. (U.S.EPA, 2000; Behera et al., 2013). As shown in Fig. 5, subsequent to their emission or secondary formation, gaseous acids and bases can condense on existing aerosol particles or fog/cloud droplets and can then contribute to aerosol acidity.

295




Figure 5. Schematic of sources (red text) and conditions of acidity in different aqueous aerosol particles (green text) together with microphysical and chemical processes that are able to influence the acidity of tropospheric aerosols (Fig. created after Raes et al. (2000) and McMurry (2015)) The dashed gray line represents an aerosol number size distribution.

300 Acidic and alkaline aerosol components are also (i) primarily emitted by anthropogenic and natural processes (Zhang et al., 2015) or secondarily formed in aqueous aerosol solutions or at their interface (see Sect. 4 and 5 below). Important anthropogenic primary sources of acidic and alkaline aerosols (see Fig. 5) are urban combustion aerosols and agricultural aerosols, including e.g. agricultural ammonia from livestock farming. Important natural primary sources of acidic and alkaline aerosols are sea spray, desert dust, biomass burning and volcanic emissions. Besides the secondary acid formation in the gas-
 305 phase, in-cloud oxidation of SO_2 contributes more than 50% globally to sulfate aerosol mass formation (Alexander et al., 2009) (see Sect. 4.1 for details). Thus, the aqueous-phase formation of sulfate from the oxidation of SO_2 is the largest source of acidity in the atmosphere. However, besides sulfate, other acidic components are also secondarily formed in aqueous aerosols



such as nitrate, chloride, formate, acetate, and oxalate (see (Chebbi and Carlier, 1996; Spataro and Ianniello, 2014; Ervens, 2015; Zhang et al., 2015; Kawamura and Bikkina, 2016)).

310 In the past, emissions of SO₂ in industrialized countries were the predominant cause of strong acidification of aerosol particles, cloud droplets and precipitation, typically known as the acid rain phenomenon (Adriano and Johnson, 1989; Seinfeld and Pandis, 2006). However, due to strongly reduced anthropogenic sulfur emissions in some parts of the world, a reduction in cloud/fog acidity has been observed over recent decades (see Pye et al. (2020)). As a consequence of the changing acid and base sources, the composition of continental aerosol particles and cloud/fog/rain droplets will most likely continue to evolve

315 toward compositions observed pre-industrially in rural continental areas, e.g. in North America and Western Europe. These environments are characterized by higher contribution of organic acids and chloride (less acid displacement) and less sulfate/nitrate mass (see precipitation composition data compiled by Vet et al. (2014)). In such a future environment, natural acidity sources become a much more important source for the acidity of tropospheric cloud/fog/rain droplets. On the other hand, no significant changes are expected for the acidity of marine droplets, except downwind of continents. Their main acidity

320 and alkalinity sources, such as the emission of DMS, marine NH₃ and sea salt particles containing chloride and base cations, are not expected to change significantly. However, it should be mentioned that the impact of climate change including higher temperatures and ocean acidification and related changes in the ocean biochemistry, may unequally affect the emission of DMS in different regions. The effects of climate change on DMS emission patterns are still under debate due to the complex interactions of marine biochemistry and atmosphere-ocean interactions (Six et al., 2013; Gypens and Borges, 2014; Dani and

325 Loreto, 2017; Hopkins et al., 2020).

4 Interactions of acidity and chemical processes: Inorganic systems

In this section, the feedbacks between particle/droplet acidity and key inorganic chemical subsystems, the sulfur(IV) oxidation and tropospheric halogen chemistry are discussed in detail.

4.1 Acidity and Sulfur Oxidation

330 In addition to its reaction with OH in the gas phase, SO₂ is oxidized via heterogeneous and multiphase reactions in clouds, fog, or aerosol particles to form particulate sulfate. Sulfate is a major component of PM_{2.5}, especially in areas affected by emissions from burning coal or other sulfur-containing fossil fuels (Attwood et al., 2014). Because sulfate lifetime is on the order of days (Barth et al., 2000), sulfate contributes to regional haze and acid deposition, as well as local air pollution.

Once in the aqueous phase, SO₂ is hydrated and undergoes acid-base equilibrium to form other S(IV) species, bisulfite (HSO₃⁻)

335 (pK_{a,R-1} = 1.9) and sulfite (SO₃²⁻) (pK_{a,R-2} = 7.2). The hydration of SO₂ upon uptake alone, according to (R-1) already leads to the release of acidity:






S(IV) oxidation in the aqueous phase to form S(VI) species (sulfate (SO_4^{2-}), bisulfate (HSO_4^-), and sulfuric acid (H_2SO_4)) leads to further acidification. S(IV) oxidation can take place via a number of chemical pathways, many of which are pH-sensitive (Fig. 6). As a result of the equilibrium reactions described by R-1 and R-2, the effective solubility of SO_2 in aqueous solutions increases rapidly with increasing pH (see Eq. 2a). Partly for this reason, as well as because of their relatively small liquid water content ($\sim 10^{-9} \text{ cm}^3 \text{ cm}^{-3}$), sulfate formation in aerosols is generally believed to be less significant than in clouds and fog (Schwartz, 1986). Only S(VI) formation in the gas-phase and in clouds is included in most large-scale atmospheric chemistry models. Globally, in-cloud formation is thought to be the dominant sulfate production pathway ($\sim 60\%$), particularly over the oceans (generally $>75\%$) (Barth et al., 2000; Barrie et al., 2001; Manktelow et al., 2007; Alexander et al., 2009; Faloon, 2009; Alexander et al., 2012). However, there is evidence that significant sulfate formation also occurs in polluted urban areas, during periods of high aerosol surface area and few clouds (Hering and Friedlander, 1982; Wang et al., 2014; He et al., 2018a). This suggests that aerosol chemistry is also an important source of sulfate under some conditions.

In the aqueous phase, S(VI) species exist in acid-base equilibrium according to:



Since sulfuric acid is a very strong acid ($K_{a,R-4} \cong 1000 \text{ mol L}^{-1}$ at 298 K (Graedel and Weschler, 1981)), almost no unionized H_2SO_4 exists in aqueous solution and HSO_4^- is significant only at $\text{pH} < 3$. As a consequence, the conversion of S(IV) to S(VI) in the aqueous phase increases the acidity of the cloud or aerosol particle not only by the initial acidification through the SO_2 reaction with water, but additionally through the dissociation of sulfuric acid. Some S(IV) oxidation reactions have other acidic byproducts such as halous acid species HX (with $X = \text{Cl}, \text{Br}$) or HONO, and thus may contribute additional acidity to the aerosol (Fig. 6). Figure 6 illustrates that sulfate oxidation under urban haze conditions can significantly contribute to the acidification of aerosols. After a short period of chemical processing, aerosols are expected to reach $\text{pH} 4.5$ or lower. Particularly for haze particles with initial pH conditions above 4, a fast acidification can be modeled as a consequence of the higher initial S(IV) oxidation rates under less acidic conditions.

In the next subsections, we outline the major S(IV) oxidation pathways, their sensitivity to the pH of the aqueous medium, and their potential to alter pH via the formation of acid products.




365 **Figure 6.** S(IV) oxidation rates for Beijing winter haze conditions (following (Cheng et al., 2016)). Shown are initial rates (top), aerosol pH after 10 s of reaction at the initial rates (center), and S(IV) oxidation rates after 10s of reaction (bottom). Total S(IV) rate with and without taking into account ionic strength at the maximum reported limit is shown. Rates used were those recommended in this text.



370 4.2 S(IV) oxidation through O₃, H₂O₂, ROOH, and HOX (with X = Cl, Br, I)

Due to the pH-dependent partitioning of S(IV) species and hence solubility of SO₂, most S(IV) oxidation mechanisms are highly pH-dependent. However, S(IV) oxidation by H₂O₂, is only weakly pH dependent. At pH values typical of cloud water (pH = 2 - 7 (Pye et al., 2020)), S(IV) oxidation by H₂O₂ is thought to dominate sulfate production (Faloona, 2009) although other oxidants can be important at higher pH values or if H₂O₂ is depleted (e.g., Shen et al. (2012)). In-cloud S(IV) oxidation
375 by H₂O₂ proceeds via reaction with HSO₃⁻, followed by addition of H⁺ (R-5/6a).



380

Therefore, the intrinsic reaction rate decreases rapidly with increasing pH above pH 2 (McArdle and Hoffmann, 1983). This is balanced by the fact that the effective SO₂ solubility increases with increasing pH. As a result, the overall rate is relatively independent of pH, above pH ~ 1.5. The rate expression for S(VI) formation by S(IV) + H₂O₂ is given by McArdle and Hoffmann (1983); Lind et al. (1987); and Gunz and Hoffmann (1990):

385

$$R_{\text{H}_2\text{O}_2} = k_{6a} \frac{[\text{H}^+][\text{HSO}_3^-]}{1 + K_5[\text{H}^+]} [\text{H}_2\text{O}_2] \quad (6a)$$

with a recommended temperature-dependent rate constant $k_{6a} = 7.45 \times 10^7 \exp\left(-4000\left(\frac{1}{T} - \frac{1}{298}\right)\right) \text{ L mol}^{-1} \text{ s}^{-1}$ and $K_5 = 13 \text{ mol L}^{-1}$.

390 Recently, Liu et al. (2020) investigated S(VI) formation by S(IV) + H₂O₂ in a flow reactor under aqueous aerosol conditions (pH = 2.5, high ionic strength, and 73-90% relative humidity) and in the presence of malonic acid. This study revealed that, under concentrated aqueous-aerosol conditions, the S(VI) formation rate can be significantly increased compared to dilute aqueous conditions like those in clouds. The study demonstrated that ionic strength and general acid catalysis promotes faster S(VI) formation via R-6b. This additional pathway is expected to contribute to S(VI) missing from model simulations of severe
395 haze episodes (Hering and Friedlander (1982); Wang et al. (2014); He et al. (2018a)).

The rate expression given by Liu et al. (2020) is as follows:

$$R_{\text{H}_2\text{O}_2} = (k + k_{HX}[\text{HX}][\text{H}^+]^{-1})K_{a1}H_{\text{SO}_2}p_{\text{SO}_2}H_{\text{H}_2\text{O}_2}p_{\text{H}_2\text{O}_2} \quad (\text{pH} > 2) \quad (6b)$$



400 with the following ionic strength dependencies of the reaction rate constant, Henry's law constants and dissociation constants (see Liu et al. (2020) and references therein).

$$k: \quad \log\left(\frac{k}{k_{I=0}}\right) = 0.36 \cdot I - \frac{1.018\sqrt{I}}{1+1.018\sqrt{I}} \quad (I_{\max} = 5 \text{ molal}) \quad (6c)$$

$$H_{H_2O_2}: \quad \frac{H_{H_2O_2}}{H_{H_2O_2}^{I=0}} = 1 - 1.414 \cdot 10^{-3} \cdot I^2 + 0.121 \cdot I \quad (I_{\max} = 5 \text{ molal}) \quad (6d)$$

$$405 \quad H_{SO_2}: \quad \frac{H_{SO_2}}{H_{SO_2}^{I=0}} = \left(\frac{22.3}{T} - 0.0997\right) \cdot I \quad (I_{\max} = 6 \text{ molal}) \quad (6e)$$

$$K_{a1}^*: \quad \log\left(\frac{K_{a1}^*}{K_{a1}^{I=0}}\right) = 0.5 \cdot \sqrt{I} - 0.31 \cdot I \quad (I_{\max} = 6 \text{ molal}) \quad (6f)$$

and

$$K_{a2}^*: \quad \log\left(\frac{K_{a2}^*}{K_{a2}^{I=0}}\right) = 0.5 \cdot \sqrt{I} - 0.36 \cdot I \quad (I_{\max} = 6 \text{ molal}) \quad (6g)$$

In Eq. 6b, $k = k' \cdot 1.3 \times 10^{-2} \cdot e^{1960\left(\frac{1}{T} - \frac{1}{298}\right)} \cdot 6.6 \times 10^{-8} \cdot e^{1500\left(\frac{1}{T} - \frac{1}{298}\right)}$ (reaction rate constant of proton-catalyzed pathway
 410 R-6a), k_{HX} (overall reaction rate constant of the catalysis pathway of a general acid HX (R-6b), $k_{\text{malonic acid}} = 5.61 \times 10^5 \text{ mol}^2 \text{ kg}^{-2} \text{ s}^{-1}$ (at $I = 3.9 \text{ mol kg}^{-1}$), $k_{\text{malonate}} = 1.32 \times 10^5 \text{ mol}^2 \text{ kg}^{-2} \text{ s}^{-1}$ (at $I = 6.6 \text{ mol kg}^{-1}$)),
 $K_{a1} = 1.3 \times 10^{-2} \cdot e^{1960\left(\frac{1}{T} - \frac{1}{298}\right)}$ (thermodynamic dissociation constant of R-5), $H_{SO_2} = 1.23 \cdot e^{3145.3\left(\frac{1}{T} - \frac{1}{298}\right)}$ (Henry's law
 constant of SO_2) and $H_{H_2O_2} = 1.3 \times 10^5 \cdot e^{7297.1\left(\frac{1}{T} - \frac{1}{298}\right)}$ (Henry's law constant of H_2O_2). Furthermore, p_{SO_2} and $p_{H_2O_2}$
 415 represent partial pressure of SO_2 and H_2O_2 in the gas phase, respectively. Note, the kinetic of the study by Liu et al. (2020) has
 been determined for NaCl/NaNO₃-malonate/malonic acid mixtures only which could restrict their applicability. Hence, further
 investigations for other aerosol composition mixtures (e.g., considering ammonium-sulfate salts and other general acids), lower
 pH conditions and higher ionic strengths are definitely needed to provide even more advanced rate expressions for concentrated
 aqueous-aerosol conditions.

420 Organic hydroperoxides (ROOH) can also oxidize HSO_3^- by a similar mechanism to H_2O_2 , although at lower rates (Graedel
 and Goldberg, 1983; Lind et al., 1987; Drexler et al., 1991). The oxidation of HSO_3^- by methylhydroperoxide, CH_3OOH , has
 methanol as a product, with the overall reaction given as (Lind et al., 1987):



with a third-order rate law:

$$425 \quad R_{CH_3OOH} = k_7 [HSO_3^-][CH_3OOH][H^+] \quad (7)$$



with $k_7 = 1.7 \times 10^7 \exp\left(-3800\left(\frac{1}{T} - \frac{1}{298}\right)\right) \text{ L}^2 \text{ mol}^{-2} \text{ s}^{-1}$.

The S(IV) oxidation rate for peroxyacetic acid is faster (Lind et al., 1987), and produces acetic acid as a byproduct, thereby further increasing the acidity of the aqueous phase:



430 with a third-order rate law:




with $k_8 = 5.6 \times 10^7 \exp\left(-3990\left(\frac{1}{T} - \frac{1}{298}\right)\right) \text{ L}^2 \text{ mol}^{-2} \text{ s}^{-1}$.

The aerosol- and gas-phase abundances of organic hydroperoxides are poorly constrained, so S(IV) oxidation by ROOH may be more important than previously thought in aerosols containing secondary organic material (Ye et al., 2018b; Dovrou et al., 2019; Wang et al., 2019a). Organosulfates have been proposed as minor products of the S(IV) + ROOH reactions with secondary organic material, with further implications for aerosol pH (Wang et al., 2019a).

In contrast to S(IV) oxidation by H_2O_2 , the oxidation of S(IV) by reaction with O_3 becomes faster with increasing pH. Since S(VI) formation contributes to acidification of the aerosol, these processes are therefore potentially self-limiting, depending on the buffering capacity of the aqueous medium (Fig. 6).



Each S(IV) species reacts with O_3 , leading to a composite rate expression of:



Here, F_i is an empirically determined factor accounting for the effect of ionic strength, I , on the rate. Lagrange et al. (1994) explored the effects of ionic strength on the oxidation of S(IV) by O_3 (up to 4 mol L^{-1}) and found that $F = 1.59 \pm 0.3$ for NaCl and $F = 3.71 \pm 0.7$ for Na_2SO_4 . The rate constant for oxidation of SO_3^{2-} by O_3 ($k_{9a} = 1.5 \times 10^9 \exp\left(-5280\left(\frac{1}{T} - \frac{1}{298}\right)\right) \text{ L mol}^{-1} \text{ s}^{-1}$) is over three orders of magnitude larger than the rate constant for $\text{O}_3 + \text{HSO}_3^-$ ($k_{9b} = 3.7 \times 10^5 \exp\left(-5530\left(\frac{1}{T} - \frac{1}{298}\right)\right) \text{ L mol}^{-1} \text{ s}^{-1}$) (Hoffmann and Calvert, 1985), which is more than ten times the rate constant for the reaction of O_3 with $\text{SO}_2 \cdot \text{H}_2\text{O}$ ($k_{9c} = 2.4 \times 10^4 \text{ L}^2 \text{ mol}^{-2} \text{ s}^{-1}$) when the respective maximum values are compared (Hoffman, 1986). Therefore, the overall rate of S(IV) oxidation by O_3 increases rapidly with increasing pH, and is most important above pH 5-6 (Chameides, 1984; Calvert et al., 1985; Turnock et al., 2019).

Sulfate can also form via reaction of S(IV) with O_3 on the surface of alkaline aerosols, e.g. freshly emitted sea salt aerosols and some mineral dust aerosols (Sievering et al., 1992; Chameides and Stelson, 1993; Zhang and Carmichael, 1999; Li et al., 2006; Wu et al., 2011; Yu et al., 2017; Zhang et al., 2018b). At pH values typical of fresh sea salt aerosol (pH ≈ 8), the S(IV)



loss rate constant for oxidation by O_3 in these aerosols is 10^5 times larger than in-cloud oxidation by H_2O_2 , more than making up for their lower liquid water content (Sievering et al., 1992; Chameides and Stelson, 1993). However, like other $S(IV) + O_3$ mechanisms, these processes are potentially self-limiting, as noted above.


460 Reactions of $S(IV)$ with hypohalous acids ($HOBr$, $HOCl$, and HOI ; see reactions R-17/ R-18) contribute to sulfate formation in the marine boundary layer (Vogt et al., 1996; von Glasow et al., 2002a; Chen et al., 2016). These reactions act as a sink for reactive halogens by converting them to their acidic form (e.g., $HOBr \rightarrow HBr$, see Sect. 4.8 for further details) (Chen et al., 2016).

4.3 Free radical pathways for $S(IV)$ oxidation

465 The hydroxyl radical (OH) can oxidize $S(IV)$ in the aqueous phase through a radical pathway involving SO_3^- , SO_5^- , HSO_5^- , and SO_4^- . This process is more likely to be important in cloud water than in aqueous aerosol due to the higher liquid water content of clouds and the relatively lower OH concentration in aqueous aerosols (Herrmann et al., 2010; McNeill, 2015). The high concentrations of organic material in aerosols can quench of radical and triplet species (Herrmann et al., 2010; McNeill, 2015; Wang et al., 2020). Furthermore, the reaction of OH with SO_3^{2-} is somewhat faster than that of OH with HSO_3^-
470 ($k = 4.6 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$ vs. $2.7 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$) (Buxton et al., 1996). This, along with the pH dependence of the water solubility of SO_2 , suggests that $S(IV)$ oxidation by OH is more efficient at higher pH and in clouds (and is potentially self-limiting). The production of SO_4^- via this reaction pathway couples $S(IV)$ oxidation to organosulfate production (Perri et al., 2010), although this is a minor pathway (McNeill et al., 2012).

Laboratory studies have demonstrated sulfate production on the surface of acidic aerosols, via direct electron transfer from
475 HSO_3^- to O_2 , followed by a free-radical chain oxidation of bisulfite to sulfate (Hung and Hoffmann, 2015), however the significance of this pathway is not confirmed by field and modeling studies (Shao et al., 2019). Catalytic oxidation of $S(IV)$ by NO_3 (Exner et al., 1992; Rudich et al., 1998; Feingold et al., 2002), also believed to take place via a free radical mechanism, may be important in the remote troposphere. Recent experimental studies suggest that photolysis of particulate nitrate and hydrolysis of NO_2 to form nitrate and $HONO$ (Li et al., 2018) may accelerate oxidation of $S(IV)$ under Beijing conditions by
480 generating NO_2 and OH radicals (Gen et al., 2019). However, the consumption of OH radicals by organic constituents present in aerosols were ignored in this study likely leading to an overestimation of the effect.

Another suggested $S(IV)$ oxidation pathway is the reaction of excited triplet states of photosensitizers (PS^*) with $S(IV)$ species (see R-10). This pathway potentially involves produced sulfur-containing radicals and/or excited transient species (see e.g., Wang et al. (2020) and Loeff et al. (1993)). Currently, it is also discussed as potential $S(IV)$ oxidation pathway under polluted
485 aerosol conditions (Wang et al., 2020).



The exact reaction pathway is still uncertain particularly with respect to the involved sulfur-containing radicals or excited transient species. Some studies (Loeff et al., 1993; Wang et al., 2020) already determined chemical reaction rate constants for



certain PS* species such as acetophenone, flavone, xanthone, 4-(benzoyl)benzoic acid and anthraquinone-1-sulfonate. The
490 second-order reaction rate constants of PS* with S(IV) species measured in the laboratory are between 6.0×10^7 and
 1.0×10^9 mol L⁻¹ s⁻¹. Kinetic measurements of the reactive PS* quenching by S(IV) using of ambient filter extracts taken during
Chinese winter haze conditions revealed a rate constant of 1.3×10^8 mol L⁻¹ s⁻¹ (Wang et al. (2020)). Note, the kinetic
investigations of Wang et al. (2020) assumed that the initial reaction step is the rate-limiting step in this reaction sequence and
the reaction rate constant is pH-independent. So, based on Wang et al. (2020), the rate expression is as follows:

$$495 \quad R_{PS^*} = k_{10}[PS^*][S(IV)] \quad (10)$$

Due to the presently strong uncertainties in the existing kinetic data and mechanistic understanding of R-10, a recommendation
of a proper kinetic reaction rate constant is rather difficult. Thus, we preliminary recommend the chemical rate constant of
 $k_{10} = 1.3 \times 10^8$ mol L⁻¹ s⁻¹. Finally, it should be noted that great care is needed for estimating the rate of R-10 because of lacking
knowledge about the present PS* concentrations in ambient aerosols and cloud droplets and the very rapid quenching and
500 deactivation triplet species by water, dissolved oxygen and organic/inorganic aerosol constituents which might lead to very
low PS* concentrations which can strongly limit or inhibit this pathway similarly to the S(IV) oxidation by free radical which
can be effectively scavenged by particle constituents other than S(IV) as outline above in the present section.

4.4 S(IV) oxidation catalyzed by transition metal ions

The oxidation of S(IV) by O₂ as catalyzed by transition metal ions (TMI, mainly Fe(III) and Mn(II); see R-11/R-12)
505 (Humphreys, 1964; Martin and Hill, 1987a; Martin and Hill, 1987b; Brandt and van Eldik, 1995; Alexander et al., 2009; Harris
et al., 2013) is an efficient pathway for S(VI) formation, especially under conditions where photochemistry is limited, e.g.
wintertime at high latitudes (Simpson et al., 2019).



510 The solubility and speciation of the TMI (Pye et al., 2020), as well as the reaction rates, all depend on pH. As primary
pollutants, TMI concentrations are higher in aerosols than in cloud water, but this effect is limited by the pH-dependent
solubility of the active species. The TMI-S(IV) reactions (R-11/R-12) are also reported to be inhibited by ionic strength (Martin
and Hill, 1987a; Martin and Hill, 1987b), although this dependence is only known under the relatively dilute conditions which
are accessible in bulk solutions. This introduces considerable additional uncertainty to estimates of the aerosol-phase TMI
515 catalyzed S(IV) oxidation rate.

TMI-mediated S(IV) oxidation has been proposed to proceed through radical intermediates (Grgić and Berčić, 2001), at least
for pH > 3.6 (Martin et al., 1991). A detailed discussion of the mechanisms can be found in Brandt and van Eldik (1995). A
pH-dependent synergistic effect has been reported when multiple transition metal ions are present in solution (Ibusuki and
Takeuchi, 1987; Martin and Good, 1991; Harris et al., 2013). Martin et al. (1991) observed that water-soluble organic material
520 inhibits Fe(III)-catalyzed S(IV) oxidation for pH ≥ 5. Given this pH range, the effect is not expected to be significant for



atmospheric aerosols, although interactions with organics, for example complexation with oxalate, may impact TMI chemistry in other ways (e.g., Okochi and Brimblecombe (2002); Passananti et al. (2016)).

Given the current focus on sulfate formation in atmospheric aerosols, our recommendations for kinetics of S(IV) oxidation by TMI favor studies which included the ionic strength and pH effects. For Fe(III)-catalyzed S(IV) oxidation, the expression from Martin and Hill (1987b) and Martin et al. (1991) is as follows:

$$R_{\text{Fe},10a} = \begin{cases} \frac{k_{11a}[\text{Fe(III)}][\text{S(IV)}]10^{-2\sqrt{I}/(1+\sqrt{I})}}{[\text{H}^+](1+K_{11}[\text{S(VI)})^{2/3}]} & \text{for pH} < 3.6 \\ k_{11b}[\text{Fe(III)}]^2[\text{S(IV)}] & \text{for } 3.6 \leq \text{pH} \leq 5 \\ k_{11c}[\text{S(IV)}] & \text{for } 5 < \text{pH} \leq 6 \\ k_{11d}[\text{S(IV)}] & \text{for pH} > 6 \end{cases} \quad (11a)$$

Here, $k_{11a} = 6 \text{ s}^{-1}$, $K_{11} = 150 (\text{mol L}^{-1})^{-2/3}$, $k_{11b} = 10^9 \text{ L}^2 \text{ mol}^{-2} \text{ s}^{-1}$, $k_{11c} = 10^{-3} \text{ s}^{-1}$ and $k_{11d} = 10^{-4} \text{ s}^{-1}$. However, the dependence of Eq. 11a on ionic strength (I) is only known up to 1 mol L^{-1} and unfortunately the rate law is valid for a limited range of conditions only ($[\text{Fe}^{3+}] > 10^{-7} \text{ mol L}^{-1}$, $[\text{S(IV)}] < 10^{-5} \text{ mol L}^{-1}$, $[\text{S(VI)}] < 10^{-4} \text{ mol L}^{-1}$, $I < 10^{-2} \text{ mol L}^{-1}$). Moreover, note that the ionic strength effect was verified at $\text{pH} = 2$ and $T = 25 \text{ }^\circ\text{C}$ only. Additionally, the study implied that the effect of higher S(IV) and S(VI) concentrations may be more important than the ionic strength effect (see Martin et al. (1991) for details). Due to the limited range of conditions where the expression of Martin and Hill (1987b) and (Martin et al., 1991) are valid and existing gaps in the understanding of this reaction, thus, we recommend the rate expression by (Hoffmann and Calvert, 1985).

$$R_{\text{Fe},11b} = k_{10e}[\text{Fe(III)}][\text{SO}_3^{2-}] \quad (\text{for pH} < 5) \quad (11b)$$

with $k_{11e} = 1.2 \times 10^6 \text{ L mol}^{-1} \text{ s}^{-1}$.

The rate for Mn(II)-catalyzed S(IV) oxidation from Martin and Hill (1987a) is recommended:

$$R_{\text{Mn}} = \begin{cases} k_{12a}[\text{Mn(II)}][\text{S(IV)}] & \text{for S(IV)} < 10^{-4} \text{ mol L}^{-1} \\ k_{12b}[\text{Mn}^{2+}]^2 & \text{for S(IV)} > 10^{-4} \text{ mol L}^{-1} \end{cases} \quad (12)$$

where $k_{12a} = k_{12a,0}10^{-4.07\sqrt{I}/(1+\sqrt{I})} \text{ L mol}^{-1}\text{s}^{-1}$ and $k_{12b} = k_{12b,0}10^{-4.07\sqrt{I}/(1+\sqrt{I})} \text{ L mol}^{-1}\text{s}^{-1}$ with $k_{12a,0} = 10^3 \text{ L m}^{-1}\text{s}^{-1}$ and $k_{12b,0} = 680 \text{ L mol}^{-1}\text{s}^{-1}$. Note that, while Martin and Hill (1987b) and Martin and Hill (1987a) observed strong inhibition with increasing ionic strength, k_{12a} is only reported for ionic strength up to 1 mol L^{-1} . Overall, TMI-catalyzed reactions are still not very well understood, and further studies of these reactions particularly under aerosol conditions are needed.

A synergistic effect has been reported in laboratory studies when Fe(III) and Mn(II) are both present in solution (Martin, 1984; Ibusuki and Takeuchi, 1987; Martin and Good, 1991; Grgić et al., 1992), but more work must be done to reconcile the rates of R-13 from those studies with single-ion studies, and the effect of ionic strength is not known.





550 The recommended rate of R-13 is from Ibusuki and Takeuchi (1987), who investigated the effect as a function of pH and temperature:

$$R_{\text{TMI-Syn}} = \begin{cases} k_{13a}[\text{H}^+]^{-0.74}[\text{Mn(II)}][\text{Fe(III)}][\text{S(IV)}] & \text{for } 2.6 \leq \text{pH} \leq 4.2 \\ k_{13b}[\text{H}^+]^{0.67}[\text{Mn(II)}][\text{Fe(III)}][\text{S(IV)}] & \text{for } 4.2 < \text{pH} \leq 6.5 \end{cases} \quad (13)$$

where $k_{13a} = 3.72 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$ and $k_{13b} = 2.51 \times 10^{13} \text{ L mol}^{-1} \text{ s}^{-1}$.

For the sake of completeness, a more comprehensive literature overview on reaction rate constants related to TMI-catalyzed S(IV) oxidation kinetics is given in Radojevic (1992) and Brandt and van Eldik (1995).

555

4.5 NO₂ and HNO₄

NO₂ can oxidize HSO₃⁻ in the aqueous phase (Lee and Schwartz, 1983) through adduct formation, followed by decomposition, to eventually form SO₃⁻ and the weak acid HONO. The thermodynamic driving force for this process is small (Spindler et al., 2003). The reaction favors basic conditions and therefore is unlikely to be significant for most atmospheric aerosols, and self-limiting. Early studies Lee and Schwartz (1983) reported relatively high reaction rates which decreased rapidly with decreasing pH. Spindler et al. (2003) demonstrated, based on coupled gas- and aqueous phase measurements together with the direct measurement of NO₂ in aqueous solution, that the reaction between NO₂ and S(IV) proceeds first by an adduct formation equilibrium (R-14a and R-14b) followed by the adduct's unimolecular decomposition (R-15a and R-15b) to products nitrite/HONO and SO₃⁻.



570 This mechanism was invoked to explain the formation of 'artifact HONO' in a wet denuder when both NO₂ and SO₂ are present in the gas phase. The kinetic data of Spindler et al. (2003) were experimentally determined by measuring NO₂ in aqueous solution in a laser photolysis-broadband optical absorption experimental set-up and newly kinetically analyzed for the present review. The measurement was performed at pH = 4.5 and pH = 10 to investigate either the HSO₃⁻ or the fully deprotonated form SO₃²⁻. From the T-dependent rate constants in Table S2, of the forward (k_{14a} , k_{14b}) and backward reaction (k_{-14a} , k_{-14b}) the equilibrium constants (K_{14a} , K_{14b}) were calculated and the Arrhenius expressions were derived as follows.

575



At pH 10.0:

$$k_{14a}(T) = (1.4 \pm 0.2) 10^7 \text{ L mol}^{-1} \text{ s}^{-1} \quad (288 \text{ K} \leq T \leq 328 \text{ K})$$

$$k_{-14a}(T) = (3.5 \pm 0.5) 10^6 \exp[-(2440 \pm 710) \text{ K} / T] \text{ s}^{-1}$$

580 $K_{14a}(T) = (1.9 \pm 15) \exp[-(-2700 \pm 1600) \text{ K} / T] \text{ L mol}^{-1}$

At pH 4.5:

$$k_{14b}(T) = (8.5 \pm 1.9) 10^{12} \exp[-(4670 \pm 2010) \text{ K} / T] \text{ L mol}^{-1} \text{ s}^{-1}$$

$$k_{-14b}(T) = (3.8 \pm 0.5) 10^7 \exp[-(3560 \pm 680) \text{ K} / T] \text{ s}^{-1}$$

585 $K_{14b}(T) = (2.2 \pm 0.1) 10^5 \exp[-(2270 \pm 150) \text{ K} / T] \text{ L mol}^{-1} \quad (298 \text{ K} \leq T \leq 328 \text{ K})$

Finally, from the measurements of ‘artifact HONO’ in the Spindler et al. (2003) publication the unimolecular rate of decomposition for the adduct was determined as $k_{15a}(T) = (8.4 \pm 0.1) 10^{-3} \text{ s}^{-1}$ ($T = 298 \text{ K}$).

The most significant difference between the results of Spindler et al. (2003) and earlier studies is that the mechanism identified by Spindler et al. (2003) (adduct formation with a slow adduct decomposition) limits the potential for S(VI) formation via this mechanism under environmental conditions. Here, from the viewpoint of aqueous-phase thermochemistry, it should also be noted that such high rate constants for a prompt bimolecular reaction with a concerted single electron transfer from HSO_3^- to NO_2 would not be feasible. The one-electron reduction potentials of $\text{NO}_{2(\text{aq})}$ and $\text{HSO}_{3(\text{aq})}^-$ are very similar with $E^\circ(\text{HSO}_3^-) = 0.84 \text{ V}$ vs. NHE (Huie and Neta, 1984; Wardman, 1989) and $E^\circ(\text{NO}_2^-/\text{NO}_2) = 1.04 \pm 0.02 \text{ V}$ vs. NHE (Armstrong et al., 2013).

590

595

The oxidation of S(IV) by NO_2 in aerosol water was previously proposed to be important during wintertime haze episodes in Beijing (Cheng et al., 2016; Wang et al., 2016). The significance of this S(IV) oxidation pathway rests on (a) the hypothesis that aerosols in Beijing have unusually high pH (Wang et al., 2016), which is not supported by thermodynamic models, and (b) the mechanism and relatively fast kinetic parameters of earlier studies by (Lee and Schwartz, 1983) and Clifton et al. (1988) without considering the more recent findings of Spindler et al. (2003).

600

Recent isotopic studies provide further evidence that this reaction is not important in Beijing (Au Yang et al., 2018; He et al., 2018a; Shao et al., 2019; Li et al., 2020a) in line with the aforementioned mechanistic and thermodynamic considerations.

The importance of the $\text{NO}_2 + \text{HSO}_3^-$ reaction has also been highlighted for fogs in China with $\text{pH} > 5$ (Xue et al., 2016; Xue et al., 2019). However, as with the aerosol aqueous chemistry, this sulfate production pathway should be self-limiting due to its production of H^+ .

605


Peroxynitric acid (HNO_4), a product of the gas-phase reaction of HO_2 and NO_2 , also oxidizes HSO_3^- , primarily in cloud water, with a rate constant of $3.3 \times 10^5 \text{ L mol}^{-1} \text{ s}^{-1}$ (Amels et al., 1996; Warneck, 1999; Dentener et al., 2002). The reaction rate increases with increasing aqueous pH due to increased solubility of S(IV) and HNO_4 . Besides the acidifying effect of S(IV) to S(VI) conversion, the reaction yields nitric acid (HNO_3) as an acidic byproduct. The significance of this pathway depends on gas-phase HO_x and NO_x levels and the relative abundance of other competing S(IV) oxidants.

610



4.6 Overall S(IV) oxidation considerations

To compare the potential atmospheric relevance of the different S(IV) to S(VI) conversion pathways with respect to different environmental and acidity regimes in aerosols, haze and clouds, initial S(IV) oxidation rates of the different pathways discussed up to here were calculated. Figure 7 shows the resulting calculated S(IV) oxidation rates of these reaction pathways in mol L⁻¹ s⁻¹ for continental urban haze and rural aerosol conditions as well as continental urban and rural cloud conditions. These rates were calculated with the rate expressions from the subsections above (Eq.s 6a, 7, 8, 9, 10, 11b, 12) and are based on the typical conditions as summarized in Table 1. For the NO₂, kinetic rates were calculated applying the pseudo-steady-state approximation ($k_{\text{PSSA,HSO}_3^-} = 1.3 \times 10^1 \text{ L mol}^{-1} \text{ s}^{-1}$, $k_{\text{PSSA,SO}_3^{2-}} = 2.7 \times 10^2 \text{ L mol}^{-1} \text{ s}^{-1}$). For HNO₄, the reaction rate was calculated with a rate constant of $3.3 \times 10^5 \text{ L mol}^{-1} \text{ s}^{-1}$ (Amels et al., 1996; Warneck, 1999; Dentener et al., 2002). For Fe(III) and Mn(II), the rate expressions by Hoffmann and Calvert (1985) and Martin and Hill (1987a) were applied, respectively. Note that the synergistic rates of Ibusuki and Takeuchi (1987) (Eq. 13) were not used due to the still large uncertainties of this oxidation pathway.



625 **Figure 7.** Calculated S(IV) oxidation rates of different reaction pathways in $\text{mol L}^{-1} \text{s}^{-1}$ for urban winter haze (a) and rural aerosol (b) conditions as well as urban (c) and rural (d) cloud conditions at 298 K. Applied conditions are given in Table 1 and the rate expressions used were those given in this text. The atmospherically relevant acidity range in the different cases is marked in yellow.

For diluted cloud conditions, the S(IV) oxidation by dissolved H_2O_2 , O_3 , HNO_4 and potentially the iron-catalyzed pathway are the most important oxidation pathways (see Fig. 7c and 7d). The reaction with dissolved H_2O_2 is the major oxidation pathway under acidic cloud conditions. Under less acidic cloud conditions ($\text{pH} > 5$), the other reaction pathways are able to contribute

630



significantly to the S(VI) formation. Moreover, Fig. 7 displays that the oxidation rates of other oxidants such as NO₂, excited triplet states of photosensitizers (PS*) and organic hydroperoxides (CH₃COOH, CH₃C(O)OOH) are unimportant under cloud conditions mainly because of their low in-cloud concentrations.

635 Differently, under more concentrated haze and deliquesced aerosol conditions, the molar concentrations of TMIs are significantly higher. Thus, the contributions of TMI-catalyzed S(IV) oxidation pathways are elevated against cloud conditions. From the calculation output in Fig. 7a and 7b, it can be seen that the S(IV) oxidation by dissolved H₂O₂ is still predominant below pH ≤ 3. However already at quite low acidity conditions with pH ≈ 3.5, the TMI-catalyzed pathways can become the main oxidation route for S(IV). Note that the synergistic rate of Ibusuki and Takeuchi (1987) ((Eq. 13) were not included in
640 the current study, so even higher contributions of TMI-catalyzed S(IV) oxidation pathways can be possible. Besides the TMI-catalyzed S(IV) oxidation pathways, also S(IV) oxidations by dissolved HNO₄ and O₃ as well as, to some extent, PS* can be important under polluted haze and rural aerosol conditions when pH is above pH > 5, respectively. Importantly, the current comparison clearly shows that the NO₂-driven oxidation route even under very high NO_x conditions (66 ppb) applied in the urban haze case still remains of minor importance. Only by the combination of applying unusually high aerosol pH
645 values, artificially low H₂O₂ concentrations and unrealistically fast kinetic parameters from earlier studies by Clifton et al. (1988), NO₂ rates can fall into the range of other key oxidants discussed here.

In conclusion, the outcomes of this overall comparison are in agreement with findings of isotope field investigations (see e.g., Harris et al. (2013); Au Yang et al. (2018); He et al. (2018a); Shao et al. (2019); Li et al. (2020a)) which have implicated that mainly H₂O₂, O₃, and TMI-catalyzed pathways are responsible for the S(IV) to S(VI) conversion in atmospheric aqueous-
650 phase cloud and aerosol solutions. However, due to the uncertainties still existing with regard to kinetics and mechanisms further acidity-dependent investigations are warranted.

4.7 Sequestering of S(IV) as HMS


HSO₃⁻ or SO₃²⁻ can react with a variety of aldehydes to form hydroxyalkylsulfonates (Olson and Hoffmann, 1989). Of particular interest has been S(IV) reaction with HCHO to produce hydroxymethanesulfonate (HMS, HOCH₂SO₃⁻) (Munger et al., 1986).
655 The formation of HMS is strongly dependent on drop acidity, increasing rapidly at higher pH values due to increased partitioning of S(IV) to HSO₃⁻ and SO₃²⁻ (Rao and Collett, 1995). Furthermore, the reaction rate increases with increasing pH due to the fact that the rate coefficient for SO₃²⁻ ($k = 2.5 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$) is more than four orders of magnitude higher than that for HSO₃⁻ ($k = 790 \text{ L mol}^{-1} \text{ s}^{-1}$) (Boyce and Hoffmann, 1984; Olson and Hoffmann, 1989). At pH values > 6, HMS formation becomes so fast that it can limit aqueous sulfate production in large droplets where mass transport limits SO₂ uptake from the gas phase (Reilly et al., 2001). Since oxidation of HMS is slow (Hoigne et al., 1985; Kok et al., 1986; Barlow et al., 1997a, b), its formation effectively protects S(IV) from oxidation to S(VI) by non-radical oxidants such as H₂O₂, O₃ and others. Whiteaker and Prather (2003) demonstrated the utility of HMS measurements in single particles as a tracer for fog processing. Recent field and modeling studies have suggested that HMS production may also be an important contributor to fine particle sulfur content under polluted haze conditions (Moch et al., 2018; Song et al., 2019b; Ma et al., 2020; Moch et al., 2020).



665 Sulphur in particles may exist in the form of other sulphonates ($R-C-SO_3^-$) besides organosulphates ($R-C-O-SO_3^-$) (Le Breton et al., 2018; Brüggemann et al., 2020).

4.8 Acid-driven production of tropospheric reactive halogens: Multiphase halogen activation

670 Of the many acid-catalyzed reactions in the atmosphere, the acid-catalyzed formation of reactive halogens (Br, Cl, I) in the troposphere has the potential to render acidity as an influencer of the oxidative capacity of the atmosphere, although its influence has yet to be fully quantified. Reactive halogens and halogen reservoir species are of the form Br_y ($= Br + 2 Br_2 + HOBr + BrO + HBr + BrNO_2 + BrNO_3 + IBr + BrCl$), Cl_y ($= Cl + 2 Cl_2 + HOCl + ClO + HCl + ClNO_2 + ClNO_3 + ICl + BrCl + ClOO + OClO + 2 Cl_2O_2$), and I_y ($I + 2 I_2 + HOI + IO + OIO + HI + HIO_3 + INO + INO_2 + INO_3 + 2 I_2O_2 + 2 I_2O_3 + 2 I_2O_4$). Tropospheric reactive halogens can impact the oxidation capacity of the atmosphere by acting as an effective sink for ozone (O_3), e.g. bromine explosion events in the Arctic, and nitrogen oxides ($NO_x = NO + NO_2$) and by influencing the partitioning of HO_x ($= OH + HO_2$) (Oltmans et al., 1989; Simpson et al., 2015; Schmidt et al., 2016; Sherwen et al., 2016; Hoffmann et al., 2019a). Reactive halogens also directly impact the lifetime of reduced trace gases such as methane (CH_4) and non-methane volatile organic compounds (VOCs), dimethylsulfide (DMS), and mercury in the atmosphere (Barnes et al., 2006; Saiz-Lopez and von Glasow, 2012; Ariya et al., 2015; Simpson et al., 2015). Sources of tropospheric reactive halogens include oxidation of organohalogens (e.g., CH_3Br and CH_3I) (Saiz-Lopez et al., 2012b; Saiz-Lopez and von Glasow, 2012), deposition of ozone to the ocean surface to yield HOI and I_2 (Carpenter et al., 2013), release from sea salt aerosols (Parrella et al., 2012; Schmidt et al., 2016; Sherwen et al., 2017) and, to a minor extent, transport from the stratosphere (Schmidt et al., 2016; Wang et al., 2019b). Liberation of halogens to their reactive form via acid-catalyzed reactions on sea salt aerosols (see Fig. 8) is the largest source of reactive bromine in the troposphere (Vogt et al., 1996; von Glasow et al., 2002a; Pechtl et al., 2007; Pechtl and von Glasow, 2007; Parrella et al., 2012; Chen et al., 2017). As shown in Fig. 8, the formation and processing of reactive halogens strongly depends on the aqueous-phase conditions, i.e. the LWC and the acidity of the solution.



690 **Figure 8: Simplified scheme of the reactive halogen chemistry and their differences between diluted less acidic cloud conditions and more concentrated and acidic aerosol conditions. Differences in the chemical rates and uptake fluxes are indicated by lighter and thicker arrows.**

Formation of reactive halogens (Br, Cl, I) from sea salt aerosols proceeds in pristine environments via the uptake of hypohalous acid species (HOX, where X = Br, Cl, or I) from the gas phase (von Glasow et al., 2002b) or in more polluted environments via the hydrolysis of N₂O₅ forming ClNO₂ (Finlayson-Pitts et al., 1989; Roberts et al., 2009; Sarwar et al., 2014) as well as via
 695 the hydrolysis of XNO₃ forming HOX (Schmidt et al., 2016; Hoffmann et al., 2019b), see reactions 16a-16c:



where X = Br, Cl, or I. If two different halogens are involved, Y denotes the second halogen atom. The formed XX or XY
 700 species then either reacts further or partitions to the gas phase, where it is photolyzed and participates in gas-phase oxidation chemistry to ultimately regenerate HOX or XNO₃. Non-linear reactive halogen production proceeds via uptake of one molecule HOX or XNO₃, carrying one halogen atom, yielding two halogen atoms released back to the gas phase (see Fig. 8). Note this is an acid-driven process which consumes H⁺ in the aqueous-particle phase without recycling it and also one halogen anion Y⁻ is consumed. When Br is a participant, this auto-catalytic reaction cycle can lead under high bromide concentrations to so-called bromine explosion events characterized by high concentrations of BrO (Evans et al., 2003) resulting from the gas-phase
 705 reaction of Br with O₃.



Changing atmospheric acidity due to changes in anthropogenic emissions of acid precursor gases may influence the formation of reactive halogens via reaction (R-16a) (Keene et al., 1998). However, lower acidity conditions might also result in stronger aqueous-phase partitioning of hydrogen halides which might partly compensate for the reduced acidity effect via reaction (R-16a). Changes in sulfur dioxide (SO₂) may contribute to sources or sinks of reactive halogens. The formation of sulfate, from the oxidation of SO₂, is typically the largest source of acidity in the atmosphere (see Sect. 3). However, reactions of HOX with dissolved S(IV) (HSO₃⁻ + SO₃²⁻) in aqueous aerosols can convert halogens to their less-reactive acid form (HX) via R-17 and R-18 (Fogelman et al., 1989; Troy and Margerum, 1991; von Glasow et al., 2002a; Chen et al., 2017; Liu and Abbatt, 2020). Here, especially the reaction with HOI can be very significant (Pechtl and von Glasow, 2007; Bräuer et al., 2013; Hoffmann et al., 2019a).



The rate expression for S(VI) formation by S(IV) + HOX is given by:

$$R_{\text{HOX},1} = k_{17,\text{HOX}}[\text{HSO}_3^-][\text{HOX}] \quad (17)$$

$$R_{\text{HOX},2} = k_{18,\text{HOX}}[\text{SO}_3^{2-}][\text{HOX}] \quad (18)$$

with recommended rate constants for HOCl of $k_{17,\text{HOCl}} = 2.8 \times 10^5 \text{ L mol}^{-1} \text{ s}^{-1}$ (Liu and Abbatt, 2020) and $k_{18,\text{HOCl}} = 7.6 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$ (Fogelman et al., 1989), and for HOBr of $k_{17,\text{HOBr}} = 2.6 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$ (Liu and Abbatt, 2020) and $k_{18,\text{HOBr}} = 5.0 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$ (Troy and Margerum, 1991), respectively. Unfortunately, reaction rate constants for HOI with dissolved S(IV) (HSO₃⁻ + SO₃²⁻) have not been measured yet. However, following the augmentation of Pechtl et al. (2007), the reaction rate constants of HOI with HSO₃⁻ and SO₃²⁻ should be even faster than the reaction rate constants of HOCl and HOBr or is likely diffusion-limit controlled.

Finally, the overall impact of changes in anthropogenic emissions of SO₂ or other acid-gas precursors on tropospheric reactive halogen production remains unknown. Because of the impact of reactive halogens on the radiative forcing of the powerful greenhouse gas ozone (Saiz-Lopez et al., 2012a) as well as aerosol particle composition (Hoffmann et al., 2016; Lee et al., 2019), their chemistry can be of crucial importance for climate predictions. Therefore, more laboratory investigations, chamber studies and accompanied modelling efforts are needed to determine chemical reaction rate constant of crucial halogen processes, such as the oxidation of S(IV) by HOI, and to better characterize the overall reactive cycling of halogens including its sensitivity to aerosol particle and cloud acidity.

4.9 Discussion and outlook: atmospheric multiphase chemistry of inorganic species

Multiple reactive pathways for the conversion of S(IV) to S(VI) have been discussed here. Many of these processes are limited in atmospheric aerosols by acidic conditions and the presence of particle-phase organics, which quench highly reactive radical and triplet species. Studies from the past four decades have shown that, under polluted conditions, such as found in urban areas worldwide or in the North China Plain (NCP), only relatively stable oxidants or TMI catalysis may lead to the required rate of



S(IV) to S(VI) conversion to explain the observed S(VI) budgets (Jacob and Hoffmann, 1983; Chameides, 1984; Saxena and
740 Seigneur, 1987; Seigneur and Saxena, 1988; Pandis et al., 1992; Amels et al., 1996; Berglund and Elding, 1996). That being
said, our understanding of atmospheric multiphase sulfate production, especially in the aerosol phase, is still incomplete,
despite more than a century of studies on aqueous sulfur oxidation. S(IV) conversion explaining the aerosol sulfate budgets
encountered today, especially under urban or semi-urban polluted conditions still need further elucidation from the basic
aqueous-phase processes to concrete field measurements. This includes the role of acidity in these processes which could be
745 decisive to whether or not a process can really be important in the environment.

Areas of focus should include:

- a) Laboratory studies of S(IV) oxidation by all pathways under atmospheric aerosol conditions, i.e. in aerosol flow tube
reactors, to assess the impact of high ionic strength and other factors specific to the aerosol phase
- b) Advanced sulfur-isotope measurements of ambient aerosol and cloud water samples to identify driving sulfur oxidation
750 pathways under various atmospheric conditions
- c) Advanced knowledge of TMI-catalyzed S(IV) oxidation pathways, including investigation of synergy effects and the role
of other metal catalysts present in aqueous atmospheric solutions besides Fe and Mn. The impact of acidity and ionic
strength on both the speciation of TMIs, i.e. their presence in free and complexed form, and the specific chemical reaction
rates of single-TMIs have to be studied.
- 755 d) Kinetic and mechanistic investigations on other potential oxidants, especially comparatively stable oxidants such as
ROOHs and HOI
- e) Investigations of pH-dependent in-situ formation of key S(IV) oxidants such as H₂O₂ and ROOH resulting from TMI-
HO_x-DOM chemistry.

5 Interactions of acidity and chemical processes: Organic systems

760 Acidity in aerosol particles can strongly enhance secondary organic aerosol (SOA) formation (Jang and Kamens, 2001; Jang
et al., 2002; Jang et al., 2003; Iinuma et al., 2004; Jang et al., 2004; Liggió and Li, 2006; Surratt et al., 2007b). These early
observations triggered immense research interest in investigating aqueous-phase reactions leading to the accumulation of
organic particle constituents. These so-called ‘accretion reactions’ are often acid-driven, or even acid-catalyzed. In the
following, the most important organic compound families and the influence of acidity on their aqueous-phase chemistry are
765 discussed. In this section we discuss the role of acidity on the gas-particle partitioning of semi-volatile organic compounds
through its influence on hydration of carbonyls and dicarbonyls. We then discuss in detail the impact of acidity on the
multiphase oxidation of organic material. Oxidative organic chemistry can be influenced by acidity because this influences the
reactant speciation, such as in acid and diacid oxidations by radicals and non-radical oxidants such as dissolved ozone. Finally,
we discuss accretion reactions.

770



5.1 Acidity and hydration of carbonyl compounds

775 Carbonyl compounds (i.e., aldehydes or ketones), omnipresent in the tropospheric gas and aqueous phase, result from primary emissions or are secondary oxidation products. The photolysis of carbonyl compounds can be important for both their degradation in the troposphere and gas-phase oxidant production. Water-soluble carbonyl compounds may partition into the aqueous phase of deliquesced aerosols and cloud/fog droplets. Once in the aqueous phase, carbonyl compounds can undergo hydration, leading to conversion of carbonyl group into gem-diol moieties. As hydration processes are typically acid- or base-catalyzed, the acidity of an aqueous solution can affect the hydration and consequently all other processes linked to it. With regard to phase partitioning, the hydration equilibria increase the effective partitioning of the carbonyl-containing compound towards the aqueous phase (Sumner et al., 2014). Moreover, compared to the carbonyl group, the diol functionality is

780 photochemically inactive. Thus, partitioning to the aqueous phase and subsequent hydration can, in part, protect carbonyl compounds from photolysis and shut off possible photochemistry of the carbonyl group (George et al., 2015; Herrmann et al., 2015; McNeill and Canonica, 2016). However, hydrated aldehydes are often characterized by a somewhat lower reactivity with radical oxidants such as OH compared to the unhydrated carbonyl species (Schuchmann and von Sonntag, 1988).

785 This sub-section summarizes the present knowledge on the acidity dependence of carbonyl hydration constants, and implications for the chemical conversions of carbonyl compounds in atmospheric aqueous media.

5.1.1 The influence of acidity on hydration constants and its implications

The reversible hydration and dehydration of the carbonyl group of an aldehyde or ketone in the aqueous phase is illustrated in Fig. 9 (Bell and Darwent, 1950; Bell, 1966; Ogata and Kawasaki, 1970; Lowry and Richardson, 1976).




Figure 9: General mechanism of the acid- (A) or base- (B) catalyzed formation of diols resulting from the hydration of the carbonyl group.



Simple aldehydes such as formaldehyde (Zavitsas et al., 1970; Li et al., 2011; Rivlin et al., 2015), acetaldehyde (Ahrens and
795 Strehlow, 1965; Kuschel and Polarz, 2010) and glyoxal (Liggio et al., 2005a; Loeffler et al., 2006) tend to self-oligomerize,
e.g. via hemiacetal formation or aldol condensation, which further influences the hydration equilibrium. .

The ratio of the hydrated and dehydrated fraction under equilibrium conditions is described by equilibrium constant K_{hyd} ,
defined as follows for a dilute aqueous solution:

$$800 \quad K_{\text{hyd}} = \frac{[\text{diol compound}]}{[\text{carbonyl compound}]} = \frac{k_{\text{hyd}}}{k_{\text{dehyd}}} \quad (19)$$

Where k_{hyd} is the rate constant for hydration and k_{dehyd} is the rate constant for dehydration. In general, the hydration constants
 K_{hyd} decrease with decreasing electron-withdrawing power of the substituent in a substituted organic acid (Clayden et al.,
2012). The equilibrium constants of simple aldehydes or ketones generally show no pH dependence but are dependent on
temperature.

805 For most carbonyls, the hydration reaction with H_2O under neutral conditions is slow. In the presence of hydrogen ions,
hydroxyl ions, undissociated acid molecules and anion bases, the hydration reaction proceeds faster. The overall hydration rate
considering all acid and base dependencies can be calculated by means of Eq. 20 (Ogata and Kawasaki, 1970; Lowry and
Richardson, 1976):

$$k_{\text{hyd}} = k_0 + k_{\text{H}^+}[\text{H}_3\text{O}^+] + k_{\text{OH}^-}[\text{OH}^-] + \left(k_a + k_b \frac{[\text{B}]}{[\text{HA}]}\right) \quad (20)$$


810 The catalytic constants of the hydration rate in equation (11) are described as follows: k_0 for the solvent influence, k_{H^+} for the
effect of the H_3O^+ ion, k_{OH^-} as influence of the OH^- ion, k_a and k_b as general acid or base contribution (Ogata and Kawasaki,
1970). An overview of the acid or base catalytic constants for the hydration of formaldehyde and acetaldehyde by a few
different organic acids is presented in Table 2. As can be seen from the data compiled in Table 2, the presence of acids clearly
influences the hydration rate of the carbonyl group.

815 Experimentally determined values for K_{hyd} and our recommended values are presented for several atmospherically relevant
simple aldehydes, ketones and α -oxocarboxylic acids in the Supporting Information Tables S3 and S4. The general trend in
 K_{hyd} is: glyoxal (2nd hydration) > formaldehyde > methylglyoxal (CHO-group hydration) > glyoxylic acid > glyoxal
(1st hydration) > glyoxylate > glycolaldehyde > pyruvic acid > biacetyl > acetaldehyde > propanal > butanal > pivaldehyde
> pyruvate > acetone. These data are discussed in more detail in the Supporting Information.

820 It is important to note that the hydrolysis of simple aldehydes and ketones and dicarbonyls is unaffected by pH. For
multifunctional carbonyl compounds, the hydration equilibrium constant of the carbonyl group is strongly influenced by the
electronic effects of the adjacent group. The hydration of carbonyl groups in compounds that also contain pH sensitive
moieties, such as α -oxocarboxylic acids, is highly influenced by the acidity of the surrounding environment.


Besides the hydration and dissociation equilibria, condensation (dimerization or polymerization) equilibria as well as keto-
825 enol-equilibria could influence these compounds (Fig. 10). The equilibria are related by $K_{\text{hyd},1} \times K_{\text{diss},2} = K_{\text{diss},1} \times K_{\text{hyd},2}$, while
the apparent dissociation constant is given by

$$K_{Hyd} = \frac{[H^+] \times K_{diss,1} + K_{hyd,2} \times K_{diss,1}}{[H^+] + K_{diss,1}} \quad (21)$$



830 **Figure 10:** Scheme describing the four equilibria of pyruvic acid, a representative α -keto-carboxylic acid, in aqueous solution.

An overview of the determined K_{hyd} values for atmospherically relevant α -oxocarboxylic acids is given in Table S4 and the existing data are outlined for each of the listed chemical compounds separately in the Supplement. The most prominent α -oxocarboxylic acid compounds in the atmosphere are glyoxylic acid and pyruvic acid. Special emphasis in the recent literature was put on pyruvic acid. The recent data on pyruvic acid is summarized in Fig. 11.



835

Figure 11: Measured pH dependency of the apparent hydration equilibria (see Eq. 21) of pyruvic acid in aqueous solution. Experimental data are shown from the literature (Pocker et al., 1969; Reed Harris et al., 2014; Lopalco et al., 2016; Rapf et al., 2017a; Rapf et al., 2017b).



As shown in Fig. 11, K_{hyd} for pyruvic acid increases rapidly with decreasing pH for $\text{pH} < 3$. Note that the formation of the
840 hydrated pyruvic acid (2,2-dihydroxypropanoic acid) is also dependent on the water concentration (Pocker et al., 1969; Maron
et al., 2011) – which may have implications for aqueous aerosol chemistry.

The impact of acidity and its feedback on the hydration, as well as their impact on the photochemistry of pyruvic acid have
been examined by spectroscopic investigations performed at TROPOS. These investigations have shown that the molar
absorption coefficient spectra of pyruvic acid are rather different under low and high acidity conditions. Measured absorption
845 coefficient spectra of pyruvic acid at $\text{pH} = 0$ and $\text{pH} = 9$ (Fig. 12) shows higher absorption coefficients under $\text{pH} = 9$
conditions.




Fig. 12. Measured UV absorption coefficient spectra of pyruvic acid in water under acidic ($\text{pH} = 0$, red line) and alkaline
($\text{pH} = 9$, blue line) conditions.

850

At 300 nm wavelength, the measured absorption coefficient is about 4 times larger at $\text{pH} = 9$ than at $\text{pH} = 0$. Under high pH
conditions ($\text{pH} = 9$), a large fraction of pyruvic acid is present in its unhydrated form and, consequently, higher absorption
coefficients are observed compared to very acidic conditions, where pyruvic acid is mainly present in its photochemically
inactive hydrated form. This difference has implications for photochemistry of pyruvic acid which will become less efficient
855 in more acidic solutions compared to less acidic ones. Such effects should be implemented into aerosol liquid water chemistry
models. Finally, hydration processes can be characterized by both temperature and acidity dependencies particularly for
 α -/ β -keto-carboxylic acids such as pyruvic acid. These dependencies need to be included in future models to be able to
accurately investigate their impact on the partitioning and occurring multiphase chemistry. For that reason, more laboratory
studies are needed to extend the available database for other atmospherically-relevant functionalized α -/ β -keto-carboxylic



860 acids. In concrete terms, more studies appear desirable for glyoxylic acid, mesoxalic acid, oxalacetic acid and oxalglycolic acid.

5.2 pH-sensitive organic accretion reactions

Organic accretion reactions are considered to be a source of high-molecular-weight organic material in atmospheric aerosols, playing a key role in the formation of secondary organic aerosol material (Barsanti and Pankow, 2004, 2005, 2006). These reactions are typically multistep, bond-forming reactions, and are highly pH-sensitive. Many organic accretion reactions are acid- or base-driven or, in some cases, even catalyzed. Others require acidity, e.g. in the form of the participation of a proton (H^+) in the reaction, but are not technically ‘acid catalyzed’ because the proton is incorporated into the formed reaction product (e.g., ring-opening of epoxides, cf. Sect. 5.3)). Examples of atmospherically important accretion reactions include (i) aldol condensation, (ii) hemiacetal and acetal formation, and (iii) esterification of carboxylic acids which will be treated in the following subsections.

The current kinetic and mechanistic knowledge on tropospheric accretion reactions has been summarized not too long ago in a review by Herrmann et al. (2015) and a book by Barker et al. (2016). Accordingly, the present subsection only briefly outlines the mechanisms and emphasizes their dependence on acidity. For further specific details on organic accretion reactions please see Herrmann et al. (2015), Zhao et al. (2016) and references therein.

5.2.1 Aldol condensation and ammonium catalyzed reactions

Overview

The Aldol (short form of ‘aldehyde alcohol’) condensation is a carbon-carbon-bond formation requiring the participation of an enolizable carbonyl compound (e.g., ketones or aldehydes with an α -hydrogen) (Loudon, 1995). Under acidic conditions, the enol reacts with a protonated carbonyl compound to form the aldol addition product. The product may be dehydrated to form the aldol condensation product, a conjugated enone compound. The acid- or base-catalyzed nature of the enol formation, as well as the role of the protonated carbonyl reactant under acidic conditions, make this family of reactions pH-sensitive. This pathway has been suggested as a source of light-absorbing secondary organic material (i.e., brown carbon) in atmospheric aerosols (Laskin et al., 2015; Nozière et al., 2015), which has been discussed in more detail for acetaldehyde, glyoxal as well as methylglyoxal (Laskin et al., 2015; Nozière et al., 2015), which was discussed in more detail for acetaldehyde, glyoxal as well as methylglyoxal (Nozière and Esteve, 2005; Nozière et al., 2007; Nozière and Esteve, 2007; Nozière and Cordova, 2008; De Haan et al., 2009a; Shapiro et al., 2009; Bones et al., 2010; Sareen et al., 2010; Li et al., 2011; Yu et al., 2011; Kampf et al., 2012; Nguyen et al., 2012; Laskin et al., 2015; Lin et al., 2015; Maxut et al., 2015; Nozière et al., 2015; Van Wyngarden et al., 2015; Aiona et al., 2017; Rodriguez et al., 2017). Several studies which focused on sulfuric acid-catalyzed aldol formation have shown that this chemistry occurs efficiently only under strongly acidic conditions (Duncan et al., 1999; Kane



et al., 1999; Imamura and Akiyoshi, 2000; Nozière and Riemer, 2003; Esteve and Noziere, 2005; Noziere and Esteve, 2005; Liggio and Li, 2006; Noziere et al., 2006; Casale et al., 2007; Noziere and Esteve, 2007; Krizner et al., 2009).

Surface films

895 Van Wyngarden et al. (2015) reported on the formation of surface films from H₂SO₄-propanal mixtures with or without glyoxal and/or methylglyoxal. These films tended to form faster when the acidity was increased up to 48 wt % H₂SO₄, but with an acidity of 76 wt % H₂SO₄, the film formation slowed down or even stop in all mixtures except propanal/glyoxal.

Mechanistic and kinetic considerations

900 Yasmeen et al. (2010) suggested that the favorable mechanism under acidic conditions pH < 3.5 is the acetal formation, while the aldol condensation only occur at a pH = 4-5, which is in contrast to the above-mentioned conditions. Ammonium-catalyzed or amine-catalyzed aldol condensation proceeds under higher, but still acidic, pH values typical for tropospheric aerosol particles (Noziere and Cordova, 2008; De Haan et al., 2009a; Noziere et al., 2010; Sareen et al., 2010; Li et al., 2011; Sedehi et al., 2013; Powelson et al., 2014; Aiona et al., 2017). The pH-dependent rate constants for aldol condensation reactions of glyoxal and methylglyoxal with ammonium sulfate and amines have been further investigated in several studies (Noziere et al., 2009; Sareen et al., 2010; Yu et al., 2011; Kampf et al., 2012; Sedehi et al., 2013; Powelson et al., 2014; Yi et al., 2018).
905 Noziere et al. (2009) observed that the pathway via iminium ion is faster at higher pH, whereas the aldol condensation is favored at lower pH values, which suggests a pH dependency incorporation of N-containing products. The results of pH dependency appear to be contradictory. On the one hand, Sareen et al. (2010) reported an enhanced product formation by decreasing the pH and concluded an acid-catalyzed aldol formation of the light-absorbing product at 280 nm. On the other
910 hand, Yu et al. (2011) reported an exponential increase of the formation rate of condensation products, e.g. imidazole by increasing the pH and concluded an ammonium-catalyzed reaction. Similarly, Kampf et al. (2012) observed a higher production rate of the imidazole bicycle with increasing pH values. Furthermore, Sedehi et al. (2013), showed a strong pH-dependence with an increasing reaction rate proportional to the concentration of the deprotonated amine or in other words an increase of the pH value. Nevertheless, the pH-dependent character of the reaction of ammonium sulfate or amine reaction
915 with glyoxal is stronger than for methylglyoxal. A study by Yi et al. (2018), describes an acceleration of the pH-dependent reaction of ammonium sulfate or amine in the presence of glycolaldehyde, whereas no cyclic compounds (e.g. imidazole) were formed in this reaction. Powelson et al. (2014), Grace et al. (2019) and (Li et al., 2019) reported the formation of heterocyclic compounds under similar conditions. Hawkins et al. (2018) reported an increasing formation of pyrazine-based chromophores in an aqueous mixture containing methylglyoxal and ammonium sulfate by increasing the pH from 2 to 9. This indicates that
920 the nitrogen nucleophile is more important than the acid-catalyzed aldol condensation, which is consistent with the observation of Kampf et al. (2012), Kampf et al. (2016) and Yi et al. (2018). A theoretical analysis of glyoxal condensation in the presence of ammonia conduct by Tuguldurova et al. (2019) describes two different imidazole formation pathways: the imine pathway and the aminoethanetriol pathway. These authors reported that the imine concentrations are very low due to the high-energy barriers for imine formation. Although a pH decrease due to amino alcohol dehydration leads to higher imine concentrations,
925 it also leads to higher ammonium cation formation, which is another difficulty for ammonium addition to the carbonyl group.



The second proposed pathway, which includes the intermediate aminoethanetriol, has a lower energy barrier and appears to be kinetically favourable due to the higher concentrations. Finally, imidazole formation is determined by the glyoxal concentration, the ratio of glyoxal/ (amine or ammonium), the composition of the solvent, and the pH value. All in all, aldol condensations are today generally regarded as demanding to drastically acidic conditions to be really important in particle and multiphase chemistry.

5.2.2 Hemiacetal and acetal formation

Hemiacetal and acetal formation are the addition of an organic molecule containing either one or two hydroxyl groups (e.g. alcohols) to a carbonyl compound leading to the formation of one(hemiacetal) or two (acetal) ether-type C–O–C bonds. This type of accretion reaction is significant for aqueous secondary organic aerosol formation involving glyoxal, methylglyoxal, acetaldehyde, formaldehyde, and other common atmospheric carbonyl-containing compounds (Schweitzer et al., 1998; Tobias and Ziemann, 2000; Jang et al., 2002; Kalberer et al., 2004; Hastings et al., 2005; Liggio et al., 2005b, a; Loeffler et al., 2006; Zhao et al., 2006; De Haan et al., 2009a; De Haan et al., 2009b; Shapiro et al., 2009; Sareen et al., 2010; Yasmeeen et al., 2010; Li et al., 2011). Hemiacetal formation is initiated by protonation of a carbonyl group, followed by nucleophilic addition of the alcohol (Loudon, 1995). After the deprotonation of the attacking alcohol, the hemiacetal is formed. Promoted by acidity, the hemiacetal can react further to a full acetal, by protonation of the alcohol group of the hemiacetal to eliminate water again under the formation of a carbocation. This carbocation can react in a subsequent reaction with another alcohol molecule to form the full acetal by deprotonation of the hydroxyl group. Hemiacetal and acetal formation are reversible. In addition to the aldol condensation product, Liggio et al. (2005a, 2005b); and Liggio and Li (2006) reported on an acetal formation during the reactive uptake of glyoxal and pinonaldehyde on acidic aerosols. It has been reported by De Haan et al. (2009b) that glyoxal is more prone to undergo the acetal formation, while methylglyoxal reacts mainly by the aldol condensation reaction mentioned above, whereas the contribution of oligomer formation was strongly dependent on the relative humidity and hence the particulate water concentration. Holmes and Petrucci (2006, 2007) observed the formation of hemiacetals in the oligomerization process of levoglucosan induced by the Fenton chemistry. Noziere et al. (2009, 2010) observe a pH dependent ammonium-catalyzed acetal formation from glyoxal and acetaldehyde. The hydration and the subsequent acetal formation involving methylglyoxal is strongly dependent on the pH value and occur at a pH < 3.5 (Yasmeeen et al., 2010). Maxut et al. (2015) investigated also the ammonium catalyzed imidazole formation with glyoxal in neutral aqueous solution concluded that the contribution of acetal oligomer formation pathway is small. Grace et al. (2019) referred to the study by De Haan et al. (2011) and Kampf et al. (2016), who reported that aldol condensation type reactions are more important than acetal or hemiacetal formation under atmospheric conditions. In summary, hemiacetal and acetal formation requires acidic conditions, but the contribution of this reaction pathway is small compared to aldol formation under atmospheric conditions.



5.2.3 Esterification of carboxylic acids

Esterification is a reversible, acid- or base-catalyzed condensation reaction of carboxylic acids and hydroxyl group containing molecules under the formation of an C(O)–O–C type bond (Ingold, 1969; Larson and Weber, 1994). The acid-catalyzed mechanism can be described as follows: The carbonyl group of the undissociated carboxylic acid can be protonated under acidic conditions to form a carbocation. The carbocation then is subject to a nucleophilic attack by a hydroxyl group-containing molecule. The resulting intermediate further reacts by tautomerization (proton shift in the molecule), which subsequently decays in an equilibrium reaction to a protonated ester and a water molecule (Loudon, 1995).

The base-catalyzed mechanism includes the reaction of a carboxylate group (resulting from the deprotonation of carboxylic acid group) and a hydroxyl group-containing molecule, such as alcohol. First, a proton transfer from the alcohol to the carboxylate occurs. Second, the deprotonated hydroxyl group reacts in a nucleophilic attack with the carbon atom of the carboxylic acid forming a metastable intermediate, which subsequently decays to an ester molecule and a hydroxide ion in an equilibrium reaction. In addition to being pH-sensitive, esterification reactions are also strongly dependent on the water content. The majority of esters are hydrolyzed in the presence of water. Both the formation and the hydrolysis of esters are slow processes under tropospheric conditions. Moreover, the hydrolysis rate of esters will increase with increasing acidity (Mabey and Mill, 1978). Altieri et al. (2006) and (2008) reported the esterification mechanism occur in the cloud processing of pyruvic acid (pH = 2.7-3.1) and methylglyoxal (pH = 4.2-4.5). The oxidation leads to carboxylic acids and proceeds through α - or β -hydroxy acid to esters or oligoesters, similarly to the proposed mechanisms for oligomers in the aerosol phase (Gao et al., 2004; Tolocka et al., 2004; Surratt et al., 2006a; Surratt et al., 2007a). Since then, ester formation by oxidation of organic matter in the troposphere has been the subject of many laboratory investigations (Hamilton et al., 2006; Surratt et al., 2006b; Szmigielski et al., 2007; Altieri et al., 2008; Galloway et al., 2009; Zhang et al., 2011; Birdsall et al., 2013; Kristensen et al., 2013; Strollo and Ziemann, 2013; Claffin and Ziemann, 2019; Mekic et al., 2019) and field studies (Raja et al., 2008; Raja et al., 2009; Kristensen et al., 2013). The work from (Birdsall et al., 2013) suggested that esterification by condensation of carboxylic acids with hydroxyl group containing molecules is not efficient enough to explain the oligoesters under realistic aerosol acidities. In a recent study by Zhao et al. (2019), heterogeneous oxidation processes near the gas-particle interface open up a further formation pathway of ester-like structures, namely the dimerization of organic oxygen-containing radicals leads dominantly to ester formation. In summary, it should be noted that this accretion reaction in the atmospheric aerosol-phase depends more on the hygroscopicity, than the acidity of the aerosols (Zhao et al., 2006; De Haan et al., 2009a), since the hydrolysis competes with the ester formation.


5.3 Epoxide reactions

Isoprene -derived epoxides

In the last decade, acid-catalyzed ring-opening reactions of epoxides (see Fig. 13) in aqueous aerosols have emerged as a significant source of secondary organic aerosol material. In the aqueous phase, protonation of the epoxide by an acid (H_3O^+ or



NH_4^+ (Minerath et al., 2008; Minerath and Elrod, 2009; Eddingsaas et al., 2010; Nguyen et al., 2014; Noziere et al., 2018))
990 occurs in concert with nucleophilic addition. Typically, the participating nucleophiles are H_2O , HSO_4^- , and SO_4^{2-} , although
amines (Stropoli and Elrod, 2015) and alcohols (Surratt et al., 2010; Piletic et al., 2013) can also add. Nucleophilic attack by
 H_2O results in hydrolysis and polyol formation, thus explaining the presence of isoprene-derived tetrols in particles (Kourtchev
et al., 2005; Xia and Hopke, 2006; Liang et al., 2012; Zhang et al., 2013). The hydrolysis of epoxides catalyzed by NH_4^+ can
only be important in less acidic aerosol solutions due to the orders of magnitude lower rate coefficients (Noziere et al., 2018).
995 Addition of HSO_4^- or SO_4^{2-} to the protonated epoxide in the aerosol phase is a more efficient pathway for organosulfate (OS)
formation than radical mechanisms (McNeill et al., 2012; Schindelka et al., 2013). While the formation of polyols via
hydrolysis of epoxides may be acid catalyzed (Eddingsaas et al., 2010), OS formation can consume H^+ (Riva et al., 2019;
Brüggemann et al., 2020) (e.g. see Fig. 13).



1000

Figure 13. Schematic of the OS and polyol formation via acid-catalyzed ring-opening reactions of epoxides.

Isoprene epoxydiol (IEPOX), a photooxidation product of isoprene (Paulot et al., 2009; Surratt et al., 2010), is calculated to
contribute 34% of global SOA mass (Lin et al., 2012), and 28% of organic aerosol mass in the SE USA (Marais et al., 2016).
1005 The reactive uptake of IEPOX to aqueous media is strongly pH-dependent, with the reactive uptake coefficient decreasing
rapidly with increasing pH for $\text{pH} > 1$ (Gaston et al., 2014). For this reason, the rate of IEPOX SOA formation is slow in cloud
water (McNeill, 2015), but given the relatively large liquid water content of clouds and the relatively large water solubility of
IEPOX, it could be significant in more acidic cloud droplets ($\text{pH} 3\text{-}4$) (Tsui et al., 2019).

1010 *Terpene-derived epoxides*

In regions with lower isoprene but higher monoterpene emissions, e.g., in the boreal forests, monoterpene-derived OSs formed
via different proposed pathways can also contribute to SOA mass in atmospheric aerosols (see Brüggemann et al. (2020) for
details). Their importance for SOA is still not well characterized. Formation of monoterpene-derived OS has been observed in



1015 chamber experiments and measured in field samples (Iinuma et al., 2004; Iinuma et al., 2007; Ye et al., 2018b; Brüggemann et al., 2019; Cui et al., 2019). However, there are only a few measurements of monoterpene-derived OSs in boreal forests areas.

OS formation via acid-catalyzed ring-opening reactions of several monoterpene epoxides (β -pinene oxide, limonene oxide, and limonene dioxide) has been kinetically and mechanistically investigated (Cortes and Elrod, 2017). Investigations demonstrated that monoterpene epoxides react faster than IEPOX in aqueous solution and might even react in less acidic
1020 solutions. However, this study also revealed that the formed OS compounds are not long-lived compounds under aqueous aerosol conditions and may quickly react further mainly through hydrolysis. Therefore, Cortes and Elrod (2017) concluded that other OS formation mechanisms, than the acid-catalyzed ring-opening mechanism of monoterpene epoxides, are needed to explain the formation of more long-lived OS from monoterpenes. In agreement with these findings, recent chamber studies on the OS formation from α -pinene oxidation (Duporte et al., 2016; Duporte et al., 2020) showed that the OS yield, including
1025 the subsequent formation of OS dimers and trimers, decreases with increasing relative humidity. Furthermore, these studies revealed that effective formation rates of OS from α -pinene are 2 orders of magnitude higher under very acidic aerosol conditions and that the OS formation under slightly acidic aerosols conditions is limited. Further sensitivity studies showed a strong dependency of the OS formation on the available sulfate supporting an acid-catalyzed processing of monoterpene epoxides yielding OS. However, it should be noted that regions with high monoterpene emissions are usually not associated
1030 with high sulfate aerosol loadings and quite acidic aerosols, hence, their contribution to SOA might be limited and important only in mixed environments.

Other epoxides

Other atmospheric epoxides have been proposed to contribute to SOA formation, including 2-methyl-3-buten-2-ol (MBO)
1035 (Mael et al., 2015), methacrylic acid epoxide (MAE) (Lin et al., 2013; Birdsall et al., 2014), and epoxides from toluene oxidation (Baltaretu et al., 2009; McNeill et al., 2012). However, none of these species have the relatively large gas-phase production rate and water solubility of IEPOX, so they probably lead to small SOA mass contributions.

5.4 Oxidation reactions of acids and diacids

Acidity changes the speciation of dissociating organic compounds in the atmospheric aqueous phase. More specifically, acidity
1040 decreases the degree of dissociation for organic acids, i.e. lowers the fraction of a compound in its deprotonated form. The protonated and deprotonated forms of a dissociating compound are characterized by different molecular properties (e.g., different bond-dissociation energies (BDEs)). Therefore, key aqueous-phase oxidants, such as the radicals OH, NO₃ or the non-radical oxidant O₃, may react via different possible reaction pathways and kinetics with the protonated and deprotonated forms. Accordingly, acidity can strongly affect the chemical processing of dissociating organic compounds.



1045 Within this subsection, the potential effect of acidity on the chemical processing of dissociating organic compounds in atmospheric aqueous solutions is summarized. The discussion will focus primarily on acids and their respective anions, however, acidity may influence reactivity and partitioning for any dissociating species (including, e.g., imidazoles or phenols).

5.4.1 Reaction pathways of dissociating organic compounds with different oxidants

Similar to the gas phase radical oxidants such as OH and NO₃ can react with dissociating organic compounds via H-abstraction. 1050 Oxidation of dissociated organic compounds may also proceed through an electron transfer reaction (ETR). For unsaturated organic compounds, radical addition to the C=C double bond represents a third possible reaction pathway. Overviews on atmospheric aqueous-phase radical oxidants are available by Buxton et al. (1988); Herrmann (2003); Herrmann et al. (2010); Herrmann et al. (2015).

In Fig. 14, the three types of radical-initiated reactions are schematically displayed for carboxylic acids, the most prominent 1055 dissociating organic compound class in tropospheric aerosols and clouds. The H-abstraction related reactivity of an organic molecule strongly depends on the BDE of the abstractable hydrogen atoms. Carbon-hydrogen bonds (C-H) are typically characterized by lower BDEs (e.g., BDE = 410 ± 5 kJ mol⁻¹ for acetone (Herrmann, 2003)) than other bonds such as oxygen-hydrogen bonds (e.g., O-H in acids, BDE = 445 kJ mol⁻¹ (Luo, 2002)). However, please note that the given BDEs are gas-phase BDEs and that BDEs can be slightly altered by an aqueous solvent. As a consequence of the weaker carbon-hydrogen bonds, 1060 the H-abstraction reaction is currently expected to predominantly proceed at the carbon chain of dissociating organic compounds and not on the hydroxyl group, e.g. of the acid group. The H-abstraction leads to carbon-centered radicals that further react with oxygen leading to the formation of peroxy-radicals and subsequently to functionalized organic compounds with possibly changed dissociation properties. Dissociated organic compounds can also react with radical oxidants via ETR in the aqueous phase, e.g. by removing an electron from a deprotonated and ionized acid group. Particularly for more selective 1065 radical oxidants such as NO₃ (or others such as Cl₂⁻, Br₂⁻), ETR is often preferred over H-abstraction. The reaction rate constants of NO₃ for ETRs are generally larger than those for H-abstraction. Overall, the different contributions of ETR and H-abstraction pathways modify also the product distributions as a function of pH.


The third mentioned pathway of radical oxidants, the addition reaction, occurs for unsaturated aliphatic and aromatic 1070 compounds. This reaction type is typically the fastest radical reaction pathway and proceeds almost at the aqueous-phase diffusion limit (see Sect. 5.4.2), except for double bonds where the electron density is strongly reduced by electron-drawing substituents, such as halogen atoms.




H-Abstraction



Electron-Transfer-Reaction (ETR)



Addition



1075 **Figure 14.** Schematic of the initial reaction steps for the most important radical oxidation pathways of dissociating organic compounds, exemplified for carboxylic acids.

Besides the radical oxidation reactions, dissociating organic compounds can be also oxidized by ozone. In aqueous solutions, the decomposition of ozone is strongly affected by the acidity due to its strong chemical interaction with the water matrix (see Herrmann et al. (2015) and references therein). Ozone is known to be an electrophilic and selective oxidant for organic
1080 compounds, with particular selectivity for C=C double bonds. Therefore, ozone reacts primarily with both unsaturated aliphatic compounds and aromatic compounds. O₃ is also known to react slowly with saturated aliphatic compounds such as hydrated organic acids and carbonyl compounds. Rate constants for these reactions have recently been compiled in Herrmann et al. (2015).

Similar to radical oxidants, ozone reactions are expected to proceed via (i) H-abstraction (e.g. from the hydrated carbonyl
1085 groups of carboxylic acids, see Schöne and Herrmann (2014)), (ii) addition onto C=C double bonds (e.g., in case of unsaturated aliphatic compounds and aromatic compounds, see Mvula and von Sonntag (2003) and Leitzke and von Sonntag (2009)) and, finally, (iii) ETR (see Mvula and von Sonntag (2003)). However, current knowledge of the above-mentioned ozone oxidation mechanisms remains quite limited. An overview of proposed oxidation mechanisms of aqueous-phase ozone can be found, e.g. in (Hoigne and Bader, 1983a, b; Mvula and von Sonntag, 2003; Leitzke and von Sonntag, 2009; von Sonntag and von
1090 Gunten, 2012; Schöne and Herrmann, 2014) and references therein.



5.4.2 Comparison of kinetic data of dissociated and undissociated organic compounds

To examine the effect of acidity on the chemical processing of dissociating organic compounds, kinetic data for their oxidation by OH, NO₃ and O₃ have been newly compiled for the present review following several published review articles and data compilations (Buxton et al., 1988; Neta et al., 1988; Ross et al., 1998; Herrmann, 2003; Herrmann et al., 2010; Herrmann et al., 2015; Bräuer et al., 2019). These data are presented in Tables S5-S7 and Fig. S2-S5 in the Supporting Information. It should be noted that the Tables and Figures in the Supporting Information only show kinetic data for dissociating organic compounds where data for both their protonated and deprotonated form are available. For comparing differences in the kinetic reactivity data of protonated and deprotonated organic compounds, a reactivity ratio κ_R has been calculated. The calculated ratios are defined as the quotient of the kinetic reaction rate constants of deprotonated and protonated form (see below) with the respective oxidant (OH, NO₃, O₃):

$$\kappa_R(i) = \frac{k_{deprotonated}^{298K}}{k_{protonated}^{298K}} \quad (i = \text{OH, NO}_3, \text{O}_3) \quad (22)$$

In brief, values of $\kappa_R(i)$ above 1 imply that reaction of the deprotonated form will proceed more readily than the reaction of the protonated acid. Furthermore, in case of an acid, a $\kappa_R(i)$ above 1 implies that at higher pH, with an increased abundance of the deprotonated form, the overall reaction rate of a compound will be increased, i.e. oxidations are favored under decreasing acidic conditions (cf. Sect. 5.4.3).

5.4.2.1 OH radical oxidations


Overall, for OH reactions, the impact of acidity on the chemical kinetics is often quite small and crucial only for some specific compounds. Thus, with respect to OH reactions, acidity will mostly alter the lifetime of dissociating compounds mainly because of its impact on the partitioning and the consequently affected aqueous-phase concentrations and not so much because of changes in the OH kinetics (see also Sect. 5.4.3). Figure 15 shows the calculated ratios ($\kappa_R(\text{OH})$) of OH reactions with several dissociating organic compounds. The calculated reactivity ratios are typically close to unity, i.e. a similar reactivity exists for the undissociated molecule and its corresponding anion. Larger ratios are calculated for a few small carboxylic and dicarboxylic acids. For formic acid and malonic acid, the $\kappa_R(\text{OH})$ ratios are larger than 10, and for oxalic acid, they are larger than 100. For mono- and dicarboxylic acids, Fig. 15 shows decreasing ratios with increasing carbon chain. For larger carboxylic acids, calculated ratios are scattered around unity. This result indicates that the reaction mechanism of the carboxylic acid and the corresponding carboxylate is similar for the larger acids and H-abstraction is the dominant reaction pathway. With longer carbon chain, the impact of the acid functionality decreases and, thus, almost no acidity dependence exists. This also seems to be true for functionalized carboxylic acids. On the other hand, the substantially larger ratios of the smaller carboxylic acids demonstrate a much faster OH degradation of the smaller carboxylates compared to their protonated acids. This implies that the carboxylate group and their steric effects on the surrounding C-H bonds can facilitate an easier H-abstraction causing higher H-abstraction rate constants for smaller carboxylates compared to corresponding protonated acids. Besides, former



studies have shown that the ETR pathway contributes minor. Schuchmann et al. (1995) reported a contribution of less than 5%. Thus, the ETR pathway should be not responsible for the reactivity difference.

Acetic acid and acetate show the lowest OH reactivity among the considered unsubstituted monocarboxylic acids, with
1125 $1.70 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$ and $7.30 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$ (Chin and Wine, 1994), respectively. The weakest bond H-atoms in these molecules are part of the methyl-group. Those primary C-H bonds are much stronger than secondary or ternary C-H bonds. This explains why acetic acid is less reactive towards OH compared to higher non-substituted primary C-H bonds than monocarboxylic acids containing more CH₂ groups, or secondary C-H bonds. This also explains why the reactivity difference of acetic acid/ acetate is more distinct and the impact of the carboxylate group is higher compared to longer chain acids.

1130 Moreover, its worthy to mention that the comparison of kinetic differences of protonated and deprotonated carboxylic acids given in Fig. 15 is more comprehensive compared to the work of Zhao et al. (2016). In Zhao et al. (2016), the OH oxidation kinetics of formic, glyoxylic, pyruvic acid, lactic acid, malic acid, oxalic acid has been compared. Based limited number of kinetic data and, particularly, because of the selected compounds, this study concluded that the oxidation of carboxylate forms is much more rapid compared to that of free carboxylic acid, indicating an acidity dependence in the reactivity of carboxylic
1135 acids. However, from the present study, it can be concluded that this not true, except for smaller organic acids that are characterized by higher $\kappa_R(OH)$ ratios, due to their special structure properties, such as less or even no abstractable carbon-bonded H-atoms as in case of oxalic acid. Hence, an acidity dependence in the reactivity exists only for these smaller carboxylic acids and almost no acidity dependence exists for other carboxylic acids with a longer carbon chain. Therefore, the statements of Zhao et al. (2016) regarding the pH dependence in the reactivity of saturated carboxylic acids are by far too overgeneralized.



1140

Figure 15. Calculated reactivity ratios $\kappa_R(OH)$ of different dissociating organic compounds. The $\kappa_R(OH)$ ratio of the dianion and protonated diacid is indicated by (di). The applied aqueous-phase OH reaction rate constants are provided in Table S5 in the SI. Different colors indicate different compounds classes (such as unsubstituted saturated monoacids (red), unsubstituted unsaturated monoacids (purple), unsubstituted saturated diacids (blue), substituted saturated monoacids (green), substituted saturated diacids (brown), phenols (light green), aromatic acids (pink), imidazoles (orange)). The dotted bars mark the ratio of the dianions.

1145



The rate constants for unsaturated aliphatic organic acids (protonated and deprotonated) reacting with OH are quite high, in the range of about 10^9 - 10^{10} L mol⁻¹ s⁻¹. The available kinetic data indicate that the considered protonated C4 unsaturated acids (methyl crotonic acid, methacrylic acid) react almost exclusively via OH-addition because of the high bond strengths for the C-H bond in the methyl and the O-H bond in the carboxyl groups. For the smaller protonated unsaturated acids, the protonated acid group likely inhibits the OH-addition because the -COOH group lowers the electron density of the double bond and, therefore, leads to lower reaction constants. However, it can be concluded from the available kinetic dataset that the kinetic acidity effect on unsaturated aliphatic organic acids reacting mostly via OH addition generally should be small.

Aromatic compounds may also dissociate in case of side chains. Both protonated and deprotonated aromatic compounds are characterized mostly by very high OH reaction rate constants up to about 10^{10} L mol⁻¹ s⁻¹ due to the preference of OH radicals to add onto the aromatic ring (Adams et al., 1965). The probability of an H-abstraction from the OH group or an ETR with deprotonated acid groups is minor. Therefore, the calculated reactivity ratios of OH reactions with aromatic compounds are around unity (0.7-2.7). This implies there is only a small kinetic acidity effect for OH reactions with aromatic compounds. A special class of aromatic compounds considered in Fig. 15 are imidazoles. In contrast to the acids that have been the primary focus of discussion, the partitioning of imidazoles, as a base, increases with higher acidity and, therefore, greater aqueous-phase chemical processing is feasible under more acidic particle conditions. Reaction rate constants of protonated imidazoles (Rao et al., 1975; Felber et al., 2019) are approximately a factor of 2 higher than their deprotonated forms (see Fig. 15). This indicates a minor kinetic effect of acidity.


5.4.2.2 NO₃ radical oxidations

In comparison to the OH radical, the NO₃ radical is commonly characterized by a lower reactivity, especially for plain H-abstraction reactions (Herrmann and Zellner, 1998; Herrmann et al., 2015). However, this could be compensated through the high reactivity of NO₃ in single electron transfer reactions. As specifically for the acids, the compiled kinetic data demonstrate that NO₃ radical reaction rate constants for the reactions with undissociated and dissociated acids are often quite different (see Fig. 16). Reaction rate constants of NO₃ radical with saturated protonated aliphatic mono- and di-carboxylic acids are typically in the range of $k = 10^4$ - 10^6 L mol⁻¹ s⁻¹, where the higher values correspond to rate constants for reactions of functionalized acids. In contrast, saturated deprotonated aliphatic mono- and di-carboxylic acids are oxidized by NO₃ with rate constants typically in the range of $k = 10^6$ - 10^8 L mol⁻¹ s⁻¹. Accordingly, the calculated reactivity ratios $\kappa_R(\text{NO}_3)$ (see Fig. 16) are often above 10 and in some cases up to 10^4 . As a consequence, acidity is a very important parameter when the reactivity of NO₃ in the atmospheric aqueous phase is to be described.

Compared to saturated aliphatic acids, unsaturated aliphatic acids show higher NO₃ reactions rate constants ($k = 10^7$ - 10^8 L mol⁻¹ s⁻¹). This is due to a possible addition of the NO₃ radical to the carbon double bond, which proceeds faster than the H-abstraction. $\kappa_R(\text{NO}_3)$ of unsaturated acids, such as acrylic and methacrylic acid, is about 6.4 and 1.8, respectively



(see Fig 16). The NO_3 addition reaction on the C=C double bond is more important than the ETR for both the undissociated and dissociated acid. Hence, the calculated reactivity ratios are smaller compared to NO_3 reactions of saturated acids.



1180

Figure 16: Calculated reactivity ratios $\kappa_R(\text{NO}_3)$ of different carboxylic acids. The $\kappa_R(\text{NO}_3)$ ratio of the dianion and protonated diacid is indicated by the add-on (di) behind the acid name. The applied aqueous-phase NO_3 reaction rate constants are provided in Table S6 in the SI. Different colors indicate different compounds classes (such as unsubstituted saturated monoacids (red), unsubstituted unsaturated monoacids (purple), unsubstituted saturated diacids (blue), substituted saturated monoacids (green), substituted saturated diacids (brown)). The dotted bars mark the ratio of the dianions.

1185

Overall, it can be concluded for NO_3 in aqueous atmospheric systems, particularly clouds conditions, that acidity can substantially affect the chemical NO_3 -initiated processing of organic compounds. Less acidic conditions will enhance the degradation of dissociating compounds via NO_3 because of more rapid oxidation and increased partitioning (see Sect. 5.4.3 for further details). This acidity effect could be important in urban mixed regimes, where higher NO_x regimes mix with marine, continental dust or soil aerosols which are typically less acidic. However, due to of the sparse kinetic database for NO_3 radical reactions in the aqueous phase, more kinetic and mechanistic laboratory investigations are needed with special emphasis on acidity effects.

1190




5.4.2.3 O₃ oxidations

In comparison to OH, ozone (O₃) is commonly known to be an electrophilic and very selective oxidant for organic compounds, covering a very wide range of reactivities. It should be noted here that lower ozone rate constants might be compensated by much higher concentrations of ozone compared to OH (Tilgner and Herrmann, 2010; Schöne and Herrmann, 2014). Comparison of the O₃ oxidation rates for protonated and deprotonated forms of dissociating compounds (Fig. 17) shows that O₃ oxidation kinetics depend significantly on acidity, especially for phenolic compounds. For saturated carboxylic acids, carboxylates demonstrate roughly a factor of 10 higher reactivity towards O₃ as compared to the protonated acids. This higher reactivity can be explained by the higher electron-withdrawing properties of the carboxylate. Therefore, BDEs of the carbon-bonded H-atoms are smaller and more easily abstractable. Furthermore, ETR can also occur.

As expected, compared to saturated carboxylic acids, unsaturated carboxylic acids have significantly higher reactivities with O₃, i.e. more than 4 orders of magnitude higher reaction rate constants. In case of unsaturated carboxylic acids, addition to the C=C double bond will establish an important reaction pathway for both the protonated and the deprotonated unsaturated acid. Nevertheless, the calculated reactivity ratios $\kappa_R(O_3)$ (see Fig. 17) show that the deprotonated unsaturated acid reacts more rapidly (1.3-25 times faster) with ozone than its protonated form. Possible reasons could be the same as for saturated organic acids. The deprotonation likely leads to a reduction in the electron density at the carbon-carbon double bond enabling an easier O₃ addition, i.e. a more rapid oxidation. From inductive effect theory, it is known that the COOH group is electron-withdrawing and COO⁻ is electron-donating. Thus, the obtained behavior is feasible.

A comparison of the kinetic data of maleic and fumaric acids demonstrates that the molecular structure (symmetry/bonds) strongly affects ozone reactivity. These isomers are characterized by different physical and chemical properties (dipole moments, pK_a, reactivity). The differences in the molecular structure lead to a higher O₃ reactivity of fumaric acid. The C=C double bond in a fumaric acid molecule is less shielded from the two acids groups which simplifies O₃ addition onto the double bond. Thus, a 6 times higher reactivity of the protonated fumaric acid results compared to maleic acid.

Similar $\kappa_R(O_3)$ values are also found for aromatic acids containing unsaturated carbon side chains, e.g. cinnamic acid. On the other hand, for hydroxylated acids such as p-hydroxy benzoic acid, significantly higher reactivity differences are found with decreasing acidity. Under highly acidic conditions (pH = 2), p-hydroxy benzoic acid shows a reaction rate constant with O₃ of $2.0 \cdot 10^2 \text{ L mol}^{-1} \text{ s}^{-1}$ (Beltran et al., 2000) whereas it reacts much more rapidly ($6.4 \cdot 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$, (Beltran et al., 2000)) under alkaline conditions (pH = 9). This increase can be explained by increasing deprotonation of the hydroxy group leading to formation of the phenolate form and a higher contribution of the ETR with decreasing acidity. The significantly higher reactivity of the fully deprotonated form implies that the oxidation rate at a pH of 6 is still dominated by the reaction of O₃ with the fully deprotonated form.



1225 **Figure 17: Calculated $\kappa_R(O_3)$ of different dissociating organic compounds. The $\kappa_R(O_3)$ ratio of the dianion and protonated diacid is indicated by the add-on (di) behind the acid name. The applied aqueous-phase O_3 reaction rate constants are provided in Table S7 in the SI. Different colors indicate different compounds classes (substituted saturated monoacids (green), unsaturated monoacids (purple), unsaturated diacids (brown), phenols (light green), aromatic acids (blue)).**

1230 Rather high acidity dependencies of the reactivity data exist for phenolic compounds, too (see Fig. 17), with a large increase of the ozone reaction rate of up to six orders of magnitude. However, it should be mentioned that the present kinetic literature data are based on extrapolations (see Hoigne and Bader (1983b) for details). Due to the huge reactivity difference, the ozone oxidation of phenol can be dominated by the reaction with phenolate even at neutral or slightly acidic conditions. Moreover, charge transfer to ozone can lead to the formation of OH radical (Mvula and von Sonntag, 2003) which can initiate further oxidation reactions. Thus, less acidic conditions can enhance the aqueous oxidation of phenolic compounds by dissolved ozone and additionally promote further oxidation due to the initiated OH chemistry.

1235 Overall, the present O_3 kinetic data analyses have demonstrated the crucial role of acidity for ozonolysis processes and, hence, the chemical processing of dissociating compounds in tropospheric aqueous solutions. The possible formation of OH radicals following initial ozone reactions can further enhance the oxidation capacity of the atmospheric aqueous phase. Further laboratory measurements and modelling studies are urgently needed to improve current knowledge.




5.4.3 Overall considerations for the oxidation of dissociating species and the role of acidity

1240 The overall reaction rate of a compound at a given pH depends on both the individual reaction rate constants (see above) and
the degree of dissociation of the compound (which, in turn, is determined by its pK_a value(s)). The rate constants of the
individual free acids and their dissociated anions represent only extreme values, and the overall processing rate for a weak acid
or base will often fall between these two values, depending on the pH and the pK_a . In order to illustrate this point, Fig. 18
shows the dependence of the overall reaction rate constant through a typical tropospheric pH range of 0 to 9 for a few selected
1245 mono- and diacids. The overall reaction rate constant is calculated by means of the individual reaction rate constants of the
protonated and deprotonated forms and their respective speciation fraction. Please note that the overall second order reaction
rate constants consider the dissociation speciation of the carboxylic acids but not their effective solubility. Thus, the overall
chemical reaction rate will depend on both the aqueous oxidant concentration and on the total aqueous compounds
concentration. The latter largely depend on the microphysical conditions present.

1250 Briefly, Fig. 18 demonstrates that the overall reaction rate constant can be largely pH dependent. This is particularly true for
compounds with large reactivity ratios, i.e. those for which the anion is more reactive than the unionized form. For such
compounds, the overall rate constant typically increases with increasing pH and more efficient oxidation can be expected under
less acidic conditions. In view of decreasing inorganic acid aerosol content, together with decreasing acidity in clouds in some
parts of the world (Pye et al., 2020), this might imply both stronger partitioning and more efficient oxidation of organic acids
1255 (lower chemical aqueous-phase lifetime) in the troposphere under future conditions.

Overall, based on a compiled kinetic dataset for oxidation by OH, NO_3 and O_3 of both protonated and deprotonated organic
compounds, investigations of the reactivity ratio κ_R showed that, for OH reactions, the impact of acidity on the chemical
kinetics is often quite small and only important for some specific compounds. For NO_3 reactions, particularly in cloud droplets,
acidity can substantially affect the chemical NO_3 -initiated processing of organic compounds and less acidic conditions will
1260 enhance the degradation of dissociating compounds via NO_3 because of more rapid oxidation from an increased likelihood for
Electron-Transfer-Reactions (ETRs). Furthermore, the present O_3 kinetic data analyses demonstrate the crucial role of acidity
for ozonolysis processes, especially for phenolic compounds.



1265 **Figure 18.** Overall condensed-phase second order rate constant k_{2nd} as a function of pH of selected mono- (top) and dicarboxylic (bottom) acids for different radical (OH/NO₃) and non-radical (O₃) oxidants.



6 Atmospheric implications

In the review of Pye et al. (2020), a detailed compilation of acidity data measured in cloud and fog water around the globe showed decreasing trends across North America and Europe, mainly driven by a decreased sulfate and nitrate aerosol content due to reduced anthropogenic emissions of SO₂ and NO_{x,y}. The reduction of fossil fuel combustion emissions in a changing world and its related feedback on acidity will have several implications on chemistry-related topics discussed in the present review. As a similar trend in acidity of aqueous aerosols particles has not yet been widely predicted by thermodynamic models and as observations of such a trend for aerosol particles are scarce (see Pye et al. (2020)), this section will mainly focus on implications of changes in the acidity of cloud and fog on multiphase chemistry.

As a result of reductions in anthropogenic emissions of acid precursors in many western industrialized countries, the relative contributions of other sources to the acidification in fog and cloud droplets will continue to grow in importance over the next few decades unless emissions of ammonia from agricultural fertilization are simultaneously reduced. Other direct and indirect acid sources are, for example, (1) the emission of dimethyl sulfide (DMS), (2) emission of SO₂ from volcanic activity, (3) sea spray aerosol related emission of HCl/Cl⁻ and (4) the emission and secondary multiphase formation of organic acids such as formic and oxalic acid.

On the one hand, at pH ranges between 4 and 6, weaker acids tend to partition into less acidic cloud/fog waters more effectively and, thus, contribute more substantially to acidity in less acidic droplet waters. On the other hand, less acidic cloud and fog water pH values are in the typical range of the pK_a values of weak acids such as acetic acid (pK_a = 4.75), so that they can efficaciously buffer acidification by stronger acids in this acidity range (see Sect. 2.2). As a consequence, the increased aqueous-phase partitioning enables higher chemical processing rates of weak acids such as SO₂, HONO and organic carboxylic acids. Both lower acidity and stronger buffering can support faster S(IV) to S(VI) conversions due to the higher efficiency of other chemical pathways such as ozone oxidation (Li et al., 2020b) and therefore reduce the tropospheric lifetime (and in-cloud lifetime) of SO₂. Therefore, under future conditions with a lower overall SO₂ budget, the increased secondary sulfate mass formation probabilities may compensate at least partly the reduced sulfate formation potential. In the case of organic acids, the increased in-cloud partitioning allows them greater opportunities for chemical processing, leading to higher formation yields of functionalized organic acids which tend to partition even stronger towards the aqueous phase of particles and droplets. Hence, higher in-cloud SOA formation yields can be expected as a consequence of the lower acidification of cloud and fog waters by anthropogenic sulfate.

Having affected in-cloud chemistry processes, the decreasing SO₂ budget will also presumably influence the isoprene-related SOA formation, particularly the OS formation. Here, several projection studies (Pye et al., 2013; Marais et al., 2016; Budisulistiorini et al., 2017; He et al., 2018b; Zhang et al., 2018a) have proposed a reduced IEPOX-derived SOA formation under reduced SO₂ emissions. Also, studies for the SE-US (Pye et al., 2013; Marais et al., 2016; Budisulistiorini et al., 2017) implied that a reduction of SO₂ by 25-48% led to reduction of the IEPOX-derived SOA formation of about 35-70%. This effect is mainly related to the changes in aerosol acidity but could be further modulated by the resulting changes in particle viscosity



and phase separation that result from extensive conversion of inorganic to organic sulfur expected with declines in SO₂ (Riva et al., 2019). For the Pearl River Delta region, a reduction of ~ 45% of the IEPOX-derived SOA was reported by He et al. (2018b) due to an aerosol sulfate reduction by 25%. Finally, all studies clearly demonstrated that a SO₂ emission decline in polluted regions could significantly lower the isoprene-related SOA. Similar effects can also be assumed for other acid-catalyzed or acidity-dependent processes.

Another chemical subsystem that will be likely affected by reduced anthropogenic acid precursor emissions in the future is TMI solubilization (see Pye et al. (2020) and references therein). The smaller possible acidification of aqueous interfacial layers on crustal aerosols can lower the acid-driven solubilization of TMI, particularly in regions where dust particles are mixed with urban pollutants. The decreased formation of soluble and, hence, bioavailable TMIs can (1) cause lower nutrient inputs into oceans impacting the ocean biological activity there, (2) decrease the chemical HO_x radical cycling in both aqueous particles and droplets and (3) might also affect the TMI-related S(IV) oxidation.

Decreasing atmospheric acidity may also impact the acidity-driven production of reactive halogens, with potential implications for ozone and OH. Observations of sea-salt aerosol bromide and chloride deficits over the northeastern Pacific Ocean revealed that depletions in bromide and chloride relative to seawater were correlated with particle acidity (see e.g., Newberg et al. (2005)).

In order to explore the expected changes in the tropospheric multiphase chemistry and their overarching impacts in a changing environment, further field measurements, laboratory studies and accompanied modeling are needed to both monitor occurring changes and improve air quality and climate model predictions.



1320 7 Conclusions and outlook

In the present review, we have outlined different aspects and chemical subsystems where acidity affects multiphase chemistry and, in turn, acidity is affected by tropospheric multiphase chemistry. Although many advances have been made in our understanding of acidity-driven and acid-catalyzed chemical processes, there are still many open issues which need to be addressed in order to further advance our understanding of the complex role acidity in the atmosphere. Besides the implications
1325 caused by changing acidity conditions in the atmosphere (cf. Sect. 6), the present review has also identified chemistry-related research targets and needs for further investigation that are outlined below. Specifically, these chemistry-related future research needs and objectives are:

- (1) Advance our understanding of the activation mechanisms and multiphase chemical processing of reactive halogen species in different acidity environments and quantify their presence in and above the tropospheric boundary layer
- 1330 (2) Undertake more sophisticated field, laboratory and model investigations on the importance of different S(IV) oxidation pathways under various acidity conditions in different environments, particularly under conditions typical of aerosols (lower pH, high ionic strength)
- (3) Advance our understanding of aqueous-phase organic accretion reactions including their dependence on acidity in order to assess their role for the secondary aerosol formation in clouds and aqueous particles
- 1335 (4) Perform advanced kinetic and mechanistic studies on acidity-dependencies of aqueous-phase organic oxidations
- (5) Quantify the role of organosulfates (OS)/ organonitrates (ON)/ nitrooxy organosulfates (NOS) as potential acidity reservoir or sink and characterize the role of acidity for their formation and fate in aerosols
- (6) Investigate the impact of particle acidity in the formation and early growth of tropospheric nanometer-sized particles from highly-oxidized molecules (HOMs)
- 1340 (7) Improve size/time-resolved cloud/fog measurement techniques and develop in situ measurements techniques for directly determining aerosol acidity, burdens which still limits our knowledge about the impact of acidity on the multiphase chemistry in different droplet and aerosol sizes and their feedbacks on chemical composition

(1) While much effort has been devoted to investigations of reactive halogen chemistry in pristine open ocean regions and
1345 partly coastal areas (see Sect. 4.8 for details), the impact of reactive halogens on atmospheric chemistry in developing countries is less examined. Especially in developing economies such as China and India, where a substantial amount of the air pollution is related to coal combustion and other biomass burning, a significant fraction of the aerosol matter consists of halogens related to such sources (Goetz et al., 2018; Gani et al., 2019). Further studies are needed to investigate the role of halogen chemistry in strongly polluted environments which are characterized by very acidic particles compared to marine environments. Under
1350 very polluted conditions, high acidity linked with high $\text{NO}_{x,y}$ can cause active halogen radical chemistry that might influence the tropospheric cleaning capacity. However, the role of multiphase halogen chemistry in such environments is still not well investigated. For example, recent studies of Wang et al. (2019b) have demonstrated that current models are not able to



reproduce high observed Cl_2 in daytime in continental regimes. Furthermore, recent model studies of Zhu et al. (2019) also reported overestimates of free-tropospheric BrO during the extratropical winter–spring. Moreover, our understanding of halogen activation processes in Arctic regions needs further improvement. Arctic regions are undergoing unprecedented climate changes, most likely with substantial changes in the aerosol composition that can affect the aerosol acidity and consequently halogen activation processes. So, further research is needed to focus on how Arctic climate changes will impact halogen chemistry in this dynamic and sensitive environment.

(2) Comparisons of model findings with field measurements in polluted regions have shown that current models often underestimate the S(VI) formation rates or cannot reproduce the findings of sulfur-isotope measurements regarding the responsible oxidation pathways. Hence, the current chemical kinetic and mechanistic understanding of the S(IV) to S(VI) conversion processes, including their acidity dependency, is still incomplete to adequately predict the budgets of S(VI) in cloud, fog, and especially aqueous aerosol conditions. As multiphase chemistry models rely on detailed acidity-dependent kinetic and mechanistic knowledge, further laboratory studies are indispensable to improve model predictions of S(VI), particularly under conditions of high ionic strength (e.g. aerosol chamber or aerosol flow tube studies).

(3) Non-oxidative aqueous organic chemical process, such as accretion reactions (aldol condensation, hemiacetal as well as acetal formation and the esterification of carboxylic acids), are affected by acidity/basicity of the solution and are expected to be important formation pathways of aqSOA. However, the potential role of such acidity-related processes in tropospheric aqueous solutions is still not fully explored yet since mechanistic and particularly kinetic data on acid-catalyzed accretion reactions in aerosols are still sparse (see Herrmann et al. (2015)). So, these acidity-related processes should be a key objective of future laboratory and chamber studies towards a better representation of such processes in detailed chemistry mechanisms and models.

(4) More advanced kinetic and mechanistic studies on acidity-dependencies of aqueous-phase oxidations of dissociating organic compounds, such as functionalized carboxylic acids, are needed to better describe such processes in future multiphase models and to, finally, elucidate their impacts. Compared to the huge number of functionalized organic dissociating organic compounds that are formed by various multiphase reaction pathways, the body of investigated aqueous-phase oxidations are still rather small. Furthermore, existing estimation methods for aqueous-phase kinetic reaction rate constants are either characterized by large uncertainties particularly for functionalized compounds (Bräuer et al., 2019) or don't exist because of the sparse dataset available. Even though the present review showed that acidity could play an important role for ozonolysis processes of dissociating compounds in tropospheric aqueous solutions, nevertheless, the kinetic and mechanistic knowledge of such oxidation processes including the possible formation of OH radicals is currently rather limited. Further kinetic and mechanistic laboratory investigation are urgently needed to minimize the current enormous knowledge limitations and uncertainties, for example, with regards to the production of organic mono- and dicarboxylic acids as well as their functionalized derivatives.

(5) Organosulfates (OSs) are ubiquitous constituents of atmospheric aerosol particles that not only contribute substantially to OM, but may also bind a considerable portion of the sulfate content of atmospheric particles (Riva et al., 2019; Brüggemann



et al., 2020). For this reason, these compounds have the potential to reduce the free sulfate, and consequently the H^+ formation potential, with major implications for aerosol acidity. Overall, OSs may be a temporary acidity reservoir due to the binding and release of sulfate, or a sink of H^+ if H^+ is incorporated into the OS and unavailable for further processing. While progress has been achieved in the understanding of OS formation pathways in the last decade, the scientific understanding of their chemical processing in aqueous aerosols is still uncertain. In order to elucidate the role of OSs to act as crucial acidity reservoir/buffer in atmospheric particles, more detailed knowledge of their chemical processing is highly desirable. Thus, the chemical transformations of OSs through hydrolysis and oxidations by atmospheric radicals (e.g., OH, NO_3 , Cl etc.) require more kinetic and mechanistic laboratory investigations as reaction data are usually not yet available. Additionally, similar investigations on the formation and transformation must be performed for organonitrates (ONs) and nitrooxy organosulfates (NOSs) which could also potentially act as nitrate/sulfate reservoirs, and thus affect the acidity.

(6) Historically, sulfuric acid has been considered the driver of new particle formation events (e.g. Sipila et al. (2010)). However, several studies (see Lee et al. (2019) and references therein) have shown that the gas-phase oxidation of biogenic and anthropogenic VOCs, via autoxidation reactions, can produce highly-oxidized organic molecules (HOMs). HOMs are characterized by extremely low saturation vapor pressures and can effectively condense on nanometer-sized aerosol, contributing there to the early growth of particles. Due to their high degree of oxygenation, including several chemical functional groups (e.g., peroxide, hydroperoxide, carbonyl, and carboxylic acid) that are sensitive to acidity, they can undergo chemical reactions there which can enhance their uptake and contribution to the particle growth. As a result, one possible chemical pathway would be the formation of HOOS (highly oxidized OSs). Having first been reported by Mutzel et al. (2015), they potentially serve as an example to better explain the early growth of freshly formed particles. Such acidic aerosols might provide a chemical environment where extremely low volatility compounds can condense and react, leading to the formation of HOOSs. Elucidating the role of acidity for the formation of SOA in such small particles and their importance for early nanoparticle growth will be a crucial objective for future field and chamber studies.

(7) Because aerosols are ubiquitous in the atmosphere, and clouds cover 60% of the Earth, understanding their contribution to tropospheric composition and how it is evolving, is crucial. Here, the present review has shown the role of acidity in determining aqueous chemistry and vice versa, yet several issues demand more advanced field, kinetic laboratory, chamber and modeling studies. Armed with this new fundamental knowledge, better predictions can be made of aerosol and cloud/fog chemistry on tropospheric oxidizing capacity, air quality, climate, and human health. For example, to better characterize the effect of varying acidity on chemical processing and the identification of potential changes due to changing anthropogenic emissions, further advances in measuring the chemical composition of cloud/fog in a both size- and time-resolved manner is needed. The lack of measurement and analytical techniques for directly determining aerosol acidity in situ is even more urgent.



Data availability. All data used for the figures are provided in the Tables and supplement.

Supplement. The supplement related to this article is available online at:

1420 The supplement includes additional documentation on available hydration equilibrium constants K_{hyd} and compiled kinetic data of OH, NO₃ and ozone reactions of dissociating organic compounds. In addition, figures further displaying differences in reaction rate constants of protonated and deprotonated organic compounds are shown. Data used as input, to create the partitioning and speciation plots (Fig. 1 and 2) are available in the supplement.

1425 *Author contributions.* VFM and HH coordinated the manuscript preparation and AT was the lead author. HOTP and AN designed the overall scope of this study. AT, TS, BA, MB, JLC, KLF, HH, AN, HOTP, and VFM wrote the manuscript. All authors prepared text, figures, and tables in collaboration.

Competing interests. The authors declare no conflicts of interest.

1430

Disclaimer. The U.S. Environmental Protection Agency through its Office of Research and Development collaborated in the research described here. The research has been subjected to Agency administrative review and approved for publication but may not necessarily reflect official Agency policy. The views expressed in this article are those of the authors and do not necessarily represent the views or policies of the U.S. Environmental Protection Agency.

1435

Acknowledgements. We thank the EPA for funding and hosting a workshop on the State of Acidity in the Atmosphere: Particles and Clouds. We thank Ken Elstein, Brooke Hemming, and Randa Boykin for their assistance during the workshop. This work has received funding from the European Union's Horizon 2020 research and innovation program through the EUROCHAMP-2020 Infrastructure Activity under grant agreement no. 730997. Support for TROPOS-ACD by EFRE
1440 (Contract 3000582010) is acknowledged.

Financial support. We thank the EPA for funding and hosting the workshop *The State of Acidity in the Atmosphere: Particles and Clouds* and Ken Elstein, Brooke Hemming, and Randa Boykin for their assistance during the workshop. AN was supported by the project PyroTRACH (ERC-2016-COG) funded by H2020-EU.1.1. – Excellent Science – European Research Council
1445 (ERC), project ID 726165. The work by JC was supported by grant number NSF-AGS- 1650786.



References

- Adams, G. E., Boag, J. W., Currant, J., and Michael, B. D.: Absolute rate constants for the reaction of the hydroxyl radical with organic compounds, in: Pulse Radiolysis, edited by: Ebert, M., Keene, J. P., Swallow, A. J., and Baxendale, J. H., Academic Press, New York, 131-143, 1965.
- Adriano, D. C., and Johnson, A. H.: Acidic precipitation, Vol. 2: Biological and ecological effects, Springer-Verlag New York, Inc., New York, NY, USA, 1989.
- Ahrens, M. L., and Strehlow, H.: Acid Catalyzed Hydration of Acetaldehyde, *Discuss. Faraday Soc.*, 112-120, <https://doi.org/10.1039/DF9653900112>, 1965.
- Aiona, P. K., Lee, H. J., Leslie, R., Lin, P., Laskin, A., Laskin, J., and Nizkorodov, S. A.: Photochemistry of Products of the Aqueous Reaction of Methylglyoxal with Ammonium Sulfate, *ACS Earth Space Chem.*, 1, 522-532, <https://doi.org/10.1021/acsearthspacechem.7b00075>, 2017.
- Alexander, B., Park, R. J., Jacob, D. J., and Gong, S. L.: Transition metal-catalyzed oxidation of atmospheric sulfur: Global implications for the sulfur budget, *J. Geophys. Res.-Atmos.*, 114, D02309, <https://doi.org/10.1029/2008jd010486>, 2009.
- Alexander, B., Allman, D. J., Amos, H. M., Fairlie, T. D., Dachs, J., Hegg, D. A., and Sletten, R. S.: Isotopic constraints on the formation pathways of sulfate aerosol in the marine boundary layer of the subtropical northeast Atlantic Ocean, *J. Geophys. Res.-Atmos.*, 117, D06304, <https://doi.org/10.1029/2011jd016773>, 2012.
- Ali, H. M., Iedema, M., Yu, X. Y., and Cowin, J. P.: Ionic strength dependence of the oxidation of SO₂ by H₂O₂ in sodium chloride particles, *Atmos. Environ.*, 89, 731-738, <https://doi.org/10.1016/j.atmosenv.2014.02.045>, 2014.
- Altieri, K. E., Carlton, A. G., Lim, H. J., Turpin, B. J., and Seitzinger, S. P.: Evidence for oligomer formation in clouds: reactions of isoprene oxidation products, *Environ. Sci. Technol.*, 40, 4956-4960, <https://doi.org/10.1021/es052170n>, 2006.
- Altieri, K. E., Seitzinger, S. P., Carlton, A. G., Turpin, B. J., Klein, G. C., and Marshall, A. G.: Oligomers formed through in-cloud methylglyoxal reactions: Chemical composition, properties, and mechanisms investigated by ultra-high resolution FT-ICR mass spectrometry, *Atmos. Environ.*, 42, 1476-1490, <https://doi.org/10.1016/j.atmosenv.2007.11.015>, 2008.
- Amels, P., Elias, H., Gotz, U., Steingens, U., and Wannowius, K. J.: Sulfur(IV) Oxidation, in: Heterogeneous and Liquid Phase Processes. Transport and Chemical Transformation of Pollutants in the Troposphere, edited by: Warneck, P., Springer, Berlin, Heidelberg, 77-96, <https://doi.org/10.1007/978-3-642-61445-3>, 1996.
- Ariya, P. A., Amyot, M., Dastoor, A., Deeds, D., Feinberg, A., Kos, G., Poulain, A., Ryjkov, A., Semeniuk, K., Subir, M., and Toyota, K.: Mercury physicochemical and biogeochemical transformation in the atmosphere and at atmospheric interfaces: a review and future directions, *Chem. Rev.*, 115, 3760-3802, <https://doi.org/10.1021/cr500667e>, 2015.
- Armstrong, D. A., Huie, R. E., Lymar, S., Koppenol, W. H., Merényi, G., Neta, P., Stanbury, D. M., Steenken, S., and Wardman, P.: Standard electrode potentials involving radicals in aqueous solution: inorganic radicals, *Bioinorg. React. Mech.*, 9, <https://doi.org/10.1515/irm-2013-0005>, 2013.
- Attwood, A. R., Washenfelder, R. A., Brock, C. A., Hu, W., Baumann, K., Campuzano-Jost, P., Day, D. A., Edgerton, E. S., Murphy, D. M., Palm, B. B., McComiskey, A., Wagner, N. L., de Sá, S. S., Ortega, A., Martin, S. T., Jimenez, J. L., and Brown, S. S.: Trends in sulfate and organic aerosol mass in the Southeast U.S.: Impact on aerosol optical depth and radiative forcing, *Geophys. Res. Lett.*, 41, 7701-7709, <https://doi.org/10.1002/2014gl061669>, 2014.
- Au Yang, D., Bardoux, G., Assayag, N., Laskar, C., Widory, D., and Cartigny, P.: Atmospheric SO₂ oxidation by NO₂ plays no role in the mass independent sulfur isotope fractionation of urban aerosols, *Atmos. Environ.*, 193, 109-117, <https://doi.org/10.1016/j.atmosenv.2018.09.007>, 2018.
- Baker, A. R., Kanakidou, M., Nenes, A., Myriokefalitakis, S., Croot, P. L., Duce, R. A., Gao, Y., Guieu, C., Ito, A., Jickells, T. D., Mahowald, N. M., Middag, R., Perron, M. M. G., Sarin, M. M., Shelley, R., and Turner, D. R.: Changing atmospheric acidity as a modulator of nutrient deposition and ocean biogeochemistry, *Sci. Adv.*, in review, 2020.
- Baltaretu, C. O., Lichtman, E. I., Hadler, A. B., and Elrod, M. J.: Primary atmospheric oxidation mechanism for toluene, *J. Phys. Chem. A*, 113, 221-230, <https://doi.org/10.1021/jp806841t>, 2009.
- Bao, L. F., Matsumoto, M., Kubota, T., Sekiguchi, K., Wang, Q. Y., and Sakamoto, K.: Gas/particle partitioning of low-molecular-weight dicarboxylic acids at a suburban site in Saitama, Japan, *Atmos. Environ.*, 47, 546-553, <https://doi.org/10.1016/j.atmosenv.2009.09.014>, 2012.
- Barker, J. R., Steiner, A. L., and Wallington, T. J.: Advances in Atmospheric Chemistry, *Advances in Atmospheric Chemistry, Volume 1*, World Scientific, 608 pp., <https://doi.org/10.1142/10216>, 2016.



- Barlow, S., Buxton, G. V., Murray, S. A., and Salmon, G. A.: Free-radical-induced oxidation of hydroxymethanesulfonate in aqueous solution. 1. A pulse radiolysis study of the reactions of (OH)-O-center dot and SO₄ center dot-, *J. Chem. Soc. Faraday T.*, 93, 3637-3640, <https://doi.org/10.1039/a703267f>, 1997a.
- 1500 Barlow, S., Buxton, G. V., Murray, S. A., and Salmon, G. A.: Free-radical-induced oxidation of hydroxymethanesulfonate in aqueous solution. 2. A pulse and steady-state radiolysis study of oxygenated solutions, *J. Chem. Soc. Faraday T.*, 93, 3641-3645, <https://doi.org/10.1039/a703266h>, 1997b.
- Barnes, I., Hjorth, J., and Mihalopoulos, N.: Dimethyl sulfide and dimethyl sulfoxide and their oxidation in the atmosphere, *Chem. Rev.*, 106, 940-975, <https://doi.org/10.1021/cr020529+>, 2006.
- 1505 Barrie, L. A., Yi, Y., Leaitch, W. R., Lohmann, U., Kasibhatla, P., Roelofs, G. J., Wilson, J., McGovern, F., Benkovitz, C., Melieres, M. A., Law, K., Prospero, J., Kritz, M., Bergmann, D., Bridgeman, C., Chin, M., Christensen, J., Easter, R., Feichter, J., Land, C., Jeuken, A., Kjellstrom, E., Koch, D., and Rasch, P.: A comparison of large-scale atmospheric sulphate aerosol models (COSAM): overview and highlights, *Tellus B*, 53, 615-645, <https://doi.org/10.1034/j.1600-0889.2001.530507.x>, 2001.
- Barsanti, K. C., and Pankow, J. F.: Thermodynamics of the formation of atmospheric organic particulate matter by accretion reactions - Part 1: aldehydes and ketones, *Atmos. Environ.*, 38, 4371-4382, <https://doi.org/10.1016/j.atmosenv.2004.03.035>, 2004.
- Barsanti, K. C., and Pankow, J. F.: Thermodynamics of the formation of atmospheric organic particulate matter by accretion reactions - 2. Dialdehydes, methylglyoxal, and diketones, *Atmos. Environ.*, 39, 6597-6607, <https://doi.org/10.1016/j.atmosenv.2005.07.056>, 2005.
- 1515 Barsanti, K. C., and Pankow, J. F.: Thermodynamics of the formation of atmospheric organic particulate matter by accretion reactions - Part 3: Carboxylic and dicarboxylic acids, *Atmos. Environ.*, 40, 6676-6686, <https://doi.org/10.1016/j.atmosenv.2006.03.013>, 2006.
- Barth, M. C., Rasch, P. J., Kiehl, J. T., Benkovitz, C. M., and Schwartz, S. E.: Sulfur chemistry in the National Center for Atmospheric Research Community Climate Model: Description, evaluation, features, and sensitivity to aqueous chemistry, *J. Geophys. Res.-Atmos.*, 105, 1387-1415, <https://doi.org/10.1029/1999jd900773>, 2000.
- 1520 Behera, S. N., Sharma, M., Aneja, V. P., and Balasubramanian, R.: Ammonia in the atmosphere: a review on emission sources, atmospheric chemistry and deposition on terrestrial bodies, *Environ. Sci. Pollut. Res.*, 20, 8092-8131, <https://doi.org/10.1007/s11356-013-2051-9>, 2013.
- Bell, R. P., and Darwent, B. D.: The Kinetics of the Hydration of Acetaldehyde, *T. Faraday Soc.*, 46, 34-41, <https://doi.org/10.1039/tf9504600034>, 1950.
- 1525 Bell, R. P.: The Reversible Hydration of Carbonyl Compounds, *Adv. Phys. Org. Chem.*, 4, 1-29, [https://doi.org/10.1016/S0065-3160\(08\)60351-2](https://doi.org/10.1016/S0065-3160(08)60351-2), 1966.
- Bell, R. P., and Evans, P. G.: Kinetics of the dehydration of methylene glycol in aqueous solution, *Proc. R. Soc. Lond. A*, 291, 297-323, <https://doi.org/10.1098/rspa.1966.0097>, 1966.
- 1530 Beltran, F. J., Garcia-Araya, J. F., Rivas, F. J., Alvarez, P., and Rodriguez, E.: Kinetics of competitive ozonation of some phenolic compounds present in wastewater from food processing industries, *Ozone-Sci. Eng.*, 22, 167-183, <https://doi.org/10.1080/01919510008547218>, 2000.
- Berglund, J., and Elding, L. I.: Transition Metal Ions in Atmospheric Waters, in: *Heterogeneous and Liquid Phase Processes: Laboratory Studies Related to Aerosols and Clouds*, edited by: Warneck, P., Springer Berlin Heidelberg, Berlin, Heidelberg, 97-131, https://doi.org/10.1007/978-3-642-61445-3_4, 1996.
- 1535 Birdsall, A. W., Zentner, C. A., and Elrod, M. J.: Study of the kinetics and equilibria of the oligomerization reactions of 2-methylglyceric acid, *Atmos. Chem. Phys.*, 13, 3097-3109, <https://doi.org/10.5194/acp-13-3097-2013>, 2013.
- Birdsall, A. W., Miner, C. R., Mael, L. E., and Elrod, M. J.: Mechanistic study of secondary organic aerosol components formed from nucleophilic addition reactions of methacrylic acid epoxide, *Atmos. Chem. Phys.*, 14, 12951-12964, <https://doi.org/10.5194/acp-14-12951-2014>, 2014.
- 1540 Bones, D. L., Henricksen, D. K., Mang, S. A., Gonsior, M., Bateman, A. P., Nguyen, T. B., Cooper, W. J., and Nizkorodov, S. A.: Appearance of strong absorbers and fluorophores in limonene-O₃ secondary organic aerosol due to NH₄⁺-mediated chemical aging over long time scales, *J. Geophys. Res.-Atmos.*, 115, <https://doi.org/10.1029/2009jd012864>, 2010.
- 1545 Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V. M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S. K., Sherwood, S., Stevens, B., and Zhang, X. Y.: Clouds and Aerosols, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the*



- Intergovernmental Panel on Climate Change, edited by: Stocker, T. F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgle, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- 1550 Boyce, S. D., and Hoffmann, M. R.: Kinetics and mechanism of the formation of hydroxymethanesulfonic acid at low pH, *J. Phys. Chem.*, 88, 4740-4746, <https://doi.org/10.1021/j150664a059>, 1984.
- Brandt, C., and van Eldik, R.: Transition-Metal-Catalyzed Oxidation of Sulfur(IV) Oxides - Atmospheric-Relevant Processes and Mechanisms, *Chem. Rev.*, 95, 119-190, <https://doi.org/10.1021/cr00033a006>, 1995.
- 1555 Bräuer, P., Tilgner, A., Wolke, R., and Herrmann, H.: Mechanism development and modelling of tropospheric multiphase halogen chemistry: The CAPRAM Halogen Module 2.0 (HM2), *J. Atmos. Chem.*, 70, 19-52, <https://doi.org/10.1007/s10874-013-9249-6>, 2013.
- Bräuer, P., Mouchel-Vallon, C., Tilgner, A., Mutzel, A., Böge, O., Rodigast, M., Poulain, L., van Pinxteren, D., Wolke, R., Aumont, B., and Herrmann, H.: Development of a protocol for the auto-generation of explicit aqueous-phase oxidation schemes of organic compounds, *Atmos. Chem. Phys.*, 19, 9209-9239, <https://doi.org/10.5194/acp-19-9209-2019>, 2019.
- 1560 Brüggemann, M., van Pinxteren, D., Wang, Y., Yu, J. Z., and Herrmann, H.: Quantification of known and unknown terpenoid organosulfates in PM10 using untargeted LC-HRMS/MS: contrasting summertime rural Germany and the North China Plain, *Environ. Chem.*, 16, 333-346, <https://doi.org/10.1071/EN19089>, 2019.
- Brüggemann, M., Xu, R., Tilgner, A., Kwong, K. C., Mutzel, A., Poon, H. Y., Otto, T., Schaefer, T., Poulain, L., Chan, M. N., and Herrmann, H.: Organosulfates in Ambient Aerosol: State of Knowledge and Future Research Directions on Formation, Abundance, Fate, and Importance, *Environ. Sci. Technol.*, 54, 3767-3782, <https://doi.org/10.1021/acs.est.9b06751>, 2020.
- 1565 Budisulistiorini, S. H., Nenes, A., Carlton, A. G., Surratt, J. D., McNeill, V. F., and Pye, H. O. T.: Simulating Aqueous-Phase Isoprene-Epoxydiol (IEPOX) Secondary Organic Aerosol Production During the 2013 Southern Oxidant and Aerosol Study (SOAS), *Environ. Sci. Technol.*, 51, 5026-5034, <https://doi.org/10.1021/acs.est.6b05750>, 2017.
- Buxton, G. V., Greenstock, C. L., Helman, W. P., and Ross, A. B.: Critical-Review of Rate Constants for Reactions of Hydrated Electrons, Hydrogen-Atoms and Hydroxyl Radicals ($\cdot\text{OH}/\cdot\text{O}$) in Aqueous-Solution, *J. Phys. Chem. Ref. Data*, 17, 513-886, <https://doi.org/10.1063/1.555805>, 1988.
- 1570 Buxton, G. V., McGowan, S., Salmon, G. A., Williams, J. E., and Wood, N. D.: A study of the spectra and reactivity of oxysulphur-radical anions involved in the chain oxidation of S(IV): A pulse and γ -radiolysis study, *Atmos. Environ.*, 30, 2483-2493, [https://doi.org/10.1016/1352-2310\(95\)00473-4](https://doi.org/10.1016/1352-2310(95)00473-4), 1996.
- 1575 Calvert, J. G., Lazrus, A., Kok, G. L., Heikes, B. G., Walega, J. G., Lind, J., and Cantrell, C. A.: Chemical Mechanisms of Acid Generation in the Troposphere, *Nature*, 317, 27-35, <https://doi.org/10.1038/317027a0>, 1985.
- Cappa, C. D., Lovejoy, E. R., and Ravishankara, A. R.: Evidence for liquid-like and nonideal behavior of a mixture of organic aerosol components, *P. Natl. Acad. Sci. USA*, 105, 18687-18691, <https://doi.org/10.1073/pnas.0802144105>, 2008.
- 1580 Carpenter, L. J., MacDonald, S. M., Shaw, M. D., Kumar, R., Saunders, R. W., Parthipan, R., Wilson, J., and Plane, J. M. C.: Atmospheric iodine levels influenced by sea surface emissions of inorganic iodine, *Nat. Geosci.*, 6, 108-111, <https://doi.org/10.1038/Ngeo1687>, 2013.
- Casale, M. T., Richman, A. R., Elrod, M. J., Garland, R. M., Beaver, M. R., and Tolbert, M. A.: Kinetics of acid-catalyzed aldol condensation reactions of aliphatic aldehydes, *Atmos. Environ.*, 41, 6212-6224, <https://doi.org/10.1016/j.atmosenv.2007.04.002>, 2007.
- 1585 Chameides, W. L.: The Photochemistry of a Remote Marine Stratiform Cloud, *J. Geophys. Res.-Atmos.*, 89, 4739-4755, <https://doi.org/10.1029/JD089iD03p04739>, 1984.
- Chameides, W. L., and Stelson, A. W.: Aqueous-Phase Chemical Processes in Deliquescent Sea-Salt Aerosols - a Mechanism That Couples the Atmospheric Cycles of S and Sea-Salt, *J. Geophys. Res.-Atmos.*, 98, 9051-9054, <https://doi.org/10.1029/93jd00310>, 1993.
- 1590 Charlson, R. J., Schwartz, S. E., Hales, J. M., Cess, R. D., Coakley, J. A., Hansen, J. E., and Hofmann, D. J.: Climate Forcing by Anthropogenic Aerosols, *Science*, 255, 423, <https://doi.org/10.1126/science.255.5043.423>, 1992.
- Chebbi, A., and Carlier, P.: Carboxylic acids in the troposphere, occurrence, sources, and sinks: A review, *Atmos. Environ.*, 30, 4233-4249, [https://doi.org/10.1016/1352-2310\(96\)00102-1](https://doi.org/10.1016/1352-2310(96)00102-1), 1996.
- 1595 Chen, Q., Schmidt, J. A., Shah, V., Jaegle, L., Sherwen, T., and Alexander, B.: Sulfate production by reactive bromine: Implications for the global sulfur and reactive bromine budgets, *Geophys. Res. Lett.*, 44, 7069-7078, <https://doi.org/10.1002/2017gl073812>, 2017.



- Chen, Q. J., Geng, L., Schmidt, J. A., Xie, Z. Q., Kang, H., Dachs, J., Cole-Dai, J., Schauer, A. J., Camp, M. G., and Alexander, B.: Isotopic constraints on the role of hypohalous acids in sulfate aerosol formation in the remote marine boundary layer, *Atmos. Chem. Phys.*, 16, 11433-11450, <https://doi.org/10.5194/acp-16-11433-2016>, 2016.
- 1600 Cheng, Y., Zheng, G., Wei, C., Mu, Q., Zheng, B., Wang, Z., Gao, M., Zhang, Q., He, K., Carmichael, G., Poschl, U., and Su, H.: Reactive nitrogen chemistry in aerosol water as a source of sulfate during haze events in China, *Sci. Adv.*, 2, e1601530, <https://doi.org/10.1126/sciadv.1601530>, 2016.
- Chin, M., and Wine, P.: A temperature-dependent competitive kinetics study of the aqueous-phase reactions of OH radicals with formate, formic acid, acetate, acetic acid, and hydrated formaldehyde, in: *Aquatic and Surface Photochemistry*, edited by: Helz, G., Zepp, R., and Crosby, D., CRC Press, Boca Raton, FL, 85-96, 1994.
- 1605 Claffin, M. S., and Ziemann, P. J.: Thermal desorption behavior of hemiacetal, acetal, ether, and ester oligomers, *Aerosol Sci. Technol.*, 53, 473-484, <https://doi.org/10.1080/02786826.2019.1576853>, 2019.
- Clayden, J., Greeves, N., and Warren, S.: *Organic Chemistry*, 2 ed., Oxford University Press, Oxford, 2012.
- Clegg, S. L., and Seinfeld, J. H.: Thermodynamic Models of Aqueous Solutions Containing Inorganic Electrolytes and Dicarboxylic Acids at 298.15 K. 1. The Acids as Nondissociating Components, *J. Phys. Chem. A*, 110, 5692-5717, <https://doi.org/10.1021/jp056149k>, 2006.
- 1610 Clifton, C. L., Altstein, N., and Huie, R. E.: Rate constant for the reaction of nitrogen dioxide with sulfur(IV) over the pH range 5.3-13, *Environ. Sci. Technol.*, 22, 586-589, <https://doi.org/10.1021/es00170a018>, 1988.
- Collett, J. L., Hoag, K. J., and Rao, X.: Internal acid buffering in San Joaquin Valley fog drops and its influence on aerosol processing, *Atmos. Environ.*, 33, 4833-4847, [https://doi.org/10.1016/S1352-2310\(99\)00221-6](https://doi.org/10.1016/S1352-2310(99)00221-6), 1999.
- 1615 Cortes, D. A., and Elrod, M. J.: Kinetics of the Aqueous Phase Reactions of Atmospherically Relevant Monoterpene Epoxides, *J. Phys. Chem. A*, 121, 9297-9305, <https://doi.org/10.1021/acs.jpca.7b09427>, 2017.
- Cui, T., Green, H. S., Selleck, P. W., Zhang, Z., O'Brien, R. E., Gold, A., Keywood, M., Kroll, J. H., and Surratt, J. D.: Chemical Characterization of Isoprene- and Monoterpene-Derived Secondary Organic Aerosol Tracers in Remote Marine Aerosols over a Quarter Century, *ACS Earth Space Chem.*, 3, 935-946, <https://doi.org/10.1021/acsearchspacechem.9b00061>, 2019.
- 1620 Dani, K. G. S., and Loreto, F.: Trade-Off Between Dimethyl Sulfide and Isoprene Emissions from Marine Phytoplankton, *Trends Plant. Sci.*, 22, 361-372, <https://doi.org/10.1016/j.tplants.2017.01.006>, 2017.
- Dassios, K. G., and Pandis, S. N.: Evaporation of ammonium nitrate aerosol and the effect of organic coatings, *J. Aerosol Sci.*, 29, S983-S984, [https://doi.org/10.1016/S0021-8502\(98\)90674-7](https://doi.org/10.1016/S0021-8502(98)90674-7), 1998.
- 1625 De Haan, D. O., Corrigan, A. L., Smith, K. W., Stroik, D. R., Turley, J. J., Lee, F. E., Tolbert, M. A., Jimenez, J. L., Cordova, K. E., and Ferrell, G. R.: Secondary organic aerosol-forming reactions of glyoxal with amino acids, *Environ. Sci. Technol.*, 43, 2818-2824, <https://doi.org/10.1021/es803534f>, 2009a.
- De Haan, D. O., Corrigan, A. L., Tolbert, M. A., Jimenez, J. L., Wood, S. E., and Turley, J. J.: Secondary Organic Aerosol Formation by Self-Reactions of Methylglyoxal and Glyoxal in Evaporating Droplets, *Environ. Sci. Technol.*, 43, 8184-8190, <https://doi.org/10.1021/es902152t>, 2009b.
- 1630 De Haan, D. O., Hawkins, L. N., Kononenko, J. A., Turley, J. J., Corrigan, A. L., Tolbert, M. A., and Jimenez, J. L.: Formation of nitrogen-containing oligomers by methylglyoxal and amines in simulated evaporating cloud droplets, *Environ. Sci. Technol.*, 45, 984-991, <https://doi.org/10.1021/es102933x>, 2011.
- 1635 Deguillaume, L., Leriche, M., Desboeufs, K., Mailhot, G., George, C., and Chaumerliac, N.: Transition metals in atmospheric liquid phases: Sources, reactivity, and sensitive parameters, *Chem. Rev.*, 105, 3388-3431, <https://doi.org/10.1021/cr040649c>, 2005.
- Dentener, F., Williams, J., and Metzger, S.: Aqueous phase reaction of HNO₄: The impact on tropospheric chemistry, *J. Atmos. Chem.*, 41, 109-134, <https://doi.org/10.1023/A:1014233910126>, 2002.
- 1640 Dovrou, E., Rivera-Rios, J. C., Bates, K. H., and Keutsch, F. N.: Sulfate Formation via Cloud Processing from Isoprene Hydroxyl Hydroperoxides (ISOPOOH), *Environ. Sci. Technol.*, 53, 12476-12484, <https://doi.org/10.1021/acs.est.9b04645>, 2019.
- Drexler, C., Elias, H., Fecher, B., and Wannowius, K. J.: Kinetic Investigation of Sulfur(IV) Oxidation by Peroxo Compounds R-OOH in Aqueous-Solution, *Fresen. J. Anal. Chem.*, 340, 605-615, <https://doi.org/10.1007/Bf00321521>, 1991.



- 1645 Duncan, J. L., Schindler, L. R., and Roberts, J. T.: Chemistry at and near the Surface of Liquid Sulfuric Acid: A Kinetic, Thermodynamic, and Mechanistic Analysis of Heterogeneous Reactions of Acetone, *J. Phys. Chem. B*, 103, 7247-7259, <https://doi.org/10.1021/jp991322w>, 1999.
- Duporte, G., Flaud, P. M., Geneste, E., Augagneur, S., Pangui, E., Lamkaddam, H., Gratien, A., Doussin, J. F., Budzinski, H., Villenave, E., and Perraudin, E.: Experimental Study of the Formation of Organosulfates from alpha-Pinene Oxidation. Part I: Product Identification, Formation Mechanisms and Effect of Relative Humidity, *J. Phys. Chem. A*, 120, 7909-7923, <https://doi.org/10.1021/acs.jpca.6b08504>, 2016.
- 1650 Duporte, G., Flaud, P. M., Kammer, J., Geneste, E., Augagneur, S., Pangui, E., Lamkaddam, H., Gratien, A., Doussin, J. F., Budzinski, H., Villenave, E., and Perraudin, E.: Experimental Study of the Formation of Organosulfates from alpha-Pinene Oxidation. 2. Time Evolution and Effect of Particle Acidity, *J. Phys. Chem. A*, 124, 409-421, <https://doi.org/10.1021/acs.jpca.9b07156>, 2020.
- 1655 Eddingsaas, N. C., VanderVelde, D. G., and Wennberg, P. O.: Kinetics and products of the acid-catalyzed ring-opening of atmospherically relevant butyl epoxy alcohols, *J. Phys. Chem. A*, 114, 8106-8113, <https://doi.org/10.1021/jp103907c>, 2010.
- Ervens, B., Herckes, P., Feingold, G., Lee, T., Collett, J. L., and Kreidenweis, S. M.: On the drop-size dependence of organic acid and formaldehyde concentrations in fog, *J. Atmos. Chem.*, 46, 239-269, <https://doi.org/10.1023/A:1026393805907>, 2003.
- 1660 Ervens, B.: Modeling the processing of aerosol and trace gases in clouds and fogs, *Chem. Rev.*, 115, 4157-4198, <https://doi.org/10.1021/cr5005887>, 2015.
- Esteve, W., and Noziere, B.: Uptake and reaction kinetics of acetone, 2-butanone, 2,4-pentanedione, and acetaldehyde in sulfuric acid solutions, *J. Phys. Chem. A*, 109, 10920-10928, <https://doi.org/10.1021/jp051199a>, 2005.
- 1665 Evans, M. J., Jacob, D. J., Atlas, E., Cantrell, C. A., Eisele, F., Flocke, F., Fried, A., Mauldin, R. L., Ridley, B. A., Wert, B., Talbot, R., Blake, D., Heikes, B., Snow, J., Walega, J., Weinheimer, A. J., and Dibb, J.: Coupled evolution of BrOx-ClOx-HOx-NOx chemistry during bromine-catalyzed ozone depletion events in the arctic boundary layer, *J. Geophys. Res.-Atmos.*, 108, 8368, <https://doi.org/10.1029/2002jd002732>, 2003.
- Exner, M., Herrmann, H., and Zellner, R.: Laser-Based Studies of Reactions of the Nitrate Radical in Aqueous Solution, *Ber. Bunsenges. Phys. Chem.*, 96, 470-477, <https://doi.org/10.1002/bbpc.19920960347>, 1992.
- 1670 Faloon, I.: Sulfur processing in the marine atmospheric boundary layer: A review and critical assessment of modeling uncertainties, *Atmos. Environ.*, 43, 2841-2854, <https://doi.org/10.1016/j.atmosenv.2009.02.043>, 2009.
- Fang, T., Guo, H., Zeng, L., Verma, V., Nenes, A., and Weber, R. J.: Highly Acidic Ambient Particles, Soluble Metals, and Oxidative Potential: A Link between Sulfate and Aerosol Toxicity, *Environ. Sci. Technol.*, 51, 2611-2620, <https://doi.org/10.1021/acs.est.6b06151>, 2017.
- 1675 Feingold, G., Frost, G. J., and Ravishankara, A. R.: Role of NO₃ in sulfate production in the wintertime northern latitudes, *J. Geophys. Res.-Atmos.*, 107, 4640, <https://doi.org/10.1029/2002jd002288>, 2002.
- Felber, T., Schaefer, T., and Herrmann, H.: OH-Initiated Oxidation of Imidazoles in Tropospheric Aqueous-Phase Chemistry, *J. Phys. Chem. A*, 123, 1505-1513, <https://doi.org/10.1021/acs.jpca.8b11636>, 2019.
- 1680 Finlayson-Pitts, B. J., Ezell, M. J., and Pitts, J. N.: Formation of chemically active chlorine compounds by reactions of atmospheric NaCl particles with gaseous N₂O₅ and ClONO₂, *Nature*, 337, 241-244, <https://doi.org/10.1038/337241a0>, 1989.
- Fogelman, K. D., Walker, D. M., and Margerum, D. W.: Nonmetal redox kinetics: hypochlorite and hypochlorous acid reactions with sulfite, *Inorg. Chem.*, 28, 986-993, <https://doi.org/10.1021/ic00305a002>, 1989.
- Fountoukis, C., and Nenes, A.: ISORROPIA II: a computationally efficient thermodynamic equilibrium model for K⁺ - Ca²⁺ - Mg²⁺ - NH₄⁺ - Na⁺ - SO₄²⁻ - NO₃⁻ - Cl⁻ - H₂O aerosols, *Atmos. Chem. Phys.*, 7, 4639-4659, <https://doi.org/10.5194/acp-7-4639-2007>, 2007.
- 1685 Fuzzi, S., Baltensperger, U., Carslaw, K., Decesari, S., Denier van der Gon, H., Facchini, M. C., Fowler, D., Koren, I., Langford, B., Lohmann, U., Nemitz, E., Pandis, S., Riipinen, I., Rudich, Y., Schaap, M., Slowik, J. G., Spracklen, D. V., Vignati, E., Wild, M., Williams, M., and Gilardoni, S.: Particulate matter, air quality and climate: lessons learned and future needs, *Atmos. Chem. Phys.*, 15, 8217-8299, <https://doi.org/10.5194/acp-15-8217-2015>, 2015.
- 1690 Galloway, M. M., Chhabra, P. S., Chan, A. W. H., Surratt, J. D., Flagan, R. C., Seinfeld, J. H., and Keutsch, F. N.: Glyoxal uptake on ammonium sulphate seed aerosol: reaction products and reversibility of uptake under dark and irradiated conditions, *Atmos. Chem. Phys.*, 9, 3331-3345, <https://doi.org/10.5194/acp-9-3331-2009>, 2009.



- 1695 Gani, S., Bhandari, S., Seraj, S., Wang, D. S., Patel, K., Soni, P., Arub, Z., Habib, G., Hildebrandt Ruiz, L., and Apte, J. S.: Submicron aerosol composition in the world's most polluted megacity: the Delhi Aerosol Supersite study, *Atmos. Chem. Phys.*, 19, 6843-6859, <https://doi.org/10.5194/acp-19-6843-2019>, 2019.
- Gao, S., Ng, N. L., Keywood, M., Varutbangkul, V., Bahreini, R., Nenes, A., He, J., Yoo, K. Y., Beauchamp, J. L., Hodyss, R. P., Flagan, R. C., and Seinfeld, J. H.: Particle Phase Acidity and Oligomer Formation in Secondary Organic Aerosol, *Environ. Sci. Technol.*, 38, 6582-6589, <https://doi.org/10.1021/es049125k>, 2004.
- 1700 Gaston, C. J., Riedel, T. P., Zhang, Z., Gold, A., Surratt, J. D., and Thornton, J. A.: Reactive uptake of an isoprene-derived epoxydiol to submicron aerosol particles, *Environ. Sci. Technol.*, 48, 11178-11186, <https://doi.org/10.1021/es5034266>, 2014.
- Gen, M. S., Zhang, R. F., Huang, D. D., Li, Y. J., and Chan, C. K.: Heterogeneous SO₂ Oxidation in Sulfate Formation by Photolysis of Particulate Nitrate, *Environ. Sci. Tech. Lett.*, 6, 86-91, <https://doi.org/10.1021/acs.estlett.8b00681>, 2019.
- George, C., Ammann, M., D'Anna, B., Donaldson, D. J., and Nizkorodov, S. A.: Heterogeneous Photochemistry in the Atmosphere, *Chem. Rev.*, 115, 4218-4258, <https://doi.org/10.1021/cr500648z>, 2015.
- 1705 Goetz, J. D., Giordano, M. R., Stockwell, C. E., Christian, T. J., Maharjan, R., Adhikari, S., Bhave, P. V., Praveen, P. S., Panday, A. K., Jayarathne, T., Stone, E. A., Yokelson, R. J., and DeCarlo, P. F.: Speciated online PM_{2.5} from South Asian combustion sources – Part 1: Fuel-based emission factors and size distributions, *Atmos. Chem. Phys.*, 18, 14653-14679, <https://doi.org/10.5194/acp-18-14653-2018>, 2018.
- Grace, D. N., Sharp, J. R., Holappa, R. E., Lugos, E. N., Sebold, M. B., Griffith, D. R., Hendrickson, H. P., and Galloway, M. M.: Heterocyclic Product Formation in Aqueous Brown Carbon Systems, *ACS Earth Space Chem.*, 3, 2472-2481, <https://doi.org/10.1021/acsearthspacechem.9b00235>, 2019.
- 1710 Graedel, T. E., and Weschler, C. J.: Chemistry within Aqueous Atmospheric Aerosols and Raindrops, *Rev. Geophys.*, 19, 505-539, <https://doi.org/10.1029/RG019i004p00505>, 1981.
- Graedel, T. E., and Goldberg, K. I.: Kinetic-Studies of Raindrop Chemistry .1. Inorganic and Organic Processes, *J. Geophys. Res.-Oceans*, 88, 865-882, <https://doi.org/10.1029/JC088iC15p10865>, 1983.
- 1715 Grgić, I., Hudnik, V., Bizjak, M., and Levec, J.: Aqueous S(IV) oxidation—II. Synergistic effects of some metal ions, *Atmos. Environ.*, 26, 571-577, [https://doi.org/10.1016/0960-1686\(92\)90170-p](https://doi.org/10.1016/0960-1686(92)90170-p), 1992.
- Grgić, I., and Berčić, G.: A simple kinetic model for autoxidation of S(IV) oxides catalyzed by iron and/or manganese ions, *J. Atmos. Chem.*, 39, 155-170, <https://doi.org/10.1023/a:1010638902653>, 2001.
- 1720 Gunz, D. W., and Hoffmann, M. R.: Atmospheric chemistry of peroxides: A review, *Atmos. Environ.*, 24, 1601-1633, [https://doi.org/10.1016/0960-1686\(90\)90496-a](https://doi.org/10.1016/0960-1686(90)90496-a), 1990.
- Guo, H., Weber, R. J., and Nenes, A.: High levels of ammonia do not raise fine particle pH sufficiently to yield nitrogen oxide-dominated sulfate production, *Sci. Rep.*, 7, 12109, <https://doi.org/10.1038/s41598-017-11704-0>, 2017.
- 1725 Gypens, N., and Borges, A. V.: Increase in dimethylsulfide (DMS) emissions due to eutrophication of coastal waters offsets their reduction due to ocean acidification, *Front. Mar. Sci.*, 1, <https://doi.org/10.3389/fmars.2014.00004>, 2014.
- Hamilton, J. F., Lewis, A. C., Reynolds, J. C., Carpenter, L. J., and Lubben, A.: Investigating the composition of organic aerosol resulting from cyclohexene ozonolysis: low molecular weight and heterogeneous reaction products, *Atmos. Chem. Phys.*, 6, 4973-4984, <https://doi.org/10.5194/acp-6-4973-2006>, 2006.
- 1730 Harris, E., Sinha, B., van Pinxteren, D., Tilgner, A., Fomba, K. W., Schneider, J., Roth, A., Gnauk, T., Fahlbusch, B., Mertes, S., Lee, T., Collett, J., Foley, S., Borrmann, S., Hoppe, P., and Herrmann, H.: Enhanced role of transition metal ion catalysis during in-cloud oxidation of SO₂, *Science*, 340, 727-730, <https://doi.org/10.1126/science.1230911>, 2013.
- Hastings, W. P., Koehler, C. A., Bailey, E. L., and De Haan, D. O.: Secondary Organic Aerosol Formation by Glyoxal Hydration and Oligomer Formation: Humidity Effects and Equilibrium Shifts during Analysis, *Environ. Sci. Technol.*, 39, 8728-8735, <https://doi.org/10.1021/es050446l>, 2005.
- 1735 Hawkins, L. N., Welsh, H. G., and Alexander, M. V.: Evidence for pyrazine-based chromophores in cloud water mimics containing methylglyoxal and ammonium sulfate, *Atmos. Chem. Phys.*, 18, 12413-12431, <https://doi.org/10.5194/acp-18-12413-2018>, 2018.
- 1740 He, P. Z., Alexander, B., Geng, L., Chi, X. Y., Fan, S. D., Zhan, H. C., Kang, H., Zheng, G. J., Cheng, Y. F., Su, H., Liu, C., and Xie, Z. Q.: Isotopic constraints on heterogeneous sulfate production in Beijing haze, *Atmos. Chem. Phys.*, 18, 5515-5528, <https://doi.org/10.5194/acp-18-5515-2018>, 2018a.



- He, Q. F., Ding, X., Fu, X. X., Zhang, Y. Q., Wang, J. Q., Liu, Y. X., Tang, M. J., Wang, X. M., and Rudich, Y.: Secondary Organic Aerosol Formation From Isoprene Epoxides in the Pearl River Delta, South China: IEPOX- and HMML-Derived Tracers, *J. Geophys. Res.-Atmos.*, 123, 6999-7012, <https://doi.org/10.1029/2017jd028242>, 2018b.
- 1745 Hering, S. V., and Friedlander, S. K.: Origins of Aerosol Sulfur Size Distributions in the Los-Angeles Basin, *Atmos. Environ.*, 16, 2647-2656, [https://doi.org/10.1016/0004-6981\(82\)90346-8](https://doi.org/10.1016/0004-6981(82)90346-8), 1982.
- Herrmann, H., and Zellner, R.: Reactions of NO₃-Radicals in Aqueous Solution, in: *The Chemistry of Free Radicals: N-Centered Radicals*, edited by: Alfassi, Z. B., Wiley, Chichester, 291-343, 1998.
- Herrmann, H.: Kinetics of aqueous phase reactions relevant for atmospheric chemistry, *Chem. Rev.*, 103, 4691-4716, <https://doi.org/10.1021/cr020658q>, 2003.
- 1750 Herrmann, H., Hoffmann, D., Schaefer, T., Brauer, P., and Tilgner, A.: Tropospheric aqueous-phase free-radical chemistry: radical sources, spectra, reaction kinetics and prediction tools, *ChemPhysChem*, 11, 3796-3822, <https://doi.org/10.1002/cphc.201000533>, 2010.
- Herrmann, H., Schaefer, T., Tilgner, A., Styler, S. A., Weller, C., Teich, M., and Otto, T.: Tropospheric aqueous-phase chemistry: kinetics, mechanisms, and its coupling to a changing gas phase, *Chem. Rev.*, 115, 4259-4334, <https://doi.org/10.1021/cr500447k>, 2015.
- 1755 Hoffmann, E. H., Tilgner, A., Schrodner, R., Brauer, P., Wolke, R., and Herrmann, H.: An advanced modeling study on the impacts and atmospheric implications of multiphase dimethyl sulfide chemistry, *Proc. Natl. Acad. Sci. USA*, 113, 11776-11781, <https://doi.org/10.1073/pnas.1606320113>, 2016.
- Hoffmann, E. H., Tilgner, A., Vogelsberg, U., Wolke, R., and Herrmann, H.: Near-Explicit Multiphase Modeling of Halogen Chemistry in a Mixed Urban and Maritime Coastal Area, *ACS Earth Space Chem.*, 3, 2452-2471, <https://doi.org/10.1021/acsearthspacechem.9b00184>, 2019a.
- Hoffmann, E. H., Tilgner, A., Wolke, R., and Herrmann, H.: Enhanced Chlorine and Bromine Atom Activation by Hydrolysis of Halogen Nitrates from Marine Aerosols at Polluted Coastal Areas, *Environ. Sci. Technol.*, 53, 771-778, <https://doi.org/10.1021/acs.est.8b05165>, 2019b.
- 1765 Hoffmann, E. H., Schrödner, R., Tilgner, A., Wolke, R., and Herrmann, H.: CAPRAM reduction towards an operational multiphase halogen and dimethyl sulfide chemistry treatment in the chemistry transport model COSMO-MUSCAT(5.04e), *Geosci. Model Dev.*, 13, 2587-2609, <https://doi.org/10.5194/gmd-13-2587-2020>, 2020.
- Hoffmann, M. R., and Calvert, J. G.: Chemical transformation modules for eulerian acid deposition models, v2, the aqueous-phase chemistry, EPA/600/3-85/017, US Environmental Protection Agency, Research Triangle Park, NC, 1985.
- 1770 Hoigne, J., and Bader, H.: Rate Constants of Reactions of Ozone with Organic and Inorganic-Compounds in Water .1. Non-Dissociating Organic-Compounds, *Water Res.*, 17, 173-183, [https://doi.org/10.1016/0043-1354\(83\)90098-2](https://doi.org/10.1016/0043-1354(83)90098-2), 1983a.
- Hoigne, J., and Bader, H.: Rate Constants of Reactions of Ozone with Organic and Inorganic-Compounds in Water .2. Dissociating Organic-Compounds, *Water Res.*, 17, 185-194, [https://doi.org/10.1016/0043-1354\(83\)90099-4](https://doi.org/10.1016/0043-1354(83)90099-4), 1983b.
- 1775 Hoigne, J., Bader, H., Haag, W. R., and Staehelin, J.: Rate Constants of Reactions of Ozone with Organic and Inorganic-Compounds in Water .3. Inorganic-Compounds and Radicals, *Water Res.*, 19, 993-1004, [https://doi.org/10.1016/0043-1354\(85\)90368-9](https://doi.org/10.1016/0043-1354(85)90368-9), 1985.
- Holmes, B. J., and Petrucci, G. A.: Water-Soluble Oligomer Formation from Acid-Catalyzed Reactions of Levoglucosan in Proxies of Atmospheric Aqueous Aerosols, *Environ. Sci. Technol.*, 40, 4983-4989, <https://doi.org/10.1021/es060646c>, 2006.
- Holmes, B. J., and Petrucci, G. A.: Oligomerization of levoglucosan by Fenton chemistry in proxies of biomass burning aerosols, *J. Atmos. Chem.*, 58, 151-166, <https://doi.org/10.1007/s10874-007-9084-8>, 2007.
- 1780 Hopkins, F. E., Nightingale, P. D., Stephens, J. A., Moore, C. M., Richier, S., Cripps, G. L., and Archer, S. D.: A meta-analysis of microcosm experiments shows that dimethyl sulfide (DMS) production in polar waters is insensitive to ocean acidification, *Biogeosciences*, 17, 163-186, <https://doi.org/10.5194/bg-17-163-2020>, 2020.
- Huie, R. E., and Neta, P.: Chemical behavior of sulfur trioxide(1-) (SO₃⁻) and sulfur pentoxide(1-) (SO₅⁻) radicals in aqueous solutions, *J. Phys. Chem.*, 88, 5665-5669, <https://doi.org/10.1021/j150667a042>, 1984.
- 1785 Humphreys, E. M.: Atherosclerosis in the Coronary Arteries of Rats, *J. Atheroscler. Res.*, 4, 416-434, [https://doi.org/10.1016/s0368-1319\(64\)80026-0](https://doi.org/10.1016/s0368-1319(64)80026-0), 1964.
- Hung, H. M., and Hoffmann, M. R.: Oxidation of Gas-Phase SO₂ on the Surfaces of Acidic Microdroplets: Implications for Sulfate and Sulfate Radical Anion Formation in the Atmospheric Liquid Phase, *Environ. Sci. Technol.*, 49, 13768-13776, <https://doi.org/10.1021/acs.est.5b01658>, 2015.
- 1790



- Hyslop, N. P.: Impaired visibility: the air pollution people see, *Atmos. Environ.*, 43, 182-195, <https://doi.org/10.1016/j.atmosenv.2008.09.067>, 2009.
- 1795 Ibusuki, T., and Takeuchi, K.: Sulfur-Dioxide Oxidation by Oxygen Catalyzed by Mixtures of Manganese(II) and Iron(III) in Aqueous-Solutions at Environmental Reaction Conditions, *Atmos. Environ.*, 21, 1555-1560, [https://doi.org/10.1016/0004-6981\(87\)90317-9](https://doi.org/10.1016/0004-6981(87)90317-9), 1987.
- Inuma, Y., Böge, O., Gnauk, T., and Herrmann, H.: Aerosol-chamber study of the α -pinene/O₃ reaction: influence of particle acidity on aerosol yields and products, *Atmos. Environ.*, 38, 761-773, <https://doi.org/10.1016/j.atmosenv.2003.10.015>, 2004.
- Inuma, Y., Müller, C., Berndt, T., Böge, O., Claeys, M., and Herrmann, H.: Evidence for the Existence of Organosulfates from β -Pinene Ozonolysis in Ambient Secondary Organic Aerosol, *Environ. Sci. Technol.*, 41, 6678-6683, <https://doi.org/10.1021/es070938t>, 2007.
- 1800 Imamura, T., and Akiyoshi, H.: Uptake of acetone into sulfuric-acid solutions, *Geophys. Res. Lett.*, 27, 1419-1422, <https://doi.org/10.1029/1999gl011137>, 2000.
- Ingold, C.: *Structure and Mechanism in Organic Chemistry*, Cornell University Press, Ithaca, N.Y., 1969.
- Ip, H. S. S., Huang, X. H. H., and Yu, J. Z.: Effective Henry's law constants of glyoxal, glyoxylic acid, and glycolic acid, *Geophys. Res. Lett.*, 36, L01802, <https://doi.org/10.1029/2008gl036212>, 2009.
- 1805 Jacob, D. J., and Hoffmann, M. R.: A dynamic model for the production of H⁺, NO₃⁻, and SO₄²⁻ in urban fog, *J. Geophys. Res.-Atmos.*, 88, <https://doi.org/10.1029/JC088iC11p06611>, 1983.
- Jacob, D. J., Munger, J. W., Waldman, J. M., and Hoffmann, M. R.: The H₂SO₄-HNO₃-NH₃ System at High Humidities and in Fogs .1. Spatial and Temporal Patterns in the San-Joaquin-Valley of California, *J. Geophys. Res.-Atmos.*, 91, 1073-1088, <https://doi.org/10.1029/JD091iD01p01073>, 1986.
- 1810 Jang, M., and Kamens, R. M.: Atmospheric secondary aerosol formation by heterogeneous reactions of aldehydes in the presence of a sulfuric acid aerosol catalyst, *Environ. Sci. Technol.*, 35, 4758-4766, <https://doi.org/10.1021/es010790s>, 2001.
- Jang, M., Czoschke, N. M., Lee, S., and Kamens, R. M.: Heterogeneous atmospheric aerosol production by acid-catalyzed particle-phase reactions, *Science*, 298, 814-817, <https://doi.org/10.1126/science.1075798>, 2002.
- 1815 Jang, M., Carroll, B., Chandramouli, B., and Kamens, R. M.: Particle growth by acid-catalyzed heterogeneous reactions of organic carbonyls on preexisting aerosols, *Environ. Sci. Technol.*, 37, 3828-3837, <https://doi.org/10.1021/es021005u>, 2003.
- Jang, M., Czoschke, N. M., and Northcross, A. L.: Atmospheric organic aerosol production by heterogeneous acid-catalyzed reactions, *ChemPhysChem*, 5, 1647-1661, <https://doi.org/10.1002/cphc.200301077>, 2004.
- 1820 Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., Dunlea, J., Huffman, J. A., Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimojo, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of Organic Aerosols in the Atmosphere, *Science*, 326, 1525, <https://doi.org/10.1126/science.1180353>, 2009.
- 1825 Kalberer, M., Paulsen, D., Sax, M., Steinbacher, M., Dommen, J., Prevot, A. S., Fisseha, R., Weingartner, E., Frankevich, V., Zenobi, R., and Baltensperger, U.: Identification of polymers as major components of atmospheric organic aerosols, *Science*, 303, 1659-1662, <https://doi.org/10.1126/science.1092185>, 2004.
- 1830 Kampf, C. J., Jakob, R., and Hoffmann, T.: Identification and characterization of aging products in the glyoxal/ammonium sulfate system – implications for light-absorbing material in atmospheric aerosols, *Atmos. Chem. Phys.*, 12, 6323-6333, <https://doi.org/10.5194/acp-12-6323-2012>, 2012.
- Kampf, C. J., Waxman, E. M., Slowik, J. G., Dommen, J., Pfaffenberger, L., Praplan, A. P., Prevot, A. S. H., Baltensperger, U., Hoffmann, T., and Volkamer, R.: Effective Henry's Law Partitioning and the Salting Constant of Glyoxal in Aerosols Containing Sulfate, *Environ. Sci. Technol.*, 47, 4236-4244, <https://doi.org/10.1021/es400083d>, 2013.
- 1835 Kampf, C. J., Filippi, A., Zuth, C., Hoffmann, T., and Opatz, T.: Secondary brown carbon formation via the dicarbonyl imine pathway: nitrogen heterocycle formation and synergistic effects, *Phys. Chem. Chem. Phys.*, 18, 18353-18364, <https://doi.org/10.1039/C6CP03029G>, 2016.



- 1840 Kane, S. M., Timonen, R. S., and Leu, M.-T.: Heterogeneous Chemistry of Acetone in Sulfuric Acid Solutions: Implications for the Upper Troposphere, *J. Phys. Chem. A*, 103, 9259-9265, <https://doi.org/10.1021/jp9926692>, 1999.
- Kawamura, K., and Bikkina, S.: A review of dicarboxylic acids and related compounds in atmospheric aerosols: Molecular distributions, sources and transformation, *Atmos. Res.*, 170, 140-160, <https://doi.org/10.1016/j.atmosres.2015.11.018>, 2016.
- 1845 Keene, W. C., and Galloway, J. N.: Organic Acidity in Precipitation of North-America, *Atmos. Environ.*, 18, 2491-2497, [https://doi.org/10.1016/0004-6981\(84\)90020-9](https://doi.org/10.1016/0004-6981(84)90020-9), 1984.
- Keene, W. C., Sander, R., Pszenny, A. A. P., Vogt, R., Crutzen, P. J., and Galloway, J. N.: Aerosol pH in the marine boundary layer: A review and model evaluation, *J. Aerosol Sci.*, 29, 339-356, [https://doi.org/10.1016/S0021-8502\(97\)10011-8](https://doi.org/10.1016/S0021-8502(97)10011-8), 1998.
- Kok, G. L., Gitlin, S. N., and Lazrus, A. L.: Kinetics of the Formation and Decomposition of Hydroxymethanesulfonate, *J. Geophys. Res.-Atmos.*, 91, 2801-2804, <https://doi.org/10.1029/JD091iD02p02801>, 1986.
- 1850 Koop, T., Bookhold, J., Shiraiwa, M., and Pöschl, U.: Glass transition and phase state of organic compounds: dependency on molecular properties and implications for secondary organic aerosols in the atmosphere, *Phys. Chem. Chem. Phys.*, 13, 19238-19255, <https://doi.org/10.1039/c1cp22617g>, 2011.
- Kourtchev, I., Ruuskanen, T., Maenhaut, W., Kulmala, M., and Claeys, M.: Observation of 2-methyltetrols and related photo-oxidation products of isoprene in boreal forest aerosols from Hyytiälä, Finland, *Atmos. Chem. Phys.*, 5, 2761-2770, <https://doi.org/10.5194/acp-5-2761-2005>, 2005.
- 1855 Kristensen, K., Enggrob, K. L., King, S. M., Worton, D. R., Platt, S. M., Mortensen, R., Rosenoern, T., Surratt, J. D., Bilde, M., Goldstein, A. H., and Glasius, M.: Formation and occurrence of dimer esters of pinene oxidation products in atmospheric aerosols, *Atmos. Chem. Phys.*, 13, 3763-3776, <https://doi.org/10.5194/acp-13-3763-2013>, 2013.
- Krizner, H. E., De Haan, D. O., and Kua, J.: Thermodynamics and Kinetics of Methylglyoxal Dimer Formation: A Computational Study, *J. Phys. Chem. A*, 113, 6994-7001, <https://doi.org/10.1021/jp903213k>, 2009.
- 1860 Kroll, J. H., Ng, N. L., Murphy, S. M., Varutbangkul, V., Flagan, R. C., and Seinfeld, J. H.: Chamber studies of secondary organic aerosol growth by reactive uptake of simple carbonyl compounds, *J. Geophys. Res.*, 110, <https://doi.org/10.1029/2005jd006004>, 2005.
- Kurz, J. L.: Hydration of acetaldehyde. I. Equilibrium thermodynamic parameters, *J. Am. Chem. Soc.*, 89, 3524-3528, <https://doi.org/10.1021/ja00990a032>, 1967.
- 1865 Kurz, J. L., and Coburn, J. I.: The Hydration of Acetaldehyde. II. Transition-State Characterization, *J. Am. Chem. Soc.*, 89, 3528-3537, <https://doi.org/10.1021/ja00990a600>, 1967.
- Kuschel, A., and Polarz, S.: Effects of primary and secondary surface groups in enantioselective catalysis using nanoporous materials with chiral walls, *J. Am. Chem. Soc.*, 132, 6558-6565, <https://doi.org/10.1021/ja1017706>, 2010.
- 1870 Lagrange, J., Pallares, C., Wenger, G., and Lagrange, P.: Electrolyte effects on aqueous atmospheric oxidation of sulphur dioxide by hydrogen peroxide, *Atmos. Environ.*, 27, 129-137, [https://doi.org/10.1016/0960-1686\(93\)90342-V](https://doi.org/10.1016/0960-1686(93)90342-V), 1993.
- Lagrange, J., Pallares, C., and Lagrange, P.: Electrolyte effects on aqueous atmospheric oxidation of sulphur dioxide by ozone, *J. Geophys. Res.-Atmos.*, 99, 14595-14600, <https://doi.org/10.1029/94JD00573>, 1994.
- 1875 Larson, R. A., and Weber, E. J.: *Reaction Mechanisms in Environmental Organic Chemistry*, Lewis Publishers CRC Press, Boca Raton, FL, 1994.
- Laskin, A., Laskin, J., and Nizkorodov, S. A.: Chemistry of Atmospheric Brown Carbon, *Chem. Rev.*, 115, 4335-4382, <https://doi.org/10.1021/cr5006167>, 2015.
- 1880 Le Breton, M., Wang, Y., Hallquist, Å. M., Pathak, R. K., Zheng, J., Yang, Y., Shang, D., Glasius, M., Bannan, T. J., Liu, Q., Chan, C. K., Percival, C. J., Zhu, W., Lou, S., Topping, D., Wang, Y., Yu, J., Lu, K., Guo, S., Hu, M., and Hallquist, M.: Online gas- and particle-phase measurements of organosulfates, organosulfonates and nitrooxy organosulfates in Beijing utilizing a FIGAERO ToF-CIMS, *Atmos. Chem. Phys.*, 18, 10355-10371, <https://doi.org/10.5194/acp-18-10355-2018>, 2018.
- Lee, S.-H., Gordon, H., Yu, H., Lehtipalo, K., Haley, R., Li, Y., and Zhang, R.: New Particle Formation in the Atmosphere: From Molecular Clusters to Global Climate, *J. Geophys. Res.-Atmos.*, 124, 7098-7146, <https://doi.org/10.1029/2018jd029356>, 2019.
- 1885 Lee, Y.-N., and Schwartz, S. E.: Kinetics of Oxidation of Aqueous Sulfur(IV) by Nitrogen Dioxide, in: *Precipitation Scavenging, Dry Deposition, and Resuspension. Volume 1: Precipitation Scavenging*, edited by: Pruppacher, H. R., Semonin, R. G., and Slinn, W. G., Elsevier, New York, Amsterdam, Oxford, 453-470, 1983.
- Leitzke, A., and von Sonntag, C.: Ozonolysis of Unsaturated Acids in Aqueous Solution: Acrylic, Methacrylic, Maleic, Fumaric and Muconic Acids, *Ozone-Sci. Eng.*, 31, 301-308, <https://doi.org/10.1080/01919510903041354>, 2009.



- 1890 Lelieveld, J., Evans, J. S., Fnais, M., Giannadaki, D., and Pozzer, A.: The contribution of outdoor air pollution sources to premature mortality on a global scale, *Nature*, 525, 367-371, <https://doi.org/10.1038/nature15371>, 2015.
- Li, J., Zhang, Y. L., Cao, F., Zhang, W., Fan, M., Lee, X., and Michalski, G.: Stable Sulfur Isotopes Revealed a Major Role of Transition-Metal Ion-Catalyzed SO₂ Oxidation in Haze Episodes, *Environ. Sci. Technol.*, 54, 2626-2634, <https://doi.org/10.1021/acs.est.9b07150>, 2020a.
- 1895 Li, J., Zhu, C., Chen, H., Fu, H., Xiao, H., Wang, X., Herrmann, H., and Chen, J.: A More Important Role for the Ozone-S(IV) Oxidation Pathway Due to Decreasing Acidity in Clouds, *J. Geophys. Res.-Atmos.*, 125, <https://doi.org/10.1029/2020jd033220>, 2020b.
- Li, L., Chen, Z. M., Zhang, Y. H., Zhu, T., Li, J. L., and Ding, J.: Kinetics and mechanism of heterogeneous oxidation of sulfur dioxide by ozone on surface of calcium carbonate, *Atmos. Chem. Phys.*, 6, 2453-2464, <https://doi.org/10.5194/acp-6-2453-2006>, 2006.
- 1900 Li, L., Hoffmann, M. R., and Colussi, A. J.: Role of Nitrogen Dioxide in the Production of Sulfate during Chinese Haze-Aerosol Episodes, *Environ. Sci. Technol.*, 52, 2686-2693, <https://doi.org/10.1021/acs.est.7b05222>, 2018.
- Li, Z., Schwier, A. N., Sareen, N., and McNeill, V. F.: Reactive processing of formaldehyde and acetaldehyde in aqueous aerosol mimics: surface tension depression and secondary organic products, *Atmos. Chem. Phys.*, 11, 11617-11629, <https://doi.org/10.5194/acp-11-11617-2011>, 2011.
- 1905 Li, Z., Nizkorodov, S. A., Chen, H., Lu, X., Yang, X., and Chen, J.: Nitrogen-containing secondary organic aerosol formation by acrolein reaction with ammonia/ammonium, *Atmos. Chem. Phys.*, 19, 1343-1356, <https://doi.org/10.5194/acp-19-1343-2019>, 2019.
- Liang, L., Engling, G., Duan, F., Cheng, Y., and He, K.: Characteristics of 2-methyltetrols in ambient aerosol in Beijing, China, *Atmos. Environ.*, 59, 376-381, <https://doi.org/10.1016/j.atmosenv.2012.05.052>, 2012.
- 1910 Liggio, J., Li, S.-M., and McLaren, R.: Heterogeneous Reactions of Glyoxal on Particulate Matter: Identification of Acetals and Sulfate Esters, *Environ. Sci. Technol.*, 39, 1532-1541, <https://doi.org/10.1021/es048375y>, 2005a.
- Liggio, J., Li, S.-M., and McLaren, R.: Reactive uptake of glyoxal by particulate matter, *J. Geophys. Res.-Atmos.*, 110, <https://doi.org/10.1029/2004JD005113>, 2005b.
- 1915 Liggio, J., and Li, S.-M.: Reactive uptake of pinonaldehyde on acidic aerosols, *J. Geophys. Res.-Atmos.*, 111, <https://doi.org/10.1029/2005JD006978>, 2006.
- Liljestrand, H. M.: Average Rainwater Ph, Concepts of Atmospheric Acidity, and Buffering in Open Systems, *Atmos. Environ.*, 19, 487-499, [https://doi.org/10.1016/0004-6981\(85\)90169-6](https://doi.org/10.1016/0004-6981(85)90169-6), 1985.
- 1920 Limbeck, A., Kraxner, Y., and Puxbaum, H.: Gas to particle distribution of low molecular weight dicarboxylic acids at two different sites in central Europe (Austria), *J. Aerosol Sci.*, 36, 991-1005, <https://doi.org/10.1016/j.jaerosci.2004.11.013>, 2005.
- Lin, G., Penner, J. E., Sillman, S., Taraborrelli, D., and Lelieveld, J.: Global modeling of SOA formation from dicarbonyls, epoxides, organic nitrates and peroxides, *Atmos. Chem. Phys.*, 12, 4743-4774, <https://doi.org/10.5194/acp-12-4743-2012>, 2012.
- 1925 Lin, P., Laskin, J., Nizkorodov, S. A., and Laskin, A.: Revealing Brown Carbon Chromophores Produced in Reactions of Methylglyoxal with Ammonium Sulfate, *Environ. Sci. Technol.*, 49, 14257-14266, <https://doi.org/10.1021/acs.est.5b03608>, 2015.
- Lin, Y. H., Zhang, H., Pye, H. O., Zhang, Z., Marth, W. J., Park, S., Arashiro, M., Cui, T., Budisulistiorini, S. H., Sexton, K. G., Vizuete, W., Xie, Y., Lueken, D. J., Piletic, I. R., Edney, E. O., Bartolotti, L. J., Gold, A., and Surratt, J. D.: Epoxide as a precursor to secondary organic aerosol formation from isoprene photooxidation in the presence of nitrogen oxides, *Proc. Natl. Acad. Sci. USA*, 110, 6718-6723, <https://doi.org/10.1073/pnas.1221150110>, 2013.
- 1930 Lind, J. A., Lazrus, A. L., and Kok, G. L.: Aqueous phase oxidation of sulfur(IV) by hydrogen peroxide, methylhydroperoxide, and peroxyacetic acid, *J. Geophys. Res.-Atmos.*, 92, <https://doi.org/10.1029/JD092iD04p04171>, 1987.
- Liu, T., and Abbatt, J. P. D.: An Experimental Assessment of the Importance of S(IV) Oxidation by Hypohalous Acids in the Marine Atmosphere, *Geophys. Res. Lett.*, 47, <https://doi.org/10.1029/2019gl086465>, 2020.
- 1935 Liu, T., Clegg, S. L., and Abbatt, J. P. D.: Fast oxidation of sulfur dioxide by hydrogen peroxide in deliquesced aerosol particles, *Proc. Natl. Acad. Sci. USA*, 117, 1354-1359, <https://doi.org/10.1073/pnas.1916401117>, 2020.
- Loeff, I., Rabani, J., Treinin, A., and Linschitz, H.: Charge transfer and reactivity of $n\pi^*$ and $\pi\pi^*$ organic triplets, including anthraquinonesulfonates, in interactions with inorganic anions: A comparative study based on classical Marcus theory, *J. Am. Chem. Soc.*, 115, 8933-8942, <https://doi.org/10.1021/ja00073a007>, 1993.



- 1940 Loeffler, K. W., Koehler, C. A., Paul, N. M., and De Haan, D. O.: Oligomer formation in evaporating aqueous glyoxal and methyl glyoxal solutions, *Environ. Sci. Technol.*, 40, 6318-6323, <https://doi.org/10.1021/es060810w>, 2006.
- Lopalco, A., Douglas, J., Denora, N., and Stella, V. J.: Determination of pKa and Hydration Constants for a Series of alpha-Keto-Carboxylic Acids Using Nuclear Magnetic Resonance Spectrometry, *J. Pharm. Sci.*, 105, 664-672, <https://doi.org/10.1002/jps.24539>, 2016.
- 1945 Loudon, G. M.: *Organic Chemistry*, 3 ed., Benjamin/Cummings, Redwood City, 1995.
- Lowry, T. H., and Richardson, K. S.: *Mechanism and Theory in Organic Chemistry*, Haper & Row, Publishers, New York, Hagerstown, San Francisco, London, 1976.
- Luo, Y. R.: *Handbook of Bond Dissociation Energies in Organic Compounds*, Taylor and Francis, CRC Press, Boca Raton, FL., <https://doi.org/10.1201/9781420039863>, 2002.
- 1950 Ma, T., Furutani, H., Duan, F., Kimoto, T., Jiang, J., Zhang, Q., Xu, X., Wang, Y., Gao, J., Geng, G., Li, M., Song, S., Ma, Y., Che, F., Wang, J., Zhu, L., Huang, T., Toyoda, M., and He, K.: Contribution of hydroxymethanesulfonate (HMS) to severe winter haze in the North China Plain, *Atmos. Chem. Phys.*, 20, 5887-5897, <https://doi.org/10.5194/acp-20-5887-2020>, 2020.
- Maaß, F., Elias, H., and Wannowius, K. J.: Kinetics of the oxidation of hydrogen sulfite by hydrogen peroxide in aqueous solution: ionic strength effects and temperature dependence, *Atmos. Environ.*, 33, 4413-4419, [https://doi.org/10.1016/S1352-2310\(99\)00212-5](https://doi.org/10.1016/S1352-2310(99)00212-5), 1999.
- 1955 Mabey, W., and Mill, T.: Critical-Review of Hydrolysis of Organic-Compounds in Water under Environmental-Conditions, *J. Phys. Chem. Ref. Data*, 7, 383-415, <https://doi.org/10.1063/1.555572>, 1978.
- Mael, L. E., Jacobs, M. I., and Elrod, M. J.: Organosulfate and nitrate formation and reactivity from epoxides derived from 2-methyl-3-buten-2-ol, *J. Phys. Chem. A*, 119, 4464-4472, <https://doi.org/10.1021/jp510033s>, 2015.
- 1960 Manktelow, P. T., Mann, G. W., Carslaw, K. S., Spracklen, D. V., and Chipperfield, M. P.: Regional and global trends in sulfate aerosol since the 1980s, *Geophys. Res. Lett.*, 34, <https://doi.org/10.1029/2006gl028668>, 2007.
- Marais, E. A., Jacob, D. J., Jimenez, J. L., Campuzano-Jost, P., Day, D. A., Hu, W., Krechmer, J., Zhu, L., Kim, P. S., Miller, C. C., Fisher, J. A., Travis, K., Yu, K., Hanisco, T. F., Wolfe, G. M., Arkinson, H. L., Pye, H. O. T., Froyd, K. D., Liao, J., and McNeill, V. F.: Aqueous-phase mechanism for secondary organic aerosol formation from isoprene: application to the southeast United States and co-benefit of SO₂ emission controls, *Atmos. Chem. Phys.*, 16, 1603-1618, <https://doi.org/10.5194/acp-16-1603-2016>, 2016.
- 1965 Maron, M. K., Takahashi, K., Shoemaker, R. K., and Vaida, V.: Hydration of pyruvic acid to its geminal-diol, 2,2-dihydroxypropanoic acid, in a water-restricted environment, *Chem. Phys. Lett.*, 513, 184-190, <https://doi.org/10.1016/j.cplett.2011.07.090>, 2011.
- 1970 Martin, L. R.: Kinetic studies of sulfite oxidation in aqueous solution., in: SO₂, NO and NO₂ oxidation mechanisms: Atmospheric consideration, Acid Precipitation Series, edited by: Calvert, J. G., Butterworth, Boston, 63-100, 1984.
- Martin, L. R., and Hill, M. W.: The effect of ionic strength on the manganese catalyzed oxidation of sulfur(IV), *Atmos. Environ.*, 21, 2267-2270, [https://doi.org/10.1016/0004-6981\(87\)90361-1](https://doi.org/10.1016/0004-6981(87)90361-1), 1987a.
- Martin, L. R., and Hill, M. W.: The iron catalyzed oxidation of sulfur: Reconciliation of the literature rates, *Atmos. Environ.* (1967), 21, 1487-1490, [https://doi.org/10.1016/0004-6981\(67\)90100-x](https://doi.org/10.1016/0004-6981(67)90100-x), 1987b.
- 1975 Martin, L. R., and Good, T. W.: Catalyzed oxidation of sulfur dioxide in solution: The iron-manganese synergism, *Atmos. Environ.*, 25, 2395-2399, [https://doi.org/10.1016/0960-1686\(91\)90113-1](https://doi.org/10.1016/0960-1686(91)90113-1), 1991.
- Martin, L. R., Hill, M. W., Tai, A. F., and Good, T. W.: The Iron Catalyzed Oxidation of Sulfur(IV) in Aqueous-Solution - Differing Effects of Organics at High and Low pH, *J. Geophys. Res.-Atmos.*, 96, 3085-3097, <https://doi.org/10.1029/90jd02611>, 1991.
- 1980 Maxut, A., Nozière, B., Fenet, B., and Mechakra, H.: Formation mechanisms and yields of small imidazoles from reactions of glyoxal with NH₄⁺ in water at neutral pH, *Phys. Chem. Chem. Phys.*, 17, 20416-20424, <https://doi.org/10.1039/C5CP03113C>, 2015.
- 1985 McArdle, J. V., and Hoffmann, M. R.: Kinetics and mechanism of the oxidation of aqated sulfur dioxide by hydrogen peroxide at low pH, *J. Phys. Chem.*, 87, 5425-5429, <https://doi.org/10.1021/j150644a024>, 1983.
- McMurry, P. H.: AEROSOLS | Observations and Measurements, in: *Encyclopedia of Atmospheric Sciences (Second Edition)*, edited by: North, G. R., Pyle, J., and Zhang, F., Academic Press, Oxford, 53-65, <https://doi.org/10.1016/B978-0-12-382225-3.00048-7>, 2015.



- McNeill, K., and Canonica, S.: Triplet state dissolved organic matter in aquatic photochemistry: reaction mechanisms, substrate scope, and photophysical properties, *Environ. Sci. Process. Impacts*, 18, 1381-1399, <https://doi.org/10.1039/c6em00408c>, 2016.
- McNeill, V. F., Woo, J. L., Kim, D. D., Schwier, A. N., Wannell, N. J., Sumner, A. J., and Barakat, J. M.: Aqueous-phase secondary organic aerosol and organosulfate formation in atmospheric aerosols: a modeling study, *Environ. Sci. Technol.*, 46, 8075-8081, <https://doi.org/10.1021/es3002986>, 2012.
- McNeill, V. F.: Aqueous organic chemistry in the atmosphere: sources and chemical processing of organic aerosols, *Environ. Sci. Technol.*, 49, 1237-1244, <https://doi.org/10.1021/es5043707>, 2015.
- McNeill, V. F.: Atmospheric Aerosols: Clouds, Chemistry, and Climate, *Annu. Rev. Chem. Biomol. Eng.*, 8, 427-444, <https://doi.org/10.1146/annurev-chembioeng-060816-101538>, 2017.
- Mekic, M., Liu, J., Zhou, W., Loisel, G., Cai, J., He, T., Jiang, B., Yu, Z., Lazarou, Y. G., Li, X., Brigante, M., Vione, D., and Gligorovski, S.: Formation of highly oxygenated multifunctional compounds from cross-reactions of carbonyl compounds in the atmospheric aqueous phase, *Atmos. Environ.*, 219, <https://doi.org/10.1016/j.atmosenv.2019.117046>, 2019.
- Meng, Z., Seinfeld, J. H., and Saxena, P.: Gas/Aerosol Distribution of Formic and Acetic Acids, *Aerosol Sci. Technol.*, 23, 561-578, <https://doi.org/10.1080/02786829508965338>, 2007.
- Meskhidze, N., Chameides, W. L., and Nenes, A.: Dust and pollution: A recipe for enhanced ocean fertilization?, *J. Geophys. Res.-Atmos.*, 110, D03301, <https://doi.org/10.1029/2004JD005082>, 2005.
- Minerath, E. C., Casale, M. T., and Elrod, M. J.: Kinetics Feasibility Study of Alcohol Sulfate Esterification Reactions in Tropospheric Aerosols, *Environ. Sci. Technol.*, 42, 4410-4415, <https://doi.org/10.1021/es8004333>, 2008.
- Minerath, E. C., and Elrod, M. J.: Assessing the Potential for Diol and Hydroxy Sulfate Ester Formation from the Reaction of Epoxides in Tropospheric Aerosols, *Environ. Sci. Technol.*, 43, 1386-1392, <https://doi.org/10.1021/es8029076>, 2009.
- Moch, J. M., Dovrou, E., Mickley, L. J., Keutsch, F. N., Cheng, Y., Jacob, D. J., Jiang, J. K., Li, M., Munger, J. W., Qiao, X. H., and Zhang, Q.: Contribution of Hydroxymethane Sulfonate to Ambient Particulate Matter: A Potential Explanation for High Particulate Sulfur During Severe Winter Haze in Beijing, *Geophys. Res. Lett.*, 45, 11969-11979, <https://doi.org/10.1029/2018gl079309>, 2018.
- Moch, J. M., Dovrou, E., Mickley, L. J., Keutsch, F. N., Liu, Z., Wang, Y., Dombek, T. L., Kuwata, M., Budisulistiorini, S. H., Yang, L., Decesari, S., Paglione, M., Alexander, B., Shao, J., Munger, J. W., and Jacob, D. J.: Global Importance of Hydroxymethanesulfonate in Ambient Particulate Matter: Implications for Air Quality, *J. Geophys. Res.-Atmos.*, 125, <https://doi.org/10.1029/2020jd032706>, 2020.
- Munger, J. W., Tiller, C., and Hoffmann, M. R.: Identification of Hydroxymethanesulfonate in Fog Water, *Science*, 231, 247, <https://doi.org/10.1126/science.231.4735.247>, 1986.
- Mutzel, A., Poulain, L., Berndt, T., Iinuma, Y., Rodigast, M., Böge, O., Richters, S., Spindler, G., Sipilä, M., Jokinen, T., Kulmala, M., and Herrmann, H.: Highly Oxidized Multifunctional Organic Compounds Observed in Tropospheric Particles: A Field and Laboratory Study, *Environ. Sci. Technol.*, 49, 7754-7761, <https://doi.org/10.1021/acs.est.5b00885>, 2015.
- Mvula, E., and von Sonntag, C.: Ozonolysis of phenols in aqueous solution, *Org. Biomol. Chem.*, 1, 1749-1756, <https://doi.org/10.1039/b301824p>, 2003.
- Nah, T., Guo, H. Y., Sullivan, A. P., Chen, Y. L., Tanner, D. J., Nenes, A., Russell, A., Ng, N. L., Huey, L. G., and Weber, R. J.: Characterization of aerosol composition, aerosol acidity, and organic acid partitioning at an agriculturally intensive rural southeastern US site, *Atmos. Chem. Phys.*, 18, 11471-11491, <https://doi.org/10.5194/acp-18-11471-2018>, 2018.
- Nenes, A., Krom, M. D., Mihalopoulos, N., Van Cappellen, P., Shi, Z., Bougiatioti, A., Zampas, P., and Herut, B.: Atmospheric acidification of mineral aerosols: a source of bioavailable phosphorus for the oceans, *Atmos. Chem. Phys.*, 11, 6265-6272, <https://doi.org/10.5194/acp-11-6265-2011>, 2011.
- Nenes, A., Pandis, S. N., Kanakidou, M., Russell, A., Song, S., Vasilakos, P., and Weber, R. J.: Aerosol acidity and liquid water content regulate the dry deposition of inorganic reactive nitrogen, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2020-266>, 2020a.
- Nenes, A., Pandis, S. N., Weber, R. J., and Russell, A.: Aerosol pH and liquid water content determine when particulate matter is sensitive to ammonia and nitrate availability, *Atmos. Chem. Phys.*, 20, 3249-3258, <https://doi.org/10.5194/acp-20-3249-2020>, 2020b.
- Neta, P., Huie, R. E., and Ross, A. B.: Rate Constants for Reactions of Inorganic Radicals in Aqueous-Solution, *J. Phys. Chem. Ref. Data*, 17, 1027-1284, <https://doi.org/10.1063/1.555808>, 1988.



- Newberg, J. T., Matthew, B. M., and Anastasio, C.: Chloride and bromide depletions in sea-salt particles over the northeastern Pacific Ocean, *J. Geophys. Res.-Atmos.*, 110, n/a-n/a, <https://doi.org/10.1029/2004jd005446>, 2005.
- Nguyen, T. B., Lee, P. B., Updyke, K. M., Bones, D. L., Laskin, J., Laskin, A., and Nizkorodov, S. A.: Formation of nitrogen- and sulfur-containing light-absorbing compounds accelerated by evaporation of water from secondary organic aerosols, *J. Geophys. Res.-Atmos.*, 117, n/a-n/a, <https://doi.org/10.1029/2011jd016944>, 2012.
- Nguyen, T. B., Coggon, M. M., Bates, K. H., Zhang, X., Schwantes, R. H., Schilling, K. A., Loza, C. L., Flagan, R. C., Wennberg, P. O., and Seinfeld, J. H.: Organic aerosol formation from the reactive uptake of isoprene epoxydiols (IEPOX) onto non-acidified inorganic seeds, *Atmos. Chem. Phys.*, 14, 3497-3510, <https://doi.org/10.5194/acp-14-3497-2014>, 2014.
- Noziere, B., and Esteve, W.: Organic reactions increasing the absorption index of atmospheric sulfuric acid aerosols, *Geophys. Res. Lett.*, 32, L03812, <https://doi.org/10.1029/2004gl021942>, 2005.
- Noziere, B., Voisin, D., Longfellow, C. A., Friedli, H., Henry, B. E., and Hanson, D. R.: The uptake of methyl vinyl ketone, methacrolein, and 2-methyl-3-butene-2-ol onto sulfuric acid solutions, *J. Phys. Chem. A*, 110, 2387-2395, <https://doi.org/10.1021/jp0555899>, 2006.
- Noziere, B., and Esteve, W.: Light-absorbing aldol condensation products in acidic aerosols: Spectra, kinetics, and contribution to the absorption index, *Atmos. Environ.*, 41, 1150-1163, <https://doi.org/10.1016/j.atmosenv.2006.10.001>, 2007.
- Noziere, B., and Cordova, A.: A kinetic and mechanistic study of the amino acid catalyzed aldol condensation of acetaldehyde in aqueous and salt solutions, *J. Phys. Chem. A*, 112, 2827-2837, <https://doi.org/10.1021/jp7096845>, 2008.
- Noziere, B., Dziedzic, P., and Cordova, A.: Products and kinetics of the liquid-phase reaction of glyoxal catalyzed by ammonium ions (NH_4^+), *J. Phys. Chem. A*, 113, 231-237, <https://doi.org/10.1021/jp8078293>, 2009.
- Noziere, B., Dziedzic, P., and Cordova, A.: Inorganic ammonium salts and carbonate salts are efficient catalysts for aldol condensation in atmospheric aerosols, *Phys. Chem. Chem. Phys.*, 12, 3864-3872, <https://doi.org/10.1039/b924443c>, 2010.
- Noziere, B., Fache, F., Maxut, A., Fenet, B., Baudouin, A., Fine, L., and Ferronato, C.: The hydrolysis of epoxides catalyzed by inorganic ammonium salts in water: kinetic evidence for hydrogen bond catalysis, *Phys. Chem. Chem. Phys.*, 20, 1583-1590, <https://doi.org/10.1039/c7cp06790a>, 2018.
- Nozière, B., and Riemer, D. D.: The chemical processing of gas-phase carbonyl compounds by sulfuric acid aerosols: 2,4-pentanedione, *Atmos. Environ.*, 37, 841-851, [https://doi.org/10.1016/S1352-2310\(02\)00934-2](https://doi.org/10.1016/S1352-2310(02)00934-2), 2003.
- Nozière, B., Dziedzic, P., and Córdova, A.: Formation of secondary light-absorbing “fulvic-like” oligomers: A common process in aqueous and ionic atmospheric particles?, *Geophys. Res. Lett.*, 34, L21812, <https://doi.org/10.1029/2007gl031300>, 2007.
- Nozière, B., Kalberer, M., Claeys, M., Allan, J., D’Anna, B., Decesari, S., Finessi, E., Glasius, M., Grgić, I., Hamilton, J. F., Hoffmann, T., Iinuma, Y., Jaoui, M., Kahnt, A., Kampf, C. J., Kourtev, I., Maenhaut, W., Marsden, N., Saarikoski, S., Schnelle-Kreis, J., Surratt, J. D., Szidat, S., Szmigielski, R., and Wisthaler, A.: The Molecular Identification of Organic Compounds in the Atmosphere: State of the Art and Challenges, *Chem. Rev.*, 115, 3919-3983, <https://doi.org/10.1021/cr5003485>, 2015.
- Ogata, Y., and Kawasaki, A.: Equilibrium additions to carbonyl compounds, in: *The Chemistry of the Carbonyl Group*, edited by: Zabicky, J., PATAI'S Chemistry of Functional Groups, Interscience Publishers, London, New York, Sydney, Toronto, 2-59, <https://doi.org/10.1002/9780470771228.ch1>, 1970.
- Okochi, H., and Brimblecombe, P.: Potential trace metal-organic complexation in the atmosphere, *ScientificWorldJournal*, 2, 767-786, <https://doi.org/10.1100/tsw.2002.132>, 2002.
- Olson, T. M., and Hoffmann, M. R.: Hydroxyalkylsulfonate formation: Its role as a S(IV) reservoir in atmospheric water droplets, *Atmos. Environ.*, 23, 985-997, [https://doi.org/10.1016/0004-6981\(89\)90302-8](https://doi.org/10.1016/0004-6981(89)90302-8), 1989.
- Oltmans, S. J., Schnell, R. C., Sheridan, P. J., Peterson, R. E., Li, S. M., Winchester, J. W., Tans, P. P., Sturges, W. T., Kahl, J. D., and Barrie, L. A.: Seasonal Surface Ozone and Filterable Bromine Relationship in the High Arctic, *Atmos. Environ.*, 23, 2431-2441, [https://doi.org/10.1016/0004-6981\(89\)90254-0](https://doi.org/10.1016/0004-6981(89)90254-0), 1989.
- Pandis, S. N., Seinfeld, J. H., and Pilinis, C.: Heterogeneous sulfate production in an urban fog, *Atmos. Environ.*, 26, 2509-2522, [https://doi.org/10.1016/0960-1686\(92\)90103-r](https://doi.org/10.1016/0960-1686(92)90103-r), 1992.
- Parrella, J. P., Jacob, D. J., Liang, Q., Zhang, Y., Mickley, L. J., Miller, B., Evans, M. J., Yang, X., Pyle, J. A., Theys, N., and Van Roozendaal, M.: Tropospheric bromine chemistry: implications for present and pre-industrial ozone and mercury, *Atmos. Chem. Phys.*, 12, 6723-6740, <https://doi.org/10.5194/acp-12-6723-2012>, 2012.



- 2090 Passananti, M., Vinatier, V., Delort, A. M., Mailhot, G., and Brigante, M.: Siderophores in Cloud Waters and Potential Impact on Atmospheric Chemistry: Photoreactivity of Iron Complexes under Sun-Simulated Conditions, *Environ. Sci. Technol.*, 50, 9324-9332, <https://doi.org/10.1021/acs.est.6b02338>, 2016.
- Paulot, F., Crouse, J. D., Kjaergaard, H. G., Kurten, A., St Clair, J. M., Seinfeld, J. H., and Wennberg, P. O.: Unexpected epoxide formation in the gas-phase photooxidation of isoprene, *Science*, 325, 730-733, <https://doi.org/10.1126/science.1172910>, 2009.
- 2095 Paulot, F., Wunch, D., Crouse, J. D., Toon, G. C., Millet, D. B., DeCarlo, P. F., Vigouroux, C., Deutscher, N. M., González Abad, G., Notholt, J., Warneke, T., Hannigan, J. W., Warneke, C., de Gouw, J. A., Dunlea, E. J., De Mazière, M., Griffith, D. W. T., Bernath, P., Jimenez, J. L., and Wennberg, P. O.: Importance of secondary sources in the atmospheric budgets of formic and acetic acids, *Atmos. Chem. Phys.*, 11, 1989-2013, <https://doi.org/10.5194/acp-11-1989-2011>, 2011.
- Pechtl, S., Schmitz, G., and von Glasow, R.: Modelling iodide-iodate speciation in atmospheric aerosol: Contributions of inorganic and organic iodine chemistry, *Atmos. Chem. Phys.*, 7, 1381-1393, <https://doi.org/10.5194/acp-7-1381-2007>, 2007.
- 2100 Pechtl, S., and von Glasow, R.: Reactive chlorine in the marine boundary layer in the outflow of polluted continental air: A model study, *Geophys. Res. Lett.*, 34, L11813, <https://doi.org/10.1029/2007GL029761>, 2007.
- Perri, M. J., Lim, Y. B., Seitzinger, S. P., and Turpin, B. J.: Organosulfates from glycolaldehyde in aqueous aerosols and clouds: Laboratory studies, *Atmos. Environ.*, 44, 2658-2664, <https://doi.org/10.1016/j.atmosenv.2010.03.031>, 2010.
- Piletic, I. R., Edney, E. O., and Bartolotti, L. J.: A computational study of acid catalyzed aerosol reactions of atmospherically relevant epoxides, *Phys. Chem. Chem. Phys.*, 15, 18065-18076, <https://doi.org/10.1039/C3CP52851K>, 2013.
- 2105 Pitzer, K. S.: Activity Coefficients in Electrolyte Solutions, CRC Press, Boca Raton, 542 pp., 1991.
- Pocker, Y., Meany, J. E., Nist, B. J., and Zadorojny, C.: Reversible Hydration of Pyruvic Acid I. Equilibrium Studies, *J. Phys. Chem.*, 73, 2879-2882, <https://doi.org/10.1021/j100843a015>, 1969.
- 2110 Pope, C. A., and Dockery, D. W.: Health Effects of Fine Particulate Air Pollution: Lines that Connect, *J. Air Waste Manage.*, 56, 709-742, <https://doi.org/10.1080/10473289.2006.10464485>, 2012.
- Pöschl, U.: Atmospheric aerosols: Composition, transformation, climate and health effects, *Angew. Chem. Int. Edit.*, 44, 7520-7540, <https://doi.org/10.1002/anie.200501122>, 2005a.
- Pöschl, U.: Atmospheric aerosols: composition, transformation, climate and health effects, 44, 7520-7540, <https://doi.org/10.1002/anie.200501122>, 2005b.
- 2115 Powelson, M. H., Espelien, B. M., Hawkins, L. N., Galloway, M. M., and De Haan, D. O.: Brown carbon formation by aqueous-phase carbonyl compound reactions with amines and ammonium sulfate, *Environ. Sci. Technol.*, 48, 985-993, <https://doi.org/10.1021/es4038325>, 2014.
- Pye, H. O., Pinder, R. W., Piletic, I. R., Xie, Y., Capps, S. L., Lin, Y. H., Surratt, J. D., Zhang, Z., Gold, A., Luecken, D. J., Hutzell, W. T., Jaoui, M., Offenberg, J. H., Kleindienst, T. E., Lewandowski, M., and Edney, E. O.: Epoxide pathways improve model predictions of isoprene markers and reveal key role of acidity in aerosol formation, *Environ. Sci. Technol.*, 47, 11056-11064, <https://doi.org/10.1021/es402106h>, 2013.
- 2120 Pye, H. O. T., Nenes, A., Alexander, B., Ault, A. P., Barth, M. C., Clegg, S. L., Collett Jr, J. L., Fahey, K. M., Hennigan, C. J., Herrmann, H., Kanakidou, M., Kelly, J. T., Ku, I. T., McNeill, V. F., Riemer, N., Schaefer, T., Shi, G., Tilgner, A., Walker, J. T., Wang, T., Weber, R., Xing, J., Zaveri, R. A., and Zuend, A.: The acidity of atmospheric particles and clouds, *Atmos. Chem. Phys.*, 20, 4809-4888, <https://doi.org/10.5194/acp-20-4809-2020>, 2020.
- 2125 Radojevic, M.: SO₂ and NO_x Oxidation Mechanisms in the Atmosphere, in: Atmospheric Acidity: Sources, Consequences and Abatement, edited by: Radojevic, M., and Harrison, R. M., Environmental Management Series, Springer, Netherlands, 587, 1992.
- 2130 Raes, F., Van Dingenen, R., Vignati, E., Wilson, J., Putaud, J. P., Seinfeld, J. H., and Adams, P.: Formation and cycling of aerosols in the global troposphere, *Atmos. Environ.*, 34, 4215-4240, [https://doi.org/10.1016/S1352-2310\(00\)00239-9](https://doi.org/10.1016/S1352-2310(00)00239-9), 2000.
- Raja, S., Raghunathan, R., Yu, X. Y., Lee, T. Y., Chen, J., Kommalapati, R. R., Murugesan, K., Shen, X., Qingzhong, Y., Valsaraj, K. T., and Collett, J. L.: Fog chemistry in the Texas-Louisiana Gulf Coast corridor, *Atmos Environ* 42, 2048-2061, 2008.
- 2135 Raja, S., Raghunathan, R., Kommalapati, R. R., Shen, X. H., Collett, J. L., and Valsaraj, K. T.: Organic composition of fogwater in the Texas-Louisiana gulf coast corridor, *Atmos. Environ.*, 43, 4214-4222, <https://doi.org/10.1016/J.Atmosenv.2009.05.029>, 2009.



- Rao, P. S., Simic, M., and Hayon, E.: Pulse-Radiolysis Study of Imidazole and Histidine in Water, *J. Phys. Chem.*, 79, 1260-1263, <https://doi.org/10.1021/j100580a006>, 1975.
- 2140 Rao, X., and Collett, J. L., Jr.: Behavior of S(IV) and Formaldehyde in a Chemically Heterogeneous Cloud, *Environ. Sci. Technol.*, 29, 1023-1031, <https://doi.org/10.1021/es00004a024>, 1995.
- Rapf, R. J., Dooley, M. R., Kappes, K., Perkins, R. J., and Vaida, V.: pH Dependence of the Aqueous Photochemistry of alpha-Keto Acids, *J. Phys. Chem. A*, 121, 8368-8379, <https://doi.org/10.1021/acs.jpca.7b08192>, 2017a.
- Rapf, R. J., Perkins, R. J., Carpenter, B. K., and Vaida, V.: Mechanistic Description of Photochemical Oligomer Formation from Aqueous Pyruvic Acid, *J. Phys. Chem. A*, 121, 4272-4282, <https://doi.org/10.1021/acs.jpca.7b03310>, 2017b.
- 2145 Reed Harris, A. E., Ervens, B., Shoemaker, R. K., Kroll, J. A., Rapf, R. J., Griffith, E. C., Monod, A., and Vaida, V.: Photochemical kinetics of pyruvic acid in aqueous solution, *J. Phys. Chem. A*, 118, 8505-8516, <https://doi.org/10.1021/jp502186q>, 2014.
- Reilly, J. E., Rattigan, O. V., Moore, K. F., Judd, C., Eli Sherman, D., Dutkiewicz, V. A., Kreidenweis, S. M., Husain, L., and Collett, J. L.: Drop size-dependent S(IV) oxidation in chemically heterogeneous radiation fogs, *Atmos. Environ.*, 35, 5717-5728, [https://doi.org/10.1016/S1352-2310\(01\)00373-9](https://doi.org/10.1016/S1352-2310(01)00373-9), 2001.
- 2150 Riva, M., Chen, Y., Zhang, Y., Lei, Z., Olson, N. E., Boyer, H. C., Narayan, S., Yee, L. D., Green, H. S., Cui, T., Zhang, Z., Baumann, K., Fort, M., Edgerton, E., Budisulistiorini, S. H., Rose, C. A., Ribeiro, I. O., RL, E. O., Dos Santos, E. O., Machado, C. M. D., Szopa, S., Zhao, Y., Alves, E. G., de Sa, S. S., Hu, W., Knipping, E. M., Shaw, S. L., Duvoisin Junior, S., de Souza, R. A. F., Palm, B. B., Jimenez, J. L., Glasius, M., Goldstein, A. H., Pye, H. O. T., Gold, A., Turpin, B. J., Vizuete, W., Martin, S. T., Thornton, J. A., Dutcher, C. S., Ault, A. P., and Surratt, J. D.: Increasing Isoprene Epoxydiol-to-Inorganic Sulfate Aerosol Ratio Results in Extensive Conversion of Inorganic Sulfate to Organosulfur Forms: Implications for Aerosol Physicochemical Properties, *Environ. Sci. Technol.*, 53, 8682-8694, <https://doi.org/10.1021/acs.est.9b01019>, 2019.
- Rivlin, M., Eliav, U., and Navon, G.: NMR studies of the equilibria and reaction rates in aqueous solutions of formaldehyde, *J. Phys. Chem. B*, 119, 4479-4487, <https://doi.org/10.1021/jp513020y>, 2015.
- 2160 Roberts, J. M., Osthoff, H. D., Brown, S. S., Ravishankara, A. R., Coffman, D., Quinn, P., and Bates, T.: Laboratory studies of products of N₂O₅ uptake on Cl⁻ containing substrates, *Geophys. Res. Lett.*, 36, <https://doi.org/10.1029/2009GL040448>, 2009.
- Rodriguez, A. A., de Loera, A., Powelson, M. H., Galloway, M. M., and De Haan, D. O.: Formaldehyde and Acetaldehyde Increase Aqueous-Phase Production of Imidazoles in Methylglyoxal/Amine Mixtures: Quantifying a Secondary Organic Aerosol Formation Mechanism, *Environ. Sci. Technol. Lett.*, 4, 234-239, <https://doi.org/10.1021/acs.estlett.7b00129>, 2017.
- Author: NIST: Standard reference database 40: NDRL/NIST solution kinetics database vers. 3.0, 1998.
- Rudich, Y., Talukdar, R. K., and Ravishankara, A. R.: Multiphase chemistry of NO₃ in the remote troposphere, *J. Geophys. Res.-Atmos.*, 103, 16133-16143, <https://doi.org/10.1029/98jd01280>, 1998.
- 2170 Rusumdar, A. J., Wolke, R., Tilgner, A., and Herrmann, H.: Treatment of non-ideality in the SPACCIM multiphase model – Part 1: Model development, *Geosci. Model Dev.*, 9, 247-281, <https://doi.org/10.5194/gmd-9-247-2016>, 2016.
- Rusumdar, A. J., Tilgner, A., Wolke, R., and Herrmann, H.: Treatment of non-ideality in the SPACCIM multiphase model – Part 2: Impacts on the multiphase chemical processing in deliquesced aerosol particles, *Atmos. Chem. Phys.*, 20, 10351-10377, <https://doi.org/10.5194/acp-20-10351-2020>, 2020.
- 2175 Saiz-Lopez, A., Lamarque, J. F., Kinnison, D. E., Tilmes, S., Ordóñez, C., Orlando, J. J., Conley, A. J., Plane, J. M. C., Mahajan, A. S., Sousa Santos, G., Atlas, E. L., Blake, D. R., Sander, S. P., Schauffler, S., Thompson, A. M., and Brasseur, G.: Estimating the climate significance of halogen-driven ozone loss in the tropical marine troposphere, *Atmos. Chem. Phys.*, 12, 3939-3949, <https://doi.org/10.5194/acp-12-3939-2012>, 2012a.
- Saiz-Lopez, A., Plane, J. M., Baker, A. R., Carpenter, L. J., von Glasow, R., Martin, J. C., McFiggans, G., and Saunders, R. W.: Atmospheric chemistry of iodine, *Chem. Rev.*, 112, 1773-1804, <https://doi.org/10.1021/cr200029u>, 2012b.
- 2180 Saiz-Lopez, A., and von Glasow, R.: Reactive halogen chemistry in the troposphere, *Chem. Soc. Rev.*, 41, 6448-6472, <https://doi.org/10.1039/c2cs35208g>, 2012.
- Sareen, N., Schwier, A. N., Shapiro, E. L., Mitroo, D., and McNeill, V. F.: Secondary organic material formed by methylglyoxal in aqueous aerosol mimics, *Atmos. Chem. Phys.*, 10, 997-1016, <https://doi.org/10.5194/acp-10-997-2010>, 2010.
- 2185 Sarwar, G., Simon, H., Xing, J., and Mathur, R.: Importance of tropospheric ClNO₂ chemistry across the Northern Hemisphere, *Geophys. Res. Lett.*, 41, 4050-4058, <https://doi.org/10.1002/2014GL059962>, 2014.



- Saxena, P., and Seigneur, C.: On the oxidation of SO₂ to sulfate in atmospheric aerosols, *Atmos. Environ.* (1967), 21, 807-812, [https://doi.org/10.1016/0004-6981\(87\)90077-1](https://doi.org/10.1016/0004-6981(87)90077-1), 1987.
- Schindelka, J., Iinuma, Y., Hoffmann, D., and Herrmann, H.: Sulfate radical-initiated formation of isoprene-derived organosulfates in atmospheric aerosols, *Faraday Discuss.*, 165, 237-237, <https://doi.org/10.1039/c3fd00042g>, 2013.
- 2190 Schmidt, J. A., Jacob, D. J., Horowitz, H. M., Hu, L., Sherwen, T., Evans, M. J., Liang, Q., Suleiman, R. M., Oram, D. E., Le Breton, M., Percival, C. J., Wang, S., Dix, B., and Volkamer, R.: Modeling the observed tropospheric BrO background: Importance of multiphase chemistry and implications for ozone, OH, and mercury, *J. Geophys. Res.-Atmos.*, 121, 11819-11835, <https://doi.org/10.1002/2015jd024229>, 2016.
- 2195 Schöne, L., and Herrmann, H.: Kinetic measurements of the reactivity of hydrogen peroxide and ozone towards small atmospherically relevant aldehydes, ketones and organic acids in aqueous solutions, *Atmos. Chem. Phys.*, 14, 4503-4514, <https://doi.org/10.5194/acp-14-4503-2014>, 2014.
- Schuchmann, M. N., and von Sonntag, C.: The rapid hydration of the acetyl radical. A pulse radiolysis study of acetaldehyde in aqueous solution, *J. Am. Chem. Soc.*, 110, 5698-5701, <https://doi.org/10.1021/ja00225a019>, 1988.
- 2200 Schuchmann, M. N., Schuchmann, H.-P., and von Sonntag, C.: Oxidation of Hydroxymalonic Acid by OH Radicals in the Presence and in the Absence of Molecular Oxygen. A Pulse-Radiolysis and Product Study, *J. Phys. Chem.*, 99, 9122-9129, <https://doi.org/10.1021/j100022a026>, 1995.
- Schwartz, S.: Mass-transport considerations pertinent to aqueous phase reactions of gases in liquid-water clouds, in: *Chemistry of Multiphase Atmospheric Systems. NATO ASI Series (Series G: Ecological Sciences)*, vol 6., edited by: Jaeschke, W., Springer, Berlin, 425-471, https://doi.org/10.1007/978-3-642-70627-1_16, 1986.
- 2205 Schweitzer, F., Magi, L., Mirabel, P., and George, C.: Uptake Rate Measurements of Methanesulfonic Acid and Glyoxal by Aqueous Droplets, *J. Phys. Chem. A*, 102, 593-600, <https://doi.org/10.1021/jp972451k>, 1998.
- Sedehe, N., Takano, H., Blasic, V. A., Sullivan, K. A., and De Haan, D. O.: Temperature- and pH-dependent aqueous-phase kinetics of the reactions of glyoxal and methylglyoxal with atmospheric amines and ammonium sulfate, *Atmos. Environ.*, 77, 656-663, <https://doi.org/10.1016/j.atmosenv.2013.05.070>, 2013.
- 2210 Seigneur, C., and Saxena, P.: A theoretical investigation of sulfate formation in clouds, *Atmos. Environ.* (1967), 22, 101-115, [https://doi.org/10.1016/0004-6981\(88\)90303-4](https://doi.org/10.1016/0004-6981(88)90303-4), 1988.
- Seinfeld, J. H., and Pandis, S. N.: *Atmospheric chemistry and physics: from air pollution to climate change*, John Wiley & Sons, Inc., Hoboken, xxviii + 1203 pp. pp., 2006.
- Seinfeld, J. H.: Tropospheric chemistry and composition | Aerosols/Particles, in: *Encyclopedia of Atmospheric Sciences* (Second Edition), edited by: North, G. R., Pyle, J., and Zhang, F., Academic Press, Oxford, 182-187, <https://doi.org/10.1016/B978-0-12-382225-3.00438-2>, 2015.
- 2215 Seinfeld, J. H., Bretherton, C., Carslaw, K. S., Coe, H., DeMott, P. J., Dunlea, E. J., Feingold, G., Ghan, S., Guenther, A. B., Kahn, R., Kraucunas, I., Kreidenweis, S. M., Molina, M. J., Nenes, A., Penner, J. E., Prather, K. A., Ramanathan, V., Ramaswamy, V., Rasch, P. J., Ravishankara, A. R., Rosenfeld, D., Stephens, G., and Wood, R.: Improving our fundamental understanding of the role of aerosol-cloud interactions in the climate system, *Proc. Natl. Acad. Sci. USA*, 113, 5781, <https://doi.org/10.1073/pnas.1514043113>, 2016.
- 2220 Shao, J. Y., Chen, Q. J., Wang, Y. X., Lu, X., He, P. Z., Sun, Y. L., Shah, V., Martin, R. V., Philip, S., Song, S. J., Zhao, Y., Xie, Z. Q., Zhang, L., and Alexander, B.: Heterogeneous sulfate aerosol formation mechanisms during wintertime Chinese haze events: air quality model assessment using observations of sulfate oxygen isotopes in Beijing, *Atmos. Chem. Phys.*, 19, 6107-6123, <https://doi.org/10.5194/acp-19-6107-2019>, 2019.
- 2225 Shapiro, E. L., Szprengiel, J., Sareen, N., Jen, C. N., Giordano, M. R., and McNeill, V. F.: Light-absorbing secondary organic material formed by glyoxal in aqueous aerosol mimics, *Atmos. Chem. Phys.*, 9, 2289-2300, <https://doi.org/10.5194/acp-9-2289-2009>, 2009.
- 2230 Shen, X., Lee, T., Guo, J., Wang, X., Li, P., Xu, P., Wang, Y., Ren, Y., Wang, W., Wang, T., Li, Y., Carn, S. A., and Collett, J. L.: Aqueous phase sulfate production in clouds in eastern China, *Atmos. Environ.*, 62, 502-511, <https://doi.org/10.1016/j.atmosenv.2012.07.079>, 2012.
- 2235 Sherwen, T., Schmidt, J. A., Evans, M. J., Carpenter, L. J., Grossmann, K., Eastham, S. D., Jacob, D. J., Dix, B., Koenig, T. K., Sinreich, R., Ortega, I., Volkamer, R., Saiz-Lopez, A., Prados-Roman, C., Mahajan, A. S., and Ordóñez, C.: Global impacts of tropospheric halogens (Cl, Br, I) on oxidants and composition in GEOS-Chem, *Atmos. Chem. Phys.*, 16, 12239-12271, <https://doi.org/10.5194/acp-16-12239-2016>, 2016.



- Sherwen, T., Evans, M. J., Carpenter, L. J., Schmidt, J. A., and Mickley, L. J.: Halogen chemistry reduces tropospheric O₃ radiative forcing, *Atmos. Chem. Phys.*, 17, 1557-1569, <https://doi.org/10.5194/acp-17-1557-2017>, 2017.
- Shi, Z., Bonneville, S., Krom, M. D., Carslaw, K. S., Jickells, T. D., Baker, A. R., and Benning, L. G.: Iron dissolution kinetics of mineral dust at low pH during simulated atmospheric processing, *Atmos. Chem. Phys.*, 11, 995-1007, <https://doi.org/10.5194/acp-11-995-2011>, 2011.
- 2240 Sievering, H., Boatman, J., Gorman, E., Kim, Y., Anderson, L., Ennis, G., Luria, M., and Pandis, S.: Removal of Sulfur from the Marine Boundary-Layer by Ozone Oxidation in Sea-Salt Aerosols, *Nature*, 360, 571-573, <https://doi.org/10.1038/360571a0>, 1992.
- 2245 Simpson, W., Law, K., Schmale, J., Pratt, K., Arnold, S., Mao, J., Alexander, B., Anenberg, S., Baklanov, A., Bell, D., Brown, S., Creamean, J., de Boer, G., DeCarlo, P., Descari, S., Elleman, R., Flynn, J., Fochesatto, J., Ganzenfeld, L., Gilmour, I., Griffin, R., Järvi, L., Kaspari, S., Konstantinov, P., Murphy, J., Petäjä, T., Pye, H., Raut, J.-C., Roberts, T., Shiraiwa, M., Stutz, J., Thomas, J., Thornton, J., Wagstrom, K., Weber, R., Webley, P., and Williams, B.: Alaskan Layered Pollution And Chemical Analysis (ALPACA) White Paper, 84, <https://alpaca.community.uaf.edu/files/2018/2011/ALPACA-whitepaper-2030Nov2018.pdf>, 2019.
- 2250 Simpson, W. R., Brown, S. S., Saiz-Lopez, A., Thornton, J. A., and Glasow, R.: Tropospheric halogen chemistry: sources, cycling, and impacts, *Chem. Rev.*, 115, 4035-4062, <https://doi.org/10.1021/cr5006638>, 2015.
- Sipila, M., Berndt, T., Petaja, T., Brus, D., Vanhanen, J., Stratmann, F., Patokoski, J., Mauldin, R. L., 3rd, Hyvarinen, A. P., Lihavainen, H., and Kulmala, M.: The role of sulfuric acid in atmospheric nucleation, *Science*, 327, 1243-1246, <https://doi.org/10.1126/science.1180315>, 2010.
- 2255 Six, K. D., Kloster, S., Ilyina, T., Archer, S. D., Zhang, K., and Maier-Reimer, E.: Global warming amplified by reduced sulphur fluxes as a result of ocean acidification, *Nature Clim. Change*, 3, 975-978, <https://doi.org/10.1038/nclimate1981>, 2013.
- Smith, S. J., Pitcher, H., and Wigley, T. M. L.: Global and regional anthropogenic sulfur dioxide emissions, *Global and Planetary Change*, 29, 99-119, [https://doi.org/10.1016/S0921-8181\(00\)00057-6](https://doi.org/10.1016/S0921-8181(00)00057-6), 2001.
- 2260 Smith, S. J., van Aardenne, J., Klimont, Z., Andres, R. J., Volke, A., and Delgado Arias, S.: Anthropogenic sulfur dioxide emissions: 1850–2005, *Atmos. Chem. Phys.*, 11, 1101-1116, <https://doi.org/10.5194/acp-11-1101-2011>, 2011.
- Song, S., Nenes, A., Gao, M., Zhang, Y., Liu, P., Shao, J., Ye, D., Xu, W., Lei, L., Sun, Y., Liu, B., Wang, S., and McElroy, M. B.: Thermodynamic Modeling Suggests Declines in Water Uptake and Acidity of Inorganic Aerosols in Beijing Winter Haze Events during 2014/2015–2018/2019, *Environ. Sci. Tech. Lett.*, 752-760, <https://doi.org/10.1021/acs.estlett.9b00621>, 2019a.
- 2265 Song, S. J., Gao, M., Xu, W. Q., Sun, Y. L., Worsnop, D. R., Jayne, J. T., Zhang, Y. Z., Zhu, L., Li, M., Zhou, Z., Cheng, C. L., Lv, Y. B., Wang, Y., Peng, W., Xu, X. B., Lin, N., Wang, Y. X., Wang, S. X., Munger, J. W., Jacob, D. J., and McElroy, M. B.: Possible heterogeneous chemistry of hydroxymethanesulfonate (HMS) in northern China winter haze, *Atmos. Chem. Phys.*, 19, 1357-1371, <https://doi.org/10.5194/acp-19-1357-2019>, 2019b.
- Spataro, F., and Ianniello, A.: Sources of atmospheric nitrous acid: State of the science, current research needs, and future prospects, *J. Air Waste Manag. Assoc.*, 64, 1232-1250, <https://doi.org/10.1080/10962247.2014.952846>, 2014.
- 2270 Spindler, G., Hesper, J., Brüggemann, E., Dubois, R., Müller, T., and Herrmann, H.: Wet annular denuder measurements of nitrous acid: laboratory study of the artefact reaction of NO₂ with S(IV) in aqueous solution and comparison with field measurements, *Atmos. Environ.*, 37, 2643-2662, [https://doi.org/10.1016/S1352-2310\(03\)00209-7](https://doi.org/10.1016/S1352-2310(03)00209-7), 2003.
- 2275 Stockdale, A., Krom, M. D., Mortimer, R. J., Benning, L. G., Carslaw, K. S., Herbert, R. J., Shi, Z., Myriokefalitakis, S., Kanakidou, M., and Nenes, A.: Understanding the nature of atmospheric acid processing of mineral dusts in supplying bioavailable phosphorus to the oceans, *Proc. Natl. Acad. Sci. USA*, 113, 14639-14644, <https://doi.org/10.1073/pnas.1608136113>, 2016.
- Strollo, C. M., and Ziemann, P. J.: Products and mechanism of secondary organic aerosol formation from the reaction of 3-methylfuran with OH radicals in the presence of NO_x, *Atmos. Environ.*, 77, 534-543, <https://doi.org/10.1016/J.Atmosenv.2013.05.033>, 2013.
- 2280 Stropoli, S. J., and Elrod, M. J.: Assessing the Potential for the Reactions of Epoxides with Amines on Secondary Organic Aerosol Particles, *J. Phys. Chem. A*, 119, 10181-10189, <https://doi.org/10.1021/acs.jpca.5b07852>, 2015.
- Sumner, A. J., Woo, J. L., and McNeill, V. F.: Model analysis of secondary organic aerosol formation by glyoxal in laboratory studies: the case for photoenhanced chemistry, *Environ. Sci. Technol.*, 48, 11919-11925, <https://doi.org/10.1021/es502020j>, 2014.
- 2285



- Surratt, J. D., Murphy, S. M., Kroll, J. H., Ng, N. L., Hildebrandt, L., Sorooshian, A., Szmigielski, R., Vermeylen, R., Maenhaut, W., Claeys, M., Flagan, R. C., and Seinfeld, J. H.: Chemical Composition of Secondary Organic Aerosol Formed from the Photooxidation of Isoprene, *J. Phys. Chem. A*, 110, 9665-9690, <https://doi.org/10.1021/jp061734m>, 2006a.
- 2290 Surratt, J. D., Murphy, S. M., Kroll, J. H., Ng, N. L., Hildebrandt, L., Sorooshian, A., Szmigielski, R., Vermeylen, R., Maenhaut, W., Claeys, M., Flagan, R. C., and Seinfeld, J. H.: Chemical composition of secondary organic aerosol formed from the photooxidation of isoprene, *J. Phys. Chem., A*, 110, 9665-9690, <https://doi.org/10.1021/Jp061734m>, 2006b.
- Surratt, J. D., Kroll, J. H., Kleindienst, T. E., Edney, E. O., Claeys, M., Sorooshian, A., Ng, N. L., Offenberg, J. H., Lewandowski, M., Jaoui, M., Flagan, R. C., and Seinfeld, J. H.: Evidence for Organosulfates in Secondary Organic Aerosol, *Environ. Sci. Technol.*, 41, 517-527, <https://doi.org/10.1021/es062081q>, 2007a.
- 2295 Surratt, J. D., Lewandowski, M., Offenberg, J. H., Jaoui, M., Kleindienst, T. E., Edney, E. O., and Seinfeld, J. H.: Effect of acidity on secondary organic aerosol formation from isoprene, *Environ. Sci. Technol.*, 41, 5363-5369, <https://doi.org/10.1021/es0704176>, 2007b.
- Surratt, J. D., Chan, A. W., Eddingsaas, N. C., Chan, M., Loza, C. L., Kwan, A. J., Hersey, S. P., Flagan, R. C., Wennberg, P. O., and Seinfeld, J. H.: Reactive intermediates revealed in secondary organic aerosol formation from isoprene, *Proc. Natl. Acad. Sci. USA*, 107, 6640-6645, <https://doi.org/10.1073/pnas.091114107>, 2010.
- 2300 Szmigielski, R., Surratt, J. D., Vermeylen, R., Szmigielska, K., Kroll, J. H., Ng, N. L., Murphy, S. M., Sorooshian, A., Seinfeld, J. H., and Claeys, M.: Characterization of 2-methylglyceric acid oligomers in secondary organic aerosol formed from the photooxidation of isoprene using trimethylsilylation and gas chromatography/ion trap mass spectrometry, *J. Mass Spectrom.*, 42, 101-116, <https://doi.org/10.1002/Jms.1146>, 2007.
- 2305 Tanner, P. A., and Law, P. T.: Organic acids in the atmosphere and bulk deposition of Hong Kong, *Water Air Soil Poll.*, 142, 279-297, <https://doi.org/10.1023/A:1022063925972>, 2003.
- Teich, M., van Pinxteren, D., and Herrmann, H.: A one year study of functionalised medium-chain carboxylic acids in atmospheric particles at a rural site in Germany revealing seasonal trends and possible sources, *J. Atmos. Chem.*, 76, 115-132, <https://doi.org/10.1007/s10874-019-09390-5>, 2019.
- 2310 Tilgner, A., and Herrmann, H.: Radical-driven carbonyl-to-acid conversion and acid degradation in tropospheric aqueous systems studied by CAPRAM, *Atmos. Environ.*, 44, 5415-5422, <https://doi.org/10.1016/j.atmosenv.2010.07.050>, 2010.
- Tobias, H. J., and Ziemann, P. J.: Thermal Desorption Mass Spectrometric Analysis of Organic Aerosol Formed from Reactions of 1-Tetradecene and O₃ in the Presence of Alcohols and Carboxylic Acids, *Environ. Sci. Technol.*, 34, 2105-2115, <https://doi.org/10.1021/es9907156>, 2000.
- 2315 Tolocka, M. P., Jang, M., Ginter, J. M., Cox, F. J., Kamens, R. M., and Johnston, M. V.: Formation of Oligomers in Secondary Organic Aerosol, *Environ. Sci. Technol.*, 38, 1428-1434, <https://doi.org/10.1021/es035030r>, 2004.
- Troy, R. C., and Margerum, D. W.: Non-metal redox kinetics: hypobromite and hypobromous acid reactions with iodide and with sulfite and the hydrolysis of bromosulfate, *Inorg. Chem.*, 30, 3538-3543, <https://doi.org/10.1021/ic00018a028>, 1991.
- Tsui, W. G., Woo, J. L., and McNeill, V. F.: Impact of Aerosol-Cloud Cycling on Aqueous Secondary Organic Aerosol Formation, *Atmosphere*, 10, 666-678, <https://doi.org/10.3390/atmos10110666>, 2019.
- 2320 Tuguldurova, V. P., Fateev, A. V., Poleshchuk, O. K., and Vodyankina, O. V.: Theoretical analysis of glyoxal condensation with ammonia in aqueous solution, *Phys. Chem. Chem. Phys.*, 21, 9326-9334, <https://doi.org/10.1039/c8cp07270a>, 2019.
- Turnock, S. T., Mann, G. W., Woodhouse, M. T., Dalvi, M., O'Connor, F. M., Carslaw, K. S., and Spracklen, D. V.: The Impact of Changes in Cloud Water pH on Aerosol Radiative Forcing, *Geophys. Res. Lett.*, 46, 4039-4048, <https://doi.org/10.1029/2019gl082067>, 2019.
- 2325 U.S.EPA: Air Quality Research Subcommittee. Atmospheric Ammonia: Sources and Fate. A Review of Ongoing Federal Research and Future Needs, U.S.EPA, Washington D.C., <http://www.esrl.noaa.gov/csd/AQRS/reports/ammonia.pdf>, 2000.
- Urbansky, E. T., and Schock, M. R.: Understanding, Deriving, and Computing Buffer Capacity, *J. Chem. Edu.*, 77, 1640, <https://doi.org/10.1021/ed077p1640>, 2000.
- 2330 van Pinxteren, D., and Herrmann, H.: Determination of functionalised carboxylic acids in atmospheric particles and cloud water using capillary electrophoresis/mass spectrometry, *J. Chromatogr. A*, 1171, 112-123, <https://doi.org/10.1016/j.chroma.2007.09.021>, 2007.
- Van Wyngarden, A. L., Perez-Montano, S., Bui, J. V., Li, E. S., Nelson, T. E., Ha, K. T., Leong, L., and Iraci, L. T.: Complex chemical composition of colored surface films formed from reactions of propanal in sulfuric acid at upper troposphere/lower stratosphere aerosol acidities, *Atmos. Chem. Phys.*, 15, 4225-4239, <https://doi.org/10.5194/acp-15-4225-2015>, 2015.
- 2335



- Vasilakos, P., Russell, A., Weber, R., and Nenes, A.: Understanding nitrate formation in a world with less sulfate, *Atmos. Chem. Phys.*, 18, 12765-12775, <https://doi.org/10.5194/acp-18-12765-2018>, 2018.
- 2340 Vet, R., Artz, R. S., Carou, S., Shaw, M., Ro, C.-U., Aas, W., Baker, A., Bowersox, V. C., Dentener, F., Galy-Lacaux, C., Hou, A., Pienaar, J. J., Gillett, R., Forti, M. C., Gromov, S., Hara, H., Khodzher, T., Mahowald, N. M., Nickovic, S., Rao, P. S. P., and Reid, N. W.: A global assessment of precipitation chemistry and deposition of sulfur, nitrogen, sea salt, base cations, organic acids, acidity and pH, and phosphorus, *Atmos. Environ.*, 93, 3-100, <https://doi.org/10.1016/j.atmosenv.2013.10.060>, 2014.
- Vogt, R., Crutzen, P. J., and Sander, R.: A mechanism for halogen release from sea-salt aerosol in the remote marine boundary layer, *Nature*, 383, 327-330, <https://doi.org/10.1038/383327a0>, 1996.
- 2345 von Glasow, R., Sander, R., Bott, A., and Crutzen, P. J.: Modeling halogen chemistry in the marine boundary layer 2. Interactions with sulfur and the cloud-covered MBL, *J. Geophys. Res.-Atmos.*, 107, ACH 2-1-ACH 2-12, <https://doi.org/10.1029/2001jd000943>, 2002a.
- von Glasow, R., Sander, R., Bott, A., and Crutzen, P. J.: Modeling halogen chemistry in the marine boundary layer 1. Cloud-free MBL, *J. Geophys. Res.-Atmos.*, 107, ACH 9-1-ACH 9-16, <https://doi.org/10.1029/2001jd000942>, 2002b.
- 2350 von Schneidmesser, E., Monks, P. S., Allan, J. D., Bruhwiler, L., Forster, P., Fowler, D., Lauer, A., Morgan, W. T., Paasonen, P., Righi, M., Sindelarova, K., and Sutton, M. A.: Chemistry and the Linkages between Air Quality and Climate Change, *Chem. Rev.*, 115, 3856-3897, <https://doi.org/10.1021/acs.chemrev.5b00089>, 2015.
- von Sonntag, C., and von Gunten, U.: *Chemistry of Ozone in Water and Wastewater Treatment: From Basic Principles to Applications*, IWA Publishing, 2012.
- 2355 Wang, G., Zhang, R., Gomez, M. E., Yang, L., Levy Zamora, M., Hu, M., Lin, Y., Peng, J., Guo, S., Meng, J., Li, J., Cheng, C., Hu, T., Ren, Y., Wang, Y., Gao, J., Cao, J., An, Z., Zhou, W., Li, G., Wang, J., Tian, P., Marrero-Ortiz, W., Secrest, J., Du, Z., Zheng, J., Shang, D., Zeng, L., Shao, M., Wang, W., Huang, Y., Wang, Y., Zhu, Y., Li, Y., Hu, J., Pan, B., Cai, L., Cheng, Y., Ji, Y., Zhang, F., Rosenfeld, D., Liss, P. S., Duce, R. A., Kolb, C. E., and Molina, M. J.: Persistent sulfate formation from London Fog to Chinese haze, *Proc. Natl. Acad. Sci. USA*, 113, 13630-13635, <https://doi.org/10.1073/pnas.1616540113>, 2016.
- 2360 Wang, S., Zhou, S., Tao, Y., Tsui, W. G., Ye, J., Yu, J. Z., Murphy, J. G., McNeill, V. F., Abbatt, J. P. D., and Chan, A. W. H.: Organic Peroxides and Sulfur Dioxide in Aerosol: Source of Particulate Sulfate, *Environ. Sci. Technol.*, 53, 10695-10704, <https://doi.org/10.1021/acs.est.9b02591>, 2019a.
- Wang, X., Jacob, D. J., Eastham, S. D., Sulprizio, M. P., Zhu, L., Chen, Q., Alexander, B., Sherwen, T., Evans, M. J., Lee, B. H., Haskins, J. D., Lopez-Hilfiker, F. D., Thornton, J. A., Huey, G. L., and Liao, H.: The role of chlorine in global tropospheric chemistry, *Atmos. Chem. Phys.*, 19, 3981-4003, <https://doi.org/10.5194/acp-19-3981-2019>, 2019b.
- 2365 Wang, X., Gemayel, R., Hayeck, N., Perrier, S., Charbonnel, N., Xu, C., Chen, H., Zhu, C., Zhang, L., Wang, L., Nizkorodov, S. A., Wang, X., Wang, Z., Wang, T., Mellouki, A., Riva, M., Chen, J., and George, C.: Atmospheric Photosensitization: A New Pathway for Sulfate Formation, *Environ. Sci. Technol.*, 54, 3114-3120, <https://doi.org/10.1021/acs.est.9b06347>, 2020.
- 2370 Wang, Y. X., Zhang, Q. Q., Jiang, J. K., Zhou, W., Wang, B. Y., He, K. B., Duan, F. K., Zhang, Q., Philip, S., and Xie, Y. Y.: Enhanced sulfate formation during China's severe winter haze episode in January 2013 missing from current models, *J. Geophys. Res.-Atmos.*, 119, 10425-10440, <https://doi.org/10.1002/2013jd021426>, 2014.
- Wardman, P.: Reduction Potentials of One-Electron Couples Involving Free Radicals in Aqueous Solution, *J. Phys. Chem. Ref. Data*, 18, 1637-1755, <https://doi.org/10.1063/1.555843>, 1989.
- 2375 Warneck, P.: The relative importance of various pathways for the oxidation of sulfur dioxide and nitrogen dioxide in sunlit continental fair weather clouds, *Phys. Chem. Chem. Phys.*, 1, 5471-5483, <https://doi.org/10.1039/A906558J>, 1999.
- Waxman, E. M., Elm, J., Kurtén, T., Mikkelsen, K. V., Ziemann, P. J., and Volkamer, R.: Glyoxal and Methylglyoxal Setschenow Salting Constants in Sulfate, Nitrate, and Chloride Solutions: Measurements and Gibbs Energies, *Environ. Sci. Technol.*, 49, 11500-11508, <https://doi.org/10.1021/acs.est.5b02782>, 2015.
- 2380 Weber, R. J., Guo, H., Russell, A. G., and Nenes, A.: High aerosol acidity despite declining atmospheric sulfate concentrations over the past 15 years, *Nat. Geosci.*, 9, 282-285, <https://doi.org/10.1038/ngeo2665>, 2016.
- Whiteaker, J. R., and Prather, K. A.: Hydroxymethanesulfonate as a tracer for fog processing of individual aerosol particles, *Atmos. Environ.*, 37, 1033-1043, [https://doi.org/10.1016/S1352-2310\(02\)01029-4](https://doi.org/10.1016/S1352-2310(02)01029-4), 2003.
- 2385 Wu, L. Y., Tong, S. R., Wang, W. G., and Ge, M. F.: Effects of temperature on the heterogeneous oxidation of sulfur dioxide by ozone on calcium carbonate, *Atmos. Chem. Phys.*, 11, 6593-6605, <https://doi.org/10.5194/acp-11-6593-2011>, 2011.



- Xia, X., and Hopke, P. K.: Seasonal variation of 2-methyltetrols in ambient air samples, *Environ. Sci. Technol.*, 40, 6934-6937, <https://doi.org/10.1021/es060988i>, 2006.
- Xue, J., Yuan, Z., Griffith, S. M., Yu, X., Lau, A. K., and Yu, J. Z.: Sulfate Formation Enhanced by a Cocktail of High NO_x, SO₂, Particulate Matter, and Droplet pH during Haze-Fog Events in Megacities in China: An Observation-Based Modeling Investigation, *Environ. Sci. Technol.*, 50, 7325-7334, <https://doi.org/10.1021/acs.est.6b00768>, 2016.
- 2390 Xue, J., Yu, X., Yuan, Z., Griffith, S. M., Lau, A. K. H., Seinfeld, J. H., and Yu, J. Z.: Efficient control of atmospheric sulfate production based on three formation regimes, *Nat. Geosci.*, 12, 977-982, <https://doi.org/10.1038/s41561-019-0485-5>, 2019.
- Yasmeen, F., Sauret, N., Gal, J. F., Maria, P. C., Massi, L., Maenhaut, W., and Claeys, M.: Characterization of oligomers from methylglyoxal under dark conditions: a pathway to produce secondary organic aerosol through cloud processing during
2395 nighttime, *Atmos. Chem. Phys.*, 10, 3803-3812, <https://doi.org/10.5194/acp-10-3803-2010>, 2010.
- Ye, C., Liu, P., Ma, Z., Xue, C., Zhang, C., Zhang, Y., Liu, J., Liu, C., Sun, X., and Mu, Y.: High H₂O₂ concentrations observed during haze periods during the winter in Beijing: Importance of H₂O₂ oxidation in sulfate formation, *Environ. Sci. Technol. Lett.*, 5, 757-763, <https://doi.org/10.1021/acs.estlett.8b00579>, 2018a.
- Ye, J. H., Abbatt, J. P. D., and Chan, A. W. H.: Novel pathway of SO₂ oxidation in the atmosphere: reactions with
2400 monoterpene ozonolysis intermediates and secondary organic aerosol, *Atmos. Chem. Phys.*, 18, 5549-5565, <https://doi.org/10.5194/acp-18-5549-2018>, 2018b.
- Yi, Y., Cao, Z., Zhou, X., Xue, L., and Wang, W.: Formation of aqueous-phase secondary organic aerosols from glycolaldehyde and ammonium sulfate/amines: A kinetic and mechanistic study, *Atmos. Environ.*, 181, 117-125, <https://doi.org/10.1016/j.atmosenv.2018.03.021>, 2018.
- 2405 Yu, G., Bayer, A. R., Galloway, M. M., Korshavn, K. J., Fry, C. G., and Keutsch, F. N.: Glyoxal in aqueous ammonium sulfate solutions: products, kinetics and hydration effects, *Environ. Sci. Technol.*, 45, 6336-6342, <https://doi.org/10.1021/es200989n>, 2011.
- Yu, Z., Jang, M., and Park, J.: Modeling atmospheric mineral aerosol chemistry to predict heterogeneous photooxidation of SO₂, *Atmos. Chem. Phys.*, 17, 10001-10017, <https://doi.org/10.5194/acp-17-10001-2017>, 2017.
- 2410 Zaveri, R. A.: A new method for multicomponent activity coefficients of electrolytes in aqueous atmospheric aerosols, *J. Geophys. Res.-Atmos.*, 110, D02201, <https://doi.org/10.1029/2004jd004681>, 2005.
- Zavitsas, A. A., Coffiner, M., Wiseman, T., and Zavitsas, L. R.: Reversible Hydration of Formaldehyde - Thermodynamic Parameters, *J. Phys. Chem.*, 74, 2746-2750, <https://doi.org/10.1021/j100708a003>, 1970.
- Zhang, H., Surratt, J. D., Lin, Y. H., Bapat, J., and Kamens, R. M.: Effect of relative humidity on SOA formation from
2415 isoprene/NO photooxidation: enhancement of 2-methylglyceric acid and its corresponding oligoesters under dry conditions, *Atmos. Chem. Phys.*, 11, 6411-6424, <https://doi.org/10.5194/acp-11-6411-2011>, 2011.
- Zhang, R., Wang, G., Guo, S., Zamora, M. L., Ying, Q., Lin, Y., Wang, W., Hu, M., and Wang, Y.: Formation of urban fine particulate matter, *Chem. Rev.*, 115, 3803-3855, <https://doi.org/10.1021/acs.chemrev.5b00067>, 2015.
- Zhang, Y., and Carmichael, G. R.: The Role of Mineral Aerosol in Tropospheric Chemistry in East Asia—A Model Study, *J. Appl. Meteor.*, 38, 353-366, [https://doi.org/10.1175/1520-0450\(1999\)038%3C0353:TROMAI%3E2.0.CO;2](https://doi.org/10.1175/1520-0450(1999)038%3C0353:TROMAI%3E2.0.CO;2), 1999.
- 2420 Zhang, Y., Chen, Y., Lambe, A. T., Olson, N. E., Lei, Z., Craig, R. L., Zhang, Z., Gold, A., Onasch, T. B., Jayne, J. T., Worsnop, D. R., Gaston, C. J., Thornton, J. A., Vizuete, W., Ault, A. P., and Surratt, J. D.: Effect of the Aerosol-Phase State on Secondary Organic Aerosol Formation from the Reactive Uptake of Isoprene-Derived Epoxydiols (IEPOX), *Environ. Sci. Technol. Lett.*, 5, 167-174, <https://doi.org/10.1021/acs.estlett.8b00044>, 2018a.
- 2425 Zhang, Y., Tong, S., Ge, M., Jing, B., Hou, S., Tan, F., Chen, Y., Guo, Y., and Wu, L.: The formation and growth of calcium sulfate crystals through oxidation of SO₂ by O₃ on size-resolved calcium carbonate, *RSC Adv.*, 8, 16285-16293, <https://doi.org/10.1039/c8ra02050g>, 2018b.
- Zhang, Z.-S., Engling, G., Chan, C.-Y., Yang, Y.-H., Lin, M., Shi, S., He, J., Li, Y.-D., and Wang, X.-M.: Determination of isoprene-derived secondary organic aerosol tracers (2-methyltetrols) by HPAEC-PAD: Results from size-resolved aerosols in
2430 a tropical rainforest, *Atmos. Environ.*, 70, 468-476, <https://doi.org/10.1016/j.atmosenv.2013.01.020>, 2013.
- Zhao, J., Levitt, N. P., Zhang, R., and Chen, J.: Heterogeneous Reactions of Methylglyoxal in Acidic Media: Implications for Secondary Organic Aerosol Formation, *Environ. Sci. Technol.*, 40, 7682-7687, <https://doi.org/10.1021/es060610k>, 2006.
- Zhao, R., Lee, A. K. Y., Wang, C., Wania, F., Wong, J. P. S., Zhou, S., and Abbatt, J. P. D.: The Role of Water in Organic Aerosol Multiphase Chemistry: Focus on Partitioning and Reactivity, in: *Advances in Atmospheric Chemistry*, edited by:



- 2435 Barker, J. R., Steiner, A. L., and Wallington, T. J., *Advances in Atmospheric Chemistry, Volume 1*, World Scientific, Singapore, 95-184, https://doi.org/10.1142/9789813147355_0002, 2016.
Zhao, Z., Tolentino, R., Lee, J., Vuong, A., Yang, X., and Zhang, H.: Interfacial Dimerization by Organic Radical Reactions during Heterogeneous Oxidative Aging of Oxygenated Organic Aerosols, *J. Phys. Chem. A*, 123, 10782-10792, <https://doi.org/10.1021/acs.jpca.9b10779>, 2019.
- 2440 Zheng, G., Su, H., Wang, S., Andreae, M. O., Poschl, U., and Cheng, Y.: Multiphase buffer theory explains contrasts in atmospheric aerosol acidity, *Science*, 369, 1374-1377, <https://doi.org/10.1126/science.aba3719>, 2020.
Zhu, L., Jacob, D. J., Eastham, S. D., Sulprizio, M. P., Wang, X., Sherwen, T., Evans, M. J., Chen, Q., Alexander, B., Koenig, T. K., Volkamer, R., Huey, L. G., Le Breton, M., Bannan, T. J., and Percival, C. J.: Effect of sea salt aerosol on tropospheric bromine chemistry, *Atmos. Chem. Phys.*, 19, 6497-6507, <https://doi.org/10.5194/acp-19-6497-2019>, 2019.
- 2445 Zuend, A., Marcolli, C., Luo, B. P., and Peter, T.: A thermodynamic model of mixed organic-inorganic aerosols to predict activity coefficients, *Atmos. Chem. Phys.*, 8, 4559-4593, <https://doi.org/10.5194/acp-8-4559-2008>, 2008.



2450 **Table 1: Composition conditions applied for the calculation of the S(IV) oxidation rates of different reaction pathways for urban haze and rural aerosol conditions as well as urban and rural cloud conditions (bottom) at 298 K.**

Reactant	Concentration			
	Urban haze	Rural aerosol	Urban cloud	Rural cloud
SO ₂ / ppb	40.0 [#]	1.0 [†]	5.0 [∞]	1.0 [†]
H ₂ O ₂ / ppb	0.1 [‡]	0.1 [†]	1.0 [∞]	0.1 [†]
O ₃ / ppb	1.0 [#]	30.0 [†]	50.0 [∞]	30.0 [†]
NO ₂ / ppb	66.0 [#]	1.0 [†]	10.0 [†]	1.0 [†]
HNO ₄ / ppb	0.01 [†]	0.001 [†]	0.01 [†]	0.001 [†]
CH ₃ OOH / ppb	0.1 [†]	0.1 [†]	0.1 [†]	0.1 [†]
CH ₃ C(O)OOH / ppb	0.1 [†]	0.1 [†]	0.1 [†]	0.1 [†]
Fe(III) / mol L ⁻¹	1.1×10 ⁻³ [#]	1.0×10 ⁻³ [∞]	1.0×10 ⁻⁵ [§]	1.0×10 ⁻⁶ [§]
Mn(II) / mol L ⁻¹	2.55×10 ⁻³ [#]	1.0×10 ⁻⁴ [∞]	1.0×10 ⁻⁶ [§]	1.0×10 ⁻⁷ [§]
PS* / mol L ⁻¹	1.9×10 ⁻¹¹	6.3×10 ⁻¹¹	1.2×10 ⁻¹²	3.6×10 ⁻¹³
Ionic strength mol L ⁻¹	1.0 [†]	1.0 [†]	1.0×10 ^{-4†}	1.0×10 ^{-4†}

Remarks: [∞] based on Seinfeld and Pandis (2006), [#] based on Cheng et al. (2016), [§] estimated from data given in Deguillaume et al. (2005), [‡] based on (Ye et al., 2018a), [†] estimated, daytime particle/cloud mean concentrations based on simulations using CAPRAM mechanism (Bräuer et al., 2019; Hoffmann et al., 2020). Further simulation details are given in the SI.



Table 2: Influence of acidity on the hydration rate of formaldehyde and acetaldehyde in buffered solutions.

Acidic species	Acid dissociation constant of the catalyst acid-base pair K_a / (unitless)	Acid catalytic constant k_a / $L\ mol^{-1}\ s^{-1}$	Base catalytic constant k_b / $L\ mol^{-1}\ s^{-1}$
<i>Formaldehyde*</i>			
H_3O^+	55.5	2.7	0.0051
Formic acid	1.77×10^{-4}	0.070	0.013
Phenylacetic	4.88×10^{-5}	-	0.015
Acetic acid	1.75×10^{-5}	0.043	0.022
Trimethylacetic acid	8.9×10^{-6}	0.025	0.022
Water	1.8×10^{-16}	0.0051	1600
<i>Acetaldehyde#</i>			
H_3O^+	55.5	930	0.00014
Formic acid	1.77×10^{-4}	1.74	0.065
Phenylacetic	4.88×10^{-5}	0.91	0.054
Acetic acid	1.75×10^{-5}	0.47	0.157
Trimethylacetic acid	9.4×10^{-6}	0.33	0.161
Water	1.8×10^{-16}	0.00014	8×10^4
*Bell and Evans (1966), #Kurz (1967); Kurz and Coburn (1967); Ogata and Kawasaki (1970)			