



The impact of large-scale circulation on daily fine particulate matter (PM_{2.5}) over major populated regions of China in winter

Zixuan Jia¹, Ruth M. Doherty¹, Carlos Ordóñez², Chaofan Li^{3,6}, Oliver Wild⁴, Shipra Jain¹, Xiao Tang⁵

¹School of GeoSciences, University of Edinburgh, Edinburgh, UK

5 ²Departamento de Física de la Tierra y Astrofísica, Facultad de Ciencias Físicas, Universidad Complutense de Madrid, Madrid, Spain

³Center for Monsoon System Research, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China

⁴Lancaster Environment Centre, Lancaster University, Lancaster, UK

⁵LAPC, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China

10 ⁶College of Earth and Planetary Sciences, University of the Chinese Academy of Sciences, Beijing, China

Correspondence to: Zixuan Jia (Z.Jia-6@sms.ed.ac.uk)

Abstract. The influence of large-scale circulation on daily PM_{2.5} variability through its direct effect on key regional meteorological variables is examined over three major populated regions of China: Beijing–Tianjin–Hebei (BTH), the Yangtze River Delta (YRD), and the Pearl River Delta (PRD). In BTH, a shallow East Asian trough curbs northerly cold and dry air from the Siberian High, enhancing PM_{2.5} pollution levels. Weak southerly winds in eastern and southern China, associated with a weakened Siberian High, suppress horizontal dispersion, contributing to air pollution accumulation over YRD. In PRD, weak southerly winds and precipitation deficits over southern China are conducive to high PM_{2.5} pollution levels. To account for these dominant large-scale circulation – PM_{2.5} relationships, we propose three new circulation-based indices for predicting different levels of air pollution based on regional PM_{2.5} concentrations in each region: a 500 hPa geopotential height-based index for BTH, a sea level pressure-based index for YRD and an 850 hPa meridional wind-based index for PRD. These three indices can effectively distinguish clean days from heavily polluted days in these regions, assuming variation is solely due to meteorology. We also find that including the most important regional meteorological variable in each region improves the performance of the circulation-based indices in predicting daily PM_{2.5} concentrations on the regional scale. These results are beneficial to understanding and forecasting the occurrence of heavily polluted PM_{2.5} days in BTH, YRD and PRD from a large-scale perspective.

15
20
25

1 Introduction

Over the last few decades, rapid economic progress and urbanization in China have raised a number of environmental challenges. These include sharp increases in the atmospheric concentrations of particulate matter with an aerodynamic diameter of 2.5 μm or less (PM_{2.5}), which are of utmost concern for public health (Xu et al., 2013; Zheng et al., 2015). Episodes of haze and smog pollution with high levels of PM_{2.5}, in particular during winter, have become common in the most developed and highly populated city clusters in China (Zhang et al., 2007; Chan and Yao, 2008; Zhang et al., 2014).

30



Although emissions of pollutant precursors strongly influence air pollution levels, meteorology also plays a major role in air quality variability and trends through a combination of transport, transformation and deposition processes (e.g., Wang et al., 2009; Cheng et al., 2019). For instance, the extreme haze event in January 2013 in Beijing when the maximum instantaneous

35 $PM_{2.5}$ value exceeded $500 \mu\text{g}/\text{m}^3$, one of the worst air pollution events on record in China, has been attributed to unfavourable atmospheric dispersion conditions with weak surface winds and high humidity (Wang et al., 2014; Yang et al., 2015). In contrast, during winter and spring in 2015, $PM_{2.5}$ concentrations were much lower at most Chinese monitoring stations because of more favourable atmospheric dispersion conditions compared to those of the previous year (Wang et al., 2016).

40 While regional meteorological conditions are known to strongly influence air pollution levels, the responses of $PM_{2.5}$ concentrations to different meteorological variables are complex (e.g., Tai et al., 2010; Barmadimos et al., 2012; Dawson et al., 2014; Han et al. 2016; Garrido-Perez et al., 2017, 2021). Key processes and the relevant regional meteorological variables influencing $PM_{2.5}$ levels have been identified in recent studies. These processes include: a) secondary aerosol formation and hygroscopic growth associated with high relative humidity (RH; Sun et al., 2013; Wang et al., 2014), b) wet

45 deposition due to precipitation (Koch et al., 2003; Tai et al., 2010), c) horizontal dispersion of polluted air under high wind speeds (WSPD; Wang et al., 2012; Zhang et al., 2014), and d) vertical ventilation and dilution of the boundary layer via mechanically generated turbulence associated with strong wind shear (WSHR; Wang et al., 2018, 2019a) and via thermodynamically generated turbulence as measured by inversion intensity (INV; Zhao et al., 2013; Wang et al., 2014). Specifically, high RH, weak WSPD, strong INV and weak WSHR have been found to contribute to the accumulation and

50 growth of pollutants in a shallow and stable boundary layer over the North China Plain (e.g., Wang et al., 2014; Ge et al., 2019). However, RH can also be associated with precipitation and therefore removal of aerosols by wet deposition (Zhu et al., 2012; Leung et al., 2018), and may also be an indicator of air masses from different origins.

These key regional meteorological factors have been found to be affected by circulation patterns at larger scales over different regions of the globe (Tai et al., 2012; Garrido-Perez et al., 2017; Pei et al., 2018). Prominent large-scale circulation

55 patterns over China during winter include the East Asian winter monsoon (EAWM; Chang et al., 2006; Wang and Chen, 2010) and El Niño-Southern Oscillation (ENSO; Wang et al., 2000; Zhang et al., 2017). The EAWM resulting from the development of the cold-core Siberian High system is mainly characterised by dry cold low-level northerlies along the East Asian coast, the mid-tropospheric East Asian trough and the upper-tropospheric westerly jet stream (Jhun and Lee, 2004; Li and Yang, 2010; Wang and Lu, 2017). The EAWM has a significant impact on China's regional meteorological conditions,

60 including air temperature, wind speed, RH and atmospheric stability (Jeong and Park, 2017; Wang et al., 2019b), and hence influences $PM_{2.5}$ levels as noted above. ENSO, as the dominant mode of global ocean-atmosphere interaction, also substantially modulates regional meteorological conditions in China, through changes in atmospheric circulation patterns. The regional meteorological variables affected include RH and precipitation over southeastern China, and wind speed over northern China (Sun et al., 2018; He et al., 2019).



65 Previous studies of how the large-scale wintertime circulation modulates air quality in China through its effect on regional
meteorology have been primarily focused on Beijing and the North China Plain, the regions with the most severe PM_{2.5}
pollution (e.g., Wang et al., 2014, 2019b; Zhang et al., 2019). Broader regions in northern and southern China also show
clear relationships between PM_{2.5} concentrations and the EAWM intensity (e.g., Jeong and Park, 2017), aerosol optical depth
and the position of the Siberian High (Jia et al., 2015), as well as the number of haze days and ENSO intensity (e.g., He et
70 al., 2019). However, the major city clusters in northern, eastern and southern China, i.e., Beijing–Tianjin–Hebei (BTH), the
Yangtze River Delta (YRD) and the Pearl River Delta (PRD), respectively, have been considered jointly only in a few
studies (e.g., Leung et al., 2018; Hou et al., 2019). Furthermore, most of the existing large-scale circulation indices, such as
the EAWM indices (Wang et al., 2010), the Siberian High index (Wu and Wang, 2002) and the Haze Weather Index (Cai et
al., 2017), have been proposed for the North China Plain. Consequently, they do not reflect the link between the large-scale
75 circulation and PM_{2.5} levels over YRD and PRD. Indeed, Leung et al. (2018) found using PM_{2.5} observations between June
2014 and May 2017 that different dominant large-scale circulation modes could explain annual PM_{2.5} variability in BTH,
YRD and PRD.

Understanding the impact of the large-scale circulation on PM_{2.5} air quality in these three major populated regions of China
during winter, therefore, requires consideration of regional differences in the dominant large-scale circulation features. In
80 order to understand and predict the occurrence of days with high PM_{2.5} concentrations, it is critical to investigate the
relationship between the large-scale circulation and PM_{2.5} levels on daily timescales. This study examines the dominant
large-scale circulation-PM_{2.5} relationships separately for BTH, YRD and PRD during winter, and further proposes novel
circulation-based indices to explain the day-to-day variability of PM_{2.5} levels in each region. We first explore the relationship
of daily PM_{2.5} concentrations with specific regional meteorological variables across BTH, YRD and PRD (Section 3). We
85 then identify the dominant large-scale circulation associated with heavily polluted days for the three regions through its
effect on the most important regional meteorological variables, and propose specific circulation-based indices for these three
regions (Section 4). Furthermore, we assess the performance of these circulation-based indices in distinguishing different
levels of air pollution (Section 5) and examine the joint effect of the circulation-based indices and regional meteorology on
the day-to-day variability of PM_{2.5} (Section 6). Finally, Section 7 summarises the main results.

90 **2 Data and Methodology**

We use daily meteorological data from the ECMWF Reanalysis 5 (ERA5; Hersbach et al., 2020) on a 30 km × 30 km grid,
including zonal wind at 300 hPa, 900 hPa and 1000 hPa (U300, U900, U1000); meridional wind at 850 hPa, 900 hPa and
1000 hPa (V850, V900, V1000); geopotential height at 500 hPa (Z500); air temperature at 900 hPa and 1000 hPa; RH at 900
hPa and 1000 hPa; sea level pressure (SLP) and sea surface temperatures (SSTs). Hourly data are used to calculate daily
95 averages for 450 days during the five winters from 1st Dec 2013 – 28th Feb 2014 to 1st Dec 2017 – 28th Feb 2018 (hereafter
referred to as DJF 2013–2017). Daily precipitation is from the Global Precipitation Climatology Project (GPCP; Huffman et
al. 2001) 1° daily precipitation product. These meteorological fields are used to investigate both the large-scale circulation



features and regional meteorological conditions modulating $PM_{2.5}$ concentrations. Four meteorological fields representing relevant processes affecting $PM_{2.5}$ in the boundary layer (RH, WSPD, WSHR and INV) are evaluated, following Ge et al. (2019). RH and WSPD are used at 1000 hPa. Wind shear, WSHR, is calculated as

$$WSHR = \sqrt{(U_{900} - U_{1000})^2 + (V_{900} - V_{1000})^2}, \quad (1)$$

Inversion intensity, INV, is calculated as

$$INV = \theta_{v,900 \text{ hPa}} - \theta_{v,1000 \text{ hPa}}, \quad (2)$$

where θ_v is virtual potential temperature and the subscripts 900 and 1000 hPa specify the vertical levels at which θ_v is evaluated from air temperature and RH.

The six-year long high-resolution Chinese air quality reanalysis dataset (CAQRA; Kong et al., 2021) is the latest long-term air quality reanalysis for China. It contains surface fields of conventional pollutants, including $PM_{2.5}$, at high spatial (15 km×15 km) and temporal (1 h) resolution for the period 2013–2018. This dataset has been developed by assimilating pollutant concentrations from over 1000 surface air quality monitoring sites from the China National Environmental Monitoring Centre. CAQRA has been validated against independent datasets, yielding a good performance in reproducing the magnitude and variability of surface air pollutants in China on a regional scale (Kong et al., 2021). We use $PM_{2.5}$ hourly concentrations from this dataset to calculate daily averages for the same time period as the daily meteorological data (DJF 2013–2017, 450 days). $PM_{2.5}$ concentrations show a decreasing trend over the period of analysis, consistent with the primary emission reductions and $PM_{2.5}$ concentration decreases reported by many previous studies (e.g., Li et al., 2019, 2020). Therefore, to eliminate the influence of changing anthropogenic emissions, the daily $PM_{2.5}$ data are de-trended by removing the linear trend from the Dec 2013–Feb 2018 (1550 days) time series. To understand how meteorology drives clean vs. polluted conditions in a consistent way, percentile thresholds of the de-trended daily $PM_{2.5}$ data are used. We choose the 10th percentile (p10) of $PM_{2.5}$ concentrations as the clean threshold and the 90th percentile (p90) of $PM_{2.5}$ concentrations as the heavily polluted threshold. We then group all the days below p10 and above p90 and classify them as clean or heavily polluted days (45 days each).

Statistical significance is assessed at the 95% confidence level throughout this paper, unless otherwise stated. The effective numbers of degrees of freedom are calculated in order to assess the significance of correlations considering the effect of temporal autocorrelation (Allen and Smith 1994; Hu et al. 2017). A non-parametric bootstrap resampling method is used to assess the significance of differences between meteorological variables under heavily polluted and average conditions, as these variables do not necessarily follow normal distributions. This bootstrap resampling method generates random samples of meteorological variables for the whole period of analysis. Each random sample comprises 45 days, i.e., the total number of heavily polluted days. Then the difference between the mean of each sample and all the data is calculated. This procedure is repeated 10,000 times to create a random distribution of meteorological variable differences. Following this, differences calculated for heavily polluted days are compared with the distribution of meteorological variable differences. The



130 differences calculated for heavily polluted days are considered significantly negative or positive (at 95% confidence level) when they are below or above the 2.5% and 97.5% tails, respectively.

3 Influence of regional meteorological variables on daily PM_{2.5} variability

We first identify the meteorologically coherent regions representing BTH, YRD and PRD by searching for reanalysis grid cells where the daily PM_{2.5} concentrations are highly correlated ($r \geq 0.7$) with those in the grid cells corresponding to Beijing, Shanghai and Guangzhou, respectively (Fig. 1). This accounts for the regional nature of PM_{2.5} pollution and provides a more robust result than using the closest grid cells containing the cities or some arbitrary rectangular regions as previous studies have done (e.g., Leung et al., 2018; Hou et al., 2019). Daily regional PM_{2.5} concentrations are then calculated by averaging the data over these three homogeneous regions. Note that, as the 90th percentiles (p90) of daily average PM_{2.5} differ for the three regions, heavily polluted days defined on p90 correspond to concentrations > 97 $\mu\text{g}/\text{m}^3$ for BTH, > 110 $\mu\text{g}/\text{m}^3$ for YRD and > 68 $\mu\text{g}/\text{m}^3$ for PRD. The value of p90 PM_{2.5} is higher in YRD than in BTH, because the smaller size of YRD is more representative of a coherent urban environment (Fig. 1). For consistency, the gridded meteorological fields described in Section 2 are averaged over the same regions to construct daily regional meteorological variables.

Figure 2 shows the lagged relationship of daily regional PM_{2.5} concentrations with specific regional meteorological variables in these three homogeneous regions for the entire DJF 2013–2017 period. There are positive correlations for INV and negative correlations for WSHR and WSPD with PM_{2.5} concentrations for all three regions. This occurs even when daily PM_{2.5} concentrations are lagged by a few days. This suggests that high PM_{2.5} days are associated with poor vertical ventilation (increased INV and reduced WSHR) and reduced horizontal dispersion (weak WSPD) for several days preceding the high PM_{2.5} levels. In particular, WSPD is the variable with the highest correlation with PM_{2.5} concentrations in YRD, appearing for a one-day lag ($r = -0.43$) (Fig. 2b). Unlike the other three variables considered, the relationship between RH and PM_{2.5} concentration varies across BTH, YRD and PRD. A positive correlation is seen between RH and PM_{2.5} concentrations for BTH, with the highest value at zero lag ($r = 0.66$) (Fig. 2a). This highlights the general contrast between clean, dry air reaching BTH from the northwest and more polluted, humid air reaching BTH from central and eastern China. However, RH is negatively correlated with PM_{2.5} concentrations in the other two regions, with larger correlations in PRD than in YRD. The high correlations in PRD persist over the previous four days (with the highest value of $r = -0.52$ for a two-day lag) (Fig. 2c). This reflects the association of high RH with cleaner oceanic air and precipitation, and hence wet deposition in PRD (e.g., Zhu et al., 2012; Jeong and Park, 2017). RH is the meteorological variable presenting the highest correlation value with PM_{2.5} concentrations over both BTH and PRD. These results are consistent with previous findings of the different patterns in PM_{2.5}–RH relationships over northern and southern China (Wang et al., 2014; Leung et al., 2018). Consequently, RH on the same day ($r = 0.66$), WSPD one day before ($r = -0.43$) and RH two days before ($r = -0.52$) are identified as the most important regional meteorological variables contributing to the day-to-day variability of PM_{2.5} concentrations over BTH, YRD and PRD, respectively. Among the second most relevant meteorological variables, WSPD



and INV stand out for BTH and PRD, respectively, with absolute correlation coefficients close to 0.5 for some time lags. Following previous analyses (e.g., Tai et al., 2010, 2012; Leung et al., 2018; Ge et al., 2019), we now investigate how the relationships between PM_{2.5} concentrations and the most important regional meteorological variables described above and considering the same time lags are caused by common association with large-scale circulation systems.

4 Modulation of daily PM_{2.5} by the large-scale circulation

Using ERA-5 reanalysis data for DJF 2013-17, we find the wintertime large-scale circulation over East Asia is dominated by the Siberian High as seen from the high SLP values centred over northwestern Mongolia (Fig. 3a). The Siberian High induces northerly near-surface winds along its eastern edge, which bring cold, clean air to northern and central China as indicated by negative V850 values (Fig. 3b). This northerly near-surface flow is also associated with the middle tropospheric East Asian trough, characterised by low Z500 values over Northeast China as seen in Figure 3c. Over eastern and southern China, wet and warm southerly winds blow from the South China Sea (Fig. 3b), bringing precipitation (Fig. 3d).

Previous studies have introduced a variety of large-scale circulation indices to characterise atmospheric circulation in East Asia. Here we apply three commonly used EAWM indices (*I_{Yang}* (V850): Yang et al., 2002; *I_{Sun}* (Z500): Sun and Li, 1997; *I_{Jhun}* (U300): Jhun and Lee, 2004) and a widely used Siberian High index (*I_{SH}*; Wu and Wang, 2002) to test their relationship with daily PM_{2.5} concentrations separately for the three meteorologically coherent regions using reanalysis data (Table S1). We reverse *I_{Yang}* and *I_{Sun}* by multiplying them by -1 so that a high index value represents a strong EAWM. The three EAWM indices have been selected because they reflect the circulation characteristics of the EAWM in the lower, middle and upper troposphere, respectively (e.g., Wang et al., 2019b). Linear correlations of all three EAWM indices with the daily PM_{2.5} concentrations for the whole period of analysis are significant (at 99% confidence level) only for BTH (r ranging from -0.54 to -0.36), whereas absolute correlation coefficients do not exceed 0.12 for YRD and PRD. This suggests that these three typical EAWM indices do not capture well the relationship between the large-scale circulation and daily PM_{2.5} concentrations over the YRD and PRD regions used in this study. The Siberian High index (*I_{SH}*) is significantly correlated with daily PM_{2.5} concentrations for all three regions, although the correlations are not strong (r ranging from -0.19 to -0.13). Considering the moderate correlations found for YRD and PRD, we further investigate the influence of large-scale circulation on daily PM_{2.5} variability through its direct effect on the most important regional meteorological variables identified separately for the three regions. For this purpose, we first examine the dominant large-scale circulation features associated with heavily polluted days for each region, then identify the correlation patterns of daily PM_{2.5} concentrations with these circulation variables for the whole period of analysis and define circulation-based indices separately for the three regions. These analyses will be carried out considering the same time lags as those for the most important regional meteorological variables identified in Section 3. The daily meteorological reanalysis data are normalised by subtracting the means of individual variables and dividing by their standard deviations to yield fields with zero means and unit variance before calculating these indices.



195 4.1 Beijing–Tianjin–Hebei (BTH)

As shown in Figure 2a, the strongest correlations between daily PM_{2.5} concentrations and regional meteorological variables over BTH are found for RH with no time lag. In this section, we examine circulation variables during heavily polluted days (PM_{2.5} above p90; daily PM_{2.5} concentrations > 97 µg/m³ for BTH) over this region. Figure 4 shows the average composites of circulation variables (SLP, V850 and Z500) for heavily polluted days over BTH (upper panels), along with the difference (lower panels) between heavily polluted days and the winter (DJF) mean (as displayed in Figure 3) during 2013–2017. Heavily polluted days are characterised by a weak and eastward-extended Siberian High, weak northerly winds at 850 hPa over North China, and a shallow East Asian trough at 500 hPa, reflecting a weak EAWM circulation (Jia et al., 2015; Ge et al., 2019). Following these results, we calculate daily correlations of the PM_{2.5} concentrations with SLP, V850 and Z500 for the whole period of analysis to assess to what extent the observed circulation anomalies can be used to represent the day-to-day variability of PM_{2.5}. The resulting circulation-PM_{2.5} correlation patterns are displayed in Fig. 5. The daily PM_{2.5} concentrations for BTH have negative correlations with SLP over mainland China (and positive correlations centred over Japan, Fig. 5a), positive correlations with V850 over eastern China (Fig. 5b), and positive correlations with Z500 centred over Northeast China (Fig. 5c), in accord with the observed departures of heavily polluted days from the winter mean.

Based on these circulation-PM_{2.5} correlation patterns, we now select broad regions (yellow rectangles in Figure 5) which represent the highest correlations with PM_{2.5} concentrations in BTH and then construct spatial averages of the daily meteorological fields over these regions. The area-weighted averages of daily normalised SLP, V850 and Z500 show significant correlations with daily PM_{2.5} concentrations in BTH (at 99% confidence level), especially for Z500 ($r = 0.67$), followed by V850 ($r = 0.59$) and SLP ($r = 0.54$) (Table 1). Note that these correlations are stronger than those using the EAWM indices and the Siberian High index from the literature (see Table S1). We therefore use Z500 averaged over Northeast China, Korea and the Sea of Japan [118–139°E, 33–50°N] (rectangle in Figure 5c) to build a Z500-based index for BTH (I_{Z500_BTH}) for all days in DJF 2013–17). I_{Z500_BTH} is calculated as the mean of daily normalised Z500 in that region with a reversed sign (eq. 3) so that negative values of I_{Z500_BTH} indicate a shallow East Asian trough.

$$I_{Z500_BTH} = - \overline{Z500 (33^\circ - 50^\circ N, 118^\circ - 139^\circ E)} \quad (3)$$

I_{Z500_BTH} is significantly correlated both with PM_{2.5} concentrations ($r = -0.67$ in Table 2) and with RH ($r = -0.64$ in Table 2) in BTH on daily time scales. These results point to a shallow East Asian trough as the dominant large-scale circulation pattern favouring high PM_{2.5} concentrations and high RH in BTH. The shallow East Asian trough in the middle troposphere inhibits the invasion of northerly cold air from the rear of the trough to northern and central China, yielding southerly wind anomalies (Figs. 4e–f), as found in other studies (e.g., Zhang et al., 2014). This anomalous warm and humid air from the south therefore creates appropriate conditions for the accumulation and possibly the growth of fine aerosols and also suppresses the southward transport of aerosols away from BTH (see positive correlations for RH and negative correlations for WSPD in Fig. 2a).



4.2 Yangtze River Delta (YRD)

As shown in Figure 2b, the correlations between daily PM_{2.5} concentrations and regional meteorological variables over YRD are highest for the most important regional meteorological variable (WSPD) when daily PM_{2.5} concentrations are lagged by one day. Hence, in this section, we focus on the circulation variables (SLP, V850 and Z500) one day before heavily polluted days over this region. Heavily polluted days in YRD (PM_{2.5} above p90; daily PM_{2.5} concentrations > 110 µg/m³) are mainly characterised by reduced SLP over eastern China, indicating a weak Siberian High (Figs. 6a and 6d), and a shallow East Asian trough with positive Z500 anomalies centred over Japan (Figs. 6c and 6f). This weakened intensity of the Siberian High is associated with a northerly wind anomaly over both North and South China, as well as a significant southerly wind anomaly over Northeast China and Japan (Fig. 6e). The northerly wind anomaly implies a weakening of the winter mean southerly wind over southern China and a strengthening of the winter mean northerly wind over northern China (Figs. 6b, 6e). This different pattern in southern versus northern China is further supported by the daily WSPD-PM_{2.5} correlation features for the whole period of analysis, where daily PM_{2.5} concentrations in YRD are negatively correlated with WSPD over southern China and positively correlated over northern China (Fig.S1). Furthermore, the daily PM_{2.5} concentrations for YRD have negative correlations with SLP centred over northeast China (Fig. 7a), negative correlations with V850 over both southern China and northern China (and positive correlations over northeast China and Japan, Fig. 7b), and positive correlations with Z500 centred over northwest China (Fig. 7c). These circulation-PM_{2.5} correlation patterns for the whole period of analysis are consistent with the circulation anomalies shown for heavily polluted days in Figure 6.

We then identify the regions with the highest correlations of area-weighted average daily normalised meteorological fields with daily PM_{2.5} concentrations in YRD. Among these three meteorological fields (i.e., SLP, V850, Z500), for the regions that show the highest correlations with PM_{2.5} concentrations in YRD (yellow rectangles in Fig 7), SLP is found to have the highest correlation ($r = -0.33$) (Table 1). We therefore use SLP averaged over Northeast China [30–49°N, 111–131°E] (rectangle in Figure 7a) to build a normalised SLP-based index for YRD (I_{SLP_YRD}) for all days in DJF 2013-17 (eq. 4). Negative values of I_{SLP_YRD} indicate a weak Siberian High.

$$I_{SLP_YRD} = \overline{SLP(30^{\circ}-49^{\circ}N, 111^{\circ}-131^{\circ}E)} \quad (4)$$

I_{SLP_YRD} is significantly correlated both with PM_{2.5} concentrations ($r = -0.33$ in Table 2) and with WSPD ($r = 0.29$ in Table 2) in YRD on daily time scales. This suggests a weakened Siberian High as the dominant large-scale circulation pattern contributing to higher concentrations of PM_{2.5} and reduced WSPD in YRD. The associated reduction in the southerly wind reported above for southern and eastern China together with reduced WSPD implies a greater suppression of horizontal dispersion, contributing to air pollution accumulation over YRD. Moreover, strengthened northerly winds in northern China may lead to southward transport of aerosols emitted from sources over northern China to YRD, as also indicated by previous studies (Li et al., 2012; Jeong and Park, 2017).



260 We repeated the analysis above to examine the sensitivity to different time lags. The observed circulation anomaly patterns without a lag resemble those found for one-day lag, although they are displaced to the east because of the eastward movement of synoptic systems in the midlatitudes (Fig. S2). The region that shows the highest correlations with PM_{2.5} concentrations in YRD on the SLP-PM_{2.5} correlation pattern is slightly less significant without a lag, again with an eastward shift (Fig. S3).

265 4.3 Pearl River Delta (PRD)

In contrast to BTH and YRD, the highest correlations of daily PM_{2.5} concentrations over PRD with the two most important regional meteorological variables (RH and INV) persist when PM_{2.5} is lagged by several days (Fig. 2c). As the maximum correlations are found with a lag of two days, we examine composites of two circulation variables (SLP and V850) and precipitation two days before the occurrence of heavily polluted days over PRD (PM_{2.5} above p90; daily PM_{2.5} concentrations > 68 µg/m³) (Fig. 8). These are mainly characterised by reduced SLP centred over northern China and increased SLP over southwestern China, weak southerly winds at 850 hPa over South China, as well as precipitation deficits over southern China. Correlation patterns of PM_{2.5} with the same fields (Fig. 9) for the whole period of analysis further support these circulation anomalies for heavily polluted days. Daily PM_{2.5} concentrations over PRD have negative correlations with SLP over northern China (and positive correlations over southern China, Fig. 9a), negative correlations with V850 over South China and the South China Sea (Fig. 9b), as well as with precipitation over southern China (Fig. 9c). There are also negative correlations between daily PM_{2.5} concentrations and SSTs over the central and eastern equatorial Pacific (and positive correlations over the western equatorial Pacific), as well as negative correlations for SLP over the western North Pacific (Fig. S4). These circulation-PM_{2.5} correlation features display characteristic ENSO-related patterns over the Pacific and East Asia (e.g., Wang et al., 2000). La Niña events are associated with warm SSTs in the western Pacific and cold SSTs in the central and eastern equatorial Pacific, reduced SLP over the western North Pacific and descending motion on the northwestern flank of this reduced SLP. The opposite relationships are seen for El Niño (Fig. S4). This anomalous subsidence with suppressed precipitation (Fig. 9) has been found to play a major role in high PM_{2.5} concentrations over southern China (e.g., He et al., 2019; Sun et al., 2018). We also find that more than 80% (37 out of 45) of heavily polluted days in PRD are in La Niña years, considered here as those when the Niño 3.4 index (area-weighted averages of SSTs anomaly over 5°S –5°N, 120° –170°W) is less than -0.5. Nonetheless, these results should be treated with caution because of the relatively short time series considered (only 5 winters with PM_{2.5} data).

Comparing the correlations of the area-weighted average daily normalised meteorological fields with daily PM_{2.5} concentrations, V850 is found to have the highest value ($r = -0.43$ in Table 1), followed by SLP and precipitation ($r < 0.4$) over the regions showing the highest correlation with PM_{2.5} concentrations in PRD (yellow rectangles in Figure 9). We therefore build a normalised daily V850-based index for PRD (I_{V850_PRD}) by averaging V850 over the region of South China and the South China Sea [100–118°E, 10–22°N] (rectangle in Fig. 9b) (eq. 5). Negative values of I_{V850_PRD} indicate weak southerly winds over South China.



$$I_{V850_PRD} = \overline{V850 (10^{\circ} - 22^{\circ}N, 100^{\circ} - 118^{\circ}E)} \quad (5)$$

295 Weak southerly winds over southern China as the dominant large-scale circulation pattern are associated with greater polluted continental flow and precipitation deficits under weak cleaner oceanic winds (Figs. 8e-f) that are conducive to air pollution over PRD via reduced wet deposition. Consequently, I_{V850_PRD} is not only negatively correlated with $PM_{2.5}$ concentrations ($r = -0.43$ in Table 2) but also positively correlated with regional RH in PRD ($r = 0.64$ in Table 2). The anomalous subsidence yielding precipitation deficits over southern China is also associated with a shallow and stable
300 boundary layer where the vertical dilution capacity of the lower atmosphere reduces (see negative correlations for RH and positive correlations for INV in Fig. 2c). Overall, the observed circulation patterns for smaller and zero lag are broadly similar to those found for a two-day lag (Fig. S5), although the $V850$ - $PM_{2.5}$ correlations weaken as the lag is reduced (Fig. S6).

5 Performance of circulation-based indices for differing air pollution levels

305 Our analyses confirm that the proposed circulation-based indices are significantly correlated with the most important regional meteorological variables and the $PM_{2.5}$ concentrations on daily time scales during DJF 2013-17. The correlations are significant at the 99% confidence level (Table 2). To further examine the performance of circulation-based indices for distinguishing different levels of air quality, we show the distributions of I_{Z500_BTH} , I_{SLP_YRD} and I_{V850_PRD} for several percentile thresholds of daily $PM_{2.5}$: above p90 (heavily polluted), p50-90 (moderately polluted), p10-50 (moderately clean) and below
310 p10 (clean) (Fig. 10). Note that the sample size for moderate events is larger than for heavily polluted/clean events and also that daily $PM_{2.5}$ concentrations are lagged by one and two days in the case of YRD and PRD, respectively, for consistency with the previous analysis.

For BTH, the average value of I_{Z500_BTH} with associated 95% confidence intervals are: $I_{Z500_BTH} = -1.04 \pm 0.20$ for heavily polluted days, $I_{Z500_BTH} = -0.28 \pm 0.10$ for moderately polluted days, $I_{Z500_BTH} = 0.35 \pm 0.10$ for moderately clean days and
315 $I_{Z500_BTH} = 0.83 \pm 0.23$ for clean days (Fig. 10a). The values of I_{Z500_BTH} for these four categories differ (i.e., the confidence intervals do not overlap) at the 95% confidence level and I_{Z500_BTH} can distinguish between different levels of air quality, not just extreme heavily polluted or clean conditions. Ge et al. (2019) used a Siberian High index (I_{SH} ; Wu and Wang, 2002), which we tested in Section 4, and a potential vorticity based EAWM index (I_{PV} ; Huang et al., 2016) to distinguish different $PM_{2.5}$ pollution levels in Beijing. They found that I_{SH} can effectively distinguish clean days (daily $PM_{2.5}$ concentrations ≤ 75
320 $\mu\text{g}/\text{m}^3$) from polluted days (daily $PM_{2.5}$ concentrations $\geq 75 \mu\text{g}/\text{m}^3$), but could not distinguish between moderate and severe (daily concentrations $PM_{2.5} \geq 150 \mu\text{g}/\text{m}^3$) $PM_{2.5}$ pollution. The I_{PV} index exhibited the reverse problem. This shows that I_{Z500_BTH} performs better than existing circulation indices, both in capturing the relationship between the dominant large-scale circulation and daily $PM_{2.5}$ concentrations (Tables S1 and 2) and in distinguishing pollution levels in BTH (Fig. 10a). In the case of YRD (Fig. 10b), I_{SLP_YRD} can effectively distinguish heavily polluted days ($I_{SLP_YRD} = -0.32 \pm 0.28$) from clean days
325 ($I_{SLP_YRD} = 0.51 \pm 0.19$). However, differences are not significant between heavily and moderately polluted days ($I_{SLP_YRD} = -$



0.19 ± 0.11) and are not highly significant between clean and moderately clean days ($I_{SLP_YRD} = 0.24 \pm 0.11$). For PRD (Fig. 10c), I_{V850_PRD} can distinguish well between heavily polluted days ($I_{V850_PRD} = -0.31 \pm 0.16$), moderately clean days ($I_{V850_PRD} = 0.22 \pm 0.10$) and clean days ($I_{V850_PRD} = 0.83 \pm 0.19$), but not between heavily polluted and moderately polluted days ($I_{V850_PRD} = -0.28 \pm 0.09$).

330 To further illustrate the relationships between the dominant large-scale circulation, as represented by these circulation-based indices, and the severity of PM_{2.5} pollution at daily timescales, we show the joint frequency distributions of daily values of circulation-based indices compared to daily PM_{2.5} concentrations (Fig. 11). We show the linear relationship between each respective index and PM_{2.5} concentrations, as given in Table 2, with higher PM_{2.5} concentrations and smaller (negative) index values on heavily polluted days, and vice versa. Moderately polluted days (PM_{2.5} above p50; daily PM_{2.5} concentrations > 43
335 µg/m³ for BTH, > 59 µg/m³ for YRD, > 39 µg/m³ for PRD) tend to occur when the circulation-based indices are negative. This is more often the case for heavily polluted days (PM_{2.5} above p90; daily PM_{2.5} concentrations > 97 µg/m³ for BTH, > 110 µg/m³ for YRD, > 68 µg/m³ for PRD), in particular for BTH where 98% (44 of 45) of those days have negative values of I_{Z500_BTH} compared to 66% (119 of 180) of moderately polluted days (p50-90 PM_{2.5}). However, there is no such apparent distinction in the other two regions, since around 62% of both heavily and moderately polluted days in YRD have negative
340 values of I_{SLP_YRD} , and 70% of these days in PRD have negative values of I_{V850_PRD}). Alternatively, 51% (23 of 45), 16% (7 of 45), 13% (6 of 45) of heavily polluted days in BTH, YRD and PRD, respectively, occur when circulation-based indices are below -1.

By contrast, moderately clean days (PM_{2.5} below p50) and, to a greater extent, clean days (PM_{2.5} below p10; daily PM_{2.5} concentrations < 16 µg/m³ for BTH, < 29 µg/m³ for YRD, < 15 µg/m³ for PRD) tend to occur when circulation-based indices
345 are positive. 91% (41 of 45), 80% (36 of 45), 89% (40 of 45) of clean days in BTH, YRD and PRD have positive indices values. As expected, this tendency is even more pronounced for larger values of the indices, as 93% (52 of 56), 88% (37 of 42) and 93% (37 of 40) of days with I_{Z500_BTH} , I_{SLP_YRD} and I_{V850_PRD} exceeding 1 are classified as moderately clean or clean. The share of days with positive values of the circulation indices generally decreases with increasing PM_{2.5} pollution levels for all three regions, especially for PRD where the percentage of days with positive values of I_{V850_PRD} decreases from 89% of
350 clean days to only 61% (110 of 180) of moderately clean days. The results of the analyses conducted so far show that the daily circulation-based indices proposed in this study can capture most of the day-to-day variability of PM_{2.5} and also identify days with different pollution levels, although with poorer performance for YRD than for the other two regions.

6 Joint effect of large-scale circulation and regional meteorology

The relatively moderate correlation between daily circulation-based index and daily PM_{2.5} concentrations in YRD reflects the
355 complex mix of factors affecting the day-to-day variability of this pollutant. We have also found that regional meteorological variables (the most relevant ones identified in Section 3) influence the PM_{2.5} concentrations for the three regions (e.g., $r = 0.66$ for RH in BTH, $r = -0.43$ for WSPD in YRD and $r = -0.52$ for RH in PRD). On the other hand, there are significant correlations between the circulation-based indices and the most relevant regional meteorological variables in each region,



indicating that the effect of circulation on $PM_{2.5}$ occurs through modulation of the regional meteorology. The relationship
360 between the daily circulation-based index and the most important daily regional meteorological variable is weaker in YRD (r
 $= 0.29$) than for the other two regions ($r = -0.64$ for BTH; $r = 0.64$ for PRD) (Table 2). This shows that the daily circulation-
based index is not solely capable of capturing the regional meteorological variability driving day-to-day $PM_{2.5}$ changes in
YRD.

While there is some co-variation of the large-scale circulation with the regional meteorology, they can be combined to
365 reproduce the day-to-day variability of $PM_{2.5}$ with improved performance. We therefore build multiple regression models
including a linear combination of the most important regional meteorological field and the large-scale circulation index in
each region (Table 3). The inclusion of regional meteorology explains more of the variance in the $PM_{2.5}$ concentrations for
all three regions ($R^2(I_{Z500_BTH} + RH) = 0.54$, $R^2(I_{SLP_YRD} + WSPD) = 0.23$, $R^2(I_{V850_PRD} + RH) = 0.30$) than the large-scale
circulation index alone ($R^2(I_{Z500_BTH}) = 0.45$, $R^2(I_{SLP_YRD}) = 0.11$, $R^2(I_{V850_PRD}) = 0.18$). However, if we consider the regional
370 meteorological variable alone we see that its relationship with daily $PM_{2.5}$ concentrations explains more of the variance than
the large-scale circulation variable for the YRD and PRD regions. Hence, compared to a linear model on the most relevant
regional meteorological field, these multiple models do not bring major improvements for YRD and PRD, where the
increase in explained variance is relatively small (0.18 vs. 0.23 for YRD and 0.27 vs. 0.30 for PRD). As expected, the signs
of the regression coefficients for the most important regional meteorological field and the large-scale circulation index
375 (Table S2) are consistent with those of their respective correlation coefficients with $PM_{2.5}$.

7 Discussion and conclusions

This study investigates the modulation of daily $PM_{2.5}$ concentrations by regional meteorological conditions and large-scale
circulation in three major populated regions of China during winter. Using a new high-resolution Chinese air quality
reanalysis dataset, major regions associated with BTH, YRD and PRD are identified where daily $PM_{2.5}$ concentrations are
380 spatially coherent. For these three regions, we find that the regional meteorological variables most correlated with daily
 $PM_{2.5}$ concentrations are different: RH on the same day for BTH ($r = 0.66$), WSPD one day before for YRD ($r = -0.43$), and
RH two days before for PRD ($r = -0.52$). We identify the dominant large-scale circulation patterns associated with heavily
polluted days ($PM_{2.5}$ above $p90$) considering the same time lags. Based on these dominant large-scale circulation features, we
propose three new circulation-based indices that can be used both to explain the day-to-day variability of the $PM_{2.5}$
385 concentrations and to predict the occurrence of heavily polluted days and clean days ($PM_{2.5}$ below $p10$) in each region. The
proposed circulation indices capture the relationship between the dominant large-scale circulation and daily $PM_{2.5}$
concentrations better than existing EAWM indices (Yang et al., 2002; Sun and Li, 1997; Jhun and Lee, 2004) and the
Siberian High index (Wu and Wang, 2002). They improve on the capability of circulation-based indices (e.g., Wu and Wang,
2002; Huang et al., 2016) to distinguish $PM_{2.5}$ pollution levels in BTH, and provide the first daily circulation-based indices
390 specifically for YRD and PRD. Furthermore, the inclusion of regional meteorology slightly improves the performance of the



large-scale circulation-based indices to predict the evolution of the regional PM_{2.5} concentrations in these regions on daily time scales.

Specifically, in BTH, heavily polluted days preferentially occur on days with a weak and eastward-extended Siberian High, weak northerly winds at 850 hPa over North China, and a shallow East Asian trough at 500 hPa. Of these meteorological variables, we find that the shallow East Asian trough, represented as the negative area-weighted daily averages of Z500 centred over Northeast China, has the strongest relationship with both PM_{2.5} concentrations ($r = -0.67$) and RH ($r = -0.64$). This suggests a strong contribution of the warm, humid air from the south and weak transport of northerly cold, dry air associated with the shallow East Asian trough to air pollution accumulation in BTH. Hence the intensity of the trough can be used as an indicator of PM_{2.5} levels. The 500 hPa geopotential height-based index for BTH (I_{Z500_BTH}) can distinguish between different levels of PM_{2.5} pollution well, not just between heavily polluted and clean conditions. On average, PM_{2.5} pollution levels generally increase as the daily values of I_{Z500_BTH} turn from positive to negative. Furthermore, the inclusion of RH ($R^2(I_{Z500_BTH} + \text{RH}) = 0.54$) explains more daily PM_{2.5} variability than the simpler model I_{Z500_BTH} ($R^2(I_{Z500_BTH}) = 0.45$).

In YRD, heavily polluted days are characterised by the following circulation anomalies: a weak Siberian High, strong northerly winds over North China and weak southerly winds over South China, as well as a shallow East Asian trough with positive Z500 anomalies centred over Japan. A weak Siberian High, represented as the area-weighted daily averages of SLP centred over Northeast China, shows the largest correlation with PM_{2.5} concentrations ($r = -0.33$) and WSPD ($r = 0.29$), and hence is the dominant circulation variable used to build an index of PM_{2.5} levels. This reflects the relationship between weak southerly winds over southern China, associated with a weak Siberian High, and poor horizontal dispersion of polluted air in YRD. The sea level pressure-based index for YRD (I_{SLP_YRD}) does not perform as well as I_{Z500_BTH} in distinguishing between moderately clean (p10-50 PM_{2.5}) and clean days or between moderately polluted (p50-90 PM_{2.5}) and heavily polluted days. The inclusion of WSPD ($R^2(I_{SLP_YRD} + \text{WSPD}) = 0.23$) explains more daily PM_{2.5} variability than I_{SLP_YRD} ($R^2(I_{SLP_YRD}) = 0.11$).

In PRD, heavily polluted days are characterised by La Niña-related circulation patterns: reduced SLP centred over northern China and increased SLP over southwestern China and south-eastern Asia, weak southerly winds at 850 hPa over southern China, as well as a precipitation deficit over southern China. Weak southerly winds over southern China are identified as the dominant large-scale circulation anomaly, as seen from the largest correlations of the area-weighted daily averages of V850 over southern China and the South China Sea with PM_{2.5} concentrations ($r = -0.43$) and RH ($r = 0.64$) in the region. This illustrates the influence of flow from more polluted continental regions and of precipitation deficits under weak humid southerly winds on PM_{2.5} pollution through reduced wet deposition in PRD. The 850 hPa meridional wind-based index for PRD (I_{V850_PRD}) can distinguish between moderately clean and clean days better than between moderately polluted and heavily polluted days. The inclusion of RH ($R^2(I_{V850_PRD} + \text{RH}) = 0.30$) explains considerably more daily PM_{2.5} variability than I_{V850_PRD} ($R^2(I_{V850_PRD}) = 0.18$).



These results demonstrate the benefits of considering the large-scale circulation for air quality studies over China. The novel
425 circulation indices proposed here explain a considerable fraction of the day-to-day variability of winter PM_{2.5} over major
metropolitan regions and can be combined with regional meteorological fields to improve our capability to predict the
variability of this pollutant. Although the circulation indices explain less variance than the most relevant regional
meteorological fields for YRD and PRD, we expect weather prediction and climate models to better represent these features
of the large-scale circulation than regional meteorological fields such as surface wind speed and RH. There are however two
430 limitations inherent in this work. Firstly, the relationships between atmospheric circulation and daily PM_{2.5} concentrations
may not be linear, as assumed in this study. Although we have improved the explained daily variability of PM_{2.5} by linearly
combining the most important regional meteorological field and the large-scale circulation index, non-linear models that
account for the covariance of meteorological fields (e.g., Barmadimos et al., 2011, 2012, Garrido-Perez et al., 2021) or
dimensionality reduction techniques such as principal component analysis (e.g., Tai et al., 2012; Shen et al., 2015; Leung et
435 al., 2018) merit further consideration. In addition, these large-scale relationships are based on only five winters of data,
because high spatiotemporal coverage of air pollution measurements are only available in China from 2013. Hence, whilst
our results are encouraging (e.g., we find that more than 80% of heavily polluted days in PRD occur in La Niña years), the
robustness of these results need to be verified using longer-term data. Despite these limitations, this study has shown that
dominant large-scale circulation features influence both regional meteorological processes and the PM_{2.5} concentrations in
440 BTH, YRD and PRD on a timescale of days, and can be used to construct indices for distinguishing clean days from heavily
polluted days in these major populated regions of China. These results are beneficial to understanding and forecasting the
occurrence of air pollution episodes in the three regions from a large-scale perspective.

Acknowledgements

We acknowledge the use of data from ERA-5 ([https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-pressure-](https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-pressure-levels?tab=overview)
445 [levels?tab=overview](https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-pressure-levels?tab=overview)), GPCP (<https://rda.ucar.edu/datasets/ds728.3/#!/description>) and CAQRA
(<http://www.en.scidb.cn/en/detail?dataSetId=696756084735475712&dataSetType=personal&version=V1#>). Oliver Wild
and Ruth M. Doherty thank the Natural Environment Research Council (NERC) for funding under grants NE/N006925/1,
NE/N006976/1 and NE/N006941/1. Carlos Ordóñez thanks the Spanish Ministerio de Economía y Competitividad [grant
number RYC-2014-15036]. Chaofan Li thanks the National Key Research and Development Program of China (Grant No.
450 2018YFA0606501).



455 References

- Allen, M. R., & Smith, L. A.: Investigating the origins and significance of low-frequency modes of climate variability, *Geophysical Research Letters*, 21(10), 883-886, <https://doi.org/10.1029/94GL00978>, 1994.
- Barnpadimos, I., Hueglin, C., Keller, J., Henne, S., & Prévôt, A. S. H.: Influence of meteorology on PM₁₀ trends and variability in Switzerland from 1991 to 2008, *Atmospheric Chemistry and Physics*, 11(4), 1813-1835, <https://doi.org/10.5194/acp-11-1813-2011>, 2011.
- 460 Barnpadimos, I., Keller, J., Oderbolz, D., Hueglin, C., & Prévôt, A. S. H.: One decade of parallel fine (PM_{2.5}) and coarse (PM₁₀-PM_{2.5}) particulate matter measurements in Europe: trends and variability, *Atmospheric Chemistry and Physics*, 12(7), 3189-3203, <https://doi.org/10.5194/acp-12-3189-2012>, 2012.
- Cai, W., Li, K., Liao, H., Wang, H., & Wu, L.: Weather conditions conducive to Beijing severe haze more frequent under climate change, *Nature Climate Change*, 7(4), 257-262, <https://doi.org/10.1038/nclimate3249>, 2017.
- 465 Chan, C. K., & Yao, X.: Air pollution in mega cities in China, *Atmospheric environment*, 42(1), 1-42, <https://doi.org/10.1016/j.atmosenv.2007.09.003>, 2008.
- Chang, C. P., Wang, Z., & Hendon, H.: The Asian winter monsoon, In *The Asian Monsoon* (pp. 89-127), Springer, Berlin, Heidelberg, 2006.
- 470 Cheng, J., Su, J., Cui, T., Li, X., Dong, X., Sun, F., ... & He, K.: Dominant role of emission reduction in PM_{2.5} air quality improvement in Beijing during 2013–2017: a model-based decomposition analysis, *Atmospheric Chemistry and Physics*, 19(9), 6125-6146, <https://doi.org/10.5194/acp-19-6125-2019>, 2019.
- Dawson, J. P., Bloomer, B. J., Winner, D. A., & Weaver, C. P.: Understanding the meteorological drivers of US particulate matter concentrations in a changing climate, *Bulletin of the American Meteorological Society*, 95(4), 521-532, <https://doi.org/10.1175/BAMS-D-12-00181.1>, 2014.
- 475 Garrido-Perez, J. M., Ordóñez, C., & Garcia-Herrera, R.: Strong signatures of high-latitude blocks and subtropical ridges in winter PM₁₀ over Europe, *Atmospheric Environment*, 167, 49-60, <https://doi.org/10.1016/j.atmosenv.2017.08.004>, 2017.
- Garrido-Perez, J. M., García-Herrera, R., & Ordóñez, C.: Assessing the value of air stagnation indices to reproduce PM₁₀ variability in Europe, *Atmospheric Research*, 248, 105258, <https://doi.org/10.1016/j.atmosres.2020.105258>, 2021.
- 480 Ge, W., Yin, Y., Wright, J. S., Huang, W., Jia, B., Wang, Y., & Yang, Z.: Links Between the Large-Scale Circulation and Daily Air Quality Over Central Eastern China During Winter, *Journal of Geophysical Research: Atmospheres*, 124(13), 7147-7163., <https://doi.org/10.1029/2018JD030154>, 2019.
- Han, L., Zhou, W., & Li, W.: Fine particulate (PM_{2.5}) dynamics during rapid urbanization in Beijing, 1973–2013, *Scientific reports*, 6(1), 1-5, <https://doi.org/10.1038/srep23604>, 2016.
- 485



- He, C., Liu, R., Wang, X., Liu, S. C., Zhou, T., & Liao, W.: How does El Niño-Southern Oscillation modulate the interannual variability of winter haze days over eastern China?, *Science of The Total Environment*, 651, 1892-1902, <https://doi.org/10.1016/j.scitotenv.2018.10.100>, 2019.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., ... & Thépaut, J. N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146(730), 1999-2049, 490
<https://doi.org/10.1002/qj.3803>, 2020.
- Hou, X., Zhu, B., Kumar, K. R., & Lu, W.: Inter-annual variability in fine particulate matter pollution over China during 2013–2018: Role of meteorology, *Atmospheric Environment*, 214, 116842, <https://doi.org/10.1016/j.atmosenv.2019.116842>, 2019.
- 495 Hu, J., Emile-Geay, J., & Partin, J.: Correlation-based interpretations of paleoclimate data—where statistics meet past climates, *Earth and Planetary Science Letters*, 459, 362-371, <https://doi.org/10.1016/j.epsl.2016.11.048>, 2017.
- Huang, W., Wang, B., & Wright, J. S.: A potential vorticity-based index for the East Asian winter monsoon, *Journal of Geophysical Research: Atmospheres*, 121, 9382–9399, <https://doi.org/10.1002/2016JD025053>, 2016.
- Huffman, G. J., Adler, R. F., Morrissey, M. M., Bolvin, D. T., Curtis, S., Joyce, R., ... & Susskind, J.: Global precipitation at 500
one-degree daily resolution from multisatellite observations, *Journal of hydrometeorology*, 2(1), 36-50, [https://doi.org/10.1175/1525-7541\(2001\)002<0036:GPAODD>2.0.CO;2](https://doi.org/10.1175/1525-7541(2001)002<0036:GPAODD>2.0.CO;2), 2001.
- Jeong, J. I., & Park, R. J.: Winter monsoon variability and its impact on aerosol concentrations in East Asia, *Environmental Pollution*, 221, 285-292, <https://doi.org/10.1016/j.envpol.2016.11.075>, 2017.
- Jhun, J. G., & Lee, E. J.: A new East Asian winter monsoon index and associated characteristics of the winter 505
monsoon, *Journal of Climate*, 17(4), 711-726, [https://doi.org/10.1175/1520-0442\(2004\)017<0711:ANEAWM>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<0711:ANEAWM>2.0.CO;2), 2004.
- Jia, B., Wang, Y., Yao, Y., & Xie, Y.: A new indicator on the impact of large-scale circulation on wintertime particulate matter pollution over China, *Atmospheric Chemistry and Physics*, 15(20), 11919-11929, <https://doi.org/10.5194/acp-15-11919-2015>, 2015.
- 510 Kong, L., Tang, X., Zhu, J., Wang, Z., Li, J., Wu, H., ... & Carmichael, G. R.: A 6-year-long (2013–2018) high-resolution air quality reanalysis dataset in China based on the assimilation of surface observations from CNEMC, *Earth System Science Data*, 13(2), 529-570, <https://doi.org/10.5194/essd-13-529-2021>, 2021.
- Koch, D., Park, J., & Del Genio, A.: Clouds and sulfate are anticorrelated: A new diagnostic for global sulfur models, *Journal of Geophysical Research: Atmospheres*, 108(D24), <https://doi.org/10.1029/2003JD003621>, 2003.
- 515 Leung, D. M., Tai, A. P., Mickley, L. J., Moch, J. M., Donkelaar, A. V., Shen, L., & Martin, R. V.: Synoptic meteorological modes of variability for fine particulate matter (PM_{2.5}) air quality in major metropolitan regions of China, *Atmospheric Chemistry and Physics*, 18(9), 6733-6748, <https://doi.org/10.5194/acp-18-6733-2018>, 2018.
- Li, Y., & Yang, S.: A dynamical index for the East Asian winter monsoon, *Journal of Climate*, 23(15), 4255-4262, <https://doi.org/10.1175/2010JCLI3375.1>, 2010.



- 520 Li, M., Huang, X., Zhu, L., Li, J., Song, Y., Cai, X., & Xie, S.: Analysis of the transport pathways and potential sources of
PM₁₀ in Shanghai based on three methods, *Science of the Total Environment*, 414, 525-534,
<https://doi.org/10.1016/j.scitotenv.2011.10.054>, 2012.
- Li, K., Jacob, D. J., Liao, H., Zhu, J., Shah, V., Shen, L., ... & Zhai, S.: A two-pollutant strategy for improving ozone and
particulate air quality in China, *Nature Geoscience*, 12(11), 906-910, <https://doi.org/10.1038/s41561-019-0464-x>,
525 2019.
- Pei, L., Yan, Z., Sun, Z., Miao, S., & Yao, Y.: Increasing persistent haze in Beijing: potential impacts of weakening East
Asian winter monsoons associated with northwestern Pacific sea surface temperature trends, *Atmospheric
Chemistry and Physics*, 18(5), 3173-3183, <https://doi.org/10.5194/acp-18-3173-2018>, 2018.
- Shen, L., Mickley, L. J., & Tai, A. P.: Influence of synoptic patterns on surface ozone variability over the eastern United
530 States from 1980 to 2012, *Atmospheric Chemistry and Physics*, 15(19), 10925-10938, <https://doi.org/10.5194/acp-15-10925-2015>, 2015.
- Sun, B. M., & Li, C. Y.: Relationship between the disturbances of East Asian trough and tropical convective activities in
boreal winter, *Chin. Sci. Bull.*, 42, 500-504, <https://doi.org/10.1360/csb1997-42-5-500>, 1997.
- Sun, Y., Wang, Z., Fu, P., Jiang, Q., Yang, T., Li, J., & Ge, X.: The impact of relative humidity on aerosol composition and
535 evolution processes during wintertime in Beijing, China, *Atmospheric Environment*, 77, 927-934,
<https://doi.org/10.1016/j.atmosenv.2013.06.019>, 2013.
- Sun, J., Li, H., Zhang, W., Li, T., Zhao, W., Zuo, Z., et al.: Modulation of the ENSO on winter aerosol pollution in the
eastern region of China, *Journal of Geophysical Research: Atmospheres*, 123, 11,952-11,969,
<https://doi.org/10.1029/2018JD028534>, 2018.
- 540 Tai, A. P., Mickley, L. J., & Jacob, D. J.: Correlations between fine particulate matter (PM_{2.5}) and meteorological variables
in the United States: Implications for the sensitivity of PM_{2.5} to climate change, *Atmospheric environment*, 44(32),
3976-3984, <https://doi.org/10.1016/j.atmosenv.2010.06.060>, 2010.
- Tai, A. P., Mickley, L. J., Jacob, D. J., Leibensperger, E. M., Zhang, L., Fisher, J. A., & Pye, H. O. T.: Meteorological modes
of variability for fine particulate matter (PM_{2.5}) air quality in the United States: implications for PM_{2.5} sensitivity to
545 climate change, *Atmospheric Chemistry and Physics*, 12(6), 3131-3145, <https://doi.org/10.5194/acp-12-3131-2012>,
2012.
- Wang, F. K.: Confidence interval for the mean of non-normal data, *Quality and Reliability Engineering International*, 17(4),
257-267, <https://doi.org/10.1002/qre.400>, 2001.
- Wang, L., & Chen, W.: How well do existing indices measure the strength of the East Asian winter monsoon?, *Advances in
550 Atmospheric Sciences*, 27(4), 855-870, <https://doi.org/10.1007/s00376-009-9094-3>, 2010.
- Wang, L., & Lu, M. M.: The East Asian winter monsoon, In *The global monsoon system: research and forecast* (pp. 51-61),
https://doi.org/10.1142/9789813200913_0005, 2017.



- Wang, W., Primbs, T., Tao, S., & Simonich, S. L. M.: Atmospheric particulate matter pollution during the 2008 Beijing Olympics, *Environmental science & technology*, 43(14), 5314-5320, <https://doi.org/10.1021/es9007504>, 2009.
- 555 Wang, L., Xu, J., Yang, J., Zhao, X., Wei, W., Cheng, D., ... & Su, J.: Understanding haze pollution over the southern Hebei area of China using the CMAQ model, *Atmospheric Environment*, 56, 69-79, <https://doi.org/10.1016/j.atmosenv.2012.04.013>, 2012.
- Wang, B., Wu, R., & Fu, X.: Pacific–East Asian teleconnection: how does ENSO affect East Asian climate?, *Journal of Climate*, 13(9), 1517-1536, [https://doi.org/10.1175/1520-0442\(2000\)013<1517:PEATHD>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<1517:PEATHD>2.0.CO;2), 2000.
- 560 Wang, X., Wang, K., & Su, L.: Contribution of atmospheric diffusion conditions to the recent improvement in air quality in China, *Scientific reports*, 6(1), 1-11, <https://doi.org/10.1038/srep36404>, 2016.
- Wang, L., Zhang, N., Liu, Z., Sun, Y., Ji, D., & Wang, Y.: The influence of climate factors, meteorological conditions, and boundary-layer structure on severe haze pollution in the Beijing–Tianjin–Hebei region during January 2013, *Advances in Meteorology*, 2014, 685971, <https://doi.org/10.1155/2014/685971>, 2014.
- 565 Wang, X., Dickinson, R. E., Su, L., Zhou, C., & Wang, K.: PM_{2.5} pollution in China and how it has been exacerbated by terrain and meteorological conditions, *Bulletin of the American Meteorological Society*, 99(1), 105-119, <https://doi.org/10.1175/BAMS-D-16-0301.1>, 2018.
- Wang, L., Wang, H., Liu, J., Gao, Z., Yang, Y., Zhang, X., ... & Huang, M.: Impacts of the near-surface urban boundary layer structure on PM_{2.5} concentrations in Beijing during winter, *Science of The Total Environment*, 669, 493-504, <https://doi.org/10.1016/j.scitotenv.2019.03.097>, 2019a.
- 570 Wang, C., An, X., Zhang, P., Sun, Z., Cui, M., & Ma, L.: Comparing the impact of strong and weak East Asian winter monsoon on PM_{2.5} concentration in Beijing, *Atmospheric Research*, 215, 165-177, <https://doi.org/10.1016/j.atmosres.2018.08.022>, 2019b.
- Wu, B., & Wang, J.: Winter arctic oscillation. Siberian High and East Asian Monsoon, *geophysical research letters*, 29(4), <https://doi.org/10.1029/2002GL015373>, 2002.
- 575 Xu, P., Chen, Y., & Ye, X.: Haze, air pollution, and health in China, *The Lancet*, 382(9910), 2067, [https://doi.org/10.1016/S0140-6736\(13\)62693-8](https://doi.org/10.1016/S0140-6736(13)62693-8), 2013.
- Yang, S., Lau, K. M., & Kim, K. M.: Variations of the East Asian jet stream and Asian–Pacific–American winter climate anomalies, *Journal of Climate*, 15(3), 306-325, [https://doi.org/10.1175/1520-0442\(2002\)015<0306:VOTEAJ>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<0306:VOTEAJ>2.0.CO;2), 2002.
- 580 Yang, Y., Liu, X., Qu, Y., Wang, J., An, J., Zhang, Y., & Zhang, F.: Formation mechanism of continuous extreme haze episodes in the megacity Beijing, China, in January 2013, *Atmospheric Research*, 155, 192-203, <https://doi.org/10.1016/j.atmosres.2014.11.023>, 2015.
- Zhang, Q., Streets, D. G., He, K., & Klimont, Z.: Major components of China’s anthropogenic primary particulate emissions, *Environmental Research Letters*, 2(4), 045027, <https://doi.org/10.1088/1748-9326/2/4/045027>, 2007.
- 585



- Zhang, R., Min, Q., & Su, J.: Impact of El Niño on atmospheric circulations over East Asia and rainfall in China: Role of the anomalous western North Pacific anticyclone, *Science China Earth Sciences*, 60(6), 1124-1132, <https://doi.org/10.1007/s11430-016-9026-x>, 2017.
- 590 Zhang, R. H., Li, Q., & Zhang, R.: Meteorological conditions for the persistent severe fog and haze event over eastern China in January 2013, *Science China Earth Sciences*, 57(1), 26-35, <https://doi.org/10.1007/s11430-013-4774-3>, 2014.
- Zhang, G., Gao, Y., Cai, W., Leung, L. R., Wang, S., Zhao, B., ... & Gao, H.: Seesaw haze pollution in North China modulated by the sub-seasonal variability of atmospheric circulation, *Atmospheric Chemistry and Physics*, 19(1), 565-576, <https://doi.org/10.5194/acp-19-565-2019>, 2019.
- 595 Zhao, X. J., Zhao, P. S., Xu, J., Meng, W., Pu, W. W., Dong, F., ... & Shi, Q. F.: Analysis of a winter regional haze event and its formation mechanism in the North China Plain, *Atmospheric Chemistry and Physics*, 13(11), 5685-5696, <https://doi.org/10.5194/acp-13-5685-2013>, 2013.
- Zheng, S., Pozzer, A., Cao, C. X., & Lelieveld, J.: Long-term (2001–2012) concentrations of fine particulate matter (PM_{2.5}) and the impact on human health in Beijing, China, *Atmospheric Chemistry and Physics*, 15(10), 5715-5725, <https://doi.org/10.5194/acp-15-5715-2015>, 2015.
- 600 Zhu, J., Liao, H., & Li, J.: Increases in aerosol concentrations over eastern China due to the decadal-scale weakening of the East Asian summer monsoon, *Geophysical Research Letters*, 39(9), <https://doi.org/10.1029/2012GL051428>, 2012.

605

610

615




Figure 1: Spatial correlation coefficients of daily mean $\text{PM}_{2.5}$ concentrations with those in (a) Beijing, (b) Shanghai and (c) Guangzhou during DJF 2013–2017. Regions where correlations are higher than 0.7 are selected to represent the Beijing–Tianjin–Hebei (BTH), Yangtze River Delta (YRD) and Pearl River Delta (PRD) regions, separately.


620

625

630

635


640



645 **Figure 2: Lagged correlations between daily mean PM_{2.5} concentrations and regional meteorological variables including relative**
humidity (RH; red bars), wind speed (WSPD; green bars), vertical wind shear (WSHR; yellow bars) and inversion intensity (INV;
blue bars) over (a) BTH, (b) YRD and (c) PRD during DJF 2013–2017. Black horizontal dashed lines indicate the 95% confidence
level using the two-tailed Student’s *t*-test.

650


655



660 **Figure 3: Winter mean (a) sea level pressure (SLP; hPa), (b) 850-hPa meridional wind (V850; m s⁻¹), (c) 500-hPa geopotential height (Z500; m, shading) and 850 hPa wind (arrows), and (d) precipitation (mm day⁻¹) during DJF 2013–2017.**

665


670



675 **Figure 4: Average (a) SLP (hPa), (b) V850 (m s^{-1}), (c) Z500 (m, shading) and 850 hPa wind (m s^{-1} , vector) on heavily polluted days**
(24-h $\text{PM}_{2.5}$ above the regional 90th percentile), and difference (heavily polluted days minus winter mean) for (d) SLP, (e) V850, (f)
680 Z500 and 850 hPa wind during DJF 2013–2017 over BTH. For V850 (b, e), blue regions represent northerlies and red regions
represent southerlies. Dotted regions in panels d–f mark statistically significant differences at the 95% level (determined through a
bootstrap resampling method). Grey shading represents the BTH region.

680

685




690 **Figure 5:** Correlation coefficients of daily $\text{PM}_{2.5}$ concentrations in BTH with (a) SLP, (b) V850 and (c) Z500 during DJF 2013–2017
(dotted regions indicate significant correlations at the 95% level from the two-tailed Student's t -test). Grey shading represents the
BTH region. The broad region presenting the highest correlation with BTH is marked by a yellow rectangle in each panel. The
region used for the definition of a circulation-based index (eq. 3) is marked by a yellow thick rectangle in panel c.

695

700


705



710 **Figure 6:** Average (a) SLP (hPa), (b) V850 (m s^{-1}), (c) Z500 (m, shading) and 850 hPa wind (m s^{-1} , vector) one day before heavily polluted days (24-h $\text{PM}_{2.5}$ above the regional 90th percentile), and difference (one day before heavily polluted days minus winter mean) for (d) SLP, (e) V850, (f) Z500 and 850 hPa wind during DJF 2013–2017 over YRD. For V850 (b, e), blue regions represent northerlies and red regions represent southerlies. Dotted regions in panels d–f mark statistically significant differences at the 95% level (determined through a bootstrap resampling method). Grey shading represents the YRD region.

715

720



725 **Figure 7: Correlation coefficients of daily $PM_{2.5}$ concentrations in YRD with one day before (a) SLP, (b) V850 and (c) Z500 during**
730 **DJF 2013–2017 (dotted regions indicate significant correlations at the 95% level from the two-tailed Student’s t -test). Grey**
735 **shading represents the YRD region. The broad region presenting the highest correlation with YRD is marked by a yellow**
740 **rectangle in each panel. The region used for the definition of a circulation-based index (eq. 4) is marked by a yellow thick**
745 **rectangle in panel a.**


730

735


740

745

750



755 **Figure 8:** Average (a) SLP (hPa), (b) V850 (m s^{-1}) and (c) precipitation (mm day^{-1}) two days before heavily polluted days, and
760 difference (two days before heavily polluted days minus winter mean) for (d) SLP, (e) V850, (f) precipitation during DJF 2013–
765 2017 over PRD. For V850 (b, e), blue regions represent northerlies and red regions represent southerlies. Dotted regions in panels
4d–f mark statistically significant differences at the 95% level (determined through a bootstrap resampling method). Grey shading
represents the PRD region.



770 **Figure 9: Correlation coefficients of daily $PM_{2.5}$ concentrations in PRD with two days before (a) SLP, (b) V850 and (c)**
precipitation during DJF 2013–2017 (dotted regions indicate significant correlations at the 95% level from the two-tailed Student's
***t*-test). Grey shading represents the PRD region. The broad region presenting the highest correlation with PRD is marked by a**
yellow rectangle in each panel. The region used for the definition of a circulation-based index (eq. 5) is marked by a yellow thick
rectangle in panel b.


775

780

785

790


795



800 **Figure 10:** Frequency distributions of circulation-based indices for different percentile thresholds of daily mean $\text{PM}_{2.5}$
 concentrations over (a) BTH, (b) YRD and (c) PRD during DJF 2013–2017. $\text{PM}_{2.5}$ concentration data are lagged by one and two
 days with respect to the circulation indices for YRD and PRD, respectively. The vertical lines and shading represent the averages
 and the associated 95% confidence intervals, respectively. Averages are calculated using Tukey's trimean (e.g., Ge et al., 2019):
 $\bar{X} = \frac{1}{4}(Q1 + 2Q2 + Q3)$, where $Q1$ is the lower quartile, $Q2$ is the median, and $Q3$ is the upper quartile. The confidence intervals
 805 for these averages are estimated by using bootstrap resampling (e.g., Wang, 2001). This method generates samples by randomly
 choosing daily values of circulation-based indices (resampling with replacement) and then calculating the Tukey's trimean. This
 process is repeated 10,000 times to get robust replicates of the mean. Ultimately, the lower and upper limits of the 95% confidence
 intervals are calculated as the values corresponding to the 2.5th and 97.5th percentiles.

810

815



820 **Figure 11:** Joint distributions of circulation-based indices against detrended daily $PM_{2.5}$ concentrations for different percentile thresholds (colour coded), including the corresponding linear fits with 95% prediction intervals, over (a) BTH, (b) YRD and (c) PRD during DJF 2013–2017. $PM_{2.5}$ concentrations data are lagged by one and two days behind the circulation indices in the case of YRD and PRD, respectively.

825

830

835

840



845

Table 1: Correlation of the area-weighted averages of daily normalized circulation variables over the regions marked by yellow rectangles in Figures 5, 7 and 9 with daily PM_{2.5} concentrations over BTH, YRD and PRD, respectively, during DJF 2013–2017. All correlation values are significant at the 99% confidence level. The highest correlation (absolute value) for each region is shown in bold.

Correlation coefficient (450 days)	V850	Z500	SLP	Precipitation
PM _{2.5} (BTH)	0.59	0.67	0.54	
PM _{2.5} (YRD)	0.25	0.21	-0.33	
PM _{2.5} (PRD)	-0.43		0.36	-0.29

850

855

860

865

870

875

880

885



Table 2: Correlation of circulation-based indices defined in this study (equations 3–5) with daily PM_{2.5} concentrations over BTH, YRD, PRD, and with the most important regional meteorological variable in each region during DJF 2013–2017. All correlations are significant at the 99% confidence level.

Correlation coefficient (450 days)	<i>I</i>_{Z500_BTH}	<i>I</i>_{SLP_YRD}	<i>I</i>_{V850_PRD}
PM_{2.5}	-0.67	-0.33	-0.43
RH	-0.64		0.64
WSPD		0.29	

890

895

900

905

910

915

920

925



930 **Table 3: Linear relationship (explained variance) of daily PM_{2.5} concentrations in BTH, YRD and PRD, with the circulation-based**
index of Table 2, the most important regional meteorological field in each region and the linear combination of both during DJF
2013–2017. All the linear relationships are significant at the 99% confidence level.

935

BTH	<i>I</i>_{Z500_BTH}	RH	<i>I</i>_{Z500_BTH + RH}
<i>R</i>²	0.45	0.44	0.54
YRD	<i>I</i>_{SPL_YRD}	WSPD	<i>I</i>_{SPL_YRD + WSPD}
<i>R</i>²	0.11	0.18	0.23
PRD	<i>I</i>_{V850_PRD}	RH	<i>I</i>_{V850_PRD + RH}
<i>R</i>²	0.18	0.27	0.30

940

945