<u>Comments for all Reviewers:</u> Thank you for taking the time to review this manuscript and for providing helpful and thoughtful comments. These suggestions have helped improve the quality of the manuscript. Below are responses (in green text) to the reviewer comments (in black text).

Response to Reviewer 1:

This manuscript describes CI-TOF-MS measurements of gaseous sulfur compounds emitted during a controlled, induced bloom experiment. The authors sought to determine if additional sulfur compounds, besides DMS, are emitted to the atmosphere, as well as the relative amounts. This is important for aerosol and cloud formation in the marine atmosphere. The manuscript reads well and the results are robust. The manuscript should be published after the following comments have been addressed.

Specific comments:

Lines 95-96: Missing current Lennartz references. For example (this alone might be good enough to reference):

Lennartz, S. T., Marandino, C. A., von Hobe, M., Andreae, M. O., Aranami, K., Atlas, E., Berkelhammer, M., Bingemer, H., Booge, D., Cutter, G., Cortes, P., Kremser, S., Law, C. S., Marriner, A., Simó, R., Quack, B., Uher, G., Xie, H., and Xu, X.: Marine carbonyl sulfide (OCS) and carbon disulfide (CS₂): a compilation of measurements in seawater and the marine boundary layer, Earth Syst. Sci. Data, 12, 591–609, https://doi.org/10.5194/essd-12-591-2020, 2020.

Authors' response: Thank you for bringing this to our attention. We have added the suggested Lennartz reference to line 93 of the revised manuscript.

Section 2.3: is there a reason why the authors chose not to use an equilibrator?

Authors' response: The authors thank the reviewer for this comment. Headspace equilibrators, such as those coupled to PTR-MS instruments like in Wohl et al. (2019) are a great way to assess dissolved VOC concentrations in seawater. For the SeaSCAPE experiment, multiple analytical goals made the usage of cryo purge and trap preferable, namely measurements of dissolved and particulate DMSP which required overnight reactions to yield measurable DMS. In addition, we sought to follow established protocols of dissolved DMS measurements, requiring close attention to water agitation which can disrupt phytoplankton cell membranes and can increase DMS concentrations significantly. For these reasons, purge and trap was chosen, however the suggestion to use an equilibrator is appreciated and will be strongly considered for future experimentation.

Figure 1b: are the individual peak shapes determined by injection of individual standards?

Authors' response: The individual peak shape for DMS was determined by a compressed gas standard. Other individual peak shapes in Figure 1b were determined by the processing software (Tofware v3.1.2, TOFWERK). Inputs into the software are a mass calibration, defined peak shape, width, and baseline, and a reference mass spectrum. Peaks are then found using a smoothed second-derivative approach and peak fits are applied.

Line 191: how were LODs determined?

Authors' response: Limits of detection were determined based on equation 1 in Bertram et al. (2011) for a signal-to-noise ratio of 3. This information has now been included in line 189-191, reading: "Limits of detection for DMS, MeSH, and benzothiazole at 1 minute averaging time were 0.20 ± 0.49 ppt, 1.5 ± 0.25 ppt, and 0.42 ± 0.14 ppt, respectively, calculated according to equation 1 in Bertram et al. (2011) for a signal-to-noise ratio of 3."

Line 201: why does MeSH have a water vapor dependence, but DMS does not? Have the authors considered an isotope dilution method to solve this sensitivity problem during the measurement?

Authors' response: We think that the water vapor dependence observed for MeSH and not observed for DMS could be, at least in part, due to conversion on inlet surfaces. This has been observed previously in MeSH measurements (Perraud et al., 2016). We have used isotope dilution methods in past field studies and agree this could be a useful method to confirm our calibrations from this experiment going forward.

Lines 212-220: Did the authors try using Teflon fittings? And Sulfinert® coated metals?

Authors' response: We used a mixture of both Teflon and stainless steel fittings connecting the ISV to the Vocus inlet during SeaSCAPE. Stainless steel was used on the ISV because the ISV was used for both gas and aerosol sampling. We recognize that for experiments only measuring gases, we would benefit from a fully Teflon inlet manifold. We did not try Sulfinert® coated metals in these experiments.

Line 284: I am not sure why this should be surprising if you don't have oxidation products – even if MSAM is outgassing, if no DMSO2 is formed then why would the authors expect a correlation?

Authors' response: The intention here was to suggest that given MSAM was a large emission in the Arabian Sea, especially relative to DMS, and was proposed to be formed from biogenic outgassing and not an oxidation product (Edtbauer et al., 2020), we were surprised to not see a

more significant MSAM emission. This suggests that MSAM was formed via a mechanism active in the Arabian Sea, but not in our study using coastal seawater from La Jolla, California.

We have changed the text from "Methane sulfonamide (MSAM) measured on average less than 1 ppt during the experiment and showed no positive correlation with the DMS oxidation product, DMSO₂ (Fig. S6). This is surprising as MSAM was recently measured at mixing ratios of up to 60 ppt, corresponding to up to 33% of DMS, in upwelling areas of the Arabian Sea and was correlated with DMSO₂ (r = 0.8) (Edtbauer et al., 2020)." to "Methane sulfonamide (MSAM) measured on average less than 1 ppt during the experiment. In contrast, Edtbauer et al. (2020) measured MSAM at mixing ratios of up to 60 ppt, corresponding to up to 33% of DMS, in upwelling areas of the Arabian Sea and suggested MSAM had a direct oceanic emission source. Results from our study suggest that the pathway producing MSAM in the Arabian Sea was not active in our experiments utilizing coastal seawater from Southern California."

Lines 304-306: this is only true if the atmospheric loss is the same or negligible. Is it true here that it is assumed to be negligible?

Authors' response: Since we are measuring from a low-oxidant headspace, continuously purged with zero air, we are assuming atmospheric loss is negligible. We have updated the language in lines 300-303 to clarify this, reading "Given that the Henry's law constants and diffusion constants for DMS and MeSH are roughly the same, and assuming atmospheric loss is negligible in the ISV, we expect the measured molar ratio of DMS:MeSH in the headspace to reflect the dissolved concentration ratio in the seawater."

Line 363: why would the authors assume the lifetimes to be the same?

Authors' response: Thank you for this note. You bring up a good point that the lifetimes of DMSP and DMS in the water are variable across the ocean, as shown in Table 1 of Vila-Costa et al., (2008). We have removed from line 363 the text reading "assuming the lifetimes of DMS and DMSP_t in the water are the same" and instead have replaced it with our reasoning for using this DMS:DMSP_t ratio. We have added the following text: "Further, since we do not have direct measurements of DMSP and DMS cycling, we will use the waterside DMS:DMSP_t ratio to estimate the DMSP_t to DMS conversion efficiency in seawater (Galí et al., 2018, 2021)."

Lines 366-370: how does this explanation account for bacterial cleavage?

Authors' response: As written, lines 366-370 do not explicitly account for bacterial cleavage. In this section, we propose that the low waterside DMS:DMSP_t observed in this study is either a result of low DMS production from DMSP or a large DMS loss. Low DMS production from DMSP could be due to the phytoplankton community present in this study (diatom-dominant) or from reduced bacterial cleavage of DMSP to form DMS. Given the strong correlation between DMS:MeSH and methionine aminopeptidase activity (Fig. 5h), it is possible that protein

degradation provided the primary carbon and sulfur sources to bacteria and the DMSP to DMS conversion pathway was not the most important metabolism.

Line 426: when I look at Figure 5, it does not seem like bacterial sulfur demand is the most important determinant.

Authors' response: Thank you for this comment. While there is not as strong of a correlation between DMS:MeSH and bacterial sulfur demand as there is between DMS:MeSH and other variables plotted in Figure 5, we think that the temporal dynamics of bacterial sulfur demand in relation to available sulfur (and the composition of available sulfur sources) is what makes bacterial sulfur demand important for controlling DMS:MeSH. This can be seen in Fig. 6c where there are two stages of the experiment separated by the vertical dashed line, where the high DMS:MeSH values are observed in the latter stage when the estimated assimilated sulfur exceeds the bacterial sulfur demand. We recognize how difficult it is to parse out these complex relationships in large-scale experiments like this one and as such have changed the text in line 382 from "leading to a significant increase in measured DMS:MeSH" to "likely responsible for part of the significant DMS:MeSH increase."

Supplemental material, paragraph starting at line 109: since this correlation looks so good, I wonder if there is anything with which salinity correlates that could be interesting for the ratio (but was not tested against the ratio).

Authors' response: None of the waterside variables measured (FDOM, chlorophyll-a, dissolved oxygen, and water temperature) or other sulfur-containing ions correlate with DMS:MeSH above $R^2 = 0.55$. The strongest correlation observed is between DMS:MeSH and day of the bloom ($R^2 = 0.64$). As discussed in the SI, the salinity decline throughout the bloom was likely a result of additions of ultrapure water to the wave channel meant to maintain the water level, shown by the correlation between DMS:MeSH and time. While the correlation between DMS:MeSH and salinity is interesting, we do not think that the fluctuation in salinity was enough to cause the changes in DMS:MeSH, meaning the observation is reflective of a correlation and not causation (Salgado et al., 2014).

Response to Reviewer 2:

The paper entitled "Marine gas-phase sulphur emissions during an induced phytoplankton bloom" by Kilgour and co-workers presents wind tunnel measurements of the emissions of marine reduced sulphur species during phytoplankton blooms.

The paper is well written and discusses two important points.

Firstly, the authors report that during the pre-bloom period non-DMS species contribute significantly to the reduced sulphur budget. Methanethiol and benzothiazole were found to be the largest contributors to non-DMS sulphur emissions. The authors discuss the implications of this new finding for the marine sulphur budget, in particular the implications benzothiazole emission on new particle formation in marine environments.

Secondly the authors propose that the ratio DMS:MeSH is driven primarily by methionine aminopeptidase which catalyses cleavage of amino acids from proteins and peptides. The propose that salinity is not the main driving force for the DMS:MeSH ratio.

The paper falls within the scope of atmospheric chemistry and physics and is well written. I recommend the paper for acceptance subject to minor revisions.

Minor revisions suggested:

• The authors cite an unpublished work Franklin et al. 2021. I am not sure what the journal policy in this matter is, but it may be good if the paper can be deposited as a preprint in a repository so that it becomes accessible to the reader.

Authors' response: Franklin et al. (2021) has been accepted in the journal Environmental Science and Technology. We have updated the reference with the following doi: 10.1021/acs.est.1c04422

• The authors also cite and AGU abstract, which is not a peer reviewed source to support one of their points Kiene et al. 2017. This is not ideal. Have the authors checked whether the relevant data has been published since, possible under a different title or is available in some repository.

Authors' response: Thank you for noting this. We are not aware of where this data has been published and peer-reviewed and as a result, we have removed references to it in the manuscript.

• The Sander reference for Henry's law constants appears to be incomplete Atmos. Chem. Phys., 15, 4399–4981, 2015 doi:10.5194/acp-15-4399-2015

Authors' response: Thank you for catching this mistake. We have updated the reference to accurately cite this paper.

References

Bertram, T. H., Kimmel, J. R., Crisp, T. A., Ryder, O. S., Yatavelli, R. L. N., Thornton, J. A., Cubison, M. J., Gonin, M., and Worsnop, D. R.: A field-deployable, chemical ionization time-of-flight mass spectrometer, Atmospheric Meas. Tech., 4, 1471–1479, https://doi.org/10.5194/amt-4-1471-2011, 2011.

Edtbauer, A., Stönner, C., Pfannerstill, E. Y., Berasategui, M., Walter, D., Crowley, J. N., Lelieveld, J., and Williams, J.: A new marine biogenic emission: methane sulfonamide (MSAM), dimethyl sulfide (DMS), and dimethyl sulfone (DMSO<sub>2</sub>) measured in air over the Arabian Sea, Atmospheric Chem. Phys., 20, 6081–6094, https://doi.org/10.5194/acp-20-6081-2020, 2020.

Galí, M., Levasseur, M., Devred, E., Simó, R., and Babin, M.: Sea-surface dimethylsulfide (DMS) concentration from satellite data at global and regional scales, Biogeosciences, 15, 3497–3519, https://doi.org/10.5194/bg-15-3497-2018, 2018.

Galí, M., Lizotte, M., Kieber, D. J., Randelhoff, A., Hussherr, R., Xue, L., Dinasquet, J., Babin, M., Rehm, E., and Levasseur, M.: DMS emissions from the Arctic marginal ice zone, Elem. Sci. Anthr., 9, 00113, https://doi.org/10.1525/elementa.2020.00113, 2021.

Perraud, V., Meinardi, S., Blake, D. R., and Finlayson-Pitts, B. J.: Challenges associated with the sampling and analysis of organosulfur compounds in air using real-time PTR-ToF-MS and offline GC-FID, Atmospheric Meas. Tech., 9, 1325–1340, https://doi.org/10.5194/amt-9-1325-2016, 2016.

Salgado, P., Kiene, R., Wiebe, W., and Magalhães, C.: Salinity as a regulator of DMSP degradation in Ruegeria pomeroyi DSS-3, J. Microbiol., 52, 948–954, https://doi.org/10.1007/s12275-014-4409-1, 2014.

Vila-Costa, M., Kiene, R. P., and Simí, R.: Seasonal variability of the dynamics of dimethylated sulfur compounds in a coastal northwest Mediterranean site, Limnol. Oceanogr., 53, 198–211, https://doi.org/10.4319/lo.2008.53.1.0198, 2008.

Wohl, C., Capelle, D., Jones, A., Sturges, W. T., Nightingale, P. D., Else, B. G. T., and Yang, M.: Segmented flow coil equilibrator coupled to a proton-transfer-reaction mass spectrometer for measurements of a broad range of volatile organic compounds in seawater, Ocean Sci., 15, 925–940, https://doi.org/10.5194/os-15-925-2019, 2019.

Marine gas-phase sulfur emissions during an induced phytoplankton bloom

Delaney B. Kilgour¹, Gordon A. Novak¹, Jon S. Sauer², Alexia N. Moore², Julie Dinasquet³, Sarah Amiri², Emily B. Franklin⁴, Kathryn Mayer², Margaux Winter⁵, Clare K. Morris³, Tyler Price², Francesca
Malfatti^{3,6}, Daniel R. Crocker², Christopher Lee³, Christopher D. Cappa⁷, Allen H. Goldstein^{4,8}, Kimberly A. Prather^{2,3}, and Timothy H. Bertram¹

¹Department of Chemistry, University of Wisconsin – Madison, Madison, WI 53706, USA
²Department of Chemistry and Biochemistry, University of California, San Diego, La Jolla, CA 92093, USA
³Scripps Institution of Oceanography, University of California, San Diego, La Jolla, CA 92037, USA
⁴Department of Civil and Environmental Engineering, University of California, Berkeley, CA 94720, USA
⁵Department of Chemistry and Chemical Biology, Harvard University, Cambridge, MA 02138, USA
⁶Department of Life Sciences, Universita' degli Studi di Trieste, Trieste, Italy 34127
⁷Department of Civil and Environmental Engineering, University of California, Davis, CA 95616, USA
⁸Department of Environmental Science, Policy and Management, University of California, Berkeley, CA 94720, USA

15 *Correspondence to*: Timothy H. Bertram (timothy.bertram@wisc.edu)

Abstract. The oxidation of dimethyl sulfide (DMS; CH₃SCH₃), emitted from the surface ocean, contributes to the formation of Aitken mode particles and their growth to cloud condensation nuclei (CCN) sizes in remote marine environments. It is not clear whether other, less commonly measured marine-derived, sulfur-containing gases share similar dynamics to DMS and contribute to secondary marine aerosol formation. Here, we present measurements of gas-phase volatile organosulfur

- 20 molecules taken with a Vocus proton transfer reaction high resolution time-of-flight mass spectrometer during a mesocosm phytoplankton bloom experiment using coastal seawater. We show that DMS, methanethiol (MeSH; CH₃SH), and benzothiazole (C_7H_5NS) account for on average over 90% of total gas-phase sulfur emissions, with non-DMS sulfur sources representing 36.8 ± 7.7% of sulfur emissions during the first nine days of the experiment in the pre-bloom phase prior to major biological growth, before declining to 14.5 ± 6.0% in the latter half of the experiment when DMS dominates during the bloom
- and decay phases. The molar ratio of DMS to MeSH during the pre-bloom phase (DMS:MeSH = 4.60 ± 0.93) was consistent with the range of previously calculated ambient DMS to MeSH sea-to-air flux ratios. As the experiment progressed, the DMS to MeSH emission ratio increased significantly, reaching 31.8 ± 18.7 during the bloom and decay. Measurements of dimethylsulfoniopropionate (DMSP), heterotrophic bacteria, and enzyme activity in the seawater suggest the DMS:MeSH ratio is a sensitive indicator of the bacterial sulfur demand and the composition and magnitude of available sulfur sources in
- 30 seawater. The evolving DMS:MeSH ratio and the emission of a new aerosol precursor gas, benzothiazole, have important implications for secondary sulfate formation pathways in coastal marine environments.

1 Introduction

The ocean accounts for the largest natural source of sulfur to the atmosphere, primarily as dimethyl sulfide (DMS; CH₃SCH₃)

- 35 (Andreae, 1990; Simó, 2001). Current estimates for oceanic DMS emissions range between 17.6–34.4 Tg S yr⁻¹ (Lana et al., 2011), compared to anthropogenic DMS emission estimates of 2.20 Tg S yr⁻¹ (Lee and Brimblecombe, 2016). DMS has been shown to impact the production rate of secondary marine aerosol (SMA), the concentration of cloud condensation nuclei (CCN), and Earth's radiation budget by altering cloud properties (Bates et al., 1992; Carpenter et al., 2012; Charlson et al., 1987; Lana et al., 2011).
- 40

DMS is primarily produced in seawater following the bacterial cleavage of the algal metabolite dimethylsulfoniopropionate (DMSP) (Challenger and Simpson, 1948). DMSP is present in both particulate (DMSP_p) and dissolved (DMSP_d) forms, where DMSP_p consists of phytoplankton intracellular DMSP, and DMSP_d consists of the dissolved pool in the seawater (Kiene and Linn, 2000a). DMSP_p concentrations in coastal seawater span a large range, from 5 to >300 nM, dependent on bloom dynamics,

- whereas DMSP_d is often present in lower concentrations (1-25 nM) and has a turnover rate of 1-129 nM d⁻¹ (Kiene et al., 2000; Kiene and Linn, 2000a). During blooms of DMSP-rich phytoplankton and in some colder waters, total DMSP (DMSP_t; DMSP_d + DMSP_p) can significantly exceed these ranges (Kiene et al., 2019; Kiene and Linn, 2000a; Kwint and Kramer, 1996). Isotopic labeling experiments using the ³⁵S isotope show wide variability in the DMS yield from DMSP_d (3-50%) (Carpenter et al., 2012), but the yield is commonly estimated as 10% (Kiene and Linn, 2000a). This results in waterside DMS concentrations in
- 50 the range of 1–7 nM globally, with higher values in the summer and in bloom conditions (Kiene et al., 2000; Kiene and Linn, 2000b; Lana et al., 2011). Once produced, DMS in seawater can be transformed by bacterial or photochemical processes, or converted to non-volatile sulfur, resulting in a DMS lifetime on the order of a few days in seawater (Flöck and Andreae, 1996; Kiene et al., 2000; Kiene and Linn, 2000b; Lawson et al., 2020). Approximately 10% of the dissolved DMS ventilates to the atmosphere, where it is oxidized by the hydroxyl radical (OH), halogen radicals (Cl and BrO), and nitrate radical (NO₃) to
- 55 form lower volatility products, including sulfur dioxide (SO₂) with yields ranging between 30% and 100%, and methanesulfonic acid (MSA) (Carpenter et al., 2012; Chen et al., 2018; Faloona, 2009; Lawson et al., 2020). Atmospheric SO₂ is further oxidized to sulfuric acid (H₂SO₄) and sulfate (SO₄²⁻), which can lead to new particle formation, while MSA primarily contributes to particle growth (Carpenter et al., 2012).
- 60 Methanethiol (MeSH; CH₃SH) has also been observed in marine environments, although there are fewer measurements compared to DMS and the MeSH emission rate is thought to be a small fraction of the DMS emission rate. MeSH is the major DMSP_d product (~75% yield) and is formed when bacteria demethylate or demethiolate DMSP_d (Kiene, 1996; Kiene and Linn, 2000b). However, existing measurements of dissolved MeSH concentrations are significantly smaller than collocated dissolved DMS concentrations (<1.8 nM versus <6 nM) (Kettle et al., 2001). This is a result of its reaction with dissolved organic matter
- and its rapid incorporation into bacterial cells where it is used to form methionine (Flöck and Andreae, 1996; Kiene, 1996;

Kiene et al., 1999). This leads to a dissolved MeSH lifetime on the order of minutes to an hour, which is considerably shorter than that of DMS (Lawson et al., 2020). Although the fraction of MeSH that ventilates to the atmosphere is poorly constrained, it serves as an additional source of reduced sulfur to the marine atmosphere and has a faster reaction rate with OH at 298 K ($3.3 \times 10^{-11} \text{ cm}^3$ molecules⁻¹s⁻¹) compared to that of DMS with OH at 298 K ($4.8 \times 10^{-12} \text{ cm}^3$ molecules⁻¹s⁻¹ via H-abstraction

70 and 1.7 x 10⁻¹² cm³ molecules⁻¹s⁻¹ via OH addition) (Atkinson et al., 2004), suggesting MeSH could also contribute to marine boundary layer (MBL) SO₂ and sulfate aerosol.

The emission ratio of DMS to MeSH (DMS:MeSH) is a sensitive indicator of DMSP_t production and degradation pathways, as well as the lifetimes of DMS and MeSH in seawater. The cleavage pathway that produces DMS is in competition with the demethylation/demethiolation pathway that produces MeSH. The favored pathway is a function of both the concentration of DMSP_t and the bacterial sulfur demand (Carpenter et al., 2012; Kiene et al., 2000; Kiene and Linn, 2000a; Vila et al., 2004). Low bacterial sulfur demand and high DMSP_t concentrations promote DMS production, while high bacterial sulfur demand and high DMSP_t concentrations promote DMS production, while high bacterial sulfur demand and low DMSP_t concentrations promote MeSH production (Carpenter et al., 2012; Kiene et al., 2000; Kiene et al., 2000; Kiene and Linn, 2000a).

75

80 Fluctuations in the available sulfur pool and bacterial sulfur demand can translate to significant variability in waterside measurements of DMS:MeSH over the open ocean. In upwelling regions of the Atlantic Ocean, waterside DMS:MeSH has been shown to range between 1 and 30 (Kettle et al., 2001). Measurements made in the Baltic Sea, Kattegat/Skagerrak, and North Sea have shown waterside DMS:MeSH of 16, 20, and 6, respectively (Leck and Rodhe, 1991). More recently, measurements in the subarctic northeast Pacific Ocean showed waterside DMS:MeSH ranged between 2 - 5.3 resulting in a calculated average DMS:MeSH sea to air flux ratio of 6 (Kiene et al., 2017). In the southwest Pacific Ocean, the reported DMS:MeSH flux ratio varied between 3 and 7 as estimated by the nighttime concentration accumulation method (Lawson et al.)

al., 2020). The Henry's law constants and diffusion constants in water at 298 K for DMS (5.6 × 10⁻³ mol m⁻³ Pa⁻¹; 1.217 × 10⁻⁵ cm² s⁻¹) and MeSH (3.8 × 10⁻³ mol m⁻³ Pa⁻¹; 1.556 × 10⁻⁵ cm² s⁻¹) are similar (Gharagheizi, 2012; Sander, 2015), implying the dissolved concentration ratio in the seawater is directly related to the emission ratio. Periods of low DMS:MeSH suggest that
MeSH could impact oxidative capacity of the MBL by providing a significant source of reduced sulfur to the atmosphere.

Other sulfur species, including dimethyl disulfide, carbon disulfide, and carbonyl sulfide, have previously been measured in the seawater in highly productive regions, though in significantly smaller quantities than DMS (Kettle et al., 2001; Leck and Rodhe, 1991; Lennartz et al., 2020). Recently, a previously unobserved biogenic marine volatile sulfur molecule, methane

95 sulfonamide, was measured in the gas-phase near an upwelling region of the Arabian Sea at mixing ratios up to 33% of DMS (Edtbauer et al., 2020). Additionally, the recent discovery of the DMS oxidation product hydroperoxymethyl thioformate (HPMTF) has prompted researchers to reexamine our understanding of the sulfur cycle (Veres et al., 2020). The combination of these findings raises questions regarding whether organosulfur molecules emitted in smaller quantities than DMS are important to the sulfur budget and contribute to sulfate aerosol and CCN in the marine atmosphere.

Here we report measurements of gas-phase volatile organosulfur molecules, with specific focus on DMS, MeSH, and a marine sulfur-containing molecule not reported prior to this experiment, benzothiazole (Franklin et al., 2021). These measurements were made during a mesocosm bloom experiment in a low-oxidant wave channel at the Scripps Institution of Oceanography in La Jolla, California. We examine how the distribution of emitted gas-phase sulfur molecules evolves as a function of bloom stage and provide insight into biological and environmental controls on the production and loss processes of these gases.

105

2 Methods

2.1 Scripps Institution of Oceanography Wave Channel and Mesocosm Experiment

The experiment was conducted at the Scripps Institution of Oceanography Hydraulics Laboratory wave channel as part of the Center for Aerosol Impacts on Chemistry of the Environment's intensive Sea Spray Chemistry And Particle Evolution (SeaSCAPE) experiment in July and August 2019. The collaborative study aimed to determine the impacts of biological activity, oxidative aging, and photochemistry on the emission of marine trace gases, the production of nascent sea spray aerosol, and the composition of secondary marine aerosol. Here, we present analysis of gas-phase sulfur species from the third of three phytoplankton blooms. This part of the mesocosm experiment lasted 21 days, where day 0 marks the time when the wave channel was filled with seawater from the Pacific Ocean, pumped directly from below Ellen Browning Scripps Memorial

Pier (herein Scripps Pier) in La Jolla, CA. Details of the wave channel setup and wave-breaking mechanism have been described elsewhere (Prather et al., 2013) and additional detail on this collaborative study is provided in the supplementary information (S1). A phytoplankton bloom was induced through a series of f/2 and f/20 growth medium and silicate additions. Details and timing of nutrient additions and perturbations to the mesocosm system are in the supplementary information (S2 and Table S1).

125

Due to the large volume of the wave channel, it is challenging to fully clean the headspace of background trace gases. As a result, all gas-phase measurements were made from an isolated sampling vessel (ISV) (Fig. S1) that circulated wave channel seawater using a peristaltic pump, providing a water residence time of 29 minutes. Its headspace was continuously purged with zero air (air residence time of 5 minutes) from a zero air generator (Sabio 1001), providing a headspace with low NOx, O₃, and background VOC. For this work, the ISV was sampled at 100 standard cubic centimeters per minute (sccm) through an approximately 2.5 m 0.25" O.D. PFA tube. While the ISV and wave channel were illuminated with fluorescent lights during

approximately 2.5 m 0.25" O.D. PFA tube. While the ISV and wave channel were illuminated with fluorescent lights during gas-phase measurements, these do not mimic the solar spectrum reaching the ocean's surface, providing a key difference between this work and referenced work studying gaseous emissions in the ambient environment.

¹²⁰

2.2 PTR-ToF-MS Measurements of Reduced Sulfur Compounds

- 130 A Vocus proton transfer reaction time-of-flight mass spectrometer (PTR-ToF-MS) (TOFWERK, Aerodyne, Inc.) was deployed to measure gas-phase volatile sulfur molecules. The Vocus instrument has a high resolving power ($m/\Delta m > 5000$) and 1–2 orders of magnitude improved sensitivity over prior low-pressure PTR-ToF-MS instruments, allowing detection of the subppt level gases observed in this study (Krechmer et al., 2018).
- Mass spectra were collected from 19–500 m/Q and saved at 1 Hz time resolution. Peak fitting and integration were completed in Tofware v3.1.2 (TOFWERK). The Vocus instrument parameters used in the study are as follows: The big segmented quadrupole (BSQ) voltage was 275 V, acting as a high-pass band filter to reduce the ion transmission of low mass (<35 m/Q) ions (Krechmer et al., 2018). The focusing ion-molecule reactor (FIMR) was operated at a high reduced field strength (E/N = 143 Td) with a pressure of 1.5 mbar, axial electric field gradient of 41.5 V cm⁻¹, and was heated to 100 °C. The high reduced field strength lessened reagent ion clustering and increased fragmentation of some ions.

Measurements of the ISV headspace were taken for approximately one hour at 9 am and one hour at 2 pm each day, and daily averages were calculated as the average over the total two-hour measurement period. Instrument background signals were determined approximately 8 times daily by overflowing the Vocus inlet with zero air from the same zero air generator that
provided air to the ISV headspace. Daily average background signals were used for background correction. Calibration factors for DMS and MeSH were determined by diluting a gas standard (5.08 ppm ± 5% DMS, Praxair; 6.111 ppm ± 5% MeSH, Airgas) into zero air. The benzothiazole calibration factor was measured using a syringe pump to inject dilute solutions of benzothiazole (96%, Sigma-Aldrich) in cyclohexane (99.5%, Sigma-Aldrich) into zero air carrier gas flow. The dry sensitivities to DMS, MeSH, and benzothiazole are 3.0, 1.0, and 5.8 cps ppt⁻¹, respectively. Other sulfur-containing species
(listed in Table S2) were quantified using the DMS sensitivity, as the proton transfer rate constant for DMS is similar to the proton transfer rate constants for other sulfur-containing species (Sekimoto et al., 2017). All molecules were identified and quantified by their protonated ion (MH⁺). Identifications of non-calibrated large mass species (>100 g mol⁻¹) in Table S2 were provided by thermal desorption two-dimensional gas chromatography coupled with electron ionization time-of-flight mass

155 2.3 Waterside Measurements at Wave Channel

spectrometry (TD-GCxGC-ToF-MS) (Franklin et al., 2021).

The following waterside variables were measured continuously for indication of bloom progression: fluorescent dissolved organic matter (FDOM) and chlorophyll-a (ECO-Triplet-BBFL2; Sea-Bird Scientific), dissolved oxygen and water temperature (SBE 63 Optical Dissolved Oxygen Sensor; Sea-Bird Scientific), and salinity (SBE 37 SI MicroCAT; Sea-Bird Scientific).

Bulk water samples collected from the wave channel were used for daily measurements of the following: heterotrophic bacteria abundance measured with flow cytometry (Gasol and Del Giorgio, 2000), bacterial productivity determined by radiolabeled leucine incorporation (Kirchman et al., 1985; Azam and Smith, 1992; Simon and Azam, 1989), phytoplankton enumeration determined by the Utermöhl method (Edler and Elbrächter, 2010) and dissolved DMS, DMSP_p, and DMSP_d measured by a

165 home-built purge and trap system (Wurl, 2009) coupled to a chemical ionization mass spectrometer with benzene cluster cation reagent ions (Fig. S2) (Lavi et al., 2018; Kim et al., 2016). More information on these methods is described in the supplementary information (S3 and S4).

3 Results and Discussion

3.1 Vocus PTR-ToF-MS Characterization of Organosulfur Molecules

- 170 Krechmer et al. (2018) previously characterized the Vocus performance in a lab setting and Li et al. (2020b, a) have described its abilities in forest sites. What follows is the first description of the instrument's capabilities for studying marine trace gases, which comprise a unique subset of VOCs that are often emitted in smaller quantities than in forest or urban environments. In this manuscript we focus on organosulfur molecules.
- Twenty-eight sulfur-containing ions were detected in the mass spectrum (Table S2). In addition to ions corresponding to the molecules DMS, MeSH, and benzothiazole, these included ions such as C₃H₆SH⁺, C₂H₆S₂H⁺, C₄H₈SO₂H⁺, C₁₀H₁₆SH⁺, and C₁₁H₁₆SH⁺. A sample mass spectrum highlighting sulfur-containing ions and the high-resolution fit around DMS is in Fig. 1. Several ions are detected at the unit masses of important marine gases (Fig. 1b), with ions at m/Q 49 (the m/Q corresponding to MeSH) including CClHH⁺, HO₃⁺, and CH₄O₂H⁺, and ions at m/Q 63 (the m/Q corresponding to DMS) including H₂CO₃H⁺, H₂N₂O₂H⁺, and C₂H₆O₂H⁺. Previous open ocean measurements of DMS and MeSH have been reported at unit mass m/Q 63
- and m/Q 49, respectively (Lawson et al., 2020). In this work with coastal seawater in an indoor laboratory setting, MeSH constituted $73.9 \pm 12.9\%$ of m/Q 49 and DMS constituted $76.8 \pm 18.0\%$ of m/Q 63. Therefore, the high resolution of the Vocus ensured accurate quantitative measurements of DMS and MeSH.



185 Figure 1: (a) Sample mass spectrum corresponding to an ISV headspace measurement. Peaks highlighted in red contain sulfur. (b) High-resolution fit around DMS (C₂H₆SH⁺). For this mass spectrum, DMS makes up 72.2% of the total ion current at m/Q 63 with C₂H₆O₂H⁺ being the second largest peak.

In Fig. S3 and Fig. S4, we show calibration curves for DMS, MeSH, and benzothiazole, and observed fragments of these molecules. Limits of detection_for DMS, MeSH, and benzothiazole at 1 minute averaging time were 0.20 ± 0.49 ppt, 1.5 ± 0.25 ppt, and 0.42 ± 0.14 ppt, respectively, calculated according to equation 1 in (Bertram et al., (2011) for a signal-to-noise ratio of 3. While observations of DMS, MeSH, and benzothiazole in this study were well above the instrument's limits of detection, open ocean measurements of DMS and MeSH in non-bloom conditions are on the order of tens of ppt, making the Vocus with its low limits of detection an ideal instrument to use in such conditions.

195

Krechmer et al. (2018) demonstrated the Vocus sensitivity to a number of non-sulfur containing VOCs is independent of relative humidity due to the high water vapor mixing ratio in the focusing ion-molecule reactor causing the relative humidity in the small volume of sample air to have limited effects on ion-molecule reactions in the Vocus. In Fig. 2 we show the sensitivity to DMS is humidity-independent, while a small humidity-dependence exists for MeSH between 2.7 × 10⁻⁴ kg m⁻³
and 0.018 kg m⁻³. The upper half of these values (0.009–0.018 kg m⁻³), corresponding to roughly 40–80% relative humidity at 25 °C, are commonly observed over mid-latitude oceans (Liu et al., 1991). DMS signal was within the standard deviation for

all absolute humidity values tested, but MeSH signal was strongly anticorrelated with absolute humidity ($R^2 = 0.96$) and decreased 20% across the absolute humidity range tested.



Figure 2: Vocus PTR-ToF-MS signal as a function of absolute humidity for a constant flow of (a) DMS (measured at m/Q 63.0263) and (b) MeSH (measured at m/Q 49.0107) in humidified zero air. Absolute humidity values between 0.009 and 0.018 kg m⁻³ are commonly measured in the MBL in mid-latitude oceans (Liu et al., 1991).

- 210 The decrease in MeSH with absolute humidity is likely a result of conversion on inlet surfaces in addition to humiditydependent changes in ion chemistry. It has been well-documented in the literature that MeSH oxidation to dimethyl disulfide (DMDS; $C_2H_6S_2$) can occur on metal surfaces (Perraud et al., 2016). We used stainless steel fittings on the tubing and inlet so MeSH loss and DMDS production was a possibility to consider in our system. MeSH and DMDS (measured as $C_2H_6S_2H^+$) were tightly correlated ($R^2 = 0.98$) in MeSH calibrations (done after SeaSCAPE), with DMDS representing roughly 12% of
- 215 the MeSH signal. This is likely a result of conversion in the inlet or in the calibration standard. However, there was no correlation ($R^2 = 0.017$) between DMDS and MeSH during SeaSCAPE, suggesting that the on average 7 ppt of DMDS measured during SeaSCAPE has a seawater source and cannot solely be a result of inlet conversion of MeSH (Fig. S5).

3.2 Gas-Phase Sulfur Budget during Bloom

Figures 3a and 3b depict the progression of the bloom in the wave channel and the effects of perturbations listed in Table S1

- 220 through waterside measurements including chlorophyll-a, heterotrophic bacteria, DMSP_t, and fluorescent dissolved organic matter (FDOM). Prior to the first nutrient addition on day 2, the mean chlorophyll-a concentration was $0.80 \pm 0.08 \ \mu g \ L^{-1}$, mean heterotrophic bacteria abundance was $2.97 \times 10^9 \pm 1.27 \times 10^9$ cells L⁻¹, and mean DMSP_t was 51.2 ± 20.7 nM. DMSP_t and heterotrophic bacteria had a small peak on days 5 and 6, respectively, but chlorophyll-a remained low at less than $1.25 \ \mu g \ L^{-1}$ until day 7. Chlorophyll-a began to rise on day 7, reaching 2.93 $\mu g \ L^{-1}$, indicative of the start of a small bloom. This in situ
- 225 bloom was enhanced through the addition of a 300-gallon tank of seawater containing healthy biomass, dominated by diatoms and with chlorophyll-a measuring 43.8 μ g L⁻¹, on day 9, causing significant responses in chlorophyll-a, DMSP_t, and heterotrophic bacteria. This added seawater was collected the same way as the water in the wave channel but on a different date. It was immediately spiked with f/2 nutrients and left outside for four days until the phytoplankton bloom reached the exponential growth phase at which time it was added to the wave channel. DMSP_t and chlorophyll-a peaked approximately
- 230 one day after the tank addition at values of 224 nM and 25.9 μ g L⁻¹, respectively. Heterotrophic bacteria had a small peak corresponding to the chlorophyll-a peak before reaching its maximum concentration four days after DMSP_t and chlorophyll-a at 1.3 x 10¹⁰ cells L⁻¹. Chlorophyll-a concentrations were sustained at elevated values around 7 μ g L⁻¹ following the peak, while heterotrophic bacteria measurements showed a local maximum on day 18.



Figure 3: (a) Time series of (a) 2-hour average chlorophyll-a and daily heterotrophic bacteria concentrations in the wave channel, (b) daily DMSPt and FDOM, (c) DMS, MeSH, benzothiazole, and the sum of other detectable sulfur ions in the mass spectrum in absolute concentration, and (d) fractional contribution of DMS, MeSH, benzothiazole, and the sum of other detectable sulfur ions in the mass spectrum to the total measured gas-phase sulfur budget by the Vocus. Fractional contribution is calculated by mixing ratio. The wave channel was filled on day 0 and nutrients were added on days 2.2 and 3.2. The tank of productive seawater and more nutrients were added on day 8.9, shown by the vertical line in Fig. 3. More details on these changes are recorded in Table S1.

The waterside measurements in Fig. 3a and Fig. 3b provide context for understanding the gas-phase sulfur emissions displayed in Fig. 3c and Fig. 3d. The organosulfur molecules studied are DMS, MeSH, benzothiazole, and "other S", where "other S" corresponds to the sum of 25 other detectable sulfur-containing ions in the mass spectrum. The other S signal was distributed

- among ions, with ions corresponding to DMDS and larger sulfur-containing compounds ($C_4H_8SO_2H^+$, $C_{10}H_{16}SH^+$, $C_{11}H_{16}SH^+$) present with appreciable signal throughout the bloom. Only three ions in the other S signal showed a correlation of $R^2 > 0.5$ with either DMS, MeSH, or benzothiazole. Known fragments of DMS, MeSH, and benzothiazole were not included in the other S signal as they are not unique molecules. Initial concentrations of DMS, MeSH, benzothiazole, and other S in the ISV headspace at the start of the experiment when seawater was first added were 545, 97, 41, and 141 ppt, respectively. DMS and
- 250 MeSH increased from the beginning of the bloom, with DMS peaking at 5690 ppt on day 13 and MeSH peaking at 274 ppt on day 11. These values are significantly higher than what is routinely measured over the open ocean (Lawson et al., 2020; Leck and Rodhe, 1991), likely owing to multiple additions of concentrated nutrients that induced the intense phytoplankton bloom and the gas equilibration time in the ISV. Other S and benzothiazole peaked earlier in the experiment on day 7 when some other anthropogenic gases, including benzophenone and naphthalene, peaked (Franklin et al., 2021). The other S signal from
- this day was primarily driven by contributions from $C_4H_8SO_2H^+$, which could be indicative of the molecule sulfolane, and $C_{10}H_{16}SH^+$, and $C_{11}H_{16}SH^+$.

Benzothiazole is a reduced sulfur molecule measured in the ISV headspace that contributed significantly to the gas-phase sulfur budget during the bloom. Since the Henry's law constant for benzothiazole in water is three to four orders of magnitude 260 higher than that of DMS and MeSH, it is possible that the emission ratio of benzothiazole to DMS and MeSH is uniquely sensitive to the water and air flow rates in the ISV (Sander, 2015). Benzothiazole has both biological and anthropogenic sources. It is naturally produced by the γ -Proteobacteria, *Pseudomonas fluorescens* (Le Bozec and Moody, 2009), and the Actinobacteria, *Micrococcus* sp. (Schulz-Bohm et al., 2017), both of which can be found in seawater. Benzothiazole (C_7H_5NS) belongs to a class of structurally similar molecules called benzothiazoles, which are a group of high production volume 265 chemicals found in wastewater and urban runoff (De Wever and Verachtert, 1997; Hidalgo-Serrano et al., 2019). Based on analysis in Franklin et al. (2021) where the benzothiazole molecule was (1) consistently observed in significant quantities in the dissolved, gas, and aerosol phases, (2) the gas-phase molecule displayed temporal behavior similar to a group of other anthropogenic gases, and (3) anthropogenic benzothiazole tracer species were observed, we attribute its source as primarily anthropogenic from the presence of pollutants in coastal waters. The two benzothiazole measurements, here by the Vocus 270 PTR-ToF-MS and in Franklin et al. (2021) by TD-GCxGC-ToF-MS, differ in absolute quantities but are highly correlated (R² = 0.91) (Franklin et al., 2021). Additionally, Franklin et al. (2021) showed the oxidation of benzothiazole in a potential aerosol mass-oxidation flow reactor produced increasing amounts of secondary aerosol and SO₂ from 2.9-4.7 days of equivalent aging. As a result, we suspect benzothiazole may be important in coastal regions, but is not expected to be a significant sulfur

275

source in open ocean regions.

We find little evidence for emissions of DMS oxidation products. Dimethyl sulfoxide (DMSO) did not have a signal-to-noise ratio above three on any day of the bloom. Dimethyl sulfone (DMSO₂) was present through day 6 prior to the rise in chlorophyll-a, then decayed to below the detection limit (Fig. S6). The observed small concentrations of DMS oxidation products relative to DMS are expected as the mesocosm experiment was conducted in a low-oxidant indoor environment.

- 280 Methane sulfonamide (MSAM) measured on average less than 1 ppt during the experiment. In contrast, Edtbauer et al. (2020) (Edtbauer et al., 2020a)measured MSAM at mixing ratios of up to 60 ppt, corresponding to up to 33% of DMS, in upwelling areas of the Arabian Sea and suggested MSAM had a direct oceanic emission source. Results from our study suggest that the pathway producing MSAM in the Arabian Sea was not active in our experiments utilizing coastal seawater from Southern California. and showed no positive correlation with the DMS oxidation product, DMSO₂ (Fig. S6). This is surprising as MSAM
 285 was recently measured at mixing ratios of up to 60 ppt corresponding to up to 33% of DMS in upwelling areas of the Arabian
- Sea and was correlated with DMSO₂ (r = 0.8) (Edtbauer et al., 2020).

In Fig. 3c, we show that non-DMS sulfur sources comprise a larger fraction (36.8 ± 7.64%) of the gas-phase sulfur budget in the pre-bloom period prior to the peak in chlorophyll-a on day 10. The contribution of non-DMS to total sulfur in this period is higher than in previous measurements, such as those of DMS, MeSH, DMDS, and carbon disulfide reported by Leck and Rodhe (1991), where waterside non-DMS/total sulfur was on average 10, 6, and 16% in the Baltic Sea, Kattegat/Skagerrak, and North Sea, respectively, and average chlorophyll-a was 0.5–1.9 µg L⁻¹. When comparing the same sulfur molecules as in Leck and Rodhe (1991), non-DMS represents 18.7 ± 3.4% of total sulfur for the same period. This is within the range of observations in the North Sea and suggests that this difference is either driven by a subset of sulfur molecules perhaps unique to a coastal environment or the measurement technique. Another possible cause for the high observed non-DMS sulfur is that the distribution and magnitude of organosulfur emissions may have been altered through the process of water collection, transport, and wave channel filling, when the water temperature increased 4 °C during the first two days. When chlorophyll-a was low and after the peak in bulk water heterotrophic bacterial abundance on day 15, DMS alone accounted for approximately 87% of the sulfur budget. This is within the range of previous measurements of DMS contribution to total sulfur made by Leck

300 and Rodhe (1991).

305

3.3 The DMS:MeSH Molar Ratio

In the remaining sections of the paper, we focus our analysis of the gas-phase sulfur budget on DMS and MeSH, as they collectively accounted for $84 \pm 8.1\%$ of the total sulfur budget. Further, since their production and loss are inherently linked, measurements of the molar ratio of DMS:MeSH provides unique perspective on waterside sulfur chemistry. Given that the Henry's law constants and diffusion constants for DMS and MeSH are roughly the same, and assuming atmospheric loss is negligible in the ISV, we expect the measured molar ratio of DMS:MeSH in the headspace to reflect the dissolved concentration ratio in the seawater. In this work, DMS and MeSH increased at approximately the same rate from day 1 to day 9 of the bloom shown by their similar slopes in Fig. 4a. This resulted in a relatively constant DMS:MeSH of 4.6 ± 0.9 during

the first 9 days, shown in Fig. 4b. This value is consistent with the range of previously reported waterside concentration and

- 310 sea-to-air fluxes of DMS:MeSH measured over the open ocean during non-bloom conditions (Kettle et al., 2001; Leck and Rodhe, 1991) and during a modest bloom (chlorophyll-a < $3 \mu g L^{-1}$) (Lawson et al., 2020). The stable DMS:MeSH value is unexpected as it was sustained through three nutrient additions and the addition of a tank containing highly productive seawater (Table S1). Following the peak in MeSH on day 11, DMS:MeSH began to increase considerably. The peak in DMS:MeSH on day 13 was driven by a maximum in DMS concentration relative to declining MeSH concentrations, representing a DMS
- 315 increase of 3340 ppt and MeSH decrease of 150 ppt from day 11 to 13. Additionally, the period of increasing DMS:MeSH from day 10–13 likely reflects the substantially shorter waterside lifetime of MeSH compared to DMS. Maximum DMS concentrations occurred 3 days after peak DMSP_t, whereas maximum MeSH concentrations occurred 1 day after peak DMSP_t, suggesting quick turnover of DMSP_d to form MeSH. The DMS:MeSH peak on day 17 was driven by low MeSH concentrations (35 ppt MeSH) relative to DMS (2490 ppt DMS). The episodic DMS:MeSH variations around days 13 and 17 are likely the
- 320 results of external perturbations to the wave channel water (Table S1) which affected water mixing and algal cell lysis. Despite these two external factors, DMS:MeSH increased significantly from the initial to final day of the experiment, reaching values higher than what have been observed in previous oceanic studies. Thus, increases in the DMS:MeSH ratio might reflect the biological dynamics induced in our controlled system.



325 Figure 4: (a) Time series of DMS and MeSH in the ISV, where error bars indicate the standard deviation over the 1 Hz measurements for the day. (b) Time series of DMS:MeSH, where error bars indicate the error propagation from (a). Vertical line represents the addition of highly productive seawater and more nutrients.

3.3.1 Observed Correlations of DMS:MeSH with Seawater Properties

- 330 In what follows, we explore in detail the factors that control the DMS:MeSH emission ratio. Properties with the potential to impact the waterside production and loss of DMS and MeSH are examined, including chlorophyll-a, FDOM, wave channel water temperature, dissolved oxygen, and salinity. Later, we focus on a subset of these measurements that provide insight into the transformation of precursor sulfur molecules to DMS and MeSH, namely DMSP_t, bacterial sulfur demand, and methionine aminopeptidase activity. Regressions of these variables against DMS:MeSH are shown in Fig. 5.
- 335

340

Chlorophyll-a, serving as a metric for phytoplankton biomass, and thus, intracellular DMSP (DMSP_p) is expected to trend with total available sulfur and influence production of DMS and MeSH (Galí et al., 2015). FDOM is expected to positively correlate with DMS:MeSH due to both MeSH loss by reaction with DOM (Lawson et al., 2020) and its impact on DMS production. As the concentration and chemical complexity of FDOM increases during a bloom, the available sulfur compounds are also expected to increase as there is a release of sulfur-rich amino acids in addition to DMSP (Pinhassi et al., 2005; Meon and Kirchman, 2001). An excess of available sulfur will favor DMS production, leading to an increase in DMS:MeSH, evidenced by the weak ($R^2 = 0.24$) positive correlation in Fig. 5b. Temperature and dissolved oxygen may influence DMS:MeSH through their relationship with bacterial growth rates and DMSP conversion (Kiene and Linn, 2000b, a). Both have weak correlations with DMS:MeSH in Fig. 5c and Fig. 5d. While the strong anticorrelation between DMS:MeSH and salinity in this experiment

345 is in accordance with prior experiments constraining the DMSP demethylation/demethiolation pathways as a function of salinity (Magalhães et al., 2012; Salgado et al., 2014), we argue that the observed anticorrelation in this experiment is simply a correlation and not suggestive of a salinity control. This observation is discussed further in the supplementary information (S5). The remaining sections of the paper will focus specifically on the relationship between bacterial sulfur demand, DMSP_t, and methionine, as this is expected to modulate the fate of DMSP (Kiene et al., 2000).



Figure 5: Regressions of DMS:MeSH versus waterside variables measured in the wave channel: (a) chlorophyll-a, (b) FDOM, (c) wave channel water temperature, (d) dissolved oxygen, (e) salinity, (f) DMSP_t, (g) bacterial sulfur demand (using a cellular C:S ratio in bacteria of 248 (Cuhel et al., 1982)), and (h) methionine aminopeptidase activity. The color of the marker indicates the day of the bloom. Panel a uses two hour averages, panels b, c, d, and e use minute averages, and panels f, g, and h use daily averages.

355 3.3.2 Biological Influences on DMS:MeSH

360

DMSP is a precursor for both MeSH and DMS. DMSP_t ranged from 15.3 to 224 nM (Fig. 6a), similar to concentrations observed in other phytoplankton blooms (Galí et al., 2015). Taking chlorophyll-a as a proxy for phytoplankton biomass, the ratio of DMSP_t:chlorophyll-a suggests that the bloom was largely dominated by low DMSP producers after the first few days (Fig. S7), in accordance with the observed large diatom population (Fig. S8) (Dani and Loreto, 2017; McParland and Levine, 2019).

Further, assuming the lifetimes of DMS and DMSP_t in the water are the same, since we do not have direct measurements of DMSP and DMS cycling, we will use the waterside DMS:DMSP_t ratio can be used to estimate the DMSP_t to DMS conversion efficiency in the seawater_(Galí et al., 2018, 2021). While typically considered to be around 10% (Kiene and Linn, 2000b;
Lizotte et al., 2012; Vila-Costa et al., 2008), waterside DMS:DMSP_t was much lower in this study, ranging between 0.38% and 8.30% (average 2.88%) (Fig. S7). This suggests there was low DMS production from DMSP, which could be a result of diatoms dominating the experiment while other taxa capable of directly producing DMS from DMSP (such as dinoflagellates and haptophytes) being less abundant (Lizotte et al., 2012; Stefels and van Boekel, 1993). Additionally, low DMS:DMSP_t could be representative of a significant DMS loss either through ventilation or biotic or abiotic transformations in the seawater.



370

Figure 6: (a) Time series of heterotrophic bacteria and DMSPt. (b) Time series of calculated bacterial sulfur demand (cellular C:S in bacteria of 86-248) and estimated assimilated sulfur assuming 25% of DMSPt is assimilated by bacteria (Fagerbakke et al., 1996; Cuhel et al., 1982; Kiene and Linn, 2000a). (c) DMS:MeSH is low in the pre-bloom at the beginning of the experiment, before increasing significantly driven by different sources of available sulfur. Vertical line represents the addition of productive seawater and additional nutrients to the wave channel.

375

The dynamics between bacterial sulfur demand and available sulfur sources are important for regulating the fate of DMSP and therefore the DMS:MeSH ratio (Kiene et al., 2000). Bacterial sulfur demand was calculated using measured bacterial productivity, assuming lower and upper limits on cellular C:S ratios in bacteria of 86 (Fagerbakke et al., 1996) and 248 (Cuhel

et al., 1982; Kiene and Linn, 2000a), and assuming 25% of sulfur from DMSPt is assimilated by bacteria (Pinhassi et al., 2005).
 The comparison between assimilated sulfur from DMSPt and bacterial sulfur demand (Fig. 6b) shows that at the beginning of the experiment during the pre-bloom stage, bacterial sulfur demand exceeded sulfur available from DMSP, suggesting that all the DMSP was channeled toward the demethylation pathway and the formation of MeSH, and that other sulfur sources

complemented the bacterial demand. After the peak of the bloom, the assimilated sulfur from DMSPt exceeded the bacterial

385 sulfur demand (assuming a cellular C:S ratio of 248), suggesting that bacteria produced DMS from the excess sulfur source in the latter half of the experiment, <u>likely responsible for part of the leading to a</u> significant increase in measured DMS:MeSH (Fig.6c). This preference toward DMSP cleavage at the end of this bloom is likely due to an increase in the amount and chemical complexity of dissolved organic matter and the presence of other forms of available sulfur often observed at the end of phytoplankton blooms (Pinhassi et al., 2005). Since DMSP_t represents a small fraction of bioavailable carbon throughout the bloom (<1%), other existing sulfur sources in the carbon pool may be more easily accessible to bacteria than DMSP.</p>

The preferential assimilation of other sulfur sources than DMSP is further supported by strong correlations between DMS:MeSH and aminopeptidase activities (Fig. S9), particularly methionine aminopeptidase ($R^2 = 0.82$) (Fig. 5h). Aminopeptidases catalyze cleavage of amino acids from proteins and peptides (Taylor, 1993). This suggests that protein degradation, or even direct methionine assimilation may provide additional sulfur sources to bacteria.

Taken together, the increasing trajectory of DMS:MeSH throughout the experiment reflects changes in bacterial sulfur demand and the availability of other organosulfur molecules in the system. The pre-bloom stage of this experiment where DMS:MeSH was low and stable (4.60 ± 0.93) follows ambient conditions, such as those observed in Lawson et al. (2020) (Table 1), and the significantly higher (31.8 ± 18.7) ratio observed in the bloom and decay stage in this work are likely the product of an intense induced phytoplankton bloom and water mixing conditions not usually observed in the open ocean, but could be reflective of intense blooms in coastal environments.

405

400

Measurement	DMS:MeSH	Notes	Reference	
Airside	4.60 ± 0.93	Mesocosm experiment;	This work	
		Pre-bloom	-	
	31.8 ± 18.7	Mesocosm experiment;		
		Bloom and decay		
Airside	3–33	Southwest Pacific Ocean	Lawson et al. (2020)	
Flux	3–7			
Waterside	2-5.3	Subarctic Northeast Pacific Ocean	Kiene et al. (2017)	
Waterside	1–30	Atlantic Ocean	Kettle et al. (2001)	
Waterside	16.4 (mean)	Baltic Sea	Leck and Rodhe (1991)	
	19.7 (mean)	Kattegat-Skagerrak		
	6.1 (mean)	North Sea		

Table 1. DMS: MeSH measurements from this work and previous studies.

420 4 Conclusions and Outlook

During an induced phytoplankton bloom on coastal seawater, non-DMS organosulfur molecules accounted for on average 37% of the total gas-phase sulfur budget in the pre-bloom stage when chlorophyll-a was low, representative of ambient conditions in a typical coastal environment. The ratio of DMS:MeSH increased significantly during the phytoplankton bloom, likely due to the interaction between several variables influencing the molecules' production and loss processes in the seawater.

- 425 DMS:MeSH was primarily sensitive to bacterial sulfur demand and the chemical composition and magnitude of available sulfur sources during the bloom. The low DMS:MeSH measured during the pre-bloom at the beginning of the experiment and which is more representative of average in situ conditions suggests MeSH can have a significant impact on atmospheric oxidative capacity and secondary sulfate formation in coastal environments given that it reacts with the hydroxyl radical seven times faster than DMS and has an expected unit yield of SO₂. This finding, combined with the significant emission of
- 430 benzothiazole and substantial concentrations of other sulfur gases observed in this experiment, suggest pathways to secondary sulfate formation in a coastal environment warrant further study. A more complete understanding of coastal emissions of gaseous precursors to sulfate aerosol will improve model estimates of cloud formation and radiative balance in the marine environment.
- Data Availability: Seawater measurements and Vocus PTR-ToF-MS measurements of DMS, MeSH, benzothiazole, and total 435 other sulfur will be made available at https://minds.wisconsin.edu/handle/1793/76304.

Supplement: Wave channel and mesocosm details, additional methods descriptions, DMS:MeSH salinity discussion, and supporting tables and figures.

440

450

Author Contributions: Conceptualization: DBK, GAN, JSS, KM, CL, CDC, AHG, KAP, THB; Investigation: DBK, GAN, JSS, ANM, JD, SA, EBB, MW, CKM, TP, FM, DRC, THB; Data Analysis: DBK, GAN, JSS, ANM, JD, SA, EBB, CKM, TP, FM, DRC, THB; Writing: DBK, JSS, JD, THB; All authors reviewed and edited the paper.

445 **Competing interests:** The authors declare that they have no conflict of interest.

Acknowledgments: We would like to thank the Center for Aerosol Impacts on Chemistry of the Environment (an NSF Chemical Innovation Center, (CHE-1801971)) and the full team which participated in the SeaSCAPE campaign for their support of this work. We specifically thank Michael Alves for his contribution of the initial benzothiazole observation and identification at SeaSCAPE and related support of this publication.

References

Andreae, M. O.: Ocean-atmosphere interactions in the global biogeochemical sulfur cycle, Mar. Chem., 30, 1–29, https://doi.org/10.1016/0304-4203(90)90059-L, 1990.

Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe,
 J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume I - gas phase reactions of Ox, HOx, NOx and SOx species, Atmospheric Chem. Phys., 4, 1461–1738, https://doi.org/10.5194/acp-4-1461-2004, 2004.

Azam, F. and Smith, D. C.: A simple, economical method for measuring bacterial protein synthesis rates in seawater using 3H-leucine, Mar. Microb. Food Webs, 6, 107–114, 1992.

Bates, T. S., Lamb, B. K., Guenther, A., Dignon, J., and Stoiber, R. E.: Sulfur emissions to the atmosphere from natural sources, J. Atmospheric Chem., 14, 315–337, https://doi.org/10.1007/BF00115242, 1992.

Bertram, T. H., Kimmel, J. R., Crisp, T. A., Ryder, O. S., Yatavelli, R. L. N., Thornton, J. A., Cubison, M. J., Gonin, M., and Worsnop, D. R.: A field-deployable, chemical ionization time-of-flight mass spectrometer, Atmospheric Meas. Tech., 4, 1471–1479, https://doi.org/10.5194/amt-4-1471-2011, 2011.

Carpenter, L. J., Archer, S. D., and Beale, R.: Ocean-atmosphere trace gas exchange, Chem. Soc. Rev., 41, 6473, https://doi.org/10.1039/c2cs35121h, 2012.

Challenger, F. and Simpson, M. I.: 320. Studies on biological methylation. Part XII. A precursor of the dimethyl sulphide evolved by Polysiphonia fastigiata. dimethyl-2-carboxyethylsulphonium hydroxide and its salts, J. Chem. Soc. Resumed, 1591, https://doi.org/10.1039/jr9480001591, 1948.

Charlson, R. J., Lovelock, J. E., Andreae, M. O., and Warren, S. G.: Oceanic phytoplankton, atmospheric sulphur, cloud albedo and climate, Nature, 326, 655–661, https://doi.org/10.1038/326655a0, 1987.

Chen, Q., Sherwen, T., Evans, M., and Alexander, B.: DMS oxidation and sulfur aerosol formation in the marine troposphere: a focus on reactive halogen and multiphase chemistry, Atmospheric Chem. Phys., 18, 13617–13637, https://doi.org/10.5194/acp-18-13617-2018, 2018.

Cuhel, R. L., Taylor, C. D., and Jannasch, H. W.: Assimilatory Sulfur Metabolism in Marine Microorganisms: Sulfur
 Metabolism, Protein Synthesis, and Growth of Alteromonas luteo-violaceus and Pseudomonas halodurans During Perturbed
 Batch Growtht, APPL Env. MICROBIOL, 43, 9, 1982.

Dani, K. G. S. and Loreto, F.: Trade-Off Between Dimethyl Sulfide and Isoprene Emissions from Marine Phytoplankton, Trends Plant Sci., 22, 361–372, https://doi.org/10.1016/j.tplants.2017.01.006, 2017.

De Wever, H. and Verachtert, H.: Biodegradation and toxicity of benzothiazoles, Water Res., 31, 2673–2684, 480 https://doi.org/10.1016/S0043-1354(97)00138-3, 1997.

Edler, L. and Elbrächter, M.: The Utermöhl method for quantitative phytoplankton analysis, in: Microscopic and Molecular Methods for Quantitative Phytoplankton Analysis, 13–20, 2010.

Edtbauer, A., Stönner, C., Pfannerstill, E. Y., Berasategui, M., Walter, D., Crowley, J. N., Lelieveld, J., and Williams, J.: A new marine biogenic emission: methane sulfonamide (MSAM), DMS and DMSO measured in air over the Arabian Sea,
Gases/Field Measurements/Troposphere/Chemistry (chemical composition and reactions), https://doi.org/10.5194/acp-2019-1021, 2020.

Fagerbakke, K., Heldal, M., and Norland, S.: Content of carbon, nitrogen, oxygen, sulfur and phosphorus in native aquatic and cultured bacteria, Aquat. Microb. Ecol., 10, 15–27, https://doi.org/10.3354/ame010015, 1996.

Faloona, I.: Sulfur processing in the marine atmospheric boundary layer: A review and critical assessment of modeling uncertainties, Atmos. Environ., 43, 2841–2854, https://doi.org/10.1016/j.atmosenv.2009.02.043, 2009.

Flöck, O. R. and Andreae, M. O.: Photochemical and non-photochemical formation and destruction of carbonyl sulfide and methyl mercaptan in ocean waters, Mar. Chem., 54, 11–26, https://doi.org/10.1016/0304-4203(96)00027-8, 1996.

Franklin, E. B., Alves, M. R., Moore, A. N., Kilgour, D. B., Novak, G. A., Mayer, K., Sauer, J. S., Weber, R. J., Dang, D., Winter, M., Lee, C., Cappa, C. D., Bertram, T. H., Prather, K. A., Grassian, V. H., and Goldstein, A. H.: Atmospheric Benzothiazoles in a Coastal Marine Environment, Environ. Sci. Technol., https://doi.org/10.1021/acs.est.1c04422, 2021.

495

Galí, M., Devred, E., Levasseur, M., Royer, S.-J., and Babin, M.: A remote sensing algorithm for planktonic dimethylsulfoniopropionate (DMSP) and an analysis of global patterns, Remote Sens. Environ., 171, 171–184, https://doi.org/10.1016/j.rse.2015.10.012, 2015.

Galí, M., Levasseur, M., Devred, E., Simó, R., and Babin, M.: Sea-surface dimethylsulfide (DMS) concentration from satellite data at global and regional scales, Biogeosciences, 15, 3497–3519, https://doi.org/10.5194/bg-15-3497-2018, 2018.

Galí, M., Lizotte, M., Kieber, D. J., Randelhoff, A., Hussherr, R., Xue, L., Dinasquet, J., Babin, M., Rehm, E., and Levasseur, M.: DMS emissions from the Arctic marginal ice zone. Elem. Sci. Anthr.. 9. 00113. https://doi.org/10.1525/elementa.2020.00113, 2021.

Gasol, J. M. and Del Giorgio, P. A.: Using flow cytometry for counting natural planktonic bacteria and understanding the structure of planktonic bacterial communities, Sci. Mar., 64, 197–224, https://doi.org/10.3989/scimar.2000.64n2197, 2000.

Gharagheizi, F.: Determination of Diffusion Coefficient of Organic Compounds in Water Using a Simple Molecular-Based Method, Ind. Eng. Chem. Res., 51, 2797–2803, https://doi.org/10.1021/ie201944h, 2012.

Hidalgo-Serrano, M., Borrull, F., Marcé, R. M., and Pocurull, E.: Presence of benzotriazoles, benzothiazoles and benzenesulfonamides in surface water samples by liquid chromatography coupled to high-resolution mass spectrometry, Sep. 510
Sci. PLUS, 2, 72–80, https://doi.org/10.1002/sscp.201800140, 2019.

Kettle, A. J., Rhee, T. S., Hobe, M. von, Poulton, A., Aiken, J., and Andreae, M. O.: Assessing the flux of different volatile sulfur gases from the ocean to the atmosphere, J. Geophys. Res. Atmospheres, 106, 12193–12209, https://doi.org/10.1029/2000JD900630, 2001.

Kiene, R. P.: Production of methanethiol from dimethylsulfoniopropionate in marine surface waters, Mar. Chem., 54, 69–83, https://doi.org/10.1016/0304-4203(96)00006-0, 1996.

Kiene, R. P. and Linn, L. J.: Distribution and turnover of dissolved DMSP and its relationship with bacterial production and dimethylsulfide in the Gulf of Mexico, Limnol. Oceanogr., 45, 849–861, https://doi.org/10.4319/lo.2000.45.4.0849, 2000a.

Kiene, R. P. and Linn, L. J.: The fate of dissolved dimethylsulfoniopropionate (DMSP) in seawater: tracer studies using 35S-DMSP, Geochim. Cosmochim. Acta, 64, 2797–2810, https://doi.org/10.1016/S0016-7037(00)00399-9, 2000b.

520 Kiene, R. P., Linn, L. J., González, J., Moran, M. A., and Bruton, J. A.: Dimethylsulfoniopropionate and Methanethiol Are Important Precursors of Methionine and Protein-Sulfur in Marine Bacterioplankton, Appl. Environ. Microbiol., 65, 4549– 4558, https://doi.org/10.1128/AEM.65.10.4549-4558.1999, 1999.

Kiene, R. P., Linn, L. J., and Bruton, J. A.: New and important roles for DMSP in marine microbial communities, J. Sea Res., 43, 209–224, https://doi.org/10.1016/S1385-1101(00)00023-X, 2000.

525 Kiene, R. P., Nowinski, B., Esson, K., Preston, C., Marin, R., Birch, J., Scholin, C., Ryan, J., and Moran, M. A.: Unprecedented DMSP Concentrations in a Massive Dinoflagellate Bloom in Monterey Bay, CA, Geophys. Res. Lett., 46, 12279–12288, https://doi.org/10.1029/2019GL085496, 2019.

Kim, M. J., Zoerb, M. C., Campbell, N. R., Zimmermann, K. J., Blomquist, B. W., Huebert, B. J., and Bertram, T. H.: Revisiting benzene cluster cations for the chemical ionization of dimethyl sulfide and select volatile organic compounds, Atmospheric Meas. Tech., 9, 1473–1484, https://doi.org/10.5194/amt-9-1473-2016, 2016.

530

Kirchman, D., K'Nees, E., and Hodson, R.: Leucine Incorporation and Its Potential as a Measure of Protein Synthesis by Bacteria in Natural Aquatic Systemst, APPL Env. MICROBIOL, 49, 9, 1985.

Krechmer, J., Lopez-Hilfiker, F., Koss, A., Hutterli, M., Stoermer, C., Deming, B., Kimmel, J., Warneke, C., Holzinger, R., Jayne, J., Worsnop, D., Fuhrer, K., Gonin, M., and de Gouw, J.: Evaluation of a New Reagent-Ion Source and Focusing Ion–
Molecule Reactor for Use in Proton-Transfer-Reaction Mass Spectrometry, Anal. Chem., 90, 12011–12018, https://doi.org/10.1021/acs.analchem.8b02641, 2018.

Kwint, R. and Kramer, K.: Annual cycle of the production and fate of DMS and DMSP in a marine coastal system, Mar. Ecol. Prog. Ser., 134, 217–224, https://doi.org/10.3354/meps134217, 1996.

Lana, A., Bell, T. G., Simó, R., Vallina, S. M., Ballabrera-Poy, J., Kettle, A. J., Dachs, J., Bopp, L., Saltzman, E. S., Stefels,
J., Johnson, J. E., and Liss, P. S.: An updated climatology of surface dimethlysulfide concentrations and emission fluxes in the global ocean, Glob. Biogeochem. Cycles, 25, https://doi.org/10.1029/2010GB003850, 2011.

Lavi, A., Vermeuel, M. P., Novak, G. A., and Bertram, T. H.: The sensitivity of benzene cluster cation chemical ionization mass spectrometry to select biogenic terpenes, Atmospheric Meas. Tech., 11, 3251–3262, https://doi.org/10.5194/amt-11-3251-2018, 2018.

545 Lawson, S. J., Law, C. S., Harvey, M. J., Bell, T. G., Walker, C. F., de Bruyn, W. J., and Saltzman, E. S.: Methanethiol, dimethyl sulfide and acetone over biologically productive waters in the southwest Pacific Ocean, Atmos Chem Phys, 18, 2020.

Le Bozec, L. and Moody, C. J.: Naturally Occurring Nitrogen–Sulfur Compounds. The Benzothiazole Alkaloids, Aust. J. Chem., 62, 639, https://doi.org/10.1071/CH09126, 2009.

Leck, C. and Rodhe, H.: Emissions of marine biogenic sulfur to the atmosphere of northern Europe, J. Atmospheric Chem., 12, 63–86, https://doi.org/10.1007/BF00053934, 1991.

Lee, C.-L. and Brimblecombe, P.: Anthropogenic contributions to global carbonyl sulfide, carbon disulfide and organosulfides fluxes, Earth-Sci. Rev., 160, 1–18, https://doi.org/10.1016/j.earscirev.2016.06.005, 2016.

Lennartz, S. T., Marandino, C. A., von Hobe, M., Andreae, M. O., Aranami, K., Atlas, E., Berkelhammer, M., Bingemer, H., Booge, D., Cutter, G., Cortes, P., Kremser, S., Law, C. S., Marriner, A., Simó, R., Quack, B., Uher, G., Xie, H., and Xu, X.:
Marine carbonyl sulfide (OCS) and carbon disulfide (CS₂): a compilation of measurements in seawater and the marine boundary layer, Earth Syst. Sci. Data, 12, 591–609, https://doi.org/10.5194/essd-12-591-2020, 2020.

Li, H., Canagaratna, M. R., Riva, M., Rantala, P., Zhang, Y., Thomas, S., Heikkinen, L., Flaud, P.-M., Villenave, E., Perraudin, E., Worsnop, D., Kulmala, M., Ehn, M., and Bianchi, F.: Source identification of atmospheric organic vapors in two European pine forests: Results from Vocus PTR-TOF observations, Gases/Field Measurements/Troposphere/Chemistry (chemical composition and reactions), https://doi.org/10.5194/acp-2020-648, 2020a.

Li, H., Riva, M., Rantala, P., Heikkinen, L., Daellenbach, K., Krechmer, J. E., Flaud, P.-M., Worsnop, D., Kulmala, M., Villenave, E., Perraudin, E., Ehn, M., and Bianchi, F.: Terpenes and their oxidation products in the French Landes forest: insights from Vocus PTR-TOF measurements, Atmospheric Chem. Phys., 20, 1941–1959, https://doi.org/10.5194/acp-20-1941-2020, 2020b.

565 Liu, W. T., Tang, W., and Niiler, P. P.: Humidity Profiles over the Ocean, American Meteorological Society, 1991.

Lizotte, M., Levasseur, M., Michaud, S., Scarratt, M. G., Merzouk, A., Gosselin, M., Pommier, J., Rivkin, R. B., and Kiene, R. P.: Macroscale patterns of the biological cycling of dimethylsulfoniopropionate (DMSP) and dimethylsulfide (DMS) in the Northwest Atlantic, Biogeochemistry, 110, 183–200, https://doi.org/10.1007/s10533-011-9698-4, 2012.

Magalhães, C., Salgado, P., Kiene, R. P., and Bordalo, A. A.: Influence of salinity on dimethyl sulfide and methanethiol formation in estuarine sediments and its side effect on nitrous oxide emissions, Biogeochemistry, 110, 75–86, https://doi.org/10.1007/s10533-011-9690-z, 2012.

McParland, E. L. and Levine, N. M.: The role of differential DMSP production and community composition in predicting variability of global surface DMSP concentrations, Limnol. Oceanogr., 64, 757–773, https://doi.org/10.1002/lno.11076, 2019.

Meon, B. and Kirchman, D. L.: Dynamics and molecular composition of dissolved organic material during experimental phytoplankton blooms, Mar. Chem., 75, 185–199, https://doi.org/10.1016/S0304-4203(01)00036-6, 2001.

Perraud, V., Meinardi, S., Blake, D. R., and Finlayson-Pitts, B. J.: Challenges associated with the sampling and analysis of organosulfur compounds in air using real-time PTR-ToF-MS and offline GC-FID, Atmospheric Meas. Tech., 9, 1325–1340, https://doi.org/10.5194/amt-9-1325-2016, 2016.

Pinhassi, J., Simó, R., González, J. M., Vila, M., Alonso-Sáez, L., Kiene, R. P., Moran, M. A., and Pedrós-Alió, C.:
Dimethylsulfoniopropionate Turnover Is Linked to the Composition and Dynamics of the Bacterioplankton Assemblage during a Microcosm Phytoplankton Bloom, Appl. Environ. Microbiol., 71, 7650–7660, https://doi.org/10.1128/AEM.71.12.7650-7660.2005, 2005.

Prather, K. A., Bertram, T. H., Grassian, V. H., Deane, G. B., Stokes, M. D., DeMott, P. J., Aluwihare, L. I., Palenik, B. P., Azam, F., Seinfeld, J. H., Moffet, R. C., Molina, M. J., Cappa, C. D., Geiger, F. M., Roberts, G. C., Russell, L. M., Ault, A.
P., Baltrusaitis, J., Collins, D. B., Corrigan, C. E., Cuadra-Rodriguez, L. A., Ebben, C. J., Forestieri, S. D., Guasco, T. L., Hersey, S. P., Kim, M. J., Lambert, W. F., Modini, R. L., Mui, W., Pedler, B. E., Ruppel, M. J., Ryder, O. S., Schoepp, N. G., Sullivan, R. C., and Zhao, D.: Bringing the ocean into the laboratory to probe the chemical complexity of sea spray aerosol, Proc. Natl. Acad. Sci., 110, 7550–7555, https://doi.org/10.1073/pnas.1300262110, 2013.

Salgado, P., Kiene, R., Wiebe, W., and Magalhães, C.: Salinity as a regulator of DMSP degradation in Ruegeria pomeroyi DSS-3, J. Microbiol., 52, 948–954, https://doi.org/10.1007/s12275-014-4409-1, 2014.

Sander, R.: Compilation of Henry's law constants (version 4.0) for water as solvent, Atmospheric Chem. Phys., 15, 4399–4981, https://doi.org/10.5194/acp-15-4399-2015, 2015.

Schulz-Bohm, K., Martín-Sánchez, L., and Garbeva, P.: Microbial Volatiles: Small Molecules with an Important Role in Intraand Inter-Kingdom Interactions, Front. Microbiol., 8, 2484, https://doi.org/10.3389/fmicb.2017.02484, 2017.

595 Sekimoto, K., Li, S.-M., Yuan, B., Koss, A., Coggon, M., Warneke, C., and de Gouw, J.: Calculation of the sensitivity of proton-transfer-reaction mass spectrometry (PTR-MS) for organic trace gases using molecular properties, Int. J. Mass Spectrom., 421, 71–94, https://doi.org/10.1016/j.ijms.2017.04.006, 2017.

Simó, R.: Production of atmospheric sulfur by oceanic plankton: biogeochemical, ecological and evolutionary links, Trends Ecol. Evol., 16, 287–294, https://doi.org/10.1016/S0169-5347(01)02152-8, 2001.

600 Simon, M. and Azam, F.: Protein content and protein synthesis rates of planktonic marine bacteria, Mar. Ecol. Prog. Ser., 51, 201–213, https://doi.org/10.3354/meps051201, 1989.

Stefels, J. and van Boekel, W. H. M.: Production of DMS from dissolved DMSP in axenic cultures of the marine phytoplankton species Phaeocystis sp., Mar. Ecol. Prog. Ser., 97, 11–18, 1993.

Taylor, A.: Aminopeptidases: structure and function, FASEB J., 7, 290–298, https://doi.org/10.1096/fasebj.7.2.8440407, 1993.

- 605 Veres, P. R., Neuman, J. A., Bertram, T. H., Assaf, E., Wolfe, G. M., Williamson, C. J., Weinzierl, B., Tilmes, S., Thompson, C. R., Thames, A. B., Schroder, J. C., Saiz-Lopez, A., Rollins, A. W., Roberts, J. M., Price, D., Peischl, J., Nault, B. A., Møller, K. H., Miller, D. O., Meinardi, S., Li, Q., Lamarque, J.-F., Kupc, A., Kjaergaard, H. G., Kinnison, D., Jimenez, J. L., Jernigan, C. M., Hornbrook, R. S., Hills, A., Dollner, M., Day, D. A., Cuevas, C. A., Campuzano-Jost, P., Burkholder, J., Bui, T. P., Brune, W. H., Brown, S. S., Brock, C. A., Bourgeois, I., Blake, D. R., Apel, E. C., and Ryerson, T. B.: Global airborne sampling
- 610 reveals a previously unobserved dimethyl sulfide oxidation mechanism in the marine atmosphere, Proc. Natl. Acad. Sci., 201919344, https://doi.org/10.1073/pnas.1919344117, 2020.

Vila, M., Simó, R., Kiene, R. P., Pinhassi, J., González, J. M., Moran, M. A., and Pedrós-Alió, C.: Use of Microautoradiography Combined with Fluorescence In Situ Hybridization To Determine Dimethylsulfoniopropionate Incorporation by Marine Bacterioplankton Taxa, Appl. Environ. Microbiol., 70, 4648–4657, https://doi.org/10.1128/AEM.70.8.4648-4657.2004, 2004.

615 Vila-Costa, M., Kiene, R. P., and Simí, R.: Seasonal variability of the dynamics of dimethylated sulfur compounds in a coastal northwest Mediterranean site, Limnol. Oceanogr., 53, 198–211, https://doi.org/10.4319/lo.2008.53.1.0198, 2008.

Wurl, O. (Ed.): Practical Guidelines for the Analysis of Seawater, CRC Press, https://doi.org/10.1201/9781420073072, 2009.

Supplemental Information for:

Marine gas-phase sulfur emissions during an induced phytoplankton bloom

5 Delaney B. Kilgour¹, Gordon A. Novak¹, Jon S. Sauer², Alexia N. Moore², Julie Dinasquet³, Sarah Amiri², Emily B. Franklin⁴, Kathryn Mayer², Margaux Winter⁵, Clare K. Morris³, Tyler Price², Francesca Malfatti^{3,6}, Daniel R. Crocker², Christopher Lee³, Christopher D. Cappa⁷, Allen H. Goldstein^{4,8}, Kimberly A. Prather^{2,3}, and Timothy H. Bertram¹

¹Department of Chemistry, University of Wisconsin – Madison, Madison, WI 53706, USA

- ²Department of Chemistry and Biochemistry, University of California, San Diego, La Jolla, CA 92093, USA
 ³Scripps Institution of Oceanography, University of California, San Diego, La Jolla, CA 92037, USA
 ⁴Department of Civil and Environmental Engineering, University of California, Berkeley, CA 94720, USA
 ⁵Department of Chemistry and Chemical Biology, Harvard University, Cambridge, MA 02138, USA
 ⁶Department of Life Sciences, Universita' degli Studi di Trieste, Trieste, Italy 34127
- ⁷Department of Civil and Environmental Engineering, University of California, Davis, CA 95616, USA ⁸Department of Environmental Science, Policy and Management, University of California, Berkeley, CA 94720, USA

Correspondence to: Timothy H. Bertram (timothy.bertram@wisc.edu)

S1 SIO Wave Channel

Seawater used in the wave channel was pumped from below the Ellen Browning Scripps Memorial Pier in La Jolla, CA (32-

- 30 52'00" N, 117-15'21" W) between 08:00 and 13:00 PST on July 23, 2019. The seawater intake was located 300 m offshore and 2 m above the ocean floor. Large pieces of organisms, such as seaweeds, were filtered out prior to the water being stored in a gravity flume on the pier. Water from the gravity flume was pumped (Grundfos UNILIFT AP12.40.04.A1) through a fire hose and a 100-micron nitex nylon mesh (Flystuff; Cat# 57-103) to remove bulk sediment before being transported in 300 gallon tanks to the SIO wave channel five minutes away. Prior to being pumped into the wave channel, the water was filtered
- 35 again using a 50-micron nitex nylon mesh (Flystuff; Cat# 57-106) to prevent zooplankton from entering the wave channel. This process was continued until the glass-walled wave channel (33 m length x 0.5 m width x 1 m height) was filled with approximately 13,000 L of seawater. To simulate the wave-breaking mechanism of the ocean, waves were generated by a paddle and forced to break over an inclined ramp in the channel, simulating a beach (Prather et al., 2013; Wang et al., 2015).

S2 Mesocosm Experiment Nutrient Additions and Perturbations

40 A record of nutrient additions and changes that affected the wave channel seawater and bloom progression are outlined in Table S1 below. "f Medium" nutrient additions, diluted by 2 (f/2) or 20 (f/20) from the original formulation were added to stimulate algae growth (Guillard and Ryther, 1962). The phytoplankton tows and outdoor tank addition added healthy phytoplankton biomass grown under natural sunlight to increase the likelihood a bloom occurred.

S3 Dissolved DMSP and DMS Measurements

45 S3.1 Sample Preparation

For waterside DMSP and DMS measurements, sample preparation and cryogenic purge and trap methods closely followed Wurl (2009). All tubing, syringes, and containers were rigorously cleaned daily with 0.1% HCl, ethanol, propanol, and Milli-Q (Millipore) water for sample and standard preparation. Samples of seawater for DMSP and DMS analysis were gathered from the wave channel by siphoning seawater ~5 cm below the water surface through ¼" ID PTFE tubing into a 250 mL PTFE

- 50 media bottle which was filled to overflow before closure. For DMSP_p analysis, labeled as sample 1 in Equations 1-3, a ~0.1 g pellet of high-purity NaOH (Sigma Aldrich) was added directly to 5 mL of unfiltered seawater in a 15 mL headspace vial which was then immediately crimp capped with a PTFE septum. For DMSP_d, labeled as sample 2, a 5 mL seawater aliquot was gently syringe filtered using a 0.7 μm Whatman syringe filter into a 15 mL headspace vial after which a ~0.1 g pellet of NaOH was added before sealing. For DMS analysis, labeled as sample 3, seawater was syringe filtered (0.7 μm) and analyzed
- 55 less than an hour after collection. Both DMSP_d and DMSP_p vials were left to rest overnight in a dark location to allow cleavage of DMSP to DMS before purge and trap analysis the following day. External standards of dissolved DMS (Sigma Aldrich) were freshly prepared and analyzed daily ranging from 0.1 to 5 nM. For samples of DMSP which exceeded the external

calibration, replicate samples were used to prepare dilutions in the calibration range. A sample CI-TOFMS chromatogram of dissolved organosulfur compounds and the external DMS calibration are in Fig. S2.

60 S3.2 Cryogenic Purge and Trap

Two 22 gauge needles were used to pierce the headspace vial septa with one needle positioned at the bottom of the vial and the second in the vial headspace. At a defined start time, headspace vials were bubbled with 80 sccm UHP helium (Praxair) regulated by mass flow controller (MKS) for 4 minutes. Purged gases leaving through the second needle into PTFE tubing were directed through a Nafion drier (MD-050) to remove water vapor and then passed through a Teflon loop immersed in

65 liquid nitrogen. At the end of the purging cycle, a 6 port valve (Vici - 6UWE) was switched and the trapping loop was removed from the dewar. A carrier flow of 20 sccm helium then passed through the trapping loop, eluting the trapped gases to the analyzer inlet as the loop equilibrated to room temperature. Elution gases were diluted by 2.0 slpm N₂ before analysis.

S3.3 Chemical Ionization Time-of-Flight Mass Spectrometry (CI-TOFMS) of Dissolved Sulfur Compounds

Eluted gases were measured on a chemical ionization time-of-flight mass spectrometer using benzene cluster cation ionization
which has been detailed previously (Lavi et al., 2018; Kim et al., 2016). Briefly, ~300 ppm benzene vapor was generated by passing 10 sccm N₂ vapor over a cylinder of liquid benzene and diluted with excess nitrogen to 1.9 slpm. Benzene vapor was then passed through a 20mCi Po-210 alpha source which generates benzene cluster cations. Two inline critical orifices (O'Keefe) controlled flow of benzene cluster cation vapor and analyte gases which were drawn at 1.8 slpm into an ion molecule reactor (IMR) which was held at 60 Torr, 50 V, and 50 °C. Resultant ions were directed into an electrodynamic ion funnel
which focuses and gently declusters the ion beam into a RF-only transfer quadrupole before detection by a lower resolving power (m/Δm ≈ 1200) time-of-flight mass spectrometer operating at 10 Hz. Peak area for m/Q 62 [C₂H_oS]⁺ was analyzed in the Tofware graphic user interface for Igor Pro 7. Calculation of concentrations of DMSP_p and DMSP_d were as follows using

the m/z 62 ion peak area from each sample:

80	$[DMSP_p] = Sample \ 1 - Sample \ 2 - Sample \ 3$	(1)
	$[DMSP_d] = Sample \ 2 - Sample \ 3$	(2)
	[DMS] = Sample 3	(3)

S4 Bacterial Abundance and Production Measurements and Phytoplankton Enumeration

Samples for bacterial abundance (1.0 mL) were fixed with electron microscopy grade glutaraldehyde (5% v/v) and stored at -

85 80°C after being stored at 4°C for 15 minutes and flash frozen in liquid nitrogen. Counts were performed on a FACSCanto II flow cytometer (Becton Dickinson) after SYBRGreen I staining according to Gasol and Del Giorgio (2000).

Bacterial productivity was measured by [H³]-leucine incorporation (Kirchman et al., 1985) modified for microcentrifugation (Azam and Smith, 1992). Triplicate 1.7 mL aliquots were incubated with [H³]-leucine (20 nM final concentration) for 1 hour.

Samples with 100% trichloroacetic acid added prior to $[H^3]$ -leucine addition served as blanks. Leucine incorporation was 90 converted to carbon production assuming 3.1 kg C (mol leucine)⁻¹ (Simon and Azam, 1989).

The range of bacterial sulfur demand was estimated from bacterial productivity using the cellular C:S ratios of 86 (Fagerbakke et al., 1996) and 248 as a range (Cuhel et al., 1982). Bacterial carbon demand was estimated considering an average bacterial growth efficiency of 0.3 (del Giorgio and Cole, 1998).

95

Phytoplankton functional groups were enumerated by allowing 50 mL of collected seawater to settle into a settlement chamber using the Utermöhl method (Edler and Elbrächter, 2010) and analyzed with an Olympus IX-71 inverted microscope.

S5 DMS:MeSH Correlation with Salinity

100 It has been shown previously that DMS:MeSH may be controlled by seawater salinity (Salgado et al., 2014; Magalhães et al., 2012). Controlled experiments in which salt and DMSP were added to water samples and headspace gases were measured with gas chromatography, showed more DMS accumulated at low salinities and more MeSH accumulated at high salinities. The authors hypothesized that more DMSP is stored in cells at high salinity, causing the DMSP demethylation/demethiolation pathway to be favored over the DMSP cleavage pathway (Magalhães et al., 2012; Salgado et al., 2014). This suggests that 105 DMS:MeSH should be inversely correlated with salinity.

Figure 5 shows a strong inverse correlation ($R^2 = 0.72$) between DMS:MeSH and salinity measurements that roughly follows linearly with days of the bloom. The salinity decline during the bloom is primarily due to the additions of ultrapure water to the wave channel meant to maintain the water level and outweigh the influence of evaporation. The change in salinity and DMSP concentrations over the 21-day bloom (1 psu; <150 nM) were significantly smaller than the salinity range and DMSP 110 concentrations tested in controlled experiments reported in the literature (30 psu; <500 µM) (Salgado et al., 2014). A plot of DMS:MeSH as a function of salinity based on data in Salgado et al. (2014) showed the largest rate of change occurred below

15 psu salinity for these experiments, lower values than observed in the wave channel. Further, the small change in salinity observed in the wave channel is within the natural fluctuations off of Scripps Pier in summer months (Automated Shore

115 Stations, 2021). As a result, we hypothesize that the observed correlation between DMS:MeSH and salinity during SeaSCAPE is simply a correlation and not indicative of a causal relationship between salinity and DMS:MeSH.

Day	Action
7/23/2019 08:00-13:00 (Day 0)	Flume fill
7/25/2019 17:00 (Day 2.2)	Nutrient addition – f/20 growth media and f/40 sodium metasilicate
7/26/2019 17:47 (Day 3.2)	Nutrient addition - additional silicates to bring up to f/20 sodium
	metasilicate
7/28/2019 17:00 (Day 5.2)	Outdoor tank – 300 gallons of freshly collected seawater (filtered in same
	manner as wave channel water) were spiked with f/2 growth media and
	silicates. Tank was placed outside in partial shade with a mesh screen.
8/1/2019 10:30 (Day 8.9)	<i>Outdoor tank addition</i> – Outdoor tank added to wave flume via buckets.
8/1/2019 10:45 (Day 8.9)	Nutrient addition - Additional growth media and silicates, to bring total
	concentration of both up to f/2
8/5/2019 8:15-10:30 (Day 12.8 -	Wall cleaning - Removed algal growth from flume walls so that light
12.9)	could reach interior
8/9/2019 17:00 (Day 17.2)	Mixing pumps – Placed on the floor of the flume to better mix organic
	material
8/13/2019 17:00 (Day 21.2)	Measurements end

Table S2: Resolvable sulfur-containing ions in the Vocus PTR-ToF-MS mass spectrum present at signal-to-noise ratio above 3 during the campaign. Excluding DMS (C₂H₆SH⁺), MeSH (CH₄SH⁺), and benzothiazole (C₇H₅NSH⁺), all were included in the "Other Sulfur" signal.

Ion	Exact Mass	Ion Identification	Early Bloom (Days 1-9)	Late Bloom
			Mean $\pm 1\sigma$ (ppt)	(Days 10-21)
				Mean $\pm 1\sigma$ (ppt)
H_3S^+	34.995		0.11 ± 0.059	0.35 ± 0.20
CH ₄ SH ⁺	49.0107	MeSH	140. ± 44.6	98.6 ± 82.3
$C_2H_6SH^+$	63.0263	DMS	651 ± 297	$2.08 \text{ x } 10^3 \pm 1.28 \text{ x } 10^3$
HNOSH ⁺	63.9852		0.43 ± 0.19	0.84 ± 0.41
SO_2H^+	64.9692		0.21 ± 0.12	0.25 ± 0.16
C3H6SH ⁺	75.0263		3.9 ± 5.0	6.5 ± 4.6
$C_2H_5NSH^+$	76.0216		0.15 ± 0.45	0.25 ± 0.28
CS_2H^+	76.9514		0.10 ± 0.073	0.055 ± 0.021
CHO ₂ SH ⁺	77.977		0.19 ± 0.16	0.37 ± 0.17
C ₃ H ₇ OS ⁺	91.0212		0.63 ± 1.6	1.3 ± 2.3
$C_2H_6S_2H^+$	94.9984	DMDS	3.9 ± 3.0	9.4 ± 4.3
$C_2H_6O_2SH^+$	95.0161	DMSO ₂	$25 \pm 30.$	3.1 ± 8.0
CH ₅ NO ₂ SH ⁺	96.0114	MSAM	0.47 ± 0.61	0.49 ± 0.40
C ₂ H ₆ NO ₂ S ⁺	108.011		0.095 ± 0.19	0.22 ± 0.19
C5H8NS ⁺	114.037		0.70 ± 1.4	0.89 ± 1.2
$C_4H_8SO_2H^+$	121.032		56 ± 52	26 ± 17
C2H6S3H ⁺	126.9704		0.090 ± 0.18	0.16 ± 0.40
C7H5NSH ⁺	136.0215	Benzothiazole	112 ± 67.3	146 ± 47.3
C7H5NOSH ⁺	152.0165	2(3H)-	0.86 ± 0.52	1.2 ± 0.15
		Benzothiazolone		
C9H14SH ⁺	155.0889		0.73 ± 0.91	0.89 ± 1.3
C5H2O4SH ⁺	158.975		0.089 ± 0.16	0.021 ± 0.037
C9H20SH ⁺	161.1358		0.16 ± 0.25	0.60 ± 0.72
C5H7O4S ⁺	163.006		0.65 ± 1.0	0.72 ± 0.36
$C_7H_5NS_2H^+$	167.9936		0.72 ± 0.082	0.38 ± 0.11
$C_{10}H_{16}SH^+$	169.1045		8.3 ± 3.2	5.6 ± 0.90

C9H7NOSH ⁺	178.0321	2-acetyl- benzothiazole	0.79 ± 0.31	0.97 ± 0.18
$C_9H_6O_2SH^+$	179.0161		0.062 ± 0.11	0.20 ± 0.086
$C_{11}H_{16}SH^+$	181.105		7.5 ± 2.8	4.8 ± 0.97



Figure S1: Picture of the isolated sampling vessel (ISV) used for gas sampling. Water enters the ISV through the water entry port after being pumped from the wave channel using a peristaltic pump and Tygon tubing. The water entry port is located on the far PTFE disk of the ISV, where the curved fitting allows for water to be introduced in a plunging jet motion. Zero air enters the ISV through a ¼'' stainless steel Swagelok bulkhead fitting next to the water inlet. The near side of the ISV shows four ¼'' stainless steel Swagelok bulkhead fittings arranged in a semicircle at the top of the PTFE disk. These were used for gas sampling. All gas tubing

165 Swagelok bulkhead fittings arranged in a semicircle at the top of the PTFE disk. These were used for gas sampling. All gas tubing used in the experiment is ¹/₄" O.D. PFA. Water drains back into the wave channel through the 1" water outlet port located half way up the PTFE disk. The ISV is made of borosilicate glass with O.D., thickness, and length of 400 mm, 6 mm, and 73.66 cm respectively, leading to an ISV volume of 0.0871 m³. In operation, the ISV was approximately half filled with water (0.0436 m³). The zero air flow rate through the headspace was 8-10 slpm (average residence time of 5 minutes), and water flow rate was 1.5 slpm (water residence time 29 minutes).



185 Figure S2: (a) Typical cryogenic purge and trap CI-TOFMS chromatogram of SeaSCAPE seawater featuring multiple organosulfur peaks. (b) Example daily DMS external calibration of cryogenic purge and trap CI-TOFMS.



Figure S3: Example calibration curves for DMS, MeSH, and benzothiazole diluted in dry zero air. Average dry sensitivities for DMS, MeSH, and benzothiazole are 3.0, 1.0, and 5.8 cps ppt⁻¹, respectively.



230 Figure S4: (a) DMS, (b) MeSH, and (c) benzothiazole fragments present in calibrations are displayed. The fragment ions are not included in the "Other Sulfur" ions, as they are not unique molecules.





Figure S5: There exists no correlation between MeSH and DMDS during SeaSCAPE, despite a small dependence of DMDS on MeSH during post-SeaSCAPE calibrations.



Figure S6: (a) Time series of DMSO₂. Correlations between (b) DMSO₂ and (c) methane sulfonamide (MSAM) with DMS are weak and represent minor contributors to the total "Other S" signal. DMSO is not detectable above signal-to-noise ratio 3 for any day of SeaSCAPE. (d) MSAM and DMSO₂ are also weakly anti-correlated (r = -0.48), in contrast to what was observed in the Arabian Sea (Edtbauer et al., 2020).

285



Figure S7: Time series of DMSP_t:chlorophyll-a, suggesting the bloom was dominated by a community of low DMSP producers, and DMS_w:DMSP_t, showing a low conversion efficiency of DMSP_t to DMS.



Figure S8: Time series in experiment day of the relative abundance of diatom aggregates, diatoms, and other taxa. Diatoms were dominant after day six.



Figure S9: Regression between methionine, leucine, and serine aminopeptidase activity with DMS:MeSH, suggesting more protein degradation with higher DMS:MeSH.

References

375

Automated Shore Stations: https://sccoos.org/data/autoss/, last access: 19 May 2021.

Azam, F. and Smith, D. C.: A simple, economical method for measuring bacterial protein synthesis rates in seawater using 3H-leucine, Mar. Microb. Food Webs, 6, 107–114, 1992.

355 Cuhel, R. L., Taylor, C. D., and Jannasch, H. W.: Assimilatory Sulfur Metabolism in Marine Microorganisms: Sulfur Metabolism, Protein Synthesis, and Growth of Alteromonas luteo-violaceus and Pseudomonas halodurans During Perturbed Batch Growtht, APPL Env. MICROBIOL, 43, 9, 1982.

Edler, L. and Elbrächter, M.: The Utermöhl method for quantitative phytoplankton analysis, in: Microscopic and Molecular Methods for Quantitative Phytoplankton Analysis, 13–20, 2010.

- 360 Edtbauer, A., Stönner, C., Pfannerstill, E. Y., Berasategui, M., Walter, D., Crowley, J. N., Lelieveld, J., and Williams, J.: A new marine biogenic emission: methane sulfonamide (MSAM), DMS and DMSO measured in air over the Arabian Sea, Gases/Field Measurements/Troposphere/Chemistry (chemical composition and reactions), https://doi.org/10.5194/acp-2019-1021, 2020.
- Fagerbakke, K., Heldal, M., and Norland, S.: Content of carbon, nitrogen, oxygen, sulfur and phosphorus in native aquatic and cultured bacteria, Aquat. Microb. Ecol., 10, 15–27, https://doi.org/10.3354/ame010015, 1996.

Gasol, J. M. and Del Giorgio, P. A.: Using flow cytometry for counting natural planktonic bacteria and understanding the structure of planktonic bacterial communities, Sci. Mar., 64, 197–224, https://doi.org/10.3989/scimar.2000.64n2197, 2000.

del Giorgio, P. A. and Cole, J. J.: BACTERIAL GROWTH EFFICIENCY IN NATURAL AQUATIC SYSTEMS, Annu. Rev. Ecol. Syst., 29, 503–541, https://doi.org/10.1146/annurev.ecolsys.29.1.503, 1998.

370 Guillard, R. R. L. and Ryther, J. H.: STUDIES OF MARINE PLANKTONIC DIATOMS: I. CYCLOTELLA NANA HUSTEDT, AND DETONULA CONFERVACEA (CLEVE) GRAN., Can. J. Microbiol., 8, 229–239, https://doi.org/10.1139/m62-029, 1962.

Kim, M. J., Zoerb, M. C., Campbell, N. R., Zimmermann, K. J., Blomquist, B. W., Huebert, B. J., and Bertram, T. H.: Revisiting benzene cluster cations for the chemical ionization of dimethyl sulfide and select volatile organic compounds, Atmospheric Meas. Tech., 9, 1473–1484, https://doi.org/10.5194/amt-9-1473-2016, 2016.

Kirchman, D., K'Nees, E., and Hodson, R.: Leucine Incorporation and Its Potential as a Measure of Protein Synthesis by Bacteria in Natural Aquatic Systemst, APPL Env. MICROBIOL, 49, 9, 1985.

Lavi, A., Vermeuel, M. P., Novak, G. A., and Bertram, T. H.: The sensitivity of benzene cluster cation chemical ionization mass spectrometry to select biogenic terpenes, Atmospheric Meas. Tech., 11, 3251–3262, https://doi.org/10.5194/amt-11-3251-2018, 2018.

- Magalhães, C., Salgado, P., Kiene, R. P., and Bordalo, A. A.: Influence of salinity on dimethyl sulfide and methanethiol formation in estuarine sediments and its side effect on nitrous oxide emissions, Biogeochemistry, 110, 75–86, https://doi.org/10.1007/s10533-011-9690-z, 2012.
- Prather, K. A., Bertram, T. H., Grassian, V. H., Deane, G. B., Stokes, M. D., DeMott, P. J., Aluwihare, L. I., Palenik, B. P.,
 Azam, F., Seinfeld, J. H., Moffet, R. C., Molina, M. J., Cappa, C. D., Geiger, F. M., Roberts, G. C., Russell, L. M., Ault, A.
 P., Baltrusaitis, J., Collins, D. B., Corrigan, C. E., Cuadra-Rodriguez, L. A., Ebben, C. J., Forestieri, S. D., Guasco, T. L.,

Hersey, S. P., Kim, M. J., Lambert, W. F., Modini, R. L., Mui, W., Pedler, B. E., Ruppel, M. J., Ryder, O. S., Schoepp, N. G., Sullivan, R. C., and Zhao, D.: Bringing the ocean into the laboratory to probe the chemical complexity of sea spray aerosol, Proc. Natl. Acad. Sci., 110, 7550–7555, https://doi.org/10.1073/pnas.1300262110, 2013.

390 Salgado, P., Kiene, R., Wiebe, W., and Magalhães, C.: Salinity as a regulator of DMSP degradation in Ruegeria pomeroyi DSS-3, J. Microbiol., 52, 948–954, https://doi.org/10.1007/s12275-014-4409-1, 2014.

Simon, M. and Azam, F.: Protein content and protein synthesis rates of planktonic marine bacteria, Mar. Ecol. Prog. Ser., 51, 201–213, https://doi.org/10.3354/meps051201, 1989.

Wang, X., Sultana, C. M., Trueblood, J., Hill, T. C. J., Malfatti, F., Lee, C., Laskina, O., Moore, K. A., Beall, C. M.,
McCluskey, C. S., Cornwell, G. C., Zhou, Y., Cox, J. L., Pendergraft, M. A., Santander, M. V., Bertram, T. H., Cappa, C. D.,
Azam, F., DeMott, P. J., Grassian, V. H., and Prather, K. A.: Microbial Control of Sea Spray Aerosol Composition: A Tale of
Two Blooms, ACS Cent. Sci., 1, 124–131, https://doi.org/10.1021/acscentsci.5b00148, 2015.

Wurl, O. (Ed.): Practical Guidelines for the Analysis of Seawater, CRC Press, https://doi.org/10.1201/9781420073072, 2009.