

Influence of ~~the Change in~~ Total Ozone Column (TOC) on the Occurrence of Tropospheric Ozone Depletion Events (ODEs) in the Antarctic

Le Cao^{1,*}, Linjie Fan^{1,2,*}, Simeng Li¹, and Shuangyan Yang²

¹Key Laboratory for Aerosol-Cloud-Precipitation of China Meteorological Administration, Nanjing University of Information Science and Technology, Nanjing 210044, China

²Key Laboratory of Meteorological Disaster, Ministry of Education (KLME)/Joint International Research Laboratory of Climate and Environmental Change (ILCEC)/Collaborative Innovation Center on Forecast and Evaluation of Meteorological Disasters (CIC-FEMD), Nanjing University of Information Science and Technology, Nanjing 210044, China

*These authors contributed equally to this work.

Correspondence: L. Cao

(le.cao@nuist.edu.cn)

Abstract. The occurrence of the tropospheric ozone depletion events (ODEs) in the Antarctic can be influenced by ~~the change in many factors such as the~~ Total Ozone Column (TOC). In this study, we ~~combined~~ analyzed the observational data obtained from ground observation stations ~~with and used~~ two numerical models (TUV and KINAL), to ~~figure out~~ discover the relationship between the TOC ~~change~~ and the occurrence of ODEs in the Antarctic. A sensitivity analysis was also performed on ~~the change in~~ ozone and major bromine species (BrO, HOBr and HBr) to find out key photolysis reactions determining the impact on the occurrence of tropospheric ODEs brought by ~~the change in~~ TOC. From the analysis of the observational data and the numerical results, we suggested that the occurrence frequency of ODEs in the Antarctic ~~seems negatively correlated with the variation of TOC is negatively associated with TOC, after screening out the impact on ODEs caused by solar zenith angle (SZA). This negative impact of TOC on the occurrence of ODEs was suggested to be exerted through altering the solar radiation reaching the ground surface and changing the rates of photolysis reactions.~~ Moreover, major ODE accelerating reactions (i.e., photolysis of ozone, H₂O₂ and HCHO) and decelerating reactions (i.e., photolysis of BrO and HOBr), which heavily control the start of ODEs, were also identified. ~~It was~~ We found that when TOC ~~varies~~ decreases, the major ODE accelerating reactions speed up significantly, ~~while~~. In contrast, the major ODE decelerating reactions are only slightly affected, ~~thus leading to the negative dependence of the ODE occurrence on the change in~~. As a result of the different impacts of TOC on photolysis reactions, the occurrence of ODEs depends negatively on TOC.


1 Introduction

Ozone is a short-lived trace gas in the atmosphere, with about 90% located in the stratosphere and 10% in the troposphere (Seinfeld and Pandis, 2006; Akimoto, 2016). In the stratosphere, ozone plays a role in absorbing the ultraviolet (UV) radiation from the sun, thus protecting the lives on the earth. In contrast, ozone in the troposphere is a pollutant. It causes eye irrita-

tions and disorders of the lung function of human beings at a high concentration (Lippmann, 1991). Moreover, ozone in the troposphere also acts as a greenhouse gas, contributing to the global warming (Seinfeld and Pandis, 2006). It was suggested by Toumi et al. (1996) that the tropospheric ozone mostly originates from three sources: downward entrainment from the stratosphere, photochemical reactions occurring in the troposphere, and the vertical convection. Thus, the amount of ozone in the troposphere can be affected by many factors such as the variation of the stratospheric ozone.

Ozone in polar regions is always a focus of the scientific community. Due to the special geographical location and the unique environment, polar regions are also called “natural laboratory” of the earth (Heinemann, 2008). Moreover, because polar regions especially the Antarctic are hardly affected by anthropogenic activities, the climate of polar regions is capable of reflecting the global change of the climate (Prather and Jaffe, 1990). In 1980s, ~~a~~an extraordinary event, i.e., ozone hole, was found occurring over the Antarctic (Farman et al., 1985). This event refers to a continuous decline in the total ozone amount over the Antarctic during the springtime of every year. Because the majority of ozone in the atmosphere resides in the stratosphere, the Antarctic ozone hole mostly represents a depletion of the stratospheric ozone. After the discovery of the ozone hole, large efforts were made to reveal the reasons causing the emergence of this event, such as ~~figuring out~~discovering the role of chlorofluorocarbons (CFCs) from human activities (Molina and Rowland, 1974; Bedjanian and Poulet, 2003), heterogeneous reactions on the surface of PSCs (polar stratospheric clouds) and the photolysis of ClO dimer, i.e., ClOOC1 at polar night (Finlayson-Pitts and Pitts, 1999; Brasseur and Solomon, 2005). Features of the ozone loss were also revealed using total column measurements from ground-based stations in the Antarctic (Kuttiapurath et al., 2010).

Similar to the ozone hole phenomenon representing a depletion of the stratospheric ozone, in the 1980s, an ozone depletion event (ODE) was also observed in the troposphere of polar regions (Oltmans, 1981). It was ~~firstly~~first reported that during the springtime of Arctic, the surface ozone drops from a normal level (~ 40 ppb) to less than 1 ppb within a few hours or 1-2 days. After that, the tropospheric ODE was also reported occurring in the coastal areas of Antarctic (Kreher et al., 1997; Frieß et al., 2004; Wagner et al., 2007). Subsequent studies suggested that the tropospheric ODE is a common phenomenon that occurs in the atmospheric boundary layer during the springtime of both the Arctic and the Antarctic. It was also reported by Roscoe and Roscoe (2006) that the tropospheric ODE occurs in the Antarctic since as early as the 1950s. Following studies suggested that the occurrence of the tropospheric ODE is ~~related to~~driven by an auto-catalytic reaction cycle involving bromine species at polar sunrise during polar spring, as follows (Simpson et al., 2007):



This bromine-involved reaction cycle includes heterogeneous reactions occurring on substrates such as the snow-/ice-covered ground surface and the suspended aerosols. Through reaction cycle (I), bromide ions (Br^-) are activated from the substrates ; ~~and the total bromine amount in the troposphere is thus elevated. The bromine released to the atmosphere and then released into~~

50 the atmosphere in the form of Br₂. In the presence of sunlight, Br₂ is photolyzed to be Br atoms, which then consumes ozone
near the ground, leading to the occurrence of the tropospheric ODEs. ~~This ozone-depleting cycle is thus called~~ Thus, the net
effect of reaction cycle (I) is converting the surface ozone in polar regions into O₂. Meanwhile, due to the activation of bromine
ions from the substrates, the total bromine amount in the troposphere is also exponentially elevated, which is thus called as
“bromine explosion mechanism” (Platt and Janssen, 1995; Platt and Lehrer, 1997; Wennberg, 1999). Therefore, necessary
55 conditions required for the occurrence of ODEs include the existence of substrates such as the snow-/ice-covered surfaces and
the suspended aerosols and the presence of sunlight (Lehrer et al., 2004).

Apart from the bromine chemistry, the occurrence of the tropospheric ODE was also found to be determined by many factors
such as :- (1) Temperature. Tarasick and Bottenheim (2002) examined historical ozonesonde records at three Canadian stations
over the time period 1966-2000. They suggested that a low temperature (<-20°C) is probably a necessary condition for the
60 occurrence of ODEs, because heterogeneous reactions that activate bromide from substrates and the formation of frost flowers
are favored under this cold condition. However, in a later analysis of ozone data obtained from a transpolar drift, Bottenheim
et al. (2009) found the temperature well above -20°C during the most persistent ~~ozone depletion~~ ODE period over the Arctic
Ocean. It was also reported by Koo et al. (2012) that there is no ~~evidence in observations~~ observational evidence for the
threshold value of temperature for the occurrence of ODEs. Instead, they suggested the variability of temperature potentially
65 an important factor for the ~~depletion of ozone~~ occurrence of ODEs. (2) Passing of pressure systems. By analyzing the values
of ozone mixing ratio and meteorological parameters from balloon sondes during the 1994 Polar Sunrise Experiment (PSE94),
Hopper et al. (1998) suggested that the occurrence of ODEs in the Arctic is strongly correlated with high-pressure systems. This
dependence of the ozone decline on pressure systems was also confirmed by Jacobi et al. (2010) who proposed that mesoscale
synoptic systems are able to transport air masses with low ozone mixing ratio to the observational site, leading to the detection
70 of ODEs at the Arctic coastal stations. It was also suggested by Boylan et al. (2014) that the transport caused by synoptic
patterns acts as the major factor for the occurrence of ODEs at Barrow, Alaska, rather than the change in local meteorological
parameters. In contrast, Jones et al. (2006) analyzed the observational data of ozone and meteorological parameters obtained at
Halley station in coastal Antarctica, and they ~~figured out~~ found that in Antarctica, the occurrence of ODEs is highly associated
with low pressure systems, denoting the remarkable differences in the atmospheric system between the Arctic and the Antarctic.
75 (3) Formation of fresh sea ice. Based on sea ice maps obtained from satellite detection, Bottenheim et al. (2009) ~~figured out~~
identified regions of the Arctic Ocean as the origin of the tropospheric ozone depletion. In these regions, open leads, polynyas
and fresh sea ice are frequently formed, which favors the release of bromine and thus the depletion of the tropospheric ozone.
These ODE-originating regions proposed by Bottenheim et al. (2009) are also consistent with the “cold spots” discovered in a
previous study of Bottenheim and Chan (2006) where the depletion of the tropospheric ozone possibly initiates and develops.
80 The connection between the ODE occurrence and the formation of fresh sea ice was also identified by Jones et al. (2006) by
revealing that the air masses causing rapid ~~ozone depletion~~ ODEs at Halley station originate in a region where a large amount
of fresh sea ice is formed. (4) Other factors. ODEs were also found to be impacted by the presence of mixed-phase clouds in
the boundary layer due to the cloud-top radiative cooling (Hu et al., 2011) and regional climate variability such as the Western
Pacific (WP) teleconnection pattern (Koo et al., 2014).

85 Although there exist many studies discussing the determining factors for the occurrence of the tropospheric ODE, the impact caused by the ~~change in the~~ stratospheric ozone on the occurrence of the tropospheric ODE has not been thoroughly investigated yet. In previous studies, most of often, ozone ~~change in the~~ stratosphere and the tropospheric ODEs were investigated separately. However, ODEs can be strongly influenced by the ~~change in the~~ stratospheric ozone. For example, the ~~ozone-lacking air in the boundary layer can be partly replenished by the ozone-rich air due to the downward entrainment from the lower stratosphere~~ (Kuang et al., 2017). However, because of the strong stability of the polar boundary layer during ODEs, ~~this influence might be minor. Moreover, the~~ variation of the stratospheric ozone would ~~also~~ lead to a change in the solar radiation reaching the ground surface, thus affecting the rates of photolysis reactions in the troposphere. As a result, the lifetimes of many atmospheric constituents in the troposphere and the occurrence frequency of the tropospheric ODEs can be altered. ~~But it is still unclear whether the change in~~ However, the exact mechanism of how the stratospheric ozone ~~will foster or retard the occurrence of the tropospheric ODE~~ affects the tropospheric ODEs is still unclear to the present. Therefore, in the present ~~this~~ study, we combined the observational data from ground observation stations with two numerical models, to ~~figure out the impacts~~ discover the impact on the occurrence of tropospheric ODEs in the Antarctic brought about by the ~~change in the~~ total ozone amount including the stratospheric ozone. A concentration sensitivity analysis was also performed to reveal photolysis reactions heavily responsible for this impact on the tropospheric ODE.

100 The structure of the manuscript is as follows. In Sect. 2, observational data analyzed in the present study are described. The numerical models and the governing equations are also presented in this section. In Sect. 3, results obtained from the analysis of the observational data and the computations of the numerical models are shown and discussed. At last, in Sect. 4, conclusions achieved in this study are summarized, and studies to be made in the future are also prospected.

2 Description of the Observational Data and the Numerical Method

105 In the present study, we first analyzed the observational data of ozone obtained from ground observation stations, and then tried to ~~figure out~~ discover the relationship between the ~~variation of the~~ total ozone column (TOC) and the occurrence frequency of tropospheric ODEs. Then we ~~took the Halley station as an example, and~~ applied two numerical models, TUV model and KINAL box model, to capture the temporal ~~change in~~ variations of ozone and major bromine species during ODEs ~~under the conditions of the Halley station~~. Sensitivity tests were also performed in these models to ~~figure out the influence caused by the change in the total ozone column (TOC)~~ discover the influence exerted by TOC on the time ~~variation~~ variations of these air constituents (i.e., ozone and bromine species). Then the most influential photolysis reactions were identified through a concentration sensitivity analysis on ~~the change in~~ these air constituents.

110

2.1 Observational Data of Ozone

Two different types of ozone data were used in the present study. One is the total ozone column (TOC) obtained from ground-based observations. This kind of data mainly indicates the total ozone amount in a vertical column extending from the ground surface to the top of the atmosphere. Because approximately 90% of ozone in the atmosphere resides in the stratosphere, ~~the~~

115

~~change in~~ TOC can heavily reflect the ~~variation amount~~ of the stratospheric ozone. The other type of the ozone data is the near-surface ozone mixing ratio recorded at ground observation stations, and it can partly represent the ozone concentration in the atmospheric boundary layer. The details of these ozone data are given below.

120 2.1.1 Total Ozone Column (TOC)

The TOC data used in this study were obtained from World Ozone and Ultraviolet Radiation Data Center of Canada (WOUDC, <https://woudc.org/home.php>) for all the registered stations in the Antarctic. These TOC data were observed using a Dobson instrument (Balis et al., 2007), and cover a time span from the year 2000 to 2016. The time resolution of these data is 1 day. After filtering out stations that possess only out-dated or incomplete data, we picked ~~up out~~ Halley station (75.52°S, 26.73°W), which
125 has the most complete TOC data, ~~as an example to indicate the typical temporal change in TOC for the present investigation.~~
Moreover, TOC data from Faraday-Vernadsky (FAD) station (65.25°S, 64.27°W) were also used for comparison. The FAD station is located in the Antarctic Peninsula area, which is to the northwest of the Halley station. Thus, TOC detected at the FAD station is more capable of reflecting the conditions of the Weddell Sea, which is between the Halley station and the FAD station. In order to guarantee the representativeness of the ~~data from this station, we compared the time series of TOC~~
130 data from these stations, we also compared the TOC recorded at the Halley station with that obtained at a station nearby, Belgrano II (77.88°S, 34.63°W) (see Tab. S1 in the supplementary material). It can be seen from Tab. S1 that the correlation coefficients between the TOCs obtained at the Halley station and ~~other stations~~ the Belgrano II station mostly possess a value above 0.9, indicating that the TOC ~~data~~ obtained at the Halley station can represent the typical TOC variation surrounding Halley. Moreover, we also validated the observed TOC obtained at the FAD station using the observations from Marambio
135 station (64.24°S, 56.62°W), which is located on the northeast side of the Antarctic Peninsula, and we found the correlation between these two data high (also see Tab. S1 in the supplementary material), which ensures the validity of the TOC observed at the FAD station. In addition, we also calculated the correlation coefficient between TOCs observed at the Halley station and the ~~Faraday-Vernadsky station might be caused by the special geographical location of the Faraday-Vernadsky station, which is out of the scope of the present study.~~
140 FAD station, and the value of the correlation coefficient mostly resides in a range of
0.3-0.8. The difference between the observed TOCs at these two stations might be caused by atmospheric dynamics.

2.1.2 Surface Ozone

After choosing the Halley station as an example ~~representing the typical TOC change in the Antarctic for the present investigation,~~
we adopted the surface ozone data at the Halley station from World Data Center for Greenhouse Gases (WDCGG, <https://gaw.kishou.go.jp>),
~~and the.~~ The surface ozone data used in this study can also be found in Section “Code and data availability” of this paper and
145 from the data archive provided by World Data Centre for Reactive Gases (WDCRG, <https://www.gaw-wdcr.org>). These data have also been used in many previous investigations such as that of Kumar et al. (2021). The time span of the provided surface ozone data ~~is between~~ for the Halley station is from the year 2007 ~~and 2013.~~ to 2019, and we adopted the data belonging to the
years 2007-2013 in the present study. The time resolution of the surface data is 1 hour. Because the tropospheric ODE, which

we focused on in the present study, mostly occurs in the springtime of every year, we thus only analyzed the surface ozone data during the springtime of the Halley station (from Sept. 1 to Nov. 30).

We then ~~calculated the occurrence frequency of the tropospheric ODE~~ picked out time points representing the occurrence of tropospheric ODEs at the Halley station for each month, based on the daily ~~data of the~~ surface ozone. However, at present, the definition of the ODE occurrence is still under debate. In some previous studies (Tarasick and Bottenheim, 2002; Koo et al., 2012), the occurrence of ODEs was recognized by the surface ozone mixing ratio. When the surface ozone drops to lower than 20 ppb, it is called partial ODEs. Moreover, when the surface ozone declines to a level lower than 10 ppb, it is called major or severe ODEs. In contrast, the ODEs can also be judged by the ~~change in variation of~~ the ozone mixing ratio. This method was suggested by Bian et al. (2018) to indicate ~~extreme events uncommon variations of the surface ozone~~ in polar regions. In the present study, we ~~used the method suggested by Bian et al. (2018) and picked up the time point~~ picked out the time points representing the occurrence of ODEs from the observational data ~~when it based on the method of Bian et al. (2018), when the~~ instantaneous ozone at these time points fulfills the following criterion,

$$[O_3]_i - \overline{[O_3]} < -\alpha \cdot \sigma, \quad (1)$$

where $[O_3]_i$ is the instantaneous ozone ~~data~~ at the i -th time point, and $\overline{[O_3]}$ is the mean ozone value ~~of this over a~~ month. σ in Eq. (1) is the standard deviation. ~~After picking up the time points representing the tropospheric ODEs from the observational data, we calculated the frequency of ODEs belonging to each month during the springtime of the years 2007–2013. The constant~~ α in Eq. (1) is set to 1.5 in the present study so that many partial ODEs during the springtime can also be identified. By using Eq. (1), we defined the occurrence of ODEs as the period when the surface ozone drops remarkably instead of the time when ozone is lower than a level. The identified ODEs using this selection criterion are shown in Fig. S1 in the supplementary material, and from the results we feel the method identifying ODEs used in the present study acceptable.

2.2 Numerical Methods

Two numerical models, a radiation model (TUV, Tropospheric Ultraviolet and Visible) (Madronich and Flocke, 1997, 1999) and a chemical box model (KINAL, KInetic aNALysis of reaction mechanics) (Turányi, 1990), were used in this study. The TUV model was used to estimate the photolytic rates of atmospheric constituents in the troposphere of the Antarctic. The KINAL box model was used to capture the ~~concentration change in variations of~~ these constituents such as ozone in the boundary layer over time. KINAL was also used to compute the sensitivity of these constituents to each photolysis reaction in the chemical mechanism.

2.2.1 TUV model

The TUV model (Tropospheric Ultraviolet and Visible Radiation Model), provided by NCAR (National Center for Atmospheric Research), is able to calculate the tropospheric photo-dissociation coefficient (Madronich and Flocke, 1997, 1999), based on input parameters such as the total ozone column (TOC). The vertical ozone profile assumed in the model is taken from the

180 US standard atmosphere [1976 \(Krueger and Minzner, 1976\)](#). A total of 112 photolysis reactions are implemented in the TUV model.

The photolytic rate constant j_p (unit: s^{-1}) for each photolysis reaction is calculated by the TUV model as follows,

$$j_p = \int_0^{\infty} \sigma(\lambda) \Phi(\lambda) F(\lambda) d\lambda. \quad (2)$$

In Eq. (2), $\sigma(\lambda)$ represents the absorption cross section at the wavelength λ . $\Phi(\lambda)$ denotes the photolytic quantum yield. F in
185 Eq. (2) is the actinic flux, and it is determined by many factors such as the presence of clouds and ~~the change in TOC~~ [TOC](#). [The input parameters of the TUV model are listed in Tab. S2 of the supplementary material, among which TOC and the date vary on different days.](#) A 4-stream discrete ordinate method (van Oss and Spurr, 2002) with a step length of 1 nm is implemented in TUV, calculating the photolytic rate constants.

Thus, in the present study, we implemented the observed TOC and other weather conditions into the TUV model, to estimate
190 the actinic flux F reaching the boundary layer and the rates of photolysis reactions for different time periods. Then we used a chemical box model, KINAL, to capture the ~~concentration change in~~ [temporal evolution of](#) ozone and bromine species in the process of the tropospheric ODE under different photolytic conditions. By doing that, the influence of the ~~change in the~~ total amount of ozone, i.e., [TOC](#), on the occurrence of the tropospheric ODE can be revealed.

2.2.2 KINAL model

195 After obtaining the photolytic dissociation rates of many atmospheric constituents using the TUV model, we then applied the chemical box model, KINAL (KINetic aNALysis of reaction mechanics) (Turányi, 1990) to capture the temporal evolution of chemical species such as ozone and many bromine species. Moreover, sensitivities of these chemical species to each photolysis reaction in the chemical mechanism were also computed using KINAL.

The governing equation describing the temporal evolution of chemical species in the KINAL model is as follows (Turányi,
200 1990):

$$\frac{dc}{dt} = f(c, k) + E, \quad (3)$$

with the initial condition $c|_{t=0} = c_0$, where c represents a vector of chemical species concentrations. k in Eq. (3) is a vector of rate constants of chemical reactions and t denotes time. E indicates the near-surface source emissions, and $\frac{dc}{dt}$ is the temporal evolution of chemical species such as ozone. In the present study, a chemical mechanism including the bromine
205 and chlorine chemistry was adopted from previous box model studies (Cao et al., 2014, 2016a,c; Zhou et al., 2020), and the reaction rate constants were updated with the latest chemical kinetic data (Atkinson et al., 2006). There are in total 49 species and 141 chemical reactions included in the latest version of the chemical mechanism, which are listed in Tab. ~~S2~~ [S3](#) of the supplementary material. Among these reactions, there are 23 photolysis reactions of which the rates are associated with ~~the change in~~ [TOC](#), and these photolysis reactions are listed in Tab. 1 along with the reaction numbers in the mechanism. Among
210 these photolysis reactions, a part of them can enhance the occurrence of ODEs, while the others will retard it.

Table 1. Listing of photolysis reactions in the chemical mechanism of the KINAL model, of which the rates vary with TOC. The reaction numbers correspond to those listed in Tab. [S2-S3](#) of the supplements.

Reaction Number	Reaction
(SR1)	$\text{O}_3 + h\nu \rightarrow \text{O}(^1\text{D}) + \text{O}_2$
(SR6)	$\text{Br}_2 + h\nu \rightarrow 2\text{Br}$
(SR7)	$\text{BrO} + h\nu \xrightarrow{\text{O}_2} \text{Br} + \text{O}_3$
(SR11)	$\text{HOBr} + h\nu \rightarrow \text{Br} + \text{OH}$
(SR57)	$\text{H}_2\text{O}_2 + h\nu \rightarrow 2\text{OH}$
(SR58)	$\text{HCHO} + h\nu \xrightarrow{2\text{O}_2} 2\text{HO}_2 + \text{CO}$
(SR59)	$\text{HCHO} + h\nu \rightarrow \text{H}_2 + \text{CO}$
(SR60)	$\text{C}_2\text{H}_4\text{O} + h\nu \rightarrow \text{CH}_3\text{O}_2 + \text{CO} + \text{HO}_2$
(SR61)	$\text{CH}_3\text{O}_2\text{H} + h\nu \rightarrow \text{OH} + \text{HCHO} + \text{HO}_2$
(SR62)	$\text{C}_2\text{H}_5\text{O}_2\text{H} + h\nu \rightarrow \text{C}_2\text{H}_5\text{O} + \text{OH}$
(SR74)	$\text{HNO}_3 + h\nu \rightarrow \text{NO}_2 + \text{OH}$
(SR75)	$\text{NO}_2 + h\nu \xrightarrow{\text{O}_2} \text{NO} + \text{O}_3$
(SR76)	$\text{NO}_3 + h\nu \xrightarrow{\text{O}_2} \text{NO}_2 + \text{O}_3$
(SR77)	$\text{NO}_3 + h\nu \rightarrow \text{NO} + \text{O}_2$
(SR88)	$\text{BrONO}_2 + h\nu \rightarrow \text{NO}_2 + \text{BrO}$
(SR89)	$\text{BrNO}_2 + h\nu \rightarrow \text{NO}_2 + \text{Br}$
(SR91)	$\text{PAN} + h\nu \rightarrow \text{NO}_2 + \text{CH}_3\text{CO}_3$
(SR135)	$\text{BrCl} + h\nu \rightarrow \text{Br} + \text{Cl}$
(SR136)	$\text{Cl}_2 + h\nu \rightarrow 2\text{Cl}$
(SR137)	$\text{ClO} + h\nu \rightarrow \text{Cl} + \text{O}_3$
(SR138)	$\text{HOCl} + h\nu \rightarrow \text{Cl} + \text{OH}$
(SR139)	$\text{ClONO}_2 + h\nu \rightarrow \text{Cl} + \text{NO}_3$
(SR140)	$\text{OCIO} + h\nu \rightarrow \text{ClO} + \text{O}_3$

In the KINAL model, it is also assumed that bromide stored in the ice/snow-covered ground surface is inexhaustible. As a result, the rates of heterogeneous reactions such as $\text{HOBr}_{(\text{gas})} + \text{H}^+_{(\text{liquid})} + \text{Br}^-_{(\text{liquid})} \rightarrow \text{Br}_{2(\text{gas})} + \text{H}_2\text{O}_{(\text{liquid})}$, which are responsible for the bromine explosion, only depend on the availability of HOBr in the atmosphere in model calculations.

~~Moreover, the wind speed~~ The rates of these heterogeneous reactions, v , are estimated as:

$$v = -\frac{d[\text{HOBr}]}{dt} = \frac{d[\text{Br}_2]}{dt} = \frac{v_d}{L} [\text{HOBr}]. \quad (4)$$

Table 2. Initial atmospheric composition in the boundary layer of the Antarctic (ppm = parts per million, ppb = parts per billion, ppt = parts per trillion) (Piot, 2007), and the prescribed intensities of emission fluxes from the ground surface (units: molec. cm⁻² s⁻¹) (Hutterli et al., 2004; Riedel et al., 2005; Jones et al., 2011), assumed in the model.

Species	Mixing Ratio	Emissions	Species	Mixing Ratio	Emissions
O ₃	25 ppb	-	C ₂ H ₆	0.4 ppb	-
Br ₂	0.3 ppt	-	C ₂ H ₄	50 ppt	-
HBr	0.01 ppt	-	C ₂ H ₂	300 ppt	-
CH ₄	1.7 ppm	-	C ₃ H ₈	0.2 ppb	-
CO ₂	371 ppm	-	NO	2 ppt	1.6 × 10 ⁷
CO	50 ppb	-	NO ₂	8 ppt	1.6 × 10 ⁷
HCHO	500 ppt	9.0 × 10 ⁹	HONO	-	1.6 × 10 ⁷
CH ₃ CHO	500 ppt	-	H ₂ O ₂	-	1.0 × 10 ⁹
H ₂ O	800 ppm	-			

In Eq. (4), [HOBr] and [Br₂] are the concentrations of HOBr and Br₂ in the boundary layer, respectively. *L* is the boundary layer height, and *v_d* is the dry deposition velocity of HOBr at the ice/snow surface. The estimation of *v_d* depends on the values of three resistances (i.e., *r_a*, *r_b* and *r_c*), which are associated with the wind speed and the roughness of the ground surface (Seinfeld and Pandis, 2006). The wind speed was assumed as 8 m s⁻¹ (Beare et al., 2006), which is a typical wind speed in polar regions. In addition, the roughness of the ice/snow surface is set to 10⁻⁵ m (Stull, 1988), and the height of the polar boundary layer is assumed as 200 m because the typical thickness of the boundary layer in polar regions is about 100-500 m (Simpson et al., 2007; Anderson and Neff, 2008). Details of the parameterization of the heterogeneous reactions can be found in previous publications (Lehrer et al., 2004; Cao et al., 2014, 2016b). Aside from this parameterization, we also assumed that the loss of chemical species caused by dry deposition is equivalent to the flux brought by the entrainment from the free atmosphere into the boundary layer, which is similar to the treatment of Michalowski et al. (2000). By doing that, in the absence of chemistry, the concentrations of chemical species are able to maintain in the balance of the dry deposition and the entrainment. The initial atmospheric composition used in the model is listed in Tab. 2, which represents a typical air composition in the Antarctic (Piot, 2007). Constant emission fluxes from the ground surface were also prescribed in the model according to observations (Hutterli et al., 2004; Riedel et al., 2005; Jones et al., 2011), and the prescribed intensities of the emission fluxes are also presented in Tab. 2.

2.2.3 Concentration Sensitivity Analysis

After obtaining the temporal evolution of ozone and major bromine species, relative concentration sensitivities of these species to different photolysis reactions in the chemical mechanism were computed to reveal the dependence of these species on each

photolysis reaction of the mechanism. The relative concentration sensitivity S_{ij} is calculated by:

$$S_{ij} = \frac{\partial \ln c_i}{\partial \ln k_j} = \frac{k_j}{c_i} \frac{\partial c_i}{\partial k_j}, \quad (5)$$

which shows the importance of the j -th reaction for the concentration change in the i -th chemical species. In Eq. (5), i is the index of chemical species, and j is the index of chemical reactions in the mechanism. c_i is the concentration of the i -th species, and k_j is the rate constant of the j -th reaction. S_{ij} , an element of the relative sensitivity matrix, indicates the change in the i -th species concentration resulted from a small perturbation in the j -th reaction rate. The evaluation of the concentration sensitivity is helpful for discovering the importance of specific reactions in the chemical mechanism for the concentration change in of the focused species.

In the following section, the computational results are presented and discussed.

3 Results and Discussions

In this section, we first show the ~~relationship between the TOC change and the occurrence frequency of the tropospheric~~
~~ODE connection between TOCs detected at two monitoring stations (i.e., Halley station and FAD station) and the tropospheric~~
~~ODE observed~~ at the Halley station ~~based on the observational data~~. Later, we ~~presented~~ present the computational results of ODEs for ~~the year 2013~~ a time period of Sept. 29 - Oct 8, 2008 as an example to show the time ~~variations~~ series of ozone and major bromine compounds during ODEs. The depletion rate of ozone and the temporal ~~change in~~ evolution of bromine species under different ~~weather~~ conditions were then displayed to indicate the influence caused by ~~the TOC change~~ TOC on the ozone depletion and the bromine activation. At last, a concentration sensitivity analysis was performed to see which photolysis reactions playing important roles in the connection between ~~the TOC change~~ TOC and the occurrence of ODEs.

3.1 Relationship between the ~~TOC Change~~ TOCs and the Tropospheric ODE

The time series of the ~~monthly-averaged TOC~~ daily TOCs detected at the Halley station and the ~~occurrence frequency of~~
~~ODEs~~ Faraday-Vernadsky (FAD) station as well as the surface ozone of Halley during the springtime of years 2007-2013 are
 presented in Fig. 1. From the temporal evolution of the surface ozone, we found that at Halley, the occurrence frequency of
ODEs in November is substantially lower than that in September or October. Moreover, by comparing the surface ozone of
Halley with the TOC detected at Halley, we did not find any obvious correlation between them, except that the ODEs occur
more frequently in a relatively low TOC condition. However, from the comparison between the surface ozone of Halley and
the TOC detected at the FAD station, we found that the ODEs observed at Halley usually followed a decline in TOC at the FAD
station (see the marks in Fig. 1). It suggests that the decrease of TOC surrounding FAD possibly favors the occurrence of ODEs
at the Halley station. As the FAD station is located to the northwest of the Halley station and near the Weddell Sea, the TOC
detected at this station is more capable of reflecting conditions of the Weddell Sea. Thus, we suggest the possible mechanism
as that the decline in TOC over the area of the Weddell Sea favors the tropospheric ozone depletion in this region. Then the
ozone-lacking air was transported from the sea to the Halley station, leading to the detection of ODEs at this site. Thus, there

265 exists a lag time between the TOC decline observed at the FAD station and the detection of ODE at the Halley station, and the
length of the lag time depends on the weather conditions during that period. In previous studies, the source of ODEs observed
at Halley has also been discussed by Jones et al. (2006), who found that air masses causing rapid ODEs of Halley originated in
the southern Weddell Sea. Our findings are consistent with the conclusions of Jones et al. (2006).

In order to further clarify the ~~correlation between the change in TOC and~~ role of TOC in affecting the occurrence of ODEs,
270 we then took the year ~~2013-2008~~ as an example and used the models (i.e., TUV and KINAL) with the input ~~data of TOC~~
~~belonging to the year 2013 (220.03 DU on average)~~ of TOC observed at the FAD station. The reason we chose the year ~~2013 is~~
~~that it is the latest year with abundant springtime surface ozone data.~~ 2008 is that the TOC variation at the FAD station in this
year is more stable than those in other years (see Fig. S2 in the supplements).

3.2 Temporal Behavior of Ozone and Bromine Species during ODEs ~~for the Year 2013~~ in October, 2008

275 The temporal ~~change in profiles of~~ ozone and bromine species ~~under the conditions of the year 2013 with an average TOC~~
~~of 220.03 DU is simulated by models implementing the daily variation of TOC observed at the FAD station from Sept. 29~~
~~to Oct. 8, 2008 are~~ shown in Fig. 2. We chose this period because during this time a significant drop in TOC from 250 DU
to 131 DU was observed at the FAD station (see Fig. S2 in the supplements). From the temporal behavior of these chemical
species, we can better understand the interconversion of bromine species and the reasons causing the depletion of ozone in
280 the troposphere. Because similar results have been shown and discussed in previous publications (Cao et al., 2014, 2016a), we
only describe it briefly here.

It can be seen in Fig. 2 that the whole process of the ODE can be divided into four time stages, according to the types
of bromine species in the troposphere. The first stage is the beginning of ODEs, in which the concentration of bromine is
low, and ozone is only slightly depleted. This time stage would last for about ~~65~~ 65 days in the present simulation (see Fig. 2).
285 Subsequently, in the second time stage, Br₂ released from substrates such as the ice/snow-covered surface due to the bromine
explosion mechanism is continuously photolyzed, forming Br atoms. The Br atoms then react with ozone and form BrO:



A part of BrO then gets oxidized and converted to HOBr:



290 Thus, in the second time stage, the major bromine species are BrO and HOBr. In contrast, the concentration of Br is almost
zero due to the presence of ozone. Under this high-HOBr environment, a large amount of bromide is activated from substrates,
so that ozone is rapidly consumed by the massive amount of bromine in the atmosphere. Thus, this second time stage is also
the key time period that the majority of the tropospheric ozone is depleted, and this time stage lasts for about 1 day, in which
the majority of ozone is depleted with a rate of 0.5-1.0 ppb hr⁻¹.

295 When the mixing ratio of ozone drops to less than 5 ppb, this event enters the third stage. In this time stage, the depletion
of ozone continues, but the formations of BrO and HOBr are retarded, because of the low ozone in the atmosphere. The major



Figure 1. (Continued...)




Figure 1. Time series of TOCs belonging to the Halley station and the Faraday-Vernadsky (FAD) station as well as the surface ozone detected at the Halley station during the springtime of years 2007-2013 (the observational data of the surface ozone for October and November in the year 2012 are missing). The green-shaded areas in the figure indicate the periods identified as the occurrence of ODEs at Halley in the present study, and the red-shaded areas represent the significant decline in TOC at FAD, which might be associated with the occurrence of ODEs at Halley.

bromine species at this time is Br, formed by the photo-decomposition of BrO and HOBr in the atmosphere. Then the last stage comes, in which ozone in the troposphere is almost completely consumed. At this time stage, BrO and HOBr are all photolyzed to Br, thus not existing in the atmosphere. Meanwhile, the formed Br is eliminated by aldehydes and HO_x free radicals in the atmosphere, and is converted into HBr. As a result, at this last stage, a complete depletion of ozone in the troposphere is




Figure 2. Variations-Temporal profiles of ozone and major bromine species with time during the tropospheric ODE under the conditions of the year-2013, from Sept. 9 to Oct. 8, 2008.

achieved, and the major bromine species is HBr, which is in accordance with previous observations (Langendörfer et al., 1999).

3.3 Impact of the Change in TOC on the Occurrence of ODEs

We then compared the model results for different months of the springtime of 2008. The adopted time periods representing
 305 the three months for the present investigation are Sept. 1-Sept. 10, Sept. 29-Oct. 8 and Nov. 1-Nov. 10, respectively. The TOC
variations in these three months can be found in Fig. S2 of the supplementary material. However, it should be noted that aside
 from the change-difference in TOC, the value of SZA also varies between different-these months, which may significantly
 affect the radiation fluxes reaching the boundary layer and thus the rates of photolysis reactions.

The temporal change-in-profiles of ozone during ODEs under the conditions of different months belonging to the year-2013
 310 is-of 2008 are shown in Fig. 3. It is seen that compared with the situation in September, the decline of ozone in November is
 delayed. Moreover, the depletion rate in November was also found to be lower than that in September. It suggests that under
 the weather conditions of November 2013-2008, the occurrence of ODEs is more difficult to achieve, compared with that in
 September 2013-2008. This result is also consistent with the change-in-the-relatively lower occurrence frequency of ODEs
observed in November shown above. However, as mentioned above, although TOC increases from September to November in
 315 2013differs between these months, it does not guarantee that the occurrence of ODEs is negatively dependent on the variation




Figure 3. Variations-Temporal profiles of ozone with time during ODEs for TOC in different months of the springtime of 2013, 2008 and the simulated surface ozone during ODEs.

of TOC difference in the ODE occurrence between these simulations is caused by the use of different TOCs, because SZA also decreases varies, which may heavily enhance-influence the radiation fluxes reaching the ground surface -and thus the occurrence frequency of ODEs. Therefore, we designed two sensitivity tests in the present study, to discover the role of SZA and TOC in affecting the ODEs separately.

320 In order to clarify the role of SZA in determining the occurrence of ODEs, we designed a sensitivity test by replacing the input TOC variation in September and November simulations with that belonging to October. By performing this test, we were able to discover the influence on the occurrence of ODEs solely by SZA. Figure 4 shows the temporal evolution of the surface ozone in this sensitivity test. It can be seen that when applying a same TOC variation in these simulations, the ozone depletion in September occurs remarkably earlier than that in November. It denotes that the decline of SZA leads to a retardation of ODEs. Thus, the ozone depletion is more difficult to achieve when SZA becomes smaller. The reasons for this
 325 positive association between SZA and the occurrence frequency of ODEs would be discussed in a later context.

After clarifying the role of SZA, we continued to discover the role of TOC in affecting ODEs. We thus implemented different temporal profiles of TOC in the October simulation. As mentioned above, in October, TOC at the FAD station drops sharply from 250 DU in Sept. 29 to 131 DU in Oct. 4. In this sensitivity test, we first assumed that the TOC keeps as a constant 250 DU
 330 instead of dropping, and we named this simulation scenario as the “constant” scenario. Next, we performed another simulation in which the TOC increases sharply from 250 DU to 350 DU from the beginning of the simulation rather than decreasing, and




Figure 4. Temporal evolution of ozone during ODEs in different months of the springtime of 2008, using a same TOC temporal profile.

this scenario was named as “increase” in the following context. Thus, in these simulation scenarios, the TOC value for the year 2008 level in the original scenario is the lowest, while the TOC value for 2013 level in the “increase” scenario is the highest. In contrast, the values of SZA and other meteorological parameters are mostly similar in these scenarios. By doing that, we were able to figure find out the impact on the occurrence of ODEs brought about only by the change in exerted by only TOC using TUV and KINAL models.

The time variations temporal profiles of ozone and major bromine species are presented in Fig. 5. It can be seen in Fig. 5(a) that when TOC in October is lower (i.e., the situation in 2008 the original simulation), the depletion of ozone is accelerated and the depletion rate becomes higher than those for in the other two years scenarios. It denotes that it takes less time for the ozone to be completely depleted in 2008 than in the other two years due to the lower TOC in the October of 2008 under a lower TOC. Moreover, it can be found from Fig. 5(b)-(d) that the peaks of BrO, HOBr and HBr in October 2008 the original simulation scenario occur earlier than those in October 2013 the other two scenarios. In addition, it can be seen from Fig. 5(d) that the total amount of bromine in the atmosphere (i.e., in the form of HBr) at the end of ODEs for 2008 in the original scenario is slightly higher than that for 2013 those in the other two scenarios. Thus, from the model results, the decrease of TOC favors enhances the occurrence of the tropospheric ODEs and the bromine release. This simulation result also partly supports our previous suggestion about the negative dependence of the ODE occurrence frequency on the TOC change that the decline in TOC possibly favors the tropospheric ozone depletion in the analysis of the observational data.




Figure 5. Variations-Time series of (a) ozone, (b) BrO, (c) HOBr and (d) HBr with time during ODEs under the conditions of October for 2008, implementing three different years-TOC profiles (i.e., original, “constant” and “increase”).

The mechanism we proposed is that when TOC decreases, a larger amount of solar radiation would reach the troposphere, leading to an acceleration of a part of photo-chemical reactions associated with the ozone depletion and the bromine activation in the troposphere. As a result, the formation of major bromine species such as BrO and HOBr as well as the bromine activation become faster, and the occurrence of ODEs is also accelerated.

However, it is still unclear through which photolysis reactions the change-in-variation of TOC deeply affects the occurrence of tropospheric ODEs. Thus, we continued to analyze the photo-chemical reactions using the concentration sensitivity analysis, as presented below.




Figure 6. Relative sensitivities of (a) ozone, (b) BrO, (c) HOBr and (d) HBr to photolysis reactions on Day 6:55.8, which resides in the time period when the strongest ozone depletion occurs.

355 3.4 Sensitivities of Ozone and Major Bromine Species to Photolysis Reactions

In the present study, the impact on ODEs caused by the change in TOC in the models is exerted through 23 photolysis reactions, which are listed in Tab. 1. In order to figure out discover which photolysis reactions are the most important ones during the ODE process, we performed a concentration sensitivity analysis on the change in ozone and major bromine species for the

~~springtime of 2013, simulation of October 2008 (i.e., the simulation presented in Sect. 3.2),~~ so that the dependence of ozone

360 and bromine species on these photolysis reactions can be revealed.

The relative concentration sensitivities of ozone and major bromine species (i.e., BrO, HOBr and HBr) to all the 23 photolysis reactions on Day 6.55.8, which resides in the second time stage of ODEs when the strongest ozone depletion occurs (see Fig. 2), are shown in Fig. 6. From Fig. 6(a), it can be seen that Reactions (SR7) and (SR11):




365



possess the largest positive sensitivities for the mixing ratio of ozone. It means that the rate increase of these two photolysis reactions leads to an elevation of the ozone value during ODEs and thus a retardation of ODEs. These two reactions are thus named “major ODE decelerating reactions” in the following context. The reason for the delay impact on ODEs brought by
370 Reaction (SR7) is that in this reaction, BrO is photolyzed, forming Br. As a result, the formation of HOBr by the oxidation of BrO is decelerated due to the reduction of the available BrO. Thus, the heterogeneous bromine activation process, i.e., bromine explosion mechanism that HOBr participates in, gets retarded, leading to a slow down of the bromine activation and the ozone depletion. Apart from that, additional ozone is also formed through Reaction (SR7). With respect to Reaction (SR11), HOBr is photo-decomposed through this reaction. Thus, the heterogeneous bromine activation also gets suppressed by the strengthening
375 of this reaction, resulting in a delay of the ozone depletion.

In contrast, Reactions (SR1), (SR57), and (SR58),



380



are largely negatively correlated with the change in ozone (see Fig. 6a). It denotes that when these three reactions speedup, the level of ozone drops, which represents an acceleration of the tropospheric ODE. These three reactions are thus named “major ODE accelerating reactions”. For Reaction (SR1), it is not surprising as this reaction is the direct photolysis of the tropospheric
385 ozone. Moreover, Reaction (SR1) also serves as a major formation pathway of OH radicals. OH radicals are important for the occurrence of ODEs and the bromine explosion mechanism, as they are involved in Reaction (SR18):



In Reaction (SR18), not only the Br atoms are generated by the conversion from the released Br₂, but also the formation of HOBr is strengthened, leading to an acceleration of the bromine activation. As a result, an enhancement of Reaction (SR1) would lead to a speedup of the ozone depletion. Regarding Reaction (SR57), the occurrence of this reaction also strengthens the occurrence of ODEs, because this reaction also forms OH radicals, which are critical for the bromine explosion mechanism as mentioned above. With respect to Reaction (SR58), it possesses the most negative ozone sensitivity (see Fig. 6a), which means that this reaction heavily controls the depletion of ozone. It is because this reaction reinforces the formation of HO₂. As HO₂ is the key oxidant for the formation of HOBr through reaction:



the enhancement of Reaction (SR58) thus favors the release of bromine and the depletion of ozone.

From Fig. 6(b)-(d), we can see that the sensitivities of major bromine species (i.e., BrO, HOBr and HBr) to photolysis reactions mostly have an opposite sign, compared with those corresponding to ozone (shown in Fig. 6a). It is because the bromine species in the atmosphere are mostly responsible for the ozone depletion in the troposphere during ODEs. Within these photolysis reactions, Reactions (SR7) and (SR11) have the largest negative sensitivities as they strongly decelerate the bromine explosion mechanism. In contrast, Reactions (SR1), (SR57) and (SR58) exert a positive impact on the change in bromine species. It is because the speedup of these three reactions can reinforce the bromine explosion mechanism as mentioned above, thus leading to a positive dependence of these bromine containing compounds on these three reactions.

The results of the sensitivity analysis help to explain the impact on the occurrence of ODEs exerted by ~~the change in TOC~~ in SZA in previous discussions. First, in the comparison of the ~~simulation~~ results corresponding to different months of ~~the springtime of 2013 (see 2008 (see Fig. 3 and Fig. 4 in Sect. 3.3),~~ it was found that the tropospheric ODE is more difficult to achieve in November than in September ~~of the same year due to a smaller SZA in November.~~ From the sensitivity analysis, we ~~attributed were able to discover~~ the reason for the retardant of ODEs ~~in November to be caused by~~ the smaller SZA ~~in November than that in September.~~ Due to the shift in SZA, solar radiation with all the wavelengths that reaches the ground surface is strengthened in November. Consequently, although the TOC value in November is higher than that in September, both the major ODE accelerating reactions (i.e., Reactions (SR1), (SR57) and (SR58)) and the major ODE decelerating reactions (i.e. Reactions (SR7) and (SR11)) are promoted in November. From Fig. 6(a), it can be seen that the ozone level during ODEs is more sensitive to the major ODE decelerating reactions than the major ODE accelerating reactions. As a result, the outcome of the SZA decline in November is that the occurrence of ODEs is retarded. In this situation, ~~the change in TOC~~ only plays a minor role in affecting the occurrence of ODEs.

In contrast, in the comparison of the ODE occurrence belonging to October ~~of different years (see using different TOC profiles (see Fig. 5 in Sect. 3.3),~~ because the values of SZA are ~~mostly~~ similar in these scenarios, the change in ODEs is mainly determined by the difference in TOC between these simulations. In a lower TOC environment, the intensity of the solar radiation reaching the atmospheric boundary layer, especially the ultraviolet radiation in a wavelength range of 200-320 nm (i.e., UV-B and UV-C), is significantly enhanced, as ozone has strong absorption bands in 200-320 nm (i.e., Hartley bands). Moreover, solar radiation in 320-350 nm also gets moderately elevated under a low TOC condition, because of the absorption

bands of ozone in 320-350 nm with a vibrational structure (i.e., Huggins bands). On the contrary, solar radiation in other wavelength ranges would not be significantly affected by the decrease of TOC. In this situation, the major ODE accelerating reactions, i.e., photolysis of ozone, H_2O_2 and HCHO in the boundary layer, are remarkably promoted. The reasons are as follows: (1) The photolysis of ozone in the troposphere that forms $\text{O}(^1\text{D})$ depends heavily on the strength of the solar radiation in 295-360 nm (Akimoto, 2016). Thus, the decrease in TOC would significantly accelerate the photolysis of ozone in the troposphere, i.e., Reaction (SR1). (2) The absorption cross section of H_2O_2 decreases monotonically from the wavelength of 190 nm to 350 nm (Vaghjiani and Ravishankara, 1989), which overlaps the wavelength range that the TOC change strongly affects. Thus, the photo-decomposition of H_2O_2 (i.e., Reaction (SR57)) also gets strongly promoted when TOC decreases. (3) The absorption spectrum of HCHO spreads from 260 nm to 360 nm, with many vibrational structures (Rogers, 1990). As a result, the decline of TOC substantially enhances the photolysis of HCHO (i.e., Reaction (SR58)).

On the contrary, for the major ODE decelerating reactions, i.e., photolysis of BrO and HOBr , their rates are only slightly influenced by the decrease of TOC. It is because that BrO has an absorption spectrum in the range of 290-380 nm, peaking at approximately 330 nm (Wilmouth et al., 1999). The strengthening of the solar radiation especially in 200-320 nm due to the decline of TOC thus only exerts a small impact on the photolysis of BrO (i.e., Reaction (SR7)). Regarding HOBr , its photolysis rate relies more on the strength of the UV-A radiation (i.e., in 320-400 nm) reaching the boundary layer, as it has a broad absorption spectrum between 200 and 400 nm (Burkholder et al., 2015). Therefore, the influence caused by ~~the TOC change~~ TOC on the photolysis of HOBr is also weak.

Hence, it can be concluded that when TOC decreases, the rates of major ODE accelerating reactions (i.e., Reactions (SR1), (SR57) and (SR58)) significantly increase, while the rates of major ODE decelerating reactions (i.e., Reactions (SR7) and (SR11)) are hardly changed. Consequently, the ODEs are accelerated under a low TOC condition, and vice versa, resulting in a negative association between ~~the change of~~ TOC and the occurrence frequency of ODEs as presented above.

4 Conclusions and Future Work

In this study, we investigated the connection between the ~~change in the~~ total ozone column (TOC) and the occurrence of the tropospheric ozone depletion events (ODEs) in the Antarctic. Photolysis reactions dominating this connection were also identified using a concentration sensitivity analysis. Major conclusions achieved in the present study are as follows.

Based on the analysis of the observational data belonging to the years 2007-2013, we suggested that the decrease ~~in TOC~~ of TOC surrounding the Faraday-Vernadsky (FAD) station possibly favors the occurrence of the tropospheric ODEs at the Halley station. Then, the model results with the ~~input data representing the conditions of October belonging to different years~~ implementation of different TOC profiles also indicate that the occurrence of ODEs would be accelerated when TOC decreases. Moreover, key photolysis reactions that dominate the production and the consumption of ozone during ODEs, i.e., major ODE accelerating reactions and major ODE decelerating reactions, were also ~~figured out~~ discovered. It was found that when TOC ~~changes~~ varies, the rates of major ODE accelerating reactions are substantially altered, while the rates of major ODE

decelerating reactions mostly remain unchanged, leading to the negative association between ~~the TOC change~~ TOC and the
455 occurrence frequency of ODEs.

Improvements can be made to the present study. For instance, many other factors that are able to influence the occurrence of ODEs such as the type of the sea ice and the existence of frost flowers should be considered in the future work. Unfortunately, currently we are still lack of these observational data. Aside from that, a study for Arctic conditions should also be conducted, so that the conclusions obtained in the present study can be compared and verified, which is attributed to a future publication.

460 *Code and data availability.* The observational data used in this study and the source code of the models as well as the computational results shown in this article can be acquired from the link <https://faculty.nuist.edu.cn/caole/en/kyxm/72647/content/17580.htm#kyxm> (Cao, 2022).

Acknowledgements. The authors wish to thank the financial support by the National Natural Science Foundation of China (Grant No. 41705103). The numerical calculations in this paper have been done on the high performance computing system in the High Performance Computing Center, Nanjing University of Information Science ~~&~~ and Technology.

465 *Author contributions.* Le Cao conceived the idea of the article and extended the KINAL model. Linjie Fan processed the observational data, performed the computations, and wrote the paper with Le Cao together. Simeng Li revised the chemical mechanism and Shuangyan Yang gave valuable suggestions on the improvement of the manuscript. All the authors listed have read and approved the final manuscript.

Competing interests. The authors declare no conflict of interest.

References

- 470 Akimoto, H.: Atmospheric Reaction Chemistry, Springer Atmospheric Sciences, Springer Japan, 2016.
- Anderson, P. S. and Neff, W. D.: Boundary layer physics over snow and ice, *Atmospheric Chemistry and Physics*, 8, 3563–3582, <https://doi.org/10.5194/acp-8-3563-2008>, <https://acp.copernicus.org/articles/8/3563/2008/>, 2008.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., Troe, J., and Subcommittee, I.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume II - gas phase reactions of organic species, *Atmospheric*
475 *Chemistry and Physics*, 6, 3625–4055, <https://doi.org/10.5194/acp-6-3625-2006>, 2006.
- Balis, D., Kroon, M., Koukouli, M. E., Brinksma, E. J., Labow, G., Veefkind, J. P., and McPeters, R. D.: Validation of Ozone Monitoring Instrument total ozone column measurements using Brewer and Dobson spectrophotometer ground-based observations, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/https://doi.org/10.1029/2007JD008796>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2007JD008796>, 2007.
- 480 Beare, R., Macvean, M., Holtslag, A., Cuxart, J., Esau, I., Golaz, J.-C., Jimenez, M., Khairoutdinov, M., Kosovic, B., Lewellen, D., Lund, T., Lundquist, J., McCabe, A., Moene, A., Noh, Y., Raasch, S., and Sullivan, P.: An intercomparison of large-eddy simulations of the stable boundary layer, *Boundary Layer Meteorol.*, 118, 247–272, <https://doi.org/10.1007/s10546-004-2820-6>, <http://dx.doi.org/10.1007/s10546-004-2820-6>, 2006.
- Bedjanian, Y. and Poulet, G.: Kinetics of Halogen Oxide Radicals in the Stratosphere, *Chemical Reviews*, 103, 4639–4656, <https://doi.org/10.1021/cr0205210>, <https://doi.org/10.1021/cr0205210>, PMID: 14664627, 2003.
- 485 Bian, L., Ye, L., Ding, M., Gao, Z., Zheng, X., and Schnell, R.: Surface Ozone Monitoring and Background Concentration at Zhongshan Station, Antarctica, *Atmospheric and Climate Sciences*, 08, 1–14, <https://doi.org/10.4236/acs.2018.81001>, 2018.
- Bottenheim, J. W. and Chan, E.: A trajectory study into the origin of spring time Arctic boundary layer ozone depletion, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/https://doi.org/10.1029/2006JD007055>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2006JD007055>, 2006.
- 490 Bottenheim, J. W., Netcheva, S., Morin, S., and Nghiem, S. V.: Ozone in the boundary layer air over the Arctic Ocean: measurements during the TARA transpolar drift 2006–2008, *Atmos. Chem. Phys.*, 9, 4545–4557, <https://doi.org/10.5194/acp-9-4545-2009>, 2009.
- Boylan, P., Helmig, D., Staebler, R., Turnipseed, A., Fairall, C., and Neff, W.: Boundary layer dynamics during the Ocean-Atmosphere-Sea-Ice-Snow (OASIS) 2009 experiment at Barrow, AK, *Journal of Geophysical Research: Atmospheres*, 119, 2261–2278, <https://doi.org/https://doi.org/10.1002/2013JD020299>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2013JD020299>, 2014.
- 495 Brasseur, G. and Solomon, S.: *Aeronomy of the Middle Atmosphere: Chemistry and Physics of the Stratosphere and Mesosphere*, Springer, 2005.
- Burkholder, J., Sander, S., Abbatt, J., Barker, J., Huie, R., Kolb, C., Kurylo, M., Orkin, V., Wilmouth, D., and Wine, P.: *Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies*, Evaluation Number 18, Tech. rep., <https://doi.org/10.13140/RG.2.1.2504.2806>, 2015.
- 500 Cao, L.: The observational data and the source code of the models as well as the computational results for “Influence of Total Ozone Column (TOC) on the Occurrence of Tropospheric Ozone Depletion Events (ODEs) in the Antarctic”, NUIST Information Platform [code and data set], available at: <https://faculty.nuist.edu.cn/caole/en/kyxm/72647/content/17580.htm#kyxm>, last access: 14 February 2022, 2022.
- Cao, L., Sihler, H., Platt, U., and Gutheil, E.: Numerical analysis of the chemical kinetic mechanisms of ozone depletion and halogen release
505 in the polar troposphere, *Atmos. Chem. Phys.*, 14, 3771–3787, <https://doi.org/10.5194/acp-14-3771-2014>, 2014.

- Cao, L., He, M., Jiang, H., Grosshans, H., and Cao, N.: Sensitivity of the Reaction Mechanism of the Ozone Depletion Events during the Arctic Spring on the Initial Atmospheric Composition of the Troposphere, *Atmosphere*, 7, 124, 2016a.
- Cao, L., Platt, U., and Gutheil, E.: Role of the boundary layer in the occurrence and termination of the tropospheric ozone depletion events in polar spring, *Atmospheric Environment*, 132, 98–110, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2016.02.034>, <https://www.sciencedirect.com/science/article/pii/S1352231016301479>, 2016b.
- 510 Cao, L., Wang, C., Mao, M., Grosshans, H., and Cao, N.: Derivation of the reduced reaction mechanisms of ozone depletion events in the Arctic spring by using concentration sensitivity analysis and principal component analysis, *Atmos. Chem. Phys.*, 16, 14 853–14 873, <https://doi.org/10.5194/acp-16-14853-2016>, 2016c.
- Farman, J. C., Gardiner, B. G., and Shanklin, J. D.: Large losses of total ozone in Antarctica reveal seasonal ClO_x/NO_x interaction, *Nature*, 315, 207–210, <https://doi.org/10.1038/315207a0>, 1985.
- 515 Finlayson-Pitts, B. and Pitts, J.: *Chemistry of the Upper and Lower Atmosphere: Theory, Experiments and Applications*, Academic Press, 1999.
- Frieß U., Hollwedel, J., König-Langlo, G., Wagner, T., and Platt, U.: Dynamics and chemistry of tropospheric bromine explosion events in the Antarctic coastal region, *J. Geophys. Res.*, 109, <https://doi.org/10.1029/2003JD004133>, 2004.
- 520 Heinemann, G.: The polar regions: a natural laboratory for boundary layer meteorology a review, *Meteorologische Zeitschrift*, 17, 589–601, <https://doi.org/10.1127/0941-2948/2008/0327>, <http://dx.doi.org/10.1127/0941-2948/2008/0327>, 2008.
- Hopper, J. F., Barrie, L. A., Silis, A., Hart, W., Gallant, A. J., and Dryfhout, H.: Ozone and meteorology during the 1994 Polar Sunrise Experiment, *J. Geophys. Res.*, 103, 1481–1492, <https://doi.org/10.1029/97JD02888>, 1998.
- Hu, X.-M., Zhang, F., Yu, G., Fuentes, J. D., and Wu, L.: Contribution of mixed-phase boundary layer clouds to the termination of ozone depletion events in the Arctic, *Geophys. Res. Lett.*, 38, L21 801, <https://doi.org/10.1029/2011GL049229>, 2011.
- 525 Hutterli, M. A., McConnell, J. R., Chen, G., Bales, R. C., Davis, D. D., and Lenschow, D. H.: Formaldehyde and hydrogen peroxide in air, snow and interstitial air at South Pole, *Atmospheric Environment*, 38, 5439–5450, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2004.06.003>, <https://www.sciencedirect.com/science/article/pii/S1352231004005175>, antarctic Atmospheric Chemistry: ISCAT 2000, 2004.
- 530 Jacobi, H.-W., Morin, S., and Bottenheim, J. W.: Observation of widespread depletion of ozone in the springtime boundary layer of the central Arctic linked to mesoscale synoptic conditions, *J. Geophys. Res.*, 115, <https://doi.org/10.1029/2010JD013940>, 2010.
- Jones, A. E., Anderson, P. S., Wolff, E. W., Turner, J., Rankin, A. M., and Colwell, S. R.: A role for newly forming sea ice in springtime polar tropospheric ozone loss? Observational evidence from Halley station, Antarctica, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/https://doi.org/10.1029/2005JD006566>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2005JD006566>, 2006.
- 535 Jones, A. E., Wolff, E. W., Ames, D., Bauguutte, S. J.-B., Clemitchaw, K. C., Fleming, Z., Mills, G. P., Saiz-Lopez, A., Salmon, R. A., Sturges, W. T., and Worton, D. R.: The multi-seasonal NO_y budget in coastal Antarctica and its link with surface snow and ice core nitrate: results from the CHABLIS campaign, *Atmospheric Chemistry and Physics*, 11, 9271–9285, <https://doi.org/10.5194/acp-11-9271-2011>, <https://acp.copernicus.org/articles/11/9271/2011/>, 2011.
- Koo, J.-H., Wang, Y., Kurosu, T. P., Chance, K., Rozanov, A., Richter, A., Oltmans, S. J., Thompson, A. M., Hair, J. W., Fenn, M. A., Weinheimer, A. J., Ryerson, T. B., Solberg, S., Huey, L. G., Liao, J., Dibb, J. E., Neuman, J. A., Nowak, J. B., Pierce, R. B., Natarajan, M., and Al-Saadi, J.: Characteristics of tropospheric ozone depletion events in the Arctic spring: analysis of the ARCTAS, ARCPAC, and ARCIONS measurements and satellite BrO observations, *Atmos. Chem. Phys.*, 12, 9909–9922, <https://doi.org/10.5194/acp-12-9909-2012>, 2012.
- 540

- Koo, J.-H., Wang, Y., Jiang, T., Deng, Y., Oltmans, S. J., and Solberg, S.: Influence of climate variability on near-surface ozone depletion events in the Arctic spring, *Geophysical Research Letters*, 41, 2582–2589, <https://doi.org/https://doi.org/10.1002/2014GL059275>, 2014.
- Kreher, K., Johnston, P. V., Wood, S. W., Nardi, B., and Platt, U.: Ground-based measurements of tropospheric and stratospheric BrO at Arrival Heights, Antarctica, *Geophys. Res. Lett.*, 24, 3021–3024, <https://doi.org/10.1029/97GL02997>, 1997.
- Krueger, A. J. and Minzner, R. A.: A mid-latitude ozone model for the 1976 U.S. Standard Atmosphere, *Journal of Geophysical Research* (1896-1977), 81, 4477–4481, <https://doi.org/https://doi.org/10.1029/JC081i024p04477>, 1976.
- Kuang, S., Newchurch, M. J., Johnson, M. S., Wang, L., Burris, J., Pierce, R. B., Eloranta, E. W., Pollack, I. B., Graus, M., de Gouw, J., Warneke, C., Ryerson, T. B., Markovic, M. Z., Holloway, J. S., Pour-Biazar, A., Huang, G., Liu, X., and Feng, N.: Summertime tropospheric ozone enhancement associated with a cold front passage due to stratosphere-to-troposphere transport and biomass burning: Simultaneous ground-based lidar and airborne measurements, *Journal of Geophysical Research: Atmospheres*, 122, 1293–1311, <https://doi.org/https://doi.org/10.1002/2016JD026078>, 2017.
- Kumar, P., Kuttippurath, J., Gathen, P., Petropavlovskikh, I., Johnson, B., McClure-Begley, A., Cristofanelli, P., Bonasoni, P., Barlasina, M., and Sánchez, R.: The Increasing Surface Ozone and Tropospheric Ozone in Antarctica and Their Possible Drivers, *Environmental Science & Technology*, 55, <https://doi.org/10.1021/acs.est.0c08491>, 2021.
- Kuttippurath, J., Goutail, F., Pommereau, J.-P., Lefèvre, F., Roscoe, H. K., Pazmiño, A., Feng, W., Chipperfield, M. P., and Godin-Beekmann, S.: Estimation of Antarctic ozone loss from ground-based total column measurements, *Atmospheric Chemistry and Physics*, 10, 6569–6581, <https://doi.org/10.5194/acp-10-6569-2010>, <https://acp.copernicus.org/articles/10/6569/2010/>, 2010.
- Langendörfer, U., Lehrer, E., Wagenbach, D., and Platt, U.: Observation of filterable bromine variabilities during Arctic tropospheric ozone depletion events in high (1 hour) time resolution, *J. Atmos. Chem.*, 34, 39–54, <https://doi.org/10.1023/A:1006217001008>, 1999.
- Lehrer, E., Hönninger, G., and Platt, U.: A one dimensional model study of the mechanism of halogen liberation and vertical transport in the polar troposphere, *Atmos. Chem. Phys.*, 4, 2427–2440, <https://doi.org/10.5194/acp-4-2427-2004>, 2004.
- Lippmann, M.: Health effects of tropospheric ozone, *Environ. Sci. Technol.*, 25, 1954–1962, <https://doi.org/10.1021/es00024a001>, 1991.
- Madronich, S. and Flocke, S.: Theoretical Estimation of Biologically Effective UV Radiation at the Earth’s Surface, in: *Solar Ultraviolet Radiation*, edited by Zerefos, C. S. and Bais, A. F., pp. 23–48, Springer Berlin Heidelberg, Berlin, Heidelberg, 1997.
- Madronich, S. and Flocke, S.: The Role of Solar Radiation in Atmospheric Chemistry, pp. 1–26, Springer Berlin Heidelberg, Berlin, Heidelberg, https://doi.org/10.1007/978-3-540-69044-3_1, 1999.
- Michalowski, B. A., Francisco, J. S., Li, S.-M., Barrie, L. A., Bottenheim, J. W., and Shepson, P. B.: A computer model study of multiphase chemistry in the Arctic boundary layer during polar sunrise, *J. Geophys. Res. Atmos.*, 105, 15 131–15 145, <https://doi.org/10.1029/2000JD900004>, 2000.
- Molina, M. J. and Rowland, F. S.: Stratospheric sink for chlorofluoromethanes: chlorine atom-catalysed destruction of ozone, *Nature*, 249, 810–812, 1974.
- Oltmans, S. J.: Surface ozone measurements in clean air, *J. Geophys. Res.*, 86, 1174–1180, <https://doi.org/10.1029/JC086iC02p01174>, 1981.
- Piot, M.: Modeling Halogen Chemistry during Ozone Depletion Events in Polar Spring: A Model Study, Ph.D. thesis, University of Heidelberg, Germany, 2007.
- Platt, U. and Janssen, C.: Observation and role of the free radicals NO₃, ClO, BrO and IO in the troposphere, *Faraday Discuss.*, 100, 175–198, <https://doi.org/10.1039/FD9950000175>, 1995.
- Platt, U. and Lehrer, E.: Arctic tropospheric ozone chemistry, ARCTOC, no. 64 in Air pollution research report, European Commission Directorate-General, Science, Research and Development, Luxembourg, 1997.

- Prather, M. and Jaffe, A. H.: Global impact of the Antarctic ozone hole: Chemical propagation, *Journal of Geophysical Research: Atmospheres*, 95, 3473–3492, <https://doi.org/https://doi.org/10.1029/JD095iD04p03473>, 1990.
- 585 Riedel, K., Allan, W., Weller, R., and Schrems, O.: Discrepancies between formaldehyde measurements and methane oxidation model predictions in the Antarctic troposphere: An assessment of other possible formaldehyde sources, *Journal of Geophysical Research: Atmospheres*, 110, <https://doi.org/https://doi.org/10.1029/2005JD005859>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2005JD005859>, 2005.
- Rogers, J. D.: Ultraviolet absorption cross sections and atmospheric photodissociation rate constants of formaldehyde, *The Journal of Physical Chemistry*, 94, 4011–4015, <https://doi.org/10.1021/j100373a025>, <https://doi.org/10.1021/j100373a025>, 1990.
- 590 Roscoe, H. K. and Roscoe, J.: Polar tropospheric ozone depletion events observed in the International Geophysical Year of 1958, *Atmospheric Chemistry and Physics*, 6, 3303–3314, <https://doi.org/10.5194/acp-6-3303-2006>, <https://acp.copernicus.org/articles/6/3303/2006/>, 2006.
- Seinfeld, J. and Pandis, S.: *Atmos. Chem. Phys.: from air pollution to climate change*, A Wiley-Interscience publications, Wiley, 2006.
- Simpson, W. R., von Glasow, R., Riedel, K., Anderson, P., Ariya, P., Bottenheim, J., Burrows, J., Carpenter, L. J., Frieß, U., Goodsite, M. E., Heard, D., Hutterli, M., Jacobi, H.-W., Kaleschke, L., Neff, B., Plane, J., Platt, U., Richter, A., Roscoe, H., Sander, R., Shepson, P., Sodeau, J., Steffen, A., Wagner, T., and Wolff, E.: Halogens and their role in polar boundary-layer ozone depletion, *Atmos. Chem. Phys.*, 7, 4375–4418, <https://doi.org/10.5194/acp-7-4375-2007>, 2007.
- 595 Stull, R. B.: *An Introduction to Boundary Layer Meteorology*, Kluwer Academic Publishers, The Netherlands, 1988.
- Tarasick, D. W. and Bottenheim, J. W.: Surface ozone depletion episodes in the Arctic and Antarctic from historical ozonesonde records, *Atmos. Chem. Phys.*, 2, 197–205, <https://doi.org/10.5194/acp-2-197-2002>, 2002.
- 600 Toumi, R., Haigh, J. D., and Law, K. S.: A tropospheric ozone-lightning climate feedback, *Geophysical Research Letters*, 23, 1037–1040, <https://doi.org/https://doi.org/10.1029/96GL00944>, 1996.
- Turányi, T.: KINAL - a program package for kinetic analysis of reaction mechanisms., *Comput. Chem.*, 14, 253–254, 1990.
- Vaghjiani, G. L. and Ravishankara, A. R.: Absorption cross sections of CH₃OOH, H₂O₂, and D₂O₂ vapors between 210 and 365 nm at 297 K, *Journal of Geophysical Research: Atmospheres*, 94, 3487–3492, <https://doi.org/https://doi.org/10.1029/JD094iD03p03487>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/JD094iD03p03487>, 1989.
- 605 van Oss, R. F. and Spurr, R. J.: Fast and accurate 4 and 6 stream linearized discrete ordinate radiative transfer models for ozone profile retrieval, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 75, 177–220, [https://doi.org/https://doi.org/10.1016/S0022-4073\(01\)00246-1](https://doi.org/https://doi.org/10.1016/S0022-4073(01)00246-1), <https://www.sciencedirect.com/science/article/pii/S0022407301002461>, 2002.
- Wagner, T., Ibrahim, O., Sinreich, R., Frieß, U., von Glasow, R., and Platt, U.: Enhanced tropospheric BrO over Antarctic sea ice in mid winter observed by MAX-DOAS on board the research vessel Polarstern, *Atmos. Chem. Phys.*, 7, 3129–3142, <https://doi.org/10.5194/acp-7-3129-2007>, 2007.
- 610 Wennberg, P.: Atmospheric chemistry: Bromine explosion, *Nature*, 397, 299–301, <https://doi.org/10.1038/16805>, 1999.
- Wilmouth, D. M., Hanisco, T. F., Donahue, N. M., and Anderson, J. G.: Fourier Transform Ultraviolet Spectroscopy of the A $2\pi/3/2 \leftarrow \Psi 2\pi/3/2$ Transition of BrO, *The Journal of Physical Chemistry A*, 103, 8935–8945, <https://doi.org/10.1021/jp991651o>, <https://doi.org/10.1021/jp991651o>, 1999.
- 615 Zhou, J., Cao, L., and Li, S.: Influence of the Background Nitrogen Oxides on the Tropospheric Ozone Depletion Events in the Arctic during Springtime, *Atmosphere*, 11, <https://doi.org/10.3390/atmos11040344>, <https://www.mdpi.com/2073-4433/11/4/344>, 2020.