

Limitations of assuming internal mixing between different aerosol species: a case study with sulfate geoengineering simulations

Daniele Visioni¹, Simone Tilmes², Charles Bardeen², Michael Mills², Douglas G. MacMartin¹, Ben Kravitz^{3,4}, and Jadwiga Richter⁵

¹Sibley School of Mechanical and Aerospace Engineering, Cornell University, Ithaca, NY, USA

²Atmospheric Chemistry, Observations, and Modeling Laboratory, National Center for Atmospheric Research, Boulder CO, USA

³Department of Earth and Atmospheric Sciences, Indiana University, Bloomington, IN, USA

⁴Atmospheric Sciences and Global Change Division, Pacific Northwest National Laboratory, Richland, WA, USA

⁵Climate and Global Dynamics Laboratory, National Center for Atmospheric Research, Boulder CO, USA

Correspondence: Daniele Visioni (daniele.visioni@cornell.edu)

Abstract. Simulating the complex aerosol microphysical processes in a comprehensive Earth System Model can be very computationally intensive and therefore many models utilize a modal approach, where aerosol size distributions are represented by observations-derived lognormal functions, and internal mixing between different aerosol species [within an aerosol mode](#) is often assumed. This approach has been shown to yield satisfactory results in a large array of applications, but there may

5 be cases where the simplification in this approach may produce some shortcomings. In this work we show specific conditions under which the current approximations used in some modal approaches might yield [some](#)-incorrect answers. Using results from the Community Earth System Model v1 (CESM1) Geoengineering Large Ensemble (GLENS) project, we analyze the effects in the troposphere of a continuous increasing load of sulfate aerosols in the stratosphere, with the aim of counteracting the surface warming produced by non-mitigated increasing greenhouse gases concentration between 2020-2100. We show
10 that the simulated results pertaining to the evolution of sea salt and dust aerosols in the upper troposphere are not realistic due to internal mixing assumptions in the modal aerosol treatment, which in this case reduces the size, and thus the settling velocities, of those particles and ultimately changes their mixing ratio below the tropopause. The unnatural increase of these aerosol species affects, in turn, the simulation of upper tropospheric ice formation, resulting in an increase in ice clouds that is not due to any meaningful physical mechanisms. While we show that this does not significantly affect the overall
15 results of the simulations, we point to some areas where results should be interpreted with care in modeling simulations using similar approximations: in particular, the evolution of upper tropospheric clouds when large amount of sulfate is present in the stratosphere, as after a large explosive volcanic eruption or in similar stratospheric aerosol injection cases. Finally, we suggest that this could be avoided if sulfate aerosols in the coarse mode, the predominant species in these situations, are treated separately from other aerosol species in the model.

A comprehensive representation of aerosol processes in Earth System models is crucial for a variety of reasons. Aerosols are one of the main short-term forcing agents in the climate system, and uncertainties in the estimate of their overall forcing effects are still quite large Boucher et al. (2013). Directly, they scatter incoming solar radiation, thus influencing surface temperatures, and also absorb both solar radiation and outgoing planetary radiation, locally increasing air temperatures, modifying circulation patterns, and affecting meteorology. Indirectly, they affect the climate by modifying cloud cover, acting as cloud condensation nuclei (CCN) and ice nuclei (IN), and changing clouds' physical and radiative properties (also through local atmospheric heating). Lastly, once they are deposited to the surface, they affect land and ice albedo through the melting of snow and ice. Aerosols at the surface also contribute to air pollution (in particular with particulate matter (PM) below $2.5\text{ }\mu\text{m}$ and $10\text{ }\mu\text{m}$ in diameter, Ayala et al. (2012)) and may affect soils and ecosystems through acid deposition (Vet et al. (2014)).

30

Atmospheric aerosols may have different sizes, ranging from $0.001\text{ }\mu\text{m}$ to $100\text{ }\mu\text{m}$ in diameter, and their characteristics (number concentration, mass, shape, chemical composition and other physical properties) may change through emission (from natural or anthropogenic sources), nucleation (defined as the formation of new particles), coagulation (defined as the combination of existing aerosol particles, decreasing their number concentration but leaving the overall mass unaltered), condensational growth of chemical species in vapor form (such as H_2SO_4 , NH_3 , HNO_3 and volatile organics gases) on existing particles, gas-phase and aqueous-phase chemistry, water uptake (Ghan and Zaveri (2007)) and their removal through gravitational settling (dry deposition), in-cloud scavenging (defined as the removal of aerosol particles by precipitation particles), below-cloud scavenging (defined as the capture of aerosol particles by precipitating droplets, Feng (2009)). They can, of course, be composed of different chemical species: the main components are usually sea salt, mineral dust, black carbon, organic matter, nitrate, ammonium and sulfate. Wang et al. (2020) provide a recent overview of all aerosol-related processes: this list of processes can give an idea of how challenging it can be to represent aerosol processes correctly in climate models, both due to the uncertainties in each process and in their combination, and because of the high computational burden necessary to correctly reproduce the known parts of each process. For this reason, most processes need to treat aerosols in parametrized ways with different complexities, ranging from the least (bulk model, considering all aerosols as described by a single mean radius and standard deviation) to the most complex (sectional model, considering a large number of "bins" of different sizes for each species where the aerosols can grow or shrink). An intermediate approach is the modal method: it works on the assumption that the actual size distribution of the aerosols can be represented by the combination of multiple log-normal functions with fixed standard deviations that are based on observations. Furthermore, a modal approach to aerosol microphysics must also make assumptions about the internal and external mixing of aerosol species (Fassi-Fihri et al. (1997)): internal mixing is defined as the modeling of an aerosol particle as a mixing of the constituent species (Clarke et al. (2004)), either as an homogeneous mix or as a coated sphere containing a solid core and coated by a liquid exterior. More simply put, it assumes that different species of aerosols in every gridbox are present in the same proportion, and that their physical characteristics can be described by the same average size distribution. External mixing is defined as the treatment of particle populations as composed of different species with

55 distinct compositions, and no assumption needs to be made with regards to the localized proportion of the various species, as they are treated differently (Riemer et al., 2019). Some discussion and comparisons of the application of sectional and modal microphysics modules in climate models can be found in Weisenstein et al. (2007); Kokkola et al. (2009); Kleinschmitt et al. (2017), and more specifically for sulfate geoengineering applications in Laakso et al. (2021).

56 A modal treatment of aerosols in climate models has been shown to successfully reproduce aerosol measurements in various occasions. In this work, in particular, we focus on the Community Atmosphere Model version 5.0 (CAM5) and its implementation in the Community Earth System Model (CESM), using the Modal Aerosol Module with 3 modes (MAM3, Liu et al. (2012)). This approach has been shown to correctly reproduce tropospheric aerosols in a baseline climate (see Liu et al. (2012); Samset et al. (2014)), and also to correctly reproduce the evolution of the stratospheric sulfate aerosol layer in the extreme case of explosive volcanic eruptions (Robock (2000)), as detailed by Mills et al. (2016, 2017) for the Whole Atmosphere Community Climate Model (WACCM) version of CESM1, with a high top (140 km) and 70 vertical layers. This model has been used for climate simulations of Stratospheric Aerosol Intervention (SAI), a form of climate engineering that has been proposed (Budyko, 1978; Crutzen, 2006) (Budyko, 1969; Crutzen, 2006) as a way to temporarily reduce global surface temperatures by mimicking the cooling effect of volcanic eruptions through injecting SO_2 in the stratosphere. In both the case of volcanic eruptions and for proposed artificial injections, once SO_2 reaches the stratosphere, it would oxidize, eventually forming gaseous H_2SO_4 , which would then either nucleate forming new sulfate aerosol particles of $\text{H}_2\text{SO}_4\text{-H}_2\text{O}$ or condense onto existing particles (already existing particles could also coagulate, resulting in larger particles with the same overall mass). While this process transforms all SO_2 into aerosols with an e-folding time of around 1 month (Mills et al. (2017)), the produced aerosols tend to remain in the stratosphere for one year or more (Visioni et al. (2018b)), and are removed through large-scale stratospheric circulation, that moves them poleward, or by gravitational settling or stratosphere-troposphere exchange, crossing the tropopause. Once in the troposphere, they are quickly removed through dry or wet deposition (Kremser et al. (2016)).

75 Climate engineering simulations with CESM1(WACCM) in the Geoengineering Large Ensemble (GLENS; Tilmes et al. (2018a)) have shown that it would be possible to maintain global surface temperatures at 2010-2029 levels even under a scenario where emissions of greenhouse gases continue unabated, by increasing the amount of SO_2 injected throughout the century. This technique may be able to reduce some of the harmful climatic effects produced by the temperature increase ((Tilmes et al. (2018a); Kravitz et al. (2019))), by reducing the amount of incoming sunlight, but would not be a perfect solution. In fact, the atmosphere and surface would be impacted in various ways by the produced aerosols, for example, the chemical composition of the stratosphere and the dynamical response produced by the local stratospheric heating (Richter et al. (2017); Tilmes et al. (2018b)), which in turn can influence precipitation (Simpson et al. (2019)) and the high-latitude seasonal cycle of temperature (Jiang et al. (2019)). Once the aerosols are deposited to the surface they may also affect soils; however, considering the much larger amount of tropospheric sulfate aerosols produced by both natural and human activities, those settling down from the stratosphere would have a marginal impact everywhere except in some pristine areas (Visioni et al. (2020b)). Other processes that may be affected by SAI include aerosol interactions with cirrus clouds, which are a key component of the radiation balance. Water clouds at lower altitudes have a net cooling effect because they reflect solar radiation (see for instance Yi et al. (2017)); on the other hand, the effect of cirrus clouds made of ice crystals and produced in the upper

troposphere by supercooled water particles is harder to determine, but is widely understood to be positive (i.e., it produces a net warming at the surface) (Fusina et al. (2007)) due to their trapping of outgoing planetary radiation. Kuebbeler et al. (2012) and Visioni et al. (2018a) found in two different climate models (ECHAM-HAM5 and ULAQ-CCM, using a modal and a sectional aerosol approach, respectively) that the change in vertical temperature gradient resulting from the stratospheric heating would reduce the formation of tropical cirrus ice clouds by less than 10%, thus contributing to the surface cooling. This effect is tied to the fact that, in both models, the amount of water vapor reaching the upper troposphere and necessary for the cloud formation is directly tied to the available turbulent kinetic energy, which is a function of the vertical temperature gradient, and is therefore a purely dynamical effect. Cirisan et al. (2013) on the other hand investigated the possible microphysical effect of the increased sulfate load on the cirrus cloud formation, but found no significant impact due to the much larger size of the aerosols from the stratosphere compared to those already present in the troposphere by other activities.

In this study, as an example to illustrate some of the shortcomings of the modal aerosol treatment in MAM3, we use the GLENS simulations to explore the effects of large amount of sulfate aerosols in the stratosphere on tropospheric aerosol concentrations and on cirrus ice cloud formation in CESM1(WACCM). In the following sections, we will briefly describe how aerosol microphysics and cirrus ice formation are parameterized in Section 2, then discuss how aerosols in the upper troposphere change in the simulations of interest in Section 3 and how that affects upper tropospheric ice in Section 4. Finally, possible radiative effects at the top of the atmosphere of the identified changes will be discussed in Section 5.

105 2 Model description

In this section we will briefly describe the simulations used in this work, and then describe the components of the model that will be of use in our analyses.

The Geoengineering Large Ensemble (GLENS, Tilmes et al. (2018a)) is an ensemble of simulations performed with CESM1 (WACCM), with all simulations using surface emissions from the Representative Concentration Pathway 8.5 (RCP) scenario. 110 21 ensemble members are available for the period 2010-2030 under RCP8.5 (hereafter, termed Baseline). From each of these, in 2020 a scenario is simulated where SO_2 is injected at four locations: 30° N and S, with injections at 23 km altitude, and 15° N and S at 25 km altitude. Each year, an algorithm (described by Kravitz et al. (2017)) determines the amount of SO_2 to be injected 115 at each location in order to maintain the mean surface temperature, the inter-hemispheric surface temperature gradient, and the equator-to-pole temperature gradient at their 2020 values in the presence of growing greenhouse gas concentrations. All the simulations of SAI are extended to 2100 (hereafter, termed GLENS), and 4 ensemble members of the Baseline cases (without SAI) are run to at least the year 2097. Please note that, compared to the original simulations described in Tilmes et al. (2018a), one more ensemble member is available for both GLENS and Baseline (from 20 to 21 and from 3 to 4, respectively).

2.1 The Modal Aerosol Model in CESM

120 The Modal Aerosol Model (MAM) has been first described by Liu et al. (2012), and therein evaluated for tropospheric aerosol loads. Some modifications have been made for including interactive stratospheric aerosols, described in depth by Mills et al. (2016) and therein validated in the case of the Pinatubo 1991 eruption.

125 For CESM1(WACCM) climate simulations, the 3-mode version (MAM3) is used, with aerosol species divided in three different lognormal modes of fixed width, named Aitken (dry diameter size range between 0.015 and 0.05 μm), Accumulation (between 0.05 and 0.3 μm) and Coarse mode (between 0.80 and 3.65 μm ; all estimates from Liu et al. (2012)), going from the smallest to the largest (in this manuscript, we will always refer to the names of the modes with a capitalized letter to avoid ambiguity). Sulfate particles can grow by condensation (from H_2SO_4 vapor condensing on existing particles, locally maintaining particle numbers but increasing mass) or coagulation (locally reducing particle numbers but maintaining mass). When, by either process, the tail of the distribution of the particles in one of the modes grows to a size that would nominally be in the 130 size range for the larger mode, the particles are transferred to the larger mode. This is done, as detailed by Easter et al. (2004), by defining a lower and upper limit for the dry diameter in each mode, and transferring part of the local number concentration to the larger mode when the threshold is surpassed. In the stratosphere the sulfate particles can also shrink due to evaporation, thus allowing for a Coarse-to-Accumulation mode transfer.

135 Compared to the more computationally-demanding MAM version with 7 aerosol modes (MAM7), in MAM3 all aerosol species are considered internally mixed within each of the three modes, thus sharing composition and size distribution. The mass for the single species has to be conserved, both globally and locally, and thus can only change in each gridbox if particles are moved from one gridbox to the other, either because of air mass movement, or because of gravitational settling or other 140 tropospheric removal processes. Liu et al. (2012) justify this approach by noting that the sources of different aerosol types are geographically separated, thus unlikely to affect each other in the simplified version used for long climate applications. Coarse mode sulfate aerosols are, in quiescent conditions, also scarce in the troposphere: in the AeroCom multi-model mean (Textor et al. (2006)) they were determined to be less than 2% of all sulfate aerosols, with a preponderance of particles in the Accumulation mode in the troposphere. The assumption by Liu et al. (2012) would therefore hold in the background atmosphere. We 145 will show in the next section that the presence of the Coarse mode sulfate particles produced by SAI fundamentally breaks this assumption, unnaturally modifying the size and quantity of non-sulfate aerosol species in the upper troposphere.

2.2 The formation of cirrus ice clouds in CESM

Cirrus clouds play an important part in the radiation budget, but have been generally represented poorly in general circulation models (GCMs) for a variety of reasons, amongst them a poor horizontal resolution which fails to capture the scale needed to represent some of the processes and a large spread in the ice water content simulated by models (Jiang et al. (2012)). There 150 are two processes that can produce ice crystals in the upper troposphere: homogeneous nucleation of ice crystals from sulfate aerosols and heterogeneous immersion freezing of mineral dust. Normally, homogeneous freezing is assumed to be the

dominant process, although there are instances where this is not the case (Knopf and Koop (2006); Cziczo et al. (2013)). In the atmospheric model used in CESM1(WACCM), the Community Atmospheric Model version 5 (CAM5), both processes are present and we will briefly describe both below.

155

The process of homogeneous freezing is based on the assumption that only sulfate particles in the Aitken mode work as ice nuclei (IN), using the portion of the Aitken mode particles with radii greater than 0.1 microns (Liu and Penner (2005); Liu et al. (2007)). Other works have used all available sulfate modes for homogeneous freezing (Shi et al. (2015)), and this would clearly affect the results. The process of homogeneous freezing is assumed to happen only when clouds are present, together with a probability distribution from Kärcher and Burkhardt (2008) that determines when the supersaturation is above the threshold for homogeneous freezing to happen. This means that homogeneous freezing can happen only when local vertical velocities are high (Kärcher and Lohmann (2002)). Those velocities are determined following Morrison and Pinto (2005) as

$$w_{sub} = \sqrt{\frac{2}{3} TKE} \quad (1)$$

with the turbulent kinetic energy (TKE) determined using a steady state energy balance as in Bretherton and Park (2009).

165

For heterogeneous freezing, only Coarse mode dust particles are assumed to be available IN (although the exclusion of soot particles and other particles as inefficient ice nuclei has been debated; see Kärcher and Burkhardt (2008)). Given the internal mixing assumption in MAM3, the number of available dust nuclei is determined as a fraction of the overall amount of Coarse mode particles given the mass of dust and the overall aerosol mass

$$N_d = \left(\frac{m_d}{m_d + m_{ss}} \right) \times N_c \quad (2)$$

170

where m_d is the mass of dust in the Coarse mode, m_{ss} is the mass of sea salt in the Coarse mode, and N_c is the total number of particles in the Coarse mode. This approach assumes a negligible amount of Coarse mode sulfate in the upper tropical stratosphere in the denominator of the fraction. Another shortcoming of the heterogeneous freezing description in this version of WACCM, as already discussed by Mills et al. (2017), is that when heterogeneous freezing occurs, IN that have nucleated to form ice particles are not removed from the available reservoir, thus allowing for too many particles to be formed via heterogeneous freezing.

175

This approach to the microphysical modeling of cirrus clouds in CESM has been discussed recently by Maloney et al. (2019), comparing it with a more complex approach using the coupled Community Aerosol and Radiation Model for Atmospheres (CARMA) sectional scheme, and comparing both with measurements from the National Aeronautics and Space Administration Airborne Tropical Tropopause Experiment (ATTREX 3) and with observations from the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) onboard the CALIPSO satellite. They found that while the CAM5 approach was capable of correctly representing the annual average cloud fraction profile in the tropics (20°N to 20°S), it tended to underrepresent the cirrus fraction in the tropical tropopause layer. Similar results were found previously also by Bardeen et al. (2013).

3 Simulated tropospheric aerosols

The geographical distribution of tropospheric aerosols in the unperturbed atmosphere depends mainly on the surface sources 185 of the aerosols or aerosol precursors; once the aerosols are produced in the atmosphere or are directly emitted, they can be affected by long range transport, upward currents and sediment through gravitational settling or scavenging. An in-depth analyses of sources and sinks of atmospheric aerosols can be found, for instance, in Lamarque et al. (2010). We focus here on sulfate, dust and sea salt, as their changes will be of interest when analyzing the effects on cirrus formation. Sulfate particles 190 are mostly formed through surface emission of SO_2 , which can be oxidized to form sulfuric acid and then sulfate particles by condensation in the smaller (Aitken) mode. For this reason, Coarse mode sulfate particles near the surface are much less frequent than dust or sea salt, as shown in Fig. 1a-[SO4a](#) as compared to 1-Da and 1-SSa. For the latter two, their concentration decreases with increasing altitude as the scavenging processes reduce their number, whereas there is an additional sulfate layer 195 in the stratosphere (the Junge layer), discovered and discussed by Junge et al. (1961). This happens because of the presence of various sources of sulfate aerosols in the stratosphere, even without considering volcanic aerosols from explosive volcanic 200 eruptions. In particular, surface emissions of carbonyl sulfide (OCS) and dimethyl sulfide (DMS), which are light, well mixed gases that may reach the stratosphere where they are oxidized, forming sulfate aerosols (Vet et al. (2014)). Meteoric sulfur also plays a part in the formation of the layer (see Gómez Martín et al. (2017)). The quantity of sulfate aerosols produced in the stratosphere with an artificial injection of SO_2 would be larger by some orders of magnitude compared to the amount in the quiescent Junge layer; the full stratospheric distribution in the case of GLENS has been described elsewhere (i.e., 205 Kravitz et al. (2019)), so here we simply report that at 50 hPa, between 30°N and 30°S, the average mass concentration for the last 20 years of simulation is 163 $\mu\text{g}/\text{kg}$, compared to 0.4 $\mu\text{g}/\text{kg}$ in the unperturbed case; while this simulation considered high cooling ($\sim 4^\circ\text{C}$ reduction in global mean temperature), even much smaller amounts of SAI would lead to substantially 210 more stratospheric sulfate than the unperturbed case. This magnitude is driven by the long lifetime of the produced aerosol particles in the stratosphere (around 12 months, Visoni et al. (2018b)). Once the particles cross the tropopause, the various removal processes strongly reduce the lifetime, driving the concentration down. This is visible already in Fig. 1a-[SO4a](#), with the tropopause clearly visible as a change in concentration. Nonetheless, the average concentration of sulfate under SAI is 215 much larger even in the upper troposphere (Fig. 1b-[SO4b](#)) and only returns close to the Baseline levels close to the surface, where other sulfate sources are predominant (Fig. 1e-[SO4c](#)).

Fig. 1 also shows the behavior of dust and sea salt in the Coarse mode in the same altitude-latitude region (those are 220 the only other aerosol species considered in this version of CESM1 in the Coarse mode: black carbon is only found in the Accumulation mode). Various behaviors can be found depending on the altitude of analyses for the GLENS case. We focus here on the uppermost troposphere, right below the tropopause (black boxes in Fig. 1) and on a layer below, in the upper troposphere (green boxes). In the upper level, both dust (Fig. 1-Db) and sea salt (Fig. 1-SSb) present a behavior that is much 225 more similar to that of sulfate: a large, initial increase in concentration in the first 5 to 10 years followed by either a constant evolution or by a small decrease (from 0.1 to 0.2 $\mu\text{m}/\text{kg}$ for dust in the upper layer, from 0.06 to 0.2 $\mu\text{m}/\text{kg}$ and from 0.9 to 0.17 $\mu\text{m}/\text{kg}$ for sea salt in the upper and lower layer). Only dust in the lower layer (Fig. 1-Dc) does not show such an abrupt 230

initial increase. On the other hand, in the Baseline case the dust mixing ratios show a decrease by 25% over the 80 years of analyses in the lower level. There is no immediate physical mechanism by which such changes would be observed in aerosol species independent of sulfate; therefore, a more in-depth analysis is necessary.

220 One possible explanation would be a change in the surface emissions of these species. However, the vertical profiles of tropical concentration for the three species seem to exclude that (Fig. 1, all three d) panels and Fig. 3). This is further confirmed by the analyses of the overall burdens shown in Fig. 2. For these, no initial abrupt change can be identified, and the time evolution shows the opposite behavior from that found in Fig. 1, with a minor reduction in GLENS compared to Baseline. The behavior of the Baseline case is consistent with previous projections: Mahowald and Luo (2003) predicted a reduction in overall dust 225 emission with increasing GHGs due to higher precipitation and more surface moisture produced by the warmer air; thus the precipitation decrease in GLENS (Cheng et al. (2019)) would exacerbate differences between dust burden in GLENS and the baseline. Struthers et al. (2013) on the other hand projected a small increase in sea salt aerosols mostly as a result of increased surface wind speeds. Having excluded that changes in surface emissions are the cause of the abrupt increase identified, we further investigate the smaller aerosol modes (Fig. 3). In this case, we see a much closer agreement of the values in GLENS to those 230 of the initial baseline conditions for sea salt and dust: this indicates that the initial aim of the GLENS simulations, which was to maintain the state of the climate as close as possible to 2010-2029 conditions, also helps with not modifying surface emission sources of those species. The time evolution of the globally averaged quantities for these species and modes also doesn't show the same abrupt change as that found in Fig. 1 (see Fig. S1 and S2). So clearly the cause of the rapid change in concentration of the Coarse mode must be the result of the very rapid increase in SO_4 aerosol concentration by more than 4 orders of magnitude.

235

The solution to this conundrum can be found by analyzing the behavior of the simulated radius of the particles in the three modes (Fig. 4). In MAM3, all aerosol species within an aerosol mode are assumed to be internally mixed. The simplified assumption that different aerosols peak at different locations in the lower atmosphere is reasonable for the baseline scenario (Fig. 1 for the vertical distribution, and Fig. 2 for the spatial distribution). The parameterization requires that in each gridbox, 240 all aerosols in one mode are treated as the same in terms of their size distribution, so for each mode, only one mode diameter and one number concentration is used in the online calculations (since the geometric standard deviation is constant throughout the atmosphere). The mass concentrations of the different species are then calculated using a reference density for each species (Liu et al. (2012)). This information is necessary to calculate the gravitational settling velocities (at each layer), which follow Seinfeld and Pandis (2016), where the equation of a free-falling spherical particle in a fluid which has reached terminal velocity 245 (which is done in less than 1×10^{-6} for particles of diameter of $\simeq 1 \mu\text{m}$) is shown to be:

$$v_t = \left(\frac{4gD_p C_c \rho_p}{3C_D \rho} \right)^{1/2} \quad (3)$$

where D_p is the diameter of the particle, ρ_p is the density of the particle, C_c is the slip-correction factor that accounts for non-continuum effects (dependent on the diameter of the particles), and C_D is the empirical drag coefficient (dependent on the Reynolds number R_e).




Figure 1. Mass concentration (in μg per kg of air) of Coarse mode species (Sulfate (SO4), Dust (D) and Sea Salt (SS), from top to bottom). On the left, panels a) indicate the Baseline (2020-2029) zonal mean concentration for the respective species, with the black and green boxes indicating the areas considered in the averages for the central panels (panels b and c). The uppermost limit of the green boxes and the lowermost limit of the black boxes coincide. Panels b) and c) for each species show the annual evolution of the concentration in the black (panels b) and green (panels c) boxes, with lighter lines showing the single ensemble realizations, and the thicker lines show the ensemble mean; red lines indicate the Baseline simulations, while blue lines indicate the GLENS simulations. Black dashed lines in the b) and c) panels indicate the periods of analyses for the a) and d) panels, respectively. On the right, panels ed) show the vertical profiles in both cases (Baseline, in red, and GLENS, in blue) for the period 2080-2099. Black dashed The black lines in the e) and d) panels indicate the periods of analyses for in the ab) and d) panels, respectively; while the same green lines in the d) panels indicate the altitude of analyses in the b) and c) panels.




Figure 2. Zonally averaged total column burden of dust (a) and sea salt (c) (considering all modes) in 2010-2029 (red, dashed line), 2080-2099 in Baseline (red line) and 2080-2099 in GLENS (blue line). Evolution of the globally integrated dust (b) and sea salt (d) burden (Tg/yr). Lighter lines show the single ensemble realizations, while the thicker lines show the ensemble mean.




Figure 3. Mass concentration (in μg per kg of air) of Sulfate, Dust and Sea Salt in the Aitken (smallest) and Accumulation (intermediate) mode, between 30°N and 30°S . Lighter lines show the single ensemble realizations, while the thicker lines show the ensemble mean.

250 The initial identified changes can thus be explained this way. The SO_4 formed in the stratosphere is the predominant form of aerosols in the stratosphere, deciding the radius of the particles at those altitudes. Most of these particles are in the Coarse mode, therefore changing the radius considered by the model in those grid-boxes. Once the sulfate aerosols cross the tropopause (whether by gravitational settling, mostly in the tropics, or by large scale circulation, at higher latitudes) they are added to the particle distribution already present in the upper troposphere. Due to the small amount of Coarse mode aerosols there, and due 255 to the internal mixing assumption in MAM3, the dust and sea salt aerosols already present are “forced” to reduce in size (and to conserve mass, to increase in number concentration, see Fig. S3). This produces a drop in the overall size by around 25% within 3 years in the upper troposphere. Because of the strong dependence of gravitational settling velocities on size, the result is a drop in these velocities that, for dust aerosols, can be estimated to be over 40%. This is not a direct output of the model, but we estimate it from Eq. 3 using the approach of Seinfeld and Pandis (2016). The clear drop in v_t can then explain the initial 260 identification of increase in non-sulfate aerosol species: reducing settling deposition abruptly directly affects the concentration by decreasing removal, especially in regions where contributions from below are scarce. And the reason why this is only visible in the upper troposphere is that, further below, the preponderance of pre-existing, background Coarse mode aerosol reduces the strength of this effect, not changing the radii. The preponderance of other removal mechanisms (in-cloud scavenging) also reduces the effect of the gravitational settling changes.

265 **4 Effect on cirrus clouds formation**

We have shown that the simulated changes in dust and sea salt in the upper troposphere in GLENS using CESM1(WACCM), are non-physical in nature. However, by looking at the concentration of the same species in the Baseline conditions it is clear that the overall presence of those aerosols at the altitudes of analyses is small, and therefore would have a negligible effect both 270 when looking at the direct effects (for instance, on the radiative fluxes, which would be overshadowed by the effects of the sulfate aerosols in the stratosphere) and when eventually looking at surface effects. We show in this section, however, that their change does influence the simulation of ice clouds formation in the upper troposphere. Visioni et al. (2021) showed that some noticeable changes in cloud cover are present in the GLENS simulations, and that when separating the contribution of different 275 types of clouds, most of these changes were attributable to changes in high clouds, that are defined in CESM1(WACCM) as all clouds formed between the altitude of 400 and 50 hPa. Furthermore, these changes were not as noticeable in other performed simulations where the same cooling as GLENS was achieved using a solar constant reduction approach, and when together with that the stratospheric heating produced by the aerosols was imposed, without the presence of the aerosol themselves. This would point towards a contribution to the changes in high cloud cover in GLENS by some effects connected to the aerosol 280 themselves. In particular, this points to some changes in the freezing processes that produce ice crystals in the upper troposphere, given that at those altitudes there is much less water than in the lower troposphere, and most of that is in the form of ice rather than liquid droplets (see Fig. S5).




Figure 4. Wet radius (in μm) of all three modes in the Baseline case ([2010-2029], a-b-c-d) and in GLENS ([2080-2099], e-f-g-h). g) Vertical distribution of aerosol size in all three modes between 30°N and 30°S . Black boxes in panels c-f) indicate the area analyzed in the average for panel h), where the temporal evolution in Baseline (red) and GLENS (blue) is shown. Lighter lines show the single ensemble realizations, while the thicker lines show the ensemble mean. i) Estimate of the changes in gravitational settling velocities following Seinfeld and Pandis (2016).

As we detailed in Section 2.2, there are two types of processes that may lead to the formation of ice crystals at those altitudes: the spontaneous freezing of small aqueous sulfate droplets at high relative humidity rates (Chen et al. (2000)), known as homogeneous freezing, and the freezing of water droplets mediated by the presence of insoluble aerosol particles (ice nuclei, IN), known as heterogeneous nucleation (Diehl and Wurzler (2004)), which can happen at lower relative humidity conditions and higher temperatures (Knopf and Koop (2006)). Due to the difficulties in measuring the amount of ice crystals in the upper troposphere, and to the challenges of representing the processes in models, there are plenty of uncertainties over the predominance of one of the two formation processes over the other. In CESM, only sulfate particles in the Aitken mode can act as the substrate over which homogeneous freezing can take place; this is generally understood to be correct, as too large particles would be increasingly harder to freeze (Chen et al. (2000)). Since Aitken mode particles do not change in GLENS (see Fig. 3a), changes in homogeneous freezing due to that can be excluded, as predicted already by Cirisan et al. (2013). However, changes in homogeneous freezing may happen as a result of dynamical changes in the local vertical velocities that determine the amount of water vapor in the upper troposphere. Since the processes that determine those local vertical velocities happen at a much smaller scale compared to the resolution of climate model, they are usually parameterized in climate models as a function 295 of vertical stability (see Eq. 2.2). Both previous studies analyzing the response of ice clouds in a SAI scenario, Kuebbeler et al. (2012) and Visioni et al. (2018a), used climate models with a similar parameterization as CESM, and noticed a reduction in ice cloud coverage because the warming of the stratosphere, combined with a cooling of the surface, produced a reduction in the vertical temperature gradients and thus in the turbulent kinetic energy used to determine the velocities. A similar process can then be expected in CESM. Lastly, there would be no physical reason why SAI would change heterogeneous freezing in ice 300 clouds. However, given what we identified in the previous section related to dust aerosols in the upper troposphere, an influence on freezing processes cannot be excluded.

In Fig. 5 we show an analyses of the changes in the amount of ice in clouds under GLENS. Two particularly different behaviors can be noted when separating the effects at low latitudes (30°N-30°S) versus the effects noted elsewhere. At low latitudes, the dynamical changes produced by the different vertical stabilities dominate, resulting in a slow decrease in ice concentration 305 in line with predictions discussed by Kuebbeler et al. (2012) and Visioni et al. (2018a) (which is not a surprise, considering the very similar parameterization of sub-grid vertical velocities). At higher latitudes, where there are fewer changes in the vertical temperature gradient (since the stratosphere warms much less, see Richter et al. (2017)), the predominant effect is the sudden increase in IN, resulting from the simplified aerosol treatment in the Coarse mode. The increase in ice formation the model “sees” is therefore not physical in origin, but simply an artifact of the microphysical parameterization and thus should not be 310 treated as a physical side-effect to SAI. A clue as to the two different mechanisms at play can be determined by observing the rate of change at low versus high latitudes. At low latitudes, the changes happen gradually as the stratospheric sulfate load (and thus the warming) increases. At high altitudes, the in-cloud ice changes are very abrupt, much more similar to the aerosol changes identified previously in this work. Further confirmation can be derived by observing the differences between the full GLENS simulations and the simulations described by Visioni et al. (2021) with stratospheric heating imposed, but no aerosols. 315 By comparing the two (see bottom panels of Fig. 5), the effect of the presence of the aerosols can be separated from the dynamical ones, and indeed this confirms the different nature of the ice changes, since the two simulations (aerosols+stratospheric

heating and stratospheric heating alone) show no differences at low latitudes, whereas they are as different as GLENS-Baseline at high latitudes.

Further information would be gained by observing the changes in heterogeneous versus homogeneous freezing in these 320 simulations. However, output fields that separate the two processes have not been saved in the original simulations from GLENS. Therefore, a complete analysis is not possible using the whole ensemble and time period. Noting that the main, unexplained processes (with our current understanding of ice formation) happen in the very first decade of the simulations, for the following analyses we have re-run the first 21 years of one of the ensemble members with the same configuration (thus 325 bit-by-bit, the result is the same) but retaining information related to the two different upper tropospheric freezing processes. The results in Fig. 6 confirm our previous observations. For homogeneous freezing, very few changes are initially observable at all latitudes, whereas large changes are observable in the concentration of what the model considers to be IN for heterogeneous freezing, sharply in the first few years. This increase is much larger than the increase in Coarse mode dust aerosols identified in Fig. 1, where the overall concentration was doubled, while here the increase is much larger, at more than 10 times the amount 330 in the first two years. To partially explain this change we refer to the equation used by the model to determine the fraction of the available dust in the Coarse mode, which is used as IN for heterogeneous freezing (Eq. 2.2). The available number of dust particles that can be used by the model depends on the overall amount of Coarse mode particles, but is weighted by which fraction, locally, of aerosol is dust compared to sea salt. This formula (which in normal conditions gives satisfactorily 335 results, see Maloney et al. (2019)) presumes a lack of sulfate in the upper troposphere. In our case however, as we have shown, the amount of Coarse mode sulfate is actually by far preponderant in the upper troposphere. So while the number of Coarse mode particles N_C grows (see Fig. S3), the fraction of all aerosols that is considered does not. This is a separated, but similar, problem as the one noted for the behavior of the aerosols. We tested how sensitive this assumption is by performing an identical simulation to the one that gave us information about homogeneous and heterogeneous freezing, but modifying the code so that 340 Eq. 2.2 becomes

$$N_d = \left(\frac{m_d}{m_d + m_{ss} + m_s} \right) \times N_C \quad (4)$$

thus correctly accounting for the mass of sulfate when considering which fraction of the number of Coarse mode particles 345 locally present is made of dust aerosols. We show the results of these sensitivity simulation in Fig. 6 using dotted lines: as expected, this doesn't change the homogeneous freezing processes, but it does change the amount of heterogeneous IN, and reduces the non-physical increase in cloud ice identified in Fig. 5 at high latitudes.




Figure 5. Baseline concentration of cloud ice number concentration (a, number of particles per cm^{-3}) and fractional occurrence (b) in the period 2010-2030. In the panels c-e) and f-h), the yearly evolution of the differences between GLENS and Baseline (in the 2010-2030 period, as for the panels above) are shown for the three latitudinal boxes separately. In the small panels below, the effect of the changes produced by the aerosols are isolated by plotting the difference between GLENS and the simulations with surface cooling and stratospheric heating but no aerosols.




Figure 6. Evolution of the occurrence of homogeneous freezing in clouds (a), the concentration of what the model considers to be IN for homogeneous freezing (Aitken mode sulfate) (b) and the concentration of IN for heterogeneous freezing (coarse mode dust) (c) in GLENS in the first 21 years of simulation, for the three latitudinal band already considered in Fig. 5. On the right, the column mean between 50 and 400 hPa is considered (thick lines), together with the case described in the text with a sensitivity test for Eq. 2.2 (dotted lines). In panel f) the ice number concentration shown in Fig. 5 is compared with that for the sensitivity experiment.

5 Radiative effects

345 Here we investigate whether the changes shown in the previous section produce some signal in the modeled radiative fluxes at the top of the atmosphere, as that would be important in order to determine how significant the effect is. To do so we use the method suggested by Ghan (2013) to separate both the direct radiative effect of the aerosols and the effect of cloud changes produced by the aerosols. This is possible here as the radiative output of CESM is comprised of both the full radiative fluxes (\mathbf{F}_{ALL}) at the top of the atmosphere (TOA), that consider all effects from aerosols, clouds and atmospheric composition, but
 350 also of the diagnosed Clear Sky radiative fluxes (where the radiative transfer model calculates the radiative fluxes as if no clouds were present, \mathbf{F}_{CLEAR}), Clean Sky radiative fluxes (where the radiative fluxes are calculated as if no aerosols were present, \mathbf{F}_{CLEAN}) and their combination ($\mathbf{F}_{CLEAN,CLEAR}$). This allows a separation of the contributions of the added aerosols and their effects for both the longwave (LW) and shortwave (SW) components: the direct forcing produced by the aerosols direct interaction with the radiation ($\Delta\mathbf{F}_D$) can be defined as

$$355 \quad \Delta\mathbf{F}_D = (\mathbf{F}_{ALL}^{GLENS} - \mathbf{F}_{CLEAN}^{GLENS}) - (\mathbf{F}_{ALL}^{BASELINE} - \mathbf{F}_{CLEAN}^{BASELINE}) \quad (5)$$

and the forcing produced by the changes in clouds ($\Delta\mathbf{C}$) can be defined as

$$\Delta\mathbf{F} = (\mathbf{F}_{CLEAN}^{GLENS} - \mathbf{F}_{CLEAN,CLEAR}^{GLENS}) - (\mathbf{F}_{CLEAN}^{BASELINE} - \mathbf{F}_{CLEAN,CLEAR}^{BASELINE}) \quad (6)$$

As Ghan (2013) notes, normally these fluxes are just an estimation of the real components of the forcing, as to properly estimate them it would be necessary to maintain surface and tropospheric temperatures fixed between the two cases to avoid changes
 360 in the radiative emissions of the troposphere, which would be at different temperature in the two cases. In our case, however, the GLENS simulations have been performed on purpose to maintain similar global surface temperatures as the Baseline 2010-2030 period: some small regional temperature differences are still present (Tilmes et al. (2018a)), but overall this can be considered a minor factor and the estimated forcing quite robust if we are comparing the radiative fluxes always against the 2010-2030 period. In the case of cloud changes, this further assures that we are not counting effects produced by the surface
 365 warming, which might also locally modify cloud cover.

In Fig. 7 we show the global evolution of $\Delta\mathbf{F}_D$ and $\Delta\mathbf{C}$ in GLENS. The aerosol direct radiative effect is linear, negative in the shortwave (implying a cooling) as they reflect incoming sunlight and ~~negative~~ positive in the longwave (implying a warming) as they absorb and re-emit longwave radiation, with an overall negative effect, as expected. We show the latitudinal breakdown of the fluxes in panels 7b-c normalized by the stratospheric AOD produced by the SO_2 injections for the initial
 370 period (we pick 2026-2035 to avoid the very first few years when the algorithm that determines the SO_2 injections is still converging, see Kravitz et al. (2017)) and the last twenty years (2081-2100). A partial non-linearity can be identified between the two periods. This can be explained by the slight increase in stratospheric sulfate aerosol during the whole period of analyses (see Visioni et al. (2020a), the effective radius in the stratosphere in GLENS grows from $0.4 \mu\text{m}$ to over $0.5 \mu\text{m}$ as a result of the increasing injection rates resulting in more coagulation of SO_2 with pre-existing particles). Larger aerosols scatter slightly less
 375 efficiently, but they absorb more LW radiation. The changes in upper tropospheric aerosols described before do not influence these radiative fluxes, as they are, in proportion, negligible compared to the stratospheric increase. Looking at the $\Delta\mathbf{C}$ fluxes,

however, we can see that the behavior of the LW ΔC shows a very different behavior compared with the SW. This effect is positive in the first $\simeq 30$ years, and then negative afterwards, whereas the SW forcing is always positive. The SW forcing is easily explainable as directly connected to the presence of the aerosols above: if less sunlight reaches the troposphere, because 380 it is partially reflected in the stratosphere, the same clouds would appear as to be less reflective, hence the positive sign of the SW (cloud masking). This doesn't apply to the LW, as globally the surface temperatures are the same, but can be explained by noting that the main contributor to LW trapping by clouds is upper tropospheric ice (Fusina et al. (2007)). Since, in the first decades, mid-latitudinal ice clouds increase as shown in Fig. 5 because of "more" dust aerosols acting as IN, they would trap more outgoing LW radiation. When tropical ice clouds start decreasing because of the dynamical mechanism produced 385 by stratospheric heating, the positive bump is erased by the negative forcing (less ice clouds, more LW radiation escaping to space) already analyzed by Visioni et al. (2018a). This is further confirmed by the latitudinal breakdown of ΔC in panels 7e-f, where the (normalized by stratospheric AOD) forcing in the LW goes from positive, in the first 10 years, to negative in the last 20.

390 6 Conclusions

In this work we have identified the presence of some weaknesses in the 3-mode modal approach (MAM3) used in CESM1(WACCM) when a large presence of aerosols settles down from the stratosphere, as under stratospheric sulfate aerosol injection, which results in some artificial changes to cirrus clouds. MAM3 only separates the species in three modes depending on their size, by treating all aerosol species as the same (internal mixing assumption). When a large amount of sulfate is produced (mostly 395 in the coarse, largest mode) in the stratosphere following the injection of SO_2 in large quantities, the particles slowly descend upon the troposphere, where they are quickly removed. However, the size of the Coarse mode particles is different (smaller) compared to that of the Coarse mode particles already present in the troposphere, whose source is at the surface: therefore, MAM3 "sees" an abrupt decrease in all aerosol size in each gridbox in the upper troposphere for the coarse mode, resulting in smaller settling velocities for aerosol species that would not be otherwise affected in the real world. This effect results in an 400 increase in the mass of particles of all species at those altitudes even if there would be no natural causes for it to happen. The effect is small, and its direct effect on the cooling produced by the stratospheric aerosols negligible. However, the unnatural addition of dust in the upper troposphere results in more particles that the model can use as solid ice nuclei for the freezing of ice particles in clouds. This effect is particularly evident at mid-and-high latitudes, where the low relative humidity and lack of aerosols make other ice formation processes more scarce in normal conditions. The formation of these ice crystals, as 405 simulated by the model, does produce a noticeable change in the cloud forcing, especially in the first years, when the effect from the incorrect presence of the dust aerosols is large compared to other, dynamical effects that tend to reduce the amount of ice crystals in the upper troposphere when a warming of the stratosphere is present.




Figure 7. Global mean evolution of TOA radiative fluxes in the LW (orange), SW (green) and Net (black) for the forcing directly produced by the aerosols ΔF_d (a) and by the clouds as changed by the presence of the aerosols (ΔC) (d). In panels b-c and e-f, respectively, the latitudinal breakdown of the global fluxes in two periods (2026-2035 and 2081-2100) is shown for the various components. On the right of each panel, the global mean in the considered period is shown with a triangle of the same color, and the value on the right. Lighter lines show the single ensemble realizations, while the thicker lines show the ensemble mean.

Given the set-up of the GLENS experiments, this effect is counteracted by the presence of the algorithm which determines
410 how much SO_2 is needed every year to counteract the effect of the increased emissions in RCP8.5: even if the radiative im-
balance were to be large, the algorithm would just prescribe more SO_2 to be injected, therefore resulting, overall, in the same
415 global mean temperatures as if the error in the ice cloud formation was not there. In the sensitivity simulations produced for
Section 4, which reduced the amount of ice clouds incorrectly formed by the model, the cumulative amount of SO_2 injected
420 in the first 20 years was 131 Tg- SO_2 (6.2 Tg- SO_2 /year), whereas the same period in the default simulation had a cumulative
amount of SO_2 of 154 Tg- SO_2 (7.3 Tg- SO_2 /year): considering that, aside from the change in Eq. 2.2, the two simulations
were otherwise the same, we can assume that this difference arises from the changes in ice clouds as simulated by the model.
Overall then, if we assume this effect is constant throughout the whole simulation period, it would account for an cumulative
425 injected 88 Tg- SO_2 , in the entire century: while this amount might seem large, it accounts for less than two years of anthro-
pogenic emissions of SO_2 at present (Visioni et al. (2020b)). This, furthermore, assumes a very large injection to overcome a
considerable amount of warming (over 4 °) over the entire century. More moderate mitigation scenarios would require far less
430 cumulative injections of SO_2 (for instance, see Tilmes et al. (2020)).

This doesn't imply that the effect can be ignored, but suggests that going forward when using a modal approach to aerosol
435 microphysics, simulations where large amount of sulfate is present in the stratosphere should treat sulfate aerosols in a separate
coarse mode that is not internally mixed with the other species compared (dust, sea salt, black carbon, etc.). A similar approach
has already been used to include an additional primary carbon mode in MAM4 (Liu et al., 2016) in order to account for pro-
cesses that affect the microphysical properties of primary carbonaceous aerosols in the atmosphere. There could be various
applications where this observation might be useful: SAI is one example, but the simulation of explosive volcanic eruptions is
another case where it could be useful: for instance, Schmidt et al. (2018) used the same CESM1(WACCM) model described
here to estimate the global volcanic radiative forcing in the last 45 years, and ran into a similar observation as found in this
440 paper, but did not find an explanation for it. The mechanism that produced, in their simulations, an increase in ice particles in
the upper troposphere is definitely the same we have encountered here, and would explain some of the LW forcing changes di-
agnosed in their simulations with a similar method (see Fig. 4 and 5, Schmidt et al. (2018)). More generally, it would be crucial
to properly represent the upper troposphere in the case of volcanic eruptions to verify the influence of volcanic eruptions on ice
clouds as observed by some studies in the observational record (see for instance Friberg et al. (2015)). Future modeling efforts
aimed at better understanding the climatic effect of volcanic eruptions, such as the Volcanic Model Intercomparison Project
(Clyne et al. (2021)) or the Interactive Stratospheric Aerosol Model Intercomparison Project (Timmreck et al. (2018)) should
take this into account and consider how that might affect some of their results, since models with similar modal approaches are
present in both.

440

Lastly, this observation would also be crucial in studies that aim to combine sulfate injections with the artificial seeding of
upper tropospheric ice clouds with solid nuclei, to increase the size of the ice crystal and make them sediment faster (cloud
seeding, Gasparini et al. (2020)). Cao et al. (2017), for instance, proposed a combination of the two methods to stabilize global

temperatures and precipitation: such simulations, performed with CESM1(WACCM) or any model with a similar microphysical
445 approach would not give meaningful results. Our study shows that more care should be given to make sure that the climate
models used for simulating sulfate geoengineering application are not applied outside of the parameters for which the models
will give reliable answers. Most of the time, models are also compared against volcanic events to make sure they are properly
simulating the results, but not all effects are immediately visible in those cases as compared to a long term geoengineering
simulation, and a comparison of cloud changes after a volcanic eruption may be complicated by various factors (Friberg et al.
450 (2015)).

Data availability. All simulations analyzed in this work are available online. All GLENS simulations data are available at <https://doi.org/10.5065/D6JH3JXX>. Data from the stratospheric heating simulations used for Fig. 5 are available at <https://ecommons.cornell.edu/handle/1813/76871>.

Author contributions. DV performed the analysis and wrote the manuscript. ST, CB and MJM contributed to the analysis and with the
455 interpretation of MAM3 and ice formation processes results. DGM, BK and YR offered helpful suggestions on the manuscript and analysis.

Competing interests. The authors declare no competing interests

Acknowledgements. Support for DGM was provided by the National Science Foundation through agreement CBET-1818759. Support for
DV was provided by the Atkinson Center for a Sustainable Future at Cornell University. Support for BK was provided in part by the National
Sciences Foundation through agreement CBET-1931641, the Indiana University Environmental Resilience Institute, and the Prepared for
460 Environmental Change Grand Challenge initiative. The Pacific Northwest National Laboratory is operated for the U.S. Department of Energy
by Battelle Memorial Institute under contract DE-AC05-76RL01830. The CESM project is supported primarily by the National Science
Foundation.

References

- Ayala, A., Brauer, M., Mauderly, J. L., and Samet, J. M.: Air pollutants and sources associated with health effects, *Air Quality, Atmosphere & Health*, 5, 151–167, 2012.
- 465 Bardeen, C. G., Gettelman, A., Jensen, E. J., Heymsfield, A., Conley, A. J., Delanoë, J., Deng, M., and Toon, O. B.: Improved cirrus simulations in a general circulation model using CARMA sectional microphysics, *Journal of Geophysical Research: Atmospheres*, 118, 11,679–11,697, [https://doi.org/https://doi.org/10.1002/2013JD020193](https://doi.org/10.1002/2013JD020193), <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2013JD020193>, 2013.
- 470 Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S., Sherwood, S., Stevens, B., and Zhang, X.: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, in: Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by IPCC, IPCC, https://www.ipcc.ch/site/assets/uploads/2018/02/WG1AR5_Chapter07_FINAL-1.pdf, 2013.
- 475 Bretherton, C. S. and Park, S.: A New Moist Turbulence Parameterization in the Community Atmosphere Model, *Journal of Climate*, 22, 3422 – 3448, <https://doi.org/10.1175/2008JCLI2556.1>, <https://journals.ametsoc.org/view/journals/clim/22/12/2008jcli2556.1.xml>, 2009.
- Budyko, M. I.: The effect of solar radiation variations on the climate of the Earth, *Tellus*, 21, 611–619, [https://doi.org/https://doi.org/10.1111/j.2153-3490.1969.tb00466.x](https://doi.org/10.1111/j.2153-3490.1969.tb00466.x), <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.2153-3490.1969.tb00466.x>, 1969.
- 480 Budyko, M. I.: The Climate of the Future, American Geophysical Union, <https://doi.org/10.1002/9781118665251.ch7>, <http://dx.doi.org/10.1002/9781118665251.ch7>, 1978.
- Cao, L., Duan, L., Bala, G., and Caldeira, K.: Simultaneous stabilization of global temperature and precipitation through cocktail geoengineering, *Geophysical Research Letters*, 44, 7429–7437, [https://doi.org/https://doi.org/10.1002/2017GL074281](https://doi.org/10.1002/2017GL074281), <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2017GL074281>, 2017.
- 485 Chen, Y., DeMott, P. J., Kreidenweis, S. M., Rogers, D. C., and Sherman, D. E.: Ice Formation by Sulfate and Sulfuric Acid Aerosol Particles under Upper-Tropospheric Conditions, *Journal of the Atmospheric Sciences*, 57, 3752 – 3766, [https://doi.org/10.1175/1520-0469\(2000\)057<3752:IFBSAS>2.0.CO;2](https://doi.org/10.1175/1520-0469(2000)057<3752:IFBSAS>2.0.CO;2), https://journals.ametsoc.org/view/journals/atsc/57/22/1520-0469_2000_057_3752_ifbsas_2.0.co_2.xml, 2000.
- 490 Cheng, W., MacMartin, D. G., Dagon, K., Kravitz, B., Tilmes, S., Richter, J. H., Mills, M. J., and Simpson, I. R.: Soil Moisture and Other Hydrological Changes in a Stratospheric Aerosol Geoengineering Large Ensemble, *Journal of Geophysical Research: Atmospheres*, 124, 12 773–12 793, <https://doi.org/10.1029/2018JD030237>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018JD030237>, 2019.
- Cirisan, A., Spichtinger, P., Luo, B. P., Weisenstein, D. K., Wernli, H., Lohmann, U., and Peter, T.: Microphysical and radiative changes in cirrus clouds by geoengineering the stratosphere, *Journal of Geophysical Research: Atmospheres*, 118, 4533–4548, [https://doi.org/https://doi.org/10.1002/jgrd.50388](https://doi.org/10.1002/jgrd.50388), <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/jgrd.50388>, 2013.
- 495 Clarke, A. D., Shinozuka, Y., Kapustin, V. N., Howell, S., Huebert, B., Doherty, S., Anderson, T., Covert, D., Anderson, J., Hua, X., Moore II, K. G., McNaughton, C., Carmichael, G., and Weber, R.: Size distributions and mixtures of dust and black carbon aerosol in Asian outflow: Physiochemistry and optical properties, *Journal of Geophysical Research: Atmospheres*, 109, [https://doi.org/https://doi.org/10.1029/2003JD004378](https://doi.org/10.1029/2003JD004378), <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2003JD004378>, 2004.

- 500 Clyne, M., Lamarque, J.-F., Mills, M. J., Khodri, M., Ball, W., Bekki, S., Dhomse, S. S., Lebas, N., Mann, G., Marshall, L., Niemeier, U., Poulain, V., Robock, A., Rozanov, E., Schmidt, A., Stenke, A., Sukhodolov, T., Timmreck, C., Toohey, M., Tummon, F., Zanchettin, D., Zhu, Y., and Toon, O. B.: Model physics and chemistry causing intermodel disagreement within the VolMIP-Tambora Interactive Stratospheric Aerosol ensemble, *Atmospheric Chemistry and Physics*, 21, 3317–3343, <https://doi.org/10.5194/acp-21-3317-2021>, <https://acp.copernicus.org/articles/21/3317/2021/>, 2021.
- 505 Crutzen, P. J.: Albedo Enhancement by Stratospheric Sulfur Injections: A Contribution to Resolve a Policy Dilemma?, *Climatic Change*, 77, 211–220, <https://doi.org/10.1007/s10584-006-9101-y>, <http://dx.doi.org/10.1007/s10584-006-9101-y>, 2006.
- 510 Cziczo, D. J., Froyd, K. D., Hoose, C., Jensen, E. J., Diao, M., Zondlo, M. A., Smith, J. B., Twohy, C. H., and Murphy, D. M.: Clarifying the Dominant Sources and Mechanisms of Cirrus Cloud Formation, *Science*, 340, 1320–1324, <https://doi.org/10.1126/science.1234145>, <https://science.sciencemag.org/content/340/6138/1320>, 2013.
- 515 Diehl, K. and Wurzler, S.: Heterogeneous Drop Freezing in the Immersion Mode: Model Calculations Considering Soluble and Insoluble Particles in the Drops, *Journal of the Atmospheric Sciences*, 61, 2063 – 2072, [https://doi.org/10.1175/1520-0469\(2004\)061<2063:HDFITI>2.0.CO;2](https://doi.org/10.1175/1520-0469(2004)061<2063:HDFITI>2.0.CO;2), https://journals.ametsoc.org/view/journals/atsc/61/16/1520-0469_2004_061_2063_hdfiti_2.0.co_2.xml, 2004.
- 520 Easter, R. C., Ghan, S. J., Zhang, Y., Saylor, R. D., Chapman, E. G., Laulainen, N. S., Abdul-Razzak, H., Leung, L. R., Bian, X., and Zaveri, R. A.: MIRAGE: Model description and evaluation of aerosols and trace gases, *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/https://doi.org/10.1029/2004JD004571>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2004JD004571>, 2004.
- 525 Fassi-Fihri, A., Suhre, K., and Rosset, R.: Internal and external mixing in atmospheric aerosols by coagulation: Impact on the optical and hygroscopic properties of the sulphate-soot system, *Atmospheric Environment*, 31, 1393 – 1402, [https://doi.org/https://doi.org/10.1016/S1352-2310\(96\)00341-X](https://doi.org/https://doi.org/10.1016/S1352-2310(96)00341-X), <http://www.sciencedirect.com/science/article/pii/S135223109600341X>, 1997.
- 530 Feng, J.: A size-resolved model for below-cloud scavenging of aerosols by snowfall, *Journal of Geophysical Research: Atmospheres*, 114, <https://doi.org/https://doi.org/10.1029/2008JD011012>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2008JD011012>, 2009.
- 535 Friberg, J., Martinsson, B. G., Sporre, M. K., Andersson, S. M., Brenninkmeijer, C. A. M., Hermann, M., van Velthoven, P. F. J., and Zahn, A.: Influence of volcanic eruptions on midlatitude upper tropospheric aerosol and consequences for cirrus clouds, *Earth and Space Science*, 2, 285–300, <https://doi.org/https://doi.org/10.1002/2015EA000110>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2015EA000110>, 2015.
- Fusina, F., Spichtinger, P., and Lohmann, U.: Impact of ice supersaturated regions and thin cirrus on radiation in the midlatitudes, *Journal of Geophysical Research: Atmospheres*, 112, n/a–n/a, <https://doi.org/10.1029/2007JD008449>, <http://dx.doi.org/10.1029/2007JD008449>, d24S14, 2007.
- 540 Gasparini, B., McGraw, Z., Storelvmo, T., and Lohmann, U.: To what extent can cirrus cloud seeding counteract global warming?, *Environmental Research Letters*, 15, 054 002, <https://doi.org/10.1088/1748-9326/ab71a3>, <https://doi.org/10.1088%2F1748-9326%2Fab71a3>, 2020.
- Ghan, S. J.: Technical Note: Estimating aerosol effects on cloud radiative forcing, *Atmospheric Chemistry and Physics*, 13, 9971–9974, <https://doi.org/10.5194/acp-13-9971-2013>, <https://acp.copernicus.org/articles/13/9971/2013/>, 2013.
- 545 Ghan, S. J. and Zaveri, R. A.: Parameterization of optical properties for hydrated internally mixed aerosol, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/https://doi.org/10.1029/2006JD007927>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2006JD007927>, 2007.

- Gómez Martín, J. C., Brooke, J. S. A., Feng, W., Höpfner, M., Mills, M. J., and Plane, J. M. C.: Impacts of meteoric sulfur in the Earth's atmosphere, *Journal of Geophysical Research: Atmospheres*, 122, 7678–7701, [https://doi.org/https://doi.org/10.1002/2017JD027218](https://doi.org/10.1002/2017JD027218), <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2017JD027218>, 2017.
- 540 Jiang, J., Cao, L., MacMartin, D. G., Simpson, I. R., Kravitz, B., Cheng, W., Visioni, D., Tilmes, S., Richter, J. H., and Mills, M. J.: Stratospheric Sulfate Aerosol Geoengineering Could Alter the High-Latitude Seasonal Cycle, *Geophysical Research Letters*, 46, 14 153–14 163, <https://doi.org/10.1029/2019GL085758>, 2019.
- Jiang, J. H., Su, H., Zhai, C., Perun, V. S., Del Genio, A., Nazarenko, L. S., Donner, L. J., Horowitz, L., Seman, C., Cole, J., Gettelman, A., Ringer, M. A., Rotstain, L., Jeffrey, S., Wu, T., Brient, F., Dufresne, J.-L., Kawai, H., Koshiro, T., Watanabe, M., LÉcuyer, T. S., Volodin, 545 E. M., Iversen, T., Drange, H., Mesquita, M. D. S., Read, W. G., Waters, J. W., Tian, B., Teixeira, J., and Stephens, G. L.: Evaluation of cloud and water vapor simulations in CMIP5 climate models using NASA "A-Train" satellite observations, *Journal of Geophysical Research: Atmospheres*, 117, [https://doi.org/https://doi.org/10.1029/2011JD017237](https://doi.org/10.1029/2011JD017237), <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2011JD017237>, 2012.
- 550 Junge, C. E., Chagnon, C. W., and Manson, J. E.: A World-wide Stratospheric Aerosol Layer, *Science*, 133, 1478–1479, <https://doi.org/10.1126/science.133.3463.1478-a>, <https://science.sciencemag.org/content/133/3463/1478.2>, 1961.
- Kleinschmitt, C., Boucher, O., Bekki, S., Lott, F., and Platt, U.: The Sectional Stratospheric Sulfate Aerosol module (S3A-v1) within the LMDZ general circulation model: description and evaluation against stratospheric aerosol observations, *Geoscientific Model Development*, 10, 3359–3378, 2017.
- 555 Knopf, D. A. and Koop, T.: Heterogeneous nucleation of ice on surrogates of mineral dust, *Journal of Geophysical Research: Atmospheres*, 111, [https://doi.org/https://doi.org/10.1029/2005JD006894](https://doi.org/10.1029/2005JD006894), <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2005JD006894>, 2006.
- Kokkola, H., Hommel, R., Kazil, J., Niemeier, U., Partanen, A.-I., Feichter, J., and Timmreck, C.: Aerosol microphysics modules in the framework of the ECHAM5 climate model - intercomparison under stratospheric conditions, *Geoscientific Model Development*, 2, 97–112, 2009.
- 560 Kravitz, B., Lamarque, J.-F., Tribbia, J. J., Tilmes, S., Vitt, F., Richter, J. H., MacMartin, D. G., and Mills, M. J.: First Simulations of Designing Stratospheric Sulfate Aerosol Geoengineering to Meet Multiple Simultaneous Climate Objectives, *Journal of Geophysical Research: Atmospheres*, 122, 12,616–12,634, <https://doi.org/10.1002/2017jd026874>, 2017.
- Kravitz, B., MacMartin, D. G., Tilmes, S., Richter, J. H., Mills, M. J., Cheng, W., Dagon, K., Glanville, A. S., Lamarque, J.-F., Simpson, 565 I. R., Tribbia, J., and Vitt, F.: Comparing Surface and Stratospheric Impacts of Geoengineering With Different SO₂ Injection Strategies, *Journal of Geophysical Research: Atmospheres*, 124, 7900–7918, <https://doi.org/10.1029/2019JD030329>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019JD030329>, 2019.
- Kremser, S., Thomason, L. W., von Hobe, M., Hermann, M., Deshler, T., Timmreck, C., Toohey, M., Stenke, A., Schwarz, J. P., Weigel, R., Fueglistaler, S., Prata, F. J., Vernier, J.-P., Schlager, H., Barnes, J. E., Antuña-Marrero, J.-C., Fairlie, D., Palm, M., Mahieu, E., Notholt, J., Rex, M., Bingen, C., Vanhellemont, F., Bourassa, A., Plane, J. M. C., Klocke, D., Carn, S. A., Clarisse, L., Trickl, T., Neely, R., 570 James, A. D., Rieger, L., Wilson, J. C., and Meland, B.: Stratospheric aerosol—Observations, processes, and impact on climate, *Reviews of Geophysics*, 54, 278–335, [https://doi.org/https://doi.org/10.1002/2015RG000511](https://doi.org/10.1002/2015RG000511), <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2015RG000511>, 2016.
- Kuebbeler, M., Lohmann, U., and Feichter, J.: Effects of stratospheric sulfate aerosol geo-engineering on cirrus clouds, *Geophysical Research Letters*, 39, <https://doi.org/10.1029/2012GL053797>, <http://dx.doi.org/10.1029/2012GL053797>, I23803, 2012.

- 575 Kärcher, B. and Burkhardt, U.: A cirrus cloud scheme for general circulation models, *Quarterly Journal of the Royal Meteorological Society*, 134, 1439–1461, <https://doi.org/10.1002/qj.301>, <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.301>, 2008.
- Kärcher, B. and Lohmann, U.: A parameterization of cirrus cloud formation: Homogeneous freezing of supercooled aerosols, *Journal of Geophysical Research: Atmospheres*, 107, AAC 4–1–AAC 4–10, <https://doi.org/10.1029/2001JD000470>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2001JD000470>, 2002.
- 580 Laakso, A., Niemeier, U., Visioni, D., Tilmes, S., and Kokkola, H.: Dependency of the impacts of geoengineering on the stratospheric sulfur injection strategy part 1: Intercomparison of modal and sectional aerosol module, *Atmospheric Chemistry and Physics Discussions*, 2021, 1–36, <https://doi.org/10.5194/acp-2021-526>, <https://acp.copernicus.org/preprints/acp-2021-526/>, 2021.
- Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Lioussse, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., 585 Riahi, K., and van Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application, *Atmospheric Chemistry and Physics*, 10, 7017–7039, <https://doi.org/10.5194/acp-10-7017-2010>, <https://acp.copernicus.org/articles/10/7017/2010/>, 2010.
- Liu, X. and Penner, J. E.: Ice nucleation parameterization for global models, *Meteorologische Zeitschrift*, 14, 499–514, <https://doi.org/10.1127/0941-2948/2005/0059>, <http://dx.doi.org/10.1127/0941-2948/2005/0059>, 2005.
- 590 Liu, X., Penner, J. E., Ghan, S. J., and Wang, M.: Inclusion of Ice Microphysics in the NCAR Community Atmospheric Model Version 3 (CAM3), *Journal of Climate*, 20, 4526 – 4547, <https://doi.org/10.1175/JCLI4264.1>, <https://journals.ametsoc.org/view/journals/clim/20/18/jcli4264.1.xml>, 2007.
- Liu, X., Easter, R. C., Ghan, S. J., Zaveri, R., Rasch, P., Shi, X., Lamarque, J.-F., Gettelman, A., Morrison, H., Vitt, F., Conley, A., Park, S., Neale, R., Hannay, C., Ekman, A. M. L., Hess, P., Mahowald, N., Collins, W., Iacono, M. J., Bretherton, C. S., Flanner, M. G., and Mitchell, 595 D.: Toward a minimal representation of aerosols in climate models: description and evaluation in the Community Atmosphere Model CAM5, *Geoscientific Model Development*, 5, 709–739, <https://doi.org/10.5194/gmd-5-709-2012>, <https://gmd.copernicus.org/articles/5/709/2012/>, 2012.
- Liu, X., Ma, P.-L., Wang, H., Tilmes, S., Singh, B., Easter, R. C., Ghan, S. J., and Rasch, P. J.: Description and evaluation of a new four-mode version of the Modal Aerosol Module (MAM4) within version 5.3 of the Community Atmosphere Model, *Geoscientific Model Development*, 9, 505–522, 2016.
- 600 Mahowald, N. M. and Luo, C.: A less dusty future?, *Geophysical Research Letters*, 30, <https://doi.org/10.1029/2003GL017880>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2003GL017880>, 2003.
- Maloney, C., Bardeen, C., Toon, O. B., Jensen, E., Woods, S., Thornberry, T., Pfister, L., Diskin, G., and Bui, T. P.: An Evaluation of the 605 Representation of Tropical Tropopause Cirrus in the CESM/CARMA Model Using Satellite and Aircraft Observations, *Journal of Geophysical Research: Atmospheres*, 124, 8659–8687, <https://doi.org/10.1029/2018JD029720>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018JD029720>, 2019.
- Mills, M. J., Schmidt, A., Easter, R., Solomon, S., Kinnison, D. E., Ghan, S. J., Neely III, R. R., Marsh, D. R., Conley, A., Bardeen, C. G., and Gettelman, A.: Global volcanic aerosol properties derived from emissions, 1990–2014, using CESM1(WACCM), *Journal of Geophysical Research: Atmospheres*, 121, 2332–2348, <https://doi.org/10.1002/2015JD024290>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2015JD024290>, 2016.
- Mills, M. J., Richter, J. H., Tilmes, S., Kravitz, B., Macmartin, D. G., Glanville, A. A., Tribbia, J. J., Lamarque, J. F., Vitt, F., Schmidt, A., Gettelman, A., Hannay, C., Bacmeister, J. T., and Kinnison, D. E.: Radiative and chemical response to interactive strato- 610

- spheric sulfate aerosols in fully coupled CESM1(WACCM), *Journal of Geophysical Research: Atmospheres*, 122, 13,061–13,078, <https://doi.org/10.1002/2017JD027006>, 2017.
- 615 Morrison, H. and Pinto, J. O.: Mesoscale Modeling of Springtime Arctic Mixed-Phase Stratiform Clouds Using a New Two-Moment Bulk Microphysics Scheme, *Journal of the Atmospheric Sciences*, 62, 3683 – 3704, <https://doi.org/10.1175/JAS3564.1>, <https://journals.ametsoc.org/view/journals/atsc/62/10/jas3564.1.xml>, 2005.
- Richter, J. H., Tilmes, S., Mills, M. J., Tribbia, J. J., Kravitz, B., Macmartin, D. G., Vitt, F., and Lamarque, J. F.: Stratospheric dynamical response and ozone feedbacks in the presence of SO₂ injections, *Journal of Geophysical Research: Atmospheres*, 122, 12,557–12,573, 620 <https://doi.org/10.1002/2017JD026912>, 2017.
- Riemer, N., Ault, A. P., West, M., Craig, R. L., and Curtis, J. H.: Aerosol Mixing State: Measurements, Modeling, and Impacts, *Reviews of Geophysics*, 57, 187–249, [https://doi.org/https://doi.org/10.1029/2018RG000615](https://doi.org/10.1029/2018RG000615), <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018RG000615>, 2019.
- Robock, A.: Volcanic eruptions and climate, *Reviews of Geophysics*, 38, 191–219, <https://doi.org/10.1029/1998RG000054>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/1998RG000054>, 2000.
- 625 Samset, B. H., Myhre, G., Herber, A., Kondo, Y., Li, S.-M., Moteki, N., Koike, M., Oshima, N., Schwarz, J. P., Balkanski, Y., Bauer, S. E., Bellouin, N., Berntsen, T. K., Bian, H., Chin, M., Diehl, T., Easter, R. C., Ghan, S. J., Iversen, T., Kirkevåg, A., Lamarque, J.-F., Lin, G., Liu, X., Penner, J. E., Schulz, M., Seland, Ø., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., and Zhang, K.: Modelled black carbon radiative forcing and atmospheric lifetime in AeroCom Phase II constrained by aircraft observations, *Atmospheric Chemistry and Physics*, 630 14, 12 465–12 477, <https://doi.org/10.5194/acp-14-12465-2014>, <https://acp.copernicus.org/articles/14/12465/2014/>, 2014.
- Schmidt, A., Mills, M. J., Ghan, S., Gregory, J. M., Allan, R. P., Andrews, T., Bardeen, C. G., Conley, A., Forster, P. M., Gettelman, A., Portmann, R. W., Solomon, S., and Toon, O. B.: Volcanic Radiative Forcing From 1979 to 2015, *Journal of Geophysical Research: Atmospheres*, 123, 12 491–12 508, <https://doi.org/10.1029/2018JD028776>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018JD028776>, 2018.
- 635 Seinfeld, J. H. and Pandis, S. N.: *Atmospheric chemistry and physics: from air pollution to climate change*, John Wiley & Sons, 2016.
- Shi, X., Liu, X., and Zhang, K.: Effects of pre-existing ice crystals on cirrus clouds and comparison between different ice nucleation parameterizations with the Community Atmosphere Model (CAM5), *Atmospheric Chemistry and Physics*, 15, 1503–1520, <https://doi.org/10.5194/acp-15-1503-2015>, <https://acp.copernicus.org/articles/15/1503/2015/>, 2015.
- Simpson, I., Tilmes, S., Richter, J., Kravitz, B., MacMartin, D., Mills, M., Fasullo, J., and Pendergrass, A.: The regional hydroclimate 640 response to stratospheric sulfate geoengineering and the role of stratospheric heating, *Journal of Geophysical Research: Atmospheres*, p. 2019JD031093, <https://doi.org/10.1029/2019JD031093>, <https://onlinelibrary.wiley.com/doi/abs/10.1029/2019JD031093>, 2019.
- Struthers, H., Ekman, A. M. L., Glantz, P., Iversen, T., Kirkevag, A., Seland, O., Martensson, E. M., Noone, K., and Nilsson, E. D.: Climate-induced changes in sea salt aerosol number emissions: 1870 to 2100, *Journal of Geophysical Research: Atmospheres*, 118, 670–682, <https://doi.org/https://doi.org/10.1002/jgrd.50129>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/jgrd.50129>, 2013.
- 645 Textor, C., Schulz, M., Guibert, S., Kinne, S., Balkanski, Y., Bauer, S., Berntsen, T., Berglen, T., Boucher, O., Chin, M., Dentener, F., Diehl, T., Easter, R., Feichter, H., Fillmore, D., Ghan, S., Ginoux, P., Gong, S., Grini, A., Hendricks, J., Horowitz, L., Huang, P., Isaksen, I., Iversen, I., Kloster, S., Koch, D., Kirkevåg, A., Kristjansson, J. E., Krol, M., Lauer, A., Lamarque, J. F., Liu, X., Montanaro, V., Myhre, G., Penner, J., Pitari, G., Reddy, S., Seland, Ø., Stier, P., Takemura, T., and Tie, X.: Analysis and quantification of the diversities of aerosol life cycles within AeroCom, *Atmospheric Chemistry and Physics*, 6, 1777–1813, <https://doi.org/10.5194/acp-6-1777-2006>, 650 <https://acp.copernicus.org/articles/6/1777/2006/>, 2006.

- Tilmes, S., Richter, J. H., Kravitz, B., Macmartin, D. G., Mills, M. J., Simpson, I. R., Glanville, A. S., Fasullo, J. T., Phillips, A. S., Lamarque, J. F., Tribbia, J., Edwards, J., Mickelson, S., and Ghosh, S.: CESM1(WACCM) stratospheric aerosol geoengineering large ensemble project, *Bulletin of the American Meteorological Society*, pp. 2361–2371, <https://doi.org/10.1175/BAMS-D-17-0267.1>, 2018a.
- Tilmes, S., Richter, J. H., Mills, M. J., Kravitz, B., MacMartin, D. G., Garcia, R. R., Kinnison, D. E., Lamarque, J. F., Tribbia, J., and Vitt, F.:
 655 Effects of Different Stratospheric SO₂ Injection Altitudes on Stratospheric Chemistry and Dynamics, *Journal of Geophysical Research: Atmospheres*, 123, 4654–4673, <https://doi.org/10.1002/2017JD028146>, 2018b.
- Tilmes, S., MacMartin, D. G., Lenaerts, J. T. M., van Kampenhout, L., Muntjewerf, L., Xia, L., Harrison, C. S., Krumhardt, K. M., Mills, M. J., Kravitz, B., and Robock, A.: Reaching 1.5 and 2.0C global surface temperature targets using stratospheric aerosol geoengineering, *Earth System Dynamics*, 11, 579–601, <https://doi.org/10.5194/esd-11-579-2020>, <https://esd.copernicus.org/articles/11/579/2020/>, 2020.
- Timmreck, C., Mann, G. W., Aquila, V., Hommel, R., Lee, L. A., Schmidt, A., Brühl, C., Carn, S., Chin, M., Dhomse, S. S., Diehl, T., English, J. M., Mills, M. J., Neely, R., Sheng, J., Toohey, M., and Weisenstein, D.: The Interactive Stratospheric Aerosol Model Intercomparison Project (ISA-MIP): motivation and experimental design, *Geoscientific Model Development*, 11, 2581–2608, <https://doi.org/10.5194/gmd-11-2581-2018>, <https://gmd.copernicus.org/articles/11/2581/2018/>, 2018.
- Vet, R., Artz, R. S., Carou, S., Shaw, M., Ro, C.-U., Aas, W., Baker, A., Bowersox, V. C., Dentener, F., Galy-Lacaux, C., Hou, A.,
 665 Pienaar, J. J., Gillett, R., Forti, M. C., Gromov, S., Hara, H., Khodzher, T., Mahowald, N. M., Nickovic, S., Rao, P., and Reid, N. W.: A global assessment of precipitation chemistry and deposition of sulfur, nitrogen, sea salt, base cations, organic acids, acidity and pH, and phosphorus, *Atmospheric Environment*, 93, 3–100, <https://doi.org/10.1016/j.atmosenv.2013.10.060>, <https://www.sciencedirect.com/science/article/pii/S1352231013008133>, a global assessment of precipitation chemistry and deposition of sulfur, nitrogen, sea salt, base cations, organic acids, acidity and pH, and phosphorus, 2014.
- Visioni, D., Pitari, G., Di Genova, G., Tilmes, S., and Cionni, I.: Upper tropospheric ice sensitivity to sulfate geoengineering, *Atmospheric Chemistry and Physics*, <https://doi.org/10.5194/acp-18-14867-2018>, 2018a.
- Visioni, D., Pitari, G., Tuccella, P., and Curci, G.: Sulfur deposition changes under sulfate geoengineering conditions: Quasi-biennial oscillation effects on the transport and lifetime of stratospheric aerosols, *Atmospheric Chemistry and Physics*, <https://doi.org/10.5194/acp-18-2787-2018>, 2018b.
- Visioni, D., MacMartin, D. G., Kravitz, B., Lee, W., Simpson, I. R., and Richter, J. H.: Reduced poleward transport due to stratospheric heating under stratospheric aerosols geoengineering, *Geophysical Research Letters*, n/a, e2020GL089470, <https://doi.org/10.1029/2020GL089470>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2020GL089470>, e2020GL089470 2020a.
- Visioni, D., Slessarev, E., MacMartin, D., Mahowald, N. M., Goodale, C. L., and Xia, L.: What goes up must come down: impacts of deposition in a sulfate geoengineering scenario, *Environmental Research Letters*, 15, <http://iopscience.iop.org/10.1088/1748-9326/ab94eb>, 2020b.
- Visioni, D., MacMartin, D. G., and Kravitz, B.: Is Turning Down the Sun a Good Proxy for Stratospheric Sulfate Geoengineering?, *Journal of Geophysical Research: Atmospheres*, n/a, e2020JD033952, <https://doi.org/10.1029/2020JD033952>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2020JD033952>, e2020JD033952 2020JD033952, 2021.
- Wang, H., Easter, R. C., Zhang, R., Ma, P.-L., Singh, B., Zhang, K., Ganguly, D., Rasch, P. J., Burrows, S. M., Ghan, S. J., Lou, S., Qian, Y., Yang, Y., Feng, Y., Flanner, M., Leung, L. R., Liu, X., Shrivastava, M., Sun, J., Tang, Q., Xie, S., and Yoon, J.-H.: Aerosols in the E3SM Version 1: New Developments and Their Impacts on Radiative Forcing, *Journal of Advances in Modeling Earth Sys-*

- tems, 12, e2019MS001851, <https://doi.org/https://doi.org/10.1029/2019MS001851>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019MS001851>, e2019MS001851 2019MS001851, 2020.
- 690 Weisenstein, D. K., Penner, J. E., Herzog, M., and Liu, X.: Global 2-D intercomparison of sectional and modal aerosol modules, *Atmospheric Chemistry and Physics*, 7, 2339–2355, 2007.
- Yi, B., Rapp, A. D., Yang, P., Baum, B. A., and King, M. D.: A comparison of Aqua MODIS ice and liquid water cloud physical and optical properties between collection 6 and collection 5.1: Cloud radiative effects, *Journal of Geophysical Research: Atmospheres*, 122, 4550–4564, <https://doi.org/https://doi.org/10.1002/2016JD025654>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016JD025654>, 2017.
- 695