

Measurement report: Photochemical production and loss rates of formaldehyde and ozone across Europe

Clara M. Nussbaumer¹, John N. Crowley¹, Jan Schuladen¹, Jonathan Williams^{1,2}, Sascha Hafermann¹, Andreas Reiffs¹, Raoul Axinte¹, Hartwig Harder¹, Cheryl Ernest^{1,a}, Anna Novelli^{1,b}, Katrin Sala¹, Monica Martinez¹, Chinmay Mallik^{1,c}, Laura Tomsche^{1,d}, Christian Plass-Dülmer³, Birger Bohn⁴, Jos Lelieveld^{1,2}, and Horst Fischer¹

¹Max Planck Institute for Chemistry, Department of Atmospheric Chemistry, 55128 Mainz, Germany

²Climate and Atmosphere Research Center, The Cyprus Institute, Nicosia, Cyprus

³German Meteorological Service, Meteorological Observatory Hohenpeissenberg (MOHp), 83282 Hohenpeissenberg, Germany

⁴Institute of Energy and Climate Research, IEK-8: Troposphere, Forschungszentrum Jülich GmbH, 52428 Jülich, Germany

^anow at: Department of Neurology, University Medical Center of the Johannes Gutenberg University Mainz, 55131 Mainz, Germany

^bnow at: ⁴

^cnow at: Department of Atmospheric Science, Central University of Rajasthan, Rajasthan 305817, India

^dnow at: German Aerospace Center, Institute of Atmospheric Physics, 82234 Weßling-Oberpfaffenhofen, Germany

Correspondence: Clara M. Nussbaumer (clara.nussbaumer@mpic.de)

Abstract. Various atmospheric sources and sinks regulate the abundance of tropospheric formaldehyde (HCHO) which is an important trace gas impacting the HO_x (\equiv HO₂ + OH) budget and the concentration of ozone (O₃). In this study, we present the formation and destruction terms of ambient HCHO and O₃ calculated from in-situ observations of various atmospheric trace gases measured at three different sites across Europe during summer time. These include a coastal site in Cyprus in the scope of the Cyprus Photochemistry Experiment (CYPHEX) in 2014, a mountain site in Southern Germany as part of the Hohenpeissenberg Photochemistry Experiment (HOPE) in 2012 and a forested site in Finland where measurements were performed during the Hyytiälä United Measurements of Photochemistry and Particles (HUMPPA) campaign in 2010. We show that at all three sites formaldehyde production from the OH oxidation of methane (CH₄), acetaldehyde (CH₃CHO), isoprene (C₅H₈) and methanol (CH₃OH) can almost completely balance the observed loss via photolysis, OH oxidation and dry deposition. Ozone chemistry is clearly controlled by nitrogen oxides (NO_x \equiv NO + NO₂) that includes O₃ production from NO₂ photolysis and O₃ loss via the reaction with NO. Finally, we use the HCHO budget calculations to determine whether net ozone production is limited by the availability of VOCs (VOC limited regime) or NO_x (NO_x limited regime). At the mountain site in Germany O₃ production is VOC limited, whereas it is NO_x limited at the coastal site in Cyprus. The forested site in Finland is in the transition regime.

Formaldehyde (HCHO) is an important atmospheric trace gas, which provides insight into various photochemical processes taking place in the earth's atmosphere. It has both anthropogenic sources, such as industrial and vehicle emissions, and natural sources including for example biomass burning or VOC precursors, with natural sources dominating in remote locations (Luecken et al., 2018; Anderson et al., 2017; Stickler et al., 2006; Wittrock et al., 2006; Lowe and Schmidt, 1983). The majority of these HCHO sources is secondary and due to its short lifetime, atmospheric transport of HCHO from primary (direct) emissions (e.g. biomass burning or industry) to remote locations can be mostly neglected (Fortems-Cheiney et al., 2012; Vigouroux et al., 2009; Anderson et al., 2017). Loss processes of HCHO include deposition, reaction with OH and photolysis yielding mainly HO₂, CO and H₂ (Anderson et al., 2017). HCHO production paths are more diverse and include oxidation processes of almost any volatile organic compound (VOC) including acetone (CH₃COCH₃), methane (CH₄), acetaldehyde (CH₃CHO), methanol (CH₃OH), isoprene (C₅H₈), methyl hydroperoxide (CH₃OOH), ethene (C₂H₄) (these selected species are included in this study due to the availability of measurement data) and many more, the majority of which are initiated by the OH radical during the day. (Stickler et al., 2006; Wittrock et al., 2006). Net production processes of formaldehyde therefore influence the HO_x (HO_x ≡ OH + HO₂) budget which in turn controls the atmospheric oxidizing capacity (Luecken et al., 2018). This includes the regulation of the atmospheric ozone (O₃) abundance, a trace gas with adverse health effects for humans, animals and plants leading to cardiovascular and respiratory diseases and the loss of life expectancy (Nuvolone et al., 2018; Lippmann, 1989). It is therefore important to understand the processes influencing and contributing to HCHO and O₃ formation and loss processes in the earth's atmosphere (see also Figure 1 and 2 for an overview of the reactions considered in this study).

Previous studies have investigated the processes contributing to HCHO production from secondary sources. Palmer et al. (2003) identified isoprene, methane and methanol to be the main HCHO precursor over the United States of America contributing by over 80 % in the GEOS-CHEM model (Goddard Earth Observing System global 3-D model of tropospheric chemistry). Anderson et al. (2017) evaluated HCHO concentrations in the tropical western pacific and found methane and acetaldehyde to be the main precursors of HCHO based on box model simulations. Fried et al. (2011) identified methane to be the main precursor of HCHO in remote regions based on model simulations and measurements during the campaign INTEX-B (Intercontinental Transport Experiment-Phase B) in 2006. Sumner et al. (2001) investigated the HCHO budget at a forest in Pellston, Michigan (U.S.A.) based on observations in the scope of PROPHET (Program for Research on Oxidants: Photochemistry, Emission and Transport) in 1998. They identified isoprene to be the main HCHO precursor with around 80 %. Dienhart et al. (2021) investigated the relationship between OH reactivity and HCHO production rates during the shipborne campaign AQABA (Air Quality and Climate Change in the Arabian Basin) around the Arabian Peninsula in 2017 which they found to be highest in polluted areas, suggesting a high diversity of HCHO precursors. Kaiser et al. (2015) studied OH reactivities and HCHO concentrations in the Po Valley based on zeppelin measurements during the research campaign PEGASOS (Pan-European Gas-AeroSols Climate Interaction Study) in 2012 in comparison with model simulations and attributed discrepancies to possible primary HCHO emissions from agriculture.

Tropospheric ozone chemistry is dependent on the O₃ precursors NO_x (NO_x ≡ NO + NO₂) and volatile organic compounds (VOC). Depending on the ambient concentrations of NO_x and VOC, net ozone formation can either be NO_x or VOC limited. A
50 NO_x limitation is usually dominant for low NO_x concentrations in which increasing NO_x leads to an increase in O₃ formation. For high NO_x concentrations, ozone formation is usually VOC limited and an increase in ambient NO_x reduces O₃ formation through loss of HO_x as OH is converted to HNO₃ (Pusede et al., 2015; Nussbaumer and Cohen, 2020). Consequently, changes in ambient NO_x concentrations can either increase or decrease O₃ or - at a transition between both regimes - have only a weak
55 net effect on O₃ production. In urban environments, the chemical regime can be characterized using the weekend effect which describes the ozone response to decreasing NO_x emissions on weekends as reported by Pusede and Cohen (2012), Nussbaumer and Cohen (2020), Pires (2012), Wang et al. (2014) and many more (Levitt and Chock, 1976; Seguel et al., 2012; Sadanaga et al., 2011). Another measure for identifying the prevailing chemical regime is the ratio of HCHO and NO₂ which has been determined via satellite measurements in various studies (Sillman, 1995; Jin et al., 2020; Martin et al., 2004; Duncan et al.,
60 2010; Jin and Holloway, 2015). Sillman (1995) initially suggested a threshold of 0.28 for the ratio HCHO/NO_y (NO_y is here the sum of NO_x, HNO₃, peroxyacetylnitrates and alkyl nitrates), below which chemistry is VOC limited, based on model simulations for Lake Michigan and the northeast corridor (United States of America). Martin et al. (2004) investigated the ratio of the HCHO/NO₂ column based on satellite observations and found a threshold of 1. This is in agreement with findings from Duncan et al. (2010) who suggested a threshold of 1 for a VOC limited regime, a threshold of 2 for a NO_x limited regime and a transition in between both using remote sensing. Schroeder et al. (2017) present in-situ measurements of HCHO and NO₂
65 for determining the dominant regime and points out that exact thresholds are geographically variable due to locally different atmospheric composition and ambient conditions, such as VOC variety or humidity.

In this study, we evaluate the formaldehyde and ozone budget during the field experiment CYPHEX (Cyprus Photochemistry Experiment) which took place in July 2014 at a coastal site in Cyprus (Ineia) based on in-situ trace gas observations of NO, NO₂, O₃, OH, HO₂, CH₄, CH₃OH (methanol), C₅H₈ (isoprene), CH₃CHO (acetaldehyde), CH₃COCH₃ (acetone), CH₃OOH
70 (methyl hydroperoxide), C₂H₄ (ethene), CH₃SCH₃ (DMS) and HCHO (Derstroff et al., 2017; Meusel et al., 2016; Mallik et al., 2018). We compare the results with two other field campaigns in central and northern Europe which are the Hohenpeißenberg Photochemistry Experiment (HOPE 2012, Novelli et al. (2017)) at a mountain site in Germany and the Hyytiälä United Measurements of Photochemistry and Particles (HUMPPA 2010, Williams et al. (2011)) at a boreal forest site in Finland. Only few studies have evaluated the HCHO budget and mainly through model simulations. To our knowledge there is only one study by
75 Sumner et al. (2001) that has previously presented HCHO budget calculations from in-situ trace gas observations in the United States of America in 1998. We are first to present HCHO budget calculations from in-situ measurements across Europe and show that in all three locations HCHO production can be predominantly accounted for by the oxidation of methane, methanol, acetaldehyde and isoprene.

2 Observations and methods

80 2.1 HCHO chemistry calculations

Figure 1 shows an overview of the main production and loss processes for formaldehyde which we consider in this study and for which measurements were obtained during the field campaign CYPHEX (see Section 2.3.1 for further details). The relationships we derive in this study are based on boundary layer conditions and we therefore assume no relevant intrusion from higher altitudes. Acetone and methane can form methyl radicals (CH_3) through oxidation by OH or photolysis which are subsequently oxidized to methyl peroxy radicals (CH_3O_2) by molecular oxygen (O_2). Another pathway yielding CH_3O_2 is the OH oxidation of acetaldehyde which forms $\text{CH}_3\text{C}(\text{O})$ in the first step which can then be oxidized to $\text{CH}_3\text{C}(\text{O})\text{O}_2$ when O_2 is present. $\text{CH}_3\text{C}(\text{O})\text{O}_2$ yields CH_3O_2 through reaction with nitric oxide (NO) or the hydroperoxyl radical (HO_2) via CH_3CO_2 . The CH_3O_2 yield from the reaction of $\text{CH}_3\text{C}(\text{O})\text{O}_2$ with HO_2 is approximately 50 % (k_c). Other reactions pathways result in the formation of $\text{CH}_3\text{C}(\text{O})\text{OH}$ (k_a) and $\text{CH}_3\text{C}(\text{O})\text{OOH}$ (k_b) (IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation, 2019). We calculate the overall fraction $\alpha_{\text{CH}_3\text{CHO}}$ of acetaldehyde oxidation that results in CH_3O_2 formation via Equation (1).

$$\alpha_{\text{CH}_3\text{CHO}} = \frac{k_{\text{CH}_3\text{C}(\text{O})\text{O}_2+\text{NO}} \times [\text{NO}] + k_c \times [\text{HO}_2]}{k_{\text{CH}_3\text{C}(\text{O})\text{O}_2+\text{NO}} \times [\text{NO}] + k_a \times [\text{HO}_2] + k_b \times [\text{HO}_2] + k_c \times [\text{HO}_2]} \quad (1)$$

Additionally, methyl hydroperoxide (CH_3OOH) forms CH_3O_2 via OH oxidation (60 %) (or reacts directly to HCHO via photolysis or OH-initiated oxidation (40 %)) (IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation, 2007). CH_3O_2 can then either react with HO_2 to form CH_3OOH or yield HCHO through reaction with NO or OH via CH_3O (Anderson et al., 2017; Stickler et al., 2006; Lowe and Schmidt, 1983; Fittschen et al., 2014). The importance of CH_3O_2 loss via OH in remote locations has been recently shown in several studies and the reaction primarily yields HCHO (Lightfoot et al., 1992; Assaf et al., 2017, 2016; Fittschen et al., 2014; Yan et al., 2016). For simplification, we assume this yield to be 100 % which slightly increases the uncertainty of the calculation but is negligible given the small fraction of CH_3O_2 that reacts with OH (< 10 % for CYPHEX). In our study, we use the rate constant $k_{\text{CH}_3\text{O}_2+\text{OH}}$ recommended by the IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation (2017). Table S1 of the Supplement gives an overview of all rate constants used in this study, most of which were taken from IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation (2021). The fraction of CH_3O_2 that forms HCHO ($\alpha_{\text{CH}_3\text{O}_2}$) is dependent on the ambient concentrations of HO_2 , NO and OH and can be calculated via Equation (2). CH_3O_2 loss via self-reaction is negligibly small and therefore not included. The reaction with NO_2 forming $\text{CH}_3\text{O}_2\text{NO}_2$ can also be excluded due to its thermal instability in the boundary layer.

$$\alpha_{\text{CH}_3\text{O}_2} = \frac{k_{\text{CH}_3\text{O}_2+\text{NO}} \times [\text{NO}] + k_{\text{CH}_3\text{O}_2+\text{OH}} \times [\text{OH}]}{k_{\text{CH}_3\text{O}_2+\text{NO}} \times [\text{NO}] + k_{\text{CH}_3\text{O}_2+\text{OH}} \times [\text{OH}] + k_{\text{CH}_3\text{O}_2+\text{HO}_2} \times [\text{HO}_2]} \quad (2)$$

Isoprene oxidation results in the formation of HCHO through the intermediate products methyl vinyl ketone and methacrolein as described by Wolfe et al. (2016). The HCHO yield from isoprene (α_{Isoprene}) is dependent on the ambient NO concentration




Figure 1. Overview of the chemical and photolytic reactions which lead to HCHO production and loss considering the trace gases measured during the field experiment CYPHEX. Black and bold font identifies the species which contribute mainly (~80 %) to HCHO formation according to the findings in this study. For a better overview, we have omitted intermediate steps with a 100 % yield which are instead described in the main text.

and varies between around 30 % for low NO_x and 60 % for high NO_x (RO_2 react primarily with NO) (Atkinson et al., 2006; Palmer et al., 2003; Sumner et al., 2001). We calculate the HCHO yield via Equation (3) and estimate $[\text{HO}_2] \approx [\text{RO}_2]$ as suggested by Sumner et al. (2001). This assumption is justified when looking at O_3 production ($\text{P}(\text{O}_3)$) terms. $\text{P}(\text{O}_3)$ can either be calculated via the photolytic reaction of NO_2 as presented in Section 2.2 or via the reaction of HO_2 or RO_2 with NO. We equate the two terms, using the rate constant of the reaction of NO and CH_3O_2 as estimate for the reaction of NO and RO_2 , and calculate RO_2 . We show the diurnal profiles of HO_2 and calculated RO_2 in Figure S1a of the Supplement.

Conversely, we calculate $\text{P}(\text{O}_3)$ for both cases, equating RO_2 to HO_2 which show close agreement and can be seen in Figure S1b of the Supplement. This is also confirmed by findings from Crowley et al. (2018) (presented in Figure 9) based on model simulations of HO_2 and RO_2 during HUMPPA. The yield can vary between 34 % (in the absence of NO) and 57 % (when NO chemistry dominates the fate of the RO_2 formed) according to Equation (3) which was originally determined experimentally by Miyoshi et al. (1994). We discuss the effects of the threshold values in Section 3.1. $\text{C}_5\text{H}_8(\text{OH})\text{O}_2$ is the peroxy radical resulting from isoprene oxidation and has six relevant isomers which can undergo multiple reactions yielding HCHO and many other products (Wennberg et al., 2018; Schwantes et al., 2020). Additionally, the formation of HCHO from isoprene does not occur instantaneously (as for example from methane or other VOC), but is likely time-dependent. However, the consideration of this time-dependent formation as well as the detailed evaluation of the reactions paths from each peroxy radical isomer is beyond

the scope of this study and we therefore adapt the methodology presented by Sumner et al. (2001) to estimate the HCHO
 125 production from isoprene.

$$\alpha_{Isoprene} = 0.34 + 0.23 \times \left(\frac{k_{C_5H_8(OH)O_2+NO} \times [NO]}{k_{C_5H_8(OH)O_2+NO} \times [NO] + k_{C_5H_8(OH)O_2+HO_2} \times [HO_2] + k_{C_5H_8(OH)O_2+RO_2} \times [RO_2]} \right) \quad (3)$$

Methanol reacts with OH yielding HCHO via CH₂OH and CH₃O and following oxidation by O₂ (Anderson et al., 2017; Stickler et al., 2006). Ethene is a precursor to HCHO through OH oxidation or ozonolysis (Alam et al., 2011; Atkinson et al., 2006). A potential source of HCHO in marine environments is dimethyl sulfide (DMS) via OH oxidation (Ayers et al., 1997; 130 Urbanski et al., 1997). Terpenes such as limonene or α -/ β -pinene emitted from plants can additionally be HCHO sources as described by Lee et al. (2006).

Reactions (R1) - (R3) present the chemical loss processes of HCHO through OH oxidation and two different photolysis pathways. In addition, HCHO dry deposition - the uptake of HCHO by the earth's surface - plays a role in HCHO loss, particularly during the night (Anderson et al., 2017; Possanzini et al., 2002; Sumner et al., 2001; Wesely and Hicks, 2000). 135 While HCHO loss can also occur via wet deposition as for example described by Seyfioglu et al. (2006) or via liquid-phase reactions in cloud droplets as shown by Franco et al. (2021), this study investigates summertime campaigns without significant precipitation.



We will show in the scope of this work that reactions of methane, acetaldehyde, methanol and isoprene with OH almost completely account for HCHO production in the environments considered in this paper across Europe. We have highlighted these pathways in Figure 1 in black and bold font. Equations (4) and (5) show the calculation of the basic production $P(HCHO)_{basic}$ (compared to the reactions shown in Figure 1) and the loss $L(HCHO)$ terms. k values represent the rate coefficients, $j(HCHO)$ is the summed photolysis frequency for reactions (R2) and (R3), v_d describes the dry deposition velocity in cm s⁻¹ and BLH is the boundary layer height in cm.

$$\begin{aligned} P(HCHO)_{basic} &= P_{CH_4+OH} + P_{CH_3CHO+OH} + P_{CH_3OH+OH} + P_{C_5H_8+OH} \\ &= [OH] \times ([CH_4] \times k_{CH_4+OH} \times \alpha_{CH_3O_2} + [CH_3CHO] \times k_{CH_3CHO+OH} \times \alpha_{CH_3O_2} \times \alpha_{CH_3CHO} \\ &\quad + [CH_3OH] \times k_{CH_3OH+OH} + [C_5H_8] \times k_{C_5H_8+OH} \times \alpha_{Isoprene}) \end{aligned} \quad (4)$$




Figure 2. Overview of the chemical and photolytic reactions which lead to O₃ production and loss.

$$\begin{aligned}
 L(HCHO) &= L_{HCHO+OH} + L_{HCHO+h\nu} + L_{deposition} \\
 &= [HCHO] \times \left([OH] \times k_{HCHO+OH} + j(HCHO) + \frac{v_d(HCHO)}{BLH} \right) \quad (5)
 \end{aligned}$$

Changes in the HCHO concentration are represented by Equation (6) which includes production and loss from Equations (4) -
 150 (5), a transport term T(HCHO) such as advection or entrainment (can be positive or negative) and a term for primary emissions (Fischer et al., 2019).

$$\frac{d[HCHO]}{dt} = P(HCHO)_{basic} - L(HCHO) + T(HCHO) + P_{emission} \quad (6)$$

2.2 O₃ chemistry calculations

Figure 2 presents the main processes contributing to O₃ formation and loss. The only significant chemical source of tropo-
 155 spheric O₃ is the photolysis of nitrogen dioxide (NO₂) which is converted to NO and O(³P) under the influence of sunlight. The reaction of O(³P) with molecular oxygen (O₂) subsequently yields O₃ (Jacob, 1999). NO₂ in turn is generated from the oxidation of NO by ozone or peroxy radicals (HO₂, RO₂) (Pusede et al., 2015).

Ozone loss processes include the reaction with NO forming NO₂, conversion with OH or HO₂ to HO₂ and OH, respectively,
 160 deposition processes and photolysis. O₃ photolysis yields O(¹D) which is deactivated to O(³P) (and then O₃) via collision with nitrogen (N₂) and oxygen (O₂). O₃ loss occurs when O(¹D) reacts with water to form OH. The fraction of this reaction is presented by α_{O¹D} as shown in Equation (7) (Bozem et al., 2017). The reaction of O₃ and NO₂ forming NO₃ could potentially

yield a net loss of O_3 when being photolyzed back to NO_x . However, only around 10 % of the NO_3 photolysis leads to NO formation (which results in a O_3 net loss) (Stockwell and Calvert, 1983). Additionally, the reaction is likely negligible during the day as O_3 reacts much more rapidly with NO.

$$165 \quad \alpha_{O^1D} = \frac{k_{O^1D+H_2O} \times [H_2O]}{k_{O^1D+N_2} \times [N_2] + k_{O^1D+O_2} \times [O_2] + k_{O^1D+H_2O} \times [H_2O]} \quad (7)$$

Equations (8) and (9) present the calculations for O_3 production and loss.

$$P(O_3) = P_{NO_2+h\nu} = [NO_2] \times j(NO_2) \quad (8)$$

$$\begin{aligned} L(O_3) &= L_{O_3+NO} + L_{O_3+OH} + L_{O_3+HO_2} + L_{O_3+h\nu} + L_{deposition} \\ &= [O_3] \times \left([NO] \times k_{O_3+NO} + [OH] \times k_{O_3+OH} + [HO_2] \times k_{O_3+HO_2} + j(O^1D) \times \alpha_{O^1D} + \frac{v_d(O_3)}{BLH} \right) \end{aligned} \quad (9)$$

As for HCHO, net changes in O_3 are represented by production, loss and a transport term (either positive or negative) as shown
170 in Equation (10).

$$\frac{d[O_3]}{dt} = P(O_3) - L(O_3) + T(O_3) \quad (10)$$

2.3 Field experiments

We have analyzed trace gases and further measurement parameters in regard to the HCHO and O_3 budget at three different measurement sites across Europe which are located in Cyprus (CYPHEX campaign 2014), Southern Germany (HOPE cam-
175 paign 2012) and Finland (HUMPPA campaign 2010). Their geographic locations are shown in Figure 3. We provide details on each campaign in the following sections. Please note that all times are in UTC (coordinated universal time). The time difference between 12:00 UTC and mean local noon are UTC + 128 min for Cyprus, UTC + 44 min for Germany and UTC + 96 min for Finland (Fischer et al., 2019).

2.3.1 CYPHEX campaign 2014

180 The Cyprus Photochemistry Experiment (CYPHEX) took place in Ineia, Cyprus in July and August 2014. The measurement site was situated on a hilltop 650 m above sea level (34.96 °N, 32.38 °E) in a remote location with low population in the surrounding areas. The distance to the coastline of the Mediterranean Sea was approximately 10 km in the North and in the West. A detailed description of the measurement site can be found in Derstroff et al. (2017), Meusel et al. (2016) and Mallik et al. (2018). In this study, we consider the campaign days for which the trace gas measurements were available simultaneously
185 which is the time period 22 - 31 July 2014. NO and NO_2 were measured via photolytic chemiluminescence (detector from ECO




Figure 3. Geographic locations of the measurement sites included in this analysis. CYPHEX: 34.96 °N, 32.38 °E, 650 m a.s.l. (above sea level), UTC + 128 min (mean local time); HOPE: 47.80 °N, 11.01 °E, 980 m a.s.l., UTC + 44 min; HUMPPA: 61.85°N, 24.28 °E, 181 m a.s.l., UTC + 96 min.

Physics CLD 790 SR, Dürnten Switzerland, photolytic converter from Droplet Measurement Technologies, Boulder, USA) with a total uncertainty of 20 and 30 % and a detection limit of 5 and 20 pptv, respectively (Hosaynali Beygi et al., 2011; Tadic et al., 2020). O₃ was measured via UV photometry (O₃-Analyzer model 49, Thermo Environment Instruments, USA) with a detection limit of 2 ppbv and a total uncertainty of 5 %. OH and HO₂ were measured via laser-induced fluorescence spectroscopy with the custom-built HORUS (HydrOxyl Radical measurement Unit based on fluorescence Spectroscopy) instrument (accuracy 28.5 and 36 % & detection limit 1×10^6 molec cm⁻³ and 0.8 pptv, respectively) (Marno et al., 2020; Novelli et al., 2014). HCHO was measured via the method of Hantzsch with a commercial instrument (Aero-Laser model AL 4021, Garmisch Partenkirchen, Germany) with a detection limit of 38 pptv and a total uncertainty of 16 % (Kormann et al., 2003). C₂H₄ and CH₄ were determined via gas chromatography flame ionization detection (GC 5000 VOC, AMA instruments, Ulm, Germany). CH₄ measurements had a detection limit of 20 ppbv and a total uncertainty of 2 % and C₂H₄ measurements had a detection limit of 1 - 8 pptv and a total uncertainty of 10 % (Sobanski et al., 2016; Mallik et al., 2018). Isoprene was measured via gas chromatography mass spectrometry (MSD 5973, Agilent Technologies GmbH, Böblingen, Deutschland) with a detection limit of 1 pptv and a total uncertainty of 14.5 % (Derstroff et al., 2017). Oxygenated VOC (OVOC) were measured via proton-transfer-reaction time-of-flight mass spectrometry (PTR-TOF-MS) (Ionicon Analytik GmbH, Innsbruck, Austria) (CH₃OH: 242 pptv limit of detection (LOD), 37 % total uncertainty (TU); CH₃COCH₃: 97 ppt_v, LOD, 10 % TU; CH₃CHO: 85 pptv LOD, 22 % TU; DMS: 18 pptv LOD, 12 % TU; the TU is 4 - 7 % higher for a relative humidity below 25 %) (Veres et al., 2013; Graus et al., 2010). CH₃OOH was measured via high-performance liquid chromatography with a detection limit of 25 pptv and a total

uncertainty of 9%. Photolysis frequencies were determined with a single monochromator spectral radiometer (Meterologic Consult GmbH, Königstein, Germany) with a total uncertainty of around 10%. The photolysis frequencies for acetaldehyde $j(\text{CH}_3\text{CHO})$ and formaldehyde $j(\text{HCHO})$ were determined via parameterizations based on $j(\text{NO}_2)$ and $j(\text{O}^1\text{D})$ according to Equation (S1) with the coefficients presented in Table S2 of the Supplement. The latter were derived from least-squares fits to photolysis frequencies from a large set of spectroradiometer measurements at Jülich, Germany (Bohn et al., 2008) under all weather conditions and were originally derived for the HUMPPA campaign. In this work more recent quantum yields for the HCHO photolysis as recommended by IUPAC (2013) were used with an estimated uncertainty of 20%. An example for the performance of the parameterization is shown in Figure S2 of the Supplement. An overview of all measured trace gases including the measurement uncertainty and the time resolution of the data used in this study can be found in Table S3 of the Supplement. For the point-by-point calculations, the data were interpolated to the OH time stamp with a 4-minutes time resolution. All stated detection limits refer to the time resolution shown in Table S3 of the Supplement.

2.3.2 HOPE campaign 2012

The Hohenpeißenberg Photochemistry Experiment (HOPE) took place from June to September 2012 in Hohenpeißenberg, Germany at the Global Atmospheric Watch Meteorological Observatory (47.80°N, 11.01°E). The measurement location was situated on a hilltop 980 m above sea level in a remote and vegetated area. More details on the campaign and the site location can be found in Novelli et al. (2017). O_3 , HCHO, OH and HO_2 were measured with the same methods as described for the CYPHEX campaign in Section 2.3.1. NO and NO_2 were measured via (photolysis) chemiluminescence by the German Weather Service with an estimated uncertainty of 10%. CH_4 was measured via GC-FID. Isoprene and OVOC were determined via a custom built GC (Agilent Technologies) FID/MS system (Novelli et al., 2017; Werner et al., 2013). $j(\text{NO}_2)$ and $j(\text{O}^1\text{D})$ were measured with filter radiometers (Meterologic Consult GmbH, Königstein, Germany) with an uncertainty of 10% (Bohn et al., 2008). $j(\text{HCHO})$ was determined via parameterization (Section 2.3.1). BLH measurements were not performed and instead adopted from Fischer et al. (2019) which are 1500 m for daytime ($j(\text{NO}_2) > 10^{-3} \text{ s}^{-1}$) and 200 m for nighttime ($j(\text{NO}_2) < 10^{-3} \text{ s}^{-1}$). These values were derived from BLH measurements at other locations during summertime including the CYPHEX and the HUMPPA measurement site, and a site in central Germany situated at a comparable altitude. We assume that this estimate increases the uncertainty from 20% (BLH measurements) to 30%. We determined the deposition velocity v_d from the HCHO nighttime loss as previously performed by Fischer et al. (2019) for H_2O_2 . The loss rate coefficient k_d was determined from the [HCHO] decrease from 21:00 - 01:30 UTC divided by the average HCHO concentration $[\text{HCHO}]_{av}$ in this time interval. Multiplication with the nighttime boundary layer height (BLH) (200 m) then yielded the deposition velocity according to Equation (11). The HCHO loss via deposition during nighttime is independent of the BLH (see Equation (5) and (11)).

$$v_d = \frac{k_{d,night} \times BLH_{night}}{x} = \frac{\frac{d[\text{HCHO}]}{dt}}{[\text{HCHO}]_{av} \times x} \times BLH_{night} \quad (11)$$

Please note that v_d derived this way represents a lower limit to the nighttime loss rate as HCHO could be formed from NO_3 and O_3 chemistry (for example from ozone and isoprene) at night (Crowley et al., 2018). The factor x considers the inconsistent




Figure 4. The determination of the deposition velocity is based on the HCHO nighttime loss - here exemplarily shown for one night during the campaign HOPE (2012) in Hohenpeißenberg, Southern Germany.

235 mixing of the boundary layer at night for which x was 2, assuming a linear gradient between the top and the bottom of the boundary layer. During the day, x equaled 1 (Shepson et al., 1992; Fischer et al., 2019). As we determined the loss rate from the nighttime decrease in HCHO, the daytime deposition velocity was twice the nighttime deposition velocity according to Equation (11) which gives $v_d(\text{day}) = 0.94 \text{ cm s}^{-1}$ and $v_d(\text{night}) = 0.47 \text{ cm s}^{-1}$. Literature values for daytime deposition velocities range between 0.36 and 1.5 cm s^{-1} and for nighttime between 0.18 and 0.65 cm s^{-1} (Sumner et al., 2001; Stickler et al.,

240 2007; DiGangi et al., 2011; Ayers et al., 1997). We therefore consider our calculation to yield reasonable estimates.

We have performed this calculation for nine nights. Figure 4 exemplarily shows the nighttime loss of HCHO for one night during the HOPE campaign. Red data points represent the HCHO mixing ratios, black color highlights the data points which we included in our analysis (nighttime HCHO loss) and the cyan line is the linear fit of these data points. An overview of the nights thus analyzed can be found in Figure S3b of the Supplement. The uncertainty of each calculation results from the single uncertainties of $\frac{d[HCHO]}{dt}$, $[HCHO]_{av}$ and the nighttime BLH. The uncertainty of $\frac{d[HCHO]}{dt}$ is composed of the HCHO measurement uncertainty (16 %) and the uncertainty of the fit (30 % upper limit) with the latter dominating. The uncertainty of $[HCHO]_{av}$ is based on the HCHO measurement uncertainty and the HCHO averaging (20 % upper limit). Again, the uncertainty of the averaging prevails over the measurement uncertainty. The uncertainty of the BLH is 30 %. Gaussian error propagation gives an overall uncertainty of $\sqrt{30\%^2 + 30\%^2 + 20\%^2} = 47\%$ from the calculation. Besides the uncertainty resulting from the calculation,

250 an additional uncertainty arises from the atmospheric variability which describes ambient, instrumentally independent variations of the considered trace gases and parameters caused by for example atmospheric turbulence. The uncertainty from the atmospheric variability represented by the 1σ standard deviation of the mean over the considered nine nights is 54 % which

exceeds the uncertainty from the calculation. We therefore estimate the uncertainty of v_d to be 54 %. Please note that in our uncertainty analysis we consider the arising statistical errors but not the systematic errors which are not quantifiable but could potentially increase the overall uncertainty. An overview of all uncertainties and the time resolution of the measured trace gases can be found in Table S3 of the Supplement. Absolute H₂O concentrations were estimated from the measured relative humidity via the Magnus formula over water. Please note that some trace gases were not available simultaneously or only for a short overlap. We therefore present the averaged diurnal profiles.

2.3.3 HUMPPA campaign 2010

The Hyytiälä United Measurements of Photochemistry and Particles (HUMPPA) campaign took place in July and August 2010 at SMEAR II (Station for Measuring Ecosystem-Atmosphere Relation) in Hyytiälä, Finland (61.85°N, 24.28°E, 181 m above sea level). The site is located in a remote area in a boreal forest. A detailed description of the campaign can be found in Williams et al. (2011). The measurement methods for NO_x, O₃, HCHO and CH₄ were the same as presented for the CYPHEX campaign. CH₃OH was measured via ColdTrap PTR-MS with a detection limit of around 50 pptv (integration time of 5 mins). Acetaldehyde was measured via PTR-MS with a detection limit of 50 pptv (integration time of 6 mins) (Williams et al., 2011). Isoprene was measured via GC (GC 6890A, Agilent Technologies) coupled to a Mass Selective Detector (MSD 5973 inert, Agilent Technologies) with an uncertainty of around 15 %. Highly constrained box-model simulations which were in fair agreement with experimental data where available for OH and HO₂ as described by Crowley et al. (2018) with an uncertainty in the order of 30 - 40 %. $j(\text{NO}_2)$ and $j(\text{O}^1\text{D})$ were measured with filter radiometers (Meteorologie Consult GmbH, Königstein, Germany) with an uncertainty of around 10 %; $j(\text{HCHO})$ was determined via parameterization (in analogy to CYPHEX and HOPE). H₂O was measured with an infrared light absorption analyser (URAS 4 H₂O, Hartmann & Braun, Frankfurt am Main, Germany). The boundary layer height was measured by radio soundings as presented by Ouwersloot et al. (2012) ranging from 200 m during nighttime (here $j(\text{NO}_2) < 10^{-3} \text{ s}^{-1}$) to 1500 m during daytime (here $j(\text{NO}_2) < 10^{-3} \text{ s}^{-1}$) (Fischer et al., 2019). The deposition velocity was determined in analogy to the HOPE campaign based on the nighttime HCHO loss on the basis of 14 nights and was 0.85 cm s⁻¹ during the day and 0.43 cm s⁻¹ during the night. The uncertainty of the calculation results from the single uncertainties of $\frac{d[\text{HCHO}]}{dt}$ (16 %), $[\text{HCHO}]_{av}$ (33 %) and the nighttime BLH (20 %). Gaussian error propagation gives an overall uncertainty of 42 % from the calculation. The atmospheric variability equals 54 %. We therefore estimate the total uncertainty of the deposition velocity to be 54 %. An overview of all considered nights and the according HCHO loss is presented in Figure S3c of the Supplement. The deposition velocity of ozone was adopted from Rannik et al. (2012) for the time period W (week) 25 - 34 of the year and a relative humidity below 70 % which is 0.491 cm s⁻¹ for daytime and 0.069 cm s⁻¹ for nighttime. A modeling study by Emmerichs et al. (2021) presents average values in the same order of magnitude (~0.2 - 0.3 cm s⁻¹ for July and August). Again, uncertainties and time resolution are shown in Table S3 of the Supplement. Similar to the HOPE campaign, some trace gases were not available simultaneously and we therefore present the averaged diurnal profiles.

3.1 Net HCHO production during CYPHEX 2014

HCHO concentrations during the CYPHEX campaign ranged between 0.3 and 1.9 ppb_v (1.1 ± 0.4 ppbv on average) with a maximum in the diel cycle during morning hours (04:00 UTC) and a minimum in the afternoon (15:00 UTC). Please note the mean local time difference stated in Section 2.3 and Figure 3. The temporal development of HCHO concentrations during the campaign and the respective $j(\text{NO}_2)$ values (to illustrate the daily cycle) are shown in Figure S3a of the Supplement. Figure S4a of the Supplement presents the diurnal average of HCHO including its rate of change $\frac{d[\text{HCHO}]}{dt}$. The uncertainty is dominated by the atmospheric variability which is on average 27 % for daytime HCHO.

Figure 5 shows the time series of (a) the HCHO production terms, (b) the HCHO loss terms and (c) the net HCHO production during the research campaign CYPHEX in Cyprus in July 2014. We calculated the HCHO production terms for all measured gas-phase precursors shown in Figure 1. The individual production terms are shown in Figure S5 of the Supplement. The total measurement uncertainties were determined via gaussian error propagation and were 28 % for HCHO production (31 % for only considering the OH oxidation of methane, methanol, isoprene and acetaldehyde) and 26 % for HCHO loss. Table S4 of the Supplement provides an overview of all calculated uncertainties. We present two example step-by-step calculations via gaussian error propagation in Equations (S2)-(S8) of the Supplement. All other calculations were made accordingly. We have neglected the uncertainties for the HCHO yield from isoprene, $\alpha_{\text{CH}_3\text{O}_2}$ and $\alpha_{\text{CH}_3\text{CHO}}$ and instead present a sensitivity study for these parameters later in this section. Dark green colors show the overall HCHO production as sum of all single production terms. All production terms show a diurnal cycle which follows the course of the photolysis frequency $j(\text{NO}_2)$ which we show in Figure 5d. Photolysis reactions do not take place after sunset and oxidation reactions are dependent on the abundance of OH radicals which is low during nighttime. Therefore, HCHO production approaches zero during nighttime. Due to data gaps a full diurnal profile was only available for July 23, July 28 and July 30 where the overall HCHO production reached maxima between 0.6 and 0.8 ppbv h⁻¹. The reactions of methane, methanol, isoprene and acetaldehyde with OH dominated the HCHO production processes. This is additionally illustrated in Figure 6 which presents the daily average share (including day- and nighttime values) of each production term based on a balance to the overall loss rate. HCHO production was therefore dominated by the reaction of methane and OH (almost a third), followed by the oxidation of acetaldehyde contributing around 15 % and the OH oxidations of isoprene and methanol, both contributing around 14 %. These four species together represented 75 % of the overall HCHO production required to balance the sinks. The production through OH oxidation of methyl hydroperoxide and dimethyl sulfide and through the reaction of isoprene and ozone contributed by around 1 - 2 % each. The remaining species each yielded less than 1 % of the overall HCHO. Less than 20 % was unaccounted for ("rest" in Figure 6). This part also includes HCHO production from terpenes via oxidation through OH or O₃. The yields from these reactions vary greatly in the literature. Considering the yields from OH oxidation suggested by Lee et al. (2006) from laboratory investigations, limonene, β - and α -pinene would account for 3, 2 and 1 % of the overall HCHO production, respectively. The isoprene yield is limited to a value between 34 and 57 %. The lower limit would give a HCHO production from isoprene of 11 % and the upper limit would yield 19 %. The value for $\alpha_{\text{CH}_3\text{O}_2}$ can theoretically be between 0 and 100 % but is likely situated at the upper end due to the




Figure 5. Temporal development of (a) HCHO production terms, (b) HCHO loss terms and (c) net HCHO from July 22 to July 31, 2014 during the research campaign CYPHEX in Cyprus. The NO₂ photolysis frequency $j(\text{NO}_2)$ is shown in (d) as illustration of the diurnal cycle. All times are in UTC.

availability of NO. As an example, a 20 % decrease in $\alpha_{\text{CH}_3\text{O}_2}$ would give a HCHO production from CH₃O₂ of 38 % (compare to 47 % for the calculated $\alpha_{\text{CH}_3\text{O}_2}$). A 20 % increase would yield 57 % on average and decrease the "rest" to less than 10 %. For $\alpha_{\text{CH}_3\text{CHO}}$, a 20 % decrease and increase would give a HCHO yield from acetaldehyde of 12 % and 18 %, respectively,




Figure 6. Chemical production terms of HCHO during CYPHEX including daily averages of all data.

assuming a constant $\alpha_{CH_3O_2}$. Please note that the uncertainty of the absolute values used to create this pie chart is dominated by the atmospheric and diurnal variability of the single terms and is in the order of 100 %.

HCHO loss was determined from the reaction with OH and photolysis. Red colors in Figure 5b show the overall calculated HCHO loss. Similarly to HCHO production, HCHO loss via OH and photolysis only plays a role during daytime. At night, dry deposition is the only HCHO loss mechanism and the deposition velocity v_d can be determined from the nighttime decrease of ambient HCHO concentrations as shown by Sumner et al. (2001). However, we often observed increasing HCHO concentrations during nighttime, particularly before sunrise. The reason for this nighttime HCHO increase is not yet fully understood. The role of deposition processes at elevated altitudes is not yet fully understood but is likely influenced by flows along mountain slopes as well as horizontal advection on hilltops. For hilltops, incoming air through advection likely originated from areas without deposition (too high above the ground) (Derstroff et al., 2017). Please note that the effect of deposition processes could also be counteracted by a nighttime HCHO source, such as terpene oxidation by ozone or advection. The determined value for the deposition can therefore be seen as a lower estimate Crowley et al. (2018). As we do not observe net loss of HCHO at night during the CYPHEX campaign, we estimate dry deposition to be negligible. Figure 5c shows the overall HCHO production and loss rates as well as the difference between both, which we refer to as net HCHO production, in black color. Production and loss showed a good balance with values of ± 0.2 ppbv h⁻¹. On most days, HCHO loss prevailed over HCHO production based on measured precursors.

In order to better account for diurnal changes we investigated the daily cycle of HCHO production and loss rates based on hourly averages over all measurement days. Figure 7 shows the diurnal profiles of (a) the main HCHO production terms as determined above, (b) the HCHO loss terms and (c) net HCHO production. The solid lines represent the hourly averages of the point-by-point calculation of the production and loss terms. The uncertainty is composed of the measurement uncertainty and the atmospheric variability, the latter demonstrated by the 1 σ error shades. For HCHO production, the measurement uncertainty was 31 % and the daytime atmospheric variability was 40 % (61 % for day and night). For HCHO loss, the uncertainties were 26 % and 34 % (57 %), respectively. In both cases, the overall uncertainty was dominated by the atmospheric variability. The dashed lines show the hourly production and loss terms calculated from the hourly averaged trace gas concentrations.




Figure 7. Diurnal profiles of (a) HCHO production $P(\text{HCHO})$, (b) HCHO loss $L(\text{HCHO})$ and (c) HCHO net terms during the CYPHEX campaign 2014. Solid lines show the hourly averaged HCHO production and loss terms based on the point-by-point calculation of simultaneous parameter measurements. The uncertainty from the atmospheric variability is represented by the 1σ error shades. For the dashed lines, all measurement parameters were averaged first, followed by the calculation of hourly HCHO production and loss terms. The uncertainty is based on the atmospheric variability of all parameters and exemplarily shown by the error bar for one point for each production and loss term. The gray dashed line represents the diurnal solar elevation angle at the measurement site as determined by help of the NOAA Solar Calculator (2021).

The uncertainty from the atmospheric variability was 22 % (34 %) for HCHO production and 25 % (52 %) for HCHO loss, exemplarily indicated by an error bar for one point for each production and loss term in Figure 7. Both methods have advantages and disadvantages. The point-by-point calculation allows for a simultaneous consideration of all measurement parameters. Potential atmospheric changes or incidents such as winds, precipitation or rare primary emissions events are reflected by all parameters as they are monitored at the same time. On the other hand, this method reduces the amount of data points used as only one single missing species prevents the calculation of the overall term. In order to increase the number of results we have interpolated the values for the isoprene yield (data coverage 81 %), $\alpha_{\text{CH}_3\text{CHO}}$ and $\alpha_{\text{CH}_3\text{O}_2}$ (data coverage 64 %). Calculating production and loss terms from the hourly averaged trace gas concentrations allows for the inclusion of all measured parameters but could potentially increase the uncertainty of the estimate depending on the duration and overlap of the single measurements. However in this case, the data were limited to one week in July with mainly simultaneous measurement of all parameters and we therefore do not expect large uncertainties. The similarity of the results as shown in Figure 7 indicates that both methods provide reasonable results.

For HCHO production, daily maxima of around 0.4 - 0.45 ppbv h^{-1} were reached between 9:00 and 10:00 UTC which was coincident with the maximum of the photolysis frequency $j(\text{NO}_2)$. HCHO production from methane dominated the overall production term with a maximum of close to 0.2 ppbv h^{-1} , followed by acetaldehyde, isoprene and methanol. HCHO loss peaked around 9:00 UTC with a value of 0.5 ppbv h^{-1} with around 55 % contribution from photolysis and 45 % from OH oxidation. Figure 7c shows that the calculated production term for HCHO can almost completely balance the loss which, assuming that the major loss processes are well constrained, leads to the conclusion that HCHO production can be approximated




Figure 8. Diurnal profiles of (a) ozone production $P(\text{O}_3)$, (b) ozone loss $L(\text{O}_3)$ and (c) ozone net terms during the CYPHEX campaign 2014. Solid lines show the hourly averaged O_3 production and loss terms based on the point-by-point calculation of simultaneous parameter measurements. The uncertainty from the atmospheric variability is represented by the 1σ error shades. For the dashed lines, all measurement parameters were averaged first, followed by the calculation of hourly O_3 production and loss terms. The uncertainty is mostly dominated by the atmospheric variability of all parameters and exemplarily shown by the error bar for one point for each production and loss term. Please find the temporal development of production and loss terms throughout the campaign in Figure S8 of the Supplement.

by OH oxidation of methane, acetaldehyde, isoprene and methanol. This is in line with the HCHO rate of change presented in Figure S4a of the Supplement which fluctuated evenly around zero. Therefore, transport processes and primary emissions during CYPHEX can likely be excluded.

3.2 Net O_3 production during CYPHEX 2014

Ozone varied between 46 and 104 ppbv during the CYPHEX campaign (70 ± 13 ppbv on average) with a diel maximum at 04:00 UTC and a minimum at 15:00 UTC. The time series of O_3 concentrations are presented in Figure S6a of the Supplement. The diel mean including the rate of change $d\text{O}_3/dt$ can be found in Figure S7a of the Supplement. The uncertainty is dominated by the atmospheric variability which is on average 16% (1σ) for O_3 .

Figure 8 shows the diurnal cycle of the production and loss rates for ozone, analogous to Figure 7. Solid lines show the hourly averaged point-by-point calculations of the O_3 production and loss terms. The 1σ error shades present the atmospheric variability which was 59% for O_3 production and 37% for O_3 daytime loss (44% for day and night). The measurement uncertainties were 32% and 16%, respectively, and the overall uncertainty was therefore dominated by the atmospheric variability. Dashed lines show the O_3 production and loss terms based on the hourly averaged measurement parameters. The uncertainty resulting from the averaging of the individual parameters is exemplarily shown by the error bar for one point and is similar to the atmospheric variability obtained by the point-by-point method (57% for $P(\text{O}_3)$ and 35% (49%) for $L(\text{O}_3)$). Figure 8a shows O_3 production represented by the photolysis of NO_2 . Figure 8b shows the O_3 loss terms. O_3 loss through reaction with NO was dominant, followed by photolysis. The loss via OH and HO_2 was negligibly small. O_3 net production during the CYPHEX campaign was therefore clearly dominated by nitrogen oxides chemistry. Figure 8c shows net O_3 production.

O₃ production and loss were similar throughout the day with peak values of 4 - 7 ppbv h⁻¹ between 5:00 and 10:00 UTC. Net O₃ production was between -1 and 1 ppbv h⁻¹. Please note that we have excluded the NO₂ data on July 24 between 13:15 and 16:15 UTC due to a singular high concentration event. The large spike in NO₂ (O₃ titration of NO) is likely the result of sampling air impacted by the exhaust from a diesel generator which provided on site power and was located about 200 m from the containers housing the instruments. We show O₃ net production including all NO₂ data points in Figure S9a of the Supplement. O₃ production is directly proportional to the NO₂ concentration according to Equation (8) explaining the large afternoon production peak. We show the diel profile of NO₂ concentrations with and without the afternoon peak in Figure S9b of the Supplement. Production and loss terms for O₃ were balanced and the O₃ rate of change fluctuated evenly around zero (Figure S7a of the Supplement) suggesting that the diel variability was likely not impacted by transport processes.

3.3 Comparison with HOPE 2012 and HUMPPA 2010

3.3.1 Net HCHO production

We have shown in Sections 3.1 and 3.2 that the hourly averaging of measurement parameters and subsequent calculation of production and loss terms yielded reliable results during the CYPHEX campaign. For HUMPPA and HOPE, we have pursued this approach regarding the calculations of HCHO and O₃ production and loss terms, as the simultaneous data availability for all measurement parameters was too low for a point-by-point analysis.

HCHO mixing ratios during the HOPE campaign varied between 0.1 and 3.2 ppbv (1.1 ± 0.5 ppbv on average) with a maximum in the diel cycle at 15:00 UTC and a minimum at 06:00 UTC. The average mixing ratio was similar to the average mixing ratio during the CYPHEX campaign, but the variability was higher which is likely due to the longer time period of available data which was about four weeks for HOPE (compared to a bit more than one week for CYPHEX) including e.g. larger temperature variations. HCHO mixing ratios in Hyytiälä during the HUMPPA campaign ranged between 0.03 and 5.7 ppbv, with an average of 0.4 ± 0.5 ppbv. The variability is high because of biomass burning events in Russia which were detected at the site in Finland on July 26 - 30 and August 9 as described by Williams et al. (2011). The large HCHO peaks detected on these days can also be seen in Figure S3c of the Supplement. Figure S4b and S4c of the Supplement present the diurnal cycles for HCHO including the rate of change $\frac{d[HCHO]}{dt}$ during HOPE and HUMPPA. The uncertainty is dominated by the atmospheric variability represented by the 1 σ standard deviation which is on average 44 % for HCHO for HOPE and 106 % for HUMPPA.

Figure 9 shows the HCHO production and loss rates for the two campaigns. For all cases, the uncertainty was dominated by the atmospheric variability (1 σ) which was 42 % for daytime HCHO production and 40 % for daytime HCHO loss in Hohenpeißenberg and 38 % and 77 % in Hyytiälä, respectively. The measurement uncertainty was around 30 to 40 %. We present the atmospheric variability by the error bars in Figure 9. For better clarity, we only show one error bar for each term. An overview of all calculated uncertainties can be found in Table S4 of the Supplement. HCHO production during HOPE is shown in Figure 9a. The maximum production rate of 0.48 ppbv h⁻¹ was reached between 11:00 and 12:00 UTC and comparison to j(NO₂) shows good agreement with local noon. The production of HCHO was dominated by the oxidation of acetaldehyde which contributed to peak production by around 50 %, followed by methane, isoprene and methanol. We show a pie chart




Figure 9. Diurnal profiles of HCHO production $P(\text{HCHO})$ and loss $L(\text{HCHO})$ terms during the HOPE campaign 2012 (top row: (a) $P(\text{HCHO})$, (b) $L(\text{HCHO})$ and (c) net HCHO) and the HUMPPA campaign 2010 (bottom row: (d) $P(\text{HCHO})$, (e) $L(\text{HCHO})$ and (f) net(HCHO)).

415 representing the contribution of the single HCHO production terms during HOPE in Figure S10a of the Supplement. HCHO
 loss is shown in Figure 9b. The maximum loss rate of HCHO was 0.46 ppbv h^{-1} . During the day, HCHO loss was dominated by
 photolysis and oxidation while at nighttime, deposition was the main loss path for formaldehyde in Hohenpeißenberg. Figure 9c
 shows HCHO net production during HOPE. The calculated production of HCHO slightly prevailed over its loss. At 11:00 UTC,
 HCHO loss was $>95\%$ of HCHO production. Overall absolute loss and production terms were very similar compared to the
 420 results obtained for the site in Cyprus. The main difference is the composition of the HCHO production which was dominated
 by the oxidation of acetaldehyde in Hohenpeißenberg and by the oxidation of methane in Cyprus. HCHO production and
 loss during the HUMPPA campaign are shown in Figure 9d - f. HCHO production reached a peak value of 0.15 ppbv h^{-1} at
 12:00 UTC. Methane and acetaldehyde contributed to the overall production by approximately equal parts followed by isoprene
 and methanol. Figure S10b shows the share of the individual HCHO production terms during HUMPPA. Axinte (2016) showed
 425 that the contribution from terpene oxidation to HCHO production was small (1 - 2 % each) which is in line with our findings
 for the CYPHEX campaign. HCHO loss was dominated by photolysis during the day and by dry deposition at night. Figure
 9f shows that HCHO production and loss were in good agreement throughout the day ($\sim 90\%$) apart for the morning hours

06:00 - 07:00 UTC (09:00 - 10:00 UTC) when HCHO production prevailed over its loss by around 2 to 3 times indicating a missing "loss" term, most likely due to dilution with HCHO-poor air during the rise of the planetary boundary layer in the early morning hours (accompanied by a peak in the HCHO rate of change as shown in Figure S4b of the Supplement). Overall production and loss terms for HCHO were around 3 times smaller compared to the values obtained for the sites in Germany and Cyprus. We also observed smaller concentrations of CH₄, OH radicals and HCHO. It is notable that HCHO production was slightly higher than the loss for both the HUMPPA and the HOPE campaign. This is also reflected by peaks in the rate of change (0.1 - 0.2 ppbv h⁻¹) in the morning/midday hours (Figure S4b and c of the Supplement) and could indicate a transport effect from areas with lower HCHO concentration, e.g. entrainment, according to Equation (6). For HUMPPA, this idea is supported when excluding the data impacted by biomass burning which we show in Figure S11a of the Supplement. It can be seen that HCHO production prevailed over its loss suggesting a missing loss term. The difference was highest in the morning hours with approximately 0.1 ppbv h⁻¹ and decreased throughout the day indicating e.g. a vertical dilution from higher, HCHO-poor altitudes. For comparison, Figure S11b shows the HCHO production and loss terms when only considering data impacted by biomass burning. Due to high HCHO mixing ratios the loss terms were substantially higher compared to its production. In periods influenced by biomass burning, the highest HCHO yield was from acetaldehydhe (averaged diel mixing ratios of 1.1 ppbv compared to 0.6 ppbv without biomass burning); when excluding biomass burning, the yield from methane was slightly higher.

3.3.2 Net O₃ production

Ozone concentrations in Hohenpeißenberg were on average 44 ± 11 ppbv with a minimum of 5 and a maximum of 97 ppbv. The campaign averaged, diel profile displayed a peak of 48 ppbv at 16:00 UTC and a minimum of 39 ppbv at 7:00 UTC. In Hyytiälä, ozone concentrations were between 20 and 76 ppbv and 42 ± 11 ppbv on average. A diel peak of 49 ppbv was reached between 15:00 and 16:00 UTC and a minimum of 34 ppbv at 5:00 UTC. The temporal development of ozone concentrations during the campaigns and the diurnal averages can be found in Figure S6 and S7 of the Supplement. The uncertainty is dominated by the atmospheric variability (1σ) which is on average 24 % for O₃ for HOPE and 23 % for HUMPPA.

Figure 10 shows ozone production and loss terms for the research campaigns HOPE and HUMPPA. The overall uncertainty was again dominated by the atmospheric variability (1σ) which was 58 % for daytime O₃ production and 90 % for daytime O₃ loss in Hohenpeißenberg and 44 % and 51 %, respectively, in Hyytiälä. In comparison, the measurement uncertainty regarding overall production and loss was between 10 and 15 %. A detailed overview of the uncertainties can be found in Table S4 of the Supplement. We adopted the dry deposition velocity for ozone in Hyytiälä from Rannik et al. (2012). No literature values for ozone dry deposition in Hohenpeißenberg were available. Therefore, we applied the same values as for the HUMPPA campaign which increased the uncertainty of the analysis. However, the fraction of ozone dry deposition of the overall loss was small. For the applied deposition velocity, dry deposition contributed by around 4 % during daytime. For a change in v_d of ± 100 %, the fraction varied between 2 and 7 %. We therefore assume a maximum additional uncertainty of 5 % resulting from this estimate. Ozone production in Hohenpeißenberg reached peak values of 38 ppbv h⁻¹. In contrast, ozone loss showed a maximum of only 21 ppbv h⁻¹ most of which was due to the reaction with NO. In Hyytiälä, ozone production was 6.7 ppbv h⁻¹ at its diurnal




Figure 10. Diurnal profiles of O_3 production $P(O_3)$ and loss $L(O_3)$ terms during the HOPE campaign 2012 (top row: (a) $P(O_3)$, (b) $L(O_3)$ and (c) net O_3) and the HUMPPA campaign 2010 (bottom row: (d) $P(O_3)$, (e) $L(O_3)$ and (f) net O_3).

maximum while ozone loss reached 5.1 ppbv h^{-1} and was mainly composed of the loss via NO, followed by dry deposition. The differences between production and loss terms could again indicate a transport effect according to Equation (10) as described for HCHO. The rate of change for O_3 showed peak values ($2 - 3 \text{ ppbv h}^{-1}$) during morning/midday hours which was not observed
465 for the CYPHEX campaign (Figure S7 of the Supplement). O_3 production and loss terms during HUMPPA and CYPHEX were similar while the values were significantly higher during HOPE. O_3 production in Hohenpeißenberg was almost one order of magnitude higher compared to the other sites which was likely due to the higher ambient NO_x concentrations. The net O_3 production (the difference between O_3 production and loss) at each site could give a hint regarding the dominant chemical ozone regime. For HOPE, net O_3 production was significantly above zero with diurnal maximum values of around 20 ppbv h^{-1}
470 at 10:00 UTC which could indicate a VOC limitation. In contrast, net O_3 production was close to zero for CYPHEX and the ozone regime was more likely NO_x limited. We will discuss the dominant chemical ozone regime in detail in the following section.

3.4 Chemical regime

Various methods exist for determining the prevailing chemical ozone regime (i.e. the net efficiency of production or loss), one of which is the ratio of formaldehyde and nitrogen dioxide. We have calculated the diel HCHO/NO₂ ratios for all three measurement sites which is shown in Figure 11a. The HCHO/NO₂ ratio was highest with 8.0 ± 2.4 in Cyprus, followed by 1.4 ± 0.7 in Hyytiälä and 0.7 ± 0.2 in Hohenpeißenberg. According to findings by Martin et al. (2004) and Duncan et al. (2010) these values indicate a dominant NO_x limited regime during CYPHEX and a dominant VOC limited regime during HOPE. The regime during HUMPPA was likely transitioning between both limitations. We show two additional approaches for determining the present chemical regime in Figure 11. In Figure 11b, we present the daytime averages of $\alpha_{CH_3O_2}$ versus the NO concentration. All campaigns show a linear correlation. Daytime average NO concentrations during CYPHEX ranged from 1 to 45 pptv accompanied by an increase in $\alpha_{CH_3O_2}$ from around 55 to 75 %, giving a slope of 5.54 ppb_v^{-1} . The slope for the increase in $\alpha_{CH_3O_2}$ with NO is approximately half for the HUMPPA campaign with a value of 2.52 ppb_v^{-1} . The NO concentration ranged from 22 to 76 pptv along with an increase in $\alpha_{CH_3O_2}$ from 73 to 87 %. Finally for the HOPE campaign, NO concentrations and its range were highest with values between 60 and 350 pptv. At the same time, $\alpha_{CH_3O_2}$ showed the smallest increase by only 3 percentage points. The resulting slope was smallest with a value of 0.11 ppb_v^{-1} . $\alpha_{CH_3O_2}$ indicates the share of CH₃O₂ that forms HCHO, predominantly through the reaction with NO. The competing reaction is the conversion with HO₂ to CH₃OOH. A high value for $\alpha_{CH_3O_2}$ which is not or only little responsive to a changing NO concentration indicates a VOC limited regime while in a NO_x limited regime, small changes in ambient NO have a large effect on the HCHO formation from CH₃O₂. In analogy, we show the HCHO yield from isoprene according to Equation (3) versus NO concentrations in Figure 11c which we suggest as an indicator for the present chemical regime, too. For the CYPHEX campaign, the isoprene yield was most responsive to ambient NO concentrations with a slope of 1.61 ppb_v^{-1} indicating a NO_x limitation. In contrast, the isoprene yield during the HOPE campaign was almost non-responsive to changing NO indicating a VOC limitation (slope of 0.05 ppb_v^{-1}). Although specialized instrumentation is still necessary to measure NO, OH and HO₂, these methods to determine the dominant chemical regime only require the knowledge of a small number of trace gas concentrations and the ambient temperature.

4 Conclusions

In this study, we have analyzed the photochemical processes contributing to formaldehyde and ozone production and loss across Europe based on in-situ trace gas observations during three different stationary field campaigns in Cyprus (CYPHEX 2014), Germany (HOPE 2012) and Finland (HUMPPA 2010). Very consistently across all sites, we found that formaldehyde loss can be predominantly accounted for by the production via OH oxidation of methane, acetaldehyde, isoprene and methanol. Formaldehyde loss is represented by photolysis, OH oxidation and to a small part by dry deposition. Ozone chemistry is mainly controlled by nitrogen oxides. The production can be described by NO₂ photolysis and the loss is mainly a function of NO reduction and to a smaller part of photolysis and dry deposition. We found a good agreement between O₃ production and loss in Cyprus and Finland, while the production was approximately double its loss in Southern Germany. Finally, we have presented




Figure 11. Determination of the dominant chemical regime via (a) the HCHO/NO₂ ratio (colored areas indicate the dominant chemical regime according to Duncan et al. (2010) which are blue (HCHO/NO₂ > 2) for a NO_x limitation, green (HCHO/NO₂ < 1) for a VOC limitation and yellow (1 < HCHO/NO₂ < 2) for the transition), (b) the fraction of methyl peroxy radicals forming HCHO and (c) the HCHO yield from isoprene.

several different approaches for determining the prevalent chemical regime which included the HCHO/NO₂ ratio as well as the fraction of CH₃O₂ forming HCHO and the HCHO yield from isoprene in dependence on the ambient NO concentration. We identify a VOC limited regime during the HOPE campaign in Germany and a NO_x limited regime during the CYPHEX campaign in Cyprus, whereas chemistry during the HUMPPA campaign in Finland was likely at a transition point.

510 While ongoing research on HCHO photochemical processes has continuously widened the contributors to possible production paths and the complexity of calculations and models, we show that the consideration of only four precursor VOC is capable of almost completely representing the HCHO production term at various sites across Europe. We encourage to widen HCHO budget calculations based on in-situ trace gas observations to more sites worldwide as a simple, but effective tool to monitor photochemical processes and air quality, including the dominant chemical regime.

515 *Data availability.* Data measured during the research campaigns CYPHEX, HOPE and HUMPPA are available to all scientists and will be uploaded to <https://keeper.mpg.de/>.

Author contributions. HF had the idea. CMN and HF designed the study. CMN analyzed the data and wrote the manuscript. JS, JNC and BB provided J-values. CPD provided trace-gas measurements for HOPE. JNC provided OH and HO₂ model data for HUMPPA. JW provided VOC data. HH, CE, AN, KT, MM, CM and LT provided OH and HO₂ data for CYPHEX and HOPE. HCHO data were measured and provided by AR and RA. SH provided CH₃OOH data. NO_x and O₃ data were obtained from HF.

Competing interests. The authors have no competing interests to declare.

Acknowledgements. We acknowledge all researchers and supporting personnel who participated in the CYPHEX campaign 2014, the HOPE campaign 2012 and the HUMPPA campaign 2010. We would like to thank Pekka Rantala (University of Helsinki) for providing the CH₃CHO data for HUMPPA. This work was supported by the Max Planck Graduate Center with the Johannes Gutenberg-Universität Mainz (MPGC).

525 *Financial support.* The article processing charges for this open-access publication were covered by the Max Planck Society.

References

- Alam, M. S., Camredon, M., Rickard, A. R., Carr, T., Wyche, K. P., Hornsby, K. E., Monks, P. S., and Bloss, W. J.: Total radical yields from tropospheric ethene ozonolysis, *Physical Chemistry Chemical Physics*, 13, 11 002–11 015, <https://doi.org/10.1039/C0CP02342F>, 2011.
- Anderson, D., Nicely, J., Wolfe, G., Hanisco, T., Salawitch, R., Canty, T., Dickerson, R., Apel, E., Baidar, S., Bannan, T., Blake, N. J., Chen, D., Dix, B., Fernandez, R. P., Hall, S. R., Hornbrook, R. S., Huey, L. G., Josse, B., Jöckel, P., Kinnison, D. E., Koenig, T. K., Breton, M. L., Marécal, V., Morgenstern, O., Oman, L. D., Pan, L. L., Percival, C., Plummer, D., Revell, L. E., Rozanov, E., Saiz-Lopez, A., Stenke, A., Sudo, K., Tilmes, S., Ullmann, K., Volkamer, R., Weinheimer, A. J., , and Zeng, G.: Formaldehyde in the tropical western Pacific: Chemical sources and sinks, convective transport, and representation in CAM-Chem and the CCM1 models., *Journal of Geophysical Research: Atmospheres*, 122, 11,201–11,226, <https://doi.org/10.1002/2016JD026121>, 2017.
- 530 Assaf, E., Song, B., Tomas, A., Schoemaeker, C., and Fittschen, C.: Rate constant of the reaction between CH₃O₂ radicals and OH radicals revisited, *The Journal of Physical Chemistry A*, 120, 8923–8932, <https://doi.org/10.1021/acs.jpca.6b07704>, 2016.
- Assaf, E., Sheps, L., Whalley, L., Heard, D., Tomas, A., Schoemaeker, C., and Fittschen, C.: The reaction between CH₃O₂ and OH radicals: product yields and atmospheric implications, *Environmental science & technology*, 51, 2170–2177, <https://doi.org/10.1021/acs.est.6b06265>, 2017.
- 540 Atkinson, R., Baulch, D., Cox, R., Crowley, J., Hampson, R., Hynes, R., Jenkin, M., Rossi, M., Troe, J., and Subcommittee, I.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume II—gas phase reactions of organic species, *Atmospheric chemistry and physics*, 6, 3625–4055, <https://doi.org/10.5194/acp-6-3625-2006>, 2006.
- Axinte, R. D.: The oxidation photochemistry and transport of hydrogen peroxide and formaldehyde at three sites in Europe: trends, budgets and 3-D model simulations, Ph.D. thesis, <https://doi.org/10.25358/openscience-3313>, 2016.
- 545 Ayers, G., Gillett, R., Granek, H., De Serves, C., and Cox, R.: Formaldehyde production in clean marine air, *Geophysical research letters*, 24, 401–404, <https://doi.org/10.1029/97GL00123>, 1997.
- Bohn, B., Corlett, G. K., Gillmann, M., Sanghavi, S., Stange, G., Tensing, E., Vrekoussis, M., Bloss, W., Clapp, L., Kortner, M., Dorn, H.-P., Monks, P. S., Platt, U., Plass-Dulmer, C., Mihalopoulos, N., Heard, D. E., Clemmshaw, K. C., Meixner, F. X., Prevot, A. S. H., and Schmitt, R.: Photolysis frequency measurement techniques: results of a comparison within the ACCENT project, *Atmospheric chemistry and physics*, 8, 5373–5391, <https://doi.org/10.5194/acp-8-5373-2008>, 2008.
- 550 Bozem, H., Butler, T. M., Lawrence, M. G., Harder, H., Martinez, M., Kubistin, D., Lelieveld, J., and Fischer, H.: Chemical processes related to net ozone tendencies in the free troposphere, *Atmospheric Chemistry and Physics*, 17, 10 565–10 582, <https://doi.org/10.5194/acp-17-10565-2017>, 2017.
- Crowley, J. N., Pouvesle, N., Phillips, G. J., Axinte, R., Fischer, H., Petäjä, T., Nölscher, A., Williams, J., Hens, K., Harder, H., Novelli, M. M.-H. A., Kubistin, D., Bohn, B., and Lelieveld, J.: Insights into HO_x and RO_x chemistry in the boreal forest via measurement of peroxyacetic acid, peroxyacetic nitric anhydride (PAN) and hydrogen peroxide, *Atmospheric chemistry and physics*, 18, 13 457–13 479, <https://doi.org/10.5194/acp-18-13457-2018>, 2018.
- 555 Derstroff, B., Hüser, I., Bourtsoukidis, E., Crowley, J. N., Fischer, H., Gromov, S., Harder, H., Janssen, R. H., Kesselmeier, J., Lelieveld, J., Mallik, C., Martinez, M., Novelli, A., Parchatka, U., Phillips, G. J., Sander, R., Sauvage, C., Schuladen, J., Stönnner, C., Tomsche, L., and Williams, J.: Volatile organic compounds (VOCs) in photochemically aged air from the eastern and western Mediterranean, *Atmospheric Chemistry and Physics*, 17, 9547–9566, <https://doi.org/10.5194/acp-17-9547-2017>, 2017.
- 560

- Dienhart, D., Crowley, J. N., Harder, H., Hottmann, B., Martinez, M., Parchatka, U., Pfannerstill, E. Y., Rohloff, R., Schuladen, J., Tadic, I., Tauer, S., Wang, N., Williams, J., Lelieveld, J., and Fischer, H.: Measurement report: Observation-based formaldehyde production rates and their relation to OH reactivity during AQABA, *Atmospheric Chemistry and Physics*, <https://doi.org/10.5194/acp-2021-304>, 2021.
- 565 DiGangi, J. P., Boyle, E. S., Karl, T., Harley, P., Turnipseed, A., Kim, S., Cantrell, C., Maudlin Iii, R. L., Zheng, W., Flocke, F., Hall, S. R., Ullmann, K., Nakashima, Y., Paul, J. B., Wolfe, G. M., Desai, A. R., Kajii, Y., Guenther, A., and Keutsch, F. N.: First direct measurements of formaldehyde flux via eddy covariance: implications for missing in-canopy formaldehyde sources, *Atmospheric Chemistry and Physics*, 11, 10 565–10 578, <https://doi.org/10.5194/acp-11-10565-2011>, 2011.
- Duncan, B. N., Yoshida, Y., Olson, J. R., Sillman, S., Martin, R. V., Lamsal, L., Hu, Y., Pickering, K. E., Retscher, C., Allen, D. J., and
570 Crawford, J. H.: Application of OMI observations to a space-based indicator of NO_x and VOC controls on surface ozone formation, *Atmospheric Environment*, 44, 2213–2223, <https://doi.org/10.1016/j.atmosenv.2010.03.010>, 2010.
- Emmerichs, T., Kerkweg, A., Ouwersloot, H., Fares, S., Mammarella, I., and Taraborrelli, D.: A revised dry deposition scheme for land–atmosphere exchange of trace gases in ECHAM/MESSEy v2. 54, *Geoscientific Model Development*, 14, 495–519, <https://doi.org/10.5194/gmd-14-495-2021>, 2021.
- 575 Fischer, H., Axinte, R., Bozem, H., Crowley, J. N., Ernest, C., Gilge, S., Hafermann, S., Harder, H., Hens, K., Janssen, R. H., Königstedt, R., Kubistin, D., Mallik, C., Martinez, M., Novelli, A., Parchatka, U., Plass-Dülmer, C., Pozzer, A., Regelin, E., Reiffs, A., Schmidt, T., Schuladen, J., and Lelieveld, J.: Diurnal variability, photochemical production and loss processes of hydrogen peroxide in the boundary layer over Europe, *Atmospheric chemistry and physics*, 19, 11 953–11 968, <https://doi.org/10.5194/acp-19-11953-2019>, 2019.
- Fittschen, C., Whalley, L. K., and Heard, D. E.: The reaction of CH₃O₂ radicals with OH radicals: a neglected sink for CH₃O₂ in the remote
580 atmosphere, <https://doi.org/10.1021/es502481q>, 2014.
- Fortems-Cheiney, A., Chevallier, F., Pison, I., Bousquet, P., Saunois, M., Szopa, S., Cressot, C., Kurosu, T., Chance, K., and Fried, A.: The formaldehyde budget as seen by a global-scale multi-constraint and multi-species inversion system, *Atmospheric Chemistry and Physics*, 12, 6699–6721, <https://doi.org/10.5194/acp-12-6699-2012>, 2012.
- Franco, B., Blumenstock, T., Cho, C., Clarisse, L., Clerbaux, C., Coheur, P.-F., De Mazière, M., De Smedt, I., Dorn, H.-P., Emmerichs, T.,
585 Fuchs, H., Gkatzelisand, G., Griffith, D. W. T., Gromov, S., Hannigan, J. W., Hase, F., Hohaus, T., Jones, N., Kerkweg, A., Kiendler-Scharr, A., Lutsch, E., Mahieu, E., Novelli, A., Ortega, I., Paton-Walsh, C., Pommier, M., Pozzer, A., Reimer, D., Rosanka, S., Sander, R., Schneider, M., Strong, K., Tillmann, R., Van Roozendae, M., Vereecken, L., Vigouroux, C., Wahner, A., and Taraborrelli, D.: Ubiquitous atmospheric production of organic acids mediated by cloud droplets, *Nature*, 593, 233–237, <https://doi.org/10.1038/s41586-021-03462-x>, 2021.
- 590 Fried, A., Cantrell, C., Olson, J., Crawford, J. H., Weibring, P., Walega, J., Richter, D., Junkermann, W., Volkamer, R., Sinreich, R., Heikes, B. G., O’Sullivan, D., Blake, D. R., Blake, N., Meinardi, S., Apel, E., Weinheimer, A., Knapp, D., Perring, A., Cohen, R. C., Fuelberg, H., Shetter, R. E., Hall, S. R., Ullmann, K., Brune, W. H., Mao, J., Ren, X., Huey, L. G., Singh, H. B., Hair, J. W., Riemer, D., Diskin, G., and Sachse, G.: Detailed comparisons of airborne formaldehyde measurements with box models during the 2006 INTEX-B and MILAGRO campaigns: potential evidence for significant impacts of unmeasured and multi-generation volatile organic carbon compounds,
595 *Atmospheric Chemistry and Physics*, 11, 11 867–11 894, 2011.
- Graus, M., Müller, M., and Hansel, A.: High resolution PTR-TOF: quantification and formula confirmation of VOC in real time, *Journal of the American Society for Mass Spectrometry*, 21, 1037–1044, <https://doi.org/10.1016/j.jasms.2010.02.006>, 2010.

- Hosaynali Beygi, Z., Fischer, H., Harder, H., Martinez, M., Sander, R., Williams, J., Brookes, D., Monks, P., and Lelieveld, J.: Oxidation photochemistry in the Southern Atlantic boundary layer: unexpected deviations of photochemical steady state, *Atmospheric Chemistry and Physics*, 11, 8497–8513, <https://doi.org/10.5194/acp-11-8497-2011>, 2011.
- IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation: Data Sheet HOx VOC34, <http://iupac.pole-ether.fr>, "accessed on 2021-07-05", 2007.
- IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation: Data Sheet HOx VOC95, <http://iupac.pole-ether.fr>, "accessed on 2021-06-09", 2017.
- IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation: Data Sheet HOx VOC54, <http://iupac.pole-ether.fr>, "accessed on 2021-06-09", 2019.
- IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation: Evaluated Kinetic Data, <http://iupac.pole-ether.fr>, "accessed on 2021-07-05", 2021.
- Jacob, D. J.: *Introduction to atmospheric chemistry*, Princeton University Press, 1999.
- Jin, X. and Holloway, T.: Spatial and temporal variability of ozone sensitivity over China observed from the Ozone Monitoring Instrument, *Journal of Geophysical Research: Atmospheres*, 120, 7229–7246, <https://doi.org/10.1002/2015JD023250>, 2015.
- Jin, X., Fiore, A., Boersma, K. F., Smedt, I. D., and Valin, L.: Inferring Changes in Summertime Surface Ozone–NO_x–VOC Chemistry over US Urban Areas from Two Decades of Satellite and Ground-Based Observations, *Environmental science & technology*, 54, 6518–6529, <https://doi.org/10.1021/acs.est.9b07785>, 2020.
- Kaiser, J., Wolfe, G., Bohn, B., Broch, S., Fuchs, H., Ganzeveld, L. N., Gomm, S., Häsel, R., Hofzumahaus, A., Holland, F., Jäger, J., Li, X., Lohse, I., Lu, K., Prévôt, A. S. H., Rohrer, F., Wegener, R., Wolf, R., Mentel, T. F., Kiendler-Scharr, A., Wahner, A., and Keutsch, F. N.: Evidence for an unidentified non-photochemical ground-level source of formaldehyde in the Po Valley with potential implications for ozone production, *Atmospheric chemistry and physics*, 15, 1289–1298, <https://doi.org/10.5194/acp-15-1289-2015>, 2015.
- Kormann, R., Fischer, H., Reus, M. d., Lawrence, M., Brühl, C., Kuhlmann, R. v., Holzinger, R., Williams, J., Lelieveld, J., Warneke, C., J. d. G., Heland, J., Ziereis, H., and Schlager, H.: Formaldehyde over the eastern Mediterranean during MINOS: Comparison of airborne in-situ measurements with 3D-model results, *Atmospheric Chemistry and Physics*, 3, 851–861, <https://doi.org/10.5194/acp-3-851-2003>, 2003.
- Lee, A., Goldstein, A. H., Kroll, J. H., Ng, N. L., Varutbangkul, V., Flagan, R. C., and Seinfeld, J. H.: Gas-phase products and secondary aerosol yields from the photooxidation of 16 different terpenes, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2006JD007050>, 2006.
- Levitt, S. B. and Chock, D. P.: Weekday-weekend pollutant studies of the Los Angeles basin, *Journal of the Air Pollution Control Association*, 26, 1091–1092, 1976.
- Lightfoot, P. D., Cox, R. A., Crowley, J. N., Destriau, M., Hayman, G. D., Jenkin, M. E., Moortgat, G. K., and Zabel, F.: Organic peroxy radicals: kinetics, spectroscopy and tropospheric chemistry, *Atmospheric Environment. Part A. General Topics*, 26, 1805–1961, [https://doi.org/10.1016/0960-1686\(92\)90423-I](https://doi.org/10.1016/0960-1686(92)90423-I), 1992.
- Lippmann, M.: Health effects of ozone a critical review, *Japca*, 39, 672–695, <https://doi.org/10.1080/08940630.1989.10466554>, 1989.
- Lowe, D. C. and Schmidt, U.: Formaldehyde (HCHO) measurements in the nonurban atmosphere, *Journal of Geophysical Research: Oceans*, 88, 10 844–10 858, <https://doi.org/10.1029/JC088iC15p10844>, 1983.

- Luecken, D., Napelenok, S., Strum, M., Scheffe, R., and Phillips, S.: Sensitivity of ambient atmospheric formaldehyde and ozone to precursor species and source types across the United States, *Environmental science & technology*, 52, 4668–4675, <https://doi.org/10.1021/acs.est.7b05509>, 2018.
- Mallik, C., Tomsche, L., Bourtsoukidis, E., Crowley, J. N., Derstroff, B., Fischer, H., Hafermann, S., Hueser, I., Javed, U., Kessel, S., Lelieveld, J., Martinez, M., Meusel, H., Novelli, A., Phillips, G. J., Pozzer, A., Reiffs, A., Sander, R., Taraborrelli, D., Sauvage, C., Schuladen, J., Su, H., Williams, J., and Harder, H.: Oxidation processes in the eastern Mediterranean atmosphere: evidence from the modelling of HO_x measurements over Cyprus, *Atmospheric chemistry and physics*, 18, 10 825–10 847, <https://doi.org/10.5194/acp-18-10825-2018>, 2018.
- Marno, D., Ernest, C., Hens, K., Javed, U., Klimach, T., Martinez, M., Rudolf, M., Lelieveld, J., and Harder, H.: Calibration of an airborne HO_x instrument using the All Pressure Altitude-based Calibrator for HO_x Experimentation (APACHE), *Atmospheric Measurement Techniques*, 13, 2711–2731, <https://doi.org/10.5194/amt-13-2711-2020>, 2020.
- Martin, R. V., Fiore, A. M., and Van Donkelaar, A.: Space-based diagnosis of surface ozone sensitivity to anthropogenic emissions, *Geophysical Research Letters*, 31, <https://doi.org/10.1029/2004GL019416>, 2004.
- Meusel, H., Kuhn, U., Reiffs, A., Mallik, C., Harder, H., Martinez, M., Schuladen, J., Bohn, B., Parchatka, U., Crowley, J. N., Fischer, H., Tomsche, L., Novelli, A., Hoffmann, T., Janssen, R. H. H., Hartogensis, O., Pikridas, M., Vrekoussis, M., Bourtsoukidis, E., Weber, B., Lelieveld, J., Williams, J., Pöschl, U., Cheng, Y., and Su, H.: Daytime formation of nitrous acid at a coastal remote site in Cyprus indicating a common ground source of atmospheric HONO and NO, *Atmospheric Chemistry and Physics*, 16, 14 475–14 493, <https://doi.org/10.5194/acp-16-14475-2016>, 2016.
- Miyoshi, A., Hatakeyama, S., and Washida, N.: OH radical-initiated photooxidation of isoprene: An estimate of global CO production, *Journal of Geophysical Research: Atmospheres*, 99, 18 779–18 787, <https://doi.org/10.1029/94JD01334>, 1994.
- NOAA Solar Calculator: NOAA Solar Calculator, <https://gml.noaa.gov/grad/solcalc/>, "accessed on 2021-07-08", 2021.
- Novelli, A., Hens, K., Tatum Ernest, C., Kubistin, D., Regelin, E., Elste, T., Plass-Dülmer, C., Martinez, M., Lelieveld, J., and Harder, H.: Characterisation of an inlet pre-injector laser-induced fluorescence instrument for the measurement of atmospheric hydroxyl radicals, *Atmospheric measurement techniques*, 7, 3413–3430, <https://doi.org/10.5194/amt-7-3413-2014>, 2014.
- Novelli, A., Hens, K., Tatum Ernest, C., Martinez, M., Nölscher, A. C., Sinha, V., Paasonen, P., Petäjä, T., Sipilä, M., Elste, T., Plass-Dülmer, C., Phillips, G. J., Kubistin, D., Williams, J., Vereecken, L., Lelieveld, J., and Harder, H.: Estimating the atmospheric concentration of Criegee intermediates and their possible interference in a FAGE-LIF instrument, *Atmospheric Chemistry and Physics*, 17, 7807–7826, <https://doi.org/10.5194/acp-17-7807-2017>, 2017.
- Nussbaumer, C. M. and Cohen, R. C.: The Role of Temperature and NO_x in Ozone Trends in the Los Angeles Basin, *Environmental Science & Technology*, 54, 15 652–15 659, <https://doi.org/10.1021/acs.est.0c04910>, 2020.
- Nuvolone, D., Petri, D., and Voller, F.: The effects of ozone on human health, *Environmental Science and Pollution Research*, 25, 8074–8088, <https://doi.org/10.1007/s11356-017-9239-3>, 2018.
- Ouwensloot, H., Vilà-Guerau de Arellano, J., Nölscher, A., Krol, M., Ganzeveld, L., Breitenberger, C., Mammarella, I., Williams, J., and Lelieveld, J.: Characterization of a boreal convective boundary layer and its impact on atmospheric chemistry during HUMPPA-COPEC-2010, *Atmospheric Chemistry and Physics*, 12, 9335–9353, <https://doi.org/10.5194/acp-12-9335-2012>, 2012.
- Palmer, P. I., Jacob, D. J., Fiore, A. M., Martin, R. V., Chance, K., and Kurosu, T. P.: Mapping isoprene emissions over North America using formaldehyde column observations from space, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2002JD002153>, 2003.

- Pires, J. C. M.: Ozone weekend effect analysis in three European urban areas, *CLEAN–Soil, Air, Water*, 40, 790–797, <https://doi.org/10.1002/clen.201100410>, 2012.
- Possanzini, M., Di Palo, V., and Cecinato, A.: Sources and photodecomposition of formaldehyde and acetaldehyde in Rome ambient air, *Atmospheric Environment*, 36, 3195–3201, [https://doi.org/10.1016/S1352-2310\(02\)00192-9](https://doi.org/10.1016/S1352-2310(02)00192-9), 2002.
- 675 Pusede, S. and Cohen, R.: On the observed response of ozone to NO_x and VOC reactivity reductions in San Joaquin Valley California 1995–present, *Atmospheric Chemistry and Physics*, 12, 8323–8339, <https://doi.org/10.5194/acp-12-8323-2012>, 2012.
- Pusede, S. E., Steiner, A. L., and Cohen, R. C.: Temperature and recent trends in the chemistry of continental surface ozone, *Chemical reviews*, 115, 3898–3918, <https://doi.org/10.1021/cr5006815>, 2015.
- 680 Rannik, Ü., Altimir, N., Mammarella, I., Bäck, J., Rinne, J., Ruuskanen, T., Hari, P., Vesala, T., and Kulmala, M.: Ozone deposition into a boreal forest over a decade of observations: evaluating deposition partitioning and driving variables, *Atmospheric Chemistry and Physics*, 12, 12 165–12 182, <https://doi.org/10.5194/acp-12-12165-2012>, 2012.
- Sadanaga, Y., Sengen, M., Takenaka, N., and Bandow, H.: Analyses of the ozone weekend effect in Tokyo, Japan: regime of oxidant (O₃+NO₂) production, *Aerosol and Air Quality Research*, 12, 161–168, <https://doi.org/10.4209/aaqr.2011.07.0102>, 2011.
- 685 Schroeder, J. R., Crawford, J. H., Fried, A., Walega, J., Weinheimer, A., Wisthaler, A., Müller, M., Mikoviny, T., Chen, G., Shook, M., Blake, D. R., and Tonnesen, G. S.: New insights into the column CH₂O/NO₂ ratio as an indicator of near-surface ozone sensitivity, *Journal of Geophysical Research: Atmospheres*, 122, 8885–8907, <https://doi.org/10.1002/2017JD026781>, 2017.
- Schwantes, R. H., Emmons, L. K., Orlando, J. J., Barth, M. C., Tyndall, G. S., Hall, S. R., Ullmann, K., Clair, J. M. S., Blake, D. R., Wisthaler, A., and V. Bui, T. P.: Comprehensive isoprene and terpene gas-phase chemistry improves simulated surface ozone in the southeastern US, *Atmospheric Chemistry and Physics*, 20, 3739–3776, <https://doi.org/10.5194/acp-20-3739-2020>, 2020.
- 690 Seguel, R. J., S., R. G. E. M., and G., M. A. L.: Ozone weekend effect in Santiago, Chile, *Environmental Pollution*, 162, 72–79, <https://doi.org/10.1016/j.envpol.2011.10.019>, 2012.
- Seyfioglu, R., Odabasi, M., and Cetin, E.: Wet and dry deposition of formaldehyde in Izmir, Turkey, *Science of the Total Environment*, 366, 809–818, <https://doi.org/10.1016/j.scitotenv.2005.08.005>, 2006.
- 695 Shepson, P. B., Bottenheim, J. W., Hastie, D. R., and Venkatram, A.: Determination of the relative ozone and PAN deposition velocities at night, *Geophysical research letters*, 19, 1121–1124, <https://doi.org/10.1029/92GL01118>, 1992.
- Sillman, S.: The use of NO_y, H₂O₂, and HNO₃ as indicators for ozone-NO_x-hydrocarbon sensitivity in urban locations, *Journal of Geophysical Research: Atmospheres*, 100, 14 175–14 188, <https://doi.org/10.1029/94JD02953>, 1995.
- Sobanski, N., Tang, M., Thieser, J., Schuster, G., Pöhler, D., Fischer, H., Song, W., Sauvage, C., Williams, J., Fachinger, J., Berkes, F., Hoor, P., Platt, U., Lelieveld, J., and Crowley, J. N.: Chemical and meteorological influences on the lifetime of NO₃ at a semi-rural mountain site during PARADE, *Atmospheric Chemistry and Physics*, 16, 4867–4883, <https://doi.org/10.5194/acp-16-4867-2016>, 2016.
- 700 Stickler, A., Fischer, H., Williams, J., De Reus, M., Sander, R., Lawrence, M., Crowley, J., and Lelieveld, J.: Influence of summertime deep convection on formaldehyde in the middle and upper troposphere over Europe, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2005JD007001>, 2006.
- 705 Stickler, A., Fischer, H., Bozem, H., Gurk, C., Schiller, C., Martinez-Harder, M., Kubistin, D., Harder, H., Williams, J., Eerdeken, G., Yassaa, N., Ganzeveld, L., Sander, R., and Lelieveld, J.: Chemistry, transport and dry deposition of trace gases in the boundary layer over the tropical Atlantic Ocean and the Guyanas during the GABRIEL field campaign, *Atmospheric Chemistry and Physics*, 7, 3933–3956, <https://doi.org/10.5194/acp-7-3933-2007>, 2007.

- Stockwell, W. R. and Calvert, J. G.: The mechanism of NO₃ and HONO formation in the nighttime chemistry of the urban atmosphere, *Journal of Geophysical Research: Oceans*, 88, 6673–6682, <https://doi.org/10.1029/JC088iC11p06673>, 1983.
- 710 Sumner, A. L., Shepson, P. B., Couch, T. L., Thornberry, T., Carroll, M. A., Sillman, S., Pippin, M., Bertman, S., Tan, D., Faloona, I., Brune, W., Young, V., Cooper, O., Moody, J., , and Stockwell, W.: A study of formaldehyde chemistry above a forest canopy, *Journal of Geophysical Research: Atmospheres*, 106, 24 387–24 405, <https://doi.org/10.1029/2000JD900761>, 2001.
- Tadic, I., Crowley, J. N., Dienhart, D., Eger, P., Harder, H., Hottmann, B., Martinez, M., Parchatka, U., Paris, J.-D., Pozzer, A., Rohloff, R., Schuladen, J., Shenolikar, J., Tauer, S., Lelieveld, J., and Fischer, H.: Net ozone production and its relationship to nitrogen oxides and volatile organic compounds in the marine boundary layer around the Arabian Peninsula, *Atmospheric Chemistry and Physics*, 20, 6769–6787, <https://doi.org/10.5194/acp-20-6769-2020>, 2020.
- 715 Urbanski, S. P., Stickel, R. E., Zhao, Z., and Wine, P. H.: Mechanistic and kinetic study of formaldehyde production in the atmospheric oxidation of dimethyl sulfide, *Journal of the Chemical Society, Faraday Transactions*, 93, 2813–2819, <https://doi.org/10.1039/A701380I>, 1997.
- 720 Veres, P. R., Faber, P., Drewnick, F., Lelieveld, J., and Williams, J.: Anthropogenic sources of VOC in a football stadium: Assessing human emissions in the atmosphere, *Atmospheric Environment*, 77, 1052–1059, <https://doi.org/10.1016/j.atmosenv.2013.05.076>, 2013.
- Vigouroux, C., Hendrick, F., Stavrakou, T., Dils, B., Smedt, I. D., Hermans, C., Merlaud, A., Scolas, F., Senten, C., Vanhaelewyn, G., Fally, S., Carleer, M., Metzger, J.-M., Müller, J.-F., Roozendael, M. V., and Mazière, M. D.: Ground-based FTIR and MAX-DOAS observations of formaldehyde at Réunion Island and comparisons with satellite and model data, *Atmospheric Chemistry and Physics*, 9, 9523–9544, <https://doi.org/10.5194/acp-9-9523-2009>, 2009.
- 725 Wang, Y., Hu, B., Ji, D., Liu, Z., Tang, G., Xin, J., Zhang, H., Song, T., Wang, L., Gao, W., Wang, X., and Wang, Y.: Ozone weekend effects in the Beijing–Tianjin–Hebei metropolitan area, China, *Atmospheric chemistry and physics*, 14, 2419–2429, <https://doi.org/10.5194/acp-14-2419-2014>, 2014.
- 730 Wennberg, P. O., Bates, K. H., Crouse, J. D., Dodson, L. G., McVay, R. C., Mertens, L. A., Nguyen, T. B., Praske, E., Schwantes, R. H., Smarte, M. D., St Clair, J. M., Teng, A. P., Zhang, X., and Seinfeld, J. H.: Gas-phase reactions of isoprene and its major oxidation products, *Chemical reviews*, 118, 3337–3390, <https://doi.org/10.1021/acs.chemrev.7b00439>, 2018.
- Werner, A., Plass-Dülmer, C., Englert, J., Michl, K., and Tensing, E.: VOCs at Hohenpeißenberg during HOPE 2012, in: EGU General Assembly Conference Abstracts, pp. EGU2013–8175, 2013.
- 735 Wesely, M. and Hicks, B.: A review of the current status of knowledge on dry deposition, *Atmospheric environment*, 34, 2261–2282, [https://doi.org/10.1016/S1352-2310\(99\)00467-7](https://doi.org/10.1016/S1352-2310(99)00467-7), 2000.
- Williams, J., Crowley, J., Fischer, H., Harder, H., Martinez, M., Petäjä, T., Rinne, J., Bäck, J., Boy, M., Maso, M. D., Hakala, J., Kajos, M., Keronen, P., Rantala, P., Aalto, J., Aaltonen, H., Paatero, J., Vesala, T., Hakola, H., Levula, J., Pohja, T., Herrmann, F., Auld, J., Mesarchaki, E., Song, W., Yassaa, N., Nolscher, A., Johnson, A. M., Custer, T., Sinha, V., Thieser, J., Pouvesle, N., Taraborrelli, D., Tang, M. J., Bozem, H., Hosaynali-Beygi, Z., Axinte, R., Oswald, R., Novelli, A., Kubistin, D., Hens, K., Javed, U., Trawny, K., Breitenberger, C., Hidalgo, P. J., Ebben, C. J., Geiger, F. M., Corrigan, A. L., Russell, L. M., Ouwersloot, H. G., de Arellano, J. V.-G., Ganzeveld, L., Vogel, A., Beck, M., Bayerle, A., Kampf, C. J., Bertelmann, M., Kollner, F., Hoffmann, T., Valverde, J., Gonzalez, D., Riekkola, M.-L., Kulmala, M., and Lelieveld, J.: The summertime Boreal forest field measurement intensive (HUMPPA-COPEC-2010): an overview of meteorological and chemical influences, *Atmospheric Chemistry and Physics*, 11, 10 599–10 618, <https://doi.org/10.5194/acp-11-10599-2011>, 2011.

- 745 Wittrock, F., Richter, A., Oetjen, H., Burrows, J. P., Kanakidou, M., Myriokefalitakis, S., Volkamer, R., Beirle, S., Platt, U., and Wagner, T.: Simultaneous global observations of glyoxal and formaldehyde from space, *Geophysical Research Letters*, 33, <https://doi.org/10.1029/2006GL026310>, 2006.
- Wolfe, G., Kaiser, J., Hanisco, T., Keutsch, F., De Gouw, J., Gilman, J., Graus, M., Hatch, C., Holloway, J., Horowitz, L., Lee, B. H., Lerner, B. M., Lopez-Hilfiker, F., Mao, J., Marvin, M. R., Peischl, J., Pollack, I. B., Roberts, J. M., Ryerson, T. B., Thornton, J. A., Veres, 750 P. R., and Warneke, C.: Formaldehyde production from isoprene oxidation across NO_x regimes, *Atmospheric chemistry and physics*, 16, 2597–2610, <https://doi.org/10.5194/acp-16-2597-2016>, 2016.
- Yan, C., Kocevskaja, S., and Krasnoperov, L. N.: Kinetics of the reaction of CH₃O₂ radicals with OH studied over the 292–526 K temperature range, *The Journal of Physical Chemistry A*, 120, 6111–6121, <https://doi.org/10.1021/acs.jpca.6b04213>, 2016.