

Ground-based Investigation of HOx and Ozone Chemistry in Biomass Burning Plumes in Rural Idaho

Andrew J. Lindsay¹, Daniel C. Anderson^{1,*,**}, Rebecca A. Wernis^{2,3}, Yutong Liang³, Allen H. Goldstein^{2,3}, Scott C. Herndon⁴, Joseph R. Roscioli⁴, Christoph Dyroff⁴, Ed C. Fortner⁴, Philip L. Croteau⁴, Francesca Majluf⁴, Jordan E. Krechmer⁴, Tara I. Yacovitch⁴, Walter B. Knighton⁵, Ezra C. Wood¹

¹Department of Chemistry, Drexel University, Philadelphia, PA, USA

²Department of Civil and Environmental Engineering, University of California Berkeley, Berkeley, CA, USA

³Department of Environmental Science, Policy and Management, University of California Berkeley, Berkeley, CA, USA

⁴Aerodyne Research Inc., Billerica, MA, USA

⁵Department of Chemistry and Biochemistry, Montana State University, Bozeman, MT, USA

*Now at: [University of Maryland, Baltimore County, Baltimore](#), MD, USA

**Now at: NASA Goddard Space Flight Center, Greenbelt, MD, USA

Correspondence to: [Ezra C. Wood \(ew456@drexel.edu\)](mailto:Ezra C. Wood (ew456@drexel.edu))

Abstract. Ozone (O₃), a potent greenhouse gas that is detrimental to human health, is typically found in elevated concentrations within biomass burning (BB) smoke plumes. The radical species OH, HO₂, and RO₂ (known collectively as ROx) have central roles in the formation of secondary pollutants including O₃ but are poorly characterized for BB plumes. We present measurements of total peroxy radical concentrations ([XO₂] ≡ [HO₂] + [RO₂]) and additional trace-gas and particulate matter measurements from McCall, Idaho during August 2018. There were five distinct periods in which BB smoke impacted this site. During BB events, O₃ concentrations were enhanced as evidenced by ozone enhancement ratios ($\Delta O_3 / \Delta CO$) that ranged up to 0.06 ppbv ppbv⁻¹. [XO₂] was similarly elevated during some BB events. Overall, quantified instantaneous ozone production rates (P(O₃)) were minimally impacted by the presence of smoke as NO_x enhancements were minimal. Measured XO₂ concentrations were compared to zero-dimensional box modeling results to evaluate the effectiveness of the Master Chemical Mechanism (MCM) and GEOS-Chem mechanisms overall and during periods of BB influence. The models consistently over-estimated XO₂ with the base MCM and GEOS-Chem XO₂ predictions high by an average of 28% and 20%, respectively. One period of BB influence had distinct measured enhancements of 15 pptv XO₂ that were not reflected in the model output, likely due to the presence of unmeasured HOx sources. To our knowledge, this is the first BB study featuring peroxy radical measurements.

1 Introduction

Unprecedented wildfire activity has been observed in recent years. For example, Brazil's Amazon rainforest wildfires in 2019 (Cardil et al., 2020) and Australia's bush fires in 2019-2020 (Yu et al., 2020) were both marked by historically high amounts of land burned. The United States has also observed several staggering wildfire seasons characterized by hundreds of fatalities and tens of thousands of homes destroyed. For example, the 2018 season included California's deadliest fire ever (the Camp Fire) responsible for 85 deaths (CAL-FIRE, 2021), burned a total of 3.5 Mha of land, and elicited government spending of 3.1 billion dollars for fire suppression (NIFC, 2019). Due to climate change, the impacts associated with wildfire are projected to worsen. Many regions, including much of the Western US, are expected to be warmer and drier causing increases in fuel aridity. These conditions are conducive to increases in land area burned, extended fire seasons, and frequent extreme fires (Spracklen et al.,

Deleted: Universities Space Research Association, Columbia...

Deleted: Andrew J. Lindsay (ajl384@drexel.edu)

Deleted: 25

Deleted: only slightly

Deleted: and overall

Deleted: agreed within 31%

Deleted: an

Deleted: , quite likely nitrous acid (HONO)


50 2009;Yue et al., 2014;Yue et al., 2013;Goss et al., 2020). As a result, the amount of land area burned in the Western US is expected to increase by 24 – 124% by the mid-21st century (Yue et al., 2014;Yue et al., 2013).

Wildfire smoke degrades air quality both locally and far downwind. Wildfire smoke is associated with respiratory and cardiovascular health risks (Liu et al., 2015;Reid et al., 2016;DeFlorio-Barker et al., 2019;Stowell et al., 2019). These health risks will result in modest increases of respiratory illness hospitalizations as wildfire smoke will more frequently impact densely populated areas (Liu, 2016). Smoke from biomass burning (BB), a term that includes wildfires, changes in composition as it is transported downwind. The initial composition is based on direct BB emissions: particulate matter (PM), NO_x, and organic compounds spanning a wide range of volatilities. These emissions then transform through photochemical reactions that form secondary pollutants such as ozone (O₃) and secondary organic aerosol (SOA). The formation of O₃ within BB smoke is of concern because O₃ poses additional health concerns and is a prominent greenhouse gas. While the large quantities of PM within BB smoke are primarily responsible for BB smoke health concerns (Liu et al., 2015;DeFlorio-Barker et al., 2019;Reid et al., 2016), health risks from O₃ exposure include lung irritation, decreases in lung and cardiac function, higher susceptibility to respiratory infection, and early mortality (Bell et al., 2006;Park et al., 2004;Jerrett et al., 2009;Turner et al., 2016;Silva et al., 2013). Tropospheric ozone is the third most important anthropogenic greenhouse gas after CO₂ and CH₄. Increases in tropospheric ozone between the pre-industrial era and the present, from 1750 to 2011, have accounted for 0.40 W m⁻² of radiative forcing. For perspective, CH₄ and CO₂ increases are responsible for 0.48 W m⁻² and 1.82 W m⁻² of radiative forcing, respectively (Stocker et al., 2013).

Mitigating O₃ pollution will be increasingly challenging as background O₃ concentrations are increasing in some regions. Long-term aircraft observations across the Northern Hemisphere indicate average increases of tropospheric O₃ of 5% per decade (Gaudel et al., 2020). More specifically, Western US O₃ concentrations have been increasing 0.41 ppbv yr⁻¹ despite declining concentrations in the Eastern US (Cooper et al., 2012). As biomass burning accounts for an estimated 3.5% of global tropospheric O₃ formation (Jaffe and Wigder, 2012), this source of O₃ will be more important with projected wildfire activity (Jacob and Winner, 2009;Yue et al., 2015). Cities in the western US, which often have existing O₃ pollution issues, have their air quality exacerbated by BB smoke. For example, a study focused on eight predominantly western US cities found wildfire smoke to correlate with 19% of exceedances to the previous 75 ppbv O₃ NAAQS (National Ambient Air Quality Standard) standard despite smoke only being present 4.1% of the total days studied between May and the end of September (Gong et al., 2017). This NAAQS O₃ standard is in terms of maximum 8-hour daily concentration and has been set to a more stringent standard of 70 ppbv since 2015.

The correlation between O₃ concentrations and BB smoke is not completely straightforward. Depleted O₃ concentrations can be found in freshly emitted smoke plumes due to NO emissions reacting with background O₃. Aged plumes have also been observed with depleted O₃. This has been attributed to meteorological conditions (Wentworth et al., 2018), NO_x sequestration via peroxy acetyl nitrate (PAN) formation (Alvarado et al., 2010), and significant PM emissions that attenuate sunlight and limit photochemistry (Verma et al., 2008). Ozone depletion is further exemplified by the wide variation in reported ozone enhancement ratios ($\Delta O_3/\Delta CO$). While most reported ratios are positive indicating net O₃ formation, negative $\Delta O_3/\Delta CO$ values have also been reported and indicate depletion. $\Delta O_3/\Delta CO$ ratios in temperate and boreal forest (those typical of the United States) are on average 0.018, 0.15, and 0.22 ppbv ppbv⁻¹ for smoke plumes aged < 2 days, 2-5 days, and > 5 days, respectively (Jaffe and Wigder, 2012). The wide variation in net O₃ production within BB smoke is related to several factors including fire dynamics, the extent of emissions, and meteorological conditions. Each of these factors influence the underlying ROx (“ROx” = OH, HO₂, and RO₂) chemistry that controls oxidation processes and secondary pollutant formation.

ROx chemistry has rarely been studied within BB plumes. The peroxy radicals HO₂ and RO₂ (R = organic group) oxidize nitric oxide (NO) to form NO₂ (R1 and R2). During the day, the resulting NO₂ is converted to O₃ by photolysis (R3 and R4):



95 Within these smoke plumes, the concentration and composition of ROx species depends on fire emissions, photochemical conditions, and smoke age. BB emissions include direct HOx precursors, but individual emission factors and ratios are highly variable between fires. Fire dynamics alone have a significant effect on emissions. Flaming conditions have greater combustion efficiencies and are characterized by smaller emission ratios of VOCs and PM, while reactive nitrogen emissions are dominated by HONO, NO, and NO₂ (Burling et al., 2010; Roberts et al., 2020). Smoldering fire conditions have greater VOC emission ratios, PM emissions, but lower NOx emissions (Yokelson et al., 1996). Different fuel sources, such as unique tree species, shrubs, grasses, and crops, have unique emission profiles (Koss et al., 2018). Direct HOx precursor emissions of formaldehyde (HCHO), acetaldehyde (CH₃CHO), and nitrous acid (HONO) are greater than those from more typical urban combustion sources. HONO within BB plumes is short lived with a chemical lifetime ranging 10 to 30 minutes due to photolysis. This rapid photolysis yields OH and NO and has caused HONO to act as the dominant ROx source in some freshly emitted smoke plumes (Peng et al., 2020; Yokelson et al., 2009). Ozone formation in young BB plumes is, in almost all cases, initially NOx-saturated (VOC-limited) but transitions to being NOx-limited as the NOx is photochemically processed to nitric acid and organic nitrates (Alvarado et al., 2015; Müller et al., 2016; Folkins et al., 1997). In particular, NO₂ can be efficiently sequestered in the form of peroxyacetyl nitrate (PAN) due to the importance of the acetaldehyde emissions and the relative importance of the CH₃C(O)OO radical. The subsequent thermal decomposition of PAN into air masses in which ozone production is NOx-limited can potentially lead to sustained O₃ formation far downwind of a fire. BB emissions also include unique VOCs that are typically unaccounted for by chemical mechanisms employed by models. For instance, the importance of furanoids for model predictions of secondary pollution formation has only recently been studied (Müller et al., 2016; Coggon et al., 2019; Decker et al., 2019; Salvador et al., 2021).

115 For the best understanding of the composition and concentration of radicals, direct measurements and improved models are necessary. Measurements of any ROx compounds within BB plumes are rare. Only one study has reported direct measurements of OH in a BB smoke plume, which found that freshly emitted BB plumes (22-43 minutes aged) had OH concentrations five times greater than that of background air (Yokelson et al., 2009). Our understanding of ROx chemistry in BB smoke has historically relied on calculations and models (Mason et al., 2001; Hobbs et al., 2003; de Gouw et al., 2006; Akagi et al., 2012; Liu et al., 2016; Müller et al., 2016; Parrington et al., 2013). Elevated OH concentrations have been suggested for freshly emitted smoke (Hobbs et al., 2003; Akagi et al., 2012), while especially low OH concentrations were calculated for aged plumes of ~4 days in western Canada (de Gouw et al., 2006). There are few studies that have focused on peroxy radical chemistry within BB studies and it appears there have been no direct measurements in BB smoke. Model suggested peroxy radical concentrations have exhibited a wide range of values (Mason et al., 2001; Parrington et al., 2013; Liu et al., 2016; Baylon et al., 2018), in some cases reaching unrealistically high values ($[\text{HO}_2 + \text{RO}_2] \gg 200 \text{ pptv}$ for wildfires in Nova Scotia, Canada as presented in Parrington et al. (2013)).

125 This manuscript focuses on smoke observations collected in McCall, Idaho in the Pacific Northwest – a region particularly prone to wildfire – as part of the joint NCAR WE-CAN (Western Wildfire Experiment for Cloud Chemistry, Aerosol Absorption, and Nitrogen) and NOAA FIREX (Fire Influence on Regional to Global Environments Experiment) study. Increases in wildfire activity are anticipated for parts of this region, including much of Idaho, because of climate change (Halofsky et al., 2020). Presented are possibly the first measurements of total peroxy radicals in biomass burning plumes, enabling a unique investigation

130 into the impacts of biomass burning on photochemistry and ozone production and the accuracy of commonly used atmospheric chemistry models.

2 Methods

2.1 Campaign Description


135 Measurements were collected in McCall, Idaho during the WE-CAN/FIREX 2018 campaign. McCall (elevation ~1.5 km) is a rural town in Valley County, Idaho approximately 160 km north of Boise, Idaho (Fig. 1) within the West Mountains of Idaho. While much of the local area is used for cattle grazing, the town attracts tourists year-round for outdoor recreation due to the presence of mountains, surrounding forests, and a large lake.

140 The McCall field site included two mobile laboratories and one building. Each were outfitted with instrumentation for gas- and particle-phase measurements. The largest mobile lab was the Aerodyne Mobile Laboratory (AML) (Herndon et al., 2005). The AML split its time between stationary sampling in McCall and mobile measurements in other parts of Idaho and surrounding states. Unlike the AML, the second mobile lab, known as the Miniature Aerodyne Mobile Lab (minAML), was permanently stationed at the McCall site. Our analyses focus on the date ranges 16-18 August and 21-24 August when the AML was stationed at the McCall site.

2.2 ECHAMP

145 ECHAMP (Ethane CHEMical AMPlifier) is a chemical amplification-based instrument that was used to measure total peroxy radical concentrations ($[\text{HO}_2] + [\text{RO}_2]$, or simply $[\text{XO}_2]$). ECHAMP was stationed within the minAML and sampled on a 2-minute time base. The sampling and calibration methods have been described in detail elsewhere (Anderson et al., 2019; Wood et al., 2017) and only a brief summary including details specific to the McCall deployment is described here.

150 ECHAMP ‘amplifies’ each sampled XO_2 molecule into a greater number of NO_2 molecules. The enhancement in NO_2 concentration from ‘amplification’ is then divided by an amplification factor to determine $[\text{XO}_2]$. Amplification is achieved by mixing sampled air with elevated concentrations of NO and C_2H_6 to take advantage of a radical propagation scheme (Reactions 1-2 and 5-8). Since these reactions can proceed multiple times, each XO_2 produces up to 20 NO_2 molecules. There are two sampling channels: in the amplification channel there is an immediate addition of C_2H_6 and NO , whereas in the background channel the C_2H_6 addition is delayed to avoid Reaction 5, and all peroxy radicals are converted into HONO (Wood et al., 2017). The NO_2 within the two channels are transported to and measured by respective Cavity Attenuated Phase Shift (CAPS) NO_2 monitors (Kebabian et al., 2005).



160 Ambient air is first sampled at 2.5 SLPM into a ¼” PFA tee and immediately diluted with 0.8 SLPM O_2 . The sampled air then flows through 10 cm of 0.635 cm OD PFA tubing protruding out of an inlet box. This ECHAMP inlet box is a weatherproof container with dimensions of 39 cm × 44 cm × 16 cm and was mounted 3 m above ground level on scaffolding. The sampled air

165 then entered a glass cross that was internally coated with halocarbon wax. Connected orthogonally from the sampling lines were two 15.2 cm long (0.635 cm OD) reaction chambers. The reaction chambers each subsampled at a flow rate of 0.83 lpm. The remaining flow rate of 1.64 SLPM traveled by a Vaisala HMP60 probe that measured temperature and humidity. The reaction chambers included additions of NO, C₂H₆, and N₂. In amplification mode, 20 sccm of 39.3 ppmv NO in N₂ and 50 sccm of 25% C₂H₆ in N₂ were added at the beginning of the reaction chamber, while 50 sccm of N₂ is added in a downstream position. In the background mode, the locations for the N₂ and C₂H₆ additions are reversed. The final concentrations of NO and C₂H₆ after reaction chambers were 0.827 ppmv and 1.32%, respectively. These reaction chambers alternate in one-minute intervals between background and amplification modes leading to the overall two-minute sampling time. The resulting mixtures from both channels were transported in approximately 23 m of their respective tubing to the minAML that housed both CAPS monitors.

175 Dilution of sampled air with O₂ is a new addition to ECHAMP and is similar in some ways to its use by the perCIMS method (Hornbrook et al., 2011). The elevated O₂ concentration (40%) increases the ratio of the rate of the propagation reaction R8 (C₂H₃O + O₂ → CH₃CHO + HO₂) to the rate of ethyl nitrite formation (C₂H₃O + NO + M → C₂H₅ONO + M) which is a termination reaction. Dilution also dries the sampled air leading to lower and less variable relative humidity. This is beneficial as the amplification factor decreases with increasing RH (Anderson et al., 2019; Wood et al., 2017). The humidity-dependence of the amplification factor is due to humidity dependent reactions of HO₂ and the HO₂-H₂O adduct reacting with NO to form HNO₃ (Reichert et al., 2003; Butkovskaya et al., 2009) and HO₂ wall losses (Mihele and Hastie, 1998; Reichert et al., 2003).

180 ECHAMP was calibrated to the methyl peroxy radical CH₃O₂ over a range of relative humidities (RH) using the CH₃I photolysis method as described in Anderson et al. (2019) six times at McCall. The calibrant was prepared by mixing humidified ZA with trace amounts of CH₃I from a permeation source. The resulting mixture flows to a quartz tube where CH₃I is photolyzed at 254 nm by an O₃-free Hg lamp. The resulting CH₃O₂ calibrant is then added in excess flow to the ECHAMP inlet. To quantify the CH₃O₂ produced, both reaction chambers were initially operated in background mode and the CH₃O₂ source modulated on and off by alternating the flow between UV photolysis cell and a by-pass chamber. An improvement over the prior version of this calibration method was the elimination of dead-volume in the chamber. Further details regarding this calibration technique including recent improvements are provided in the supplement. Calibrations were also conducted using the H₂O photolysis method (Anderson et al., 2019). Unfortunately, inconsistent results were obtained, and at the end of the project we discovered that the quartz photolysis cell was broken. Therefore, calibrations using the H₂O photolysis method were disregarded and only the CH₃I calibrations were used. Based on the uncertainties in the individual calibration points, the variability among individual calibration points, and uncertainties regarding sampling losses, we ascribe an uncertainty of 34% (2σ) to the measurements. See SI Sect. S1 for more information.

Deleted: 27

Deleted: the

2.3 Additional Measurements

195 Onboard the AML, Quantum Cascade Tunable Infrared Laser Direct Absorption Spectrometers (QC-TILDAS, Aerodyne Research Inc.) (McManus et al., 2015) were used to measure 1) NO, NO₂; 2) CO, N₂O, H₂O; 3) HCHO, HCOOH; 4) CH₄, C₂H₆; and 5) HCN. Ozone was measured by a 2B-Tech UV absorption instrument. While particulate matter and VOCs can positively interfere with photometric O₃ measurements (Huntzicker and Johnson, 1979; Long et al., 2021), comparison to a separate Ox measurement revealed minimal interferences in the 2B-Tech O₃ observations (see Sect. S2 of SI). VOC measurements were made by an ARI Vocus (proton-transfer-reaction high-resolution time-of-flight (PTR-HR-ToF) mass spectrometer) (Krechmer et al., 2018). Measured VOCs include isoprene, acetaldehyde, acetone, acetic acid, benzene, toluene, C₂-benzenes, C₃-benzenes, methanol, total monoterpenes, and the sum of methyl vinyl ketone (MVK) and methacrolein (MACR). BB-related VOCs measured

Deleted: AML-based instrumentation is listed here. Aerodyne Research Inc. (ARI)

Deleted: instruments

Deleted: .

Deleted: O₃

210 with the Vocus include furan, methyl furan, furfural, the sum of methyl furfural and catechol, and guaiacol. Chemically-resolved
measurements of particulate matter mass concentrations were made by an ARI Soot Particle Aerosol Mass Spectrometer (SP-
AMS) (Canagaratna et al., 2007). Additional particle-phase measurements were made with an ARI Aerosol Chemical Speciation
Monitor (ACSM) permanently stationed at the McCall site building (Ng et al., 2011).

215 Onboard the minAML, NO₂ was measured with a CAPS monitor and VOCs with the Berkeley Comprehensive Thermal
Desorption Aerosol Gas Chromatograph (cTAG) (Wernis et al., 2021). cTAG measures concentrations of VOCs, intermediate
volatility organic compounds and semi-volatile organic compounds spanning an alkane-equivalent volatility range from C₅ to
C₃₀ every hour via pre-concentration followed by thermal desorption and gas chromatography-time-of-flight mass spectrometry
(GC-TOFMS). This manuscript uses isoprene, speciated monoterpenes, 2-methyl-3-buten-2-ol (MBO), styrene, benzene and
toluene measurements taken by cTAG.

220 Meteorological measurements were made both on the AML and permanently at the McCall site. Temperature, wind speed,
and wind direction were collected by a 3-D R.M. Young (Model 81000RE) sonic anemometer stationed permanently at the McCall
site at a height of 10 m. Additional wind was measured with a 2D R.M. Young (Model 81000RE) sonic anemometer mounted to
the AML rooftop and corrected for speed and truck orientation with data from a Hemisphere GPS compass (model Vector V103).
225 [Temperature, RH, and wind data are shared in the supporting information \(Fig. S3\). Daily maximum temperatures ranged 22 °C
and 28 °C while minimum temperatures ranged 4 °C to 13 °C.](#) Solar irradiance was measured by a permanently stationed ARISense
air quality sensor system (Cross et al., 2017). This was used to derive photolysis frequencies of interest, such as J_{NO_2} , by scaling
measured irradiance to outputs from the National Center for Atmospheric Research (NCAR) Tropospheric Ultraviolet and Visible
(TUV) radiation model. This process for deriving photolysis frequencies is described in greater detail in the supplement.


230 For most calculations and chemistry analyses, measured concentrations of other compounds were synchronized to the 2-
minute ECHAMP time scale. This was achieved by averaging greater frequency measurements and linearly interpolating lower
frequency measurements. Most measurements fall in the former category and were measured at a 1 Hz sampling rate. VOCs from
cTAG were obtained at an hourly rate and were therefore linearly interpolated.

2.4 Smoke Events


235 Time periods impacted by smoke were identified with observations of the biomass burning tracers HCN, CH₃CN, organic
aerosol (OA), and CO. HCN was used as the primary tracer for BB smoke. Nitriles are commonly used as tracers for BB, and HCN
is particularly useful in the absence of nearby vehicle sources. Emissions of HCN are essentially inert within BB plumes with an
atmospheric lifetime of 2 to 4 months (Li et al., 2000). HCN emission ratios are dependent on the biomass burning fuel type (Koss
et al., 2018; Coggon et al., 2016) and can vary with fire dynamics (Roberts et al., 2020). Figure 2 presents McCall site observations
and indicates smoke-impacted time periods when ECHAMP was actively sampling and the AML was present. The lowest
240 concentrations of HCN, CH₃CN, CO, and OA were all observed on 24 August, suggesting the air sampled up until then was always
somewhat affected by BB emissions. We experienced 5 distinct smoke-impacted periods evident by clear enhancements of HCN
along with organic PM and CO. The periods of greatest smoke influence (red shaded regions in Fig. 2) were identified by sustained
periods of HCN concentrations greater than 1 ppbv. [General smoke presence \(tan shaded regions in Fig. 2\) was identified before
and after each significant smoke period when background smoke tracer concentrations remained elevated compared to stable
245 background air.](#) There were distinct smoke periods in the early evenings of 16 August and 17 August. These occurred from 18:48
MDT to 21:55 MDT on 16 Aug and 15:27 MDT to 18:15 MDT on 17 Aug. While appreciable enhancements in CO were observed
during both of these periods, a significant OA enhancement of ~20 μg m⁻³ was only observed during 17 August. For the period 21

to 24 August, smoke events occurred at earlier times. Periods of significant smoke influence on 22 August, 23 August, and 24 August were from 12:55 MDT to 16:34 MDT, 12:26 MDT to 17:07 MDT, and 7:42 MDT to 15:00 MDT, respectively. The 22 August smoke event had minimal enhancement of OA. The smoke event on 23 August was quite distinct with the greatest HCN concentrations observed for the entire campaign. After this 23 August significant smoke period highlighted in Fig. 2, concentrations of HCN, CO, and OA decreased but remained at levels above background concentrations until the 24 August event. Following the 24 August significant smoke period, CO and OA decreased to their lowest observed concentrations indicating that we were sampling an air mass with minimal smoke influence.

We identify the sources of the observed smoke by pairing NOAA Air Resources Laboratory HYbrid Single Particle Lagrangian Integrated Trajectory (HYSPLIT) (Stein et al., 2015) model back trajectories with satellite detected wildfire locations (Lindaas et al., 2017; Rogers et al., 2020). Panel A of Fig. 1 shows all wildfires detected for the date range we focus our analyses on. Three well documented fires are indicated: the Mesa Fire, the Rattlesnake Creek Fire, and the Rabbit Foot Fire. The Mesa Fire, located 38.3 km southwest of McCall, was responsible for burning 34,700 acres (~14,000 hectares) and started 26 July 2018 (FWAC, 2021c). The Rattlesnake Creek Fire started 23 July, 2018, burned over 8,000 acres (>3,300 hectares) (FWAC, 2021b). The Rabbit Foot Fire began on 2 August 2018, burned 36,000 acres (~15,000 hectares), that was not contained until November (FWAC, 2021a). Panel B of Fig. 1 shows 17 August as a representative smoke-impacted day. Shown are bihourly HYSPLIT 48 hour-long back trajectories initialized from the McCall site at 10 m above ground level using archived Global Data Assimilation System (GDAS) 1-degree meteorological data. Based on fire tracer observations (see Fig. 2), the 17 August significant smoke period started at ~15:30 MDT and persisted until 18:15 MDT. This is consistent with the 16:00 MDT and 18:00 MDT back trajectories that show air sampled during that period was transported past an active fire approximately 60 km southwest of the McCall, Idaho site. Though an uncontrolled portion of the Mesa Fire may have also contributed to these smoky conditions, the HYSPLIT trajectories would suggest earlier smoke influence possibly starting near 14:00 MDT. The smoke likely traveled between 3-5 hours to get to the McCall site from the unspecified active fire for the 17 August example shown. Figures showing HYSPLIT trajectories for other smoke influenced days are included in the SI (Sect. S4). The likely sources and smoke ages for other events are detailed here. The 16 August smoke event was likely sourced from the same unspecified wildfire ranging in age between 4-10 hours. The smoke on 22 August was likely 12-18 hours or longer and sourced from an unspecified fire located east between the McCall site and the Rabbit Foot Fire. The 23 August smoke was likely sourced from the southwest from another unspecified fire at an age of 12-18 hours, though the Mesa Fire may have contributed here. The smoky conditions of the morning of 24 August were likely from wildfires in central Oregon aged 18-30 hours until especially clean air was sourced from Northern Oregon beginning at approximately 12:00 MDT.



280 **Figure 1** The McCall, Idaho site is indicated by the red star and wildfire locations indicated by orange circles. Panel A shows detected wildfires between 15 August 2018 and 24 August 2018. Panel B includes bihourly HYSPLIT back trajectories for the representative smoke impacted day of 17 August. Shown here are active wildfires of 15 August through 17 August, as back trajectories end 24 hours back. Fire locations were retrieved using the NOAA Hazard Mapping System (HMS) (NOAA, 2021).



285 **Figure 2** Time series of smoke tracers. Tan shading represents smoke presence, while red shading signifies significant smoke periods. Significant smoke periods are defined by HCN concentrations greater than 1 ppbv. This smoke-defining concentration is indicated by the purple dashed line.

2.5 Calculations

Ozone enhancement ratios $\Delta O_3/\Delta CO$ from smoke influence were determined for the most distinct BB events of 16, 17, and 24 August. For the 17 and 24 August events, the ozone enhancement ratio was determined using the York bi-variate linear regression method (York et al., 2004) using a continuous section of O_3 and CO data that includes 60 minutes of background air, a transitional smoke period (tan shaded regions in Fig. 2), and 60 minutes of significant smoke period data (red shaded regions in Fig. 2). Enhancements in NO_2 were typically under 0.2 ppbv and so the difference between considering ΔOx ($Ox = O_3 + NO_2$) and ΔO_3 was negligible. The linear regressions are included in the supplement (Fig. S8). $\Delta O_3/\Delta CO$ for the 16 August event was determined using Eq. (1) with O_3 and CO data collected during a stable period at the start of the significant smoke period and a stable background prior to smoke presence.

$$\Delta O_3/\Delta CO = ([O_3]_{Smoke} - [O_3]_{Background}) / ([CO]_{Smoke} - [CO]_{Background}) \quad (1)$$

This event had a temporary depletion in O_3 by ~20 ppbv for the start of smoke significance, then returned to near background levels of O_3 . $\Delta O_3/\Delta CO$ values were not calculated for the remaining 22 August and 23 August smoke events. These events had less distinct O_3 enhancements and occurred at times that O_3 increased during non-smoky time periods.

The gross instantaneous Ox ($[Ox] \equiv [O_3] + [NO_2]$) production rate ($P(Ox)$, often referred to as $P(O_3)$), is the rate at which NO is converted to NO_2 by reaction with HO_2 or RO_2 (Eq. 2). Since we measure the sum of HO_2 and RO_2 , we calculate $P(Ox)$ with Eq. (3) using ECHAMP XO_2 and AML-based QC-TILDAS NO measurements. Noting that the differences between k_{HO_2+NO} and most k_{RO_2+NO} rate constants are small (Anderson et al., 2019), we use an effective rate constant (k_{eff}) equal to k_{HO_2+NO} . $P(Ox)$ from box model results were calculated using Eq. (2) and the model HO_2 and speciated RO_2 concentrations.

$$P(Ox) = k_{HO_2+NO}[HO_2][NO] + \sum k_{RO_2(i)+NO}[RO_2(i)][NO] \quad (2)$$

$$P(Ox) = k_{eff}[XO_2][NO] \quad (3)$$

Instantaneous ROx production rates $P(ROx)$ from measured compounds were calculated using Eq. (4). Each term represents a compound that undergoes photolysis to produce two ROx radicals, with compound-specific photolysis rate constants (frequencies) indicated by the J variables. Our calculated values for $P(ROx)$ are limited by the lack of measurements for HONO, which is the dominant HOx source in freshly emitted BB smoke (Peng et al., 2020). While HONO has a lifetime of ~20 minutes during daylight hours, dark plume conditions and possible photochemical formation on aerosol particles may lead to sustained HONO concentrations. Ozonolysis of measured alkenes had minimal contribution to daytime $P(ROx)$ values and was therefore omitted from this calculation but is included in model predictions.

$$P(ROx) = 2J_{O^1D}[O_3] \frac{k_{O^1D+H_2O}[H_2O]}{k_{O^1D+H_2O}[H_2O] + k_{O^1D+N_2}[N_2] + k_{O^1D+O_2}[O_2]} + 2J_{HCHO}[HCHO] + 2J_{CH_3CHO}[CH_3CHO] + 2J_{CH_3COCH_3}[CH_3COCH_3] \quad (4)$$

2.6 Zero-Dimensional Modeling

The Framework for 0-D Atmospheric Modeling (F0AM) box model (v3.2) (Wolfe et al., 2016) was used to evaluate ECHAMP XO_2 measurements and further investigate BB impacts on instantaneous chemistry. F0AM simulations were conducted for the ECHAMP 2-minute time basis for dates in which AML concomitant measurements were present. Modeling was conducted separately for the two date ranges of interest of 16 to 18 August and 21 to 24 August then combined for analysis. We primarily focus on results acquired by employing a subset of the Master Chemical Mechanism (MCM) version 3.3.1 (Saunders et al.,

Deleted: each

Deleted:

Deleted: .

Deleted: Regressions were typically conducted using a continuous section of O_3 and CO data starting with 60 minutes of background air, followed by a transitional smoke period (tan shaded regions in Fig. 2), then 60 minutes of significant smoke period data (red shaded regions in Fig. 2). Since the 24 August event was unique in that the BB influenced period was followed by very clean background air, this regression was conducted on the initial BB influenced air up until the first 60 minutes of the clean background period.

Deleted: The

Deleted: of

Deleted: uniquely

Deleted: without regression by comparing a

Deleted: to a

Deleted: $Ox = O_3 + NO_2$

Deleted: 1

Deleted: using

Deleted: 2

Deleted: 1

Deleted: 1

Deleted: 2

Deleted: 3

Deleted: 3

2003; Jenkin et al., 2003; Jenkin et al., 2015) that included only the relevant chemical species in order to avoid unnecessary reactions and improve model time consumption. [The model was constrained with all available measurements \(see SI for full list\), including concentrations of ozone, formaldehyde, acetaldehyde, acetone, isoprene, speciated monoterpenes, MVK, MACR, and MBO. While F0AM allows for total NO_x to be constrained, we instead constrained NO and NO₂ individually. Model results obtained using total NO_x constraints led to nearly identical daytime XO₂ predictions but with unrealistic nighttime XO₂ values.](#) The base MCM mechanism, referred to as “MCM-base” from here on, was augmented for two additional F0AM simulations. First, the MCM-base was expanded by including additional chemistry for BB-related VOCs (referred to as “MCM-BBVOC”) of furan, methyl furan, furfural, methyl furfural, and guaiacol by manually adding the relevant chemical reactions to the MCM as detailed by Coggon et al. (2019). Second, a mechanism referred to as “MCM-BBVOC-het” included heterogeneous chemistry for HO₂ loss on organic aerosols in addition to the previously detailed BB VOC chemistry. The heterogeneous loss rates are dependent on the predicted [HO₂] values, organic aerosol surface area concentration, an uptake coefficient (γ), and mean molecular speed (Tang et al., 2014). Aerosol surface area concentrations were unmeasured and instead calculated from mass concentration measurements by applying a specific surface area. The default specific surface area was set to 4 m² g⁻¹. This setting falls slightly below the typical values measured for an urban environment of Tokyo, Japan (Hatoya et al., 2016). The default uptake coefficient was 0.20 as recommended by Jacob (2000). We explored the sensitivity of model results to both the specific surface area and uptake coefficient parameters by varying settings. We also share F0AM results acquired using the GEOS-Chem chemical mechanism. This version of the GEOS-Chem mechanism uses version 9-02 (Mao et al., 2013) with isoprene chemistry updates (Marais et al., 2016; Fisher et al., 2016; Travis et al., 2016; Kim et al., 2015). A small first-order dilution was implemented for all model experiments so that all compounds would have 24-hour lifetimes in order to prevent unreasonable [accumulation of secondary species with background concentrations for all unmeasured compounds set to 0 ppbv](#) (Wolfe et al., 2016). [Concentrations of unmeasured species were also set to 0 ppbv for the model results presented in this manuscript. Minimal changes in model results were observed for additional simulations that included a “spin-up” period in order to determine initial concentrations of unmeasured compounds.](#) As mentioned earlier, HONO is a particularly important RO_x precursor in BB plumes. In addition to it not being measured during this study, the zero-dimensional models utilized cannot be expected to accurately predict HONO concentrations since a portion of the HONO in the sampled air masses was undoubtedly emitted directly by the smoke. Furthermore, there are no HONO formation processes in the chemical mechanisms besides its homogenous formation from the reaction of OH with NO (i.e., there are no heterogeneous formation mechanisms). A complete description of our model setup, including observational constraints and uncertainties, is provided in the SI ([see Sect. S7](#)).

Deleted: This

Deleted:

Deleted: buildup

Deleted: with . b

3 Results and Discussion

3.1 Smoke Influence on ozone and its precursors




Figure 3 Time-series of O_3 , NO , NO_2 , total peroxy radicals (XO_2), and aldehydes ($HCHO$ and CH_3CHO). Modeled J_{NO_2} frequencies and calculated parameters of $P(Ox)$ and $P(ROx)$ are also provided. The red shading indicates periods of smoke influence, while somewhat smoky periods are shaded tan. $P(Ox)$ is shown in both 16-minute averages (blue circles) and 2-minute data (grey points). The dashed J_{NO_2} trace is entirely simulated by NCAR TUV model whereas the rest of the data is derived from ARISense solar irradiance measurements.

Deleted: $P(ROx)$ is speciated by reaction of $O(^1D)$ with H_2O , $HCHO$ photolysis, CH_3CHO photolysis, and acetone photolysis colored by red, blue, purple, and yellow respectively. ...




Figure 4 Correlation between O_3 and smoke tracer HCN for all observations between 9:00 and 22:00 MDT. Data points are colored by date collected.

400 Figure 3 shows site observations of O_3 and relevant measurements of NO_x , total peroxy radicals, and aldehydes. Typically O_3 increased from ~ 20 ppbv overnight up to daily maximum concentrations between 50 and 60 ppbv, with the greatest O_3 concentrations of near 70 ppbv observed on 23 August. NO_x concentrations were consistently low with typical daytime $[NO]$ and $[NO_2]$ values below 0.1 and 1.0 ppbv, respectively, suggesting O_3 production in the NO_x limited regime (Sillman et al., 1990). Ozone production is further discussed in Sect. 3.2. XO_2 typically reached daily maximum concentrations between 40 and 60 pptv.

405 Distinct increases in $[O_3]$ and $[XO_2]$ were observed during some BB events. The 17 August event had the clearest impact, with $[O_3]$ increasing ~ 10 ppbv and $[XO_2]$ increasing ~ 15 pptv over 45 minutes resulting in the maximum $[XO_2]$ observed of near 70 pptv for the entire campaign. The 16 August and 24 August smoke events had distinct $[O_3]$ increases but with less notable change in $[XO_2]$. For 24 August, smoke impacted the site throughout the morning and persisted into the afternoon. Ozone concentrations were ~ 53 ppbv for over an hour until smoky conditions dispersed. Ozone concentrations then decreased and remained near 46

410 ppbv, demonstrating a 7 ppbv elevation associated with the presence of smoke. No distinct impact on XO_2 was observed for this event. The smoke impact on 16 August was unique as $[O_3]$ was depleted by ~ 10 ppbv upon smoke arrival, returned to near background concentrations, then was again depleted by ~ 10 ppbv as smoke exited. Enhancements in $[NO_2]$ during these periods of depleted O_3 were small – only 0.5 and 1.0 ppbv – thus $[Ox]$ was depleted as well. This variation in $[O_3]$ and smoke tracers was likely due to different regions of this smoke plume impacting the site at different times. The edges of the plume may have been

415 sampled at the beginning and end of the smoke period, and the plume center sampled during the period with the greatest O_3 concentrations. XO_2 responded similarly to O_3 and was depleted by ~ 10 pptv upon the arrival and departure of smoke. The 23

August event was particularly smoky with the greatest smoke tracer concentrations of HCN, CO, and organic PM as well as the highest concentrations of O₃ for the entire campaign.

420 For the entire campaign, there is a positive correlation between daytime O₃ (and Ox) concentrations and smoke tracer
HCN (Fig. 4) with the highest values for both observed on 23 August. Enhancements in O₃ from smoke influence are highly
variable due to O₃ precursor emissions and photochemistry conditions, and there is also considerable variability in BB HCN
emission ratios as mentioned in Sect. 2.4. Clusters of data points at HCN concentrations below 0.75 ppbv are observed for the 18
August and 24 August data sets. For 18 August, there was no distinct BB influenced period and a minimal range in [HCN]. This
led to the cluster of 18 August data points with [O₃] near 60 ppbv. The 24 August data cluster near 50 ppbv [O₃] captures the stable
425 period after [O₃] is depleted by ~7 ppbv upon smoke departure followed by a slow build in concentration. Smoke age also plays a
role in this correlation plot. Based on literature trends where $\Delta O_3/\Delta CO$ values increase with age until an eventual plateau (Jaffe
and Wigder, 2012; Baker et al., 2016), young smoke plumes are likely to have smaller O₃ enhancements relative to smoke tracers
like HCN compared to aged plumes. A similar figure with O₃ plotted against CO but for times specific to the arrival or departure
of smoke is shown in the SI (Fig. S8).

430 Daily maximum P(ROx) values calculated from measured compounds ranged from 0.45 to 0.65 pptv s⁻¹ (Fig. 3). P(ROx)
was dominated by HCHO photolysis and the reaction of O(¹D) (from O₃ photolysis) with water vapor. ROx production from
photolysis of acetaldehyde and acetone were of minor importance. Aldehydes were enhanced for some of the BB events. Typically,
HCHO ranged between 1.5 and 4.5 ppbv, and BB events led to enhancements of near 2 ppbv for the 16, 17, and 24 August events.
This led to distinct impacts on P(ROx) and is most evident for the 24 August event. Distinct enhancements in O₃ for the 17 August
435 and 24 August events similarly affected P(ROx). While measurements of [XO₂] generally reflect P(ROx) trends, this was not the
case for the 17 August event. As XO₂ increased by ~27% upon the arrival of the smoke-affected air mass, P(ROx) from measured
compounds increased by at most 5%. Changes in NOx were mostly negligible during this period, though [NO] remained below
0.1 ppbv indicating that small changes in [NO] could have had a large impact on [XO₂]. Increases in both [O₃] and [HCHO] of
~20% did not contribute to a significant increase in P(ROx) due to a ~30% decrease in [H₂O] (see Fig. S16 in the supplement) and
440 a ~10% decrease in photolysis frequencies. The sudden increase in measured [XO₂] when there were only small changes in P(ROx),
NOx, and VOCs suggests the prominence of unmeasured ROx sources – most likely HONO. Peng et al. (2020) suggested HONO
enhancement ratios ($\Delta HONO/\Delta CO$) of ~0.1 pptv ppbv⁻¹ for Western US wildfire BB plumes aged 3 hours, the likely age of our
smoke plume sampled here, causing HONO photolysis to remain a significant ROx source even after 3 hours of aging. On average,
HONO accounted for >90% and 50% of P(ROx) in 30 minute aged plumes and 3 hour aged plumes, respectively (Peng et al.,
445 2020).

Deleted: S12

3.2 Ozone Production




Figure 5 P(Ox) results derived from 2 min XO₂ observations. Panel A shows the diurnal profile of all calculated P(Ox) values. The data points are colored by smoke tracer HCN. The green and blue traces represent 15-minute median values of P(Ox) as determined using measurements and model results, respectively. The provided model results were acquired with F0AM using the MCM-BBVOX mechanism. Panel B shows the variation of P(Ox) with NO. Data is colored by P(ROx) and sized by [HCN].

We describe the extent of ozone formation for the 16 August, 17 August and 24 August BB influenced periods using the commonly used $\Delta O_3/\Delta CO$ metric. These values depict the O₃ produced in transit to the McCall site while accounting for plume dilution or overall smoke influence of the site air sampled. $\Delta O_3/\Delta CO$ values were -0.02, 0.06 and 0.03 ppbv ppbv⁻¹ for the 16, 17 and 24 August smoke events, respectively. These calculated values fall within the wide variability and range of literature $\Delta O_3/\Delta CO$ values for boreal and temperate forest fire smoke plumes aged less than two days, including numerous examples of ozone depletion for aged plumes (Jaffe and Wigder, 2012). Though the smoke was likely sourced from the same wildfire for the 16 August and 17 August events (section 2.4), we observe O₃ depletion on 16 August and O₃ enhancement on 17 August. $\Delta O_3/\Delta CO$ values were not calculated for the 22 August and 23 August smoke events as we were unable to attribute the observed increases of O₃ to smoke influence as they occurred at the same time that O₃ usually increased during non-smoky time periods as mentioned in the section 2.5. Ox enhancement ratios are not presented but differed insignificantly from O₃ enhancement ratios as NO₂ concentrations were much lower than O₃ concentrations (see Fig. 3).

Instantaneous O₃ production rates are calculated using NO and XO₂ concentrations (Fig. 3). Gaps in P(Ox) are due to measurement gaps in XO₂ when ECHAMP was offline for calibrations and diagnostic tests. The highest P(Ox) values occurred on 17 and 18 August during non-smoky periods between 10:00 MDT and 12:00 MDT, reaching formation rates slightly greater than 8 ppbv hr⁻¹. For the entire campaign, median P(Ox) peaked at 11:00 MDT at 5.8 ppbv hr⁻¹. As NO concentrations were low and rarely exceeded 1 ppbv, changes in [NO] had a near-linear impact on P(Ox). Noisy P(Ox) periods, such as the entire afternoon of 16 August, are mainly attributed to the atmospheric variability of and measurement precision for NO. Overall, there is little

- Deleted:** five
- Deleted:** sampled air masses with
- Deleted:** -0.02, 0.06, 0.17, 0.25, and 0.03 ppbv ppbv⁻¹ for the 16, 17, 22, 23, and 24
- Deleted:** are generally consistent with
- Deleted:** result
- Deleted:** for boreal and temperate forest, with the older smoke plumes having greater $\Delta O_3/\Delta CO$ values
- Deleted:** The youngest smoke plumes observed were on August 16 and 17 and were both likely aged <10 hours, whereas the plumes of August 22, 23, and 24 were each likely aged >12 hours. While $\Delta O_3/\Delta CO$ values were generally greater for the aged plumes, the 24 August event was unique in that the 0.03 ppbv ppbv⁻¹ $\Delta O_3/\Delta CO$ value was most similar to the youngest plumes of August 16 and 17. This wide variability is not surprising given the range of $\Delta O_3/\Delta CO$ values observed in many studies, including numerous examples of ozone depletion for aged plumes (Jaffe and Wigder, 2012).
- Deleted:** are
- Deleted:** essentially the same
- Deleted:** Due to the negligible changes in [NOx] observed in BB-affected air masses,

correlation between P(Ox) and smoke tracers. However, elevated P(Ox) during the 17 August event is somewhat evident. The ~27% increase in XO₂ and near constant value for NO led to this temporary increase in P(Ox). P(Ox) increased from ~2.5 to 8.9 ppb hr⁻¹ during the transition from background air to significant smoke, remained elevated for 34 minutes, and then returned to near background P(Ox) rates. The overall lack of impact of BB influence on P(Ox) is further depicted in the P(Ox) diurnal cycle of Fig. 5. Modeled P(Ox) results for the same time period are also presented in Fig. 5a with the green median trend. These model results were acquired using F0AM with the MCM-BBVOC mechanism. Modeled P(Ox) is consistently greater than measured values with the greatest discrepancy occurring in the 7:45 to 8:15 MDT period. This difference is due to modeled [XO₂] being greater than measured [XO₂]. While we present results acquired using four unique chemical mechanisms, model predicted P(Ox) was always greater than measurements, though within the combined uncertainties.

Figure 5b shows the relationship between P(Ox), [NO], and P(ROx). For P(ROx) values above 0.4 ppt s⁻¹, P(Ox) increases almost linearly with [NO] up until at least 400 ppt, consistent with ozone formation being NO_x-limited. The vast majority of NO concentrations are below 400 ppt. P(Ox) at lower P(ROx) values (less than 0.2 ppt/s) exhibits much more noise and are typically below 2 ppb/hr.

3.3 Model Evaluation




Figure 6 Time series of modeled OH, HO₂, speciated RO₂, and OH reactivity (OHR). These results were acquired using the MCM-BBVOC mechanism. Periods of smoke are shaded as per Fig. 2. Measured XO₂ is included as black markers for comparison. Legend categories for XO₂ and OHR are mostly straightforward. Note that the isoprene XO₂ category includes several RO₂ species produced from isoprene oxidation.

Deleted: P(Ox) is slightly higher during the afternoon and evening smoke influenced periods compared to non-smoke periods. ...

Deleted:


Deleted: presented here

Deleted: 3


Deleted: 3

Deleted: 3

Deleted: 2



520 **Figure 7** Comparison of all daytime XO_2 observations (9:00 - 21:00 MDT) to modeled XO_2 .



525 **Figure 8** Modeled ROx production and termination for 16, 17, and 18 August. The results provided were modeled with F0AM using the MCM-BBVOc-het mechanism. Net rates are provided for reversible processes. ROx termination $L(ROx)$ and ROx production $P(ROx)$ are separated into several straightforward categories as represented in their respective legends. Heterogeneous HO_2 loss is represented by the $HO_2 + AER$ category. $P(ROx)$ derived from direct measurements, as presented in Fig. 3, is shown for comparison and is represented by the blue markers. The modeled results are shown in 10-minute intervals and were averaged from the 2-minute basis, while [measurement-based](#) $P(ROx)$ is shown in 2-minute intervals.

530 To investigate our understanding of photochemistry in biomass burning plumes we conducted zero-dimensional modeling constrained by our available measurements excluding ECHAMP XO₂ observations. Fig. 6 shows a time-series of model predicted HO₂, speciated RO₂, OH, and OH reactivity acquired using the MCM 3.3.1 mechanism with added BB VOC chemistry (“MCM-BBVOCC”). Additional results included in the SI were acquired using the three other mechanisms: MCM-base, MCM-BBVOCC-het, and the base GEOS-Chem mechanism (Figures S11-S13 in the supplement). Model [OH] (Fig. 6) had daily peak values ranging from 0.10 to 0.15 pptv (2 – 3 x 10⁶ molecules cm⁻³). The MCM-base [OH] results were typically higher but generally agreed within 5% of the other two altered MCM mechanisms. Daytime GEOS-Chem OH concentrations were typically ~75% higher than the base MCM prediction. OH predictions were generally unaffected by smoke influence for all mechanisms, though upon smoke arrival on 17 August [OH] decreased from ~0.12 pptv to 0.05 pptv for the MCM-BBVOCC mechanism with similar results observed with the other mechanisms. Modeled XO₂ comprises HO₂ (typically ~45-50%), CH₃O₂ (~20-25%), and the remaining portion a combination of CH₃CO₃, RO₂ derived from isoprene oxidation, and other organic peroxy radicals. Measured XO₂, as included in Fig. 3, is overlaid on Fig. 6. Daytime modeled [XO₂] (between 9:00 to 21:00 MDT) using the MCM-base, MCM-BBVOCC, MCM-BBVOCC-het, and GEOS-chem mechanisms are consistently greater than measured [XO₂] on average by 28%, 34%, 27%, and 20%, respectively (Fig. 7), all of which are within the combined 2σ measurement uncertainty (34%) and predicted model uncertainty (25%), though there are periods in which model values are greater than measured XO₂ by nearly a factor of two (i.e., the afternoon of 18 August). The inclusion of BB chemistry (MCM-BBVOCC) increased XO₂ predictions (compared to MCM-base results), and the inclusion of heterogeneous HO₂ uptake (MCM-BBVOCC-het) led to slight decreases (compared to MCM-BBVOCC results). The rapid increase in [XO₂] observed on 17 August was not captured under any model conditions. As discussed in Sect. 2.6, this discrepancy from 17 August is likely at least partially due to the impact of HONO, which was not measured and unlikely properly accounted for by the zero-dimensional model. Observed XO₂ on 23 August was high compared to all other days observed, while XO₂ modeled is only somewhat greater than other days. Comparison between daytime XO₂ observations and model results (Fig. 7) was determined using the York bi-variate linear regression method (York et al., 2004). Both measured and modeled [XO₂] are low at night (below 10 pptv). Model XO₂ predictions with all chemical mechanisms agreed within 10%.

545 The OH reactivities from the MCM-BBVOCC mechanism ranged from 5 to 25 s⁻¹ (Fig. 6). These OH reactivities are divided into several categories based on direct measurements and an ‘other’ category for non-measured model outputs. The ‘other’ category was often the greatest category contributing as much as 40 percent of the total reactivity at times. Incorporating heterogeneous losses to the BB VOC mechanism (MCM-BBVOCC-het results) had nearly no effect on OH reactivity, and removing BB chemistry (MCM-base results) led to a smaller range in OH reactivity – from 5 to 20 s⁻¹. The GEOS-CHEM OH reactivities were lowest ranging 3 to 14 s⁻¹ due to this mechanism having a limited number of reactions causing fewer measured compounds to be constrained. The BB VOC category shown (catechol, furan, methyl furan, furfural, methyl furfural, and guaiacol) was often the greatest contributor to the portion of OH reactivity attributed to measured compounds. BB VOC values were typically 2 to 5 s⁻¹ accounting for 10 to 30% of measured reactivity. This category played a lesser role in the MCM-base mechanism as only one measured compound was included - catechol – leading to at most 10 percent of measured reactivity. Reactivity from carbonyls (HCHO, CH₃CHO, acetone, and methyl ethyl ketone (MEK)) typically contributed 15 to 20% of the measured reactivity. The inorganic category typically accounted for 10% of measured reactivity, or 1 to 3 s⁻¹, and was dominated by CO. The four biogenic categories included were isoprene, monoterpenes (measured monoterpenes sorted into alpha pinene, beta pinene, and limonene), MVK/MACR (methyl vinyl ketone and methacrolein), and MBO (2-methyl-3-buten-2-ol). These categories typically account for between 30 to 50% of the OH reactivity attributed to measured species. Other measured reactivity categories were minor and included aromatics (benzene, toluene, C2 benzenes, C3 benzenes, and phenol), alkanes (CH₄, C₂H₆, and ethyne), and acids (formic acid and acetic acid). The ‘other’ category had greatest contributions from carbonyl compounds. Changes in OH reactivity were

Deleted: S9-S11

Deleted: results

Deleted: 31

Deleted: 27

Deleted: .

Deleted: Model-predicted XO₂ tends to follow the same pattern as observations, although the rapid increase in [XO₂] observed on 17 August was not captured under any model conditions. ...

Deleted: (9:00 to 21:00 MDT)

Deleted: Both measured and modeled [XO₂] at night were low (below 10 pptv) during this period.

Deleted: broken down

Deleted: The relative role a category plays in O₃ formation is represented by its relative OH reactivity contribution.

Deleted: had

Deleted: measured

Deleted: measured

590 observed for some smoke periods. Subtle changes were noted during the 16 and 17 August events due to changes in the inorganic, carbonyl, biogenic, and BB VOC portions. A noticeable [decrease](#) in reactivity [from 10 to <5 s⁻¹](#) occurred on 24 August upon smoke [departure](#). Decreases in nearly all reactivity categories contributed with notable contributions due to depletion in aldehyde concentrations and [CO]. This period sustained the lowest reactivities for the entire data set while also having the lowest concentrations of smoke tracers.

595 Figure 8 shows modeled ROx production and termination acquired using MCM-BBVOC-het mechanism for the period of 16 August through 18 August. ROx production is sorted into several categories, and only net rates are provided for reversible processes (e.g., net decomposition of peroxy nitrates). The sum of [measurement-based P\(ROx\)](#), [comprising the reaction of O\(¹D\)](#) with [H₂O](#) and [the photolysis of HCHO, CH₃CHO, and acetone](#) is overlaid in Fig. 8 (shown speciated in Fig. 3) and typically accounts for between 50-60% of total modeled values. For modeled P(ROx), O(¹D) + H₂O and HCHO photolysis both contribute ~25% of predicted ROx production during midday. Photolysis of carbonyls accounts for most of the remaining modeled daytime P(ROx), though only a fraction (less than 15%) [of this category](#) is from [the measured carbonyl](#) compounds [acetaldehyde and acetone](#). [Unmeasured carbonyls account for the rest](#), with [methylglyoxal and glycolaldehyde accounting for 26% and 8%](#), respectively. [Predicted methylglyoxal concentrations were typically between 0.4 and 0.6 ppbv. These concentrations are about an order of magnitude greater than those measured at mountaintop sites](#) (Mitsuishi et al., 2018; Kawamura et al., 2013) [but lower than those observed at a suburban site in China](#) (Liu et al., 2020) [and in biomass burning plumes observed in the Amazon](#) (Kluge et al., 2020). [Methylglyoxal is largely formed from the oxidation of MVK and MACR, themselves oxidation products of isoprene. A BB VOC related dicarbonyl, 4-oxo-2-pentenal \(listed as C5DICARB within MCM\), which is formed from methyl furan oxidation, accounted for 13% of P\(ROx\) from carbonyl photolysis.](#) Photolysis of acids and alcohols contribute up to ~10% of modeled P(ROx). Nighttime P(ROx) is typically ~0.1 pptv s⁻¹ and is primarily from alkene ozonolysis (> 80%) and the reaction of alkenes with NO₃. Net formation of peroxy RO₂NO₂, mainly PAN, was the dominant modeled ROx sink from the morning until ~ 12:00 MDT most days. Given that [neither PAN nor the acetyl peroxy radical \(CH₃CO₃\) were](#) directly measured and that a zero-dimensional model cannot be expected to accurately model PAN concentrations due to [its long lifetime](#), this result is highly uncertain. Midday ROx termination was dominated by ROx self-reactions with ~75% of L(ROx) from HO₂ + RO₂ and the remainder of L(ROx) almost entirely from HO₂ + HO₂. Reactions between RO₂ and other RO₂ played a comparatively minor role (<5% L(ROx)). Other L(ROx) categories consisted of reactions of RO₂ + NO, RO + NO₂, and OH + NO₂. [Heterogeneous uptake of HO₂ generally had small contributions to L\(ROx\) but at times of elevated PM concentrations such as the BB-influenced periods 17, 23, and 24 August, accounted for up to ~10% of total ROx termination \(see section 3.4 for further discussion\). Results from the 21 August through 24 August period \(see Fig. S17 in SI\) are similar to the results presented above for 16 – 18 August, though the unmeasured portion of P\(ROx\) is smaller for the former. This resulted from the considerably smaller concentrations of MVK, MACR, isoprene and BB VOCs measured during this period. The smaller portions for the carbonyls and ‘alcohols, acids’ categories led to smaller P\(ROx\) totals that peaked near 0.7 pptv s⁻¹ rather than the calculated 1.2 pptv s⁻¹ for the period shown in Fig. 8.](#)

620 3.4 Model Sensitivity

There are important fundamental limitations to how well a zero-dimensional model can describe the McCall measurements. Concentrations of several radical precursors such as nitrous acid, glycolaldehyde, methylglyoxal, and glyoxal were not constrained by measurements and were instead determined by the model. For days in which dilute biomass burning plumes arrived suddenly, it is unrealistic to expect that the model can accurately determine the concentrations of these compounds which depend on the history of the air mass. XO₂ predictions are sensitive to several model inputs. Since some secondary compounds'

Deleted: change

Deleted: with a decrease by about 50%

Deleted: arrival

Deleted: with the measured reactivity decreased from 10 to <5 s⁻¹

Deleted: measured

Deleted: sources consists of

Deleted: reaction

Deleted: O₃

Deleted: ,

Deleted: of

Deleted: – the greatest contributions are from u

Deleted:

Deleted: 19

Deleted: 13

Deleted: the

Deleted: of PAN

Deleted: .

645 concentrations build up in the model and can drive significant ROx production, we explore the sensitivity of model XO₂ predictions to the first-order dilution rate constant applied to all compounds. Shortening the dilution lifetime from 24-hr to 6-hr (the minimum value suggested by Wolfe et al. (2016) decreases model XO₂ predictions and reduces the GEOS-Chem XO₂ overprediction from 20% to 2% (Fig. S15 in SI). The most important unmeasured radical precursors that are affected by this dilution are methylglyoxal, glycolaldehyde, and 4-oxo-2-pentenal due to their collective contribution to P(ROx). We also investigate model XO₂ sensitivity to NOx. Increasing NOx inputs by 50% leads to peak daily XO₂ predictions by roughly 10%.

650 As the 15 pptv increase in [XO₂] observed on 17 August is not captured by model simulations, we include HONO as an additional model constraint in additional GEOS-Chem simulations (see Sect. S8 of SI). Constrained HONO concentrations were determined by the product of selected ΔHONO/ΔCO values to the measured CO mixing ratios during BB periods. To achieve a similar ~15 pptv XO₂ enhancement as measured during the 17 August BB event, a HONO enhancement ratio of near 3 pptv ppbv⁻¹ is required, which provides an additional 0.15 - 0.60 ppbv HONO throughout the BB period. This ΔHONO/ΔCO value is 30 times larger than observed by Peng et al. (2020) for similarly aged plumes. While this value is likely unrealistic, larger ΔHONO/ΔCO have been reported by Peng et al. (2020). Other unmeasured ROx precursors were likely present and at least partially responsible for the elevated XO₂ concentrations observed.

655 Model sensitivity to heterogeneous HO₂ uptake was also investigated. The introduction of heterogeneous losses of HO₂ had overall minimal impacts (see Fig. 7 MCM-BBVOC-het results) even though a fairly high HO₂ uptake coefficient of 0.2 was used (Abbatt et al., 2012). Heterogeneous losses decreased modeled [OH] and [XO₂] by an average of 3.4% and 2.9%, respectively, though the impact is more evident for smoke periods with elevated PM. Our analysis of HO₂ heterogeneous uptake is limited by the uncertainty in the HO₂ uptake coefficient and the specific surface area parameter. To investigate the sensitivity of model results to these parameters, these parameters were varied for several GEOS-Chem model simulations (see Figures S9 and S10 in SI). We focus our heterogeneous chemistry sensitivity tests on a 40 min period during the 17 August BB event in which OA concentrations were 30 μg m⁻³. Inclusion of heterogeneous chemistry with standard parameter settings (γ = 0.2) leads to a 11% decrease in [HO₂], whereas use of a much higher and likely unrealistic HO₂ uptake coefficient of 0.5 resulted in a 25% HO₂ decrease. A similar HO₂ decrease results from using a higher specific surface area of 10 m² g⁻¹. Use of a smaller HO₂ uptake coefficient of 0.02 led to a nearly negligible decrease in [HO₂]. Constraining the OA concentration at 100 μg m⁻³ for the same period - much higher than actually observed - leads to heterogeneous [HO₂] loss of 3% for an uptake coefficient of 0.02 and 30% for an uptake coefficient of 0.2. The only conditions in which heterogeneous loss of HO₂ to BB smoke would appear to be important would be for less dilute plumes ([OA] > 100 μg m⁻³) and a high uptake coefficient (γ > 0.2).

4 Conclusions

675 Peroxy radicals were measured during the FIREX 2018 campaign in McCall, Idaho in order to better characterize ROx chemistry and study O₃ formation within BB plumes. There were five distinct BB influenced periods that were identified using smoke tracers, primarily HCN. HYSPLIT back trajectories were paired with satellite data to suggest smoke sources and plume age. Most smoke periods had distinct enhancements in O₃ with enhancement ratios of up to 0.06 ppbv ppbv⁻¹ ΔO₃/ΔCO. Zero-dimensional box model results for XO₂ were generally greater than measured XO₂. These simulated results were acquired with F0AM using a variety of chemical mechanisms - GEOS-Chem, MCM, and two expanded versions of MCM. All model iterations followed the general trends observed for [XO₂] measurements, though a measured 15 pptv XO₂ enhancement during a 17 August BB event was not captured by any model iteration. This includes simulated results acquired using expanded versions of the MCM that had additional BB VOC chemistry and heterogeneous HO₂ and OH losses intended to better capture BB influence.

Deleted: 1

Deleted: leads to

Deleted: addition

Deleted: on average

Deleted: . For example, the inclusion of this heterogeneous chemistry led to a 11% decrease in [HO₂] during a 40 min period of the 17 August smoke period that had PM concentrations above 30 μg m⁻³ (Fig. S7 in the supplement). These results suggest that for less dilute BB plumes where PM concentrations exceed 100 μg m⁻³ (Zhou et al., 2017), heterogeneous HO₂ uptake is likely a major HO₂ sink and thus has a large impact on HOx chemistry. The modeled L(ROx) breakdown in Fig. 8 (and Fig. S13 in the supplement) shows heterogeneous HO₂ uptake to generally be of minimal importance but at times of elevated PM concentrations, such as BB influenced periods 17, 23, and 24 August, make up to ~10% of total ROx termination.

Deleted: we varied the HO₂ uptake coefficient

Deleted: Use

Deleted: in HO₂ for the same 17 August BB period

Deleted: These results suggest that for less dilute BB plumes where PM concentrations exceed 100 μg m⁻³, heterogeneous HO₂ uptake is likely a major HO₂ sink and thus has a large impact on HOx chemistry.

Formatted: Font color: Text 1

Deleted: positive

Deleted: 2

Deleted: 2

Heterogeneous losses overall had minimal impact on XO₂ even though BB smoke periods with elevated organic PM levels often near 30 μg m⁻³ led to decreases in [HO₂] by 10%. Heterogeneous chemistry was investigated using a variety of HO₂ uptake coefficients and aerosol specific surface areas. At the greatest settings, these variables led to decreases in HO₂ of at most 25% and 25%, respectively, during a period with PM above 30 μg m⁻³. Heterogeneous HO₂ uptake is likely a major HO₂ sink and thus has a large impact on HO_x chemistry for less dilute BB plumes. Both measured and modeled XO₂ concentrations were used to calculate P(Ox). The presence of smoke had an overall negligible impact on P(Ox) as NO_x enhancements were minimal.

Quantification of P(ROx) suffered from lack of constraints for several compounds including HONO, methylglyoxal, and glycolaldehyde. HONO photolysis likely contributed to the enhanced concentrations in [XO₂] measured during the 17 August BB period, though a rather large ΔHONO/ΔCO value near 3.0 ppbv ppbv⁻¹ is necessary for a similar XO₂ enhancement. An additional gas-phase process in need of validation by measurements is the role of PAN formation as a ROx sink during the morning. The importance of unmeasured ROx precursors was especially sensitive to the first-order dilution rate constant applied to all unmeasured species. Finally, the role of heterogeneous HO₂ uptake as a ROx sink would benefit from more direct measurements of particle size distribution and knowledge of HO₂ uptake coefficients. Heterogeneous HO₂ uptake was minimal for the dilute BB plumes studied here, and would appear to only be important in less dilute BB plumes if the uptake coefficient is relatively high (e.g., 0.2).

Data Availability

Peroxy radical measurements are available at <https://doi.org/10.26023/CY1Q-OT7V-G80R>. Other supporting measurements are available at https://data.eol.ucar.edu/master_lists/generated/we-can/ and upon request.

Supplement

Author contributions

AJL and ECW wrote the manuscript. All authors discussed results and commented on the manuscript. AJL, ECW, and DCA were responsible for XO₂ measurements. JRR, CD, and SCH operated and were responsible for the O₃ instrument and the TILDAS instruments for trace gases. FM, JEK, and WBK contributed with VOCUS measurements. RAW, YL, and AHG contributed with cTAG VOC measurements. EF and PLC were responsible for particle phase measurements. TIY and SCH coordinated the ARI data analysis.

Competing Interests

The authors declare that they have no conflict of interest.

Acknowledgements

This work was supported by NOAA/ACCC grants NA16OAR4310105 to Drexel University, NA16OAR4310104 to Aerodyne Research, Inc., and NA16OAR4310107 to UC Berkeley.

- Deleted: only slightly impacted
- Deleted: Both modeled and measured
- Deleted: , and
- Deleted: was quite likely responsible
- Deleted: for
- Deleted: A
- Deleted: es
- Deleted: include the role of methyl glyoxal and glycolaldehyde as ROx precursors and
- Deleted: dominant
- Deleted: While h
- Deleted: considered
- Deleted: this ROx sink likely plays a more
- Deleted: role
- Deleted: M

We profoundly thank Michele Crester and the Brundage Mountain Company for the use of the Activity Barn in McCall, Idaho as our sampling location. We acknowledge the work of additional scientists Leonid Nichman and Nathan Kreisberg for their operation and support of instruments in the field. We thank Eben Cross of QuantAQ, Inc. in Somerville, MA for providing ARISense solar irradiance data. We also acknowledge Jason Curry, Connor Daube, and Salvador Cartagena of Aerodyne Research Inc. for their general assistance including the transport of the mobile laboratories.

References:

- 755 Abbatt, J., Lee, A., and Thornton, J.: Quantifying trace gas uptake to tropospheric aerosol: recent advances and remaining challenges, *Chemical Society Reviews*, 41, 6555-6581, 2012.
- Akagi, S. K., Craven, J., Taylor, J., McMeeking, G., Yokelson, R., Burling, I., Urbanski, S., Wold, C., Seinfeld, J., and Coe, H.: Evolution of trace gases and particles emitted by a chaparral fire in California, *Atmospheric Chemistry and Physics*, 12, 1397-1421, 2012.
- 765 Alvarado, M., Logan, J., Mao, J., Apel, E., Riemer, D., Blake, D., Cohen, R., Min, K.-E., Perring, A., and Browne, E.: Nitrogen oxides and PAN in plumes from boreal fires during ARCTAS-B and their impact on ozone: an integrated analysis of aircraft and satellite observations, *Atmospheric Chemistry and Physics*, 10, 9739-9760, 2010.
- Alvarado, M., Lonsdale, C., Yokelson, R., Akagi, S. K., Coe, H., Craven, J., Fischer, E., McMeeking, G., Seinfeld, J., and Soni, T.: Investigating the links between ozone and organic aerosol chemistry in a biomass burning plume from a prescribed fire in California chaparral, *Atmospheric Chemistry and Physics*, 15, 6667-6688, 2015.
- 770 Anderson, D. C., Pavelec, J., Daube, C., Herndon, S. C., Knighton, W. B., Lerner, B. M., Roscioli, J. R., Yacovitch, T. I., and Wood, E. C.: Characterization of ozone production in San Antonio, Texas, using measurements of total peroxy radicals, *Atmospheric Chemistry and Physics*, 19, 2845-2860, 2019.
- Baker, K., Woody, M., Tonnesen, G., Hutzell, W., Pye, H., Beaver, M., Pouliot, G., and Pierce, T.: Contribution of regional-scale fire events to ozone and PM_{2.5} air quality estimated by photochemical modeling approaches, *Atmospheric Environment*, 140, 539-554, 2016.
- 775 Baylon, P., Jaffe, D., Hall, S., Ullmann, K., Alvarado, M., and Lefer, B.: Impact of biomass burning plumes on photolysis rates and ozone formation at the Mount Bachelor Observatory, *Journal of Geophysical Research: Atmospheres*, 123, 2272-2284, 2018.
- Bell, M. L., Peng, R. D., and Dominici, F.: The exposure-response curve for ozone and risk of mortality and the adequacy of current ozone regulations, *Environmental health perspectives*, 114, 532-536, 2006.
- 780 Burling, I., Yokelson, R. J., Griffith, D. W., Johnson, T. J., Veres, P., Roberts, J., Warneke, C., Urbanski, S., Reardon, J., and Weise, D.: Laboratory measurements of trace gas emissions from biomass burning of fuel types from the southeastern and southwestern United States, *Atmospheric Chemistry and Physics*, 10, 11115-11130, 2010.
- 785 Butkovskaya, N., Rayez, M.-T., Rayez, J.-C., Kukui, A., and Le Bras, G.: Water vapor effect on the HNO₃ yield in the HO₂+NO reaction: experimental and theoretical evidence, *The Journal of Physical Chemistry A*, 113, 11327-11342, 2009.
- CAL-FIRE Camp Fire: <https://www.fire.ca.gov/incident/?incident=75d4fe80-f18a-4a4a-9a37-4b564c5f6014>, access: 18 August, 2021.
- 790 Canagaratna, M., Jayne, J., Jimenez, J., Allan, J., Alfarra, M., Zhang, Q., Onasch, T., Drewnick, F., Coe, H., and Middlebrook, A.: Chemical and microphysical characterization of ambient aerosols with the aerodyne aerosol mass spectrometer, *Mass spectrometry reviews*, 26, 185-222, 2007.
- Cardil, A., De-Miguel, S., Silva, C. A., Reich, P. B., Calkin, D., Brancalion, P. H., Vibrans, A. C., Gamarra, J. G., Zhou, M., and Pijanowski, B. C.: Recent deforestation drove the spike in Amazonian fires, *Environmental Research Letters*, 15, 121003, 2020.
- Coggon, M. M., Veres, P. R., Yuan, B., Koss, A., Warneke, C., Gilman, J. B., Lerner, B. M., Peischl, J., Aikin, K. C., and Stockwell, C. E.: Emissions of nitrogen-containing organic compounds from the burning of herbaceous and arboraceous

- 795 biomass: Fuel composition dependence and the variability of commonly used nitrile tracers, *Geophysical Research Letters*, 43, 9903-9912, 2016.
- Coggon, M. M., Lim, C. Y., Koss, A. R., Sekimoto, K., Yuan, B., Gilman, J. B., Hagan, D. H., Selimovic, V., Zarzana, K. J., and Brown, S. S.: OH chemistry of non-methane organic gases (NMOGs) emitted from laboratory and ambient biomass burning smoke: evaluating the influence of furans and oxygenated aromatics on ozone and secondary NMOG formation, *Atmospheric Chemistry and Physics*, 19, 14875-14899, 2019.
- 800 Cooper, O. R., Gao, R. S., Tarasick, D., Leblanc, T., and Sweeney, C.: Long-term ozone trends at rural ozone monitoring sites across the United States, 1990–2010, *Journal of Geophysical Research: Atmospheres*, 117, 2012.
- Cross, E. S., Williams, L. R., Lewis, D. K., Magoon, G. R., Onasch, T. B., Kaminsky, M. L., Worsnop, D. R., and Jayne, J. T.: Use of electrochemical sensors for measurement of air pollution: correcting interference response and validating measurements, *Atmospheric Measurement Techniques*, 10, 3575-3588, 2017.
- 805 de Gouw, J. A., Warneke, C., Stohl, A., Wollny, A., Brock, C., Cooper, O., Holloway, J., Trainer, M., Fehsenfeld, F., and Atlas, E. L.: Volatile organic compounds composition of merged and aged forest fire plumes from Alaska and western Canada, *Journal of Geophysical Research: Atmospheres*, 111, 2006.
- Decker, Z. C., Zarzana, K. J., Coggon, M., Min, K.-E., Pollack, I., Ryerson, T. B., Peischl, J., Edwards, P., Dubé, W. P., and Markovic, M. Z.: Nighttime chemical transformation in biomass burning plumes: a box model analysis initialized with aircraft observations, *Environmental science & technology*, 53, 2529-2538, 2019.
- 810 DeFlorio-Barker, S., Crooks, J., Reyes, J., and Rappold, A. G.: Cardiopulmonary effects of fine particulate matter exposure among older adults, during wildfire and non-wildfire periods, in the United States 2008–2010, *Environmental health perspectives*, 127, 037006, 2019.
- 815 Fisher, J. A., Jacob, D. J., Travis, K. R., Kim, P. S., Marais, E. A., Miller, C. C., Yu, K., Zhu, L., Yantosca, R. M., and Sulprizio, M. P.: Organic nitrate chemistry and its implications for nitrogen budgets in an isoprene-and monoterpene-rich atmosphere: constraints from aircraft (SEAC4RS) and ground-based (SOAS) observations in the Southeast US, *Atmospheric chemistry and physics*, 16, 5969, 2016.
- Folkens, I., Wennberg, P., Hanisco, T., Anderson, J., and Salawitch, R.: OH, HO₂, and NO in two biomass burning plumes: Sources of HO_x and implications for ozone production, *Geophysical Research Letters*, 24, 3185-3188, 1997.
- 820 FWAC Rabbit Foot Fire: <https://www.fireweatheravalanche.org/wildfire/incident/51954/idaho/rabbit-foot-fire>, access: 5 July, 2021a.
- FWAC Rattlesnake Creek Fire: <https://www.fireweatheravalanche.org/wildfire/incident/48778/idaho/rattlesnake-creek-fire>, access: 18 August, 2021b.
- 825 FWAC Mesa Fire: <https://www.fireweatheravalanche.org/wildfire/incident/50013/idaho/mesa-fire>, access: 8 August, 2021c.
- Gaudel, A., Cooper, O. R., Chang, K.-L., Bourgeois, I., Ziemke, J. R., Strode, S. A., Oman, L. D., Sellitto, P., Nédélec, P., and Blot, R.: Aircraft observations since the 1990s reveal increases of tropospheric ozone at multiple locations across the Northern Hemisphere, *Science advances*, 6, eaba8272, 2020.
- 830 Gong, X., Kaulfus, A., Nair, U., and Jaffe, D. A.: Quantifying O₃ Impacts in Urban Areas Due to Wildfires Using a Generalized Additive Model, *Environ Sci Technol*, 51, 13216-13223, 10.1021/acs.est.7b03130, 2017.
- Goss, M., Swain, D. L., Abatzoglou, J. T., Sarhadi, A., Kolden, C. A., Williams, A. P., and Diffenbaugh, N. S.: Climate change is increasing the likelihood of extreme autumn wildfire conditions across California, *Environmental Research Letters*, 15, 094016, 2020.
- 835 Halofsky, J. E., Peterson, D. L., and Harvey, B. J.: Changing wildfire, changing forests: the effects of climate change on fire regimes and vegetation in the Pacific Northwest, USA, *Fire Ecology*, 16, 4, 2020.
- Hatoya, K., Okuda, T., Funato, K., and Inoue, K.: On-line measurement of the surface area concentration of aerosols in Yokohama, Japan, using the diffusion charging method, *Asian Journal of Atmospheric Environment*, 10, 1-12, 2016.

- 840 Herndon, S. C., Jayne, J. T., Zahniser, M. S., Worsnop, D. R., Knighton, B., Alwine, E., Lamb, B. K., Zavala, M., Nelson, D. D., and McManus, J. B.: Characterization of urban pollutant emission fluxes and ambient concentration distributions using a mobile laboratory with rapid response instrumentation, *Faraday discussions*, 130, 327-339, 2005.
- Hobbs, P. V., Sinha, P., Yokelson, R. J., Christian, T. J., Blake, D. R., Gao, S., Kirchstetter, T. W., Novakov, T., and Pilewskie, P.: Evolution of gases and particles from a savanna fire in South Africa, *Journal of Geophysical Research: Atmospheres*, 108, 2003.
- 845 Hornbrook, R., Crawford, J., Edwards, G., Goyea, O., Mauldin Iii, R., Olson, J., and Cantrell, C.: Measurements of tropospheric HO₂ and RO₂ by oxygen dilution modulation and chemical ionization mass spectrometry, *Atmospheric Measurement Techniques*, 4, 735, 2011.
- Huntzicker, J. J., and Johnson, R. L.: Investigation of an ambient interference in the measurement of ozone by ultraviolet absorption photometry, *Environmental Science & Technology*, 13, 1414-1416, 1979.
- Jacob, D. J.: Heterogeneous chemistry and tropospheric ozone, *Atmospheric Environment*, 34, 2131-2159, 2000.
- 850 Jacob, D. J., and Winner, D. A.: Effect of climate change on air quality, *Atmospheric environment*, 43, 51-63, 2009.
- Jaffe, D. A., and Wigder, N. L.: Ozone production from wildfires: A critical review, *Atmospheric Environment*, 51, 1-10, 2012.
- Jenkin, M., Saunders, S., Wagner, V., and Pilling, M.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part B): tropospheric degradation of aromatic volatile organic compounds, *Atmospheric Chemistry and Physics*, 3, 181-193, 2003.
- 855 Jenkin, M., Young, J., and Rickard, A.: The MCM v3. 3.1 degradation scheme for isoprene, *Atmospheric Chemistry and Physics*, 15, 11433, 2015.
- Jerrett, M., Burnett, R. T., Pope III, C. A., Ito, K., Thurston, G., Krewski, D., Shi, Y., Calle, E., and Thun, M.: Long-term ozone exposure and mortality, *New England Journal of Medicine*, 360, 1085-1095, 2009.
- 860 Kawamura, K., Okuzawa, K., Aggarwal, S., Irie, H., Kanaya, Y., and Wang, Z.: Determination of gaseous and particulate carbonyls (glycolaldehyde, hydroxyacetone, glyoxal, methylglyoxal, nonanal and decanal) in the atmosphere at Mt. Tai, *Atmospheric Chemistry and Physics*, 13, 5369-5380, 2013.
- Kebabian, P. L., Herndon, S. C., and Freedman, A.: Detection of nitrogen dioxide by cavity attenuated phase shift spectroscopy, *Analytical chemistry*, 77, 724-728, 2005.
- 865 Kim, P. S., Jacob, D. J., Fisher, J. A., Travis, K., Yu, K., Zhu, L., Yantosca, R. M., Sulprizio, M., Jimenez, J. L., and Campuzano-Jost, P.: Sources, seasonality, and trends of southeast US aerosol: an integrated analysis of surface, aircraft, and satellite observations with the GEOS-Chem chemical transport model, 2015.
- Kluge, F., Hüneke, T., Knecht, M., Lichtenstern, M., Rotermund, M., Schlager, H., Schreiner, B., and Pfeilsticker, K.: Profiling of formaldehyde, glyoxal, methylglyoxal, and CO over the Amazon: Normalised excess mixing ratios and related emission factors in biomass burning plumes, *Atmos. Chem. Phys. Discuss*, 1-37, 2020.
- 870 Koss, A. R., Sekimoto, K., Gilman, J. B., Selimovic, V., Coggon, M. M., Zarzana, K. J., Yuan, B., Lerner, B. M., Brown, S. S., and Jimenez, J. L.: Non-methane organic gas emissions from biomass burning: identification, quantification, and emission factors from PTR-ToF during the FIREX 2016 laboratory experiment, *Atmospheric Chemistry and Physics*, 18, 3299-3319, 2018.
- 875 Krechmer, J., Lopez-Hilfiker, F., Koss, A., Hutterli, M., Stoermer, C., Deming, B., Kimmel, J., Warneke, C., Holzinger, R., and Jayne, J.: Evaluation of a new reagent-ion source and focusing ion-molecule reactor for use in proton-transfer-reaction mass spectrometry, *Analytical chemistry*, 90, 12011-12018, 2018.
- Li, Q., Jacob, D. J., Bey, I., Yantosca, R. M., Zhao, Y., Kondo, Y., and Notholt, J.: Atmospheric hydrogen cyanide (HCN): Biomass burning source, ocean sink?, *Geophysical research letters*, 27, 357-360, 2000.

- 880 Lindaas, J., Farmer, D. K., Pollack, I. B., Abeleira, A., Flocke, F., Roscioli, R., Herndon, S., and Fischer, E. V.: Changes in ozone and precursors during two aged wildfire smoke events in the Colorado Front Range in summer 2015, *Atmospheric Chemistry and Physics*, 17, 10691-10707, 2017.
- Liu, J.: *Exposure and Health Impacts of PM_{2.5} from Wildfires in the Western US for the Present-day (2004-2009) and in the Future (2046-2051)*, Yale University, 2016.
- 885 Liu, J., Li, X., Li, D., Xu, R., Gao, Y., Chen, S., Liu, Y., Zhao, G., Wang, H., and Wang, H.: Observations of glyoxal and methylglyoxal in a suburban area of the Yangtze River Delta, China, *Atmospheric Environment*, 238, 117727, 2020.
- Liu, J. C., Pereira, G., Uhl, S. A., Bravo, M. A., and Bell, M. L.: A systematic review of the physical health impacts from non-occupational exposure to wildfire smoke, *Environmental research*, 136, 120-132, 2015.
- 890 Liu, X., Zhang, Y., Huey, L., Yokelson, R., Wang, Y., Jimenez, J., Campuzano-Jost, P., Beyersdorf, A., Blake, D., and Choi, Y.: Agricultural fires in the southeastern US during SEAC4RS: Emissions of trace gases and particles and evolution of ozone, reactive nitrogen, and organic aerosol, *Journal of Geophysical Research: Atmospheres*, 121, 7383-7414, 2016.
- Long, R. W., Whitehill, A., Habel, A., Urbanski, S., Halliday, H., Colón, M., Kaushik, S., and Landis, M. S.: Comparison of ozone measurement methods in biomass burning smoke: an evaluation under field and laboratory conditions, *Atmospheric Measurement Techniques*, 14, 1783-1800, 2021.
- 895 Mao, J., Paulot, F., Jacob, D. J., Cohen, R. C., Crouse, J. D., Wennberg, P. O., Keller, C. A., Hudman, R. C., Barkley, M. P., and Horowitz, L. W.: Ozone and organic nitrates over the eastern United States: Sensitivity to isoprene chemistry, *Journal of Geophysical Research: Atmospheres*, 118, 11,256-211,268, 2013.
- Marais, E. A., Jacob, D. J., Jimenez, J. L., Campuzano-Jost, P., Day, D. A., Hu, W., Krechmer, J., Zhu, L., Kim, P. S., and Miller, C. C.: Aqueous-phase mechanism for secondary organic aerosol formation from isoprene: application to the southeast United States and co-benefit of SO₂ emission controls, 2016.
- 900 Mason, S. A., Field, R. J., Yokelson, R. J., Kochivar, M. A., Tinsley, M. R., Ward, D. E., and Hao, W. M.: Complex effects arising in smoke plume simulations due to inclusion of direct emissions of oxygenated organic species from biomass combustion, *Journal of Geophysical Research: Atmospheres*, 106, 12527-12539, 2001.
- 905 McManus, J., Zahniser, M., Nelson, D., Shorter, J., Herndon, S., Jervis, D., Agnese, M., McGovern, R., Yacovitch, T., and Roscioli, J.: Recent progress in laser-based trace gas instruments: performance and noise analysis, *Applied Physics B*, 119, 203-218, 2015.
- Mihele, C., and Hastie, D.: The sensitivity of the radical amplifier to ambient water vapour, *Geophysical research letters*, 25, 1911-1913, 1998.
- 910 Mitsubishi, K., Iwasaki, M., Takeuchi, M., Okochi, H., Kato, S., Ohira, S.-I., and Toda, K.: Diurnal variations in partitioning of atmospheric glyoxal and methylglyoxal between gas and particles at the ground level and in the free troposphere, *ACS Earth and Space Chemistry*, 2, 915-924, 2018.
- Müller, M., Anderson, B. E., Beyersdorf, A. J., Crawford, J. H., Diskin, G. S., Eichler, P., Fried, A., Keutsch, F. N., Mikoviny, T., and Thornhill, K. L.: In situ measurements and modeling of reactive trace gases in a small biomass burning plume, *Atmos. Chem. Phys*, 16, 3813-3824, 2016.
- 915 Ng, N. L., Herndon, S. C., Trimborn, A., Canagaratna, M. R., Croteau, P., Onasch, T. B., Sueper, D., Worsnop, D. R., Zhang, Q., and Sun, Y.: An Aerosol Chemical Speciation Monitor (ACSM) for routine monitoring of the composition and mass concentrations of ambient aerosol, *Aerosol Science and Technology*, 45, 780-794, 2011.
- NIFC Suppression Costs: https://www.nifc.gov/fireInfo/fireInfo_documents/SuppCosts.pdf, access: 1 November 2020, 2019.
- NOAA Hazard Mapping System Fire and Smoke Product: <https://www.ospo.noaa.gov/Products/land/hms.html#data>, access: 24 June 2021, 2021.
- 920 Park, S. K., O'Neill, M. S., Vokonas, P. S., Sparrow, D., and Schwartz, J.: Effects of air pollution on heart rate variability: the VA normative aging study, *Environmental health perspectives*, 113, 304-309, 2004.

- Parrington, M., Palmer, P., Lewis, A., Lee, J., Rickard, A., Carlo, P. D., Taylor, J., Hopkins, J., Punjabi, S., and Oram, D.: Ozone photochemistry in boreal biomass burning plumes, *Atmospheric Chemistry and Physics*, 13, 7321-7341, 2013.
- 925 Peng, Q., Palm, B. B., Melander, K. E., Lee, B. H., Hall, S. R., Ullmann, K., Campos, T., Weinheimer, A., Apel, E., and Hornbrook, R. S.: HONO Emissions from Western US Wildfires Provide Dominant Radical Source in Fresh Wildfire Smoke, *Environmental Science & Technology*, 2020.
- Reichert, L., Andres Hernandez, M., Stöbener, D., Burkert, J., and Burrows, J.: Investigation of the effect of water complexes in the determination of peroxy radical ambient concentrations: Implications for the atmosphere, *Journal of Geophysical Research: Atmospheres*, 108, ACH 4-1-ACH 4-16, 2003.
- 930 Reid, C. E., Brauer, M., Johnston, F. H., Jerrett, M., Balmes, J. R., and Elliott, C. T.: Critical review of health impacts of wildfire smoke exposure, *Environmental health perspectives*, 124, 1334-1343, 2016.
- Roberts, J. M., Stockwell, C. E., Yokelson, R. J., de Gouw, J., Liu, Y., Selimovic, V., Koss, A. R., Sekimoto, K., Coggon, M. M., and Yuan, B.: The nitrogen budget of laboratory-simulated western US wildfires during the FIREX 2016 Fire Lab study, *Atmospheric Chemistry and Physics*, 20, 8807-8826, 2020.
- 935 Rogers, H. M., Ditto, J. C., and Gentner, D. R.: Evidence for impacts on surface-level air quality in the northeastern US from long-distance transport of smoke from North American fires during the Long Island Sound Tropospheric Ozone Study (LISTOS) 2018, *Atmospheric Chemistry and Physics*, 20, 671-671, 2020.
- Salvador, C. M. G., Tang, R., Priestley, M., Li, L., Tsiligiannis, E., Le Breton, M., Zhu, W., Zeng, L., Wang, H., and Yu, Y.: Ambient nitro-aromatic compounds—biomass burning versus secondary formation in rural China, *Atmospheric Chemistry and Physics*, 21, 1389-1406, 2021.
- 940 Saunders, S. M., Jenkin, M. E., Derwent, R., and Pilling, M.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part A): tropospheric degradation of non-aromatic volatile organic compounds, 2003.
- Sillman, S., Logan, J. A., and Wofsy, S. C.: The sensitivity of ozone to nitrogen oxides and hydrocarbons in regional ozone episodes, *Journal of Geophysical Research: Atmospheres*, 95, 1837-1851, 1990.
- 945 Silva, R. A., West, J. J., Zhang, Y., Anenberg, S. C., Lamarque, J.-F., Shindell, D. T., Collins, W. J., Dalsoren, S., Faluvegi, G., and Folberth, G.: Global premature mortality due to anthropogenic outdoor air pollution and the contribution of past climate change, *Environmental Research Letters*, 8, 034005, 2013.
- Spracklen, D. V., Mickle, L. J., Logan, J. A., Hudman, R. C., Yevich, R., Flannigan, M. D., and Westerling, A. L.: Impacts of climate change from 2000 to 2050 on wildfire activity and carbonaceous aerosol concentrations in the western United States, *Journal of Geophysical Research: Atmospheres*, 114, 2009.
- 950 Stein, A., Draxler, R. R., Rolph, G. D., Stunder, B. J., Cohen, M., and Ngan, F.: NOAA's HYSPLIT atmospheric transport and dispersion modeling system, *Bulletin of the American Meteorological Society*, 96, 2059-2077, 2015.
- Stocker, T. F., Qin, D., Plattner, G.-K., Alexander, L. V., Allen, S. K., Bindoff, N. L., Bréon, F.-M., Church, J. A., Cubasch, U., and Emori, S.: Technical summary, in: *Climate change 2013: the physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, 33-115, 2013.
- 955 Stowell, J. D., Geng, G., Saikawa, E., Chang, H. H., Fu, J., Yang, C.-E., Zhu, Q., Liu, Y., and Strickland, M. J.: Associations of wildfire smoke PM_{2.5} exposure with cardiorespiratory events in Colorado 2011–2014, *Environment international*, 133, 105151, 2019.
- 960 Tang, M., Cox, R., and Kalberer, M.: Compilation and evaluation of gas phase diffusion coefficients of reactive trace gases in the atmosphere: volume 1. Inorganic compounds, *Atmospheric Chemistry and Physics*, 14, 9233-9247, 2014.
- Travis, K. R., Jacob, D. J., Fisher, J. A., Kim, P. S., Marais, E. A., Zhu, L., Yu, K., Miller, C. C., Yantosca, R. M., and Sulprizio, M. P.: Why do models overestimate surface ozone in the Southeast United States?, *Atmospheric Chemistry and Physics*, 16, 13561-13577, 2016.

- 965 Turner, M. C., Jerrett, M., Pope III, C. A., Krewski, D., Gapstur, S. M., Diver, W. R., Beckerman, B. S., Marshall, J. D., Su, J., and Crouse, D. L.: Long-term ozone exposure and mortality in a large prospective study, *American journal of respiratory and critical care medicine*, 193, 1134-1142, 2016.
- Verma, S., Worden, J., Pierce, B., Jones, D., Al-Saadi, J., Boersma, F., and Bowman, K.: Ozone production in boreal fire smoke plume as observed by the tropospheric emission spectrometer and the ozone monitoring instrument, *AGUSM*, 2008, A31A-03, 2008.
- 970 Wentworth, G. R., Aklilu, Y.-a., Landis, M. S., and Hsu, Y.-M.: Impacts of a large boreal wildfire on ground level atmospheric concentrations of PAHs, VOCs and ozone, *Atmospheric Environment*, 178, 19-30, 2018.
- Wernis, R. A., Kreisberg, N. M., Weber, R. J., Liang, Y., Jayne, J., Hering, S., and Goldstein, A. H.: Development of an In Situ Dual-Channel Thermal Desorption Gas Chromatography Instrument for Consistent Quantification of Volatile, Intermediate Volatility and Semivolatile Organic Compounds, *Atmos. Meas. Tech. Discuss.*, 2021, 1-30, <https://doi.org/10.5194/amt-2021-156>, in review, 2021.
- 975 Wolfe, G. M., Marvin, M. R., Roberts, S. J., Travis, K. R., and Liao, J.: The framework for 0-D atmospheric modeling (F0AM) v3. 1, *Geoscientific Model Development*, 9, 3309, 2016.
- Wood, E. C., Deming, B. L., and Kundu, S.: Ethane-based chemical amplification measurement technique for atmospheric peroxy radicals, *Environmental Science & Technology Letters*, 4, 15-19, <https://doi.org/10.1021/acs.estlett.6b00438>, 2017.
- 980 Yokelson, R. J., Griffith, D. W., and Ward, D. E.: Open-path Fourier transform infrared studies of large-scale laboratory biomass fires, *Journal of Geophysical Research: Atmospheres*, 101, 21067-21080, 1996.
- Yokelson, R. J., Crouse, J., DeCarlo, P., Karl, T., Urbanski, S., Atlas, E., Campos, T., Shinozuka, Y., Kasputin, V., and Clarke, A.: Emissions from biomass burning in the Yucatan, *Atmospheric Chemistry and Physics*, 9, 5785, 2009.
- 985 York, D., Evensen, N. M., Martinez, M. L., and De Basabe Delgado, J.: Unified equations for the slope, intercept, and standard errors of the best straight line, *American Journal of Physics*, 72, 367-375, 2004.
- Yu, P., Xu, R., Abramson, M. J., Li, S., and Guo, Y.: Bushfires in Australia: a serious health emergency under climate change, *The Lancet Planetary Health*, 4, e7-e8, 2020.
- Yue, X., Mickley, L. J., Logan, J. A., and Kaplan, J. O.: Ensemble projections of wildfire activity and carbonaceous aerosol concentrations over the western United States in the mid-21st century, *Atmospheric Environment*, 77, 767-780, 2013.
- 990 Yue, X., Mickley, L. J., and Logan, J. A.: Projection of wildfire activity in southern California in the mid-twenty-first century, *Climate dynamics*, 43, 1973-1991, 2014.
- Yue, X., Mickley, L., Logan, J., Hudman, R., Martin, M. V., and Yantosca, R.: Impact of 2050 climate change on North American wildfire: consequences for ozone air quality, *Atmospheric Chemistry and Physics*, 15, 10033-10055, 2015.
- 995