

~~The nitrate radical (NO_3) oxidation of alpha-pinene is a significant source of s~~Secondary organic aerosol and organic nitrogen yields from the nitrate radical (NO_3) oxidation of alpha-pinene from various RO_2 fates ~~the under simulated ambient nighttime conditions~~

5 Kelvin H. Bates,^{1,2} Guy J.P. Burke,¹ James D. Cope,¹ and Tran B. Nguyen¹

¹Department of Environmental Toxicology, University of California Davis, Davis CA 95616, USA


²Center for the Environment, Harvard University, Cambridge, MA 02138, USA

Correspondence to: Tran B. Nguyen (tbn@ucdavis.edu)

Abstract. The reaction of α -pinene with NO_3 is an important sink of both α -pinene and NO_3 at night in regions with mixed
10 biogenic and anthropogenic emissions; however, there is debate on its importance for secondary organic aerosol (SOA) and
reactive nitrogen budgets in the atmosphere. Previous experimental studies have generally observed low or zero SOA
formation, often due to excessive $[\text{NO}_3]$ conditions. Here, we characterize the SOA and organic nitrogen formation from α -
pinene + NO_3 as a function of nitrooxy peroxy ($n\text{RO}_2$) radical fates with HO_2 , NO , NO_3 , and RO_2 in an atmospheric chamber.
We show that SOA yields are not small when the $n\text{RO}_2$ fate distribution in the chamber mimics that in the atmosphere, and the
15 formation of pinene nitrooxy hydroperoxide (PNP) and related organonitrates in the ambient can be reproduced. Nearly all
SOA from α -pinene + NO_3 chemistry derives from the $n\text{RO}_2$ + RO_2 pathway, which alone has an SOA mass yield of
~~56~~5(\pm 7)%. Molecular composition analysis shows that particulate nitrates are a large (60-70%) portion of the SOA, and that
dimer formation is the primary mechanism of SOA production from α -pinene + NO_3 under simulated nighttime conditions.
~~We estimate an average $n\text{RO}_2$ + $n\text{RO}_2 \rightarrow \text{ROOR}$ branching ratio of 18%.~~ Synergistic dimerization between $n\text{RO}_2$ and RO_2
20 derived from ozonolysis and OH oxidation also contribute to SOA formation, and should be considered in models. We report
a 58 (\pm 20)% molar yield of PNP from the $n\text{RO}_2$ + HO_2 pathway. Applying these laboratory constraints to model simulations
of summertime conditions observed in the Southeast United States (where 80% of α -pinene is lost via NO_3 oxidation, leading
to 20% $n\text{RO}_2$ + $n\text{RO}_2$ and 45% $n\text{RO}_2$ + HO_2), we estimate yields of 113% SOA and 79% particulate nitrate by mass, and 26%
PNP by mole, from α -pinene + NO_3 in the ambient. These results suggest that α -pinene + NO_3 significantly contributes to the
25 SOA budget, and likely constitutes a major removal pathway of reactive nitrogen from the nighttime boundary layer in mixed
biogenic/anthropogenic areas.

1 Introduction

Monoterpenes ($C_{10}H_{16}$) are a major class of biogenic hydrocarbons. Although less abundant than isoprene in terms of absolute emission flux of non-methane hydrocarbons, they have a disproportionate importance for the formation of



Scheme 1: The NO_3 -initiated oxidation of α -pinene produces a nitrated RO_2 radical (nRO_2). The major isomer is shown.

secondary organic aerosol (SOA), accounting for half of the total fine aerosol globally (Zhang et al., 2018), and for nitrogen oxide (NO , NO_2 , NO_3) sequestration through the formation of gaseous and particle-phase organic nitrates (Pye et al., 2015). Thus, monoterpene chemistry plays a prevailing role in aerosol-climate interactions and atmospheric air quality. Of the monoterpenes, α -pinene is the most abundant globally (Sindelarova et al., 2014). This is especially notable over boreal coniferous forests where the α -pinene emission flux alone can overtake isoprene and the combined flux of all other monoterpenes during the summer season (Hakola et al., 2003). The atmospheric abundance, fast reaction rates, and nighttime emission profile of α -pinene conspire for it to dominate the fate of the nitrate radical (NO_3) in the dark, and also to play a significant role in the daytime (Ayres et al., 2015). The reaction of α -pinene + NO_3 , thus, is one of the most prevalent reactions observed in the summer in mixed biogenic-anthropogenic sites, such as the Southeastern United States.

NO_3 reacts with α -pinene by addition to the double bond, mainly producing a nitrooxy alkyl radical in the tertiary position, which is rapidly converted to a nitrooxy peroxy radical (nRO_2) upon collisions with molecular oxygen (**Scheme 1**). Prominent field observations of a monoterpene nitrooxy hydroperoxide ($(O_2NO)ROOH$) further suggest that the nRO_2 reacts with HO_2 significantly (Nguyen et al., 2015). However, laboratory research on α -pinene + NO_3 has so far not demonstrated nitrooxy hydroperoxide formation, except for one experiment during the FIXCIT chamber studies where the pinene nitrooxy hydroperoxide (PNP) was abundantly produced from experiments using ppm-level concentrations of formaldehyde to produce dark HO_2 via the slow reaction of $CH_2O + NO_3 \rightarrow HO_2 + HNO_3$ (Nguyen et al., 2014a). Although we now appreciate that the nRO_2 from α -pinene can form monoterpene nitrooxy hydroperoxide, its absolute yield is unknown; furthermore, such high concentrations of formaldehyde in the exploratory FIXCIT experiment will affect SOA formation, and thus, aerosol yields were not extractable.

Determinations of SOA yields of α -pinene + NO_3 in the literature have also suffered from high uncertainty, although it has generally been accepted that the yields are lower than those of many other monoterpenes and, thus, often not considered for modeling SOA and organic nitrogen formation (Pye et al., 2015; Ayres et al., 2015). Fry et al. (2014) reported an SOA yield of exactly zero from this reaction from experiments using N_2O_5 as an NO_3 precursor. Hallquist et al. (1999), Moldanova et al. (2000), Bell et al. (2021), and Mutzel et al. (2021) also reported low mass yields (0.3–7%, 0.3–6.9%, 3–11%, and 5.9–

6.4% respectively, depending on precursor concentrations) with reactions performed similarly. Spittler et al. (2006) performed the experiment with slow and minute introductions of N_2O_5 and reported higher mass yields, although they depend on whether the seed particle chosen was ammonium sulfate (9%) or organic (16%). Nah et al. (2016) and Kurtén et al. (2017) also observed minimal SOA formation (mass yields of 3.6% and <1%, respectively) using $\text{NO}_2 + \text{O}_3$ as a source of NO_3 , and formaldehyde to promote HO_2 chemistry. From these observations, it is clear that chamber reaction conditions are highly influential in the observed SOA yields, and that previous studies may have each probed different $n\text{RO}_2$ fates. Thus, a systematic investigation of how $n\text{RO}_2$ fates dictate reaction outcomes will enable reconciliation of past results and accurate representation of this reaction in atmospheric models. The high initial NO_3 concentrations (tens of ppb) used in some previous studies, derived from the decomposition of N_2O_5 , will cause $n\text{RO}_2 + \text{NO}_3$ to dominate in the chamber, when it is negligible in the field ($[\text{NO}_3]$ is persistently at or below the detection limit of 1 pptv in the rural Southeast United States; Ayres et al., 2015). Kurtén Kjaergaard and coworkers further illuminated the role of alkoxy radical scission in SOA formation, and predicted low SOA yields from the α -pinene + NO_3 reaction when the RO_2 radical is reduced to RO via bimolecular reaction with NO_3 , RO_2 , and even HO_2 (Kurtén et al., 2017).

Given the high relevance of the α -pinene + NO_3 reaction, it is critical to place tighter constraints on how this reaction contributes to SOA and organic nitrogen in the ambient environment. In light of the emerging appreciation for the importance of RO_2 radical fate in designing chamber experiments (Nguyen et al, 2014a; Xu et al., 2019; Boyd et al., 2015; Teng et al., 2017; Crounse et al., 2013), we re-investigate this reaction to probe the SOA yield and organic nitrate formation from α -pinene + NO_3 from each relevant $n\text{RO}_2$ reaction channel. While a chamber experiment may never truly replicate the field, and ours certainly are no exception, the $n\text{RO}_2$ fate distribution in this work was designed to approach those expected in the ambient nighttime (Ayres et al., 2015; Romer et al., 2018), including any reaction synergies that may occur (Kenseth et al., 2018; Inomata, 2021). Finally, relatively little information is available for the $n\text{RO}_2$ compared to their hydroxylated counterparts; this work also constrains the rate coefficients and branching ratios of the α -pinene $n\text{RO}_2$ through a combination of chamber reactions and modeling. We demonstrate a new HO_2 formation route in the dark chamber that does not require carbon inputs, and thus enables SOA yields to be more accurately measured when probing the $\text{RO}_2 + \text{HO}_2$ pathway from the α -pinene + NO_3 reaction.

2 Methods

2.1 Chamber reactions

Experiments were performed in a 10 m³ FEP Teflon atmospheric chamber in the dark ~~in the dark~~ and at a temperature of ~22 °C, consistent with the average nighttime temperatures in the Southeastern United States in the summer (Hidy et al., 2014). Experiments were performed dry to reduce uncertainty from variable wall loss corrections, ~~to~~ ~~and~~ avoid rapid hydrolysis of tertiary nitrates (Vasquez et al., 2021) ~~and the effects of gas-particle partitioning of H_2O_2 and N_2O_5 , and because relative~~

95 **humidity (RH)** is not expected to influence radical reactions. Temperature and RH were monitored continuously by a
 membrane probe (Vaisala Inc.) calibrated with saturated salt solutions. NO_x and O₃ mixing ratios were quantified with
 commercial chemiluminescence (Thermo 42i) and photometric (Thermo 49i) analyzers. The chemiluminescence analyzer was
 calibrated with a NO₂ primary standard (200 ppmv, Air Liquide) diluted to desired concentrations using ultra-high-purity
 (UHP) N₂ and zeroed with UHP N₂. The photometric analyzer was cross calibrated for O₃, produced by an ozone generator
 100 (A2Ozone Inc.) from ultra zero air (Air Liquide), with a Fourier-transform infrared spectrometer (FT-IR, Shimadzu Scientific
 Inst. Inc., IR-Tracer 100) using a 10 m gas cell. α -Pinene mixing ratios were quantified with a gas chromatograph coupled to
 a flame ionization detector (GC-FID) using a PLOT-Q column (Agilent Inc.) and a custom pneumatically-injected method.
 The GC-FID was calibrated with a NIST-traceable 40 ppmv α -pinene primary standard in N₂ (Matheson Gas) diluted to several
 concentrations with UHP N₂ using calibrated mass flow controllers (SEC, Horiba Inst. Inc.). The mass flow controllers were
 105 calibrated using a primary flow calibrator (AP Buck, Inc.).

Table 1. Initial conditions for chamber experiments performed in this work. Controls to subtract SOA yields from background chemistry (not listed here) were conducted for experiments 1-11 and 17-25 using identical conditions but omitting of NO₂.

Expt	Type	pinene (ppb)	H ₂ O ₂ (ppm)	O ₃ (ppb)	NO ₂ (ppb)	NO (ppb)	N ₂ O ₅ (ppb)	Mixing time (min)	seed area ($\mu\text{m}^2/\text{cm}^3$)	seed mass ($\mu\text{g}/\text{m}^3$)
1	RO ₂ fate	27	0	53	44	0	0	60	645	23.6
2	RO ₂ fate	27	0	99	73	0	0	60	662	24.3
3	RO ₂ fate	27	4	50	149	0	0	70	645	24.3
4	RO ₂ fate	27	8	53	149	0	0	90	631	23.2
5	RO ₂ fate	27	3.5	96	225	0	0	120	700	25.8
6	RO ₂ fate	27	9	39	99	0	0	120	650	25.2
7	RO ₂ fate	27	2	43	125	0	0	60	666	25.7
8	RO ₂ fate	27	8	103	145	0	0	120	695	27.2
9	RO ₂ fate	54	2	45	120	0	0	60	615	24.5
10	RO ₂ fate	130	2	43	121	0	0	60	660	26.6
11	RO ₂ fate	27	2	98	86	0	0	120	9300	392
12	RO ₂ fate	27	0	0	0	0	31	0	7260	252
13	RO ₂ fate	27	0	0	0	0	200	0	7960	330
14	RO ₂ fate	60	0	0	11	83	110	0	5330	214
15	RO ₂ fate	60	0	0	240	110	85	0	5010	221
16	RO ₂ fate	45	0	0	565	65	80	0	4380	205
17	seed area	27	4	102	72	0	0	75	0	0
18	seed area	27	4	97	71	0	0	75	73	2.79
19	seed area	27	4	98	73	0	0	75	78	2.9
20	seed area	27	4	99	75	0	0	75	143	5.3
21	seed area	27	4	98	74	0	0	75	335	11.5
22	seed area	27	4	102	69	0	0	75	714	26.8
23	seed area	27	4	97	68	0	0	75	1460	51.1
24	seed area	42	4	99	71	0	0	75	2920	101

25	seed area	27	4	101	82	0	0	120	7400	304
26	filter	165	0	0	0	0	145	0	0	0
27	filter	100	10	520	175	0	0	60	0	0

Initial conditions for all chamber experiments are described in **Table 1**. The chamber was cleaned prior to each experiment by continually flushing with custom-filtered zero air, quantified to be below the detection limits of all available analyzers, for at least 12 hours (>7 full air exchanges). Two types of experiments were conducted: those using O₃ + NO₂ as an NO₃ source, and those using N₂O₅ as an NO₃ source. For the former, variable concentrations of NO₂ (diluted from 2042 ppm ±1% in N₂, Praxair) and O₃ were injected to initiate the formation of low levels of NO₃, according to the desired *n*RO₂ fate. H₂O₂ (50 wt.% in H₂O, Aldrich) was then injected by flowing 4 L min⁻¹ of ultra zero air through a bubbler warmed to 40 °C in a water bath. After the inorganic gas-phase reactants were introduced, the chamber was allowed to mix for 1-2 h, during which time secondary formation of NO₃ (from O₃ + NO₂) and HO₂ (from NO₃ + H₂O₂) could proceed. Seed particles were introduced to the chamber during mixing by atomizing a solution of 0.06 M ammonium sulfate ((NH₄)₂SO₄, ≥99%, Aldrich) through a ²¹⁰Po neutralizer. Lastly, liquid standards of α-pinene (Sigma Aldrich, >99%) were injected with gas-tight syringes into an airtight glass bulb, and introduced to the chamber by a 4 L min⁻¹ flow of ultra zero air. The α-pinene reacts quickly and thus was mixed rapidly from pulsed injections of high pressure ultra zero air (100 psi) for 2 min in order to initiate the reaction. The ozonolysis of α-pinene occurs concurrently with its NO₃-initiated oxidation with this experiment design. Ozonolysis of α-pinene produces OH, which reacts with H₂O₂ to be an additional dark formation source of HO₂, as well as with α-pinene itself. Control reactions were performed to accompany each experiment listed in **Table 1** that includes ozone, using the same conditions minus NO₂, in order to subtract out the SOA and other product formation from the purely ozonolytic reaction.

For experiments that used N₂O₅ as the NO₃ source, injections of α-pinene and other desired inorganic reactants, including seed particles, NO₂, and NO (200 ppm ±1% in N₂, Praxair), were conducted first. The reactions were then initiated by the rapid injection of gas-phase N₂O₅, which was previously evaporated into an evacuated 500 mL glass bulb to the desired pressure and backfilled to room pressure with N₂. N₂O₅ was synthesized according to Claflin and Ziemann (2018), verified using FT-IR, and stored in the dark at -20°C prior to use. In these experiments, the decomposition equilibria of N₂O₅ ⇌ NO₂ + NO₃ was manipulated via injections of NO₂ in order to slow NO₃ formation and thus control the *n*RO₂ fate.

During experiments, mixing ratios of oxygenated gas-phase organics were quantified with a custom-built triple-quadrupole chemical ionization mass spectrometer (CIMS) using CF₃O⁺ as the reagent ion. Instrumental details, including humidity-dependent calibration methods, have been described in detail previously (Crounse et al., 2006; St. Clair et al., 2010; Nguyen et al., 2014b; Praske et al., 2015). The CIMS detects PNP and other polar analytes predominantly without fragmentation as clusters with CF₃O⁺. Although authentic standards of PNP are not available for direct calibration in CIMS, the analytical sensitivity of synthesized organic nitrates of different carbon length and neighboring groups in the CF₃O⁺ CIMS were found to be different from each other by a factor of 20-30% (Lee et al., 2014; Teng et al., 2017); thus, sensitivity of PNP and pinene hydroxy-nitrate were assumed to be the same as isoprene hydroxynitrates with 30% uncertainty.

A scanning mobility particle sizer (SMPS), comprised of an electrostatic classifier (TSI 3080) and a condensation
140 particle counter (TSI 3772), was used to measure particle size distributions between 15 nm and 670 nm. Control experiments
monitoring the dark decay of ammonium sulfate seed aerosol concentrations in the chamber were used to determine diameter-
dependent particle wall loss rates (Schwantes et al., 2019), which were then used to correct experiment particle concentrations.
Calculations of particle mass from measured aerodynamic diameter assume a density of 1.2 g cm⁻³.

2.2 Kinetic modelling

145 We use a kinetic model (**Mech. S1**) to simulate gas-phase chemistry for each experiment in the environmental
chamber. The mechanism uses reaction parameters from the JPL Chemical Kinetics and Photochemical Data Evaluation
(Burkholder et al., 2019), and is run on Matlab (MathWorks, Inc.). We also include reactions of α -pinene with OH, O₃, and
NO₃, and isomer-specific reactions of the subsequently produced peroxy radicals with HO₂, NO, NO₃, and other RO₂ radicals
(individually represented). Product yields from each pathway, along with rates of RO₂ + RO₂ reactions, are adjusted to fit the
150 experimental data. We initialize simulations with the inorganic species listed in **Table 1**, and allow the model to run for the
allotted mixing time before instantaneously adding α -pinene. The model does not include any wall deposition of vapors or
gas-particle interactions, and is used only to estimate the concentrations of gas-phase species and the contributions of each
peroxy radical reactive pathway.

2.3 SOA composition analysis by high-resolution mass spectrometry (HRMS)

155 SOA were collected for composition analysis using Omnipore hydrophilic Teflon filters (0.2 μ m diameter pore,
Millipore Corp.) that is compatible with polar and non-polar organics. The filters were gently extracted using LC-MS grade
acetonitrile (Optima, Fisher Scientific) to mass concentrations on the order of 100 μ g/mL, depending on the experiment, by
several ultrasound pulses of duration 1 s in order to limit cavitation in the ultrasonic bath that may alter analyte compositions.
Filter extracts were directly infused into a linear-trap-quadrupole (LTQ) Orbitrap XL mass spectrometer (Thermo Instrument
160 Corp.) using positive and negative ion mode electrospray ionization (ESI) at 4 kV spray voltage and a mass resolving power
of 60,000 $m/\Delta m$ at m/z 400. An external calibration was performed in both ion modes using commercial mass standards in the
range of 100 – 2000 m/z (Pierce™ LTQ ESI Positive and Pierce™ LTQ ESI Negative calibration solutions, Fisher Scientific),
and the data were recalibrated until the mass accuracy obtained from standard solutions was < 1 ppm.

The data analysis was performed similarly to our previous works (Nguyen et al., 2010; Nguyen et al., 2011). Briefly,
165 the raw data were de-convoluted using Decon2LS (freeware from Pacific Northwest National Laboratory), and background
subtracted for peaks present in the solvent. The m/z peaks were assigned to molecular formulas (C_cH_hO_oN_n) based on a custom
Matlab script that applies Lewis and Senior rules (Kind and Fiehn, 2007) and a Kendrick Mass Defect analysis (base CH₂;
Roach et al., 2011) that have been demonstrated on SOA mixtures. The prevalent ionization mechanism for this specific analyte
mixture was found to be sodium cluster formation (M+Na⁺) in the positive mode, which occurs preferentially for carbonyls
170 (Kruve et al., 2013), and nitrate cluster formation (M+NO₃⁻) in the negative mode, which is efficient for organic nitrates,

alcohols, and other functional groups (Sisco and Forbes, 2015; Mathis and McCord, 2005). The nitrate anion, prominently detected in the mass analyzer at m/z 61.988, was not purposefully introduced but likely formed either from in-source fragmentation of organic nitrates or from HNO_3 produced during the hydrolysis of tertiary organonitrates from the aqueous LC solvents, and fortuitously acted as a reagent ion for chemical-assisted electrospray. While peak heights correlate well with concentration in direct-infusion ESI HRMS when the analyte matrix is similar (Chan et al., 2020), the correlation coefficients are unknown for each analyte in the mixture; thus, the HRMS data is qualitative. Furthermore, due to the labile $-\text{ONO}_2$ groups, the organonitrate observations from HRMS likely represent a lower limit.

3 Results and discussion

3.1 SOA and PNP yields from different $n\text{RO}_2$ fates

To investigate the dependence of aerosol and gaseous PNP yields on the $n\text{RO}_2$ reaction partner, we performed a series of environmental chamber experiments (experiments 1-16, **Table 1**) with various starting conditions designed to isolate or




Figure 1. Modelled (solid lines) and measured (points) concentrations of key reactive species and products in three example chamber experiments. Timing of chamber injections are demarcated at top. Experiment time of 0 corresponds to the beginning of α -pinene oxidation. Pie charts show the percent contribution of each α -pinene oxidative pathway (α -pinene + O_3 in red, α -pinene + OH in grey and α -pinene + NO_3 speciated by subsequent RO_2 reaction partner: orange for NO_3 , blue for RO_2 , green for HO_2).

maximize the contributions of each RO₂ reaction pathway. Example time profiles from three representative experiments are shown in **Figure 1**. *n*RO₂ + NO₃, *n*RO₂ + NO, and *n*RO₂ + RO₂ chemistry were isolated in experiments using N₂O₅ as the NO₃ source. Using excess N₂O₅ as a source causes a rapid initial spike of NO₃, making the *n*RO₂ + NO₃ pathway **dominant** **easy-to** **isolate** in Experiment 13 (**Fig. 1**, left). The *n*RO₂ + NO pathway is similarly easy to isolate with excess NO added before N₂O₅ injection, as in Experiment 15 (**Fig. S1**). *n*RO₂ + *n*RO₂ chemistry can be maximized by injecting excess α-pinene prior to N₂O₅ addition; the NO₃ is thus dominantly consumed by reaction with α-pinene, leaving the subsequently produced *n*RO₂ to react with each other, as in Experiment 26 (**Fig. 1**, middle), for which we calculate that 79% of *n*RO₂ reacted with other *n*RO₂.

*n*RO₂ + HO₂ chemistry is more difficult to isolate due to the scarcity of clean HO_x sources in the dark. Here, we describe a method **whereby reaction with HO₂ represents a majority of the *n*RO₂ fate** **that can achieve a domination of *n*RO₂ + HO₂ chemistry** without additional carbon inputs, initiated by the slow production of NO₃ in situ via NO₂ + O₃, which will introduce three reaction partners for α-pinene (NO₃, O₃, OH). Alkene ozonolysis has been used previously to produce dark OH in chamber experiments (Xu et al., 2019); Leveraging the ozonolysis of α-pinene already occurring in the chamber, we can amplify HO₂ production by injecting excess H₂O₂ prior to α-pinene to scavenge OH. Thus, the OH + H₂O₂ → HO₂ + H₂O reaction simultaneously produces HO₂ and suppresses the side chemistry of α-pinene + OH. This HO₂ source is relevant to the nighttime atmosphere, as ozonolysis ~~is~~ always occurring with NO₃ reaction in the ambient due to the major source chemistry from NO₂ + O₃.

The added H₂O₂ provides an additional benefit – its reaction with NO₃ is a source of HO₂ prior to α-pinene injection (**Fig. S12**). The H₂O₂ + NO₃ → HO₂ + HNO₃ reaction has been estimated by Burrows, Tyndall, and Moortgat (1985) to have an upper limit of <2×10⁻¹⁵ cm³ molec⁻¹ s⁻¹ – too slow for atmospheric relevance, but sufficient to produce significant HO₂ in chambers when H₂O₂ is in excess. The rate coefficient of this reaction was further constrained from its upper limit based on the ratio of *k*_{CH₂O+OH}/*k*_{CH₂O+NO₃} (Burkholder et al., 2019), resulting in a rate coefficient of 1.1×10⁻¹⁶ cm³ molec⁻¹ s⁻¹ used in our simulations. The formation of PNP was not adequately reproduced if HO₂ is assume to originate from H₂O₂ + OH alone, i.e., omitting the H₂O₂ + NO₃ reaction in the kinetic model; the yield of PNP would need to be unphysically high to reconcile the difference.

The α-pinene + NO₃ reaction is modeled to be the major α-pinene sink in experiments initiated by NO₂ + O₃ (e.g., 73% in Experiment 22, **Fig.**




Figure 2. (a) α-pinene loss and (b) SOA formation in ‘simulated nighttime’ Experiment 22 (blue) and the ozonolysis control experiment (red). The difference (green) shows the contribution from α-pinene + NO₃.

215 **2, Table 2).** Even so, the background chemistry from α -pinene + O_3 and α -pinene + OH (from the ozone control experiments without NO_2) are substantial sources of SOA and require careful subtraction (**Fig. 2**). Caveats to this approach include: (1) in the controls, ozone and OH are larger sinks for α -pinene due to a lack of competition from NO_3 ; thus, a larger fraction of α -pinene produces SOA from ozonolysis in the control compared to the experiment and the subsequent subtraction obtains a lower-limit SOA yield; (2) synergistic reactions between RO_2 intermediates from the different oxidation pathways are not possible to isolate, and contributed roughly 20% to the analytical signal from the SOA composition analysis (**Section 3.3**); however, it is now appreciated that these synergies also occur in the ambient and are not realistic to ignore in laboratory and modeling studies (Kenseth et al., 2018; Inomata, 2021).




Figure 3. Negative ion CIMS mass spectrum after 1 h of oxidation under ‘simulated nighttime’ conditions, showing PNP, PHN, and another organic nitrogen product at m/z 314. Pinonaldehyde is observed at m/z 253, and pinonic acid at m/z 269.

230 These ‘simulated nighttime’ experiments (e.g. Experiment 20, **Fig. 1**, right) are termed as such because they provide an atmospherically relevant balance of reactive pathways ~~in which fractional contributions are comparable to those on summer nights in the Southeast United States~~ (60–80% α -pinene + NO_3 , 20–40% α -pinene + O_3 ; 30–50% nRO_2 + HO_2 , 30–50% nRO_2 + RO_2) comparable to those on summer nights in the Southeast United States (based on measurements reported in Ayres et al., 2015 and Romer et al., 2018; see Section 4). This is corroborated by gas phase data from the CIMS (**Fig. 3**), which show that NO_3 -initiated products are the major volatile products formed in the simulated nighttime experiments. Furthermore, these experiments are able to reproduce gaseous organonitrates observed during the Southern Oxidant and Aerosol Study (SOAS; Carlton et al., 2018), including PNP – a major monoterpene semivolatile at the SOAS site in rural Alabama – and an unknown compound at m/z 314 that has also been observed in the ambient with similar time profiles to PNP (**Fig. S32**).

245 Product yield results from different nRO_2 reaction pathways are shown in **Table 2**, along with the modeled contributions of each nRO_2 reaction pathway. We observe low SOA formation in experiments maximizing nRO_2 + NO_3 (3(\pm 3)% mass yield, Experiment 13) and nRO_2 + NO (12(\pm 7)%, Experiment 15). This is consistent with previous observations of 0–16% SOA mass yields from N_2O_5 -initiated experiments (in which the nRO_2 + NO_3 pathway tends to dominate), where the larger yields were obtained only with slow continuous introductions of N_2O_5 (Bell et al., 2021; Fry et al., 2014; Hallquist et al., 1999; Moldanova et al., 2000; Mutzel et al., 2021; Spittler et al., 2006). We also observe no correlation between SOA yields in ‘simulated nighttime’ experiments and the fractional contribution of the nRO_2 + HO_2 pathway (**Fig. S43**), suggesting no SOA production from this pathway, consistent with Kurten et al. (2017) and Nah et al. (2016). This result conforms to

expectations as $n\text{RO}_2 + \text{HO}_2$ produces either the RO radical (Iyer et al., 2018) leading to the volatile pinonaldehyde (Kurtén et al., 2017) or the hydroperoxide, which has high enough volatility to be observed by the CIMS. We do, however, observe high yields of SOA in experiments targeting $n\text{RO}_2 + n\text{RO}_2$ chemistry (up to 82% mass yield, Experiment 26). This includes both N_2O_5 -initiated experiments with excess α -pinene and $\text{O}_3 + \text{NO}_2 + \text{H}_2\text{O}_2$ -initiated ‘simulated nighttime’ experiments, during which we measured SOA mass yields of 19–55%.

Table 2. Peroxy radical pathway contributions and measured experimental outcomes for experiments performed in this work.

Expt	Modelled								Measured	
	$X(\text{HO}_2)_i$ (ppt)	$X(\text{NO}_3)_i$ (ppt)	$X(\text{RO}_2)_{\text{max}}$ (ppb)	pinene + NO_3 (%)	$n\text{RO}_2$ + NO_3 (%)	$n\text{RO}_2$ + NO (%)	$n\text{RO}_2$ + HO_2 (%)	$n\text{RO}_2$ + RO_2 (%)	SOA mass yield (%) ^a	PNP molar yield (%)
1	0	259	2.9	60	11	0	0	89	26.9	0
2	0	542	4.7	85	31	0	0	69	28.9	0
3	73	160	3.1	95	19	0	37	44	18.3	17
4	121	114	1.9	89	10	0	65	25	9.2	31
5	120	303	4.0	99	37	0	41	21	2.5	19
6	97	56	0.94	74	5	0	70	25	5.1	30
7	34	152	3.3	90	17	0	18	65	22	9
8	147	151	2.0	89	11	0	66	23	10.5	32
9	37	159	4.0	69	6	0	41	53	22.1	17
10	35	151	4.6	46	2	0	22	76	23.2	9
11	88	277	3.1	81	16	0	23	61	26.4	10
12	0	0	6.0	100	41	0	0	59	67	0
13	0	0	7.4	100	95	0	0	5	3	0
14	0	0	5.6	100	8	53	0	39	43	0
15	0	0	0.01	100	0	100	0	0	12	0
16	0	0	3.0	100	7	59	0	33	33	0
17	113	194	2.1	75	10	0	45	45	0	20
18	111	186	2.0	75	10	0	45	45	4	21
19	112	191	2.1	75	10	0	44	45	5.7	21
20	114	195	2.1	76	11	0	44	45	21	26
21	113	192	2.1	76	10	0	45	45	21.9	25
22	112	188	2.0	73	10	0	45	45	19.9	25
23	110	181	1.9	71	9	0	45	45	19	21
24	112	189	2.3	60	6	0	46	48	21.6	20
25	110	178	2.0	75	10	0	44	46	23.5	18
26	0	0	23	100	21	0	0	79	82	0
27	295	504	5.5	64	12	0	41	47	55	^b

^aozonolysis-corrected (see Table S1); ^bnot measured

Across all experiments, the modeled fractional contribution of the $n\text{RO}_2 + \text{RO}_2$ pathway correlates strongly with SOA yield (Fig. 4a). An error-weighted linear regression (York et al., 2004) of

260 ozonolysis-corrected SOA yield against the modeled $n\text{RO}_2 + \text{RO}_2$ contribution for seeded experiments suggests a $67.58(\pm 6)\%$ mass yield ($R^2 = 0.54$) from this pathway alone. With such a large number of experiments spanning a range of different pathway contributions, however, we can improve upon this simple fit by performing a

265 multivariate linear regression to estimate SOA yields from all pathways, with coefficients limited to be ≥ 0 . This results in a predicted SOA mass yield from the $n\text{RO}_2 + \text{RO}_2$ pathway identical to results from the simple linear estimate within error: $56.65(\pm 7.9)\%$. SOA mass yields from the other pathways are not significantly

270 different from zero: $13.1(\pm 11)\%$ for $n\text{RO}_2 + \text{NO}$, $4.3(\pm 8.7)\%$ from $n\text{RO}_2 + \text{NO}_3$, $0(\pm 5)\%$ from $n\text{RO}_2 + \text{HO}_2$. Results from this regression analysis, which gives an R^2 of 0.6662 against measured SOA yields, are shown in Figure 4b. Repeating the regression analysis for all α -pinene sinks in the ‘simulated nighttime’ experiments, including the

275 ozonolysis and OH pathways as additional independent variables, also gives coefficients indistinguishable within error (0% for α -pinene + OH, 1.58% for α -pinene + O_3).

By inherently treating individual reactive pathways as independent variables, these regression analyses cannot separate the possible contributions of interactions between multiple pathways, e.g. from synergistic dimerization between $n\text{RO}_2$ and ozonolysis-derived RO_2 (see Section 3.3). The reported coefficients may therefore misrepresent what each pathway

280 on its own would contribute to SOA formation without such synergy. However, because the analysis was performed on experiments predominantly designed to replicate the reactive pathways experienced by α -pinene under ambient nighttime conditions, we expect the coefficients reported above to represent reasonable estimates of each pathway’s contribution to SOA formation in the nighttime atmosphere, including any synergistic reactive pathways that occur in ambient conditions. Conditions which deviate substantially from the nighttime atmosphere and therefore lack the same synergistic reactions (e.g.

285 experiments isolating single reactive pathways) may measure different SOA yields. Among the experiments performed here, those with high $n\text{RO}_2 + \text{RO}_2$ contributions without any ozonolysis (Experiments 12, 14, and 26) exhibited some of the highest measured SOA yields – higher than the regression model would predict – suggesting perhaps that the $n\text{RO}_2 + n\text{RO}_2$ pathway on its own results in even higher SOA yields while $n\text{RO}_2 + \text{other RO}_2$ pathways have lower yields. Without knowing the relative rates of various $n\text{RO}_2 + \text{RO}_2$ reactions, though, we cannot sufficiently constrain these differences, and




Figure 4. Measured (non-ozonolysis-corrected) SOA mass yields from each experiment as a function of the modelled contribution of $n\text{RO}_2 + n\text{RO}_2$ (color top) and of the predicted SOA yield from the sum of each pathway contribution (x axis bottom). Dotted lines denote error-weighted ordinary least squares fits with intercepts constrained to zero ($R^2 = 0.54$ top, 0.62 bottom). Error bars denote uncertainty from SMPS measurements.

290 additional regression analyses including interaction terms between the reactive pathways did not yield statistically robust results.

These fits regression analyses also intrinsically depend on kinetic model parameters such as bimolecular RO₂ reaction rates, some of which are uncertain. While we are unable to fully quantify these rates, we find that certain ratios between rates are constrained by our experimental outcomes. For example, the negligible SOA yield in the high-NO₃ Experiment 13 suggests
295 that *n*RO₂ + RO₂ chemistry cannot play a major role in that experiment. This can only be achieved if the bulk rate coefficient for *n*RO₂ + RO₂ reactions (*k*_{RO₂+RO₂}) is approximately two orders of magnitude slower than that of *n*RO₂ + NO₃ reactions (*k*_{RO₂+NO₃}). We therefore set the bulk *k*_{RO₂+RO₂} to 1×10⁻¹³ cm³ molecule⁻¹ s⁻¹ (with the corresponding bulk *k*_{RO₂+NO₃} equal to 1×10⁻¹¹ cm³ molecule⁻¹ s⁻¹ (a value higher than most reported *k*_{RO₂+NO₃}, although few measurements exist for these reactions). The PNP yields in ‘simulated nighttime’ experiments also provide a constraint on the relative rates of the *n*RO₂ + RO₂ and *n*RO₂ +
300 HO₂ reactions. RO₂ + HO₂ rate coefficients are, in general, are better constrained than those of RO₂ + RO₂ or RO₂ + NO₃ reactions; we use the parameterization in Wennberg et al. (2018), which gives 1.7×10⁻¹¹ cm³ molecule⁻¹ s⁻¹, similar to the measured value for pinene-derived radicals of 2.1×10⁻¹¹ cm³ molecule⁻¹ s⁻¹ from Boyd et al. (2003). Increasing *k*_{RO₂+RO₂} in our simulations causes a decrease in the contribution of RO₂ + HO₂ chemistry, but this decrease is only plausible until the measured PNP yield is 100%, which occurs at a *k*_{RO₂+RO₂} value slightly above 1×10⁻¹² cm³ molecule⁻¹ s⁻¹. The substantial yield of
305 pinonaldehyde instead of PNP from *n*RO₂ + HO₂ calculated by Kurtén et al. (2017) suggests that *k*_{RO₂+RO₂} must be well below 1×10⁻¹² cm³ molecule⁻¹ s⁻¹ to keep the PNP yield well below 100%). ~~but we note that this rate likely represents a weighted mean of different rates for the different *n*RO₂ isomers (Orlando and Tyndall, 2012), and that it remains uncertain. Substantially higher *k*_{RO₂+RO₂} on the order of 10⁻¹⁰ cm³ molecule⁻¹ s⁻¹, which has been suggested elsewhere for 1,3,5 trimethylbenzenepinene (Berndt et al., 2018), would necessitate a much higher *k*_{RO₂+HO₂} rate in ‘simulated nighttime’ experiments, a value that is better~~
310 constrained (Boyd et al., 2003).

Based on these constraints, a value of 1×10⁻¹³ cm³ molecule⁻¹ s⁻¹ represents our optimal fit and a value of 1×10⁻¹² cm³ molecule⁻¹ s⁻¹ represents an upper limit for the bulk *k*_{RO₂+RO₂} of α-pinene *n*RO₂. A higher value would require implausibly high *k*_{RO₂+NO₃} values in order to maintain dominance of the *n*RO₂ + NO₃ pathway in Experiment 13, and would require implausibly high *k*_{RO₂+HO₂} values in order to maintain a measured PNP branching ratio below the maximum of 100%. Our best estimate is
315 reasonably within the broad range of measured *k*_{RO₂+RO₂} for other peroxy radicals, particularly given that rate coefficients tend to decrease with size and degree of substitution at the radical site (Orlando and Tyndall, 2012). It is smaller, however, than some recent measurements of the *k*_{RO₂+RO₂} of peroxy radicals derived from α-pinene ozonolysis (1×10⁻¹² to 1×10⁻¹¹ cm³ molecule⁻¹ s⁻¹), highlighting the uncertainty in reaction rates of this type (Zhao et al., 2018; Berndt et al., 2018b). A slower bulk rate of 1×10⁻¹⁴ cm³ molecule⁻¹ s⁻¹, which would still be consistent with our constraints on relative rate ratios, would scale
320 the contribution of *n*RO₂ + RO₂ chemistry in the ‘simulated nighttime’ experiments by a factor of 0.67, which in turn would require scaling SOA yields from this pathway up by 50%. A faster bulk *k*_{RO₂+RO₂} of 1×10⁻¹² cm³ molecule⁻¹ s⁻¹ would scale the *n*RO₂ + RO₂ contribution in the same experiments by a factor of 1.16, which in turn would require scaling the *n*RO₂ + RO₂

SOA yield down by 14%. Despite our constraints, the bulk $k_{\text{RO}_2+\text{RO}_2}$ remains highly uncertain, and is also likely to differ between $n\text{RO}_2$ isomers for self- and cross-reactions (Orlando and Tyndall, 2012).

To constrain the sequestration of reactive nitrogen under ambient conditions from the α -pinene + NO_3 reaction, we also quantify PNP yields in our simulated nighttime experiments. **Figure 5** (top) shows measured PNP molar yields as a function of the $n\text{RO}_2$ + HO_2 fate contribution. Regression analysis (York et al., 2004) suggests a branching ratio for PNP formation of $58(\pm 2)\%$ from $n\text{RO}_2$ + HO_2 ($R^2 = 0.88$), which we implement in our kinetic model (**Fig. 5**, bottom). Including the uncertainty in the CIMS calibration increases the uncertainty bounds to $58(\pm 20)\%$. The bulk yield likely represents a combination of different branching ratios to PNP formation from the different $n\text{RO}_2$ isomers. We hypothesize that the secondary (minor) $n\text{RO}_2$ produces exclusively PNP in its reaction with HO_2 , which would imply a branching ratio of 37% for PNP formation from the tertiary (major) $n\text{RO}_2$ + HO_2 , assuming an initial branching of 65:35 major:minor isomers from α -pinene + NO_3 (Jenkin et al., 1997; Saunders et al., 2003). Our measured bulk PNP yield is somewhat higher than the 30% estimated in the FIXCIT studies (Kurtén et al., 2017), which used similar instrumentation but slightly different chamber conditions (formaldehyde instead of H_2O_2 and slow addition of α -pinene to minimize RO_2 + RO_2 chemistry). Some of this discrepancy might be explained by an overestimate of the contribution of $n\text{RO}_2$ + HO_2 chemistry in the FIXCIT experiment, where they assumed that all NO_3 produced by O_3 + NO_2 reacted with α -pinene and that all $n\text{RO}_2$ reacted with HO_2 (Kurtén et al., 2017).

We also detect α -pinene hydroxy nitrate (PHN), a product of $n\text{RO}_2$ + $n\text{RO}_2$ chemistry, and α -pinene dinitrate (PDN), a product of $n\text{RO}_2$ + NO or $n\text{RO}_2$ + NO_3 chemistry, by $\text{CF}_3\text{O}^\bullet$ CIMS. Based on similar regression analyses of observations and modeling results (**Fig. S54**), we estimate a bulk PHN yield from $n\text{RO}_2$ + $n\text{RO}_2$ of $11.7(\pm 3.3)\%$. However, because the major $n\text{RO}_2$ isomer (with the tertiary peroxy radical) is unable to donate an α -hydrogen for PHN formation, a bulk yield of 11.8% represents a 34% branching ratio from the minor (secondary) $n\text{RO}_2$ isomer (assuming an initial ratio of 65:35




Figure 5. Measured α -pinene nitrooxy-hydroperoxide (PNP) yields as a function of the modelled contribution of the $n\text{RO}_2$ + HO_2 pathway (top), and measured PNP at the end of each experiment as a function of the predicted PNP based on the simple kinetic model. Dotted lines denote error-weighted ordinary least squares fits with intercepts constrained to zero ($R^2 = 0.88$ top, 0.74 bottom; slope = 0.58 top, 1.0 bottom). Error bars denote uncertainty from CIMS measurements.

major:minor isomers from α -pinene + NO_3 ; Jenkin et al., 1997; Saunders et al., 2003). While observed PDN correlates with $n\text{RO}_2 + \text{NO}_3$ chemistry, its low signal and uncertain CIMS sensitivity makes quantification difficult.

3.2 SOA yield dependence on seed surface area

SOA from gas-particle partitioning tends to exhibit a strong dependence on seed aerosol surface area, because of competition between particle surfaces and chamber walls (Schwantes et al., 2019; Zhang et al., 2014; Zhang et al., 2015). This effect can cause an underestimation of SOA yields in atmospheric chambers if insufficient seed aerosol is used. To quantify this effect and ensure that RO_2 fate experiments (#1-16) were initiated with sufficient seed aerosol, we performed an additional set of experiments (#17-25) under ‘simulated nighttime’ conditions to investigate




Figure 6. Measured ozonolysis-corrected SOA yields from seed area experiments (#17-25) plotted against initial seed surface area. Error bars denote experimental uncertainty from SMPS measurements.

the dependence of SOA yield on seed surface area (**Fig. 6**). No SOA formation was observed in the absence of seed aerosol, indicating that α -pinene + NO_3 with simulated nighttime RO_2 fates does not nucleate on its own at the low precursor concentrations used in these experiments (and the even lower concentrations present in the atmosphere). This is in contrast to our ozonolysis control experiments where nucleation was observed, in agreement with other accounts (Burkholder et al., 2007; Hoppel et al., 2001; Inomata, 2021; Takeuchi et al., 2019). The lack of nucleation is consistent with the expectation that the α -pinene + NO_3 reaction dominates the nighttime experiments over ozonolysis. SOA yields reached a maximum when initial seed surface area reached $143 \mu\text{m}^2 \text{cm}^{-3}$ (equivalent under experimental conditions to $5.3 \mu\text{g m}^{-3}$). This threshold is at least an order of magnitude lower than those measured by Schwantes et al. (2019) for SOA from isoprene OH oxidation under high-NO conditions and by Zhang et al. (2014) for SOA from toluene oxidation. Although vapor wall losses may still cause a decrease in measured SOA yields regardless of seed surface area (Krechmer et al., 2020), we assume here that these effects are minimal at the high seed surface areas ($> 600 \mu\text{m}^2 \text{cm}^{-3}$) used in our experiments to investigate RO_2 fate. ~~suggesting much lower volatility for the α -pinene + NO_3 products.~~ However, true SOA yields may be higher than reported if vapor wall losses are considered.

Filters collected following SOA formation in Experiments 26 (high $n\text{RO}_2 + n\text{RO}_2$), 27 (simulated nighttime), and the ozonolysis control (without NO_2) for Experiment 27 were analyzed by HRMS to determine the SOA molecular composition (**Figure 7**). As expected, the mass spectrum from the 'simulated nighttime' experiment (#27, **Fig. 7B**) exhibit substantial overlap with those from the N_2O_5 -initiated experiment in which the $n\text{RO}_2 + n\text{RO}_2$ pathway dominated (#26, **Fig. 7D**),
390 corroborating the results in **Section 3.1** that suggested the $n\text{RO}_2 + n\text{RO}_2$ pathway is responsible for most SOA formation from α -pinene + NO_3 chemistry. The SOA from $n\text{RO}_2 + n\text{RO}_2$ chemistry is dominated by compounds containing 17-20 carbon atoms, suggesting that dimerization is the primary mechanism by which $n\text{RO}_2 + n\text{RO}_2$ reactions lead to SOA formation.

Masses in the 'simulated nighttime' SOA that are also observed in the $n\text{RO}_2 + n\text{RO}_2$ control experiment (#26) account for 39% of the negative mode peak signal and 29% of the positive mode peak signal in the SOA from the 'simulated nighttime' experiment. Negative mode may overestimate the contributions from dinitrates dimers from $n\text{RO}_2 + n\text{RO}_2$, whereas positive mode may underestimate them. In the positive mode, it was also found that 34% of the overall peaks signal in the 'simulated nighttime' experiment can also be found observed from ozonolysis chemistry (including OH reaction), 14% of peak signal is are found in both ozonolysis and $n\text{RO}_2 + n\text{RO}_2$ chemistry, and 24% of overall peaks signal is are completely unique to the nighttime chemistry. SOA from α -pinene ozonolysis is dominated by oxygenated monomers in the negative mode and lacks
400 organonitrate functionalities, making it easy to distinguish from the mass spectrum of NO_3 -oxidized SOA. The peaks from α -pinene ozonolysis can thus be subtracted off from the 'simulated nighttime' SOA mass spectrum, leaving a spectrum attributable to α -pinene + NO_3 under 'simulated nighttime' conditions (**Fig. 7C**).





Figure 7. Mass spectra of filter samples from Experiments 26 (D) and 27 (B), along with the ozonolysis-only control for Experiment 27 (A) and the signal difference between B and A (C). Peaks **uniquely** attributable to α -pinene ozonolysis alone are shown in red, while those **uniquely** attributable to α -pinene + NO_3 followed by RO_2 + RO_2 chemistry are shown in blue with all other peaks in black. Pie charts at right denote the fraction of α -pinene reacting via each pathway for a given experiment. Formation mechanisms for the key dimer species (a-d) identified in the bottom panels are shown in Schemes 2-3.

The signal-weighted composition of SOA from the ‘simulated nighttime’ reaction (**Fig. 7B**), which includes background ozonolysis, is 60–70% nitrogen-containing organics with 1–3 nitrate groups, as determined in both negative and positive mode (where organic nitrogen species are not enhanced). Of the organic nitrogen, it appears that 2N species are the most abundant and comprise ~30% of the SOA on average. This suggests dimerization from nRO_2 + nRO_2 to form C_{14-20} compounds is highly important to SOA formation. Previous analyses of the molecular composition of SOA from α -pinene + NO_3 have also found a dominant contribution of dimers in the particle phase (Bell et al., 2021; Takeuchi et al., 2019). The 0N, 1N, and 3N species may also be dimers (4N were not highly observed), as each of the expected RO_2 s from all channels can have 0–2 N (**Fig. 8**). There is also a clear population of trimers from α -pinene SOA that is observed here (**Fig. 7**, m/z centered around 700, with associated carbons C_{21} - C_{30}) and elsewhere (Claflin and Ziemann, 2018; Romonosky et al., 2017). **This that**

may suggest RO₂s can react with neutral compounds to propagate RO₂ radicals that then terminate with another RO₂·, or may alternatively be attributable to second-generation accretion reactions of dimers (especially those still containing a double bond,



Scheme 2. Proposed mechanisms for the formation of stable products (red) from α -pinene + NO₃ oxidation, including nitrooxy-hydroperoxide (PNP), α -pinene hydroxynitrate (PHN), pinene dinitrate (PDN), the oxidation product observed by CIMS at m/z 314, and pinonaldehyde, as well as α -pinene-derived peroxy radicals (n RO₂ α - ϕ , blue) that may contribute to particle-phase dimers (**Fig. 8**) following the dominant secondary addition of NO₃ to α -pinene. “X” stands in for any of RO₂, HO₂, NO, or NO₃ as bimolecular reaction partners.

which would react rapidly with NO₃) with RO₂ monomers.

The data show that SOA formation from α -pinene + NO₃ originates from a varied cohort of distinct n RO₂ isomers. In **Scheme 2** we suggest multiple pathways involving bimolecular reactions, β -scission, and intramolecular isomerization that can produce different C₇-C₁₀ n RO₂ isomers following the initial reaction of α -pinene with NO₃. These pathways are consistent with RO₂ chemistry previously suggested from theoretical (Kurtén et al., 2017) or lab studies (Xu et al., 2019) of similar systems. Some reactions of n RO₂ with NO, NO₃, or HO₂ shown in **Scheme 2** propagate n RO₂ radicals due to unimolecular decomposition rather than forming stable products; this implies that dimer-containing SOA from n RO₂ can originate from all n RO₂ fate pathways. **Figure 8C** shows the most abundantly observed dimers in the SOA from n RO₂ + n RO₂ chemistry (labeled peaks a-d in **Fig. 7D**), which include C₂₀H₃₀O₄ (α - α , **Fig. 7D.a**; presumably following the loss of 2 HONO from dimer C₂₀H₃₂N₂O₈ in the ESI source), C₁₇H₂₆N₂O₁₀ (δ - α , **Fig. 7D.b**), C₂₀H₃₂N₂O₁₀ (β - β , **Fig. 7D.c**), and C₂₀H₃₁N₃O₁₃ (ϵ - α , **Fig. 7D.d**). Bell et al. (2021) also observed C₂₀H₃₂N₂O₈ and C₂₀H₃₂N₂O₁₀ among the most abundant compounds in SOA from α -pinene + NO₃ chemistry in which n RO₂ + n RO₂ reactions were thought to dominate.

Dimers from the ozonolysis control experiments are notably absent from the ‘simulated nighttime’ experiment. This is likely because the peroxy radicals from α -pinene ozonolysis, which would have reacted with each other to dimerize in the
435 ozonolysis-only experiment, instead react with the more abundant $n\text{RO}_2$ radicals from α -pinene + NO_3 in the ‘simulated nighttime’ experiment. This dimerization from cross-reactions of peroxy radicals from both oxidation pathways is responsible for the $\sim 24\%$ additional HRMS signals in the α -pinene + NO_3 SOA spectrum (**Fig. 7C**, black peaks) but not in the $n\text{RO}_2$ + $n\text{RO}_2$ SOA spectrum (**Fig. 7D**). **Figure 8D** show how RO_2 s formed from ozonolysis (**i-iii**; Iyer et al., 2021), initially via the vinylhydroperoxide channel, and OH reactions (**1-2**; Xu et al., 2019) can couple with $n\text{RO}_2$ s to produce synergistic dimers
440 during nighttime oxidation of α -pinene. This type of synergy is missing when lab studies cleanly isolate reaction pathways, but may be an important part of SOA formation in the ambient as reactions occur simultaneously (**Berndt et al., 2018a & b**; **Kenseth et al., 2018**; **Zhao et al., 2018**; **McFiggans et al., 2019**; **Inomata, 2021** **Teng et al., 2017**). These results underscore the importance of conducting chamber experiments under atmospherically relevant conditions.

While we do not seek to represent each individual $n\text{RO}_2$ and dimerization pathway in our kinetic model, we adjust
445 the product yields from the first-generation $n\text{RO}_2$ + $n\text{RO}_2$ reactions to provide a reasonable estimate of dimer formation. The observed combined positive and negative mode peak signal attributable to C_{14-20} compounds (presumed dimers) in $n\text{RO}_2$ + $n\text{RO}_2$ SOA (Experiment 27, **Fig. 6D**) constitutes 40-60% of the total signal. To produce this 40-60% of the ~~56~~65% SOA mass yield estimated from $n\text{RO}_2$ + $n\text{RO}_2$ chemistry in **Section 3.1** using dimers, assuming a mean dimer molar mass of 485 g mol^{-1} (the average of the positive mode and negative mode peak-signal-weighted mean), requires a molar yield of 0.168 dimer
450 molecules from the $n\text{RO}_2$ + $n\text{RO}_2$ reaction (0.089 per $n\text{RO}_2$). This bulk effective yield represents an average across the various $n\text{RO}_2$ + $n\text{RO}_2$ isomer permutations; branching ratios from individual $n\text{RO}_2$ + $n\text{RO}_2$ isomer reactions may differ substantially, but in our kinetic model we simply apply the bulk effective yield to all peroxy radical self- and cross-reactions and scale down the pinonaldehyde-forming pathway accordingly. This 168% dimer yield is somewhat higher than has previously been measured in other systems (Orlando and Tyndall, 2012), including from α -pinene ozonolysis (Zhao et al., 2018), but fits with
455 recent evidence suggesting that increased molecular size and functionalization can significantly increase dimer branching ratios (Molteni et al., 2019; Berndt et al., 2018). However, our estimated dimer yield is highly uncertain for a wide range of reasons, including (a) variations in compound-specific sensitivities within the ESI HRMS, which may accentuate the dimer contribution as ESI efficiency has been shown to increase with molecular size (Kenseth et al., 2020); (b) sensitivity to the uncertain estimates of RO_2 fate, which depend on poorly-constrained RO_2 + RO_2 and RO_2 + NO_3 rates (see **Section 3.1**); (c) the
460 possibility of particle-phase reactions, or reactions during filter extraction, which may form or break dimers; and (d) lack of consideration of trimers, which account for 14% of overall peak intensity and may form from dimers, but are also subject to even greater uncertainty from ESI sensitivity. For all these reasons, we caution that the dimer yield estimated here is only a best guess meant to fit our kinetic model, and that further investigation is needed with more quantitative methodology.




Figure 8. Structures and molecular formulas for (A) $n\text{RO}_2$ from NO_3 -initiated chemistry (Scheme 2), (B) RO_2 from OH or O_3 chemistry (Xu et al., 2019; Iyer et al., 2021), (C) prominent dimers observed from $n\text{RO}_2$ self/cross reactions, and (D) dimers from synergistic $n\text{RO}_2 + \text{RO}_2$ couplings observed in the ‘simulated nighttime’ reactions. Dimers with chemical formulas written in pink are uniquely observed in the ‘simulated nighttime’ experiments; those marked with a * have moderate-to-high signal ($40 > \text{S/N}$).

465 4 Conclusions and atmospheric relevance

Contrary to previous chamber experimental results, we have shown here that the reaction of α -pinene with NO_3 can form high mass yields of SOA ($>21\%$, at ~ 30 ppbv of α -pinene) under RO_2 fate branching ratios designed to mimic those of summer nights in the Southeast United States. Much of this SOA originates from the self- and cross-reactions of nitrated peroxy radicals ($n\text{RO}_2 + n\text{RO}_2$); we estimate that this pathway alone has an SOA mass yield of ~~56~~⁶⁵%, with an associated dimer branching ratio of ~ 16 ⁸%, while SOA formation from other $n\text{RO}_2$ pathways is negligible within uncertainty. This hypothesis is also supported by the dominance of dimers in the ozonolysis-subtracted mass spectra of SOA collected from ‘simulated nighttime’ experiments, the diversity of which suggests that many different peroxy radical rearrangements and dimerization channels contribute to SOA formation from $n\text{RO}_2 + n\text{RO}_2$ chemistry.

The magnitude of α -pinene concentrations in the Southeast United States will be lower than that used in our chamber; thus, we model the α -pinene + NO_3 chemistry at nighttime concentrations measured during the SOAS campaign in rural Alabama in the summer (1 ppb α -pinene, 1 ppb other terpenes, 20 ppb ozone, 1 ppt NO_3 , and 2 ppt HO_2 ; Ayres et al., 2015;

Romer et al., 2018) and assuming steady-state for $n\text{RO}_2$. At atmospheric concentrations, our kinetic model predicts that 80% of α -pinene reacts with NO_3 . Of the $n\text{RO}_2$ formed, approximately 20% is predicted to react with α -pinene-derived $n\text{RO}_2$ or o_3RO_2 at $k_{\text{RO}_2+\text{RO}_2}$ to $1 \times 10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. Combined with our measured 5665% SOA mass yield from $n\text{RO}_2 + \text{RO}_2$, this suggests an overall SOA mass yield of 113% from α -pinene + NO_3 chemistry under the ambient conditions of SOAS. This SOA yield corresponds to a 79% particulate nitrate yield for compounds that did not hydrolyze. The mass of the $-\text{ONO}_2$ group alone represents 13% of the SOA mass. A rough calculation predicts that the nighttime reaction of 1 ppbv ($\sim 5.5 \mu\text{g}/\text{m}^3$) α -pinene may produce $\sim 0.5665 \mu\text{g}/\text{m}^3$ SOA ($\sim 0.3844 \mu\text{g}/\text{m}^3$ of which is particulate nitrates) through its reaction with NO_3 . These calculations neglect contributions of cross reactions between $n\text{RO}_2$ with and other terpene RO_2 , estimated to comprise another 20% of the $n\text{RO}_2$ fate, which may also lead to SOA formation. Our model excludes the contribution of other non-terpene RO_2 from precursors such as isoprene, which, while minor at night, may still contribute to RO_2 reactivity, and will tend to depress SOA yields due to the smaller dimers formed in those reactions (McFiggans et al., 2019). Although the yields in this work would predict α -pinene + NO_3 represents a significant fraction of aerosol and particulate nitrate mass observations during SOAS, previous modeling efforts did not track the SOA or organic nitrates from α -pinene + NO_3 due to belief that this reaction does not produce aerosol (Pye et al., 2015; Ayres et al., 2015). Atmospheric models should therefore include SOA formation from α -pinene + NO_3 , ideally with a dependence on $n\text{RO}_2$ fate.

Since HRMS analysis suggests that most compounds in α -pinene + NO_3 SOA retain their organonitrate functionality, they will become a permanent sink of NO_x following particle deposition. The α -pinene + NO_3 chemistry can also have important implications for reactive nitrogen budgets in the gas phase. The high yield of PNP (58%) from $n\text{RO}_2 + \text{HO}_2$ will temporarily sequester NO_y , which may become a permanent sink when the PNP undergoes deposition (Nguyen et al., 2015) or retains its nitrate functionality following further oxidation. Approximately 45% of $n\text{RO}_2$ is expected to react with HO_2 during the conditions of SOAS; thus, the nighttime reaction of 1 ppbv α -pinene is expected to produce ~ 260 pptv of volatile PNP. During SOAS, PNP is observed with maximum concentrations of ~ 30 pptv, possibly limited by its atmospheric deposition and transport, among other fates (Nguyen et al., 2015). Together, both particle and gas-phase organonitrate formation and deposition from α -pinene + NO_3 chemistry may amount to a substantial removal pathway for atmospheric NO_x .

Our results also highlight the necessity of performing chamber experiments under conditions that more closely match those in the ambient atmosphere. Measured SOA yields depend heavily on the reactive fate of RO_2 in chamber experiments. Previous experiments using N_2O_5 as a source of NO_3 to study SOA formation from α -pinene + NO_3 likely observed low mass yields due to the dominance of $n\text{RO}_2 + \text{NO}_3$ reactions, from which we observe little-to-no SOA formation. Similarly, experiments that maximize $n\text{RO}_2 + \text{HO}_2$ may observe primarily volatile products. Only in experiments designed to allow sufficient $n\text{RO}_2 + n\text{RO}_2$ reactions was SOA formation observed. Furthermore, HRMS analysis shows that a large fraction of the compounds in SOA formed under ‘simulated nighttime’ conditions are not observed in either ozonolysis-only or $n\text{RO}_2 + n\text{RO}_2$ -only SOA, suggesting that synergistic reaction pathways can enhance SOA formation when multiple oxidation channels occur simultaneously. Because ozonolysis and NO_3 oxidation occur together at night in the atmosphere, chamber experiments

510 isolating one or the other will necessarily be biased in their representation of nighttime SOA from α -pinene and potentially other volatile hydrocarbons.

Code and data availability. Chamber experiment data are available online at the Index of Chamber Atmospheric Research in the United States (ICARUS; <https://icarus.ucdavis.edu/>). The chemical mechanism for kinetic modelling can be found in the
515 Supplement (Mech. S1).

Author contributions. T. B. N., K. H. B. and G. J. P. B. designed the chamber experiments. K. H. B. and G. J. P. B. performed the chamber experiments. K. H. B. performed the kinetic modelling. J. D. C. assisted with chemical syntheses and experiments. T. B. N. and K. H. B. prepared the manuscript.

520
Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This work was funded by the US National Oceanic and Atmospheric Administration.

525 **References**

- Ayres, B.; Allen, H.; Draper, D.; Brown, S.; Wild, R.; Jimenez, J.; Day, D.; Campuzano-Jost, P.; Hu, W.; Gouw, J. d., Organic nitrate aerosol formation via NO_3 + biogenic volatile organic compounds in the southeastern United States. *Atmos. Chem. Phys.*, 15 (23), 13377-13392, 2015
- Bell, D. M., Wu, C., Bertrand, A., Graham, E., Schoonbaert, J., Giannoukos, S., Baltensperger, U., Prevot, A. S. H., Riipinen, I., El Haddad, I., and Mohr, C.: Particle-phase processing of α -pinene NO_3 secondary organic aerosol in the dark, *Atmos. Chem. Phys. Discuss.* [preprint], DOI: 10.5194/acp-2021-379, in review, 2021.
- Berndt, T., Scholz, W., Mentler, B., Fischer, L., Herrmann, H., Kulmala, M. and Hansel, A.: Accretion product formation from self- and cross-reactions of RO_2 radicals in the atmosphere. *Angew. Chem. Int. Ed.*, 57, 3820-3824, DOI: 10.1002/anie.201710989, 2018.
- 535 Boyd, A. A., Flaud, P.-M., Daugey, N., and Lesclaux, R.: Rate constants for RO_2 + HO_2 reactions measured under a large excess of HO_2 , *J. Phys. Chem. A*, 107, 818–821, 2003.

- Boyd, C.; Sanchez, J.; Xu, L.; Eugene, A. J.; Nah, T.; Tuet, W.; Guzman, M. I.; Ng, N., Secondary organic aerosol formation from the β -pinene + NO₃ system: effect of humidity and peroxy radical fate. *Atmos. Chem. Phys.*, 15 (13), 7497-7522, 2015.
- 540 Burkholder, J. B., Baynard, T., Ravishankara, A. R., and Lovejoy, E. R.: Particle nucleation following the O₃ and OH initiated oxidation of α -pinene and β -pinene between 278 and 320 K, *J. Geophys. Res.*, 112, D10216, DOI:10.1029/2006JD007783, 2007.
- Burkholder, J. B., S. P. Sander, J. Abbatt, J. R. Barker, C. Cappa, J. D. Crounse, T. S. Dibble, R. E. Huie, C. E. Kolb, M. J. Kurylo, V. L. Orkin, C. J. Percival, D. M. Wilmouth, and P. H. Wine "Chemical Kinetics and Photochemical Data for Use
545 in Atmospheric Studies, Evaluation No. 19," JPL Publication 19-5, Jet Propulsion Laboratory, Pasadena, 2019. <http://jpldataeval.jpl.nasa.gov>.
- Burrows, J. P., Tyndall, G. S., and Moortgat, G. K.: Absorption spectrum of NO₃ and kinetics of the reactions of NO₃ with NO₂, Cl, and several stable atmospheric species at 298 K. *J. Phys. Chem.*, 89 (22), 4848-4856, DOI: 10.1021/j100268a038, 1985.
- 550 Carlton, A. G., de Gouw, J., Jimenez, J. L., Ambrose, J. L., Attwood, A. R., Brown, S., Baker, K. R., Brock, C., Cohen, R. C., Edgerton, S., Farkas, C. M., Farmer, D., Goldstein, A. H., Gratz, L., Guenther, A., Hunt, S., Jaeglé, L., Jaffe, D. A., Mak, J., McClure, C., Nenes, A., Nguyen, T. K., Pierce, J. R., de Sa, S., Selin, N. E., Shah, V., Shaw, S., Shepson, P. B., Song, S., Stutz, J., Surratt, J. D., Turpin, B. J., Warneke, C., Washenfelter, R. A., Wennberg, P. O., and Zhou, X. Synthesis of the Southeast Atmosphere Studies: Investigating Fundamental Atmospheric Chemistry Questions. *Bull. Am. Met. Soc.* 99, 3, 547-567, DOI: 10.1175/BAMS-D-16-0048.1, 2018.
- 555 Chan, L. K.; Nguyen, K. Q.; Karim, N.; Yang, Y.; Rice, R. H.; He, G.; Denison, M. S.; Nguyen, T. B., Relationship between the molecular composition, visible light absorption, and health-related properties of smoldering woodsmoke aerosols. *Atmos. Chem. Phys.*, 20, 539-559, DOI: 10.5194/acp-20-539-2020, 2020.
- Clafflin, M. S.; Ziemann, P. J., Identification and quantitation of aerosol products of the reaction of β -pinene with NO₃ radicals
560 and implications for gas-and particle-phase reaction mechanisms. *J. Phys. Chem. A* 2018, 122 (14), 3640-3652.
- Crounse, J. D.; McKinney, K. A.; Kwan, A. J.; and Wennberg, P. O.: Measurement of Gas-Phase Hydroperoxides by Chemical Ionization Mass Spectrometry. *Anal. Chem.*, 78 (19), 6726-6732, DOI: 10.1021/ac0604235, 2006.
- Crounse, J. D.; Nielsen, L. B.; Jørgensen, S.; Kjaergaard, H. G.; Wennberg, P. O., Autoxidation of organic compounds in the atmosphere. *J. Phys. Chem. Lett.*, 4 (20), 3513-3520, 2013.
- 565 Fry, J. L.; Draper, D. C.; Barsanti, K. C.; Smith, J. N.; Ortega, J.; Winkler, P. M.; Lawler, M. J.; Brown, S. S.; Edwards, P. M.; Cohen, R. C., Secondary Organic Aerosol Formation and Organic Nitrate Yield from NO₃ Oxidation of Biogenic Hydrocarbons. *Environ. Sci. Technol.*, 48 (20), 11944-11953, 2014.

- Hakola, H.; Tarvainen, V.; Laurila, T.; Hiltunen, V.; Hellén, H.; Keronen, P., Seasonal variation of VOC concentrations above a boreal coniferous forest. *Atmos. Environ.*, 37 (12), 1623-1634, 2003
- 570 Hallquist, M.; Waengberg, I.; Ljungstroem, E.; Barnes, I.; Becker, K.-H., Aerosol and Product Yields from NO₃ Radical-Initiated Oxidation of Selected Monoterpenes. *Environ. Sci. Technol.*, 33 (4), 553-559, 1999.
- Hidy, G. M.; Blanchard, C. L.; Baumann, K.; Edgerton, E.; Tanenbaum, S.; Shaw, S.; Knipping, E.; Tombach, I.; Jansen, J.; Walters, J., Chemical climatology of the southeastern United States, 1999-2013. *Atmos. Chem. Phys.* 2014, 14 (21), 11893-11914.
- 575 Hoppel, W., Fitzgerald, J., Frick, G., Caffrey, P., Pasternack, L., Hegg, D., Gao, S., Leaitch, R., Shantz, N., Cantrell, C., Albrechtinski, T., Ambrusko, J., Sullivan, W.: Particle formation and growth from ozonolysis of α -pinene, *J. Geophys. Res.*, 106 (D21), 27603–27618, DOI: 10.1029/2001JD900018, 2001.
- Inomata, S.: New particle formation promoted by OH reactions during α -pinene ozonolysis. *ACS Earth Space Chem*, Just Accepted, DOI: 10.1021/acsearthspacechem.1c00142, 2021.
- 580 Iyer, S., Reiman, H., Møller, K. H., Rissanen, M. P., Kjaergaard, H. G., and Kurtén, T.: Computational Investigation of RO₂ + HO₂ and RO₂ + RO₂ Reactions of Monoterpene Derived First-Generation Peroxy Radicals Leading to Radical Recycling. *J. Phys. Chem. A*, 122 (49), 9542-9552, DOI: 10.1021/acs.jpca.8b09241, 2018.
- Iyer, S., Rissanen, M. P., Valiev, R., Barua, S., Krechmer, J. E., Thornton, J., Ehn, M., and Kurtén, T.: Molecular mechanism for rapid autoxidation in α -pinene ozonolysis. *Nature. Comm.*, 12, 878, DOI: 10.1038/s41467-021-21172-w, 2021.
- 585 Jenkin, M. E.; Saunders, S. M.; Pilling, M. J. The Tropospheric Degradation of Volatile Organic Compounds: A Protocol for Mechanism Development. *Atmos. Environ.*, 31, 81–104, 1997.
- Kenseth, C. M.; Huang, Y.; Zhao, R.; Dalleska, N. F.; Hethcox, J. C.; Stoltz, B. M.; Seinfeld, J. H., Synergistic O₃ + OH oxidation pathway to extremely low-volatility dimers revealed in β -pinene secondary organic aerosol. *Proc. Nat'l. Acad. Sci.*, 115 (33), 8301-8306, 2018.
- 590 Kind, T.; Fiehn, O., Seven Golden Rules for heuristic filtering of molecular formulas obtained by accurate mass spectrometry. *Bmc Bioinformatics*, 8, 105; DOI: 10.1186/1471-2105-8-105, 2007.
- Kruve, A.; Kaupmees, K.; Liigand, J.; Oss, M.; Leito, I., Sodium adduct formation efficiency in ESI source. *J. Mass Spectrom.*, 48 (6), 695-702, 2013.
- Kurtén, T.; Møller, K. H.; Nguyen, T. B.; Schwantes, R. H.; Misztal, P. K.; Su, L.; Wennberg, P. O.; Fry, J. L.; Kjaergaard, H.
- 595 G., Alkoxy Radical Bond Scissions Explain the Anomalously Low Secondary Organic Aerosol and Organonitrate Yields from α -Pinene + NO₃. *J. Phys. Chem. Lett.*, 8, 13, 2826–2834, 2017.

- Lee, L., Teng, A. P., Wennberg, P. O., Crounse, J. D., and Cohen, R. C. On rates and mechanism of OH and O₃ reactions with isoprene-derived hydroxy nitrates. *J. Phys. Chem. A*, 118 (9), 1622 – 1637, 2014.
- Mathis, J. A.; McCord, B. R., The analysis of high explosives by liquid chromatography/electrospray ionization mass spectrometry: multiplexed detection of negative ion adducts. *Rapid Comm. Mass Spectrom.*, 19 (2), 99-104, 2005.
- Moldanova, J. and Ljungström, E.: Modelling of particle formation from NO₃ oxidation of selected monoterpenes. *J. Aerosol Sci.*, 31 (11), 1317-1333, DOI: 10.1016/S0021-8502(00)00041-0, 2000.
- Molteni, U., Simon, M., Heinritz, M., Hoyle, C. R., Bernhammer, A.-K., Bianchi, F., Breitenlechner, M., Brilke, S., Dias, A., Duplissy, J., Frege, C., Gordon, H., Heyn, C., Jokinen, T., Kurten, A., Lehtipalo, K., Makhmutov, V., Petäjä, T., Pieber, S. M., Praplan, A. P., Schobesberger, S., Steiner, G., Stozhkov, Y., Tomé, A., Tröstl, J., Wagner, A. C., Wagner, R., Williamson, C., Yan, C., Baltensperger, U., Curtius, J., Donahue, N., Hansel, A., Kirkby, J., Kulmala, M., Worsnop, D. R., and Dommen, J. Formation of highly oxygenated organic molecules from α -pinene ozonolysis: chemical characteristics, mechanism, and kinetic model development. *ACS Earth Space Chem*, 3 (5), 873-883, DOI: 10.1021/acsearthspacechem.9b00035, 2019.
- Mutzel, A., Zhang, Y., Böge, O., Rodigast, M., Kolodziejczyk, A., Wang, X., and Herrmann, H.: Importance of secondary organic aerosol formation of α -pinene, limonene, and m-cresol comparing day- and nighttime radical chemistry, *Atmos. Chem. Phys.*, 21, 8479–8498, DOI: 10.5194/acp-21-8479-2021, 2021.
- Nah, T.; Sanchez, J.; Boyd, C. M.; Ng, N. L., Photochemical Aging of α -pinene and β -pinene Secondary Organic Aerosol formed from Nitrate Radical Oxidation. *Environ. Sci. Technol.*, 50 (1), 222-231, 2016.
- Nguyen, T. B.; Bateman, A. P.; Bones, D. L.; Nizkorodov, S. A.; Laskin, J.; Laskin, A., High-resolution mass spectrometry analysis of secondary organic aerosol generated by ozonolysis of isoprene. *Atmos. Environ.*, 44 (8), 1032-1042, 2010.
- Nguyen, T. B.; Laskin, J.; Laskin, A.; Nizkorodov, S. A., Nitrogen containing organic compounds and oligomers in secondary organic aerosol formed by photooxidation of isoprene. *Environ. Sci. Technol.*, 45, 6908-6918, 2011.
- Nguyen, T. B.; Crounse, J. D.; Schwantes, R. H.; Teng, A. P.; Bates, K. H.; Zhang, X.; St. Clair, J. M.; Brune, W. H.; Tyndall, G. S.; Keutsch, F. N.; Seinfeld, J. H.; Wennberg, P. O., Overview of the Focused Isoprene eXperiment at the California Institute of Technology (FIXCIT): mechanistic chamber studies on the oxidation of biogenic compounds. *Atmos. Chem. Phys.*, 14 (24), 13531-13549, 2014a.
- Nguyen, T. B., Coggon, M. M., Bates, K. H., Zhang, X., Schwantes, R. H., Schilling, K. A., Loza, C. L., Flagan, R. C., Wennberg, P. O., and Seinfeld, J. H.: Organic aerosol formation from the reactive uptake of isoprene epoxydiols (IEPOX) onto non-acidified inorganic seeds, *Atmos. Chem. Phys.*, 14, 3497–3510, DOI: 10.5194/acp-14-3497-2014, 2014b.

- Nguyen, T. B.; Crounse, J. D.; Teng, A. P.; St. Clair, J. M.; Paulot, F.; Wolfe, G. M.; Wennberg, P. O., Rapid deposition of oxidized biogenic compounds to a temperate forest. *Proc. Nat'l. Acad. Sci.*, 112 (5), E392–E401, 2015
- Orlando, J. J., and Tyndall, G. S.: Laboratory studies of organic peroxy radical chemistry: an overview with emphasis on recent issues of atmospheric significance, *Chem. Soc. Rev.*, 41, 6294-6317, DOI: 10.1039/c2cs35166h, 2012.
- 630 Praske, E., Crounse, J. D., Bates, K. H., Kurtén, T., Kjaergaard, H. G., Wennberg, P. O. Atmospheric fate of methyl vinyl ketone: peroxy radical reactions with NO and HO₂. *J. Phys. Chem. A*, 119 (19), 4562-4572. DOI: 10.1021/jp5107058, 2015.
- Pye, H. O.; Luecken, D. J.; Xu, L.; Boyd, C. M.; Ng, N. L.; Baker, K. R.; Ayres, B. R.; Bash, J. O.; Baumann, K.; Carter, W. P., Modeling the current and future roles of particulate organic nitrates in the southeastern United States. *Environ. Sci. Technol.*, 49 (24), 14195-14203, 2015.
- 635 Roach, P. J.; Laskin, J.; Laskin, A., Higher-Order Mass Defect Analysis for Mass Spectra of Complex Organic Mixtures. *Anal. Chem.*, 83 (12), 4924-4929, 2011
- Romer, P. S., Duffey, K. C., Wooldridge, P. J., Edgerton, E., Baumann, K., Feiner, P. A., Miller, D. O., Brune, W. H., Koss, A. R., de Gouw, J. A., Misztal, P. K., Goldstein, A. H., and Cohen, R. C.: Effects of temperature-dependent NO_x emissions on continental ozone production, *Atmos. Chem. Phys.*, 18, 2601–2614, DOI: 10.5194/acp-18-2601-2018, 2018.
- 640 Romonosky, D. E., Li, Y., Shiraiwa, M., Laskin, A., Laskin, J., and Nizkorodov, S. A.: Aqueous photochemistry of secondary organic aerosol of α -pinene and α -humulene oxidized with ozone, hydroxyl radical, and nitrate radical. *J. Phys. Chem. A*, 121 (6), 1298-1309, DOI: 10.1021/acs.jpca.6b10900, 2017.
- Saunders, S. M.; Jenkin, M. E.; Derwent, R. G.; Pilling, M. J. Protocol for the Development of the Master Chemical Mechanism, MCM v3 (Part A): Tropospheric Degradation of Non-Aromatic Volatile Organic Compounds. *Atmos. Chem. Phys.*, 3, 161–180, 2003.
- 645 Schwantes, R. H., Charan, S. M., Bates, K. H., Huang, Y., Nguyen, T. B., Mai, H., Kong, W., Flagan, R. C., and Seinfeld, J. H.: Low-volatility compounds contribute significantly to isoprene secondary organic aerosol (SOA) under high-NO_x conditions, *Atmos. Chem. Phys.*, 19, 7255–7278, DOI: 10.5194/acp-19-7255-2019, 2019.
- Sindelarova, K.; Granier, C.; Bouarar, I.; Guenther, A.; Tilmes, S.; Stavrakou, T.; Müller, J.-F.; Kuhn, U.; Stefani, P.; Knorr, W., Global data set of biogenic VOC emissions calculated by the MEGAN model over the last 30 years. *Atmos. Chem. Phys.*, 14 (17), 9317-9341, 2014
- Sisco, E.; Forbes, T. P., Rapid detection of sugar alcohol precursors and corresponding nitrate ester explosives using direct analysis in real time mass spectrometry. *Analyst*, 140 (8), 2785-2796, 2015.

- 655 Spittler, M.; Barnes, I.; Bejan, I.; Brockmann, K. J.; Benter, T.; Wirtz, K., Reactions of NO₃ radicals with limonene and α -pinene: Product and SOA formation. *Atmos. Environ.*, 40 (Suppl. 1), S116-S127, DOI: 10.1016/j.atmosenv.2005.09.093, 2006.
- St. Clair, J. M.; McCabe, D. C.; Crounse, J. D.; Steiner, U.; and Wennberg, P. O.: Chemical Ionization Tandem Mass Spectrometer for the *In Situ* Measurement of Methyl Hydrogen Peroxide. *Rev. Sci. Instrum.*, 81, 094102, DOI: 10.1063/1.3480552, 2010.
- 660 Takeuchi, M. and Ng, N. L.: Chemical composition and hydrolysis of organic nitrate aerosol formed from hydroxyl and nitrate radical oxidation of α -pinene and β -pinene, *Atmos. Chem. Phys.*, 19, 12749–12766, DOI: 10.5194/acp-19-12749-2019, 2019.
- Teng, A. P.; Crounse, J. D.; Wennberg, P. O., Isoprene peroxy radical dynamics. *J. Amer. Chem. Soc.*, 139 (15), 5367-5377, 2017.
- 665 Vasquez, K. T., Crounse, J. D., Schulze, B. C., Bates, K. H., Teng, A. P., Xu, L., and Wennberg, P. O. Rapid hydrolysis of tertiary isoprene nitrate efficiently removes NO_x from the atmosphere. *Proc. Nat'l Acad. Sci.*, 117 (52) 33011-33016. DOI: 10.1073/pnas.2017442117, 2020.
- Wennberg, P. O., Bates, K. H., Crounse, J. D., Dodson, L. G., McVay, R. C., Mertens, L. A., Nguyen, T. B., Praske, E., Schwantes, R. H., Smarte, M. D., St. Clair, J. M., Teng, A. P., Zhang, X., and Seinfeld, J. The gas-phase oxidation of isoprene and its first-generation products. *Chem. Rev.*, 118 (7), 3337-3390, DOI: 10.1021/acs.chemrev.7b00439, 2018.
- 670 Xu, L.; Møller, K. H.; Crounse, J. D.; Otkjær, R. V.; Kjaergaard, H. G.; Wennberg, P. O., Unimolecular reactions of peroxy radicals formed in the oxidation of α -pinene and β -pinene by hydroxyl radicals. *J. Phys. Chem. A*, 123 (8), 1661-1674, 2019
- 675 York, D.; Evensen, N.; Martinez, M.; Delgado, J. Unified Equations for the Slope, Intercept, and Standard Errors of the Best Straight Line. *Am. J. Phys.*, 72, 367-375, DOI: 10.1119/1.1632486, 2004.
- Zhang, H.; Yee, L. D.; Lee, B. H.; Curtis, M. P.; Worton, D. R.; Isaacman-VanWertz, G.; Offenberg, J. H.; Lewandowski, M.; Kleindienst, T. E.; Beaver, M. R., Monoterpenes are the largest source of summertime organic aerosol in the southeastern United States. *Proc. Nat'l Acad. Sci.*, 115 (9), 2038-2043, 2018.
- 680 Zhang, X., Cappa, C. D., Jathar, S. H., McVay, R. C., Ensberg, J. J., Kleeman, M. J., and Seinfeld, J. H.: Influence of vapor wall loss in laboratory chambers on yields of secondary organic aerosol, *Proc. Nat'l Acad. Sci.*, 111, 5802–5807, 2014.
- Zhang, X., Schwantes, R. H., McVay, R. C., Lignell, H., Coggon, M. M., Flagan, R. C., and Seinfeld, J. H.: Vapor wall deposition in Teflon chambers, *Atmos. Chem. Phys.*, 15, 4197–4214, DOI: 10.5194/acp-15-4197-2015, 2015.

- 685 Zhao, J., Ortega, J., Chen, M., McMurry, P. H., and Smith, J. N.: Dependence of particle nucleation and growth on high-molecular-weight gas-phase products during ozonolysis of α -pinene, *Atmos. Chem. Phys.*, 13, 7631–7644, DOI: 10.5194/acp-13-7631-2013, 2013.
- Zhao, Y., Thornton, J. A., and Pye, H. O. T.: Quantitative constraints on autoxidation and dimer formation from direct probing of monoterpene-derived peroxy radical chemistry, *Proc. Nat'l. Acad. Sci.*, 115 (48), 12142-12147, DOI: 10.1073/pnas.1812147115, 2018.