Long Range Prediction and the Stratosphere

2

- 3 Adam A. Scaife^{1,2}, Mark P. Baldwin³, Amy H. Butler⁴, Andrew J. Charlton-Perez⁵, Daniela
- 4 I.V. Domeisen^{6,7}, Chaim I. Garfinkel⁸, Steven C. Hardiman¹, Peter Haynes⁹, Alexey Yu
- 5 Karpechko¹⁰, Eun-Pa Lim¹¹, Shunsuke Noguchi^{12,13}, Judith Perlwitz¹⁴, Lorenzo Polvani^{15,16},
- 6 Jadwiga H. Richter¹⁷, John Scinocca¹⁸, Michael Sigmond¹⁸, Theodore G. Shepherd⁵, Seok-
- 7 Woo Son¹⁹, and David W.J. Thompson²⁰.
- 8 ¹ Met Office Hadley Centre for Climate Prediction and Research, Exeter, U.K.
- 9 ² College of Engineering, Mathematics and Physical Sciences, Exeter University, Exeter, U.K.
- ³ Department of Mathematics and Global Systems Institute, University of Exeter, Exeter, U.K.
- ⁴ NOAA Chemical Sciences Laboratory (CSL), Boulder, CO, USA.
- ⁵ Department of Meteorology, University of Reading, Reading, U.K.
- 13 ⁶ ETH Zurich, Zurich, Switzerland.
- ⁷ University of Lausanne, Lausanne, Switzerland

15

- ⁸ Fredy and Nadine Herrmann Institute of Earth Sciences, Hebrew University of Jerusalem,
- 17 Jerusalem, Israel
- ⁹ Department of Applied Mathematics and Theoretical Physics, University of Cambridge, Cambridge,
- 19 UK
- 20 ¹⁰ Finnish Meteorological Institute, Helsinki, Finland
- 21 ¹¹ Bureau of Meteorology, Melbourne, Australia
- 22 ¹² Research Center for Environmental Modeling and Application, Japan Agency for Marine-Earth
- 23 Science and Technology (JAMSTEC), Yokohama, Japan
- 24 ¹³ Meteorological Research Institute, Japan Meteorological Agency, Tsukuba, Japan
- 25 ¹⁴ NOAA Physical Sciences Laboratory (PSL), Boulder, CO, USA
- 26 ¹⁵ Columbia University, Department of Applied Physics and Applied Mathematics, U.S.A.
- 27 ¹⁶ Department of Earth and Environmental Sciences, New York City, NY, U.S.A.
- 28 ¹⁷ Climate and Global Dynamics Laboratory, National Center for Atmospheric Research, Boulder,
- 29 Colorado, U.S.A.
- 30 ¹⁸ Canadian Centre for Climate Modelling and Analysis, Environment and Climate Change Canada,
- 31 Victoria, BC, Canada
- 32 ¹⁹ School of Earth and Environmental Sciences, Seoul National University, Seoul, Republic of Korea
- 33 ²⁰ Department of Atmospheric Science, Colorado State University, Fort Collins, Colorado, U.S.A.

34

35 Correspondence to: Adam A. Scaife (adam.scaife@metoffice.gov.uk)

Abstract. Over recent years there have been concomitant advances in the development of stratosphere resolving numerical models, our understanding of stratosphere-troposphere interaction and the extension of long-range forecasts to explicitly include the stratosphere. These advances are now allowing new and improved capability in long range prediction. We present an overview of this development and show how the inclusion of the stratosphere in forecast systems aids monthly, seasonal, annual to decadal climate predictions and multidecadal projections. We end with an outlook towards the future and identify areas for improvement that could further benefit these rapidly evolving predictions.

44

45

46

47

48

49

50 51

52

53

54 55

56

57 58

59

60 61

62

63

64

65

36

37

38

39


40 41

42

43

1 Introduction

Daily weather fluctuations are thought to have a deterministic predictability horizon of around two weeks due to the sensitivity of the evolution of the atmospheric state to small errors in initial conditions (Lorenz 1969) - the so-called 'butterfly effect'. Recent estimates (Leung et al., 2020; Domeisen et al., 2018) as well as tests of the predictability of midlatitude daily weather using the latest global prediction models (Zhang et al., 2019; Son et al., 2020) produce similar estimates for this predictability limit. However, this does not preclude skilful forecasts of the statistics (most notably the average) of conditions at long range beyond this timescale (e.g. Shukla 1981). This predictability owes its existence to slowly varying predictable components of the climate system in the ocean, and in some cases the atmosphere, as well as externally forced changes such as volcanic or solar variability effects (e.g. Kushnir et al., 2019). Some of the more prominent examples of stratospheric variability such as sudden stratospheric warmings and their subsequent impact on the stratosphere and the troposphere (Baldwin et al., 2021), or the quasi-biennial oscillation and its associated teleconnections (Scaife et al., 2014a), have been shown to be predictable out to timescales well beyond the traditional two-week predictability horizon from initial tropospheric conditions alone. Other examples involve stratospheric pathways for teleconnections originating in the troposphere or ocean (e.g., Schwartz and Garfinkel 2017, Byrne et al. 2019). On longer timescales, boundary forcing, for example from composition changes such as ozone depletion and recovery allows the stratosphere to provide relatively slowly varying conditions to guide the turbulent troposphere and hence provide long range predictability (e.g., Thompson et al. 2011). The relative importance of stratospheric initial conditions to boundary conditions decreases with lead time as shown in the schematic in Figure 1.



Timescale

Figure 1: Schematic representation of the role of the stratosphere in long range

Figure 1: Schematic representation of the role of the stratosphere in long range prediction showing the transition from initial condition predictability in the atmosphere (blue) and the ocean (green), to boundary condition predictability at longer timescales (orange). Individual mechanisms involving the stratosphere are labelled in black showing their approximate contributions to forecast variance at different lead times.

The extension of long-range prediction systems to explicitly include representation of the stratosphere follows many years of development of stratosphere resolving general circulation models (GCMs). By the late 20th century many leading centres for climate research had started to include the stratosphere in versions of their GCMs (Pawson et al., 2000; Gerber et al., 2012). Much of the early model development was motivated by the discovery of the ozone hole in the 1980s (Farman et al., 1985) and the need for simulations of ozone depletion and potential recovery of the ozone hole following the 1987 Montreal Protocol, which required atmospheric models that represented both the atmospheric dynamics and chemistry of stratospheric ozone depletion (Molina and Rowland 1974; Crutzen 1974). In most cases this was achieved by adding further quasi-horizontal layers to the domain of existing climate models to extend their representation of the atmosphere to the stratopause or beyond (e.g. Rind et al 1988; Beagley et al., 1997; Swinbank et al., 1998; Sassi et al., 2002), while also incorporating key radiative (e.g. Fels et al., 1985), chemical (e.g. Steil et al., 1998) and dynamical (e.g. Scaife et al., 2000) processes.

The early development of so called 'high top' climate models, which represent the whole depth of the stratosphere, in general preceded the discovery of the main body of evidence that the variability of the stratosphere is not only affected by, but also interacts with the lower atmosphere and surface climate. Pioneering early studies suggested that the stratosphere might have direct effects on the troposphere and surface climate (e.g. Labitzke 1965; Boville 1984; Kodera et al., 1990, 1995; Haynes et al., 1991; Perlwitz and Graf 1995). In subsequent years, as reliable observational records lengthened and large enough samples of stratospheric variability were amassed it was unequivocally demonstrated that stratospheric variability precedes important tropospheric changes in the extratropics (Baldwin and Dunkerton 1999, 2001). There was debate about causality and whether the stratosphere really does affect the atmosphere below (e.g. Plumb and Semeniuk 2003). However, experiments where the stratosphere is perturbed in numerical models show changes in surface climate and reproduce similar patterns of response at the surface to those found in real world observations (e.g. Polvani and Kushner 2002; Norton et al., 2003; Scaife et al., 2006; Joshi et al., 2006; Scaife and Knight 2008; Hitchcock and Haynes 2016, White et al., 2020). These involve changes to planetary scale waves and also baroclinic

eddies in the troposphere that are consistent with changes in baroclinicity near the tropopause (Kushner and Polvani 2004; Song and Robinson 2004; Wittman et al., 2004, 2007; Scaife et al., 2012; Domeisen et al., 2013; Hitchcock and Simpson 2014; White et al., 2020). Importantly, as we discuss below, the same mechanisms also appear to be at work across a broad range of timescales (Kidston et al., 2015).

In recent years, motivated by the evidence of surface effects of stratospheric variability in the midlatitudes, the high-top model configurations used for stratospheric research were incorporated into leading prediction systems. Improved vertical resolution was already known to improve the atmospheric data assimilation of satellite instrument observations whose sensitivity was often heavily weighted towards stratospheric altitudes. This also provided initial stratospheric conditions for sets of retrospective forecasts, some of which were internationally coordinated (e.g. Butler et al., 2016; Tompkins et al., 2017). A growing number of operational systems are now producing regular ensembles of predictions at lead times of months or years with coupled ocean-atmosphere models that extend to the stratopause or beyond; for example at Environment Canada (Merryfield et al., 2013), the Met Office in the UK (MacLachlan et al., 2014), the German Weather Service DWD (Baehr et al., 2015), the Japan Meteorological Agency (Takaya et al., 2017) and the European Centre for Medium Range Weather Forecasts (Johnson et al., 2019). In the following sections we document the emerging impacts and benefits of this new capability for surface climate predictions at monthly, seasonal, and annual to decadal lead times starting with the shorter-range, initial condition cases and ending with the longerrange boundary-condition cases.

2 The stratosphere and monthly prediction

The best-established phenomenon that gives rise to predictability of surface climate from the stratosphere is the tropospheric circulation changes that follow strong and weak conditions in the stratospheric polar vortex (Baldwin and Dunkerton 1999, 2001). For example, weak vortex conditions such as those found in a sudden stratospheric warming (SSW, Baldwin et al., 2021) are typically followed by a weakening and southward shift of the tropospheric mid-latitude jet stream (see e.g. Kidston et al., 2015 and references therein) and thus the negative polarity of the North Atlantic Oscillation (NAO), Arctic Oscillation (AO) and Northern Annular Mode (NAM). These fluctuations also show a tendency to vacillate between strong westerly and weak (SSW) states on subseasonal timescales (Kuroda and Kodera 2001; Hardiman et al., 2020a). The changes in the troposphere persist roughly as long as those in the lower stratosphere, and last for around two months (Baldwin and Dunkerton 2001; Baldwin et al., 2003; Hitchcock et al., 2013; Son et al., 2020; Domeisen 2019). The impacts on surface climate also include changes in the frequency of extremes of temperature and rainfall (Scaife et al., 2008; King et al., 2019; Cai et al., 2016; Domeisen et al., 2020b).

Although *major* SSW events, involving a complete reversal of the zonal flow in the mid stratosphere, are rare in the southern hemisphere (Wang et al., 2020; Jucker et al., 2021), variations of the Antarctic polar vortex are likewise followed by similar signatures in the underlying tropospheric flow, in this case via the Southern Annular Mode (SAM). Weakening of the vortex is typically followed by a negative shift in the SAM and associated changes in rainfall and near surface temperature (Thompson et al., 2005; E. Lim et al., 2018, 2019, 2021, Rao et al., 2020e). These changes in Southern Hemisphere circulation typically take longer to reach the surface than their Northern Hemisphere counterparts (Graverson and Christiansen 2003), perhaps due to the stronger stratospheric polar vortex and weaker wave driving in the southern hemisphere, but they are nonetheless better predicted by improving stratospheric resolution of forecast models (Roff et al., 2011). The timescale of weeks for the predictability of sudden warmings is limited by the predictability of weather patterns in the troposphere which might trigger SSW events (e.g. Mukougawa et al., 2005; Taguchi 2016; Garfinkel and Schwarz 2017; Jucker and Reichler 2018; Lee et al., 2020a). However, if we add this timescale to the timescale of a month or more for the persistence of lower stratospheric anomalies and their surface effects (e.g.

Baldwin et al., 2003; Butler et al., 2019), we arrive at the conclusion that on these occasions at least,

initial conditions in the atmosphere can provide predictability well beyond the usual two-week horizon

for daily weather in either hemisphere.

Predictability of the atmosphere at monthly lead times is also known to originate in part from the Madden Julian Oscillation (MJO) in the troposphere and its teleconnection to the extratropics (e.g. Vitart 2017). The circulation pattern associated with the MJO resembles a poleward and eastward propagating Rossby wave with centres of action over the Pacific and extending into the Atlantic sector where it also maps strongly onto the North Atlantic Oscillation. The lead time of around 10 days for the impact of a change in the MJO to appear in the extratropical flow (e.g. Cassou 2008; Lin et al., 2009) is also consistent with the timescale for poleward propagation of Rossby waves (e.g. Scaife et al., 2017). It turns out that this tropospheric MJO teleconnection on monthly timescales also interacts with the stratosphere (Garfinkel and Schwartz 2017). The MJO teleconnection to the North Pacific affects the region most strongly associated with tropospheric precursors to SSW events, and consistent with this, SSWs in the observational record have tended to follow certain MJO phases. The subsequent weak vortex anomaly then propagates down to the troposphere (Garfinkel et al 2012), where it may strengthen and prolong any existing negative NAO signal that is directly linked to the MJO via the troposphere (Schwartz and Garfinkel 2017, 2020; Barnes et al., 2019).

In addition to the interaction of the MJO with the extratropical stratosphere, a further, completely different link between the stratosphere and the MJO has recently been uncovered which modulates MJO amplitude and persistence in the troposphere via the phase of the Quasi-Biennial Oscillation (QBO) in the tropical lower stratosphere (Liu et al., 2014; Yoo and Son 2016; Martin et al., 2021). In this case, easterly phases of the QBO appear to energise the MJO compared to westerly QBO phases, likely due to changes in temperature and hence static stability close to the tropopause (Hendon and Abhik 2018; Martin et al., 2019) with a potential contribution of cloud-radiation feedbacks (Son et al., 2017, see Martin et al., 2021 for a review). This modulation of the MJO is in turn important for predictability as it gives rise to higher monthly prediction skill of the MJO and its surface teleconnections during the easterly phase of the QBO (Marshall et al., 2017; Abhik and Hendon 2019; Y. Lim et al., 2019).

The traditional view of stratosphere-troposphere interaction involves upward propagation of planetary scale Rossby waves (Charney and Drazin 1961), but this linear theory applies equally well to downward propagation. Harnik and Lindzen (2001) and Perlwitz and Harnik (2003) identified a possible source of downward propagating planetary waves in the form of reflecting surfaces in the winter stratosphere. Examples of specific reflection events, showing upwards and then downward propagation have since been observed (e.g. Kodera et al., 2008; Harnik 2009; Kodera and Mukougawa 2017; Mukougawa et al., 2017; Matthias and Kretschmer 2020). These results suggest that the details of the stratospheric circulation such as regions of negative vertical wind shear could be important for the formation of reflecting conditions (Perlwitz and Shaw 2013) and may yet provide a further mechanism by which the stratosphere can affect the troposphere (Domeisen et al., 2019; Butler et al., 2019).

Following studies demonstrating enhanced tropospheric predictability after SSW events in individual climate models (e.g. Kuroda 2008; Mukougawa et al 2009; Marshall and Scaife 2010; Sigmond et al. 2013), subseasonal forecast systems which explicitly represent the stratosphere in the climate system were developed and implemented at operational prediction centres worldwide. It is often difficult to demonstrate significant increases in overall skill (e.g. Richter et al. 2020a) but routinely produced ensembles of subseasonal predictions show that both stratospheric variability and its subsequent tropospheric signature are predictable at monthly lead times (Domeisen et al. 2020a, 2020b). The strongest surface impacts occur if the polar vortex in the lower stratosphere is in a weakened state at the time of the SSW (Karpechko et al., 2017) and there appears to be a roughly linear relationship between the strength of these lower stratospheric anomalies and the tropospheric response (e.g. Runde et al. 2016; White et al. 2020 and see Baldwin et al. 2019 for a review). We should note however that

there is no one-to-one correspondence between stratospheric variability and tropospheric events, and some prominent examples of sudden stratospheric warmings are followed by differing tropospheric anomalies (e.g. Charlton-Perez et al., 2018; Knight et al., 2020; Butler et al., 2020; Rao et al., 2020a). Nevertheless, the canonical response is seen in the majority (~70%) of cases and periods of intense wintertime stratospheric variability are important windows of opportunity to provide skilful monthly forecasts (Mariotti et al., 2020; Tripathi et al., 2015a).



200

201

202

203204

205

206

207

208

209

210211

212

213214

215

216217

218219

220

221

222

223

224225

226

227

228

198

193

194

195

196 197




Figure 2: Monthly forecasts prior to the 2018 sudden stratospheric warming and severe cold event over northern Europe. Forecast Polar Cap Index (a) and February sea level pressure anomalies (b). Ensemble mean anomalies are shown for the average of forecasts initialised between 8th and 22nd of January 2018 relative to hindcasts over the 1993-2016 period using the Met Office Hadley Centre GloSea prediction system (MacLachlan et al., 2015). Sea level pressure is measured in hPa and Polar Cap Index is the geopotential height anomaly (m) averaged over 65N to the North Pole.

These forecast systems are now important tools for national meteorological and hydrological services to monitor impending stratospheric variability and associated surface impacts in real time. Recent extreme examples illustrate the importance of this activity. In February 2018 a major SSW occurred and was followed by a strong negative NAO-like pattern at the surface with easterly wind anomalies over Europe and multiple cold air outbreaks over the following weeks, including extreme snowfall across northern Europe (Figure 2, Karpechko et al., 2018; Knight et al., 2020; Rao et al., 2020a) and an abrupt end to Iberian drought in Southern Europe (Ayarzagueña et al., 2018). Studies of monthly ensemble predictions of this event with operational stratosphere resolving systems showed that the stratospheric event was predictable at least 2 weeks in advance (Figure 2) and that the ensemble forecasts indicated increased likelihood of cold surface conditions for several weeks after the event (Karpechko 2018; Butler et al., 2020; Statnaia et al., 2020; Rao et al., 2020a). Again, as in the analysis of previous events, there was also a strong association with the MJO entering Phase 7 with increased convection in the West Pacific (cf. Garfinkel and Schwartz 2017) in the 2018 event. Finally, we should also note that cases of monthly forecasts where the stratosphere plays an important role are not restricted to winters with sudden stratospheric warmings and periods when the stratospheric polar vortex is above normal strength also provide opportunities for skilful monthly forecasts (Tripathi et al., 2015b; Scaife et al., 2016). In this case an opposite but symmetric surface response results, with strong positive NAO. A very recent example occurred in February 2020, when, following an extremely strong polar vortex (Hardiman et al., 2020b; Lee et al., 2020b; Lawrence et al., 2020; Rao et al., 2021a), the tropospheric jet in the Atlantic sector strengthened, and the associated increased storminess and rainfall in this case resulted in UK monthly rainfall reaching a new record high (Davies et al., 2021).

3 The stratosphere and seasonal prediction

230 Prior to the advent of dynamical forecast systems which explicitly represent the stratosphere, seasonal forecasts using empirical relationships and statistical methods were proposed. These relied on the prior 231 state of the polar vortex and other predictable factors such as the OBO that are known to have links to 232 surface climate (Thompson et al., 2002; Charlton et al., 2003; Christiansen et al., 2005; Boer and 233 Hamilton 2008). In some cases they indicated additional predictability that was absent in existing 234 235 operational forecast systems, providing further evidence of predictability involving the stratosphere and further motivating the extension of dynamical forecast systems to properly represent the stratosphere. 236 Similar empirical forecast studies continue, and although they cannot provide evidence of predictability 237 that is as strong as from GCM experiments based on fundamental physical principles, they do continue 238 239 to be useful to indicate sources of predictability that need to be properly represented in comprehensive forecast systems (e.g. Folland et al., 2012; Wang et al., 2017; Hall et al., 2017; Byrne and Shepherd 240 241 2018).

242

243244

245

246247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263264

265

266

267

268

269

270271

Following the introduction of dynamical seasonal forecast systems with a good representation of the stratosphere, clear links between successful seasonal prediction of the North Atlantic Oscillation, the closely related Arctic Oscillation and the state of the stratospheric polar vortex have been identified in forecast output (e.g. Scaife et al., 2014b; Stockdale et al., 2015; Jia et al., 2017). Similar signals are also seen in the southern hemisphere in relation to predictability of the Southern Annular Mode (Seviour et al., 2014; Byrne et al., 2019; Lim et al., 2021). Statistically significant increases in overall skill directly attributable to the inclusion of the stratosphere in prediction systems is sometimes difficult to demonstrate (e.g. Butler et al., 2016), especially given that other factors such as horizontal resolution and physical parametrizations are often simultaneously changed. Nevertheless, the body of evidence now weighs heavily in favour of predictability of the NAO and SAM from the stratospheric polar vortex and from analyses showing reduced surface prediction skill in the absence of stratospheric variability (e.g. Hardiman et al 2011; Sigmond et al., 2013; Scaife et al., 2016).

A second clear example of seasonal predictability originating in the stratosphere is the Quasi-Biennial Oscillation (QBO). The QBO has such inherently long timescales that it persists for several months in seasonal forecasts from initial atmospheric conditions alone and its regularity means that it can be predicted from simple composites of earlier cycles. Nevertheless, a growing number of numerical models used in seasonal forecast systems can now simulate and predict the oscillation within climate forecasts (Garfinkel et al., 2018; Richter et al., 2020b; Stockdale et al., 2021) with the aid of forcing from parametrized non-orographic gravity waves, and there is skill in predicting QBO phase changes at lead times of a few months (e.g. Pohlman et al., 2013, Scaife et al., 2014a). The surface impact of the QBO is also well established and has stood the test of time since it was first identified in the 1970s (Ebdon 1975; Thompson et al., 2002; Anstey and Shepherd 2014; Gray et al 2018). Yet again this response projects closely onto the North Atlantic Oscillation (and hence the Arctic Oscillation/Northern Annular Mode) and the Southern Annular Mode. The favoured mechanism involves refraction of vertically propagating Rossby waves in the lower stratosphere (Holton and Tan 1980), although other pathways may also be involved (e.g. Inoue et al., 2011; Yamazaki et al., 2020; Rao et al., 2020b, 2021b). The observed magnitude of the QBO teleconnection is also large enough to provide seasonal predictability of surface climate (Boer and Hamilton 2008) but its modelled amplitude at the surface appears to be under-represented in current operational prediction systems and models (Scaife et al., 2014b; Garfinkel et al. 2018; O'Reilly et al. 2019; Rao et al. 2020b; Anstey et al. 2021).

In addition to the stratosphere acting as a source of predictability, other mechanisms by which the stratosphere plays a role in seasonal predictions involve a pathway for global scale teleconnections. These often originate in the tropics where the longer timescales of coupled ocean-atmosphere variability such as the El Niño Southern Oscillation (ENSO, L'Heureux et al. 2020) provide a predictable source of low frequency variability. Effects on the extratropics can occur by tropical excitation of anomalous Rossby waves which propagate polewards but also upwards into the stratosphere, as in the case of ENSO (Manzini et al., 2006; Domeisen et al., 2019), giving two pathways for extratropical influence

(Butler et al., 2014; Kretschmer et al., 2021). These highly predictable tropical sources of climate variability alter the strength and position of the stratospheric polar vortex in the extratropics as well as the frequency of SSWs (Polvani et al., 2017) and these are followed by changes in the seasonal westerly jets in the troposphere and surface climate via the North Atlantic Oscillation (Ineson and Scaife 2009; Cagnazzo and Manzini 2009) or the Southern Annular Mode (Byrne et al., 2019). As might be expected, both the QBO and ENSO teleconnections are best represented in seasonal forecast systems which contain a well resolved stratosphere (Butler et al., 2016). We note that new examples of the stratosphere acting as a conduit for seasonal teleconnections are still being uncovered (Hurwitz et al., 2012, Woo et al., 2015). For example, the Indian Ocean Dipole (IOD) received little attention in this context until the recent record event of late 2019, when it appears to have driven an extreme winter strengthening of the northern hemisphere stratospheric polar vortex. This strengthening took many weeks to decay, giving rise to extreme yet highly predictable conditions in the stratosphere and around the Atlantic sector in late boreal winter (Hardiman et al., 2020b; Lee et al., 2020b). The same event was also implicated in extreme changes in the polar vortex and the near SSW in the southern hemisphere (Rao et al., 2020e); an event that itself likely helped to drive the extreme summer conditions and wildfires over Australia that year (Lim et al., 2021).

279

280

281

282

283284

285

286

287 288

289

290

291 292

293

294

295

296

297

298299

300

301

302

303 304

305

306 307

308

309

310

311

312313

314

315316

317

318

319 320

321 322

323

324

325

326327

Apparent links between Arctic sea ice and seasonal winter climate in the mid latitudes have also been suggested to be mediated by the stratosphere, with increased Rossby wave activity and a weakening of the stratospheric polar vortex in response to reduced sea ice, especially in the Barents-Kara Sea (Jaiser et al., 2013; Kim et al. 2014; King et al., 2016; Kretschmer et al., 2016). Some studies also reproduced surface signals in response to sea ice anomalies in seasonal forecasts of particular years that are in apparent agreement with observational estimates (e.g. Balmaseda et al., 2010; Orsolini et al., 2012). However, recent updates to observational records show a weakening of these apparent effects (Blackport and Screen 2020) and significant non-stationarity (Kolstad and Screen 2019). Subsequent modelling studies with larger samples of simulations have provided mixed results (Zhang et al 2018; Dai and Song 2020; Smith et al 2021), and some have argued that the atmospheric response to sea ice is weak and that while the sensitivity to Barents-Kara sea ice may be stronger, the stratospheric response in particular is highly variable (McKenna et al. 2017). While there may well be a longer-term effect via the stratosphere from sea ice decline (Sun et al., 2015; Screen and Blackport 2019; Kretschmer et al., 2020), sensitivity of the response to the background state complicates the issue (Labe et al., 2019; Smith et al. 2017), as do possible confounding influences from the tropics (Warner et al. 2020) and to date there is no clear consensus for strong enough year-to-year effects to provide significant seasonal predictability.

Other proposed teleconnections acting via the stratosphere have been found in observations but remain to be confirmed with successful reproduction in physically based climate models. A prominent example involves a proposed link between Eurasian snow amounts and the stratosphere, followed by a return influence on the NAO and surface climate. In this case, enhanced snow cover or depth is associated with high pressure over north Eurasia, an increase in the flux of Rossby wave activity into the stratosphere and a subsequent weakening of the stratospheric polar vortex, followed by the expected negative shift in the NAO and AO (Cohen and Entekhabi 1999, Cohen and Jones 2011; Cohen et al., 2014; Furtado et al., 2015). However, the strength of this link in climate models and seasonal predictions is modest (Fletcher et al., 2009; Riddle et al., 2013; Tyrrell et al., 2018, 2019) and does not agree with apparent links to the AO in observations (Kretschmer et al., 2016; Garfinkel et al., 2020) even when model mean state biases are corrected (Tyrrell et al., 2020). It has also been suggested that teleconnections to snow are non-stationary or non-causal and there is continued debate about its long-term robustness (Peings et al., 2013; Henderson et al., 2018).

In summary, a number of mechanisms by which the stratosphere acts to provide seasonal predictability either by acting directly as a source of predictable variability (e.g. the QBO, SSWs), or as a conduit for teleconnections (e.g. ENSO, MJO, IOD) have now been established in observations and have been

confirmed using climate model simulations based on first principles. These operate in seasonal forecast systems, albeit with remaining errors such as the weakness of the QBO connection to surface climate. Meanwhile, other mechanisms involving the stratosphere (for example the response to snow cover variations) have been proposed based on apparent observed relationships, but until we have agreement between these observations and theory (model simulations), scientists remain sceptical of whether they represent actual sources of seasonal predictability and these remain topics of current research.

4 The stratosphere and annual to decadal prediction

In recent years, initialised predictions on longer timescales were developed on the premise of multiyear memory in the ocean (e.g. Smith et al 2007), and following the development pathway mapped out by seasonal forecasts in the past, these are now being run operationally to produce real time multi-model forecasts (Smith et al., 2013). Kushnir et al. (2019) mapped out this operational development of annual to decadal predictions and highlighted a number of sources of predictability, some of which involve the stratosphere (Figure 3), but not all of which are fully represented in climate prediction systems.




Figure 3. Sources of annual to decadal predictability, some of which involve the stratosphere through the response to external forcing, internal atmospheric dynamics, or ozone chemistry changes. After Kushnir et al., 2019.

Despite common misconceptions, not all annual to decadal predictability stems from the ocean. Indeed, it has been clearly demonstrated that multiyear predictability of the QBO exists in current decadal predictions systems out to lead times of several years (Pohlman et al., 2013; Scaife et al., 2014a). This offers the prospect of a stratospheric contribution to multiyear predictability of the extratropics through the teleconnection with the Arctic Oscillation (Anstey and Shepherd 2014, Gray et al., 2018) and to tropical predictability through links to the MJO (e.g. Martin et al., 2021) and wider tropical climate variability (Haynes et al., 2021).

Although it is more important on multidecadal timescales (see below), external forcing of the stratosphere can also act as a source of decadal predictability. Forced climate signals from changes in greenhouse gases or stratospheric effects such as ozone depletion occur on a much longer timescale than the lead time of decadal forecasts but their contribution to the skill of predictions is not trivial. For example, it is not immediately obvious whether the slow changes from multidecadal forced signals would simply be swamped by unpredictable internal variability on decadal timescales, rendering long term external forcing changes useless for decadal predictions. However, this is not the case and long-term forcing is now known to be an important source of decadal prediction skill (Smith et al., 2019, 2020).

353

354

355

356357

358

359

360

361

362

363

364

365

366367

368

369 370

371

372

373

374375

376

377378

379380

381

382

383

384

385

386 387

388

389 390

391

392

393 394

395396

397

398

399 400

401

External forcing involving the stratosphere on shorter timescales is also important for annual to decadal predictions. The stratosphere has long been known to be influenced by volcanic eruptions, particularly in the case of tropical volcanic eruptions which are powerful enough to inject significant quantities of sulphur dioxide into the atmosphere. Here it reacts with water to form sulphuric acid and persists in aerosol form, leading to predictable multiyear global surface cooling, tropical stratospheric warming and an intensification of the westerly stratospheric polar vortex in the extratropics (Robock and Mao 1992). Although the sample of observed events is limited, modelling studies have reproduced an observed post-eruption intensification of the westerly winds in the stratosphere and some impacts on the surface Arctic Oscillation. However, generations of models have struggled to reproduce the twoyear persistence of volcanic effects seen in observations and the observed magnitude of the effect on the winter AO (e.g. Stenchikov et al., 2006; Marshall et al., 2009; Charlton-Perez et al., 2013, Bittner et al., 2016). In addition to these changes in the atmosphere, the intensification of stratospheric westerlies and hence Arctic Oscillation also combines with surface cooling of the ocean to generate predictable changes in the Atlantic meridional overturning circulation (Reichler et al., 2012) which can extend the volcanic influence to decadal timescales (Swingedouw et al., 2015). Finally, although the mechanism is debated, there is also evidence of a multiyear effect of tropical volcanic eruptions on ENSO, presumably requiring the persistent radiative forcing that arises through the long residence time of volcanic products, particularly sulphate aerosols, in the stratosphere. This reportedly increases the frequency of El Nino events by a factor of two in the years following volcanic eruptions (Adams et al., 2003), again suggesting an important source of multiannual predictability via the stratosphere.

A second source of multiannual predictability from external forcing originates from solar variability and in particular the 11-year solar activity cycle. Although a number of alternative mechanisms have been proposed (see Gray et al., 2010 for a review), the established mechanism for surface effects via the stratosphere is the change in the polar vortex that results from changes in upper stratospheric heating over the course of each cycle between solar minimum and solar maximum. Atmospheric wave-mean flow interactions amplify the initial radiatively driven change and drive its descent to the troposphere (Kodera and Kuroda 2002; Marsh et al 2007; Ineson et al., 2011; Givon et al., 2021), where changes in the extratropical jets result in a negative (positive) Arctic Oscillation pattern following solar minimum (maximum). There is also evidence that it contributes to interannual prediction skill (Dunstone et al., 2016) and an interesting aspect that has emerged in recent years is the integrating effect of the ocean on solar induced changes in the NAO via interannual persistence of ocean heat content anomalies which leads to a lag of around 3 years ($\pi/2$ cycles) in the peak response, as would be expected if the ocean is integrating the effects of a periodic solar forcing (Scaife et al., 2013; Gray et al., 2013; Andrews et al., 2015; Thiéblemont et al., 2015). However, debate continues as to whether the solar signal is indeed large enough to be detectable in observations in the presence of large internal tropospheric variability (Chiodo et al., 2019).

Perhaps the longest known timescale for predictability from initial conditions, which also involves the stratosphere, is the interaction of Atlantic Multidecadal Variability (AMV) with the stratospheric circulation. The Atlantic has followed pronounced multidecadal variations over the last century (Mann et al., 1995) and these variations are predictable out to years ahead (Hermanson et al., 2014). Some

studies link these variations to the stratosphere and the NAO/AO (Reichler et al., 2012; Omrani et al., 2014). Indeed, the pronounced multidecadal increase in the surface NAO between the 1960s and 1990s is strongly coupled to changes in the strength of the stratospheric polar night jet (Scaife et al., 2005). Although current models simulate weak coupling between the AMV and the free atmosphere, this coupling appears to increase with model resolution (Lai et al., 2021) suggesting that the links between AMV, the stratosphere and the NAO offer potential for improved decadal scale prediction involving the stratosphere.

The currently recognised role of the stratosphere in decadal forecasts of surface climate again appears mainly via the impact on annular modes and, in the northern hemisphere, the North Atlantic Oscillation. Indeed, while current decadal prediction systems are now able to produce skilful predictions of variations in the NAO on multiyear lead times (Smith et al., 2019, 2020; Athanassiadis et al., 2020), much work is still needed to attribute these variations to external forcing or internal variability and to understand the interaction between boundary and initial conditions which blurs the simple distinction between the two. These new results are important because they indicate new-found decadal predictability of events like the high NAO of the 1990s which yielded a run of mild but wet and stormy winters in northern Europe and the eastern USA. These winters are well known to have caused significant impact for example on the insurance sector (Leckebusch et al., 2007) and coincided with the longest observed absence of SSW events (Pawson and Naujokat 1999; Domeisen 2019). Given the indications of coupled stratosphere-troposphere variations on decadal timescales (Scaife et al., 2005; Omrani et al., 2014; Garfinkel et al., 2017; Woo et al., 2015), understanding the role of the stratosphere in extratropical decadal predictions needs further investigation.

5 The stratosphere and multidecadal projection

The importance of the stratosphere for climate projections on multidecadal timescales was generally recognised before its role in predictions on shorter timescales. This is in part a legacy of the early development of stratosphere-troposphere models for ozone depletion studies described in the introduction. On these longer timescales, coupling between stratospheric composition, thermal structure and atmospheric circulation gives rise to surface climate predictability.

Perhaps the best-known case for the stratosphere affecting multidecadal projections of surface climate is the influence of ozone depletion on the southern annular mode (SAM; Thompson and Solomon 2002, 2005; McLandress et al., 2011; Polvani et al., 2011; Son et al., 2018) where decreasing ozone in the late 20th century led to a strengthened pole-to-equator temperature gradient, a stronger stratospheric polar vortex and a shift to strong positive SAM phases at the surface. In this case, studies again show the importance of stratospheric resolution to generate the full response, consistent with a genuine downward influence (Karpechko et al., 2008). The associated poleward shift in the tropospheric jet is connected to a delay in the spring breakdown of the stratospheric polar vortex (Byrne et al., 2017) and delivered significant and prolonged changes in rainfall across many regions of the southern hemisphere (Kang et al., 2011; Purich and Son 2012). Implementation of the Montreal Protocol in 1987 and subsequent reductions in the rate of ozone depletion mean that recovery of the ozone layer is now expected over the coming decades and the reversible effects of this on the surface climate form an important element of current multidecadal projections (Thompson et al., 2011; Previdi and Polyani 2014; Solomon et al., 2016; Banarjee et al., 2020, Zambri et al., 2021) where they are expected to play an important role alongside other changes in the southern stratosphere due to continuing increases in greenhouse gases (Son et al., 2009; Barnes et al., 2012), some of which occur via the stratospheric polar vortex in a similar way to those from ozone depletion and recovery (Ceppi and Shepherd, 2019).

The more limited effects of ozone depletion in the northern hemisphere meant that the role of the stratosphere in multidecadal projections took longer to become established. Some early studies found

potential amplification of positive Arctic Oscillation trends under climate change when the stratosphere was included (Shindell et al., 2001). However, this was not borne out in later studies as simulations with other fully coupled ocean-troposphere-stratosphere models suggested weakening of the stratospheric polar vortex (e.g. Huebener et al., 2007). Subsequent studies with multiple models also indicated a southward shift in the polar night jet with weakening high latitude winds and strengthening subtropical winds (Scaife et al., 2012; Manzini et al., 2014). These changes result from increased atmospheric wave driving of the winds which can overwhelm the cooling effect of greenhouse gases (Karpechko and Manzini 2012) and can lead to important differences in future surface climate, for example in regional rainfall in areas typically affected by the stratosphere via the Arctic Oscillation and NAO (Scaife et al., 2012). There is still significant uncertainty due to the diversity of modelled stratospheric responses to greenhouse gas increases (Manzini et al., 2014, Simpson et al., 2018, Zappa and Shepherd 2017), and it has proved difficult to identify any clear change in the frequency of sudden stratospheric warmings (Ayarzagüena et al., 2018, 2020; Rao et al., 2020c). This is perhaps due to the competition between strengthening latitudinal temperature gradients near the tropopause and enhanced meridional overturning in the mid stratosphere. There is also strong inherent unpredictable variability from decade to decade in the frequency of SSW occurrence (Butchart et al., 2000; McLandress and Shepherd 2009).

Other aspects of future climate change where the stratosphere plays a role have also been identified, for example, in the debate over the response to future levels of Arctic sea ice. In this case it seems that the response of the mid-latitude circulation involves a negative shift in the Arctic Oscillation (Screen et al., 2018; Zappa et al., 2018; McKenna et al., 2018). This could again be amplified by interaction with the stratosphere as some studies suggest that the stratospheric response is necessary for a large surface response (Kim et al., 2014), while others highlight that the stratospheric interaction is sensitive to the regional pattern of sea ice decline (McKenna et al., 2018), and still others show evidence of non-linear stratospheric, and stratosphere-mediated surface response (Manzini et al., 2018), coincident with the time when the Barents and Kara seas become ice-free (Kretschmer et al., 2020). Furthermore, studies also indicate that the surface climate response to sea ice decline depends systematically on the phase of the stratospheric QBO (Labe et al., 2019).

Although it is much less certain than anthropogenic climate change, there have also been suggestions of a multidecadal decline of external solar irradiance which can impact multidecadal climate projections via the stratosphere. Previous multidecadal solar minima, so called 'grand minima', have occurred in sunspot records and have been connected to the Little Ice Age period around the end of the 17th century using proxy and other data (Owens et al., 2017). Given recent weak amplitude 11 year solar cycles, there are now suggestions of a future solar 'grand minimum' where the 11 year cycle described above could become muted or even absent for a prolonged period (Lockwood et al., 2010). In this case, the upper stratospheric cooling in the tropics and summer hemisphere can change the meridional temperature gradient in a similar fashion to the 11 year cycle (Maycock et al., 2015) and leads to a negative shift in the AO and the NAO, and hence affects regional climate (Ineson et al., 2015). However, in this case it appears that while regional changes could be significant, they are generally much smaller than the surface warming due to anticipated levels of anthropogenic greenhouse gases (Anet et al., 2013; Ineson et al., 2015; Maycock et al., 2015).

Finally, we note that although low frequency variability in teleconnections is observed (e.g. Garfinkel et al., 2019), it is often unclear whether this is a systematic variation or simply due to sampling variability of an underlying stationary process (Jain et al., 2018). Nevertheless, there is growing evidence for systematic climate change in some of the teleconnections by which the stratosphere enables surface predictability. Under future climate change it appears that some of the teleconnections discussed above may *strengthen* in amplitude. For example, the strength of ENSO induced anomalies in the extratropical Atlantic/European sector increases in future climate projections (Müller and Roeckner 2006; Fereday et al., 2020). Similarly, recent analyses suggest that the MJO teleconnection to the

extratropics increases in amplitude under climate change (Samarasinghe et al., 2020). The same is also true of the extratropical effects of the stratospheric QBO, where in this case, the amplitude of the teleconnection in composite anomalies doubles under future climate change (Rao et al., 2020d) despite the QBO itself becoming weaker (Richter et al., 2020c).

6 Outlook

Long range prediction has evolved quickly in recent years (Merryfield et al., 2017; 2020, Butler et al., 2019; Meehl et al., 2021) and this rapid development is due in part to the improved representation of stratospheric processes and stratospheric initial conditions in ensemble prediction systems. The longrange forecast community originally focused on predictability from initial ocean conditions and this remains the primary source of long-range predictability, for example from ENSO, but some of these long-range prediction systems contained poor representations of the stratosphere. In the meantime, those working in parallel on climate modelling of the stratosphere were rarely involved in initialised long-range prediction, instead being driven primarily by the ozone depletion problem. Knowledge exchange across fields is important in science and precursors to a new paradigm often occur when a topic is investigated by researchers from outside the field (Kuhn 1970). The crossover and collaboration between long range prediction and stratospheric research communities is no exception and the interaction between these communities has yielded rapid progress and new insights. Examples where initial atmospheric conditions can provide predictability beyond the usually assumed limit have been demonstrated, particularly for the extratropics but also for the tropics, and we now know that in some situations, for example when sudden stratospheric warmings occur, the initial conditions in the stratosphere can have more impact than initial conditions in the ocean (Thompson et al 2002; Scaife and Knight 2008; Polvani et al 2017). This suggests that initial atmospheric conditions in the stratosphere are likely to be more important for long range forecasts than previously assumed (Mukougawa et al., 2005, 2009; Stockdale et al., 2015; Noguchi et al., 2016, 2020a; Choi and Son 2019; O'Reilly et al., 2019; Nie et al., 2019), not least because the overturning and breaking of Rossby waves in the stratosphere is followed by long lived atmospheric anomalies due to synoptic scale eddy feedbacks that prolong the effects in the troposphere (Kunz and Greatbatch 2013Kang et al., 2011; White et al., 2020). More research on the initial conditions in the stratosphere might therefore help to reveal potential for further improvements in prediction skill.

A notable simplification to understanding the role of the stratosphere, at least in extratropical long-range predictions, is its apparently seamless mechanism across different timescales and different phenomena. Following the early ground-breaking studies showing surface impacts of stratospheric variability (e.g. Labitzke 1965, Boville 1984) and a multitude of studies on individual teleconnections between the stratosphere and surface climate, the projection of stratospheric variability onto the Arctic Oscillation/North Atlantic Oscillation/Annular Mode circulation patterns across timescales and hemispheres is now well established (see the review by Kidston et al., 2015). This suggests that similar coupling processes occur between the stratosphere and troposphere from months to decades and these processes lead to some of the most intense extratropical climate extremes, in winter in the northern hemisphere and in late spring/early summer in the southern hemisphere (Karpechko et al., 2018; Fereday et al., 2012, Kautz et al., 2019; Domeisen and Butler 2020). While studies point to changes in upper tropospheric baroclinicity and tropospheric eddy feedbacks as crucial in these teleconnections, a full mechanistic understanding of how this occurs is still lacking.

Some, but not all, leading forecast systems now include a well resolved stratosphere with reasonable representation of relevant processes such as the body force from sub-grid orographic and non-orographic gravity waves. However, many outstanding problems remain. Although their number is increasing, only a subset of current GCMs have the ability to simulate a realistic QBO beyond its decay from initial conditions and it seems that all GCMs have problems with the fidelity of modelled QBO

546 teleconnections, which are either too weak or absent altogether (Scaife et al., 2014a; Kim et al., 2020; Anstey et al., 2021). Even the relatively well studied ENSO teleconnection via the stratosphere to the 547 extratropics still has outstanding questions, such as whether the northern hemisphere stratosphere 548 exhibits more SSW events during the La Niña phase (Butler and Polvani 2011; Song and Son 2012). 549 550 This is not generally reproduced in modelling systems (Garfinkel et al., 2012) but occurred in the recent 551 La Niña winter of 2020/2021. Similarly, while the increased monthly predictability from the MJO during the easterly phase of the QBO has been detected in monthly forecast experiments, the QBO-552 MJO connection does not persist in longer predictions and simulations with current models (Kim et al., 553 2020). Research and model development on stratosphere-troposphere interaction, including tropical 554 555 effects (Noguchi et al., 2020b), will no doubt lead to further progress in resolving this issue (Haynes et al., 2021). 556

Errors in the modelled climatological mean climate are inevitably present to varying degrees in even the latest climate models. The common protocol of running a set of retrospective predictions to allow these mean biases to be estimated and hence subtracted from real time predictions may well correct for much of this error. However, the degree to which biases have a nonlinear, state dependent impact on the predictions is not fully understood. In some contexts, the nonlinear impacts of biases may be minimal (Karpechko et al., 2021) while others show sensitivity (Sigmond et al., 2008, 2010) and increases of prediction skill occur under certain background conditions, for example during Easterly QBO phases (Taguchi 2018). Other processes generally omitted from long range predictions include interactive variations of ozone and other trace gases. Although reports of impacts and benefits have varied, it is thought that surface signals on interannual timescales come mainly from dynamical rather than chemical changes (Seviour et al., 2014; Harari et al., 2019). Nevertheless, some studies suggest detectable effects from interannual variability of ozone and it may be that ozone fluctuations could help to amplify surface signals (Karpechko et al., 2014; Son et al., 2013; Smith and Polyani 2014; Oehrlein et al., 2020; Hendon et al., 2020), providing a further area for future development. Given that the cost of full atmospheric chemistry schemes remains computationally expensive, it seems likely that simple parametrizations of ozone chemistry (e.g. Monge-Sanz et al., 2021) would be valuable in this context.

We end with a pointer to an issue that has now been found to affect long-range predictions from monthly to seasonal to decadal and multidecadal timescales, particularly in the extratropics. So called 'perfect model studies', which test the ability of models to predict their own ensemble members, are now known to underestimate the true predictability of climate in some regions, particularly around the Atlantic basin, and so models are better at predicting real world variations than they are at predicting themselves. This so-called 'Signal to Noise Paradox' (Scaife and Smith 2018) is at first surprising, because perfect model prediction scores are often assumed to represent an upper (rather than lower) limit for prediction skill of the real world. The problem can be understood in terms of unrealistically weak ensemble mean predictions (e.g. Eade et al., 2014) but whether the stratosphere is involved directly in the cause of this problem remains to be seen (Saito et al., 2017; Stockdale et al., 2015), as it initially appears in the troposphere rather than the stratosphere in long range forecasts (Domeisen et al., 2020a). Nevertheless, the unrealistically weak amplitude of ensemble mean predictions may well have the same root cause as the weaker than observed amplitude of modelled teleconnections to the stratosphere discussed in this review, including, for example, the under-representation of the surface impact of the QBO. Resolving this problem will therefore likely amplify these signals, provide greater levels of prediction skill, and further strengthen the role of the stratosphere in long range predictions of surface climate.

Author Contributions

557

558

559560

561

562563

564

565566

567

568

569

570 571

572

573

574

575

576

577

578579

580

581

582

583

584

585

586 587

588

589

592

AAS wrote the draft manuscript. All other co-authors contributed relevant references and input to revisions and editing of the manuscript. SWS helped produce Figure 1.

Acknowledgements

- 593 AAS and SCH were supported by the Met Office Hadley Centre Climate Programme funded by BEIS
- and Defra. MPB was supported by the Natural Environment Research Council (grant number
- NE/M006123/1). JHR was supported by the Regional and Global Model Analysis (RGMA) component
- of the Earth and Environmental System Modeling Program of the U.S. Department of Energy's Office
- of Biological & Environmental Research (BER) via NSF Interagency Agreement 1844590. SN was
- supported by the Japan Society for the Promotion of Science (KAKENHI, Grant Number: 19K14798).
- 599 EPL was supported by the Australian government's National Environmental Science Program Phase 2
- and the Victorian Water and Climate Initiative Phase 2. SWS was supported by the National Research
- Foundation of Korea (NRF) grant funded by the South Korean government (Ministry of Science and
- 602 ICT) (2017R1E1A1A01074889). DWJT is supported by the US National Science Foundation Climate
- and Large-Scale Dynamics program. Support from the Swiss National Science Foundation through
- projects PP00P2_170523 and PP00P2_198896 to D.D. is gratefully acknowledged. CIG was supported
- by a European Research Council starting Grant under the European Union Horizon 2020 research and
- 606 innovation program (Grant agreement 677756).

608 References

607

- Abhik, S. and Hendon, H.H.: Influence of the QBO on the MJO during coupled model multiweek
- forecasts. Geophysical Research Letters, 46(15), 9213-9221, 2019.
- Adams, B. J., Mann, M. and Ammann, C. Proxy evidence for an El Niño-like response to volcanic
- 612 forcing. Nature, 426, 274–278, 2003.
- Andrews, M., Knight, J. & Gray, L. A simulated lagged response of the North Atlantic Oscillation to
- the solar cycle over the period 1960–2009. Env. Res. Lett. 10, L054022, 2015.
- Anet, J. G., et al.: Impact of a potential 21st century "grand solar minimum" on surface temperatures
- and stratospheric ozone, Geophys. Res. Lett., 40, 4420–4425, doi:10.1002/grl.50806, 2003.
- Anstey, J.A. and Shepherd, T.G.: High-latitude influence of the Quasi-Biennial Oscillation. Quart. J.
- 618 Roy. Meteor. Soc., 140, 1–21, 2014.
- Anstey, J.A., Simpson, I.R., Richter, J.H., Naoe, H., Taguchi, M., Serva, F., et al.: Teleconnections of
- the Quasi-Biennial Oscillation in a multi-model ensemble of QBO-resolving models. Q J R Meteorol
- 621 Soc, 1–26, 2021.
- 622 Ayarzagüena, B., L.M. Polvani, U. Langematz, and CCMI co-authors: No robust evidence of future
- changes in major stratospheric sudden warmings: A multi-model assessment from CCMI, Atmos.
- 624 Chem. Phys., 18, 11277-11287, 2018.
- 625 Ayarzagüena, B., Barriopedro, D., Garrido-Perez, J. M., Abalos, M., de la Cámara, A., García-
- Herrera, R., et al.: Stratospheric connection to the abrupt end of the 2016/2017 Iberian drought.
- 627 Geophysical Research Letters, 45, 12, 639–12, 646, 2018.
- 628 Ayarzagüena et al.: Uncertainty in the response of stratospheric sudden warmings and stratosphere-
- troposphere coupling to quadrupled CO2 concentrations in CMIP6 models, J. Geophys. Res., 126, 6,
- 630 e2019JD032345, 2020.
- Baehr, J., Fröhlich, K., Botzet, M., Domeisen, D. I. V., Kornblueh, L., Notz, D., et al.: The prediction
- of surface temperature in the new seasonal prediction system based on the MPI-ESM coupled climate
- 633 model. Climate Dynamics, 44(9-10), 2723–2735, 2015.
- Baldwin, M. P., and Dunkerton, T. J.: Propagation of the Arctic Oscillation from the stratosphere to
- 635 the troposphere, J. Geophys. Res., 104 (D24), 30937–30946, 1999.

- Baldwin, M.P. and T.J. Dunkerton: Stratospheric Harbingers of Anomalous Weather Regimes.
- 637 Science, 294, 5542, 581-584, 2001.
- Baldwin M.P., David B. Stephenson, David W. J. Thompson, Timothy J. Dunkerton, Andrew
- J. Charlton, Alan O'Neill. Stratospheric Memory and Skill of Extended-Range Weather Forecasts.
- 640 Science, 301, 636-640, 2003.
- Baldwin M.P., et al.: 100 Years of Progress in Understanding the Stratosphere and Mesosphere. Met.
- 642 Monographs, 59, 27.1-27.62, 2019.
- Baldwin, M. P., Ayarzagüena, B., Birner, T., Butchart, N., Butler, A. H., Charlton-Perez, A. J., et al.:
- Sudden stratospheric warmings. Reviews of Geophysics, 59, e2020RG000708, 2021.
- Balmaseda, M.A., Ferranti, L., Molteni, F. and Palmer, T.N.: Impact of 2007 and 2008 Arctic ice
- anomalies on the atmospheric circulation: Implications for long-range predictions. Q.J.R. Meteorol.
- 647 Soc., 136: 1655-1664, 2010.
- Barnes, E.A, N.W. Barnes and L.M. Polvani: Delayed Southern Hemisphere climate change induced
- by stratospheric ozone recovery, as projected by the CMIP5 models, J. Climate, 27, 852-867, 2014.
- Barnes, E. A., Samarasinghe, S. M., Ebert-Uphoff, I., & Furtado, J. C.: Tropospheric and stratospheric
- 651 causal pathways between the MJO and NAO. Journal of Geophysical Research: Atmospheres, 124,
- 652 9356–9371, 2019.
- Beagley, S. R., J. de Grandpre', J. N. Koshyk, N. A. McFarlane, and T. G. Shepherd. Radiative-
- dynamical climatology of the first-generation Canadian middle atmosphere model. Atmosphere-
- 655 Ocean 35, 293-331, 1997.
- Bittner, M., Timmreck, C., Schmidt, H., Toohey, M., and Krüger, K.: The impact of wave-mean flow
- 657 interaction on the Northern Hemisphere polar vortex after tropical volcanic eruptions, J. Geophys.
- 658 Res. Atmos., 121, 5281–5297, 2016.
- Blackport, R. and J.A. Screen. Weakened evidence for mid-latitude impacts of Arctic warming.
- 660 Nature. Clim. Change, 10, 1065-1666, 2020.
- Boer, G.J., Hamilton, K. QBO influence on extratropical predictive skill. *Clim Dyn* **31**, 987–1000,
- 662 2008.
- Boville, B. A.: The Influence of the Polar Night Jet on the Tropospheric Circulation in a GCM, J.
- 664 Atm. Sci., 41(7), 1132-1142, 1984.
- Butchart, N., Austin, J., Knight, J. R., Scaife, A. A., & Gallani, M. L.: The Response of the
- 666 Stratospheric Climate to Projected Changes in the Concentrations of Well-Mixed Greenhouse Gases
- from 1992 to 2051, Journal of Climate, 13(13), 2142-2159, 2000.
- Butler, A.H., A. Charlton-Perez, D.I.V. Domeisen, I.R. Simpson, and J. Sjoberg, The predictability of
- Northern Hemisphere final stratospheric warmings and their surface impacts, Geophys. Res. Lett., 46,
- 670 10578-10588, 2019.
- Butler A.H., Arribas A., Athanassiadou M., Baehr J., Calvo N., Charlton-Perez A., Deque M.,
- Domeisen D.I.V., Fröhlich K., Hendon H., Imada Y., Ishii M., Iza M., Karpechko A.Y., Kumar A.,
- Maclachlan C., Merryfield W.J., Muller W.A., O'Neill A., Scaife A.A., Scinocca, J., Sigmond M.,
- 674 Stockdale T.N., Yasuda T.: The Climate-system Historical Forecast Project: Do stratosphere-
- 675 resolving models make better seasonal climate predictions in boreal winter? Quart. J. Roy. Met. Soc.,
- 676 142, 1413-1427, doi:10.1002/qj.2743, 2016.
- Butler, A.H., Z.D. Lawrence, S.H. Lee, S.P. Lillo, and C.S. Long: Differences between the 2018 and
- 678 2019 stratospheric polar vortex split events, Quar. Jour. Roy. Met. Soc., 1-19, 2020.

- 679 Butler, A.H., A. Charlton-Perez, D.I.V. Domeisen, C. Garfinkel, E.P. Gerber, P. Hitchcock, A.-Yu
- 680 Karpechko, A.C. Maycock, M. Sigmond, I. Simpson, S.-W. Son. Sub-seasonal Predictability and the
- Stratosphere- Chapter 11, The Gap Between Weather and Climate Forecasting, p. 223-241, 2019.
- Butler, A.H., L.M. Polvani and C. Deser: Separating the stratospheric and tropospheric pathways of El
- Niño-Southern Oscillation teleconnections, Environ. Res. Lett., 9, 024014, 2014.
- Butler A.H. and L.M. Polvani: El Niño, La Niña, and stratospheric sudden warmings: A re-evaluation
- in light of the observational record, Geophys. Res. Lett., 38, L13807, 2011.
- Byrne, N.J., Shepherd, T.G., Woollings, T. and Plumb, R.A.: Non-stationarity in Southern
- 687 Hemisphere climate variability associated with the seasonal breakdown of the stratospheric polar
- 688 vortex. J. Clim., 30, 7125–7139, 2017.
- Byrne, N. J., and T. G. Shepherd: Seasonal persistence of circulation anomalies in the Southern
- Hemisphere stratosphere and its implications for the troposphere. J. Clim., 31, 3467–3483,
- 691 doi:10.1175/JCLI-D-17-0557.1, 2018.
- Byrne, N.J., Shepherd, T.G. and Polichtchouk, I.: Subseasonal-to-seasonal predictability of the
- 693 Southern Hemisphere eddy-driven jet during austral spring and early summer. J. Geophys. Res., 124,
- 694 6841–6855, 2019.
- 695 Cai, M., Y. Yu, Y. Deng, H.M. van den Dool, R. Ren, S. Saha, X. Wu, J. Huang: Feeling the pulse of
- the stratosphere: an emerging opportunity for predicting continental-scale cold-air outbreaks 1 month
- 697 in advance. Bull. Am. Meteorol. Soc., 97, pp. 1475-1489, 2016.
- 698 Cassou, C.: Intraseasonal interaction between the Madden–Julian Oscillation and the North Atlantic
- 699 Oscillation. Nature 455, 523–527, 2008.
- 700 Christiansen B.: Downward propagation and statistical forecast of the near-surface weather. J.
- 701 Geophys. Res. Atmos., 110, 2005.
- Ceppi, P. and Shepherd, T.G.: The role of the stratospheric polar vortex for the austral jet response to
- 703 greenhouse gas forcing. Geophys. Res. Lett., 46, 6972–6979, 2019.
- Charlton, A.J., O'Neill, A., Stephenson, D.B., Lahoz, W.A. and Baldwin, M.P.: Can knowledge of the
- state of the stratosphere be used to improve statistical forecasts of the troposphere?. Q.J.R. Meteorol.
- 706 Soc., 129: 3205-3224, 2003.
- 707 Charlton-Perez AJ, et al.: On the lack of stratospheric dynamical variability in low-top versions of the
- 708 CMIP5 models, J. Geophys. Res. Atmos, 118, 2494–2505, 2013.
- 709 Charlton-Perez, A. J., Ferranti, L. and Lee, R. W.: The influence of the stratospheric state on North
- 710 Atlantic weather regimes. Quarterly Journal of the Royal Meteorological Society, 144 (713). pp.
- 711 1140-1151. ISSN 1477-870X, 2018.
- 712 Chiodo, G., Oehrlein, J., Polvani, L.M. et al.: Insignificant influence of the 11-year solar cycle on the
- North Atlantic Oscillation. Nat. Geosci. 12, 94–99, 2019.
- 714 Choi, J., and S.-W. Son: Stratospheric initial condition for skillful surface prediction in the ECMWF
- 715 model, Geophysical Research Letters, 21, 12556-12564, 2019.
- Cohen J. and D. Entekhabi: Eurasian snow cover variability and northern hemisphere climate
- 717 variability. Geophys. Res. Lett., 26, 345-348, 1999.
- 718 Cohen, J., and Jones, J.: A new index for more accurate winter predictions. Geophysical Research
- 719 Letters, 38, L21701, 2011.
- 720 Cohen J., Furtado JC, Jones J, Barlow M, Whittleston D, Entekhabi D.: Linking Siberian snow cover
- to precursors of stratospheric variability. J Clim., 27(14), 5422–5432, 2014.

- 722 Crutzen P.J.: Estimates of Possible Variations in Total Ozone Due to Natural Causes and Human
- 723 Activities. Ambio, 3, No. 6, 201-210, 1974.
- Dai, A., Song, M.: Little influence of Arctic amplification on mid-latitude climate. Nat. Clim. Chang.
- 725 10, 231–237, 2020.
- Davies, P.A., McCarthy, M., Christidis, N., Dunstone, N., Fereday, D., Kendon, M., Knight, J.R.,
- Scaife, A.A. and Sexton, D.: The wet and stormy UK winter of 2019/2020. Weather,
- 728 https://doi.org/10.1002/wea.3955, 2021.
- 729 Domeisen, D. I. V., Sun, L., & Chen, G.: The role of synoptic eddies in the tropospheric response to
- 730 stratospheric variability. Geophysical Research Letters, 40, 1–5, 2013.
- 731 Domeisen, D. I. V., Badin, G., & Koszalka, I. M.: How Predictable Are the Arctic and North Atlantic
- Oscillations? Exploring the Variability and Predictability of the Northern Hemisphere. Journal of
- 733 Climate, 31(3), 997–1014, 2018.
- Domeisen, D. I. V.: Estimating the Frequency of Sudden Stratospheric Warming Events From Surface
- 735 Observations of the North Atlantic Oscillation. Journal of Geophysical Research-Atmospheres,
- 736 124(6), 3180–3194, 2019.
- 737 Domeisen, D. I. V., Garfinkel, C. I., & Butler, A. H.: The Teleconnection of El Niño Southern
- Oscillation to the Stratosphere. Reviews of Geophysics, 57(1), 5–47, 2019.
- 739 Domeisen, D. I. V., & Butler, A. H.: Stratospheric drivers of extreme events at the Earth's surface.
- 740 Communications Earth & Environment, 1–8, 2020.
- 741 Domeisen, D.I.V., A.H. Butler, A.J. Charlton-Perez, B. Ayarzaguena, M.P. Baldwin, E. Dunn-
- 742 Sigouin, J.C. Furtado, C.I. Garfinkel, P. Hitchcock, A. Yu. Karpechko, H. Kim, J. Knight, A.L. Lang,
- 743 E.-P. Lim, A. Marshall, G. Roff, C. Schwartz, I.R. Simpson, S.-W. Son, M. Taguchi: The role of
- stratosphere-troposphere coupling in sub-seasonal to seasonal prediction. 1. Predictability in the
- 745 Stratosphere, J. Geophys. Res., 125, e2019JD030920, 2020a.
- 746 Domeisen, D.I.V., A.H. Butler, A.J. Charlton-Perez, B. Ayarzaguena, M.P. Baldwin, E. Dunn-
- 747 Sigouin, J.C. Furtado, C.I. Garfinkel, P. Hitchcock, A. Yu. Karpechko, H. Kim, J. Knight, A.L. Lang,
- 748 E.-P. Lim, A. Marshall, G. Roff, C. Schwartz, I.R. Simpson, S.-W. Son, M. Taguchi: The role of
- stratosphere-troposphere coupling in sub-seasonal to seasonal prediction. 2. Predictability arising
- from stratosphere-troposphere coupling, J. Geophys. Res., 125, e2019JD030923, 2020b.
- 751 Douville H.: Stratospheric polar vortex influence on Northern Hemisphere winter climate variability.
- 752 Geophys. Res. Lett., 36, L18703, 2009.
- 753 Eade R., D. Smith, A.A. Scaife and E. Wallace. Do seasonal to decadal climate predictions
- underestimate the predictability of the real world? Geophys. Res. Lett, 5620–5628, 2014.
- 755 Farman, J., Gardiner, B. & Shanklin, J.: Large losses of total ozone in Antarctica reveal seasonal
- 756 ClOx/NOx interaction. Nature 315, 207–210, 1985.
- 757 Fels S.B.: Radiative–Dynamical Interactions in the Middle Atmosphere. Advances in Geophysics, 28,
- 758 Part A, 277-300, 1985.
- 759 Fereday D. R. et al.: Seasonal forecasts of northern hemisphere winter 2009/10. Environ. Res. Lett. 7,
- 760 034031, 2012.
- 761 Fereday D. et al.: Tropical rainfall drives stronger future ENSO-NAO teleconnection in CMIP5
- 762 models. Geophys. Res. Lett., 47, e2020GL088664, 2020.

- Fletcher, C. G., S. C. Hardiman, P. J. Kushner, and J. Cohen: The dynamical response to snow cover
- perturbations in a large ensemble of atmospheric GCM integrations, J. Climate, 22(5), 1208-1222,
- 765 doi:10.1175/2008JCLI2505.1, 2009.
- Furtado, J.C., Cohen, J.L., Butler, A.H. et al.: Eurasian snow cover variability and links to winter
- 767 climate in the CMIP5 models. Clim Dyn 45, 2591–2605, 2015.
- 768 Garfinkel, C.I., C. Schwartz, D.I.V. Domeisen, S.-W. Son, A.H. Butler, and I.P. White, Extratropical
- atmospheric predictability from the Quasi-Biennial Oscillation in subseasonal forecast models, Jour.
- 770 Geophys. Res., 123 (15), 7855-7866, 2018.
- Garfinkel, C. I., Schwartz, C., Butler, A. H., Domeisen, D. I. V., Son, S.-W., & White, I. P.:
- 772 Weakening of the Teleconnection From El Niño-Southern Oscillation to the Arctic Stratosphere Over
- the Past Few Decades: What Can Be Learned From Subseasonal Forecast Models? Journal of
- 774 Geophysical Research-Atmospheres, 124(14), 7683–7696, 2019.
- 775 Garfinkel, C.I. and C. Schwartz: MJO-related tropical convection anomalies lead to more accurate
- stratospheric vortex variability in subseasonal forecast models, GRL, doi:10.1002/2017GL074470,
- 777 2017.
- 778 Garfinkel, C. I., Son, S.-W., Song, K., Aquila, V., and Oman, L. D.: Stratospheric variability
- contributed to and sustained the recent hiatus in Eurasian winter warming, Geophys. Res. Lett., 44,
- 780 374–382, 2017.
- 781 Garfinkel C.I., C. Schwartz, D. I. P. Domeisen, S-W Son, A. H. Butler, I. P. White: Extratropical
- atmospheric predictability from the Quasi-Biennial Oscillation in subseasonal forecast models , JGR,
- 783 doi: 10.1029/2018JD028724.., 123 (15), 7855-7866, 2018.
- Garfinkel, C. I., A.H. Butler, D. W. Waugh, M. M. Hurwitz, L. M. Polvani: Why might stratospheric
- 785 sudden warmings occur with similar frequency in El Nino and La Nina winters?, J. Geophys. Res.
- 786 Atmos., 117, D19106, doi:10.1029/2012JD017777, 2012.
- 787 Garfinkel, C. I., Feldstein, S. B., Waugh, D. W., Yoo, C., and Lee, S.: Observed connection between
- stratospheric sudden warmings and the Madden-Julian Oscillation, Geophys. Res. Lett., 39, L18807,
- 789 2012.
- 790 Gerber, E. P., A. Butler, N. Calvo, A. Charlton-Perez, M. Giorgetta, E. Manzini, J. Perlwitz, L. M.
- 791 Polvani, F. Sassi, A. A. Scaife, T. A. Shaw, S.-W. Son and S. Watanabe: Assessing and
- 792 Understanding the Impact of Stratospheric Dynamics and Variability on the Earth System. Bull.
- 793 Amer. Met. Soc., 93, 845-859, 2012.
- Givon, Y., Garfinkel, C.I. and White, I.: Transient extratropical response to solar ultraviolet radiation
- in the Northern Hemisphere winter. Journal of Climate, 34(9), pp.3367-3383, 2021.
- 796 Graversen, R. G., and B. Christiansen: Downward propagation from the stratosphere to the
- troposphere: A comparison of the two hemispheres. J. Geophys. Res. Atmos., 108, 4780, 2003.
- 798 Gray, L. J., et al.: Solar influences on climate, Rev. Geophys., 48, RG4001, 2010.
- 799 Gray, L. J., Scaife, A. A., Mitchell, D. M., Osprey, S., Ineson, S., Hardiman, S., Butchart, N., Knight,
- 300 J., Sutton, R., and Kodera, K.: A lagged response to the 11 year solar cycle in observed winter
- Atlantic/European weather patterns, J. Geophys. Res. Atmos., 118, 13,405–13, 420, 2013.
- 802 Gray, L. J., Anstey, J. A., Kawatani, Y., Lu, H., Osprey, S., and Schenzinger, V.: Surface impacts of
- the Quasi-Biennial Oscillation, Atmos. Chem. Phys., 18, 8227–8247, 2018.

- Hall, R. J., Scaife, A. A., Hanna, E., Jones, J. M., & Erdélyi, R.: Simple statistical probabilistic
- forecasts of the winter NAO. Weather and Forecasting, 32(4), 1585–1601, 2017.
- Harari, O., C.I. Garfinkel, O. Morgenstern, G. Zeng, S. Tilmes, D. Kinnison, M. Deushi, P. Jockel, A.
- Pozzer, and F. M. O'Connor, Influence of Arctic Stratospheric Ozone on Surface Climate in CCMI
- 808 models, ACP, doi: 10.5194/acp-2018-1031, 2019.
- Hardiman, Steven C., Adam A. Scaife, Nick. J. Dunstone, and Lin Wang: Subseasonal vacillations in
- the winter stratosphere, Geophys. Res. Lett., 47, e2020GL087766, 2020a.
- Hardiman, Steven C., Nick J. Dunstone, Adam A. Scaife, Doug M. Smith, Jeff R. Knight, Paul
- Davies, Martin Claus, and Richard J. Greatbatch (2020b), Predictability of European winter 2019/20:
- 813 Indian Ocean dipole impacts on the NAO, Atmos. Sci. Lett. 2020, e1005, 2020b.
- Hardiman, Steven C., Neal Butchart, Andrew J. Charlton-Perez, Tiffany A. Shaw, Hideharu Akiyoshi,
- Andreas Baumgaertner, Slimane Bekki, Peter Braesicke, Martyn Chipperfield, Martin Dameris,
- Rolando R. Garcia, Martine Michou, Steven Pawson, Eugene Rozanov, and Kiyotaka Shibata:
- 817 Improved predictability of the troposphere using stratospheric final warmings, J. Geophys. Res., 116,
- 818 D18113, 11 PP., 2011.
- Harnik N.. Observed stratospheric downward reflection and its relation to upward pulses of wave
- 820 activity. J. Geophys. Res., 114, Article D08120, 2009.
- Harnik, N. and R.S. Lindzen. The effect of reflecting surfaces on the vertical structure and variability
- of stratospheric planetary waves. J. Atmos. Sci., 58, 2872-2894, 2001.
- Haynes, P. H., McIntyre, M. E., Shepherd, T. G., Marks, C. J., & Shine, K. P.: On the "Downward
- 824 Control" of Extratropical Diabatic Circulations by Eddy-Induced Mean Zonal Forces, *Journal of*
- 825 *Atmospheric Sciences*, 48(4), 651-678, 1991.
- Haynes, P., Hitchcock, P., Hitchman, M., Yoden, S., Hendon, H., Kiladis, G., Kodera, K., Simpson,
- 827 I.: The influence of the stratosphere on the tropical troposphere. J. Meteor. Soc. Japan, 2021.
- Haynes, P., Hitchcock, P., Hitchman, M., Yoden, S., Hendon, H., Kiladis, G., Kodera, K., Simpson,
- 829 I.: The influence of the stratosphere on the tropical troposphere. J. Meteor. Soc. Japan, (accepted
- subject to final small changes), 2021.
- Henderson GR, Peings Y, Furtado JC, Kushner PJ: Snow-atmosphere coupling in the northern
- 832 hemisphere. Nat Clim Change 8(11), 954–963, 2018
- Hendon, H. H., and S. Abhik: Differences in vertical structure of the Madden-Julian oscillation
- associated with the quasi-biennial oscillation. Geophys. Res. Lett., 45, 4419–4428, 2018.
- Hendon, H. H., E. -P. Lim, and S. Abhik: Impact of Interannual Ozone Variations on the Downward
- 836 Coupling of the 2002 Southern Hemisphere Stratospheric Warming. J. Geophys. Res. Atmos., 1–16,
- 837 doi:10.1029/2020JD032952, 2020.
- Hermanson L., R. Eade, N.H. Robinson, M.B. Andrews, J.R. Knight, A.A. Scaife and D.M. Smith,
- Forecast cooling of the Atlantic subpolar gyre and associated impacts. Geophys. Res. Lett., 41, 5167-
- 840 5174, 2014.
- Hitchcock, P., Shepherd, T.G., Taguchi, M., Yoden, S. and Noguchi, S.: Lower-stratospheric radiative
- damping and Polar-night Jet Oscillation events. J. Atmos. Sci., 70, 1391–1408, 2013.
- Hitchcock, P., & Simpson, I. R.: The Downward Influence of Stratospheric Sudden Warmings,
- *Journal of the Atmospheric Sciences*, 71(10), 3856-3876, 2014.

- Hitchcock, P., Haynes, P.H.: Stratospheric control of planetary waves. Geophys. Res. Lett., 43,
- 846 11,884–11,892, doi:10.1002/2016GL071372, 2016.
- Holton, J.R., H.-C. Tan. The influence of the equatorial quasi-biennial oscillation on the global
- 848 circulation at 50 mb. J. Atmos. Sci., 37, 2200-2208, 1980.
- Honda, M., Inoue, J., and Yamane, S.: Influence of low Arctic sea-ice minima on anomalously cold
- 850 Eurasian winters, Geophys. Res. Lett., 36, L08707, doi:10.1029/2008GL037079, 2009.
- Huebener H, Cubasch U, Langematz U, Spangehl T, Niehörster F, Fast I and Kunze M.: Ensemble
- climate simulations using a fully coupled ocean–troposphere–stratosphere general circulation
- 853 modelPhil. Trans. R. Soc. A.3652089–2101, 2007.
- Hurwitz, M. M., Newman, P. A., and Garfinkel, C. I.: On the influence of North Pacific sea surface
- temperature on the Arctic winter climate, *J. Geophys. Res.*, 117, D19110, 2012.
- 856 Ineson S. and Scaife A.A.: The role of the stratosphere in the European climate response to El Nino.
- 857 Nature Geoscience, 2, 32-36, 2009.
- 858 Ineson S., A. A. Scaife, J.R. Knight, J.C. Manners, N.J. Dunstone, L.J. Gray and J.D. Haigh: Solar
- Forcing of Winter Climate Variability in the Northern Hemisphere. Nat. Geosci., 4, 753-757, 2011.
- 860 Ineson, S., Maycock, A., Gray, L. et al.: Regional climate impacts of a possible future grand solar
- 861 minimum. *Nat Commun* **6,** 7535, 2015.
- 862 Inoue, M., M. Takahashi, H. Naoe: Relationship between the stratospheric quasi-biennial oscillation
- and tropospheric circulation in northern autumn. J. Geophys. Res. Atmos., 116, 2011.
- Jain, S., Scaife, A.A. & Mitra, A.K.: Skill of Indian summer monsoon rainfall prediction in multiple
- seasonal prediction systems. *Clim Dyn* **52**, 5291–5301, 2019.
- Jaiser R, Dethloff K, Handorf D. Stratospheric response to arctic sea ice retreat and associated
- planetary wave propagation changes. Tellus A Dyn Meteorol Oceanogr 65(1):19375, 2013.
- Jia, L., Yang, X., Vecchi, G., Gudgel, R., Delworth, T., Fueglistaler, S., Lin, P., Scaife, A. A.,
- 869 Underwood, S., and Lin, S.: Seasonal Prediction Skill of Northern Extratropical Surface Temperature
- 870 Driven by the Stratosphere, Journal of Climate, 30(12), 4463-4475, 2017.
- Johnson, S. J., Stockdale, T. N., Ferranti, L., Balmaseda, M. A., Molteni, F., Magnusson, L., Tietsche,
- 872 S., Decremer, D., Weisheimer, A., Balsamo, G., Keeley, S. P. E., Mogensen, K., Zuo, H., and Monge-
- 873 Sanz, B. M.: SEAS5: the new ECMWF seasonal forecast system, Geosci. Model Dev., 12, 1087–
- 874 1117, 2019.
- Joshi, M. M., A. J. Charlton, and A. A. Scaife: On the influence of stratospheric water vapor changes
- on the tropospheric circulation. Geophys. Res. Lett., 33, L09806, 2006.
- Jucker, M., & Reichler, T.: Dynamical precursors for statistical prediction of stratospheric sudden
- warming events. Geophysical Research Letters, 45, 13,124–13,132, 2018.
- Jucker, M., Reichler, T., & Waugh, D. W.: How frequent are Antarctic sudden stratospheric warmings
- in present and future climate? *Geophysical Research Letters*, 48, e2021GL093215, 2021.
- Kang, I.S., Kug, JS., Lim, MJ. et al: Impact of transient eddies on extratropical seasonal-mean
- predictability in DEMETER models. Clim Dyn 37, 509–519, 2011.
- 883 Kang, S.M., L.M. Polvani, J.C. Fyfe and M. Sigmond: Impact of Polar Ozone Depletion on
- Subtropical Precipitation, Science, 332, 951-954, 2011.

- 885 Karpechko A., N.P. Gillett, G.J. Marshall and A.A. Scaife: Stratospheric influence on circulation
- changes in the Southern Hemisphere troposphere in coupled climate models. Geophys. Res. Lett.,
- 887 L20806, 2008.
- 888 Karpechko, A. Y., and Manzini, E.: Stratospheric influence on tropospheric climate change in the
- Northern Hemisphere, J. Geophys. Res., 117, D05133, doi:10.1029/2011JD017036, 2012.
- 890 Karpechko, A. Yu., J. Perlwitz, and E. Manzini: A model study of tropospheric impacts of the Arctic
- 891 ozone depletion 2011, J. Geophys. Res. Atmos., 119, 7999–8014, 2014
- 892 Karpechko A. Yu., P. Hitchcock, D. H.W. Peters, A. Schneidereit: Predictability of downward
- propagation of sudden stratospheric warmings, Quart. J. Roy. Meteor. Soc., v. 143, 704, 1459-1470,
- 894 2017.
- 895 Karpechko A. Y., A. Charlton-Perez, M. Balmaseda, N. Tyrrell, and F. Vitart: Predicting sudden
- stratospheric warming 2018 and its climate impacts with a multimodel ensemble, Geophys. Res. Lett.,
- 897 45, 13538–13546, 2018.
- 898 Karpechko A. Yu.: Predictability of Sudden Stratospheric Warmings in the ECMWF Extended-Range
- 899 Forecast System, Mon. Wea. Rev., 146(4), 1063–1075, doi: 10.1175/MWR-D-17-0317.1, 2018.
- Warpechko, AY, Tyrrell, NL, Rast, S.: Sensitivity of QBO teleconnection to model circulation biases.
- 901 Q J R Meteorol Soc. 147, 2147–2159, 2021.
- 902 Kautz, L. A., Polichtchouk, I., Birner, T., Garny, H., and Pinto, J. G.: Enhanced extended-range
- predictability of the 2018 late-winter Eurasian cold spell due to the stratosphere, Q. J. Roy.
- 904 Meteorol.Soc., 146, 1040–1055, 2019.
- 905 Kidston, J., A. A. Scaife, S. C. Hardiman, D. M. Mitchell, N. Butchart, M. P. Baldwin and L. J. Gray.
- 906 Stratospheric influence on tropospheric jet streams, storm tracks and surface weather, Nature
- 907 Geoscience, 8, 433–440, 2015.
- 908 Kim, BM., Son, SW., Min, SK. et al.: Weakening of the stratospheric polar vortex by Arctic sea-ice
- 909 loss. Nat Comm., 5, 4646, 2014.
- 910 Kim, H., Caron, J. M., Richter, J. H., & Simpson, I. R.: The lack of QBO-MJO connection in CMIP6
- 911 models. Geophysical Research Letters, 47, e2020GL087295, 2020.
- 912 King, M.P., Hell, M. & Keenlyside, N. Investigation of the atmospheric mechanisms related to the
- autumn sea ice and winter circulation link in the Northern Hemisphere. Clim. Dyn., 46, 1185–1195,
- 914 2016.
- 915 King, A. D., A.H. Butler, M. Jucker, N.O. Earl, and I. Rudeva, Observed relationships between
- 916 sudden stratospheric warmings and European climate extremes, J. Geophys. Res., 124, 13943-13961,
- 917 2019.
- 918 Kodera, K. and Kuroda, Y. Dynamical response to the solar cycle. J. Geophys. Res. 107, 4749, 2002.
- 919 Kolstad, E. W., & Screen, J. A.. Non-stationary relationship between autumn Arctic sea ice and the
- 920 winter North Atlantic Oscillation. Geophysical Research Letters, 46, 7583–7591, 2019.
- 921 Knight J. et al.: Predictability of European winters 2017/2018 and 2018/2019: Contrasting Influences
- from the Tropics and Stratosphere. Atm. Sci. Lett., e1009, doi.org/10.1002/asl.1009, 2020.
- 923 Kodera, K., Yamazaki, K., Chiba, M., Shibata, K.: Downward propagation of upper stratospheric
- mean zonal wind perturbation to the troposphere. Geophys. Res. Lett., 17, 9, 0094-8276, 1990.

- 925 Kodera, K.: On the origin and nature of the interannual variability of the winter stratospheric
- circulation in the northern hemisphere, J. Geophys. Res., 100 (D7), 14077–14087, 1995.
- 927 Kodera, K., Mukougawa, H., and Itoh, S.: Tropospheric impact of reflected planetary waves from the
- 928 stratosphere, Geophys. Res. Lett., 35, L16806, doi:10.1029/2008GL034575, 2008.
- 929 Kretschmer, M., Coumou, D., Donges, J.F., and Runge, J.: Using causal effect networks to analyze
- 930 different Arctic drivers of midlatitude winter circulation. *J. Clim.*, 29, 4069–4081, 2016.
- 931 Kretschmer, M., Zappa, G. and Shepherd, T.G.: The role of Barents-Kara sea ice loss in projected
- 932 polar vortex changes. Wea. Clim. Dyn., 1, 715–730, 2020.
- 933 Kretschmer, M., Adams, S.V., Arribas, A., Prudden, R., Robinson, N., Saggioro, E. and Shepherd,
- T.G.: Quantifying causal pathways of teleconnections. *Bull. Amer. Meteor. Soc.*, in press, doi:
- 935 10.1175/BAMS-D-20-0117.1, 2021.
- 836 Kuhn, T. S.: The Structure of Scientific Revolutions. University of Chicago Press, 1970.
- 937 Kunz, T., & Greatbatch, R. J.: On the Northern Annular Mode Surface Signal Associated with
- 938 Stratospheric Variability, *Journal of the Atmospheric Sciences*, 70(7), 2103-2118, 2013.
- 839 Kuroda, Y.: Role of the stratosphere on the predictability of medium-range weather forecast: A case
- study of winter 2003–2004. Geophysical Research Letters, 35, L19701, 2008.
- Kushnir, Y., Scaife, A.A., Arritt, R. et al.: Towards operational predictions of the near-term climate.
- 942 Nature Clim. Change 9, 94–101, 2019.
- Kushner P.J. and L.M. Polvani: Stratosphere-troposphere coupling in a relatively simple AGCM: The
- 944 role of eddies, J. Climate, 17, 629-639, 2004.
- Labe, Z., Peings, Y., & Magnusdottir, G.: The effect of QBO phase on the atmospheric response to
- projected Arctic sea ice loss in early winter. Geophysical Research Letters, 46, 7663–7671, 2019.
- Labitzke, K.: On the Mutual Relation between Stratosphere and Troposphere during Periods of
- 948 Stratospheric Warmings in Winter, J. Appl. Met. Climatol., 4(1), 91-99, 1965.
- 949 Lai, W. K. M., Robson, J. I., Wilcox, L. J., & Dunstone, N. Mechanisms of Internal Atlantic
- 950 Multidecadal Variability in HadGEM3-GC3.1 at Two Different Resolutions, Journal of Climate,
- 951 2021.
- Lawrence, Z. D., Perlwitz, J., Butler, A. H., Manney, G. L., Newman, P. A., Lee, S. H., & Nash, E.
- 953 R.: The remarkably strong Arctic stratospheric polar vortex of winter 2020: Links to record-breaking
- Arctic Oscillation and ozone loss. Journal of Geophysical Research: Atmospheres, 125,
- 955 e2020JD033271, 2020.
- Leckebusch, G. C., Ulbrich, U., Fröhlich, L., and Pinto, J. G.: Property loss potentials for European
- 957 midlatitude storms in a changing climate, Geophys. Res. Lett., 34, L05703, 2007.
- 958 Lee, SH, Charlton-Perez, AJ, Furtado, JC, Woolnough, SJ. Representation of the Scandinavia-
- Greenland pattern and its relationship with the polar vortex in S2S forecast models. Q J R Meteorol
- 960 *Soc.* 146: 4083–4098, 2020a.
- 961 Lee, S.H., Z.D. Lawrence, A.H. Butler, and A. Karpechko. Seasonal forecasts of the exceptional
- Northern Hemisphere winter of 2020, Geophys. Res. Lett., 47, e2020GL090328, 2020b.
- Leung, T.Y., Leutbecher, M., Reich, and Shepherd, T.G.. Impact of the mesoscale range on error
- growth and the limits to atmospheric predictability. J. Atmos. Sci., 77, 3769–3779, 2020.

- 965 L'Heureux, M.L., Levine, A.F.Z., Newman, M., Ganter, C., Luo, J.-J., Tippett, M.K. and Stockdale,
- 966 T.N.: ENSO Prediction. In El Niño Southern Oscillation in a Changing Climate (eds M.J. McPhaden,
- 967 A. Santoso and W. Cai), 2020.
- Lim, E.-P., H. H. Hendon, and D. W. J. Thompson: Seasonal evolution of stratosphere-troposphere
- oupling in the Southern Hemisphere and implications for the predictability of surface climate. J.
- 970 Geophys. Res. Atmos., 123, 12,002–12,016, 2018.
- Lim, E.-P., H. H. Hendon, G. Boschat, D. Hudson, D. W. J. Thompson, A. J. Dowdy, and J. Arblaster:
- Australian hot and dry extremes induced by weakening of the stratospheric polar vortex. Nature
- 973 Geoscience, 12 (11), 896-901, 2019.
- 2019 Lim, E.-P., and Coauthors: The 2019 Southern Hemisphere polar stratospheric warming and its
- 975 impacts. BAMS. 102 (6), E1150-E1171, 2021.
- 276 Lim, Y., S.-W. Son, A. G. Marshall, H. H. Hendon, and K.-H. Seo: Influence of the QBO on MJO
- prediction skill in the subseasonal-to-seasonal prediction models, Climate Dynamics, 53, 1681-1695,
- 978 2019.
- 279 Lin, H., Brunet, G., & Derome, J.: An Observed Connection between the North Atlantic Oscillation
- and the Madden–Julian Oscillation, *Journal of Climate*, 22(2), 364-380, 2009.
- 981 Lockwood, M.: Solar change and climate: An update in the light of the current exceptional solar
- 982 minimum, Proc. R. Soc. A, 466, 303–329, 2010.
- Lorenz, E.N.: The predictability of a flow which possesses many scales of motion. Tellus, 21: 289-
- 984 307, 1969.
- 985 Manzini, E., Giorgetta, M. A., Esch, M., Kornblueh, L., & Roeckner, E.: The Influence of Sea Surface
- Temperatures on the Northern Winter Stratosphere: Ensemble Simulations with the MAECHAM5
- 987 Model, Journal of Climate, 19(16), 3863-3881, 2006.
- 988 Manzini, E., Karpechko, A. Y., and Kornblueh, L.: Nonlinear Response of the Stratosphere and the
- North Atlantic-European Climate to Global Warming, Geophys. Res. Lett., 45, 4255–4263, 2018.
- 990 Mariotti, A., C. Baggett, E. A. Barnes, E. Becker, A.H. Butler, D.C. Collins, P.A. Dirmeyer, L.
- 991 Ferranti, N.C. Johnson, J. Jones, B. P. Kirtman, A.L. Lang, A. Molod, M. Newman, A.W. Robertson,
- 992 S. Schubert, Waliser, D.E. and J. Albers, Windows of opportunity for skillful forecasts subseasonal to
- 993 seasonal and beyond, BAMS, 101, E608-E625, 2020.
- 994 Marsh, D. R., Garcia, R. R., Kinnison, D. E., Boville, B. A., Sassi, F., Solomon, S. C., and Matthes,
- 995 K.: Modeling the whole atmosphere response to solar cycle changes in radiative and geomagnetic
- 996 forcing, J. Geophys. Res., 112, D23306, 2007.
- 997 Marshall A. and A.A. Scaife: Improved predictability of stratospheric sudden warming events in an
- 998 AGCM with enhanced stratospheric resolution. J. Geophys. Res., 115, D16114, 2010.
- 999 Marshall, A.G., Hendon, H.H., Son, S.W. and Lim, Y.: Impact of the quasi-biennial oscillation on
- predictability of the Madden–Julian oscillation. Climate Dynamics, 49(4), pp.1365-1377, 2017.
- Martin, Z., Wang, S., Nie, J., & Sobel, A.: The Impact of the QBO on MJO Convection in Cloud-
- 1002 Resolving Simulations, Journal of the Atmospheric Sciences, 76(3), 669-688, 2019.
- 1003 Martin, Z., Son, SW., Butler, A. et al.: The influence of the quasi-biennial oscillation on the Madden-
- Julian oscillation. Nat Rev Earth Environ 2, 477–489, 2021.
- 1005 Matthias, V., & Kretschmer, M.: The Influence of Stratospheric Wave Reflection on North American
- 1006 Cold Spells, Monthly Weather Review, 148(4), 1675-1690, 2020.

- Maycock, A. C., Ineson, S., Gray, L. J., Scaife, A. A., Anstey, J. A., Lockwood, M., Butchart, N.,
- Hardiman, S. C., Mitchell, D. M., and Osprey, S. M.: Possible impacts of a future grand solar
- minimum on climate: Stratospheric and global circulation changes, J. Geophys. Res. Atmos., 120,
- 1010 9043-9058, 2015.
- 1011 McKenna, C.M., Bracegirdle, T.J., Shuckburgh, E.F., Haynes, P.H., Joshi, M.M.: Arctic sea-ice loss
- in different regions leads to contrasting Northern Hemisphere impacts. Geophys. Res. Lett., 45, 945–
- 1013 954, 2018.
- 1014 MacLachlan, C., Arribas, A., Peterson, K.A., Maidens, A., Fereday, D., Scaife, A.A., Gordon, M.,
- 1015 Vellinga, M., Williams, A., Comer, R.E., Camp, J., Xavier, P. and Madec, G.: Global Seasonal
- forecast system version 5 (GloSea5): a high-resolution seasonal forecast system. Q.J.R. Meteorol.
- 1017 Soc., 141: 1072-1084, 2015.
- Mann, M. E., J. Park, and R. S. Bradley. Global interdecadal and century-scale climate oscillations
- 1019 during the past 5 centuries, Nature, 378, 266 270, 1995.
- McLandress, C. and Shepherd, T.G.: Impact of climate change on stratospheric sudden warmings as
- simulated by the Canadian Middle Atmosphere Model. J. Clim., 22, 5449–5463, 2009.
- 1022 McLandress, C., Shepherd, T.G., Scinocca, J.F., Plummer, D.A., Sigmond, M., Jonsson, A.I. and
- Reader, M.C.: Separating the dynamical effects of climate change and ozone depletion: Part 2.
- 1024 Southern Hemisphere Troposphere. J. Clim., 24, 1850–1868, 2011.
- Meehl G. et al.: Initialised Earth System Prediction from Subseasonal to Decadal Timescales. Nat.
- 1026 Rev. Earth. Environ. 2, 340–357, 2021.
- Merryfield, W.J., J. Baehr, L. Batté, E.J. Becker, A.H. Butler, et al., Current and emerging
- developments in subseasonal to decadal prediction, BAMS, 101, E869–E896, 2020.
- Merryfield, W. J., F. J. Doblas-Reyes, L. Ferranti, J.-H. Jeong, Y. J. Orsolini, R. I. Saurral, A. A.
- Scaife, M. A. Tolstykh, and M. Rixen. Advancing climate forecasting. EOS, 98, 17-21, 2017.
- 1031 Merryfield, W. J., Lee, W., Boer, G. J., Kharin, V. V., Scinocca, J. F., Flato, G. M., Ajayamohan, R.
- S., Fyfe, J. C., Tang, Y., & Polavarapu, S.: The Canadian Seasonal to Interannual Prediction System.
- Part I: Models and Initialization, Monthly Weather Review, 141(8), 2910-2945, 2013.
- 1034 Mechoso, C.R., A. Molod, A. O'Neill, R.B. Pierce, W.J. Randel, R.B. Rood, F. Wu: The GCM-
- 1035 Reality Intercomparison Project for SPARC (GRIPS): Scientific Issues and Initial Results. Bull.
- 1036 Amer. Met. Soc., 81, 781-796, 2000.
- Molina, M., Rowland, F.: Stratospheric sink for chlorofluoromethanes: chlorine atom-catalysed
- 1038 destruction of ozone. Nature 249, 810–812, 1974.
- Monge-Sanz, B., et al.: A stratospheric prognostic ozone for seamless Earth System Models:
- performance, impacts and future. Atmos. Chem. Phys. Disc., doi: 10.5194/acp-2020-1261, 2021.
- Müller, W. A., and Roeckner, E.: ENSO impact on midlatitude circulation patterns in future climate
- change projections, Geophys. Res. Lett., 33, L05711, 2006.
- Mukougawa, H., Sakai, H., & Hirooka, T.: High sensitivity to the initial condition for the prediction
- of stratospheric sudden warming. Geophysical Research Letters, 32, L17806, 2005.
- Mukougawa, H., Hirooka, T., and Kuroda, Y.: Influence of stratospheric circulation on the
- 1046 predictability of the tropospheric Northern Annular Mode, Geophys. Res. Lett., 36, L08814,
- doi:10.1029/2008GL037127, 2009.

- Mukougawa, H., Noguchi, S., Kuroda, Y., Mizuta, R., & Kodera, K.: Dynamics and predictability of
- downward-propagating stratospheric planetary waves observed in March 2007. Journal of the
- 1050 Atmospheric Sciences, 74(11), 3533–3550, 2017.
- Nie Y. et al.: Stratospheric initial conditions provide seasonal predictability of the North Atlantic and
- 1052 Arctic Oscillations. Env. Res. Lett., 14, 3, 2019.
- Noguchi, S., Mukougawa, H., Kuroda, Y., Mizuta, R., Yabu, S., & Yoshimura, H.: Predictability of
- the stratospheric polar vortex breakdown: An ensemble reforecast experiment for the splitting event in
- January 2009. Journal of Geophysical Research: Atmospheres, 121, 3388–3404, 2016.
- Noguchi, S., Kuroda, Y., Mukougawa, H., Mizuta, R., & Kobayashi, C.: Impact of satellite
- observations on forecasting sudden stratospheric warmings. Geophysical Research Letters, 47,
- 1058 e2019GL086233, 2020a.
- Noguchi, S., Kuroda, Y., Kodera, K., & Watanabe, S.: Robust enhancement of tropical convective
- activity by the 2019 Antarctic sudden stratospheric warming. Geophysical Research Letters, 47,
- 1061 e2020GL088743, 2020b.
- Oehrlein J., G. Chiodo and L.M. Polvani: The effect of interactive ozone chemistry on weak and
- strong stratospheric polar vortex events, Atmos. Chem. Phys., 20, 10531-10544, 2020.
- Omrani, NE., Keenlyside, N.S., Bader, J. et al.: Stratosphere key for wintertime atmospheric response
- to warm Atlantic decadal conditions. Clim Dyn 42, 649–663, 2014.
- 1066 O'Reilly CH, Weisheimer A, Woollings T, Gray LJ, MacLeod D: The importance of stratospheric
- initial conditions for winter North Atlantic Oscillation predictability and implications for the signal-
- to-noise paradox. Q J R Meteorol Soc 145(718):131–146, 2019.
- Orsolini YJ, Senan R, Benestad RE, Melsom A. Autumn atmospheric response to the 2007 low Arctic
- sea ice extent in coupled ocean–atmosphere hindcasts. Clim Dyn 38:2437–2448, 2012.
- 1071 Owens, M.J., Mike Lockwood, Ed Hawkins, Ilya Usoskin, Gareth S. Jones, Luke Barnard, Andrew
- 1072 Schurer and John Fasullo: The Maunder minimum and the Little Ice Age: an update from recent
- reconstructions and climate simulations. J. Space Weather Space Clim., 7 (2017) A33, 2017.
- Pawson, S. and B. Naujokat: The cold winters of the middle 1990s in the northern lower stratosphere.
- 1075 J. Geophys. Res., 104, 14 209–14 222, 1999.
- 1076 Pawson, S., Kodera, K., Hamilton, K., Shepherd, T. G., Beagley, S. R., Boville, B. A., Farrara, J. D.,
- Fairlie, T. D. A., Kitoh, A., Lahoz, W. A., Langematz, U., Manzini, E., Rind, D. H., Scaife, A. A.,
- 1078 Shibata, K., Simon, P., Swinbank, R., Takacs, L., Wilson, R. J., Al-Saadi, J. A., Amodei, M., Chiba,
- 1079 M., Coy, L., de Grandpré, J., Eckman, R. S., Fiorino, M., Grose, W. L., Koide, H., Koshyk, J. N., Li,
- D., Lerner, J., Mahlman, J. D., McFarlane, N. A., Mechoso, C. R., Molod, A., O'Neill, A., Pierce, R.
- 1081 B., Randel, W. J., Rood, R. B., & Wu, F.: The GCM-Reality Intercomparison Project for SPARC
- 1082 (GRIPS): Scientific Issues and Initial Results, Bulletin of the American Meteorological Society, 81(4),
- 1083 781-796, 2000.
- 1084 Peings, Y. E. Brun, V. Mauvais, H. Douville: How stationary is the relationship between Siberian
- snow and Arctic Oscillation over the 20th century? Geophys. Res. Lett., 40, 183-188, 2013.
- Perlwitz, J., and N. Harnik: Observational evidence of a stratospheric influence on the troposphere by
- planetary wave reflection, J. Clim., 16, 3011–3026, 2003.
- Perlwitz, J., & Graf, H.: The Statistical Connection between Tropospheric and Stratospheric
- 1089 Circulation of the Northern Hemisphere in Winter, *Journal of Climate*, 8(10), 2281-2295, 1995.

- 1090 Plumb, R. A., and Semeniuk, K.: Downward migration of extratropical zonal wind anomalies, J.
- 1091 Geophys. Res., 108, 4223, doi:10.1029/2002JD002773, D7, 2003.
- 1092 Polvani L.M. and P.J. Kushner: Tropospheric response to stratospheric perturbations in a relatively
- simple general circulation model, Geophys. Res. Lett., 29, no. 7 (2002) Liu C., B. Tian, K.-F. Li, G.L.
- Manney, N.J. Livesey, Y.L. Yung, D.E. Waliser. Northern Hemisphere mid-winter vortex-
- displacement and vortex-split stratospheric sudden warmings: influence of the Madden-Julian
- Oscillation and Quasi-Biennial Oscillation. J. Geophys. Res. Atmos., 119, 12,599-12,620, 2014.
- 1097 Polvani, L.M., D.W. Waugh, G.J.P. Correa and S.-W. Son: Stratospheric ozone depletion: the main
- driver of 20th Century atmospheric circulation changes in the Southern Hemisphere, J. Climate, 24,
- 1099 795-812, 2011.
- 1100 Polvani L.M., L. Sun, A.H. Butler, J.H. Richter and C. Deser: Distinguishing stratospheric sudden
- warmings from ENSO as key drivers of wintertime climate variability over the North Atlantic and
- 1102 Eurasia, J. Clim., 30, 1959-1969, 2017.
- 1103 Polvani, L. M., and Kushner, P. J., Tropospheric response to stratospheric perturbations in a relatively
- simple general circulation model, Geophys. Res. Lett., 29(7), doi:10.1029/2001GL014284, 2002.
- Polvani, L.M., L. Sun, A.H. Butler, J.H. Richter and C. Deser: Distinguishing stratospheric sudden
- warmings from ENSO as key drivers of wintertime climate variability over the North Atlantic and
- 1107 Eurasia, J. Climate, 30, 1959-1969, 2017.
- 1108 Previdi, M. and L.M. Polvani: Climate System Response to Stratospheric Ozone Depletion and
- 1109 Recovery, Quart. J. Roy. Meteor. Soc., 140, 2401-2419, 2014.
- Purich, A., and S.-W. Son: Impact of Antarctic ozone depletion and recovery on Southern Hemisphere
- precipitation, evaporation and extreme changes, Journal of Climate, 25, 3145-3154, 2012.
- 1112 Rao, J., C. I. Garfinkel, and I. P. White: Predicting the downward and surface influence of the
- 1113 February 2018 and January 2019 sudden stratospheric warming events in subseasonal to seasonal
- 1114 (S2S) models, Journal of Geophysical Research: Atmospheres, 125, 2020a
- Rao, J., C. I. Garfinkel, I. P. White, C. Schwartz: How does the Quasi-Biennial Oscillation affect the
- boreal winter tropospheric circulation in CMIP5/6 models?, Journal of Climate, 33, 8975-8996,
- 1117 2020b.
- 1118 Rao, J. and C. I. Garfinkel: CMIP5/6 Models Project Little Change in the Statistical Characteristics of
- 1119 Sudden Stratospheric Warmings in the 21st Century, Environmental Research Letters, 2020c.
- 1120 Rao, J., C. I. Garfinkel, and I. P. White: Projected strengthening of the extratropical surface impacts
- of the stratospheric Quasi Biennial Oscillation, Geophysical Research Letters, 47, e2020GL089149,
- 1122 2020d.
- Rao, J., C. I. Garfinkel, I. P. White, C. Schwartz: The Southern Hemisphere Minor Sudden
- 1124 Stratospheric Warming in September 2019 and its predictions in S2S Models, Journal of Geophysical
- 1125 Research: Atmospheres, 125, e2020JD032723, doi:10.1029/2020JD032723, 2020e.
- 1126 Rao, J. and Garfinkel, C.I.: The Strong Stratospheric Polar Vortex in March 2020 in Sub-Seasonal to
- 1127 Seasonal Models: Implications for Empirical Prediction of the Low Arctic Total Ozone Extreme.
- Journal of Geophysical Research: Atmospheres, 126(9), p.e2020JD034190., 2021a.
- Rao, J., Garfinkel, C. I., & White, I. P. Development of the Extratropical Response to the Stratospheric
- 1130 Quasi-Biennial Oscillation. Journal of Climate, 34(17), 7239-7255, 2021. Reichler, T., Kim, J.,
- 1131 Manzini, E., & Kröger, J. A stratospheric connection to Atlantic climate variability. Nature
- 1132 Geoscience, 5(11), 783–787, 2012.

- Richter, J. H., Pegion, K., Sun, L., Kim, H., Caron, J. M., Glanville, A., LaJoie, E., Yeager, S., Kim,
- W. M., Tawfik, A., & Collins, D.: Subseasonal Prediction with and without a Well-Represented
- Stratosphere in CESM1, Weather and Forecasting, 35(6), 2589-2602, 2020a.
- 1136 Richter, J. H., Anstey, J. A., Butchart, N., Kawatani, Y., Meehl, G. A., Osprey, S., & Simpson, I.
- 1137 R.: Progress in simulating the quasi-biennial oscillation in CMIP models. Journal Geophysical
- 1138 Research: Atmospheres, 125, e2019JD032362, 2020b.
- Richter, JH, Butchart, N, Kawatani, Y, et al.: Response of the Quasi-Biennial Oscillation to a
- warming climate in global climate models. Q J R Meteorol Soc.; 1–29, 2020c.
- 1141 Riddle, E.E., A.H. Butler, J.C. Furtado, J.L. Cohen, and A. Kumar, CFSv2 ensemble prediction of the
- wintertime Arctic Oscillation, Climate Dynamics, 41, 1099-1116, 2013.
- Rind, D., Suozzo, R., Balachandran, N. K., Lacis, A., & Russell, G.. The GISS global climate-middle
- atmosphere model. Part I: Model structure and climatology. Journal of the Atmospheric Sciences, 45,
- 1145 329–370, 1988.
- 1146 Roff, G., D. W. J. Thompson, and H. Hendon: Does increasing model stratospheric resolution
- improve extended-range forecast skill? Geophys. Res. Lett., 38, L05809, 2011.
- Runde, T., Dameris, M., Garny, H., & Kinnison, D. E.: Classification of stratospheric extreme events
- according to their downward propagation to the troposphere. Geophysical Research Letters, 43(12),
- 1150 6665–6672, 2016.
- 1151 Saito, N., Maeda, S., Nakaegawa, T., Takaya, Y., Imada, Y., & Matsukawa C.: Seasonal predictability
- of the North Atlantic Oscillation and zonal mean fields associated with stratospheric influence in
- JMA/MRI-CPS2. Scientific Online Letters on the Atmosphere, 13, 209–213, 2017.
- Samarasinghe, S. M., Connolly, C., Barnes, E. A., Ebert-Uphoff, I., & Sun, L.: Strengthened causal
- connections between the MJO and the North Atlantic with climate warming. Geophysical Research
- 1156 Letters, 48, e2020GL091168, 2021.
- Sassi, F., Garcia, R. R., Boville, B. A., and Liu, H.: On temperature inversions and the mesospheric
- 1158 surf zone, J. Geophys. Res., 107(D19), 4380, 2002.
- 1159 Scaife A.A., T. Spangehl, D. Fereday, U. Cubasch, U. Langematz, H. Akiyoshi, S. Bekki, P.
- Braesicke, N. Butchart, M. Chipperfield, A. Gettelman, S. Hardiman, M. Michou, E. Rozanov and
- 1161 T.G. Shepherd. Climate Change and Stratosphere-Troposphere Interaction. Clim. Dyn., 38, Page
- 1162 2089-2097, 2012.
- Scaife A.A., N. Butchart, C.D. Warner, D. Stainforth, W.A. Norton and J. Austin. Realistic Quasi-
- Biennial Oscillations in a simulation of the global climate. Geophys. Res. Let. 27, 3481-3484, 2000.
- Scaife A.A., J.R. Knight, G.K. Vallis, C.K. Folland. A stratospheric influence on the winter NAO and
- North Atlantic surface climate. Geophys. Res. Let., 32, L18715, 2005.
- Scaife A.A. and J.R. Knight. Ensemble simulations of the cold European winter of 2005/6. Q. J. R.
- 1168 Meteorol. Soc., 134, 1647-1659, 2008.
- 1169 Scaife A.A., C.K. Folland, L. Alexander, A. Moberg and J.R. Knight: European climate extremes and
- the North Atlantic Oscillation. J. Clim., 21, 72-83, 2008.
- 1171 Scaife, A. A., et al.: Predictability of the quasi-biennial oscillation and its northern winter
- teleconnection on seasonal to decadal timescales, Geophys. Res. Lett., 41, 1752–1758, 2014a.
- 1173 Scaife, A. A., et al.: Skillful long-range prediction of European and North American winters,
- 1174 Geophys. Res. Lett., 41, 2514–2519, 2014b.

- 1175 Scaife A.A., A.-Yu. Karpechko, M.P. Baldwin, A. Brookshaw, A.H. Butler, R. Eade, M. Gordon, C.
- 1176 MacLachlan, N. Martin, N. Dunstone and D. Smith: Seasonal winter forecasts and the stratosphere.
- 1177 Atm. Sci. Lett., 2016.
- 1178 Schwartz, C. and C.I. Garfinkel: Relative Roles of the MJO and Stratospheric Variability in North
- Atlantic and European Winter Climate . J. Geoph. Res, doi: 10.1002/2016JD025829, 2017.
- 1180 Schwartz, C. and C.I. Garfinkel: Troposphere-stratosphere coupling in subseasonal-to-seasonal
- models and its importance for a realistic extratropical response to the Madden-Julian Oscillation.
- Journal of Geophysical Research: Atmospheres, 125(10), p.e2019JD032043, 2020.
- Screen, J.A., Deser, C., Smith, D.M. et al.: Consistency and discrepancy in the atmospheric response
- to Arctic sea-ice loss across climate models. *Nature Geosci* **11**, 155–163, 2018.
- 1185 Screen, J. A. and Blackport, R.: How robust is the atmospheric response to projected Arctic sea ice
- loss across climate models?. Geophysical Research Letters, 46, 11406–11415, 2019.
- 1187 Seviour, W. J. M., S. C. Hardiman, L. J. Gray, N. Butchart, C. MacLachlan, and A. A. Scaife: Skillful
- seasonal prediction of the Southern Annular Mode and Antarctic ozone. J. Clim., 27, 7462–7474,
- 1189 doi:10.1175/JCLI-D-14-00264.1, 2014.
- 1190 Shukla, J.: Dynamical Predictability of Monthly Means. J. Atmos. Sci., 38(12), 2547-2572, 1981.
- Shindell, D. T., Schmidt, G. A., Miller, R. L., and Rind, D.: Northern hemisphere winter climate
- response to greenhouse gas, ozone, solar, and volcanic forcing, J. Geophys. Res., 106(D7), 7193–
- 1193 7210, 2001.
- 1194 Sigmond, M., Scinocca, J.F., Kharin, V.V. and Shepherd, T.G.. Enhanced seasonal forecast skill
- following stratospheric sudden warmings. Nature Geosci., 6, 98–102, 2013.
- 1196 Simpson, I. R., Hitchcock, P., Seager, R., Wu, Y., & Callaghan, P.: The Downward Influence of
- 1197 Uncertainty in the Northern Hemisphere Stratospheric Polar Vortex Response to Climate Change.
- 1198 Journal of Climate, 31(16), 6371–6391., 2018.
- 1199 Sigmond, M., J. F. Scinocca and P. J. Kushner: The impact of the stratosphere on tropospheric climate
- 1200 change. Geophys. Res. Lett., 35, L12706, 2008.
- 1201 Sigmond, M. and J.F. Scinocca: The influence of basic state on the Northern Hemisphere circulation
- response to climate change. J. Clim., 23, 1434-1446, , 2010.
- 1203 Sigmond M., Scinocca J.F., Kharin V.V., Shephgerd T.G.. Enhanced seasonal forecast skill following
- stratospheric sudden warmings. Nature Geoscience, 6, 98-102, 2013.
- Shaw T.A. and J. Perlwitz. The life cycle of Northern Hemisphere downward wave coupling between
- the stratosphere and troposphere. J. Clim., 26, 1745-1763, 2013
- 1207 Smith K.L. and L M Polvani. The surface impacts of Arctic stratospheric ozone anomalies. Environ.
- 1208 Res. Lett. 9 074015, 2014.
- 1209 Solomon, S., Ivy, D. J., Kinnison, D., Mills, M. J., Neely, R. R., & Schmidt, A.: Emergence of healing
- in the Antarctic ozone layer. Science, 353(6296), 269–274, 2016.
- Son, S.W., L.M. Polvani, D.W. Waugh and CCMVal co-authors: The impact of stratospheric ozone
- recovery on the Southern Hemisphere westerly jet, Science, 320, 1486-1489, 2008.
- 1213 Son, S.-W., Purich, A., Hendon, H. H., Kim, B.-M., and Polyani, L. M.: Improved seasonal forecast
- using ozone hole variability?, Geophys. Res. Lett., 40, 6231–6235, 2013.
- 1215 Son, S-W., Y. Lim, C. Yoo, H. H. Hendon, and J. Kim: Stratospheric control of Madden-Julian
- 1216 Oscillation, Journal of Climate, 30, 1909-1922, 2017.

- Son. S-W, B-R. Han, C.I. Garfinkel, and 26 others: Tropospheric jet response to Antarctic ozone
- depletion: An update with Chemistry-Climate Model Initiative (CCMI) models, Environmental
- 1219 Research Letters, 13, 054024, 2018.
- Son, S.-W., H. Kim, K. Song, S.-W. Kim, P. Martineau, Y.-K. Hyun, and Y. Kim: Extratropical
- prediction skill of the subseasonal-to-seasonal (S2S) prediction models, Journal of Geophysical
- 1222 Research Atmosphere, 125, e2019JD031273, 2020.
- Song, K. and Son, S.: Revisiting the ENSO–SSW Relationship, *Journal of Climate*, 31(6), 2133-2143,
- **1224** 2018.
- 1225 Statnaia, I., A. Yu. Karpechko, H. Järvinen: Mechanisms and predictability of Sudden Stratospheric
- Warming in winter 2018, Weather and Climate Dynamics, 2020.
- Steil, B., Dameris, M., Brühl, C. et al.: Development of a chemistry module for GCMs: first results of
- a multiannual integration. Annales Geophysicae 16, 205–228, 1998.
- 1229 Stenchikov, G., Hamilton, K., Stouffer, R. J., Robock, A., Ramaswamy, V., Santer, B., and Graf, H.-
- 1230 F.: Arctic Oscillation response to volcanic eruptions in the IPCC AR4 climate models, J. Geophys.
- 1231 Res., 111, D07107, 2006.
- 1232 Stockdale, T. N., Molteni, F. and Ferranti, L. (2015), Atmospheric initial conditions and the
- predictability of the Arctic Oscillation. Geophys. Res. Lett., 42: 1173–1179. doi:
- 1234 10.1002/2014GL062681.
- Stockdale et al.: Multi-Model Predictions of the Quasi-Biennial Oscillation. Quart. J. Roy. Met. Soc.,
- in press, 2021.
- Sun, L., Deser, C., & Tomas, R. A.: Mechanisms of Stratospheric and Tropospheric Circulation
- Response to Projected Arctic Sea Ice Loss, Journal of Climate, 28(19), 7824-7845, 2015.
- Swinbank, R., Douglas, C.S., Lahoz, W.A., O'neill, A. and Heaps, A.: Middle atmosphere variability
- in the UK Meteorological Office Unified Model. Q.J.R. Meteorol. Soc., 124: 1485-1525, 1998.
- Swingedouw, D., Pablo Ortega, Juliette Mignot, Eric Guilyardi, Valérie Masson-Delmotte, Paul G.
- 1242 Butler, Myriam Khodri, Roland Séférian: Bidecadal North Atlantic ocean circulation variability
- 1243 controlled by timing of volcanic eruptions. Nature Communications, 2015; 6: 6545, 2015.
- 1244 Taguchi, M.: Seasonal Winter forecasts of the northern stratosphere and troposphere: Results from
- JMA seasonal hindcast experiments. Journal of the Atmospheric Sciences, 75 (3), 827-840, 2018.
- 1246 Taguchi, M.: Predictability of Major Stratospheric Sudden Warmings: Analysis Results from JMA
- 1247 Operational 1-Month Ensemble Predictions from 2001/02 to 2012/13, Journal of the Atmospheric
- 1248 Sciences, 73(2), 789-806, 2016.
- Takaya, Y., T. Yasuda, Y. Fujii, S. Matsumoto, T. Soga, H. Mori, M. Hirai, I. Ishikawa, H. Sato, A.
- 1250 Shimpo, M. Kamachi, T. Ose: Japan Meteorological Agency/Meteorological Research Institute-
- 1251 Coupled Prediction System version 1 (JMA/MRI-CPS1) for operational seasonal forecasting, Clim.
- 1252 Dyn., 1–2, 313–333, 2017.
- 1253 Thiéblemont, R., Matthes, K., Omrani, NE. et al.: Solar forcing synchronizes decadal North Atlantic
- 1254 climate variability. Nat. Comm. 6, 8268, 2015.
- 1255 Thompson, D.W.J., Susan Solomon: Interpretation of Recent Southern Hemisphere Climate Change.
- 1256 Science, 296, Issue 5569, pp. 895-899, 2002.
- Thompson, D.W.J., M.P. Baldwin, and J. M. Wallace: Stratospheric connection to Northern
- Hemisphere wintertime weather: Implications for prediction. J. Climate, 15, 1421-1428, 2002.

- 1259 Thompson, D. W. J., M. P. Baldwin, and S. Solomon: Stratosphere–troposphere coupling in the
- 1260 Southern Hemisphere. J. Atmos. Sci., 62, 708–715, 2005.
- 1261 Thompson, D., Solomon, S., Kushner, P. et al.: Signatures of the Antarctic ozone hole in Southern
- Hemisphere surface climate change. Nature Geosci 4, 741–749 (2011).
- Tompkins, A. M., Ortiz De Zárate, M. I., Saurral, R. I., Vera, C., Saulo, C., Merryfield, W. J.,
- 1264 Sigmond, M., Lee, W., Baehr, J., Braun, A., Butler, A., Déqué, M., Doblas-Reyes, F. J., Gordon, M.,
- 1265 Scaife, A. A., Imada, Y., Ishii, M., Ose, T., Kirtman, B., Kumar, A., Müller, W. A., Pirani, A.,
- 1266 Stockdale, T., Rixen, M., & Yasuda, T.: The Climate-System Historical Forecast Project: Providing
- Open Access to Seasonal Forecast Ensembles from Centers around the Globe, Bulletin of the
- 1268 *American Meteorological Society*, *98*(11), 2293-2301, 2017.
- 1269 Tripathi, O.P., M. Baldwin, A. Charlton-Perez, M. Charron, S. Eckermann, E. Gerber, G. Harrison, D.
- Jackson, B. Kim, Y. Kuroda, A. Lang, C. Lee, S. Mahmood, R. Mizuta, G. Roff, M. Sigmond, S-W.
- Son: The predictability of the extra-tropical stratosphere on monthly timescales and its impact on the
- skill of tropospheric forecasts, Q. J. R. Meteorol. Soc., 141, 987-1003, 2015a.
- 1273 Tripathi, O., Charlton-Perez, A., Sigmond, M., and Vitart, F: Enhanced long-range forecast skill in
- boreal winter following stratospheric strong vortex conditions. Environ. Res. Lett. 10, 104007, 2015b
- 1275 Tyrrell, N. L., A. Yu. Karpechko and P. Raisanen: The Influence of Eurasian Snow Extent on the
- Northern Extratropical Stratosphere in a QBO Resolving Model, J. Geophys. Res., 123, 1, 315-328,
- **1277** 2018.
- 1278 Tyrrell N. L., Karpechko A. Y., Uotila P., Vihma T.: Atmospheric circulation response to anomalous
- 1279 Siberian forcing in October 2016 and its long-range predictability. Geophys. Res. Lett., 46, 2800–
- 1280 2810, 2019.
- 1281 Tyrrell, N., A. Yu. Karpechko and S. Rast: Siberian snow forcing in a dynamically bias-corrected
- 1282 model, J. Climate, v. 33, 10455–10467, 2020.
- 1283 Vitart, F.: Madden—Julian Oscillation prediction and teleconnections in the S2S database. Q.J.R.
- 1284 Meteorol. Soc, 143: 2210-2220, 2017.
- 1285 Wang, L., Ting, M. & Kushner, P.J.: A robust empirical seasonal prediction of winter NAO and
- 1286 surface climate. Sci Rep 7, 279, 2017.
- Wang L. et al.: What chance of a sudden stratospheric warming in the southern hemisphere? Environ.
- 1288 Res. Lett., 15, 104038, 2020.
- Warner, J. L., Screen, J. A., & Scaife, A. A.: Links between Barents-Kara sea ice and the extratropical
- atmospheric circulation explained by internal variability and tropical forcing. Geophysical Research
- 1291 Letters, 47, e2019GL085679, 2020.
- White, I., C. I. Garfinkel, E. P. Gerber, M. Jucker, P. Hitchcock, and J. Rao: The generic nature of the
- tropospheric response to sudden stratospheric warmings, Journal of Climate, 33, 5589-5610, 2020.
- 1294 Wittman, L.M. Polvani, R.K. Scott and A.J. Charlton: Stratospheric influence on baroclinic lifecycles
- and its connection to the Arctic Oscillation, Geophys. Res. Lett., 31, L16113, 2004.
- 1296 Wittman, A.J. Charlton and L.M. Polvani: The effect of lower stratospheric shear on baroclinic
- instability, J. Atmos. Sci., 64, 479-496, 2007.
- 1298 Woo, SH., Sung, MK., Son, SW. et al.: Connection between weak stratospheric vortex events and the
- 1299 Pacific Decadal Oscillation. *Clim Dyn* **45**, 3481–3492 (2015).
- Yamazaki, K., Nakamura, T., Ukita, J., and Hoshi, K.: A tropospheric pathway of the stratospheric
- quasi-biennial oscillation (QBO) impact on the boreal winter polar vortex, Atmos. Chem. Phys., 20,
- 1302 5111–5127, 2020.

- 1303 Yoo, C., and Son, S.-W.: Modulation of the boreal wintertime Madden-Julian oscillation by the
- stratospheric quasi-biennial oscillation, Geophys. Res. Lett., 43, 1392–1398, 2016.
- Zappa, G. and Shepherd, T.G.: Storylines of atmospheric circulation change for European regional
- 1306 climate impact assessment. J. Clim., 30, 6561–6577, 2017.
- Zappa, G., Pithan, F. and Shepherd, T.G.: Multimodel evidence for an atmospheric circulation
- response to Arctic sea ice loss in the CMIP5 future projections. *Geophys. Res. Lett.*, 45, 1011–1019,
- 1309 2018.
- Zhang, F., Sun, Y. Q., Magnusson, L., Buizza, R., Lin, S., Chen, J., & Emanuel, K.: What Is the
- Predictability Limit of Midlatitude Weather?, J. Atm. Sci., 76(4), 1077-1091, 2019.
- Zhang, P., Wu, Y., Simpson, I. R., Smith, K. L., Zhang, X., De, B., & Callaghan, P.: A stratospheric
- pathway linking a colder Siberia to Barents-Kara Sea sea ice loss. Science Advances, 4(7), eaat6025,
- 1314 2018.