



Interaction between aerosol and thermodynamic stability within the PBL during the wintertime over the North China Plain: Aircraft observation and WRF-Chem simulation

Hao Luo¹, Li Dong², Yichen Chen^{3,4*}, Yuefeng Zhao^{5*}, Delong Zhao^{3,4}, Mengyu Huang^{3,4}, Deping Ding^{3,4}, Jiayuan Liao¹, Tian Ma¹, Maohai Hu^{2*}, Yong Han^{1,6*}

¹Advanced Science & Technology of Atmospheric Physics Group (ASAG), School of Atmospheric Sciences, Sun Yat-sen University & Southern marine Science and Engineering Guangdong Laboratory, Zhuhai 519082, China

²School of Electronic and Optical Engineering, Nanjing University of Science and Technology, Nanjing 210094, China

³Beijing Weather Modification Office, Beijing 100089, China

10 ⁴Beijing Key Laboratory of Cloud, Precipitation and Atmospheric Water Resources (LCPW), Beijing Meteorological Bureau, Beijing 100089, China

⁵School of Physics and Electronics, Shandong Normal University, Jinan 250014, China

⁶Key Laboratory of Tropical Atmosphere-Ocean System (Sun Yat-sen University), Ministry of Education, Zhuhai 519082, China

15 * Correspondence to: Yichen Chen (chenyichen@bj.cma.gov.cn); Yuefeng Zhao (yuefengzhao@sdu.edu.cn); Maohai Hu (hmh@njtu.edu.cn) & Yong Han (hany66@mail.sysu.edu.cn)

Abstract. Aerosol-planetary boundary layer (PBL) interaction has been proposed as a key mechanism for stabilizing the atmosphere and exacerbating surface air pollution. Although the understanding of this process has progressed enormously, its magnitude and impact remain uncertain and vary widely concerning aerosol types, vertical distributions, synoptic conditions, etc. In this study, our primary interest is to distinguish the aerosol-PBL interaction of absorbing and scattering aerosols under contrasting synoptic patterns and aerosol vertical distributions. Detailed in-situ aircraft (KingAir-350) measurements and online coupled model Weather Research and Forecasting with Chemistry (WRF-Chem) simulations are explored over the North China Plain (NCP). Furthermore, a long-term PBL stability trend from 1980 to 2020 over the NCP is also investigated. The aircraft measurements and surface observations show that the surface air pollution over the Baoding City on 3 January is heavier than that on 4 January, 2020. In addition, the aerosols are restricted to the low layer on 3 January, whereas the aerosols mix more homogeneous upwards on 4 January. Thereupon, we focus on the two days with distinct synoptic circumstances, PBL stability, and aerosol vertical distributions over the NCP. According to the WRF-Chem modelling, the synoptic pattern over the Baoding City differs between the two days. The prevailing wind direction is opposite with a southwest wind on 3 January and a northeast wind on 4 January. The results indicate that the synoptic condition may affect the PBL thermal structure, thus affecting the aerosol vertical distribution. Additionally, the sensitive numerical experiments reveal that the light-absorbing and light-scattering aerosols have different effects on altering the PBL thermal structure. The inhibition effect of scattering aerosols on the PBL appears to be independent of the aerosol height distribution and solely depends on its concentration. However, aerosol-PBL feedback of absorbing aerosols is highly



dependent on its vertical distribution. Our analysis highlights that we should principally concentrate on controlling the
emissions of scattering aerosols under the stable stratification while cooperating to control the emissions of scattering and
absorbing aerosols in an unstable stratification. Moreover, the long-term inter-annual variation in PBL stability shows a
strong correlation with the East Asian Winter Monsoon, which seems to be valuable in determining which pollutants to
target in different monsoon years and attaining more precise air pollution control. Based on the numerical simulations and
observational constraints, a concept scheme description has been concluded to deepen our recognition of the interactions
between thermodynamic stability and aerosols within the PBL over the NCP region.

1 Introduction

Ambient air pollution has been one of the major environmental issues in China, particularly in the highly populated and
industrialized areas, such as the North China Plain (NCP) (Chan and Yao, 2008; Sun et al., 2014; Zhang, Q., Zheng, Y. et al.,
2019a; Fan et al., 2020; Luo et al., 2021). Severe and persistent air pollution episodes commonly break out in the presence of
high aerosol emissions and unfavorable synoptic circumstances (e.g. low wind speed, high moisture, and stable stratification),
posing great threats to human health (Che et al., 2019; Zhang, X. et al., 2019b). The interaction of aerosols with the planetary
boundary layer (PBL), which is regarded to be a critical process for stabilizing the atmosphere and worsening surface air
pollution, has been widely explored in the context of aerosol weather and climate effects (Li, Z. et al., 2017a; Su et al., 2020;
Hung et al., 2021).

Aerosols contribute to considerable uncertainties in interpreting and quantifying the Earth's radiative budget and
hydrological cycles through aerosol-radiation interactions (ARI) and aerosol-cloud interactions (ACI) (Rosenfeld et al., 2014;
Luo et al., 2019; Zhao et al., 2019). Despite the great progress in observational and numerical studies of ARI and ACI over
the recent decades, correctly quantifying the aerosol radiative effect (ARE) on the weather and climate systems remains a
challenge. The principal reason for this challenge is the inadequate understanding of strong variations in aerosol types,
loadings, and vertical distributions, as well as the complex mechanisms among large-scale synoptic patterns, local-scale
planetary boundary layer (PBL) structures and AREs (Wang, H. et al., 2015; Li, Z. et al., 2017a; Huang et al., 2018; Su et al.,
2020).

The thermodynamic stability of the PBL dictates the planetary boundary layer height (PBLH), thereby dominating the
vertical dissipation of surface pollutants to some extent (Zhang, Q., Ma, X. et al., 2009; Zhang, W. et al., 2018; Su et al.,
2020). Aerosols, in turn, have vital feedback on the stability of the PBL, depending on their properties, particularly those of
light-absorbing aerosols (e.g., black, brown and organic carbon) (Huang et al., 2018; Menon et al., 2002; Wang, Z. et al.,
2021). The feedback mechanisms between aerosol and PBL are critical in modulating air pollution outbreaks (Ding et al.,
2013; Ding et al., 2016; Huang et al., 2018; Ma et al., 2020). This feedback process can be divided into two aspects, relating
to surface cooling and atmospheric heating through ARE. On the one hand, the suspended aerosols in the atmosphere
attenuate the incident solar radiation, leading to an overall cooling of the surface, diminishing surface sensible heat flux, and



impeding the PBL development, which is known as the umbrella effect (Ma et al., 2020; Shen and Zhao, 2020). On the other hand, light-absorbing aerosols can trap solar energy and heat the atmosphere, which has different effects depending on the aerosol vertical distribution. For one thing, the absorbing aerosols in the low layer of the atmosphere strongly heat the near-surface layer to form a well-mixed PBL, which is recognized as the stove effect (Ma et al., 2020). For another, the absorbing aerosols in the upper layer of the atmosphere strengthen the PBL inversion intensity and weaken its development, which is noticed as the dome effect (Ding et al., 2016).

In addition, the large-scale synoptic patterns disturb the sensitivity of the aerosol-PBL feedback mechanism. PBLH is not only influenced by the local-scale surface properties (e.g., surface cover types, sensible heat fluxes, latent heat fluxes, etc.), but also regulated by large-scale synoptic forcing (Hu et al., 2016; Miao et al., 2020). Synoptic conditions can alter the PBL thermodynamic stability through cold or warm advection, thus affecting the aerosol vertical dispersion (Zhang, J. et al., 2012; Ye et al., 2016; Miao et al., 2019). Furthermore, due to the prevailing south winds, aerosols emitted in the southern regions can be carried to NCP via cross-region transport, and the increased aerosols eventually modulate the PBL thermodynamic stability (Du et al., 2020; Ma et al., 2020). On the inter-annual scale, previous studies reported that the wintertime air quality is closely connected with the East Asian Winter Monsoon (EAWM), and the strengthening (weakening) of EAWM is typically associated with changes in representative circulation patterns, which can improve (worsen) regional air quality (Niu et al., 2010; Zhang, Y. et al., 2016).

According to the aforementioned efforts, most of them primarily concentrated on the aerosol-PBL interactions neglecting the simultaneous influences of the synoptic condition, the aerosol type and vertical distribution. With numerical simulations and observational constraints, the roles of synoptic pattern, PBLH, aerosol type and vertical distribution in aerosol-PBL interactions warrant further investigation. To address this issue, this study attempts to analyze the effects of synoptic forcing and aerosol types on the PBL thermodynamic stability, aerosol vertical distribution, and aerosol-PBL interaction by using in-situ aircraft measurements, surface observations, and on-line coupled model Weather Research and Forecasting with Chemistry (WRF-Chem) simulations. On the 3rd and 4th of January, 2020, two days with distinct synoptic conditions, PBL stability, and aerosol vertical distributions over NCP are analyzed. Five parallel numerical experiments are conducted to investigate the AREs of scattering and absorbing aerosols under different aerosol vertical distributions. Moreover, the long-term inter-annual variation in PBL stability from 1980 to 2020 over the NCP region is examined, and its driving factors are figured out. The conclusions appear to be beneficial in determining which pollutants to target in different weather conditions and achieving more precise controls of air pollution.

The remainder of this paper is organized as follows. Section 2 describes the data and methods used in this study. Section 3 presents and discusses the aerosol-PBL interactions, as well as the factors of synoptic forcing and aerosol types. The conclusions and summary are provided in section 4.

2 Data and Methods

2.1 Flight experiments and observational data

2.1.1 Intensity aircraft observation experiments

To study the vertical distributions of aerosol concentrations and meteorological conditions, the intensity aircraft observation experiments (aircraft model: KingAir-350, which is displayed in Fig. 1e) were carried out to measure the vertical profiles on January 3 and 4, 2020. The terrain height and three-dimensional flight routes are shown in Fig. 1 during the study periods. The flight routes on the two days were almost identical, which is conducive to the comparative analysis. The KingAir-350 took off and climbed to about 3000 m over the Shahe airport (40.1° N, 116.3° E), which is located in Beijing City. Afterward, the aircraft proceeded towards the southwest for about 150 km, arriving in Baoding City and performing vertical measurements ranging from around 650 m to 3000 m.




Figure 1: WRF-Chem domains with the terrain height and flight routes. (a) The outermost domain (d01) and the two embedded domains (d02 and d03); (b) the innermost domain (d03); (c-d) the three-dimensional flight routes on January 3 and 4, 2020, respectively; (e) the KingAir-350 aircraft. The black dots and brown boxes in panel (b) show the locations of the air quality monitoring sites (AQMS) and automatic weather stations (AWS), respectively. The red and blue lines in panel (b) represent the horizontal flight routes on January 3 and 4, 2020, respectively.



115 In this study, the aerosol number concentrations and size distributions were measured by a passive cavity aerosol
 spectrometer probe (PCASP) mounted on the wingtip of the KingAir-350 aircraft. PCASP detected aerosol size distributions
 ranging from 0.1 to 3.0 μm at 1 Hz. Given the substantial noise of the first channel, we considered all the spectral bands
 except the first one, i.e. 0.11 to 3.0 μm , in this study (Twohy et al., 2005). In addition, an integrated meteorological
 measurement system (AIMMS) was employed to measure the aircraft location and ambient meteorological conditions (e.g.
 120 temperature, wind, etc.). The detailed information of aircraft observation is summarized in Table 1.

Table 1: Information of the KingAir-350 aircraft flight times and instruments.

	Date	Take-off time	Landing time
Flight time	January 3, 2020	1600 LST	1830 LST
	January 4, 2020	1530 LST	1820 LST
	Instrument	Objects	Manufacturer
Observation instrument	PASCP-100X	Aerosol number concentration, size distribution, etc.	Droplet Measurement Technologies
	AIMMS-20	Location, temperature, wind speed, etc.	Advantech Research Inc.

2.1.2 Ground-based observations

Surface meteorological observations at three ground-based automatic weather stations (AWS) in Beijing City were
 collected for model meteorological validation, which includes the hourly datasets of surface pressure, 2 m temperature, 10 m
 125 wind speed, etc. In addition, air quality observations, including hourly ground-based $\text{PM}_{2.5}$ concentration measurements at
 six air quality monitoring sites (AQMS) in Baoding City, were used in this study to validate the model performance. The
 observational air pollutant data was collected from the China National Environmental Monitoring Center
 (<http://www.cnemc.cn/>). Ground-based observations on January 3 and 4, 2020 were applied in this study, and the
 locations of AWS and AQMS are shown in Fig. 1b.

130 2.2 Reanalysis data

The data of isobaric temperature taken from the European Centre for Medium-Range Weather Forecasts (ECMWF)
 ERA5 reanalysis dataset was used to summarize the long-term thermal structure of PBL in North China Plain. The
 temperature data of seven isobaric surfaces between 1000 hPa and 850 hPa were considered to be within the PBL. The sea
 level pressure from the ERA5 reanalysis product was used to calculate and describe the intensity of the EAWM and Siberian
 135 High (SH), which are demonstrated to be the aspects that vigorously govern the PBL thermal structure over the NCP in this



study. Here, the monthly average data during the wintertime (December, January, February) from 1980 to 2020 with a grid of $0.25^\circ \times 0.25^\circ$ was adopted.

2.3 WRF-Chem model

In this study, we simulated the aerosol concentrations and meteorological conditions by using the WRF-Chem model version 3.9, which is an online-coupled three-dimensional Eulerian chemical transport model considering complex physical and chemical processes (Grell et al., 2005). The three nesting domains with horizontal resolutions of 27, 9 and 3 km, respectively, are shown in Fig. 1a. There were 29 vertical layers extended from the ground to the top pressure of 50 hPa in the model, with more than 15 layers located below 3 km to fully describe the vertical structure of the PBL. The simulations were carried out from 0000 UTC on January 2 to 1800 UTC on January 4, 2020, and the first 16 hours were set as the model spin-up time.

The 6-hour National Centers for Environmental Prediction (NCEP) global final (FNL) reanalysis fields with a spatial resolution of $1^\circ \times 1^\circ$ were input for the model initial and lateral boundary meteorological conditions. Anthropogenic emissions were adopted from the Multi-resolution Emission Inventory for China (MEIC) developed by Tsinghua University (<http://meicmodel.org>), whose emission sources were classified into five sectors: industrial process, power plants, residential combustion, on-road mobile sources, and agricultural activities (Li, M., Liu, H. et al., 2017b; Zheng et al., 2018). The Model of Emissions of Gases and Aerosols from Nature (MEGAN) was used online in the simulation to calculate the biogenic emissions (Guenther et al., 2006).

The model physical configurations we used in the simulation included the Morrison double-moment microphysics scheme (Morrison et al., 2005), the RRTMG radiation scheme (Iacono et al., 2008), the Yonsei University (YSU) boundary layer scheme (Noh et al., 2003) and the Noah land surface scheme (Ek et al., 2003). Model for Simulating Aerosol Interactions and Chemistry (MOSAIC) was applied as the sectional aerosol scheme, with aerosols specified as 8 size sections (bins) ranging from 39 nm to 10 μm (Zaveri et al., 2008).

To investigate the interactions between aerosol and PBL thermodynamic stability over the NCP, five parallel numerical experiments were conducted as presented in Table 2. In the control experiment (denoted as the EXP_Ctrl), the anthropogenic emissions were performed at the normal level, and the aerosol optical properties were calculated at each time step and then coupled with the short- and long-wave radiative transfer model. In the EXP_WoF, the emissions were the same as the EXP_Ctrl, except that ARE was not included. In the other three experiments, ARE was taken into consideration, but the emissions varied. Specifically, the BC emission was subtracted in the EXP_WFexBC, the BC emission was 20 times higher in the EXP_WF20BC, and the total aerosol emissions were strengthened to 20 times in the EXP_WF20Aer.



Table 2: Descriptions of five parallel WRF-Chem numerical experiments.

Experiment	Description
EXP_Ctrl	Control experiment, with the aerosol radiative feedback.
EXP_WoF	Without any aerosol radiative feedback.
EXP_WFexBC	With the aerosol feedback excluding the contribution of BC.
EXP_WF20BC	With the aerosol feedback including 20 times the BC emission.
EXP_WF20Aer	With the aerosol feedback including 20 times the total aerosol emission.

2.4 Statistical methods for model validation

Correlation coefficient (R), mean bias (MB), normalized mean bias (NMB), mean error (ME) and root mean square error (RMSE) were applied to assess the WRF-Chem model veracity in simulating meteorological parameters and air pollutants against the ground-based observations with the following equations (Itahashi et al., 2015; Granella et al., 2021):

$$R = \frac{\sum_{i=1}^n (S_i - \bar{S})(O_i - \bar{O})}{\sqrt{\sum_{i=1}^n (S_i - \bar{S})^2 \sum_{i=1}^n (O_i - \bar{O})^2}}, \quad (1)$$

$$\text{MB} = \frac{1}{n} \sum_{i=1}^n (S_i - O_i), \quad (2)$$

$$\text{NMB} = \frac{\sum_{i=1}^n (S_i - O_i)}{\sum_{i=1}^n (O_i)}, \quad (3)$$

$$\text{ME} = \frac{1}{n} \sum_{i=1}^n |S_i - O_i|, \quad (4)$$

$$\text{RMSE} = \sqrt{\frac{1}{n} \sum_{i=1}^n (S_i - O_i)^2}, \quad (5)$$

where S_i and O_i are the simulated and observed parameters, respectively, n is the total number of the values used for validation, and \bar{S} and \bar{O} are the averages of the simulation and observation, respectively.

3 Results and Discussion

3.1 Validations of meteorological parameters and aerosol concentrations

Considering that the weather conditions play important roles in pollutant transport, aerosol dispersion, and chemical reactions in the atmosphere, it is crucial to validate the simulated meteorological parameters with the observations. Fig. 2 shows the validation of meteorological parameters between the modeled and observed results, and model evaluations suggest



that the WRF-Chem model is able to simulate the weather characteristics in the NCP region. The simulated temporal series and magnitudes of temperature, pressure and wind speed generally agree well with the ground-based and vertical aircraft observations. The accuracy of simulation during the daytime is superior to that at night, which is instrumental in the intention of discussing the PBL thermodynamic stability and ARE in the daytime.

190 The statistical evaluations of model performance during the daytime from 1000LST to 1800LST on January 3 and 4, 2020 are concluded in Table 3. Underestimation of temperature exists in both the surface level and vertical profile, though the R between the ground-based observation and EXP_Ctrl is 0.87 that indicates the model well describes the temperature variation. The surface pressure is well simulated with an MB of 0.81 and a small NMB of 0.08%. Slight overestimation of wind speed (0.40 m s^{-1}) in the simulation might be due to the unresolved topography in the WRF model, which is also
 195 demonstrated in the previous studies (Jiménez et al., 2013; Li, M., Wang, T. et al., 2021a).

Table 3: Statistical evaluations of the WRF-Chem performance between ground-based observation and EXP_Ctrl during the daytime from 1000LST to 1800LST on January 3 and 4, 2020.

Variable	Observation	EXP_Ctrl	R^a	MB ^a	NMB ^a	ME ^a	RMSE ^a
Temperature ($^{\circ}\text{C}$)	3.25	2.37	0.87	-0.87	-26.90%	0.95	1.11
Pressure (hPa)	1021.95	1022.76	0.42	0.81	0.08%	1.44	1.81
Wind speed (m s^{-1})	1.14	1.54	0.29	0.40	6.97%	0.49	0.64
PM _{2.5} ($\mu\text{g m}^{-3}$)	104.77	99.86	0.79	-4.91	4.69%	17.64	20.31

^a R : correlation coefficient; MB: mean bias; NMB: normalized mean bias; ME: mean error; RMSE: root mean square error.




Figure 2: Validation of meteorological parameters. (a-c) Time series of 2m temperature, surface pressure, and 10m wind speed in Beijing City between the modeling (EXP_Ctrl) and ground-based observation, respectively; (d-e) vertical profiles of temperature and wind speed in Baoding City between the modeling (EXP_Ctrl) and aircraft observation, respectively. The shaded areas indicate the error bars (standard deviation).



205 In addition, the EXP_Ctrl modeled surface $\text{PM}_{2.5}$ mass concentrations in Baoding City also compare well with the ground-based measurements, especially during the daytime (Fig. 2). The statistical validations indicate an R of 0.79, a total MB of $-4.91 \mu\text{g m}^{-3}$, and an NMB of 4.69% during the daytime (Table 3). Both simulation and observation display a high level of air pollution on 3 January, and good air quality on 4 January 4. Therefore, in this study, we consider that the WRF-Chem simulation is in line with the observation and can capture the general distribution and variation in air pollutants.




Figure 3: Validation of $\text{PM}_{2.5}$ concentration between the modeling (EXP_Ctrl) and ground-based observation. The shaded areas indicate the error bars (standard deviation).

3.2 Aerosol vertical characteristics and PBL thermal structures

215 Despite the large-scale weather condition, the local-scale PBL thermal structure is also reported as a critical aspect that substantially affects the surface aerosol loading (Zhang, Q., Ma, X. et al., 2009). The PBL thermodynamic stability dictates the PBLH, thereby dominating the vertical dissipation of surface pollutants to some extent. Therefore, the detailed synoptic condition, PBL thermal structure, and their influences on the aerosol vertical characteristics are discussed in the following subsections.

3.2.1 Vertical thermal structure and synoptic condition

220 The temporal evolution of the EXP_Ctrl simulated temperature profile is presented in Fig. 4, which reflects the variation in the thermal structure and stability during the study period. The results indicate that the thermal structures differ significantly between the two days. It is noticed that the temperature on 3 January is warmer than that on 4 January, suggesting different wintertime synoptic characteristics. Furthermore, the average lapse rate below 1.5 km is lower on 3 January, with lapse rate values of $3.62 \text{ }^{\circ}\text{C km}^{-1}$ and $5.38 \text{ }^{\circ}\text{C km}^{-1}$ during the aircraft observation periods on 3 and 4 January, respectively.




Figure 4: Temporal evolution of the EXP_Ctrl simulated temperature profile over the Baoding City. The pink and blue dashed boxes correspond to the aircraft observation periods on 3 and 4 January, respectively.

The thermal structure of the atmosphere is closely related to and highly sensitive to the synoptic condition over the NCP, especially during the wintertime (Yang et al., 2016; Miao et al., 2019). Figs. 5a and 5b display the surface synoptic patterns during the aircraft observation periods on 3 and 4 January, respectively. In the adjacent regions of the Baoding City, the wind fields are practically opposite in these two periods, and the contrast is also intuitively shown in Fig. 5c with the vertical profile difference in the meridional wind speed. The prevailing surface wind direction on 3 January is southwest, while on 4 January is northeast. The south wind at the low layer warms the atmosphere below 1.5 km on 3 January, whereas the north wind leads to a cooling effect on 4 January, thus there appears a large temperature difference manifested in Fig. 4. Furthermore, the south wind warms the atmosphere, particularly the aloft layer at about 1 km altitude, leading to a stable thermal stratification below 1 km and a low PBLH on 3 January.




Figure 5: The EXP_Ctrl simulated synoptic patterns during the study period. (a-b) the surface wind, temperature, and pressure field during the aircraft observation periods on 3 and 4 January, respectively; (c) the temporal evolution of the meridional wind speed over the Baoding City. The green dots in (a-b) denote the location of the Baoding City. The pink and blue dashed boxes in (c) correspond to the aircraft observation periods on 3 and 4 January, respectively.

3.2.2 Observed and simulated aerosol vertical distributions

As demonstrated in Fig. 6, contrasting aerosol vertical distributions are observed with aircraft during the daytimes of January 3 and 4, 2020. The accumulation mode particles (0.1-1 μm) dominate the aerosol number concentrations during the study period, which is primarily emitted by anthropogenic pollutants, such as BC aerosol loadings (Wu, Y. et al., 2017). On 3 January, larger amounts of aerosol particles are constrained to the low layer of the atmosphere (below 650 m), while the pollutants decrease sharply above 650 m and gradually tend to a stable low value. On 4 January, the aerosol number

concentrations below 650 m are lower than that on 3 January, but higher when the height is above 1000 m. This contrast is attributed to the differences in atmospheric stability (Su et al., 2020). The great difference in lapse rate shown in Fig. 4 leads to the disparity in atmospheric stability and aerosol dispersion ability. On 4 January, the more unstable thermal stratification promotes the dispersion of the near-surface pollutants, and meanwhile, exacerbating the pollution on the elevated layer.




Figure 6: Aircraft observations of aerosol vertical distributions. (a) the aerosol number concentrations on 3 and 4 January, the dashed and solid lines denote the observations over Beijing and Baoding, respectively; (b-c) the aerosol size distributions on 3 and 4 January, respectively.

The temporal evolution of the EXP_Ctrl simulated vertical profiles of total aerosol number concentrations and BC mass concentrations are shown in Fig. 7. Both profiles have similar features, indicating that anthropogenic fine particles like BC, are the dominant pollutants, which is consistent with the particle size observations exhibited in Figs. 6b and 6c. The diurnal variation in the aerosol profile is strongly driven by the evolution of PBLH, which is well examined in the previous studies (Qu et al., 2017; Huang et al., 2018). The pollutants are well constrained to the near-surface by the stable PBL at nighttime, while the vertical distribution of aerosol particles is more homogenous by turbulent mixing in the afternoon. In addition, on 4 January, the vertical mixing of aerosols is more homogenous, transporting aerosols to a higher layer, which implies a more unstable thermal stratification and a higher PBLH.




Figure 7: Temporal evolution of the EXP_Ctrl simulated aerosol vertical distributions over the Baoding City. (a) the aerosol number concentration (particle diameter: 0.15-2.5 μm); (b) the BC concentration. The pink and blue dashed boxes correspond to the aircraft observation periods on 3 and 4 January, respectively.

The contrasting aerosol vertical distributions between the two days are interpreted from two aspects. From the perspective of aerosol regional transport, the air quality is remarkably altered by the synoptic condition (Zhang, Y. et al., 2016; An et al., 2019). The accumulation of pollutants over NCP is contributed by regional transport from the polluted southwest region via the northward movement of the air mass, whereas the cessation of pollution over NCP is caused by prevailing northern air mass and then the pollution hotspot moves to the south (Luo et al., 2021). Therefore, the observed and simulated high surface pollution loading on 3 January is externally affected by the unfavorable synoptic condition, which is consistent with the south wind demonstrated in Fig. 5. From the perspective of aerosol vertical distribution, the surface air pollution loading is efficiently driven by the thermal structure and stability of the PBL (Miao et al., 2020; Li, Q. et al., 2021b). On 3 January, the prevailing surface south wind leads to a stable stratification, restraining aerosol vertical dispersion



and finally exacerbating the surface air pollution. Besides, the north wind condition on 4 January has the opposite effect, which reduces the surface aerosol loading.

280 In conclusion, the stable stratification related by the prevailing surface south wind limits the PBL development and exacerbates the surface air pollution by restraining the aerosols to the low layer. Conversely, the unstable stratification generated by surface north wind promotes the PBL turbulent mixing and lightens the surface aerosol concentration via vertical diffusion.

3.3 Aerosol and PBL thermodynamic stability interactions

285 3.3.1 ARE on PBL thermal structure

PBL structure modulates the aerosol vertical distribution by turbulent mixing, and inversely, the suspended aerosols may also modify the PBL thermal structure by ARE to some extent. Fig. 8 presents the temperature variation affected by ARE, and it is noticeable that light-absorbing aerosols heat the atmosphere while light-scattering aerosols contribute to a cooling effect. On 3 January, aerosols are constrained to the low layer during the daytime, and the overall ARE results in a warming effect below 1 km. The conspicuous heating suggests that the leading role of absorptive BC aerosols on the variation in PBL thermal structure when coexisting with other types of scattering aerosols on 3 January. On 4 January, due to the strong turbulence mixing, aerosols are carried to the aloft layer. The overall ARE results in a cooling effect below 0.6 km and a warming effect between 0.6 and 1.5 km. When the aerosols are transported to a high altitude, the absorbing and scattering particles inhibit the incident solar radiation from reaching the low layer and exhibit a cooling effect, while the absorbing aerosols heat the upper layer. The contrasting aerosol vertical distribution caused by the varied PBL thermal structure between the two daytimes results in different AREs, therefore, it is quite crucial to examine the aerosol vertical structure when evaluating the ARE.

290

295




Figure 8: The temporal evolution of temperature profile variation influenced by aerosol radiative effect (ARE). (a) ARE by all the aerosols (EXP_Ctrl – EXP_WoF); (b) ARE by BC (EXP_Ctrl – EXP_WFexBC); (c) ARE by other aerosols (EXP_WFexBC – EXP_WoF). The pink and blue dashed boxes correspond to the aircraft observation periods on 3 and 4 January, respectively.



The simulated ARE modify the thermal structure with a temperature variation of less than 0.5 °C shown in Fig. 8, to amplify the signal of ARE when discussing its interactions with PBL, we further performed two experiments, one with strong overall aerosol loading (EXP_WF20Aer) and the other only with strong BC aerosol loading (EXP_WF20BC). The simulated thermal structures are presented in Fig. 9. The amplified simulations reveal a temperature variation of less than 2.5 °C, which is consistent with the altered magnitude of aerosol emissions. ARE by only absorptive BC aerosols heat the low layer more remarkable when compared with the role of overall aerosols, suggesting that the coexists of absorbing and scattering aerosols partially conceal the signal of ARE. This result is owing to the opposite optical roles of the two aerosol types in the low layer, as well as the strong inhibition of the incident solar radiation by the aerosols in the upper layer.




Figure 9: The temporal evolution of temperature profile with (a) high aerosol emissions (EXP_WF20Aer) and (c) high BC emissions (EXP_WF20Aer); the temporal evolution of temperature variation profile influenced by (b) high scattering aerosol loadings (EXP_WF20Aer – EXP_WF20BC) and (d) high absorbing aerosol loadings (EXP_WF20BC – EXP_Ctrl). The pink and blue dashed boxes correspond to the aircraft observation periods on 3 and 4 January, respectively.

3.3.2 ARE on the PBLH, lapse rate, and surface pollution

Since ARE can modulate the PBL thermal structure, the related impacts caused by the thermal variation are discussed here. Fig. 10 shows the time series of the simulated lapse rate, PBLH, and surface carbon monoxide (CO) concentration under five parallel experiments.

Generally, on 3 January, the PBLH is lower and the surface air pollution is heavier than those on 4 January. Influenced by the southwest winds, the warmer air mass can be guided to NCP (Fig. 5a), enhancing the thermal stability and restraining



the growth of the PBL (Fig. 10b). On the one hand, from the local-scale view, the lower PBLH constrains the air pollution to the near-surface; on the other hand, from the regional transport view, the southwest prevailing winds can transport the pollutants emitted from upstream plain regions to NCP, which further worsen the surface air quality (Fig. 10c). However, the relatively higher PBLH and the northeast winds on 4 January are able to reduce the surface air pollution and inversely carry the pollutants to the higher altitude with the strong turbulent mixing. The disparate synoptic conditions regulate the vertical aerosol structures and further affect the ARE and other related variables.

The distinct aerosol types contribute to varied AREs when different aerosol vertical distributions are dominant. The lapse rate and PBLH are closely associated with the vertical thermal structure. A warmer low layer and a colder upper layer induce a higher lapse rate and PBLH, while a colder low layer and a warmer upper layer decrease the lapse rate and PBLH. On 3 January, the pollutants are restricted to the low layer, the absorbing and scattering aerosols play the opposite roles. The absorbing aerosols heat the low layer and invigorate the turbulent mixing (stove effect), leading to a higher lapse rate and PBLH. On the contrary, the scattering aerosols cool the low layer and inhibit the growth of the PBL (umbrella effect), causing a lower lapse rate and PBLH (Figs. 10a and 10b). On 4 January, the aerosol vertical profile differs from that on 3 January, with a huge number of pollutants being carried to the high altitude by the unstable stratification. The absorbing and scattering aerosols both prevent the solar radiation to the low layer (umbrella effect), while the absorbing aerosols also heat the upper layer (dome effect), generating a lower lapse rate and PBLH (Figs. 10a and 10b).

The variations in lapse rate and PBLH further alter the surface air pollution by the thermodynamic and dynamic effect, and high PBLH ordinarily lightens surface air pollution. Due to the CO is a relatively stable pollutant that is hardly interactions with other pollutants and the main sources of CO are primary emissions from anthropogenic pollution and open biomass burning (Chi et al., 2013), we use CO concentration to evaluate the variation in surface air pollution caused by the ARE in this study. The surface CO concentration in the EXP_WF20Aer is divided by a factor of 20 to match other experiments since the emissions are magnified 20 times. The results presented in Fig. 10c indicate that the surface air pollution variation is closely related to the PBLH variation and driven by the aerosol vertical dispersion. When the stable stratification confines the absorbing aerosols to the low layer, the stove effect destroys the stability, enhancing the thermal convection and decreasing the surface air pollution. The absorbing aerosols in the upper layer or the scattering aerosols in both the low and upper layers stabilize the atmosphere, deteriorating the surface air quality.




Figure 10: The temporal evolution of the (a) lapse rate within 1.5 km, (b) planetary boundary layer height (PBLH), and (c) surface CO concentration simulated by the five parallel experiments. The upward or downward arrows indicate an increase or decrease compared to the EXP_Ctrl.



3.3.3 ARE under different PBL thermodynamic stabilities

It is noticeable that different aerosol vertical distributions between the two days contribute to distinct AREs due to the PBL thermal structure and stability differences. In particular, the absorptive BC aerosols have both stove and dome effects, which are sensitive to the PBL thermal structure. Here, we further analyze the spatial distribution of atmospheric stability during the daytime of the aircraft observation periods on 3 January. The atmospheric stability is divided into two categories following the method developed by Wolyn and McKee (1989): a stable stratification known as the deep stable layer (DSL) and an unstable stratification known as the non-deep stable layer (non-DSL). The DSL is identified supposing that 65% of the lowest 1.5 km has a lapse rate of $2.5\text{ }^{\circ}\text{C km}^{-1}$ or less, and the DSL definition examines a layer much deeper than a typical nocturnal inversion (Wolyn and McKee, 1989). The spatial distributions of the DSL and non-DSL are displayed in Fig. 11.





Figure 11: The spatial distributions of the deep stable layer (DSL) and non-deep stable layer (non-DSL) during aircraft observation periods on 3 January over the NCP region.

The impacts of PBL thermodynamic stability on the BC ARE are compared between DSL and non-DSL conditions demonstrated in Fig. 12. Under the non-DSL condition, the BC dome effect is dominant, the BC aerosols in the upper layer absorb the solar radiation, stabilize the PBL and decrease the PBLH (Fig. 12a), and further increase the surface air pollution (Fig. 12c). Under the DSL condition, the BC stove effect plays the dominant role, the strong BC concentrations in the low layer contribute to the development of the PBL and increase the PBLH (Fig. 12b), and further decrease the surface air pollution (Fig. 12d).



370 **Figure 12: Scatter plots of the correlations between EXP_Ctrl and EXP_WF20BC for (a-b) PBLH and (c-d) CO concentration under non-DSL condition and DSL condition, respectively. The PBLHs lower than 500 m are discarded to avoid the incorrect simulation.**

3.4 Long-term variation in PBL thermodynamic stability over NCP

The linkage between synoptic pattern and PBL thermal structure is analyzed from a climatological view, which is critical for the examination of aerosol vertical distribution and ARE. The potential impact of inter-annual climate variability on the PBL thermodynamic stability is explored in this study over the NCP during the wintertime of 1980-2013. Following the method developed by Wu and Wang (2002), the index of SH and the index of EAWM are used to analyze the possible factors that influence the PBL thermal structure over the NCP. The index of SH and EAWM are the two variables describing the meridional wind intensity over the coast of East Asia, which are crucial to the PBL thermal structure. The PBL lapse rate is calculated between 1000 hPa and 850 hPa.

Fig. 13 suggests that the relationships between SH index, EAWM index, and PBL lapse rate are highly correlated. Weak EAWM and SH with the prevailing south wind stabilize the PBL and constrain the aerosols to the near-surface, while



the strong EAWM and SH play the opposite role. The result indicates that the inter-annual variability of the EAWM and SH can influence surface air quality, aerosol vertical distribution, and ARE through its reflection on the intensity of PBL thermodynamic stability over the NCP region.




Figure 13: (a) Time series of the standardized anomaly of the wintertime boundary layer lapse rate (LR), the index of Siberian High (SH), and the index of East Asian Winter Mooson (EAWM) from 1980 to 2020. The scatter plots of the correlations between (b) ESWM and LR, as well as between (c) SH and LR.




390 4 Conclusions and Summary

In this study, the complex relationships between large-scale synoptic patterns, local PBL thermal structures, aerosol vertical distributions, and AREs of different aerosol types are systematically investigated by the combined aircraft observations, surface measurements, reanalysis data, and WRF-Chem simulations. The in-situ aircraft observations were carried out over the Beijing and Baoding Cities during the daytimes on January 3 and 4, 2020, and the numerical simulations were performed over NCP from 3 to 4 January. By and large, the meteorological variables and air pollutants are well simulated when validated by the observations.

Observations show that the surface air pollution over the Baoding City on 3 January is heavier than that on 4 January, and the aerosols are constrained to the low layer on 3 January while the aerosols mix more homogeneous vertically on 4 January. The numerical simulations suggest that the synoptic pattern over the Baoding City differs between the two days, and the prevailing wind direction is opposite with a southwest wind on 3 January and a northeast wind on 4 January. The synoptic condition may affect the PBL thermal structure, thus affecting the vertical dispersion of aerosols. The south winds flow the warm and polluted air to the NCP and stabilize the PBL, hence exacerbating the surface air pollution. The north winds flow the cold and clean air to the NCP, establishing an unstable stratification and transporting the pollutants to the upper layer.

The suspended aerosols may also modify the PBL thermodynamic stability through ARE to some extent, and the synoptic condition can modulate the sensitivity of PBL-aerosol feedback by influencing the PBL thermal structure and aerosol vertical distribution. The interactions among the synoptic pattern, PBL thermal structure, and aerosol vertical distribution are demonstrated in Fig. 14.



410 **Figure 14: The diagram of the interactions among the synoptic pattern, PBL thermal structure and aerosol vertical distribution.**



The sensitive numerical experiments reveal that the absorbing and scattering aerosols have different effects when modifying the PBL thermal structure, and the AREs are also influenced by the aerosol vertical distribution. Aerosol-PBL feedback of absorbing aerosols is highly dependent on its vertical distribution. In the case of stable weather patterns under the influence of the south wind, the absorbing aerosols concentrated in the low layer have a strong radiative heating effect on the atmosphere (stove effect). This stove effect disturbs the stratification stability of the low layer, promotes the development of the PBL in coordination with the surface sensible heat flux, and transports heat and pollutants to the upper layer through turbulence. In the case of unstable weather patterns under the influence of the north wind, the absorbing aerosols in the upper atmosphere heat the atmosphere (dome effect) and inhibit the development of the PBL, which further leads to the accumulation of surface pollutants. The inhibition effect of scattering aerosols on the PBL is independent of the aerosol height distribution but only depends on its concentration. Scattering aerosols reflect solar radiation back to the space (umbrella effect), thus weakening the surface sensible heat flux and upward heat transfer, inhibiting the development of the PBL and further accumulating surface pollutants. In summary, the umbrella, stove and dome effects of the scattering and absorbing aerosols under different synoptic patterns are illustrated in Fig. 15. The conclusions enlighten that we should mainly control the emissions of scattering aerosols under the stable stratification, and cooperate to control the emissions of scattering and absorbing aerosols in an unstable stratification, especially to prevent the accumulation of surface pollutants caused by the dome effect of absorbing aerosols.




Figure 15: The illustration of the umbrella, stove, and dome effects of the scattering and absorbing aerosols under different synoptic patterns. The red plus and blue minus signs indicate the aerosol heating and dimming effect, respectively. The upward and downward arrows denote the increase and decrease in PBLH caused by ARE, respectively.

In addition, the factors affecting the wintertime PBL stability over the NCP area are further analyzed from the climatological perspective, which is mainly controlled by the intensity of EAWM and SH. A strong winter monsoon creates an unstable PBL stratification, whereas a weak winter monsoon stabilizes the PBL. Combined with the umbrella, stove, and dome effects of scattering and absorbing aerosols under different synoptic conditions, this finding aids us in determining which pollutants to target in different monsoon years and achieving more precise air pollution control.



It should be noted that only two days of pollution episodes are examined in this study. While the model simulations produce evidence that is compatible with the observations and substantiate the observational analysis, additional measurements should be carried out in the near future.

Code/Data availability

440 The surface observational air pollutant data are collected from <http://www.cnemc.cn>. The meteorological data taken from the ERA5 reanalysis dataset are obtained from <https://www.ecmwf.int/en/forecasts/datasets/reanalysis-datasets/era5>. The NCEP-FNL global reanalysis dataset used in the WRF-Chem model is available from <https://rda.ucar.edu/datasets/ds083.2>. The MEIC anthropogenic emission inventories are available from www.meicmodel.org, and for more information, please contact Q. Zhang (qiangzhang@tsinghua.edu.cn). The WRF-Chem model version 3.9 is available from
445 https://www2.mmm.ucar.edu/wrf/users/download/get_source.html. The aircraft data and surface meteorological data are available from the corresponding author. The codes that support the findings of this study are available from the corresponding author upon reasonable request.

Author Contributions

Y. H. designed research and edited the paper; H. L., L. D., Y. C., Y. Z., M. H., and Y. H. performed research and analyzed
450 data; D. Z., M. H., D. D., J. L., and T. M. participated in the discussions. H. L. wrote the manuscript.

Competing Interests

The authors declare they have no conflict of interest.

Acknowledgements

This work was jointly supported by the National Natural Science Foundation of China (Grant Nos. 41775026, 42027804,
455 41075012 and 40805006).

References

An, Z., Huang, R.-J., Zhang, R., Tie, X., Li, G., Cao, J., Zhou, W., Shi, Z., Han, Y., Gu, Z., and Ji, Y.: Severe haze in northern China: A synergy of anthropogenic emissions and atmospheric processes, *Proceedings of the National Academy of Sciences*, 116, 8657-8666, 10.1073/pnas.1900125116, 2019.



- 460 Chan, C. K., and Yao, X.: Air pollution in mega cities in China, *Atmospheric Environment*, 42, 1-42,
 10.1016/j.atmosenv.2007.09.003, 2008.
- Che, H., Gui, K., Xia, X., Wang, Y., Holben, B. N., Goloub, P., Cuevas-Agulló, E., Wang, H., Zheng, Y., Zhao, H., and
 Zhang, X.: Large contribution of meteorological factors to inter-decadal changes in regional aerosol optical depth,
Atmos. Chem. Phys., 19, 10497-10523, 10.5194/acp-19-10497-2019, 2019.
- 465 Chi, X., Winderlich, J., Mayer, J. C., Panov, A. V., Heimann, M., Birmili, W., Heintzenberg, J., Cheng, Y., and Andreae, M.
 O.: Long-term measurements of aerosol and carbon monoxide at the ZOTTO tall tower to characterize polluted and
 pristine air in the Siberian taiga, *Atmos. Chem. Phys.*, 13, 12271-12298, 10.5194/acp-13-12271-2013, 2013.
- Ding, A. J., Fu, C. B., Yang, X. Q., Sun, J. N., Petäjä, T., Kerminen, V. M., Wang, T., Xie, Y., Herrmann, E., Zheng, L. F.,
 Nie, W., Liu, Q., Wei, X. L., and Kulmala, M.: Intense atmospheric pollution modifies weather: a case of mixed
 470 biomass burning with fossil fuel combustion pollution in eastern China, *Atmos. Chem. Phys.*, 13, 10545-10554,
 10.5194/acp-13-10545-2013, 2013.
- Ding, A. J., Huang, X., Nie, W., Sun, J. N., Kerminen, V.-M., Petäjä, T., Su, H., Cheng, Y. F., Yang, X.-Q., Wang, M. H.,
 Chi, X. G., Wang, J. P., Virkkula, A., Guo, W. D., Yuan, J., Wang, S. Y., Zhang, R. J., Wu, Y. F., Song, Y., Zhu, T.,
 Zilitinkevich, S., Kulmala, M., and Fu, C. B.: Enhanced haze pollution by black carbon in megacities in China,
 475 *Geophysical Research Letters*, 43, 2873-2879, 10.1002/2016GL067745, 2016.
- Du, H., Li, J., Wang, Z., Dao, X., Guo, S., Wang, L., Ma, S., Wu, J., Yang, W., Chen, X., and Sun, Y.: Effects of Regional
 Transport on Haze in the North China Plain: Transport of Precursors or Secondary Inorganic Aerosols, *Geophysical
 Research Letters*, 47, e2020GL087461, 10.1029/2020GL087461, 2020.
- Ek, M. B., Mitchell, K. E., Lin, Y., Rogers, E., Grunmann, P., Koren, V., Gayno, G., and Tarpley, J. D.: Implementation of
 480 Noah land surface model advances in the National Centers for Environmental Prediction operational mesoscale Eta
 model, *Journal of Geophysical Research: Atmospheres*, 108, 10.1029/2002JD003296, 2003.
- Fan, H., Zhao, C., and Yang, Y.: A comprehensive analysis of the spatio-temporal variation of urban air pollution in China
 during 2014–2018, *Atmospheric Environment*, 220, 117066, 10.1016/j.atmosenv.2019.117066, 2020.
- Granella, F., Aleluia Reis, L., Bosetti, V., and Tavoni, M.: COVID-19 lockdown only partially alleviates health impacts of
 485 air pollution in Northern Italy, *Environmental Research Letters*, 16, 035012, 10.1088/1748-9326/abd3d2, 2021.
- Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., and Eder, B.: Fully coupled “online”
 chemistry within the WRF model, *Atmospheric Environment*, 39, 6957-6975, 10.1016/j.atmosenv.2005.04.027, 2005.
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene
 emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181-
 490 3210, 10.5194/acp-6-3181-2006, 2006.
- Hu, X.-M., Li, X., Xue, M., Wu, D., and Fuentes, J. D.: The Formation of Barrier Winds East of the Loess Plateau and Their
 Effects on Dispersion Conditions in the North China Plains, *Boundary-Layer Meteorology*, 161, 145-163,
 10.1007/s10546-016-0159-4, 2016.



- Huang, X., Wang, Z., and Ding, A.: Impact of Aerosol-PBL Interaction on Haze Pollution: Multiyear Observational
 495 Evidences in North China, *Geophysical Research Letters*, 45, 8596-8603, 10.1029/2018GL079239, 2018.
- Hung, W.-T., Lu, C.-H., Alessandrini, S., Kumar, R., and Lin, C.-A.: The impacts of transported wildfire smoke aerosols on
 surface air quality in New York State: A multi-year study using machine learning, *Atmospheric Environment*, 259,
 118513, 10.1016/j.atmosenv.2021.118513, 2021.
- Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., and Collins, W. D.: Radiative forcing by long-
 500 lived greenhouse gases: Calculations with the AER radiative transfer models, *Journal of Geophysical Research:
 Atmospheres*, 113, 10.1029/2008JD009944, 2008.
- Itahashi, S., Hayami, H., and Uno, I.: Comprehensive study of emission source contributions for tropospheric ozone
 formation over East Asia, *Journal of Geophysical Research: Atmospheres*, 120, 331-358, 10.1002/2014JD022117, 2015.
- Jiménez, P. A., Dudhia, J., González-Rouco, J. F., Montávez, J. P., García-Bustamante, E., Navarro, J., Vilà-Guerau de
 505 Arellano, J., and Muñoz-Roldán, A.: An evaluation of WRF's ability to reproduce the surface wind over complex terrain
 based on typical circulation patterns, *Journal of Geophysical Research: Atmospheres*, 118, 7651-7669,
 10.1002/jgrd.50585, 2013.
- Li, M., Liu, H., Geng, G., Hong, C., Liu, F., Song, Y., Tong, D., Zheng, B., Cui, H., Man, H., Zhang, Q., and He, K.:
 Anthropogenic emission inventories in China: a review, *National Science Review*, 4, 834-866, 10.1093/nsr/nwx150,
 510 2017b.
- Li, M., Wang, T., Xie, M., Li, S., Zhuang, B., Fu, Q., Zhao, M., Wu, H., Liu, J., Saikawa, E., and Liao, K.: Drivers for the
 poor air quality conditions in North China Plain during the COVID-19 outbreak, *Atmospheric Environment*, 246,
 118103, 10.1016/j.atmosenv.2020.118103, 2021a.
- Li, Q., Zhang, H., Cai, X., Song, Y., and Zhu, T.: The impacts of the atmospheric boundary layer on regional haze in North
 515 China, *npj Climate and Atmospheric Science*, 4, 9, 10.1038/s41612-021-00165-y, 2021b.
- Li, Z., Guo, J., Ding, A., Liao, H., Liu, J., Sun, Y., Wang, T., Xue, H., Zhang, H., and Zhu, B.: Aerosol and boundary-layer
 interactions and impact on air quality, *National Science Review*, 4, 810-833, 10.1093/nsr/nwx117, 2017a.
- Luo, H., Han, Y., Lu, C., Yang, J., and Wu, Y.: Characteristics of Surface Solar Radiation under Different Air Pollution
 Conditions over Nanjing, China: Observation and Simulation, *Advances in Atmospheric Sciences*, 36, 1047-1059,
 520 10.1007/s00376-019-9010-4, 2019.
- Luo, H., Han, Y., Cheng, X., Lu, C., and Wu, Y.: Spatiotemporal Variations in Particulate Matter and Air Quality over China:
 National, Regional and Urban Scales, *Atmosphere*, 12, 43, 10.3390/atmos12010043, 2021.
- Ma, Y., Ye, J., Xin, J., Zhang, W., Vilà-Guerau de Arellano, J., Wang, S., Zhao, D., Dai, L., Ma, Y., Wu, X., Xia, X., Tang,
 G., Wang, Y., Shen, P., Lei, Y., and Martin, S. T.: The Stove, Dome, and Umbrella Effects of Atmospheric Aerosol on
 525 the Development of the Planetary Boundary Layer in Hazy Regions, *Geophysical Research Letters*, 47,
 e2020GL087373, 10.1029/2020GL087373, 2020.



- Menon, S., Hansen, J., Nazarenko, L., and Luo, Y.: Climate Effects of Black Carbon Aerosols in China and India, *Science*, 297, 2250-2253, 10.1126/science.1075159, 2002.
- Miao, Y., Liu, S., and Huang, S.: Synoptic pattern and planetary boundary layer structure associated with aerosol pollution during winter in Beijing, China, *Science of The Total Environment*, 682, 464-474, 10.1016/j.scitotenv.2019.05.199, 2019.
- Miao, Y., Che, H., Zhang, X., and Liu, S.: Integrated impacts of synoptic forcing and aerosol radiative effect on boundary layer and pollution in the Beijing–Tianjin–Hebei region, China, *Atmos. Chem. Phys.*, 20, 5899-5909, 10.5194/acp-20-5899-2020, 2020.
- Morrison, H., Curry, J. A., and Khvorostyanov, V. I.: A New Double-Moment Microphysics Parameterization for Application in Cloud and Climate Models. Part I: Description, *Journal of Atmospheric Sciences*, 62, 1665, 10.1175/jas3446.1, 2005.
- Niu, F., Li, Z., Li, C., Lee, K.-H., and Wang, M.: Increase of wintertime fog in China: Potential impacts of weakening of the Eastern Asian monsoon circulation and increasing aerosol loading, *Journal of Geophysical Research: Atmospheres*, 115, 10.1029/2009JD013484, 2010.
- Noh, Y., Cheon, W. G., Hong, S. Y., and Raasch, S.: Improvement of the K-profile Model for the Planetary Boundary Layer based on Large Eddy Simulation Data, *Boundary-Layer Meteorology*, 107, 401, 10.1023/a:1022146015946, 2003.
- Qu, Y., Han, Y., Wu, Y., Gao, P., and Wang, T.: Study of PBLH and Its Correlation with Particulate Matter from One-Year Observation over Nanjing, Southeast China, *Remote Sensing*, 9, 668, 10.3390/rs9070668, 2017.
- Rosenfeld, D., Sherwood, S., Wood, R., and Donner, L.: Climate Effects of Aerosol-Cloud Interactions, *Science*, 343, 379-380, 10.1126/science.1247490, 2014.
- Shen, L., and Zhao, C.: Dominance of Shortwave Radiative Heating in the Sea-Land Breeze Amplitude and its Impacts on Atmospheric Visibility in Tokyo, Japan, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD031541, 10.1029/2019JD031541, 2020.
- Su, T., Li, Z., Li, C., Li, J., Han, W., Shen, C., Tan, W., Wei, J., and Guo, J.: The significant impact of aerosol vertical structure on lower atmosphere stability and its critical role in aerosol–planetary boundary layer (PBL) interactions, *Atmos. Chem. Phys.*, 20, 3713-3724, 10.5194/acp-20-3713-2020, 2020.
- Sun, Y., Jiang, Q., Wang, Z., Fu, P., Li, J., Yang, T., and Yin, Y.: Investigation of the sources and evolution processes of severe haze pollution in Beijing in January 2013, *Journal of Geophysical Research: Atmospheres*, 119, 4380-4398, 10.1002/2014JD021641, 2014.
- Twohy, C. H., Petters, M. D., Snider, J. R., Stevens, B., Tahnk, W., Wetzel, M., Russell, L., and Burnet, F.: Evaluation of the aerosol indirect effect in marine stratocumulus clouds: Droplet number, size, liquid water path, and radiative impact, *Journal of Geophysical Research: Atmospheres*, 110, 10.1029/2004JD005116, 2005.
- Wang, H., Shi, G. Y., Zhang, X. Y., Gong, S. L., Tan, S. C., Chen, B., Che, H. Z., and Li, T.: Mesoscale modelling study of the interactions between aerosols and PBL meteorology during a haze episode in China Jing–Jin–Ji and its near



- surrounding region – Part 2: Aerosols' radiative feedback effects, *Atmos. Chem. Phys.*, 15, 3277-3287, 10.5194/acp-15-3277-2015, 2015.
- Wang, Z., Huang, X., Ding, K., Ren, C., Cao, L., Zhou, D., Gao, J., and Ding, A.: Weakened Aerosol-PBL Interaction During COVID-19 Lockdown in Northern China, *Geophysical Research Letters*, 48, e2020GL090542, 10.1029/2020GL090542, 2021.
- Wolyn, P. G., and McKee, T. B.: Deep Stable Layers in the Intermountain Western United States, *Monthly Weather Review*, 117, 461-472, 10.1175/1520-0493(1989)117<0461:dsliti>2.0.co;2, 1989.
- Wu, B., and Wang, J.: Winter Arctic Oscillation, Siberian High and East Asian Winter Monsoon, *Geophysical Research Letters*, 29, 3-1-3-4, 10.1029/2002GL015373, 2002.
- Wu, Y., Wang, X., Tao, J., Huang, R., Tian, P., Cao, J., Zhang, L., Ho, K. F., Han, Z., and Zhang, R.: Size distribution and source of black carbon aerosol in urban Beijing during winter haze episodes, *Atmos. Chem. Phys.*, 17, 7965-7975, 10.5194/acp-17-7965-2017, 2017.
- Yang, X., Zhao, C., Guo, J., and Wang, Y.: Intensification of aerosol pollution associated with its feedback with surface solar radiation and winds in Beijing, *Journal of Geophysical Research: Atmospheres*, 121, 4093-4099, 10.1002/2015JD024645, 2016.
- Ye, X., Song, Y., Cai, X., and Zhang, H.: Study on the synoptic flow patterns and boundary layer process of the severe haze events over the North China Plain in January 2013, *Atmospheric Environment*, 124, 129-145, 10.1016/j.atmosenv.2015.06.011, 2016.
- Zaveri, R. A., Easter, R. C., Fast, J. D., and Peters, L. K.: Model for Simulating Aerosol Interactions and Chemistry (MOSAIC), *Journal of Geophysical Research: Atmospheres*, 113, 10.1029/2007JD008782, 2008.
- Zhang, J. P., Zhu, T., Zhang, Q. H., Li, C. C., Shu, H. L., Ying, Y., Dai, Z. P., Wang, X., Liu, X. Y., Liang, A. M., Shen, H. X., and Yi, B. Q.: The impact of circulation patterns on regional transport pathways and air quality over Beijing and its surroundings, *Atmos. Chem. Phys.*, 12, 5031-5053, 10.5194/acp-12-5031-2012, 2012.
- Zhang, Q., Ma, X., Tie, X., Huang, M., and Zhao, C.: Vertical distributions of aerosols under different weather conditions: Analysis of in-situ aircraft measurements in Beijing, China, *Atmospheric Environment*, 43, 5526-5535, 10.1016/j.atmosenv.2009.05.037, 2009.
- Zhang, Q., Zheng, Y., Tong, D., Shao, M., Wang, S., Zhang, Y., Xu, X., Wang, J., He, H., Liu, W., Ding, Y., Lei, Y., Li, J., Wang, Z., Zhang, X., Wang, Y., Cheng, J., Liu, Y., Shi, Q., Yan, L., Geng, G., Hong, C., Li, M., Liu, F., Zheng, B., Cao, J., Ding, A., Gao, J., Fu, Q., Huo, J., Liu, B., Liu, Z., Yang, F., He, K., and Hao, J.: Drivers of improved PM_{2.5} air quality in China from 2013 to 2017, *Proceedings of the National Academy of Sciences*, 116, 24463-24469, 10.1073/pnas.1907956116, 2019a.
- Zhang, W., Guo, J., Miao, Y., Liu, H., Song, Y., Fang, Z., He, J., Lou, M., Yan, Y., Li, Y., and Zhai, P.: On the Summertime Planetary Boundary Layer with Different Thermodynamic Stability in China: A Radiosonde Perspective, *Journal of Climate*, 31, 1451-1465, 10.1175/jcli-d-17-0231.1, 2018.



- 595 Zhang, X., Xu, X., Ding, Y., Liu, Y., Zhang, H., Wang, Y., and Zhong, J.: The impact of meteorological changes from 2013 to 2017 on PM_{2.5} mass reduction in key regions in China, *Science China Earth Sciences*, 62, 1885-1902, 10.1007/s11430-019-9343-3, 2019b.
- Zhang, Y., Ding, A., Mao, H., Nie, W., Zhou, D., Liu, L., Huang, X., and Fu, C.: Impact of synoptic weather patterns and inter-decadal climate variability on air quality in the North China Plain during 1980–2013, *Atmospheric Environment*, 124, 119-128, 10.1016/j.atmosenv.2015.05.063, 2016.
- 600 Zhao, C., Yang, Y., Fan, H., Huang, J., Fu, Y., Zhang, X., Kang, S., Cong, Z., Letu, H., and Menenti, M.: Aerosol characteristics and impacts on weather and climate over the Tibetan Plateau, *National Science Review*, 7, 492-495, 10.1093/nsr/nwz184, 2019.
- Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., Qi, J., Yan, L., Zhang, Y., Zhao, H., Zheng, Y., He, K., and Zhang, Q.: Trends in China's anthropogenic emissions since 2010 as the consequence of clean air actions, *Atmos. Chem. Phys.*, 18, 14095-14111, 10.5194/acp-18-14095-2018, 2018.
- 605