

Cloud droplet formation at the base of tropical convective clouds: closure between modeling and measurement results of ACRIDICON-CHUVA

Ramon Campos Braga¹, Barbara Ervens², Daniel Rosenfeld³, Meinrat O. Andreae^{4,5}, Jan-David Förster¹, Daniel Fütterer⁶, Lianet Hernández Pardo¹, Bruna A. Holanda¹, Tina Jurkat - Witschas⁶, Ovid O. Krüger¹, Oliver Lauer¹, Luiz A. T. Machado^{1,7}, Christopher Pöhlker¹, Daniel Sauer⁶, Christiane Voigt^{6,8}, Adrian Walser⁶, Manfred Wendisch⁹, Ulrich Pöschl¹, and Mira L. Pöhlker¹

¹Multiphase Chemistry Department, Max Planck Institute for Chemistry, 55128 Mainz, Germany

²Université Clermont Auvergne, CNRS, SIGMA Clermont, Institut de Chimie de Clermont-Ferrand, 63000 Clermont-Ferrand, France

³Institute of Earth Sciences, The Hebrew University of Jerusalem, 9190401 Jerusalem, Israel.

⁴Biogeochemistry Department, Max Planck Institute for Chemistry, 55128 Mainz, Germany

⁵Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA 92037, USA

⁶Institute of Atmospheric Physics, German Aerospace Center (DLR), 82234 Oberpfaffenhofen, Germany

⁷National Institute for Space Research (INPE), 12227-010 São José Dos Campos, Brazil

⁸Johannes Gutenberg University Mainz, 55099 Mainz, Germany

⁹Faculty of Physics and Earth Sciences, Leipzig Institute for Meteorology, University of Leipzig, 04103 Leipzig, Germany

Correspondence: Mira L. Pöhlker (m.pohlker@mpic.de) and Barbara Ervens (barbara.ervens@uca.fr)

Abstract.

Aerosol-cloud interactions contribute to the large uncertainties in current estimates of climate forcing. We investigated the effect of aerosol particles on cloud droplet formation by model calculations and aircraft measurements over the Amazon and over the western tropical Atlantic during the ACRIDICON-CHUVA campaign in September 2014. On the HALO research aircraft, cloud droplet number concentrations (N_d) were measured near the base of clean and polluted growing convective cumuli using a cloud combination probe (CCP) and a cloud and aerosol spectrometer (CAS-DPOL). An adiabatic parcel model was used to perform cloud droplet number closure studies for flights in differently polluted air masses. Model input parameters included aerosol size distributions measured with an ultra-high sensitive aerosol spectrometer (UHSAS), in combination with a condensation particle counter (CPC). Updraft velocities (w) were measured with a boom-mounted Rosemount probe. Over the continent, the aerosol size distributions were dominated by accumulation mode particles, and good agreement between measured and modeled N_d values was obtained (deviations $\lesssim 10\%$) assuming an average hygroscopicity of $\kappa \sim 0.1$, which is consistent with Amazonian biomass burning and secondary organic aerosol. Above the ocean, fair agreement was obtained assuming an average hygroscopicity of $\kappa \sim 0.2$ (deviations $\lesssim 16\%$) and further improvement was achieved assuming different hygroscopicities for Aitken and accumulation mode particles ($\kappa_{\text{Ait}} = 0.8$, $\kappa_{\text{acc}} = 0.2$; deviations $\lesssim 10\%$), which may reflect secondary marine sulfate particles. Our results indicate that Aitken mode particles and their hygroscopicity can be important for droplet formation at low pollution levels and high updraft velocities in tropical convective clouds.

1 Introduction

Aerosol-cloud-interactions represent one of the largest uncertainties in our current understanding of the Earth's climate system, according to the latest report by the Intergovernmental Panel on Climate Change (IPCC, 2021). Aerosols have a strong effect on cloud properties since cloud droplets form on cloud condensation nuclei (CCN) by condensation of water vapor. The uptake of water vapor by aerosol particles and the subsequent activation and growth of cloud droplets is described by the Köhler theory (Köhler, 1936), relates the water vapor saturation ratio (s) to the water activity in the aqueous solution (a_w), Raoult term), which is the size and composition dependencies of the droplet's solute effect, as well as the increase in equilibrium water vapor pressure due to droplet's surface curvature Kelvin-term (K_c (Kelvin term)). The Raoult effect (solute term) is commonly parameterized using the hygroscopicity parameter κ with values ranging from ~ 0.1 to ~ 0.9 for single components of atmospheric aerosol particles (Petters and Kreidenweis, 2007). The κ values of ambient aerosol particles are in the range of $\sim 0.09 < \kappa < 0.18$ for the complex mixtures of organic aerosols and $0.1 < \kappa < 0.3$ for biomass burning aerosol (e.g., Andreae and Rosenfeld, 2008; Carrico et al., 2010; Engelhart et al., 2012).

~~The Amazon Basin is a unique region to test our understanding of aerosol-cloud interactions in shallow and deep convective clouds due to large variability in aerosol concentration during the dry and wet seasons (e.g., Artaxo, 2002; Andreae et al., 2004; Pöhlker et al., 2018). The properties and dynamics of clouds over this pristine rain forest region can be fundamentally changed by anthropogenic emissions (e.g., Roberts et al., 2003; Rosenfeld et al., 2008; Reutter et al., 2009; Pöhlker et al., 2018). To explore aerosol-cloud interactions, cloud microstructure and precipitation-forming processes above the Amazon rain forest, the ACRIDICON-CHUVA (Aerosol, Cloud, Precipitation, and Radiation Interactions and Dynamics of Convective Cloud Systems—Cloud processes of the main precipitation systems in Brazil: A contribution to cloud resolving modeling and to the Global Precipitation Measurements) campaign with the HALO (High Altitude Long Range Research) aircraft took place during the dry season in September 2014 (Wendisch et al., 2016).~~

Many CCN closure studies in various parts of the world have been performed to improve our understanding of the relationship between aerosol properties and their ability to form cloud droplets (e.g., Rissler et al., 2004; Broekhuizen et al., 2006; Wang et al., 2008; Ervens et al., 2010). Such closure studies compare the predicted CCN number concentration (N_{CCN}) according to Köhler theory based on particle size and composition (hygroscopicity) to results from CCN measurements, i.e. for equilibrium conditions at different supersaturations in CCN counters. Much fewer studies compare predicted ($N_{d,p}$) and measured cloud droplet number concentrations ($N_{d,m}$) ('cloud droplet number closure'), since often direct measurements or estimates of the updraft velocities (w) near cloud base are not available. $N_{d,p}$ at cloud bases is commonly calculated in adiabatic cloud models based on the hygroscopic growth of CCN particles with a prescribed κ and w . These models simulate the expansion and cooling of air, the resulting changes in relative humidity, and the condensational growth of cloud droplets (Reutter et al., 2009; Ervens et al., 2010) (Reutter et al., 2009; Ervens et al., 2010; Leaitch et al., 2010). Updraft velocity is often used as a fitting parameter to match $N_{d,m}$ (e.g., Anttila et al., 2012). Other studies used w distributions to yield-predict a range of $N_{d,p}$ (Chuang et al., 2000; Peng et al., 2005; Meskhidze et al., 2005; Hsieh et al., 2009). Previous cloud droplet

50 number closure studies suggested that w is one of the most poorly constrained parameters leading to large uncertainties in the predictions of $N_{d,p}$ (e.g., Conant et al., 2004; Fountoukis et al., 2007).

~~Braga et al. (2017a) compared N_d at cloud bases measured during the~~ The Amazon Basin is a unique region to test our understanding of aerosol-cloud interactions in shallow and deep convective clouds due to large variability in aerosol concentration during the dry and wet seasons (e.g., Artaxo, 2002; Andreae et al., 2004; Pöhlker et al., 2016, 2018). The properties and dynamics of clouds over this pristine rain forest region can be fundamentally changed by anthropogenic emissions (e.g., Roberts et al., 2003; Rosenfeld
55 . To explore aerosol-cloud interactions, cloud microstructure and precipitation-forming processes above the Amazon rain forest, the ACRIDICON-CHUVA ~~campaign to CCN number concentrations to those predicted based on traditional parameterizations for~~ (Aerosol, Cloud, Precipitation, and Radiation Interactions and Dynamics of Convective Cloud Systems - Cloud processes of tHe main precipitation systems in Brazil: A contribUtion to cloud resolVing modeling and to the GlobAl Precipitation
60 Measurements) campaign with the HALO (High Altitude Long Range Research) aircraft took place during the dry season in September 2014 (Wendisch et al., 2016).

The focus of this study is to describe a closure analysis based on the $N_{d,m}$ as a function of N_d and at cloud bases of convective clouds and $wN_{d,p}$ (Twomey, 1959). The current study represents an extension of this work, in which we explicitly ~~simulate~~ calculated from an adiabatic parcel model (Feingold and Heymsfield, 1992; Ervens et al., 2005) that is based on other
65 independently measured properties. To this end, measurements of droplet concentrations at cloud bases performed during the ACRIDICON-CHUVA campaign were used. Furthermore, our calculations explicitly simulated the condensational growth of aerosol particles from below cloud base up to a height of several meters above the level, at which they grow to cloud droplet sizes. We ~~compare~~ compared $N_{d,m}$ at cloud bases of convective clouds in different air masses with $N_{d,p}$ predicted by an adiabatic ~~parcel model,~~ using in situ measurements of aerosol size distributions as model input. Model studies are performed to-

70 The closure analysis was performed separately for two cloud probes (Cloud droplet number concentrations and size distributions were measured by a Cloud Combination Probe – Cloud Droplet Probe, CCP-CDP, and Cloud and Aerosol Spectrometer with Depolarization, CAS-DPOL) mounted onboard HALO (Wendisch et al., 2016; Voigt et al., 2017). This was performed to verify our methodology using two types of instruments to measure number concentrations of droplets with different particle inlet characteristics and uncertainties. Our study was performed to explore the sensitivities of effective particle hygroscopic-
75 ity (κ), aerosol particle number concentration (N_a), and w to N_d . The results reveal the sensitivity of N_d to κ , N_a and w for measurements over the Amazon basin during the dry season and over Atlantic Ocean.

2 Aircraft Measurements

The measurements were performed aboard the High Altitude Long Range aircraft (HALO), a modified business jet G550 (manufactured by Gulfstream, Savannah, USA). In situ meteorological and avionics data, such as the vertical velocity, were
80 obtained at 1 Hz from the BAasic HALO Measurement And Sensor System (BAHAMAS). A boom-mounted Rosemount model 858 AJ air velocity probe was used to measure the updraft velocity with BAHAMAS, measuring in a range of $0.1 \text{ m s}^{-1} \leq w \leq 6 \text{ m s}^{-1}$. The uncertainties in measured w are $\Delta w < 0.2 \text{ m s}^{-1}$ for $w < 5 \text{ m s}^{-1}$ and $\Delta w \approx 0.25 \text{ m s}^{-1}$ for $w > 5 \text{ m s}^{-1}$.

Further details on the uncertainties of w measurements are described by Mallaun et al. (2015). The measurements took place over the Amazon Basin and over the western tropical Atlantic in September 2014 during the ACRIDICON–CHUVA campaign
85 (Wendisch et al., 2016).

Figure 1a shows the measurement region for the flights analyzed in this study (the flights are labelled with 'AC' and a running number, in agreement with the naming e.g., (Wendisch et al., 2016)). The region of cloud base measurements is indicated by circles for each flight. The measurement strategy was developed such that measurements were made within at most 10 minutes and 60 km from each other. This was performed to assure that droplet measurements at cloud base pertain to the same air mass
90 as the aerosol measurements below cloud base. A *conceptual* representation of the cloud profiling, including flight legs below and within cloud base, is shown for measurements during flight AC19 in Fig. 1b. For the present study, the flight legs below and at cloud base are of primary relevance. Such flight legs, during which the relevant aerosol and cloud microphysical data were obtained, are distinguished by different colors. During flight AC19 the profiling of the marine shallow cumulus clouds was conducted up to an altitude of 4.3 km; details on the air mass origin and the aerosol properties during this flight can be
95 found in Section 2.1. Aerosol properties were investigated during the flight leg below cloud base, which had a length of about 19 km at an altitude of ~ 450 m above sea level (asl). Cloud microphysical properties of the marine shallow cumulus clouds (Atlantic ocean) were investigated during the flight leg near cloud base, which had a length of ~ 60 km at an altitude at ~ 600 m asl. A similar strategy was applied for in-land flights. Convective cumuli formed in very polluted environments (arc of deforestation) directly above the Amazonian deforestation arc during flight AC07. Less polluted clouds were found farther
100 away from the deforestation fires over the tropical rain forest (remote Amazon) during flights AC09 and AC18.

2.1 Aerosol size distribution below cloud base

Aerosol size distributions were measured using an Ultra-High Sensitivity Aerosol Spectrometer (UHSAS; Droplet Measurement Technologies, Inc., Longmont, CO, USA) (Cai et al., 2008; Moore et al., 2021). The UHSAS combines a high-power infrared laser ($\lambda = 1054$ nm) and a large solid angle range in a side-ways direction for the detection of light scattered by
105 individual particles (Andreae et al., 2018). The aircraft instrument measures particles in the diameter size range between ~~100~~
92 nm and 600 nm. The instrument is mounted in an under-wing canister. The sampled air is entering the instrument by a forward facing diffusor inlet, and the airflow is reduced by a second inlet to approximately isokinetic conditions. The measured
particle ~~diameter~~diameters can be assumed to be close to their dry diameters due to heating effects (Chubb et al., 2016). The UHSAS was calibrated with monodisperse polystyrene latex (PSL) spheres of known size. Typical uncertainties of UHSAS
110 measurements are both 15 % in diameter and concentration (Cai et al., 2008; Moore et al., 2021).

The total particle number concentration in the size range of ~ 10 nm to ~ 500 nm (N_{CN}) below cloud base were measured using the Aerosol Measurement System (AMETYST)~~);~~; the uncertainty of these measurements is estimated to be 10 % (Andreae et al., 2018). N_{CN} was measured by a butanol-based condensation particle counter (CPCs, modified Grimm CPC 5.410 by Grimm Aerosol Technik, Ainring, Germany) with a flow of 0.6 L min^{-1} . Particle losses in the sampling lines have been
115 estimated and taken into account with the particle loss calculator by von der Weiden et al. (2009). Typical uncertainties of CPC measurements are on the order of ~ 10 % (~~Petzold et al., 2011~~)(Petzold et al., 2011; Andreae et al., 2018).




Figure 1. a) HALO flight tracks during the ACRIDICON–CHUVA experiment, color-coded for the different flights. Circles indicate the region of aerosol and cloud measurements. The average aerosol particle concentration measured below cloud bases during flights AC07, AC09, AC18, and AC19 were 2417 cm^{-3} , 737 cm^{-3} , 809 cm^{-3} and 428 cm^{-3} , respectively. b) Cloud profiling maneuvers during flight AC19 above the Atlantic Ocean near the Amazon River delta shown as three-dimensional profiles corresponding to the two dimensional profile in panel a). Relevant flight segments - particularly legs below cloud base and within cloud base, as well as cloud penetrations above cloud base - are emphasized by color-coding. c) Aerosol size distributions for each flight as used in this study.

The geometric mean of the aerosol size distribution and N_{CN} below cloud were calculated. The mean aerosol size distribution was fitted by one-modal-one-modal lognormal distributions. The integral of the fit for the aerosol size distribution should be similar to N_{CN} if mainly accumulation mode particles are present. This was fulfilled for AC07, AC09 and AC18, but not for AC19 (Tables S1-S4). For this latter flight, the integrated number concentration of the monomodal lognormal fit made up approximately half of the total N_{CN} . This discrepancy led to the assumption that a significant number concentration of particles in the size range of Aitken mode particles were present during AC19, but not captured by the UHSAS measurements. Consequently, a bimodal ASD shape was inferred. The geometric parameters for the lognormal distribution assumed for measurements during Flight-AC19 were based on averages of bimodal aerosol size distributions measured above the ocean in previous studies (Figure S4) (Wex et al., 2016; Quinn et al., 2017; Gong et al., 2019). Other shapes of marine aerosol size distributions, e.g. as reported by Leaitch et al. (2010), were not considered for our lognormal fit because they were not

[in agreement with the measured UHSAS data](#). The resulting shape of the two modes based on literature data was weighted by the difference between UHSAS and CPC measurements (Table S4). The number concentrations of all fitted aerosol size distributions were normalized to the measured N_{CN} . The variability of the aerosol number size distributions was calculated by the standard deviation on average $\sim 10\%$ and up to $\sim 20\%$ for very clean conditions. As a conservative approach $\sim 20\%$ was used in our model sensitivity study to take into account the impact of this variability on cloud droplet number concentration (Section 4.2). All concentrations are reported for normalized atmospheric conditions (~~Corrected~~[corrected](#) for standard conditions (STP): $T = 273.15^\circ\text{C}$ and $p = 1013.25\text{ mbar}$).

2.2 Cloud droplet measurements at cloud base

Cloud droplet number concentrations and size distributions were measured by a Cloud Combination Probe – Cloud Droplet Probe (CCP-CDP) and by a Cloud and Aerosol Spectrometer with Depolarization (CAS-DPOL) mounted onboard HALO (Wendisch et al., 2016; Voigt et al., 2017). Cloud droplet number size distributions (DSDs) between $3\ \mu\text{m}$ and $50\ \mu\text{m}$ in diameter were measured at a temporal resolution of 1 s by the CAS-DPOL and CCP-CDP probes (Baumgardner et al., 2011; Voigt et al., 2010, 2011; Kleine et al., 2018; Wendisch and Brenguier, 2013). These probes have different measurement characteristics such as particle inlet, sampling area of detection, size sensitivities etc. The CCP-CDP is an open-path instrument that detects forward-scattered laser light from cloud particles as they pass through the CDP detection area (Lance et al., 2010). CAS-DPOL collects forward-scattered light to determine particle size and number that pass the sampling area centered in an inlet shaft that guides the airflow. CCP-CDP and CAS-DPOL has similar values of uncertainty ($\sim 10\%$) in the sample area. However, particle velocities in the sampling tube may be modified by the CAS tube when compared to the open path instruments (like CCP-CDP). This results in an additional uncertainty in the droplet number concentration measured by CAS-DPOL. During the ACRIDICON-CHUVA campaign the resulting uncertainty in the droplet concentration measured by CCP-CDP and CAS-DPOL were $\sim 10\%$ and $\sim 21\%$, respectively (Braga et al., 2017a).

For cloud base measurements, each probe DSD spectrum represented 1 s of flight path (covering between 70 m to 120 m of horizontal distance for the aircraft speed at cloud bases). We refer in the current study to the measurements closest to cloud base as 'cloud base' measurements, even if the actual cloud base might have been slightly below this altitude of measurements (Section 3.2.2 and Figure 2). [The cloud base measurements were selected based on the videos recorded by the HALO cockpit forward-looking camera](#). From the DSDs, the droplet number concentrations were derived by size integration. Braga et al. (2017a) showed that both probes were in agreement within their uncertainty range for probe DSDs ($\pm \sim 16\%$). The overall systematic errors in the cloud probe integrated water content with respect to a King type hot-wire device are $\sim 6\%$ for CAS-DPOL and $\sim 21\%$ for CCP-CDP. A positive bias of $\sim 20\%$ was found for CAS-DPOL droplet concentration in comparison with those measured with CCP-CDP for cloud passes with cloud droplet effective radius $< 7\ \mu\text{m}$ (mostly measured at cloud bases). Cloud passes were defined for conditions, under which the number droplet concentration (i.e., particles with diameter larger than $3\ \mu\text{m}$) exceeded 20 cm^{-3} . This criterion was applied to avoid cloud passes well mixed with subsaturated environment air ($\text{RH} < 100\%$) and counts of haze particles, typically found at cloud edges. Additional details about the cloud probes measurements at cloud bases used in this study can be found in Tables S5-S6.

3 Methodology

3.1 Probability matching method (PMM): Pairing measured updraft velocities (w) and droplet number concentrations ($N_{d,m}$)

The thermal instability in the boundary layer promotes the formation of clouds consisting of regions with updrafts and down-
165 drafts. At cloud bases, the variability in vertical velocities and droplet concentration is high due to air turbulence. Clouds
develop in updrafts, and during their vertical development the continued movement as a turbulent eddy adds a large random
component to the relationship of w with $N_{d,m}$. These intrinsic characteristics of clouds reduce the confidence that a measured
 w in the cloud led to the simultaneously measured $N_{d,m}$. Such inconsistencies often result in poor correlations of w and $N_{d,m}$.
As w is highly variable in clouds and is measured independently from $N_{d,m}$, we apply the “Probability Matching Method”
170 (PMM, Haddad and Rosenfeld (1997)) to statistically determine the most probable combinations of $N_{d,m}$ and w values using
the same percentiles of their occurrence. The PMM analysis is based on the assumption that these two related variables increase
monotonically with each other. Measured $N_{d,m}$ and w values were sorted in ascending order and the most likely w value was
assigned to $N_{d,m}$ for each of the four flights. To avoid biases caused by outlier measurements, $N_{d,m}$ above the 97.5th and below
the 2.5th percentile were removed. Only cloud passes with positive vertical velocities (i.e., updrafts) were considered in the
175 analysis. Furthermore, we take into account only data of non-precipitating clouds, typically from cumulus humilis and cumulus
mediocris clouds. Braga et al. (2017a) have shown that the PMM can be used to find the best agreement between measured and
estimated N_d at cloud base as a function of w . In the current study, PMM analysis is used to compare the $N_{d,m}$ at cloud base and
its assigned w with $N_{d,p}$ at a constant w in the model. [Figure 2 shows an example of measured \$w\$ at cloud bases and estimated
 \$w\$ based on the PMM analysis \(\$w_{PMM}\$ \). The figure shows that \$w_{PMM}\$ are in well agreement with measurements at cloud bases.
180 \[Furthermore, for cloud passes in which the values of \\$w\\$ are negative realistic \\$w\\$ are estimated based on PMM.\]\(#\)](#)

3.2 Adiabatic cloud parcel model

3.2.1 Model description and simulations

The adiabatic parcel model describes the growth of aerosol particles by water vapor uptake on a moving mass grid (Feingold
and Heymsfield, 1992; Ervens et al., 2005). The air parcel is described to rise with a constant w below and inside of the
185 cloud. Saturation with respect to water vapor in the air parcel is calculated based on the standard thermodynamic equations
for adiabatic conditions as a function of w and particle properties (N_a , particle sizes and hygroscopicity) (Pruppacher and
Klett, 1997). It is assumed that the aerosol particles are internally mixed with identical hygroscopicity (κ) of all particles.
This assumption was made based on previous sensitivity studies that have shown that for marine and aged continental air
masses internal mixtures are suitable approximations (Ervens et al., 2010). [We note that \$\kappa\$ is regarded here as an effective
190 \[parameter, encompassing all factors that affect water uptake.\]\(#\) Simulations are performed up to a height of 70 m above the
level of predicted maximum supersaturation. The initial conditions for the model simulations are summarized in Tables S1-
S4. Particles that exceed a diameter of 3 \$\mu\text{m}\$ are defined as droplets; this definition allows a direct comparison of \$N_{d,p}\$ and](#)




Figure 2. Time series of droplet size distribution measured by CCP-CDP [top], number concentration of droplets (N_d) [middle], measured vertical velocities w , and estimated w based on PMM method [bottom]. The measurements were performed during flight AC19 above the Atlantic Ocean.

$N_{d,m}$. Collision/coalescence processes are not considered as we restrict our analysis to heights near cloud base where droplets are relatively small and the cloud droplet size distribution is narrow. Under such conditions, collision-coalescence is likely negligible (Shaw et al., 1998; Xue et al., 2008; Rosenfeld, 2018; Braga et al., 2017b). Sensitivity studies are performed for the observed ranges of w and $N_d \pm 40\%$ and assumed range of $0.02 \leq \kappa \leq 1$ to identify parameter ranges and combinations for which droplet closure can be achieved.

3.2.2 Determination of in-cloud height to compare $N_{d,m}$ and $N_{d,p}$

The cloud base measurements were performed at approximately constant altitude during each research flight and were selected based on the videos recorded by the HALO cockpit forward-looking camera. However, this height these measurements might represent different levels in relation to cloud base and to the level of maximum supersaturation, depending on at cloud bases, which depends on the updraft velocity and turbulence in cloud. In order to determine the height at which $N_{d,m}$ and $N_{d,p}$ should

be compared, the measured liquid water content (LWC) was compared to the simulated LWC using the aerosol size distribution for the different flights together with w measured at cloud base and assumed hygroscopicity of $\kappa = 0.1$.

205 Under adiabatic conditions, $N_{d,p}$ is predicted to be approximately constant at ~ 20 m above the level of the maximum supersaturation S_{max} (Fig. S5). Figure 3 shows the ~~frequency of measured LWC and the modeled LWC at different heights. At ~ 20 m above cloud base the LWC measured with the highest frequency and the modeled LWC is the same. For this reason the model results at 20 m above cloud base are compared to the measured cloud droplet number concentrations in the scope of this study.~~

210 ~~Predicted LWC g m^{-3} as a function of height above the level of S_{max} left axis and w (lines, color-coded by $w \text{ m s}^{-1}$) for flights a) AC07, b) AC09, c) AC18, d) AC19. The vertical bars indicate the number of cloud passes (with a temporal resolution of 1 s) as a function of the measured LWC by CAS-DPOL and CCP-CDP in cloud (right axis). The dashed line denotes the level of 20 m above predicted maximum supersaturation, at which $N_{d,p}$ is predicted (Section 3.2.2).~~

215 ~~Figure 3 shows the values of predicted liquid water content (LWC) from the same LWC based on the simulations as a function of height above S_{max} for the four flights. Overlaid on the model results (colored lines) are the frequencies of measured LWC by the cloud probes near cloud base (white bars). The measured LWC represents the cumulative mass size distribution. For AC09, AC18 and AC19, all flights, the model predictions match the most frequently measured LWC in most of cases match the minimum LWC measured at ~ 20 m above the S_{max} level. This height level might represent slightly different absolute heights above the surface and the level of saturation estimated by the model (RH = 100%) (Fig. S6). For AC07, the LWC frequency distribution is very flat and leads to ambiguity of the height in cloud where predicted and measured LWC coincide.~~ However, since we focus our discussion in the following section on the comparison of $N_{d,m}$ and $N_{d,p}$, we perform our analysis for based on model predictions at a height of 20 m above S_{max} ~~for all flights, as above this height cloud droplet number is not predicted to change.~~

220

4 Results and Discussion

225 4.1 Constraining aerosol hygroscopicity (κ) based on N_d and w

Figure 4 shows the range of $N_{d,m}$ as a function of w as determined by the PMM (Section 3.1); the symbols indicate $N_{d,m}$ from CAS-DPOL (black diamonds) and CCP-CDP (black triangles). The lines in Fig. 4a-d represent model predictions for the assumption of $\kappa = 0.05$, $\kappa = 0.1$, $\kappa = 0.3$ and $\kappa = 0.6$ for up to 38 w values for each flight, covering the measured w range. Figure 4e shows results of additional simulations for Flight AC19 (marine conditions) assuming $\kappa = 0.6$ and $\kappa = 0.8$ for aerosol particles from Aitken ($d < 70$ nm) and $\kappa = 0.1$ and $\kappa = 0.2$ for aerosol particles from accumulation ($d \geq 70$ nm) mode sizes, respectively. For all flights, $N_{d,m}$ values are reasonably reproduced by the model assuming a particle hygroscopicity of $0.05 < \kappa < 0.3$; $N_{d,p}$ are closer to the measured values from CCP-CDP assuming a slightly lower κ , whereas $N_{d,m}$ from CAS-DPOL indicate a slightly higher κ . However, these deviations are within the uncertainty range of the cloud probe measurements, i.e., $\sim 10\%$ and $\sim 21\%$ for CCP-CDP and CAS-DPOL, respectively (Braga et al., 2017a).

230




Figure 3. Predicted LWC [$g\ m^{-3}$] as a function of height above the level of S_{max} [left axis] and w (lines, color-coded by w [$m\ s^{-1}$]) for flights a) AC07, b) AC09, c) AC18, d) AC19. The vertical bars indicate the number of cloud passes (with a temporal resolution of 1 s) as a function of the measured LWC by CAS-DPOL and CCP-CDP in cloud (right axis). The dashed line denotes the level of 20 m above predicted maximum supersaturation, at which $N_{d,p}$ is predicted (Section 3.2.2).

235 Figure 4 shows that the agreement between measured and predicted cloud droplet number concentration is obtained for low w during all flights. However, the value of w , above which the model predictions deviate from measurements varies among the flights: For continental clouds as encountered during AC07, AC18 and AC09, the model results agree well with observations for $w \lesssim 2.5 \text{ m s}^{-1}$. At higher w , $N_{d,m}$ shows a much stronger increase with w than predicted by the model. For AC19, i.e., above the ocean, this trend is even obvious for $w \gtrsim 0.5 \text{ m s}^{-1}$. The statistical analysis based on bias, root mean square error (RMSE),
240 and mean absolute error (MAE) from the closure analysis are shown in Tables S7-S18. This analysis suggests that the use of two probes to perform the closure does not have a large effect on the inferred value of κ . We find best agreement, quantified by the smallest absolute bias and RMSE, for all cases for single κ values of $0.05 \leq \kappa \leq 0.2$. The deviations between $N_{d,m}$ from CCP-CDP and CAS-DPOL ($\sim 21\%$ on average) reinforce the advantage of duplicate measurements for the closure analysis. The use of a single cloud probe might lead to a biased κ estimate based on the data set of each cloud probe separately. ~~In our cases, analysis of the CCP-CDP data only might result in conclusions on the 'best' κ at the upper end of the κ whereas the CAS-DPOL data rather suggest κ values at the lower end.~~
245 The consideration of both cloud probes shows the uncertainty in N_d measurements and therefore the uncertainty range of κ and/or N_d values for N_d closure. Therefore, we base our conclusions in the following on the statistical analysis of all data from both probes together (Tables S9, S12, S15 and S18).

The results in Fig. 4 imply that the assumption of an internally mixed aerosol population with moderate hygroscopicity ($\kappa \sim 0.1$) is justified to reproduce $N_{d,m}$ for flights AC07, AC09 and AC18 for wide ranges of updraft speeds ($0.1 \text{ m s}^{-1} \leq w \leq 2.5 \text{ m s}^{-1}$). This κ value has been suggested previously for comparable air masses during the dry season in the Amazon Basin (e.g., Pöhlker et al., 2016, 2018). In these prior studies, κ was constrained based on size-resolved CCN measurements and measurements of the aerosol chemical composition, dominated by an aged organic fraction. Our results ~~confirm that this κ value are in agreement with previous studies.~~
250 The value range is representative of internally mixed aerosol particle populations during the dry season in the Amazon Basin, which are influenced by fresh and aged biomass burning aerosol from Amazon and Africa.

~~The systematic bias in $N_{d,p}$ suggests that the adiabatic model might not be suitable for high updraft conditions. Under such conditions, entrainment of additional aerosol particles from air masses surrounding the cloud could explain the observed trend in $N_{d,m}$ which would not be captured by the model. However, while entrainment of biomass burning aerosol may be possible, we do not have any quantitative information on such processes.~~
260

While also particles of different hygroscopicities and activation thresholds depending on w might explain the trends in Fig. 4a-c, there is no indication of higher hygroscopicity of smaller accumulation mode aerosol particles during the Amazonian dry season (e.g., Pöhlker et al., 2016, 2018). In air masses of different origin, aerosol particles would likely not only exhibit different chemical composition and hygroscopicity but also large variability in their particle number concentrations. Given the relatively
265 small standard deviations in the measured N_a (Tables S1 - S4), we are confident that the sampled aerosol populations did not have large variability in their composition. ~~While the hygroscopicity of particles could possibly change near cloud base, e.g., due to dissolution of soluble compounds, this effect would result in the opposite trend as predicted, i.e., a higher $N_{d,p}$ at low w when particles dissolve over longer time scales. The resulting curves of $N_{d,p}$ as a function of w would be even flatter than shown in Fig. 4, as opposed to the steep increase in $N_{d,m}$. Thus, a significant role of such composition effects can likely be excluded.~~




Figure 4. Cloud droplet number concentration (N_d) as a function of updraft velocity near cloud base of convective clouds during flights: a) AC07, b) AC09, c) AC18, d) and e) AC19. The measured updraft velocities are based on the “probability matching method” (PMM) using the same percentiles for updraft velocity and $N_{d,m}$ (Section 3.1). The black diamonds and triangles represent $N_{d,m}$ near cloud base from the CAS-DPOL and CCP-CDP probes, respectively. Measurement uncertainties, indicated by error bars, are $\sim 21\%$ and $\sim 10\%$ for CAS-DPOL and CCP-CDP data (Braga et al. (2017a)). The colored lines in panels a) - d) show $N_{d,p}$ assuming a single κ value for both modes (labeled on the left). Panel e) shows $N_{d,p}$ based on simulations assuming different values of κ for Aitken and accumulation mode particles during flight AC19.

270 ~~The~~The chemical composition of Aitken mode particles often differs significantly from that of accumulation mode particles, which are more aged and internally mixed (e.g., [Wex et al. \(2016\)](#); [Pöhlker et al. \(2018\)](#) ([Wex et al., 2016](#); [Pöhlker et al., 2018](#))), and thus continental Aitken mode particles usually exhibit a lower hygroscopicity than accumulation mode particles (McFiggans et al., 2006).

The air masses below cloud encountered during flight AC19 were mostly impacted by marine air, as supported by prior
275 back trajectory analysis (Section S1 and Holanda et al. (2020)) and exhibited a bi-modal aerosol size distribution with low $N_{d,m}$ (Figure 1c). For this flight, the cloud droplet closure is worse as compared to the reasonable agreement for the other three cases. Not only is the absolute difference between $N_{d,m}$ and $N_{d,p}$ relatively larger (Figure 4d), but also the trend of $N_{d,m}$ with w cannot be well reproduced: While at $w < 0.5 \text{ m s}^{-1}$, the range of $N_{d,p}$ agrees well with $N_{d,m}$, above this threshold the model strongly underestimates the droplet number concentration even for $\kappa = 0.3$ (Figure 4d). Assuming $\kappa = 0.6$ only slightly
280 increases $N_{d,p}$ as compared to the results for $\kappa = 0.3$. This trend shows that $N_{d,p}$ is rather insensitive to κ if particles are very hygroscopic ($\kappa \gtrsim \kappa > 0.3$). While all κ values lead to reasonable agreement at low w , none of the model results can reproduce the strongly increased $N_{d,m}$ with w . Therefore, we conclude that the simplifying assumptions made in the model, i.e., identical hygroscopicities across both aerosol modes, may not be appropriate.

The measured aerosol size distribution during flights AC19 differed significantly from the other ones (Figure 1c) because of
285 (i) low N_a , and (ii) a distinct Aitken mode (mean diameter 37 nm) that comprised $\sim 47\%$ of the particle number concentration. At such low N_a , the maximum supersaturation in the clouds is relatively high so that at sufficiently high w , Aitken mode particles (diameter $\lesssim 70 \text{ nm}$) may be activated into cloud droplets and contribute to N_d ([Pöhlker et al., 2021](#) (?)). Highly hygroscopic Aitken mode particles over the ocean may reflect secondary marine sulfate aerosols (Andreae and Raemdonck, 1983).

To account for different hygroscopicities in Aitken and accumulation modes, we performed further sensitivity analyses using combinations of $\kappa = 0.1$ and 0.6 for the two modes (Figure 4e). It is obvious that the choice of κ for the Aitken mode (κ_{Ait}) does not affect $N_{d,p}$ for $w \lesssim 1 \text{ m s}^{-1}$ in the presence of very hygroscopic accumulation mode particles ($\kappa_{\text{acc}} = 0.6$) or below
290 $w \lesssim 0.5 \text{ m s}^{-1}$ with $\kappa_{\text{acc}} = 0.1$, respectively. Even assuming rather extreme values of $\kappa_{\text{Ait}} = 0.8$ cannot fully reproduce the large increase in N_d at $w \gtrsim 1.5 \text{ m s}^{-1}$ as observed by the CAS probes; assuming very hygroscopic Aitken mode and less
295 hygroscopic accumulation mode particles can approximately reproduce the trend in $N_{d,m}$ from the CDP.

Varying κ_{acc} from 0.1 to 0.6 leads to a large increase of $N_{d,p}$ at all w . The corresponding change in $N_{d,p}$ by increasing κ_{Ait} is much smaller. The reason for this relatively smaller sensitivity of $N_{d,p}$ to κ_{Ait} is the fact that the supersaturation in the cloud is mostly controlled by the droplet growth on accumulation mode particles. The sensitivity of $N_{d,p}$ formed on Aitken mode particles to κ_{acc} is slightly larger if $\kappa_{\text{acc}} = 0.1$ as compared to $\kappa_{\text{acc}} = 0.6$, because in the latter case the supersaturation
300 is efficiently suppressed preventing a higher number of Aitken mode particles from activating. Overall we can conclude that assuming different κ values for accumulation and Aitken mode leads to a better representation of the observed trends of $N_{d,m}$ with w (Tables S16 and S17). However, in the absence of more information on the particle hygroscopicity we cannot state with certainty that the assumptions of the two κ values are appropriate for this aerosol population. Figure 4d clearly shows that the simplified assumption of a single κ is not appropriate to infer $N_{d,p}$ for low aerosol loading and when the particle number

305 concentrations of the accumulation and Aitken modes are comparable. By using a single κ value, we cannot reproduce the observed continuously strong increase of $N_{d,m}$ for the whole w range. Instead we predict a smaller increase at $w \sim 1 \text{ m s}^{-1}$, i.e., a flattening of the curve.

In general, the observed trends of N_d with w for flights AC07, AC09 and AC18 confirm results from previous sensitivity studies that have shown that with increasing w , changes in N_d become small and, thus, sensitivity of N_d to κ and w decreases
310 (Ervens et al., 2005; Fountoukis et al., 2007; Reutter et al., 2009). In these studies, it was demonstrated that the sensitivity of N_d to N_a becomes small when nearly all particles are activated ('aerosol-limited regime'). For these simulations, either only an accumulation mode was considered, or N_d closure studies were performed for situations with low w and/or fairly small Aitken mode particles ($< 40 \text{ nm}$) that were not predicted to activate.

Anttila and Kerminen (2007) showed in a model study focusing only on Aitken mode particles that N_d is highly sensitive
315 to the chemical composition of Aitken mode particles. In our recent model study, we systematically explored the extent to which the presence of an Aitken mode might significantly affect N_d as a function of updraft velocity (Pöhlker et al., 2021). In that study, we show that the sensitivities of $N_{d,p}$ are different to the properties (N_a , κ) of accumulation and Aitken mode particles, respectively. Generally, we find that $N_{d,m}$ is not highly sensitive to Aitken mode particle properties in the presence of a dominant accumulation mode, which is in agreement to our results in Figures 4 and S7.

320 4.2 Influence of aerosol number concentration (N_a) on predicted N_d

The measurements of N_a were associated with uncertainties of $\pm \sim 20\%$ (Section 2.1). In order to account for this uncertainty and possible fluctuation in N_a at cloud base, $N_{d,m}$ and $N_{d,p}$ are compared for all flights, using N_a (Figure 1), reduced and increased by 20%, 30% and 40% as model input, respectively. Figure 5 shows the comparison of measured and predicted N_d assuming the uncertainty range of N_a . Figure 4a-d shows the $N_{d,p}$ with the values of κ that are within the uncertainty range of
325 cloud probes measurements. The green lines show the model results for $\kappa = 0.1$ for flights AC07, AC09 and AC18 and $\kappa = 0.2$ for flight AC19, which show the smallest absolute bias and RMSE to the measured data (Tables S9, S12, S15 and S18); the other lines denote $N_{d,p}$ using the higher and lower input N_a . For flights AC07, AC09 and AC18, $N_{d,p}$ is within the range of $N_{d,m}$ for the assumed model parameter space. Also the curves for $N_{d,p}$ as a function of w exhibit a similar shape as predicted for a variation in κ . The agreement between measurements and model results decreases with increasing w . However, unlike the $N_{d,p}$
330 curves for high κ that level off at high w when all particles are activated, an increase in N_a leads to continuously higher N_d as the aerosol-limited regime is not yet reached.

Using different κ values for the two modes leads to a better representation of the N_d trend with w , i.e., the shape of the curve can be fairly reproduced for different combinations of separate values of κ_{acc} and κ_{ait} . However, N_d is systematically underestimated which suggests that in addition uncertainties in N_a are important for N_d closure and likely more important
335 than those in κ . In fact, the closure results in Fig. 5 show that $N_{d,m}$ can be reproduced well by the model over wide w ranges if the variability in N_a of $\pm 20\%$ and the uncertainty in $N_{d,m}$ is taken into account, and an average hygroscopicity of $\kappa = 0.1$ is assumed for the aerosol in the Amazon basin during the dry season and $\kappa = 0.2$ for that in the western tropical Atlantic. However, also the assumption of two different κ values leads to good N_d closure for the bimodal ASD as observed during

flight AC19 (Fig. 5e). These two different assumptions on the mode hygroscopicities result in ambiguous conclusions on the importance of knowledge of κ values for Aitken mode particles contribute to N_d (Pöhlker et al., 2021)(?). Generally, for both sets of simulations, i.e., varying N_a or κ , the agreement between model and measurements is best at low w . At $w \gtrsim 2 \text{ m s}^{-1}$, the $N_{d,p}$ curves flatten suggesting a decreasing sensitivity of $N_{d,p}$ above this w threshold. Unlike the trends in $N_{d,p}$ for different κ values, the lines for different N_a keep diverging with increasing w . Thus, the sensitivity of $N_{d,p}$ to N_a is predicted to remain high, and nearly independent of w which may point to conditions near the aerosol-limited regime.

4.3 Sensitivities of N_d predictions to w , N_a and κ

The sensitivities of cloud drop number concentrations to hygroscopicity (κ), N_a and w have been explored in numerous previous studies. In the following, we place our results in the context of such studies. Consistent with such previous sensitivity studies, we calculated the sensitivity ξ of $N_{d,p}$ to κ , N_a and w

$$\xi(X) = \frac{\partial \ln N_d}{\partial \ln X} \quad (\text{E.1})$$

whereas X is κ , N_a or w , respectively. The results are summarized in Figure S7. They show that $\xi(\kappa)$ is smallest as compared to $\xi(N_a)$ and $\xi(w)$. $\xi(\kappa)$ is highest for low κ as conditions, the activated fraction is smallest and thus a small change in κ might cause a significant change in N_d . κ is often termed 'effective hygroscopicity' since it is used as a parameter that reflects multiple parameters that affect the water uptake by the particles. Generally, sensitivities are high under conditions of high supersaturation which are present at high w and/or low N_a . While the sensitivities calculated for flights AC07, AC09 and AC18 follow these general trends according to N_a , the $\xi(X)$ values are higher for AC19 even though N_a was lowest for this flight. The reason for this difference is the successive activation of Aitken mode particles at high w (Pöhlker et al. (2021)(?)). A sensitivity study for the same flights has been performed previously (Cecchini et al., 2017) in which the sensitivity sensitivities of N_d and effective cloud droplet diameter to N_a and w was were explored in detail at various heights in cloud. The focus of that study was the change of the sensitivities during cloud evolution, i.e., as a function of height in cloud. In the present study, we focus on the sensitivities of N_d near cloud base, but additionally explore the importance of κ in determining N_d . Such analysis can be used to give guidance for future measurements in similar clouds on the absolute values and relative importance of the three parameters to predict N_d .

Generally, prior sensitivity studies agree in the rankings of the relative importance of hygroscopicity (κ), N_a and w , as also shown in Figure S7. Feingold (2003) has shown that N_a has the largest influence on effective radius which is indirectly related to N_d . The sensitivities to the effective radius are typically smaller than those to N_d (Pardo et al., 2019). In our recent model study, we have shown that in the transitional regime, i.e., in the parameter space between the aerosol- and updraft limited regimes, as defined by Reutter et al. (2009), N_d can be equally sensitive to κ and w (Pöhlker et al., 2021)(?). In that study, we show latter study, it was shown that with increasing N_a , the sensitivities to both parameters decrease; however, the sensitivity of N_d to w remains higher under such conditions than that to κ .

The uncertainties in updraft measurements are larger than those of hygroscopicity due to the great variability of w near cloud base. Peng et al. (2005) compared $N_{d,p}$ based on a w distribution in a range of $0.09 - 1 \text{ m s}^{-1}$ and using characteristic




Figure 5. Cloud droplet number concentration (N_d) as a function of updraft velocity near cloud base of convective clouds during flights: a) AC07, b) AC09, c) AC18, d) and e) AC19. The measured updraft velocities are based on the “probability matching method” (PMM) using the same percentiles for updraft velocity and $N_{d,m}$ (Section 3.1). The black diamond and triangle symbols represent $N_{d,m}$ near cloud base with the CAS-DPOL and CCP-CDP probes, respectively. Measurement uncertainties (indicated by error bars) are $\sim 21\%$ and $\sim 10\%$ for CAS-DPOL and CCP-CDP data (Braga et al. (2017a)). The lines show $N_{d,p}$ assuming the uncertainty range of N_a measurements, colored-coded by ΔN_a [%].

single w values. They found differences in $N_{d,p}$ on the order of $< 10\%$ for the two sets of model simulations. Meskhidze et al. (2005) performed model simulations of low-level cumuliform clouds for which a range of $0.9 \text{ m s}^{-1} \leq w \leq 2.8 \text{ m s}^{-1}$ had been observed. They concluded that parameterizations of $N_{d,p}$ should include a weighting factor for high values of w as otherwise
375 N_d might be biased high due to enhanced vertical velocity within in cloud cores as compared to cloud base.

In turbulent clouds with high w , the determination of w near cloud bases might be challenging; however, the resulting uncertainties in updraft velocity or its distributions cannot explain the discrepancies between $N_{d,m}$ and $N_{d,p}$ at high w (Figures 4 and 5). Under such conditions, the activated fraction approaches unity and any increase in w would not lead to higher N_d and improve the overall N_d closure (e.g., Hsieh et al. (2009)). Therefore under such condition, $\xi(w)$ becomes small (Figure S7b,
380 e, h, k) These previous N_d sensitivity and closure studies either considered w as a fitting parameter to obtain good closure or used w values or distributions as relatively poorly constrained parameters. The PMM analysis as applied in the current study partially overcomes these uncertainties as it provides a stronger constraint of the w and N_d pairs for the full w range (Section 3.1), as opposed to the previous studies that derived their w distributions from averaging measured updraft velocities without sorting w and $N_{d,m}$ data based on their frequency occurrence.

385 Reutter et al. (2009) termed conditions under which nearly all particles are activated into cloud droplets as ‘aerosol-limited regime’ when N_d is only dependent on N_a , and not on w . Such conditions are present at relatively low total N_a and high w , i.e., when the maximum supersaturation in the cloud is relatively high. When an increase in N_a results in an equal increase in N_d , $\xi(N_a)$ approaches unity (Figure S7c, f, i, l). The measured and predicted activated fractions for flights AC07, AC09 and AC18 reach $\geq 80\%$ at updraft velocities of $w \gtrsim 1 \text{ m s}^{-1}$ if the measured value is based on the CAS data (Figure 4). Therefore, we
390 conclude that the sensitivity of N_d to N_a is much greater than that to w under these conditions which is also reflected by the rather small increase in N_d with w at high updraft velocities.

Overall, the variability of predicted N_d due to inferred κ ranges in the present study confirm trends from previous sensitivity studies for mono-modal aerosol size distributions: The sensitivity to N_d decreases with increasing w , i.e. when the activated fraction is large and activation of additional smaller particles increases N_d only to a small extent (Figure S7 and Ervens et al.
395 (2005); Reutter et al. (2009); Cecchini et al. (2017); Pardo et al. (2019)). If low hygroscopicity limits the water vapor uptake, a small change in κ may lead to a significant change in N_d , resulting in high $\xi(\kappa)$ values. A change of κ by the same factor for highly hygroscopic particles, however, might not lead to a significant change in N_d due to the regulation of the supersaturation (‘buffering’), i.e., the efficient growth of more cloud droplets which, in turn, reduces the supersaturation. Our sensitivity study of AC19 exceeds these previous sensitivity studies that focused on monomodal aerosol size distributions. We show that the
400 uncertainties in $N_{d,p}$ become larger under conditions when Aitken mode particles contribute to N_d as only at very high w the aerosol-limited regime is reached and $\xi(\kappa)$ and $\xi(w)$ decrease. Qualitatively this was also suggested in a previous N_d closure study for marine stratocumulus clouds, where it was concluded that only the presence of an Aitken mode could explain the high $N_{d,m}$ at updraft velocities of $w \geq 1 \text{ m s}^{-1}$ (Schulze et al., 2020). Our analysis exceeds this former study as we show that the w threshold above which Aitken mode particles contribute to N_d depends on the properties (e.g., κ , N_{acc}) of the accumulation
405 mode. In addition, we show that various combinations of inferred κ_{acc} and κ_{Ait} result in similar $N_{d,p}$ and thus cannot be constrained without more detailed composition measurements. While these conclusions are drawn on a single observationally-

based case study, a more systematic analysis of parameter ranges of Aitken and accumulation mode particles is provided in our [follow-up study \(Pöhlker et al., 2021\)](#) [recent study \(?\)](#).

5 Summary and conclusions

410 Airborne measurements of cloud droplet number concentrations ($N_{d,m}$), aerosol particle size distributions and updraft velocities (w) near cloud base were performed during the ACRIDICON-CHUVA campaign in September 2014. Using an adiabatic air parcel model, the importance of aerosol particle number concentration (N_a) and [effective](#) hygroscopicity (κ) and their uncertainties on predicted cloud droplet number concentrations ($N_{d,p}$) near cloud bases of growing convective cumuli, formed over the Amazon and western Atlantic, were explored. Data from aerosol and cloud probes onboard HALO were used as
415 model input for this cloud droplet closure analysis. Model results for four different scenarios in terms of aerosol loading and size distributions, and of w confirm previously suggested values of the hygroscopicity parameter κ to reasonably predict N_d for most conditions: best N_d closure is achieved for [an effective hygroscopicity of](#) $\kappa \sim 0.1$ for the Amazon basin during the dry season using the full data set of CCP-CDP and CAS-DPOL measurements. Above the western Atlantic best N_d closure was achieved for $\kappa \sim 0.2$ applying a single κ value for both Aitken and accumulation modes; an even better representation
420 of the increase in N_d with w was obtained when moderately hygroscopic accumulation mode particles ($\kappa_{acc} = 0.2$) and highly hygroscopic Aitken mode particles ($\kappa_{ait} = 0.8$) were assumed.

While we could not further constrain the hygroscopicities of the two modes based on the available data, our results suggest that knowledge of Aitken mode particle properties is required to predict cloud droplet number concentrations in convective clouds and/or clean air masses. We conclude that in the case of a bi-modal aerosol size distribution with distinct Aitken and
425 accumulation modes as encountered during flight AC19, $N_{d,p}$ may be significantly underestimated as compared to $N_{d,p}$ for $w \gtrsim 0.5 \text{ m s}^{-1}$ if a single κ for both modes is assumed that might be only appropriate for the larger accumulation mode particles. Our results also suggest that the ratio of the number concentrations of the Aitken and accumulation modes and their κ values can influence cloud properties near cloud base differently than for one-modal aerosol size distributions. More detailed sensitivity studies of cloud properties to Aitken mode aerosol properties (N_a , κ) have been recently performed for wider parameter ranges
430 to identify conditions, under which they might affect aerosol-cloud interactions ([Pöhlker et al., 2021](#)) [\(?\)](#).

Our droplet closure study represents a complementary approach to constrain CCN hygroscopicity, in addition to previous studies in the same region, in which a similar κ range (0.1–0.35) was determined for aerosol in the Amazon Basin, and a range of $0.1 < \kappa < 0.9$ above the ocean, based on CCN measurements and detailed analysis of chemical composition (Wex et al., 2016; Thalman et al., 2017; Pöhlker et al., 2016, 2018). Our comparison between predicted and measured N_d showed
435 largest discrepancies at high updraft velocities ($w > 2.5 \text{ m s}^{-1}$), which could be possibly explained by non-adiabaticity and/or entrainment of aerosol particles near cloud bases of convective clouds.

While in previous cloud droplet number closure studies the updraft velocity was often assumed to be a major factor of uncertainty, this parameter was well constrained in the current study. Implying that higher N_d are formed in regions of higher updraft velocities, we sorted observed data of N_d and w by their frequency of occurrence ('probability matching method').

440 Using this approach, we reduced the uncertainty of w for the N_d closure. Therefore, we could largely limit our sensitivity analysis to the investigation of the importance of particle hygroscopicity and number concentration for cloud droplet number concentrations.

Variability in N_a measurements ($\sim \pm 20\%$) translate into similar differences in predicted droplet number concentration as uncertainties assuming different κ values, in particular at low w . In previous cloud droplet number closure studies, composition effects, such as slow dissolution of soluble compounds (Asa-Awuku and Nenes, 2007), reduced surface tension or variation of the water mass accommodation coefficient (Conant et al., 2004) have been inferred to explain observed droplet number concentrations. Our analysis shows that measurement uncertainties in basic aerosol properties might equally explain such differences. If particles exceed a hygroscopicity threshold (~~$\kappa > \sim 0.3$~~ $\kappa > 0.3$), predicted cloud droplet number concentration becomes very insensitive to κ when a large fraction of all particles are activated ('aerosol-limited regime'). In the presence of a distinct Aitken mode, the parameter space (w, N_a, κ) at which this regime prevails is suggested to be shifted to even higher updraft velocity regimes than in the presence of monomodal accumulation mode size distributions.

445
450

Data availability. The data used in this study can be found at <https://halo-db.pa.op.dlr.de/mission/5>.

Author contributions. RCB, BE, MLP led the analyses and the manuscript preparation. The measurements of aerosol and cloud properties were conducted and analyzed by RCB, DF, BAH, TJ, OOK, CP, DS, CV, AW, MLP. The measurements were led by MOA, MW, UP. The modeling studies were performed and interpreted by RCB, BE, DR, JDF, OL, LHP, LATM, MLP.

455

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. The ACRIDICON–CHUVA campaign was supported by the Max Planck Society (MPG), the German Science Foundation (DFG Priority Program SPP 1294), the German Aerospace Center (DLR), and a wide range of other institutional partners. BE was supported by the French National Research Agency (ANR) (grant no. ANR-17-MPGA- 0013).

460 References

- Andreae, M. O. and Raemdonck, H.: Dimethyl Sulfide in the Surface Ocean and the Marine Atmosphere: A Global View, *Science*, 221, 744–747, <https://doi.org/10.1126/science.221.4612.744>, 1983.
- Andreae, M. O. and Rosenfeld, D.: Aerosol-cloud-precipitation interactions. Part I. The nature and sources of cloud-active aerosols, *Earth-Science Reviews*, 89, 13–41, <https://doi.org/10.1016/j.earscirev.2008.03.001>, 2008.
- 465 Andreae, M. O., Rosenfeld, D., Artaxo, P., Costa, A. A., Frank, G. P., Longo, K. M., and Silva-Dias, M. A. F.: Smoking rain clouds over the Amazon., *Science (New York, N.Y.)*, 303, 1337–1342, <https://doi.org/10.1126/science.1092779>, 2004.
- Andreae, M. O., Afchine, A., Albrecht, R., Amorim Holanda, B., Artaxo, P., Barbosa, H. M., Borrmann, S., Cecchini, M. A., Costa, A., Dollner, M., Fütterer, D., Järvinen, E., Jurkat, T., Klimach, T., Konemann, T., Knote, C., Krämer, M., Krisna, T., Machado, L. A., Mertes, S., Minikin, A., Pöhlker, C., Pöhlker, M. L., Pöschl, U., Rosenfeld, D., Sauer, D., Schlager, H., Schnaiter, M., Schneider, J., Schulz, C., Spanu, A., Sperling, V. B., Voigt, C., Walser, A., Wang, J., Weinzierl, B., Wendisch, M., and Ziereis, H.: Aerosol characteristics and particle production in the upper troposphere over the Amazon Basin, *Atmospheric Chemistry and Physics*, <https://doi.org/10.5194/acp-18-921-2018>, 2018.
- 470 Anttila, T. and Kerminen, V. M.: On the contribution of Aitken mode particles to cloud droplet populations at continental background areas - A parametric sensitivity study, *Atmospheric Chemistry and Physics*, 7, 4625–4637, <https://doi.org/10.5194/acp-7-4625-2007>, 2007.
- 475 Anttila, T., Brus, D., Jaatinen, A., Hyvärinen, A. P., Kivekäs, N., Romakkaniemi, S., Komppula, M., and Lihavainen, H.: Relationships between particles, cloud condensation nuclei and cloud droplet activation during the third Pallas Cloud Experiment, *Atmospheric Chemistry and Physics*, 12, 11 435–11 450, <https://doi.org/10.5194/acp-12-11435-2012>, 2012.
- Artaxo, P.: Physical and chemical properties of aerosols in the wet and dry seasons in Rondônia, Amazonia, *Journal of Geophysical Research*, 107, 8081, <https://doi.org/10.1029/2001JD000666>, 2002.
- 480 Asa-Awuku, A. and Nenes, A.: Effect of solute dissolution kinetics on cloud droplet formation: Extended Köhler theory, *Journal of Geophysical Research Atmospheres*, 112, <https://doi.org/10.1029/2005JD006934>, 2007.
- Baumgardner, D., Brenguier, J. L., Bucholtz, A., Coe, H., DeMott, P., Garrett, T. J., Gayet, J. F., Hermann, M., Heymsfield, A., Korolev, A., Krämer, M., Petzold, A., Strapp, W., Pilewskie, P., Taylor, J., Twohy, C., Wendisch, M., Bachalo, W., and Chuang, P.: Airborne instruments to measure atmospheric aerosol particles, clouds and radiation: A cook's tour of mature and emerging technology, <https://doi.org/10.1016/j.atmosres.2011.06.021>, 2011.
- 485 Braga, R. C., Rosenfeld, D., Weigel, R., Jurkat, T., Andreae, M. O., Wendisch, M., Pöhlker, M. L., Klimach, T., Pöschl, U., Pöhlker, C., Voigt, C., Mahnke, C., Borrmann, S., Albrecht, R. I., Molleker, S., Vila, D. A., Machado, L. A. T., and Artaxo, P.: Comparing parameterized versus measured microphysical properties of tropical convective cloud bases during the ACRIDICON-CHUVA campaign, *Atmos. Chem. Phys.*, <https://doi.org/doi.org/10.5194/acp-2016-872>, 2017a.
- 490 Braga, R. C., Rosenfeld, D., Weigel, R., Jurkat, T., Andreae, M. O., Wendisch, M., Pöschl, U., Voigt, C., Mahnke, C., Borrmann, S., Albrecht, R. I., Molleker, S., Vila, D. A., Machado, L. A. T., and Grulich, L.: Further evidence for CCN aerosol concentrations determining the height of warm rain and ice initiation in convective clouds over the Amazon basin, *Atmos. Chem. Phys.*, 17, 14 433–14 456, <https://doi.org/10.5194/acp-17-14433-2017>, 2017b.
- Broekhuizen, K., Chang, R. Y., Leaitch, W. R., Li, S. M., and Abbatt, J. P.: Closure between measured and modeled cloud condensation nuclei (CCN) using size-resolved aerosol compositions in downtown Toronto, <https://doi.org/10.5194/acp-6-2513-2006>, 2006.
- 495

- Cai, Y., Montague, D. C., Mooiweer-Bryan, W., and Deshler, T.: Performance characteristics of the ultra high sensitivity aerosol spectrometer for particles between 55 and 800 nm: Laboratory and field studies, *Journal of Aerosol Science*, 39, 759–769, <https://doi.org/10.1016/j.jaerosci.2008.04.007>, 2008.
- 500 Carrico, C. M., Petters, M. D., Kreidenweis, S. M., Sullivan, A. P., McMeeking, G. R., Levin, E. J., Engling, G., Malm, W. C., and Collett, J. L.: Water uptake and chemical composition of fresh aerosols generated in open burning of biomass, *Atmospheric Chemistry and Physics*, 10, 5165–5178, <https://doi.org/10.5194/acp-10-5165-2010>, 2010.
- Cecchini, M. A., MacHado, L. A., Andreae, M. O., Martin, S. T., Albrecht, R. I., Artaxo, P., Barbosa, H. M., Borrmann, S., Fütterer, D., Jurkat, T., Mahnke, C., Minikin, A., Molleker, S., Pöhlker, M. L., Pöschl, U., Rosenfeld, D., Voigt, C., Weinzierl, B., and Wendisch, M.: Sensitivities of Amazonian clouds to aerosols and updraft speed, *Atmospheric Chemistry and Physics*, 17, 10037–10050, 505 <https://doi.org/10.5194/acp-17-10037-2017>, 2017.
- Chuang, P. Y., Collins, D. R., Pawlowska, H., Snider, J. R., Jonsson, H. H., Brenguier, J. L., Flagan, R. C., and Seinfeld, J. H.: CCN measurements during ACE-2 and their relationship to cloud microphysical properties, *Tellus, Series B: Chemical and Physical Meteorology*, 52, 843–867, <https://doi.org/10.1034/j.1600-0889.2000.00018.x>, 2000.
- Chubb, T., Huang, Y., Jensen, J., Campos, T., Siems, S., and Manton, M.: Observations of high droplet number concentrations in Southern 510 Ocean boundary layer clouds, *Atmospheric Chemistry and Physics*, 16, 971–987, <https://doi.org/10.5194/acp-16-971-2016>, 2016.
- Conant, W. C., VanReken, T. M., Rissman, T. A., Varutbangkul, V., Jonsson, H. H., Nenes, A., Jimenez, J. L., Delia, A. E., Bahreini, R., Roberts, G. C., Flagan, R. C., and Seinfeld, J. H.: Aerosol-cloud drop concentration closure in warm cumulus, *Journal of Geophysical Research D: Atmospheres*, 109, <https://doi.org/10.1029/2003JD004324>, 2004.
- Engelhart, G. J., Hennigan, C. J., Miracolo, M. A., Robinson, A. L., and Pandis, S. N.: Cloud condensation nuclei activity of fresh primary 515 and aged biomass burning aerosol, *Atmospheric Chemistry and Physics*, 12, 7285–7293, <https://doi.org/10.5194/acp-12-7285-2012>, 2012.
- Ervens, B., Feingold, G., and Kreidenweis, S. M.: Influence of water-soluble organic carbon on cloud drop number concentration, *Journal of Geophysical Research D: Atmospheres*, 110, 1–14, <https://doi.org/10.1029/2004JD005634>, 2005.
- Ervens, B., Cubison, M. J., Andrews, E., Feingold, G., Ogren, J. A., Jimenez, J. L., Quinn, P. K., Bates, T. S., Wang, J., Zhang, Q., Coe, H., Flynn, M., and Allan, J. D.: CCN predictions using simplified assumptions of organic aerosol composition and mixing state: A synthesis 520 from six different locations, *Atmospheric Chemistry and Physics*, 10, 4795–4807, <https://doi.org/10.5194/acp-10-4795-2010>, 2010.
- Feingold, G.: Modeling of the first indirect effect: Analysis of measurement requirements, *Geophysical Research Letters*, <https://doi.org/10.1029/2003GL017967>, 2003.
- Feingold, G. and Heymsfield, A. J.: Parameterizations of condensational growth of droplets for use in general circulation models, *Journal of the Atmospheric Sciences*, 49, 2325–2342, [https://doi.org/10.1175/1520-0469\(1992\)049<2325:POCGOD>2.0.CO;2](https://doi.org/10.1175/1520-0469(1992)049<2325:POCGOD>2.0.CO;2), 1992.
- 525 Fountoukis, C., Nenes, A., Meskhidze, N., Bahreini, R., Conant, W. C., Jonsson, H., Murphy, S., Sorooshian, A., Varutbangkul, V., Brechtel, F., Flagan, R. C., and Seinfeld, J. H.: Aerosol-cloud drop concentration closure for clouds sampled during the International Consortium for Atmospheric Research on Transport and Transformation 2004 campaign, *Journal of Geophysical Research Atmospheres*, 112, <https://doi.org/10.1029/2006JD007272>, 2007.
- Gong, X., Wex, H., Müller, T., Wiedensohler, A., Höhler, K., Kandler, K., Ma, N., Dietel, B., Schiebel, T., Möhler, O., and Stratmann, 530 F.: Characterization of aerosol properties at Cyprus, focusing on cloud condensation nuclei and ice-nucleating particles, *Atmospheric Chemistry and Physics*, 19, 10883–10900, <https://doi.org/10.5194/acp-19-10883-2019>, 2019.
- Haddad, Z. S. and Rosenfeld, D.: Optimality of Empirical Z-R Relations, *Quarterly Journal of the Royal Meteorological Society*, 123, 1283–1293, <https://doi.org/10.1002/qj.49712354107>, 1997.

- Holanda, B. A., Pöhlker, M. L., Walter, D., Saturno, J., Sörgel, M., Ditas, J., Ditas, F., Schulz, C., Aurélio Franco, M., Wang, Q., Donth, T.,
535 Artaxo, P., Barbosa, H. M., Borrmann, S., Braga, R., Brito, J., Cheng, Y., Dollner, M., Kaiser, J. W., Klimach, T., Knote, C., Krüger, O. O.,
Fütterer, D., t.V. Lavric, J., Ma, N., MacHado, L. A., Ming, J., Morais, F. G., Paulsen, H., Sauer, D., Schlager, H., Schneider, J., Su, H.,
Weinzierl, B., Walser, A., Wendisch, M., Ziereis, H., Zöger, M., Pöschl, U., Andreae, M. O., and Pöhlker, C.: Influx of African biomass
burning aerosol during the Amazonian dry season through layered transatlantic transport of black carbon-rich smoke, *Atmospheric Chem-
istry and Physics*, 20, 4757–4785, <https://doi.org/10.5194/acp-20-4757-2020>, 2020.
- 540 Hsieh, W. C., Nenes, A., Flagan, R. C., Seinfeld, J. H., Buzorius, G., and Jonsson, H.: Parameterization of cloud droplet
size distributions: Comparison with parcel models and observations, *Journal of Geophysical Research Atmospheres*, 114,
<https://doi.org/10.1029/2008JD011387>, 2009.
- IPCC: *Climate Change 2021 - The Physical Science Basis*, Cambridge University Press, cambridge edn., 2021.
- Kleine, J., Voigt, C., Sauer, D., Schlager, H., Scheibe, M., Jurkat-Witschas, T., Kaufmann, S., Kärcher, B., and Anderson, B. E.:
545 In Situ Observations of Ice Particle Losses in a Young Persistent Contrail, *Geophysical Research Letters*, 45, 13,553–13,561,
<https://doi.org/10.1029/2018GL079390>, 2018.
- Köhler, H.: The nucleus in and the growth of hygroscopic droplets, *Transactions of the Faraday Society*, 32, 1152–1161, 1936.
- Lance, S., Brock, C. A., Rogers, D., and Gordon, J. A.: Water droplet calibration of the Cloud Droplet Probe (CDP) and in-
flight performance in liquid, ice and mixed-phase clouds during ARCPAC, *Atmospheric Measurement Techniques*, 3, 1683–1706,
550 <https://doi.org/10.5194/amt-3-1683-2010>, 2010.
- Leaitch, W. R., Lohmann, U., Russell, L. M., Garrett, T., Shantz, N. C., Toom-Sauntry, D., Strapp, J. W., Hayden, K. L., Marshall, J.,
Wolde, M., Worsnop, D. R., and Jayne, J. T.: Cloud albedo increase from carbonaceous aerosol, *Atmospheric Chemistry and Physics*, 10,
7669–7684, <https://doi.org/10.5194/acp-10-7669-2010>, 2010.
- Mallaun, C., Giez, A., and Baumann, R.: Calibration of 3-D wind measurements on a single-engine research aircraft, *Atmospheric Measure-
ment Techniques*, 8, 3177–3196, <https://doi.org/10.5194/amt-8-3177-2015>, 2015.
- 555 McFiggans, G., P. Artaxo, Baltensperger, U., Coe, H., Facchini, M. C., Feingold, G., Fuzzi, S., Gysel, M., Laaksonen, A., Lohmann, U.,
Mentel, T. F., Murphy, D. M., O’Dowd, C. D., Snider, J., and Weingartner, E.: The effect of physical and chemical aerosol properties on
warm cloud droplet activation, *Atmos. Chem. Phys.*, 6, 2593–2649, <https://doi.org/10.5194/acp-6-2593-2006>, 2006.
- Meskhidze, N., Nenes, A., Conant, W. C., and Seinfeld, J. H.: Evaluation of a new cloud droplet activation parameteriza-
560 tion with in situ data from CRYSTAL-FACE and CSTRIFE, *Journal of Geophysical Research D: Atmospheres*, 110, 1–10,
<https://doi.org/10.1029/2004JD005703>, 2005.
- Moore, R. H., Wiggins, E. B., Ahern, A. T., Zimmerman, S., Montgomery, L., Campuzano Jost, P., Robinson, C. E., Ziemba, L. D., Win-
stead, E. L., Anderson, B. E., Brock, C. A., Brown, M. D., Chen, G., Crosbie, E. C., Guo, H., Jimenez, J. L., Jordan, C. E., Lyu, M.,
Nault, B. A., Rothfuss, N. E., Sanchez, K. J., Schueneman, M., Shingler, T. J., Shook, M. A., Thornhill, K. L., Wagner, N. L., and
565 Wang, J.: Sizing Response of the Ultra-High Sensitivity Aerosol Size Spectrometer (UHSAS) and Laser Aerosol Spectrometer (LAS)
to Changes in Submicron Aerosol Composition and Refractive Index, *Atmospheric Measurement Techniques Discussions*, 2021, 1–36,
<https://doi.org/10.5194/amt-2021-21>, 2021.
- Pardo, L. H., MacHado, L. A. T., Cecchini, M. A., and Gácita, M. S.: Quantifying the aerosol effect on droplet size distribution at cloud top,
Atmospheric Chemistry and Physics, 19, 7839–7857, <https://doi.org/10.5194/acp-19-7839-2019>, 2019.
- 570 Peng, Y., Lohmann, U., and Leaitch, W. R.: Importance of vertical velocity variations in the cloud droplet nucleation process of marine
stratus clouds, *Journal of Geophysical Research Atmospheres*, 110, 1–13, <https://doi.org/10.1029/2004JD004922>, 2005.

- Petters, M. D. and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud\condensation nucleus activity, *Atmos. Chem. Phys.*, pp. 1961–1971, <https://doi.org/10.5194/acp-7-1961-2007>, 2007.
- 575 Petzold, A., Marsh, R., Johnson, M., Miller, M., Sevcenco, Y., Delhaye, D., Ibrahim, A., Williams, P., Bauer, H., Crayford, A., Bachalo, W. D., and Raper, D.: Evaluation of Methods for Measuring Particulate Matter Emissions from Gas Turbines, *Environmental Science & Technology*, 45, 3562–3568, <https://doi.org/10.1021/es103969v>, PMID: 21425830, 2011.
- Pöhlker, M. L., Pöhlker, C., Ditas, F., Klimach, T., De Angelis, I. H., Araújo, A., Brito, J., Carbone, S., Cheng, Y., Chi, X., Ditz, R., Gunthe, S. S., Kesselmeier, J., Könemann, T., Lavrič, J. V., Martin, S. T., Mikhailov, E., Moran-Zuloaga, D., Rose, D., Saturno, J., Su, H., Thalman, R., Walter, D., Wang, J., Wolff, S., Barbosa, H. M., Artaxo, P., Andreae, M. O., and Pöschl, U.: Long-term observations of cloud condensation nuclei in the Amazon rain forest - Part 1: Aerosol size distribution, hygroscopicity, and new model parametrizations for CCN prediction, *Atmospheric Chemistry and Physics*, 16, 15 709–15 740, <https://doi.org/10.5194/acp-16-15709-2016>, 2016.
- 580 Pöhlker, M. L., Ditas, F., Saturno, J., Klimach, T., Hrabě De Angelis, I., Araújo, A. C., Brito, J., Carbone, S., Cheng, Y., Chi, X., Ditz, R., Gunthe, S. S., Holanda, B. A., Kandler, K., Kesselmeier, J., Könemann, T., Krüger, O. O., Lavric, J. V., Martin, S. T., Mikhailov, E., Moran-Zuloaga, D., Rizzo, L. V., Rose, D., Su, H., Thalman, R., Walter, D., Wang, J., Wolff, S., Barbosa, H. M., Artaxo, P., Andreae, M. O., Pöschl, U., and Pöhlker, C.: Long-term observations of cloud condensation nuclei over the Amazon rain forest - Part 2: Variability and characteristics of biomass burning, long-range transport, and pristine rain forest aerosols, *Atmospheric Chemistry and Physics*, <https://doi.org/10.5194/acp-18-10289-2018>, 2018.
- 585 Pöhlker, M. L., Zhang, M., Campos Braga, R., Krüger, O. O., Pöschl, U., and Ervens, B.: Aitken mode particles as CCN in aerosol- and updraft-sensitive regimes of cloud droplet formation, *Atmospheric Chemistry and Physics*, 21, 11 723–11 740, <https://doi.org/10.5194/acp-21-11723-2021>, <https://acp.copernicus.org/articles/21/11723/2021/>, 2021.
- 590 Pruppacher, H. and Klett, J. D.: *Microphysics of Clouds and Precipitation*, Kluwer Academic Publishers, second edn., 1997.
- Quinn, P. K., Coffman, D. J., Johnson, J. E., Upchurch, L. M., and Bates, T. S.: Small fraction of marine cloud condensation nuclei made up of sea spray aerosol, *Nature Geoscience*, 10, 674–679, <https://doi.org/10.1038/ngeo3003>, 2017.
- Reutter, P., Su, H., Trentmann, J., Simmel, M., Rose, D., Gunthe, S. S., Wernli, H., Andreae, M. O., and Pöschl, U.: Aerosol- and updraft-limited regimes of cloud droplet formation: influence of particle number, size and hygroscopicity on the activation of cloud condensation nuclei (CCN), *Atmospheric Chemistry and Physics*, 9, 7067–7080, <https://doi.org/10.5194/acp-9-7067-2009>, 2009.
- 595 Rissler, J., Swietlicki, E., Zhou, J., Roberts, G., Andreae, M. O., Gatti, L. V., and Artaxo, P.: Physical properties of the sub-micrometer aerosol over the Amazon rain forest during the wet-to-dry season transition - comparison of modeled and measured CCN concentrations, *Atmospheric Chemistry and Physics*, 4, 2119–2143, <https://doi.org/10.5194/acp-4-2119-2004>, 2004.
- 600 Roberts, G. C., Nenes, A., Seinfeld, J. H., and Andreae, M. O.: Impact of biomass burning on cloud properties in the Amazon Basin - art. no. 4062, *Journal of Geophysical Research - Atmospheres*, 108, 4062, <https://doi.org/10.1029/2001JD000985>, 2003.
- Rosenfeld, D.: Cloud-Aerosol-Precipitation Interactions Based of Satellite Retrieved Vertical Profiles of Cloud Microstructure, in: *Remote Sensing of Aerosols, Clouds, and Precipitation*, pp. 129–152, Elsevier, <https://doi.org/10.1016/B978-0-12-810437-8.00006-2>, 2018.
- Rosenfeld, D., Lohmann, U., Raga, G. B., O’Dowd, C. D., Kulmala, M., Fuzzi, S., Reissell, A., and Andreae, M. O.: Flood or drought: how do aerosols affect precipitation?, *Science (New York, N.Y.)*, 321, 1309–1313, <https://doi.org/10.1126/science.1160606>, 2008.
- 605 Schulze, B. C., Charan, S. M., Kenseth, C. M., Kong, W., Bates, K. H., Williams, W., Metcalf, A. R., Jonsson, H. H., Woods, R., Sorooshian, A., Flagan, R. C., and Seinfeld, J. H.: Characterization of Aerosol Hygroscopicity Over the Northeast Pacific Ocean: Impacts on Prediction of CCN and Stratocumulus Cloud Droplet Number Concentrations, *Earth and Space Science*, 7, <https://doi.org/10.1029/2020EA001098>, 2020.

- 610 Shaw, R. A., Reade, W. C., Collins, L. R., and Verlinde, J.: Preferential Concentration of Cloud Droplets by Turbulence: Effects on the Early Evolution of Cumulus Cloud Droplet Spectra, *Journal of the Atmospheric Sciences*, 55, 1965–1976, [https://doi.org/10.1175/1520-0469\(1998\)055<1965:PCOCDB>2.0.CO;2](https://doi.org/10.1175/1520-0469(1998)055<1965:PCOCDB>2.0.CO;2), 1998.
- Thalman, R., De Sá, S. S., Palm, B. B., Barbosa, H. M., Pöhlker, M. L., Alexander, M. L., Brito, J., Carbone, S., Castillo, P., Day, D. A., Kuang, C., Manzi, A., Ng, N. L., Sedlacek, A. J., Souza, R., Springston, S., Watson, T., Pöhlker, C., Pöschl, U., Andreae, M. O., Artaxo, P., Jimenez, J. L., Martin, S. T., and Wang, J.: CCN activity and organic hygroscopicity of aerosols downwind of an urban region in central Amazonia: Seasonal and diel variations and impact of anthropogenic emissions, *Atmospheric Chemistry and Physics*, 17, 11 779–11 801, <https://doi.org/10.5194/acp-17-11779-2017>, 2017.
- Twomey, S.: The nuclei of natural cloud formation part II: The supersaturation in natural clouds and the variation of cloud droplet concentration, *Geofisica Pura e Applicata*, 43, 243–249, <https://doi.org/10.1007/BF01993560>, 1959.
- 620 Voigt, C., Schumann, U., Jurkat, T., Schäuble, D., Schlager, H., Petzold, A., Gayet, J. F., Krämer, M., Schneider, J., Borrmann, S., Schmale, J., Jessberger, P., Hamburger, T., Lichtenstern, M., Scheibe, M., Gourbeyre, C., Meyer, J., Kübbeler, M., Frey, W., Kalesse, H., Butler, T., Lawrence, M. G., Holzäpfel, F., Arnold, F., Wendisch, M., Döpelheuer, A., Gottschaldt, K., Baumann, R., Zöger, M., Sölch, I., Rautenhaus, M., and Dörnbrack, A.: In-situ observations of young contrails - Overview and selected results from the CONCERT campaign, *Atmospheric Chemistry and Physics*, 10, 9039–9056, <https://doi.org/10.5194/acp-10-9039-2010>, 2010.
- 625 Voigt, C., Schumann, U., Jessberger, P., Jurkat, T., Petzold, A., Gayet, J. F., Krmer, M., Thornberry, T., and Fahey, D. W.: Extinction and optical depth of contrails, *Geophysical Research Letters*, 38, <https://doi.org/10.1029/2011GL047189>, 2011.
- Voigt, C., Schumann, U., Minikin, A., Abdelmonem, A., Afchine, A., Borrmann, S., Boettcher, M., Buchholz, B., Bugliaro, L., Costa, A., Curtius, J., Dollner, M., Dörnbrack, A., Dreiling, V., Ebert, V., Ehrlich, A., Fix, A., Forster, L., Frank, F., and Zoeger, M.: ML-CIRRUS: The Airborne Experiment on Natural Cirrus and Contrail Cirrus with the High-Altitude Long-Range Research Aircraft HALO, *Bulletin of the American Meteorological Society*, 98, 271–288, <https://doi.org/10.1175/BAMS-D-15-00213.1>, 2017.
- 630 von der Weiden, S.-L., Drewnick, F., and Borrmann, S.: Particle Loss Calculator – a new software tool for the assessment of the performance of aerosol inlet systems, *Atmospheric Measurement Techniques*, 2, 479–494, <https://doi.org/10.5194/amt-2-479-2009>, <https://amt.copernicus.org/articles/2/479/2009/>, 2009.
- Wang, J., Lee, Y. N., Daum, P. H., Jayne, J., and Alexander, M. L.: Effects of aerosol organics on cloud condensation nucleus (CCN) concentration and first indirect aerosol effect, *Atmospheric Chemistry and Physics*, 8, 6325–6339, <https://doi.org/10.5194/acp-8-6325-2008>, 2008.
- 635 Wendisch, M. and Brenguier, J. L.: *Airborne measurements for environmental research: methods and instruments.*, Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, Germany, <https://doi.org/10.1002/9783527653218>, 2013.
- Wendisch, M., Pöschl, U., Andreae, M. O., Machado, L. A. T., Albrecht, R., Schlager, H., Rosenfeld, D., Martin, S. T., Abdelmonem, A., Afchine, A., Araújo, A., Artaxo, P., Aufmhoff, H., Barbosa, H. M. J., Borrmann, S., Braga, R., Buchholz, B., Cecchini, M. A., Costa, A., Curtius, J., Dollner, M., Dorf, M., Dreiling, V., Ebert, V., Ehrlich, A., Ewald, F., Fisch, G., Fix, A., Frank, F., Fütterer, D., Heckl, C., Heidelberg, F., Hüneke, T., Jäkel, E., Järvinen, E., Jurkat, T., Kanter, S., Kästner, U., Kenntner, M., Kesselmeier, J., Klimach, T., Knecht, M., Kohl, R., Kölling, T., Krämer, M., Krüger, M., Krisna, T. C., Lavric, J. V., Longo, K., Mahnke, C., Manzi, A. O., Mayer, B., Mertes, S., Minikin, A., Molleker, S., Münch, S., Nillius, B., Pfeilsticker, K., Pöhlker, C., Roiger, A., Rose, D., Rosenow, D., Sauer, D., Schnaiter, M., Schneider, J., Schulz, C., de Souza, R. A. F., Spanu, A., Stock, P., Vila, D., Voigt, C., Walser, A., Walter, D., Weigel, R., Weinzierl, B., Werner, F., Yamasoe, M. A., Ziereis, H., Zinner, T., and Zöger, M.: The ACRIDICON-CHUVA campaign: Studying
- 645

- tropical deep convective clouds and precipitation over Amazonia using the new German research aircraft HALO, *Bulletin of the American Meteorological Society*, p. 160128144638003, <https://doi.org/10.1175/BAMS-D-14-00255.1>, 2016.
- 650 Wex, H., Dieckmann, K., Roberts, G. C., Conrath, T., Izaguirre, M. A., Hartmann, S., Herenz, P., Schäfer, M., Ditas, F., Schmeissner, T., Henning, S., Wehner, B., Siebert, H., and Stratmann, F.: Aerosol arriving on the Caribbean island of Barbados: Physical properties and origin, *Atmospheric Chemistry and Physics*, 16, 14 107–14 130, <https://doi.org/10.5194/acp-16-14107-2016>, 2016.
- Xue, Y., Wang, L.-P., and Grabowski, W. W.: Growth of Cloud Droplets by Turbulent Collision–Coalescence, *Journal of the Atmospheric Sciences*, 65, 331–356, <https://doi.org/10.1175/2007JAS2406.1>, 2008.