

Dispersion of particulate matter (PM_{2.5}) from wood combustion for residential heating: Optimisation of mitigation actions based on large-eddy simulations

Tobias Wolf¹, Lasse H. Pettersson¹, Igor Esau¹

¹Nansen Environmental and Remote Sensing Center, Thormøhlens gate 47, 5006, Bergen, Norway

Correspondence to: Igor Esau (igore@nersc.no)

Abstract. Many cities in the world experience significant air pollution from residential wood combustion. Such an advection-diffusion problem as applied to geographically distributed small-scale pollution sources presently does not have a satisfactory theoretical or modelling solution. For example, statistical models do not allow for pollution accumulation in local stagnation zones – a type of phenomena that is commonly observed over complex terrain. This study applies a Parallelized Atmospheric Large-eddy simulation Model (PALM) to investigate dynamical phenomena that control variability and pathways of the atmospheric pollution emitted by wood-burning household stoves. The model PALM runs with a real spatial distribution of the pollution source and with realistic surface boundary conditions that characterize a medium-sized urban area fragmented by water bodies and hills. Such complex geography is expected to favour local air quality hazards, which makes this study of general interest. The case study here is based on winter conditions in Bergen, Norway. We investigate the turbulent diffusion of a passive scalar associated with small sized particles (PM_{2.5}) emitted by household stoves. The study considers air pollution effects that could be observed under different policy scenarios of stove replacement; modern wood stoves emit significantly less PM_{2.5} than the older ones, but replacement of stoves is costly and challenging process. To solve the advection-diffusion problem, we used the PALM model at spatial resolution of 10 m in an urban-sized modelling domain of 15 km by 15 km.

We found significant accumulation of near-surface pollution in the local stagnation zones. The ~~observed-simulated~~ concentrations were larger than ~~the~~ concentrations ~~obtained only~~ due to the local PM_{2.5} emission, thus, indicating dominant trans-boundary contribution of pollutants for other districts. We demonstrate how the source of critical pollution can be

attributed through model disaggregation of emissions from specific districts. The study reveals a decisive role of local air circulations over complex terrain that makes high-resolution modelling indispensable for adequate management of the urban

25 air quality.

This modelling study has important policy-related implications. Uneven spatial distribution of the pollutants suggests prioritizing certain limited urban districts in policy scenarios. We show that focused efforts towards stove replacement in specific areas may have dominant positive effect on the air quality in the whole municipality. The case study identifies urban districts where limited incentives would result in the strongest reduction of the population's exposure to PM2.5.

30 **1 Introduction**

Residential wood combustion in households is a significant global air polluter. Even in developed countries, e.g., in Scandinavia, wood burning for residential heating is a dominant source of air pollution in cold winter days (Savolahti et al., 2019; Kukkonen et al., 2020). A low temperature wood-burning process emits a considerable amount of particulate matter of less than 2.5 micro-meter (μm) in size. Such particles are collectively abbreviated as PM_{2.5}. In fact, residential combustion is the largest PM_{2.5} emitter in the Nordic countries, including Norway (Im et al., 2019). Studies reveal that PM_{2.5} concentrations attributed to residential wood combustion in Oslo may reach 60% of the total annual average concentrations of this pollutant (Kukkonen et al., 2020). The attributed fraction of PM_{2.5} can be even larger in smaller cities (Hedberg et al., 2006). In the short run, air pollution during cold winter days can result in reduced visibility and pervasive smoky smell in the air. The highest levels of pollution are found in the most densely populated central urban districts where wood-burning is used in apartments of multi-story houses. The PM_{2.5} concentrations exceeding $40 \mu\text{g m}^{-3}$ can lead to an increased number of health issues in exposed populations. About 2000 annual premature mortality cases have been attributed to PM_{2.5} pollution in Norway alone (Im et al., 2019). Overall, non-industrial wood combustion in Europe contributes to about 10% of the total health cost of air pollution (Brandt et al., 2013). These numbers, as well as persistent public pressure and regulations from environmental protection agencies, are pushing many countries in the world, including Nordic municipalities, to introduce policy measures fostering reduction of residential emissions.

To be effective and efficient, policy measures must be supported by science. Atmospheric pollution caused by small-scale spatially distributed emitters is a non-trivial advection-diffusion problem that does not have a general solution. It is typically dealt with using numerical modelling. However, the most common Gaussian statistical models are unable to represent pollution aggravation in local stagnation zones. This problem is related to the fact that pollution dispersion models use a pure diffusion operator to calculate the concentrations along the pollution transport pathways. The worst air quality is typically observed in atmospheric conditions of weak turbulent mixing where horizontal advection dominates over lateral diffusion. In such conditions, pockets of highly polluted air can be transported over long distances, eventually exacerbating the air quality problems in remote stagnation zones downstream. It is important to emphasize that local emission in the stagnation zones may

55 be less significant, so that, those zones are not identified by statistical models as being vulnerable to air quality hazards. Turbulence-resolving atmospheric models have been found to be a proper and general tool to study passive scalar dispersion over complex terrain (Gousseau et al., 2015; Zhong et al., 2016). Such models simulate turbulence diffusion and pollution transport on the basis of air flow dynamics. Up to now, however, realistic studies with the turbulence-resolving models were limited as they require significant computing resources. This study is perhaps one of the first that assesses air quality and
60 supports policy development for the whole urban municipality with hydrodynamical simulations that resolve all significant buildings.

Air quality is controlled by near-surface turbulent mixing and local winds. We treat small smoke particles (PM_{2.5}) as passive scalars emitted to the atmosphere at actual locations of household chimneys. Each chimney creates high PM_{2.5} concentrations
65 only in its immediate vicinity. A statistical model adds up these local emissions making the total concentrations roughly proportional to the chimney density and the local emission rates (Zhong et al., 2016). With our turbulence-resolving model, we aim to understand contributions from the dynamical component of turbulent diffusion and transport in spatial variability of the observable PM_{2.5} concentrations. In realistic urban domains such as, e.g., in Bergen, Norway, concentrations are patchy and follow intricate pattern of local wind convergence and stagnation zones (Wolf et al., 2020). Such microscale climate
70 information must be considered in air quality impact studies (Chandler, 1976), (Bai, 2018). A challenge is however related to modelling of local winds and turbulence patterns. A turbulence-resolving model must be run with a realistic inventory of household emission sources and with correct topography in a sufficiently large domain. One might argue that a dense network of low-cost meteorological and air quality sensors can do the job when it is combined with statistical models (Schneider et al., 2017) or machine learning (Venter et al., 2020). That is likely to be true. Policy makers, however, would benefit from the
75 model-based approach as it provides quantitative evaluations of the impact of proposed measures, even in the case when those measures are changing configurations and intensities of emission sources. Ultimately, as we will demonstrate in this study, the models can help to identify urban areas vulnerable to air pollution, and to justify more optimal policies complying with targeted air quality levels. Locked within a silo (i.e., vertically integrated) approach to environmental management, Nordic municipalities are reluctant to explore advanced modelling approaches (Leiren and Jacobsen, 2018). In such circumstances, a

80 realistic case study is beneficial to demonstrate the added value of turbulence-resolving modelling for air quality management.
We chose a domain of the Bergen municipality for such a prototype demonstration study.

Actionable policy scenarios must reflect the complex physio-geographical context of urban areas. Hence, the models of atmospheric chemistry and physics must include effects of local topography and spatial distribution of population and pollution sources. A specific interest for research is related to long-distance (in some local sense) pollution transport where the strongest
85 air quality effects are observed far away from the source of the emission. Such effects are typical in valleys or in complex coastal areas with significant local circulations (Fernando et al., 2010). Numerical weather prediction models with their typical geographical resolution of 1 km or coarser cannot represent this complex environment (Baklanov et al., 2017a; Køltzow et al., 2019). In addition, large buildings, roads systems and other infrastructure objects modify winds and turbulent dispersion shading some places or aggravating pollution in others (Gousseau et al., 2015). A growing body of literature suggests that
90 turbulence-resolving or at least turbulence-permitting large-eddy simulation (LES) models are needed to deal with the urban pollution issues (Baklanov et al., 2017b; Brötz et al., 2014; Grimmond et al., 2020; Kurppa et al., 2018; Stoll et al., 2020).

Fine-scale LES is a computational fluid dynamics technique that becomes affordable with growing performance of massive-parallel computers. The LES models resolve the most energetic and dispersive turbulent eddies in the stratified lower atmosphere over realistically complex surfaces.

95 The LES models have already been used to simulate turbulent flows and atmospheric pollution in several smaller urban areas (e.g., Castillo et al., 2009; Cécé et al., 2016; Gronemeier et al., 2017; Keck et al., 2014; Letzel et al., 2008; Park et al., 2015b; Resler et al., 2017a). Our study further extends LES modeling of the air quality transport and diffusion problem to a whole municipality. We run the model with real surface boundary conditions, atmospheric meteorological conditions, and actual distribution and effect of emission sources. In this configuration, the modeling results could be used in policy scenario
100 evaluation and decision-making processes. We use the Parallelize Atmospheric Large-eddy simulation Model (PALM) described by Maronga et al. (Maronga et al., 2015, 2019b) with minor own modifications as described by Wolf et al. (Wolf-Grosse et al., 2017a; Wolf et al., 2020). Furthermore, we investigate a set of plausible mitigation policy scenarios, which were proposed to reduce emissions from the residential wood combustion sources – the household stoves. The central policy measure is to push for replacement of older stoves to less polluting new stoves. This push is given through soft economic incentives –

Field Code Changed

105 a limited cash refund of 5000 NOK per stove – and unconditional hard policy stipulating a ban on the use of the older stoves
by 2021. The policy was designed primarily fire safety considerations in mind, as well as improving the air quality. However,
its effect on air quality and people’s exposure to high PM2.5 concentrations have not been quantitatively evaluated before this
study.

The manuscript has the following structure. The next section describes the context, geography, and datasets of this study. The
110 third section describes the PALM model, setup of the simulations, and our methodology. The fourth section presents the
obtained results. The fifth section provides a broader discussion with generalizations of the methodology, data usage and
potentials for policy implications. The final section summarizes the conclusions.

2 The Bergen case study: Local context and datasets

Bergen is the second largest city in Norway. Its population is more than 275,000 people. It occupies over 465 km² divided into
115 8 districts. Lower parts of hills and coastal valleys within the Bergen municipality are built up. Central urban districts and
several neighbourhoods around large shopping malls are densely populated and include high-rise residential and administrative
buildings. The central urban districts have a population of more than 75,000. They occupy a narrow Bergen valley, which
opens on both sides towards large sea inlets (fjords). Topographic sheltering (Jonassen et al., 2013) by surrounding mountains
up to 643 meters height, together with maritime boreal climate at 60.4°N, favour persistent winter-time surface temperature
120 inversions capping cold air pools in the lowest parts of relief (Lareau et al., 2013). The temperature inversions, typically about
250 meters height, can last over several days. They are robustly associated with clear sky, calm and cold weather conditions
(Wolf et al., 2014).

As many Nordic cities, Bergen has clean air brought to the city with westerlies from the Atlantic Ocean. The main local
permanent PM2.5 sources are ships in the harbor and road traffic (Wolf et al., 2020). In winter months, however, households
125 actively use wood-burning stoves as a secondary heating source. The household stoves considerably contribute to the PM2.5
pollution in Bergen in November through April. Accurate inventory of the residential PM2.5 emission is nevertheless
problematic as we will discuss later in our study. Ample use of wood-burning frequently leads to high ambient concentration
of PM2.5 that in several urban districts exceeds a threshold of 40 µgm⁻³ set by the environmental protection agency (Høiskar

et al., 2017; Wolf et al., 2020). A large fraction of PM_{2.5} emission comes out from household wood-burning stoves installed
130 before 1998, hereafter referred to as older stoves. These older stoves have much higher emissions of particulate matter per unit
volume of wood; and they are less combustion and energy efficient than more modern “clean-burning” stoves, which we will
refer to as new stoves. There is a strong anti-correlation between air quality and air temperature in Bergen (Wolf and Esau,
2014). Low air temperatures are associated with calm weather periods when residential wood-burning is enhanced and the
emitted particles are trapped in the urban canopy due to weak vertical turbulent diffusion in the strongly, often inversely
135 stratified lower atmosphere (Wolf et al., 2014). Although turbulent diffusion is weak, local near-surface micro-circulations
(local winds) develop within this highly heterogeneous urban domain (Wolf-Grosse et al., 2017a). The local winds aggravate
air pollution in convergence and stagnation zones. Thus, a general advection-diffusion problem translates in the Bergen context
into the concrete modeling task of identification of urban districts with elevated PM_{2.5} concentrations given a realistic
distribution of household stoves, emission scenarios and weather conditions.

140 This study addresses the following questions. What is the spatial distribution of PM_{2.5} concentrations in the city during critical
air pollution episodes? What is the impact of emissions in each district on the (overall) air quality in the most polluted parts of
the city? What would be the effect of replacing the existing older stoves with only clean burning (new) stoves in the entire city
or in some of its districts? These research questions were formulated by scientists, whereas policy makers and stakeholders
were less open to explore different opportunities and stove replacement strategies. The study benefits from communication
145 with the Bergen municipality and other interested parties. Many Nordic cities are suffering from similar challenges, and this
is today an active field of research (Simpson et al., 2018).

3 The model and method

3.1 The large-eddy simulation model PALM

We simulate the lower atmosphere (up to 2.2 km) over the whole Bergen municipality with the LES model PALM. PALM
150 version 5.0 (revision 3063) is an atmospheric large-eddy simulation model developed by the PALM group at the Leibnitz
University of Hannover, Germany (Maronga et al., 2015). The model solves primitive hydro- and thermo-dynamic equations
for incompressible, Boussinesq fluids. PALM explicitly resolves a part of relevant three-dimensional atmospheric turbulence

dynamics as well as turbulence-flow interactions with complex surface geometry. ~~These~~ Such features give some advantages to the LES-based approach as compared to a more traditional meso-meteorological modelling in the urban areas. Our reader shall, however, observe that we do not claim accurate simulations of details of such interactions. We only expect that PALM is robust for the dominant scales of the relief (hundreds of meters in Bergen), which are well resolved by the model grid.

Our own user-code in PALM aims to improve treatment of the complex surface and environmental conditions. This user-code includes a routine to relax the mean vertical temperature profile in the computational domain towards a given input profile. In this way, the model preserves the temperature inversion that traps the emitted pollutants within the urban canopy. The user-code allows for ~~prescribe~~ mixed-type boundary conditions for the surface temperature; the von Neuman constant flux boundary conditions are set over land, the Dirichlet constant temperature conditions – over water. Constant temperature is set for the water surface grid cells, whereas a constant heat flux is set for the land surface grid cells. Our user-code also recalculates temperature and concentrations of pollutants in ~~the~~ terrain-following coordinates system from the model output ~~provided on~~ the rectangular grid ~~coordinates~~. For this, a linear extrapolation is conducted, if the specified height above the ground is not a multiple of the grid height at the given grid-point or simply the variable at the correct grid-height is given.

Pollutants are treated as passive scalars. Chemical reactions are not simulated. Household chimneys are not resolved in the model. To compensate for variable heights of the houses and their chimneys, the user-code allows for emission at any height above the surface. The emission height is then given as input to the model as an array of numbersecontaining a number for each horizontal grid point. The passive scalar is assumed being instantaneously mixed within the entire grid cell of emission. The lateral boundary conditions do not allow for recirculation and inflow of passive scalars into the model domain through any boundary. The three first methodologies are described in Wolf-Grosse et al. (2017) and Wolf et al. (2020) and the latter has been developed for this study.

3.2 The model setup for the Bergen case study

The model setup for the Bergen case study is discussed in details in Wolf et al. (Wolf et al., 2020), where it has been applied to study air pollution from a mixture of diverse (localized and spatially distributed) emission sources. Only a short summary and important differences are presented below. The computational domain includes Bergen and parts of the surrounding

municipalities (see Fig. 1). The total domain size is 28800 x 34560 m² in the zonal and meridional directions, respectively.

The simulations run with horizontally periodic lateral boundary conditions [for wind, pressure and temperature](#). Therefore, this domain includes buffer zones used for linear interpolation between the opposing periodic boundaries of 1000 m width. The horizontal grid resolution is 10 m. It gives a mesh of almost one million grid cells at each model level – the largest achieved urban simulations so far. The vertical grid resolution is 10 m up to 660 m height. Above that the vertical grid space increases by 1% for each additional grid level. The total domain height is 160 levels or 2239 m. Thus, the model domain top is found well above the highest hill (640 m) in the area of simulations. The surface in the model is approximated with cubes of 10 x 10 x 10 m³. The topographic data for the approximation are taken from a laser-scan digital elevation model (DEM) of 1 m resolution provided by the Norwegian mapping authority (Statens Kartverk, 2018). At this resolution, DEM includes all buildings and trees in the city. DEM was delivered to us in the GeoTIFF format. We use the ArcGIS © software to process the data and to create a complete topography dataset of the required 10 m horizontal resolution. We fill in small gaps in the original DEM using the standard linear interpolation.

Water bodies have a distinct impact on boundary layer circulations, also in urban areas (Ronda et al., 2017; Wolf-Grosse et al., 2017a). This impact is accounted for in simulations by prescribing a constant negative surface heat flux of -20 W m⁻² over [the land surface grid cells](#). Constant surface temperature of 275.65 K (equal to 2.5° C) is applied over [the sea water surface grid cells](#); temperature of 273.15 K (equal to 0° C) is applied over [the freshwater bodies \(lakes\)](#). This setup reflects conditions of winter temperature inversions with radiative surface cooling and warmer sea surface temperatures as compared to the land and lake surface temperatures.

3.3 Weather scenarios

Damaging levels of air pollution are almost exclusively observed during persistent calm and cold weather episodes. We have already identified typical meteorological conditions that correspond to the pollution episodes (Wolf et al., 2014). Analysis of such conditions produced two influential meteorological scenarios, i.e., two sets of [vertical profiles: one for temperature \(from the meteorological temperature profiler in Bergen\)](#) and [another for wind \(from the ERA-Interim reanalysis\) profiles](#) that correspond to the observed high concentrations of PM2.5. The first scenario is for the most typical winter conditions with high air pollution. This scenario is determined by the south-easterly geostrophic wind, which is used to force the model simulations,

and by the fjord surface water temperature of 2.5° C, which is used as the surface boundary condition in the model. In this scenario, the local winds transport polluted air from land to the city fjord through the densely populated central urban valley.

205 The second meteorological scenario has: the easterly geostrophic wind of slightly higher speed; the fjord surface water temperature of 0° C, which is still above the freezing point for salt water. Both scenarios are described in details in ~~Wolf et al. (2019)~~ (Wolf et al., 2020) under the abbreviations ws01_wd01_ft01 and ws03_wd02_ft03, respectively. Model forcing and surface temperatures are fixed in both scenarios for the complete duration of the model runs. The runs are initialised with 12-hours long precursor runs. At this point, most meteorological parameters do not show any significant drift. Simulations with different configurations of emission sources are then conducted following the precursor runs for another 6 hours.

210 Positive turbulent sensible heat fluxes over water and negative fluxes over land are not in balance and do not cancel each other within the model domain. This imbalance, if not treated, would erode the stable stratification of the atmosphere in the upper layers of the model domain. We resolve the problem relaxing the mean temperature profile to the initially prescribed profile using nudging. Nudging corrects the temperature at every grid point and time step by a weighted difference between the actual domain-averaged temperature and that at the model initialisation. The weight is given by the relaxation time scale set to 43200 s at elevations below 400 m, which effectively gives no nudging in the lower layers of the model domain. The relaxation time scale linearly decreases till 1800 s at 600 m and higher elevations. Nudging is not applied at the first grid point above the surface.

We run PALM with real spatial configuration of small-scale emission sources in the case study domain that is provided to us
220 by ~~stove~~ the Bergen Fire Brigade (Bergen brannvesen) ~~stove~~. Overall, there were 80 506 household stoves registered in Bergen in June 2018. The new clean burning stoves constitute 45.8% or 36 864 stoves. Most of them were installed in the recent years. Since 1995, ~~installation~~ only ~~the~~ new stoves are allowed. We have aggregated and averaged the number of stoves within 3 x 3 grid cells plots (90 m²) to emulate emission dispersion immediately after exhaust from chimneys. Figure 1 (bottom) shows the geographical distribution of the resulting emission source density. Concentrations are set to zero at starting both the scenario
225 and sensitivity runs. In this study, we apply the same fixed emission height of 15 m (the 2nd model level) above the ground everywhere in the model domain as it corresponds to the most typical building height in Bergen. Plausible effects of different emission heights are considered in a sensitivity study that is included in the supplementary material.

The Bergen Fire Brigade has also provided us with stipulated emission rates per stove. The new clean wood-burning stoves have the relative emission rate of 7 g kg⁻¹. This number shall be understood as 7 g of PM2.5 per kg of burned wood. The older stoves have the relative emission rate of 30 g kg⁻¹. With the given mix of the new and older stoves, the average relative emission rate is 19.42 g kg⁻¹. To obtain the actual emission rates, one must know consumption of wood per unit time per stove and frequency of using the stoves, i.e., parameters that characterize burning habits in the city. Felius et al. (Felius et al., 2019) published an updated study of the wood-burning habits in Norway. It reveals that about 80% of households use stoves only during the afternoon and evening hours, while 90% of households still use wood-burning as a significant (primary and secondary) heating source. The typical average usage time is 4 to 10 hours. Shorter usage time (2.5 to 3.5 hours) is reported by Wyss et al. (Wyss et al., 2016) in a study of relations between indoor PM2.5 concentrations and wood-burning habits in Norwegian households. Emission rates also depend on the wood quality, wetness, and tree species in the wood supply (Hellén et al., 2008). These numbers are not available so that we estimated them using a small sampling study that is based on our social networks. We assume that each stove is used to burn approximately 11 kg wood per day, and that the wood consumption is 1.25 kg per hour (Solli et al., 2009), we arrived to 8.8 hours of wood burning, primarily between 15:00 and 24:00 local time; 90% of all aerosol exhaust comes out as PM2.5. Taken together, these assumptions and estimations result in the average PM2.5 emission rate of 0.7647 g s⁻¹ per stove. This number is about 10 times smaller than it has been assumed earlier by the Bergen Fire Brigade, but simulations with this emission rate shows good ~~correspondence-agreement between~~ computed PM2.5 concentrations ~~and to~~ the concentrations measured in the city. During the second week of February 2021, a persistent cold weather situation similar to our first scenario was observed in Bergen. The maximum simulated (observed on 11.02.2021) PM2.5 concentrations were 76.7 µg m⁻³ (81.2 µg m⁻³) at Danmarks plass, 53.4 µg m⁻³ (59.2 µg m⁻³) at Klosterhaugen in the city center, and 26.1 µg m⁻³ (18.6 µg m⁻³) at Rådalen. This agreement demonstrates reasonably good capture of the spatial variability and accumulation of PM2.5 in the scenario simulations despite the accepted simplifications, assumptions, and uncertainties.

Formatted: Superscript

4 Results

Air pollution in urban areas is patchy. As one might expect, simulations reveal higher concentrations of PM2.5 in more densely populated districts. This pattern is however significantly distorted by intricate interplay of local air circulations and turbulent

diffusion in the lower parts of the atmosphere. Although there is an overall near-surface air flow towards the fjord, pollution tends to concentrate in some stagnation zones on the way. Thus, the study allows for certain generalizations. Our results reveal that the highest PM_{2.5} concentrations are created by long-distance horizontal air advection into stagnation zones. Gradual
255 accumulation and convergence of the polluted air is more important than the local dispersion over emission sources. A considerable fraction of pollution in the stagnation zones is advected from upstream urban districts. Figure 2 shows the near-surface wind- and temperature fields in the actual boundaries of the Bergen municipality for the two pre-defined weather scenarios. This area is sufficiently distant from the model boundary zone, which is shown as a grey rectangle in Figure 1 (top panel).

260 Our simulations represent two weather scenarios. Scenario 1 has higher water temperature (Figure 2; the top panel). It represents the conditions with strong water-land surface temperature difference. Warmer water forces stronger wind convergence over the fjord and increases the outflow from the central valley. In some more general sense, it can be seen as a scenario with effective urban ventilation with local breeze. The increased wind convergence is also observed over smaller or medium-size water bodies. We found that these surface-layer breeze-like circulations, which are driven by the horizontal
265 temperature differences, are to the large degree autonomous, existing independently and in many places in opposition to the upper air winds. Scenario 2 has lower water temperature (Figure 2; the bottom panel). Weaker convergence of the near-surface winds produces different configuration of the local air flows, and therefore, different pollution pathways. This scenario describe a situation when local surface heterogeneity is not strong enough or not well organized to increase urban ventilation. Its most pronounced effect is in the formation of air stagnation in the central urban districts. Stronger upper layer winds do not
270 break this stagnation. Thus, this stagnation zone is robust, which is an important result from the citizens and stakeholders' point of view.

A combination of uneven distribution of emission sources with an intricate pattern of local air circulation ~~s-patterns~~ produces a patched picture of the pollution concentrations. ~~The goal of this study is to evaluate population exposure to the high concentrations of PM_{2.5}.~~ Our simulations identify the pathways of pollution in the surface layer. In both scenarios, the high
275 concentrations are found not only in the areas with the highest emission rates but also in other areas where local winds advect the pollutants and concentrate them in converge zones. Figure 3 presents the map of the PM_{2.5} concentrations near the surface.

It is simulated using the current distribution of new and older stoves. Both weather scenarios reveal high concentrations in cold air pools designating the effect and importance of topographic sheltering.

4.1 Implications of pollution advection across municipal districts

280 Now, we shall look at the role of long-distance pollution advection and convergence ~~in setting the PM2.5 concentrations in~~ different municipal districts. To do this, we apply a methodology of artificial pollutants prescribing emission of an independent pollution specimen at each district of choice. ~~Thus, pollution from each district remains independent and traceable over the~~ whole domain. In this way, our simulations will reveal the pollution pathways and relative contributions for each area of interest. It is perhaps not surprising that the highest concentrations are found in the central, densest urban districts as, e.g., just
285 south and upward in central Bergen valley. More detailed analysis, however, suggests that a considerable fraction of the concentration shall be attributed to advection of polluted air from other upstream districts. Bergen has eight administrative districts. Figure 4 shows the simulated near-surface concentration obtained for emission sources (stoves) that are active only in the central district (Bergenuhus) and in the most populated district (Årstad). We plot both the absolute and relative concentrations. The high concentrations in Årstad are determined by the stagnant cold air pool in a local topographical shelter
290 and the high concentration of small apartments in old houses, each having individual wood-burning stove. In addition, the local wind convergence adds to increase the concentration even in the most polluted districts. Here, 80% to 90% of the total concentrations are of the local origin within the district itself. The non-local advection adds up to 60% to the concentrations in the most central district of Bergenuhus (see Fig. 4, top panel). This district is built-up by modern high-rise buildings where stoves are not in use. Nevertheless, the PM2.5 concentrations are relatively high there. The pollution in this district is advected
295 from the upstream Årstad district following the circulation pathway along the northern slope of the Bergen valley. By contrast, emissions within the city centre have almost no impact on other districts as the area is located down-stream in the Bergen valley and next to the waterfront (Byfjorden). Scenarios 1 and 2 are very similar to each other with respect to the pollution transboundary transport and non-local contribution of district's emission sources. Scenario 2 leads to lower local contribution (not shown), while the effects on the other areas remain similar. Since we want to identify the conditions with severe impact
300 on the population, we will focus on details of the scenario 1 simulations.

Looking at the three districts located further south in the municipality, we clearly see the effect of the down-stream and -valley transport towards the city centre. The main polluter, both locally (80%) and downstream (up to 60%), is the Årstad district. The pollution trapped in the surface layer are however transported over long distances within the municipality. According to our simulations, the two southernmost districts (Fana, Ytrebygda) add up to 20% to the concentrations in the city centre, while standing for near 100% of their own local pollution with their districts. The picture is very different when western (Fyllingsdalen, Laksevåg) and eastern (Åsane, Arna) urban districts are considered. Those districts are located behind mountains and water areas looking from the central Bergen perspective. Due to this, surface layer pollution will not be transported to the central valley, thus having no impact on the concentration of air pollutants. The local concentrations in these peripheral districts are less significant, due to different local topographic steering of the near surface air masses.

4.2 Assessment of mitigation actions to reduce emissions from residential wood combustion

The most environmentally friendly clean air mitigation action is to place bans on all emissions from private wood stoves. However, the need for heating during cold winter days implies that more realistic policy actions must be considered. Moreover, active use of the wood-burning stoves could be an effective retrofitting measure helping to save electricity and fossil fuels (Felius et al., 2019). Therefore, a massive transition to new stoves is deemed necessary both from political and environmental points of view. The results of our simulations impose two important constraints on plausible retrofitting measures. First, we argue and demonstrate it with model runs that there is a tight connection between weather conditions and exceedance of the concentration thresholds in particular districts. We found that PM2.5 readings in one district, say Åsane, may tell little about the concentrations in another district or even over smaller neighborhoods. So, the measures need to be constrained by geographical boundaries, which in turn could be determined only through high resolution atmospheric simulations (or analysis of dense observations if they are available). Second, if the target is not to cross a certain concentration threshold, the measures must account for the developing local atmospheric circulations, and not so much for dispersion of the very local emissions.

The presented simulations and many additional sensitivity runs suggest that unavoidable inaccuracies and uncertainties of emissions should not be overdriven in mitigation measures. Indeed, there are many uncertainties associated with the current

usage and replacement of the stoves, which includes uncertainties related to the emission factor (Seljeskog et al., 2017). ~~Despite the many uncertainties our simulations support geographically differentiated mitigation actions. However, whatever high or low the specific emission factor is, the spatial redistribution and aggravation of pollutants is robustly simulated with stagnation (high concentration) zones found over the identified areas. In particular, the convergence zone related to the breeze inflow and~~
330 ~~land outflow is of general interest as it might be found in many coastal cities. Hence, limiting~~Limiting economic incentives for stove replacement to just certain districts ~~could be~~~~may be~~ a cost-effective measure. We however acknowledge that other, e.g., political, factors may render such an efficiency sub-optimal. In our model simulations, we can alter the configurations and rates of emission sources in specific districts to estimate effects of different mitigation actions on the local and total level of urban pollution.

335 Plausible alternatives for the total and district-wise replacements of stoves with cleaner burning technology are presented in Figure 6. We change the current distribution of the older and new stoves in two clusters of four districts each. First, we replaced all stoves in the Bergen valley (the Bergenhus, Årstad, Fana and Ytrebygda districts) with the new stoves only. The simulations result in ~~a~~ strongly improved air quality with the highest concentrations being almost eliminated. Concurrently, the moderate and low concentrations are not markedly lower determining the baseline level of pollution from the new stoves – all stoves
340 will pollute. Then, we replaced ~~the~~ stoves in other districts (Laksevåg, Fyllingsdalen, Åsane and Arna). It results in ~~to~~ further reduction of concentrations in the total accumulated concentrations of over the entire municipality.

We summarize the effect of mitigation actions in Figure 7 and Table 1. The national regulators set the concentration threshold of $40 \mu\text{g m}^{-3}$ for PM_{2.5} as the delimiter of high and low air pollution. The simulations reveal that majority (respectively 17818
345 and 17243) of the exposed households are exposed only to low air pollution in both scenarios. Nevertheless, a total of 4031 (scenario 1) and 5986 (scenario 2) households are still exposed to high air pollution under the current distribution of stoves. Switching to the new stoves in the entire city results in elimination of high air pollution exposure. We discover that a very similar effect could be obtained with more modest but better targeted efforts focussed replacements of stoves on the central Bergen valley (Bergenhus, Årstad, Fana and Ytrebygda districts). Only 2 (scenario 1) and 44 (scenario 2) households are still
350 exposed to high air pollution in the simulations. By contrast, 3848 and 5636 households respectively remain exposed when the

stoves are installed only in peripheral districts (Laksevåg, Fyllingsdalen, Åsane and Arna). That corresponds to the relative exposure reduction by 5% and 6% of households only. At the same time, the whole 95% of households will continue to be exposed to high air pollution of such policy action.

5 Discussion and conclusions

355

Despite the frequent depiction of residential wood burning as clean energy, urban air data clearly associate wood combustion with elevated concentrations of the particulate matter (PM2.5) in cities. These particles are extremely harmful to humans because their small size allows them to reach the lungs and to deliver toxic contaminants they adsorbed. The World Health Organization attributed 61000 of the premature deaths in 2010 alone in Europe to outdoor PM2.5 pollution from residential wood combustion; whereas in Denmark, estimations suggested that up to 30-40% to the premature deaths should be attributed to this type of air pollution (Cincinelli et al., 2019). Residential wood combustion has its immanent importance for air quality not only in the Nordic cities where it contributes up to 82% of the total particulate matter in Northern Sweden (Krecl et al., 2008), but also elsewhere in Europe (up to 60% in Southern Germany (Bari et al., 2009)). Even in sub-tropical cities, e.g., Santiago, Chili (Mazzeo et al., 2018), impact of residential wood combustion could be critical for urban health, so that its modelling is in high demand. Due to these factors, there is a persistent interest and concentrated efforts to include the particulate matter into an urban integrated system (UIS) modelling approach (Grimmond et al., 2020). The UIS approach combines meteorological and air quality modelling with socio-economic and policy scenarios. It has been however recognized that integration of high-resolution models in existing systems still requires further development. In this study, we described a perspective UIS approach that has emerged from our co-production efforts with local policy makers.

365

370

PALM has been proposed as a perspective model that would allow resolving urban dispersion from highly localized pollution sources (Esau et al., 2021; Heinze et al., 2017; Maronga et al., 2019b; Zhang et al., 2020). PALM simulations were evaluated in several case studies but so far, this study is the first model application for policy and meteorological scenario development at the level of the whole administrative unit – an urban municipality. In this sense, our study sets up methodology and present perspective ways to utilize the large-eddy simulation model in the UIS. This modeling is advantageous for population-weighted

Formatted: Danish

Field Code Changed

Formatted: Danish

Formatted: English (United States)

Formatted: English (United States)

375 ~~exposure estimates that are still missing the high-resolution meteorological component (Savolahti et al., 2019). We demonstrate what such a modeling component could contribute for such expose estimates (Figure 7).~~

Formatted: English (United States)

~~The UIS studies benefit from co-production with local policy actors. Our case study in Bergen was co-designed with city managers to select the most relevant scenarios and targets. Concluding our case study that prototypes the modelling support~~

380 ~~for science-policy interactions, we shall note that the The Bergen city council in Bergen~~ has banned firing of the older stoves since January, 1st, 2021. To facilitate transition to the new stoves, financial incentives of 5000 Norwegian kroner (about 500

€) per stove have been allocated. The total allocated amount is 50 million Norwegian kroner or just 25% or the amount needed to support the total replacement of the older stoves in the municipality. So, it is not surprizing that by 2021, Bergen still has

385 more than 30 000 older stoves. Our simulations however suggest that targeted incentives might be sufficient to cover the stove replacement in the most critical urban districts. We show that such a selective replacement could have a major effect on

pollution reduction below the threshold of 40 $\mu\text{g m}^{-3}$ of PM2.5 ~~concentrations~~ for almost all households in Bergen municipality. Thus, we conclude that the turbulence-resolving atmospheric modelling has a potential to contribute to the design of more

efficient policy ~~actions~~ and scenarios than those presently applied. Analysis of simulation scenarios may not only increase the effects of limited incentives, which is desired per se, but can also assure that “no one is left behind” or in other words, that all

390 households will experience positive effects of the proposed policy ~~measures-actions~~; no district ~~would~~ remain unnecessarily polluted or even with worse air quality.

~~There are available other modelling approaches and more comprehensive studies of residential wood combustion, notably, applications of the Norwegian modelling system based on the MetVed emission model, the EPISODES air quality model and~~

395 ~~AROME weather model (Grythe et al., 2019; Grythe and Lopez-Aparicio, 2020). Both the data graining (250 m) and atmospheric model resolution (1000 m) in this modelling system are much coarser than in the present study (30 m and 10 m~~

~~correspondingly). Kukkonen et al. (Kukkonen et al., 2018) presented PM2.5 simulations in Helsinki, Finland, at higher resolution (100 m) with the SILAM modelling system. This system relies on the Gaussian (down-gradient) approach and hence~~

~~cannot resolve aggregation of polluted air. Emission inventory analysis and modeling results for different systems were presented in a study of four (Oslo, Helsinki, Copenhagen and Umeå) cities (Kukkonen et al., 2020). It considers the mean~~

400 ~~concentrations and the city-scale level and does not discuss relevant meteorological conditions. Our study is less model and site specific. Its focus is on explicit modelling of the turbulent dispersion and influential meteorological scenarios. Our simulations represent moderate to strong impact of the atmospheric stability on the concentration exceedance. The leading effect of atmospheric stability have been found in other our studies (Wolf et al., 2014, 2020) as well as by other authors concerned with the urban wood combustion (Grange et al., 2013). We show that this resolution and methodological differences~~
405 ~~do matter. Although the overall effect of the stove replacement programmes on the emission reduction in the city could be moderate, the replacement in certain sensitive districts helps to dramatically reduce the population expose. In this sense, our study supports the conclusions of Lopez-Aparicio and Grythe study (Lopez-Aparicio and Grythe, 2020) in Oslo, Norway, may help better understand physical reasons responsible for low effectiveness of subsidies and administrative actions set to reduce the wood burning emissions. One Norwegian study (Lopez Aparicio and Grythe, 2020) that the current indiscriminate found~~
410 ~~no evidence that municipalities with subsidies reduce emissions faster than those without subsidies. That study concluded that the subsidy programmes are ineffective. Our study however suggests that such subsidies may have the intended effect on emission reduction but a much more modest effect on the concentration and exposure reduction as subsidies are spent in less affected areas.~~

Concluding the modelling study of urban pollution dispersion, we observe that local winds within the turbulent boundary layer
415 can transport polluted air over long distances in the urban areas. Convergence and accumulation of pollution from distributed small-scale emission sources, such as the considered household stoves, deteriorate air quality in stagnation zones where the local emissions are not sufficient to maintain such high concentrations. We simulated only the worst-case scenarios for wintertime air pollution, which are ~~that~~ associated with the coldest weather and ~~possible~~ most frequent use of wood stoves for heating. ~~Therefore, our concentrations considerably exceed the mean values given by other studies.~~ (Grythe et al., 2019;
420 Kukkonen et al., 2020). Other weather regimes ~~can also result in produce~~ less severe air pollution which, ~~might~~may however contribute ~~significantly~~ over the whole heating season. The weather conditions favourable for air quality deterioration in Bergen are considered in other our works (Wolf et al., 2020; Wolf and Esau, 2014). Two studies by Wolf et al. (Wolf-Grosse et al., 2017b; Wolf et al., 2014) specifically consider co-variability between the temperature inversions and weather regimes

in Bergen as the main control factors of the local atmospheric pollution. Emission scenario testing like the one highlighted
425 here can be an important step of such a reasoning.

The ~~applied~~ methodology of this study is experimental. ~~It and~~ requires further development ~~and integration with other physical
430 and chemical components of the urban environment. Such efforts are on the way~~ (Maronga et al., 2019b). More specifically,
the quality of the PALM simulations needs assessment against in situ observations of concentrations as well as against more
constraining laboratory and other model results (Gronemeier et al., 2017, 2021). The input data sets ought to be more
constrained too (Heldens et al., 2020; Masson et al., 2020). Uncertainty in emissions and surface boundary conditions currently
impede fine tuning of the simulations. Societal uncertainties could be also influential (Feliuss et al., 2019; Wyss et al., 2016).
Heating patterns and habits are poorly constrained. Concerns that the new stoves will be used more, and therefore retard the
effect on air pollution, have been also raised ~~by stakeholders~~. A broader study of societal effects is out of scope of this work.
Finally, the PALM model is under extensive development now to allow for more realistic and comprehensive urban
435 meteorological studies. It will include a block for improved surface boundary conditions (Maronga et al., 2019a, 2019b; Resler
et al., 2017) and connection with the regional model COSMO (Fröhlich and Matzarakis, 2020).

Code availability

The model code PALM is available from the PALM group from <https://palm.muk.uni-hannover.de/trac/wiki/doc/install>. The
model is free to download and use upon registration. The scripts for data analysis and visualization are written in MATLAB
440 and available from the Nansen Center upon request to the corresponding author.

Data availability

Availability conditions for the collection of data sets to setup and run the model and weather scenarios are described in Wolf
et al. (2020). Data from new model simulations (a large data files) for this study are available from the Nansen Center upon
request to the corresponding author.

445 **Author contributions**

Tobias Wolf was responsible for the data acquisition, setting and running the model PALM, data analysis scripts and visualization routines. He contributed to writing the manuscript and interpretation of the modelling results. Lasse Pettersson was responsible for contacts with data providers and end-users, design of scenarios, interpretation, and communication of the results. He contributed to writing the manuscript. Igor Esau was responsible to the conceptual design of the study, methodology
450 of numerical simulations and data analysis. He wrote the manuscript and connected the results of the study with the need of end-users.

Competing interests

The authors declare that they have no conflict of interests.

Special issue statement

455 This study is a contribution to a special issue “Pan-Eurasian Experiment (PEEX) – Part II” of the Atmospheric Chemistry and Physics journal.

Acknowledgements

This study is performed within the GC Rieber Climate Research Institute at the Nansen Center. Our studies of air pollution in the Bergan have capitalized on stakeholder and user perspectives from and cooperation with Ulrik Jørgensen, Sverre Østvold,
460 Nils Møllerup and Even Husby – Port of Bergen, and Eva Britt Isager and Per Vikse – Climate section, Mette Iversen and Nils-Eino Langhelle – Section for Plan and Geodata, Per Hallstein Fauske and Arve Bang, Health Care Agency, all Bergen municipality.

Financial support.

465 This research has been supported by the GC Rieber Foundations, Bergen, and strategic institute funding (RCN grant no. 218857).

The finalization of data analysis and publication were supported by the Norway Grant TO01000219 (TURBAN). The project "TURBAN - Turbulent-resolving urban modeling of air quality and thermal comfort" benefits from a research grant from Norway and Technology Agency Czech Republic (TA CR).

470 ~~The finalization of data analysis and publication were supported by the EEA grant TO01000219 (TURBAN).~~

References

- Bai, X.: Advance the ecosystem approach in cities, *Nature*, 559(7712), 7, doi:10.1038/d41586-018-05607-x, 2018.
- Baklanov, A., Korsholm, U. S., Nuterman, R., Mahura, A., Nielsen, K. P., Sass, B. H., Rasmussen, A., Zaakey, A., Kaas, E.,
475 Kurganskiy, A., Sørensen, B. and González-Aparicio, I.: Enviro-HIRLAM online integrated meteorology–chemistry
modelling system: strategy, methodology, developments and applications (v7.2), *Geosci. Model Dev.*, 10, 2971–2999,
doi:10.5194/gmd-10-2971-2017, 2017a.
- Baklanov, A., Brunner, D., Carmichael, G., Flemming, J., Freitas, S., Gauss, M., Hov, Ø., Mathur, R., Schlünzen, K. H.,
Seigneur, C. and Vogel, B.: Key Issues for Seamless Integrated Chemistry–Meteorology Modeling, *Bull. Am. Meteorol. Soc.*,
480 98(11), 2285–2292, doi:10.1175/BAMS-D-15-00166.1, 2017b.
- Bari, M. A., Baumbach, G., Kuch, B. and Scheffknecht, G.: Wood smoke as a source of particle-phase organic compounds in
residential areas, *Atmos. Environ.*, 43(31), 4722–4732, doi:10.1016/j.atmosenv.2008.09.006, 2009.
- Brandt, J., Silver, J. D., Christensen, J. H., Andersen, M. S., Bønløkke, J. H., Sigsgaard, T., Geels, C., Gross, A., Hansen, A.
B., Hansen, K. M., Hedegaard, G. B., Kaas, E. and Frohn, L. M.: Contribution from the ten major emission sectors in Europe
485 and Denmark to the health-cost externalities of air pollution using the EVA model system – an integrated modelling approach,
Atmos. Chem. Phys., 13(15), 7725–7746, doi:10.5194/acp-13-7725-2013, 2013.
- Brötz, B., Eigenmann, R., Dörnbrack, A., Foken, T. and Wirth, V.: Early-Morning Flow Transition in a Valley in Low-
Mountain Terrain Under Clear-Sky Conditions, *Boundary-Layer Meteorol.*, 152(1), 45–63, doi:10.1007/s10546-014-9921-7,
2014.
- 490 Castillo, M. C. L., Kanda, M. and Letzel, M. O.: Heat ventilation efficiency of urban surfaces using large-eddy simulation,
Annu. J. Hydraul. Eng., 53, 175–180, 2009.
- Cécé, R., Bernard, D., Brioude, J. and Zahibo, N.: Microscale anthropogenic pollution modelling in a small tropical island
during weak trade winds: Lagrangian particle dispersion simulations using real nested {LES} meteorological fields, *Atmos.*
Environ., 139, 98–112, doi:10.1016/j.atmosenv.2016.05.028, 2016.
- 495 Chandler, T. J.: *Urban Climatology and its Relevance to Urban Design.*, 1976.
- Cincinelli, A., Guerranti, C., Martellini, T. and Scodellini, R.: Residential wood combustion and its impact on urban air quality
in Europe, *Curr. Opin. Environ. Sci. Heal.*, 8, 10–14, doi:10.1016/j.coesh.2018.12.007, 2019.

- Esau, I., Bobylev, L., Donchenko, V., Gnatiuk, N., Lappalainen, H. K., Konstantinov, P., Kulmala, M., Mahura, A., Makkonen, R., Manvelova, A., Miles, V., Petäjä, T., Poutanen, P., Fedorov, R., Varentsov, M., Wolf, T., Zilitinkevich, S. and Baklanov, A.: An enhanced integrated approach to knowledgeable high-resolution environmental quality assessment, *Environ. Sci. Policy*, 122(February), 1–13, doi:10.1016/j.envsci.2021.03.020, 2021.
- Felius, L. C., Thalfeldt, M., Georges, L., Hrynyszyn, B. D., Dessen, F. and Hamdy, M.: Wood burning habits and its effect on the electrical energy demand of a retrofitted Norwegian detached house, *IOP Conf. Ser. Earth Environ. Sci.*, 352(1), doi:10.1088/1755-1315/352/1/012022, 2019.
- 505 Fernando, H. J. S. S., Zajic, D., Di Sabatino, S., Dimitrova, R., Hedquist, B. and Dallman, A.: Flow, turbulence, and pollutant dispersion in urban atmospheres, *Phys. Fluids*, 22(5), 051301, doi:10.1063/1.3407662, 2010.
- Fröhlich, D. and Matzarakis, A.: Calculating human thermal comfort and thermal stress in the PALM model system 6.0, *Geosci. Model Dev.*, 13(7), 3055–3065, doi:10.5194/gmd-13-3055-2020, 2020.
- Gousseau, P., Blocken, B., Stathopoulos, T. and van Heijst, G. J. F.: Near-field pollutant dispersion in an actual urban area: Analysis of the mass transport mechanism by high-resolution Large Eddy Simulations, *Comput. Fluids*, 114, 151–162, doi:10.1016/j.compfluid.2015.02.018, 2015.
- 510 Grange, S. K., Salmond, J. A., Trompeter, W. J., Davy, P. K. and Ancelet, T.: Effect of atmospheric stability on the impact of domestic wood combustion to air quality of a small urban township in winter, *Atmos. Environ.*, 70, 28–38, doi:10.1016/j.atmosenv.2012.12.047, 2013.
- 515 Grimmond, S., Bouchet, V., Molina, L. T., Baklanov, A., Tan, J., Schlünzen, K. H., Mills, G., Golding, B., Masson, V., Ren, C., Voogt, J., Miao, S., Lean, H., Heusinkveld, B., Hovespyan, A., Teruggi, G., Parrish, P. and Joe, P.: Integrated urban hydrometeorological, climate and environmental services: Concept, methodology and key messages, *Urban Clim.*, 33(May 2019), 100623, doi:10.1016/j.uclim.2020.100623, 2020.
- Gronemeier, T., Raasch, S. and Ng, E.: Effects of Unstable Stratification on Ventilation in Hong Kong, *Atmosphere*, 8, 168, doi:10.3390/atmos8090168, 2017.
- 520 Gronemeier, T., Sum, K., Harms, F., Leitl, B., Maronga, B. and Raasch, S.: Evaluation of the dynamic core of the PALM model system 6.0 in a neutrally stratified urban environment: comparison between LES and wind-tunnel experiments, *Geosci. Model Dev.*, 14(6), 3317–3333, doi:10.5194/gmd-14-3317-2021, 2021.
- Grythe, H. and Lopez-Aparicio, S.: *MetVed v2.0 model and application*, Oslo, Norway., 2020.
- 525 Grythe, H., Lopez-Aparicio, S., Vogt, M., Vo Thanh, D., Hak, C., Karine Halse, A., Hamer, P. and Santos, G. S.: The MetVed model: development and evaluation of emissions from residential wood combustion at high spatio-temporal resolution in Norway, *Atmos. Chem. Phys.*, 19(15), 10217–10237, doi:10.5194/acp-19-10217-2019, 2019.
- Hedberg, E., Johansson, C., Johansson, L., Swietlicki, E. and Brorström-Lundén, E.: Is Levoglucosan a Suitable Quantitative Tracer for Wood Burning? Comparison with Receptor Modeling on Trace Elements in Lycksele, Sweden, *J. Air Waste Manage. Assoc.*, 56(12), 1669–1678, doi:10.1080/10473289.2006.10464572, 2006.
- 530 Heinze, R., Moseley, C., Böske, L. N., Muppa, S. K., Maurer, V., Raasch, S. and Stevens, B.: Evaluation of large-eddy

- simulations forced with mesoscale model output for a multi-week period during a measurement campaign, *Atmos. Chem. Phys.*, 17(11), 7083–7109, doi:10.5194/acp-17-7083-2017, 2017.
- 535 Heldens, W., Burmeister, C., Kanani-Sühring, F., Maronga, B., Pavlik, D., Sühring, M., Zeidler, J. and Esch, T.: Geospatial input data for the PALM model system 6.0: Model requirements, data sources and processing, *Geosci. Model Dev.*, 13(11), 5833–5873, doi:10.5194/gmd-13-5833-2020, 2020.
- Hellén, H., Hakola, H., Haaparanta, S., Pietarila, H. and Kauhaniemi, M.: Influence of residential wood combustion on local air quality, *Sci. Total Environ.*, 393(2–3), 283–290, doi:10.1016/j.scitotenv.2008.01.019, 2008.
- Høiskar, B. A. K., Sundvor, I., Johnsrud, M., Haug, T. W. and Nr, N. P.: Tiltaksutredning for lokal luftkvalitet i Bergen., 2017.
- 540 Im, U., Christensen, J. H., Nielsen, O.-K., Sand, M., Makkonen, R., Geels, C., Anderson, C., Kukkonen, J., Lopez-Aparicio, S. and Brandt, J.: Contributions of Nordic anthropogenic emissions on air pollution and premature mortality over the Nordic region and the Arctic, *Atmos. Chem. Phys.*, 19(20), 12975–12992, doi:10.5194/acp-19-12975-2019, 2019.
- Jonassen, M. O., Ólafsson, H., Valved, A. S., Reuder, J. and Olseth, J. A.: Simulations of the Bergen orographic wind shelter, *Tellus A Dyn. Meteorol. Oceanogr.*, 65(1), 19206, doi:10.3402/tellusa.v65i0.19206, 2013.
- 545 Keck, M., Raasch, S., Letzel, M. O. and Ng, E.: First Results of High Resolution Large-Eddy Simulations of the Atmospheric Boundary Layer, *J. Heat Isl. Inst. Int.*, 9, 39–43, 2014.
- Køltzow, M., Casati, B., Bazile, E., Haiden, T. and Valkonen, T.: An NWP Model Intercomparison of Surface Weather Parameters in the European Arctic during the Year of Polar Prediction Special Observing Period Northern Hemisphere 1, *Weather Forecast.*, 34(4), 959–983, doi:10.1175/WAF-D-19-0003.1, 2019.
- 550 Krecl, P., Hedberg Larsson, E., Ström, J. and Johansson, C.: Contribution of residential wood combustion and other sources to hourly winter aerosol in Northern Sweden determined by positive matrix factorization, *Atmos. Chem. Phys.*, 8(13), 3639–3653, doi:10.5194/acp-8-3639-2008, 2008.
- Kukkonen, J., Kangas, L., Kauhaniemi, M., Sofiev, M., Aarnio, M., Jaakkola, J. J. K., Kousa, A. and Karppinen, A.: Modelling of the urban concentrations of PM_{2.5} on a high resolution for a period of 35 years, for the assessment of lifetime exposure and health effects, *Atmos. Chem. Phys.*, 18(11), 8041–8064, doi:10.5194/acp-18-8041-2018, 2018.
- 555 Kukkonen, J., López-Aparicio, S., Segersson, D., Geels, C., Kangas, L., Kauhaniemi, M., Maragkidou, A., Jensen, A., Assmuth, T., Karppinen, A., Sofiev, M., Hellén, H., Riikonen, K., Nikmo, J., Kousa, A., Niemi, J. V., Karvosenoja, N., Sousa Santos, G., Sundvor, I., Im, U., Christensen, J. H., Nielsen, O. K., Plejdrup, M. S., Klenø Nøjgaard, J., Omstedt, G., Andersson, C., Forsberg, B. and Brandt, J.: The influence of residential wood combustion on the concentrations of PM_{2.5} in four Nordic cities, *Atmos. Chem. Phys.*, 20(7), 4333–4365, doi:10.5194/acp-20-4333-2020, 2020.
- 560 Kurppa, M., Hellsten, A., Auvinen, M., Raasch, S., Vesala, T. and Järvi, L.: Ventilation and Air Quality in City Blocks Using Large-Eddy Simulation—Urban Planning Perspective, *Atmosphere (Basel)*, 9(2), 65, doi:10.3390/atmos9020065, 2018.
- Lareau, N. P., Crosman, E., Whiteman, C. D., Horel, J. D., Hoch, S. W., Brown, W. O. J. and Horst, T. W.: The Persistent Cold-Air Pool Study, *Bull. Am. Meteorol. Soc.*, 94(1), 51–63, doi:10.1175/BAMS-D-11-00255.1, 2013.
- 565 Leiren, M. D. and Jacobsen, J. K. S.: Silos as barriers to public sector climate adaptation and preparedness: insights from road

- closures in Norway, *Local Gov. Stud.*, 44(4), 492–511, doi:10.1080/03003930.2018.1465933, 2018.
- Letzel, M. O., Krane, M. and Raasch, S.: High resolution urban large-eddy simulation studies from street canyon to neighbourhood scale, *Atmos. Environ.*, 42, 8770–8784, doi:10.1016/j.atmosenv.2008.08.001, 2008.
- Lopez-Aparicio, S. and Grythe, H.: Evaluating the effectiveness of a stove exchange programme on PM2.5 emission reduction, *Atmos. Environ.*, 231(October 2019), doi:10.1016/j.atmosenv.2020.117529, 2020.
- 570 Maronga, B., Gryscha, M., Heinze, R., Hoffmann, F., Kanani-Sühring, F., Keck, M., Ketelsen, K., Letzel, M. O., Sühring, M. and Raasch, S.: The Parallelized Large-Eddy Simulation Model (PALM) version 4.0 for atmospheric and oceanic flows: model formulation, recent developments, and future perspectives, *Geosci. Model Dev.*, 8, 2515–2551, doi:10.5194/gmd-8-2515-2015, 2015.
- 575 Maronga, B., Knigge, C. and Raasch, S.: An Improved Surface Boundary Condition for Large-Eddy Simulations Based on Monin–Obukhov Similarity Theory: Evaluation and Consequences for Grid Convergence in Neutral and Stable Conditions, *Boundary-Layer Meteorol.*, 1–45, doi:10.1007/s10546-019-00485-w, 2019a.
- Maronga, B., Gross, G., Raasch, S., Banzhaf, S., Forkel, R., Heldens, W., Kanani-Sühring, F., Matzarakis, A., Mauder, M., Pavlik, D., Pfafferoth, J., Schubert, S., Seckmeyer, G., Sieker, H. and Winderlich, K.: Development of a new urban climate model based on the model PALM – Project overview, planned work, and first achievements, *Meteorol. Zeitschrift*, 28(2), 105–119, doi:10.1127/metz/2019/0909, 2019b.
- 580 Masson, V., Heldens, W., Bocher, E., Bonhomme, M., Bucher, B., Burmeister, C., de Munck, C., Esch, T., Hidalgo, J., Kanani-Sühring, F., Kwok, Y. T., Lemosu, A., Lévy, J. P., Maronga, B., Pavlik, D., Petit, G., See, L., Schoetter, R., Tornay, N., Votsis, A. and Zeidler, J.: City-descriptive input data for urban climate models: Model requirements, data sources and challenges, *Urban Clim.*, 31(December 2018), doi:10.1016/j.uclim.2019.100536, 2020.
- 585 Mazzeo, A., Huneus, N., Ordoñez, C., Orfanoz-Cheuquelaf, A., Menut, L., Mailler, S., Valari, M., Denier van der Gon, H., Gallardo, L., Muñoz, R., Donoso, R., Galleguillos, M., Osses, M. and Tolvett, S.: Impact of residential combustion and transport emissions on air pollution in Santiago during winter, *Atmos. Environ.*, 190, 195–208, doi:10.1016/j.atmosenv.2018.06.043, 2018.
- 590 Park, S.-B., Baik, J.-J. and Han, B.-S.: Large-eddy simulation of turbulent flow in a densely built-up urban area, *Environ. Fluid Mech.*, 15(2), 235–250, doi:10.1007/s10652-013-9306-3, 2015.
- Resler, J., Krč, P., Belda, M., Juruš, P., Benešová, N., Lopata, J., Vlček, O., Damašková, D., Eben, K., Derbek, P., Maronga, B. and Kanani-Sühring, F.: PALM-USM v1.0: A new urban surface model integrated into the PALM large-eddy simulation model, *Geosci. Model Dev.*, 10(10), 3635–3659, doi:10.5194/gmd-10-3635-2017, 2017.
- 595 Ronda, R. J., Steeneveld, G. J., Heusinkveld, B. G., Attema, J. J. and Holtslag, A. A. M.: Urban Finescale Forecasting Reveals Weather Conditions with Unprecedented Detail, *Bull. Am. Meteorol. Soc.*, 98(12), 2675–2688, doi:10.1175/BAMS-D-16-0297.1, 2017.
- Savolahti, M., Lehtomäki, H., Karvosenoja, N., Ville-Veikko, P., Korhonen, A., Kukkonen, J., Kupiainen, K., Kangas, L., Karppinen, A. and Hänninen, O.: Residential wood combustion in Finland: PM2.5 emissions and health impacts with and

- 600 without abatement measures, *Int. J. Environ. Res. Public Health*, 16(16), doi:10.3390/ijerph16162920, 2019.
- Schneider, P., Castell, N., Vogt, M., Dauge, F. R., Lahoz, W. A. and Bartonova, A.: Mapping urban air quality in near real-time using observations from low-cost sensors and model information, *Environ. Int.*, 106, 234–247, doi:10.1016/j.envint.2017.05.005, 2017.
- Seljeskog, M., Goile, F. and Skreiberg, Ø.: Recommended Revisions of Norwegian Emission Factors for Wood Stoves, *Energy Procedia*, 105, 1022–1028, doi:10.1016/j.egypro.2017.03.447, 2017.
- 605 Simpson, W., Law, K., Schmale, J., Pratt, K., Arnold, S. and Mao, J.: Alaskan Layered Pollution And Chemical Analysis (ALPACA) White Paper., 2018.
- Solli, C., Reenaas, M., Strømman, A. H. and Hertwich, E. G.: Life cycle assessment of wood-based heating in Norway, *Int. J. Life Cycle Assess.*, 14(6), 517–528, doi:10.1007/s11367-009-0086-4, 2009.
- 610 Statens Kartverk: Høydedata, 2018.
- Stoll, R., Gibbs, J. A., Salesky, S. T., Anderson, W. and Calaf, M.: Large-Eddy Simulation of the Atmospheric Boundary Layer, *Boundary-Layer Meteorol.*, doi:10.1007/s10546-020-00556-3, 2020.
- Venter, Z. S., Brousse, O., Esau, I. and Meier, F.: Hyperlocal mapping of urban air temperature using remote sensing and crowdsourced weather data, *Remote Sens. Environ.*, 242(March), 111791, doi:10.1016/j.rse.2020.111791, 2020.
- 615 Wolf-Grosse, T., Esau, I. and Reuder, J.: Sensitivity of local air quality to the interplay between small- and large-scale circulations: a large-eddy simulation study, *Atmos. Chem. Phys.*, 17(11), 7261–7276, doi:10.5194/acp-17-7261-2017, 2017a.
- Wolf-Grosse, T., Esau, I. and Reuder, J.: The large-scale circulation during air quality hazards in Bergen, Norway, *Tellus A Dyn. Meteorol. Oceanogr.*, 69(1), 1406265, doi:10.1080/16000870.2017.1406265, 2017b.
- Wolf, T. and Esau, I.: A proxy for air quality hazards under present and future climate conditions in Bergen, Norway, *Urban Clim.*, 10(1), doi:10.1016/j.uclim.2014.10.006, 2014.
- 620 Wolf, T., Esau, I. and Reuder, J.: Analysis of the vertical temperature structure in the Bergen valley, Norway, and its connection to pollution episodes, *J. Geophys. Res. Atmos.*, 119(18), 10,645–10,662, doi:10.1002/2014JD022085, 2014.
- Wolf, T., Pettersson, L. H. and Esau, I.: A very high-resolution assessment and modelling of urban air quality, *Atmos. Chem. Phys.*, 20(2), 625–647, doi:10.5194/acp-20-625-2020, 2020.
- 625 Wyss, A. B., Jones, A. C., Bølling, A. K., Kissling, G. E., Chartier, R., Dahlman, H. J., Rodes, C. E., Archer, J., Thornburg, J., Schwarze, P. E. and London, S. J.: Particulate Matter 2.5 Exposure and Self-Reported Use of Wood Stoves and Other Indoor Combustion Sources in Urban Nonsmoking Homes in Norway, edited by R. A. Coulombe, *PLoS One*, 11(11), e0166440, doi:10.1371/journal.pone.0166440, 2016.
- Zhang, Y., Ye, X., Wang, S., He, X., Dong, L., Zhang, N., Wang, H., Wang, Z., Ma, Y., Wang, L., Chi, X., Ding, A., Yao, M.,
630 Li, Y., Li, Q., Zhang, L. and Xiao, Y.: Large-eddy simulation of traffic-related air pollution at a very high-resolution in a mega-city: Evaluation against mobile sensors and insights for influencing factors, *Atmos. Chem. Phys. Discuss.*, (November), doi:10.5194/acp-2020-1168, 2020.
- Zhong, J., Cai, X. M. and Bloss, W. J.: Coupling dynamics and chemistry in the air pollution modelling of street canyons: A

review, *Environ. Pollut.*, 214, 690–704, doi:10.1016/j.envpol.2016.04.052, 2016.

635 Bai, X.: Advance the ecosystem approach in cities, *Nature*, 559(7712), 7, doi:10.1038/d41586-018-05607-x, 2018.

Baklanov, A., Korsholm, U. S., Nuterman, R., Mahura, A., Nielsen, K. P., Sass, B. H., Rasmussen, A., Zakey, A., Kaas, E., Kurganskiy, A., Sørensen, B. and González Aparicio, I.: Enviro HIRLAM online integrated meteorology chemistry modelling system: strategy, methodology, developments and applications (v7.2), *Geosci. Model Dev.*, 10, 2971–2999, doi:10.5194/gmd-10-2971-2017, 2017a.

640 Baklanov, A., Brunner, D., Carmichael, G., Flemming, J., Freitas, S., Gauss, M., Hov, Ø., Mathur, R., Schlünzen, K. H., Seigneur, C. and Vogel, B.: Key Issues for Seamless Integrated Chemistry–Meteorology Modeling, *Bull. Am. Meteorol. Soc.*, 98(11), 2285–2292, doi:10.1175/BAMS-D-15-00166.1, 2017b.

Brandt, J., Silver, J. D., Christensen, J. H., Andersen, M. S., Bønløkke, J. H., Sigsgaard, T., Geels, C., Gross, A., Hansen, A. B., Hansen, K. M., Hedegaard, G. B., Kaas, E. and Frohn, L. M.: Contribution from the ten major emission sectors in Europe and Denmark to the health-cost externalities of air pollution using the EVA model system—an integrated modelling approach, *Atmos. Chem. Phys.*, 13(15), 7725–7746, doi:10.5194/acp-13-7725-2013, 2013.

645 Brötz, B., Eigenmann, R., Dörnbrack, A., Foken, T. and Wirth, V.: Early Morning Flow Transition in a Valley in Low Mountain Terrain Under Clear Sky Conditions, *Boundary Layer Meteorol.*, 152(1), 45–63, doi:10.1007/s10546-014-9921-7, 2014.

Castillo, M. C. L., Kanda, M. and Letzel, M. O.: Heat-ventilation efficiency of urban surfaces using large-eddy simulation, *Annu. J. Hydraul. Eng.*, 53, 175–180, 2009.

Cécé, R., Bernard, D., Brioude, J. and Zahibo, N.: Microscale anthropogenic pollution modelling in a small tropical island during weak trade winds: Lagrangian particle dispersion simulations using real-nested {LES} meteorological fields, *Atmos. Environ.*, 139, 98–112, doi:10.1016/j.atmosenv.2016.05.028, 2016.

655 Chandler, T. J.: *Urban Climatology and its Relevance to Urban Design.*, 1976.

Felius, L. C., Thalfeldt, M., Georges, L., Hrynyszyn, B. D., Dessen, F. and Hamdy, M.: Wood burning habits and its effect on the electrical energy demand of a retrofitted Norwegian detached house, *IOP Conf. Ser. Earth Environ. Sci.*, 352(1), doi:10.1088/1755-1315/352/1/012022, 2019.

660 Fernando, H. J. S. S., Zajic, D., Di Sabatino, S., Dimitrova, R., Hedquist, B. and Dallman, A.: Flow, turbulence, and pollutant dispersion in urban atmospheres, *Phys. Fluids*, 22(5), 051301, doi:10.1063/1.3407662, 2010.

Fröhlich, D. and Matzarakis, A.: Calculating human thermal comfort and thermal stress in the PALM model system-6.0, *Geosci. Model Dev.*, 13(7), 3055–3065, doi:10.5194/gmd-13-3055-2020, 2020.

665 Grimmond, S., Bouchet, V., Molina, L. T., Baklanov, A., Tan, J., Schlünzen, K. H., Mills, G., Golding, B., Masson, V., Ren, C., Voogt, J., Miao, S., Lean, H., Heusinkveld, B., Hovespyan, A., Teruggi, G., Parrish, P. and Joe, P.: Integrated urban hydrometeorological, climate and environmental services: Concept, methodology and key messages, *Urban Clim.*, 33(May 2019), 100623, doi:10.1016/j.uelim.2020.100623, 2020.

- Gronemeier, T., Raasch, S. and Ng, E.: Effects of Unstable Stratification on Ventilation in Hong Kong, *Atmosphere*, 8, 168, doi:10.3390/atmos8090168, 2017.
- 670 Hedberg, E., Johansson, C., Johansson, L., Swietlicki, E. and Brorström-Lundén, E.: Is Levoglucosan a Suitable Quantitative Tracer for Wood Burning? Comparison with Receptor Modeling on Trace Elements in Lycksele, Sweden, *J. Air Waste Manage. Assoc.*, 56(12), 1669–1678, doi:10.1080/10473289.2006.10464572, 2006.
- Hellén, H., Hakola, H., Haaparanta, S., Pietarila, H. and Kauhaniemi, M.: Influence of residential wood combustion on local air quality, *Sci. Total Environ.*, 393(2–3), 283–290, doi:10.1016/j.scitotenv.2008.01.019, 2008.
- 675 Høiskar, B. A. K., Sundvor, I., Johnsrud, M., Haug, T. W. and Nr, N. P.: Tiltaksutredning for lokal luftkvalitet i Bergen., 2017.
- Im, U., Christensen, J. H., Nielsen, O. K., Sand, M., Makkonen, R., Geels, C., Anderson, C., Kukkonen, J., Lopez Aparicio, S. and Brandt, J.: Contributions of Nordic anthropogenic emissions on air pollution and premature mortality over the Nordic region and the Arctic, *Atmos. Chem. Phys.*, 19(20), 12975–12992, doi:10.5194/acp-19-12975-2019, 2019.
- Jonassen, M. O., Olafsson, H., Valved, A. S., Reuder, J. and Olseth, J. A.: Simulations of the Bergen orographic wind shelter, *Tellus A Dyn. Meteorol. Oceanogr.*, 65(1), 19206, doi:10.3402/tellusa.v65i0.19206, 2013.
- 680 Keck, M., Raasch, S., Letzel, M. O. and Ng, E.: First Results of High Resolution Large Eddy Simulations of the Atmospheric Boundary Layer, *J. Heat Isl. Inst. Int.*, 9, 39–43, 2014.
- Költzow, M., Casati, B., Bazile, E., Haiden, T. and Valkonen, T.: An NWP Model Intercomparison of Surface Weather Parameters in the European Arctic during the Year of Polar Prediction Special Observing Period Northern Hemisphere I, *Weather Forecast.*, 34(4), 959–983, doi:10.1175/WAF-D-19-0003.1, 2019.
- 685 Kukkonen, J., López Aparicio, S., Segersson, D., Geels, C., Kangas, L., Kauhaniemi, M., Maragkidou, A., Jensen, A., Assmuth, T., Karppinen, A., Sofiev, M., Hellén, H., Riikonen, K., Nikmo, J., Kousa, A., Niemi, J. V., Karvosenoja, N., Sousa Santos, G., Sundvor, I., Im, U., Christensen, J. H., Nielsen, O. K., Plejdrup, M. S., Klenø Nøjgaard, J., Omstedt, G., Andersson, C., Forsberg, B. and Brandt, J.: The influence of residential wood combustion on the concentrations of PM_{2.5} in four Nordic cities, *Atmos. Chem. Phys.*, 20(7), 4333–4365, doi:10.5194/acp-20-4333-2020, 2020.
- 690 Kurppa, M., Hellsten, A., Auvinen, M., Raasch, S., Vesala, T. and Järvi, L.: Ventilation and Air Quality in City Blocks Using Large Eddy Simulation—Urban Planning Perspective, *Atmosphere (Basel)*, 9(2), 65, doi:10.3390/atmos9020065, 2018.
- Lareau, N. P., Crosman, E., Whiteman, C. D., Horel, J. D., Hoch, S. W., Brown, W. O. J. and Horst, T. W.: The Persistent Cold Air Pool Study, *Bull. Am. Meteorol. Soc.*, 94(1), 51–63, doi:10.1175/BAMS-D-11-00255.1, 2013.
- 695 Leiren, M. D. and Jacobsen, J. K. S.: Silos as barriers to public sector climate adaptation and preparedness: insights from road closures in Norway, *Local Gov. Stud.*, 44(4), 492–511, doi:10.1080/03003930.2018.1465933, 2018.
- Lemos, M. C., Kirchhoff, C. J. and Ramprasad, V.: Narrowing the climate information usability gap, *Nat. Clim. Chang.*, 2(11), 789–794, doi:10.1038/nclimate1614, 2012.
- Letzel, M. O., Krane, M. and Raasch, S.: High resolution urban large eddy simulation studies from street canyon to neighbourhoood scale, *Atmos. Environ.*, 42, 8770–8784, doi:10.1016/j.atmosenv.2008.08.001, 2008.
- 700

Lopez-Aparicio, S. and Grythe, H.: Evaluating the effectiveness of a stove exchange programme on PM2.5 emission reduction, *Atmos. Environ.*, 231(October 2019), doi:10.1016/j.atmosenv.2020.117529, 2020.

705 Maronga, B., Gryschka, M., Heinze, R., Hoffmann, F., Kanani Sühling, F., Keck, M., Ketelsen, K., Letzel, M. O., Sühling, M. and Raasch, S.: The Parallelized Large Eddy Simulation Model (PALM) version 4.0 for atmospheric and oceanic flows: model formulation, recent developments, and future perspectives, *Geosci. Model Dev.*, 8, 2515–2551, doi:10.5194/gmd-8-2515-2015, 2015.

Maronga, B., Knigge, C. and Raasch, S.: An Improved Surface Boundary Condition for Large Eddy Simulations Based on Monin–Obukhov Similarity Theory: Evaluation and Consequences for Grid Convergence in Neutral and Stable Conditions, *Boundary Layer Meteorol.*, 1–45, doi:10.1007/s10546-019-00485-w, 2019a.

710 Maronga, B., Gross, G., Raasch, S., Banzhaf, S., Forkel, R., Heldens, W., Kanani Sühling, F., Matzarakis, A., Mauder, M., Pavlik, D., Pffafferoth, J., Schubert, S., Seckmeyer, G., Sicker, H. and Winderlich, K.: Development of a new urban climate model based on the model PALM—Project overview, planned work, and first achievements, *Meteorol. Zeitschrift*, 28(2), 105–119, doi:10.1127/metz/2019/0909, 2019b.

Park, S.-B., Baik, J.-J. and Han, B.-S.: Large-eddy simulation of turbulent flow in a densely built-up urban area, *Environ. Fluid Mech.*, 15(2), 235–250, doi:10.1007/s10652-013-9306-3, 2015.

715 Resler, J., Krč, P., Belda, M., Juruš, P., Benešová, N., Lopata, J., Vlček, O., Damašková, D., Eben, K., Derbek, P., Maronga, B. and Kanani Sühling, F.: PALM USM v1.0: A new urban surface model integrated into the PALM large-eddy simulation model, *Geosci. Model Dev.*, 10(10), 3635–3659, doi:10.5194/gmd-10-3635-2017, 2017.

720 Ronda, R. J., Steeneveld, G. J., Heusinkveld, B. G., Attema, J. J. and Holtslag, A. A. M.: Urban Finescale Forecasting Reveals Weather Conditions with Unprecedented Detail, *Bull. Am. Meteorol. Soc.*, 98(12), 2675–2688, doi:10.1175/BAMS-D-16-0297.1, 2017.

Schneider, P., Castell, N., Vogt, M., Dauge, F.-R., Lahoz, W. A. and Bartonova, A.: Mapping urban air quality in near real-time—using observations from low cost sensors and model information, *Environ. Int.*, 106, 234–247, doi:10.1016/j.envint.2017.05.005, 2017.

725 Seljeskog, M., Goile, F. and Skreiberg, Ø.: Recommended Revisions of Norwegian Emission Factors for Wood Stoves, *Energy Procedia*, 105, 1022–1028, doi:10.1016/j.egypro.2017.03.447, 2017.

Simpson, W., Law, K., Schmale, J., Pratt, K., Arnold, S. and Mao, J.: Alaskan Layered Pollution And Chemical Analysis (ALPACA) White Paper., 2018.

730 Solli, C., Reenaas, M., Strømman, A. H. and Hertwich, E. G.: Life cycle assessment of wood-based heating in Norway, *Int. J. Life Cycle Assess.*, 14(6), 517–528, doi:10.1007/s11367-009-0086-4, 2009.

Statens Kartverk: Høydedata, 2018.

Stoll, R., Gibbs, J. A., Salesky, S. T., Anderson, W. and Calaf, M.: Large Eddy Simulation of the Atmospheric Boundary Layer, *Boundary Layer Meteorol.*, doi:10.1007/s10546-020-00556-3, 2020.

Venter, Z. S., Brousse, O., Esau, I. and Meier, F.: Hyperlocal mapping of urban air temperature using remote sensing and crowdsourced weather data, *Remote Sens. Environ.*, 242(March), 111791, doi:10.1016/j.rse.2020.111791, 2020.

Wolf Grosse, T., Esau, I. and Reuder, J.: Sensitivity of local air quality to the interplay between small and large scale circulations: a large eddy simulation study, *Atmos. Chem. Phys.*, 17(11), 7261–7276, doi:10.5194/acp-17-7261-2017, 2017a.

Wolf Grosse, T., Esau, I. and Reuder, J.: The large scale circulation during air quality hazards in Bergen, Norway, *Tellus A Dyn. Meteorol. Oceanogr.*, 69(1), 1406265, doi:10.1080/16000870.2017.1406265, 2017b.

Wolf, T. and Esau, I.: A proxy for air quality hazards under present and future climate conditions in Bergen, Norway, *Urban Clim.*, 10(1), doi:10.1016/j.uclim.2014.10.006, 2014.

Wolf, T., Esau, I. and Reuder, J.: Analysis of the vertical temperature structure in the Bergen valley, Norway, and its connection to pollution episodes, *J. Geophys. Res. Atmos.*, 119(18), 10,645–10,662, doi:10.1002/2014JD022085, 2014.

Wolf, T., Pettersson, L. H. and Esau, I.: A very high-resolution assessment and modelling of urban air quality, *Atmos. Chem. Phys.*, 20(2), 625–647, doi:10.5194/acp-20-625-2020, 2020.

Wyss, A. B., Jones, A. C., Bølling, A. K., Kissling, G. E., Chartier, R., Dahlman, H. J., Rodes, C. E., Archer, J., Thornburg, J., Schwarze, P. E. and London, S. J.: Particulate Matter 2.5 Exposure and Self-Reported Use of Wood Stoves and Other Indoor Combustion Sources in Urban Nonsmoking Homes in Norway, edited by R. A. Coulombe, *PLoS One*, 11(11), e0166440, doi:10.1371/journal.pone.0166440, 2016.








Figure 1: *Top:* Topographic map of the Bergen municipality. The colour shading indicates elevations of the land surface area. Water surfaces are marked with blue colour shading. The black curved lines indicate district boundaries. The black rectangle indicates the full extent of the model simulation domain. Buffer zones are not included. The grey rectangle indicates the part of the model domain that is selected for the analysis. *Bottom:* Map with the locations of wood-burning stoves in Bergen. The colour shading indicates the number of households with registered stoves per 10 x 10 m² grid box (averaged over 3 x 3 boxes). Black lines indicate waterfront. Data sources are the Bergen Municipality and the Bergen Fire department.



755



760 **Figure 2:** Surface layer meteorology obtained from the PALM model runs for the scenario 1 (top) and 2 (bottom). The colour shading represents the surface air temperature at 2 m above the ground (terrain-following temperature field). Vectors represent the surface-layer wind at 10 m. Data overlay the grayscale map from Map data © 2019 Google.



765 **Figure 3: Simulated concentrations at 5 m above the ground (the surface layer concentrations) of PM_{2.5} for the scenarios 1 (top) and 2 (bottom). Vectors show the surface wind at 10 m. The simulations use the current distributions of the wood-burning stoves and the current fraction of the new and older stoves in Bergen. Data overlay the grayscale map from Map data ©2019 Google.**



770

Figure 4: The simulated surface-layer concentrations of PM_{2.5} for the scenario 1 (shown in Fig. 3, top panel) but for emissions only from stoves within the Bergenhus (top) and Årstad (bottom) districts. The simulations use the current distributions of the number and types of stoves. The blue lines show the district boundaries. Thin grey isolines indicate the fraction of the total pollution that was caused by the emissions within the respective district. Data overlay the grayscale map from Map data ©2019 Google.







Figure 5: Same as Figure 4 but for the emissions within the districts Fana and Ytregyda (top), and Laksevåg and Fyllingsdalen (bottom). Data overlay the grayscale map from Map data ©2019 Google.



780


785

Figure 6: The same as in Figure 4 but for the emissions expected when all older stoves have been replaced with the new stoves in Bergenhus, Årstad, Fana and Ytrebygda districts (top) or in Laksevåg, Fyllingsdalen, Åsane and Arna districts (bottom). The boundaries of the districts are highlighted with the blue lines. Thin grey isolines indicate the fraction of the total pollution that was caused by the respective districts, when compared to the total pollution distribution in Figure 3 (top). Data overlay the grayscale map from Map data ©2019 Google.



affected today 17818
only new 1018

affected today 4031
only new 0



affected today 17243
only new 2137

affected today 5986
only new 0

790

Figure 7: Histogram of the accumulated number of households (# properties on the y-axis given in logarithmic scale) in Bergen that exposed for different concentrations (the binned concentration intervals on the x-axis) for the scenario 1 (top) and 2 (bottom). Abbreviations are: “2018” (upper part of all columns) means the exposure from the simulations with the current mix of older and new stoves; “old stoves exch.” means a complete replacement with the new stoves in various subsets of districts (as shown in Figure 6). The impact of the complete replacement in Åsane (ÅS), Arna (AR), Fyllingsdalen (FY) and Laksevåg (LA) (Figure 6, bottom) is shown as blue columns; the same in Bergenhus (BE), Årstad (ÅR), Fana (FA) and Ytrebygda (YT) (Figure 6, top) is shown as yellow columns; the same in all districts – as green columns. The threshold of 40 µg m⁻³ for PM_{2.5} concentrations is indicated with a vertical red line.

795

800

Table 1. Summary of the total simulated effect of the suggested mitigation actions.

	Weather scenario 1: Winter temperature inversion		Weather scenario 2: Winter temperature inversion and stagnation	
	Medium	Strong	Medium	Strong
Air pollution level	Medium	Strong	Medium	Strong
PM2.5 concentration range ($\mu\text{g m}^{-3}$)	20 - 40	40 - 110	20 - 40	40 - 110
The present-day situation: the total number of exposed households	17 818	4 031	17 243	5 986
Reduction in the number of the exposed households when transition to the new stoves is completed in Åsane, Arna, Fyllingsdalen and Laksevåg districts	20%	5%	20%	6%
Reduction in the number of the exposed households when transition to the new stoves is completed in Bergenhus, Årstad, Fana and Ytrebygda districts	82%	100%	72%	99%
Reduction in the number of the exposed households when transition to the new stoves is completed in the whole Bergen municipality	94%	100%	88%	100%

805