



## Measurement report: Ice nucleating particles active $\geq -15$ °C in free tropospheric air over western Europe

Franz Conen<sup>1</sup>, Annika Einbock<sup>1</sup>, Claudia Mignani<sup>1</sup>, Christoph Hüglin<sup>2</sup>

<sup>1</sup>Department of Environmental Sciences, University of Basel, CH-4056 Basel, Switzerland

5 <sup>2</sup>Laboratory for Air Pollution / Environmental Technology, Empa, CH-8600 Dübendorf, Switzerland

Correspondence to: Franz Conen (franz.conen@unibas.ch)

**Abstract.** Ice nucleating particles (INP) initiate ice formation in supercooled clouds, typically starting at a few km above ground. However, little is known about the concentration and composition of INP in the lower free troposphere (FT). Here, we analysed INP active at  $-10$  °C (INP<sub>-10</sub>) and  $-15$  °C (INP<sub>-15</sub>) collected during FT conditions at the high-altitude observatory  
10 Jungfraujoch. We relied on continuous radon measurements to distinguish FT conditions from those influenced by the planetary boundary layer. Median concentrations in the FT were  $2.4$  INP<sub>-10</sub> m<sup>-3</sup> and  $9.8$  INP<sub>-15</sub> m<sup>-3</sup>, with a multiplicative standard deviation of  $2.0$  and  $1.6$ , respectively. A majority of INP was deactivated after exposure to  $60$  °C, thus probably originated from certain epiphytic bacteria or fungi. Subsequent heating to  $95$  °C deactivated another  $15\%$  to  $20\%$  of the initial INP, likely other types of fungal INP that might be associated with soil organic matter or with decaying leaves. Very  
15 few INP<sub>-10</sub> withstood heating to  $95$  °C, but on average  $20\%$  of INP<sub>-15</sub> in FT samples did so. This percentage doubled during Saharan dust intrusions, which had practically no influence on INP<sub>-10</sub>. Overall, the results suggest that aerosolised epiphytic microorganisms, or parts thereof, are responsible for the majority of primary ice formation in moderately supercooled clouds above western Europe.

### 1 Introduction

20 Free troposphere (FT) designates a part of the troposphere that only occasionally is in exchange with Earth's surface, whereas the planetary boundary layer (PBL) continuously exchanges particles with surface sources and sinks. Consequently, the FT integrates particle emissions over a much larger area than does the PBL. A community of airborne microorganisms, for example, is in the PBL mainly composed of organisms from within a distance of several tens of kilometers (Tignat-Perrier et al., 2019). In contrast, particle populations sampled at a high-altitude mountain station under FT conditions  
25 constitute a mixture of many sources and sinks active on a continental scale (Herrmann et al., 2015, Sun et al., 2021). A special kind of aerosol particles, so-called ice-nucleating particles (INP), is relevant for primary ice formation in mid-level clouds (Findeisen, 1938). Ice formation starts in the midlatitudes a few kilometers above ground at around  $5$  °C (Kanitz et al., 2011). However, FT air masses have been little explored in terms of the INP they carry (He et al., 2021). In FT conditions at the high-altitude observatory Jungfraujoch (3580 m a.s.l.) in the Swiss Alps, Lacher et al. (2018) found similar  
30 concentrations of INP active at  $31$  °C (INP<sub>-31</sub>) as had been reported for the FT in other regions of the world, and little seasonal variation. Later, continuous measurements by Brunner et al. (2021b) found the monthly median of INP<sub>-30</sub> in the FT background to peak in April and to be lowest in December. Main components of the INP<sub>-31</sub> population studied by Lacher et al. (2018, 2021) were mineral dust ( $\sim 58\%$ ) and sea spray particles ( $17\%$ ), whereas biological particles were less frequent ( $10\%$ ). Also, Brunner et al. (2021a, 2021b) found most INP<sub>-30</sub> at Jungfraujoch occur during Saharan dust intrusions, by far  
35 exceeding any background concentration. Yet, at more moderate supercooling (here:  $\geq 15$  °C), biological particles constitute the main part of the INP population, at least in the PBL (Suski et al., 2018; Chen et al., 2021, Testa et al., 2021). Here, we investigate whether the same holds true for the FT. Our objectives were to quantify INP active at moderate supercooling in the FT, to narrow down the likely composition of these INP and how it might change during Saharan dust (SD) intrusions.



We will also compare INP concentrations in the FT with those of air masses influenced by the PBL as presented in earlier  
40 studies.

## 2 Material and methods

### 2.1 Sampling and analysis

For our investigation we relied on archived samples of particulate matter (PM<sub>10</sub>), collected on quartz-fibre filters at the high-  
altitude observatory Jungfraujoch (07°59'02" E, 46°32'53" N, 3580 m.a.s.l.), which is part of the Swiss national air quality  
45 monitoring network. The site, PM<sub>10</sub> sampling and INP analysis have been described earlier (Conen et al., 2012). Units of air  
volume throughout this paper are normalised to standard pressure. Briefly, air was sampled at a rate of 720 m<sup>3</sup> day<sup>-1</sup> through  
a heated inlet, followed by a PM<sub>10</sub> cut-off, onto a 150 mm diameter quartz-fibre filter (Pallflex Tissuquartz 2500-UP) with  
140 mm diameter of active area. Every day at midnight a new filter was exposed to the continuous air stream for 24 hours.  
From selected filters we took 72 punches with 2 mm diameter, containing aerosol particles from a total of 10.6 m<sup>3</sup>. Each  
50 punch was placed into a 0.5 mL Eppendorf safe-lock tube and 0.1 mL ultra-pure water (Sigma-Aldrich, W4502-1L) was  
added. The tubes were then immersed in a cooling bath and exposed to a temperature ramp from -5 °C to -15 °C (0.3 °C  
min<sup>-1</sup>). After each 1 °C step the number of frozen tubes was counted by eye and the cumulative concentration of INP was  
calculated according to Vali (1971). After the first such freezing assay, the tubes were placed for 10 min in a water bath at 60  
°C before the freezing assay was repeated. It was followed by a second 10-min heat treatment at 95 °C and a third freezing  
55 assay. Probably the largest uncertainty in these assays is related to an often small number of INP assessed. The 68%  
confidence interval ( $\pm 1 \sigma$ ) of a determined INP number concentration is roughly proportional to the square root of INP in an  
assay. In this study, an assay contained particles collected from 10.6 m<sup>3</sup> of air. Hence, a concentration of 1 INP m<sup>-3</sup> was  
represented by a total of 10 or 11 INP in the assay and is associated with an uncertainty of around  $\pm 0.3$  INP m<sup>-3</sup> ( $\sqrt{11 / 11 \times$   
1 INP m<sup>-3</sup>). Two assays with punches from 5 mm wide filter-fringes that were not exposed to the sample air were analysed  
60 the same way as the samples. Both of these blanks were composed of punches from four filters each. Samples values were  
corrected for blank values, on average 7% of a sample value. We calculated by difference the number concentration of heat  
sensitive INP (INP active before any heat treatment - INP active after 60 °C treatment) and moderately heat tolerant INP  
(INP active after 60 °C treatment - INP active after 95 °C treatment). What remained active after the 95 °C treatment is  
termed heat tolerant. In one sample (06.02.2021), the assay was completely frozen at -15 °C and was repeated with smaller  
65 punches (1 mm diameter).

### 2.2 Identification of free tropospheric condition

A key issue in our study is the distinction between FT and PBL-influenced air masses. For this purpose, we routinely  
monitor radon (<sup>222</sup>Rn) at Jungfraujoch with a dual-loop two-filter detector (Whittlestone and Zadorowski, 1998). Radon is  
emitted relatively homogeneously in space and time from land surfaces. Its only sink in the atmosphere is radioactive decay.  
70 Because of its short half-life (3.82 days), the concentration difference is large between the FT and the PBL, making radon a  
useful tracer to discriminate between these two types of air masses (Chambers et al., 2016). The probability density function  
of all hourly mean radon concentrations measured over the past five years is well reproduced by the sum of two log-normal  
distributions (Fig. 1), most likely representing FT and PBL-influenced air masses (Conen et al., 2021, Brunner et al., 2021a).  
Although FT conditions prevailed 41.5% of the time, it was only 1 in 13 days in 2019 and 2020 that such conditions lasted  
75 for a full 24-hour PM<sub>10</sub> sampling period. Here, we consider a 24-hour PM<sub>10</sub> sampling period as representative of the FT  
when all 24 values of hourly mean radon concentration during the sampling period were below 1.05 Bq m<sup>-3</sup>. Ninety percent  
of all values below 1.05 Bq m<sup>-3</sup> belong to the distribution representing FT conditions (Fig. 1). Two-thirds of the FT samples  
defined that way were collected in the months of January, February, and March. We selected for INP analysis 28 such



80 samples from between January 2019 and March 2021, making sure to have samples from each month of the year for which they were available. In addition, we selected from the filter archive an additional seven samples with high PM<sub>10</sub> loads resulting from Saharan dust intrusions (SD).

85 The median radon concentration in the FT distribution is only 0.28 times that of the PBL-influenced distribution (Fig. 1), hence, PBL injections in the FT may have a median age of about seven days. Thus, the airshed from where INP in FT air masses sampled at Jungfraujoch might have originated, is roughly indicated by seven-day backward trajectories calculated by the HYSPLIT model (Fig. 2). One trajectory was calculated for each day, ending at 13:00 UTC at 500 m above ground level, which for Jungfraujoch is 1500 m above sea level in the model surface. The model was run with the following settings: meteorology: GDAS (1 degree, global), vertical motion calculation method: model vertical velocity, location coordinates: 46°32'53'' N, 7°59'2'' E.

## 90 3 Results and discussion

### 3.1 Concentration and likely composition of ice nucleating particles in free tropospheric air

95 The observed range of INP concentration in FT air masses was 1.0 m<sup>-3</sup> to 5.6 m<sup>-3</sup> for INP active at -10 °C (INP<sub>-10</sub>) and 4.1 m<sup>-3</sup> to 16.3 m<sup>-3</sup> for INP active at -15 °C (INP<sub>-15</sub>). These values are within the lower range of INP concentrations observed in other contexts (Petters and Wright, 2015). The INP sampled at Jungfraujoch during FT conditions likely originated from the northern part of western Europe and from the North Atlantic (Fig. 2). Latter influence was probably minor. McCluskey et al. (2018) found a mean of 1.1 INP<sub>-15</sub> m<sup>-3</sup> in pristine marine air masses arriving at the west coast of Ireland, one tenth of the median we found at Jungfraujoch. A majority of INP<sub>-10</sub> (83%) and INP<sub>-15</sub> (57%) in the FT were heat sensitive and lost their activity after exposure to 60 °C (Table 1). Possible contributors to this category include the bacteria *Erwinia herbicola* (Phelps et al., 1986) and *Pseudomonas* sp. (Pouleur et al., 1992), as well as heat sensitive spores of several fungal species (Table 2). These microorganisms live on plant surfaces from where they, or parts of them, are emitted to the atmosphere (Lindow et al., 1978; Lindemann et al., 1982; Hirno and Upper, 2000; Huffman et al., 2013). Further potential sources include other fungi, such as *Fusarium graminearum* (Vujanovic et al., 2012; Keller et al., 2014) and *Puccinia* sp. (Morris et al., 2013). However, we do not know whether these already lose their ice nucleation activity at 60 °C, or between 60 °C and 95 °C, as heat stress tests were done only at around 95 °C, which is why they are not included in Table 2.

105

Between 15% and 20% of INP we found in the FT were moderately heat tolerant (deactivated between 60 °C and 95 °C). This deactivation temperature matches the profile of *Mortierella alpina*, a saprophytic fungus associated with decaying leaf litter (Vasebi et al., 2019) and with soil particles (Fröhlich-Nowoisky et al., 2015; Conen and Yakutin, 2018). Other potential sources of moderately heat tolerant INP include fungal symbionts in lichen (Kieft, 1988), *Fusarium avenaceum* (Pouleur et al., 1992), and the other above-mentioned fungi of which the deactivation temperature is not clearly defined. The sum of heat sensitive and moderately heat sensitive INP<sub>-15</sub> in a FT sample was on average 80% (standard deviation = 9%) of its total INP<sub>-15</sub> content, little smaller than the combined fractions of fungal and bacterial INP<sub>-15</sub> (92%) estimated in a case study over Amazonia (Patade et al., 2021).

115 Heat tolerant INP (not deactivated at 95 °C) constituted a negligibly small fraction of all INP<sub>-10</sub>, but to INP<sub>-15</sub> they contributed as much as did the moderately heat tolerant ones. The fraction of heat tolerant INP<sub>-15</sub> should substantially increase during SD intrusions, if mineral dust, in particular K-feldspar, was their main component (Vergara-Temprado et al., 2017), an issue addressed in the next section.



### 3.2 Effect of Saharan dust intrusions

120 Mineral dust makes by far the largest contribution to INP activated at around  $-30\text{ }^{\circ}\text{C}$  at Jungfraujoch (Larcher et al., 2018,  
2021; Brunner et al., 2021a, 2021b). Most INP in airborne mineral dust are probably K-feldspars, some of which are already  
activated at temperatures above  $-10\text{ }^{\circ}\text{C}$  (Boose et al., 2016; Harrison et al., 2019). Here, we analysed seven samples collected  
during SD intrusions (Annex, Table A1), but excluded from further analysis one sample (06.02.2021), in which INP had  
substantially increased at  $-15\text{ }^{\circ}\text{C}$  after each heat treatment. A similar reaction to heating was previously observed by Boose et  
125 al. (2019) and explained by the removal of secondary organic coatings that had masked ice-nucleation active sites before the  
heat treatment. In the remaining six samples, median  $\text{PM}_{10}$  concentration was 19 times as large as in FT samples.  
Surprisingly, the median concentration of heat tolerant  $\text{INP}_{-15}$  during SD intrusions was larger by only a factor of four, as  
compared to the FT (Table 1). Hence, mineral dust cannot be the main heat tolerant  $\text{INP}_{-15}$  in the FT. Otherwise, SD  
intrusions, which consist mainly of mineral dust, should have led to an increase in the heat tolerant fraction of  $\text{INP}_{-15}$  that is  
130 roughly proportional to the increase in dust load. Since we found no peak in heat tolerant  $\text{INP}_{-15}$  during the main pollen  
season in spring, pollen and pollen fragments (Diehl et al., 2002; Pummer et al., 2012) are also an unlikely source of them  
(note: recently, Burkart et al. (2021) found convincing evidence that ice nucleation activity birch pollen is related to a  
protein. Therefore, it probably is also heat sensitive, contrarily to earlier conclusions). More likely candidates of the heat  
tolerant fraction in the FT include soil organic material stabilised through bonding to minerals surfaces against deactivation  
135 by heat (Perkins et al., 2020), or lignin, not necessarily bound to mineral surfaces (Bogler and Borduas-Dedekind, 2020).  
Such organic compounds are more abundant in fertile soils in the mid-latitudes as compared to desert soils and could explain  
the comparatively small effect of SD intrusions on heat tolerant  $\text{INP}_{-15}$  observed at Jungfraujoch.

Further, the median of heat sensitive and of moderately heat tolerant  $\text{INP}_{-15}$  was larger during SD intrusions by a factor of  
140 1.6 and 2.3, respectively (Tab. 1). Their increase may be due to a concurrent influence of the PBL with all SD intrusions, as  
suggested by enhanced radon concentrations (Annex, Table A1). Overall, intrusions of SD and the associated PBL influence  
had doubled the median concentration of  $\text{INP}_{-15}$  and doubled the average fraction of heat tolerant  $\text{INP}_{-15}$  in a sample, but had  
no discernable effect on the median concentration of  $\text{INP}_{-10}$  (Table 1).

### 3.3 Comparison with other studies in the Swiss Alps

145 Mignani et al. (2021) have made a large number of impinger-based  $\text{INP}_{-15}$  measurements ( $n = 124$ , each integrating over 20  
min) in February and March 2019 at Weissfluhjoch, a mountain station 145 km east of Jungfraujoch, and at a 909 m lower  
elevation (2671 m a.s.l.). This station was throughout the day increasingly influenced by air from the PBL (Wieder et al.,  
2021). The median of all observations, excluding those influenced by SD ( $n = 11$ ), was  $15.5\text{ m}^{-3}$  ( $s^* = 4.0\text{ m}^{-3}$ ). This value is  
about a factor of two larger than the median of the four FT samples collected at Jungfraujoch during the same months,  
150 February and March 2019 ( $8.9\text{ m}^{-3}$ , range:  $5.5\text{ m}^{-3}$  to  $13.0\text{ m}^{-3}$ ). A factor of two decrease with an increase of 1000 m in  
altitude was already observed for  $\text{INP}_{-8}$  in Switzerland (Conen et al., 2017). Similar to the present study, SD intrusions at  
Weissfluhjoch enhanced the median  $\text{INP}_{-15}$  concentration by much less than the concurrent increase in particle number  
concentration.

155 A surprising feature of  $\text{INP}_{-10}$  in FT samples is the apparent lack of seasonal variation (Fig. 3), which is in contrast to what  
we had found earlier at Jungfraujoch in filters not selected for FT conditions (Conen et al., 2015). Those filters were sampled  
between June 2012 and May 2013 and analysed by the same method as used in this study, but without the heat treatments.  
Generally, the influence of the PBL on Jungfraujoch is largest in summer and smallest in winter (Collaud Coen et al., 2011;  
Griffith et al., 2014). Hence, if samples are not explicitly selected for FT conditions, the majority of them will have at least  
160 some PBL influence, in particular during summer (Brunner et al., 2021b). Therefore, concentration of  $\text{INP}_{-10}$  on filters not



selected for FT conditions occasionally exceeded that of filters selected for FT conditions, in particular during summer (Fig. 3). However, the opposite was true for winter. Especially during January and February, the concentration of INP<sub>-10</sub> was smaller by one or two orders of magnitude in the randomly selected samples of the year 2013, as compared to the FT samples of the years 2019, 2020, and 2021. A plausible explanation for this difference could be that in 2013 a substantially larger fraction of INP<sub>-10</sub> had been activated and deposited before reaching Jungfraujoch, as compared to the later years. Indeed, mean air temperature during sampling in January and February 2013 at Jungfraujoch was -16.2 °C (s.d. 5.3 °C), which is 6.4 °C colder than during sampling in January and February 2019, 2020, and 2021 (-9.8 °C; s.d. 4.6 °C). Yet, high relative humidity and low temperature does not always result in a particularly low number concentration of INP<sub>-10</sub> recorded at Jungfraujoch. On 14.03.2021 an exceptionally strong storm from North-West passed Jungfraujoch with gusts up to 122 km h<sup>-1</sup> and a mean daily windspeed of 51.5 km h<sup>-1</sup>. Daily mean values of relative humidity and temperature were 98% and -18.8 °C, respectively. Nevertheless, we found a non-negligible concentration of INP<sub>-10</sub> (0.7 m<sup>-3</sup>) on the PM<sub>10</sub> filter from that day. Updraft velocity at Jungfraujoch can reach a six-minute mean value of 8 m s<sup>-1</sup> during northwesterly wind (Hammer et al., 2014). An INP<sub>-10</sub> activated 1350 m below Jungfraujoch at -10 °C would grow for about 3 min ((18.8 °C - 10.0 °C) / 6.5 °C km<sup>-1</sup> / 8 m s<sup>-1</sup> = 169 s) into a crystal with a fall velocity of 5 cm s<sup>-1</sup> (Fukuta and Takahashi, 1999). As such, it can still enter the heated inlet of the sampler, which takes in droplets up to a size of about 40 μm (Weingartner et al., 1999), evaporate, and let the INP<sub>-10</sub> pass the PM<sub>10</sub> cut-off to be collected on the filter. At lower windspeed, growth time increases and activated INP<sub>-10</sub> are more likely to either settle or become too large to enter the heated inlet, eventually leading to the very low concentrations, which might explain our earlier observations in 2013. In an increasingly warmer climate, seasonality in the sink term of INP active at moderately supercooled conditions above the Alps may fade.

180

In summary, the results of this study suggest that epiphytic microorganisms contribute the majority of INP to ice formation in moderately supercooled clouds above the northern part of western Europe, whereas the impact of Saharan dust is negligible at -10 °C and still limited at -15 °C.

#### Acknowledgements

185 We thank the International Foundation High Altitude Research Stations Jungfraujoch and Gornergrat (HFSJG), 3012 Bern, Switzerland, for making it possible for us to work and to operate instruments at the High-Altitude Research Station at Jungfraujoch. We thank Claudia Zellweger and Stefan Reimann at Empa for helping us with the selection of PM<sub>10</sub> filter samples and for generously sending sections of them to Basel. We are grateful to Alastair Williams and his group at ANSTO for the ongoing collaboration as the supplier and supporter of the radon detection system. The radon observations are supported by the Swiss National Science Foundation (SNSF) as a contribution to the pan-European Integrated Carbon Observation System (ICOS) (<https://www.icos-ri.eu>). MeteoSwiss is acknowledged for the uncomplicated provision of meteorological data through its data portal IDAweb, and the Air Resources Laboratory at NOAA for providing access to its HYSPLIT model.

#### Author contribution

195 FC and CM conceived the study. CH organised the particle collection and provided the samples, AE did the freezing assays. FC prepared the manuscript with contribution from all coauthors.

#### Competing interests

The authors declare that they have no conflict of interest.



## References

- 200 Bogler, S., and Borduas-Dedekind, N.: Lignin's ability to nucleate ice via immersion freezing and its stability towards physicochemical treatments and atmospheric processing, *Atmos. Chem. Phys.*, 20, 14509–14522, <https://doi.org/10.5194/acp-20-14509-2020>, 2020.
- Boose, Y., Welti, A., Atkinson, J., Ramelli, F., Danielczok, A., Bingemer, H. G., Plötze, M., Sierau, B., Kanji, Z. A., and Lohmann, U.: Heterogeneous ice nucleation on dust particles sourced from nine deserts worldwide – Part 1: Immersion freezing, *Atmos. Chem. Phys.*, 16, 15075–15095, <https://doi.org/10.5194/acp-16-15075-2016>, 2016.
- Boose, Y., Baloh, P., Plötze, M., Ofner, J., Grothe, H., Sierau, B., Lohmann, U., and Kanji, Z. A.: Heterogeneous ice nucleation on dust particles sourced from 9 deserts worldwide – Part 2: Deposition nucleation and condensation freezing, *Atmos. Chem. Phys.*, 19, 1059–1076, <https://doi.org/10.5194/acp-19-1059-2019>, 2019.
- 210 Brunner, C., Brem, B. T., Martine Coen, M., Conen, F., Hervo, M., Henne, S., Steinbacher, M., Gysel-Beer, M., and Kanji, Z. A.: The contribution of Saharan dust to the ice nucleating particle concentrations at the High Altitude Station Jungfraujoch, Switzerland, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2021-643>, 2021a.
- Brunner, C., Brem, B. T., Collaud Coen, M., Conen, F., Steinbacher, M., Gysel-Beer, M., and Kanji, Z. A.: The diurnal and seasonal variability of ice nucleating particles at the High Altitude Station Jungfraujoch (3580 m a.s.l.), Switzerland, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2021-710>, 2021b.
- 215 Burkart, J., Glatzl, J., Seifried, T. M., Bieber, P., and Grothe, H.: Subpollen particles (SPP) of birch as carriers of ice nucleating macromolecules, *Biogeosciences Discuss.*, <https://doi.org/10.5194/bg-2021-8>, 2021.
- Chambers, S. D., Williams, A. G., Conen, F., Griffith, A. D., Reimann, S., Steinbacher, M., Krummel, P. B., Steele, L. P., van der Schoot, M. V., Galbally, I. E., Molloy, S. B., and Barnes, J. E.: Towards a universal "baseline" characterisation of air masses for high- and low-altitude observing stations using radon-222, *Aerosol Air Qual. Res.*, 16, 885-899, doi: 10.4209/aaqr.2015.06.0391, 2016.
- Chen, J., Wu, Z., Chen, J., Reicher, N., Fang, X., Rudich, Y., and Hu, M.: Size-resolved atmospheric ice-nucleating particles during East Asian dust events, *Atmos. Chem. Phys.*, 21, 3491–3506, <https://doi.org/10.5194/acp-21-3491-2021>, 2021.
- 225 Collaud Coen, M., Weingartner, E., Furger, M., Nyeki, S., Prévôt, A. S., Steinbacher, M., and Baltensperger, U.: Aerosol climatology and planetary boundary influence at the Jungfraujoch analyzed by synoptic weather types, *Atmospheric Chemistry and Physics*, 11, 5931–5944, <https://doi.org/10.5194/acp-11-5931-2011>, 2011.
- Conen, F., Morris, C. E., Leifeld, J., Yakutin, M. V., and Alewell, C.: Biological residues define the ice nucleation properties of soil dust, *Atmos. Chem. Phys.*, 11, 9643–9648, doi:10.5194/acp-11-9643-2011, 2011.
- Conen, F., Henne, S., Morris, C. E., and Alewell, C.: Atmospheric ice nucleators active  $\geq -12$  °C can be quantified in PM<sub>10</sub> filters, *Atmos. Meas. Tech.*, 5, 321–327, <https://doi.org/10.5194/amt-5-321-2012>, 2012.
- 230 Conen, F., Rodríguez, S., Hüglin, C., Henne, S., Herrmann, E., Bukowiecki, N., and Alewell, C.: Atmospheric ice nuclei at the high-altitude observatory Jungfraujoch, Switzerland, *Tellus B*, 67, 25014, <https://doi.org/10.3402/tellusb.v67.25014>, 2015.
- Conen, F., Yakutin, M. V., Yttri, K. E., and Hüglin, C.: Ice nucleating particle concentrations increase when leaves fall in autumn, *Atmosphere*, 8, 202; doi:10.3390/atmos8100202, 2017
- 235 Conen, F., and Yakutin, M. V.: Soils rich in biological ice-nucleating particles abound in ice-nucleating macromolecules likely produced by fungi, *Biogeosciences*, 15, 4381–4385, <https://doi.org/10.5194/bg-15-4381-2018>, 2018.
- Conen, F., and Zimmermann, L.: Separating 'free tropospheric conditions' from those 'influenced by the planetary boundary layer', *International Foundation HFSJG Activity Report 2020*, 49-50, [https://www.hfsjg.ch/reports/2020/pdf/120\\_UniBasel\\_Conen\\_radon\\_cf.pdf](https://www.hfsjg.ch/reports/2020/pdf/120_UniBasel_Conen_radon_cf.pdf), 2021.
- 240 Daily, M. I., Tarn, M. D., Whale, T. F., and Murray, B. J.: The sensitivity of the ice-nucleating ability of minerals to heat and the implications for the heat test for biological ice nucleators, *Atmos. Meas. Tech. Discuss.*, <https://doi.org/10.5194/amt-2021-208>, 2021.
- 245 Diehl, K., Matthias-Maser, S., Jaenicke, R., and Mitra, S. K.: The ice nucleation ability of pollen Part II: Laboratory studies in immersion and contact freezing modes, *Atmos. Res.*, 61, 125–133, 2002.



- Failor, K. C., Schmale III, D. G., Vinatzer, B. A., and Monteil, C. L.: Ice nucleation active bacteria in precipitation are genetically diverse and nucleate ice by employing different mechanisms, *ISME J.*, 11, 2740–2753, <https://doi.org/10.1038/ismej.2017.124>, 2017.
- 250 Findeisen, W.: Die kolloidmeteorologischen Vorgänge bei der Niederschlagsbildung (Colloidal meteorological processes in the formation of precipitation), *Meteorol. Z.* 55, 121–133, 1938 (translated and edited by Volken, E., Giesche, A. M., Brönnimann, S., *Meteorol. Z.*, 24, doi:10.1127/metz/2015/0675, 2015.
- Fröhlich-Nowoisky, J., Hill, T. C. J., Pummer, B. G., Franc, G. D., and Pöschl, U.: Ice nucleation activity in the widespread soil fungus *Mortierella alpina*, *Biogeosciences*, 12, 1057–1071, <https://doi.org/10.5194/bg-12-1057-2015>, 2015.
- 255 Fukuta, N. and Takahashi, T.: The growth of atmospheric ice crystals: A summary of findings in vertical supercooled cloud tunnel studies, *J. Atmos. Sci.*, 56, 1963–1979, 1999.
- Griffith, A. D., Conen, F., Weingartner, E., Zimmermann, L., Chambers, S. D., Williams, A. G., and Steinbacher, M.: Surface-to-mountaintop transport characterised by radon observations at the Jungfrauoch, *Atmos. Chem. Phys.*, 14, 12763–12779, doi:10.5194/acp-14-12763-2014, 2014.
- 260 Hammer, E., Bukowiecki, N., Gysel, M., Jurányi, Z., Hoyle, C. R., Vogt, R., Baltensperger, U., and Weingartner, E.: Investigation of the effective peak supersaturation for liquid-phase clouds at the high-alpine site Jungfrauoch, Switzerland (3580 m a.s.l.), *Atmos. Chem. Phys.*, 14, 1123–1139, doi:10.5194/acp-14-1123-2014, 2014.
- Harrison, A. D., Lever, K., Sanchez-Marroquin, A., Holden, M. A., Whale, T. F., Tarn, M. D., McQuaid, J. B., and Murray, B. J.: The ice-nucleating ability of quartz immersed in water and its atmospheric importance compared to K-feldspar, *Atmos. Chem. Phys.*, 19, 11343–11361, 10.5194/acp-19-11343-2019, 2019.
- 265 He, C., Yin, Y., Wang, W., Chen, K., Mai, R., Jiang, H., Zhang, X., and Fang, C.: Aircraft observations of ice nucleating particles over the Northern China Plain: Two cases studies, *Atmos. Res.*, 248, 105242, <https://doi.org/10.1016/j.atmosres.2020.105242>, 2021.
- Herrmann, E., Weingartner, E., Henne, S., Vuilleumier, L., Bukowiecki, N., Steinbacher, M., Conen, F., Coen, M. C., Hammer, E., Jurányi, Z., Baltensperger, U., and Gysel, M.: Analysis of long-term aerosol size distribution data from Jungfrauoch with emphasis on free tropospheric conditions, cloud influence, and air mass transport, *J. Geophys. Res.*, 120, 9459–9480, <https://doi.org/10.1002/2015JD023660>, 2015.
- 270 Hill, T. C. J., DeMott, P. J., Tobo, Y., Fröhlich Nowoisky, J., Moffett, B. F., Franc, G. D., and Kreidenweis, S. M.: Sources of ice nucleating particles in soils, *Atmos. Chem. Phys.*, 16, 7195–7211, <https://doi.org/10.5194/acp-16-7195-2016>, 2016.
- 275 Hirano, S. S., and Upper, C. D.: Bacteria in the Leaf Ecosystem with Emphasis on *Pseudomonas syringae* - a Pathogen, Ice Nucleus, and Epiphyte, *Microbiol. Mol. Biol. Rev.*, 64, 624–653, 2000.
- Huffman, J. A., Prenni, A. J., DeMott, P. J., Pöhlker, C., Mason, R. H., Robinson, N. H., Fröhlich-Nowoisky, J., Tobo, Y., Després, V. R., Garcia, E., Gochis, D. J., Harris, E., Müller-Germann, I., Ruzene, C., Schmer, B., Sinha, B., Day, D. A., Andreae, M. O., Jimenez, J. L., Gallagher, M., Kreidenweis, S. M., Bertram, A. K., and Pöschl, U.: High concentrations of biological aerosol particles and ice nuclei during and after rain, *Atmos. Chem. Phys.*, 13, 6151–6164, <https://doi.org/10.5194/acp-13-6151-2013>, 2013.
- 280 Kanitz, T., Seifert, P., Ansmann, A., Engelmann, R., Althausen, D., Casaccia, C., and Rohwer, E. G.: Contrasting the impact of aerosols at northern and southern midlatitudes on heterogeneous ice formation, *Geophys. Res. Lett.*, 38, L17802, doi:10.1029/2011GL048532, 2011.
- 285 Keller, M. D., Bergstrom, G. C., and Shields, E. J.: The aerobiology of *Fusarium graminearum*, *Aerobiologia*, 30, 123–136, doi: 10.1007/s10453-013-9321-3, 2014.
- Kieft, T. L.: Ice nucleation activity in lichens, *Appl. Environ. Microbiol.*, 54, 1678–1681, 1988.
- Kunert, A. T., Pöhlker, M. L., Tang, K., Krevert, C. S., Wieder, C., Speth, K. R., Hanson, L. E., Morris, C. E., Schmale, D. G. III, Pöschl, U., and Fröhlich-Nowoisky, J.: Macromolecular fungal ice nuclei in *Fusarium*: effects of physical and chemical processing, *Biogeosciences*, 16, 4647–4659, <https://doi.org/10.5194/bg-16-4647-2019>, 2019.
- 290 Lacher, L., DeMott, P. J., Levin, E. J. T., Suski, K. J., Boose, Y., Zipori, A., Herrmann, E., Bukowiecki, N., Steinbacher, M., Gute, E., Abbatt, J. P., Lohmann, U., and Kanji, Z. A.: Background free-tropospheric ice nucleating particle concentrations at mixed-phase cloud conditions, *Journal of Geophysical Research: Atmospheres*, 123, 10,506–10,525, <https://doi.org/10.1029/2018JD028338>, 2018.
- 295 Lacher, L., Clemen, H.-C., Shen, X., Mertes, S., Gysel-Beer, M., Moallemi, A., Steinbacher, M., Henne, S., Saathoff, H., Möhler, O., Höhler, K., Schiebel, T., Weber, D., Schrod, J., Schneider, J., and Kanji, Z. A.: Sources and nature of ice-



- nucleating particles in the free troposphere at Jungfraujoch in winter 2017, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2021-415>, 2021.
- Limpert, E., Stahel, W.A., and Abbt, M.: Log-normal distributions across the sciences: Keys and clues, *BioScience*, 51, 341-352, <http://stat.ethz.ch/~stahel/lognormal/bioscience.pdf>, 2001.
- 300 Limpert, E., Burke, J., Galan, C., del Mar Trigo, M., West, J.S., and Stahel, W.A.: Data, not only in aerobiology: How normal is the normal distribution? *Aerobiologia* 2008, 24, 121–124, doi:10.1007/s10453-008-9092-4, 2008.
- Lindemann, J., Constantinidou, H. A., Barchet, W. R., and Upper, C. D.: Plants as sources of airborne bacteria, including ice nucleation-active bacteria, *Appl. Environ. Microbiol.*, 44, 1059-1063, 1982.
- Lindow, S. E., Arny, D. C., and Upper, C. D.: Distribution of ice nucleation-active bacteria on plants in nature, *Appl. Environ. Microbiol.*, 36, 831-838, 1978.
- 305 McCluskey, C. S., Ovadnevaite, J., Rinaldi, M., Atkinson, J., Belosi, F., Ceburnis, D., Marullo, S., Hill, T. C. J., Lohmann, U., Kanji, Z. A., O'Dowd, C., Kreidenweis, S. M., and DeMott, P. J.: Marine and terrestrial organic ice-nucleating particles in pristine marine to continentally influenced northeast Atlantic air masses, *J. Geophys. Res. Atmos.* 123, 6196–6212. <https://doi.org/10.1029/2017JD028033>, 2018.
- 310 Mignani, C., Wieder, J., Sprenger, M. A., Kanji, Z. A., Henneberger, J., Alewell, C., and Conen, F.: Towards parameterising atmospheric concentrations of ice-nucleating particles active at moderate supercooling, *Atmos. Chem. Phys.*, 21, 657–664, <https://doi.org/10.5194/acp-21-657-2021>, 2021.
- Morris, C. E., Sands, D. C., Glaux, C., Samsatly, J., Asaad, S., Moukahel, A. R., Gonçalves, F. L. T., and Bigg, E. K.: Urediospores of rust fungi are ice nucleation active at > -10 °C and harbor ice nucleation active bacteria, *Atmos. Chem. Phys.*, 13, 4223–4233, <https://doi.org/10.5194/acp-13-4223-2013>, 2013.
- 315 O'Sullivan, D., Murray, B. J., Ross, J. F., Whale, T. F., Price, H. C., Atkinson, J. D., Umo, N. S., and Webb, M. E.: The relevance of nanoscale biological fragments for ice nucleation in clouds, *Sci. Rep.*, 5, 8082, doi:10.1038/srep08082, 2015.
- Patade, S., Phillips, V. T. J., Amato, P., Bingemer, H. G., Burrows, S. M., DeMott, P. J., Goncalves, F. L. T., Knopf, D. A., Morris, C. E., Alwmark, C., Artaxo, P., Pöhlker, C., Schrod, J., and Weber, B.: Empirical formulation for multiple groups of primary biological ice nucleating particles from field observations over Amazonia, *J. Atmos. Sci.*, 78, 2195-2220, <https://doi.org/10.1175/JAS-D-20-0096.1>, 2021.
- 320 Perkins, R. J., Gillette, S. M., Hill, T. C. J., and DeMott, P. J.: The Labile Nature of Ice Nucleation by Arizona Test Dust, *ACS. Earth. Space. Chem.*, 4, 133-141, 10.1021/acsearthspacechem.9b00304, 2020.
- Petters, M. D. and Wright, T. P.: Revisiting ice nucleation from precipitation samples, *Geophys. Res. Lett.*, 42, 8758–8766, <https://doi.org/10.1002/2015GL065733>, 2015.
- 325 Phelps, P., Giddings, T. H., Prochoda, M., and Fall, R.: Release of cell-free ice nuclei by *Erwinia herbicola*, *J. Bacteriol.*, 167, 496-502, 1986.
- Pouleur, S., Richard, C., Martin, J.-G., and Antoun, H.: Ice nucleation activity in *Fusarium acuminatum* and *Fusarium avenaceum*, *Appl. Environ. Microbiol.*, 58, 2960–2964, 1992.
- 330 Pummer, B. G., Bauer, H., Bernardi, J., Bleicher, S., and Grothe, H.: Suspendable macromolecules are responsible for ice nucleation activity of birch and conifer pollen, *Atmos. Chem. Phys.*, 12, 2541-2550, 10.5194/acp-12-2541-2012, 2012.
- Schnell, R. C. and Vali, G.: Worldwide source of leaf derived freezing nuclei, *Nature*, 246, 212–213, 1973.
- Sun, J., Herrmann, M., Yuan, Y., Birmili, W., Collaud Coen, M., Weinhold, K., Madueno, L., Poulain, L., Tuch, T., Ries, L., Sohmer, R., Couret, C., Frank, G., Brem, B. T., Gysel-Beer, M., Ma, N., and Wiedensohler, A.: Long-term trends of black carbon and particle number concentration in the lower free troposphere in Central Europe, *Environ. Sci. Eur.*, 33, 47, <https://doi.org/10.1186/s12302-021-00488-w>, 2021.
- 335 Suski, K. J., Hill, T. C. J., Levin, E. J. T., Miller, A., DeMott, P. J., and Kreidenweis, S. M.: Agricultural harvesting emissions of ice-nucleating particles, *Atmos. Chem. Phys.*, 18, 13755–13771, <https://doi.org/10.5194/acp-18-13755-2018>, 2018.
- 340 Testa, B., Hill, T. C. J., Marsden, N., Barry, K. R., Hume, C. C., Bian, Q., Uetake, J., Hare, H., Perkins, J. R., Möhler, O., Kreidenweis, S. M., and DeMott, P. J.: Ice nucleating particle connections to regional Argentinian land surface emissions and weather during the Cloud, Aerosol, and Complex Terrain Interactions experiment, *ESSOAr*, <https://doi.org/10.1002/essoar.10506960.1>, 2021.





- 345 Tignat-Perrier, R., Dommergue, A., Thollot, A., Keuschnig, C., Magand, O., Vogel, T. M., and Larose, C.: Global airborne microbial communities controlled by landscapes and wind condition, *Sci. Rep.*, 9, 14441, <https://doi.org/10.1038/s41598-019-51073-4>, 2019.
- Vali, G.: Quantitative evaluation of experimental results on the heterogeneous freezing nucleation of supercooled liquids, *J. Atmos. Sci.*, 28, 402–409, [https://doi.org/10.1175/1520-0469\(1971\)028<0402:QEOERA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1971)028<0402:QEOERA>2.0.CO;2), 1971.
- 350 Vergara-Temprado, J., Murray, B. J., Wilson, T. W., O’Sullivan, D., Browse, J., Pringle, K. J., Ardon-Dryer, K., Bertram, A. K., Burrows, S. M., Ceburnis, D., DeMott, P. J., Mason, R. H., O’Down, C. D., Rinaldi, M., and Carslaw, K. S.: Contribution of feldspar and marine organic aerosols to global ice nucleating particle concentrations, *Atmos. Chem. Phys.*, 17, 3637–3658, [www.atmos-chem-phys.net/17/3637/2017/](http://www.atmos-chem-phys.net/17/3637/2017/), 2017.
- 355 Vasebi, Y., Mehan Llonop, M. E., Hanlon, R., Schmale III, D. G., Schnell, R. C., and Vinatzer, B. A.: Comprehensive characterization of an aspen (*Populus tremuloides*) leaf litter sample that maintained ice nucleation activity for 48 years, *Biogeosciences*, 16, 1675–1683, <https://doi.org/10.5194/bg-16-1675-2019>, 2019.
- Vujanovic, V., Goh, Y. K., and Daida, P.: Heat- and Cold-Shock Responses in *Fusarium graminearum* 3 Acetyl- and 15 Acetyl-Deoxynivalenol Chemotypes, *J. Microbiol.* 50, 97-102, doi: 10.1007/s12275-012-1381-5, 2012.
- 360 Weingartner, E., Nyeki, S., and Baltensperger, U.: Seasonal and diurnal variation of aerosol size distributions ( $10 < D < 750$  nm) at a high-alpine site (Jungfraujoch 3580 m asl), *J. Geophys. Res.*, 104, 26809–26820, <https://doi.org/10.1029/1999JD900170>, 1999.
- Wieder, J., Mignani, C., Schär, M., Roth, L., Sprenger, M., Henneberger, J., Lohmann, U., Brunner, C., and Kanji, Z. A.: Unveiling atmospheric transport and mixing mechanisms of ice nucleating particles over the Alps, (in preparation).
- Whittlestone, S. and Zahorowski, W.: Baseline radon detectors for shipboard use: Development and deployment in the First Aerosol Characterization Experiment (ACE 1), *J. Geophys. Res.*, 103, 16743–16751, 1998.

365



370 **Table 1: Median and multiplicative standard deviation of INP concentration found in the free troposphere (FT) and during**  
**Saharan dust intrusions (SD), categorised according to heat sensitivity. Use of the multiplicative standard deviation ( $s^*$ ) is**  
**appropriate because multiplicative processes determine the concentration of biological particles in the atmosphere (Limpert et al.,**  
**2001, 2008). It was estimated from the upper (q3) and lower (q1) quartiles of a distribution:  $s^* = (q3/q1)0.741$  (Limpert et al.,**  
 375 **2001). It was not estimated where the lower quartile was below detection limit ( $< 0.095$  INP m<sup>-3</sup>). Heat sensitive are designated**  
**INP that had lost their activity after exposure to 60 °C. Moderately heat tolerant are termed those INP that had retained their**  
**activity after exposure to 60 °C, but had lost it after exposure to 95 °C. Heat tolerant INP are those that were still active after both**  
**heat treatments.**

Category	INP <sub>-10</sub> (m <sup>-3</sup> )		INP <sub>-15</sub> (m <sup>-3</sup> )	
	FT	SD	FT	SD
all	2.4 <sub>2.0</sub>	2.4 <sub>1.5</sub>	9.8 <sub>1.6</sub>	20.0 <sub>1.7</sub>
heat sensitive	2.0 <sub>1.8</sub>	1.9 <sub>1.6</sub>	5.6 <sub>1.5</sub>	9.2 <sub>2.1</sub>
moderately heat tolerant	0.4 <sub>1.9</sub>	0.2 <sub>n.d.</sub>	1.5 <sub>2.8</sub>	3.5 <sub>2.8</sub>
heat tolerant	0.0 <sub>n.d.</sub>	0.1 <sub>2.8</sub>	1.6 <sub>1.6</sub>	6.2 <sub>1.5</sub>



380


385

**Table 2: Reported heat sensitivity of INP active at moderate supercooling. Indicated is the temperature range in which at least 90% of a specific type of INP active  $\geq 15$  °C was found to be deactivated, although a smaller fraction of the same INP may already have been deactivated at a lower temperature. For example, soil particles analysed by Conen and Yakutin (2018) had lost half of their INP-10 after exposure to 60 °C, but more than 98% after exposure to 95 °C. These particles are assigned a deactivation temperature between 60 °C and 95 °C. Note: studies where INP were found to be deactivated after a single heat treatment  $\sim 95$  °C are not listed because deactivation might already have happened  $< 60$  °C.**

Type	Deactivation temperature			Reference
	$< 60$ °C	60 °C - 95 °C	$> 95$ °C	
Bacteria	<i>Erwinia herbicola</i>			Phelps et al. (1986)
	<i>Pseudomonas sp.</i>			Pouleur et al. (1992)
			<i>Lysinibacillus sp.</i>	Failor et al. (2017)
Fungi	<i>Acremonium implicatum</i>			Pummer et al. (2015)
	<i>Isaria farinosa</i>			Pummer et al. (2015)
	<i>Fusarium acuminatum</i>			Kunert et al. (2019)
	<i>Fusarium langsethiae</i>			Kunert et al. (2019)
	<i>Fusarium armeniacum</i>			Kunert et al. (2019)
		Fungal symbiont in lichen		
Others		<i>Mortierella alpina</i>		Fröhlich-Nowoisky et al. (2015)
		<i>Fusarium avenaceum</i>		Pouleur et al. (1992)
	mixed	Leaf-derived ice nuclei		Schnell and Vali (1973)
organic		Soil particles		Hill et al. (2016)
		Soil particles		Conen and Yakutin (2018)
			Pollen	Diehl et al. (2002), Pummer et al. (2012)
mineral			Lignin	Bogler and Borduas-Dedekind (2020)
			K-feldspars	Daily et al. (2021)
			Montmorillonite	Conen et al. (2011)
			Illite	O'Sullivan et al. (2015)
		Kaolinite	Daily et al. (2021)	




390



395 **Figure 1: Probability density function of log-transformed hourly mean radon concentration observed during the years 2016 to 2020 at Jungfraujoch (black line). The sum of two fitted and weighed log-normal distributions (grey dotted line) closely matches the observed distribution. The fitted distributions most likely represent the distribution of free tropospheric air masses (blue continuous line) and that of air masses influenced by the planetary boundary layer (red line). Ninety percent of the values below the dashed vertical line belong to the fitted 'free troposphere' distribution (Figure adapted from Conen et al., 2021).**

400

405




410

**Figure 2.** Ensemble of seven-day backward trajectories ending at Jungfraujoch at noon local time on days for which we analysed INP in free tropospheric conditions. The trajectories were calculated by the HYSPLIT model, made available by the Air Resources Laboratory at NOAA (<https://www.ready.noaa.gov/HYSPLIT.php>).



415



420 **Figure 3. Concentration of INP<sub>-10</sub> when free tropospheric conditions prevailed at Jungfraujoch throughout a full day (open**  
**squares) and for days not selected for this criterion (crosses; data from Conen et al., 2015, see Annex, Table A2).**

425

430



435

**Table A1: Number concentration of ice nucleating particles active at -10 °C (INP<sub>-10</sub>) and at -15 °C (INP<sub>-15</sub>) at Jungfraujoch, determined for free tropospheric conditions and during Saharan dust intrusions. The same sample material was analysed three times. A first freezing assay was followed by a heat treatment (10 min at 60 °C) and second freezing assay, followed by a second heat treatment (10 min at 95 °C) and a third freezing assay. All steps were carried out in an uninterrupted operation. Blank values were subtracted from raw measured values. Three categories of INP were derived by difference between INP number concentration before and after each heat treatment, resulting in negative values where INP concentration had increased after a heat treatment. Meteorological data is from MeteoSwiss and PM<sub>10</sub> data from the Swiss air quality monitoring network (NABEL).**

Date	INP <sub>-10</sub> (m <sup>-3</sup> )			INP <sub>-15</sub> (m <sup>-3</sup> )			T <sub>mean</sub> (°C)	Fraction of hours with RH > 95%	Radon			PM <sub>10</sub> (µg m <sup>-3</sup> )
	Deactivation temperature (°C)			Deactivation temperature (°C)					min	max	mean	
	< 60	60-95	> 95	< 60	60-95	> 95			(Bq m <sup>-3</sup> )			
<i>Free tropospheric conditions</i>												
06.01.19	1.6	0.3	0.0	5.1	0.7	1.1	-13.1	0.00	0.32	0.48	0.39	0.5
26.01.19	1.1	0.3	0.1	3.8	1.5	1.6	-11.1	0.00	0.20	0.40	0.29	0.7
05.02.19	1.5	0.2	0.0	4.3	0.1	0.7	-9.6	0.00	0.30	0.62	0.46	0.8
03.03.19	3.8	1.2	0.1	10.4	1.7	1.3	-8.4	0.04	0.30	0.82	0.54	0.9
08.03.19	2.3	0.8	0.1	6.6	2.6	3.6	-13.8	0.04	0.45	0.86	0.66	0.8
27.03.19	1.2	0.2	0.0	3.9	-0.2	1.3	-8.7	0.00	0.22	0.48	0.34	1.3
07.05.19	2.2	0.1	0.1	6.3	1.8	2.0	-7.7	0.00	0.37	0.92	0.58	1.9
14.05.19	2.9	0.3	0.2	7.0	1.8	5.6	-7	0.00	0.29	1.01	0.51	2.3
03.09.19	2.6	1.6	0.0	5.6	2.9	1.6	1.3	0.00	0.35	0.75	0.48	1.4
12.09.19	1.7	1.4	0.0	3.1	3.6	4.3	1.2	0.13	0.21	0.99	0.48	1.1
16.10.19	3.3	1.1	0.0	5.3	2.2	2.1	-4.9	0.13	0.43	0.69	0.56	0.6
28.12.19	3.3	0.3	0.1	7.3	1.3	2.0	-7	0.00	0.43	0.94	0.64	0.7
30.12.19	1.6	0.4	0.0	2.9	0.1	1.1	-2.4	0.00	0.38	1.01	0.68	0.4
05.01.20	1.7	0.5	0.1	7.5	0.1	1.2	-7.5	0.00	0.26	0.78	0.43	0.5
14.01.20	4.0	0.6	0.0	10.5	1.4	1.0	-6.1	0.00	0.54	0.84	0.68	1.1
16.02.20	4.2	0.6	0.2	8.9	1.3	2.0	-0.9	0.00	0.23	0.66	0.49	0.8
20.02.20	4.9	0.7	0.0	11.4	1.6	3.3	-10.1	0.00	0.19	0.51	0.34	1.7
10.03.20	1.4	0.1	0.1	7.4	0.3	1.4	-8.4	0.08	0.16	0.96	0.45	2.6
02.05.20	0.7	0.3	0.0	1.9	1.6	1.0	-8.8	0.00	0.48	1.04	0.79	0.5
03.05.20	3.9	0.5	0.0	9.8	3.0	3.4	-9.2	0.04	0.22	0.88	0.50	0.9
07.07.20	2.4	2.2	0.1	3.9	6.1	6.3	0.3	0.00	0.50	1.00	0.73	1.3
31.10.20	1.8	0.3	0.2	4.9	2.5	1.8	-0.9	0.00	0.48	0.98	0.66	0.7
01.11.20	1.7	0.5	0.2	6.8	2.6	0.6	-1.3	0.42	0.26	0.64	0.46	0.6
22.11.20	0.9	0.5	0.0	4.4	0.9	0.7	-2.3	0.00	0.31	0.96	0.56	0.7
22.12.20	2.2	-0.1	0.0	5.5	0.1	1.8	-7	0.00	0.15	0.35	0.22	0.4
19.01.21	4.1	0.0	0.0	12.8	-0.1	1.6	-13	0.00	0.34	0.64	0.49	1.6
14.02.21	0.7	0.2	0.1	5.7	0.9	0.9	-17	0.00	0.30	0.62	0.40	0.5
14.03.21	0.5	0.5	0.1	1.5	2.3	0.4	-18.8	0.83	0.57	0.90	0.75	1.7
<i>Saharan dust intrusions</i>												
25.01.20	2.4	-0.1	0.5	5.7	1.0	2.1	-11.0	0.00	2.10	4.99	3.01	20.8
03.02.20	1.5	0.9	0.1	4.5	1.9	5.1	-5.9	0.00	1.17	2.31	1.68	8.7
21.03.20	4.8	0.3	0.1	13.9	5.1	9.4	-8.8	0.00	1.24	5.03	2.71	16.5
22.03.20	1.2	0.2	0.0	13.0	9.9	5.6	-9.2	0.00	0.46	5.14	3.60	9.6
08.11.20	2.3	-0.1	0.0	12.7	1.5	6.8	-4.3	0.00	1.37	4.06	2.36	13.8
06.02.21	1.5	1.2	0.4	-25.1	-37.8	116.0	-7.2	0.00	1.15	3.91	2.47	147.1
03.03.21	0.4	0.5	0.1	-0.2	6.8	12.4	-7.1	0.00	1.25	2.40	1.58	61.5



440 **Table A2: Number concentration of ice nucleating particles active at -10 °C (INP<sub>-10</sub>) at Jungfraujoch, determined on randomly selected PM<sub>10</sub> filter samples. Each sample was collected throughout a full calendar day, for which local mean air temperature and the fraction of hours with relative humidity above 95% is shown. The INP data are from the study by Conen et al. (2015) and meteorological data is from MeteoSwiss. Assays of INP that showed no freezing at all at -10 °C were set to a value of 0.01 INP<sub>-10</sub> m<sup>-3</sup> to be displayable on the log-scale in Figure 2.**

date	INP <sub>-10</sub> (m <sup>-3</sup> )	T <sub>mean</sub> (°C)	fraction of hours with RH > 95%	date	INP <sub>-10</sub> (m <sup>-3</sup> )	T <sub>mean</sub> (°C)	fraction of hours with RH > 95%
06.06.12	2.23	-1.7	0.17	04.12.12	1.00	-16.6	0.08
13.06.12	0.78	-8.1	0.54	07.12.12	0.36	-18.1	0.00
16.06.12	6.43	2.2	0.00	08.12.12	0.43	-21.1	0.00
23.06.12	3.61	1.2	0.00	14.12.12	0.50	-9.9	0.00
27.06.12	8.78	1.7	0.17	24.12.12	0.50	-3.7	0.00
30.06.12	6.94	4.4	0.08	28.12.12	3.19	-13.7	0.42
03.07.12	19.50	1.7	0.25	01.01.13	0.71	-10.7	0.21
08.07.12	8.78	0.7	0.08	11.01.13	0.57	-16.2	0.04
11.07.12	3.73	-1.1	0.04	19.01.13	0.09	-9.5	0.04
13.07.12	4.07	-0.9	0.17	20.01.13	0.03	-11.9	0.13
16.07.12	2.05	-5	0.25	22.01.13	0.01	-18.8	0.00
28.07.12	15.38	2	0.17	26.01.13	0.01	-14.2	0.00
01.08.12	5.50	3.2	0.17	01.02.13	0.92	-9.5	0.54
03.08.12	20.74	1.8	0.00	06.02.13	0.85	-21.3	0.96
10.08.12	2.60	-0.8	0.17	09.02.13	0.03	-26.8	0.54
15.08.12	5.95	3	0.17	11.02.13	0.16	-17.4	0.00
18.08.12	3.95	7.4	0.00	21.02.13	0.01	-20	0.13
20.08.12	6.10	5.8	0.13	26.02.13	0.01	-18	0.00
03.09.12	4.55	-0.9	0.21	04.03.13	0.93	-10.6	0.00
15.09.12	2.14	0.6	0.00	06.03.13	0.10	-10.8	0.25
19.09.12	1.80	-7	0.75	13.03.13	0.23	-15.5	0.00
21.09.12	1.39	-0.9	0.04	14.03.13	0.36	-25.4	1.00
25.09.12	12.03	-3.9	0.13	18.03.13	0.10	-16.3	0.00
30.09.12	4.68	-2.9	0.79	31.03.13	0.30	-17.9	0.00
04.10.12	2.32	-4.4	0.29	10.04.13	0.78	-11.8	0.00
11.10.12	2.05	-4.3	0.04	11.04.13	0.50	-6.3	0.00
12.10.12	3.29	-6.1	0.29	14.04.13	1.16	-2.3	0.00
16.10.12	0.64	-8	0.00	19.04.13	2.79	-7.6	0.46
23.10.12	2.14	1.8	0.00	23.04.13	0.36	-4.5	0.00
28.10.12	0.03	-16.5	0.00	29.04.13	0.43	-4.1	0.92
02.11.12	3.51	-10.6	0.00	05.05.13	1.88	-5.9	0.00
10.11.12	9.02	-7.1	0.58	17.05.13	0.86	-5.2	0.50
13.11.12	1.88	-4.3	0.00	22.05.13	0.03	-9.1	0.00
14.11.12	11.31	-2	0.00	25.05.13	0.50	-15.5	0.67
21.11.12	0.23	-5.8	0.00	28.05.13	1.80	-7	0.63
26.11.12	0.86	-6.9	0.63	29.05.13	0.30	-12.4	0.33

445