

# Chemical composition and mixing state of BC-containing particles and the implications on light absorption enhancement

Jiaxing Sun<sup>1,2</sup>, Yele Sun<sup>1,2,3</sup>, Conghui Xie<sup>1,2, a</sup>, Weiqi Xu<sup>1</sup>, Chun Chen<sup>1,2</sup>, Zhe Wang<sup>1</sup>, Lei Li<sup>4</sup>, Xubing  
5 Du<sup>4,5</sup>, Fugui Huang<sup>6</sup>, Yan Li<sup>1,2</sup>, Zhijie Li<sup>1,2</sup>, Xiaole Pan<sup>1</sup>, Nan Ma<sup>7</sup>, Wanyun Xu<sup>8</sup>, Pingqing Fu<sup>9</sup>, and Zifa  
Wang<sup>1,2,3</sup>

<sup>1</sup>State Key Laboratory of Atmospheric Boundary Layer Physics and Atmospheric Chemistry, Institute of Atmospheric Physics,  
Chinese Academy of Sciences, Beijing 100029, China

10 <sup>2</sup>College of Earth and Planetary Sciences, University of Chinese Academy of Sciences, Beijing 100049, China

<sup>3</sup>Center for Excellence in Regional Atmospheric Environment, Institute of Urban Environment, Chinese Academy of Sciences,  
Xiamen 361021, China

<sup>4</sup>Institute of Mass Spectrometer and Atmospheric Environment, Jinan University, Guangzhou, 510632, China

15 <sup>5</sup>Guangdong Provincial Engineering Research Center for On-Line Source Apportionment System of Air Pollution, Guangzhou,  
510632, China

<sup>6</sup>Guangzhou Hexin Analytical Instrument Company Limited, Guangzhou, 510530, China

<sup>7</sup>Institute for Environmental and Climate Research, Jinan University, Guangzhou 511443, China

<sup>8</sup>State Key Laboratory of Severe Weather & Key Laboratory for Atmospheric Chemistry, Institute of Atmospheric  
Composition, Chinese Academy of Meteorological Sciences, Beijing, 100081, China

20 <sup>9</sup>Institute of Surface-Earth System Science, Tianjin University, Tianjin 300072, China

<sup>a</sup>now at: State Key Joint Laboratory of Environmental Simulation and Pollution Control, College of Environmental Sciences  
and Engineering, Peking University, Beijing, 100871, China

Correspondence: Yele Sun ([sunyele@mail.iap.ac.cn](mailto:sunyele@mail.iap.ac.cn))([sunyele@mail.iap.ac.cn](mailto:sunyele@mail.iap.ac.cn))

**Abstract.** The radiative forcing of black carbon (BC) depends strongly on its mixing state in different chemical environments.

25 Here, we analyzed the chemical composition and mixing state of BC-containing particles by using a single particle aerosol mass spectrometer and investigated their impacts on light absorption enhancement ( $E_{\text{abs}}$ ) at an urban (Beijing) and a rural site (Gucheng) in North China Plain. While the BC was dominantly mixed with organic carbon (OC), nitrate and sulfate at both urban and rural sites, the rural site showed much higher fraction of BC coated with OC and nitrate (36% vs. 15—20%). Moreover, the BC mixing state evolved significantly as a function of relative humidity (RH) with largely increased coatings of OC-nitrate and nitrate at high RH levels. By linking with ~~the bulk composition of organic aerosol (OA)~~ composition, we found that the OC coated on BC comprised dominantly secondary OA in Beijing, while primary and secondary OA were similarly important in Gucheng. Furthermore,  $E_{\text{abs}}$  was highly dependent on ~~the~~ secondary inorganic aerosol coated on BC at both sites, while the coated primary OC also resulted in an  $E_{\text{abs}}$  of ~1.2 for relatively fresh BC particles at the rural site. Positive matrix factorization analysis was performed to quantify the impact of different mixing state on  $E_{\text{abs}}$ . Our results showed the small  $E_{\text{abs}}$  (1.06—1.11) for BC particles from fresh primary emissions, while the  $E_{\text{abs}}$  increased significantly above 1.3 when BC was aged rapidly with increased coatings of OC-nitrate or nitrate, and it can reach above 1.4 as sulfate was involved in BC aging.

## 1 Introduction

Black carbon (BC), also referred to soot and element carbon, contributes a substantial and positive impact on climate radiative forcing (Bond et al., 2013; IPCC, 2013). High loading of atmospheric BC could depress the development of planetary boundary layer and aggravate the haze pollution episodes (Ding et al., 2016). However, accurate estimation of light absorption and radiative forcing of BC is still challenging due to its complex emission sources (e.g., fossil fuel and biomass burning) and microphysical properties (e.g., mixing state and coating composition) (Kahnert, 2010; Vignati et al., 2010; Liu et al., 2017; Cappa et al., 2019; [Sun et al., 2022](#)). Light absorption of BC is composed of pure BC absorption and the enhanced light absorption ( $E_{\text{abs}}$ ) induced by "lensing effect", which is due to the chemical materials coated on BC (Fuller et al., 1999; Bond and Bergstrom, 2006; Lack and Cappa, 2010). In order to determine the  $E_{\text{abs}}$ , the thermodenuder (TD)-method (Liu et al., 2015;

Zhang et al., 2016), mass absorption efficiency (MAE)-method (Wang et al., 2014; Zhang et al., 2018) and Mie theoretical calculations (Liu et al., 2017) are widely used in previous studies. However, the  $E_{\text{abs}}$  in different chemical environments is considerably different, such as the negligible  $E_{\text{abs}}$  (1.06) in California while a significant light absorption enhancement (~1.8) in Kanpur (Cappa et al., 2012; Thamban et al., 2017). One explanation is due to the variety of mixing state of BC from different sources, urban and rural background sites, and aging processes (Liu et al., 2015; Liu et al., 2017; [Liu et al., 2020](#)).

Extensive studies have been conducted to characterize the mixing state of BC (Li et al., 2016b). Buseck et al. (2014) used the transmission electron microscopy (TEM) to determine the size, morphology, mixing state and chemical compositions of soot particles, and Li et al. (2016a) used the same technique for characterization of size and aging dependent mixing state of individual particles in both clean and polluted environment in China. More recently, the development of soot particle aerosol mass spectrometer (SP-AMS) is able to measure refractory BC (rBC) and coated aerosol species in real-time (Lee et al., 2015). For example, Xie et al. (2019a) found that secondary organic and inorganic components contributed mostly to the coating materials of rBC in summer, while [Liu et al. \(2019\)](#)[Liu et al. \(2019a\)](#) reported that rBC was more mixed with primary organic components (e.g., coal combustion and biomass burning) than secondary species in winter. Comparatively, more comprehensive mixing state of BC particles and the chemical compositions of coatings could be acquired by aerosol time-of-flight mass spectrometry (ATOFMS) or single particle aerosol mass spectrometer (SPA-MS) (Pratt and Prather, 2012). ~~A recent study~~[Recent studies](#) in winter in North China Plain (NCP) showed that most of BC particles were mixed with sulfate and organic carbon (OC) due to the intensive coal combustion and biomass burning emissions in heating period (Chen et al., 2020b; Wang et al., 2020), while BC particles were more mixed with nitrate in summer in Beijing (Xie et al., 2020). Although the mixing state, coating compositions and  $E_{\text{abs}}$  of BC have been widely discussed in previous studies, our understanding of BC properties in winter particularly the differences between urban and rural areas in NCP are limited.

In this study, we deployed a newly developed high resolution single particle aerosol mass spectrometer (Zhu et al., 2020), and a photoacoustic extinctions (PAX) coupled with a thermodenuder at an urban and a rural site in NCP to investigate the mixing state of BC-containing particles and its relationship with light absorption enhancement. Meanwhile, the aerosol bulk composition was measured simultaneously by a high-resolution aerosol mass spectrometer (HR-AMS). The chemical composition and mixing state of BC-containing particles are characterized, and the evolution of BC mixing state and the

driving factors are investigated. Finally, the impacts of the changes in mixing state of BC on light absorption enhancement and radiative forcing are elucidated.

## 2 Methods

### 75 2.1 Sampling sites and measurements

The measurements were conducted at an urban site from 18 October to 1 December, 2019 and a rural site from 9 December 2019 to 13 January 2020. The urban site is located at the Institute of Atmospheric Physics (IAP), Chinese Academy of Sciences (39°58'28"N, 116°22'16"E) in Beijing (BJ). The rural site of Gucheng (GC) in Hebei province is located ~120 km to the southwest of Beijing. Ambient aerosols with a flow rate of 3 L min<sup>-1</sup> were drawn into a PM<sub>2.5</sub> cyclone (Model: URG-2000-

80 30ED) and a nafion dryer, then, aerosol particles (~2 L min<sup>-1</sup>) were sampled by switching between the thermodenuder (TD) and bypass line (BP) every 15 min into a HR-AMS to measure non-refractory submicron aerosol (NR-PM<sub>1</sub>) species and a PAX (Droplet Measurement Technologies) for absorption [ $b_{\text{abs}}(\lambda)$ ] and scattering [ $b_{\text{sca}}(\lambda)$ ] coefficients at 870 nm. Note that equivalent BC (eBC) is converted from  $b_{\text{abs}}(\lambda)$  into mass concentration with a reference mass absorption efficient (MAE<sub>ref</sub>, 4.44 m<sup>2</sup> g<sup>-1</sup> in this study) (Bond and Bergstrom, 2006).

85 increased gradually from 25°C to 250 °C followed by a 15 min cooling down to 25°C.  ~~$E_{\text{abs}}$  To minimize the uncertainties due to the changes during the measurements of TD and bypass, the  $b_{\text{abs, total}}$  was then calculated obtained from the linear interpolation of measured ambient absorptions adjacent to the TD time, and the  $E_{\text{abs}}$  was then determined as the ratio of ambient BC- $b_{\text{abs, total}}$  to  $b_{\text{abs, BC pure}}$ , that was defined as the thermodenuded particle absorption ( $b_{\text{abs, total}}$ ) to that thermodenuded (considered as pure BC absorption) when temperature was higher than in TD line at  $T > 200^{\circ}\text{C}$  ( $b_{\text{abs, BC pure}}^{\circ}\text{C}$ ). Note that we did not do TD loss~~

90 ~~correction in this study. The reason is that the underestimated  $E_{\text{abs}}$  of ~14% due to the TD loss (Fig. S1) can be relatively offset by the overestimation of ~16% due to the coating materials that did not evaporate at 210°C (Ma et al., 2020; Xu et al., 2021a).~~

More detailed ~~descriptions~~ descriptions of the measurements and methods could be found in ~~our~~ previous study (Sun et al., 2021).

域代码已更改

The mixing state and chemical composition of single particles are detected by a high-resolution single particle aerosol mass spectrometer (HR-SPAMS, Hexin Analytical Instrument Co., Ltd.) was deployed independently in the same room for detecting the mixing state and chemical composition of single particles in ambient atmosphere. The aerodynamic diameters of single particles are determined from the velocity detected by two continuous laser beams (diode Nd: YAG, 532 nm). After passing through the sizing region, particles are desorbed and ionized by a pulsed Nd: YAG laser (266 nm), and the positive and negative fragments are detected by a Z-shaped bipolar time-of-flight mass spectrometer to inform the chemical compositions. A more detailed description of the SPA-MS can be found elsewhere (Li et al., 2011). Compared with the traditional SPA-MS, the new HR-SPAMS uses a new aerosol concentration sampling device and improves the transmission efficiency of coarse particles. Meanwhile, a delayed extraction technology is introduced to improve the mass resolution and increase the hit rate by a factor of 2–4 for ambient particles (Chen et al., 2020c). Also, the ion signals with high and low intensity are separated by multichannel acquisition technology and detected simultaneously, which makes the system dynamic range more than 40 times of the traditional data acquisition system and improves the detection of ions with low signals greatly (Shen et al., 2018; Zhong et al., 2021).

## 2.2 Data analysis

In total, 3-619-0383619038 and 4-655-4264655426 particles were analyzed in BeijingBJ and GuehengGC, respectively. The size and chemical composition of each single particle is informed by Computational Continuation Core (COCO) toolkit in MATLAB software. Based on the marker of  $C_n^{\pm}$  ( $n = 1, 2, 3 \dots$ ) clusters, 2-2692269 659 and 3-399-5653399565 BC-containing particles were identified in BeijingBJ and GuehengGC, respectively. Four typical sources of BC-containing particles are further identified according to the characteristic ion markers: (1) particles containing abundant signals of 39[K]<sup>+</sup>, 45[CHO<sub>2</sub>]<sup>-</sup> and 59[C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>]<sup>-</sup> or 73[C<sub>3</sub>H<sub>5</sub>O<sub>2</sub>]<sup>-</sup> with the peak areas more than 0.5% are classified as BB<sub>pure</sub> type from biomass burning (Silva et al., 1999; Healy et al., 2010); (2) particles containing abundant signals of 7[Li]<sup>+</sup>, 23[Na]<sup>+</sup>, 27[Al]<sup>+</sup>, 43[AlO]<sup>+</sup>, 80[SO<sub>3</sub>]<sup>-</sup>, 97[HSO<sub>4</sub>]<sup>-</sup> and polycyclic aromatic hydrocarbons are classified as CC<sub>pure</sub> type from coal combustion (Zhang et al., 2009; Healy et al., 2010); (3) particles containing abundant signals of 40[Ca]<sup>+</sup>, 51[V]<sup>+</sup>, 55[Mn]<sup>+</sup>, 67[VO]<sup>+</sup>, 46[NO<sub>2</sub>]<sup>-</sup>, 62[NO<sub>3</sub>]<sup>-</sup> and 79[PO<sub>3</sub>]<sup>-</sup> are classified as TR<sub>pure</sub> type from traffic emissions (Yang et al., 2017); (4) particles internally mixed

with more than one source (three sources above) are unified together and named as  $Mix_{Source}$  type. The BC-containing particles above are collectively referred to  $BC_{fresh}$ . The remain BC-containing particles are named  $BC_{aged}$  and classified by using the ART-2a algorithm with a vigilance factor of 0.75, a learning rate of 0.05 and 20 iterations (Song et al., 1999). Seven particle types are grouped and named based on two principles: (1) the particles are named BCOC when the signals of  $37[C_3H]^+$ ,  $43[C_2H_3O]^+$ ,  $51[C_4H_3]^+$  and  $63[C_5H_3]^+$  are comparable with  $C_n^+$  in positive ion area (Healy et al., 2012; Xie et al., 2020), otherwise they are named BC; (2) On the basis of (1), the particles are named  $BCOC_N$  or  $BCOC_S$  when they are only mixed significantly with nitrate ( $46[NO_2]^+$  and  $62[NO_3]^+$ ) or sulfate ( $97[HSO_4]^+$ ). Otherwise, they are named  $BCOC_{NS}$  when they present comparable peak areas of nitrate and sulfate. The more detailed names of BC-containing particle types are shown in Table S1. [According to previous studies, the coating materials on BC-containing particles measured by SPA-MS referred to chemical components that are partially or fully coated on BC \(Bond and Bergstrom, 2006; Healy et al., 2012; Pratt and Prather, 2012; Bi et al., 2015; Chen et al., 2016; Xie et al., 2020\).](#)

The sources of bulk OA [from HR-AMS measurements](#) were analyzed by positive matrix factorization (PMF) and five OA factors were identified at both urban and rural sites, including biomass burning OA (BBOA), fossil fuel-related OA (FFOA), cooking OA (COA), less oxidized oxygenated OA (LO-OOA) and more oxidized OOA (MO-OOA) in Beijing and BBOA, coal combustion OA (CCOA), hydrocarbon-like OA (HOA), OOA and aqueous-related OOA (aq-OOA) in GC [\(Table S1\)](#). The detailed PMF analysis of OA in Beijing and Gucheng are presented by [Xu et al. \(2021\)](#) [Xu et al. \(2021b\)](#) and Chen et al. (2021). The PMF analysis was also performed to identify the effect of different mixing state on  $E_{abs}$ , by inputting  $b_{abs, total}$ ,  $b_{abs, BC_{pure}}$  and 11 major types of BC-containing particles derived from HR-SPAMS. The detailed pre-treatment of error matrix [\(in supplementary\)](#), and the selection of factor solutions could be found in previous studies (Petit et al., 2014; Xie et al., 2019b). Then, the  $E_{abs}$  of each factor can be calculated as the ratio of  $b_{abs, total}$  and  $b_{abs, BC_{pure}}$  in factor  $i$ .

### 3 Results and discussion

#### 3.1 BC-containing particles at urban and rural sites

140 ~~The~~ BC-containing particles ~~on-average~~ accounted for 62% of the total particles in ~~Beijing~~BJ, lower than that in ~~Gucheng~~GC (73%) ~~and yet~~ higher than that in winter ~~Beijing~~ 2018 ~~in Beijing~~ (55%) (Xie et al., 2020), ~~likely due to the higher mass fraction of eBC in this study (9.3% vs. 6.1%).~~ Similarly, a previous winter study in Beijing also found that 60—78% of aerosol particles contained BC (Chen et al., 2020a). According to Figs. 1 and ~~S1S2~~, we found that the mass spectra of BC-containing particles at the two sites are somewhat similar which are both characterized by  $C_n^+$  ( $n = 1-7$ ),  $27[C_2H_3]^+$ ,  $37[C_3H]^+$ ,  
145  $43[C_2H_3O]^+$ ,  $51[C_4H_3]^+$ ,  $63[C_5H_3]^+$ ,  $46[NO_2]^+$ ,  $62[NO_3]^+$  and  $97[HSO_4]^+$  peaks, indicating that BC-containing particles are consistently mixed with OC, nitrate ( $NO_3$ ) and sulfate ( $SO_4$ ) at the rural and urban sites. Comparatively, ~~more than 80% of BC-containing particles are internally mixed with  $SO_4$  in Beijing while those in Gucheng accounted for less than 60%, suggesting that BC particles were more aged at the urban site.~~ ~~more than 80% of BC-containing particles were internally mixed with  $SO_4$  in BJ while those in GC accounted for less than 60%.~~ Considering the higher relative area of secondary organic  
150 ~~fragment of  $43[C_2H_3O]^+$  (Healy et al., 2010; Chen et al., 2019) in BJ (Fig. 1), we concluded that BC particles were likely more aged at the urban site. Another support is much lower primary emissions of biomass burning and coal combustion in BJ than GC (Sun et al., 2020).~~ In addition, approximately 40% of BC-containing particles was mixed with amines (e.g.,  $59[(CH_3)_3N]^+$  and  $74[(C_2H_5)_2NH_2]^+$ ) in GC, which ~~is was~~ twice ~~than~~ that in BJ. One explanation is the more nitrogen-containing compounds formed from either aqueous-phase processing due to higher RH (69% vs. 45%) or biomass burning emissions at the rural site  
155 (Zhang et al., 2012; Chen et al., 2019).

The study in Beijing was divided into two periods, i.e., non-heating period (BJ-NHP) from 28 October to 10 November and heating period (BJ-HP) from 10 November to 1 December, while the observation in ~~Gucheng~~GC was performed during heating period. As illustrated in Fig. 2a, the BC-containing particles are dominantly contributed by  $BC_N$  and  $BCOC_N$  (~20%) during BJ-NHP, indicating that BC was mainly internally mixed with nitrate at the urban site. Compared with BJ-NHP, the fractions  
160 of  $BCOC_S$  and  $Mix_{Source}$  increased significantly during BJ-HP, especially during relatively clean periods, suggesting a considerable change in BC mixing state from non-heating to heating period due to the enhanced primary emissions, e.g., coal

combustion. Previous studies showed that BC was mainly mixed with sulfate in winter in Beijing (Chen et al., 2020b; Xie et al., 2020), while this study showed a dominant mixing of BC with nitrate. This was likely due to the fact that coal fuels in Beijing were replaced by clean energies, e.g., natural gas and electricity after 2017 (Zhang et al., 2019). Indeed, the changes in nitrate concentrations were relatively small in winter in Beijing since clean air action although the sulfate concentrations showed large decreases (Zhou et al., 2019; Lei et al., 2020). Comparatively, BCOC<sub>N</sub> was the major BC-containing particle type accounting for 36% in GC, which was twice ~~than~~ that in BJ-HP indicating that BC particles were dominantly mixed with OC and nitrate in an environment with high RH and intensive primary emissions, e.g., coal combustion emissions. In addition, BB<sub>pure</sub> and TR<sub>pure</sub> showed pronounced diurnal cycles in GC (Fig. S2S3) compared with the relatively flat diurnal variations of BB<sub>pure</sub> in BJ, suggesting intensive biomass burning and diesel vehicle emissions at the rural site especially at nighttime.

### 3.2 Chemical composition and mixing state in different environment

Figure 3 shows the variations of number fractions of different BC-containing types under different RH levels in BJ and GC. Almost all BC<sub>aged</sub> types showed strong dependence on RH while the number fractions of BC<sub>fresh</sub> types decreased with increasing RH at both sites indicating that high RH environment was more favorable for BC aging (Zhang et al., 2021)(Zhang et al., 2021). Similar to BC<sub>fresh</sub>, BCOC<sub>S</sub> was the only type of aged BC showing decreased fraction as a function of RH in BJ and GC. In fact, the high correlations between BCOC<sub>S</sub> and CC<sub>pure</sub> ( $R^2 = 0.89$  and  $0.98$ , in BJ and GC, respectively) highlight that BC emitted from coal combustion could be directly mixed with OC and SO<sub>4</sub> at low RH level, and ~~evolve~~evolved towards the mixing with OC and NO<sub>3</sub> under high RH conditions. Moreover, the number fraction of BC<sub>N</sub> increased gradually as the increase of RH and dominated BC particles (~~~30%~~at RH=90–100% in BJ, ~~suggesting~~(25–30%) at RH = 70–100% in BJ. Considering the similar increases of PM (NR-PM<sub>1</sub> + eBC) as a function of RH (Figs. 3a and S4), the RH dependence of BC<sub>N</sub> suggested that the newly formed nitrate that coated on fresh BC played an important role in the formation of severe pollution in urban region. Comparatively, the number fraction of BCOC<sub>N</sub> increased the most by 43%, accounting for more than half of BC-containing particles at high RH and PM levels in GC. This result indicates that the type of BCOC<sub>N</sub> was more important to aggravate air pollution in rural area. ~~In addition, we, supported by the relatively high correlation between the number fraction of BCOC<sub>N</sub> and PM (Fig. S4). In addition, BCOC<sub>N</sub> was likely affected by the photochemical production in GC, which is supported by the~~



significantly increased number counts and fraction of  $\text{BCOC}_{\text{NS}}$  during daytime (Fig. S3). We found that the fraction of  $\text{BCOC}_{\text{NS}}$  decreased obviously as a function of RH in GC indicating the impact of the transition from photochemical production to aqueous-phase reactions on the mixing state of BC. This result also suggests ~~Considering the increased  $\text{SO}_4$  mass fraction yet the relatively stable  $E_{\text{abs}}$  at RH > 70% (Sun et al., 2021), we inferred~~ that aqueous-phase formation of sulfate at high RH level appeared not to affect BC mixing state substantially, consistent with previous ~~studies that aqueous-phase processing did not affect the total  $E_{\text{abs}}$  in GC (Sun et al., 2021; Zhang et al., 2021) study (Zhang et al., 2021)~~. However,  $\text{BCOC}_{\text{NS}}$  in BJ showed relatively stable fractions across different RH levels suggesting the different sources from the rural site.

~~Figure 3 shows~~ Figs. 3c and 3d show the evolution of mixing state of BC-containing particles during two different haze events in ~~Beijing~~BJ. During the initial stage of haze ~~episode~~case1 (P0, Fig. 3c), the contribution of  $\text{BCOC}_{\text{S}}$  started to decrease while the number fraction of  $\text{BCOC}_{\text{N}}$  increased significantly. As a consequence,  $E_{\text{abs}}$  increased rapidly from ~1.1 to 1.3 in half day. Then, the number fraction of  $\text{BC}_{\text{N}}$  increased while that of  $\text{BC}_{\text{fresh}}$  decreased during P1 period. These results indicated that fresh BC was gradually aged by mixing with nitrate during the evolution of haze episode.  $\text{BC}_{\text{N}}$  increased continually during P2 with high  $E_{\text{abs}}$  up to 1.4, and finally the mixing state of BC was stabilized as indicated by the relatively stable number fractions of most BC particle types and small changes in  $E_{\text{abs}}$ . Similar to haze ~~episode 1, the freshly emitted~~case1, BC was ~~gradually~~mixed with nitrate and OC causing high  $E_{\text{abs}}$  (up to 1.5) during P3 and P4. (Fig. 3d). As shown in ~~Fig~~Figs. 3d and S5, the sources of fine particles were dominated by fossil fuel OA and presented strong diurnal variations ~~due to the influences of mountain valley winds. As a result,  $E_{\text{abs}}$  also presented a pronounced diurnal cycle with higher values during daytime, indicating that the different air masses brought by the mountain valley winds affected BC mixing state and light absorption enhancement, consistent with changes in wind direction due to the influences of mountain valley winds (Sun et al., 2016). As a result,  $E_{\text{abs}}$  also presented a relatively consistent variation with FFOA and showed higher values at nighttime, indicating that the different chemical composition and mixing state associated with the changes in air masses due to the mountain valley winds had affected the light absorption enhancement of BC (Ding et al., 2021)~~. The number fraction of  $\text{BC}_{\text{N}}$  increased significantly during the severest polluted period (P5) associated with simultaneous increases in BBOA, indicating the mixing of BC from biomass burning with nitrate under high RH and PM levels, and hence  $E_{\text{abs}}$  was comparably high (1.35). After the P5 period, ~~(from 21:00 on 9 November to 0:00 on 10 November), the BBOA was relatively stable~~decreased slightly and FFOA

increased significantly. Correspondently, the proportion of BCOC<sub>N</sub> in BC is correspondingly higher, indicating. Similar variation could also be found during 14:00–19:00 on 8 November. These results indicated that fresh BC emitted from fossil fuel emission was likely mixed with OC and nitrate at high PM level. Different from Beijing, the evolution of BC mixing state during most haze events were similar in GC (Figs. 2e Fig. S3 and S32c), which was characterized by most significant increase in BCOC<sub>N</sub> rather BC<sub>N</sub> as the PM increased. Moreover, BCOC<sub>N</sub> particles were increased more significantly during nighttime while BCOC<sub>NS</sub> was more significant during daytime. Such differences were mainly due to the enhanced coal combustion pollutants at nighttime (Fig. S2S3) which were mixed with photochemical products during daytime.

Overall, our results suggest that the fresh BC particles from biomass burning emissions are more directly mixed with nitrate under high RH conditions, and then mixed with more sulfate during further aging. Comparatively, the fresh BC particles from coal combustion are often mixed with OC and sulfate first, and then mix further with OC and nitrate at high RH level. In addition, our results also demonstrate that the  $E_{\text{abs}}$  in GC was largely due to coal combustion emissions internally mixed with OC, consistent with our previous study showing a large impact of coal combustion emissions on  $E_{\text{abs}}$  (Sun et al., 2021) (Sun et al., 2021).

### 3.3 Effects of chemical composition on $E_{\text{abs}}$

As shown in the average positive mass spectra of total BC-containing particles (Fig. S2), the peak areas of  $C_n^+$ , OC and metal contributed more than 95% to the total peak area, while the peak areas of  $\text{NO}_3$  ( $46[\text{NO}_2^+]$  and  $62[\text{NO}_3^+]$ ) and  $\text{SO}_4$  ( $97[\text{HSO}_4^+]$ ) accounted for more than 80% in the negative mass spectra. To better characterize the relationship between chemical species and  $E_{\text{abs}}$ , we summed  $C_n^{\pm}$  ( $n = 1-5$ , accounting for more than 99% in  $C_n$ ) peak areas to represent BC and the total of  $\text{NO}_3$  and  $\text{SO}_4$  peak areas to represent secondary inorganic components coated on BC. In addition, the sum of positive peak areas except  $C_n^+$  was defined as OC + Metal to represent the OC and metal components coated on BC. These peak areas covered almost all of the chemical components coated on BC in total BC-containing particles.

Figures 4a and 4b show the relationship between peak area ratios (measured by HR-SPAMS) and mass concentration ratios (measured by HR-AMS) in BeijingBJ and GuchengGC, respectively. As the increase of  $(\text{NO}_3 + \text{SO}_4)_{\text{AMS}}/\text{eBC}$  mass concentration ratio, the  $(\text{NO}_3 + \text{SO}_4)/C_n$  peak area ratio increased first and then gradually became stable at both sites. These

235 results indicated that BC was rapidly aged and internally mixed with secondary inorganic components during the early stage of haze episode, and appeared to be fully aged when the ratio of mixing efficiency of NO<sub>3</sub> and SO<sub>4</sub> to BC reached a maximum, i.e., (NO<sub>3</sub> + SO<sub>4</sub>)<sub>AMS</sub>/eBC exceeded ~6. Different from secondary inorganic species, the peak area ratio of (OC + Metal)/C<sub>n</sub> showed a high dependence on the mass concentration ratio of POA (e.g., the sum of BBOA and FFOA in BJ and the sum of BBOA, CCOA and HOA in GC) to eBC at both sites. These results indicated that the POA co-emitted with BC was more easily internally mixed with BC than secondary inorganic components (e.g., SO<sub>4</sub> and NO<sub>3</sub>) although the mass concentration was much lower than SIA at both urban and rural sites. Moreover, the mass concentration ratio of SOA/eBC also presented a high correlation with (OC + Metal)/C<sub>n</sub> in GC (R<sup>2</sup> = 0.81) and BJ (R<sup>2</sup> = 0.81) and GC (R<sup>2</sup> = 0.95, Fig. S7). We then used multiple linear regression analysis to quantify the impacts of POA and SOA on BC-OC coated OC on BC. Our results showed that the average contribution of SOA to the coated OC was nearly twice that of POA (65% vs. 35%) in Beijing BJ, while POA and SOA contributed similarly in Gucheng GC.

$$y = 0.93 * (1 - e^{-0.62*x}) \quad (1)$$

$$y = 0.85 * (1 - e^{-0.71*x}) \quad (2)$$

$$\frac{OC+Metal}{C_n} = 0.61 * \frac{POA}{eBC} + 0.60 * \frac{SOA}{eBC} \quad (3)$$

$$\frac{OC+Metal}{C_n} = 0.79 * \frac{POA}{eBC} + 0.41 * \frac{SOA}{eBC} \quad (4)$$

250 Figures 4c and 4d show the relationship between E<sub>abs</sub> and the ratio of coating materials to C<sub>n</sub> in Beijing and Gucheng, respectively. Light absorption enhancement showed strong dependence on (NO<sub>3</sub> + SO<sub>4</sub>)/C<sub>n</sub> at both sites yet the changes in E<sub>abs</sub> appeared independent of (OC + Metal)/C<sub>n</sub>. Figures 4c and 4d show the relationship between E<sub>abs</sub> and the ratio of mixing materials to C<sub>n</sub> in BJ and GC, respectively. Light absorption enhancement showed a strong dependence on (NO<sub>3</sub> + SO<sub>4</sub>)/C<sub>n</sub> at both sites. Previous studies found that SOA played an important role in BC absorption enhancement (Liu et al., 2019b), whereas in our study, the changes in E<sub>abs</sub> seemed to be independent on (OC + metal)/C<sub>n</sub>, likely because (OC + metal)/C<sub>n</sub> was influenced by both primary and secondary factors. As shown in Fig. 4c, the ratio of (OC + Metal)/C<sub>n</sub> still presented high values at E<sub>abs</sub> = ~1, while the secondary species coated on BC were negligible. These results suggested OC and metals were likely either filled

internal void spaces of fresh BC or mainly **externallypartly** mixed with BC which did not induce light absorption enhancement at the urban site. As the progress of aging,  $E_{\text{abs}}$  increased significantly mainly due to the increased secondary coating materials.

260 Similar to **BeijingBJ**,  $E_{\text{abs}}$  also increased significantly as a function of  $(\text{NO}_3 + \text{SO}_4)/C_n$  in **GuehengGC**. The difference is the high background  $E_{\text{abs}}$  of  $\sim 1.20$  in GC when the  $(\text{NO}_3 + \text{SO}_4)/C_n$  ratio was closed to 0. **Combined-with-the-peak-area-ratio-of**  
**(OC + Metal)/ $C_n$  was about 2.5 at that time, we found that OC and metals were not only as filler materials but also**  
**Pervious**  
**study showed that  $E_{\text{abs}}$  is  $> 1$  when the non-BC material is sufficient to encapsulate the BC (Liu et al., 2017). Considering the**  
**peak area ratio of  $(\text{OC} + \text{Metal})/C_n$  was  $\sim 2.5$ , we inferred that OC and metals were not only as filler materials but also likely**

265 coated on fresh BC and induced light absorption enhancement at the rural site. After the further aging process in atmosphere,  $E_{\text{abs}}$  was mainly due to the increased secondary inorganic components coated on BC. **Considering-that-the-measurements-of**  
**BC-mixing-state-by-SPAMS-are-often-not-available-in-field-campaignsMoreover**, we predicted the  $E_{\text{abs}}$  by using Eqs. 1–6**the**  
**statistical equations in Fig. 4** with aerosol species measured by AMS. **The-predicted-We found that the estimated  $E_{\text{abs}}$  showed**  
**overall agreements with the measured values in both BJ and GC, yet the variations at specific NR-PM<sub>10</sub>/eBC. Although the**

270 **correlation was not significant (Fig. S8), the average measured and estimated  $E_{\text{abs}}$  values were much smaller (Fig. S4)-similar**  
**which are  $1.21 (\pm 0.12)$  and  $1.22 (\pm 0.18)$ , respectively in BJ, and  $1.31 (\pm 0.15)$  and  $1.25 (\pm 0.07)$ , respectively in GC. Also, the**  
**uncertainty estimated from the difference of measured and predicted was overall below 10% in both BJ and GC indicating that**  
**the approach is reasonably well to estimate  $E_{\text{abs}}$ .**

We also estimated the  $E_{\text{abs}}$  in summer 2017 using the same method and compared with the measurements by a TD-CAPS (Fig. S4S9). Our results showed that the average  $E_{\text{abs}}$  in summer was 1.24,  
275 which was close to the average about 1.2 reported by Liu et al. (2019), yet higherLiu et al. (2019a), yet lower than that in Xie et al. (2019a).

$$E_{\text{abs}} = 0.29 * \frac{\text{NO}_3 + \text{SO}_4}{C_n} + 0.02 * \frac{\text{OC} + \text{Metal}}{C_n} + 1 \quad (5)$$

$$E_{\text{abs}} = 0.15 * \frac{\text{NO}_3 + \text{SO}_4}{C_n} + 0.08 * \frac{\text{OC} + \text{Metal}}{C_n} + 1 \quad (6)$$

### 3.4 Effects of mixing state on $E_{\text{abs}}$

280 The PMF analysis is used to characterize the effects of different mixing state on  $E_{\text{abs}}$  (Fig. 5). Five and four factors were identified in BJ and GC, respectively to elaborate the influence of different mixing state on  $E_{\text{abs}}$  (Fig. 5). Note that  $E_{\text{abs}}$  was not estimated when the factor contributed negligibly to the total BC, such as Factor5 in BJ (Fig. 5c). As illustrated in Fig. 5b, Factor2 is the major type of aged BC in the urban region, accounting for more than 60% of the total BC. This factor was dominated by  $\text{BC}_\text{N}$ ,  $\text{BCOC}_\text{N}$ ,  $\text{BCOC}_\text{NS}$  and  $\text{BC}_\text{NS}$ , and presented a high  $E_{\text{abs}}$  of 1.38. Comparatively, FactorB (Fig. 5g) is the major type of aged BC in the rural area which was dominated by  $\text{BCOC}_\text{N}$ . The  $E_{\text{abs}}$  was  $\sim 1.35$  for this factor, which was comparable to that in BJ. As this factor evolved towards FactorA (Fig. 5f) after further aging and internally mixed with a large amount of sulfate,  $E_{\text{abs}}$  was increased up to 1.41. In BeijingBJ, the relatively fresh traffic emissions are dominant in Factor4 (Fig. 5d) which showed a negligible impact on light absorption enhancement ( $\sim 1.06$ ), consistent with the results in previous studies (Liu et al., 2017; Sun et al., 2021). Compared with traffic emissions, the relatively fresh biomass burning and coal combustion emissions (Factor3 Fig. 5c) comprising mainly  $\text{Mix}_{\text{source}}$  and  $\text{BCOC}_\text{S}$  showed a moderate  $E_{\text{abs}}$  ( $\sim 1.11$ ) in BJ. Although the mixed fresh primary emissions (FactorD Fig. 5i) in GC presented a relatively low  $E_{\text{abs}}$  (1.06), we found that the FactorC from coal combustion emissions and mixed with much OC and nitrate showed a much higher  $E_{\text{abs}}$  ( $\sim 1.31$ ) than that in BJ (Factor3 Fig. 5c). After more aging, more sulfate could be internally mixed with BC and enhanced the  $E_{\text{abs}}$ , such as Factor1 (Fig. 5a) with  $E_{\text{abs}}$  up to 1.42. Overall,  $E_{\text{abs}}$  shows a similar dependence on the evolution of mixing state of BC-containing particles at urban and rural sites, i.e., fresh BC particles from primary emissions (e.g., biomass burning, coal combustion and traffic) showed the small  $E_{\text{abs}}$  (1.06–1.11). At relatively high RH level, BC could be directly mixed with nitrate or OC-nitrate ( $\text{BC}_\text{N}$  and  $\text{BCOC}_\text{N}$ ), accounting for more than 60% of BC, and led to an increase in  $E_{\text{abs}}$  above 1.30; and then, the BC-containing particles were further mixed with sulfate ( $\text{BC}_\text{NS}$  and  $\text{BCOC}_\text{NS}$ ) after continuous aging in atmosphere, and resulted in the highest  $E_{\text{abs}}$  above 1.40.

290

295

300 Based on  $E_{\text{abs}}$  for each factor and its contribution to  $b_{\text{abs, BCpure}}$ , we estimated the direct radiative forcing ( $\Delta F_\text{R}$ ) caused by BC-containing particles with their mixing state pure BC at the top-of-the atmosphere (TOA) and the absorption  $\Delta F_\text{g}$  enhanced by the mixed state of BC-containing particles (Chylek and Wong, 1995; Chen and Bond, 2010). The detail descriptions for the

estimation are presented in supplementary. ~~It should be noted that BrC can also absorb light at 870 nm, leading to an overestimation of BC absorption. Considering that the contribution of BrC to the total absorption at 870 nm is typically small (< 1%) (Clarke et al., 2004; Fialho et al., 2005; Yang et al., 2009), the impact of BrC on the estimation of radiative forcing of BC is expected to be small as well.~~ As shown in Fig. 5, the  $\Delta F_R$  caused by pure BC particles is about  $+0.43 \text{ W m}^{-2}$  and  $+0.60 \text{ W m}^{-2}$  in Beijing and Gucheng, respectively. Considering the mixing state of BC,  $\Delta F_R$  could increase to  $0.59 \text{ W m}^{-2}$  and  $0.82 \text{ W m}^{-2}$  in Beijing and Gucheng, respectively. Our results demonstrated that the mixing state of BC-containing particles can have a large impact on radiative forcing estimation by up to 27%.

#### 310 4 Conclusions

HR-SPAMS, TD-PAX and HR-AMS were deployed at the urban and rural sites in North China Plain in winter 2019 to characterize the chemical composition and mixing state of BC-containing particles and their impacts on light absorption enhancement. Our results showed that BC-containing particles were primarily mixed with OC,  $\text{NO}_3$  and  $\text{SO}_4$  at both sites while the rural site showed much higher fraction of OC and nitrate coated BC (36% vs. 15—20%). The increased BC particles internally mixed with large amount of  $\text{NO}_3$  was mainly due to the effect of clean air action that reduced much more sulfate than nitrate in  $\text{PM}_{2.5}$ . The average contribution of SOA to the OC coated on BC was about twice ~~than~~ that of POA in BJ, while both of them contributed similarly to the OC coating in GC. In addition, OC and metals were likely as filler materials internally mixed with fresh BC and did not induce much light absorption enhancement at the urban site, while they were coated on fresh BC and induced light absorption enhancement ( $\sim 1.2$ ) at the rural site.

320 By ~~analyzing~~~~analysing~~ the variations of BC mixing state in different environments, we found that BC particles were primarily mixed with  $\text{NO}_3$  as the increase of RH at both urban and rural sites. In particular, the BC emitted from biomass burning was first mixed with  $\text{NO}_3$  at relatively high RH level and then mixed with both nitrate and sulfate after further aging. Comparatively, the BC emitted from coal combustion was more internally mixed with OC and  $\text{NO}_3$ , and then mixed with sulfate ( $\text{BCOC}_{\text{NS}}$ ) with similar processes. Thus, ~~higher~~~~high~~ coal combustion emissions can result in the increase ~~in the of~~  $\text{BCOC}_{\text{N}}$  fraction ~~of~~  $\text{BCOC}_{\text{N}}$  as the increase of RH at the rural site, while  $\text{BC}_{\text{N}}$  was generally the dominant BC-containing particles in Beijing.

Although BC particles presented different mixing state in different environments,  $E_{\text{abs}}$  showed a similar evolutionary dependence on the changes in mixing state, i.e., from the small  $E_{\text{abs}}$  (1.06–1.11) with fresh BC emissions to above 1.30 after [agedaging](#) and internally mixed with nitrate and OC-nitrate (BC<sub>N</sub> and BCOC<sub>N</sub>), and to above 1.40 after further aging with the sulfate involved.

330 **Data availability.** The data in this study are available from the authors upon request ([sunyele@mail.iap.ac.cn](mailto:sunyele@mail.iap.ac.cn)).

**Author contributions.** YS and JS designed the research. JS, CX, WX, CC and ZL conducted the measurements. JS, WX, CC, ZW and YL analyzed the data. CX, LL, XD, FH, XP, NM, WX, PF and ZiW reviewed and commented on the paper. JS and YS wrote the paper.

**Competing interests.** The authors declare that they have no conflict of interest.

335 **Acknowledgements.** This work was supported by the National Natural Science Foundation of China (9204430003, 42061134008).

## References

- 340 [Bi, X. H., Dai, S. H., Zhang, G. H., Qiu, N., Li, M., Wang, X. M., Chen, D. H., Peng, P. A., Sheng, G. Y., Fu, J. M., and Zhou, Z.: Real-time and single-particle volatility of elemental carbon containing particles in the urban area of Pearl River Delta region, China, Atmos. Environ., 118, 194-202, 10.1016/j.atmosenv.2015.08.012, 2015.](#)
- Bond, T. C. and Bergstrom, R. W.: Light absorption by carbonaceous particles: an investigative review, *Aerosol Sci. Technol.*, 40, 27-67, 10.1080/02786820500421521, 2006.
- Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., 345 Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., P., S. J., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: a scientific assessment, *J. Geophys. Res.-Atmos.*, 118, 5380-5552, 10.1002/jgrd.50171, 2013.
- Buseck, P. R., Adachi, K., Andras, G., Tompa, E., and Mihaly, P.: Ns-Soot: a material-based term for strongly light-absorbing carbonaceous particles, *Aerosol Sci. Technol.*, 48, 777-788, 10.1080/02786826.2014.919374, 2014.
- 350 Cappa, C. D., Onasch, T. B., Massoli, P., Worsnop, D. R., Bates, T. S., Cross, E. S., Davidovits, P., Hakala, J., Hayden, K. L., and Jobson, B. T.: Radiative absorption enhancements due to the mixing state of atmospheric black carbon, *Science*, 337, 6098, 10.1126/science.1223447, 2012.

- Cappa, C. D., Zhang, X., Russell, L. M., Collier, S., Lee, A. K. Y., Chen, C.-L., Betha, R., Chen, S., Liu, J., Price, D. J., Sanchez, K. J., McMeeking, G. R., Williams, L. R., Onasch, T. B., Worsnop, D. R., Abbatt, J., and Zhang, Q.: Light absorption by ambient black and brown carbon and its dependence on black carbon coating state for two California, USA, cities in winter and summer, *J. Geophys. Res.-Atmos.*, 124, 1550-1577, 10.1029/2018jd029501, 2019.
- 355 Chen, C., Qiu, Y., Weiqi, X., He, Y., Li, Z., Sun, J., Ma, N., Xu, W., Pan, X., Fu, P., Wang, Z., and Sun, Y.: Primary emissions and secondary aerosol processing during wintertime in rural area of North China Plain, *J. Geophys. Res.-Atmos.*, <under review>, 2021.
- 360 Chen, L., Zhang, F., Yan, P., Wang, X., Sun, L., Li, Y., Zhang, X., Sun, Y., and Li, Z.: The large proportion of black carbon (BC)-containing aerosols in the urban atmosphere, *Environ. Pollut.*, 263, 114507, 10.1016/j.envpol.2020.114507, 2020a.
- Chen, Y. and Bond, T. C.: Light absorption by organic carbon from wood combustion, *Atmos. Chem. Phys.*, 10, 1773-1787, 2010.
- Chen, Y., [Cao, J., Huang, R., Yang, F., Wang, Q., and Wang, Y.: Characterization, mixing state, and evolution of urban single particles in Xi'an \(China\) during wintertime haze days, \*Sci. Total Environ.\*, 573, 937-945, 10.1016/j.scitotenv.2016.08.151, 2016.](#)
- 365 [Chen, Y.,](#) Tian, M., Huang, R.-J., Shi, G., Wang, H., Peng, C., Cao, J., Wang, Q., Zhang, S., Guo, D., Zhang, L., and Yang, F.: Characterization of urban amine-containing particles in southwestern China: seasonal variation, source, and processing, *Atmos. Chem. Phys.*, 19, 3245-3255, 10.5194/acp-19-3245-2019, 2019.
- 370 Chen, Y., Cai, J., Wang, Z., Peng, C., Yao, X., Tian, M., Han, Y., Shi, G., Shi, Z., Liu, Y., Yang, X., Zheng, M., Zhu, T., He, K., Zhang, Q., and Yang, F.: Simultaneous measurements of urban and rural particles in Beijing – Part 1: chemical composition and mixing state, *Atmos. Chem. Phys.*, 20, 9231-9247, 10.5194/acp-20-9231-2020, 2020b.
- Chen, Y., Kozlovskiy, V., Du, X., Lv, J., Nikiforov, S., Yu, J., Kolosov, A., Gao, W., Zhou, Z., Huang, Z., and Li, L.: Increase of the particle hit rate in a laser single-particle mass spectrometer by pulse delayed extraction technology, *Atmos. Meas. Tech.*, 13, 941-949, 10.5194/amt-13-941-2020, 2020c.
- 375 Chylek, P. and Wong, J.: Effect of aerosols on global budget, *Geophys. Res. Lett.*, 22, 929-931, 1995.
- [Clarke, A. D., Shinozuka, Y., Kapustin, V. N., Howell, S., Huebert, B., Doherty, S., Anderson, T., Covert, D., Anderson, J., Hua, X., Moore, K. G., McNaughton, C., Carmichael, G., and Weber, R.: Size distributions and mixtures of dust and black carbon aerosol in Asian outflow: physiochemistry and optical properties, \*J. Geophys. Res.-Atmos.\*, 109, e2003jd004378, 10.1029/2003jd004378, 2004.](#)
- 380 Ding, A. J., Huang, X., Nie, W., Sun, J. N., Kerminen, V. M., Petäjä, T., Su, H., Cheng, Y. F., Yang, X. Q., Wang, M. H., Chi, X. G., Wang, J. P., Virkkula, A., Guo, W. D., Yuan, J., Wang, S. Y., Zhang, R. J., Wu, Y. F., Song, Y., Zhu, T., Zilitinkevich, S., Kulmala, M., and Fu, C. B.: Enhanced haze pollution by black carbon in megacities in China, *Geophys. Res. Lett.*, 43, 2873-2879, 10.1002/2016gl067745, 2016.
- 385 [Ding, S., Liu, D., Hu, K., Zhao, D., Tian, P., Wang, F., Li, R., Chen, Y., He, H., Huang, M., and Ding, D.: Optical and hygroscopic properties of black carbon influenced by particle microphysics at the top of the anthropogenically polluted boundary layer, \*Atmos. Chem. Phys.\*, 21, 681-694, 10.5194/acp-21-681-2021, 2021.](#)




- 390 [Fialho, P., Hansen, A. D. A., and Honrath, R. E.: Absorption coefficients by aerosols in remote areas: a new approach to decouple dust and black carbon absorption coefficients using seven-wavelength Aethalometer data, \*J. Aerosol. Sci.\*, \*\*36\*\*, 267-282, 10.1016/j.jaerosci.2004.09.004, 2005.](#)
- Fuller, K. A., Malm, W. C., and Kreidenweis, S. M.: Effects of mixing on extinction by carbonaceous particles, *J. Geophys. Res.*, **104**, 15941-15954, 10.1029/1998jd100069, 1999.
- 395 Healy, R. M., Hellebust, S., Kourtchev, I., Allan, A., O'Connor, I. P., Bell, J. M., Healy, D. A., Sodeau, J. R., and Wenger, J. C.: Source apportionment of PM<sub>2.5</sub> in Cork Harbour, Ireland using a combination of single particle mass spectrometry and quantitative semi-continuous measurements, *Atmos. Chem. Phys.*, **10**, 9593-9613, 10.5194/acp-10-9593-2010, 2010.
- 400 Healy, R. M., Sciare, J., Poulain, L., Kamili, K., Merkel, M., Müller, T., Wiedensohler, A., Eckhardt, S., Stohl, A., Sarda-Estève, R., McGillicuddy, E., and Connor, I. P., Sodeau, J. R., and Wenger, J. C.: Sources and mixing state of size-resolved elemental carbon particles in a European megacity: Paris, *Atmos. Chem. Phys.*, **12**, 1681-1700, 10.5194/acp-12-1681-2012, 2012.
- IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp, doi:10.1017/CBO9781107415324, 2013.
- 405 Kahnert, M.: On the discrepancy between modeled and measured mass absorption cross sections of light absorbing carbon aerosols, *Aerosol Sci. Technol.*, **44**, 453-460, 10.1080/02786821003733834, 2010.
- Lack, D. A. and Cappa, C. D.: Impact of brown and clear carbon on light absorption enhancement, single scatter albedo and absorption wavelength dependence of black carbon, *Atmos. Chem. Phys.*, **10**, 4207-4220, 10.5194/acp-10-4207-2010, 2010.
- 410 Lee, A. K. Y., Willis, M. D., Healy, R. M., Onasch, T. B., and Abbatt, J. P. D.: Mixing state of carbonaceous aerosol in an urban environment: single particle characterization using the soot particle aerosol mass spectrometer (SP-AMS), *Atmos. Chem. Phys.*, **15**, 1823-1841, 10.5194/acp-15-1823-2015, 2015.
- Lei, L., Zhou, W., Chen, C., He, Y., Li, Z., Sun, J., Tang, X., Fu, P., Wang, Z., and Sun, Y.: Long-term characterization of aerosol chemistry in cold season from 2013 to 2020 in Beijing, China, *Environ. Pollut.*, **268**, 115952, 10.1016/j.envpol.2020.115952, 2020.
- 415 Li, L., Huang, Z., Dong, J., Li, M., Gao, W., Nian, H., Fu, Z., Zhang, G., Bi, X., Cheng, P., and Zhou, Z.: Real time bipolar time-of-flight mass spectrometer for analyzing single aerosol particles, *Int. J. Mass Spectrom.*, **303**, 118-124, 10.1016/j.ijms.2011.01.017, 2011.
- Li, W., Sun, J., Xu, L., Shi, Z., Riemer, N., Sun, Y., Fu, P., Zhang, J., Lin, Y., Wang, X., Shao, L., Chen, J., Zhang, X., Wang, Z., and Wang, W.: A conceptual framework for mixing structures in individual aerosol particles, *J. Geophys. Res.-Atmos.*, **121**, 784-713, 798, 10.1002/2016jd025252, 2016a.
- 420 Li, W. J., Shao, L. Y., Zhang, D. Z., Ro, C. U., Hu, M., Bi, X. H., Geng, H., Matsuki, A., Niu, H. Y., and Chen, J. M.: A review of single aerosol particle studies in the atmosphere of East Asia: morphology, mixing state, source, and heterogeneous reactions, *J. Clean Prod.*, **112**, 1330-1349, 10.1016/j.jclepro.2015.04.050, 2016b.

- 425 Liu, D., Whitehead, J., Alfarra, M. R., Reyes-Villegas, E., Spracklen, Dominick V., Reddington, Carly L., Kong, S., Williams, Paul I., Ting, Y.-C., Haslett, S., Taylor, Jonathan W., Flynn, Michael J., Morgan, William T., McFiggans, G., Coe, H., and Allan, James D.: Black-carbon absorption enhancement in the atmosphere determined by particle mixing state, *Nat. Geosci.*, 10, 184-188, 10.1038/ngeo2901, 2017.
- Liu, D., Joshi, R., Wang, J., Yu, C., Allan, J. D., Coe, H., Flynn, M. J., Xie, C., Lee, J., Squires, F., Kotthaus, S., Grimmond, S., Ge, X., Sun, Y., and Fu, P.: Contrasting physical properties of black carbon in urban Beijing between winter and summer, *Atmos. Chem. Phys.*, 19, 6749-6769, 10.5194/acp-19-6749-2019, [20192019a](#).
- 430 [Liu, D., Zhao, D., Xie, Z., Yu, C., Chen, Y., Tian, P., Ding, S., Hu, K., Lowe, D., Liu, Q., Zhou, W., Wang, F., Sheng, J., Kong, S., Hu, D., Wang, Z., Huang, M., and Ding, D.: Enhanced heating rate of black carbon above the planetary boundary layer over megacities in summertime, \*Environ. Res. Lett.\*, 14, 124003, 10.1088/1748-9326/ab4872, 2019b.](#)
- 435 [Liu, D., Ding, S., Zhao, D., Hu, K., Yu, C., Hu, D., Wu, Y., Zhou, C., Tian, P., Liu, Q., Wu, Y., Zhang, J., Kong, S., Huang, M., and Ding, D.: Black carbon emission and wet scavenging from surface to the top of boundary layer over Beijing region, \*J. Geophys. Res.-Atmos.\*, 125, e2020JD033096, 10.1029/2020jd033096, 2020.](#)
- Liu, S., Aiken, A. C., Gorkowski, K., Dubey, M. K., Cappa, C. D., Williams, L. R., Herndon, S. C., Massoli, P., Fortner, E., and Chhabra, P. S.: Enhanced light absorption by mixed source black and brown carbon particles in UK winter, *Nat. Commun.*, 6, 8435-8435, 10.1080/02786820500421521, 2015.
- 440 [Ma, Y., Huang, C., Jabbour, H., Zheng, Z., Wang, Y., Jiang, Y., Zhu, W., Ge, X., Collier, S., and Zheng, J.: Mixing state and light absorption enhancement of black carbon aerosols in summertime Nanjing, China, \*Atmos. Environ.\*, 222, 117141, 10.1016/j.atmosenv.2019.117141, 2020.](#)
- Petit, J. E., Favez, O., Sciare, J., Canonaco, F., Croteau, P., Močnik, G., Jayne, J., Worsnop, D., and Leoz-Garziandia, E.: Submicron aerosol source apportionment of wintertime pollution in Paris, France by double positive matrix factorization (PMF2) using an aerosol chemical speciation monitor (ACSM) and a multi-wavelength Aethalometer, *Atmos. Chem. Phys.*, 14, 13773-13787, 10.5194/acp-14-13773-2014, 2014.
- Pratt, K. A. and Prather, K. A.: Mass spectrometry of atmospheric aerosols-recent developments and applications. Part II: on-line mass spectrometry techniques, *Mass Spectrom. Rev.*, 31, 17-48, 10.1002/mas.20330, 2012.
- Shen, W., Dai, X., Huang, Z., Hou, Z., Cai, W., Du, X., Zhou, Z., Li, M., and Li, L.: Improvement of the dynamic range of data acquisition system in single particle mass spectrometry, *J. Chinese Cinemas.*, 039, 331-336, 2018.
- 450 Silva, P. J., Liu, D. Y., Noble, C. A., and Prather, K. A.: Size and chemical characterization of individual particles resulting from biomass burning of local Southern California species, *Environ. Sci. Technol.*, 33, 3068-3076, 10.1021/es980544p, 1999.
- Song, X. H., Hopke, P. K., Fergenson, D. P., and Prather, K. A.: Classification of single particles analyzed by ATOFMS using an artificial neural network, ART-2A, *Anal. Chem.*, 71, 860-865, 10.1021/ac9809682, 1999.
- 455 Sun, J., Xie, C., Xu, W., Chen, C., Ma, N., Xu, W., Lei, L., Li, Z., He, Y., Qiu, Y., Wang, Q., Pan, X., Su, H., Cheng, Y., Wu, C., Fu, P., Wang, Z., and Sun, Y.: Light absorption of black carbon and brown carbon in winter in North China Plain: comparisons between urban and rural sites, *Sci. Total Environ.*, 770, [144821](#), 10.1016/j.scitotenv.2020.144821, 2021.


- 460 [Sun, J., Wang, Z., Zhou, W., Xie, C., Wu, C., Chen, C., Han, T., Wang, Q., Li, Z., Li, J., Fu, P., Wang, Z., and Sun, Y.: Measurement report: long-term changes in black carbon and aerosol optical properties from 2012 to 2020 in Beijing, China, \*Atmos. Chem. Phys.\*, 22, 561-575, 10.5194/acp-22-561-2022, 2022.](#)
- [Sun, Y., Wang, Z., Wild, O., Xu, W., Chen, C., Fu, P., Du, W., Zhou, L., Zhang, Q., Han, T., Wang, Q., Pan, X., Zheng, H., Li, J., Guo, X., Liu, J., and Worsnop, D. R.: "APEC Blue": secondary aerosol reductions from emission controls in Beijing, \*Sci. Rep.\*, 6, 20668, 10.1038/srep20668, 2016.](#)
- 465 [Sun, Y., He, Y., Kuang, Y., Xu, W., Song, S., Ma, N., Tao, J., Cheng, P., Wu, C., Su, H., Cheng, Y., Xie, C., Chen, C., Lei, L., Qiu, Y., Fu, P., Croteau, P., and Worsnop, D. R.: Chemical differences between PM<sub>1</sub> and PM<sub>2.5</sub> in highly polluted environment and implications in air pollution studies, \*Geophys. Res. Lett.\*, 47, e2019GL086288, 10.1029/2019gl086288, 2020.](#)
- Thamban, N. M., Tripathi, S. N., Moosakutty, S. P., Kuntamukkala, P., and Kanawade, V. P.: Internally mixed black carbon in the Indo-Gangetic Plain and its effect on absorption enhancement, *Atmos. Res.*, 197, 211-223, 10.1016/j.atmosres.2017.07.007, 2017.
- Vignati, E., Karl, M., Krol, M., Wilson, J., Stier, P., and Cavalli, F.: Sources of uncertainties in modelling black carbon at the global scale, *Atmos. Chem. Phys.*, 10, 2595-2611, 10.5194/acp-10-2595-2010, 2010.
- 470 Wang, Q., Cao, J., Han, Y., Wang, G., Li, G., Wang, Y., Dai, W., Zhang, R., Zhou, Y., and Psi, V.: Mixing state of black carbon aerosol in a heavily polluted urban area of China: implications for light absorption enhancement, *Aerosol Sci. Technol.*, 48, 689-697, 2014.
- 475 Wang, Q., Li, L., Zhou, J., Ye, J., Dai, W., Liu, H., Zhang, Y., Zhang, R., Tian, J., Chen, Y., Wu, Y., Ran, W., and Cao, J.: Measurement report: source and mixing state of black carbon aerosol in the North China Plain: implications for radiative effect, *Atmos. Chem. Phys.*, 20, 15427-15442, 10.5194/acp-20-15427-2020, 2020.
- 480 Xie, C., Xu, W., Wang, J., Liu, D., Ge, X., Zhang, Q., Wang, Q., Du, W., Zhao, J., and Zhou, W.: Light absorption enhancement of black carbon in urban Beijing in summer, *Atmos. Environ.*, 213, 499-504, 2019a.
- Xie, C., Xu, W., Wang, J., Wang, Q., Liu, D., Tang, G., Chen, P., Du, W., Zhao, J., Zhang, Y., Zhou, W., Han, T., Bian, Q., Li, J., Fu, P., Wang, Z., Ge, X., Allan, J., Coe, H., and Sun, Y.: Vertical characterization of aerosol optical properties and brown carbon in winter in urban Beijing, China, *Atmos. Chem. Phys.*, 19, 165-179, 10.5194/acp-19-165-2019, 2019b.
- 485 Xie, C., He, Y., Lei, L., Zhou, W., Liu, J., Wang, Q., Xu, W., Qiu, Y., Zhao, J., Sun, J., Li, L., Li, M., Zhou, Z., Fu, P., Wang, Z., and Sun, Y.: Contrasting mixing state of black carbon-containing particles in summer and winter in Beijing, *Environ. Pollut.*, 263, 10, 10.1016/j.envpol.2020.114455, 2020.
- Xu, W., Chen, C., Qiu, Y., [Li, Y., Zhang, Z., Karnezi, E., Pandis, S. N., Xie, C., Li, Z., Sun, J., Ma, N., Xu, W., Fu, P., Wang, Z., Zhu, J., Worsnop, D. R., Ng, N. L., and Sun, Y.: Organic aerosol volatility and viscosity in the North China Plain: contrast between summer and winter, \*Atmos. Chem. Phys.\*, 21, 5463-5476, 10.5194/acp-21-5463-2021, 2021a.](#)
- 490 [Xu, W., Chen, C., Qiu, Y., Xie, C., Chen, Y., Ma, N., Xu, W., Fu, P., Wang, Z., Pan, X., Zhu, J., Ng, N. L., and Sun, Y.: Size-resolved characterization of organic aerosol in the North China Plain: new insights from high resolution spectral analysis, \*Environ. Sci.: Atmos.\*, 1, 346-358, 10.1039/d1ea00025j, 2021b.](#)

- 495 Yang, J., Ma, S., Gao, B., Li, X., Zhang, Y., Cai, J., Li, M., Yao, L., Huang, B., and Zheng, M.: Single particle mass spectral signatures from vehicle exhaust particles and the source apportionment of on-line PM<sub>2.5</sub> by single particle aerosol mass spectrometry, *Sci. Total Environ.*, 593-594, 310-318, 10.1016/j.scitotenv.2017.03.099, 2017.
- [Yang, M., Howell, S. G., Zhuang, J., and Huebert, B. J.: Attribution of aerosol light absorption to black carbon, brown carbon, and dust in China - interpretations of atmospheric measurements during EAST-AIRE. \*Atmos. Chem. Phys.\*, 9, 2035-2050, 10.5194/acp-9-2035-2009, 2009.](#)
- 500 Zhang, G., Bi, X., Chan, L. Y., Li, L., Wang, X., Feng, J., Sheng, G., Fu, J., Li, M., and Zhou, Z.: Enhanced trimethylamine-containing particles during fog events detected by single particle aerosol mass spectrometry in urban Guangzhou, China, *Atmos. Environ.*, 55, 121-126, 10.1016/j.atmosenv.2012.03.038, 2012.
- Zhang, Q., Zheng, Y., Tong, D., Shao, M., Wang, S., Zhang, Y., Xu, X., Wang, J., He, H., Liu, W., Ding, Y., Lei, Y., Li, J., Wang, Z., Zhang, X., Wang, Y., Cheng, J., Liu, Y., Shi, Q., Yan, L., Geng, G., Hong, C., Li, M., Liu, F., Zheng, B., Cao, J., Ding, A., Gao, J., Fu, Q., Huo, J., Liu, B., Liu, Z., Yang, F., He, K., and Hao, J.: Drivers of improved PM<sub>2.5</sub> air quality in China from 2013 to 2017, *Proc. Natl. Acad. Sci. U. S. A.*, 116, 24463-24469, 10.1073/pnas.1907956116, 2019.
- 505 Zhang, X., Kim, H., Parworth, C. L., Young, D. E., Zhang, Q., Metcalf, A. R., and Cappa, C. D.: Optical properties of wintertime aerosols from residential wood burning in Fresno, CA: results from DISCOVER-AQ 2013, *Environ. Sci. Technol.*, 50, 1681-1690, 10.1021/acs.est.5b04134, 2016.
- 510 Zhang, Y., Wang, X., Chen, H., Yang, X., Chen, J., and Allen, J. O.: Source apportionment of lead-containing aerosol particles in Shanghai using single particle mass spectrometry, *Chemosphere*, 74, 501-507, 10.1016/j.chemosphere.2008.10.004, 2009.
- Zhang, Y., Favez, O., Canonaco, F., Liu, D., Močnik, G., Amodeo, T., Sciare, J., Prévôt, A. S. H., Gros, V., and Albinet, A.: Evidence of major secondary organic aerosol contribution to lensing effect black carbon absorption enhancement, *npj Clim. Atmos. Sci.*, 1, 1-8, 10.1038/s41612-018-0056-2, 2018.
- 515 Zhang, Y., Liu, H., Lei, S., Xu, W., Tian, Y., Yao, W., Liu, X., Liao, Q., Li, J., Chen, C., Sun, Y., Fu, P., Xin, J., Cao, J., Pan, X., and Wang, Z.: Mixing state of refractory black carbon in fog and haze at rural sites in winter on the North China Plain, *Atmos. Chem. Phys. Discuss. [preprint], in review.*, 21, 17631-17648, 10.5194/acp-21-17631-2021, 2021.
- Zhong, Q. E., Cheng, C., Wang, Z., Li, L., Li, M., Ge, D., Wang, L., Li, Y., Nie, W., Chi, X., Ding, A., Yang, S., Chen, D., and Zhou, Z.: Diverse mixing states of amine-containing single particles in Nanjing, China, *Atmos. Chem. Phys. Discuss. [preprint], in review.*, 21, 17953-17967, 10.5194/acp-2021-593, 2021.
- 520 Zhou, W., Gao, M., He, Y., Wang, Q., Xie, C., Xu, W., Zhao, J., Du, W., Qiu, Y., Lei, L., Fu, P., Wang, Z., Worsnop, D. R., Zhang, Q., and Sun, Y.: Response of aerosol chemistry to clean air action in Beijing, China: insights from two-year ACSM measurements and model simulations, *Environ. Pollut.*, 255, 113345, 10.1016/j.envpol.2019.113345, 2019.
- 525 Zhu, S., Li, L., Wang, S., Li, M., Liu, Y., Lu, X., Chen, H., Wang, L., Chen, J., Zhou, Z., Yang, X., and Wang, X.: Development of an automatic linear calibration method for high-resolution single-particle mass spectrometry: improved chemical species identification for atmospheric aerosols, *Atmos. Meas. Tech.*, 13, 4111-4121, 10.5194/amt-13-4111-2020, 2020.




**Figure 1.** The digital positive and negative mass spectra of BC-containing particles in Beijing and Gucheng. The ion height indicates its number fraction in the BC-containing dataset (i.e., the number ratio of BC-containing particles with the corresponding ion detected in mass spectra to the total BC-containing particles). The color bars represent each relative peak area corresponding to a

535 specific fraction in individual particles.



**Figure 2. (a) Relative number abundance distribution of major BC-containing particle types during different periods. Temporal variation of ambient temperature ( $T$ ), relative humidity (RH) and number fractions of BC-containing particle types in (b) Beijing and (c) Gucheng.**










Figure 3. Variations of number fractions of BC-containing particle types with RH in (a) Beijing and (b) Gucheng. **The bottom and top error bars represent 25th and 75th percentiles.** Temporal variations of **wind speed (WS)**, **wind direction (WD)**,  $E_{abs}$ , number fractions of BC-containing particle types and mass concentration of species during polluted (c) Case1 and (d) Case2 **in Beijing.**








Figure 4. Relationships between peak area ratios and mass concentration ratios: (a)  $(NO_3 + SO_4)/C_n$  vs.  $(NO_3 + SO_4)_{AMS}/eBC$  and (b)  $(OC + Metal)/C_n$  vs.  $POA/eBC$ . Relationships between  $E_{abs}$  and peak area ratios of coating materials ( $(NO_3 + SO_4)/C_n$  and  $(OC + Metal)/C_n$ ) in (c) Beijing and (d) Gucheng.  $(NO_3 + SO_4)/C_n$  and  $(OC + Metal)/C_n$  in (c) Beijing and (d) Gucheng. The triangles in (a) are the averages of different bins that are grouped according to  $(NO_3 + SO_4)_{AMS}/eBC$ . The rhombuses in (b) are the averages of different bins that are grouped according to  $POA/eBC$ . The shaded areas in (b) indicate 25th and 75th percentiles of  $(OC + Metal)/C_n$  ratios. The triangles and circles in (c) and (d) are the averages of different bins that are grouped according to  $(NO_3 + SO_4)/C_n$ .

550




Figure 5. Factor profiles and their contributions to each factor identified by the positive matrix factorization model in **BJ**Beijing and **GC**Gucheng.  $b_{abs, total}$  and  $b_{abs, BCpure}$  represent light absorption ( $Mm^{-1}$ ) of coated BC particles and pure BC, respectively. BC types e.g.,  $BB_{pure}$  in units of count.