#### Dear Reviewer,

Thank you very much for your comments on this paper. We have carefully read all comments and revised the manuscript as suggested. The following are our responses to all comments point by point. The italicized sentences are all comments, and the other sentences are the author's responses. The green sentences and words are the specific revisions. We also marked all relevant changes in the manuscript in the same way.

This study investigated the important impacts of the northeast Asian anomalous anticyclone (NAAA) on the intraseasonal variations of PM2.5 pollution in the North China Plain (NCP). The paper presents novel concepts, ideas and tools. The scientific methods and assumptions are valid and clearly outlined so that substantial conclusions are reached. The description of dataset and calculations are sufficiently complete and precise. Hence, the manuscript is recommended for publication after minor revision. Specific comments can be found as follows.

## General comments:

(1) In this study, the authors reported the air quality deterioration two day prior to the peak day of NAAA and suggested that the geopotential height anomaly and meridional wind anomaly were the main causes. However, the cause and effect between the anomalies and PM2.5 should be fully discussed. Because, increases in aerosols, especially absorbing aerosols, can also heat the air, leading to the atmospheric stagnation (e.g., Ding et al., 2016), and decrease the winds over the NCP (e.g., Lou et al., 2019). Therefore, the aerosol accumulation may also cause or at least intensify the dynamical or thermodynamical anomalies.

**Response:** Thank you so much for your suggestion.

Following your advice, we have added a discussion about the effect of  $PM_{2.5}$  on the circulation anomalies in Section 4 of the revised manuscript.

Lines 266–272: "In addition, as shown in Figs. 8–9, PM2.5 pollution occurred over the NCP one day before the peak day of the NAAA, which implies the former might exert an impact on its formation and thus form a positive feedback of PM2.5 pollution–atmospheric circulation. The increases in aerosols, especially absorbing aerosols, have been reported to heat the air, and therefore lead to the atmospheric stagnation (e.g., Ding et al., 2016) and the weaker ventilation over the NCP (e.g., Lou et al., 2019). Therefore, the PM2.5 accumulation before the peak day of the NAAA may also cause or at least intensify the dynamical or thermodynamical anomalies, which in turn might support the formation of PM2.5 pollution over the NCP. This is a potential interesting topic that deserves further investigation in the future."

## Specific comments:

1. Title: The term "air pollution" was used throughout the text including the title. But this study only focused on  $PM_{2.5}$  pollution. It should be revised for the whole text.

**Response:** Thank you for your valuable idea for this paper. We have revised this description in the whole text. So please check the revised manuscript.

2. Lines 16 and 21: "day -3" and day "day -1" should be made clearer.

**Response:** Thank you for your good suggestion. We have revised these descriptions in the manuscript.

Line 17: "... day -3 ..." -> "... day 3 prior to its peak day ..."

Line 22: "... day -1 ..." -> "... the next day ..."

3. Line 39: All sea surface temperature studies listed here are belong to ENSO impacts. It could be more accurately presented. Also, a recent study also reveals impacts of different duration of El Nino on PM<sub>2.5</sub> over China (Zeng et al., 2021).

**Response:** Thank you for your carefully check and help. We have revised this sentence as suggested, so please check the revised manuscript.

Lines 40-41: "... including El Niño (Chang et al., 2016; Jeong et al., 2018; Yu et al., 2020; Zeng et al., 2021), ..."

4. Line 43: Studies also revealed that the aerosol pollution over NCP during COVID-19 was related to NAAA, which could be included here (Ren et al., 2021).

**Response:** Thank you for your carefully check and help. We have revised this sentence as suggested, so please check the revised manuscript.

Lines 43-45: "Studies also revealed that the aerosol pollution over the NCP during COVID-19 was related to the northeast Asian anomalous anticyclone (NAAA) (Ren et al., 2021)."

5. Line 77: The  $PM_{2.5}$  data used in this study were from TAP generated using a machine learning approach and covers the period from 2000 to present. But the reanalysis data are from 1979 to present. The authors are suggested to use the longterm aerosol data for the same period as meteorological parameters in future studies (e.g., Li et al., 2021).

**Response:** Thank you for your suggestion. We will use this data for future studies as much as possible. In fact, it has been used to examine air pollution in our other papers.  $PM_{2.5}$  data provided by Yang (2020) is in excellent agreement with ground measurements, and can capture a trend of continuous increase in the mean  $PM_{2.5}$  concentrations from 1985 to 2014 in China (Li et al., 2021), even daily evolution characteristics of  $PM_{2.5}$  over the North China Plain (An et al., 2022).

6. Line 100: What do the 'e' and 'T' mean?

**Response:** We are sorry for this confusion. "e" represents the spatial pattern of EOF1, which is mentioned in Line 100 (i.e., the EOF1 spatial pattern (e)). We're sorry we didn't use italics for e.  $e^{T}$  is called the transpose of e. We have added these descriptions into the manuscript.

Line 99: "... the EOF1 spatial pattern (*e*) ..."

Line 102: "Here,  $e^T$  is called the transpose of *e*."

7. Line 125: "which is conduced to the accumulation of pollutants in the NCP" should be given after "is weaker than normal".

Response: Thank you for your carefully check and suggestion. We have revised this sentence.

Lines 127–128: "... is weaker than normal (Wang et al., 2009), which is conduced to the accumulation of pollutants in the NCP (An et al., 2020)."

8. Line 154: Please give a short description about the wavenumbers 1-10.

**Response:** Thank you for your valuable advice. We have added the description of the wavenumbers 1-10 in Section 3.1 of the revised manuscript.

Lines 153–155: "Wavenumbers 1–10 are the spectral components on wave-number domain produced by Fourier transform in the spatial domain. Among them, wavenumbers 1–2 represent ultra-long wave, wavenumbers 3–5 denote long wave, and wavenumbers 6–10 are synoptic waves."

## References

- An, X. D., Sheng, L. F., Li, C., Chen, W., Tang, Y. L., and Huangfu, J. L.: Effect of rainfall-induced diabatic heating over southern China on the formation of wintertime haze on the North China Plain, Atmos. Chem. Phys., 22, 725–738, https://doi.org/10.5194/acp-22-725-2022, 2022.
- Ding, A. J., Huang, X., Nie, W., Sun, J. N., Kerminen, V. M., Petäjä, T., et al.: Enhanced haze pollution by black carbon in megacities in China, Geophys. Res. Lett., 43, 2873–2879, https://doi.org/10.1002/2016GL067745, 2016.
- Lou, S., Yang, Y., Wang, H., Smith, S. J., Qian, Y., and Rasch, P. J.: Black carbon amplifies haze over the North China Plain by weakening the East Asian winter monsoon, Geophys. Res. Lett., 46, 452–460, https://doi.org/10.1029/2018GL080941, 2019.
- Zeng, L., Yang, Y., Wang, H., Wang, J., Li, J., Ren, L., Li, H., Zhou, Y., Wang, P., and Liao, H.: Intensified modulation of winter aerosol pollution in China by El Niño with short duration, Atmos. Chem. Phys., 21, 10745–10761, https://doi.org/10.5194/acp-21-10745-2021, 2021.

- Ren, L., Yang, Y., Wang, H., Wang, P., Chen, L., Zhu, J., and Liao, H.: Aerosol transport pathways and source attribution in China during the COVID-19 outbreak, Atmos. Chem. Phys., 21, 15431–15445, https://doi.org/10.5194/acp-21-15431-2021, 2021.
- Li, H., Yang, Y., Wang, H., Li, B., Wang, P., Li, J., and Liao, H.: Constructing a spatiotemporally coherent long-term PM2.5 concentration dataset over China during 1980–2019 using a machine learning approach, Sci. Total Environ., 765, 144263, https://doi.org/10.1016/j.scitotenv.2020.144263, 2021.

# Intraseasonal variation of the northeast Asian anomalous anticyclone and its impacts on PM<sub>2.5</sub> pollution in the North China Plain in early winter

Xiadong An<sup>1,2</sup>, Wen Chen<sup>2</sup>, Peng Hu<sup>2</sup>, Shangfeng Chen<sup>2</sup>, Lifang Sheng<sup>1,3</sup>

<sup>5</sup> <sup>1</sup>Department of Marine Meteorology, College of Oceanic and Atmospheric Sciences, Ocean University of China, Qingdao 266100, China

<sup>2</sup>Center for Monsoon System Research, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100190, China

<sup>3</sup>Ocean-Atmosphere Interaction and Climate Laboratory, Key Laboratory of Physical Oceanography, Ocean University of

10 China, Qingdao 266100, China

Correspondence to: W. Chen (chenw@mail.iap.ac.cn), L. F. Sheng (shenglf@ouc.edu.cn)

Abstract. The canonical view of the northeast Asian anomalous anticyclone (NAAA) is a crucial factor for determining poor air quality (i.e., higher PM<sub>2.5</sub> concentrations) in the North China Plain (NCP) on the interannual timescale. However, there is considerable intraseasonal variability in the NAAA in early winter (November to January), and the corresponding mechanism of its impacts on PM<sub>2.5</sub> pollution in the NCP is not well understood. Here, we find that the intraseasonal NAAA usually establishes quickly on day 3 prior to its peak day with a life span of eight days, and its evolution is closely tied to the Rossby wave from upstream (i.e., the North Atlantic). Moreover, we find that the NAAA with a westward tilt might be mainly related to the wavenumbers 3–4. Further results reveal that under this background, the probability of regional PM<sub>2.5</sub> pollution for at

- 20 least three days in the NCP is as high as 69% (80% at least two days) in NDJ period 2000–2021. In particular, air quality in the NCP tends to deteriorate on day 2 prior to the peak day and reaches a peak on the next day with a life cycle of four days. In the course of PM<sub>2.5</sub> pollution, a shallower atmospheric boundary layer and stronger surface southerly wind anomaly associated with the NAAA in the NCP appear 1 day earlier than poor air quality, which provides dynamic and thermal conditions for the accumulation of pollutants and finally occurrence of the PM<sub>2.5</sub> pollution on the following day. Furthermore,
- 25 we show that the stagnant air leading to poor air quality is determined by the special structure of temperature in the vertical direction of the NAAA, while weak ventilation conditions might be related to a rapid buildup of the NAAA. The present results quantify the impact of the NAAA on PM<sub>2.5</sub> pollution in the NCP on the intraseasonal timescale.

#### **1** Introduction

- 30 The North China Plain (NCP, 32–42°N, 110–120°E) has undergone a series of air pollution (i.e., higher PM<sub>2.5</sub> (fine particulate matter with a mass median aerodynamic diameter < 2.5 μm) concentrations) episodes, particularly in late autumn and early winter (Wang et al., 2019; Yin et al., 2021), which is recognized as significant human health risk and economic activity (Geng et al., 2021a). Nevertheless, PM<sub>2.5</sub> pollution in China has been successfully reduced (i.e., PM<sub>2.5</sub> concentrations fell by 42% between 2013 and 2018 across 74 large cities in China), thanks to comprehensive emission control in response to mounting
- 35 public health risks (Cleaner air for China, 2019). However, PM<sub>2.5</sub> concentration in this region remains the highest in the world (Jeong et al., 2021). Additionally, PM<sub>2.5</sub> pollution is not only related to emissions (i.e., its long-term trends), but also is modulated by the atmospheric circulation (i.e., short-term seasonal variability) (Yang et al., 2016; Dang and Liao, 2019). Moreover, Cai et al. (2017) found that global warming will further increase the incidence of haze days in China by reducing the wind strength.
- 40 Specific to air pollution in the NCP, previous studies have found many influence factors, including El Niño (Chang et al., 2016; Jeong et al., 2018; Yu et al., 2020; Zeng et al., 2021), Arctic sea-ice (Wang et al., 2015, Zou et al., 2020), Eurasian snow cover and soil (Zou et al., 2017; Yin et al., 2018) and climate internal variability including the Eurasian Teleconnection (Li et al., 2019) and subtropical westerly jet waveguide (An et al., 2020, 2021; Mei et al., 2021), and so on. Studies also revealed that the aerosol pollution over the NCP during COVID-19 was related to the northeast Asian anomalous anticyclone (NAAA) (Ren
- et al., 2021). As a matter of fact, the NAAA is directly related to PM<sub>2.5</sub> pollution in the NCP (Wang et al., 2020; Callahan and Mankin, 2020). Wang et al. (2009) and Song et al. (2016) found a weak East Asian trough is usually related to the NAAA, which is mainly induced by the low-frequency Rossby wave and synoptic transient eddy. As a synoptic system, the NAAA not only leads to higher temperature over East Asia by weakening East Asian trough (Song et al., 2016), but also directly modulating stagnant and ventilated conditions for air pollution in the NCP (e.g., Chang et al., 2016; Zhong et al., 2019).
  Moreover, the interannual variability of the NAAA is regulated by external factors mentioned above via atmospheric teleconnection (Yin et al., 2017; Wang et al., 2020; An et al., 2020). Therefore, the NAAA can't be ignored when studying

meteorological causes of  $PM_{2.5}$  pollution in the NCP.

Although previous studies have demonstrated that the NAAA is the decisive factor affecting interannual variation of wintertime air pollution in the NCP except emissions (An et al., 2020; Wang et al., 2020), the role of the NAAA on air pollution on the intraseasonal timescale requires further investigation. On the synoptic scale, Zhong et al. (2019) found that the NAAA also plays a crucial role in haze of the NCP in December. For the research within the intraseasonal timescale, however, the existing studies mainly focus on the analysis of some haze cases (i.e., haze cases are limited to December in the years 2014–2016) (Zhong et al., 2019), lacking a more quantitative statistical analysis and further mechanistic analysis. Therefore, this study focuses on influence of the NAAA on PM<sub>2.5</sub> pollution on the intraseasonal timescale. With the objectives as follows: to derive the characteristics of PM<sub>2.5</sub> pollution in the NCP under the background of the NAAA in November to January (NDJ) on the intraseasonal timescale; to assess the probability of the NAAA in relation to PM<sub>2.5</sub> pollution in the NCP; and to further explore physical mechanisms of the NAAA deriving meteorology conditions for PM<sub>2.5</sub> pollution in the NCP.

The rest of this study is organized as follows: Section 2 describes the data and methods used in this paper. The results of this paper are included in Sect. 3. Specifically, the NAAA events and associated weather patterns are described in Sect. 3.1.  $PM_{2.5}$  pollution in the NCP related to the NAAA is described in Sect. 3.2. Section 3.3 and 3.4 introduce the physical mechanisms of the NAAA causing  $PM_{2.5}$  pollution. The paper concludes with a brief summary and discussion in Sect. 4.

#### 2 Data and methods

#### **2.1 Data**

70

80

85

The monthly and daily reanalysis data were mainly obtained from the National Center for Environmental Prediction (NCEP)/National Center for Atmospheric Research (NCAR) Reanalysis 2 dataset (Kanamitsu et al., 2002, last access: 6 September 2021). The dataset extends from 1979 to present, with a spatial resolution of  $2.5^{\circ} \times 2.5^{\circ}$  and 17 vertical layers extending from 1000 to 10 hPa. The variables including zonal and meridional wind, and air temperature are daily data. The geopotential height is monthly and daily data. In addition, daily atmospheric boundary layer height (ABLH) with a spatial resolution of  $1.0^{\circ} \times 1.0^{\circ}$  in this study averaged from the 6-hourly dataset was taken from the fifth generation the European 75 Centre for Medium-Range Weather Forecasts (ECMWF) reanalysis (ERA5, Hersbach et al., 2018, last access: 8 October 2021).

Air quality degradation is often accompanied by high PM<sub>2.5</sub> concentration (e.g., Yang et al., 2016; Dang and Liao, 2019). Consequently, PM<sub>2.5</sub> concentration is used to describe PM<sub>2.5</sub> pollution in this study. The daily PM<sub>2.5</sub> concentration data used in this study is a near real-time air pollutant database known as Tracking Air Pollution in China (TAP, http://tapdata.org.cn/, last access: 7 October 2021). The daily TAP PM<sub>2.5</sub> concentration data extends from 2000 to present, with a spatial resolution of 10 km in China, which combines information from multiple data sources like ground observations, satellite aerosol optical depth, operational chemical transport model simulations, and other ancillary data (i.e., meteorological fields, land use data, population and elevation) (Geng et al., 2021b). According to Geng et al. (2021b), the TAP PM<sub>2.5</sub> concentration is estimated based on a two-stage machine learning model coupled with the synthetic minority oversampling technique and a tree-based gap-filling method, which has an averaged out-of-bag cross-validation R<sup>2</sup> of 0.83 for different years (Geng et al., 2021b), which is widely used in PM<sub>2.5</sub> pollution research (e.g., Geng et al., 2021a). The results from the TAP PM<sub>2.5</sub> concentration are generally consistent with the observed PM<sub>2.5</sub> concentration data during December 2014 to January 2021, which can be downloaded at website https://quotsoft.net/air/ (not shown, last access: 7 October 2021).

It is noteworthy that the anomalies of the reanalysis data in this paper were calculated using climatology covering the period 1981–2020. Especially, to remove long-term trend due to emission and a comprehensive emission control of Chinese 90 government (Cleaner air for China, 2019), PM<sub>2.5</sub> concentration anomaly was calculated based on a 3-year running climatology state.

#### 2.2 Methods

Firstly, we get a spatial pattern of the NAAA using the Empirical Orthogonal Function (EOF) based monthly mean geopotential height anomaly over domain 25°-55°N, 100°-160°E in NDJ period 1979-2021 (Figs. 1a and b). The first EOF mode (EOF1) 95 represents the NAAA (Fig. 1a), which explains a total variance of 44.2% and is well separated from the other eigenvalues as per the criterion of North et al. (1982). To obtain the typical NAAA on the intraseasonal timescale, the NDJ 8-90-day Butterworth bandpass-filtered daily geopotential height anomaly field at 500 hPa in the region 25°-55°N, 100°-160°E is projected onto the EOF1 to obtain a daily principal component (PC, hereafter) time series. Specifically, z is defined as the observed daily geopotential height anomaly field at 500 hPa, which is projected onto the EOF1 spatial pattern (e) to obtain the 100 PC time series (Fig. 1c) (Baldwin et al. 2009):

$$PC = \frac{ze}{e^T e}.$$
 (1)

## Here, $e^T$ is called the transpose of *e*.

Second, the typical NAAA events is defined as the following way (Fig. 2). First, we rank the values of PC time series in descending order to select the date with the largest PC (i.e., the peak day). If the PC values on at least three days centered on 105 the peak day all exceed one standard deviation, then this peak day is marked as day 0 of a strong NAAA event. Once a day 0 is found, no day within twenty-one days of the central date (day 0) can be defined as a strong NAAA event. This procedure prevents the algorithm from counting the same strong NAAA event repeatedly. Third, we repeat the above procedure until the values of PC don't exceed one standard deviation to guarantee that all the strong NAAA events are identified. Based on the above criterion (Fig. 2), 94 NAAA events in NDJ period 1979-2021 are selected in this study. This method is similar to that

110 of Franzke et al. (2011), who studied the Pacific-North American teleconnection. In addition, the same method was used by Song et al. (2016), who studied the intraseasonal variation of the East Asian trough in winter.

In addition, to examine the propagation of anomalous Rossby waves generating the NAAA, we calculated the horizontal stationary wave activity flux (WAF), as defined by Takaya and Nakamura (2001). Daily reanalysis data; i.e., the zonal wind, meridional wind, and anomalous geopotential height, are used to calculate the vector W.

115 
$$\boldsymbol{W} = \frac{p\cos\varphi}{2|\boldsymbol{U}|} \cdot \left( \frac{\frac{U}{a^2\cos^2\phi} \left[ \left( \frac{\partial\psi'}{\partial\lambda} \right)^2 - \psi' \frac{\partial^2\psi'}{\partial\lambda^2} \right] + \frac{V}{a^2\cos\phi} \left[ \frac{\partial\psi'}{\partial\lambda} \frac{\partial\psi'}{\partial\phi} - \psi' \frac{\partial^2\psi'}{\partial\lambda\partial\phi} \right] \right) \\ \frac{U}{a^2\cos\phi} \left[ \frac{\partial\psi'}{\partial\lambda} \frac{\partial\psi'}{\partial\phi} - \psi' \frac{\partial^2\psi'}{\partial\lambda\partial\phi} \right] + \frac{V}{a^2} \left[ \left( \frac{\partial\psi'}{\partial\phi} \right)^2 - \psi' \frac{\partial^2\psi'}{\partial\phi^2} \right] \right), \tag{2}$$

where **W** is the wave activity flux (unit: m<sup>2</sup> s<sup>2</sup>),  $\psi = \Phi/f$  is the geostrophic stream function,  $\Phi$  (unit: m) is geopotential height,  $f = 2\Omega \sin \phi$  is the Coriolis parameter, p is the normalized pressure (pressure per 1000 hPa), and a is Earth's radius.  $\lambda$ and  $\phi$  denote the longitude and latitude, respectively.  $U = (U, V)^{T}$ ; unit: m s<sup>-1</sup>) is the basic flow.

120

In addition to methods mentioned above, composite analysis is also used to explore the atmospheric circulation patterns related to the NAAA that cause NDJ PM<sub>2.5</sub> pollution in the NCP. The zonal Fourier harmonic analysis of atmospheric circulation is also undertaken to obtain the parameters of the atmospheric waves (van Loon et al., 1973).

## **3 Results**

#### 3.1 Spatial and temporal characteristics of the northeast Asian anomalous anticyclone

Figure 3 presents composite spatial distribution of atmospheric circulation for 94 NAAA events in NDJ period 1979–2021. 125 The results show that there is a remarkably positive geopotential height anomaly at 500 hPa over Northeast Asia with a strong center, i.e., about 40°N, 135°E (Fig. 3a). The NCP is located in the southwest of the NAAA, which is controlled by anomalous southeasterly wind related to the NAAA (Fig. 3c). This means that the East Asian winter monsoon in the NCP, is weaker than normal (Wang et al., 2009), which is conduced to the accumulation of pollutants in the NCP (An et al., 2020). Additionally, the warm and moisture flow from the west Pacific is advected by anomalous southeasterly wind into the NCP, favoring the 130 hygroscopic growth of pollution (Ma et al., 2014). As a result, the NCP might experience heavy  $PM_{2.5}$  pollution weather. Significantly, the maximum of the NAAA locates about 300 hPa with a vertical structure of westward-tilt from 1000 hPa to 850 hPa (Fig. 3b). The corresponding temperature anomaly is a dipole pattern at the lower (1000 hPa to 300 hPa) and high (300 hPa to 10 hPa) level. That is to say that the lower is positive and the higher is negative temperature anomaly (Fig. 3b), which might lead to a westward-tilt structure of the NAAA via thermal wind and transient eddy feedback (Song et al., 2016).

- 135 To understand the life span of the NAAA, we show the temporal evolution of standardized daily PC time series of 94 NAAA events (Fig. 4). The PC values become positive from day -4, meaning that the NAAA starts to emerge. Note that the PC index reaches its maximum on day 0. And the PC index is almost zero or even negative from day 4, which implies the extinction of the NAAA with a life span of eight days. Moreover, the 8-day life cycle of the NAAA suggests that it is enough to investigate the intraseasonal evolution and dynamics of the NAAA in the 21-day period described in section of method. The question right 140 now is where does the NAAA start?

To investigate the causes and evolution mechanism of the NAAA, horizontal wave activity flux is calculated and shown in the form of arrows in Fig. 5. Distinctly, there is a positive geopotential anomaly over the Gulf Stream on day -8 and propagates eastward along the upper-tropospheric polar front jet, which serves as a waveguide (Hoskins and Ambrizzi, 1993). On day -6 and the next two days, the Rossby wave energy reaches the region of Northeast Asia, but there is no positive geopotential

- height anomaly there. Note that the significantly positive geopotential height anomaly appears in Northeast Asia on day -3 and -2 (Fig. 5), which is an embryo of the NAAA, namely, means a rapid buildup of the NAAA. On day 0, the NAAA reaches the peak of its life cycle and wears out almost immediately on the next day (Figs. 4 and 5). There is almost no positive geopotential height anomaly in Northeast Asia on day 4. On the interannual timescale, the NAAA seems to always occupy the whole winter and sustain degradation effect on air quality in the NCP (Chang et al., 2016; An et al., 2020). On the synoptic
- 150 scale, however, the life cycle of the NAAA is just eight days. The results further suggest the necessity of studying the impact of the NAAA on PM<sub>2.5</sub> pollution in the NCP on the synoptic scale.

For a deeper understanding generation of the NAAA with a westward-tilting structure from wave theory, zonal harmonic analysis is used in this investigation. Wavenumbers 1–10 are the spectral components on wave-number domain produced by Fourier transform in the spatial domain. Among them, wavenumbers 1–2 represent ultra-long wave, wavenumbers 3–5 denote
long wave, and wavenumbers 6–10 are synoptic waves. Figure 6 compares the height–longitude cross section of zonal harmonic wave anomalies on the peak day of the NAAA, overlapped with raw geopotential height anomaly. Note that the reason why other wavenumbers (i.e., wavenumbers 5–10) are not shown in Fig. 6 is that their shape are quite different from the shape of the NAAA. From Fig. 6, we find that the shape of wavenumbers 3–4 (referred as the quasi-stationary wave) is consistent with that of the NAAA in general. The results suggest that wavenumbers 3–4 might play an important role in the generation and elimination of the NAAA (Fig. 6). The amplitudes and variances of the harmonics also support the significant roles played by the Rossby wave (Fig. 7). For instance, the amplitudes and variances of wavenumbers 3–4 are significantly greater than other wavenumbers (Fig. 7). In addition to the quasi-stationary wave characterized by wavenumbers 3–4, transient eddy feedback (2–8 days on the timescale) due to a baroclinic atmosphere also plays an important role in the development of the NAAA (i.e., contributes to rapid buildup of the NAAA) (Song et al., 2016).

## 165 3.2 The northeast Asian anomalous anticyclone in relation to variation of PM<sub>2.5</sub> pollution in the NCP

Sections 3.1 investigates the spatiotemporal characteristics and evolution mechanism of the NAAA on the intraseasonal timescale, how it relates to air quality in the NCP, and what potential conditions give rise to this regime. Figure 8 presents composite PM<sub>2.5</sub> concentration anomaly from day –4 to 4 of 51 NAAA events in NDJ period 2000 to 2021. PM<sub>2.5</sub> concentration tends to increase on day –3 and then increase rapidly since day –2. By 1 day before the peak of the NAAA, PM<sub>2.5</sub> concentration anomaly reaches a maximum and maintains on the next day. On day 2 after the peak of the NAAA, positive PM<sub>2.5</sub> concentration anomaly tends to dissipate. Generally, under the background of the NAAA, the NCP experiences heavy PM<sub>2.5</sub> pollution for four days. Similarly, we investigate evolution of PM<sub>2.5</sub> pollution in the NCP based on the TAP PM<sub>2.5</sub> concentration and observed PM<sub>2.5</sub> concentration data since 2013, respectively (not shown). And results are in line with the above conclusions drawn using TAP PM<sub>2.5</sub> concentration data since 2000, suggesting our finds are reliable despite PM<sub>2.5</sub> concentration data from machine learning by Geng et al. (2021).

Figure 9 shows daily PM<sub>2.5</sub> concentration anomaly averaged in the NCP for eight days before and after peak day of the NAAA. The results indicate a distinct evolution of PM<sub>2.5</sub> pollution compared with that of the NAAA. Clearly, PM<sub>2.5</sub> concentration begins to increase after day -4 with a peak on day -1 and then decreases gradually until zero on day 2. The NCP has gone through significant PM<sub>2.5</sub> pollution for day -2 to 1 of the peak day of the NAAA (Fig. 9), which is consistent with the conclusions from Fig. 8. Significantly, the interquartile range (specially interdecile range) of area-averaged PM<sub>2.5</sub> concentration anomaly during -2 to 1 has parts less than 0, meaning that not all of the NAAA events can cause PM<sub>2.5</sub> pollution for at least three days (i.e., day -2 to 0) in the NCP. This makes us aware that the probability of PM<sub>2.5</sub> pollution events in the NCP related to the NAAA should be further examined.

To answer the question what is the probability of  $PM_{2.5}$  pollution in the NCP caused by the NAAA in NDJ period of 2000–2021 on the intraseasonal timescale. The probability of  $PM_{2.5}$  pollution under the background of the NAAA is presented in Table 1, and the event of  $PM_{2.5}$  pollution is defined here as exceeding 0 for at least three days (i.e., day -2 to 0) for region-averaged  $PM_{2.5}$  concentration anomaly in the NCP (Table 1). The probability of the NAAA in relation to  $PM_{2.5}$  pollution for at least three days in the NCP is 69% if we start counting from 2000. This percent is 64% when we start counting from 2014. Additionally, the probability of the NAAA in relation to  $PM_{2.5}$  pollution for at least two days in the NCP is higher (i.e., 80% and 72%) than

190 at least three days. These results further illustrate meteorological factors, especially the NAAA, play a crucial role in NDJ PM<sub>2.5</sub> concentration in the NCP in spite of a decline of 42% of the annual mean PM<sub>2.5</sub> concentrations between 2013 and 2018 in China (Cleaner air for China, 2019), which is in line with results by Dang and Liao (2019). From what is mentioned above, we come to the robust conclusion that 69% of the NAAA might cause NDJ PM<sub>2.5</sub> pollution for at least three days in the NCP during the period of 2000–2021.

## 195 3.3 Why does PM<sub>2.5</sub> pollution occur in the NCP before the peak day of the NAAA

200

205

From the previous section, we see that  $PM_{2.5}$  pollution in the NCP begins to deteriorate significantly from day -2 of the peak day of the NAAA. What sort of meteorological conditions causes this observed fact of  $PM_{2.5}$  pollution. The NAAA is usually accompanied by southerly wind anomalies on its western flank, corresponding to lower ABLH and weaker surface winds (Yin et al., 2017). We therefore explore the possible meteorological conditions favouring  $PM_{2.5}$  pollution in terms of dynamics (i.e., diffusion condition) and thermodynamics (i.e., stability). In Fig. 10, the evolution of the ABLH anomaly four days before and after the peak day of the NAAA is shown. The results show that there is remarkably negative ABLH anomaly on day -3, which means a shallow atmospheric boundary layer, favourable to accumulation of pollutants. It should be noted that the ABLH reduction one day prior to the appearance of  $PM_{2.5}$  pollution, which provides sufficient time for the accumulation of pollutants so that the occurrence of  $PM_{2.5}$  pollution on the following day (Figs. 8 and 11). On day 1, the negative ABLH anomaly decreases abruptly, corresponding  $PM_{2.5}$  pollution is also lightly weakened (Figs. 8 and 11). While on the next day (i.e., day 2), there is no significantly negative ABLH anomaly, corresponding  $PM_{2.5}$  pollution also almost disappears in the NCP (Figs. 8 and 11).

Similar conclusions can be drawn from the wind field for four days before and after the peak day of the NAAA, which represents a diffusion condition for  $PM_{2.5}$  pollution (e.g., Yang et al., 2016, Liu et al., 2017). As shown in Fig. 11, the NCP is mainly controlled by anomalous southerly wind with a negative divergence anomaly (not shown), which also appears one day

(i.e., day -3) earlier than heavy PM<sub>2.5</sub> pollution in the NCP. The intensify and range of southerly wind increase significantly on the following two days (i.e., day -1 and day 0). The intensify and range of southerly wind, however, shrink rapidly on day 1 and almost disappear on day 2. This process is consistent with PM<sub>2.5</sub> pollution in the NCP except that the establishment of favourable wind field is earlier (for one day) than the occurrence of PM<sub>2.5</sub> pollution. The earlier emergence of southerly wind anomaly with a negative divergence anomaly and shallow atmospheric boundary layer together facilitate the accumulation of pollutants, leading to the happening of PM<sub>2.5</sub> pollution in the following day. And the maintenance of these two parameters leads to PM<sub>2.5</sub> pollution last for the next two days. While the later weakening and even disappearance of southerly wind anomaly and shallow atmospheric boundary layer together the day 1 of peak day of the NAAA.

Overall, both dynamic and thermodynamic conditions associated with the NAAA result in heavy PM<sub>2.5</sub> pollution in the NCP. Most importantly, PM<sub>2.5</sub> pollution in the NCP happens 1 day earlier than the peak of the NAAA, which provides a reference for prediction of PM<sub>2.5</sub> pollution in the NCP on the synoptic scale. In addition, if we take the positive geopotential height anomaly over domain 45°N–60°N, 80°E–100°E as a predictor, the potential prediction of PM<sub>2.5</sub> pollution in the NCP might be extended to five days (Fig. 5). Now what we wonder is why the favourable meteorological conditions related to the NAAA appear before the peak day of the NAAA.

To further understand physical mechanisms of the NAAA favoring occurrence of PM<sub>2.5</sub> pollution in the NCP, the vertical structure of temperature and geopotential anomaly and their evolution in position of 37°N, 115°E are shown in this section. As shown in Fig. 12, temperature anomaly features a backward tilt with height below 700 hPa, meaning that the higher temperature anomaly moves from higher to lower level from day –3 to –1, which is easier to cause a potential thermal inversion. In addition, the ABLH anomaly is significantly negative in this period (Fig. 10 and 12), corresponding negative PM<sub>2.5</sub> concentration anomaly in the NCP (Fig. 8). While above 700 hPa, it features a forward tilt with height, implying that the positive temperature

- anomaly moves form lower to higher level from day -1 to 2, which is unfavourable for the formation of a potential thermal inversion. Besides, there is significantly anomalous ascending motion in the troposphere on day -3 to -1 (not shown), which might suppress intrusions of clean air from upper levels (i.e., above 300 hPa) to the lower levels, resulting in a shallower atmospheric boundary layer (Zhong et al., 2019). However, the negative ABLH anomaly decreases rapidly from day -1 to 2, and corresponding air quality in the NCP is gradually improved (Fig. 10 and 12). The characteristics based on area-averaged temperature and ABLH anomaly in the NCP are similar to the results based on single position (not shown).
- Similarly, we check the evolution of geopotential height anomaly and meridional wind anomaly with time and height in point (37°N, 115°E). Unsurprisingly, positive geopotential height anomaly shows a sudden enhancement in the whole troposphere from day -4 to 3 (Fig. 13), which is in line with result in Fig. 4b. This means a rapid buildup of the NAAA with a sudden enhancement of anomalous southerly wind. As we all know, PM<sub>2.5</sub> pollution is closely tied to lower wind field, especially surface wind field (e.g., Yang et al., 2016; Liu et al., 2017; Yin et al., 2017). Therefore, on day -3, the NAAA rapidly builds with the sudden increase of anomalous geopotential height anomaly and southerly wind anomaly (Figs. 7 and 13), resulting in PM<sub>2.5</sub> pollution prior to the peak day of the NAAA. We can draw the same conclusion compared with an area-averaged geopotential height anomaly and 1000 hPa meridional wind anomaly in the NCP.

#### 4 Conclusions and discussion

- In this study, we investigate the characteristics and evolution mechanisms of the NAAA on the intraseasonal timescale and the associated PM<sub>2.5</sub> pollution in the NCP in NDJ. In particular, the intraseasonal NAAA has a life span of about eight days with a structure of westward tilt with height, and its evolution is closely tied to the Rossby wave from upstream (i.e., the North Atlantic). On day -8, there is significant circulation anomaly over the Gulf Stream and downstream propagation in the form of the Rossby wave. The NAAA reaches its peak on day 0 and decreases rapidly the next day. According to harmonic analysis, the NAAA with a westward tilt may be related to the wavenumbers 3-4. Additionally, the NAAA is also enhanced by the transient eddy, which can be induced by weak baroclinic atmosphere with the characteristic of vertical dipole pattern of temperature. For instance, Song et al. (2016) found that the transient eddy feedback leads to 30% of the NAAA amplification using geopotential height tendency equation.
- Further results show that 69% of the NAAA in NDJ period of 2000–2021 causes regional  $PM_{2.5}$  pollution for at least three days (80% for at least two days) in the NCP and its peak day lags occurrence of  $PM_{2.5}$  pollution for two days. The composite analysis reveals that the shallower atmospheric boundary layer and stronger surface southerly wind anomaly (weaker northerly wind) associated with the NAAA in the NCP appear one day prior to  $PM_{2.5}$  pollution, which provides dynamic and thermal conditions for the accumulation of pollutants and finally occurrence of  $PM_{2.5}$  pollution on the following day. We also find that the stagnant air and weak ventilation conditions are determined by a special vertical distribution of temperature anomaly and a rapid buildup of the NAAA.
  - It is well known that the wet deposition through scavenging by rainfall is an effective way to remove atmospheric aerosols and soluble gases (e.g., Atlas and Giam, 1988). When the NAAA appears, southern China tends to experience heavy rainfall and vice versa in the NCP (not shown) (e.g., Ma et al., 2018; An et al., 2020, 2021), which is only not conducive to the wet removal of aerosol in the NCP, but usually deteriorates air quality in the NCP via a local north–south circulation (not shown) (An et al., 2020, 2021; Mei et al., 2021).
  - In addition, as shown in Figs. 8–9, PM<sub>2.5</sub> pollution occurred over the NCP one day before the peak day of the NAAA, which implies the former might exert an impact on its formation and thus form a positive feedback of PM<sub>2.5</sub> pollution–atmospheric circulation. The increases in aerosols, especially absorbing aerosols, have been reported to heat the air, and therefore lead to the atmospheric stagnation (e.g., Ding et al., 2016) and the weaker ventilation over the NCP (e.g., Lou et al., 2019). Therefore, the PM<sub>2.5</sub> accumulation before the peak day of the NAAA may also cause or at least intensify the dynamical or

270

thermodynamical anomalies, which in turn might support the formation of  $PM_{2.5}$  pollution over the NCP. This is a potential interesting topic that deserves further investigation in the future.

In brief, the NAAA and associated meteorological parameters play a crucial role in formation of NDJ PM<sub>2.5</sub> pollution in the NCP on the intraseasonal timescale, which is slightly different from its role on wintertime PM<sub>2.5</sub> pollution in this region on the interannual timescale. For example, we can't draw a conclusion that the peak time of the NAAA lags PM<sub>2.5</sub> pollution in the NCP in NDJ on the interannual timescale. In addition, there is usually an anomalous descending motion in the NCP in NDJ on the interannual timescale (An et al., 2020), while on the intraseasonal timescale is an anomalous ascending motion in the NCP in NDJ on the interase of this study only investigates the influence of the NAAA on PM<sub>2.5</sub> pollution in the NCP in NDJ on the intraseasonal timescale. It should be noticed that cyclone anomaly in Northeast Asia, as a pattern of out-of-

phase of the NAAA (Wang et al., 2009; Song et al., 2016), might be a favorable atmospheric circulation to improve PM2.5

280

quality, which should also be studied in future.

## Code availability

Codes used in this paper are available upon request to the corresponding author.

## Data availability

285 NCEP reanalysis 2 data provided by the NOAA/OAR/ESRL PSL, Boulder, Colorado, USA, from their Web site at https://psl.noaa.gov/data/gridded/data.ncep.reanalysis2.html (last access: 6 September 2021).

Daily boundary layer height data is available from https://cds.climate.copernicus.eu/cdsapp#!/home (last access: 8 October 2021).

The hourly PM<sub>2.5</sub> concentration data can be downloaded from https://quotsoft.net/air/ and http://tapdata.org.cn/ (TAP, last access: 7 October 2021).

Author contributions: XA, LS and WC designed the experiments and carried them out. XA downloaded and analysed the reanalysis data and prepared all the figures. XD prepared the manuscript with contributions from all co-authors. LS, WC, PH, and SC revised the manuscript.

Competing interests: The authors declare that they have no conflict of interest.

## 295 Acknowledgements

This research was supported by the National Natural Science Foundation of China (grants no. 41975008 and 41675146).

#### References

An, X. D., Sheng, L. F., Liu, Q., Li, C., Gao, Y., and Li, J. P.: The combined effect of two westerly jet waveguides on heavy haze in the North China Plain in November and December 2015, Atmos. Chem. Phys., 20, 4667–4680, https://doi.org/10.5194/acp-20-4667-2020, 2020.

An, X. D., Sheng, L. F., Li, C., Chen, W., Tang, Y. L., and Huangfu, J. L.: Effect of rainfall-induced diabatic heating over southern China on the formation of wintertime haze on the North China Plain, Atmos. Chem. Phys., 22, 725–738, https://doi.org/10.5194/acp-22-725-2022, 2022.

Atlas, E. and Giam, C. S.: Ambient Concentration and Precipitation Scavenging of Atmospheric 461 Organic Pollutants, Water 305 Air Soil Poll., 38, 19–36, 1988.

Baldwin, M. P., Stephenson, D. B., and Jolliffe, I. T.: Spatial weighting and iterative projection methods for EOFs, J. Climate, 22, 234–243, https://doi.org/10.1175/2008JCLI2147.1, 2009.

Cleaner air for China, Nat. Geosci., 12, 497, https://doi.org/10.1038/s41561-019-0406-7, 2019.

Cai, W., Li, K., Liao, H., Wang, H. J., and Wu, L. X.: Weather conditions conducive to Beijing severe haze more frequent under climate change, Nat. Clim. Change, 7, 257–262, https://doi.org/10.1038/nclimate3249, 2017.

Callahan, C. W., and Mankin, J. S.: The influence of internal climate variability on projections of synoptically driven Beijing haze, Geophys. Res. Lett., 46, e2020GL088548. https://doi.org/10.1029/2020GL088548, 2020.

Chang, L., Xu, J., Tie, X., and Wu J..: Impact of the 2015 El Niño event on winter air quality in China, Sci. Rep., 6, 34275, https://doi.org/10.1038/srep34275, 2016.

315 Dang, R. and Liao, H.: Severe winter haze days in the Beijing-Tianjin-Hebei region from 1985 to 2017 and the roles of anthropogenic emissions and meteorology, Atmos. Chem. Phys., 19, 10801–10816, https://doi.org/10.5194/acp-19-10801-2019, 2019.

Ding, A. J., Huang, X., Nie, W., Sun, J. N., Kerminen, V. M., Petäjä, T., et al.: Enhanced haze pollution by black carbon in megacities in China, Geophys. Res. Lett., 43, 2873–2879, https://doi.org/10.1002/2016GL067745, 2016.

320 Franzke, C., Feldstein, S. B., and Lee, S.: Synoptic analysis of the Pacific–North American teleconnection pattern, Quart. J. Roy. Meteor. Soc., 137, 329–346, https://doi.org/10.1002/qj.768, 2011.

Geng, G., Xiao, Q., Liu, S., Liu, X., Cheng, J., Zheng, Y., Xue, T., Tong, D., Zheng, B., Peng, Y., Huang, X., He, K., and Zhang, Q.: Tracking Air Pollution in China: Near Real-Time PM<sub>2.5</sub> Retrievals from Multisource Data Fusion. Environ, Sci. Technol., 55, 12106–12115, https://doi.org/10.1021/acs.est.1c01863 (last access: 7 October 2021), 2021a.

325 Geng, G. N., Zheng, Y. X., Zhang, Q., Xue, T., Zhao, H. Y., Tong, D., Zheng, B., Li, M., Liu, F., Hong, C. P., He, K. B., and Davis, S. J.: Drivers of PM2.5 air pollution deaths in China 2002–2017, Nat. Geosci. 14, 645–650, https://doi.org/10.1038/s41561-021-00792-3, 2021b.

Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., Thépaut, J-N.: ERA5 hourly data on single levels from 1979 to present,
Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [data set], https://doi.org/10.24381/cds.adbb2d47 (last access: 8 October 2021), 2018.

Jeong, J. J., Park, R. J., Yeh, S. W.: Dissimilar effects of two El Niño types on PM2.5 concentrations in East Asia, Environ. Poll., 242, 1395–1403, https://doi.org/10.1016/j.envpol.2018.08.031, 2018.

Jeong, J. I., Park, R. J., Yeh, S. W., Roh, J. W.: Statistical predictability of wintertime PM2.5 concentrations over East Asia using simple linear regression, Sci. Total Environ., 776, 146059, https://doi.org/10.1016/j.scitotenv.2021.146059, 2021.

Kanamitsu, M., Ebisuzaki, W., Woollen, J., Yang, S-K., Hnilo, J. J., Fiorino, M., and Potter, G. L.: NCEP-DOE AMIP-II Reanalysis (R-2), Bulletin of the American Meteorological Society, 1631–1643, https://doi.org/10.1175/BAMS-83-11-16312002 (last access: 6 September 2021), 2002.

Li, Y., Sheng, L. F., Li, C., and Wang, Y. H.: Impact of the Eurasian Teleconnection on the Interannual Variability of Haze-Fog in Northern China in January, Atmosphere, 10, 113, https://doi.org/10.3390/atmos10030113, 2019.

Lou, S., Yang, Y., Wang, H., Smith, S. J., Qian, Y., and Rasch, P. J.: Black carbon amplifies haze over the North China Plain by weakening the East Asian winter monsoon, Geophys. Res. Lett., 46, 452–460, https://doi.org/10.1029/2018GL080941, 2019.

Ma, N., Zhao, C. S., Chen, J., Xu, W. Y., Yan, P., and Zhou, X. J.: A novel method for distinguishing fog and haze based on PM2.5, visibility, and relative humidity, Sci. China Earth Sci., 57, 2156–2164, https://doi.org/10.1007/s11430-014-4885-5, 2014.

345

Ma, T. J., Chen, W., Feng, J., and Wu, R. G.: Modulation effects of the East Asian winter monsoon on El Niño-related rainfall anomalies in southeastern China, Sci. Rep., 8, 12107, https://doi.org/10.1038/s41598-018-32492-1, 2018.

Mei, M., Ding, Y., Wang, Z., Liu, Y. and Zhang, Y.: Effects of the East Asian Subtropical Westerly Jet on Winter Persistent Heavy Pollution in the Beijing-Tianjin-Hebei Region, Int. J. Climatol., https://doi.org/10.1002/joc.7400, 2021.

350 North, G. R., Bell, T. L., Cahalan, R. F., and Moeng, F. J.: Sampling Errors in the Estimation of Empirical Orthogonal Functions, Mon. Wea. Rev., 110, 7, 699–706, https://doi.org/10.1175/1520-0493(1982)110<0699:SEITEO>2.0.CO;2, 1982.

Ren, L., Yang, Y., Wang, H., Wang, P., Chen, L., Zhu, J., and Liao, H.: Aerosol transport pathways and source attribution in China during the COVID-19 outbreak, Atmos. Chem. Phys., 21, 15431–15445, https://doi.org/10.5194/acp-21-15431-2021, 2021.

355 Song, L., Wang, L., Chen, W., and Zhang, Y.: Intraseasonal Variation of the Strength of the East Asian Trough and Its Climatic Impacts in Boreal Winter, J. Climate, 29, 2557–2577, https://doi.org/10.1175/JCLI-D-14-00834.1, 2016.

Takaya, K., and Nakamura, H.: A formulation of a phase independent wave-activity flux for stationary and migratory quasigeostrophic eddies on a zonally varying basic flow, J. Atmos. Sci., 58, 608–627, https://doi.org/10.1175/1520-0469(2001)058<0608:AFOAPI>2.0.CO;2, 2001.

360 van Loon, H., Jenne, R. L., and Labitzke, K.: Zonal harmonic standing waves, J. Geophys. Res., 78(21), 4463–4471, https://doi.org/10.1029/JC078i021p04463, 1973.

Wang, H., Chen, H., and Liu, J.: Arctic sea ice decline intensified haze pollution in Eastern China, Atmos. Oceanic Sci. Lett., 8(1), 1–9, 2015.

Wang, L., Chen, W., Zhou, W., and Huang, R. H.: Interannual Variations of East Asian Trough Axis at 500 hPa and its
Association with the East Asian Winter Monsoon Pathway, J. Clim., 22, 600–614, https://doi.org/10.1175/2008JCLI2295.1, 2009.

Wang, J., Zhu, Z., Qi, L., Zhao, Q., He, J., and Wang, J. X. L.: Two pathways of how remote SST anomalies drive the interannual variability of autumnal haze days in the Beijing–Tianjin–Hebei region, China, Atmos. Chem. Phys., 19, 1521–1535, https://doi.org/10.5194/acp-19-1521-2019, 2019.

370 Wang, J, Liu, Y, Ding, Y, et al.: Impacts of climate anomalies on the interannual and interdecadal variability of autumn and winter haze in North China: A review, Int. J. Climatol., 40, 4309–4325, https://doi.org/10.1002/joc.6471, 2020.

Yang, Y., Liao, H., and Lou, S.: Increase in winter haze over eastern China in recent decades: Roles of variations in meteorological parameters and anthropogenic emissions, J. Geophys. Res. Atmos., 121, 13050–13065, https://doi.org/10.1002/2016JD025136, 2016.

375 Yin, Z., Wang, H., and Chen, H.: Understanding severe winter haze events in the North China Plain in 2014: roles of climate anomalies, Atmos. Chem. Phys., 17, 1641–1651, https://doi.org/10.5194/acp-17-1641-2017, 2017.

Yin, Z. and Wang, H.: The strengthening relationship between Eurasian snow cover and December haze days in central North China after the mid-1990s, Atmos. Chem. Phys., 18, 4753–4763, https://doi.org/10.5194/acp-18-4753-2018, 2018.

Yin, Z. C., Zhou, B. T., Chen, H. P., and Li, Y. Y.: Synergetic impacts of precursory climate drivers on interannual-decadal 380 in haze pollution in North China: А review, Sci. Total Environ., 755. 143017. variations https://doi.org/10.1016/j.scitotenv.2020.143017, 2021.

Yu, X., Wang, Z., Zhang, H., He, J., and Li, Y.: Contrasting impacts of two types of El Niño events on winter haze days in China's Jing-Jin-Ji region, Atmos. Chem. Phys., 20, 10279–10293, https://doi.org/10.5194/acp-20-10279-2020, 2020.

Zeng, L., Yang, Y., Wang, H., Wang, J., Li, J., Ren, L., Li, H., Zhou, Y., Wang, P., and Liao, H.: Intensified modulation of
winter aerosol pollution in China by El Niño with short duration, Atmos. Chem. Phys., 21, 10745–10761, https://doi.org/10.5194/acp-21-10745-2021, 2021.

Zou, Y., Wang, Y., Zhang, Y., and Koo, J.-H.: Arctic sea ice, Eurasia snow, and extreme winter haze in China, Sci. Adv., 3, e1602751, https://doi.org/10.1126/sciadv.1602751, 2017.

Zou, Y., Wang, Y., Xie, Z., Wang, H., and Rasch, P. J.: Atmospheric teleconnection processes linking winter air stagnation and
 haze extremes in China with regional Arctic sea ice decline, Atmospheric Chemistry and Physics, 20(8), 4999–5017,
 https://doi.org/10.5194/acp-20-4999-2020, 2020.

Zhong, W., Yin, Z., and Wang, H.: The relationship between anticyclonic anomalies in northeastern Asia and severe haze in the Beijing–Tianjin–Hebei region, Atmos. Chem. Phys., 19, 5941–5957, https://doi.org/10.5194/acp-19-5941-2019, 2019.



Figure 1: (a) The spatial pattern of the first EOF and (b) the corresponding standardized PC series over the domain 25°N-55°N, 100°E-160°E in NDJ period 1979-2020. (c) The standardized daily PC series in NDJ period 1979-2020, red line is one standard deviation.



Figure 2: Process for selecting the NAAA events. Here, PC is daily index of the NAAA, sqsort (PC) represents PC is sorted in descending order, dimsize (PC) is the size of one-dimensional array PC.



405 Figure 3: (a) Composite geopotential height anomaly (shaded and contours; unit: m) at 500 hPa on the peak day of 94 NAAA events in NDJ period 1979 to 2021. (b) Composite longitude-height cross section of geopotential height anomaly (shaded; unit: m) and temperature anomaly (contours; unit: °C) at 37°N. (c) Composite wind vector (arrows; unit: m s<sup>-1</sup>) and wind speed (shaded; unit: m s<sup>-1</sup>) at 850 hPa. The white dots denote the 99% confidence level according to the Student's *t* test.



Figure 4: The temporal evolution of the standard PC time series for 94 NAAA events. In particular, black and blue curves, light and deep gray filling represent mean and median value, interdecile range and interquartile range, respectively.



415 Figure 5: Composite evolution of geopotential height anomaly (shaded; unit: m) and the WAF (arrows; unit: m<sup>2</sup> s<sup>2</sup>) at 500 hPa on days -8, -6, -4, -2, 0, +2, +4, +6 and +8 in 94 NAAA events since 1979. The white dots denote the 99% confidence level according to the Student's *t* test.



420 Figure 6: Cross section of zonal harmonic analysis of geopotential height anomaly. (a) wavenumber 1, (b) wavenumber 2, (c) wavenumber 3 and (d) wavenumber 4 (shaded; unit: m) and geopotential height anomaly (contours; unit: m) from 1000 hPa to 10 hPa on peak day of 94 NAAA events. The white dots denote the 99% confidence level according to the Student's *t* test.



425 Figure 7: Zonal harmonic analysis of geopotential height anomaly at 500 hPa on the peak day of 94 NAAA events since 1979. (a) Amplitude harmonics of wave 1 to wave 10, (b) variances harmonics of wavenumber 1 to 10. In particular, the thick black curve (Ob) represents zonal mean geopotential height anomaly in the peak day of 94 NAAA events.



430 Figure 8: Composite evolution of PM<sub>2.5</sub> concentration anomaly (shaded; unit: ug m<sup>-3</sup>) from day -8 to day 8 in 51 NAAA events in NDJ period 2000–2021. The white dots denote the 99% confidence level according to the Student's *t* test.



Figure 9: The same as Fig. 3 except for PM<sub>2.5</sub> concentration anomaly in 51 NAAA events in NDJ period 2000–2021.



Figure 10: The same as Fig. 5 except for the ABLH anomaly (shaded; unit: m).



440 Figure 11: The same as Fig. 5 except for anomalous wind vector (arrows; unit: m s<sup>-1</sup>) and anomalous wind speed (shaded; unit: m s<sup>-1</sup>) at 1000 hPa.



Figure 12: Composite time-height cross section of temperature anomaly (shaded; unit: m) at 37°N, 115°E from the developing stage (day –8 to day 0) to the decaying stage (day 0 to day +8) for 51 NAAA events in NDJ period 2000–2021. The white dots denote the 99% confidence level according to the Student's *t* test. Black bar charts represent the variation of the ABLH anomaly (bars; unit: m) at 37°N, 115°E from the developing stage (day –8 to day 0) to the decaying stage (day 0 to day +8) of 51 NAAA events in NDJ period 2000–2021. Especially, the coordinates on the right represent the value of the ABLH anomaly. The horizontal line denotes zero of the ABLH anomaly.



Figure 13: The same as Fig. 9, but for geopotential height anomaly (shaded; unit: m) and 1000-hPa meridional wind anomaly (bars; unit: m  $s^{-1}$ ). Positive values of meridional wind anomaly represent southerly wind anomaly.

## 455 Table 1: The probability of the NAAA in relation to PM<sub>2.5</sub> pollution in the NCP.

	The NAAA	The number of $PM_{2.5}$ pollution lasted		The number of PM <sub>2.5</sub> pollution lasted	
		two days (day $-1$ and day 0)		three days (day -2 and day 0)	
	number	number	percent	number	percent
2000.11-2021.01	51	41	80%	35	69%
2014.11-2021.01	25	18	72%	16	64%