Thanks very much for the time and efforts that you have put into reviewing the previous version of the manuscript. We really appreciate all your comments and suggestions that have enabled us to improve the manuscript. The following is a point-to-point response to the reviewer's comments. We have studied comments carefully and have made correction which we hope meet with approval. Revised portion are marked in red in the revised paper.

The paper "Two mega sand and dust storm events over northern China in March 2021: transport processes, historical ranking and meteorological drivers" by Ke Gui et al. investigates two remarkable sand and dust storm (SDSs) occurred on March 15–20, 2021 and March 27–29, 2021. The study characterizes the origins, transport processes, magnitudes of impact, and meteorological causes of these two SDS events, through satellite and ground-based observations combined with atmospheric reanalysis data. The study falls within the scope of ACP. The manuscript is well-written/structured, the presentation clear, and the language fluent. However, the submitted study is subject to major deficiencies in principal ACP evaluation criteria. Here are some of my main comments which I think will help the authors to improve their manuscript.

Response: We thank the reviewers for their professional advice. In the revised manuscript, we have changed the principal objective of this paper to emphasize aerosol optical, microphysical, and radiative properties during two mega SDS events rather than dust transport processes as in the initial manuscript, which is considered to be partially overlapping with several existing studies. After combining your suggestions and those from the first reviewer, we have made numerous improvements to the structure and content of this study, mainly including:

1) The title has been changed from "Two mega sand and dust storm events over northern China in March 2021: transport processes, historical ranking and meteorological drivers" to "*Record-breaking dust loading during two mega dust storm events over northern China in March 2021: aerosol optical/radiative properties and meteorological drivers*".

2) The major difference from the first draft is that the revised draft introduces continuous observations from a sun photometer located in the Beijing area to characterize the aerosol optical, microphysical, and radiative properties during these two SDS events. Two new sections have been introduced, including "Aerosol Optical and Microphysical Properties" in Section 3.2 and "Direct Aerosol Radiative Forcing" in Section 3.3. Please refer to the revised manuscript for details of these changes that were made.

3) The section 3.1 in the initial manuscript has been rewritten by using multi-source satellite fusion AOD, the Himawari-8 dust RGB composite images and corrected visibility observations that were not covered in several existing studies. As a result, the PM_{10} observations, individual MODIS AODs, Suomi NPP VIIRS Imagery and OMPS UVAI used in the initial manuscript have been removed from the revised manuscript.

4) In the revised manuscript we sufficiently discussed the similarities and differences with published related studies to highlight the innovative aspects of this study.

For detailed revisions, please refer to the following sections.

(1) The submitted manuscript in general presents limited novel concepts, ideas, tools, or data. An exception is the historical ranking of the dust events, although this is a secondary objective of the manuscript. To be more specific, as stated by the authors, the manuscript's principal objective is to "characterize the origins, transport processes, magnitudes of impact, and meteorological causes of these two SDS events". With respect to the most significant of the two SDSs events discussed here, the event on March 15–20, the meteorology and impact of the 3.15 SDS is discussed in Filonchyk et al., 2022 "Characteristics of the severe March 2021 Gobi Desert dust storm and its impact on air pollution in China", while the transport processes are provided by Liang et al. (2021) "Revealing the dust transport processes, magnitudes of impact, and meteorological causes" have already been discussed. A recommendation would be to focus on the second event of March 27–29, 2021 which has not been discussed so far and on the historical ranking of the dust event, which however, according to manuscript and extend of material, is interpreted as a secondary objective.

Response: We thank the reviewers for their professional advice. We regret that this study conducted some redundant analyses (some overlapping with existent studies) or similar data were used. After a careful reading of the literature mentioned by the reviewers, including but not limited to these two papers (i.e., Filonchyk et al., 2022 and Liang et al., 2021), we have revised the principal objective of this study, and also made substantial adjustments to the main structure of this study, with the aim of differentiating it from previous studies and also to highlight the innovative aspects of this study. The specific modifications are as follows:

1) The title has been changed to "*Record-breaking dust loading during two mega dust storm* events over northern China in March 2021: aerosol optical/radiative properties and meteorological drivers".

2) In the introduction section, the purpose of our study is clearly presented as follows: "To date, several studies (e.g., Liang et al., 2021; Filonchyk, 2022; Filonchyk and Peterson, 2022) have been conducted to characterize the severe SDS event in March 2021. Most of these studies have focused on investigating the evolution and transport processes of the dust plume during the 3.15 event and assess its impact on the air quality by using particulate matter (PM₁₀) concentration observations and individual satellite retrieval products. However, few studies have been carried out on the optical, microphysical, and radiative properties of aerosols during the March 2021 SDS events, which are critical to accurately assess the weather and climate effects associated with enhanced dust loadings. Furthermore, these existing studies focus on the 3.15 event, and the 3.27 event, which also has a huge impact, has not received sufficient attention. Therefore, it is essential to combine the two events to elucidate their similarities and differences in terms of dust sources, aerosol optical, microphysical, and radiative properties, and meteorological drivers.". 3) The parts (i.e., section 3.1 in the initial manuscript) that were considered to overlap with previous studies were completely removed. In the revised manuscript, a new section 3.1 has been introduced with the following main purpose:

"Several studies have been performed to reveal the transport processes of dust aerosols during the 3.15 event and their impacts on near-surface air quality using PM_{10} concentration as indicator (Liang et al., 2021; Filonchyk, 2022; Filonchyk and Peterson, 2022). However, no studies have focused on the 3.27 event and the differences between the two events have not been explored. This section will provide an overview of these two SDS events based on satellite RGB images, horizontal visibility observations, and multi-satellite fusion to reveal the similarities and differences between them from different perspectives." Please refer to the revised manuscript for details of these changes that were made.

4) To achieve the principal objective of this study (i.e., aerosol optical/radiative properties), we introduced the continuous observations from a sun photometer located in the Beijing area to characterize the aerosol optical, microphysical, and radiative properties during these two SDS events. Two new sections have been introduced, including "*Aerosol Optical and Microphysical Properties*" in Section 3.2 and "*Direct Aerosol Radiative Forcing*" in Section 3.3. Please refer to the revised manuscript for details of these changes that were made. Briefly, the following are some of the valuable conclusions obtained from this study in terms of aerosol optical, microphysical and radiative properties.

Lines 30-35 in Abstract: "Despite the shorter duration of the 3.27 event relative to the 3.15 event, sun photometer and satellite observations in Beijing recorded a larger peak AOD (~2.5) in the former than in the latter (~2.0), which was mainly attributed to the short-term intrusion of coarse-mode dust particles with larger effective radii (~1.9 μ m) and volume concentrations (~2.0 μ m³ μ m⁻²) during the 3.27 event. The direct aerosol radiative forcing (DARF) induced by dust was estimated to be -92.1 and -111.4 W m⁻² at the top of the atmosphere, -184.7 and -296.2 W m⁻² at the surface, and +92.6 and +184.8 W m⁻² in the atmosphere in Beijing during the 315 and 3.27 event, respectively."

(2) Providing the objectives of the study, it is stated by the authors that "although these two studies have strengthened our understanding of the 3.15 mega SDS event in 2021, the sources, three-dimensional evolutionary features during transport processes, historical ranking, and local meteorological anomalies of the 3.15 and 3.27 SDS events have not yet been elucidated." However, with respect to ACP evaluation criteria of "giving proper credit to related work and clearly indicate their own new/original contribution?", the significant published study of Filonchyk et al. (2022), presenting the characteristics of the severe March 2021 Gobi Desert dust storm and its impact on air pollution in China is not mentioned, and the outcomes not compared, nor discussed in terms of discrepancies, similarities – although substantial – and conclusions. A strong recommendation is to extensively discusses the similarities/differences/conclusions of the related published studies, and build on top of the previous studies.

Response: We regret that some existing studies have not been sufficiently presented and discussed to distinguish their similarities and differences from the present study. In the revised manuscript, we introduced the objectives of this study by comparing it with other similar studies, and these sentences are reintroduced below to emphasize this point.

In the introduction section: "To date, several studies (e.g., Liang et al., 2021; Filonchyk, 2022; Filonchyk and Peterson, 2022) have been conducted to characterize the severe SDS event in March 2021. Most of these studies have focused on investigating the evolution and transport processes of the dust plume during the 3.15 event and assess its impact on the air quality by using particulate matter (PM_{10}) concentration observations and individual satellite retrieval products. However, few studies have been carried out on the optical, microphysical, and radiative properties of aerosols during the March 2021 SDS events, which are critical to accurately assess the weather and climate effects associated with enhanced dust loadings. Furthermore, these existing studies focus on the 3.15 event, and the 3.27 event, which also has a huge impact, has not received sufficient attention. Therefore, it is essential to combine the two events to elucidate their similarities and differences in terms of dust sources, aerosol optical, microphysical, and radiative properties, and meteorological drivers."

In addition, based on the previous studies, we fully justify or discuss the results or conclusions of this study. For example:

Lines 355-345: "CMA station records show that, during the 3.15 event, horizontal visibility first reached a minimum on March 15 in most of NC, including Gansu, southwestern Inner Mongolia, Ningxia, northern Shaanxi, northern Shanxi, Hebei and Beijing, with the number of stations with daily mean visibility below 500 m reaching 19, due to dust plume deposition. Such remarkable contribution of dust aerosols to air quality is also supported by the results revealed by Filonchyk (2022) using PM_{10} observations. Influenced by SDS, PM_{10} concentrations in some regions of China were found to exceed 7000.0 µg m⁻³ on March 15. Similarly, Liang et al. (2021) claimed that the 3.15 SDS event was the most severe in China in the past decade, with PM_{10} concentration in Beijing reaching 6450 µg m⁻³. Our results show that on March 15, the instantaneous surface horizontal visibility was below 50m near the lower limit of the monitoring threshold at seven sites (Fig. S2)."

Lines 369-373: "Here, multi-satellite fusion provides more available retrievals and regional details by incorporating the observed dust loading at different satellite transit times than the individual satellite data sources used in the previous similarity studies (e.g., Filonchyk, 2022; Filonchyk and Peterson, 2022)."

Lines 469-472: "Generally, the instantaneous DARF on March 28 estimated in this study was stronger than similar studies previously performed during several strong SDS events, such as in Beijing in May 2017 (Filonchyk et al., 2021) and over the Indo-Gangetic Basin in May 2018 (Tiwari et al., 2019)."

Lines 498-451: "These findings are consistent with the results of Jin et al. (2022) using inverse modelling and Liang et al. (2021) using backward trajectory simulations. They revealed that wind-blown dust emissions originated from both China and Mongolia contribute to the SDS events that occur in spring 2021."

References:

- Filonchyk, M., Peterson, M. and Hurynovich, V.: Air pollution in the Gobi Desert region: Analysis of dust-storm events, Q. J. R. Meteorol. Soc., 147(735), 1097–1111, doi:10.1002/qj.3961, 2021.
- Filonchyk, M.: Characteristics of the severe March 2021 Gobi Desert dust storm and its impact on air pollution in China, Chemosphere, 287(P3), 132219, doi:10.1016/j.chemosphere.2021.132219, 2022.
- Filonchyk, M. and Peterson, M.: Development, progression, and impact on urban air quality of the dust storm in Asia in March 15 18, 2021, Urban Clim., 41, 101080, doi:10.1016/j.uclim.2021.101080, 2022.
- Jin, J., Pang, M., Segers, A., Han, W., Fang, L., Li, B., Feng, H., Lin, H. X., and Liao, H.: Inverse modeling of the 2021 spring super dust storms in East Asia, Atmos. Chem. Phys. Discuss. [preprint], https://doi.org/10.5194/acp-2021-1062, in review, 2022.
- Tiwari, S., Kumar, A., Pratap, V. and Singh, A. K.: Assessment of two intense dust storm characteristics over Indo – Gangetic basin and their radiative impacts: A case study, Atmos. Res., 228(5), 23–40, doi:10.1016/j.atmosres.2019.05.011, 2019.
- Liang, P., Chen, B., Yang, X., Liu, Q., Li, A., Mackenzie, L. and Zhang, D.: Revealing the dust transport processes of the 2021 mega dust storm event in northern China, Sci. Bull., doi:10.1016/j.scib.2021.08.014, 2021.

(3) A third point is related to the similarities in satellite and ground-based observations. With respect to ground-based in-situ observation, the PM10 data from CNEMC (http://www.cnemc.cn), have already been provided, presented and discussed in previous studies. Similarly the reanalysis dataset. Similarly, MODIS and CALIOP EO of the SDS 3.15 event. The authors could use different datasets, not implemented in previous studies, instead of datasets already used. Some suggestions of satellite-based EO datasets not used in previous studies could include the Metop IASI DOD, Sentinel 3A/B DODs, the MIDAS DOD dataset, and more.

Response: We thank the reviewers for their professional advice. We apologize for using a similar dataset to the previous study for this study. To address this issue, in the revised manuscript, we removed the datasets (including PM_{10} observations, individual MODIS AODs, Suomi NPP VIIRS Imagery and OMPS UVAI) already used in the previous studies and added some new data sources (including multi-source satellite fusion AOD, Himawari-8 dust RGB composite images and corrected visibility observations) that have not yet been covered. Moreover, continuous observations and retrievals of the optical, microphysical and radiative properties of aerosols during these two SDS events were performed using a sun photometer located in the Beijing area. In terms of the use of CALIOP data, in the revised manuscript, we separated the contribution of dust from the total extinction coefficient (see also the detailed response in the next comment), which is very different

from the previous studies (e.g., Filonchyk et al., 2021; Liang et al., 2021) that focused only on the classification or identification of dust types. In general, we cannot get information such as dust aerosol loading intensity from dust aerosol type classification, instead dust extinction coefficients can do it.

These newly added data sources are described as follows:

Lines 120-128:

"2.1.1 Combined AOD from MODIS and VIIRS

Aerosol optical depth (AOD) is the column-integrated light extinction by aerosol particles. In this study, Level 2 daily AOD at 550 nm retrieved from Moderate Resolution Imaging Spectroradiometer (MODIS) on board the Terra and Aqua satellites and from Visible Infrared Imaging Radiometer Suite (VIIRS) on board the Suomi National Polar-Orbiting Partnership (SNPP) satellite during the 3.15 and 3.27 SDS events are used to examine the downstream propagation of the dust plume. It is found that MODIS and VIIRS AOD agree well with the Aerosol Robotic Network (AERONET) observations (Hsu et al. 2019; Sayer et al. 2019). We regridded Level 2 AOD products from the three sensors, which were derived using the Dark Target (DT) and Deep Blue (DB) algorithms, respectively, to a $0.1^{\circ} \times 0.1^{\circ}$ grid and averaged among them to produce daily combined AOD. Note that only the AOD data flagged as good quality (quality flag = 2) or very good quality (quality flag = 3) was used."

Lines 200-220:

"2.1.4. Himawari-8 dust RGB composite imagery

Himawari-8 geostationary satellite operated by Japan Meteorology Agency (JMA) provides images of East Asia and Western Pacific Region at a frequency of every 10 min, day and night (Bessho et al., 2016). This allows for monitoring the source, genesis and movement of dust at high temporal resolution. Two brightness temperature (BT) differences (between 12.4 and 10.4 μ m and between 10.4 and 8.6 μ m) and the BT at 10.4 μ m are rendered to red–green–blue (RGB) beams to highlight the presence of Aeolian dust (Shimizu, 2020). To highlight the dust phenomenon, we use gamma value correction to enhance high/low brightness intensity pixel values (BYTE). The formula for such correction is

$$BYTE = \left[\frac{BT - MIN}{MAX - MIN}\right]^{\frac{1}{\Gamma}}$$
(3)

where BT is brightness temperature differences (i.e., red and green beams) or BT for blue beam, MIN and MAX represent the ranges of BT differences/BT, and Γ is the gamma value. Here, we use recommended dust RGB composite thresholds for Himawari-8 (Shimizu, 2020): MIN (MAX) is -4.0 (2.0), 0.0 (15.0), 261.0 (289.0) for red, green and blue beams, respectively, and Γ is 1.0, 2.5 and 1.0, respectively. In this study, we use Himawari-8 dust RGB composite imagery to illustrate the source, genesis and movement of dust plumes during the two SDS events."

Lines 221-267:

"2.2. Observation datasets

2.2.1. Sun photometer

A Cimel CE-318 sun photometer installed on the roof of the Chinese Academy of Meteorological Sciences (CAMS), Beijing, China (39.93°N, 116.32°E; 106 m), has been in operation since 2012. This sun photometer, which is named "Beijing-CAMS", operates in both the China Aerosol Remote Sensing Network (CARSNET) and AERONET and is calibrated at Izaña, Tenerife, Spain, together with the AERONET program (Che et al., 2009). Beijing-CAMS is the main site of CARSNET, but measurements are also uploaded to the AERONET data archive. The measurements of aerosol optical, microphysical and radiative properties at the Beijing-CAMS site can provide a reference for assessing several aspects of the dust aerosol loading intensity, evolution and its impact during the downstream transport of the dust plume.

In this study, the cloud-screened instantaneous AOD data at multiple wavelengths were calculated by using the ASTPwin software (Cimel Electronique), and extinction Ångström exponents (EAE) were calculated from the AODs for wavelengths of 440 and 870nm. The fine-mode fraction (FMF) is described as the fraction of fine-mode particles of total AOD_{440nm}. To perform a spatio-temporal synergistic analysis between satellite and ground-based observations, the AOD measurements in two adjacent channels (i.e., 440 and 675nm) from sun photometer was interpolated to 550nm for satellite retrieval, using a second-order polynomial fit to ln (AOD) vs. ln (wavelength) (Eck et al., 1999).

The aerosol microphysical properties, including volume size distributions (dV(r)/dlnr)in 22 size bins for particle radii 0.05–15 µm; the total, coarse-mode, and fine-mode volume concentrations (Volume_t, Volume_c, and Volume_f, respectively); the total, coarse-mode, and fine-mode aerosol effective radii (R_{eff_p} , R_{eff_c} , and R_{eff_f} , respectively); the absorption AOD (AAOD); SSA; and the absorption Angström exponent (AAE), were retrieved from the almucantar sky irradiance measurements in conjunction with measured spectral AOD at 440, 670, 870, and 1020nm using the algorithms of Dubovik et al. (2002, 2006).

The direct aerosol radiative forcing (DARF in $W m^{-2}$) was calculated by the radiative transfer module similar to the inversion of AERONET under the assumption of cloud-free conditions (García et al., 2008, 2012). The DARF at the top of the atmosphere (TOA) and the Earth's surface (bottom of the atmosphere, BOA) was defined as the difference in the shortwave radiative fluxes with and without aerosol effects in Eqs. (4) and (5) as follows:

$$DARF_{TOA} = F_{TOA}^{\uparrow 0} - F_{TOA}^{\uparrow}$$

$$DARF_{BOA} = F_{BOA}^{\downarrow} - F_{BOA}^{\downarrow 0} \tag{5}$$

where $F_{TOA}^{\uparrow 0}$ and $F_{BOA}^{\downarrow 0}$ denote the broadband fluxes with no aerosols at TOA and BOA, respectively. DARF_{TOA} is the reflection of solar radiation by aerosols back to space, while DARF_{BOA} indicates the combined effects of absorption and scattering of solar radiation by aerosols. The findings of García et al. (2008) show that the error for the observed solar radiation at the surface on a global scale was $+2.1 \pm 3.0\%$ for an overestimation of about $+9 \pm 12 W m^{-2}$. Subsequently, the DARF at atmosphere (DARF_{ATM}) was defined as the difference between DARF_{BOA} and DARE_{TOA} as follow:

 $DARF_{ATM} = DARF_{TOA} - DARF_{BOA}$

(6)

Defined this way, a negative value for DARF indicates aerosol cooling effects, while positive values imply warming.

2.2.2. Horizontal visibility

Horizontal visibility is closely related to air quality and can be an important indicator of the quality of the atmospheric environment in most scenarios (Gui et al., 2021). In this study, the hourly visibility observations from ~1600 national surface meteorological observation stations across NC provided from the China Meteorological Administration (CMA) during the two SDS events were used to characterize the impacts of the dust plume on air quality. In order to minimize the effect of relative humidity (RH) on horizontal visibility, RH values > 40 and < 99% were converted to the equivalent visibility in dry conditions (i.e., RH < 40%). The correction formula is expressed as VIS/VIS(dry) = 0.26 + 0.4285 log₁₀ (100 – RH) (Rosenfeld et al., 2007). The hourly visibility data accompanied by the presence of fog or precipitation were excluded from this study."

Regarding the DOD dataset recommended by the reviewer, we retained the MODIS inversion of the DOD dataset in this study, considering the limitations such as data timeliness and computational resources. Nevertheless, to our knowledge, we are the first to use the MODIS-retrieved DOD to describe these two SDS processes that occurred in northern China in March 2021. In addition, to ensure the accuracy of the long-term DOD time series, we also included a comparative analysis of the data with the long-term AERONET observations in NC at 7 sites.

Lines 169-179: "Here we compare the combined daily MODIS DOD against AERONET stations in NC (Fig. S1). It should be noted that to date, there is no valid method to derive DOD from AERONET AOD measurements. Therefore, we use coarse-mode AOD (AODc) from AERONET measurements as a proxy for DOD (Pu and Ginoux, 2018; Song et al., 2021) to compare with our DOD datasets in March for NC.

For this purpose, we use the AERONET Version 3 spectral deconvolution algorithm (SDA) daily products (Level 2.0 data). Given that AERONET SDA only provide the AODc at 500 nm, the AERONET AODc is converted to 550 nm in this study using the Ångström exponent to compare with MODIS DOD retrievals. For March dust retrievals, between 2000 and 2021, there are 121 MODIS daily mean DOD retrievals collocated with 7 AERONET sites located within NC (Fig. S1). Results showed that the MODIS DOD is in good agreement with AERONET AODc (Person correlation coefficient = 0.82), although the former generally overestimated latter in NC (root-mean-square error = 0.28)."



Figure S1: Scatter plot of the daily mean MODIS DOD against the AERONET coarse-mode AOD (AODc) retrieved at 550nm. The 1-to-1 line and linear regression line are shown by black dotted and red solid lines, respectively. The number of sites (Sites), matchups (N), Pearson correlation coefficient (R), slope, and root mean square error (RMSE) of the linear regression are indicated in the lower right of the panel.

(4) The study could be improved in terms of providing and outlining more extensively the information of the scientific methods and assumptions used, which are only briefly discussed, in terms of limitations of the implemented datasets. For instance, very few information on the quality assurance criteria is provided. With respect to CALIPSO, as QA only the CAD score is mentioned that it is implemented, while in the literature, significantly more QA filters are recommended (e.g. Tackett et al., 2018). Another point could be the particulate depolarization ratio, when observed lower to 0.3, indicates mixtures of dust and non-dust components, thus the provided CALIPSO information, considered in the study as pure-dust, overestimates the actual dust extinction coefficient. With respect to MODISretrieved DOD, which are the limitations in terms of Deep-Blue algorithm, Angström exponent, SSA and the quality of the DOD product? Which are the attenuation effects thus the limitations - of the different datasets (active and passive) and how do they compare in terms of observations? The SDSs are transported detached in extended domains over China, and how does this compare with only the surface-based in-situ? There is a wealth of datasets incorporated in the study, however, the dataset observations are not extensively spatio-temporarily inter-compared to account for and extract the closest representation and actual state of the SDSs.

Response: We thank the reviewers for their professional advice.

1) In order to completely separate the contribution of dust to the total aerosol extinction, we used the following method in the revised manuscript. In addition, with reference to similar previous studies, more quality control methods were used.

Lines 179-200: "To characterize the vertical profile of the dust plume, the Level 2 daily 532 nm aerosol profile product (05kmAPro, V4.21) that contains aerosol depolarization, backscatter, and extinction profile from CALIOP/CALIPSO (Winker et al., 2010) was used. We use the methodology in Yu et al. (2015) to derive the dust extinction profile. To reduce uncertainty, only high-quality extinction profile data with a CAD (cloud aerosol discrimination) score of between -100 and -90 were used. The aerosol profile product also provides an extinction quality control flag (Ext_QC) to indicate problematic retrievals. This study only uses layers with Ext_QC values of 0, 1, 18, and 16 (Winker et al., 2013). For each aerosol backscatter coefficient profile, we infer the ratio of dust to total backscatter (f_d) at each altitude from the following equation:

$$f_d = \frac{(\delta - \delta_{nd})(1 + \delta_d)}{(\delta_d - \delta_{nd})(1 + \delta)} \tag{2}$$

where δ is CALIOP observed particulate depolarization ratio, δ_d and δ_{nd} are a priori knowledge of depolarization ratios of dust and non-dust aerosols respectively. To account for various types of non-dust aerosols with different depolarization ratio and for the variability of dust shape and size, we follow Song et al. (2021) and use the f_d that was based on the mean of the lowest ($\delta_d = 0.30$ and $\delta_{nd} = 0.07$) and the highest ($\delta_d = 0.20$ and $\delta_{nd} =$ 0.02) dust scenario. By assuming a dust lidar ratio (LR) (i.e., extinction-to-backscatter ratio) of 40 sr at 532 nm (Yu et al., 2015), we derive dust extinction coefficient (DEC) profile from dust backscatter coefficient."

2) Regarding the uncertainty of the MODIS-retrieved DOD mentioned by the reviewer. Admittedly, the uncertainty in each input variable contributes to the uncertainty in the DOD retrievals, but it is difficult for us to distinguish between them. Therefore, to ensure the accuracy of the MODIS-retrieved DOD, we evaluated it comparatively using the inversion results of AERONET with reference to the previous studies (e.g., Pu and Ginoux, 2018; Song et al., 2021). Also, some other uncertainty descriptions were included.

Lines 161-179: "MODIS dust detection is subject to a number of uncertainties. Over land, the derived MODIS DOD here denotes the coarse-mode (aerodynamic diameters larger than 1 µm) contribution of dust only and does not include its fine-mode contribution. Estimates by Kok et al. (2017) suggest that the exclusion of submicron dust aerosol could induce around 3% underestimation of the global atmospheric dust mass load and around 15% underestimation of the global DOD. Terra and Aqua DOD values at daily and monthly scales have previously been validated with AERONET stations globally (Pu and Ginoux, 2018; Song et al., 2021). Spatially, when comparing the MODIS-derived DOD climatology with the DOD retrieved from the Cloud–Aerosol Lidar with Orthogonal Polarization (CALIOP) aboard the Cloud–Aerosol Lidar and Infrared Pathfinder Satellite Observation (CALIOP) satellite, the climatological mean of the MODIS DOD generally compares well with CALIOP (Pu and Ginoux, 2018; Song et al., 2021). Here we compare the combined daily MODIS DOD against AERONET stations in NC (Fig. S1). It should be noted that to date, there is no valid method to derive DOD from AERONET AOD measurements. Therefore, we use coarse-mode AOD (AODc) from AERONET measurements as a proxy for DOD (Pu and Ginoux, 2018; Song et al., 2021) to compare with our DOD datasets in March for NC.

For this purpose, we use the AERONET Version 3 spectral deconvolution algorithm (SDA) daily products (Level 2.0 data). Given that AERONET SDA only provide the AODc at 500 nm, the AERONET AODc is converted to 550 nm in this study using the Ångström exponent to compare with MODIS DOD retrievals. For March dust retrievals, between 2000 and 2021, there are 121 MODIS daily mean DOD retrievals collocated with 7 AERONET sites located within NC (Fig. S1). Results showed that the MODIS DOD is in good agreement with AERONET AODc (Person correlation coefficient = 0.82), although the former generally overestimated latter in NC (root-mean-square error = 0.28)."



Figure S1: Scatter plot of the daily mean MODIS DOD against the AERONET coarse-mode AOD (AODc) retrieved at 550nm. The 1-to-1 line and linear regression line are shown by black dotted and red solid lines, respectively. The number of sites (Sites), matchups (N), Pearson correlation coefficient (R), slope, and root mean square error (RMSE) of the linear regression are indicated in the lower right of the panel.

3) Regarding the limitations of different data sets

Different datasets, especially satellite retrieval products, have many limitations, such as being influenced by clouds and algorithmic assumptions. Therefore, in order to address these issues, we have introduced some new data sources and presentations in the revised manuscript. First, we used a multi-satellite (including MODIS/Terra, MODIS/Aqua and VIIRS) combined AOD to improve the coverage of the valid daily average AOD values (Fig. 4). Secondly, we introduced the dust RGB composite images from Himawari-8 to monitor the source, genesis and movement of dust at high temporal resolution (Fig.2). The dust RGB composite images are able to provide a more spatially continuous evolution of the dust plume than the satellite inversion of aerosol-related variables, as it does not need to rely on various retrieval assumptions.



Figure 2: Evolution of dust plumes (magenta) as revealed by Himawari-8 dust RGB composite images at 13:00 CST during (a–f) the 3.15 SDS event (March 15–20, 2021) and (g–i) the 3.27 SDS event (March 27–29, 2021), respectively. Overlaid on the RGB imagery is the ERA5 daily mean wind vectors at 10m.



Figure 4: Evolution of MODIS and VIIRS combined daily mean AOD during (a–f) the 3.15 event and (g–i) the 3.27 event.

4) To illustrate and extract the closest representation and actual state of the SDS, we carried out a spatial-temporal intercomparison between the multi-source satellite-retrieved AODs and ground-based sun photometer observations in Beijing.

Lines 364-369: "The accuracy of the MODIS and VIIRS instantaneous AOD values retrieved using individual algorithms (i.e., DT and DB algorithms) was confirmed by a synergistic comparison with ground-based sun photometer observations located in Beijing (i.e., Beijing-CAMS site) (Fig. 5). To complete the spatial collocation, the satellite-retrieved AOD values is represented by the average value of all the satellite product pixels within a 25 km radius (Sayer et al., 2013) around the Beijing-CAMS site. The results show that both the MODIS and VIIRS instantaneous AOD values are in high agreement with ground-based observations."



Figure 5: Daily variation of AOD at 550 nm (top row), EAE between 440 and 870 nm (middle row), and FMF (bottom row) at Beijing-CAMS site during (a-c) the 3.15 event and (d-e) the 3.27 event. The instantaneous AOD values from the MODIS and VIIRS sensors, which were derived using the Dark Target (DT) and Deep Blue (DB) algorithms, respectively, are given in the top panel.

(5) The ground-based lidar observations are a novelty however, in terms of dataset. At this point, the authors could provide more information to characterize the SDSs, including the intensive properties of dust (e.g. LR used), the pre-processing of the dataset and retrieval (e.g. Raman/Klett), the temporal averaging techniques, observed wavelength dependencies, and more.

Response: We are very grateful to the reviewers for their scientific advice, which has been fully considered in the revised version. Following your comments, we have added the following description to clarify the methods and assumptions used for the inversion of the dust extinction profiles by the AD-Net lidar observations.

Lines 278-286: "In this study, the DEC profile at 532 nm provided by an AD-Net site named "Zamynuud" were used to explore the impacts of long-range dust transport on downstream areas. Each lidar from AD-Net takes a 5-minute measurement every 15 minutes, generating an observation data file with the vertical resolution of 30m. For AD-Net, the Klett's inversion method was employed to derived the extinction coefficient, after applying a geometrical-form-factor correction (Sugimoto et al., 2003). With the assumption of external mixing of dust and spherical aerosols, the ratio (R) of contribution of dust in the extinction coefficient is calculated as follows:

$$R = \frac{(\delta_a - \delta_2)(1 + \delta_1)}{(1 + \delta_a)(\delta_1 - \delta_2)} \tag{7}$$

where δ_a is the observed aerosol depolarization ratio (ADR). δ_1 and δ_2 are ADRs of dust and air-pollution aerosols. The values of δ_1 and δ_2 were determined empirically and are 0.35 and 0.05, respectively (Sugimoto et al., 2003; Shimizu et al., 2004)."



Figure 7: Time-height evolution of dust extinction coefficient (km^{-1}) at 532 nm retrieved by ground-based Lidar (location of the site shown in Fig. 1) during the 3.15 and 3.27 events.

Although similar studies have been performed for the specific SDSs and the study domain, improvements of the present work may lead to very interesting contribution in the literature, due to the very special and diverse conditions encountered over the Eastern part of Asia. At its present state, and taking the above comments under consideration, I suggest to the journal to reject the paper. However, I would suggest the authors to go through the entire manuscript once more and build on top of the published studies and on this work, follow the suggested improvements/recommendations, and maybe focus on more novel ideas such the effects of the studied SDSs in terms of atmospheric chemistry/physics (i.e. RT, IN/CCN, dust deposition rate), and then I would encourage them to resubmit it.

Response: Once again, we sincerely thank the reviewers for these structural suggestions, which have led to a qualitative improvement of our study. Finally, we thank the reviewers for providing us with some potentially novel ideas that will be taken into account in our future studies.