

Limitations in representation of physical processes prevent successful simulation of PM_{2.5} during KORUS-AQ

Katherine R. Travis¹, James H. Crawford¹, Gao Chen¹, Carolyn E. Jordan^{1,2}, Benjamin A. Nault³, Hwajin Kim⁴, Jose L. Jimenez⁵, Pedro Campuzano-Jost⁵, Jack E. Dibb⁶, Jung-Hun Woo⁷, Younha Kim⁸, Shixian Zhai⁹, Xuan Wang¹⁰, Erin E. McDuffie¹¹, Gan Luo¹², Fangqun Yu¹², Saewung Kim¹³, Isobel J. Simpson¹⁴, Donald R. Blake¹⁴, Limseok Chang¹⁵, Michelle J. Kim¹⁶

¹NASA Langley Research Center, Hampton, VA, USA

²National Institute of Aerospace, Hampton, VA, USA

³Center for Aerosol and Cloud Chemistry, Aerodyne Research Inc. 45 Manning Road Billerica, MA, USA

⁴Department of Environmental Health Sciences, Graduate School of Public Health, Seoul National University, Seoul 08826, Korea

⁵Cooperative Institute for Research in the Environmental Sciences, University of Colorado, Boulder, Colorado, USA

⁶Earth System Research Center, University of New Hampshire, Durham, NH, USA

⁷Department of Civil and Environmental Engineering, Konkuk University, Seoul, Republic of Korea

⁸Energy, Climate, and Environment (ECE) Program, International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria

⁹John A. Paulson School of Engineering and Applied Sciences, Harvard University, Cambridge, MA, USA

¹⁰City University of Hong Kong, Kowloon, HK

¹¹Department of Energy, Environmental, and Chemical Engineering, Washington University in St. Louis, St. Louis, MO, USA

¹²Atmospheric Sciences Research Center, University at Albany, Albany, NY, USA

¹³University of California, Irvine, Irvine, CA, USA

¹⁴Department of Chemistry, University of California, Irvine, California, USA

¹⁵Air Quality Research Division, National Institute of Environmental Research, Incheon, Republic of Korea

¹⁶Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA, USA

Correspondence to: K. R. Travis, katherine.travis@nasa.gov

Abstract.

High levels of fine particulate matter (PM_{2.5}) pollution in East Asia often exceed local air quality standards. Observations from the Korea United States-Air Quality (KORUS-AQ) field campaign in May and June 2016 showed that development of extreme pollution (haze) occurred through a combination of long-range transport and favorable meteorological conditions that enhanced local production of PM_{2.5}. Atmospheric models often have difficulty simulating PM_{2.5} chemical composition during haze, which is of concern for the development of successful control measures. We use observations from KORUS-AQ to examine the ability of the GEOS-Chem chemical transport model to simulate PM_{2.5} composition throughout the campaign and identify the mechanisms driving the pollution event. In the surface levelAt the surface, the model underestimates campaign-average sulfate aerosol by -64% but overestimates nitrate aerosol by +36%. The largest underestimate in sulfate occurs during the pollution event in conditions of high relative humidity, where models typically struggle to generate the high

~~concentrations~~ elevated sulfate concentrations due to missing heterogeneous chemistry in aerosol liquid water in the polluted boundary layer. Hourly surface observations show that the model nitrate bias is driven by an overestimation of the nighttime peak. In the model, nitrate formation is limited by the supply of nitric acid, which is biased by +100% against aircraft observations. We hypothesize that this is due to a large missing sink, which we implement here as a factor of five increase in dry deposition. We show that the resulting increased deposition velocity is consistent with observations of total nitrate as a function of photochemical age. The model does not account for factors such as the urban heat island effect or the heterogeneity of the built-up urban landscape resulting in insufficient model turbulence and surface area over the study area that likely results in insufficient dry deposition. Other species such as NH_3 could be similarly affected but were not measured during the campaign. Nighttime production of nitrate is driven by NO_2 hydrolysis in the model, while observations show that unexpectedly elevated nighttime ozone (not present in the model) should result in N_2O_5 hydrolysis as the primary pathway. The model is unable to represent nighttime ozone due to an overly rapid collapse of the afternoon mixed layer and excessive titration by NO . We attribute this to missing nighttime heating driving deeper nocturnal mixing that would be expected to occur in a city like Seoul. This urban heating is not considered in air quality models run at large enough scales to treat both local chemistry and long-range transport. Key model failures in simulating nitrate, mainly overestimated daytime nitric acid, incorrect representation of nighttime chemistry, and an overly shallow and insufficiently turbulent nighttime mixed layer, exacerbate the model's inability to simulate the buildup of $\text{PM}_{2.5}$ during haze pollution. To address the underestimate in sulfate most evident during the haze event, heterogeneous aerosol uptake of SO_2 is added to the model which previously only considered aqueous production of sulfate from SO_2 in cloud water. Implementing a simple parameterization of this chemistry improves the model abundance of sulfate but degrades the SO_2 simulation implying that emissions are underestimated. We find that improving model simulations of sulfate has direct relevance to determining local vs. transboundary contributions to $\text{PM}_{2.5}$. During the haze pollution event, the inclusion of heterogeneous aerosol uptake of SO_2 decreases the fraction of $\text{PM}_{2.5}$ attributable to long-range transport from 66% to 54%. Locally-produced sulfate increased from 1% to 4625% of locally-produced $\text{PM}_{2.5}$, implying that local emissions controls would have a larger effect than previously thought. However, this additional uptake of SO_2 is coupled to the model nitrate prediction which affects the aerosol liquid water abundance and chemistry driving sulfate-nitrate-ammonium partitioning. An additional simulation of the haze pollution with heterogeneous uptake of SO_2 to aerosol and simple improvements to the model nitrate simulation results in 30% less sulfate due to 40% less nitrate and aerosol water, and results in an underestimate of sulfate during the haze event. Future studies need to better consider the impact of model physical processes such as dry deposition and nighttime boundary layer mixing on the simulation of nitrate and the effect of improved nitrate simulations on the overall simulation of secondary inorganic aerosol (sulfate+nitrate+ammonium) in East Asia. Foreign emissions are rapidly changing, increasing the need to understand the impact of local emissions on $\text{PM}_{2.5}$ in South Korea to ensure continued air quality improvements.

1 1 Introduction

South Korea enacted legislation in 2018 to address local air pollution, which ranked 13th in the world for the worst annual average fine particulate matter (PM_{2.5}) exposure levels (Energy Policy Institute, 2019). Ambient PM_{2.5} was the 5th highest risk factor for human health in South Korea in 2018, leading to over 20,000 attributable deaths (GBD, 2021). The government plans to reduce the number of days with pollution warnings (PM_{2.5} > 90 µg m⁻³ for two hours) by 50% in 2022 from the 89 that occurred in 2016 (Kim et al., 2018). The reduction of PM_{2.5} levels through policy measures relies on a thorough understanding of pollution sources and the ability of models to simulate potential control measures. Modeling studies have concluded that on average, approximately half of observed PM_{2.5} in South Korea is attributable to long-range transport from China (Lee et al., 2017; Choi et al., 2019; Jung et al., 2019; Kumar et al., 2021). Quantifying the effect of long-range transport relies on regional to global-scale models that trade-off the high resolution needed to resolve urban scales with a large enough domain to represent both the study area and upwind source regions. Estimates of long-range transport This finding is based on models that have received limited testing of their ability to simulate PM_{2.5} chemical composition, particularly during extreme pollution events. Quantifying the effect of long-range transport relies on regional to global-scale models that trade-off the high resolution needed to resolve urban scales with a large enough domain to represent both the study area and upwind source regions. This evaluation is critical as the contribution of long-range transport to PM_{2.5} in South Korea may be declining due to effective emission controls in China (Han et al., 2021), increasing the need to understand the impact of local emissions on pollution events.

Across East Asia, densely populated regions experience haze events with extremely high levels of PM_{2.5} frequently associated with periods of elevated relative humidity and low daytime mixed layer heights (An et al., 2019). These conditions are favorable for increasing gas-particle partitioning of aerosol precursors. ~~In haze, secondary~~ Secondary inorganic aerosol (secondary sulfate+nitrate+ammonium ≡ SNA) is often the dominant component of PM_{2.5} in haze, but models have difficulty SNA, particularly simulating sulfate, during these periods likely due to missing conversion of SO₂ in aqueous aerosol (Wang et al., 2014; Zheng et al., 2015a, 2015b; Shao et al., 2019). ~~More generally, the~~ The MICS-Asia multi-model comparison showed that the annual contribution of SNA to total PM_{2.5} varied by a factor of two across models, and ~~the models also~~ overpredicted the gas-particle partitioning of nitrate (Chen et al., 2019). In the global AeroCom III intercomparison, models differed in their annual concentrations of nitrate and its precursor, nitric acid, by factors of thirteen and nine, respectively (Bian et al., 2017). Models also struggle to represent organic aerosol, ~~(Zhao et al., 2016),~~ overestimating primary organic aerosol (POA) but underestimating secondary organic aerosol (SOA, ~~Zhao et al., 2016~~), possibly likely due to missing sources from anthropogenic precursors (Nault et al., ~~2020~~ 2021a). ~~This~~ The wide range of model performance in simulating PM_{2.5} composition emphasizes the urgent need for better model constraints to improve our understanding of the sources and conditions driving haze events.

~~Improving model representation of sulfate chemistry cannot be considered entirely separately from model nitrate biases. In the atmosphere, aqueous-phase chemistry is a major source of sulfate, where clouds provide the dominant source of liquid water (Herrmann et al., 2015). Recent studies have hypothesized that the high aerosol liquid water content (ALWC) associated with PM_{2.5} during extreme pollution events in East Asia allows for significant sulfate production not considered in most models (Wang et al., 2014; Zheng et al., 2015a, 2015b; Shao et al., 2019). Levels of ALWC are very sensitive to). Early modeling work suggested that this chemistry must be occurring generally in the polluted boundary layer in the United States and Europe (Kasibhatla et al., 1997). aerosol nitrate (Ge et al., 2012; Sun et al., 2018)A. The aqueous SO₂ oxidation pathway(s) for SO₂ oxidation in ALWC are uncertain in part due to poor understanding of aerosol acidity (An et al., 2019). Aerosol acidity, a key factor partially controlling nitric acid - nitrate partitioning (Guo et al., 2016). Nitrate aerosol has the greatest impact on ALWC which affects acidity (Ge et al., 2012; Sun et al., 2018). Due to this coupled nature of SNA aerosol, improving model representation of one component (i.e sulfate) cannot be considered entirely separately from the rest (nitrate + ammonium). Accurate simulation of aerosol composition and water uptake is required to interpret satellite observations of aerosol optical depth (AOD) (Saide et al., 2020) and evaluate the response to emission changes.~~

The Korea United States-Air Quality campaign (KORUS-AQ), conducted in May and June 2016 in South Korea (Crawford et al., 2021), provides an extensive set of ground and aircraft-based observations that can further constrain model simulations of the chemical and physical drivers of PM_{2.5}. The campaign included a haze event with concentrations exceeding local air quality standards, characterized by rapid buildup of SNA aerosol. Throughout KORUS-AQ, surprisingly high levels of nighttime ozone, particularly prevalent during haze, appeared to drive nighttime nitrate formation through N₂O₅ hydrolysis (Jordan et al., 2020). This was attributed to elevated nocturnal mixed layer heights (MLH). Zhai et al. (2021) found a severe model overestimate in nighttime nitrate during KORUS-AQ, implying a failure to correctly simulate these conditions. We use the GEOS-Chem chemical transport model applied at high resolution (0.25° × 0.3125°) over East Asia to investigate model representation of PM_{2.5} mass and chemical composition during KORUS-AQ. We specifically evaluate model performance during the conditions governing the development of haze pollution such as elevated relative humidity, increased SNA, and high nighttime ozone levels. We demonstrate how addressing deficiencies in model physical processes (e.g., nighttime mixing, deposition) are fundamental to the successful simulation of PM_{2.5}.

2 KORUS-AQ observations

The KORUS-AQ campaign (Crawford et al., 2021) was a joint field campaign organized by South Korea's National Institute of Environmental Research (NIER) and the United States National Aeronautics and Space Administration (NASA). KORUS-AQ included twenty flights using the NASA DC-8 aircraft from May 1 to June 9, 2016, complemented by heavily instrumented ground sites including aerosol composition at Olympic Park and the Korea Institute of Science and Technology (KIST) in Seoul. The NIER maintains the extensive AirKorea monitoring network for hourly observations of PM_{2.5} mass, ozone, and

other pollutants, with 329 sites available during the campaign, including locations near Olympic Park and KIST. There were four distinct meteorological periods during KORUS-AQ, described in Peterson et al. (2019). These included a dynamic period characterized by a series of frontal passages (Dynamic Period, May 1-16), dry, clear, and stagnant conditions (Stagnant Period, May 17-22), long-range transport and haze conditions with high humidity and extensive cloud cover (Transport/Haze Period, May 25-31), and blocking conditions limiting transport (Blocking Period, June 1-7). Details on the impact of the different meteorological periods on PM_{2.5} are provided in Jordan et al. (2020). We focus on the Seoul Metropolitan Area (SMA) with the highest density of KORUS-AQ observations and the highest PM_{2.5} levels observed by the AirKorea network during the campaign. Crawford et al. (2021) provides a full listing of all observations made during KORUS-AQ. Table 1 describes the aircraft and ground observations used in this work.

3 GEOS-Chem model

We use the GEOS-Chem chemical transport model (CTM) in version 12.7.2 (doi: 10.5281/zenodo.3701669) to simulate KORUS-AQ. The model is driven by assimilated meteorological data from the NASA Global Modeling and Assimilation Office (GMAO) Goddard Earth Observing System Forward-Processing (GEOS-FP) atmospheric data assimilation system. GEOS-FP has a native horizontal resolution of $0.25^\circ \times 0.3125^\circ$, which we apply with the nested version of GEOS-Chem (Chen et al., 2009) over East Asia ($70^\circ - 140^\circ\text{E}$, $15^\circ\text{S} - 55^\circ\text{N}$) using boundary conditions from a global simulation at $2.0^\circ \times 2.5^\circ$ with a 1-month initialization period. The model has 47 vertical layers, with the first layer centered at approximately 60 m above the surface. Model timesteps are 20 min (chemistry) and 10 min (transport) as recommended by Philip et al. (2016).

Global emissions are from the Community Emissions Database System (CEDS) inventory (Hoesly et al., 2018) overwritten by the KORUSv5 anthropogenic and shipping emissions (Woo et al., 2020) for Asia ($60^\circ - 146^\circ\text{E}$, $10^\circ\text{S} - 54^\circ\text{N}$) developed for the KORUS-AQ campaign. The translation from KORUSv5, provided using the SAPRC99 chemical mechanism, to the GEOS-Chem mechanism is given in Table S1. We apply sector-specific diurnal variation from the Multi-resolution Emission Inventory for China (MEIC) as in Miao et al. (2020) to the monthly KORUSv5 emissions. Natural emissions are from the Global Emissions Initiative (GEIA, Bouwman et al., 1997) for ammonia and from MEGANv2.1 (Guenther et al., 2012) for biogenic species. We include lightning emissions (Murray et al., 2012), biomass burning emissions (GFED4s, Werf et al., 2017), soil NO_x emissions (Hudman et al., 2012), and volcanic SO₂ emissions (Carn et al., 2015). Table 2 shows the emissions inventory for key emitted species in the nested East Asia domain for May 2016.

Model dry deposition for gas-phase species is based on the resistance-in-series scheme from Wesely (1989) as implemented by Wang et al. (1998), where species deposition is limited by aerodynamic resistance, quasi-laminar layer resistance, and canopy or surface resistance. Species with low surface resistance, such as HNO₃, are limited in their deposition velocity by aerodynamic resistance only. Aerosol deposition is from Zhang et al. (2001). The original model wet deposition scheme is

described by Liu et al. (2001) for water-soluble aerosols and Amos et al. (2012) for gases. Wet deposition includes scavenging from moist convective updrafts and rainout and washout from precipitation. We include the revised wet deposition scheme of Luo et al. (2019) that uses an empirical washout rate for nitric acid two orders of magnitude higher than the previous value and replaces the standard constant value for in-cloud condensation water content with the value calculated by the meteorological fields (GEOS-FP). GEOS-Chem uses a non-local boundary layer mixing scheme (Holtslag and Boville, 1993; Lin and McElroy, 2010) where mixing is calculated explicitly from meteorological variables provided by GEOS-FP (i.e. sensible and latent heat flux, temperature, friction velocity). The mixing height is restricted from dropping below a minimum mechanical mixing depth, defined as a function of local friction velocity (Lin and McElroy, 2010).

The GEOS-Chem HO_x-NO_x-VOC-ozone-halogen-aerosol mechanism includes improvements to PAN chemistry (Fischer et al., 2014), isoprene oxidation (Fisher et al., 2016; Travis et al., 2016; Chan Miller et al., 2017), halogen chemistry (Sherwen et al., 2016), Criegee intermediates (Millet et al., 2015), and methyl, ethyl, and propyl nitrates (Fisher et al., 2018). Heterogeneous aerosol uptake of HO₂ produces H₂O₂ (Mao et al., 2013), with a reactive uptake coefficient (γ) of 0.2 (Jacob, 2000). We implement aromatic chemistry from Yan et al. (2019) for the simulation of KORUS-AQ.

We use the model “simple scheme” for organic aerosol (OA) where OA is generated using fixed empirically derived yields from isoprene, monoterpenes, biomass burning, and anthropogenic fuel combustion (Pai et al., 2020). This scheme includes an emitted hydrophobic component (OCPO) with an assumed organic-mass-to-organic carbon (OM:OC) ratio of 1.4 that is aged to a hydrophilic oxygenated component (OCPI) with an OM:OC ratio of 2.1. Secondary organic aerosol (SOA) is a lumped product (SOAS) with a molecular weight of 150 g mol⁻¹. For comparison to observations, primary organic aerosol (POA) is defined as OCPO and SOA is the sum of OCPI and SOAS. The sulfate-nitrate-ammonium (SNA) aerosol simulation (Park, 2004) includes the addition of metal-catalyzed oxidation of SO₂ (Alexander et al., 2009), sulfur oxidation by reactive halogens (Chen et al., 2017), and improved implementation of aerosol cloud-processing and revised uptake coefficients for NO₂ (Holmes et al., 2019). Uptake of N₂O₅ on SNA includes dependence on aerosol water, organic coatings, nitrate aerosol fraction, and particulate chloride (McDuffie et al., 2018). SNA partitioning is calculated with ISORROPIA v2.2 (Pye et al., 2009). The model includes accumulation mode (SALA) and coarse mode (SALC) sea salt aerosol (Alexander et al., 2005; Jaeglé et al., 2011) and dust in four size bins (DST1 to 4) (Fairlie et al., 2010), where the first bin and 38% of the second bin are included in PM_{2.5}. The recommended definition of dry PM_{2.5} is given by Eq 1.

$$PM_{2.5} = SO_4^{2-} + NO_3^- + NH_4^+ + BC + OCPO * 1.4 + OCPI * 2.1 + SOAS + SALA + DST1 + DST2 * 0.38, \quad (1)$$

The AirKorea PM_{2.5} observations provided by NIER are obtained using the beta-ray attenuation method (BAM-1020, Table 1). We do not adjust modeled PM_{2.5} for any measurement relative humidity effects as the BAM-1020 has been shown to perform well against federal reference method monitors (Le et al., 2020).

Specific details of production of model nitric acid (HNO_3), the gas-phase precursor to aerosol nitrate ($\text{NO}_3^- = \text{pNO}_3$), are provided below as KORUS-AQ provides detailed observations of this chemistry. Reactions R1-R6 describe model production of HNO_3 from oxidation of NO_2 (R1), aqueous uptake and reaction of N_2O_5 , NO_2 , and NO_3 on aerosol (R2, R4, R5), aqueous uptake and reaction of N_2O_5 and NO_3 in cloud water (R3, R5), ~~heterogeneous halogen chemistry (Table S2),~~ and oxidation of VOCs by the nitrate radical (R6). Heterogeneous halogen chemistry is provided in Table S2. In R2, aqueous uptake and reaction of N_2O_5 with particle chloride (Cl^-) produces nitryl chloride (ClNO_2) with a yield (ϕ) of 1 on sea salt aerosol and zero on all other aerosol types.



4 Simulation of $\text{PM}_{2.5}$ during KORUS-AQ

Figure 1a shows the model simulation of daily average $\text{PM}_{2.5}$ (Eq. 1) compared to the observed average of the 15 AirKorea sites within the GEOS-Chem grid box containing the major SMA monitoring sites (KIST and Olympic Park). These two sites are in close proximity to the AirKorea monitors (Fig. 1b). Campaign average $\text{PM}_{2.5}$ is $29 \mu\text{g m}^{-3}$, but this increases to $53 \mu\text{g m}^{-3}$ during the Transport/Haze period (Table 3). The model reproduces the low $\text{PM}_{2.5}$ during the Dynamic period, the increase during the Transport/Haze period, and the variable concentrations during the Blocking period. Across the campaign, the model underestimates $\text{PM}_{2.5}$ (NMB = -15%) due to a low bias during the Stagnant period and the initial build-up during the Transport/Haze period. This model performance is similar to Choi et al. (2019) using a different GEOS-Chem configuration.

Figure 2 compares observed $\text{PM}_{2.5}$ composition against the model for the gridbox containing the KIST ground site. Speciated $\text{PM}_{2.5}$ is derived from KIST PM_1 composition and AirKorea $\text{PM}_{2.5}$ mass as described in Section S2, as the difference in composition between PM_1 and $\text{PM}_{2.5}$ is expected to be minor (Sun et al., 2020; Schlosser et al., 2022). $\text{PM}_{2.5}$ -Measured composition from the KIST HR-ToF-AMS instrument (Table 1), representative of PM_1 (Guo et al., 2021), is used to speciate daily average $\text{PM}_{2.5}$ from the AirKorea sites (Fig. 1a). Jordan et al. (2020) showed that speciated PM_1 was generally representative of $\text{PM}_{2.5}$ mass throughout KORUS-AQ, except for the Transport/Haze period when $\text{PM}_{2.5}$ significantly exceeded PM_1 . The strong correlation between $\text{PM}_{2.5}$ and PM_1 during the campaign implied growth of PM_1 to larger sizes. Dust is not a major component of $\text{PM}_{2.5}$ at the surface after May 9th, as further discussed in Section S1. Therefore PM_1 composition likely

represents the composition of $PM_{2.5}$ with the exception of a small contribution from primary aerosol species. Sun et al. (2020) showed that $PM_{2.5}$ can be up to 50% greater than PM_{10} in polluted, humid environments and the mass at sizes $>PM_{10}$ is secondary (not BC or POA). We remove BC and POA from observed $PM_{2.5}$ and scale the remaining components (SNA, SOA) to the remaining $PM_{2.5}$. The resulting speciated $PM_{2.5}$, derived from KIST PM_{10} composition and AirKorea $PM_{2.5}$ mass, is provided for each meteorological period in Table 3. Figure 2 and Table 3 include the ALWC associated with $PM_{2.5}$, calculated for the observations using the E-AIM IV thermodynamic model (Clegg and Brimblecombe, 1990; Clegg et al., 1998; Massucci et al., 1999; Wexler and Clegg, 2002; (Nault et al., 2021)), and ISORROPIAv2.2 (Pye et al., 2009) in GEOS-Chem. During KORUS-AQ, Kim et al. (2022) found that ISORROPIAv2.2 provided similar results as the E-AIM model, reproducing E-AIM pH within ~ 0.4 units.

On average, the model simulates SNA within 20%. However, this is due to compensating biases which has implications for controlling precursor species. The primary campaign average model biases are: The model underestimated sulfate (-64%), overestimated nitrate (+36%), and underestimated SOA (-43%). The excess model nitrate is the primary driver of overestimated ALWC (+82%). During the Stagnant period, the model low bias is primarily due to underestimated SOA ($-9 \mu g m^{-3}$). This may be due to missing local production from emissions of semi- and intermediate-volatility volatile organic compounds (S/IVOCs, McDonald et al., 2018) and aromatics (Nault et al., 2018), primarily attributable to solvents and vehicle emissions (Shin et al., 2013a, 2013b; Simpson et al., 2020). During the Dynamic and Blocking periods, the model $PM_{2.5}$ bias is within 20% of the observations but with overestimated nitrate and underestimated sulfate. The model severely underestimates sulfate during the Transport/Haze period ($-11 \mu g m^{-3}$, Table 3) suggesting that the model fails to reproduce the processes driving the pollution episode. As described by Jordan et al. (2020) showed, both ground and aircraft observations during KORUS-AQ showed that cloudy and humid conditions during the Transport/Haze period increased $PM_{2.5}$ through heterogeneous production of SNA and this is not included in the model.

The KORUS-AQ aircraft observations included detailed daytime (available from ~ 8 am to 4pm KST) aerosol and gas-phase observations that we use to determine the cause of model sulfate and nitrate biases and their regional extent. Model SOA biases will be the subject of future work as here they do not contribute to $PM_{2.5}$ exceedances ($50 \mu g m^{-3}$ daily average in 2016). The KORUS-AQ campaign included frequent sampling along a repeated flight pattern or “stereoroute” over the SMA up to three times a day, supplemented by less frequent flights to investigate specific source regions or transport events (Crawford et al., 2021). Figure S2 shows the high data density in the SMA compared to the rest of the study region. We use the 55 descents over Olympic Park from the SMA stereoroute to compare against the daily surface observations shown in Fig. 2.

Figure 3 shows the mean daytime aircraft profiles of sulfate and nitrate for the descents over Olympic Park below 2 km separated by the same meteorological periods as Fig. 2. The corresponding profiles for SO_2 and nitric acid are shown in Fig. S3. The model is sampled along the flight tracks and both the model and the observations are averaged to the model grid,

60 timestep and nearest vertical 0.5 km. Similar to the daily surface average, the model underestimates daytime sulfate below 2 km with the most severe bias ($-8 \mu\text{g m}^{-3}$ in the lowest altitude bin of 0.5 km) occurring during the Transport/Haze period. Unlike in the daily surface average, the model underestimates daytime nitrate below ~ 1 km with the exception of the Dynamic period ~~when nitrate is in good agreement~~. The model nitrate underestimate could be ~~partially~~ related to ~~the low~~ low model RH bias in model RH of up to ~~-83%~~ (Blocking-Stagnant period, 39-28 vs. 47-31%) below 0.5 km (Fig. S3) or overestimated mixed layer height (Oak et al., 2019). ~~Recent work suggests that If the model RH simulation was unbiased, we would expect an improved simulation of nitrate as the minimal RH bias during the Dynamic period corresponds to the best nitrate simulation (Fig. 3, Fig. S3).~~ Model aerosol dry deposition may also be too fast but this ~~effect~~ would increase model concentrations by only $\sim 10\%$ (Emerson et al., 2020).

70 There is no available measurement of $\text{PM}_{2.5}$ from the aircraft to provide ~~a similar~~ scaling from PM_1 to $\text{PM}_{2.5}$ as was done ~~in~~ for Fig. 2 and described in Section S2. However, any increase to the observed profiles of PM_1 sulfate or nitrate to account for possible growth to larger sizes would exacerbate the model underestimate of these species. The discrepancy between the model low to minimal bias against daytime aircraft nitrate observations (Fig. 3) and the overestimate against daily average nitrate at the KIST ground site (Fig. 2) implies a failure of the model to represent nighttime chemical production. We investigate the possible causes of overestimated daily average model nitrate in Section 5 and underestimated model sulfate in Section 6.

75 5 Model errors representing the nitrate diurnal cycle

The discrepancy between the model daytime ~~and vs.~~ daily average performance for nitrate demonstrates the need to evaluate the compare the model mean nitrate diurnal cycle ~~against the nitrate fraction of $\text{PM}_{2.5}$ derived from KIST observations as described in Section 4~~. Figure 4a shows the observed nitrate component of $\text{PM}_{2.5}$ calculated from PM_1 composition and $\text{PM}_{2.5}$ mass as described in Section S2. ~~that b~~ Between 6am and 6pm KST (daytime) the model bias is minimal ($< 1 \mu\text{g m}^{-3}$) while 80 the bias from 6pm to 6am KST (nighttime) is $+3 \mu\text{g m}^{-3}$. As described in Section 3, the model has a newly revised treatment of wet scavenging that significantly reduces the model nitrate and nitric acid biases present in previous model versions (Luo et al., 2019). Without this improvement, the model would have an average nighttime bias of $+7 \mu\text{g m}^{-3}$. Figure S4 shows daily precipitation in Seoul from the Korea Meteorological Administration (KMA, 2021) which ~~was~~ infrequent and negligible in the later part of the campaign. The model underestimate in total precipitation across the campaign is minimal (121 vs. 112 85 mm). Insufficient wet scavenging is unlikely to be the cause of the remaining model nitrate bias.

We perform a sensitivity test (Table 4) to determine the relative impact of daytime (R1) vs. nighttime (R2-R5, Section 3) production of HNO_3 on the model bias by shutting off the nighttime reactions. Figure 4c shows that the main model nighttime pathway is aerosol uptake of NO_2 (R4) with a small contribution from N_2O_5 hydrolysis (R2/3) in the early morning hours. Figure 4a shows that removing nighttime chemistry results in improved early morning agreement (1am to 8am KST) but the

evening overestimate (8pm to 1am KST) is less affected. Jordan et al. (2020) showed observational evidence for significant nighttime production of nitrate by N_2O_5 hydrolysis (R2). We use the removal of nighttime chemistry to hypothesize that part of the model nighttime bias is due to excess daytime HNO_3 that has not yet been lost to deposition and is converted to nitrate as conditions become thermodynamically favorable for partitioning to the aerosol-phase. The dominance of NO_2 uptake over N_2O_5 hydrolysis in the model suggests that there are additional errors in simulated nighttime chemistry.

5.1 Sensitivity of model nitrate bias to gas-phase precursors

Inorganic aerosol ammonium nitrate (NH_4NO_3) is formed by dissolution of HNO_3 , which reacts in the aqueous phase with ammonia (NH_3) to establish a thermal equilibrium with NH_4NO_3 . The conditions that favor NH_4NO_3 are generally cool and humid (i.e., nighttime) and characterized by high NH_3 and HNO_3 concentrations relative to sulfate (Guo et al., 2016). We calculate that average nighttime RH (temperature) in the SMA is ~~7469%~~ ~~(290K-291K)~~ compared to the model value of ~~7467%~~ ~~(288K-291K)~~, indicating that significant errors in RH or temperature are not the cause of nighttime biases. Overproduction of model nighttime nitrate could be due to overestimated NH_3 if this species limits NH_4NO_3 production. In South Korea, and generally East Asia, NH_4NO_3 is limited by availability of HNO_3 . This is due to high levels of NH_3 (~10 ppb) observed in East Asia, attributable to non-agricultural sources such as transportation (Song et al., 2009; Phan et al., 2013; Link et al., 2017; Sun et al., 2017; Chang et al., 2019; Lim et al., 2022). The model reproduces the expected high concentration of NH_3 with an average of 9 ppb at Olympic Park. Ibikunle et al. (2020) performed a rigorous thermodynamic assessment of KORUS-AQ observations confirming that aerosol was always sensitive to HNO_3 in polluted conditions. Nitrate-limited SNA thermodynamics were observed in similar conditions in China and successfully represented by ISORROPIA v2.2 in GEOS-Chem (Zhai et al., 2021).

Few datasets exist to further test the performance of HNO_3 - pNO_3 partitioning in the model but KORUS-AQ observations provide this opportunity. This partitioning is described by Eq. 2, where the ratio of pNO_3 to total nitrate ($\text{TNO}_3 = \text{HNO}_3$ and pNO_3), known as ϵNO_3 , is impacted by temperature, relative humidity, and aerosol composition (Guo et al., 2016, 2017).

$$\epsilon\text{NO}_3 = \frac{\text{pNO}_3}{\text{HNO}_3 + \text{pNO}_3} \quad (2)$$

Accurate simulation of ϵNO_3 is critical to regulating the deposition of TNO_3 as HNO_3 deposits more rapidly than pNO_3 (Nenes et al., 2021). Figure 5 shows ϵNO_3 as a function of RH for the observations and the model for the same domain as Fig. 3 below 1.5 km. While the model represents the increase of ϵNO_3 with RH, model ϵNO_3 is generally underestimated, particularly at lower RH (<50%). This ~~low bias in ϵNO_3 could be due to overestimated HNO_3 , as a result of underestimated ammonia, not measured during the lower RH and associated higher temperatures generally prevent excess HNO_3 (denominator of Eq. 2) from partitioning to campaign, the aerosol phase. We discuss the possibility of overestimated model HNO_3 below or errors in model temperature and RH.~~ As ϵNO_3 is underestimated in the model, excess partitioning to the aerosol-phase is not a cause

of the model nitrate overestimate shown in Fig. 2. The successful performance of ISORROPIAv2.2 during KORUS-AQ is also evident from the comparison against the E-AIM model in Kim et al. (2022).

Figure 6a shows vertical profiles of observed and modeled HNO_3 for the Olympic Park descents. The model overestimates HNO_3 in the lowest bin (0.5 km) by +1600 ppt or +100%. This high bias persists across most of the study domain except over the ocean south of 34°N (Fig. 7) where local emissions have a small impact and loss to deposition is slow. During average daytime conditions (~50% RH, 295K), model ϵNO_3 is ~0.3, indicating that while the aerosol is HNO_3 -limited, higher temperatures and low RH also prevent the excess model HNO_3 from partitioning to aerosol. A simulation turning off South Korean emissions shows that local sources contribute ~50% to model HNO_3 concentrations below 0.5 km (Fig. 6a). Thus, while model errors in emissions or chemistry could be a cause of the bias, an overestimated lifetime of HNO_3 against dry or wet deposition could also play a role. We evaluate these possibilities further in Section 5.2.

5.2 Causes of overestimated daytime HNO_3

KORUS-AQ provides aircraft and surface observations that provide additional constraints on the model HNO_3 bias of +100% described in Section 5.1. We use observations of NO_2 and OH from aircraft to evaluate whether NO_x emissions or production from R1 ($\text{NO}_2 + \text{OH}$) are overestimated. Figure 6b shows that model NO_2 is underestimated by -40% below 0.5 km. This is partially due to the expected model inability to resolve the highest observed levels of NO_2 in an urban region, illustrated by the larger standard deviation in the observations compared to the model. However, given the same emissions inventory used here (KORUSv5), a set of eight models varied in their biases for NO_x against KORUS-AQ aircraft observations from a minimal underestimate (-7%) to a large overestimate (+56%) depending on model configuration (Park et al., 2021). Thus, model biases could be due to a range of factors including underestimated emissions, inaccuracies in the emission diurnal cycle, or overestimated mixed layer heights. Errors in any of these factors that could increase model NO_2 , such as decreased mixed layer heights or (i.e., increased emissions), would be expected to increase the model overestimate of HNO_3 . Figure 6c shows that the model bias in OH is small (+20%) and well within measurement uncertainty (+32%) and therefore it is unlikely that model errors in R1 (Fig. 6d) could cause the model HNO_3 bias of +100%.

The fastest removal pathways for HNO_3 are wet and dry deposition. The model implementation of these processes is described in Section 2.3. The recently revised model wet scavenging scheme has improved annual average model simulations of HNO_3 , but the effect on HNO_3 during KORUS-AQ is limited as precipitation was infrequent after the beginning of the campaign as discussed above. Section S2-S3 further discusses the impact of this scheme on KORUS-AQ nitrate and HNO_3 but errors in wet deposition are unlikely to be the cause of overestimated model HNO_3 . Section S2-S3 also describes other possible loss pathways to dust, seasalt, or production of ClNO_2 from N_2O_5 hydrolysis (e.g. Jeong et al., 2019) that have negligible effects on the model HNO_3 and nitrate.

Previous attempts to improve model nitrate overestimates invoked an unknown sink of HNO_3 in the model (Heald et al., 2012; Weagle et al., 2018), as uncertainties in precursor emissions, the rate of N_2O_5 hydrolysis (R2/R3) or gas-phase production (R1), OH concentrations, and HNO_3 dry deposition velocity (Vd_{HNO_3}) could not explain model nitrate biases. We similarly conclude that an unknown loss process must be a main cause of the daytime model overestimate in HNO_3 and associated evening nitrate bias during KORUS-AQ that occurs as conditions become more favorable for partitioning HNO_3 to pNO_3 . This unknown loss process could be a larger underestimate in dry deposition than has been previously considered, as constraints from KORUS-AQ show that uncertainties in emissions, nighttime production (R1-R5), and wet deposition are not the cause. Heald et al. (2012) ruled out dry deposition after assuming an uncertainty of a factor of two. Here, the increase in Vd_{HNO_3} required to reproduce observed HNO_3 (Fig. 6a) is a factor of five. A similar increase in Vd_{HNO_3} was invoked by Itahashi et al. (2017) in their model study of wintertime nitrate in East Asia based on the finding from Shimadera et al. (2014) that Vd_{HNO_3} (as well as NH_3 emissions and dry deposition) were the main factors driving model nitrate performance.

The increase in Vd_{HNO_3} suggested above would result in an average value of 7.5 cm s^{-1} compared to the standard model value of 1.5 cm s^{-1} . This corresponds to a maximum midday rate of 15.4 cm s^{-1} ~~cm s^{-1}~~ compared to the original value of 3.1 cm s^{-1} (Fig S8). Deposition of HNO_3 is limited only by aerodynamic resistance (and available surface area), as it readily adheres to surfaces. While the increase to Vd_{HNO_3} we suggest here is large, this could arise from factors such as increased surface area in urban or heavily forested regions and increased vertical mixing over cities due to turbulence induced by the urban heat island effect. These factors are not accounted for in the limited existing deposition velocity measurements that have been compared against models (Nguyen et al., 2015). Increased turbulence over forested regions results in higher deposition velocities (Sievering et al., 2001; Yazbeck et al., 2021), which would also be expected in an urban environment (~~i.e.~~ Keuken et al., 1990). The model does not account for increased available surface area for deposition contributed by urban buildings, or the elevated vertical mixing over cities due to the urban heat island effect (Hong and Hong, 2016; Halios and Barlow, 2018). ~~Dry deposition rates thus may be much higher than in model parameterizations that do not include a specific treatment of the urban canopy (Cherin et al., 2015) and this is the case in GEOS Chem. The model surface roughness, an important parameter governing turbulence, is only just 0.1 m in Seoul, compared to values measured between 1 and 3 for forested or urban parts of the city (Hong and Hong, 2016). (Hong and Hong, 2016).~~

Neuman et al. (2004) derived Vd_{HNO_3} from aircraft observations of power plant plumes in eastern Texas, obtaining values between 8 and 26 cm s^{-1} , values at least four times faster than reported previously. We take a similar approach to Neuman et al. (2004) to calculate Vd_{HNO_3} from KORUS-AQ observations in the SMA using the rate equation for TNO_3 as a function of photochemical age (Fig. 8, Eq. 3).

$$\text{TNO}_3(t) = \frac{\text{NO}_x(0)}{\frac{\beta}{c}-1} (e^{-ct} - e^{-\beta t}) \quad (3)$$

85 NO_x(0) is the initial NO_x mixing ratio normalized to CO (Fig. 8, 0.24 ppbv / ppbv CO), β is the first order loss rate for TNO₃, c is the first order production rate for TNO₃ ($p\text{TNO}_3 = p\text{HNO}_3 = k_{R1}[\text{OH}]$), and TNO₃(t) is observed TNO₃ as a function of photochemical age (t). As ~~the~~ production of TNO₃ was constrained by observed OH, and ~~assuming~~ the main loss of TNO₃ (β) is from deposition of HNO₃, the unknown for TNO₃ evolution is the deposition rate. The full ~~details of this calculation are~~ provided in Section S3S4.

90

Figure 8 shows NO_x, TNO₃, and the other NO_x oxidation products of total peroxy nitrates (ΣPNs) and the sum of alkyl- and multi-functional nitrates (ΣANs) as a function of photochemical age. All species are normalized by background subtracted CO. NO_x is continuously depleted at a rate of 0.31 hr⁻¹, implying continued production of TNO₃, ΣPNs , and ΣANs . This loss rate corresponds to a lifetime of 3.2 hrs that is similar to the lifetime of 4.8 hrs for NO₂ against conversion to HNO₃ (R1) using
95 the SMA average OH of 5.2×10^{-6} molec cm⁻³. From Eq. 3, we derive a loss rate (β) of 13.9 cm s⁻¹ that best fits the observed change in TNO₃ with aging. As deposition of pNO₃ is slow, we assume that $V_{d\text{HNO}_3} = V_{d\text{TNO}_3}$. All three NO_x oxidation products (TNO₃, ΣPNs , ΣANs) exhibit similar behavior with production outpacing loss until approximately three hours of aging, where loss appears to balance production and concentrations remain relatively constant. There is likely large uncertainty in the derived photochemical ages shown in Fig 8, as the aircraft did not follow plumes as in Neuman et al. (2004). However, our derived
00 NO_x lifetime is consistent with average SMA conditions and is not affected by our choice of observed altitude range, suggesting that the aging represents true chemical processing.

Figure 8 shows that the slower value for midday $V_{d\text{HNO}_3}$ in the original model (3.1 cm s⁻¹) poorly represents observations compared to the faster value obtained in Fig. 6 (15.4 cm s⁻¹). We calculate that the original deposition rate would correspond
05 to a first order loss rate for TNO₃ of only 0.07 hr⁻¹ (assuming a 1.5 km boundary layer height) and thus observed TNO₃ should increase with photochemical age, which is not supported by the observed relationship in Fig. 8. The factor of five increase in $V_{d\text{HNO}_3}$, constrained only using observed HNO₃, implies a similar loss rate of TNO₃ as derived in Fig. 8 and leads to the observed behavior where after initial production, the normalized mixing ratio remains constant. This analysis supports the hypothesis given above, that existing observations supporting lower values for $V_{d\text{HNO}_3}$ (Nguyen et al., 2015) may
10 underrepresent deposition in regions with greater turbulence and available surface area such as in cities like Seoul. Deposition of atmospheric pollutants such as nitric acid on buildings generates ‘urban grime’ that may photolyze and produce NO_x and HONO (Baergen and Donaldson, 2013, 2016). ~~This urban grime could be a source of HONO (;~~ Zhang et al., 2016). This source and may be larger than previously thought if models underestimate nitric acid deposition.

15 Figure 4a shows the impact to the diurnal cycle of model nitrate from increasing model $V_{d\text{HNO}_3}$ by a factor of five. The rapid late afternoon /early evening increase in model nitrate (Fig. 4a) is largely resolved and ~~the~~ model ~~simulation of~~ HNO₃ is now in better agreement with aircraft observations (Fig. 6) due to a significant dampening of the HNO₃ diurnal cycle (Fig. 4b). This reduction in the HNO₃ diurnal cycle is better supported by observations of TNO₃ as discussed above. We conclude that a

key reason for the ~~overestimated-high bias in~~ daily average model nitrate ~~shown in~~ (Fig. 2) is overestimated daytime HNO₃ that ~~results in produces~~ excess ~~nighttime~~ nitrate at night when conditions become favorable (cool ~~and~~ humid) for gas to aerosol partitioning. The model overestimate is due to insufficient loss, likely underestimated dry deposition. This finding does not address possible errors ~~in in model~~ nighttime nitrate production pathways (NO₂ vs. N₂O₅) ~~), and~~ KORUS-AQ provides detailed ground observations that can be used to constrain ~~the model representation of~~ nighttime chemistry.

5.3 Errors in model nighttime production of HNO₃

Figure 4c shows that model nighttime production of HNO₃ by aerosol uptake of NO₂ (R4) is approximately twice as large as R2 (N₂O₅ hydrolysis). This contradicts the calculation from Jordan et al. (2020) that R2 is the driver of nitrate production during KORUS-AQ, particularly during the Transport/Haze period due to sufficient nighttime ozone concentrations that allow for production of the nitrate radical and N₂O₅ through R8 and R9.



Production of nitrate by N₂O₅ hydrolysis is supported by observations of ClNO₂, thought to be produced primarily by this reaction (Thornton et al., 2010). As discussed above in Section 5.2, observations of ClNO₂ at Olympic Park are elevated at night (Fig. S7). Despite recent large reductions of the uptake coefficient (γ) for NO₂ in the model (Holmes et al., 2019), NO₂ uptake still is the dominant nighttime pathway ~~for HNO₃ production in the model~~. We use observations of ozone, NO, and NO₂ at Olympic Park to determine whether errors in R7-R9 are impacting model ability to produce N₂O₅.

Figure 9 shows the mean modeled and observed diurnal cycles of ozone and NO₂ for the AirKorea sites in the model grid box (Fig. 1b) and for ozone, NO, NO₂, and NO_x at Olympic Park. Ozone might be expected to be titrated in an urban area by R7 as the mixed layer collapses in the evening, resulting in elevated NO and shutting down production of the nitrate radical (R8). This is the case in the model where nighttime ozone is <2 ppb approximately 20% of the time but this ~~never~~ occurs in the observations only twice (Fig. S9). As a result, average observed nighttime ozone is ~~24-23~~ ppb but only 13 ppb in the model (Fig. 9). The time series of observed and modeled ozone in Fig. S9 shows while the model does succeed in simulating high nighttime ozone concentrations during the Dynamic Period, characterized by higher windspeeds, ozone is incorrectly titrated at other times particularly during the buildup of the haze pollution following a frontal passage on May 24th. The implications of this excess ozone titration for the simulation of PM_{2.5} specifically during haze conditions will be further discussed in Section 6.

As shown in Fig. 9b+c, model ozone titration corresponds to excess model NO and NO₂ at night and explains the dominance of NO₂ uptake in the model over N₂O₅ hydrolysis for nighttime HNO₃ production. The model bias for NO_x is minimal during

the day, providing additional support for the level of emissions in the model, but is overestimated by ~~a factor of two~~ 50% at night. The excess model ozone titration and overestimated nighttime NO_x implies an error in nocturnal mixing. Figure 10a shows the mixed layer height (MLH) diurnal cycle measured by ceilometers at Olympic Park and Seoul National University. The aerosol gradients detected by the ceilometer to estimate MLH are less reliable at night due to the possible presence of aerosols in the residual layer (Jordan et al., 2020). We support these measurements with additional calculations of nighttime MLH from radiosonde observations of temperature and RH four times a day (Section ~~S4S5~~, Fig. S10), ~~showing that the average MLH at 3 KST could be ~300m compared to 220m in the model~~. As previously discussed in Section 5.2, in urban regions such as Seoul, the anthropogenic heat island effect and the heterogeneity of the urban land cover increase sensible heat fluxes and turbulence over non-urban areas (Halios and Barlow, 2018) and create an unstable mixed layer even at night. Min et al. (2020) showed that the nighttime mixed layer in Seoul is elevated in all seasons, and that nighttime conditions are generally unstable due to urban heat storage and anthropogenic heat release and this could explain the observed elevated nighttime MLH (Fig. 10a, Fig. S11). This effect is not captured in many meteorological models including the one used here (GEOS-CF, Section 3-), ~~instead~~ Nighttime-nighttime model sensible heat flux ~~in the model~~ is always negative (stable conditions) (Fig. 10b).

Starting at 17 KST, the model mixed layer collapses early, causing a more rapid decline in ozone than in the observations (Fig. 9a, Fig. 10a). The transition from convective daytime mixed-layer to stable nocturnal boundary layer is poorly understood (Lothon et al., 2014). The early collapse of the mixed layer has been observed in other models including the widely-used Weather Research and Forecasting (WRF) model ~~over the Baltimore-Washington, D.C. region during the NASA DISCOVER-AQ mission~~ (Hegarty et al., 2018). One possibility for the delay in this collapse is continued mixing from the last eddy of the day formed just before the sensible heat flux changes sign during the evening transition (Blay-Carreras et al., 2014). This has been hypothesized as reasons for errors in the model diurnal cycle of ozone in the ~~Southeast~~ United States (Travis and Jacob, 2019). Here, this early collapse drives excess production of nitrate from NO₂ (R4).

While addressing the shortcomings of the model mixing scheme is beyond the scope of this study, we test the sensitivity of model nitrate production to the main two problems identified above, 1) the overly rapid collapse of the afternoon mixed layer, and 2) insufficient nocturnal mixing. While model meteorology is calculated offline, mixing in the boundary layer is calculated online (Section ~~23~~), allowing us to perturb mixing parameters. We increase the nighttime MLH to 500_m (Table 4) to examine the impact on model ozone, NO, and NO₂. The effect of this change on these species is minimal (Fig. 9), similar to the findings of other model sensitivity studies that performed this same test (Oak et al., 2019; Miao et al., 2020). While the strength of model vertical mixing is sensitive to MLH, the model sensible heat flux and friction velocity have a larger impact (Holtslag and Boville, 1993), and the nighttime mixed layer will remain stable while the sensible heat flux is negative ~~regardless of MLH~~. We illustrate in Section S6 that reducing the collapse of the evening MLH without a change to the drivers of mixing (i.e., heat fluxes, friction velocity) also has negligible impact on decreasing model ozone titration (Fig. S12).

The increase in nighttime mixing in urban vs. rural regions has been addressed in the CMAQ model (Li and Rappenglueck, 2018) by using a higher value for the minimum mixing strength (eddy diffusivity) over urban areas. However, we find that this approach is insufficient to address model ozone titration without increasing the sensible heat flux to a positive value to produce an unstable mixed layer. This is illustrated in Fig. S12, where we scale the model MLH to match the profile at Olympic Park (Fig. 10a) and raise the model minimum eddy diffusivity from $0.01 \text{ m}^2 \text{ s}^{-1}$ to $1 \text{ m}^2 \text{ s}^{-1}$ over the SMA. Reducing the collapse of the evening MLH without a change to the drivers of mixing (i.e., heat fluxes, friction velocity) has negligible impact on decreasing model ozone titration (Fig. S12). In addition, the MLH at Olympic Park in the early morning hours appears inconsistent with observed ozone, likely due to the uncertainties in the measurement technique discussed above and supported by the lower values obtained from radiosonde profiles (Fig. S11). Errors in model nighttime mixing are difficult to remedy without significant revisions to the model mixing parameterizations, including implementing continued mixing from daytime eddies into the evening hours (Blay-Carreras et al., 2014) and increased parameterizing the urban heat fluxes excess sensible heat flux in urban areas (Halios and Barlow, 2018). We address the implications of these errors in-for the simulation of haze pollution events in Section 6.

00 6 Model simulation of haze buildup

The failure of models to simulate sulfate production in haze in East Asia is a current topic of intensive research and is attributable to missing sulfate production in aerosol water (Wang et al., 2014; Zheng et al., 2015a; Chen et al., 2016; Shao et al., 2019; Miao et al., 2020). ~~There~~2020). ~~There~~ has been less attention paid to the assessment of the ability of models to simulate nitrate in haze, but as nnitrate-dominated haze is a more becoming more common recent phenomenon due to the reductions in SO_2 in East Asia (Wang et al., 2020). Figure 2 and Table 3 show that the model can reproduce the increase in the nitrate component of $\text{PM}_{2.5}$ during the Transport/Haze period but overestimates absolute concentrations by ~2015%. This contributes to an 80% overestimate in ALWC. Efforts to explicitly simulate SO_2 oxidation in ALWC may be hindered by this model bias, which also impacts the rates of all other heterogeneous reactions through the increase in aerosol surface area/volume.

Figure 11a shows the hourly time series of observed and modeled nitrate at Olympic Park during the Transport/Haze period. During the haze buildup, the model initially overestimates nitrate during the day (5/24) followed by large nighttime underestimates (5/24-5/25). This is opposite to the nighttime overestimate/but daytime agreement shown in the campaign average (Fig. 4a). During the haze buildup, daytime RH remained elevated ($>50\%$, Fig. S13) and the daytime mixed layer was suppressed (Fig. S14 and Jordan et al. 2020). The model reproduces both conditions, which are favorable for SNA production. Model nitrate biases here are likely due to the errors identified in Section 5.2 (overestimated daytime HNO_3) and Section 5.3

(incorrect representation of nighttime conditions), but here the excess daytime HNO_3 in the model results in higher daytime nitrate than in the campaign average. Insufficient model sulfate during the haze event results in overestimated model pH and excess partitioning of HNO_3 to the particle phase (Guo et al., 2016). Fig. S15 shows that ϵNO_3 (the calculated fraction of TNO_3 in the aerosol phase) decreases as sulfate increases and the model sulfate bias corresponds to a difference in ϵNO_3 of ~ 0.3 .

The model underestimate of nighttime nitrate concentrations during the haze buildup must be because the rate of observed N_2O_5 hydrolysis (R2) exceeds even the erroneously high model rate of NO_2 aerosol uptake (R4). The haze buildup was characterized by a lower daytime MLH and a deeper nocturnal MLH (inferred from the lack of ozone titration) that resulted in higher nitrate production from N_2O_5 hydrolysis (Jordan et al., 2020). The model overly titrates ozone (Fig. 11c) due to insufficient nighttime mixing. We drive additional nocturnal mixing by increasing the sensible heat flux at night from slightly negative (-4 W m^{-2}) to weakly positive ($+10 \text{ W m}^{-2}$), representative of anthropogenic heat fluxes in this region (Hong and Hong, 2016; Varquez et al., 2021). To reduce the rate of R4 from overestimated NO_2 and allow for a high rate of R2, we increase the nighttime MLH over land to 300 m as suggested by the observations.

-This [sensitivity test \(Table 4\)](#) largely resolves the incorrect model ozone titration and the severe model overestimate of nighttime NO_2 on 5/23-5/24 and on 5/24-5/25 but does not remedy the early model collapse of the evening mixed layer (Fig. 11). Extending this sensitivity test past the haze buildup results in excess nighttime ozone. This may be due to the increased cloud cover during the haze buildup (Fig. S16), that could cause additional nighttime mixing over average conditions through enhancement of the urban heat island effect (Theeuwes et al., 2019). [As the meteorology in GEOS-Chem is calculated offline \(Section 3\), increasing surface sensible heat flux only impacts the boundary layer mixing parameterization but not the simulation of other meteorological fields. Future work should use a coupled system to investigate other effects of the urban heat island effect on air quality.](#)

Figure 11b shows that increased nighttime mixing allows for N_2O_5 hydrolysis (R2) to become the main nighttime pathway for HNO_3 , with a rate three times greater than NO_2 uptake (R4) in the base model. The raised mixed layer height of 300 m prevents this high rate from resulting in overestimated model nitrate. Increased model nighttime nitrate corresponds to an increase in [nighttime ALWC of 40~50%](#). We use the simulations shown in in Fig. 11 to illustrate that model errors in simulating mixed layer dynamics (overly rapid collapse of the evening mixed layer and insufficient nighttime mixing) result in errors in model chemistry. Nighttime measurements of the vertical structure of key species such as ozone, NO_2 , N_2O_5 , and HNO_3 , complemented by sensible heat flux observations, are needed to further constrain model simulations of nighttime nitrate production.

As discussed in Section 4, in addition to the above difficulties in simulating nitrate, the model fails to reproduce observed sulfate during the Transport/Haze period and this corresponds to a $15 \mu\text{g m}^{-3}$ underestimate in $\text{PM}_{2.5}$ (Table 3). Studies have

shown a strong relationship between increasing RH and conversion of gas-phase precursors to SNA in haze, indicating the occurrence of heterogeneous chemistry in ALWC (Sun et al., 2013; Liu et al., 2015; Quan et al., 2015; 2015a; Chen et al., 2016; Wu et al., 2018a). Figure 12 shows the sulfate oxidation ratio, $SOR \equiv \left(\frac{SO_4^{2-}}{SO_2 + SO_4^{2-}} \right)$ as a function of RH at Olympic Park and from aircraft observations. In the observations, SOR increases with RH, but this is missing from the model. We take the approach of Wang et al. (2014) and implement heterogeneous uptake of SO₂ on aerosol (not present in the standard model) as a function of RH according to Eq. 4,

$$k_T = \left[\frac{a}{D_g} + \frac{4}{v\gamma} \right]^{-1}, \quad (4)$$

where the mass transfer rate (k_T) at which a species is lost from the gas-phase is a function of the particle radius (a), the molecular diffusion coefficient (D_g), the mean molecular speed (v), and the reactive uptake coefficient (γ), or the probability of irreversible reaction. The value for γ depends on RH (Wang et al., 2014) according to Eq. 5.

$$\gamma = \gamma_{RH50\%} + (\gamma_{RH100\%} - \gamma_{RH50\%}) / (100\% - 50\%) \times (RH - 50\%) \quad (5)$$

The values $\gamma_{RH100\%} = 3 \times 10^{-4}$ and $\gamma_{RH50\%} = 3 \times 10^{-5}$ best fit the observations using the base model (—without the aforementioned adjustments for nitrate simulation, Table 4). These values are two orders of magnitude slower than in the original formulation of Wang et al. (2014) but similar to more recent studies (Zheng et al., 2015a; Chen et al., 2016). During the Transport/Haze period, this improves model agreement with sulfate observations at the surface (~15 $\mu\text{g m}^{-3}$ vs. Table 3: 15 $\mu\text{g m}^{-3}$) and aloft (Fig. S17). ~~average surface (Table 3, 15.4 $\mu\text{g m}^{-3}$ vs. 14.7 $\mu\text{g m}^{-3}$) and daytime aircraft (Fig. S17) sulfate observations.~~ Model agreement with daytime aircraft SO₂ observations is degraded, implying that model emissions during the Transport/Haze period are insufficient to produce both the amount of observed SO₂ and sulfate.

During the Transport/Haze period, Choi et al. (2019) estimated a contribution from transported pollution of 68%. However, the inclusion of heterogeneous uptake of SO₂ on aerosol would increase the amount of both locally produced and transported pollution, as the model attributes ~60% of SO₂ to foreign sources and ~40% to local emissions (Fig. S17). We simulate PM_{2.5} with heterogeneous conversion of SO₂ as described above, and then remove South Korean emissions in order to investigate changes to the fraction of transported pollution. Figure 13 shows the model PM_{2.5} composition for each case during the Transport/Haze period (—with an additional 15 $\mu\text{g m}^{-3}$ of PM_{2.5} in the model sensitivity with heterogeneous uptake of SO₂). In the original model, foreign transport accounts for 66% of PM_{2.5} (25 $\mu\text{g m}^{-3}$ vs 38 $\mu\text{g m}^{-3}$), but this fraction is reduced to 54% (29 $\mu\text{g m}^{-3}$ vs 53 $\mu\text{g m}^{-3}$) in the revised model as the local contribution (13 vs. 24 $\mu\text{g m}^{-3}$) makes up a greater fraction of the increase. Locally produced sulfate increases from only 1% (<1 $\mu\text{g m}^{-3}$) to 25% (6 $\mu\text{g m}^{-3}$) of local PM_{2.5}, implying that local SO₂ controls could have a greater effect on PM_{2.5} levels than previously thought. Locally produced nitrate increases from 6 $\mu\text{g m}^{-3}$ to 8 $\mu\text{g m}^{-3}$. The total amount of model nitrate (local + foreign) decreases slightly at the surface and aloft (Fig. S17) which we attribute to the impact of sulfate on reducing eNO₃ described above ~~and shown in Fig. S15 but this does not resolve the model nitrate biases described in Section 5.~~

The previous calculations only account for the missing model sulfate during the Transport/Haze period, and do not account for the incorrect model representation of nighttime nitrate production or overestimated model HNO₃. This accounts for the

dramatic increase in ALWC in Fig. 13, which is already overestimated in the original model formulation as shown in Fig 2. Given the uncertainties in revising the model nitrate simulation, we did not assess the policy implications for improving model nitrate on local vs. transported pollution. However, A simple test ~~however~~ of the haze buildup with the heterogeneous SO₂ uptake described above with the and including inclusion of a factor of five increase to V_{dHNO_3} and, increased nighttime mixing, and the addition of heterogeneous SO₂ uptake described above, results in ~40% less nitrate and ALWC, and 30% less sulfate, than in the simulation with heterogeneous SO₂ uptake alone. As a result, sulfate concentrations are 30% less than in the simulation with heterogeneous SO₂ uptake alone. Therefore, studies attempting to determine ~~estimating~~ γ to improve sulfate simulations of haze must also consider the impact of model nitrate biases on their parameterization. Follow-up work will include consideration of improvements to the model sulfate and nitrate simulation with a coupled model system such as WRF-GC (Lin et al., 2020) that is able to better simulate the urban scale as well as long-range transport.

7 Conclusions

We used aircraft and surface observations from the NIER-NASA KORUS-AQ field campaign in May and June 2016 to evaluate GEOS-Chem simulations of PM_{2.5} composition in the Seoul Metropolitan Area, including during a haze pollution event characterized by high levels of secondary inorganic aerosol. Models generally underestimate sulfate during haze and generally overestimate ~~overestimate~~ nitric acid and the gas-particle partitioning of nitric acid to aerosol and underestimate sulfate during haze events across in East Asia (An et al., 2019). This is of concern for using models to determine the fraction of PM_{2.5} pollution that can be controlled using local policy measures in South Korea, and the level to which exceedances of PM_{2.5} standards are caused by long-range transport.

The model underestimated PM_{2.5} in Seoul during the campaign (NMB = -15%) with larger errors in composition. On average, the model underestimated sulfate (-64%) and SOA (-43%) but overestimated nitrate (+36%). Models typically underestimate secondary organic aerosol (SOA, Zhao et al., 2016), and this could be due to missing sources from anthropogenic precursors (Nault et al., ~~2020~~2021a). This SOA bias will be investigated in future studies. Aircraft observations, only available during daytime hours, showed model underestimates in sulfate comparable to the bias at the surface. However, modeled nitrate was underestimated aloft, contradicting the model overestimate in the campaign average (which includes nighttime observations). Hourly surface observations showed that this was due to a model overestimate at night. During the campaign, nitrate formation was limited by the supply of nitric acid, which was overestimated against daytime aircraft observations by +100% and contributed to the model nighttime bias. Recent developments to the model wet deposition scheme have significantly improved the simulation of nitrate and nitric acid, but further improvements are unlikely to resolve the model bias.

The model overestimate in nitric acid was not due to overestimated production, insufficient loss to wet deposition, or uptake to dust or seasalt. Increasing the loss of nitric acid, implemented here as an increase in the nitric acid dry deposition velocity by a factor of five, was required to reconcile the model with observations. Aircraft observations of total nitrate ($TNO_3 = HNO_3$

and pNO₃) as a function of photochemical age support this increase. The model underestimate in deposition could be explained by missing treatment of turbulence driven by the urban heat island effect and the heterogeneity of the urban landscape, which would also increase the surface area available for deposition. Here, we only consider the effect on HNO₃, but these factors would also impact other species that readily deposit to surfaces such as NH₃, which was not measured during the campaign.

Observations of ozone, NO₂, and ClNO₂ showed that N₂O₅ hydrolysis should be the main driver of nighttime nitrate production while the model primarily produced nitrate through aerosol uptake of NO₂. The model overly titrated ozone, with an average nighttime concentration of 13 ppb compared to 24–23 ppb in the observations. This resulted in excess model NO₂ and prevented the production of N₂O₅. Observations of ozone and of the nighttime mixed layer height implied insufficient nighttime mixing and an overly rapid collapse of the afternoon mixed layer in the model. We attributed these errors to the premature shutdown of afternoon eddies and missing treatment of the urban heat island effect that typically generates a positive nighttime heat flux that is not present in the model. Nighttime measurements of the vertical structure of key species such as ozone, NO₂, N₂O₅, and HNO₃, ideally complemented by surface heat flux observations, are needed to further constrain model nighttime nitrate production, and determine the extent to which the model underestimates nighttime heating and mixing depth.

The model errors in simulating nitrate and nitric acid, mainly arising from overestimated daytime nitric acid and excess nighttime ozone titration, are exacerbated in the simulation of haze pollution. Overestimated nitric acid results in larger values of daytime nitrate during the haze buildup. This could be due to the model underestimate in sulfate as overestimated model pH would allow for increased partitioning of nitric acid to the particle phase. Nighttime nitrate in the model is underestimated during the haze buildup likely due to missing rapid N₂O₅ hydrolysis. Sensitivity simulations showed that raising the nighttime mixed layer and providing a positive nighttime sensible heat flux of +10 W m⁻² improved the model simulation of nitrate, ozone, and the allowed for nighttime production of nitrate via N₂O₅ pathway for nitrate production during haze. Previous studies have simply raised the nighttime mixed layer and found little effect on simulated pollution (Oak et al., 2019; Miao et al., 2020) but this may be due to missing nocturnal heating from anthropogenic heat release.

The underestimate in model sulfate during the KORUS-AQ haze event is typical of models that often do not include heterogeneous aerosol uptake of SO₂ (Wang et al., 2014; Zheng et al., 2015a, 2015b; Shao et al., 2019). Observations of the sulfate oxidation ratio (SOR) as a function of RH supported the need for this pathway as the strong increase in SOR with RH was not present in the model. A simple parameterization of this process increased model sulfate levels from 4 to 15 μg m⁻³ during the haze, in better agreement with observations. However, the success of this parameterization was complicated by model nitrate biases. A simulation of the haze with both improved model nitrate and heterogeneous uptake of SO₂ resulted in a 30% reduction in model sulfate over the simulation with heterogeneous uptake of SO₂, illustrating the need to consider model biases in sulfate and nitrate simultaneously. GEOS-Chem parameterizations of the urban environment are lacking and cannot be currently adjusted to robustly simulate nitrate during the campaign. However, future studies attempting to simulate sulfate in haze should consider the impact of model nitrate biases on their parameterizations. These studies require models that are

55 able to simulate a large domain to calculate long-range transport but include the detailed parameterizations of the urban environment (urban heat island effect etc.) required to successfully simulate nitrate.

60 Determining the contribution of local vs. transported PM_{2.5} is essential to the development of successful policy measures to reduce unhealthy pollution levels. Significant effort has gone into this evaluation in South Korea, but with models that have errors in PM_{2.5} composition (Choi et al., 2019; Kumar et al., 2021). The local PM_{2.5} contribution may be underestimated without including heterogeneous uptake of SO₂ on aerosol to produce sulfate during haze. Locally-produced PM_{2.5} increased from 13 to 24 µg m⁻³, decreasing the fraction of foreign pollution from 66% to 54%. Locally-produced sulfate increased from <1 µg m⁻³ to 6 µg m⁻³, implying that controls on SO₂ could have a larger impact than in model formulations without this chemistry. As a consequence of the 2013 Clean Air Action plan implemented in China, emissions of inorganic aerosol precursors have been decreasing (Zheng et al., 2018) ~~and resulting in declines in concentrations of PM_{2.5} in China have declined by approximately 5 µg m⁻³ per year from 2013–2018~~ (Zhai et al., 2019). Emission reductions in South Korea may be less rapid (Bae et al., 2021), and thus the impact of long-range transport on future PM_{2.5} pollution events could decline in the future. It is critical for models to improve representations of the interactions between physical processes and chemical production of PM_{2.5} production to support continued local air quality improvements. Follow-up studies to this work will evaluate model performance during other seasons (i.e., winter) using a model system with online meteorology to determine whether factors driving model errors in this work are occurring throughout the year.

65
70

Code Availability

The model code used in this work is available at 10.5281/zenodo.5620667.

Data Availability

75 The KORUS-AQ data archive (KORUS-AQ Science Team, 2019) includes both the aircraft and ground-based measurements from AirKorea, Olympic Park, and KIST. The precipitation data is available at: <https://www.ncdc.noaa.gov/cdo-web/datasets>. Cloud observations (RKSS ASOS station) are available here: (<http://mesonet.agron.iastate.edu/request/download.phtml>).

Author Contribution

80 The original draft preparation was completed by KRT, with review and editing by JHC, BAN, CEJ, HK, and GC. JHC, CEJ, GC, BAN, HK, and KRT contributed to project conceptualization. Modeling work was done by KRT, with additional support from SZ, XW, EM, GL and FY and formal analysis was completed by KRT and BAN. The observational data for this project was provided by BAN, HK, JLJ, PCJ, JED, MJK, SK, IJS, DRB, and LC. JHW and YK- provided the KORUS-AQ emissions.

85

Competing Interests

The authors have the following competing interests: Some authors are members of the editorial board of Atmospheric Chemistry and Physics. The peer-review process was guided by an independent editor, and the authors have also no other competing interests to declare.

90

Acknowledgements

We acknowledge Gangwoong Lee for his leadership in managing the campaign efforts at Olympic Park. We acknowledge Andrew Weinheimer for the use of his NO and NO₂ data from KORUS-AQ. We acknowledge Ron Cohen for the use of the TD-LIF data. We acknowledge Glenn Diskin for the use of DACOM CO and DLH RH data. We acknowledge Bill Brune for the use of his ATHOS OH data. We acknowledge Paul Wennberg and John Crounse for the use of their CIT-CIMS HNO₃ data. We acknowledge L. Greg Huey for the use of his SO₂ data. We acknowledge James J. Szykman for the use of ceilometer data at Olympic Park. We acknowledge Seogjo Cho for the MARGA data at Olympic Park. We acknowledge Ke Li and Yingying Yan for their help implementing aromatic chemistry in GEOS-Chem. We thank Jerome Fast, Rahul Zaveri, and David Peterson for their helpful discussions. [KRT and BAN were supported by NASA Grant 80NSSC22K0283](#). PCJ and JJJ were supported by NASA Grants 80NSSC18K0630 and 80NSSC19K0124. The GEOS-FP data used in this study/project have been provided by the Global Modeling and Assimilation Office (GMAO) at NASA Goddard Space Flight Center.

00

References

- Alexander, B., Park, R. J., Jacob, D. J., Li, Q. B., Yantosca, R. M., Savarino, J., Lee, C. C. W. and Thiemens, M. H.: Sulfate formation in sea-salt aerosols: Constraints from oxygen isotopes, *J. Geophys. Res. D: Atmos.*, 110(D10), doi:10.1029/2004JD005659, 2005.
- Alexander, B., Park, R. J., Jacob, D. J. and Gong, S.: Transition metal-catalyzed oxidation of atmospheric sulfur: Global implications for the sulfur budget, *J. Geophys. Res.*, 114(D2), D02309, 2009.
- Amos, H. M., Jacob, D. J., Holmes, C. D., Fisher, J. A., Wang, Q., Yantosca, R. M., Corbitt, E. S., Galarneau, E., Rutter, A. P., Gustin, M. S., Steffen, A., Schauer, J. J., Graydon, J. A., Louis, V. L. S., Talbot, R. W., Edgerton, E. S., Zhang, Y. and Sunderland, E. M.: Gas-particle partitioning of atmospheric Hg(II) and its effect on global mercury deposition, *Atmos. Chem. Phys.*, 12(1), 591–603, 2012.
- An, Z., Huang, R.-J., Zhang, R., Tie, X., Li, G., Cao, J., Zhou, W., Shi, Z., Han, Y., Gu, Z. and Ji, Y.: Severe haze in northern China: A synergy of anthropogenic emissions and atmospheric processes, *Proc. Natl. Acad. Sci. U. S. A.*, 116(18), 8657–8666, 2019.
- Bae, M., Kim, B.-U., Kim, H. C., Kim, J. and Kim, S.: Role of emissions and meteorology in the recent PM_{2.5} changes in China and South Korea from 2015 to 2018, *Environ. Pollut.*, 270, 116233, 2021.
- Baergen, A. M. and Donaldson, D. J.: Photochemical Renoxification of Nitric Acid on Real Urban Grime, *Environ. Sci. Technol.*, 47(2), 815–820, 2013.
- Baergen, A. M. and Donaldson, D. J.: Formation of reactive nitrogen oxides from urban grime photochemistry, *Atmos. Chem. Phys.*, 16(10), 6355–6363, 2016.
- Barlow, J. F.: Progress in observing and modelling the urban boundary layer, *Urban Climate*, 10, 216–240, 2014.
- Bain, H., Chin, M., Hauglustaine, D. A., Schulz, M., Myhre, G., Bauer, S. E., Lund, M. T., Karydis, V. A., Kucsera, T. L., Pan, X., Pozzer, A., Skeie, R. B., Steenrod, S. D., Sudo, K., Tsigaridis, K., Tsimpidi, A. P. and Tsyro, S. G.: Investigation of global particulate nitrate from the AeroCom phase III experiment, *Atmos. Chem. Phys.*, 17(21), 12911–12940, 2017.
- Blay-Carreras, E., Pardyjak, E. R., Pino, D., Alexander, D. C., Lohou, F. and Lathon, M.: Countergradient heat flux observations during the evening transition period, *Atmos. Chem. Phys.*, 14(17), 9077–9085, 2014.

25

- 30 Bouwman, A. F., Lee, D. S., Asman, W. A. H., Dentener, F. J., Van Der Hoek, K. W. and Olivier, J. G. J.: A global high-resolution emission inventory for ammonia, *Global Biogeochem. Cycles*, 11(4), 561–587, 1997.
- Carn, S. A., Yang, K., Prata, A. J. and Krotkov, N. A.: Extending the long-term record of volcanic SO₂ emissions with the Ozone Mapping and Profiler Suite nadir mapper: OMPS volcanic SO₂ measurements, *Geophys. Res. Lett.*, 42(3), 925–932, 2015.
- 35 Chang, Y., Zou, Z., Zhang, Y., Deng, C., Hu, J., Shi, Z., Dore, A. J. and Collett, J. L., Jr: Assessing Contributions of Agricultural and Nonagricultural Emissions to Atmospheric Ammonia in a Chinese Megacity, *Environ. Sci. Technol.*, 53(4), 1822–1833, 2019.
- Chan Miller, C., Jacob, D. J., Marais, E. A., Yu, K., Travis, K. R., Kim, P. S., Fisher, J. A., Zhu, L., Wolfe, G. M., Hanisco, T. F., Keutsch, F. N., Kaiser, J., Min, K.-E., Brown, S. S., Washenfelder, R. A., González Abad, G. and Chance, K.: Glyoxal yield from isoprene oxidation and relation to formaldehyde: chemical mechanism, constraints from
40 SENEX aircraft observations, and interpretation of OMI satellite data, *Atmos. Chem. Phys.*, 17(14), 8725–8738, 2017.
- Chan, Y., Evans, M. J., He, P., Holmes, C. D., Jaeglé, L., Kasibhatla, P., Liu, X., Sherwen, T., Thornton, J. A., Wang, X., Xie, Z., Zhai, S. and Alexander, B.: Heterogeneous nitrate production mechanisms in intense haze events in the North China Plain, *J. Geophys. Res.*, doi:10.1029/2021jd034688, 2021.
- 45 Chen, D., Wang, Y., McElroy, M. B., He, K., Yantosca, R. M. and Le Sager, P.: Regional CO pollution and export in China simulated by the high-resolution nested-grid GEOS-Chem model, *Atmos. Chem. Phys.*, 9(11), 3825–3839, 2009.
- Chen, D., Liu, Z., Fast, J. and Ban, J.: Simulations of sulfate–nitrate–ammonium (SNA) aerosols during the extreme haze events over northern China in October 2014, *Atmos. Chem. Phys.*, 16(16), 10707–10724, 2016.
- Chen, L., Gao, Y., Zhang, M., Fu, J. S., Zhu, J., Liao, H., Li, J., Huang, K., Ge, B., Wang, X., Lam, Y. F., Lin, C.-Y.,
50 Itahashi, S., Nagashima, T., Kajino, M., Yamaji, K., Wang, Z. and Kurokawa, J.-I.: MICS-Asia III: multi-model comparison and evaluation of aerosol over East Asia, *Atmos. Chem. Phys.*, 19(18), 11911–11937, 2019.
- Chen, Q., Schmidt, J. A., Shah, V., Jaeglé, L., Sherwen, T. and Alexander, B.: Sulfate production by reactive bromine: Implications for the global sulfur and reactive bromine budgets: Sulfur-Halogen Interactions, *Geophys. Res. Lett.*, 44(13), 7069–7078, 2017.
- 55 ~~Cherin, N., Roustan, Y., Musson-Genon, L. and Seigneur, C.: Modelling atmospheric dry deposition in urban areas using an urban canopy approach, *Geosci. Model Dev.*, 8(3), 893–910, 2015.~~
- Choi, J., Park, R. J., Lee, H.-M., Lee, S., Jo, D. S., Jeong, J. I., Henze, D. K., Woo, J.-H., Ban, S.-J., Kee, M.-D., Lim, C.-S., Park, M.-K., Shin, H. J., Cho, S., Peterson, D. and Song, C.-K.: Impacts of local vs. trans-boundary emissions from different sectors on PM_{2.5} exposure in South Korea during the KORUS-AQ campaign, *Atmos. Environ.*, 203, 196–
60 205, 2019.
- Clegg, S. L. and Brimblecombe, P.: Equilibrium partial pressures and mean activity and osmotic coefficients of 0–100% nitric acid as a function of temperature, *J. Phys. Chem.*, 94(13), 5369–5380, 1990.
- Clegg, S. L., Brimblecombe, P. and Wexler, A. S.: Thermodynamic Model of the System H⁺–NH₄⁺–SO₄²⁻–NO₃⁻–H₂O at Tropospheric Temperatures, *J. Phys. Chem. A*, 102(12), 2137–2154, 1998.
- 65 Crawford, J. H., Ahn, J.-Y., Al-Saadi, J. A., Chang, L., Emmons, L. K., Kim, J., Lee, G., Park, J.-H., Park, R. J., Woo, J. H., Song, C. K., Hong, J.-H., Hong, Y.-D., Lefer, B. L., Lee, M., Lee, T., Kim, S., Min, K.-E., Yum, S. S., Shin, H. J., Kim, Y.-W., Choi, J.-S., Park, J.-S., Szykman, J. J., Long, R. W., Jordan, C. E., Simpson, I. J., Fried, A., Dibb, J. E., Cho, S. Y. and Kim, Y. P.: The Korea–United States Air Quality (KORUS-AQ) Field Study, *Elem. Sci. Anth.*, 9(1), doi:10.1525/elementa.2020.00163, 2021.
- 70 Day, D. A., Wooldridge, P. J., Dillon, M. B., Thornton, J. A., and Cohen, R. C.: A thermal dissociation laser-induced fluorescence instrument for in situ detection of NO₂, peroxy nitrates, alkyl nitrates, and HNO₃: DETECTION OF NO₂, SPNs, EANs, AND HNO₃, *J. Geophys. Res.*, 107, ACH 4-1–ACH 4-14, doi:10.1029/2001JD000779, 2002.
- Dibb, J. E., Talbot, R. W., Scheuer, E. M., Seid, G., Avery, M. A., and Singh, H. B.: Aerosol chemical composition in Asian continental outflow during the TRACE-P campaign: Comparison with PEM-West B: AEROSOL COMPOSITION
75 IN TRACE-P ASIAN OUTFLOW, *J. Geophys. Res.*, 108, doi:10.1029/2002JD003111, 2003. Diskin, G. S.; Podolske, J. R.; Sachse, G. W.; and Slate, T. A.: "Open-Path Airborne Tunable Diode Laser Hygrometer," in Diode Lasers and Applications in Atmospheric Sensing, SPIE Proceedings 4817, A. Fried, editor, 196–204 (2002).
- Emerson, E. W., Hodshire, A. L., DeBolt, H. M., Bilsback, K. R., Pierce, J. R., McMeeking, G. R., and Farmer, D. K.:

- Revisiting particle dry deposition and its role in radiative effect estimates, *Proc. Natl. Acad. Sci USA*, 117, 26076–26082, doi:10.1073/pnas.2014761117, 2020.
- Energy Policy Institute: Air Quality Life Index: South Korea Fact Sheet. [online] Available from: https://aqli.epic.uchicago.edu/wp-content/uploads/2019/05/EPIC_SouthKorea_FactSheet_31319.pdf (Accessed 24 February 2020), 2019.
- Fairlie, T. D., Jacob, D. J., Dibb, J. E., Alexander, B., Avery, M. A., van Donkelaar, A. and Zhang, L.: Impact of mineral dust on nitrate, sulfate, and ozone in transpacific Asian pollution plumes, *Atmos. Chem. Phys.*, 10(8), 3999–4012, 2010.
- Fischer, E. V., Jacob, D. J., Yantosca, R. M., Sulprizio, M. P., Millet, D. B., Mao, J., Paulot, F., Singh, H. B., Roiger, A., Ries, L., Talbot, R. W., Dzepina, K. and Pandey Deolal, S.: Atmospheric peroxyacetyl nitrate (PAN): a global budget and source attribution, *Atmos. Chem. Phys.*, 14(5), 2679–2698, 2014.
- Fisher, J. A., Jacob, D. J., Travis, K. R., Kim, P. S., Marais, E. A., Chan Miller, C., Yu, K., Zhu, L., Yantosca, R. M., Sulprizio, M. P., Mao, J., Wennberg, P. O., Crounse, J. D., Teng, A. P., Nguyen, T. B., St. Clair, J. M., Cohen, R. C., Romer, P., Nault, B. A., Wooldridge, P. J., Jimenez, J. L., Campuzano-Jost, P., Day, D. A., Hu, W., Shepson, P. B., Xiong, F., Blake, D. R., Goldstein, A. H., Misztal, P. K., Hanisco, T. F., Wolfe, G. M., Ryerson, T. B., Wisthaler, A. and Mikoviny, T.: Organic nitrate chemistry and its implications for nitrogen budgets in an isoprene- and monoterpene-rich atmosphere: constraints from aircraft (SEAC4RS) and ground-based (SOAS) observations in the Southeast US, *Atmos. Chem. Phys.*, 16(9), 5969–5991, 2016.
- Fisher, J. A., Atlas, E. L., Barletta, B., Meinardi, S., Blake, D. R., Thompson, C. R., Ryerson, T. B., Peischl, J., Tzompa-Sosa, Z. A. and Murray, L. T.: Methyl, ethyl, and propyl nitrates: global distribution and impacts on reactive nitrogen in remote marine environments, *J. Geophys. Res.*, 123, 25, 2018.
- Ge, X., Zhang, Q., Sun, Y., Ruehl, C. R. and Setyan, A.: Effect of aqueous-phase processing on aerosol chemistry and size distributions in Fresno, California, during wintertime, *Environ. Chem.*, 9(3), 221, 2012.
- Ghosh, S., Shon, Z.-H., Kim, K.-H., Song, S.-K., Jung, K., and Kim, N.-J. (2012). Seasonal variation of PM_{2.5} and its major components in urban monitoring site. *Asian J. Atmos. Environ.*, 6(1), 23-32.
- Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K. and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Copernicus* [online] Available from: <http://dspace.mit.edu/handle/1721.1/78869> (Accessed 12 February 2018), 2012.
- Global Burden of Disease (GBD) Collaborative Network: Global Burden of Disease Study 2019 (GBD 2019) Reference Life Table, doi:10.6069/1D4Y-YQ37, 2021.
- Guo, H., Sullivan, A. P., Campuzano-Jost, P., Schroder, J. C., Lopez-Hilfiker, F. D., Dibb, J. E., Jimenez, J. L., Thornton, J. A., Brown, S. S., Nenes, A. and Weber, R. J.: Fine particle pH and the partitioning of nitric acid during winter in the northeastern United States, *J. Geophys. Res. D: Atmos.*, 121(17), 10,355–10,376, 2016.
- Guo, H., Liu, J., Froyd, K. D., Roberts, J. M., Veres, P. R., Hayes, P. L., Jimenez, J. L., Nenes, A. and Weber, R. J.: Fine particle pH and gas–particle phase partitioning of inorganic species in Pasadena, California, during the 2010 CalNex campaign, *Atmos. Chem. Phys.*, 17(9), 5703–5719, 2017.
- Guo, H., Campuzano-Jost, P., Nault, B. A., Day, D. A., Schroder, J. C., Kim, D., Dibb, J. E., Dollner, M., Weinzierl, B., and Jimenez, J. L.: The importance of size ranges in aerosol instrument intercomparisons: a case study for the Atmospheric Tomography Mission, *Atmos. Meas. Tech.*, 14, 3631–3655, doi:10.5194/amt-14-3631-2021, 2021.
- Halios, C. H. and Barlow, J. F.: Observations of the Morning Development of the Urban Boundary Layer Over London, UK, Taken During the ACTUAL Project, *Bound.-Layer Meteorol.*, 166(3), 395–422, 2018.
- Han, X., Cai, J., Zhang, M. and Wang, X.: Numerical simulation of interannual variation in transboundary contributions from Chinese emissions to PM_{2.5} mass burden in South Korea, *Atmos. Environ.*, 118440, 2021.
- Heald, C. L., Collett, J. L., Lee, T., Benedict, K. B., Schwandner, F. M., Li, Y., Clarisse, L., Hurtmans, D. R., Van Damme, M., Clerbaux, C., Coheur, P.-F., Philip, S., Martin, R. V. and Pye, H. O. T.: Atmospheric ammonia and particulate inorganic nitrogen over the United States, *Atmos. Chem. Phys.*, 12(21), 10295–10312, 2012.
- Hegarty, J. D., Lewis, J., McGrath-Spangler, E. L., Henderson, J., Scarino, A. J., DeCola, P., Ferrare, R., Hicks, M., Adams-Selin, R. D. and Welton, E. J.: Analysis of the Planetary Boundary Layer Height during DISCOVER-AQ Baltimore–Washington, D.C., with Lidar and High-Resolution WRF Modeling, *J. Appl. Meteorol. Climatol.*,

57(11), 2679–2696, 2018.

- 30 Heim, E. W., Dibb, J., Scheuer, E., Jost, P. C., Nault, B. A., Jimenez, J. L., Peterson, D., Knote, C., Fenn, M., Hair, J., Beyersdorf, A. J., Corr, C. and Anderson, B. E.: Asian dust observed during KORUS-AQ facilitates the uptake and incorporation of soluble pollutants during transport to South Korea, *Atmos. Environ.*, 117305, 2020.
- Herrmann, H., Schaefer, T., Tilgner, A., Styler, S. A., Weller, C., Teich, M. and Otto, T.: Tropospheric aqueous-phase chemistry: kinetics, mechanisms, and its coupling to a changing gas phase, *Chem. Rev.*, 115(10), 4259–4334, 2015.
- 35 Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M., Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J.-I., Li, M., Liu, L., Lu, Z., Moura, M. C. P., O'Rourke, P. R. and Zhang, Q.: Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS), *Geosci. Model Dev.*, 11(1), 369–408, 2018.
- 40 Holmes, C. D., Bertram, T. H., Confer, K. L., Graham, K. A., Ronan, A. C., Wirks, C. K. and Shah, V.: The Role of Clouds in the Tropospheric NO_x Cycle: A New Modeling Approach for Cloud Chemistry and Its Global Implications, *Geophys. Res. Lett.*, 46(9), 4980–4990, 2019.
- Holtzlag, A. a. M. and Boville, B. A.: Local Versus Nonlocal Boundary-Layer Diffusion in a Global Climate Model, *J. Clim.*, 6(10), 1825–1842, 1993.
- Hong, J.-W. and Hong, J.: Changes in the Seoul Metropolitan Area Urban Heat Environment with Residential Redevelopment, *J. Appl. Meteorol. Climatol.*, 55(5), 1091–1106, 2016.
- 45 Hudman, R. C., Moore, N. E., Mebust, A. K., Martin, R. V., Russell, A. R., Valin, L. C. and Cohen, R. C.: Steps towards a mechanistic model of global soil nitric oxide emissions: implementation and space based-constraints, *Atmos. Chem. Phys.*, 12(16), 7779–7795, 2012.
- 50 Ibikunle, I., Beyersdorf, A., Campuzano-Jost, P., Corr, C., Crounse, J. D., Dibb, J., Diskin, G., Huey, G., Jimenez, J.-L., Kim, M. J., Nault, B. A., Scheuer, E., Teng, A., Wennberg, P. O., Anderson, B., Crawford, J., Weber, R., and Nenes, A.: Fine particle pH and sensitivity to NH₃ and HNO₃ over summertime South Korea during KORUS-AQ, *Atmos. Chem. Phys. Disc.*, doi:10.5194/acp-2020-501, 2020.
- Itahashi, S., Uno, I., Osada, K., Kamiguchi, Y., Yamamoto, S., Tamura, K., Wang, Z., Kurosaki, Y. and Kanaya, Y.: Nitrate transboundary heavy pollution over East Asia in winter, *Atmos. Chem. Phys.*, 17(6), 3823–3843, 2017.
- 55 Jacob, D.: Heterogeneous chemistry and tropospheric ozone, *Atmos. Environ.*, 34(12-14), 2131–2159, 2000.
- Jaeglé, L., Quinn, P. K., Bates, T. S., Alexander, B. and Lin, J.-T.: Global distribution of sea salt aerosols: new constraints from in situ and remote sensing observations, *Atmos. Chem. Phys.*, 11(7), 3137–3157, 2011.
- Jeong, D., Seco, R., Gu, D., Lee, Y., Nault, B. A., Knote, C. J., Mcgee, T., Sullivan, J. T., Jimenez, J. L., Campuzano-Jost, P., Blake, D. R., Sanchez, D., Guenther, A. B., Tanner, D., Huey, L. G., Long, R., Anderson, B. E., Hall, S. R., Ullmann, K., Shin, H., Herndon, S. C., Lee, Y., Kim, D., Ahn, J., and Kim, S.: Integration of airborne and ground observations of nitryl chloride in the Seoul metropolitan area and the implications on regional oxidation capacity during KORUS-AQ 2016, *Atmos. Chem. Phys.*, 19, 12779–12795, <https://doi.org/10.5194/acp-19-12779-2019>, 2019.
- 60 Jeong, D., Seco, R., Gu, D., Lee, Y., Nault, B. A., Knote, C. J., Mcgee, T., Sullivan, J. T., Jimenez, J. L., Campuzano-Jost, P., Blake, D. R., Sanchez, D., Guenther, A. B., Tanner, D., Huey, L. G., Long, R., Anderson, B. E., Hall, S. R., Ullmann, K., Shin, H. J., Herndon, S. C., Lee, Y., Kim, D., Ahn, J. and Kim, S.: Integration of airborne and ground observations of nitryl chloride in the Seoul metropolitan area and the implications on regional oxidation capacity during KORUS-AQ 2016, *Atmos. Chem. Phys.*, 19(19), 12779–12795, 2019.
- 65 Jordan, C., Crawford, J. H., Beyersdorf, A. J., Eck, T. F., Halliday, H. S., Nault, B. A., Chang, L.-S., Park, J., Park, R., Lee, G., Kim, H., Ahn, J.-Y., Cho, S., Shin, H. J., Lee, J. H., Jung, J., Kim, D.-S., Lee, M., Lee, T., Whitehill, A., Szykman, J., Schueneman, M. K., Campuzano-Jost, P., Jimenez, J. L., DiGangi, J. P., Diskin, G. S., Anderson, B. E., Moore, R. H., Ziemba, L. D., Fenn, M. A., Hair, J. W., Kuehn, R. E., Holz, R. E., Chen, G., Travis, K., Shook, M., Peterson, D. A., Lamb, K. D. and Schwarz, J. P.: Investigation of factors controlling PM_{2.5} variability across the South Korean Peninsula during KORUS-AQ, *Elem Sci Anth*, 8(1), 28, 2020.
- 70 Jung, J., Lee, J., Kim, B. and Oh, S.: Seasonal variations in the NO₂ artifact from chemiluminescence measurements with a molybdenum converter at a suburban site in Korea (downwind of the Asian continental outflow) during 2015–2016, *Atmos. Environ.*, 165, 290–300, 2017.
- 75 Jung, J., Ghim, Y. S., Lyu, Y. S., Lim, Y.-J., Park, J. and Sung, M.-Y.: Quantification of regional contributions to fine

- particles at downwind areas under Asian continental outflows during winter 2014, *Atmos. Environ.*, 210, 231–240, 2019.
- 80 Kasibhatla, P., Chameides, W. L., and John, J. St.: A three-dimensional global model investigation of seasonal variations in the atmospheric burden of anthropogenic sulfate aerosols, *J. Geophys. Res.*, 102, 3737–3759, <https://doi.org/10.1029/96JD03084>, 1997.
- 85 Keuken, M. P., Bakker, F. P., Möls, J. J., Broersen, B. and Slanina, J.: Atmospheric Deposition and Conversion of Ammonium to Nitric Acid on a Historic Building: A Pilot Study, *Int. J. Environ. Anal. Chem.*, 38(1), 47–62, 1990.
- Kim, J.-H., Kim, H.-J. and Yoo, S.-H.: Public Value of Enforcing the PM_{2.5} Concentration Reduction Policy in South Korean Urban Areas, *Sustain. Sci. Pract. Policy*, 10(4), 1144, 2018.
- Kim, H., Zhang, Q., and Heo, J.: Influence of intense secondary aerosol formation and long-range transport on aerosol chemistry and properties in the Seoul Metropolitan Area during spring time: results from KORUS-AQ, *Atmos. Chem. Phys.*, 18, 7149–7168, doi: 10.5194/acp-18-7149-2018, 2018.
- 90 Kim, S., Huey, L. G., Stickel, R. E., Tanner, D. J., Crawford, J. H., Olson, J. R., Chen, G., Brune, W. H., Ren, X., Leshner, R., Wooldridge, P. J., Bertram, T. H., Perring, A., Cohen, R. C., Lefer, B. L., Shetter, R. E., Avery, M., Diskin, G., and Sokolik, I.: Measurement of HO₂ NO₂ in the free troposphere during the Intercontinental Chemical Transport Experiment–North America 2004, *J. Geophys. Res.*, 112, D12S01, doi:10.1029/2006JD007676, 2007.
- 95 Kim, Y., Park, O., Park, S. H., Kim, M., Kim, J.-J., Choi, J.-Y., Lee, D., Cho, S., and Shim, S.: PM_{2.5} pH estimation in Seoul during the KORUS-AQ campaign using different thermodynamic models, *Atmospheric Environment*, 268, 118787, doi:10.1016/j.atmosenv.2021.118787, 2022.
- Korea Meteorological Administration (KMA): KMA Weather Data Service Open MET Data Portal, [online] Available from: <https://data.kma.go.kr/resources/html/en/aowdp.html> (Accessed 29 March 2021), 2021.
- 00 KORUS-AQ Science Team: KORUS-AQ Data, NASA Langley Research Center, doi:10.5067/Suborbital/KORUSAQ/DATA01, 2019.
- Kumar, N., Park, R. J., Jeong, J. I., Woo, J.-H., Kim, Y., Johnson, J., Yarwood, G., Kang, S., Chun, S. and Knipping, E.: Contributions of International Sources to PM_{2.5} in South Korea, *Atmos. Environ.*, 118542, 2021.
- Le, T.-C., Shukla, K. K., Chen, Y.-T., Chang, S.-C., Lin, T.-Y., Li, Z., Pui, D. Y. H., and Tsai, C.-J.: On the concentration differences between PM_{2.5} FEM monitors and FRM samplers, *Atmos. Env.*, 222, 117138, doi:10.1016/j.atmosenv.2019.117138, 2020.
- 05 Lee, H.-M., Park, R. J., Henze, D. K., Lee, S., Shim, C., Shin, H.-J., Moon, K.-J. and Woo, J.-H.: PM_{2.5} source attribution for Seoul in May from 2009 to 2013 using GEOS-Chem and its adjoint model, *Environ. Pollut.*, 221, 377–384, 2017.
- 10 Li, M., Zhang, Z., Wang, T., Xie, M., Li, S., Zhuang, B. and Han, Y.: Nonlinear responses of particulate nitrate to NO_x emission controls in the megalopolises of China, , doi:10.5194/acp-2021-330, 2021.
- Lim, S., Hwang, J., Lee, M., Czimczik, C. I., Xu, X., and Savarino, J.: Robust Evidence of ¹⁴C, ¹³C, and ¹⁵N Analyses Indicating Fossil Fuel Sources for Total Carbon and Ammonium in Fine Aerosols in Seoul Megacity, *Environ. Sci. Technol.*, *acs.est.1c03903*, <https://doi.org/10.1021/acs.est.1c03903>, 2022.
- 15 Lin, H., Feng, X., Fu, T.-M., Tian, H., Ma, Y., Zhang, L., Jacob, D. J., Yantosca, R. M., Sulprizio, M. P., Lundgren, E. W., Zhuang, J., Zhang, Q., Lu, X., Zhang, L., Shen, L., Guo, J., Eastham, S. D., and Keller, C. A.: WRF-GC (v1.0): online coupling of WRF (v3.9.1.1) and GEOS-Chem (v12.2.1) for regional atmospheric chemistry modeling – Part 1: Description of the one-way model, *Geosci. Model Dev.*, 13, 3241–3265, <https://doi.org/10.5194/gmd-13-3241-2020>, 2020.
- 20 Lin, J.-T. and McElroy, M. B.: Impacts of boundary layer mixing on pollutant vertical profiles in the lower troposphere: Implications to satellite remote sensing, *Atmos. Environ.*, 44(14), 1726–1739, 2010.
- Link, M. F., Kim, J., Park, G., Lee, T., Park, T., Babar, Z. B., Sung, K., Kim, P., Kang, S., Kim, J. S., Choi, Y., Son, J., Lim, H.-J. and Farmer, D. K.: Elevated production of NH₄NO₃ from the photochemical processing of vehicle exhaust: Implications for air quality in the Seoul Metropolitan Region, *Atmos. Environ.*, 156, 95–101, 2017.
- 25 Liu, H., Jacob, D. J., Bey, I. and Yantosca, R. M.: Constraints from ²¹⁰Pb and ⁷Be on wet deposition and transport in a global three-dimensional chemical tracer model driven by assimilated meteorological fields, *J. Geophys. Res.*, 106(11), 12109–12128, 2001.
- Li, X. and Rappenglueck, B.: A study of model nighttime ozone bias in air quality modeling, *Atmos. Environ.*, 19, 2018.

- Lothon, M., Lohou, F., Pino, D., Couvreur, F., Pardyjak, E. R., Reuder, J., Vilà-Guerau de Arellano, J., Durand, P.,
30 Hartogensis, O., Legain, D., Augustin, P., Gioli, B., Lenschow, D. H., Faloona, I., Yagüe, C., Alexander, D. C.,
Angevine, W. M., Bargain, E., Barrié, J., Bazile, E., Bezombes, Y., Blay-Carreras, E., Boer, A. van de, Boichard, J.
L., Bourdon, A., Butet, A., Campistron, B., Coster, O. de, Cuxart, J., Dabas, A., Darbieu, C., Deboudt, K., Delbarre,
H., Derrien, S., Flament, P., Fourmentin, M., Garai, A., Gibert, F., Graf, A., Groebner, J., Guichard, F., Jiménez, M.
A., Jonassen, M., van den Kroonenberg, A., Magliulo, V., Martin, S., Martinez, D., Mastroiillo, L., Moene, A. F.,
35 Molinos, F., Moulin, E., Pietersen, H. P., Piguët, B., Pique, E., Román-Cascón, C., Rufin-Soler, C., Saïd, F., Sastre-
Marugán, M., Seity, Y., Steeneveld, G. J., Toscano, P., Traullé, O., Tzanos, D., Wacker, S., Wildmann, N. and
Zaldei, A.: The BLLAST field experiment: Boundary-Layer Late Afternoon and Sunset Turbulence, *Atmos. Chem.*
Phys., 14(20), 10931–10960, 2014.
- Luo, G., Yu, F. and Schwab, J.: Revised treatment of wet scavenging processes dramatically improves GEOS-Chem 12.0.0
40 simulations of surface nitric acid, nitrate, and ammonium over the United States, *Geosci. Model Dev.*, 12(8), 3439–
3447, 2019.
- Mao, J., Fan, S., Jacob, D. J. and Travis, K. R.: Radical loss in the atmosphere from Cu-Fe redox coupling in aerosols,
Atmos. Chem. Phys., 13, 509–519, 2013.
- Massucci, M., Clegg, S. L. and Brimblecombe, P.: Equilibrium Partial Pressures, Thermodynamic Properties of Aqueous and
45 Solid Phases, and Cl₂ Production from Aqueous HCl and HNO₃ and Their Mixtures, *J. Phys. Chem. A*, 103(21),
4209–4226, 1999.
- McDonald, B. C., de Gouw, J. A., Gilman, J. B., Jathar, S. H., Akherati, A., Cappa, C. D., Jimenez, J. L., Lee-Taylor, J.,
Hayes, P. L., McKeen, S. A., Cui, Y. Y., Kim, S.-W., Gentner, D. R., Isaacman-VanWertz, G., Goldstein, A. H.,
Harley, R. A., Frost, G. J., Roberts, J. M., Ryerson, T. B. and Trainer, M.: Volatile chemical products emerging as
50 largest petrochemical source of urban organic emissions, *Science*, 359(6377), 760–764, 2018.
- McDuffie, E. E., Fibiger, D. L., Dubé, W. P., Lopez-Hilfiker, F., Lee, B. H., Thornton, J. A., Shah, V., Jaeglé, L., Guo, H.,
Weber, R. J., Michael Reeves, J., Weinheimer, A. J., Schroder, J. C., Campuzano-Jost, P., Jimenez, J. L., Dibb, J.
E., Veres, P., Ebben, C., Sparks, T. L., Wooldridge, P. J., Cohen, R. C., Hornbrook, R. S., Apel, E. C., Campos, T.,
Hall, S. R., Ullmann, K. and Brown, S. S.: Heterogeneous N₂O₅ Uptake During Winter: Aircraft Measurements
55 During the 2015 WINTER Campaign and Critical Evaluation of Current Parameterizations, *J. Geophys. Res. D:*
Atmos., 123(8), 4345–4372, 2018.
- McNaughton, C. S., Clarke, A. D., Howell, S. G., Pinkerton, M., Anderson, B., Thornhill, L., Hudgins, C., Winstead, E.,
Dibb, J. E., Scheuer, E. and Maring, H.: Results from the DC-8 Inlet Characterization Experiment (DICE): Airborne
Versus Surface Sampling of Mineral Dust and Sea Salt Aerosols, *Aerosol Sci. Technol.*, 41(2), 136–159, 2007.
- 60 Miao, R., Chen, Q., Zheng, Y., Cheng, X., Sun, Y., Palmer, P. I., Shrivastava, M., Guo, J., Zhang, Q., Liu, Y., Tan, Z., Ma,
X., Chen, S., Zeng, L., Lu, K. and Zhang, Y.: Model bias in simulating major chemical components of PM_{2.5} in
China, *Atmos. Chem. Phys.*, 20(20), 12265–12284, 2020.
- Millet, D. B., Baasandorj, M., Farmer, D. K., Thornton, J. A., Baumann, K., Brophy, P., Chaliyakunnel, S., de Gouw, J. A.,
Graus, M., Hu, L., Koss, A., Lee, B. H., Lopez-Hilfiker, F. D., Neuman, J. A., Paulot, F., Peischl, J., Pollack, I. B.,
65 Ryerson, T. B., Warneke, C., Williams, B. J. and Xu, J.: A large and ubiquitous source of atmospheric formic acid,
Atmos. Chem. Phys., 15(11), 6283–6304, 2015.
- Min, J.-S., Park, M.-S., Chae, J.-H. and Kang, M.: Integrated System for Atmospheric Boundary Layer Height Estimation
(ISABLE) using a ceilometer and microwave radiometer, *Atmos. Meas. Tech.*, 13(12), 6965–6987, 2020.
- Murray, L. T., Jacob, D. J., Logan, J. A., Hudman, R. C. and Koshak, W. J.: Optimized regional and interannual variability
70 of lightning in a global chemical transport model constrained by LIS/OTD satellite data: IAV OF LIGHTNING
CONSTRAINED BY LIS/OTD, *J. Geophys. Res. D: Atmos.*, 117(D20), doi:10.1029/2012JD017934, 2012.
- Nault, B. A., Campuzano-Jost, P., Day, D. A., Schroder, J. C., Anderson, B., Beyersdorf, A. J., Blake, D. R., Brune, W. H.,
Choi, Y., Corr, C. A., Gouw, J. A. de, Dibb, J., DiGangi, J. P., Diskin, G. S., Fried, A., Huey, L. G., Kim, M. J.,
Knote, C. J., Lamb, K. D., Lee, T., Park, T., Pusede, S. E., Scheuer, E., Thornhill, K. L., Woo, J.-H. and Jimenez, J.
75 L.: Secondary organic aerosol production from local emissions dominates the organic aerosol budget over Seoul,
South Korea, during KORUS-AQ, *Atmos. Chem. Phys.*, 18(24), 17769–17800, 2018.
- Nault, B. A., Campuzano-Jost, P., Day, D. A., Jo, D. S., Schroder, J. C., Allen, H. M., Bahreini, R., Bian, H., Blake, D. R.,
Chin, M., Clegg, S. L., Colarco, P. R., Crounse, J. D., Cubison, M. J., DeCarlo, P. F., Dibb, J. E., Diskin, G. S.,

- 80 Hodzic, A., Hu, W., Katich, J. M., Kim, M. J., Kodros, J. K., Kupc, A., Lopez-Hilfiker, F. D., Marais, E. A., Middlebrook, A. M., Andrew Neuman, J., Nowak, J. B., Palm, B. B., Paulot, F., Pierce, J. R., Schill, G. P., Scheuer, E., Thornton, J. A., Tsigaridis, K., Wennberg, P. O., Williamson, C. J. and Jimenez, J. L.: Chemical transport models often underestimate inorganic aerosol acidity in remote regions of the atmosphere, *Communications Earth & Environment*, 2(1), 1–13, 2021.
- 85 Nault, B. A., Jo, D. S., McDonald, B. C., Campuzano-Jost, P., Day, D. A., Hu, W., Schroder, J. C., Allan, J., Blake, D. R., Canagaratna, M. R., Coe, H., Coggon, M. M., DeCarlo, P. F., Diskin, G. S., Dunmore, R., Flocke, F., Fried, A., Gilman, J. B., Gkatzelis, G., Hamilton, J. F., Hanisco, T. F., Hayes, P. L., Henze, D. K., Hodzic, A., Hopkins, J., Hu, M., Huey, L. G., Jobson, B. T., Kuster, W. C., Lewis, A., Li, M., Liao, J., Nawaz, M. O., Pollack, I. B., Peischl, J., Rappenglück, B., Reeves, C. E., Richter, D., Roberts, J. M., Ryerson, T. B., Shao, M., Sommers, J. M., Walega, J., Warneke, C., Weibring, P., Wolfe, G. M., Young, D. E., Yuan, B., Zhang, Q., de Gouw, J. A., and Jimenez, J. L.: Anthropogenic Secondary Organic Aerosols Contribute Substantially to Air Pollution Mortality, *Atmos. Chem. Phys.*, doi: [10.5194/acp-21-11201-2021](https://doi.org/10.5194/acp-21-11201-2021)~~[10.5194/acp-2020-914](https://doi.org/10.5194/acp-2020-914)~~, [20202021a](https://doi.org/10.5194/acp-2020-914).
- 90 Nault, B. A., Campuzano-Jost, P., Day, D. A., Jo, D. S., Schroder, J. C., Allen, H. M., Bahreini, R., Bian, H., Blake, D. R., Chin, M., Clegg, S. L., Colarco, P. R., Crounse, J. D., Cubison, M. J., DeCarlo, P. F., Dibb, J. E., Diskin, G. S., Hodzic, A., Hu, W., Katich, J. M., Kim, M. J., Kodros, J. K., Kupc, A., Lopez-Hilfiker, F. D., Marais, E. A., Middlebrook, A. M., Andrew Neuman, J., Nowak, J. B., Palm, B. B., Paulot, F., Pierce, J. R., Schill, G. P., Scheuer, E., Thornton, J. A., Tsigaridis, K., Wennberg, P. O., Williamson, C. J., and Jimenez, J. L.: Chemical transport models often underestimate inorganic aerosol acidity in remote regions of the atmosphere, *Commun Earth Environ*, 2, 93, <https://doi.org/10.1038/s43247-021-00164-0>, 2021b.
- 95 Nenes, A., Pandis, S. N., Kanakidou, M., Russell, A. G., Song, S., Vasilakos, P., and Weber, R. J.: Aerosol acidity and liquid water content regulate the dry deposition of inorganic reactive nitrogen, *Atmos. Chem. Phys.*, 21, 6023–6033, doi:10.5194/acp-21-6023-2021, 2021.
- 00 Neuman, J. A., Parrish, D. D., Ryerson, T. B., Brock, C. A., Wiedinmyer, C., Frost, G. J., Holloway, J. S. and Fehsenfeld, F. C.: Nitric acid loss rates measured in power plant plumes, *J. Geophys. Res.*, 109(D23), doi:10.1029/2004jd005092, 2004.
- 05 Nguyen, T. B., Crounse, J. D., Teng, A. P., St. Clair, J. M., Paulot, F., Wolfe, G. M. and Wennberg, P. O.: Rapid deposition of oxidized biogenic compounds to a temperate forest, *Proc. Natl. Acad. Sci. U. S. A.*, 112(5), E392–E401, 2015.
- 10 Oak, Y. J., Park, R. J., Schroeder, J. R., Crawford, J. H., Blake, D. R., Weinheimer, A. J., Woo, J.-H., Kim, S.-W., Yeo, H., Fried, A., Wisthaler, A. and Brune, W. H.: Evaluation of simulated O₃ production efficiency during the KORUS-AQ campaign: Implications for anthropogenic NO_x emissions in Korea, *Elem Sci Anth*, 7(1), 56, 2019.
- 15 Pai, S. J., Heald, C. L., Pierce, J. R., Farina, S. C., Marais, E. A., Jimenez, J. L., Campuzano-Jost, P., Nault, B. A., Middlebrook, A. M., Coe, H., Shilling, J. E., Bahreini, R., Dingle, J. H. and Vu, K.: An evaluation of global organic aerosol schemes using airborne observations, *Atmos. Chem. Phys.*, 20(5), 2637–2665, 2020.
- Park, R. J.: Natural and transboundary pollution influences on sulfate-nitrate-ammonium aerosols in the United States: Implications for policy, *J. Geophys. Res.*, 109(D15), D15204, 2004.
- 20 Park, R. J., Oak, Y. J., Emmons, L. K., Kim, C.-H., Pfister, G. G., Carmichael, G. R., Saide, P. E., Cho, S.-Y., Kim, S., Woo, J.-H., Crawford, J. H., Gaubert, B., Lee, H.-J., Park, S.-Y., Jo, Y.-J., Gao, M., Tang, B., Stanier, C. O., Shin, S. S., Park, H. Y., Bae, C. and Kim, E.: Multi-model intercomparisons of air quality simulations for the KORUS-AQ campaign, *Elementa: Science of the Anthropocene*, 9(1), doi:10.1525/elementa.2021.00139, 2021.
- Phan, N.-T., Kim, K.-H., Shon, Z.-H., Jeon, E.-C., Jung, K. and Kim, N.-J.: Analysis of ammonia variation in the urban atmosphere, *Atmos. Environ.*, 65, 177–185, 2013.
- 25 Philip, S., Martin, R. V. and Keller, C. A.: Sensitivity of chemistry-transport model simulations to the duration of chemical and transport operators: a case study with GEOS-Chem v10-01, *Geoscientific Model Development*, 9(5), 1683–1695, 2016.
- Pye, H. O. T., Liao, H., Wu, S., Mickley, L. J., Jacob, D. J., Henze, D. K. and Seinfeld, J. H.: Effect of changes in climate and emissions on future sulfate-nitrate-ammonium aerosol levels in the United States, *J. Geophys. Res.*, 114(D01205), 18, 2009.
- Qiu, X., Ying, Q., Wang, S., Duan, L., Zhao, J., Xing, J., Ding, D., Sun, Y., Liu, B., Shi, A., Yan, X., Xu, Q. and Hao, J.: Modeling the impact of heterogeneous reactions of chlorine on summertime nitrate formation in Beijing, China,

- Atmos. Chem. Phys., 19(10), 6737–6747, 2019.
- 30 Sachse, G. W., G. F. Hill, L. O. Wade, and M. G. Perry (1987), Fast-response, high precision carbon monoxide sensor using a tunable diode laser absorption technique, *J. Geophys. Res.*, 92, 2071 – 2081.
- Saide, P. E., Gao, M., Lu, Z., Goldberg, D. L., Streets, D. G., Woo, J.-H., Beyersdorf, A., Corr, C. A., Thornhill, K. L., Anderson, B., Hair, J. W., Nehrir, A. R., Diskin, G. S., Jimenez, J. L., Nault, B. A., Campuzano-Jost, P., Dibb, J., Heim, E., Lamb, K. D., Schwarz, J. P., Perring, A. E., Kim, J., Choi, M., Holben, B., Pfister, G., Hodzic, A., Carmichael, G. R., Emmons, L., and Crawford, J. H.: Understanding and improving model representation of aerosol
35 optical properties for a Chinese haze event measured during KORUS-AQ, *Atmos. Chem. Phys.*, 20, 6455–6478, doi:10.5194/acp-20-6455-2020, 2020.
- Sarwar, G., Simon, H., Bhawe, P. and Yarwood, G.: Examining the impact of heterogeneous nitryl chloride production on air quality across the United States, *Atmos. Chem. Phys.*, 12(14), 6455–6473, 2012.
- 40 Schlosser, J., Stahl, C., Sorooshian, A., Le, Y. T.-H., Jeon, K.-J., Xian, P., Jordan, C. E., Travis, K. R., Crawford, J. H., Gong, S. Y., Shin, H.-J., Song, I.-H., and Youn, J.: Evidence of haze-driven secondary production of supermicrometer aerosol nitrate and sulfate in size distribution data in South Korea, *Aerosols/Field Measurements/Troposphere/Chemistry (chemical composition and reactions)*, <https://doi.org/10.5194/acp-2021-1098>, 2022.
- 45 Shao, J., Chen, Q., Wang, Y., Lu, X., He, P., Sun, Y., Shah, V., Martin, R. V., Philip, S., Song, S., Zhao, Y., Xie, Z., Zhang, L. and Alexander, B.: Heterogeneous sulfate aerosol formation mechanisms during wintertime Chinese haze events: air quality model assessment using observations of sulfate oxygen isotopes in Beijing, *Atmos. Chem. Phys.*, 19(9), 6107–6123, 2019.
- Sherwen, T., Evans, M. J., Carpenter, L. J., Andrews, S. J., Lidster, R. T., Dix, B., Koenig, T. K., Sinreich, R., Ortega, I., Volkamer, R., Saiz-Lopez, A., Prados-Roman, C., Mahajan, A. S. and Ordóñez, C.: Iodine’s impact on tropospheric oxidants: a global model study in GEOS-Chem, *Atmos. Chem. Phys.*, 16(2), 1161–1186, 2016.
- 50 Shimadera, H., Hayami, H., Chatani, S., Morino, Y., Mori, Y., Morikawa, T., Yamaji, K. and Ohara, T.: Sensitivity analyses of factors influencing CMAQ performance for fine particulate nitrate, *J. Air Waste Manage. Assoc.*, 64(4), 374–387, 2014.
- 55 Shin, H. J., Kim, J. C., Lee, S. J. and Kim, Y. P.: Evaluation of the optimum volatile organic compounds control strategy considering the formation of ozone and secondary organic aerosol in Seoul, Korea, *Environ. Sci. Pollut. Res. Int.*, 20(3), 1468–1481, 2013a.
- Shin, H. J., Roh, S. A., Kim, J. C., Lee, S. J. and Kim, Y. P.: Temporal variation of volatile organic compounds and their major emission sources in Seoul, Korea, *Environ. Sci. Pollut. Res. Int.*, 20(12), 8717–8728, 2013b.
- 60 Sievering, H., Kelly, T., McConville, G., Seibold, C. and Turnipseed, A.: Nitric acid dry deposition to conifer forests: Niwot Ridge spruce–fir–pine study, *Atmos. Environ.*, 35(22), 3851–3859, 2001.
- Simpson, I. J., Blake, D. R., Blake, N. J., Meinardi, S., Barletta, B., Hughes, S. C., Fleming, L. T., Crawford, J. H., Diskin, G. S., Emmons, L. K., Fried, A., Guo, H., Peterson, D. A., Wisthaler, A., Woo, J.-H., Barré, J., Gaubert, B., Kim, J., Kim, M. J., Kim, Y., Knote, C., Mikoviny, T., Pusede, S. E., Schroeder, J. R., Wang, Y., Wennberg, P. O. and Zeng, L.: Characterization, sources and reactivity of volatile organic compounds (VOCs) in Seoul and surrounding
65 regions during KORUS-AQ, *Elementa*, 8(1), 37, 2020.
- Song, C. H., Park, M. E., Lee, E. J., Lee, J. H., Lee, B. K., Lee, D. S., Kim, J., Han, J. S., Moon, K. J. and Kondo, Y.: Possible particulate nitrite formation and its atmospheric implications inferred from the observations in Seoul, Korea, *Atmos. Environ.*, 43(13), 2168–2173, 2009.
- 70 Sun, J., Liu, L., Xu, L., Wang, Y., Wu, Z., Hu, M., Shi, Z., Li, Y., Zhang, X., Chen, J. and Li, W.: Key Role of Nitrate in Phase Transitions of Urban Particles: Implications of Important Reactive Surfaces for Secondary Aerosol Formation, *J. Geophys. Res. D: Atmos.*, 123(2), 1234–1243, 2018.
- Sun, K., Tao, L., Miller, D. J., Pan, D., Golston, L. M., Zondlo, M. A., Gri, R. J., Mauzerall, D. L. and Zhu, T.: Vehicle Emissions as an Important Urban Ammonia Source in the United States and China, *Environ. Sci.*, 51(4), 2472–2481, 2017.
- 75 Sun, Y., Wang, Z., Fu, P., Jiang, Q., Yang, T., Li, J., and Ge, X.: The impact of relative humidity on aerosol composition and evolution processes during wintertime in Beijing, China, *Atmospheric Environment*, 77, 927–934, <https://doi.org/10.1016/j.atmosenv.2013.06.019>, 2013.

- 80 Sun, Y., He, Y., Kuang, Y., Xu, W., Song, S., Ma, N., Tao, J., Cheng, P., Wu, C., Su, H., Cheng, Y., Xie, C., Chen, C., Lei, L., Qiu, Y., Fu, P., Croteau, P. and Worsnop, D. R.: Chemical differences between PM₁ and PM_{2.5} in highly polluted environment and implications in air pollution studies, *Geophys. Res. Lett.*, 47(5), doi:10.1029/2019gl086288, 2020.
- 85 Sun, Y., He, Y., Kuang, Y., Xu, W., Song, S., Ma, N., Tao, J., Cheng, P., Wu, C., Su, H., Cheng, Y., Xie, C., Chen, C., Lei, L., Qiu, Y., Fu, P., Croteau, P. and Worsnop, D. R.: Chemical differences between PM₁ and PM_{2.5} in highly polluted environment and implications in air pollution studies, *Geophys. Res. Lett.*, 47(5), doi:10.1029/2019gl086288, 2020.
- Theeuwes, N. E., Barlow, J. F., Teuling, A. J., Grimmond, C. S. B. and Kotthaus, S.: Persistent cloud cover over mega-cities linked to surface heat release, *npj Clim Atmos Sci*, 2(1), 15, 2019.
- 90 Travis, K. R. and Jacob, D. J.: Systematic bias in evaluating chemical transport models with maximum daily 8 h average (MDA8) surface ozone for air quality applications: a case study with GEOS-Chem v9.02, *Geosci. Model Dev.*, 12(8), 3641–3648, 2019.
- 95 Travis, K. R., Jacob, D. J., Fisher, J. A., Kim, P. S., Marais, E. A., Zhu, L., Yu, K., Miller, C. C., Yantosca, R. M., Sulprizio, M. P., Thompson, A. M., Wennberg, P. O., Crounse, J. D., St. Clair, J. M., Cohen, R. C., Laughner, J. L., Dibb, J. E., Hall, S. R., Ullmann, K., Wolfe, G. M., Pollack, I. B., Peischl, J., Neuman, J. A. and Zhou, X.: Why do models overestimate surface ozone in the Southeast United States?, *Atmos. Chem. Phys.*, 16(21), 13561–13577, 2016.
- Varquez, A. C. G., Kiyomoto, S., Khanh, D. N. and Kanda, M.: Global 1-km present and future hourly anthropogenic heat flux, *Sci Data*, 8(1), 64, 2021.
- 00 Wang, X., Jacob, D. J., Eastham, S. D., Sulprizio, M. P., Zhu, L., Chen, Q., Alexander, B., Sherwen, T., Evans, M. J., Lee, B. H., Haskins, J. D., Lopez-Hilfiker, F. D., Thornton, J. A., Huey, G. L. and Liao, H.: The role of chlorine in global tropospheric chemistry, *Atmos. Chem. Phys.*, 19(6), 3981–4003, 2019.
- Wang, Y., Jacob, D. J. and Logan, J. A.: Global simulation of tropospheric O₃-NO_x-hydrocarbon chemistry 1. Model formulation, *J. Geophys. Res.*, 103(D9), 10,713–10,725, 1998.
- Wang, Y., Zhang, Q., Jiang, J., Zhou, W., Wang, B., He, K., Duan, F., Zhang, Q., Philip, S. and Xie, Y.: Enhanced sulfate formation during China's severe winter haze episode in January 2013 missing from current models, *J. Geophys. Res. D: Atmos.*, 119(17), 10,425–10,440, 2014.
- 05 Wang, Y., Chen, Y., Wu, Z., Shang, D., Bian, Y., Du, Z., Schmitt, S. H., Su, R., Gkatzelis, G. I., Schlag, P., Hohaus, T., Voliotis, A., Lu, K., Zeng, L., Zhao, C., Alfarra, M. R., McFiggans, G., Wiedensohler, A., Kiendler-Scharr, A., Zhang, Y., and Hu, M.: Mutual promotion between aerosol particle liquid water and particulate nitrate enhancement leads to severe nitrate-dominated particulate matter pollution and low visibility, *Atmos. Chem. Phys.*, 20, 2161–2175, doi:10.5194/acp-20-2161-2020, 2020.
- 10 Weagle, C. L., Snider, G., Li, C., van Donkelaar, A., Philip, S., Bissonnette, P., Burke, J., Jackson, J., Latimer, R., Stone, E., Abboud, I., Akoshile, C., Anh, N. X., Brook, J. R., Cohen, A., Dong, J., Gibson, M. D., Griffith, D., He, K. B., Holben, B. N., Kahn, R., Keller, C. A., Kim, J. S., Lagrosas, N., Lestari, P., Khian, Y. L., Liu, Y., Marais, E. A., Martins, J. V., Misra, A., Muliane, U., Pratiwi, R., Quel, E. J., Salam, A., Segev, L., Tripathi, S. N., Wang, C., Zhang, Q., Brauer, M., Rudich, Y. and Martin, R. V.: Global Sources of Fine Particulate Matter: Interpretation of PM_{2.5} Chemical Composition Observed by SPARTAN using a Global Chemical Transport Model, *Environ. Sci. Technol.*, doi: acs.est.8b01658, 2018.
- 15 Weinheimer, A. J., Walega, J. G., Ridley, B. A., Sachse, G. W., Anderson, B. E., and Collins, J. E.: Stratospheric NO_y measurements on the NASA DC-8 during AASE II, *Geophys. Res. Lett.*, 20, 2563–2566, doi:10.1029/93GL02627, 1993.
- 20 Weinheimer, A. J., Walega, J. G., Ridley, B. A., Gary, B. L., Blake, D. R., Blake, N. J., Rowland, F. S., Sachse, G. W., Anderson, B. E., and Collins, J. E.: Meridional distributions of NO_x, NO_y, and other species in the lower stratosphere and upper troposphere during AASE II, *Geophys. Res. Lett.*, 21, 2583–2586, doi:10.1029/94GL01897, 1994.
- 25 Werf, G. R. van der, Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van Marle, M. J. E., Morton, D. C., Collatz, G. J., Yokelson, R. J. and Kasibhatla, P. S.: Global fire emissions estimates during 1997–2016, *Earth System Science Data; Katlenburg-Lindau*, 9(2), 697–720, 2017.
- Wesely, M. L.: Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical models, *Atmos. Environ.*, 23(6), 1293–1304, 1989.

- Wexler, A. S. and Clegg, S. L.: Atmospheric aerosol models for systems including the ions H^+ , NH_4^+ , Na^+ , SO_4^{2-} , NO_3^- , Cl^- , Br^- , and H_2O , *J. Geophys. Res.*, 107(D14), 4207, 2002.
- Woo, J.-H., Kim, Y., Kim, H.-K., Choi, K.-C., Eum, J.-H., Lee, J.-B., Lim, J.-H., Kim, J. and Seong, M.: Development of the CREATE Inventory in Support of Integrated Climate and Air Quality Modeling for Asia, *Sustain. Sci. Pract. Policy*, 12(19), 7930, 2020.
- Wooldridge, P. J., Perring, A. E., Bertram, T. H., Flocke, F. M., Roberts, J. M., Singh, H. B., Huey, L. G., Thornton, J. A., Wolfe, G. M., Murphy, J. G., Fry, J. L., Rollins, A. W., LaFranchi, B. W., and Cohen, R. C.: Total Peroxy Nitrates (ΣPNs) in the atmosphere: the Thermal Dissociation-Laser Induced Fluorescence (TD-LIF) technique and comparisons to speciated PAN measurements, *Atmos. Meas. Tech.*, 3, 593–607, doi:10.5194/amt-3-593-2010, 2010.
- Yan, Y., Cabrera-Perez, D., Lin, J., Pozzer, A., Hu, L., Millet, D. B., Porter, W. C. and Lelieveld, J.: Global tropospheric effects of aromatic chemistry with the SAPRC-11 mechanism implemented in GEOS-Chem version 9-02, *Geosci. Model Dev.*, 12(1), 111–130, 2019.
- Yazbeck, T., Bohrer, G., Vines, C., De Roo, F., Mauder, M. and Bakshi, B.: Effects of spatial heterogeneity of leaf density and crown spacing of canopy patches on dry deposition rates, *Agric. For. Meteorol.*, 306, 108440, 2021.
- Zhai, S., Jacob, D. J., Wang, X., Shen, L., Li, K., Zhang, Y., Gui, K., Zhao, T. and Liao, H.: Fine particulate matter ($\text{PM}_{2.5}$) trends in China, 2013–2018: separating contributions from anthropogenic emissions and meteorology, *Atmos. Chem. Phys.*, 19(16), 11031–11041, 2019.
- Zhai, S., Jacob, D. J., Brewer, J. F., Li, K., Moch, J. M., Kim, J., Lee, S., Lim, H., Lee, H. C., Kuk, S. K., Park, R. J., Jeong, J. I., Wang, X., Liu, P., Luo, G., Yu, F., Meng, J., Martin, R. V., Travis, K. R., Hair, J. W., Anderson, B. E., Dibb, J. E., Jimenez, J. L., Campuzano-Jost, P., Nault, B. A., Woo, J.-H., Kim, Y., Zhang, Q. and Liao, H.: Interpretation of geostationary satellite aerosol optical depth (AOD) over East Asia in relation to fine particulate matter ($\text{PM}_{2.5}$): insights from the KORUS-AQ aircraft campaign and seasonality, doi:10.5194/acp-2021-413, 2021.
- Zhai, S., Jacob, D. J., Wang, X., Liu, Z., Wen, T., Shah, V., Li, K., Moch, J. M., Bates, K. H., Song, S., Shen, L., Zhang, Y., Luo, G., Yu, F., Sun, Y., Wang, L., Qi, M., Tao, J., Gui, K., Xu, H., Zhang, Q., Zhao, T., Wang, Y., Lee, H. C., Choi, H., and Liao, H.: Control of particulate nitrate air pollution in China, *Nat. Geosci.*, doi:10.1038/s41561-021-00726-z, 2021.
- Zhao, B., Wang, S., Donahue, N. M., Jathar, S. H., Huang, X., Wu, W., Hao, J., and Robinson, A. L.: Quantifying the effect of organic aerosol aging and intermediate-volatility emissions on regional-scale aerosol pollution in China, *Sci Rep*, 6, 28815, doi:10.1038/srep28815, 2016.
- Zhang, L., Gong, S., Padro, J. and Barrie, L.: A size-segregated particle dry deposition scheme for an atmospheric aerosol module, *Atmos. Environ.*, 35, 549–560, 2001.
- Zhang, L., Wang, T., Zhang, Q., Zheng, J., Xu, Z. and Lv, M.: Potential sources of nitrous acid (HONO) and their impacts on ozone: A WRF-Chem study in a polluted subtropical region: MODELING HONO IN A SUBTROPICAL REGION, *J. Geophys. Res. D: Atmos.*, 121(7), 3645–3662, 2016.
- Zheng, B., Zhang, Q., Zhang, Y., He, K. B., Wang, K., Zheng, G. J., Duan, F. K., Ma, Y. L. and Kimoto, T.: Heterogeneous chemistry: a mechanism missing in current models to explain secondary inorganic aerosol formation during the January 2013 haze episode in North China, *Atmos. Chem. Phys.*, 15(4), 2031–2049, 2015a.
- Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., Qi, J., Yan, L., Zhang, Y., Zhao, H., Zheng, Y., He, K. and Zhang, Q.: Trends in China’s anthropogenic emissions since 2010 as the consequence of clean air actions, *Atmos. Chem. Phys.*, 18(19), 14095–14111, 2018.
- Zheng, G. J., Duan, F. K., Su, H., Ma, Y. L., Cheng, Y., Zheng, B., Zhang, Q., Huang, T., Kimoto, T., Chang, D., Pöschl, U., Cheng, Y. F. and He, K. B.: Exploring the severe winter haze in Beijing: the impact of synoptic weather, regional transport and heterogeneous reactions, *Atmos. Chem. Phys.*, 15(6), 2969–2983, 2015b.

Table 1. Description of the ground site and aircraft observations used in this work¹

Instrument	PI	Species	Reference ²
------------	----	---------	------------------------

Ground Observations			
<i>Korea Institute of Science and Technology (KIST)</i> ³			
Aerodyne High-Resolution Time-of-Flight Aerosol Mass Spectrometer (HR-ToF-AMS)	Hwajin Kim	OA, pNH ₄ , pNO ₃ , pSO ₄	Kim et al., 2018
Multi-angle absorption spectrometer (MAAP)	Hwajin Kim	BC	Kim et al., 2018
<i>Olympic Park</i> ⁴			
Monitor for AeRosols and Gases in ambient Air (MARGA)	Seogju Cho	SO ₂ , SO ₄ ²⁻	N/A
Chemical Ionization Mass Spectrometry (CIMS)	Saewung Kim	ClNO ₂	Slusher et al., 2004
Vaisala CL51	James Szykman	MLH	N/A
2B Tech 211, Teledyne T200U, Teledyne T500U CAPS, Aerodyne QCL	James Szykman and Andrew Whitehill	O ₃ , NO, NO ₂	N/A
Dasibi Model 2108 Oxides of Nitrogen Analyzer	NIER	O ₃ , NO ₂	N/A
BAM-1020 instruments (Met One Instruments, Inc., Grants Pass, OR, USA)	NIER	PM _{2.5}	N/A
DC8 Aircraft			
High-Resolution Time-of-Flight Aerosol Mass Spectrometer (HRTof-AMS) ⁵	Jose Jimenez	pNO ₃ , pSO ₄	Nault et al., 2018 Guo et al., 2021
Soluble Acidic Gases and Aerosol (SAGA)	Jack Dibb	Na ⁺ , Cl ⁻	Dibb et al., 2003
Caltech CIMS (CIT-CIMS)	Paul Wennberg	HNO ₃ , propene hydroxynitrate	St. Clair et al., 2010; Crounse et al., 2006
Airborne Tropospheric Hydrogen Oxides Sensor (ATHOS)	William Brune	OH	Faloona et al., 2004; Brune et al., 2020
NCAR 4-Channel chemiluminescence instrument	Andrew Weinheimer	NO, NO ₂	Weinheimer et al., 1993, 1994
Georgia Tech–Chemical Ionization Mass Spectrometer (GT-CIMS)	L. Greg Huey	SO ₂	Kim et al., 2007
Diode laser spectrometer (Differential Absorption Carbon monOxide Measurement, DACOM)	Glenn Diskin	CO	Sachse et al., 1987
Diode Laser Hygrometer measurements of H ₂ O(v) (DLH)	Glenn Diskin	RH%	Diskin et al., 2002
Thermal Dissociation–Laser-Induced Fluorescence (TD-LIF)	Ron Cohen	ΣANs, ΣPNs	Wooldridge et al., 2010; Day et al., 2002
Whole Air Sampler (WAS)	Donald Blake	propene	Simpson et al., 2020

¹For a full description of all KORUS-AQ observations, see Crawford et al., 2021.

²For specific measurement descriptions including uncertainty information, see the KORUS-AQ data archive (doi: 10.5067/Suborbital/KORUSAQ/DATA01)

³Korea Institute of Standards and Technology (KIST), 37.602°N, 127.126°E

⁴Olympic Park site in Seoul, 37.522°N, 127.124°E

⁵AMS data is written without the charge, see http://cires1.colorado.edu/jimenez-group/wiki/index.php/FAQ_for_AMS_Data_Users#Why_do_you_write_SO4_.26_NO3_and_not_SO42-_.26_NO3-.3F.

Table 2. KORUS-AQ emissions over the domain 70° to 140°E, 15°S to 55°N

May 2016 (Gg)	NO _x	CO	SO ₂	NH ₃
Natural	763 ¹	NA	143 ³	155
Biomass burning	92	7122	53	137
Fossil fuel combustion ²	1921 1920	16163	2133	1705 ⁴
Total	2776 2775	23285	2329	1997

¹Lightning, soil and fertilizer emissions

²Point, area, mobile sources, ships, aircraft from the KORUSv5 inventory

³Volcanic eruptions + degassing

⁴Includes agricultural emissions




Figure 1. a) Model simulation of PM_{2.5} during KORUS-AQ compared against the mean observations at the 15 AirKorea sites in the b) GEOS-Chem model grid-box containing Olympic Park and KIST. The gray shading shows the observed standard deviation. The correlation coefficient (R) and normalized mean bias (NMB) are inset. Map tiles by [Stamen Design](#), under [CC BY 3.0](#). Data by © OpenStreetMap contributors, under [ODbL](#).




Figure 2. Model simulation of $\text{PM}_{2.5}$ compared against observations where the fractional source contributions are calculated from KIST and applied to the mean AirKorea $\text{PM}_{2.5}$ observations from Figure 1 during the four meteorological periods. Figure values are shown in Table 3. The radius of each pie chart is scaled to the maximum value of modeled or observed $\text{PM}_{2.5}$ ($53 \mu\text{g m}^{-3}$). The blue circles show the aerosol liquid water content (ALWC) associated with $\text{PM}_{2.5}$. The sulfate-nitrate-ammonium components are bordered in black to guide the reader.

Table 3. Modeled vs. observed PM_{2.5} composition

Species	Observations (µg m ⁻³)					Model (µg m ⁻³)				
	Dynamic	Stagnant	Transport/ Haze	Blocking	Avg	Dynamic	Stagnant	Transport/ Haze	Blocking	Avg
Sulfate	3.9	3.6	14.7	5.5	6.1	1.7	1.4	4.1	2.1	2.2
Nitrate	2.4	3.4	11.2	3.1	4.5	4.2	4.0	12.9	6.2	6.1
Ammonium	1.9	2.2	8.2	2.7	3.3	1.8	1.7	5.3	2.6	2.6
SOA	6.0	14.2	11.5	8.6	9.5	3.9	4.8	10.0	5.1	5.4
POA	3.3	4.3	4.8	2.8	3.7	2.8	2.8	3.3	3.3	3.0
BC	1.2	1.7	2.2	1.3	1.5	1.0	1.2	2.1	1.5	1.3
PM _{2.5}	18.7	29.4	52.6	24.0	28.6	15.4	15.9	37.7	20.8	20.6
ALWC ¹	12.0	4.1	26.9	6.2	12.6	11.9	17.6	48.7	29.5	22.9
PM _{2.5} + H ₂ O	30.7	33.5	79.5	30.2	41.2	27.3	33.5	86.4	50.3	43.5

¹Aerosol liquid water content (ALWC) is calculated using E-AIM from temperature at KIST, the 50th percentile of RH across the AirKorea sites in Figure 1b, and the speciated PM_{2.5} components from Figure 2.

Table 4. Description of model experiments

Name	Resolution	Simulation Length	Description of changes
Base model	0.25° × 0.3125° over East Asia. Boundary conditions (BCs) from a global 2° × 2.25° simulation ¹ .	1 month initialization + KORUS-AQ period (May 1-June 9).	N/A
No nighttime production	0.25° × 0.3125° over East Asia.	KORUS-AQ period	Remove reactions R2-R5.
Old wet scavenging scheme	0.25° × 0.3125° over East Asia.	KORUS-AQ period	Remove recently implemented wet scavenging scheme (Luo et al., 2019).
5x dry deposition	0.25° × 0.3125° over East Asia.	KORUS-AQ period	Increase the deposition velocity of HNO ₃ by a factor of 5.
No local emissions	0.25° × 0.3125° over East Asia.	KORUS-AQ period	Turn off anthropogenic emissions over South Korea.
Raise nighttime PBL	0.25° × 0.3125° over East Asia.	KORUS-AQ period	Increase the nighttime MLH to 500m.
Increased nighttime mixing	0.25° × 0.3125° over East Asia.	May 23 to May 31	Increase the nighttime MLH to 300m and set nighttime sensible heat flux to 10 W m ⁻² .
Het SO ₂	0.25° × 0.3125° over East Asia.	KORUS-AQ period	Uptake of SO ₂ on aerosol with $\gamma_{RH_{100\%}} = 3 \times 10^{-4}$ and $\gamma_{RH_{50\%}} = 3 \times 10^{-5}$.

¹Boundary conditions from the base simulation are applied to all sensitivity simulations.




Figure 3. Mean vertical profiles of a) observed sulfate, b) observed nitrate, c) model sulfate, and d) model nitrate for the descents ~~into~~^{into}over Olympic Park for each meteorological period. The observations (solid lines) and model (dashed lines) are binned to the nearest 0.5 km below 2 km.




Figure 4. a) Mean hourly modeled vs. observed nitrate derived from $\text{PM}_{2.5}$ observations in the GEOS-Chem gridbox and KIST speciated composition as described in Section 4 for May 1 to June 7, 2016. The gray shading indicates the observed 25th to 75th percentile across the grid box. The model sensitivity studies are described in Section 5. b) Mean model nitric acid diurnal cycle. c) Mean model reactions that produce HNO_3 as described in Section 3.





Figure 5. Modeled and observed ϵ_{NO_3} as a function of RH below 1.5 km for the domain of Fig. 3. Median ϵ_{NO_3} as a function of equally size-binned RH is overlaid (squares). The haze buildup (5/24-5/26) is shown in gray for the observations and dark red for the model.






Figure 6. Mean vertical profiles of a) HNO_3 , b) NO_2 , c) OH , and d) production of HNO_3 (pHNO_3) for the same domain as Fig. 3 but accounting for the availability of OH , NO_2 , and HNO_3 observations. The horizontal bars show the observed and modeled standard deviations. The number of points in each altitude bin are shown in panel a). Model sensitivity simulations that are not significantly different than the base model run are plotted underneath the base model line.




Figure 7. Gridded HNO_3 from the observations a), model b), and the percent difference c) along the flight tracks at the model resolution and below 2 km. Map tiles by [Stamen Design](#), under [CC BY 3.0](#). Data by © OpenStreetMap contributors under [ODbL](#).




Figure 8. Plot of binned observations of NO_x (right axis), total nitrate ($\text{TNO}_3 = \text{HNO}_3 + \text{pNO}_3$), sum of peroxy nitrates (ΣPNs), and sum of alkyl- and multi-functional nitrates (ΣANs) (left axis for TNO_3 , ΣPNs , and ΣANs), normalized to background subtracted CO. The background CO from Nault et al. (2018) of 200 ppbv was used. The photochemical age was calculated using propene and one of its photochemical products, propene hydroxynitrate (Section S3S4). Data are binned between 0 and 5 equivalent hr between 11am to 4pm KST below 1km for the SMA (127 to 127.7°N, 37.2 to 37.7°N). The fit for NO_x (dotted gray curve) is an exponential decay, leading to a first order rate of 0.31 hr^{-1} , which represents the loss of NO_x via the production of oxidized compounds, such as TNO_3 . The best fit for $\text{TNO}_3(t)$ from Eq. 3 (dotted blue curve) includes this production and solves for first order loss, which is assumed to be equivalent to the TNO_3 deposition rate (Section S3S4). Red curves represent solutions for $\text{TNO}_3(t)$ from Eq. 3, assuming different deposition velocities (V_d) discussed in Section 5.2.




Figure 9. Mean diurnal cycle from May 1 to June 7, 2016 for a) ozone and b) NO_2 for the AirKorea sites within the GEOS-Chem gridbox (Fig. 1b) and for c) NO and d) NO_x at Olympic Park. The gray shading represents the standard deviation across the AirKorea sites. The solid gray line is the AirKorea site closest to Olympic Park, and the dashed line is the measurement from the EPA (Table 2) at Olympic Park. The sensitivity study (blue line) is described in Section 5.3.





Figure 10. a) Mean diurnal cycle for the mixed layer height (MLH) from the model and observations from May 1 to June 7, 2016, and b) sensible heat flux (HFLUX) from the model. The MLH is given for the ceilometers (CLH) at Olympic Park (black) and at Seoul National University (gray).







Figure 11. a) Transport/Haze period timeseries of modeled and observed hourly nitrate fraction of $\text{PM}_{2.5}$, b) modeled production of HNO_3 from N_2O_5 (R2) and NO_2 (R4), c) ozone, d) and NO_2 . The sensitivity studies are described in Section 6. The gray shaded regions represent ~~8pm-6pm~~ to ~~8am~~6am.






Figure 12. Sulfate oxidation ratio ($SOR = \frac{SO_4^{2-}}{SO_4^{2-} + SO_2}$) as a function of RH at Olympic Park and from aircraft below 1km for the same domain as Fig. 3 descents over Olympic Park. The squares highlight the data during the Transport/Haze period.




Figure 13. Composition of model PM_{2.5} during the Transport/Haze period. The foreign and local contributions and model sensitivity test including heterogeneous uptake of SO₂ to aerosol (Het SO₂) are calculated as discussed in Section 6. The total PM_{2.5} excluding aerosol liquid water content (ALWC) is given for each simulation.

90