

Eurodelta multi-model simulated and observed PM trends in Europe in the period of 1990-2010

Svetlana Tsyro¹, Wenche Aas², Augustin Colette³, Camilla Andersson⁴, Bertrand Bessagnet^{3,†}, Giancarlo Ciarelli⁵, Florian Couvidat³, Kees Cuvelier⁶, Astrid Manders⁷, Kathleen Mar⁸, Mihaela Mircea⁹, Noelia Otero^{8,‡}, Maria-Teresa Pay¹⁰, Valentin Raffort¹¹, Yelva Roustan¹¹, Mark R. Theobald¹², Marta G. Vivanco¹², Hilde Fagerli¹, Peter Wind^{1,13}, Gino Briganti⁹, Andrea Cappelletti⁹, Massimo D’Isidoro⁹, and Mario Adani⁹

¹Norwegian Meteorological Institute, NO-0313, Oslo, Norway

²Norwegian Institute for Air Research (NILU), Box 100, 2027 Kjeller, Norway

³INERIS, National Institute for Industrial Environment and Risks, Parc Technologique ALATA, 60550, Verneuil-en-Halatte, France

⁴Swedish Meteorological and Hydrological Institute, 60176 Norrköping, Sweden

⁵Institute for Atmospheric and Earth System Research/Physics, Faculty of Science, University of Helsinki, Finland

⁶ex European Commission, Joint Research Centre (JRC), Ispra, Italy

⁷TNO, Dept. Climate, Air and Sustainability, P.O. Box 80015, 3508 TA Utrecht, the Netherlands

⁸Institute for Advanced Sustainability Studies, Postdam, Germany

⁹ENEA, Italian National Agency for New Technologies, Energy and Sustainable Economic Development Via Martiri di Monte Sole 4, 40129 Bologna, Italy

¹⁰BSC, Barcelona Supercomputing Center, Centro Nacional de Supercomputaciòn, Nexus II Building, Jordi Girona, 29, 08034 Barcelona, Spain

¹¹CEREA, École des Ponts, EDF R&D, Île-de-France, France

¹²CIEMAT, Atmospheric Pollution Unit, Avda. Complutense, 22, 28040 Madrid, Spain

¹³Faculty of Science and Technology, University of Tromsø, Tromsø, Norway

[†]Now at European Commission, Joint Research Centre (JRC), Ispra, Italy

[‡]Now at the University of Bern, Switzerland

Correspondence: Svetlana Tsyro (svetlana.tsyro@met.no)

Abstract.

The Eurodelta-Trends multi-model experiment, aimed to assess the efficiency of emission mitigation measures in improving
5 air quality in Europe during 1990-2010, was designed to answer a series of questions regarding European pollution trends.
i.e. were there significant trends detected by observations? ~~do~~ Do the models manage to reproduce observed trends? ~~how~~ How
close is the agreement between the models and how large are the deviations from observations? In this paper, we address these
issues with respect to PM pollution. An in-depth trend analysis has been performed for PM₁₀ and PM_{2.5} for the period
10 of 2000-2010, based on results from six chemical transport models and observational data from the EMEP (Cooperative
Programme for Monitoring and Evaluation of the Long-range Transmission of Air Pollutants in Europe) monitoring network.
Given harmonization of set up and main input data, the differences in model results should mainly result from differences in

the process formulations within the models themselves, and the spread in the models simulated trends could be regarded as an indicator for modelling uncertainty.

The model ensemble simulations indicate overall decreasing trends in PM_{10} and $PM_{2.5}$, ~~with reduction from 2000 to 2010,~~
15 with the total reductions of annual mean concentrations by between 2 and ~~6-5~~ (7 for PM_{10}) $\mu g m^{-3}$ (or between 10 and 30%)
~~from 2000 to 2010~~ across most of Europe (by 0.5-2 $\mu g m^{-3}$ in Fennoscandia, north-west of Russia and Eastern Europe) during
the studied period. Compared to $PM_{2.5}$, relative PM_{10} trends are weaker due to large inter-annual variability of natural coarse
PM within the former. The changes in the concentrations of PM individual components are in general consistent with emission
20 below 30-40% over most of Europe, increasing to 50-60% in northern and eastern parts of EDT domain.

Averaged over measurement sites (26 for PM_{10} and 13 for $PM_{2.5}$), the mean ensemble simulated trends are -0.24 and -0.22
 $\mu g m^{-3} yr^{-1}$ for PM_{10} and $PM_{2.5}$, which are somewhat weaker than the observed trends of -0.35 and -0.40 $\mu g m^{-3} yr^{-1}$,
respectively, partly due to models underestimation of PM concentrations. The correspondence is better in relative PM_{10} and
 $PM_{2.5}$ trends, which are -1.7 and -2.0 % yr^{-1} from the model ensemble and -2.1 and -2.9 % yr^{-1} from the observations,
25 respectively. The observations identify significant trends (at 95% confidence level) for PM_{10} at 56 % of the sites and for
 $PM_{2.5}$ at 36% of the sites, which is somewhat less than the fractions of significant modelled trends. Further, we find somewhat
smaller spatial variability of modelled PM trends with respect to the observed ones across Europe and also within individual
countries.

The strongest decreasing PM trends and the largest number of sites with significant trends is found for the summer season,
30 according to both the model ensemble and observations. The winter PM trends are very weak and mostly insignificant. One
important reason for that is the very modest reductions and even increases in the emissions of primary PM from residential
heating in winter. It should be kept in mind that all findings regarding modeled versus observed PM trends are limited to the
regions where the sites are located.

The analysis reveals a considerable variability of the role of the individual aerosols in PM_{10} trends across European coun-
35 tries. The multi-model simulations, supported by available observations, point to decreases in SO_4^{-2} concentrations playing an
overall dominant role. Also, we see relatively large contributions of the trends of NH_4^+ and NO_3^- to PM_{10} decreasing trends in
Germany, Denmark, Poland and the Po Valley, while the reductions of primary PM emissions appears to be a dominant factor
in bringing down PM_{10} in France, Norway, Portugal, Greece and parts of the UK and Russia.

Further discussions are given with respect to emission uncertainties (including the implications of not accounting for forest
40 fires and natural mineral dust by some of the models) and the effect of inter-annual meteorological variability on the trend
analysis.

1 Introduction

The Convention on Long-range Transboundary Air Pollution (LRTAP), signed in 1979, addresses some of the major envi-
ronmental problems of the UNECE region through scientific collaboration and policy negotiation (UNECE, 2004). Parties

45 develop policies and strategies to combat the release of pollutants in the atmosphere through exchanges of information, consultation, research and monitoring. During the 1980s, 1990s and 2000s, the concentrations of particulate matter (PM) were decreasing due to the decrease of secondary inorganic aerosols (SIA) as a result of the reductions of the emissions of their gaseous precursors in order to address the acidification and eutrophication problems (Fagerli and Aas (2008), Aas et al. (2019))(Fagerli and Aas, 2008; Aas et al., 2019), mainly of SO_x due to the 1st and 2nd Sulphur Protocols, and also NO_x and NH₃ in line with the 1999 Gothenburg Protocol to Abate Acidification, Eutrophication and Ground-level Ozone (UNECE, 2004). The emissions of primary PM were not then regulated, but still were decreasing as a side-effect of the reductions of gaseous pollutants. In the end of 1990s, the issue of adverse effects of particulate pollution on human health came into focus, and in 2012, emissions of primary PM_{2.5} were included in the revised Gothenburg Protocol, stating that fine particulate matter is "the pollutant whose ambient air concentrations notoriously exceed air quality standards throughout Europe".

55 The Eurodelta-Trends (EDT) multi-model experiment, involving eight chemical transport models (CTMs), has been designed in order to better understand the evolution of air pollution and its drivers since the early 1990s. The main objective of the experiment is to assess the efficiency of air pollutant emissions mitigation measures in improving regional scale air quality in Europe. The multi-model trend analysis is a contribution to the assessment of the evolution of air pollution in the EMEP region over the 1990-2012 period coordinated by the Task Force on Monitoring and Modelling (TFMM) of EMEP (Cooperative Programme for Monitoring and Evaluation of the Long-range Transmission of Air Pollutants in Europe). The synthesis of the observational and modelling evidences of atmospheric composition and deposition change in response to actions taken by to control emissions were given in Colette et al. (2016).

A number of studies of European (and global) PM trends for the 1990s and 2000s have been performed and published ~~last years~~recently. Some studies analysed observed PM trends (e.g. ~~Guerreiro et al. (2014), Barmpadimos et al. (2012), Cusack et al. (2012)~~ EEA (2009), Crippa et al. (2016))(e.g. Guerreiro et al., 2014; Barmpadimos et al., 2012; Cusack et al., 2012; EEA, 2009; Crippa et al., 2016), including those derived from remote sensing observations (Van Donkelaar et al., 2015), whereas a limited number of ~~analysis~~analyses also included model simulations (e.g. ~~Colette et al. (2011), Mortier et al. (2020), Colette et al. (2021), Myhre et al. (2017)~~)(e.g. Colette et al., 2011; Mortier et al., 2020; Colette et al., 2021; Myhre et al., 2017). Rather a large spread of observed and modelled PM trends, both decreasing and increasing, has been reported for the period between 1998-2002 and 2008-2014. In those studies, the setup of model runs was only partly harmonised, i.e. the models in the same study used the same emissions, but otherwise different meteorology, grid resolution etc. Analysis of EMEP observed 2002-2012 trends, also performed under TFMM coordination by Colette et al. (2016), reported the median trends of $-0.35 \mu\text{g m}^{-3} \text{yr}^{-1}$ PM₁₀ and -0.29 for PM_{2.5}, resulting in the reduction over the period by -29 and -31% , respectively, with 95% probability. As we discuss in this paper, being overall consistent with the earlier trend assessments, the results presented here are believed to be more robust as they 75 rely on a multi-modelling approach.

The main science and policy questions addressed by the EDT modelling experiment are formulated in Colette et al. (2017a), in which also the design and technical specifics of the modelling exercise are described in detail. The studied period covered a 21-year time-span, from 1990 through 2010, and in total eight regional CTMs have participated. In this paper, we present the results of trend study with respect to Particulate Matter (PM) pollution in Europe. An in-depth trend analysis for PM₁₀

80 and PM_{2.5} has been performed for a period of 2000-2010, based on multi-model simulations and EMEP monitoring data. The shorter period for PM trend study than the 1990-2010 EDT period was chosen due to the lack of appropriate PM₁₀ and PM_{2.5} observations prior to 2000. Not all of the eight EDT models had resources to perform all simulations (Sec. 2.1, therefore trend analysis presented in this work are based on the results from six of the models. Also, multi-model simulated PM trends during the whole 1990-2010 period are briefly discussed here. The strength of the presented assessment is that the model ensemble
85 simulated PM trends represent more a robust estimate as compared to either of the individual models, while the multi-model simulations allowed us investigating into the variability of modelled results, obtained under the this controlled setup. Finally, the model simulations allow interpreting PM trends in term of the trends in the individual aerosols. This is a valuable contribution to better understanding the correspondence between emission changes and PM concentration levels across Europe, given the lack of observational data on PM chemical composition.

90 The paper is structured as follows: Section 2 describes the methods used, including brief information on the models, runs' setup, observations and trend calculations; Section 3 summarizes model evaluation with respect to PM; Section 4 presents emission trends; Section 5 is dedicated to PM 2000-2010 trend analysis for the whole Europe and for set of measurement sites, discusses PM seasonal trends and the relative contribution of PM components; in Section 6 we show modelled PM trends for the 1990-2010 period. Further discussion of the result is given in Section (including emission uncertainties and effect of
95 meteorological variability); and finally the main outcomes and findings can be found in Section 8.

2 Methods

2.1 Models, runs setup

The trend analysis is based on the results from six of the EDT models, namely the ones which provided a complete series of 2000-2010 simulations. Those models are CHIMERE (CHIM), EMEP MSC-W (EMEP), LOTOS-EUROS (LOTO), MATCH,
100 MINNI and Polair3D (POLR). These models, with the exception of POLR, also performed simulations for the 1990-1999 period.

A comprehensive description of the models that participated in the Eurodelta-Trends experiment, the simulations setup, input data and the overview of the computations performed is given in Colette et al. (2017a).

Briefly, the setup and input data for the EDT simulations were harmonized as far as possible. The models performed the
105 simulations on the same grid with a resolution of $0.25^\circ \times 0.4^\circ \times 0.4^\circ$ in latitude-longitude coordinates. The simulations were driven by the same meteorological input from hindcast simulations of the CORDEX project (-Jacob et al. (2014) and Stegehuis et al. (2015)) using the WRF (Weather Research and Forecast) model (Skamarock et al., 2005) at $0.44^\circ \times 0.44^\circ \times 0.44^\circ$ resolution and using boundary conditions from ERA-Interim reanalysis (Dee et al., 2011). The exceptions were LOTO and MATCH, which used ERA-Interim reanalysis downscaled respectively by RACMO2 (Van Meijgaard et al., 2012) and
110 HIRLAM (Dahlgren et al., 2016).

Furthermore, the models used the same gridded anthropogenic emissions of SO₂, NO_x, NH₃, NMVOC, CO, PM₁₀ and PM_{2.5} (-Terrenoire et al. (2015) and Bessagnet et al. (2016)). The national emissions were based on the ECLIPSE_V5 dataset,

constructed by the Greenhouse Gasses and Air pollution INteraction and Synergies (GAINS) model (Amann et al. (2011), Amann (2012), Klimont et al. (2016), Klimont et al. (2017)) and provided in SNAP (Selected Nomenclature for reporting of Air Pollutants) sectors. Spatial distribution of the national sectoral emissions ~~was~~ performed by INERIS applying auxiliary information which included road maps (for SNAP sector 7), shipping routes (for SNAP 8) and population density (for SNAP 2), the European Pollutant Release and Transfer Register (for SNAP 1, 3, 4), TNO-MACC inventory for NH₃ emissions, as well as bottom-up emission inventories for the UK and France (see details in Colette et al. (2017a) and references therein). Time changes in the spatial distribution was accounted for only for industrial emissions. Vertical distribution and temporal profiles for the emissions used in the model simulations were those used in the EMEP model standard setup (Simpson et al., 2012). The ECLIPSE_V5 emissions were available for the years 1990, 1995, 2000, 2005 and 2010, while for the intermediate years the emissions were derived through linear interpolations (Colette et al., 2017a). For temporal distribution of ECLIPSE annual emissions, the models applied the same monthly and hourly profiles based on Denier van der Gon et al. (2011); they also used the same static vertical profiles for the emissions, based Bieser et al. (2011), applied per SNAP activity sector (none of the models included explicit plume rise simulations). Regarding, chemical speciation of PM₁₀ and PM_{2.5}, the models were allowed to use their own preferred factors to split PM emission to elemental and primary organic carbon (e.g. based on Kuenen et al. (2014), or as in Simpson et al. (2012), see Summary Table in Fig. A16).

At a rather late stage of the experiment, an error was detected in the emissions of primary particulate matter from international shipping ~~and also from~~ Russia and North Africa for the period 1991-1999. Since this error was identified late in the analysis process it was not possible to re-run the simulations with corrected emissions. The additional analysis of the impact of this error carried out with the CHIMERE model showed that these errors are relatively small compared with the overall uncertainty of the model estimates and the uncertainty of the observations (see more details in Theobald et al. (2019)). Nevertheless, the main focus of this paper is on the analysis of PM trends in the course of the 2000s, i.e. the period for which model results were not affected by the emission error.

Natural emissions of biogenic VOCs, soil NO_x, sea salt and mineral dust were calculated ~~for~~ prescribed within the models individually. Online computations of windblown dust from erodible soils were performed by EMEP, LOTO and MINNI, whereas the other models included solely mineral dust from boundary conditions. ~~Forest fire emission Emissions from forest fires and volcanoes~~ were not included in the EDT simulations ~~and emissions from Italian volcanoes were only included in the EMEP and MATCH models as the main research focus was to investigate whether the models could reproduce the trends caused by anthropogenic emission changes and changes in meteorology (see discussions on possible implications of~~. Finally, the common boundary conditions, provided by the EMEP group were based mainly on a climatology of observational data (Simpson et al., 2012). Given harmonization of set up and main input data (with a few exceptions), the differences in model results should mainly result from differences in the process formulations within the models themselves.

2.2 Observations

The observations collected at the EMEP monitoring network are annually reported to the Chemical Coordinating Centre of EMEP (Tørseth et al., 2012). All submitted observational data, after routine quality and consistency control, are available in

EBAS (<http://ebas.nilu.no>). At most of the sites, 24-hourly samples were taken on a daily basis (See Table A1). Most of the sites used a gravimetric method for both size fractions, though some used monitors. The same methods are used during the whole period. Details about site locations and applied methods are found in Table A1.

150 As documented in Colette et al. (2016), the selection criteria for sites included in the trend analysis were: i) the data capture should be at least 75 % for a specific year to be counted and ii) the number of these counted years should be at least 75 % of the total number of years in the period, and had undergone visual screening tests. The datasets used in this work include yearly measurements of observed trends from respectively 26 and 13 sites of PM₁₀ and PM_{2.5} for the period 2000-2010 (Table A1 and Fig. 5, upper panel).

155 Among those 'trend-sites', PM₁₀ observations are available for all eleven years of the 2000-2010 period at 16 sites, and ~~only~~ at 4 ~~site-sites~~ for PM_{2.5} (Table A1). ~~The reason for gap years is either PM was not measured in that year, or the criterion of 75% for data coverage was not satisfied. For most of the sites with incomplete data series, 2000 is a gap-year, as PM monitoring was not started before in 2001 at those sites. In particular, the observational data is missing at rather many sites for the first trend period year, namely 2000. For the calculations of trend slopes and for models evaluation, only available observational data is used (the number of valid years used in those statistical calculations is given in Table A1). For time-series of annual and seasonal mean-~~ The other gap years are: 2009 at the Czech CZ0003R site, 2003 and 2004 at the British GB0043R, and 2009 for PM10 and , averaged over all trend-sites (Figures 4 and 8), a gap filling of the missing years was undertaken, to ensure that the same sites were used through the years 2000-2010. The gap fillings were done by averaging between the adjacent years . For the first (2000) or the last (2010) year, the missing data were filled with values from the next or previous year for PM2.5 at the
160 Swedish SE0002R (for detailed info see Table A1).
165

2.3 Trend calculation

The Mann Kendall (MK) method (~~Mann (1945)~~, Mann (1945) and Kendall (1975)) has been applied to both modelling results and observed data for identification of significant trends. The linear trends have been calculated using the Theil-Sen slope method (known to be robust to outliers), applying the probability level of 95% as a threshold for trend significance. The trend
170 calculation method used here is consistent with that in trend assessment reported in Colette et al. (2016). In addition to absolute concentration trends, relative trends have been calculated using an estimated concentration at the start of the period (i.e. the year of 2000) as reference (see Appendix A3 in Colette et al., 2016). This concentration value corresponds to PM concentration in 2000 according to the trend line, and is considered to be less sensitive to inter-annual variability than the actual observed or modelled ones.

175 A synthetic testing of the efficiency of MK methodology to identify significant trends and estimate Sen's slopes has been performed (S. Solberg, *personal comm.*, https://wiki.met.no/_media/emep/emep-experts/mannkendall_note.pdf). It showed that the chance that the MK method detects the long-term trend decreased for shorter data-series, large natural variability and relatively weak trends. ~~Further, it was found that gap-filling of data for 1-2 years did not appear to impact MK analysis results.~~ The extent to which these factors could have affected the results of our trend analysis is discussed in Sec. 7.3.
180 Furthermore, the aforementioned document also demonstrates that ~~taking into consideration (averaging)~~ averaging significant

trends only would overestimate mean absolute trends, therefore both significant and insignificant trends have been included when calculating site-average PM trends.

3 Models evaluation

Model simulated PM_{10} and $PM_{2.5}$ have been evaluated against observations at the trend-sites (26 and 13 respectively) for the 185 years from 2000 through 2010, and averaged over the measurement sites performance statistics in terms of annual mean bias and spatial correlations are summarized in Figure Fig. 1 and Tables A2 and A3 (Appendix A).

Figure 1 shows the relative biases (in %) for the individual model and the ensemble mean. The modelled PM_{10} and $PM_{2.5}$ tend to be biased low compared to the observations (marked by blue colours of different intensity). On average, the model ensemble underestimates annual mean PM_{10} by 12% and $PM_{2.5}$ by 14 % over the period 2000-2010 (rather different biases 190 for 2000 are due to fewer sites with data). PM_{10} mean relative biases for the individual models are in a range of 5-11 %, that is somewhat smaller than their biases of 5-20 % for $PM_{2.5}$ (with POLR standing out with PM_{10} bias of -31% as erroneously simulated coarse sea salt had to be excluded).

Furthermore, we find a quite moderate year-to-year variability of the model ensemble bias, namely between -7 and -18 % for PM_{10} and between -2 and -20 % for $PM_{2.5}$. This robustness in PM simulation also applies to the individual models, i.e. 195 the inter-annual bias variations are mostly within 5 % (up to 10 %). The consistency in terms of bias can be noticed between the models (e.g. smaller underestimation of PM_{10} for 2000, 2001, 2007, 2008 and 2009, whilst slightly larger underestimation for the years 2003, 2006 and 2010, characterised by elevated PM levels).

The average annual coefficients of spatial correlation (R) are 0.54 (0.41 - 0.58) for PM_{10} and 0.65 (0.58-0.72) for $PM_{2.5}$. Similar to model biases, the correlation varies only moderately between the years and the models (Tables A2 and A3). Models' 200 evaluation for the individual aerosol components and their gaseous precursors can be found in the other EDT publications (e.g. Ciarelli et al., 2019; Theobald et al., 2019).

4 Emission trends

The graphs in Figure Fig. A1 present the changes in European annual emissions, used in this work. The total emissions of aerosol gaseous precursors SO_2 , NO_x and NH_3 and primary fine and coarse PM ($PM_{2.5}$ and $PM_{10-2.5}$) are shown for the 205 whole period of EDT study, i.e. 1990-2010. The total emissions of all pollutants decrease during this period, although at different rates. From 1990 to 2010, the greatest decrease by 69 % is in SO_2 emissions, following by NO_x emissions which are decreased by 39 %. The reduction in NH_3 emissions is rather moderate 15 %. Quite considerable decrease is seen in primary PM emissions, which go down by 67 and 47 % for coarse PM and $PM_{2.5}$ respectively.

During the period of 2000-2010, which is in a focus of this publication, the total emission decreases are: 37 % for SO_2 , 17 210 % for NO_x , 6 % for NH_3 , 27 % for $PM_{2.5}$, 36 % for coarse PM, 33 % for NMVOC. For EU area, where the measurement sites

with PM observations available for the trend analysis are located, SO₂ is reduced by 24 %, NO_x by 22 %, NH₃, PM_{2.5} and coarse PM by 10 % during the same period.

Further details on emission changes across the EDT domain are provided in [Figure-Fig. A2](#), which shows the maps with annual mean trends in the emissions of primary PM and their gaseous precursors during 2000-2010 and 1990-2010. During the period of our attention 2000-2010, the emissions of SO₂ and NO_x go down in all countries, but there are many hot-spots with upward trends (also in some Eastern and south-Eastern countries for NO_x). The negative trends of SO₂ emissions are 3-7 % per-year-yr⁻¹ in most countries, exceeding 7 % /yr-yr⁻¹ in Italy, Hungary, Portugal, Ireland and parts of Sweden and Finland (below 3% /yr-yr⁻¹ in Western Balkan, Norway and Russia). NO_x emissions show a reduction of 3-5% /yr-yr⁻¹ in Central Europe and Italy, going up 5-7 % /yr-yr⁻¹ and above in Sweden, some spots in Finland, Denmark, the UK and Portugal. NO_x decreases less (by 1-3 % /yr-yr⁻¹) in Norway, parts of Spain and Eastern Europe, and increases by 1-3 % /yr-yr⁻¹ in Russia, Belarus, parts of Poland. SO₂ and NO_x emissions from international shipping decrease in the North Atlantic and the Baltic Sea, but increase in the Mediterranean Sea. Also NH₃ emissions show negative trends in most of the domain, with decrease by 0.5-3 % /yr-yr⁻¹ in most of Europe (by 3-5 % /yr-yr⁻¹ in Denmark), but they remain nearly unchanged in Scandinavia and even increase by 1-3 % /yr-yr⁻¹ in Belarus, Lithuania, Estonia and Bosnia and Herzegovina and by (0.5-1.5 % /yr-yr⁻¹) in Poland.

During 2000-2010, PM_{2.5} emissions show downward trends in Central Europe and Norway (-3-5) % /yr-yr⁻¹ and in the rest of Eastern Europe, Spain and Scandinavia (-1-3)/yr), while they go up (by 1-4 % /yr-yr⁻¹) in Italy, Poland, Denmark, Bosnia and Herzegovina, Serbia, Moldova and Turkey. Finally, the largest decrease in coarse PM emissions is in Portugal (by (3-5) % /yr-yr⁻¹), and in the UK, Belgium and parts of Central and South-Eastern Europe (by (1-5) % /yr-yr⁻¹), but there are hot-spots with 1-4 % /yr-yr⁻¹ emission increase in the latter areas. PM coarse emissions also increase in parts of Scandinavia and Finland, in the Baltic countries and in Russia (by 1-4 % /yr-yr⁻¹), whereas they change little elsewhere.

5 PM trends for the period 2000-2010

5.1 Modelled and observed European trends

Figure 2 shows the maps of mean annual trends (Sen's slopes) of PM₁₀ and PM_{2.5} over Europe for the period of 2000-2010, calculated by the ensemble of six models (mean of EMEP, CHIM, LOTO, MINNI, MATCH and POLR) and observed at EMEP sites. The trends are presented in terms of absolute (in $\mu\text{g m}^{-3} \text{yr}^{-1}$) and relative to the starting year of 2000 (% yr⁻¹) annual changes. Significant trends are represented by coloured contour maps (modelled) and triangles (observed), whereas the insignificant trends are shown as grey areas and circles respectively.

The model results over the simulation domain and the observations at the trend-sites show overall decreasing trends of PM₁₀ and PM_{2.5} levels between 2000 and 2010. The modelled mean decreasing trends vary over the studied domain from below 0.1 $\mu\text{g m}^{-3} \text{yr}^{-1}$ in northern Europe to 0.1-0.3 $\mu\text{g m}^{-3} \text{yr}^{-1}$ in the eastern parts, and to 0.3-0.5 $\mu\text{g m}^{-3} \text{yr}^{-1}$ in central Europe and most of the UK, with PM_{2.5} downward trends being just slightly smaller than those for PM₁₀. Starting from the concentration levels in 2000, the mean relative decreasing trends range mostly from 0.1 to 0.3 % yr⁻¹ for PM₁₀ and PM_{2.5}. Compared to the

distribution of absolute trends, steeper slopes of relative decreasing trends are also seen in the southern parts of Fennoscandia
245 in addition to Central Europe and the UK.

The 6-model simulated mean trends are in general comparable to the observed ones, still some discrepancies are seen in
their geographical distribution. For instance, quite strong decreasing trends for PM₁₀ and for PM_{2.5} are observed at three of
the Spanish sites, while the model ensemble ~~does not practically indicate~~ hardly indicates any significant trends over Spain. It
should be noted that the models do calculate negative PM trends for the Spanish sites (as seen in A7), but due to considerable
250 inter-annual variability most of them are not identified as significant. Furthermore, the models calculated strongest decreasing
trends of 0.5-0.7 $\mu\text{g m}^{-3} \text{yr}^{-1}$ for PM₁₀ and PM_{2.5} in Portugal and Benelux, but no measurements were available to validate
the modelled results. For Germany, the slopes of observed trends are similar or somewhat lower than the modelled, but unlike
the model results, none of the observed trends was identified as significant. In the next sections, the trends at the individual
monitoring sites will be considered more closely.

255 Figure 3 illustrates the inter-model variability in PM trend slopes, showing the Coefficient of Variability (COV) of the trends
simulated by the individual models relative to the ensemble mean (STD/ensemble mean) for PM₁₀ and PM_{2.5}. The COV is
somewhat larger for the modelled PM₁₀ trends compared to those for PM_{2.5}. This reflects larger uncertainties in modelling of
the coarse fraction of PM, which is mostly due to natural origin, i.e. sea salt and windblown dust. As shown in Table A16, the
models used different parameterisations for the source functions of natural aerosols (also some of them did not include online
260 simulations of windblown dust, but only mineral dust from boundary conditions).

The lowest spread in the modelled trends (below 20 %) appears in Central Europe (Germany, Czech Republic), and also parts
of Spain, northern regions of Italy and in the very south of Scandinavia for PM_{2.5}. Those regions correspond with the strongest
simulated PM trends. Otherwise, the COV is 20-40 % over most of Europe, increasing to 40-60 % in Poland, western and
northern Fenno-Scandia, the Baltic countries and parts of Russia, where the modelled trends are relatively low or insignificant.

265 The maps with annual mean PM₁₀ and PM_{2.5} trend slopes calculated by the individual models are provided in the Appendix.
Figures A3 and A4 show the Sen's slopes of PM₁₀ and PM_{2.5} simulated by the six models and the observed trends for the
period of 2000-2010. The significant modeled slopes are in general quite close to each other, indicating decreasing from 2000
to 2010 trends. Also the spatial variability of the Sen's slopes in the individual models' results shows much similarity, with the
strongest decreasing trends identified in Central Europe (in particular in the Benelux countries and Germany). The EMEP and
270 ~~LOTOS-LOTO~~ calculated respectively the largest and the weakest negative mean trend slopes, as well as the largest and the
smallest fraction of the modelling domain with significant PM trends, namely 45 % and 57 % grid-cells according to EMEP
and 17 and 38 % according to ~~LOTOS-LOTO~~ for respectively PM₁₀ and PM_{2.5}, with the results from the other for models lie
between those values. As most of the input and setup for the model runs were harmonized (Sec. 2.1), the differences we see here
are due to differences in model configurations and process descriptions (see Table A16), leading to different responses of the
275 models to the changes in emissions and inter-annual meteorological variability. Differences in the formulations of secondary
aerosol formations (inorganic and organic) can be pointed at as a very important reason for discrepancies in PM modelled
trends. Differences in aerosol removal, in particular wet scavenging efficiency, also play a certain role (besides LOTO and

MATCH were driven with different meteorology). Further note that the models have a different thickness of the lowest layer which affects the concentrations, removal and transport distances of primary PM and their gaseous precursors.

280 Relative to the year 2000, all the models simulate stronger trends for $PM_{2.5}$ compared to PM_{10} , as seen in [Figures-Fig. A5 and A6](#). This is to be expected as the natural contribution, which is strongly meteorology dependent, is greater in PM_{10} . The distribution patterns of relative trends from the models are in general similar to those for corresponding absolute trends. However, there is a difference between the models in the locations of their strongest simulated relative trends, namely in Central Europe (e.g. EMEP, MINNI, POLR) or in Northern Europe (e.g. CHIM, [LOFOSLOTO](#), MATCH). The fraction of the EDT
285 domain with significant PM trends simulated with the individual models ranges from 17 (LOTO) to 45 (EMEP) % for PM_{10} and from 38 (LOTO and MINNI) to 57 (EMEP) % for $PM_{2.5}$.

Figure 4 presents observed and modelled annual mean series of PM_{10} and $PM_{2.5}$ at the trend-sites for the period 2000-2010. Shown are the mean values from the 6-model ensemble (dotted curves in Fig. 4a) and from the individual models' results (Fig. 4b and c). Note that ~~for those time series, the gap filling of the missing years in observational data described in section 2.2 was~~
290 ~~applied~~in the year of 2000 is a gap-year at for 7 out of 26 sites for PM_{10} and at 8 out of 13 sites for $PM_{2.5}$, as described in Section 2.2. In particular, none of Spanish sites are included for 2000, bringing in some inconsistency in site averaged PM_{10} and $PM_{2.5}$ annual mean series.

Although they are ~~an underestimation underestimated~~ with respect to the observations, the annual mean concentrations of PM_{10} and $PM_{2.5}$ from the 6-model ensemble follow the observed year-to-year PM variations well, with a peak in 2003 and
295 a smaller one in 2006 (the years with heatwave occurrences, which facilitated enhanced photo-chemical formation of sulphate and secondary organic aerosols and inhibited aerosol wet removal). Furthermore, the observations show a trend stagnation for PM_{10} and increase of $PM_{2.5}$ towards the end of the period at the sites considered. This is not reproduced accurately by the models. A look at the individual sites reveals that the observed increase is the result of $PM_{2.5}$ going up from 2008/2009 to 2010 at 7 out of 13 sites. According to assessments of PM pollution in 2009 and 2010, presented in EMEP Status Reports 4/2011 and
300 4/2012 (www.emep.int), about half of the sites with PM measurements reported an increase in annual mean PM_{10} and $PM_{2.5}$ with respect to the year before. As documented in those reports, a 3-4 % decrease per year of PM_{10} was registered between 2008 and 2010, whereas average $PM_{2.5}$ levels were similar in 2008 and 2009 and increased by 4 % in 2010, averaged over all sites with PM data. However, large variations between monitoring sites were observed. For instance, enhanced annual mean PM_{10} , and particularly $PM_{2.5}$ levels, were reported for 2010 at Austrian, German, Swiss, and Finnish sites, which are among
305 the trend-sites included in the present trend analysis. The major reason for elevated annual PM levels is often the occurrence of winter pollution episodes (caused by stagnant conditions within a very low boundary layer and exacerbated by enhanced emissions from domestic heating), which are not always accurately modelled due to either an overestimation of mixing layer height by relatively coarse vertical resolution or/and underestimation in the emission input data.

In general, the EDT model ensemble reproduces the observed annual 2000-2010 series of PM at the trend sites quite well,
310 showing a high correlation of 0.95 for both PM_{10} and $PM_{2.5}$. Overall, the ensemble simulated PM_{10} and $PM_{2.5}$ concentrations are lower than observed values by 31 and 19 % respectively (a greater bias for PM_{10} is partly caused by the POLR model - see below). A fairly good correspondence with respect to PM year-to-year changes is seen in [Figures-Fig. 4 \(b, c\)](#) for the individual

models compared with observations (with the exception of PM_{10} concentrations from POLR having a low bias because the contribution from coarse sea salt was not accounted for). Some deviations of LOTO's results for 2003 and 2006 are probably due to a different meteorological driver used in the model runs (~~(LOTOS-EUROS was driven by ERA-interim downscaled by RACMO2-Colette see Sec. 2.1)~~). The correlation between the modelled and measured series of annual mean PM_{10} and $PM_{2.5}$ is high, with the following correlation coefficients: 0.96 and 0.93 for CHIM, 0.93 and 0.93 for EMEP, 0.77 and 0.85 for LOTO, 0.93 and 0.90 for MATCH, 0.93 and 0.88 for MINNI, and 0.70 and 0.87 for POLR, for PM_{10} and $PM_{2.5}$ respectively. These results give credibility to the results of the models and their ability to accurately simulate the changes in the PM levels due to emission changes, as well as represent the inter-annual variability due to meteorological conditions. These results also show that the model ensemble correlates better with the observations than the individual models when both PM_{10} and $PM_{2.5}$ annual series are considered.

~~Linear trends for and from the 6-model ensemble and observational data are represented by grey lines in Figure 4a. Averaged over the sites~~ Averaged over all sites (see Table 1), the mean ensemble simulated trends (the standard deviations STD are in parentheses) are -0.24 (STD=0.09) $\mu\text{g m}^{-3} \text{yr}^{-1}$ for PM_{10} and -0.21 (0.10) $\mu\text{g m}^{-3} \text{yr}^{-1}$ for $PM_{2.5}$. These are smaller compared with the observed -0.35 (STD=0.35) and -0.40 (0.38) $\mu\text{g m}^{-3} \text{yr}^{-1}$, respectively, but can be anticipated given models' underestimation of PM concentrations. The correspondence between model results and observations is better in terms of relative 2000-2010 trends (the STD are in parentheses), which are -1.7 (0.40) and -2.0 % (0.33) % yr^{-1} from the model ensemble and -2.1 (1.19) and -2.9 % (1.48) % yr^{-1} from the observations, for PM_{10} and $PM_{2.5}$ respectively.

330 5.2 PM trends at the individual sites

Figure 5 presents observed and simulated (by the 6-model ensemble) PM_{10} and $PM_{2.5}$ trend slopes for each site for the period 2000-2010. The sites at which significant trends were observed are marked with a star. The modelled significant and insignificant trends are represented respectively by dark and light blue bars.

The observed and ensemble-modelled PM_{10} and $PM_{2.5}$ trends at all sites are decreasing. Figure 5 shows quite a large variability of observed trends in mean concentrations in the trends observed at different sites, ranging between -0.08 and -0.88 $\mu\text{g m}^{-3} \text{yr}^{-1}$ for PM_{10} and between -0.05 and -1.5 $\mu\text{g m}^{-3} \text{yr}^{-1}$ for $PM_{2.5}$. Compared with the observations, ensemble-modelled trend slopes show less variability across the sites, with the Standard Deviations are of 0.09 and 0.10 $\mu\text{g m}^{-3} \text{yr}^{-1}$ versus 0.23 and 0.38 $\mu\text{g m}^{-3} \text{yr}^{-1}$ in the observations for PM_{10} and $PM_{2.5}$ respectively (Table 1). The modelled trends are mostly within -0.5 $\mu\text{g m}^{-3} \text{yr}^{-1}$, and rather poorly correlated with the observations between the trend sites.

The strongest negative PM_{10} trends were observed at three of the Spanish sites and one Austrian site (with decreases greater than 0.7 $\mu\text{g m}^{-3} \text{yr}^{-1}$), while the weakest (and mostly non-significant) trends were registered at British, Norwegian and some German sites (below -0.15 $\mu\text{g m}^{-3} \text{yr}^{-1}$). The strongest significant PM_{10} decreasing trend slopes were modelled for German and some other sites in Central Europe. For most of the Spanish sites, the model ensemble simulated PM decrease by 0.2 - 0.3 $\mu\text{g m}^{-3} \text{yr}^{-1}$, but the trends were classified as insignificant. In general, we see a similar pattern in the results for $PM_{2.5}$, with the exception that the strongest trend was both observed (-1.5 $\mu\text{g m}^{-3} \text{yr}^{-1}$) and modelled (-0.4 $\mu\text{g m}^{-3} \text{yr}^{-1}$) for Ispra

(IT0004) in the Po Valley. Uncertainties in the emission trends and spatial distribution could be one of the main reasons for the discrepancies between the model ensemble and observations (see Sec. 7 for more discussion).

The observed relative trends range from -0.5 to -4.5 % yr⁻¹ for PM₁₀ and from -0.5 to -5.2 % yr⁻¹ for PM_{2.5} (Figure Fig. 6). Also in this case, ensemble-simulated relative trends show less variability, with values between -1.0 and -2.5 % yr⁻¹. The strongest negative PM₁₀ trends (with rates of decrease greater than -3.5 % yr⁻¹) were observed at three of the Spanish sites and the Swedish one, whereas the weakest and mostly non-significant trends (under -1 % yr⁻¹) were registered at the British and some German sites. The indicated in the previous paragraph reason for model vs. observation differences applies also for relative trends, but for the latter the estimated PM at the start of the period (see 2.3) affects the results as well.

All in all, the observations show significant PM₁₀ trends at ~~14~~11 out of 26 sites and significant PM_{2.5} trends at only 5 out of 13 sites. A closer look at PM₁₀ and PM_{2.5} annual series at the individual sites (not shown) reveals that the sites where no significant trend was identified in the observations have a particularly large inter-annual variability of PM concentrations. Model ensemble results identify significant trends at more sites compared with the observations, namely at 18 sites for PM₁₀ and at 8 for PM_{2.5}. As can also be seen on the trend maps (Figure Fig. 2), the model ensemble and the observations do not always agree regarding the significance of trends at specific locations, even within the same country. For example in Spain, strong decreasing significant trends were observed at 4 out of 6 sites for PM₁₀ and at 3 out of 4 sites for PM_{2.5}, whereas the model ensemble mostly estimates non-significant trends. This is in contrast to the German sites, for which the models simulate significant and quite appreciable PM₁₀ and PM_{2.5} trends for all sites (as a result of emission reductions in the whole country), but significant observed trends are found for only ~~3~~1 out of 7 sites for PM₁₀ and for ~~1~~neither of 2 sites for PM_{2.5}. The reason for this seems to be that the trends were distorted by particular high annual mean PM concentrations in 2003, 2006 and 2010 at most of the German sites (not shown here).

Similar to Figure Fig. 5 for the model ensemble, Figure Fig. A7 presents PM₁₀ and PM_{2.5} mean trends calculated by the individual models, with only significant modelled trends shown. For any specific site, the trend slope values from the models are in general agreement (Figure Fig. A7), while there are discrepancies between the models with regards to the ~~simulation of significant~~significance levels of simulated trends. The largest number of significant PM₁₀ and PM_{2.5} trends were simulated by EMEP (23 and 14, respectively) and the smallest number by MINNI (10 and 7) (see also Table 1).

The relative trends from the individual models are compared with each other and with observed relative trends in Figure Fig. A8 for the set of trend sites.

Averaged over all sites (see Table 1), the ~~ensemble-modelled trends are~~-0.24 (ranging trends simulated with the individual models range from -0.16 to -0.33 ~~from the individual models)~~ μg m⁻³ yr⁻¹ for PM₁₀ and ~~-0.21 (ranging~~ from -0.19 to -0.26 ~~)~~ μg m⁻³ yr⁻¹ for PM_{2.5}. ~~The observed trends are stronger than the ensemble results, namely and are weaker than observed trends (-0.35 and -0.40 μg m⁻³ respectively. Better~~ yr⁻¹ ~~respectively). The agreement among the models is found for relative trends (ranging appears to be better in terms of relative trends that range~~ from -1.4 % yr⁻¹ to -2.2 % yr⁻¹ for PM₁₀ and from -1.8 % yr⁻¹ to -2.4 % yr⁻¹ for PM_{2.5} ~~) across those sites. The modelled relative trends compare quite well with the (site averages). Compared to absolute trends, those correspond better with observed trends (-2.1 and 2.9 % for and respectively)(Table 1)yr⁻¹ respectively).~~

5.3 PM seasonal trends

Figure 7 presents the maps of 2000-2010 seasonal mean trends of PM₁₀ and PM_{2.5} from the 6-model ensemble and the observations. For the winter season, the model ensemble estimates significant PM₁₀ and PM_{2.5} trends only in small areas, mostly in southern parts of Europe. The observational data do not show any significant trends for PM₁₀. For PM_{2.5}, the observations indicate quite strong significant trends at only three sites, i.e. in the north-east of Spain (also identified by the model ensemble), north Italy and south of Sweden. One probable reason for the limited number of sites with **insignificant** significant observed trends is negligible reductions and even increases in the emissions of primary PM from residential heating, most important in the winter period, which were not efficiently regulated.

For the summer period, both the model ensemble and observations estimate the strongest negative trends out of all seasons. Significant trends are simulated for most of the domain (except Northern Europe, south of Spain and most eastern parts of the domain). The number of sites with observed significant trends is also the strongest for summer, namely 12 out of 26 for PM₁₀ and 10 out of 13 for PM_{2.5}. In the spring and autumn periods, both modelled and observed trend slope values and the fraction of sites with significant trends are between those of winter and summer.

It can be noted that Ispra in northern Italy (IT0004) is the only site where significant PM_{2.5} trends were observed and **calculated** modelled for all seasons, with the exception of the modelled winter trend. For PM₁₀, the quite strong significant mean trends at four Spanish sites (ES0007, ES0008, ES0013 and ES0014) appear to be due to strong summer trends, whereas the trends are insignificant in the other seasons. Among the German sites, significant observed PM₁₀ trends are only identified at DE0001 and DE0007 and only for the spring period. The models agree with that, but also calculate significant trends for summer and autumn.

Figures 8 (a, b) present the annual series of the 6-model ensemble and observed seasonal mean trends of PM₁₀ and PM_{2.5} for the period 2000-2010, averaged over all trend sites. The values of absolute and relative trend slopes are summarized in Table 2.

Averaged over the trend-sites, the largest decrease in PM during the 2000-2010 period took place in the summer months for both PM₁₀, with the mean seasonal trend of $-0.32 \mu\text{g m}^{-3} \text{yr}^{-1}$ from the model ensemble and $-0.56 \mu\text{g m}^{-3} \text{yr}^{-1}$ from the observations, and for PM_{2.5} (-0.26 and $-0.51 \mu\text{g m}^{-3} \text{yr}^{-1}$, respectively). The weakest trends were found for the winter season from the models and observations for PM₁₀ (-0.13 and $-0.19 \mu\text{g m}^{-3} \text{yr}^{-1}$, respectively) and also for modelled PM_{2.5} ($-0.10 \mu\text{g m}^{-3} \text{yr}^{-1}$), whereas the observed PM_{2.5} trend has a minimum of $-0.27 \mu\text{g m}^{-3} \text{yr}^{-1}$ in the autumn season. The weakest winter trends are partly due to the larger amplitudes of the inter-annual changes in mean PM levels. In particular, the elevated winter levels of PM₁₀ and PM_{2.5} in 2006, and especially in 2010, contribute to reduce the mean seasonal trend.

Figure 9 presents the seasonal mean trends simulated by the individual models and the model ensemble, along with the observed trends. The graphs nicely visualize the seasonal variations of PM trend slopes discussed above. They also show quite a good correspondence between the trend seasonality from the individual models. Relative trends of PM show quite similar seasonal patterns, with the strongest trends in the summer and weaker ones in the cold seasons of 2000-2010 (**Figure Fig. 9**). For PM₁₀, observed relative trends are $-2.9 \% \text{yr}^{-1}$ in the winter period and $-3.7 \% \text{yr}^{-1}$ in the summer period; the respective

415 numbers from the model ensemble are -2.3 and -2.5 \% yr^{-1} . For $\text{PM}_{2.5}$, the observed and modelled summer trends are -3.7
and -2.9 \% yr^{-1} , whereas the weakest observed mean trend of -2.0 \% yr^{-1} was in the autumn and the weakest modelled trend
of -1.4 \% yr^{-1} was estimated for the winter period. The individual models largely agree on the seasonal profiles of the relative
trends, although some variability exists between the simulated trend slopes (similar to those for seasonal absolute trends).

5.4 Contribution of individual components to PM trends

420 PM_{10} and $\text{PM}_{2.5}$ is a complex mixture of different aerosol components originating from a variety of anthropogenic and
natural emission sources and so PM trends are basically the sum of individual trends of its constituents. Thus, for a better
understanding of the effects of emission reductions of different pollutants, it is imperative to look at the role of the individual
aerosol components in the changes of PM concentrations.

A comprehensive study of the trends for individual aerosols is beyond the scope of this paper. Besides, there is practically no
425 available observational data for individual PM components collocated with PM measurements during the period 2000-2010. In
fact, Birkenes in the south of Norway is the only site for which observational data for both PM_{10} and $\text{PM}_{2.5}$, and for secondary
inorganic aerosols (SIA) meet the required criteria for the trend study. Still, we think that for a better interpretation of PM trends
discussed in this paper, it is relevant to have a brief insight into the trends of PM components. Here, we summarize the main
results of modelled and observed trends of some PM components for 2000-2010. For more a detailed analysis of inorganic
430 gases and aerosols, the reader is referred to Ciarelli et al. (2019).

Figure A9 shows the maps of model ensemble simulated and observed annual mean 2000-2010 trends for SO_4^{-2} , NO_3^- and
 NH_4^+ aerosols. Note that due to the lack of consistent observational data sets (as pointed out above), the set of sites for SIA are
not the same [between the species](#) and also different from those used in PM trend analysis. The number of sites used here is 39,
14 and 13 for SO_4^{-2} , NO_3^- and NH_4^+ .

435 The absolute trends are all decreasing, though the rates are not directly comparable (since they are expressed as $\mu\text{g m}^{-3} (\text{S})$
 yr^{-1} and $\mu\text{g m}^{-3} (\text{N}) \text{yr}^{-1}$). The maps of relative trend slopes show the strongest trends all over Europe for SO_4^{-2} (between -2
and -4 \% yr^{-1} over most of the domain, exceeding -5 \% yr^{-1} in Spain), closely followed by NH_4^+ . For NO_3^- , the models only
estimated significant downward trends in Central European countries and Italy. The modelled trends for SIA are decreasing
over the entire domain, whereas the observations indicate significant increasing trends of SO_4^{-2} and NO_3^- at the Polish site
440 Sniezka (close to the Czech border). In addition, rather strong, though non-significant, positive trends of NO_3^- and NH_4^+ were
observed at two Dutch sites, and somewhat weaker positive trends at a few other sites. No observational datasets long enough
(or obtained with consistent analytical methods) for trend studies of carbonaceous aerosols were available at EMEP sites.
Shorter series for total carbon, available for three-four sites, show a 4-5 % decreasing trend between 2003/2004 and 2010.

In summary, the results presented here, and the analysis by Ciarelli et al. (2019), indicate that the models estimate a
445 somewhat larger than observed decrease of SO_4^{-2} in Central (also missing some positive trends) and Northern Europe and a
smaller decrease in Spain. The models appear to overestimate the observed negative trends for NO_3^- and also for NH_4^+ , though
to a smaller degree (one should keep in mind that for NO_3^- and NH_4^+ there is limited number of measurement sites covering
a limited geographic area). [It should be noted that none of the models accounts base cations \(i.e. \$\text{Na}^+\$, \$\text{K}^+\$, \$\text{Ca}^{2+}\$ and \$\text{Mg}^{2+}\$ \)](#)

450 in gas-aerosol partitioning of HNO₃ (see Table in A16). Those base cations are significant components of sea salt and mineral dust. They participate in aerosol chemistry and facilitate the formation of coarse NO₃⁻, consuming HNO₃ and thus making less of it available for NH₄NO₃ formation. As the emissions of sea salt and mineral dust strongly depend on meteorology (especially on surface wind speed), NO₃⁻ formed on the base cations (and consequently total NO₃⁻) is subject to inter-annual variability, which could weaken NO₃⁻ trends and lead to a larger fraction of insignificant trends. Thus, not including base cations in aerosol chemistry could be one reason for models' overestimating of the observed NO₃⁻ trends (see also discussion
455 in 7. Among the EDT models, MINNI and POLR did not included coarse NO₃⁻. CHIM and LOTO included NO₃⁻ formation on sea salt Na⁺, while EMEP and MATCH used constant reaction rates for coarse NO₃⁻ formation from HNO₃, irrespective of base cation availability (Table A16). However, we could not see any consistent differences in the relative trends of NO₃⁻ and NH₄⁺ between the models with and without coarse NO₃⁻ (not shown here), neither the comparison of NO₃⁻ trends from the individual models with observations at the rather limited number of sites gave conclusive results.

460 The relative contributions of SO₄⁻², NH₄⁺, NO₃⁻, total primary particulate matter (TPPM₁₀) and anthropogenic SOA (ASOA) to PM₁₀ trends in the period 2000-2010 estimated by the model ensemble are presented in ~~FigureFig.~~ 10. The maps reveal considerable variability of the role of the individual ~~aerosols in aerosol species~~ PM₁₀ trends across European countries. The decrease in SO₄⁻² concentrations (~~FigureFig.~~ 10a) played the dominating role ~~in many parts (particularly in Spain, the Baltic countries and Russia), but not in Germany, France, the UK or Norway. Germany, Denmark over most of the EDT~~
465 ~~domain, except from parts of Central Europe and Northern Italy (the Po Valley) are the regions with.~~ Namely, relatively large contributions of NO₃⁻ to PM₁₀ trends compared with the rest of the simulation domain (Figureare seen in Germany (and neighbouring parts of France, Czechia and Poland), Denmark, the Netherlands, and in the Po Valley (Fig. 10c). The reduction of NH₄⁺ levels, which includes both ammonium sulphate and ammonium nitrate, appears to be a-very-quite-an important contributor to the PM₁₀ decreasing trends, with the largest effects ~~calculated-estimated~~ for Poland, Denmark, and the Po Valley
470 (FigureFig. 10b). The reduction of primary PM emissions was according to the model ensemble simulations the dominating factor for PM₁₀ trends in France, Norway, Portugal, Greece and parts of the UK and Russia (FigurePortugal and southern parts of Balkan; as well as in many European cities (due to emission reductions from traffic and residential heating) (Fig. 10d). Finally, ASOA is also estimated to have quite a notable contribution of 3-7 % to PM₁₀ downward trends (though ASOA modelling is still associated with rather large uncertainties). The model results imply that the chemical composition of European
475 PM₁₀ has changed somewhat during the 2000-2010 period, with NO₃⁻ (and probably ASOA) becoming an increasingly important constituent compared with the other anthropogenic aerosols, i.e. SO₄⁻², NH₄⁺ and primary emitted PM (elemental and primary organic carbon, dust and metals).

The relative contributions of SO₄⁻², NH₄⁺, NO₃⁻, and ASOA to PM₁₀ trends in the period 2000-2010, as calculated by the individual models can be found in Appendix (~~FigureFig.~~ A11). ~~The models, with the exception of LOTO and POLR~~
480 Most of the models (but for POLR), agree that over most parts of Europe (though not in Germany)of the EDT domain, except from some central European countries, decreases in SO₄⁻² and concentrations were the main cause of PM₁₀ downward trends, with somewhat smaller contribution from decreasing NO₃⁻ levels. This is consistent with the emission trends shown in Figure Fig. A1. The largest emission reductions were achieved for SO_x, which explains the relatively strong trends in SO₄⁻² and

(and also appreciable trends in NH_4^+ (in the form of ammonium sulphate) trends concentrations. The reductions of NO_x and
485 NH_3 emissions from 2000 to 2010 were smaller compared with SO_4^{2-} . Thus, as the formation of ammonium sulphate was
decreasing in the 2000s, more and more NH_3 was becoming available for the formation of ammonium nitrate NH_4NO_3 .
Notably in Germany, as well as in the Benelux countries and the Po Valley, NO_3^- is estimated by the models to have the largest
contribution to the PM_{10} trends. However it should be kept in mind that in the regions influenced by mineral dust and/or sea
salt, some of nitric acid would be consumed in the formation of NO_3^- associated with base cations (as discussed above, this
490 is not fully accounted for in the EDT models), so that less NH_4NO_3 would be formed compared to what the EDT models
simulate.

Furthermore, the estimates by LOTO point to primary anthropogenic PM_{10} as the main component driving PM_{10} levels
down in a large part of the simulation domain. CHIM, MINNI and to some extent EMEP agree with the LOTO estimates for
Northern Europe and the area covering Benelux, northern parts of Germany and France, and the south of the UK. In contrast to
495 the other models, POLR estimated that NO_3^- contributed the most to the PM_{10} trends, whereas the contribution of SO_4^{2-} and
 NH_4^+ were rather moderate according to POLR in Central Europe, the UK, the Baltic countries. The modelled contributions
of ASOA to PM_{10} trends is below 5 % according to CHIM, MATCH and MINNI, whereas EMEP simulates contributions of
5-10 % and POLR 5-30 % . This variability can be explained by the different ways of handling SOA chemistry in the models.
Furthermore, somewhat weaker PM trends from LOTO could probably be explained by not including SOA chemistry in these
500 simulations. Similar results are seen with respect to the relative contributions of the individual aerosols to modelled $\text{PM}_{2.5}$
trends between 2000 and 2010 (Figure Fig. ??).

As far as natural aerosols are concerned, emissions are largely driven by meteorological conditions (e.g. by the surface
wind in the case of sea salt and windblown dust, while the air temperature controls emissions of biogenic VOCs - precursors
of biogenic secondary organic aerosol, BSOA). In addition, the generation of mineral dust is dependent on the availability
505 of erodible (snow and vegetation free) soil and its moisture (which in turn depends on precipitation frequency and amount),
whereas the temperature and salinity of sea water affect sea spray formation, though those conditions are less variable. Of
course, similar to anthropogenic aerosol, the transport and removal of the natural particles are determined by atmospheric
dynamics and precipitation. In short, year-to-year changes in the concentrations of natural aerosols are driven primarily by
inter-annual meteorological variability. Among natural aerosols, only BSOA have some dependency of formation of BSOA
510 has some dependency on anthropogenic emissions, as they BSOA can be formed from biogenic VOCs condensing on primary
organic aerosols from anthropogenic sources. Thus, BSOA production is somewhat affected by the trend in PM emissions. In
addition, as discussed above, the changes in NO_3^- formed from anthropogenic NO_x emissions are in fact dependent on the
variability of natural aerosols of sea salt and mineral dust.

Not all natural particles were calculated in a consistent way by all of the models. The missing components are: BVOC from
515 LOTO and MATCH, sea salt from POLR; and only EMEP and LOTO simulated trends of windblown dust in the modelling
domain, whereas the other models only included mineral dust from boundary conditions. Figures 10 (f, g, h) present the
computed contributions of natural aerosols estimated by the models, i.e. biogenic SOA, sea salt and mineral dust, to PM_{10}
trends, where the negative contributions (blue colours) mean increasing trends in the natural aerosols.

The model ensemble simulated decreasing BSOA trends that contribute 1-3 % of PM₁₀ decreasing trends over almost all land area (FigureFig. 10f), with the largest contribution (5-10%) in Fennoscandia and north-western Russia. The contribution of sea salt trends (derived as 3.26*sea salt Na, assuming 30.7 % sodium content in sea salt aerosols, the same as in sea water) to PM₁₀ trends is, on average, 2-5 % over land and exceeds 10 % in areas influenced more by the sea and less polluted regions (FigureFig. 10g). Comparison of the modelled sea salt trend with rather sparse observations can be found in FigureFig. A10 (a).

Furthermore, from the EMEP and LOTO results, we see contributions of 1-3 % from mineral dust to decreasing PM₁₀ trends over most of Europe (in excess of 10% in Spain, Italy), but also some negative contributions due to increasing dust trends in Greece, Portugal and south-eastern Europe and Russia (FigureFig. 10h). All in all, the inter-annual variability and increasing modelled trends for natural aerosols for some regions do not appear to have reversed the decreasing PM₁₀ trends in the 2000-2010 period (with some exceptions for windblown dust).

Model analysis of the seasonal trend of the individual PM₁₀ and PM_{2.5} components shows the strongest trends of SIA (SO₄⁻², NH₄⁺ and NO₃⁻) in summer and also in spring for NO₃⁻, while the weakest trends of all SIA are calculated for winter. On the contrary, the strongest trends for primary PM are simulated for winter and the weakest for summer.

6 PM trends in the period 1990 – 2010

As no regular measurements of PM were conducted prior to 2000, this paper mainly focus on the period 2000-2010. As far as the years prior to 2000 are concerned, we have to rely solely on model simulations to assess the effect of emission reductions on European levels of particulate pollution in the 1990s. Given that, any deep analysis of that decade is beyond the scope of the paper, but still we think it is relevant to present a multi-model assessment of PM trends during the whole 1990-2010 period, studied within the EDT framework. It should be kept in mind while looking at those results, that the emission data, in particular for PM, are much less reliable before 2000.

Figure A13 shows annual mean trends for the period 1990-2010 for PM₁₀ and PM_{2.5}, absolute and relative to 1990, produced by the ensemble of five models (all the above except POLR). Over the whole European domain, the models simulate significant decreasing PM trends. The strongest trends (0.75-1.0 μg m⁻³ yr⁻¹, or 2.5-3 % yr⁻¹) were simulated for Central Europe (extending eastward over Ukraine and European Russia for PM_{2.5}). The weakest trends of less than 0.3 μg m⁻³ yr⁻¹ (1.5-2 % yr⁻¹) are seen in Northern Europe and Russia and in Southern Europe. The rest of the domain experienced intermediate trends of 0.3-0.75 μg m⁻³ yr⁻¹ (1.5-2.5 % yr⁻¹ relative to the year 1990). Notably, the weakest decreasing trends (below 1.5 % yr⁻¹) are modelled for PM₁₀ in the southernmost parts of Mediterranean countries, which are heavily influenced by Saharan dust and so PM trends due to the reductions of anthropogenic emissions are distorted. The mean annual trends during the period of 1990-2010 are stronger compared with those for the 2000-2010 period (FigureFig. 2). This is a consequence of larger emission reductions in the 1990s compared with the 2000s. Thus, the EDT model ensemble simulated that annual mean PM₁₀ and PM_{2.5} concentrations decreased by between 5 and 15 μg m⁻³ across most of Europe (by 2-5 μg m⁻³ in the Northern Europe) from 1990 to 2010.

6.1 PM trends in European countries in 1990-2000-2010 periods

The graphs in [Figure Fig. A14](#) provide more details regarding PM₁₀ trends in individual European countries and compare the trends in the 1990s and 2000s.

555 Figure A14a shows the trends of PM₁₀ between 1990 and 2010 simulated by the five models for the individual countries and sea areas. The strongest annual mean trends, with decreases greater than $-0.6 \mu\text{g m}^{-3} \text{yr}^{-1}$, (leftmost countries in the graph) were simulated for Central European (Germany, Hungary, Czech Republic) and the Benelux countries, which were the regions with among the highest PM levels. The weakest downward trends are modelled for relatively cleaner North European (Iceland, Norway, Finland, Sweden) and Baltic countries, but also in Mediterranean countries influenced by shipping emissions and
560 African dust intrusions (rightmost countries in the graph). The models are in general agreement regarding the ranking of PM₁₀ national trends, and the spread between PM national trends calculated with the individual models is rather moderate (the mean STD between the models is $0.054 \mu\text{g m}^{-3} \text{yr}^{-1}$, varying between 0.005 and $0.104 \mu\text{g m}^{-3} \text{yr}^{-1}$ for different countries). The variation of PM_{2.5} trends across Europe is quite similar (therefore not shown here), with the only difference that the trends in the Benelux countries were the strongest.

565 Figure A14b shows for the individual countries and regions, the PM₁₀ annual trends calculated by the model ensemble for the 1990-2000 and the 2000-2010 periods separately. For most of the countries, the largest reductions of PM₁₀ levels took place in the 1990s compared with the 2000s, which is consistent with considerably larger emission reductions of PM emissions and their gaseous precursors (except from ammonia) during the first of those decades. This is especially pronounced in Central Europe, where the 1990-2000 trends were around $1 \mu\text{g m}^{-3} \text{yr}^{-1}$ compared with around $0.3 \mu\text{g m}^{-3} \text{yr}^{-1}$ in the
570 2000-2010 period. The exceptions are North-European countries, and also relatively small emitters of pollution, such as Malta, Liechtenstein, Cyprus, where PM₁₀ trends were similar during both decades.

The PM₁₀ relative trends (i.e. with respect to the starting years of 1990 and 2000) in the 1990-2000 period are also considerably stronger than those in the 2000-2010 period (not shown, or in Supplement). The model results indicate a large variability in 1990-2000 trends between the countries (from $-1.1 \% \text{yr}^{-1}$ in Central Europe to $-(0.0-0.2) \% \text{yr}^{-1}$ in Northern Europe,
575 Cyprus, Malta), whereas the 2000-2010 trends are more homogeneous across the countries, ranging between 0 and $-3 \% \text{yr}^{-1}$.

7 Discussion

7.1 Discussion of main results

The ensemble of six EDT models simulated that, from 2000 to 2010, the annual mean PM₁₀ and PM_{2.5} concentrations decreased by between 10 and 20 % over most of Europe, and respectively by up to 25 % and 30 % in Germany, the Netherlands,
580 Belgium, parts of the UK, Portugal, north/centre of Italy and large parts of Scandinavia. Notably, despite lower PM_{2.5} concentrations, the PM_{2.5} absolute downward trends appear only slightly smaller than those for PM₁₀, indicating a trend-masking role of coarse PM of natural origin. On average, we found a fair agreement between modelled and observed concentration reductions at 26 (for PM₁₀) and 13 (for PM_{2.5}) measurement sites. In the course of those 11 years, PM₁₀ and PM_{2.5} con-

centrations at the studied sites decreased respectively by 17 and 20 % according to the model ensemble and by 21 and 29 %
585 as derived from observational data. Moreover, we found a larger spatial variability of PM trends registered by observations
compared with those estimated by the model, with observed decreasing trends ranging between approximately 5 % (at British
site GB0036) and 50 % (at Swedish site SE0012). We also see some discrepancies in the geography of trends from the ob-
servations and EDT model, with the largest observed decreases (above 30 %) at the sites in Sweden, Finland and Spain (also
the Po Valley for PM_{2.5}), whereas the models simulate the strongest trends for German sites (mostly above 20 %) and do not
590 identify significant trends for Spanish sites (though 10–20 % decreases in PM₁₀ and PM_{2.5} is simulated).

Modelled PM concentrations are to a large degree determined by the emission data used and modelled PM trends reflect
the trends in national emissions. For instance, relatively strong simulated PM trends in Germany, the Benelux, the UK and
Portugal are due to considerable reductions of all gaseous precursors and primary PM in those countries (Figure Fig. A2).
Poland is among the countries with the greatest reduction of SO_x and considerable reductions in NO_x emissions from 2000
595 to 2010, but the increase in NH₃ emissions contributed to additional SIA formation during those years. Besides, the emissions
of primary PM_{2.5} went up in in Poland increased during the same period. Thus, the resulting modelled downward trends are
relatively weaker (and insignificant in parts of the country). In Northern Europe, the appreciable decrease of PM concentrations
is not only due to reductions in NO_x and primary PM_{2.5} emissions in those countries, but is also due to decreased long-range
transport from Central Europe and the UK (somewhat lessened by the increased NO_x emissions from international shipping
600 in the North and Baltic seas). For Spain, the ~~models-model ensemble~~ simulated a substantial decrease in PM concentrations
(though the PM trends were characterised as insignificant), mostly resulting from emission reductions of gaseous precursors,
while the reductions in emissions of primary PM (especially coarse PM) were relatively smaller. Only the EMEP model (and
MATCH for PM_{2.5}) simulated significant PM trends for most of Spain, whereas PM trends from the other models were found
to be insignificant due to smaller PM decreases from 2000 to 2010 or/and larger inter-annual variability (as in the results from
605 LOTO and MATCH, using different meteorology).

Furthermore, the analysis showed a considerable variability in the observed trends within the same country, which the models
could not fully reproduce. This can be due to local emissions, unaccounted for, or misrepresented spatially and temporally in
the model input. In some countries, the differences in trends could also be related to a complex topography leading to localised
pollution transport dynamics (e.g. Switzerland and Austria), unresolved by meteorological drivers.

610 As PM is a complex pollutant, consisting of different aerosol species, the concentrations and trends of PM are the result of
an intricate interplay of the effects of their direct emissions and gaseous precursors from a variety of anthropogenic and natural
sources. As discussed in ~~Section 4~~Sec. 5.4, the emissions of SO_x went down by 37 % from 2000 to 2010, resulting in the
decrease of ammonium sulphate concentrations and thus more ammonia available for reactions with nitric acid. The reduction
of NO_x emissions in the same period (17 %) was smaller than that of SO₂. Given rather moderate reductions of NH₃ emissions
615 (only 6% on average), the concentrations of ammonium nitrate decreased less compared with ammonium sulphate. The model
ensemble calculated the decrease for SO₄⁻² to be in a range of 25-45 % (45-55 % in Spain and Portugal) and for NH₄⁺ in
a range of 15-40 % over Europe from 2000 to 2010 (Figure Fig. A9, a-f). The modelled decrease of NO₃⁻ concentrations is
mostly under 30 % and the trends are insignificant in most countries. For more detailed discussion on SIA trends, we refer the

reader to the analysis published in (Ciarelli et al., 2019). In that publication, relatively moderate trends in SO_4^{-2} compared
620 with the emission reductions of SO_2 was explained by an increase in the availability of oxidant species and more efficient
pH-dependent cloud chemistry resulting from those emission reductions. (Ciarelli et al., 2019) also discusses a shift in the
thermodynamic equilibrium between HNO_3+NH_3 vs. NH_4NO_3 , favouring aerosol formation. Furthermore, the reduction of
anthropogenic VOC emissions, including aromatic hydrocarbons - precursors of SOA, by 33 %, on average, led to a decrease
in ASOA concentrations by 15-30 % from 2000-2010 (Figure Fig. A9, g, h). Finally, the emissions of both $\text{PM}_{2.5}$ and coarse
625 PM reduced, on average, over the modelled domain by 10 %, thus making primary PM an important driver of PM_{10} and $\text{PM}_{2.5}$
decreases in some European regions (not shown here).

Due to the lack of long-term observational data of PM_{10} and $\text{PM}_{2.5}$ supplemented with chemical analyses, the model results
regarding the role of the individual components in PM_{10} and $\text{PM}_{2.5}$ trends during 2000-2010 cannot be thoroughly validated.
We can only make a crude estimate, using observations of SIA and OC, which are not necessarily collocated, available at a
630 limited number of sites. The observed average trends were the strongest for organic aerosols (-3.8 \% yr^{-1} at 4 sites), followed
by NH_4^+ (-2.9 \% yr^{-1} at 13 sites), SO_4^{-2} (-2.6 \% yr^{-1} at 39 sites), and finally the weakest trends were for NO_3^- (-0.5 \% yr^{-1}
at 14 sites).

7.2 Uncertainties in emissions

As shown in the previous section, the modelled trends in PM and its components quite closely reflect emission reductions,
635 though inter-annual variability of meteorological conditions also plays an important role in PM pollution levels (see 7.3). This
means that good quality emission data is essential for accurate model simulations of the trends.

Emission estimates are associated with uncertainties due to missing or incomplete information, or limited understanding
with respect to activity data, emission factors, source locations etc. (Klimont et al., 2017).

No publication with a detailed and quantitative uncertainty estimate of the GAINS dataset used here (ECLIPSE_V5) is
640 available, but (Amann et al., 2011) and (Schöpp et al., 2005) described the treatment of uncertainties in the context of the
GAINS model. For example, for 1990, (Schöpp et al., 2005) estimated that the national total emissions used in the RAINS
integrated assessment model had an uncertainty of $\pm(6-23) \text{ \%}$ for SO_2 , $\pm(8-26) \text{ \%}$ for NO_x and $\pm(9-23 \text{ for }) \text{ \%}$ for NH_3
(95% confidence interval). However since that assessment, steps have been taken to reduce the uncertainty in the emission data
sets (Klimont et al., 2017). The European Environment Agency indicated somewhat larger uncertainties in typically top-down
645 emission estimates in the EU LRTAP inventory, namely around $\pm 10 \text{ \%}$ for SO_2 , $\pm 20 \text{ \%}$ for NO_x and $\pm 30 \text{ \%}$ for ~~(EEA, 2008).~~
Clearly, underestimation of emissions would lead to underestimation of the NH_3 and NMVOCs (EEA, 2008). Primary $\text{PM}_{2.5}$
and PM_{10} emission data is said to be of relatively higher uncertainty compared to emission estimates for the secondary PM
precursors. Clearly, uncertainties in emissions will inevitably be reflected in the uncertainties in absolute trends of PM.

Furthermore, EEA (2008) suggested that the emission trends are likely to be more accurate than the individual absolute
650 annual values, although the use of gap-filling when countries have not reported emissions for one of more years can potentially
lead to artificial trends. Regarding primary PM emissions, ECLIPSE_V5 was the first assessment of PM_{10} and $\text{PM}_{2.5}$ emis-

sions, performed using a consistent bottom-up approach across all sources and regions and, therefore, only limited comparison to other works was possible (Klimont et al., 2017).

One of the biggest sources of emissions-related uncertainty is likely to be residential wood-burning emissions of PM and VOCs (forming ASOA) (Simpson et al., 2020). Emissions of primary organic matter (POM) from residential wood burning have been known to be problematic for many years (~~Denier van der Gon et al., 2015; Simpson and Denier van der Gon, 2015~~) (Simpson et al., 2020; Denier van der Gon et al., 2015; Simpson and Denier van der Gon, 2015), with different countries accounting for, or omitting, semi-volatile compounds in different and often unknown ways. Given that wood burning for heating houses accounts for a significant percentage of European PM emissions, the lack of consistent treatment between countries has obvious implications for the reliability of any trend estimates. There is an increasing recognition that emissions of some potentially important SOA precursors, namely semi-volatile and intermediate-volatility organic compounds (SVOC, IVOC) from traffic sources, are also missing from national inventories and these can have significant impacts on ambient organic matter (OM) (Ots et al., 2016). Emissions of SVOCs and IVOCs are very dependent on e.g. the fuel and type of catalyst used in cars (Jathar et al., 2014; Platt et al., 2017), with older vehicles likely emitting substantially more than new ones, again complicating any analysis of trends. Even for the same country, condensable organics might be included or excluded differently for different sectors. Inclusion or exclusion, or the extent of inclusion of condensables, has also changed over the years, which directly affects the accuracy of trend analyses (Aas et al., 2021). It is also worth noting that the models did not account for the dependence of residential heating emissions on the outdoor temperature, i.e. they increase as it gets colder. This may lead to model underestimation of winter pollution episodes, resulting in under-predictions of annual mean PM (as for 2010, see 5.1). Finally, with respect to anthropogenic sources, assumed invariant spatial distribution of emissions (except from industrial sectors) may cause inaccuracy in modelled trends in some areas.

As far as natural emissions are concerned, biogenic VOC (BVOC) emissions estimates have also many uncertainties, both for isoprene and monoterpenes (e.g. Simpson et al., 1999; Langner et al., 2012; Messina et al., 2016). The models in this study calculate BSOA formed from the oxidation of isoprene and terpenes (CHIMERE also includes sesquiterpenes), but additionally BSOA can also be formed from the oxidation of stress-induced emissions of other VOCs that are not included in the emissions; this process is likely to be quite frequent, but can only be accounted for in speculative terms with current knowledge (Bergström et al., 2014). Beside uncertainties in emission estimates, the emission data used in the model runs omit some sources of PM. Among the omitted sources of OM are primary biological material, which can contribute e.g. 20-30% of PM₁₀ in Nordic areas in summer-early autumn (Yttri et al., 2011) (though it is likely to be much less as an annual average (Winiwarter et al., 2009)). Marine sources of OM also contribute to observed ambient OM (e.g. Spracklen et al., 2008), but the models used here have not accounted for those (some models, such as EMEP, have assumed background levels of OM which account for such diverse sources, but only in a crude way and with the same levels assumed for all years).

As described in 2.1, pollution from forest fires were not accounted for in EDT simulations mainly because of considerable uncertainties in forest fire emissions and modelling of those, but also because we aimed to look at PM trends due to emission regulation in Europe. An in-depth analysis of the effect of forest fires on PM trends is beyond the scope of the paper, but we have tested whether the discrepancies between the modelled and observed trends, in particular in terms of a relatively

larger fraction of significant trends from the model results, could be due to not including forest fire emissions in the EDT simulations. Additional simulations suggest that the effects from even large fires during the studied period (like 2010 Russian forest fires) were mostly negligible outside the regions where wild fires occurred. In fact, the pollution from major forest fires did not seem to have any large impact on simulated annual mean PM at the EDT sites in the 2000-2010 period. Therefore we are certain that not accounting for forest fires in EDT analysis did not have any significant consequences for models vs observations comparison. The same applies to not including volcano emissions in the trend simulations. For example, EMEP source-receptor calculations indicate a rather limited contribution to PM_{2.5} in European countries from volcano emissions (see for example the contributions from Italian Etna, Stromboli and Vulcano and also Eyjafjallajökull eruption in 2010 in EMEP (2012)).

7.3 Effect of inter-annual meteorological variability

As pointed out in Section 2.3, the probability of trend detection using the Mann-Kendall method decreases for shorter data-series, large natural variability and relatively weak trends. The bottom-line is that the weaker the trend is relative to the inter-annual meteorological variability, the longer the time series that is needed in order to identify a significant trend. As estimated in https://wiki.met.no/_media/emep/emep-experts/mannkendall_note.pdf, for an 11-year time series, the chances for MK methodology to detect significant trends are very small for trends of $-1\% \text{ yr}^{-1}$, with only 36% of significant trends identified by MK test will be between 92 and 100% for an inter-annual meteorological variability of just 5%, dropping down to (going down to 9% for inter-annual variability of 15%). The probability for stronger trends to be identified as significant increases, but still will be between 37 and 71% for a 10% variability and even further down to between 19 and 39% for a 15% variability. For weaker trends of -1% , the corresponding probabilities are only 36, 13 and 9% for an inter-annual meteorological variability of 5, 10, and 15% respectively.

Most of aerosol processes (some emissions, gaseous and especially heterogeneous chemistry, transport and removal) depend on the meteorological conditions. The model simulations performed in this work indicate that the relative inter-annual variability of PM concentrations due to meteorological variability is by far and large below 15% relative to the 11-year mean. It varies mostly between 5 and 10% over Central/Eastern Europe with somewhat larger variability over Northern Europe and Iberian Peninsula during 2000-2010 most of Europe, 10-12% in parts of Scandinavia and the UK, and goes up to 15-17% in the Iberian Peninsula (not shown here). This can explain the large number of sites (and areas) where the models do not estimate significant trends in the 2000-2010 period, given that average. That means that in the part of Europe, where the modelled trends are relatively strong ($-(1.5-2.5)\% \text{ yr}^{-1}$), the MK analyses has identified more significant trends (e.g. in Central and South/South-Eastern Europe). In the Iberian Peninsula, significant modelled trends are only seen in Portugal, where the PM trends are between 1 and 3%. Still, the observations indicate even more sites with non-significant PM trends quite strong ($-(2-3)\% \text{ yr}^{-1}$, but not in Spain with $-(1-2)\% \text{ yr}^{-1}$ trends. Also in southern parts of Scandinavia with PM inter-annual variability of 10-12%, PM modelled trends of $-(2-2.5)\% \text{ yr}^{-1}$ are found significant in most of modelling grid-cells. As already mentioned, compared to ensemble modelling, MK analysis could not see significant trends in PM observations at a larger number of the

trend sites. This is due to a relatively large inter-annual variability with respect to the model trend magnitudes in PM observed concentrations (e.g. at German, Austria and Swiss sites, as discussed 5.2).

In addition, we have looked at the relative effects of emission changes and inter-annual meteorological variability on PM trends by calculating the so-called normalised relative trends (NRT) introduced in Solberg et al. (2009) and also applied in
725 Colette et al. (2011). For this purpose, we used additional model results obtained from model runs with fixed 2010 emissions for the meteorological conditions 1990 to 2010 (i.e. Tier 3B as described in Colette et al. (2017a)). The effect of the emissions on PM trends was assumed to be represented by the difference in PM concentrations obtained for corresponding years in the trend runs (Tier3A) and the runs with constant emissions (Tier3B); and the inter-annual variability due to meteorological conditions was quantified by standard deviation of annual PM concentrations in the runs with constant emissions. That is to say,
730 we calculated the ratio of the difference of Sen's slopes ($PM_{Tier3A} - PM_{Tier3B}$) to $STD(PM_{Tier3B})$. The model ensemble NRT for PM_{10} and $PM_{2.5}$ are presented in [FigureFig. 11](#), where absolute NRT values greater than one indicate a larger importance of emission changes with respect to the inter-annual meteorological variability.

Figure 11 shows that the effect apparent significance of emission reduction on decreasing PM trends appears to be downsized partially masked by inter-annual meteorological variability in large part of Europe in the 2000-2010 period. It should be noted
735 that the individual EDT models have different sensitivity to meteorological variability (besides MATCH and LOTO used different meteorological drivers), which may mask the effects of emission changes. The emission reductions play a larger role in $PM_{2.5}$ trends, as PM_{10} concentrations (particularly the coarse fraction of natural origin) are more affected by variability in meteorological conditions. Evidently, the most pronounced effects of emission reductions are associated with the regions with greater emission reductions, e.g. Portugal, Benelux, some parts of South-Eastern European and the Balkan countries. These
740 results are consistent with the main conclusions from the study of PM trends in the period 1998-2007 by Colette et al. (2011). Colette et al. (2017b) arrived to somewhat different conclusions based on a different approach, namely the decomposition of the differences in EDT modelled PM concentrations in 2000 and 2010 to discriminate the role of emissions, meteorology and boundary conditions. Their analysis suggested a relatively larger on average role of emissions compared with the meteorology, though the estimated uncertainties were non-negligible. Due to different premises used by Colette et al. (2017b) and this
745 paper, discrepancies in the outcomes are to be anticipated. That is, here we compared 11-year PM trends with year-to-year PM variability due to meteorological conditions, whereas Colette et al. (2017b) looked at the difference between 2010 and 2000.

To summarise, given rather moderate reductions (and even some increases) in the emissions of some PM precursors and primary PM between 2000 and 2010, we estimate that the effect of emission decreases on 2000-2010 PM trends is roughly of the same order of magnitude as the effect of inter-annual meteorological variability. Separating the effects of emission
750 changes and meteorological variability on PM trends, we get additional insights regarding their relative roles. PM trend slopes due to emission trends ([FigureFig. 11](#)) appear to be quite similar to the total trends wherever the latter are more significant ([FigureFig. 2](#)). The remarkable difference between them is that the trends due to emissions are significant for nearly the entire domain. Model simulated PM trends due to solely inter-annual meteorological variability (not shown) are by far and large very small ($\pm 0.05 \mu\text{g m}^{-3} \text{yr}^{-1}$) and non-significant everywhere. Thus, our results suggest that the main impact of variable
755 meteorological conditions is to reduce the significance level of PM trends due to emission reductions, while the effects on PM

trend slopes are much smaller. For comparison, since the emission reduction during the 1990s were overall larger than in the 2000s, the effect of emission reductions on the decreasing PM trends is estimated to dominate meteorological variability in most of Central, Eastern and South-Eastern Europe (Figure Fig. 11).

8 Summary and outlook

760 The Eurodelta-Trends multi-model experiment, aimed to assess the efficiency of emission mitigation measures in improving air quality in Europe, was designed to answer a series of questions regarding European pollution trends in the period of 1990-2010. Among these questions are: Were there significant trends detected by observations? Do the models manage to reproduce observed trends? How close is the agreement between the models and how large are the deviations from observations? In this paper, we address these issues with respect to PM pollution.

765 An in-depth trend analysis has been performed for PM₁₀ and PM_{2.5} for the period of 2000-2010 (limited by the availability of observations), based on results from six CTMs and observational data from the EMEP monitoring network. Given harmonization of set up and main input data (with a few exceptions), the differences in model results should mainly result from differences in the process formulations within the models themselves, and the spread in the models simulated trends could be regarded as an indicator for modelling uncertainty.

770 The results of the analysis strongly indicate overall decreasing trends of annual mean PM₁₀ and PM_{2.5} concentrations between 2000 and 2010, although the trends are not characterized as significant everywhere. The model ensemble simulated mean negative trends that vary from below 0.1 $\mu\text{g m}^{-3} \text{yr}^{-1}$ in northern Europe to 0.1-0.4 $\mu\text{g m}^{-3} \text{yr}^{-1}$ in the eastern parts, and to 0.4-0.7 $\mu\text{g m}^{-3} \text{yr}^{-1}$ in central Europe and most of the UK, with PM_{2.5} negative trends being slightly weaker than those for PM₁₀, with the total reductions of annual mean concentrations by between 2 and 5 (7 for PM₁₀) $\mu\text{g m}^{-3}$ (or between 10 and 30 %) across most of Europe (by 0.5-2 $\mu\text{g m}^{-3}$ in Fennoscandia, north-west of Russia and Eastern Europe) during the studied period.

That would mean that the annual mean PM concentrations decreased by between 2 and 6-5 (7 for PM₁₀) $\mu\text{g m}^{-3}$ over Europe across most of Europe (by 0.5-2 $\mu\text{g m}^{-3}$ in Fennoscandia, north-west of Russia and Eastern Europe) during the 2000-2010 period. In relative terms, the decrease of annual mean PM₁₀ and PM_{2.5} was between 10 and 20 % over most of Europe (up to 25-30 % in Germany, the Netherlands, Belgium, parts of the UK, Portugal, north/center of Italy and large parts of Scandinavia) from 2000 to 2010. We find that the modelled PM trends are fairly consistent with emission reductions in the ECLIPSE_V5 data set used here. Among possible reasons for deviations between the ~~models and observations~~ modelled and observed PM trends are emission uncertainties, impacts of inter-annual variability in meteorological conditions (on pollutant transport and removal, secondary aerosol formation, natural PM emissions etc.), model uncertainties associated with aerosol formation and removal processes (e.g. i.e. SOA formation, cloud pH dependency of SO₄ formation, ~~thermodynamics~~ heterogeneous chemistry (including gas/aerosol partitioning of anthropogenic precursors and aerosol formation on base cations of natural origin), SO₂ and NH₃ co-deposition etc.). Not accounting for forest fires in EDT simulations should also affect the accuracy of simulated PM trends, at least in the regions of large fires, whilst this does

not appear to have a major impact on the modelled trends at the EDT sites. Furthermore, we find a fairly good general agree-
790 ment in PM trends estimated by the individual models, with the inter-model variability below 30-40 % over much of Europe
(up to 50-60 % in northern and eastern parts of EDT domain). Somewhat greater variability in the modelled PM₁₀ trends
reflects larger uncertainties in modelling of the coarse fraction of PM, which is mostly due to natural origin.

Averaged over measurement sites (26 for PM₁₀ and 13 for PM_{2.5}), the mean ensemble simulated trends are $-0.24 \mu\text{g m}^{-3} \text{yr}^{-1}$
for PM₁₀ and $-0.21 \mu\text{g m}^{-3} \text{yr}^{-1}$ for PM_{2.5}, which are somewhat weaker than the observed trends of -0.35 and -0.40
795 $\mu\text{g m}^{-3} \text{yr}^{-1}$, respectively. This is partly related to models' underestimation of PM concentrations. The correspondence be-
tween model results and observations appears better in terms of relative trends for the same period, which are -1.7 and -2.0
 $\% \text{yr}^{-1}$ from the model ensemble and -2.1 and $-2.9 \% \text{yr}^{-1}$ from the observations for PM₁₀ and PM_{2.5} respectively. We
see somewhat larger spatial variability of observed PM trends with respect to the modelled trends across Europe and within
individual countries, which could partly be explained by the uncertainties associated with national sectoral emissions and their
800 spatial distribution. In addition, the regional models have difficulties to accurately resolve pollution at some of the sites located
in the regions with complex topography. The observations identify significant trends for PM₁₀ at 56 % of the sites and for
PM_{2.5} at 36 % of the sites, which is somewhat less than those identified by the models.

The strongest decreasing trends and the largest number of sites (and larger areas) with significant trends were observed
and modelled for summer concentrations of PM₁₀ and PM_{2.5}. On the other hand for the winter season, the model ensemble
805 identifies significant PM trends for very limited areas, mostly in southern parts of Europe, whilst the observed trends are not
significant at any of the sites for PM₁₀ and only at 3 out of 14 sites for PM_{2.5}. One important reason for that is the very modest
reductions and even increases in the emissions of primary PM from residential heating in winter.

The analysis reveals a considerable variability of the role of the individual aerosols in PM₁₀ trends across European coun-
tries. The multi-model simulations, supported by available observations, point to decreases in SO₄⁻² concentrations playing
810 an overall dominant role, although with some exceptions. Namely, we see relatively large contributions of the trends of NH₄⁺
and NO₃⁻ to PM₁₀ decreasing trends in Germany, Denmark, Poland and the Po Valley, while the reductions of primary PM
emissions appears to be a dominant factor in bringing down PM₁₀ in France, Norway, Portugal, Greece and parts of the UK
and Russia.

The analysis also suggests that year-to-year variability in meteorological conditions masks decreasing PM trends due to
815 emission reductions, leading to non-significant trends in many areas and at many monitoring sites between 2000 and 2010.
Still, the role of emission reduction measures is pronounced in the regions with greater reductions, where significant trends
of PM₁₀ and PM_{2.5} are both modelled and observed. The EDT model results show that the mean annual trends during the
period of 1990-2010 were stronger compared with those in the 2000-2010 period, which is a consequence of larger emission
reductions in the 1990s compared with those in the 2000s. The EDT model ensemble estimates that annual mean PM₁₀ and
820 PM_{2.5} concentrations decreased by between 5 and 15 $\mu\text{g m}^{-3}$ across most of Europe (by 2-5 $\mu\text{g m}^{-3}$ in the Northern Europe)
from 1990 to 2021.

Data availability. Technical details of the EURODELTA project simulations that permit the replication of the experiment are available on the wiki of the EMEP Task Force on Measurement and Modelling (<https://wiki.met.no/emep/emep-experts/tfmmtrendeuodelta>, last access: 22 November 2021), which also includes ESGF links to corresponding input forcing data. The EURODELTA Trends model results are made
825 available for public use on the AeroCom server (information to gain access to the AeroCom server are available at <https://wiki.met.no/aerocom/user-server>, last access: 22 November 2021). Model input and output data are permanently stored under the /metno/aerocom-users-database/EURODELTA folder on the AeroCom Server. See Colette et al. (2017) for full terms and conditions for the use of these data.

The original data used for calculating aggregated concentrations are all available from the database infrastructure EBAS (<https://ebas.nilu.no>)

830



9 Tables and Figures

Table 1. Observed and modelled (ensemble mean and individual models), PM₁₀ and PM_{2.5} annual mean trends for the period 2000-2010, averaged over all trend-sites. The standard deviation is included in parentheses. Units are $\mu\text{g m}^{-3} \text{yr}^{-1}$ and $\% \text{yr}^{-1}$ for absolute (Abs) and relative (Rel) trends, respectively. The number of sites with significant trends identified by observations and models (Nsign) is also provided.

Parameter	Trends	Obs	ENSmean	CHIM	EMEP	LOTO	MATCH	MINNI	POLR
PM ₁₀ 26 sites	Abs	-0.35(0.23)	-0.24(0.09)	-0.22(0.09)	-0.33(0.11)	-0.23(0.11)	-0.27(0.08)	-0.24(0.10)	-0.16(0.16)
	Rel	-2.1(1.19)	-1.7(0.4)	-1.6(0.36)	-2.2(0.36)	-1.6(0.60)	-2.1(0.43)	-1.6(0.41)	-1.4(1.27)
	Nsign	14		14	23	16	20	10	14
PM _{2.5} 14 13 sites	Abs	-0.40(0.38)	-0.21(0.10)	-0.21(0.1)	-0.26(0.12)	-0.19(0.11)	-0.21(0.08)	-0.21(0.1)	-0.21(0.14)
	Rel	-2.9(1.48)	-1.6 -2.0(0.33)	-1.8(0.35)	-2.4(0.43)	-2.0(0.53)	-1.9(0.40)	-1.8(0.44)	-2.1(0.77)
	Nsign	5		9	12	8	11	7	8

Table 2. Observed (Obs) and modelled (6-model ensemble; ENS) mean seasonal trends and Standard deviations (in parentheses) for 2000-2010 at all trend-sites. Units are $\mu\text{g m}^{-3} \text{yr}^{-1}$ and $\% \text{yr}^{-1}$ for absolute (Abs) and relative (Rel) trends, respectively. The numbers of sites with significant trends are given in parentheses square brackets.

Parameter		winter	spring	summer	autumn
PM ₁₀	Obs ($\mu\text{g m}^{-3} \text{yr}^{-1}$)	-0.19(0) <u>(0.29)</u> [0]	-0.33(7) <u>(0.27)</u> [5]	-0.56(17) <u>(0.31)</u> [12]	-0.25 (7) <u>(0.26)</u> (0.25) [4]
	ENS ($\mu\text{g m}^{-3} \text{yr}^{-1}$)	-0.13(3) <u>(0.10)</u> [3]	-0.28(10) <u>(0.13)</u> [10]	-0.32(17) <u>(0.17)</u> [17]	-0.26(<u>0.13</u>) [7]
PM _{2.5}	Obs ($\mu\text{g m}^{-3} \text{yr}^{-1}$)	-0.38(3) <u>(0.51)</u> [4]	-0.42(4) <u>(0.47)</u> [4]	-0.51(10) <u>(0.34)</u> [10]	-0.27(2) <u>(0.34)</u> [2]
	ENS ($\mu\text{g m}^{-3} \text{yr}^{-1}$)	-0.10(1) <u>(0.10)</u> [1]	-0.23(3) <u>(0.15)</u> [3]	-0.26(8) <u>(0.13)</u> [8]	-0.24(7) <u>(0.14)</u> [7]
PM ₁₀	Obs ($\% \text{yr}^{-1}$)	-1.4(<u>1.7</u>)	-1.8(<u>1.2</u>)	-2.9(<u>1.0</u>)	-1.6(<u>1.8</u>)
	ENS ($\% \text{yr}^{-1}$)	-1.0(<u>0.8</u>)	-1.8(<u>0.6</u>)	-2.4(<u>0.9</u>)	-1.8(<u>0.7</u>)
PM _{2.5}	Obs ($\% \text{yr}^{-1}$)	-2.8(<u>2.2</u>)	-2.7(<u>1.7</u>)	-3.8(<u>1.5</u>)	-2.0(<u>1.9</u>)
	ENS ($\% \text{yr}^{-1}$)	-0.9(<u>1.1</u>)	-1.8(<u>0.9</u>)	-2.5(<u>0.9</u>)	-2.1(<u>0.6</u>)

	CHIM	EMEP	LOTO	MATCH	MINNI	POLR	mean
2000	-2	2	-6	-11	-4	-24	-8
2001	-4	-8	-4	-9	-8	-30	-11
2002	-7	-14	-13	-16	-14	-35	-17
2003	-8	-13	-16	-12	-12	-36	-16
2004	-7	-14	-10	-14	-9	-33	-15
2005	-7	-11	-5	-12	-7	-31	-12
2006	-6	-12	4	-8	-10	-32	-11
2007	-3	-10	-11	-10	-7	-28	-12
2008	-4	-10	-8	-11	-8	-28	-12
2009	0	-8	2	-5	-3	-27	-7
2010	-9	-16	-15	-18	-17	-35	-18
mean	-5	-10	-7	-11	-9	-31	-12

	CHIM	EMEP	LOTO	MATCH	MINNI	POLR	mean
2000	-2	-5	-6	-3	6	-2	-2
2001	-18	-21	-18	-18	-10	-22	-18
2002	-19	-25	-18	-21	-12	-26	-20
2003	-18	-20	-25	-16	-6	-25	-18
2004	-17	-23	-15	-16	-6	-23	-17
2005	-21	-24	-15	-17	-9	-22	-18
2006	-20	-24	-12	-16	-9	-25	-18
2007	-18	-25	-18	-16	-9	-22	-18
2008	-11	-17	-7	-6	1	-12	-9
2009	-7	-14	1	-4	6	-12	-5
2010	-17	-24	-14	-14	-10	-19	-16
mean	-15	-20	-13	-13	-5	-19	-14

Figure 1. Model biases (%) with respect to observations for PM₁₀ (left) and PM_{2.5} (right) for the period 2000-2010. Note: coarse sea salt is excluded in PM₁₀ from POLR.




Figure 2. Mean Sen's slopes for PM₁₀ and PM_{2.5} trends in 2000-2010: absolute (a, b) and relative (c, d) slopes calculated by the 6-model ensemble (described in Colette et al. (2017a)), Appendix A3. Modelled trends – coloured contour map (grey or white means non-significant trends) and observed trends - coloured triangles (significant) and circles (non-significant).




Figure 3. The Coefficient of Variation of PM_{10} (left) and $PM_{2.5}$ (right) trends simulated with the individual models relative to the 6-model ensemble mean for the period 2000-2010.




Figure 4. Observed and modelled (simulated with 6-model ensemble) and the individual models annual mean concentrations of PM₁₀ (upper panel) and PM_{2.5} and Sen's slopes (upper-lower panel) for the period 2000-2010, averaged over the trend sites. The 95 % confidence intervals for the period 2000-2010 observed and ensemble modelled PM concentrations are shown with shaded areas. Bottom figures also show results from The number of sites with available observations for the individual models years can be found in Table 1. (Note: PM₁₀ from POLR does not include coarse sea salt). Note: gap filling of the missing years in the observations, mostly 2000 and 2001, was undertaken to ensure that the same set of sites was used throughout the whole period; see more details in the text for explanations).




Figure 5. Observed and modelled (6-model ensemble) trend slopes ($\mu\text{g m}^{-3} \text{yr}^{-1}$) for the period 2000-2010 at the trend sites for PM_{10} (top middle) and $\text{PM}_{2.5}$ (bottom). Significant modelled trends are shown in dark blue, not-significant in light blue. Sites with significant observed-non-significant trends are indicated-represented by an-asterisk striped bars. The trend sites are shown on the map (upper panel).




Figure 6. Same as [Figure Fig. 5](#), but for relative trends ($\% \text{ yr}^{-1}$). [The trend sites are shown in Fig. 5, upper panel.](#)




Figure 7. Mean Sen's slopes for PM₁₀ and PM_{2.5} seasonal trends for 2000-2010, calculated by the 6-model ensemble (see Figure 2 for explanation).




Figure 8. Changes Observed and simulated with 6-model ensemble changes in seasonal mean PM₁₀ and PM_{2.5} concentrations in the period 2000-2010, averaged over the trend-sites. Solid line: observed; Dashed line: 6-model ensemble The 95% confidence intervals are shown with shaded areas. The number of sites with available observations for the individual years can be found in Table 2.




Figure 9. Mean relative seasonal trends in the period 2000-2010 at the trend-sites for PM₁₀ and PM_{2.5}: The trends from the observations, the individual models and the 6-model ensemble are shown.




Figure 10. Model ensemble simulated relative contribution to PM₁₀ 2000-2010 trends from anthropogenic aerosols (all 6 models): SO_4^{-2} , NH_4^+ , NO_3^- , total primary TPPM₁₀ (except POLR) and anthropogenic SOA (except LOTO), and from natural aerosols: biogenic SOA (except LOTO), sea salt (except POLR) and mineral dust particles (except MATCH). Note that a different colour scale has been used for the natural aerosols.




Figure 11. PM trends due to emission changes (upper panels) and the ratio of PM changes due to emission changes to those due to inter-annual meteorological variability (lower panels) for PM₁₀ and PM_{2.5} in the 2000-2010 period. Observed trends are shown as coloured triangles (significant) and circles (non-significant).

Appendix A: Supplementary Figures and Tables




Figure A1. Annual emissions of SO_x, NO_x, NH₃, PM_{2.5} and PM coarse (pmco) in the period 1990-2010 (all countries). Units: ktonnes.




Figure A2. Emission trends for 2000-2010 (left) and 1990-2010 (right).




Figure A3. Annual mean trends (Sen's slopes) for PM₁₀ in the period 2000-2010 as calculated by the individual models. The modelled trends are shown as coloured contour map (grey or white means non-significant trends) and the observed trends as coloured triangles (significant) and circles (non-significant). Units: $\mu\text{g m}^{-3} \text{yr}^{-1}$).




Figure A4. Same as Figure Fig. A3, but for PM_{2.5}. Units: $\mu\text{g m}^{-3} \text{ yr}^{-1}$.




Figure A5. Mean Sen's slopes relative to the starting year of 2000 ($\% \text{ yr}^{-1}$) for PM₁₀ trends in the period 2000-2010 calculated by the individual models. The modelled trends are shown as coloured contour map (grey or white means non-significant trends) and the observed trends as coloured triangles (significant) and circles (non-significant).




Figure A6. Same as Figure Fig. A5, but for PM_{2.5}. Units: % yr⁻¹.




Figure A7. Observed and modelled trend slopes ($\mu\text{g m}^{-3} \text{yr}^{-1}$) for the period 2000-2010 at the trend sites: upper two plots for PM₁₀, lower plot for PM_{2.5}. The sites are sorted by decreasing observed negative trends and the sites at which significant trends were observed are marked with an asterisk and insignificant modelled trends are shown as striped bars. Units: $\mu\text{g m}^{-3} \text{yr}^{-1}$.




Figure A8. Mean observed (black) and modelled (coloured) relative trends for PM_{10} (upper two graphs) and $PM_{2.5}$ (lower graph) in the period 2000–2010 at the individual trend-sites. ~~The sites at which significant trends were observed are marked with an asterisk and insignificant~~ Insignificant modelled trends are shown as striped bars. Units: $\% \text{ yr}^{-1}$.




Figure A9. Mean observed and modelled from 6-model ensemble Sen's trend slopes for 2000-2010 for anthropogenic aerosols: SO_4^{-2} , NO_3^- and NH_4^+ simulated by the 6-model ensemble (a-f), and for ASOA from the simulated with 5-model ensemble for ASOA (note different color scale). Left panels – absolute ($\mu\text{g m}^{-3}$ (S) and (N)) yr^{-1}) and right panels – relative ($\%$ yr^{-1}) trends. The modelled trends are shown as coloured contour map (grey or white means non-significant trends) and the observed trends as coloured triangles (significant) and circles (non-significant).




Figure A10. Mean Sen's trend slopes for 2000-2010 simulated by the 6-model ensemble for natural aerosols: (a) sea salt (observed trends also shown), (b) BSOA and (c) mineral dust. The modelled trends are shown as coloured contour map (grey or white means non-significant trends) and the observed trends as coloured triangles (significant) and circles (non-significant).




Figure A11. Relative contributions of (from left to right) SO_4^{-2} , NH_4^+ , NO_3^- and ASOA to PM_{10} trends between 2000 and 2010 calculated by (from top to bottom) CHIMERE, EMEP, LOTOS-EUROS, MATCH, MINNI and Polair3D models.




Figure A13. Annual mean Sen's slope for trends in the period 1990-2010 as calculated by the 6-model ensemble (left) for PM₁₀ and (right) PM_{2.5}. Upper panels – absolute ($\mu\text{g m}^{-3} \text{yr}^{-1}$) and lower panels – relative to 1990 ($\% \text{yr}^{-1}$).




Figure A14. Modelled PM₁₀ trends calculated for European countries ($\mu\text{g m}^{-3} \text{yr}^{-1}$): a. the individual models for the period 1990-2010, and b. the model ensemble for the periods 1990-2000 and 2000-2010 separately. The countries are ranked according to descending 1990-2010 negative trends from the EMEP model (a).




Figure A15. The ratio of PM changes due to emission changes to those due to inter-annual meteorological variability for PM_{10} and $PM_{2.5}$ for 1990-2010 (left panels) and 1990-2000 (right panels) periods.

Figure A16.

(a) Brief description of the Chemistry-Transport Models involved in the Eurodelta-Trends modeling exercise (extended version of Table S1 in Supplementary Material to Colette et al. (2017a).)

MODEL	CHIMERE	EMEP	LOTOS-EUROS	MATCH	MINNI	POLYPHEMUS
version	Modified Chimere 2013	rV4.7	v1.10.005	VSOA April 2016	V4.7	V1.9.1
operator	INERIS	MET Norway	TNO	SMHI	ENEA/Arianet S.r.l.	CEREA
Chemistry/Meteorology coupling	offline	offline	offline	offline	offline	Offline
Vertical layers	9 sigma	20 sigma	5(4 dynamic layers and a surface layer)	39 hybrid eta utilizing the meteorological model layers	16 fixed terrain following layers	9 Fixed terrain following layers
Vertical extent	500 hPa	100 hPa	5000 m	ca 5000 m (4700 – 6000 m)	10000 m	12000m
Advection scheme	(van Leer, 1984)	(Bott, 1989)	(Walcek, 2000)	Fourth order mass conserved advection scheme based on (Bott, 1989)	Blackman cubic polynomials (Yamartino, 1993)	Third-order Direct Space Time scheme (Spee, 1998) with Koren-Sweby flux limiter function
Vertical diffusion	Kz approach following (Troen and Mahrt, 1986)	Kz approach following (O'Brien, 1970) and (Jeričević et al., 2010)	Kz approach Yamartino et al (2004)	Implicit mass conservative Kz approach, see (Robertson et al., 1999) Boundary layer parameterisation as detailed in (Robertson et al., 1999) forms the basis for vertical diffusion and dry deposition	Kz approach following (Lange, 1989)	Kz approach following (Troen and Mahrt, 1986)
Depth first layer	20 m	90 m	25 m	ca 60m	40 m	40m
Surface concentration	First model level	Downscaled to 3m using dry deposition velocity and similarity theory	Downscaled to 3m	Downscaled to 3m	First model level	First model level

Figure A16.

(b) Continuation of Main Features of CTMs involved in the EDT modelling.

Biogenic VOC	MEGAN model v2.1 with high resolution spatial and temporal LAI (Yuan et al., 2011) and recomputed emissions factors based on the land-use (Guenther et al., 2006)	Based upon maps of 115 species from (Koeble and Seufert, 2001), and hourly temperature and light using (Guenther et al., 1993; Guenther et al., 1994). See (Simpson et al., 1995; Simpson et al., 2012)	Based upon maps of 115 species from (Koeble and Seufert, 2001), and hourly temperature and light using (Guenther et al., 1993; Guenther et al., 1994). See (Simpson et al., 1995; Simpson et al., 2012)	Based upon maps of 115 species from (Koeble and Seufert, 2001), and hourly temperature and light using (Guenther et al., 1993; Guenther et al., 1994). See (Simpson et al., 1995; Simpson et al., 2012)	(Simpson et al., 2012), based on hourly temperature and light.	MEGAN v2.04 (Guenther et al., 2006)	MEGAN v2.04 (Guenther et al., 2006)	MEGAN v2.04 (Guenther et al., 2006)
Forest fires	None	None	None	None	None	None	None	None
Soil-NO	MEGAN model v2.04	See in (Simpson et al., 2012)	Not used here	Not used here	None	MEGAN v2.04	MEGAN v2.04	MEGAN v2.04
Lightning	None	Monthly climatological fields, (Köhler et al., 1995)	None	None	None	None	None	None
Dust traffic suspension	None	(Denier van der Gon et al., 2010)	None	None	Not used here	None	None	None
Land-use database	GLOBCOVER (24 classes)	CCE/SEI for Europe, elsewhere GLC2000	Corine Land Cover 2000 (13 classes)	Corine Land Cover 2000 (13 classes)	CCE/SEI for Europe	Corine Land Cover 2006 (22 classes)	Corine Land Cover 2006 (22 classes)	Global Land Cover 2000 (24 classes)
Ammonia compensation points	None	None, but zero NH3 deposition over growing crops	Only for NH3 (for stomatal, external leaf surface and soil (= 0))	Only for NH3 (for stomatal, external leaf surface and soil (= 0))	None	None	None	None
Sea salt	(Monahan, 1986)	(Monahan, 1986) and (Martensson et al., 2003), see (Tsyro et al., 2011)	(Martensson et al., 2003) and (Monahan, 1986), see (Schaap et al., 2009)	(Martensson et al., 2003) and (Monahan, 1986), see (Schaap et al., 2009)	Based on parameterization by (Sofiev et al. (2011))	(Zhang K.M. et al., 2005)	(Zhang K.M. et al., 2005)	(Monahan, 1986)
Windblown Dust	No windblown dust emissions within domain Boundary conditions from EMEP	Based on Marticorena and Bergametti (1995), Marticorena et al. (1997), Alfaro and Gomes (2001), Gomes et al. (2003), and Zender et al. (2003). Boundary conditions from EMEP	Based on Marticorena and Bergametti (1995), Gomes et al. (2003) and Alfaro et al. (2004). See Schaap et al. (2009) Boundary conditions from EMEP	Based on Marticorena and Bergametti (1995), Gomes et al. (2003) and Alfaro et al. (2004). See Schaap et al. (2009) Boundary conditions from EMEP	No windblown dust emissions within domain Boundary conditions from EMEP	Within domain based on Vautard et al. (2005) Boundary conditions from EMEP	Within domain based on Vautard et al. (2005) Boundary conditions from EMEP	No windblown dust emissions within domain Boundary conditions from EMEP

Figure A16.

(c) Continuation of Main Features of CTMs involved in the EDT modelling.

Dry deposition	Resistance approach (Emberson et al., 2000a; Emberson et al., 2000b)	Resistance approach for gasses (Venkatram and Pleim, 1999) for aerosols, (Simpson et al., 2012)	Resistance approach, DEPAC3.11 for gasses, (Van Zanthen et al., 2010) and (Zhang et al., 2001) for aerosols	Resistance approach depending on aerodynamic resistance, and land use (vegetation). Similar to (Andersson et al., 2007)	Resistance model based on (Wesely, 1989)	Resistance approach for gasses (Zhang et al., 2003) and aerosols (Zhang et al., 2001)
Stomatal resistance	(Emberson et al., 2000a; Emberson et al., 2000b)	DO3SE-EMEP: (Emberson et al., 2000a; Emberson et al., 2000b); Tuovinen et al., 2004; Simpson et al., 2012)	Emberson et al., (2000a); Emberson et al., (2000b)	Simple, seasonally varying, diurnal variation of surface resistance for gasses with stomatal resistance (similar to Andersson et al., 2007)	Wesely (1989)	Zhang et al. (2003)
Wet deposition gasses	In-cloud and sub-cloud scavenging coefficients	In-cloud and sub-cloud scavenging coefficients	sub-cloud scavenging coefficient	In-cloud scavenging of some species based on Henry's law constants. Simple in-cloud and sub-cloud scavenging coefficients for other gasses.	In-cloud and sub-cloud scavenging coefficients (EMEP, 2003)	In-cloud (monodispersed raindrops with constant collection efficiency) and below cloud (Sportisse and Dubois, 2002) scavenging coefficients
Wet deposition particles	In-cloud and sub-cloud scavenging	In-cloud and sub-cloud scavenging	sub-cloud scavenging coefficient	In-cloud and sub-cloud scavenging. Similar to (Simpson et al., 2012)	In-cloud and sub-cloud scavenging coefficients	In-cloud (as for gas) and below cloud (Slinn, 1983) scavenging coefficients
Gas phase chemistry	MELCHIOR2	EmChem09 (Simpson et al., 2012)	TNO-CBM-IV	Based on EMEP (Simpson et al., 2012), with modified isoprene chemistry (Carter, 1996; Langner et al., 1998)	SAPRC99 (Carter, 1996; Carter, 2000)	CB-05 (Varwood, G. et al., 2005)
Speciation of anthropogenic PM emissions	No split applied, PPM for all primary anthropogenic PM	EMEP split of PM _{2.5} and coarse PM emissions to EC, POA, rest PPM per country/SNAP sector using fractions provided by IIASA (see Simpson et al. 2012)	No split applied, PPM for all primary anthropogenic PM	Split of PM _{2.5} emissions to EC, POA, rest PPM per country/SNAP sector using fractions by Kuenen et al. (2014), Andersson et al., (2015). No split of coarse PM emissions	PM _{2.5} are split to EC, POA and rest PPM based on Kuenen et al. (2014) per country and SNAP sectors. No split for coarse PM.	PM _{2.5} are split between EC (~ 20%), POA (~45%) and rest PPM (~35%). No split for coarse PM. Detailed speciation are applied per SNAP sector. Based on Kuenen et al. (2014)

Figure A16.

(d) Continuation of Main Features of CTMs involved in the EDT modelling. The ratio of PM changes due to emission changes to those due to inter-annual meteorological variability for and for 1990–2010 (left panels) and 1990–2000 (right panels) periods.

Cloud chemistry	Aqueous SO ₂ chemistry to form SO ₄ ²⁻ by H ₂ O ₂ , O ₃ and O ₂ ; pH dependent	Aqueous SO ₂ chemistry to form SO ₄ ²⁻ by H ₂ O ₂ , O ₃ and O ₂ ; pH dependent (Simpson et al., 2012). (N ₂ O ₅ hydrolysis to form nitrate)	Aqueous SO ₂ chemistry, particle formation, pH dependent (Banzhaf et al., 2012).	Aqueous SO ₂ chemistry to form SO ₄ ²⁻ by H ₂ O ₂ , O ₃ and O ₂	Aqueous SO ₂ chemistry to form SO ₄ ²⁻ by H ₂ O ₂ , O ₃ and O ₂ (Seinfeld and Pandis, 1998)
Ammonium nitrate equilibrium	ISORROPIA v2.1 (Nenes et al., 1999)	MARS (Binkowski and Shankar, 1995)	ISORROPIA v.2	ISORROPIA v1.7 (Nenes et al., 1998)	ISORROPIA v1.7 (Nenes et al., 1999)
Coarse nitrate	HNO ₃ condensation on coarse sea salts computed as a function of pH, condensation kinetics and accounting for HCl evaporation. Local equilibrium using ISORROPIA in reverse mode.	Two formation rates of coarse NO ₃ from HNO ₃ , for relative humidity below/above 90%	Dynamic formation on Na (Wichink Kruit et al., 2012)	None	No heterogeneous nitrate formation
SOA formation	H ₂ O (Couvidat et al., 2012) mechanism coupled with the thermodynamic model SOAP (Couvidat and Sarretet, 2015)	VBS-NPAS (Simpson et al., 2012; Bergström et al., 2012)	Not used here	SORGAM module (Schell et al., 2001)	H ₂ O (Couvidat et al., 2012)
Aerosol model	9 bins (10 nm to 10 μm)	Bulk-approach (fine and coarse modes)	Bulk-approach (2 modes)	Bulk-approach	5 bins (0.01 to 10μm)
Aerosol physics	Coagulation/condensation/nucleation. Computation of the wet diameter for each bins as a function of humidity (used for coagulation, condensation, deposition)	Not used here	Not used here	Not used here	Coagulation/Condensation

¹⁾ EC - elemental carbon, POA - primary organic aerosol, ASOA and BSOA - anthropogenic and biogenic secondary aerosol, TPPM - total primary PM (EC+POA+rest PPM)

Author contributions. ACo coordinated the EURODELTA-Trends (EDT) exercise and WA was responsible for the compilation and quality control of the observations. The following modelling teams set-up, pre-processed, ran and post-processed the simulations for each model:
835 FC, BB, MGV and ACo for CHIM; ST and PW for EMEP; AM and MS for LOTO; CA and RB for MATCH; MM, MA, GB, ACa and MD for MINNI; YR and VR for POLR. WA and CK contributed with production of figures. Additional post-processing of model output and uploading to the AeroCom server was done by KC. All of the analyses presented in this paper were carried out by ST, assisted by discussions with WA and HF and with valuable contributions from ACo, CA, AM, GC, YR, BB, MT, MGV, NO, KM, FC, and MTP. Special thanks for language proofreading go to MT.

840 *Competing interests.* No competing interest are present.

Acknowledgements. The Ineris coordination of the EURODELTA-Trends exercise was supported by the French Ministry in charge of Ecology in the context of the Task Force on Measurement and Modelling of the EMEP program of the LRTAP Convention. Meteorological forcing with the WRF model was provided by Robert Vautard and Annemiek Stegehuis from LSCE/IPSL. The CHIMERE simulations were performed using the TGCC supercomputers under GENCI computing allocation. The participation of CIEMAT was financed by the Spanish
845 Ministry of Agriculture and Fishing, Food and Environment.

Computer time for EMEP model runs was supported by the Research Council of Norway through the NOTUR project EMEP (NN2890K) for CPU and the NorStore project European Monitoring and Evaluation Programme (NS9005K) for storage of data.

Funding for the MATCH participation was jointly divided between Nordforsk through the research programme Nordic Welfair (grant no 75007), the Swedish Environmental Protection Agency through the SCAC research programme and the 2017-2018 Belmont Forum and
850 BiodivERSa joint call for research proposals, under the BiodivScen ERA-Net COFUND programme, with the funding organisations AKA (contract no 326328), ANR (ANR-18-EBI4-007), BMBF (KFZ: 01LC1810A), FORMAS (contract no's: 2018-02434, 2018-02436, 2018-02437, 2018-02438) and MICINN (APCIN; PCI2018-093149).

LOTOS-team thanks Erik van Meijgaard of the Royal Netherlands Meteorological Institute (KNMI) for providing the RACMO2 simulations that were used by LOTOS-EUROS.

855 The computing resources and the related technical support used for MINNI simulations have been provided by CRESCO/ENEAGRID High Performance Computing infrastructure and its staff. The infrastructure is funded by ENEA, the Italian National Agency for New Technologies, Energy and Sustainable Economic Development and by Italian and European research programmes (<http://www.cresco.enea.it/english>, last access: 21 December 2018). MINNI participation to this project was supported by the “Cooperation Agreement for support to international Conventions, Protocols and related negotiations on air pollution issues”, funded by the Italian Ministry for the Environment,
860 Land and Sea.

CIEMAT acknowledges the Ministry for the Ecological Transition and Demographic Challenge (MITERD) for financial support. [Giancarlo Ciarelli was supported by ADEME and the Swiss National Science Foundation \(grant no. P2EZP2_175166\).](#)

The GAINS emission trends were produced as part of the FP7 European Research Project ECLIPSE (Evaluating the Climate and Air Quality Impacts of Short-Lived Pollutants) grant no. 282688.

Table A1. Selected set of EMEP monitoring stations for PM₁₀ and PM_{2.5} trend analysis for the period 2000-2010

Sites			PM ₁₀ measurements				PM _{2.5} measurements				
Code	Name	altitude	latitude	longitude	altitude	Sampler	Frequency	N + of F Missing years	Sampler	Frequency	N + of F Missing years
AT0002R	Illmitz	117.0m	47.767	16.767	117.0m	high vol	daily	++	high vol	daily	+2000
AT0005R	Vorhegg	1020.0m	46.678	12.972	1020.0m	high vol	daily	+2000	high vol	daily	11
CH0002R	Payerne	489.0m	46.813	6.945	489.0m	high vol	daily	++	high vol	daily	11
CH0003R	Tänikon	539.0m	47.480	8.905	539.0m	high vol	daily	++	high vol	daily	++
CH0004R	Chaumont	1137.0m	47.050	6.979	1137.0m	high vol	daily	++	high vol	daily	++
CH0005R	Rigi	1031.0m	47.068	8.464	1031.0m	high vol	daily	++	high vol	daily	++
CZ0003R	Kosetice (NOAK)	535.0m	49.573	15.080	535.0m	b-attenuation ¹	hourly	+2000	b-attenuation ¹	hourly	+2000
DE0001R	Westerland	12.0m	54.926	8.310	12.0m	high vol	daily	++	high vol	daily	++
DE0002R	Waldhof	74.0m	52.802	10.759	74.0m	high vol	daily	++	high vol	daily	++
DE0003R	Schauinsland	1205.0m	47.915	7.909	1205.0m	high vol	daily	++	high vol	daily	++
DE0007R	Neuglobbow	62.0m	53.167	13.033	62.0m	high vol	daily	11	high vol	daily	++
DE0008R	Schmitzke	937.0m	50.650	10.767	937.0m	high vol	daily	++	high vol	daily	++
DE0009R	Zingst	1.0m	54.437	12.725	1.0m	high vol	daily	++	high vol	daily	++
DE0044R	Melpitz	86.0m	51.530	12.934	86.0m	high vol	daily	++	high vol	daily	++
ES0007R	Viznar	1265.0m	37.233	-3.533	1265.0m	high vol	daily	+2000	high vol	daily	+2000
ES0008R	Niembro	134.0m	43.442	-4.850	134.0m	high vol	daily	+2000	high vol	daily	+2000
ES0011R	Barcarrota	393.0m	38.476	-6.923	393.0m	high vol	daily	+2000	high vol	daily	+2000
ES0012R	Zarra	885.0m	39.086	-1.102	885.0m	high vol	daily	+2000	high vol	daily	+2000
ES0013R	Penausende	985.0m	41.283	-5.867	985.0m	high vol	daily	+2000	high vol	daily	+2000
ES0014R	Els Torms	470.0m	41.400	0.717	470.0m	high vol	daily	+2000	high vol	daily	+2000
FI0050R	Hyytiälä	181.0m	61.850	24.283	181.0m	low vol ³	daily	++	low vol ³	daily	++
GB0006R	Lough Navar	126.0m	54.443	-7.870	126.0m	TEOM FDMS	daily	++	TEOM FDMS	hourly	++
GB0036R	Harwell	137.0m	51.573	-1.317	137.0m	TEOM FDMS	daily	++	TEOM FDMS	hourly	++
GB0043R	Narberth	160.0m	51.782	-4.691	160.0m	TEOM FDMS	daily	9-2003, 2004	TEOM FDMS	hourly	++
IT0004R	Ispra	209.0m	45.800	8.633	209.0m	low vol	weekly	11	low vol	daily	+2000
NO0002R	Birkenes II	219.0m	58.389	8.252	219.0m	b-attenuation ²	daily	++	low vol	weekly	+2000
SE0012R	Aspvreten	20.0m	58.800	17.383	20.0m	b-attenuation ²	daily	+2009	TEOM	hourly	+2010

¹ Thermo FH 62 L-R, ² OPSIS SM200, ³ with 4 stage cascade impactor

Table A2. Relative bias (in %) and correlation (R) for modelled PM₁₀ with respect to available observations at 26 EDT sites for the years 2000 to 2010

Year	Nsite	CHIM		EMEP		LOTO		MATCH		MINNI		POLR*	
		Bias	CorrR	Bias	CorrR	Bias	CorrR	Bias	CorrR	Bias	CorrR	Bias	CorrR
2000	15	-2-2.0	0.64	22.1	0.58	-6-6.3	0.47	-11	0.60	-4-4.3	0.58	-24	0.62
2001	21	-4-4.4	0.41	-8-8.2	0.61	-4-4.0	0.60	-9-9.4	0.71	-8-8.3	0.48	-30	0.59
2002	22	-7-7.5	0.55	-14	0.60	-13	0.46	-16	0.63	-14	0.61	-35	0.60
2003	22	-8-8.5	0.59	-13	0.63	-16	0.40	-12	0.55	-12	0.63	-36	0.64
2004	22	-7-7.3	0.50	-14	0.64	-10-9.7	0.66	-14	0.78	-9-8.9	0.60	-33	0.64
2005	23	-7-7.2	0.55	-11	0.63	-5-5.3	0.51	-12	0.65	-7-7.0	0.62	-31	0.65
2006	21	-6-6.3	0.48	-12	0.48	44.3	0.24	-8-7.6	0.34	-10	0.52	-32	0.42
2007	21	-3-2.6	0.39	-10	0.50	-11	0.43	-10-9.8	0.61	-7-6.7	0.48	-28	0.50
2008	21	-4-3.7	0.37	-10	0.49	-8-8.23	0.44	-11	0.63	-8-7.5	0.48	-28	0.53
2009	22	0-0.4	0.53	-8-8.0	0.61	22.4	0.39	-5-5.1	0.50	-3-3.5	0.59	-27	0.57
2010	21	-9-9.4	0.58	-16	0.62	-15	0.23	-18	0.46	-17	0.60	-35	0.53

Bias - relative bias expressed in %;

* Excluding coarse sea salt

Table A3. Relative bias (in %) and correlation (R) for modelled PM_{2.5} with respect to available observations at 26-13 EDT sites for the years 2000 to 2010

Year	Nsite	CHIM		EMEP		LOTO		MATCH		MINNI		POLR	
		Bias	CorrR	Bias	CorrR	Bias	CorrR	Bias	CorrR	Bias	CorrR	Bias	CorrR
2000	75	-2-2.4	0.71	-5-5.3	0.63	-6-6.4	0.69	-3-3.1	0.64	66.1	0.71	-2-1.7	0.69
2001	12	-18	0.59	-21	0.61	-18	0.79	-18	0.70	-10	0.53	-22	0.73
2002	12	-19	0.64	-25	0.68	-18	0.73	-21	0.70	-12	0.60	-26	0.72
2003	12	-18	0.74	-20	0.72	-25	0.70	-16	0.69	-6-6.3	0.70	-25	0.74
2004	12	-17	0.61	-23	0.62	-15	0.77	-16	0.73	-6-6.2	0.54	-23	0.72
2005	13	-21	0.68	-24	0.63	-15	0.73	-17	0.69	-9-8.7	0.59	-22	0.74
2006	11	-20	0.69	-24	0.50	-12	0.65	-16	0.55	-9-9.1	0.45	-25	0.61
2007	11	-18	0.59	-25	0.52	-18	0.63	-16	0.73	-9-8.7	0.40	-22	0.69
2008	11	-11	0.60	-17	0.60	-7-6.6	0.71	-6-6.3	0.69	+1.1	0.54	-12	0.71
2009	11	-7-7.3	0.65	-14	0.61	+0.9	0.76	-4-3.8	0.64	66.1	0.57	-12	0.67
2010	10	-17	0.62	-24	0.58	-14	0.68	-14	0.66	-10	0.64	-19	0.67

Bias - relative bias expressed in %

865 References

- Transboundary acidification, eutrophication and ground level ozone in Europe: Unified EMEP model description, Tech. rep., Norwegian Meteorological Institute, 2003.
- Air quality in Europe — 2020 report, EEA Report No 09/2020, ISSN 1977-8449, European Environment Agency, Copenhagen, <https://doi.org/10.2800/786656>, 2009.
- 870 Aas, W., Mortier, A., Bowersox, V., Cherian, R., Faluvegi, G., Fagerli, H., Hand, J., Klimont, Z., Galy-Lacaux, C., Lehmann, C. M., et al.: Global and regional trends of atmospheric sulfur, *Scientific reports*, 9, 1–11, 2019.
- Aas, W., Fagerli, H., Yttri, K. E., Tsyro, S., Solberg, S., Simpson, D., Gliß, J., Mortier, A., Grøtting Wærsted, E., Brenna, H., Hjellbrekke, A., Griesfeller, J., Nyíri, A., Gauss, M., and Scheuschner, T.: Trends in observations and EMEP MSC-W model calculations 2000-2019, in: *Transboundary particulate matter, photo-oxidants, acidifying and eutrophying components. EMEP Status Report 1/2021*, pp. 83–97, 875 The Norwegian Meteorological Institute, Oslo, Norway, 2021.
- Alfaro, S. C. and Gomes, L.: Modeling mineral aerosol production by wind erosion: Emission intensities and aerosol size distributions in source areas, *Journal of Geophysical Research: Atmospheres*, 106, 18 075–18 084, 2001.
- Amann, M.: Future emissions of air pollutants in Europe-Current legislation baseline and the scope for further reductions, <http://pure.iiasa.ac.at/id/eprint/10164/1/XO-12-011.pdf>, 2012.
- 880 Amann, M., Bertok, I., Borken-Kleefeld, J., Cofala, J., Heyes, C., Höglund-Isaksson, L., Klimont, Z., Nguyen, B., Posch, M., Rafaj, P., et al.: Cost-effective control of air quality and greenhouse gases in Europe: Modeling and policy applications, *Environmental Modelling & Software*, 26, 1489–1501, 2011.
- Andersson, C., Langner, J., and Bergström, R.: Interannual variation and trends in air pollution over Europe due to climate variability during 1958–2001 simulated with a regional CTM coupled to the ERA40 reanalysis, *Tellus B: Chemical and Physical Meteorology*, 59, 77–98, 885 2007.
- Andersson, C., Bergström, R., Bennet, C., Robertson, L., Thomas, M., Korhonen, H., Lehtinen, K., and Kokkola, H.: MATCH-SALSA—Multi-scale Atmospheric Transport and CHEMistry model coupled to the SALSA aerosol microphysics model—Part 1: Model description and evaluation, *Geoscientific Model Development*, 8, 171–189, 2015.
- Banzhaf, S., Schaap, M., Kranenburg, R., Manders, A., Segers, A., Visschedijk, A., Denier van der Gon, H., Kuenen, J., Van Meijgaard, E., 890 Van Ulft, L., et al.: Dynamic model evaluation for secondary inorganic aerosol and its precursors over Europe between 1990 and 2009, *Geoscientific Model Development*, 8, 1047–1070, 2015.
- Barmadimos, I., Keller, J., Oderbolz, D., Hueglin, C., and Prévôt, A. S. H.: One decade of parallel fine ($PM_{2.5}$) and coarse ($PM_{10-PM_{2.5}}$) particulate matter measurements in Europe: trends and variability, *Atmospheric Chemistry and Physics*, 12, 3189–3203, <https://doi.org/10.5194/acp-12-3189-2012>, <https://www.atmos-chem-phys.net/12/3189/2012/>, 2012.
- 895 Beltman, J. B., Hendriks, C., Tum, M., and Schaap, M.: The impact of large scale biomass production on ozone air pollution in Europe, *Atmospheric Environment*, 71, 352–363, 2013.
- Bergström, R., Denier Van Der Gon, H., Prévôt, A. S., Yttri, K. E., and Simpson, D.: Modelling of organic aerosols over Europe (2002–2007) using a volatility basis set (VBS) framework: application of different assumptions regarding the formation of secondary organic aerosol, *Atmospheric Chemistry and Physics*, 12, 8499–8527, 2012.

- 900 Bergström, R., Hallquist, M., Simpson, D., Wildt, J., and Mentel, T. F.: Biotic stress: a significant contributor to organic aerosol in Europe?, *Atmospheric Chemistry and Physics*, 14, 13 643–13 660, <https://doi.org/10.5194/acp-14-13643-2014>, <http://www.atmos-chem-phys.net/14/13643/2014/>, 2014.
- Bessagnet, B., Pirovano, G., Mircea, M., Cuvelier, C., Aulinger, A., Calori, G., Ciarelli, G., Manders, A., Stern, R., Tsyro, S., and et al.: Presentation of the EURODELTA III intercomparison exercise – evaluation of the chemistry transport models’ performance on criteria
905 pollutants and joint analysis with meteorology, *Atmospheric Chemistry and Physics*, 16, 12 667–12 701, <https://doi.org/10.5194/acp-16-12667-2016>, 2016.
- Bieser, J., Aulinger, A., Matthias, V., Quante, M., and Van Der Gon, H. D.: Vertical emission profiles for Europe based on plume rise calculations, *Environmental Pollution*, 159, 2935–2946, 2011.
- Binkowski, F. S. and Shankar, U.: The regional particulate matter model: 1. Model description and preliminary results, *Journal of Geophysical
910 Research: Atmospheres*, 100, 26 191–26 209, 1995.
- Bott, A.: A positive definite advection scheme obtained by nonlinear renormalization of the advective fluxes, *Monthly weather review*, 117, 1006–1016, 1989.
- Carter, W. P.: Condensed atmospheric photooxidation mechanisms for isoprene, *Atmospheric Environment*, 30, 4275–4290, 1996.
- Carter, W. P.: Documentation of the SAPRC-99 chemical mechanism for VOC reactivity assessment, Contract, 92, 95–308, 2000.
- 915 Ciarelli, G., Theobald, M. R., Vivanco, M. G., Beekmann, M., Aas, W., Andersson, C., Bergström, R., Manders-Groot, A., Couvidat, F., Mircea, M., et al.: Trends of inorganic and organic aerosols and precursor gases in Europe: insights from the EURODELTA multi-model experiment over the 1990–2010 period, *Geoscientific Model Development*, 12, 4923–4954, 2019.
- Colette, A., Granier, C., Hodnebrog, Ø., Jacobs, H., Maurizi, A., Nyíri, A., Bessagnet, B., D’Angiola, A., D’Isidoro, M., Gauss, M., Meleux, F., Memmesheimer, M., Mieville, A., Rouïl, L., Russo, F., Solberg, S., Stordal, F., and Tampieri, F.: Air quality trends in Europe over
920 the past decade: a first multi-model assessment, *Atmos. Chem. Phys.*, 11, 11 657–11 678, <https://doi.org/10.5194/acp-11-11657-2011>, <https://www.atmos-chem-phys.net/11/11657/2011/>, 2011.
- Colette, A., Aas, W., Banin, L., Braban, C., Ferm, M., González Ortiz, A., Ilyin, I., Mar, K., Pandolfi, M., Putaud, J.-P., Shatalov, V., Solberg, S., Spindler, G., Tarasova, O., Vana, M., Adani, M., Almodovar, P., Berton, E., Bessagnet, B., Bohlin-Nizzetto, P., Boruvkova, J., Breivik, K., Briganti, G., Cappelletti, A., Cuvelier, K., Derwent, R., D’Isidoro, M., Fagerli, H., Funk, C., Garcia Vivanco, M., González Ortiz, A., Haeuber, R., Hueglin, C., Jenkins, S., Kerr, J., de Leeuw, F., Lynch, J., Manders, A., Mircea, M., Pay, M., Pritula, D., Putaud, J.-P.,
925 Querol, X., Raffort, V., Reiss, I., Roustan, Y., Sauvage, S., Scavo, K., Simpson, D., Smith, R., Tang, Y., Theobald, M., Tørseth, K., Tsyro, S., van Pul, A., Vidic, S., Wallasch, M., and Wind, P.: Air Pollution trends in the EMEP region between 1990 and 2012., Tech. Rep. Joint Report of the EMEP Task Force on Measurements and Modelling (TFMM), Chemical Co-ordinating Centre (CCC), Meteorological Synthesizing Centre-East (MSC-E), Meteorological Synthesizing Centre-West (MSC-W) EMEP/CCC Report 1/2016, Norwegian Institute
930 for Air Research, Kjeller, Norway, http://www.unece.org/fileadmin/DAM/env/documents/2016/AIR/Publications/Air_pollution_trends_in_the_EMEP_region.pdf, 2016.
- Colette, A., Andersson, C., Manders, A., Mar, K., Mircea, M., Pay, M.-T., Raffort, V., Tsyro, S., Cuvelier, C., Adani, M., Bessagnet, B., Bergström, R., Briganti, G., Butler, T., Cappelletti, A., Couvidat, F., D’Isidoro, M., Doumbia, T., Fagerli, H., Granier, C., Heyes, C., Klimont, Z., Ojha, N., Otero, N., Schaap, M., Sindelarova, K., Stegehuis, A. I., Roustan, Y., Vautard, R., van Meijgaard, E., Vivanco, M. G.,
935 and Wind, P.: EURODELTA-Trends, a multi-model experiment of air quality hindcast in Europe over 1990–2010, *Geoscientific Model Development*, 10, 3255–3276, <https://doi.org/10.5194/gmd-10-3255-2017>, <https://www.geosci-model-dev.net/10/3255/2017/>, 2017a.

- Colette, A., Solberg, S., Beauchamp, M., Bessagnet, B., Malherbe, L., Guerreiro, C., and Team, E.-T. M.: Long term air quality trends in Europe. Contribution of meteorological variability, natural factors and emissions, Long term air quality trends in Europe Contribution of meteorological variability, natural factors and emissions. ETC/ACM Technical Paper 2016/7, European Topic Centre on Air Pollution and Climate Change Mitigation (ETC/A, Bilthoven, The Netherlands, https://www.eionet.europa.eu/etcs/etc-atni/products/etc-atni-reports/etcacm_tp_2016_7_aqtrendseurope, 2017b.
- 940 Colette, A., Solberg, S., Aas, W., and Walker, S.-E.: Understanding Air Quality Trends in Europe, Eionet Report - ETC/ATNI 2020/8, <https://www.eionet.europa.eu/etcs/etc-atni>, 2021.
- Couvidat, F. and Sartelet, K.: The Secondary Organic Aerosol Processor (SOAP v1. 0) model: a unified model with different ranges of complexity based on the molecular surrogate approach, *Geoscientific Model Development*, 8, 1111–1138, 2015.
- 945 Couvidat, F., Debry, E., Sartelet, K., and Seigneur, C.: A hydrophilic/hydrophobic organic (H₂O) aerosol model: Development, evaluation and sensitivity analysis, *Journal of Geophysical Research: Atmospheres*, 117, 2012.
- Crippa, M., Janssens-Maenhout, G., Dentener, F., Guizzardi, D., Sindelarova, K., Muntean, M., Van Dingenen, R., and Granier, C.: Forty years of improvements in European air quality: regional policy-industry interactions with global impacts, *Atmospheric Chemistry and Physics*, 16, 3825–3841, <https://doi.org/10.5194/acp-16-3825-2016>, <https://www.atmos-chem-phys.net/16/3825/2016/>, 2016.
- 950 Cusack, M., Alastuey, A., Pérez, N., Pey, J., and Querol, X.: Trends of particulate matter (PM_{2.5}) and chemical composition at a regional background site in the Western Mediterranean over the last nine years (2002–2010), *Atmospheric Chemistry and Physics*, 12, 8341–8357, <https://doi.org/10.5194/acp-12-8341-2012>, <https://www.atmos-chem-phys.net/12/8341/2012/>, 2012.
- Dahlgren, P., Landelius, T., Kållberg, P., and Gollvik, S.: A high-resolution regional reanalysis for Europe. Part 1: Three-dimensional reanalysis with the regional High-Resolution Limited-Area Model (HIRLAM), *Quarterly Journal of the Royal Meteorological Society*, 142, 2119–2131, 2016.
- 955 Dee, D. P., Uppala, S. M., Simmons, A., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M., Balsamo, G., Bauer, d. P., et al.: The ERA-Interim reanalysis: Configuration and performance of the data assimilation system, *Quarterly Journal of the royal meteorological society*, 137, 553–597, 2011.
- 960 Denier van der Gon, H., Jozwicka, M., Hendriks, E., Gondwe, M., and Schaap, M.: Mineral dust as a component of particulate matter, Delft, the Netherlands, 2010.
- Denier van der Gon, H., Hendriks, C., Kuenen, J., Segers, A., and Visschedijk, A.: Description of current temporal emission patterns and sensitivity of predicted AQ for temporal emission patterns. EU FP7 MACC deliverable report D_D – *EMIS_{1.3}, Tech.rep., TNO*, 2011.
- Denier van der Gon, H. A. C., Bergström, R., Fountoukis, C., Johansson, C., Pandis, S. N., Simpson, D., and Visschedijk, A. J. H.: Particulate emissions from residential wood combustion in Europe - revised estimates and an evaluation, *Atmos. Chem. Physics*, 15, 6503–6519, <https://doi.org/10.5194/acp-15-6503-2015>, <http://www.atmos-chem-phys.net/15/6503/2015/>, 2015.
- EEA: Emissions of primary particles and secondary particulate matter precursors. Last modified 23 Feb 2018, Tech. rep., European Environment Agency, <https://www.eea.europa.eu/data-and-maps/indicators/emissions-of-primary-particles-and-1>, 2008.
- Emberson, L., Ashmore, M., Cambridge, H., Simpson, D., and Tuovinen, J.-P.: Modelling stomatal ozone flux across Europe, *Environmental Pollution*, 109, 403–413, 2000a.
- 970 Emberson, L., Simpson, D., Tuovinen, J., Ashmore, M., and Cambridge, H.: Towards a model of ozone deposition and stomatal uptake over Europe, MSC-W, 2000b.

- EMEP: Transboundary particulate matter, photo-oxidants, acidifying and eutrophying components, Tech. Rep. EMEP Status Report 1/2012, The Norwegian Meteorological Institute, Oslo, Norway, https://emep.int/publ/common_publications.html#975 2012, 2012.
- Fagerli, H. and Aas, W.: Trends of nitrogen in air and precipitation: Model results and observations at EMEP sites in Europe, 1980–2003, *Environmental Pollution*, 154, 448–461, 2008.
- Gomes, L., Rajot, J., Alfaro, S., and Gaudichet, A.: Validation of a dust production model from measurements performed in semi-arid agricultural areas of Spain and Niger, *Catena*, 52, 257–271, 2003.
- 980 Guenther, A., Zimmerman, P., and Wildermuth, M.: Natural volatile organic compound emission rate estimates for US woodland landscapes, *Atmospheric Environment*, 28, 1197–1210, 1994.
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmospheric Chemistry and Physics*, 6, 3181–3210, 2006.
- 985 Guenther, A. B., Monson, R. K., and Fall, R.: Isoprene and monoterpene emission rate variability: observations with eucalyptus and emission rate algorithm development, *Journal of Geophysical Research: Atmospheres*, 96, 10 799–10 808, 1991.
- Guenther, A. B., Zimmerman, P. R., Harley, P. C., Monson, R. K., and Fall, R.: Isoprene and monoterpene emission rate variability: model evaluations and sensitivity analyses, *Journal of Geophysical Research: Atmospheres*, 98, 12 609–12 617, 1993.
- 990 Guerreiro, C. B., Foltescu, V., and De Leeuw, F.: Air quality status and trends in Europe, *Atmospheric environment*, 98, 376–384, 2014.
- Jacob, D., Petersen, J., Eggert, B., Alias, A., Christensen, O. B., Bouwer, L. M., Braun, A., Colette, A., Déqué, M., Georgievski, G., et al.: EURO-CORDEX: new high-resolution climate change projections for European impact research, *Regional Environmental Change*, 14, 563–578, 2014.
- 995 Jathar, S. H., Gordon, T. D., Hennigan, C. J., Pye, H. O. T., Pouliot, G., Adams, P. J., Donahue, N. M., and Robinson, A. L.: Unspeciated organic emissions from combustion sources and their influence on the secondary organic aerosol budget in the United States, *Proceedings of the National Academy of Sciences of the United States of America*, 111, 10 473–10 478, <https://doi.org/10.1073/pnas.1323740111>, 2014.
- 1000 Jeričević, A., Kraljević, L., Grisogono, B., Fagerli, H., and Večenaj, Ž.: Parameterization of vertical diffusion and the atmospheric boundary layer height determination in the EMEP model, *Atmospheric chemistry and physics*, 10, 341–364, 2010.
- Kendall, M.: Rank correlation methods (4th edn.) Charles Griffin, San Francisco, CA, 8, 1975.
- Klimont, Z., Höglund-Isaksson, L., Heyes, C., Rafaj, P., Schoepp, W., Cofala, J., Purohit, P., Borcken-Kleefeld, J., Kupiainen, K., Kiesewetter, G., et al.: Global scenarios of air pollutants and methane: 1990–2050, 2016.

- Klimont, Z., Kupiainen, K., Heyes, C., Purohit, P., Cofala, J., Rafaj, P., Borken-Kleefeld, J., and Schöpp, W.: Global anthropogenic emissions of particulate matter including black carbon, *Atmospheric Chemistry and Physics*, 17, 8681–8723, <https://doi.org/10.5194/acp-17-8681-2017>, <https://www.atmos-chem-phys.net/17/8681/2017/>, 2017.
- Köble, R. and Seufert, G.: Novel maps for forest tree species in Europe, in: Proceedings of the 8th European symposium on the physico-chemical behaviour of air pollutants: “a changing atmosphere”, pp. 17–20, 2001.
- Köhler, I., Sausen, R., and Klenner, G.: NO_x production from lightning, the impact of NO_x emissions from aircraft upon the atmosphere at flight altitudes 8–15 km (AERONOX), edited by U. Schumann, final report to the Comm. of the Eur. Commun., Dtsch. Luft und Raumfahrt, Oberpfaffenhofen, Germany, 1995.
- Kruit, R. W. and van der Swaluw, E.: Improving the understanding of the secondary inorganic aerosol distribution over the Netherlands, Tech. rep., TNO report TNO-060-UT-2012, 2012.
- Kuenen, J., Visschedijk, A., Jozwicka, M., and Denier Van Der Gon, H.: TNO-MACC_II emission inventory; a multi-year (2003–2009) consistent high-resolution European emission inventory for air quality modelling, *Atmospheric Chemistry and Physics*, 14, 10963–10976, 2014.
- Lange, R.: Transferability of a three-dimensional air quality model between two different sites in complex terrain, *Journal of Applied Meteorology and Climatology*, 28, 665–679, 1989.
- Langner, J., Bergström, R., and Plejfel, K.: European scale modeling of sulfur, oxidized nitrogen and photochemical oxidants: Model development and evaluation for the 1994 growing season, SMHI, 1998.
- Langner, J., Engardt, M., Baklanov, A., Christensen, J. H., Gauss, M., Geels, C., Hedegaard, G. B., Nuterman, R., Simpson, D., Soares, J., Sofiev, M., Wind, P., and Zakey, A.: A multi-model study of impacts of climate change on surface ozone in Europe, *Atmos. Chem. Phys.*, 12, 10423–10440, <https://doi.org/10.5194/acp-12-10423-2012>, <http://www.atmos-chem-phys.net/12/10423/2012/>, 2012.
- Mann, H.: Non-Parametric Tests against Trend. *Econometrica*, 13, 245-259, Mantua, NJ, SR Hare, Y. Zhang, JM Wallace, and RC Francis (1997), A Pacific decadal, 1945.
- Martensson, E., Nilsson, E., de Leeuw, G., Cohen, L., and Hansson, H.-C.: Laboratory simulations and parameterization of the primary marine aerosol production, *Journal of geophysical research*, 108, AAC15–1, 2003.
- Martcorena, B. and Bergametti, G.: Modeling the atmospheric dust cycle: 1. Design of a soil-derived dust emission scheme, *Journal of geophysical research: atmospheres*, 100, 16415–16430, 1995.
- Martcorena, B., Bergametti, G., Aumont, B., Callot, Y., N'Doumé, C., and Legrand, M.: Modeling the atmospheric dust cycle: 2. Simulation of Saharan dust sources, *Journal of Geophysical Research: Atmospheres*, 102, 4387–4404, 1997.
- Messina, P., Lathiere, J., Sindelarova, K., Vuichard, N., Granier, C., Ghattas, J., Cozic, A., and Hauglustaine, D. A.: Global biogenic volatile organic compound emissions in the ORCHIDEE and MEGAN models and sensitivity to key parameters, *Atmos. Chem. Phys.*, 16, 14169–14202, <https://doi.org/10.5194/acp-16-14169-2016>, 2016.

- Monahan, E. C.: The ocean as a source for atmospheric particles, in: *The role of air-sea exchange in geochemical cycling*, pp. 129–163, Springer, 1986.
- Mortier, A., Gliß, J., Schulz, M., Aas, W., Andrews, E., Bian, H., Chin, M., Ginoux, P., Hand, J., Holben, B., et al.: Evaluation of climate model aerosol trends with ground-based observations over the last 2 decades—an AeroCom and CMIP6 analysis, *Atmospheric Chemistry and Physics*, 20, 13 355–13 378, 2020.
- Mozurkewich, M.: The dissociation constant of ammonium nitrate and its dependence on temperature, relative humidity and particle size, *Atmospheric Environment. Part A. General Topics*, 27, 261–270, 1993.
- Myhre, G., Aas, W., Cherian, R., Collins, W., Faluvegi, G., Flanner, M., Forster, P., Hodnebrog, Ø., Klimont, Z., Lund, M. T., Mülmenstädt, J., Lund Myhre, C., Olivie, D., Prather, M., Quaas, J., Samset, B. H., Schnell, J. L., Schulz, M., Shindell, D., Skeie, R. B., Takemura, T., and Tsyro, S.: Multi-model simulations of aerosol and ozone radiative forcing due to anthropogenic emission changes during the period 1990–2015, *Atmospheric Chemistry and Physics*, 17, 2709–2720, <https://doi.org/10.5194/acp-17-2709-2017>, <https://www.atmos-chem-phys.net/17/2709/2017/>, 2017.
- Nenes, A., Pandis, S. N., and Pilinis, C.: ISORROPIA: A new thermodynamic equilibrium model for multiphase multicomponent inorganic aerosols, *Aquatic geochemistry*, 4, 123–152, 1998.
- Nenes, A., Pandis, S. N., and Pilinis, C.: Continued development and testing of a new thermodynamic aerosol module for urban and regional air quality models, *Atmospheric Environment*, 33, 1553–1560, 1999.
- Ots, R., Young, D. E., Vieno, M., Xu, L., Dunmore, R. E., Allan, J. D., Coe, H., Williams, L. R., Herndon, S. C., Ng, N. L., Hamilton, J. F., Bergström, R., Di Marco, C., Nemitz, E., Mackenzie, I. A., Kuenen, J. J. P., Green, D. C., Reis, S., and Heal, M. R.: Simulating secondary organic aerosol from missing diesel-related intermediate-volatility organic compound emissions during the Clean Air for London (ClearfLo) campaign, *Atmos. Chem. Phys.*, 16, 6453–6473, <https://doi.org/10.5194/acp-16-6453-2016>, <http://www.atmos-chem-phys.net/16/6453/2016/>, 2016.
- O’Brien, J. J.: A note on the vertical structure of the eddy exchange coefficient in the planetary boundary layer, *J. atmos. Sci.*, 27, 1213–1215, 1970.
- Platt, S. M., El Haddad, I., Pieber, S. M., Zardini, A. A., Suarez-Bertoa, R., Clairotte, M., Daellenbach, K. R., Huang, R. J., Slowik, J. G., Hellebust, S., Temime-Roussel, B., Marchand, N., de Gouw, J., Jimenez, J. L., Hayes, P. L., Robinson, A. L., Baltensperger, U., Astorga, C., and Prevot, A. S. H.: Gasoline cars produce more carbonaceous particulate matter than modern filter-equipped diesel cars, *SCIENTIFIC REPORTS*, 7, <https://doi.org/10.1038/s41598-017-03714-9>, 2017.
- Robertson, L., Langner, J., and Engardt, M.: An Eulerian limited-area atmospheric transport model, *Journal of Applied Meteorology and Climatology*, 38, 190–210, 1999.
- Schaap, M., Manders, M., Hendriks, E., Cnossen, J., Segers, A., van der Gon, H. D., Jozwicka, M., Sauter, F., Velders, G., Mathijssen, J., et al.: Regional modelling of particulate matter for the Netherlands, Tech. rep., Technical Report BOP, 2008.
- Schell, B., Ackermann, I. J., Hass, H., Binkowski, F. S., and Ebel, A.: Modeling the formation of secondary organic aerosol within a comprehensive air quality model system, *Journal of Geophysical Research: Atmospheres*, 106, 28 275–28 293, 2001.

- Schöpp, W., Klimont, Z., Suutari, R., and Cofala, J.: Uncertainty analysis of emission estimates in the RAINS integrated assessment model, *Environmental Science & Policy*, 8, 601–613, 2005.
- Seinfeld John, H., Spyros, N., et al.: *Atmospheric chemistry and physics: from air pollution to climate change*, 1997.
- Simpson, D. and Denier van der Gon, H.: Problematic emissions - particles or gases?, in: *Transboundary particulate matter, photo-oxidants, acidifying and eutrophying components*. EMEP Status Report 1/2015, pp. 87–96, The Norwegian Meteorological Institute, Oslo, Norway, 2015.
- 1075 Simpson, D., Guenther, A., Hewitt, C. N., and Steinbrecher, R.: Biogenic emissions in Europe: 1. Estimates and uncertainties, *Journal of Geophysical Research: Atmospheres*, 100, 22 875–22 890, 1995.
- Simpson, D., Winiwarter, W., Börjesson, G., Cinderby, S., Ferreira, A., Guenther, A., Hewitt, C. N., Janson, R., Khalil, M. A. K., Owen, S., Pierce, T. E., Puxbaum, H., Shearer, M., Skiba, U., Steinbrecher, R., Tarrasón, L., and Öquist, M. G.: Inventorying emissions from Nature in Europe, *J. Geophys. Res.*, 104, 8113–8152, 1999.
- 1080 Simpson, D., Benedictow, A., Berge, H., Bergström, R., Emberson, L., Fagerli, H., Flechard, C., Hayman, G., Gauss, M., Jonson, J., Jenkin, M., Nyíri, A., Richter, C., Semeena, V., Tsyro, S., Tuovinen, J.-P., Valdebenito, A., and Wind, P.: The EMEP MSC-W chemical transport model – technical description, *acp*, 12, 7825–7865, <https://doi.org/10.5194/acp-12-7825-2012>, 2012.
- Simpson, D., Fagerli, H., Colette, A., van der Gon, H. D., Dore, C., Hallquist, M., Hansson, H. C., Maas, R., and Rouil, L. e. a.:
1085 How should condensables be included in PM emission inventories reported to EMEP/CLRTAP?, Report of the expert workshop on condensable organics organised by MSC-W, Gothenburg, 17-19th March 2020. EMEP/MSW Technical Report 4/2020, The Norwegian Meteorological Institute, Oslo, Norway, https://emep.int/publ/reports/2020/emep_mscw_technical_report_4_2020.pdf, 2020.
- Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Wang, W., and Powers, J. G.: A description of the advanced research WRF version 2, Tech. rep., National Center For Atmospheric Research Boulder Co Mesoscale and
1090 Microscale . . . , 2005.
- Slinn, W.: Precipitation scavenging, in *atmospheric sciences and power production-1979*, Division of Biomedical Environmental Research, US Department of Energy, Washington DC, 1983.
- Sofiev, M., Soares, J., Prank, M., deLeeuw, G., and Kukkonen, J.: A regional to global model of emission and transport of sea
1095 salt particles 20 in the atmosphere, *Journal of Geophysical Research: Atmospheres*, 116, 2011.
- Solberg, S., Jonson, J., Horalek, J., Larssen, S., and De Leeuw, F.: Assessment of ground-level ozone in EEA member countries, with a focus on long-term trends, EEA Report No7/2009, European Environment Agency, Copenhagen, 2009.
- Spee, E. J. et al.: *Numerical methods in global transport-chemistry models*, AmsterdamUvA, 1998.
- Sportisse, B. and Du Bois, L.: Numerical and theoretical investigation of a simplified model for the parameterization of below-
1100 cloud scavenging by falling raindrops, *Atmospheric Environment*, 36, 5719–5727, 2002.

- Spracklen, D. V., Arnold, S. R., Sciare, J., Carslaw, K. S., and Pio, C.: Globally significant oceanic source of organic carbon aerosol, *Geophys. Res. Lett.*, 35, <https://doi.org/10.1029/2008GL033359>, 2008.
- Stegehuis, A., Vautard, R., Ciais, P., Teuling, A., Miralles, D., and Wild, M.: An observation-constrained multi-physics WRF ensemble for simulating European mega heat waves, *Geoscientific Model Development*, 8, 2285–2298, 2015.
- 1105 Strand, A. and Hov, Ø.: A two-dimensional global study of tropospheric ozone production, *Journal of Geophysical Research: Atmospheres*, 99, 22 877–22 895, 1994.
- Terrenoire, E., Bassagnet, B., Rouil, L., Tognet, F., Pirovano, G., Letinois, L., Beauchamp, M., Colette, A., Thunis, P., Amann, M., et al.: High-resolution air quality simulation over Europe with the chemistry transport model CHIMERE, *Geoscientific Model Development*, 8, 21–42, 2015.
- 1110 Theobald, M. R., Vivanco, M. G., Aas, W., Andersson, C., Ciarelli, G., Couvidat, F., Cuvelier, K., Manders, A., Mircea, M., Pay, M.-T., Tsyro, S., Adani, M., Bergström, R., Bessagnet, B., Briganti, G., Cappelletti, A., D’Isidoro, M., Fagerli, H., Mar, K., Otero, N., Raffort, V., Roustan, Y., Schaap, M., Wind, P., and Colette, A.: An evaluation of European nitrogen and sulfur wet deposition and their trends estimated by six chemistry transport models for the period 1990–2010, *Atmospheric Chemistry and Physics*, 19, 379–405, <https://doi.org/10.5194/acp-19-379-2019>, <https://www.atmos-chem-phys.net/19/379/2019/>, 2019.
- 1115 Tørseth, K., Aas, W., Breivik, K., Fjæraa, A. M., Fiebig, M., Hjellbrekke, A.-G., Lund Myhre, C., Solberg, S., and Yttri, K. E.: Introduction to the European Monitoring and Evaluation Programme (EMEP) and observed atmospheric composition change during 1972–2009, *Atmospheric Chemistry and Physics*, 12, 5447–5481, 2012.
- Troen, I. and Mahrt, L.: A simple model of the atmospheric boundary layer; sensitivity to surface evaporation, *Boundary-Layer Meteorology*, 37, 129–148, 1986.
- 1120 Tsyro, S., Aas, W., Soares, J., Sofiev, M., Berge, H., and Spindler, G.: Modelling of sea salt concentrations over Europe: key uncertainties and comparison with observations, *Atmospheric Chemistry and Physics*, 11, 10 367–10 388, 2011.
- Tuovinen, J.-P., Ashmore, M., Emberson, L., and Simpson, D.: Testing and improving the EMEP ozone deposition module, *Atmospheric Environment*, 38, 2373–2385, 2004.
- 1125 UNECE: HANDBOOK FOR THE 1979 CONVENTION ON LONG-RANGE TRANSBOUNDARY AIR POLLUTION AND ITS PROTOCOLS, UNECE Convention on Long-range Transboundary Air Pollution, UNITED NATIONS, New York and Geneva, 2004, joint WHO/Convention Task Force on the Health Aspects of Air Pollution, 2004.
- Van Donkelaar, A., Martin, R. V., Brauer, M., and Boys, B. L.: Use of satellite observations for long-term exposure assessment of global concentrations of fine particulate matter, *Environmental health perspectives*, 123, 135–143, <https://doi.org/10.1289/ehp.1408646>, 1130 2015.
- van Leer, B.: Multidimensional explicit difference schemes for hyperbolic equations," *Sixth Symposium on Computing Methods in Applied Science and Engineering*, 1984.

- Van Meijgaard, E., Van Ulft, L., Lenderink, G., De Roode, S., Wipfler, E. L., Boers, R., and van Timmermans, R.: Refinement and application of a regional atmospheric model for climate scenario calculations of Western Europe, KVR 054/12, KVR, 1135 2012.
- Van Zanten, M., Sauter, F., RJ, W. K., Van Jaarsveld, J., and Van Pul, W.: Description of the DEPAC module: Dry deposition modelling with DEPAC_GCN2010, RIVM rapport 680180001, 2010.
- Vautard, R., Bessagnet, B., Chin, M., and Menut, L.: On the contribution of natural Aeolian sources to particulate matter concentrations in Europe: testing hypotheses with a modelling approach, *Atmospheric environment*, 39, 3291–3303, 2005.
- 1140 Venkatram, A. and Pleim, J.: The electrical analogy does not apply to modeling dry deposition of particles, *Atmospheric Environment*, 33, 3075–3076, 1999.
- Walcek, C. J.: Minor flux adjustment near mixing ratio extremes for simplified yet highly accurate monotonic calculation of tracer advection, *Journal of Geophysical Research: Atmospheres*, 105, 9335–9348, 2000.
- Wesely, M.: Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical models, *Atmospheric* 1145 *Environment*, 41, 52–63, 2007.
- Winiwarter, W., Bauer, H., Caseiro, A., and Puxbaum, H.: Quantifying emissions of primary biological aerosol particle mass in Europe, *Atmos. Environ.*, 43, 1403–1409, <https://doi.org/10.1016/j.atmosenv.2008.01.037>, 2009.
- Yamartino, R.: Nonnegative, conserved scalar transport using grid-cell-centered, spectrally constrained Blackman cubics for applications on a variable-thickness mesh, *Monthly Weather Review*, 121, 753–763, 1993.
- 1150 Yamartino, R., Strimaitis, D., and Graff, A.: Evaluation of the Concentration Fluctuation Predictive Power of the Kinematic Simulation Particle Model, in: *Air Pollution Modeling and Its Application XIV*, pp. 563–571, Springer, 2004.
- Yarwood, G., Jung, J., Whitten, G. Z., Heo, G., Mellberg, J., and Estes, M.: Updates to the Carbon Bond mechanism for version 6 (CB6), in: 9th Annual CMAS Conference, Chapel Hill, NC, pp. 11–13, 2010.
- Yttri, K. E., Simpson, D., Nøjgaard, J. K., Kristensen, K., Genberg, J., Stenström, K., Swietlicki, E., Hillamo, R., Aurela, M., 1155 Bauer, H., Offenberg, J. H., Jaoui, M., Dye, C., Eckhardt, S., Burkhardt, J. F., Stohl, A., and Glasius, M.: Source apportionment of the summer time carbonaceous aerosol at Nordic rural background sites, *Atmos. Chem. Phys.*, 11, 13 339–13 357, <https://doi.org/10.5194/acp-11-13339-2011>, <http://www.atmos-chem-phys.net/11/13339/2011/>, 2011.
- Yuan, H., Dai, Y., Xiao, Z., Ji, D., and Shanguan, W.: Reprocessing the MODIS Leaf Area Index products for land surface and climate modelling, *Remote Sensing of Environment*, 115, 1171–1187, 2011.
- 1160 Zender, C. S., Bian, H., and Newman, D.: Mineral Dust Entrainment and Deposition (DEAD) model: Description and 1990s dust climatology, *Journal of Geophysical Research: Atmospheres*, 108, 2003.
- Zhang, K. M., Knipping, E. M., Wexler, A. S., Bhave, P. V., and Tonnesen, G. S.: Size distribution of sea-salt emissions as a function of relative humidity, *Atmospheric Environment*, 39, 3373–3379, 2005.

Zhang, L., Gong, S., Padro, J., and Barrie, L.: A size-segregated particle dry deposition scheme for an atmospheric aerosol
1165 module, *Atmospheric environment*, 35, 549–560, 2001.

Zhang, L., Brook, J. R., and Vet, R.: A revised parameterization for gaseous dry deposition in air-quality models, *Atmospheric
Chemistry and Physics*, 3, 2067–2082, 2003.