

The characterization of long-range transported North American biomass burning plumes: what can a multi-wavelength Mie-Raman-polarization-fluorescence lidar provide?

Qiaoyun Hu¹, Philippe Goloub¹, Igor Veselovskii², and Thierry Podvin¹

¹Univ. Lille, CNRS, UMR 8518 - LOA - Laboratoire d'Optique Atmosphérique, F-59650 Lille, France

²Prokhorov General Physics Institute of the Russian Academy of Sciences, Moscow, Russia

Correspondence: Qiaoyun Hu (qiaoyun.hu@univ-lille.fr)

Abstract. This article presents a study of long-range transported biomass burning aerosols (BBA) originated from the North American wildfires in September 2020. The BBA plumes presented in this study were in the troposphere and underwent 1-2 weeks aging before arriving at the [observations site lidar station ATOLL \(ATmospheric Observatory of LiLle\) in northern France](#). A novel [dataset lidar-derived dataset](#), $2\alpha+3\beta+3\delta+\phi$ (α : extinction coefficient, β : backscatter coefficient, δ : particle linear depolarization ratio, *i.e.*, PLDR, ϕ : fluorescence capacity) [derived from lidar observations](#), is provided for the characterization of [long-range transported BBA](#). The fluorescence capacity [describes is an intensive aerosol parameter describing the ability of aerosols in producing fluorescence when exposed to UV excitation](#). In [the observations of BBA episode, plumes from different wildfire activities have been characterized. In the BBA plumes, we detected low PLDRs, i. e. lower our BBA observations, obvious variations in aerosol intensive parameters, reflecting the variability of BBA properties, were detected. The PLDRs varied from less than 0.03 at all wavelengths, as well as enhanced PLDRs \(\$PLDR_{355,532,1064} \approx 0.15-0.18, 0.12-0.14, 0.01-0.03\$ \) with a similar spectral dependence that had been observed in the aged BBA plumes in the upper troposphere and lower stratosphere \(Canadian smoke in 2017 and Australian smoke in 2019-2020\). Obvious variations in Angström exponent \(-0.3-1.5\), lidar ratios \(20-50 sr to 0.15-0.22 and 0.12-0.16, respectively at 355 nm, 42-90 sr at and 532 nm\) and fluorescence capacity \(- \$1.0 \times 10^{-4}\$ - \$4.0 \times 10^{-4}\$ \) are also observed during the BBA episode. Lidar ratio as low as \$24 \pm 4\$ sr \(\$50 \pm 8\$ sr\) was observed in the BBA plumes at 355 \(532\) nm on 17-18 September, which was lower than most previously observed aged BBAs. These variations are coupled with the variation of altitudes, water vapor content and wildfire events. It reflects that the properties of aged BBA particles are highly varied and depend on complex mechanisms, such as burning process and the aging process. The results also pointed out the inhomogeneity of the aging process in the BBA plumes, which means that particles in the core of the plume aged differently with those at the plume edge due to the impact of water vapor, temperatures, particle concentration and so on. These chemical and physical processes involved in BBA aging and how they could impact the particle properties are not yet well understood likely correlated with the combustion process, the lifting of BBA plumes and the conditions \(temperature, humidities, etc.\) in the aging process. In addition, our observations identified the ice crystals mixing with BBA particles, which indicates that BBA results indicate BBA could act as ice nucleating particles \(INP\) at in tropospheric conditions. The lidar fluorescence channel proves to be an efficient tool in studying the interaction of clouds and BBAs important added value in](#)

aerosol characterization and aerosol-cloud interactions studies, due to its high sensitivity. ~~Recent studies claimed that aged BBA particles are more effective INPs than they were thought. As BBAs are becoming an important atmospheric aerosols with growing global wildfires, our observations could improve our characterization about~~ With the increase of wildfires occurrence and intensity, BBAs become a more and more important atmospheric component. In this context, we show the potential of our novel lidar-derived dataset for aged BBA particles ~~and characterization and for the understanding of their importance in ice cloud formation~~ role in cloud processes.

1 Introduction

Severe wildfires caught the attention of the public and the scientific communities in ~~resent~~ recent years. Wildfires in North America increase in both frequency and intensity, due to the warming-up of the climate. Regarding ~~to~~ the global scale, the duration of fire seasons increased by about 19% from 1979 to 2013 (Jolly et al., 2015; Schoennagel et al., 2017). Wildfires can directly impact the vegetation of earth's surface, the fertility of soils, water cycle and human society (Santín and Doerr, 2016; Ditas et al., 2018; May et al., 2014; O'Dell et al., 2020). Copious amount of fire emissions, including gases and particles, were injected into the troposphere and ~~occasionally~~ occasionally into the upper troposphere and lower stratosphere (UTLS). Those emissions pose threats to the air quality and human health, because they contain mainly fine particles and some chemical compounds in wildfire emissions are toxic. They could ~~alternate~~ alter the planetary radiation budget of the planet by scattering and absorbing the incoming solar radiation and influence the cloud process by modifying cloud properties and acting as ~~INP (ice nucleation particle) and CCN (INP) and cloud condensation nuclei (CCN).~~

Wildfire emissions are composed of a mixture of gases, e.g. carbon monoxide, polycyclic aromatic hydrocarbons (PAHs) containing gas-phase hazardous air pollutants, water vapor and aerosol particles, such as black carbon (BC), organic carbon (OC), etc. In addition, water soluble ions (potassium, halides, sulfates) can also be produced by the decomposition of biomass and the subsequent condensation in the smoke plume. The OC components account for a substantial fraction, about 20% to 90% in wildfire emissions, much higher than BC. Humic-like substances, which are a complex group of high molecular weight organic compounds, contribute considerably to the mass of organic compounds and influence the light absorbing properties of aerosols (Urbanski et al., 2008; Santín and Doerr, 2016; Wu et al., 2018). For convenience, we use hereafter BBA (biomass burning aerosols) to refer to the aerosols resulting from wildfire emissions. Due to the complexity in the burning and the aging processes, the BBA properties are highly variable. The initial composition and size distribution of ~~BBA depends~~ BBAs depend on the burning materials (i.e. vegetation type, such as grassland, boreal forest, etc.) and environmental conditions (temperature, moisture, soil properties, etc.), which impact the combustion efficiency and the burning processes (i.e. flaming and smoldering). The smoldering phase, which is with lower temperature, could lead to a substantially higher conversion of burning materials to toxic compounds than the flaming phase, and produce more weakly absorbing OC aerosols. Fire emissions from high temperatures and the flaming phase tend to contain an increased proportion of BC to OC and have smaller smoke particles (Reid et al., 2005; Garofalo et al., 2019).

Fresh BBAs are fractal like aggregates of BC cores and OC coatings (China et al., 2013, 2015). During long-range trans-

port, BBAs undergo complex aging processes, including condensation, oxidation, evaporation, coagulation, compaction and humidification, humidification and so on. These processes could modify the particles' morphology, composition, optical and chemical properties and hygroscopic properties. The aging process of wildfire emissions is not yet adequately characterized. It includes a complex group of chemical reactions and physical changes, among which the gas-to-particle condensation driven by oxidation and evaporation driven by aerosol dilution, are two important and competing processes. The oxidation process can produce more organic matter with lower volatility, while it could also lead to increased volatile SOA (secondary organic aerosol (SOA)) due to fragmentation of some macromolecules. The evaporation tends to decrease the partitioning of OC aerosols as dilution shifts the gas-particle equilibrium more to the gas end. Measurements in laboratories and field campaigns do not reach a consensus on the relative balance of the two competing processes as they are controlled by many factors, such as the dilution rate, exposure to oxidants (O₃, OH radical), travel time and so on. The growth of particle size is found in both near-field and longtime aged BBAs, while the causes might not be the same. Condensational growth is the dominating factor in fresh and near-field BBAs, while coagulation growth is dominating in aged BBAs. Abel et al. (2003) reported a increase of SSA (an increase of single scattering albedo (SSA)) from about 0.84 to 0.90 in 5 hours after the emission, which is likely due to the condensation of scattering organic components on absorbing soot particles during early-stage early-stage aging. Similar changes in SSA are derived by Kleinman et al. (2020) when investigating near-field BBA plumes from western US. The aging process can result in complex BBA structures and varying properties. Dahlkötter et al. (2014) sampled BBA particles in a 3-4 days old plume from western US wildfires in the UTLS. They found that a significant proportion of the particles are BC-containing particles mixed with light-scattering materials and the mixing state is highly varying. Observations of long-ranged transported Canadian smoke in 2017 and Australian smoke in 2019-2020 in UTLS showed enhanced PLDRs (particle linear depolarization ratios (PLDR)) that had never been detected in tropospheric smoke layers (Haarig et al., 2018; Hu et al., 2019; Ohneiser et al., 2020) (Haarig et al., 2018; Hu et al., 2019; Ohneiser et al., 2020; ?). Similar signatures of PLDRs have also been observed in aged tropospheric BBAs originated from wildfires (Murayama et al., 2004; Burton et al., 2015). Researchers thought such characteristics of PLDRs are indications of complicated BBA particle morphology and effects of longtime aging. Since then, many efforts have been done-made to simulate aged BBA particles' optical properties. The simulations indicate that aerosol properties, such as the absorption and PLDRs, are strongly dependent on the morphology, mixing state and fractions of different components.

BBAs contribute to ice nucleating (IN) mainly through immersion freezing, contact freezing, deposition freezing, as well as pore condensation and freezing (Kanji et al., 2011; Umo et al., 2015; Grawe et al., 2016; Kanji et al., 2017; Umo et al., 2019) (Kanji et al., 2011; Umo et al., 2015; Grawe et al., 2016; Kanji et al., 2017; Knopf et al., 2018; Umo et al., 2019). These mechanisms are not yet well resolved and data about the onset condition of ice nucleation by BBA particles are diverse. As wildfire emissions become a more and more important atmospheric aerosol source, the role of BBAs as INP becomes increasingly important and different theories have been proposed about how the BBA impact the activation of INP to understand BBAs' IN ability. One important argument is that the amorphous state of organic matter and complicated morphology of BBAs at atmospheric conditions could influence the ability in-of water diffusion, thus causing strong perturbations in the prediction of cloud formation (Berkemeier et al., 2014) (Berkemeier et al., 2014; Knopf et al., 2018). Recently, Jahn et al. (2020) reported a

different mechanism. They claimed that the inorganic elements naturally ~~presented~~present in biomass could be transformed by combustion process to minerals, which are very ice active over a wide range of ~~temperature~~temperatures. Studies also found that the aging process could potentially increase the capability of BBA in cloud formation (Lupi and Molinero, 2014; Jahl et al., 2021). ~~The~~Therefore, the characterization of BBA properties in light scattering and cloud formation after long-range transport is an important direction to improve the representation of aerosol model and further, the accuracy of the climate model.

The 2020 California wildfires are the largest in ~~the~~ Californian modern history. They comprise about 9639 fire activities reported in the California state, located in the western coast of the United States. The fire season started from the beginning of May 2020. By the end of the year, nearly 10,000 fires had burned over 4.2 million acres, more than 4% of the state's roughly 100 million acres of land (<https://www.fire.ca.gov/incidents/2020/>). In mid-August, the first 'gigafire' spreading over 7 counties in northern California was recorded. 'Gigafire' is a level above 'megafire' and describes a blaze burning at least a million acres of land. Satellite observations indicated this fire activity caused huge BBA emissions over the middle and western US, while the plumes faded away after 1 week without being transported to other continents. In September 2020, the fire activities revived after a short pause. Fire emissions were lofted into UTLS by high winds or convective pyro-cumulonimbus, and then transported to Europe by the prevailing westerlies. ~~The most dense~~(Baars et al., 2021). The densest smoke plumes detected at the observatory ATOLL (ATmospheric Observatory of LiLle) in Lille, France, is in the period of 10-23 September. In this study, we analyzed the observations from multiple instruments and the optical properties derived from a multi-wavelength Mie-Raman polarization lidar equipped with a fluorescence channel.

2 Observations

2.1 Satellite observations – OMPS and CALIPSO

According to satellite observations, the smoke plumes detected in Lille on 10-22 September are attributed to the fire activities on 04-11 September in California and Oregon state in the USA. On 04-06 September 2020, the Creek fire started in the southwestern California. ~~The Creek fire is the most intense wildfire in September 2020.~~ On 05 September, the fires generated a pyro-cumulonimbus (~~PyCb~~PyroCb) cloud reaching about 16000 m, which might be the highest ~~PyCb~~PyroCb cloud ever (Creek Fire, 2020). The plumes from Creek fire ~~are obvious~~can be clearly seen in the earth's true color image (Figure 1(a)) and the maps of the UVAI-UV aerosol index (UVAI) from OMPS observations shown in Supplementary information (SI). The center of the lofted plumes was at (37.7N, 117.8W) on 06 September and then transported rapidly toward the northeast.

On 07-10 September, the forest in the west coast of California and Oregon started to burn, which will be called Oregon fires hereafter (Oregon Fire, 2020). The burning area was in serious drought and under the blasts of fierce winds, which lifted vast fire emissions into the atmosphere. On 09 September, the wind whipped up fires and generated a pyro-cumulus cloud. Due to cyclone activities, the plumes were trapped on the eastern Pacific ocean and west coast of the US on 07-11 September, as shown in Figure 1(b). The plumes started spreading on 12 September and arrived at in Europe on 16-18 September.

Figure 2 shows the CALIPSO observations in the transport pathway. The plumes in Figure 2(a, b, c) and in Figure 2(d, e, f) are attributed to the Creek fire and the Oregon fire, respectively. The plume origins were identified using HYSPLIT back

trajectories and OMPS UVAI maps, shown in SI. The plumes emitted by the Creek fires are mostly classified as 'elevated smoke' by the CALIPSO aerosol typing algorithm (Version 4.21), while the plumes emitted from Oregon fires are mostly detected as 'polluted dust'. The reason is that the ~~particle linear depolarization ratio~~ PLDR at 532 nm (~~PLDR₅₃₂~~) of the BBA particles from Oregon fire was higher than Creek fire. PLDR is a parameter reflecting ~~the~~ particle shape, i.e. spherical particles have PLDR of zero while morphologically complicated particles such as dust and ice crystals have high PLDRs. ~~PLDR₅₃₂~~ PLDR₅₃₂ is about 0.02–0.05 in Creek plume after 3–7 days' transport in the upper troposphere. In comparison, the PLDR₅₃₂ in Oregon plume is about 0.10–0.15 after 1–7 days' transport. The ~~difference indicates that BBA particles generated from the Creek fire and Oregon fire may have some differences in the morphology and/or other properties~~ observed differences of BBA properties in the two events may be attributed to the combustion and lifting process.

2.2 Sun/sky photometer observations near wildfire source

NASA_Ames and PNNL are two AERONET sites located near the fire activities, as shown in Figure 1(b). PNNL was mainly impacted by the Oregon smoke during 12–19 September. ~~The AOD₅₀₀ (In Figure 3, the~~ Aerosol Optical Depth at 500 nm (AOD₅₀₀) was about 0.2 and ~~AE_{340–500} (the~~ Angström exponent between 340 and 500 nm (AE_{340–500}) about 1.5 ~~in on~~ 01–08 September, before the arrival of the Oregon smoke. When the Oregon plumes arrived at PNNL on 12 September, the AOD sharply increased and the AE ~~increased~~ decreased at the same time. The AOD₅₀₀ significantly increased ~~during the period 12-19 September and the maximum and the maximal~~ AOD₅₀₀ of was about 4.0 (with, with corresponding AE_{340–500} = of 0.2), during 12-19 September. NASA_Ames is in the west of the Creek fire and was influenced firstly by the Creek fire and then by the Oregon fire. AOD₅₀₀ on 04–08 September, when Creek smoke particles dominated, was less than 1.0 (with AE_{340–500} ~1.2), because the center of the Creek plumes propagated ~~eastwardly~~ eastward and only a part of it drifted to the west. The Oregon plumes started on 09 September, as indicated by a sharp increase of AOD and a decrease of AE_{340–500}. The peak AOD₅₀₀ of 5.8, corresponding to AE_{340–500} ~0.0, was detected on 12 September in NASA_Ames. Similar to the observation at PNNL, AE_{340–500} obviously decreased during the episodes of Oregon smoke. The decrease of AE_{340–500} ~~indicated~~ indicates the increase of particle size in the Oregon BBA plumes, which is in agreement with the CALIPSO observations.

2.3 Lidar and photometer observations in Lille, France

The lidar involved in this study is a multi-wavelength Mie-Raman-polarization-fluorescence lidar, LILAS, operated at ~~Laboratoire d'optique atmosphérique (LOA)~~ ATOLL platform, University of Lille, France. The lidar system uses a Nd: YAG laser and has three emitting wavelengths, i.e., 355, 532 and 1064 nm. The backscattered lidar signals are collected at three elastic wavelengths each coupled with a parallel channel and a cross channel, and three Raman channels at 387 (~~vibration Raman of~~ nitrogen vibrational Raman of N₂ and O₂), 530 (rotational Raman of ~~nitrogen N₂ and O₂~~) and 408 nm (water vapor channel). ~~Since In~~ December 2019, the water vapor channel was replaced by a fluorescence channel centered at 466 nm (excitation wavelength: 355 nm) in order to profile atmospheric fluorescence. A full description of this configuration and the results of the feasibility test can be found in Veselovskii et al. (2020). With this configuration, we can derive the height-resolved dataset $2\alpha + 3\beta + 3\delta + \phi$ (α : extinction coefficient, β : backscattering coefficient, δ : particle linear depolarization ratio, ϕ : fluorescence

160 capacity). The ~~methodology has been presented in our previous publications (Hu et al., 2019; Veselovskii et al., 2020; ?) and will not be repeated in this paper.~~ The fluorescence signal is attributed to certain molecules that could absorb in the incident laser light and re-emit it at longer ~~wavelength~~ wavelengths. The fluorescence capacity represents the capacity of aerosol particles in producing fluorescence, hence it is linked to particle chemical compositions. Equation 1 and 2 demote respectively the fluorescence backscattering coefficient $\beta_F(z)$ and the fluorescence capacity $\phi(z)$.

$$165 \quad \beta_F(z) = \frac{C_R P_F(z) T_R(z)}{C_F P_R(z) T_F(z)} N_R(z) \sigma_R, \quad (1)$$

where $P_F(z)$ and $P_R(z)$ are the lidar signals of fluorescence and Raman channel, respectively, $\frac{C_R}{C_F}$ is the ratio of instrumental constant between the Raman and the fluorescence channel and it is determined by calibration. $N_R(z)$ is the vertically distributed number concentration of Raman scatters (i.e., N_2 and O_2) and σ_R is the Raman differential scattering cross section in the backward direction. With β_F , one can define the fluorescence capacity $\phi(z)$:

$$170 \quad \phi(z) = \frac{\beta_F(z)}{\beta_{\lambda_0}(z)}. \quad (2)$$

According to our observations, BBA and pollen particles are very efficient in producing fluorescence, while dust particles and continental aerosols are much less efficient. More information about fluorescence lidar measurements and aerosol fluorescent signatures can be found in our previous publications (Hu et al., 2019; Veselovskii et al., 2020, 2021b).

Figure 4 shows lidar observations in the period of 11-22 September 2020. Figure 4(a) and 4(b) show the range corrected
 175 lidar signal and the volume depolarization ratio at 1064 nm, respectively. The background aerosols in Lille are usually low depolarizing fine particles in the boundary layer. During the observational period, the BBA layers ~~distributing~~ distributed at 2000-14000 m and polluted dust in the boundary layer were the two aerosol species. The smoke plumes are obvious in Figure 4(a) and disappear in Figure 4(b), because BBA particles have very weak depolarization at 1064 nm. The most intense smoke plumes are observed on 11-12 September, corresponding to $AOD_{500} \approx 0.65$ and $AE_{340-500} \approx 0.70$, shown in Figure 4(c).
 180 ~~Polluted dust from west Africa arrived in Lille on 14 September and concentrated mainly in~~ Aerosols in the boundary layer on 14-16 September were strongly depolarizing. Back trajectories and modeling results indicated the cause for the enhanced depolarization could be polluted dust or transported pollen. The intrusion of polluted dust does not cause a significant decrease of $AE_{340-500}$, which can be explained by mixing with a non-negligible fraction of fine mode aerosol component, such as industrial pollution and smoke. On 17-18 September, a decrease of $AE_{340-500}$ is detected when a smoke plume appeared at
 185 6000-10000 m. The AOD_{500} and $AE_{340-500}$ on 17 September ~~is~~ are about 0.22 ± 0.02 and 0.0 ± 0.20 , respectively.

Figure 5 plots the intensive parameters in the BBA layers detected in time intervals in September 2020. The observation time and the height range of the layers are ~~resented~~ presented in Table 1. Lidar ratio (LR), i.e., the ratio between extinction coefficient and backscatter coefficient, is correlated with multiple factors, such as the particle shape, size and refractive indices. LR_{355} being lower than LR_{532} is a typical feature of aged smoke particles from North America and Siberian wildfires. The
 190 values of lidar ratios varied in the range of 20–50 sr at 355 nm and 42–90 sr at 532 nm. The spectral dependence of the PLDRs is in agreement with the aged wildfire smoke particles in the literature. The variation ranges of PLDRs are respectively

0.03–0.22, 0.02–0.16 and 0.01–0.03 at 355, 532 and 1064 nm in the investigated cases. The PLDRs on 18 and 19 September (P7 and P8) are higher than other days, with mean PLDR equals to 0.16 at 355 nm, 0.12 at 532 nm and 0.02 at 1064 nm. The fluorescence capacity varied from 1.0×10^{-4} to 4.0×10^{-4} . The $\text{BAE}_{355-532}$ (Angström exponent related to backscatter coefficient ($\text{BAE}_{355-532}$)) is in the range of 1.5–2.5. Whereas, the variation of $\text{EAE}_{355-532}$ (Angström exponent related to extinction coefficient ($\text{EAE}_{355-532}$)) is stronger. Before 17 September, the $\text{EAE}_{355-532}$ varied in the range 0.5–1.5 and after this day, it dropped to below zero. ~~Apart from the temporal variations,~~ The vertical variations in the BBA layers, indicated by the distance between the bottom and the top of each box plot in Figure 5, are also significant, such as lidar ratios ~~in~~ on 12 and 14 September, the PLDRs on 11 and 18 September. Such variations are possibly indications of the variabilities in the burning materials, combustion conditions, convective lifting and aging process. Two events corresponding to Creek fire and Oregon fire, respectively, are selected and analyzed in detail in ~~the~~ next section.

3 Case study

3.1 Case 1: 11-12 September 2020

The BBA plumes detected ~~in~~ on the night of 11-12 September 2020 are attributed to the Creek fire and have traveled about 5-7 days. Figure 6(a) shows the 4-day back trajectories overlaid with UVAI map on 8 September 2020. Two areas with intense UVAI are indicated on the map, one over the western coast of the US and the other one over the Great Lakes. The former was mostly generated by the Oregon fire, which started on 7 September. The latter was emitted by the Creek fire on 4-6 September and transported to Europe via an 'expressway'. Figure 7 shows the range corrected lidar signal (RCS) and volume linear depolarization ratio (VLDR) and ~~the relative fluorescence signal, $\frac{P_{466}(z)}{P_{387}(z)}$, on 11-12 September~~ calibrated signal ratio between 466 and 387 nm channel, i.e., $\frac{C_R P_{466}(z)}{C_F P_{387}(z)}$. The fluorescence observations are only available ~~in~~ at nighttime when there is no sunlight interference. The BBA layer was mainly distributed at 5000-10000 m and was characterized by strong fluorescence. ~~Ice particles with strong depolarization~~ In Figure 7(b), ice particles featured with increased depolarization ratios were detected within ~~the smoke layers above 8000 m~~ high smoke layers, for example, at 8000-10000 m at 17:00–18:00 UTC.

Figure 8 plots the parameters obtained from averaged observations between 22:00 UTC, 11 September and 03:00 UTC, 12 September. The extinction coefficients peaked at about 6200 m, with ~~about~~ 180 Mm^{-1} at 355 nm and 140 Mm^{-1} at 532 nm. The $\text{EAE}_{355-532}$ was about 0.5 ± 0.3 and the $\text{BAE}_{355-532}$ ~~is~~ was about 2.0 ± 0.3 . The $\text{EAE}_{355-532}$ is lower than the $\text{AE}_{340-500}$ measured at NASA_Ames when influenced by the Creek fire plumes, indicating the particle growth during the aging process. The lidar ratios, which are 40 ± 6 sr at 355 and 70 ± 11 sr at 532 nm, are typical values of aged smoke particles (Haarig et al., 2018; Hu et al., 2019; Ohneiser et al., 2020; Peterson et al., 2018). The PLDRs show strong vertical variations ~~through~~ in the BBA layer. In the core of the smoke, i.e., 5000-6000 m, the PLDRs at the three wavelengths are all below 0.03. In the ~~range of plume edge, i.e., 6500-9000 m (the plume edge),~~ m, the PLDRs at 355 ~~nm~~ and 532 nm increased significantly, with about 0.15 ± 0.03 at 355 nm and 0.12 ± 0.02 at 532 nm, ~~indicating the change of particle morphology versus altitude.~~ PLDR_{1064} shows no significant vertical variations. A slight decreasing trend ~~is observed~~ in the $\text{EAE}_{355-532}$ and a moderate

225 ~~one in BAE₃₅₅₋₅₃₂ are observed~~ when the height increases, ~~which indicate the increase of particle size. The increasing trend~~
~~of BAE₃₅₅₋₅₃₂ is stronger than EAE₃₅₅₋₅₃₂. The layer~~. BBA particles at 5000-6000 m ~~is was~~ also characterized with slightly
stronger fluorescence capacity ($\sim 3.8 \times 10^{-4} \text{ sr}^{-1} \text{ m}^{-1}$) and higher water vapor content compared to the smoke in the range of
) compared to those at 6500-9000 m. According to laboratory studies, the main fluorophores in smoke are the humic-like sub-
230 ~~stance and poly-aromatic hydrocarbons~~ Garra et al. (2015); Zhang et al. (2019) PAHs (Garra et al., 2015; Zhang et al., 2019),
which are common products of the combustion of biomass. ~~During the aging process, the formation of some SOA could also~~
~~contribute to the fluorescence signal, as well as some SOA formed in the aging process~~ (Lee et al., 2013). Therefore, temporal
and spatial variations of fluorescence capacity ~~of smoke are expectable~~ are very likely during the transport, and such variation
is ~~dynamic and controlled~~ controlled by factors, such as oxidation and evaporation of some chemical species and so on.
Figure 8(g) presents the WVMR ~~, RH and temperature~~ (water vapor mixing ratio) and RH (relative humidity) profiles measured
235 by radio sounding at ~~Beauvechain, Belgium station~~ Beauvechain station (Belgium), 120 km to ~~the lidar station in Lille~~ ATOLL
platform. The RH with respect to ice (RH_{ice}) was calculated with the radio sounding measurements (Jarraud, 2008). The
~~variation presence~~ of the BBA layer is well reflected by the ~~profile of WVMR, which is also a tracer of air mass~~ enhanced
values in the WVMR profile. It indicates that the BBA plumes were also captured by the radio sounding measurements at
~~Beauvechain~~ Beauvechain due to the plumes' large spatial coverage ~~of the plumes~~. The WVMR ~~increases from the~~ in the core
240 of the plume is obviously higher than at the plume ~~center to the~~ edge, suggesting that the WVMR is an important ~~role in the~~
aging process. ~~factor in the aging process.~~

3.2 Case 2: 17-18 September 2020

The smoke layers at 5000-9000 m on 17-18 September 2020 (Figure 9) are attributed to the Oregon fire. ~~The plumes underwent~~
~~7-12 days of transport before arriving at the observation site, as shown in~~ Figure 6(b) ~~presents the 6-day back trajectories~~
245 ~~overlaid with UVAI from OMPS on 14 September 2020.~~ The plumes covered large areas of America and ~~spread to the~~ eastern
Pacific Ocean. ~~The lidar quicklooks on 17-18 September are shown in Figure 9, and underwent 7-12 days of transport before~~
arriving at the observation site ATOLL, France.

Profiles obtained from averaged measurements between 22:30 UTC, 17 September and 03:00 UTC, 18 September are plotted in
Figure 10. ~~The extinction coefficients peaked at about 7500 m, with $80-85 \pm 9 \text{ Mm}^{-1}$ at both 355 and 532 nm~~ Compared to Case
250 1, BBA plumes in Case 2 have lower extinction coefficients, EAE₃₅₅₋₅₃₂, lidar ratios and higher PLDRs. The EAE₃₅₅₋₅₃₂
is ~~slightly negative, around~~ -0.30 ± 0.30 , obviously lower than in Case 1. ~~This feature indicates that smoke particles~~ It can be
explained by the longer travel time in Case 2 ~~are bigger than those in Case 1. The BAE₃₅₅₋₅₃₂ is comparable with that in Case 1.~~
~~Compared to Case 1, lidar ratios in Case 2 are lower, with~~, making the BBA particles bigger. Particle growth during transport
has been reported in previous aged smoke observations (Müller et al., 2007; Hu et al., 2019; Bian et al., 2020). The lidar ratios
255 are 24 ± 4 sr at 355 nm and 50 ± 8 sr at 532 nm. ~~These values are lower than many reported lidar ratios of aged smoke, lower~~
than the values in Case 1 and in most previous studies (Haarig et al., 2018; Hu et al., 2019; Ohneiser et al., 2020). ~~At the~~
355-nm, the measured lidar ratio is close to that of cirrus clouds (Haarig et al., 2016; Hu, 2018). Such low lidar ratios were
detected by Ortiz-Amezcuca et al. (2017) in ~~Raman lidar observations of~~ long-range transported North America smoke over two

European lidar stations, Leipzig ~~and Granada (Germany) and Granada (Spain)~~. The observations in Ortiz-Amezcuca et al. (2017) showed mean $EAE_{355-532}$ of 0.2-0.3, and mean lidar ratios of 23-25 sr at 355 nm and 47-51 sr at 532 nm. PLDRs in Case 2 are 0.15 ± 0.03 to 0.18 ± 0.04 at 355 nm, 0.12 ± 0.02 to 0.14 ± 0.03 at 532 nm and about 0.02 at 1064 nm, higher than those in Case 1 ~~at all, especially at shorter lidar~~ wavelengths. The fluorescence capacity varied in $2.0-3.0-2.8 \times 10^{-4}$ at 5300-8000 m, which is ~~lower than in Case 2~~ 26-47% lower than that plume core, and comparable with that in the plume edge in Case 1. ~~The BBA plumes in Case 2 traveled longer time and were diluted to a higher extent. During this process, the removal (e.g. dry and wet deposition), transformation (oxidation, dissociation) and evaporation of versatile species could lead to the decrease of the main fluorophores. Whereas, the difference in vegetation type and combustion conditions between Case 1 and 2 cannot be excluded. This decrease of fluorescence capacity may be linked to the evaporation of organic species when the BBA plumes is diluted during the transport (Hodshire et al., 2021).~~

~~A~~ In Figure 10(e), a sharp increase of $PLDR_{1064}$ to nearly 0.10 was detected at 8600 m, indicating the presence of ice crystals. At such altitude, the temperature was about -41°C to -34°C and the RH_{ice} was 100-120%, according to GDAS model data (GDAS: global data assimilation system) and radio sounding measurements. Lidar signals before and during the presence of cirrus have been averaged and shown in Figure 11. The fluorescence backscattering coefficient in the cirrus layer is as high as before the presence of ice crystals, indicating that the cirrus layer contains non-negligible BBA particles. ~~The decrease of fluorescence capacity in cirrus layer is mainly due to the enhancement of elastic scattering associated with the presence of ice crystals.~~ This observation is an indication of BBA particles acting as INP, although the possibility that ~~the~~ ice crystals formed elsewhere cannot be excluded. Considering that ~~there were almost no high clouds, in liquid or solid phase, over Lille~~ cloud coverage is quite low over northern France on 17-18 September, these ice crystals were very likely formed through deposition freezing mode or pore condensation and freezing mode (Cziczko et al., 2013; Umo et al., 2019).

4 Discussion

4.1 Optical properties: AE, PLDRs, lidar ratios and fluorescence capacity

In this study, ~~we demonstrate that the key intensive BBA optical properties, i.e. the Angström exponent ($AE_{340-500}/EAE_{355-532}$), fluorescence capacity, lidar ratios and PLDRs, have obvious vertical and temporal variations. These variations are potentially linked to complex factors, such as the fuel types, combustion conditions and aging process. Angström exponent, $AE/EAE_{355-532}$, decreased when the plumes transported from the source region to the observation site. This decrease reflects the increase of particles size during the transport, as has been observed by both in-situ and remote sensing studies. The increase of particle size in long-range transport is the result of condensational growth and coagulation of BBA particles (Müller et al., 2007; Haarig et al., 2018; Garcia et al., 2019). The spectral dependence of PLDRs and lidar ratios is also consistent with previous observations of aged North American BBA (Burton et al., 2015; Haarig et al., 2018; Hu et al., 2019). In this study, the PLDRs demonstrate a strong dependence on altitude, i.e. the higher altitude the higher PLDRs. The vertical variation of PLDRs has been discussed in previous studies. During the episode of the PLDRs of BBA originated from wildfires showed obvious variations dependent on different fire activities, different altitudes and water vapor content of the BBA layers. BBAs originating from the Oregon fire generally had~~

higher PLDRs than those from the Creek fire. Such a feature is probably due to the differences of burning vegetation type, combustion conditions and lifting process. During the episodes of transported Canadian smoke over Europe, lidar systems in Leipzig and Lille both found that the tropospheric BBA plumes ~~simultaneously detected with the UTLS plume~~ had much lower PLDRs ~~smaller than 0.10 at all wavelengths~~. Baars et al. (2019) summarized the Canadian smoke observations from different lidar stations within EARLINET and concluded that the PLDRs tend to decrease with travel time. Baars et al. (2019) explained that aged BBAs are mostly BC-containing and OC-coated aggregates and the growth of coating during aging process makes the particles more spherical. However, this argument is in conflict with the observations of Ohneiser et al. (2020), who found that in the 2020 Australian wildfires smoke layers above 18 km mostly have PLDR (at 532 nm) greater than 0.15, obviously higher than those below this altitude in than the stratosphere. Given that it takes time for the absorbing smoke plumes to raise to higher altitude by self-lifting, which is claimed by Ohneiser et al. (2020) and other researchers (Laat et al., 2012; Hu et al., 2019; Torres et al., 2020), plumes at higher altitude tend to be older. ~~mentioned that the significant PLDRs should not be simply explained by the presence of fractal BC with non-spherical OC coating and hypothesized that the OC-coated particles should be solid because they freeze or effloresce at stratospheric temperatures. This argument can explain why the PLDRs of BBAs in the stratosphere are higher than those in the troposphere. However, it cannot fully explain the sharp vertical variation of PLDRs from the core to the edge of the same BBA plume, as indicated in Case 1 of this study. Such variation indicates that the aging process is inhomogeneous in the plumes and may be sensitive to humidity, temperature and aerosol concentration. Hodshire et al. (2021) found that on the plume edges, the organic aerosol is more oxygenated compared to the plume cores. plumes simultaneously presenting in the UTLS (Haarig et al., 2018; Hu, 2018). In Case 1, the increase of PLDRs with altitude was observed in the BBA layer. This correlation between aged BBA PLDRs and altitude is likely driven by the altitude-dependent factors in the atmosphere, such as RH, temperature, etc., which influence the BBA particles' morphology in the aging process (Yu et al., 2019; Ohneiser et al., 2020; Ansmann et al., 2021). The organic aerosols, the main contributor of the BBA mass, can exist in amorphous state, in response to temperatures and RH (Koop et al., 2011; Berkemeier et al., 2014; Knopf et al., 2018). Studies demonstrated that the evaporation of condensed liquid water on the surface of BBA contributes strongly to the collapse of fractal-like aggregates into compact particles (Kütz and Schmidt-Ott, 1992; Ma et al., 2013). The organic aerosols, the main contributor of the BBA mass, can exist in amorphous state, which means that they can transform into liquid, semi-solid and solid (glassy state) in response to temperatures and humidities (Koop et al., 2011; Berkemeier et al., 2014; Knopf et al., 2018). This special property influences not only the light scattering behavior of BBA particles but also their abilities of ice nucleation. However, the current theory cannot explain the observed variability in the evolution of BBA properties concentration of BBA particles, which generally decrease during transport and aging, could also impact the evaporation and coagulation of particle aging, thus modifying the particles' morphology Hodshire et al. (2021).~~

The vertical variation of lidar ratios are also obvious in September (Figure 5), however, such variations are resulted not only from the variability of BBA properties, but also the artifacts in the calculation of the extinction-to-backscatter ratio. The calculation of extinction coefficient requires vertical smoothing, while the backscatter coefficient does not. Hence, the division of the two values could lead to some fake vertical fluctuations, especially in the range where aerosol concentration changes fast. From this aspect, the fluorescence capacity is a better parameter for aerosol typing especially in low aerosol concentrations.

The variation of lidar ratios, also the PLDRs and $EAE_{355-532}$ and fluorescence capacity, from the Creek fire to the Oregon fire is a clear indication of the variability of BBA properties originated from different fires activities. The lower lidar ratios detected in Case 2 of this study were rarely observed in previous lidar measurements. The decrease of lidar ratios in the Oregon plumes ~~is probably due to the decrease of~~ could be an indication of the decreased aerosol absorption, according to the modeling of aerosol optical properties. ~~However, other possible reasons such as particle morphology and hygroscopic effects can also impact the lidar ratio and cannot be excluded. The varied fluorescence capacity may reflect, at least to some extent, the changes of aerosol absorption, because aerosols absorb the incident radiation and re-emit the fluorescence. More observations and collocated measurements are required to explore the links between the fluorescence capacity and aerosol absorption.~~ Many efforts have been done to reproduce the light scattering properties of aged BBAs, such as absorption, PLDRs and lidar ratios, using different aerosol models. Liu and Mishchenko (2020) and Gialitaki et al. (2020) have successfully modeled the spectral PLDRs and lidar ratios observed in aged Canadian BBA plumes using near-spherical-spheroidal and Chebyshev particles. (Gialitaki et al., 2020; Liu and Mishchenko, 2020). Different scattering models have been considered to understand how aerosol composition, morphology and mixing state influence BBA absorption. Liu and Mishchenko (2018) used 11 different morphology models, ranging from bare soot to completely embedded soot mixtures, ~~and simulated several scattering parameters, such as the absorption enhancement, absorption Angström exponent and PLDRs. Liu and Mishchenko (2018) and~~ found that the absorption enhancement is affected by the particle size, morphology and the thickness of coating. ~~They also reproduced the spectral PLDRs observed by Burton et al. (2018) and concluded that multi-wavelength PLDR measurements can be used to identify the presence of morphologically complex carbonaceous particles. The results of modeling demonstrate~~ that composition and morphology of aged BBA are essential parameters that determine their properties. Moreover, the combustion and aging process, as well as the environment conditions could also contribute to the variability of BBA absorption. More efforts in the characterization of wildfire emissions and the modeling of BBA scattering are required to answer this question. The fluorescence signal brings new information for the characterization of aerosol properties. In previous studies we have shown that the fluorescence signal is a good reference for aerosol classification (Veselovskii et al., 2020, 2021b). In this study, we showed that the fluorescence capacity is correlated with the optical properties of BBA particles. As the fluorescence signal is attributed to certain chemical species, the variation of the fluorescence capacity directly reflects their abundance in BBAs, which is controlled by the combustion process, which determined their original concentration, and the aging process, consisting of complex processes, such as the oxidation, evaporation of versatile species, the condensation of gases and so on. For example, the decrease of fluorescence capacity at the plume edge in Case 1 is likely due to the evaporation of some fluorescence organic species when the BBA plume is diluted. ~~However, these studies are currently piecewise and hardly comparable, because they used different scattering models. To achieve a better characterization, more measurements and new information are needed. The idea of including laser-induced fluorescence is to extend the optical dataset and refine aerosol characterization, because with fluorescence information we could get a closer insight about the particle chemical compositions.~~

360 4.1 BBA acting as INP

Handful studies have shown that BBAs may serve as ~~a source of INP (Petters et al., 2009; McCluskey et al., 2014; Prenni et al., 2012)~~ INP (McCluskey et al., 2014; Jahn et al., 2020; Jahl et al., 2021), but existing results of ice nucleation onset conditions are highly ~~varied. The determination of the ice-active components in BBA and their mechanisms of ice nucleation are inadequate. Previous studies on ice nucleating abilities of BBA are mostly about soot particles, OC and inorganic components. Recent~~ studies based on experimental data and modeling claimed that BC is unimportant in ice cloud formation (Vergara-Temprado et al., 2018; Sel
~~-. The role the amorphous organic matter in the ice nucleating ability of BBA matter is uncertain. The phase state of organic matter influence the water diffusivity and the timescales of the phase transitions, which could modulate the pathways of ice formation (Mikhailov et al., 2009; Berkemeier et al., 2014; Wang et al., 2012; Lienhard et al., 2015; Shiraiwa et al., 2017; Price et al., 201~~ variable. Laboratory studies on organic INPs are mostly limited to pure organics, such as sucrose and levoglucosan~~ete, etc.,~~
370 while OA in ambient conditions have complex origins. As a ~~results~~ result, the conditions for initiating ice nucleation required by different organics are highly varying in literature data, which blocked us from clarifying the role of organic aerosols in ice formation (Kanji et al., 2017). ~~Recently Jahn et al. (2020) pointed out the importance of inorganic elements, which naturally existed in biomass and can be transformed into potentially ice-active minerals. Jahn et al. (2020) presented that these particles containing ice-active minerals are the major INPs in BBA because they could nucleate ice in a wide range of temperatures, up to~~ -13C. The influence of BBA aging on its ability of INP is also an important issue to be taken into account. Several studies have reported the enhancement of ice-nucleating ability of BBA after aging process, although they have not yet reached a consensus on the mechanisms (Brooks et al., 2014; Lupi and Molinero, 2014; Mahrt et al., 2020; Jahl et al., 2021). The establishment of a thorough determination the mechanism of ice nucleation by BBA particles requires more data from field measurements and laboratory measurements based on advance aerosol particle analytical techniques. Veselovskii et al. (2021a) presented that one
380 can estimate the particle concentration even in thin aerosol layers using the fluorescence signal, which offers a new opportunity for the studies of aerosol-cloud interactions. A paper about the estimation of BBA surface concentration in freshly formed cirrus layers using fluorescence measurements collected at ATOLL platform has been presented in (Veselovskii et al., 2021a).

5 Conclusions

This study reported the observations of long-range transported BBA plumes originated from wildfires in ~~West North~~ West North Amer-
385 ica in September 2020. The analysis involves data from multiple sources, including satellite measurements, ground-based sun/sky photometer and lidar and models. In September 2020, two intense fires—the Creek fire and Oregon fire, have been identified by OMPS and CALIPSO observations. After 1-2 weeks transport, these transport, the emitted BBA plumes in the troposphere were detected by LILAS, a multi-wavelength Mie-Raman polarization fluorescence lidar in ~~Lille, north of~~ Lille, north of France. The optical properties $2\alpha+3\beta+3\delta+\phi$ obtained from lidar measurements provided a good dataset for BBA particle
390 characterization. Our results demonstrated ATOLL, France. Our results confirmed the main typical features of aged BBA particles ~~that have been reported in previous research~~, i.e. the decreased Angström exponent after long-range transport, enhanced PLDRs at shorter wavelengths and lidar ratio at 532 nm greater than at 355 nm. ~~These characteristics are very likely~~

395 resulted from the growth of particle size and changes of particle morphology during the aging process, that have been reported in previous research. Moreover, our results showed strong variations of particle intensive parameters dependent on altitudes and combustion events. In the Creek plume, PLDRs increased with altitudes and from the core (PLDR_{355,532,1064} < 0.03) to (0.02) to 0.22 (0.16) at 355 (532) nm during the observations in September. Very specific lidar ratios, 24±4 sr at 355 nm and 50±8 sr at 532 nm, which are lower than most previous observations, were detected in the transported BBA plume from Oregon fire. These characteristics reflect that the BBA properties, such as morphology and absorption, are highly dependent on complex factors in the combustion (vegetation type, burning temperature, wind conditions, etc.) and aging process (humidities, temperature, etc.).
400 These variations need to be resolved and should be considered in the edge of the plume (PLDR_{355,532,1064} ≈ 0.15, 0.12, 0.01). Such characteristics suggest that the aging process is not homogeneous in the BBA plume and may dependent on environmental conditions such as humidities, particle concentration, temperature and so on. Compared to the Creek plume, the decrease of lidar ratios, EAE₃₅₅₋₅₃₂ (and AE) and fluorescence capacity ϕ in Oregon plumes, as well as the enhancement of PLDR₅₃₂ pointed out the variability of BBA properties in different combustion events. The low lidar ratios in the Oregon plumes are possible linked representation of BBAs in the climate model. The fluorescence backscattering coefficient and fluorescence capacity turned to be useful and supplementary information for BBA characterization, due to the high sensitivity to the presence of low absorbing particles fluorophores in BBAs. The fluorescence capacity derived from the fluorescence channel at 466 nm turned to be a better parameter than lidar ratio in the identification of smoke BBAs, because of its high sensitivity to low concentrations.
405 Benefitted from the fluorescence channel, lidar measurements allow us to directly detect the signature of BBA particles in ice clouds. During the smoke episode, BBA particles existing in cirrus cloud layers have been frequently detected. Our results suggest that aged BBA particles could nucleate ice crystals under atmospheric conditions in the UTLS, allows for an efficient identification of BBA particles and the quantification of fluorescent aerosols at low concentration, as has been presented in Veselovskii et al. (2020). The fluorescence capacity of future lidar LIFE (Laser-induced fluorescence explorer) at LOA will be enhanced with signal also opens the opportunity of study BBA particles inside cirrus clouds, as shown in this study and Veselovskii et al. (2021a). However, additional observations on different aerosol types and spectral fluorescence information are still needed to refine the study. A new fluorescence lidar with a more powerful laser and more fluorescence channels multiple fluorescence channels is now under development in order to obtain higher quality observations. As wildfire emissions are becoming an important atmospheric component, our observations could help improve the BBA characterization and lead to a better understanding of the BBA aging process and the role of BBA in ice nucleation advance the studies of aerosols and aerosol-cloud interactions.
420

Data availability. The data is available upon any request.

Author contributions. QH performed lidar measurements, developed the code for data analysis and wrote the manuscript. PG is the PI of the project, supervised the experiments and improved the manuscript. IV helped in data analysis and establishment of the scientific arguments.
425 TP helped perform lidar measurements and is in charge of the maintenance, automatization and operation of lidar system.

Competing interests. The authors declare no competing interests.

Acknowledgements. We acknowledge the CaPPA project (Chemical and Physical Properties of the Atmosphere), funded by the ANR (French National Research Agency) through the PIA (Programme d'Investissement d'Avenir) under contract « ANR-11-LABX-0005-01 », the Regional Council « Hauts-de-France » and the FEDER (European Funds for Regional Economic Development). We also would like to thank
430 the following organizations and individuals for the financial support and for sharing their data and expertise: Service National d'Observation PHOTONS/AERONET-EARLINET, ESA/QA4EO, AERONET (PIs of the sites involved in this study: Roy Johnson, Brent Holben, Philippe Goloub), CPER CLIMIBIO, Russian Science Foundation (project 21-17-00114) and ~~Lab-Agora~~[Agora-LAB](#).

References

- Abel, S. J., Haywood, J. M., Highwood, E. J., Li, J., and Buseck, P. R.: Evolution of biomass burning aerosol properties from an agricultural
435 fire in southern Africa, *Geophysical Research Letters*, 30, <https://doi.org/10.1029/2003GL017342>, 2003.
- Alduchov, O. A. and Eskridge, R. E.: Improved Magnus Form Approximation of Saturation Vapor Pressure, *Journal of Applied Meteorology and Climatology*, 35, 601 – 609, [https://doi.org/10.1175/1520-0450\(1996\)035<0601:IMFAOS>2.0.CO;2](https://doi.org/10.1175/1520-0450(1996)035<0601:IMFAOS>2.0.CO;2), https://journals.ametsoc.org/view/journals/apme/35/4/1520-0450_1996_035_0601_imfaos_2_0_co_2.xml, 1996.
- Ansmann, A., Riebesell, M., Wandinger, U., Weitkamp, C., Voss, E., Lahmann, W., and Michaelis, W.: Combined Raman elastic-backscatter
440 lidar for vertical profiling of moisture, aerosol extinction, backscatter, and lidar ratio, *Applied Physics B: Lasers and Optics*, 55, 18–28, <https://doi.org/https://doi.org/10.1007/BF00348608>, 1992.
- Ansmann, A., Ohneiser, K., Mamouri, R.-E., Knopf, D. A., Veselovskii, I., Baars, H., Engelmann, R., Foth, A., Jimenez, C., Seifert, P., and Barja, B.: Tropospheric and stratospheric wildfire smoke profiling with lidar: mass, surface area, CCN, and INP retrieval, *Atmospheric Chemistry and Physics*, 21, 9779–9807, <https://doi.org/10.5194/acp-21-9779-2021>, <https://acp.copernicus.org/articles/21/9779/2021/>,
445 2021.
- Baars, H., Ansmann, A., Ohneiser, K., Haarig, M., Engelmann, R., Althausen, D., Hanssen, I., Gausa, M., Pietruczuk, A., Szkop, A., Stachlewska, I. S., Wang, D., Reichardt, J., Skupin, A., Mattis, I., Trickl, T., Vogelmann, H., Navas-Guzmán, F., Haeefe, A., Acheson, K., Ruth, A. A., Tatarov, B., Müller, D., Hu, Q., Podvin, T., Goloub, P., Veselovskii, I., Pietras, C., Haeffelin, M., Fréville, P., Sicard, M., Comerón, A., Fernández García, A. J., Molero Menéndez, F., Córdoba-Jabonero, C., Guerrero-Rascado, J. L., Alados-Arboledas, L., Bortoli, D.,
450 Costa, M. J., Dionisi, D., Liberti, G. L., Wang, X., Sannino, A., Papagiannopoulos, N., Boselli, A., Mona, L., D’Amico, G., Romano, S., Perrone, M. R., Belegante, L., Nicolae, D., Grigorov, I., Gialitaki, A., Amiridis, V., Soupiona, O., Papayannis, A., Mamouri, R.-E., Nisantzi, A., Heese, B., Hofer, J., Schechner, Y. Y., Wandinger, U., and Pappalardo, G.: The unprecedented 2017–2018 stratospheric smoke event: decay phase and aerosol properties observed with the EARLINET, *Atmospheric Chemistry and Physics*, 19, 15 183–15 198, <https://doi.org/10.5194/acp-19-15183-2019>, <https://acp.copernicus.org/articles/19/15183/2019/>, 2019.
- Baars, H., Radenz, M., Floutsi, A. A., Engelmann, R., Althausen, D., Heese, B., Ansmann, A., Flament, T., Dabas, A., Trapon, D., Reitebuch, O., Bley, S., and Wandinger, U.: Californian Wildfire Smoke Over Europe: A First Example of the Aerosol Observing Capabilities of Aeolus Compared to Ground-Based Lidar, *Geophysical Research Letters*, 48, e2020GL092 194, <https://doi.org/https://doi.org/10.1029/2020GL092194>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2020GL092194>, e2020GL092194 2020GL092194, 2021.
- Berkemeier, T., Shiraiwa, M., Pöschl, U., and Koop, T.: Competition between water uptake and ice nucleation by glassy organic aerosol particles, *Atmospheric Chemistry and Physics*, 14, 12 513–12 531, <https://doi.org/10.5194/acp-14-12513-2014>, <https://acp.copernicus.org/articles/14/12513/2014/>, 2014.
- Bian, Q., Ford, B., Pierce, J. R., and Kreidenweis, S. M.: A Decadal Climatology of Chemical, Physical, and Optical Properties of Ambient Smoke in the Western and Southeastern United States, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD031 372, <https://doi.org/https://doi.org/10.1029/2019JD031372>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019JD031372>, e2019JD031372 2019JD031372, 2020.
- Brooks, S. D., Suter, K., and Olivarez, L.: Effects of chemical aging on the ice nucleation activity of soot and polycyclic aromatic hydrocarbon aerosols, *The Journal of Physical Chemistry A*, 118, 10 036–10 047, 2014.

- Burton, S., Hair, J., Kahnert, M., Ferrare, R., Hostetler, C., Cook, A., Harper, D., Berkoff, T., Seaman, S., Collins, J., et al.: Observations of the spectral dependence of linear particle depolarization ratio of aerosols using NASA Langley airborne High Spectral Resolution Lidar, *Atmospheric Chemistry and Physics*, 15, 13 453–13 473, 2015.
- Burton, S., Hostetler, C., Cook, A., Hair, J., Seaman, S., Scola, S., Harper, D., Smith, J., Fenn, M., Ferrare, R., et al.: Calibration of a high spectral resolution lidar using a Michelson interferometer, with data examples from ORACLES, *Applied optics*, 57, 6061–6075, 2018.
- China, S., Mazzoleni, C., Gorkowski, K., Aiken, A. C., and Dubey, M. K.: Morphology and mixing state of individual freshly emitted wildfire carbonaceous particles, *Nature Communications*, 4, <https://doi.org/10.1038/ncomms3122>, <https://doi.org/10.1038%2Fncomms3122>, 2013.
- China, S., Scarnato, B., Owen, R. C., Zhang, B., Ampadu, M. T., Kumar, S., Dzepina, K., Dziobak, M. P., Fialho, P., Perlinger, J. A., et al.: Morphology and mixing state of aged soot particles at a remote marine free troposphere site: Implications for optical properties, *Geophysical Research Letters*, 42, 1243–1250, 2015.
- Creek Fire: <https://www.insider.com/california-wildfires-generate-their-own-weather-via-thunderclouds-2020-9#:~:text=The%20particular%20pyrocumulonimbus%20cloud%20she,ever%20recorded%20in%20the%20US>, last accessed: 2021-05-05, 2020.
- Cziczko, D. J., Froyd, K. D., Hoose, C., Jensen, E. J., Diao, M., Zondlo, M. A., Smith, J. B., Twohy, C. H., and Murphy, D. M.: Clarifying the dominant sources and mechanisms of cirrus cloud formation, *Science*, 340, 1320–1324, 2013.
- Dahlkötter, F., Gysel, M., Sauer, D., Minikin, A., Baumann, R., Seifert, P., Ansmann, A., Fromm, M., Voigt, C., and Weinzierl, B.: The Pagami Creek smoke plume after long-range transport to the upper troposphere over Europe—aerosol properties and black carbon mixing state, *Atmospheric Chemistry and Physics*, 14, 6111–6137, 2014.
- Ditas, J., Ma, N., Zhang, Y., Assmann, D., Neumaier, M., Riede, H., Karu, E., Williams, J., Scharffe, D., Wang, Q., et al.: Strong impact of wildfires on the abundance and aging of black carbon in the lowermost stratosphere, *Proceedings of the National Academy of Sciences*, 115, E11 595–E11 603, 2018.
- Garofalo, L. A., Pothier, M. A., Levin, E. J., Campos, T., Kreidenweis, S. M., and Farmer, D. K.: Emission and evolution of submicron organic aerosol in smoke from wildfires in the western United States, *ACS Earth and Space Chemistry*, 3, 1237–1247, 2019.
- Garra, P., Maschowski, C., Liaud, C., Dieterlen, A., Trouvé, G., Le Calvé, S., Jaffrezo, J.-L., Leyssens, G., Schönnenbeck, C., Kohler, S., et al.: Fluorescence microscopy analysis of particulate matter from biomass burning: Polyaromatic Hydrocarbons as Main Contributors, *Aerosol Science and Technology*, 49, 1160–1169, 2015.
- Gialitaki, A., Tsekeri, A., Amiridis, V., Ceolato, R., Paulien, L., Kampouri, A., Gkikas, A., Solomos, S., Marinou, E., Haarig, M., et al.: Is the near-spherical shape the “new black” for smoke?, *Atmospheric Chemistry and Physics*, 20, 14 005–14 021, 2020.
- Grawe, S., Augustin-Bauditz, S., Hartmann, S., Hellner, L., Pettersson, J. B. C., Prager, A., Stratmann, F., and Wex, H.: The immersion freezing behavior of ash particles from wood and brown coal burning, *Atmospheric Chemistry and Physics*, 16, 13 911–13 928, <https://doi.org/10.5194/acp-16-13911-2016>, <https://acp.copernicus.org/articles/16/13911/2016/>, 2016.
- Haarig, M., Engelmann, R., Ansmann, A., Veselovskii, I., Whiteman, D. N., and Althausen, D.: 1064 nm rotational Raman lidar for particle extinction and lidar-ratio profiling: cirrus case study, *Atmospheric Measurement Techniques*, 9, 4269–4278, 2016.
- Haarig, M., Ansmann, A., Baars, H., Jimenez, C., Veselovskii, I., Engelmann, R., and Althausen, D.: Depolarization and lidar ratios at 355, 532, and 1064 nm and microphysical properties of aged tropospheric and stratospheric Canadian wildfire smoke, *Atmospheric Chemistry and Physics*, 18, 11 847–11 861, 2018.

- 505 Hodshire, A., Bian, Q., Ramnarine, E., Lonsdale, C., Alvarado, M., Kreidenweis, S., Jathar, S., and Pierce, J.: More than emissions and chemistry: Fire size, dilution, and background aerosol also greatly influence near-field biomass burning aerosol aging, *Journal of Geophysical Research: Atmospheres*, 124, 5589–5611, 2019.
- Hodshire, A. L., Ramnarine, E., Akherati, A., Alvarado, M. L., Farmer, D. K., Jathar, S. H., Kreidenweis, S. M., Lonsdale, C. R., Onasch, T. B., Springston, S. R., et al.: Dilution impacts on smoke aging: evidence in Biomass Burning Observation Project (BBOP) data, *Atmospheric Chemistry and Physics*, 21, 6839–6855, 2021.
- 510 Hu, Q.: Advanced aerosol characterization using sun/sky photometer and multi-wavelength Mie-Raman lidar measurements, Ph.D. thesis, Lille 1, 2018.
- Hu, Q., Goloub, P., Veselovskii, I., Bravo-Aranda, J.-A., Popovici, I. E., Podvin, T., Haefelin, M., Lopatin, A., Dubovik, O., Pietras, C., et al.: Long-range-transported Canadian smoke plumes in the lower stratosphere over northern France, *Atmospheric Chemistry and Physics*, 19, 1173–1193, 2019.
- 515 Jahl, L. G., Brubaker, T. A., Polen, M. J., Jahn, L. G., Cain, K. P., Bowers, B. B., Fahy, W. D., Graves, S., and Sullivan, R. C.: Atmospheric aging enhances the ice nucleation ability of biomass-burning aerosol, *Science Advances*, 7, eabd3440, 2021.
- Jahn, L. G., Polen, M. J., Jahl, L. G., Brubaker, T. A., Somers, J., and Sullivan, R. C.: Biomass combustion produces ice-active minerals in biomass-burning aerosol and bottom ash, *Proceedings of the National Academy of Sciences*, 117, 21 928–21 937, 2020.
- 520 Jarraud, M.: Guide to meteorological instruments and methods of observation (WMO-No. 8), World Meteorological Organisation: Geneva, Switzerland, 29, 2008.
- Jolly, W. M., Cochrane, M. A., Freeborn, P. H., Holden, Z. A., Brown, T. J., Williamson, G. J., and Bowman, D. M.: Climate-induced variations in global wildfire danger from 1979 to 2013, *Nature communications*, 6, 1–11, 2015.
- Kanji, Z. A., DeMott, P. J., Möhler, O., and Abbatt, J. P. D.: Results from the University of Toronto continuous flow diffusion chamber at ICIS 2007: instrument intercomparison and ice onsets for different aerosol types, *Atmospheric Chemistry and Physics*, 11, 31–41, <https://doi.org/10.5194/acp-11-31-2011>, <https://acp.copernicus.org/articles/11/31/2011/>, 2011.
- 525 Kanji, Z. A., Ladino, L. A., Wex, H., Boose, Y., Burkert-Kohn, M., Cziczo, D. J., and Krämer, M.: Overview of Ice Nucleating Particles, *Meteorological Monographs*, 58, 1.1–1.33, <https://doi.org/10.1175/AMSMONOGRAPHS-D-16-0006.1>, <https://doi.org/10.1175/AMSMONOGRAPHS-D-16-0006.1>, 2017.
- 530 Kleinman, L. I., Sedlacek III, A. J., Adachi, K., Buseck, P. R., Collier, S., Dubey, M. K., Hodshire, A. L., Lewis, E., Onasch, T. B., Pierce, J. R., et al.: Rapid evolution of aerosol particles and their optical properties downwind of wildfires in the western US, *Atmospheric Chemistry and Physics*, 20, 13 319–13 341, 2020.
- Knopf, D. A., Alpert, P. A., and Wang, B.: The role of organic aerosol in atmospheric ice nucleation: a review, *ACS Earth and Space Chemistry*, 2, 168–202, <https://doi.org/10.1021/acsearthspacechem.7b00120>, 2018.
- 535 Koop, T., Bookhold, J., Shiraiwa, M., and Pöschl, U.: Glass transition and phase state of organic compounds: dependency on molecular properties and implications for secondary organic aerosols in the atmosphere, *Phys. Chem. Chem. Phys.*, 13, 19 238–19 255, <https://doi.org/10.1039/C1CP22617G>, <http://dx.doi.org/10.1039/C1CP22617G>, 2011.
- Kütz, S. and Schmidt-Ott, A.: Characterization of agglomerates by condensation-induced restructuring, *Journal of aerosol science*, 23, 357–360, 1992.
- 540 Laa, A., Stein Zweers, D. C., and Boers, R.: A solar escalator: Observational evidence of the self-lifting of smoke and aerosols by absorption of solar radiation in the February 2009 Australian Black Saturday plume, *Journal of Geophysical Research: Atmospheres*, 117, 2012.

- Lee, H. J., Laskin, A., Laskin, J., and Nizkorodov, S. A.: Excitation–emission spectra and fluorescence quantum yields for fresh and aged biogenic secondary organic aerosols, *Environmental science & technology*, 47, 5763–5770, 2013.
- Lienhard, D. M., Huisman, A. J., Krieger, U. K., Rudich, Y., Marcolli, C., Luo, B., Bones, D. L., Reid, J. P., Lambe, A. T., Canagaratna, M. R.,
545 et al.: Viscous organic aerosol particles in the upper troposphere: diffusivity-controlled water uptake and ice nucleation?, *Atmospheric Chemistry and Physics*, 15, 13 599–13 613, 2015.
- Liu, L. and Mishchenko, M. I.: Scattering and Radiative Properties of Morphologically Complex Carbonaceous Aerosols: A Systematic Modeling Study, *Remote Sensing*, 10, <https://doi.org/10.3390/rs10101634>, <https://www.mdpi.com/2072-4292/10/10/1634>, 2018.
- Liu, L. and Mishchenko, M. I.: Spectrally dependent linear depolarization and lidar ratios for nonspherical smoke aerosols, *Journal of*
550 *Quantitative Spectroscopy and Radiative Transfer*, 248, 106 953, 2020.
- Lupi, L. and Molinero, V.: Does hydrophilicity of carbon particles improve their ice nucleation ability?, *The Journal of Physical Chemistry A*, 118, 7330–7337, 2014.
- Ma, X., Zangmeister, C. D., Gigault, J., Mulholland, G. W., and Zachariah, M. R.: Soot aggregate restructuring during water processing, *Journal of Aerosol Science*, 66, 209–219, 2013.
- 555 Mahrt, F., Alpert, P. A., Dou, J., Grönquist, P., Arroyo, P. C., Ammann, M., Lohmann, U., and Kanji, Z. A.: Aging induced changes in ice nucleation activity of combustion aerosol as determined by near edge X-ray absorption fine structure (NEXAFS) spectroscopy, *Environmental Science: Processes & Impacts*, 22, 895–907, 2020.
- May, A., McMeeking, G., Lee, T., Taylor, J., Craven, J., Burling, I., Sullivan, A., Akagi, S., Collett Jr, J., Flynn, M., et al.: Aerosol emissions from prescribed fires in the United States: A synthesis of laboratory and aircraft measurements, *Journal of Geophysical Research: Atmospheres*, 119, 11–826, 2014.
560
- McCluskey, C. S., DeMott, P. J., Prenni, A. J., Levin, E. J., McMeeking, G. R., Sullivan, A. P., Hill, T. C., Nakao, S., Carrico, C. M., and Kreidenweis, S. M.: Characteristics of atmospheric ice nucleating particles associated with biomass burning in the US: Prescribed burns and wildfires, *Journal of Geophysical Research: Atmospheres*, 119, 10 458–10 470, 2014.
- Mikhailov, E., Vlasenko, S., Martin, S., Koop, T., and Pöschl, U.: Amorphous and crystalline aerosol particles interacting with water vapor: conceptual framework and experimental evidence for restructuring, phase transitions and kinetic limitations, *Atmospheric Chemistry and Physics*, 9, 9491–9522, 2009.
565
- Müller, D., Mattis, I., Ansmann, A., Wandinger, U., Ritter, C., and Kaiser, D.: Multiwavelength Raman lidar observations of particle growth during long-range transport of forest-fire smoke in the free troposphere, *Geophysical Research Letters*, 34, 2007.
- Murayama, T., Müller, D., Wada, K., Shimizu, A., Sekiguchi, M., and Tsukamoto, T.: Characterization of Asian dust and Siberian smoke with multi-wavelength Raman lidar over Tokyo, Japan in spring 2003, *Geophysical Research Letters*, 31, L23 103, <https://doi.org/10.1029/2004GL021105>, 2004.
570
- O’Dell, K., Hornbrook, R. S., Permar, W., Levin, E. J., Garofalo, L. A., Apel, E. C., Blake, N. J., Jarnot, A., Pothier, M. A., Farmer, D. K., et al.: Hazardous air pollutants in fresh and aged western US wildfire smoke and implications for long-term exposure, *Environmental Science & Technology*, 54, 11 838–11 847, 2020.
- 575 Ohneiser, K., Ansmann, A., Baars, H., Seifert, P., Barja, B., Jimenez, C., Radenz, M., Teisseire, A., Floutsi, A., Haarig, M., et al.: Smoke of extreme Australian bushfires observed in the stratosphere over Punta Arenas, Chile, in January 2020: optical thickness, lidar ratios, and depolarization ratios at 355 and 532 nm, *Atmospheric Chemistry and Physics*, 20, 8003–8015, 2020.
- Oregon Fire: <https://wildfiretoday.com/2020/09/10/wildfires-have-burned-over-800-square-miles-in-oregon/>, last accessed: 2021-05-05, 2020.

- 580 Ortiz-Amezcuca, P., Guerrero-Rascado, J. L., Granados-Muñoz, M. J., Benavent-Oltra, J. A., Böckmann, C., Samaras, S., Stachlewska, I. S., Janicka, L., Baars, H., Bohlmann, S., et al.: Microphysical characterization of long-range transported biomass burning particles from North America at three EARLINET stations, *Atmospheric Chemistry and Physics*, 17, 5931–5946, 2017.
- Peterson, D. A., Campbell, J. R., Hyer, E. J., Fromm, M. D., Kablick, G. P., Cossuth, J. H., and DeLand, M. T.: Wildfire-driven thunderstorms cause a volcano-like stratospheric injection of smoke, *NPJ climate and atmospheric science*, 1, 1–8, 2018.
- 585 Petters, M. D., Parsons, M. T., Prenni, A. J., DeMott, P. J., Kreidenweis, S. M., Carrico, C. M., Sullivan, A. P., McMeeking, G. R., Levin, E., Wold, C. E., et al.: Ice nuclei emissions from biomass burning, *Journal of Geophysical Research: Atmospheres*, 114, 2009.
- Prenni, A. J., DeMott, P. J., Sullivan, A. P., Sullivan, R. C., Kreidenweis, S. M., and Rogers, D. C.: Biomass burning as a potential source for atmospheric ice nuclei: Western wildfires and prescribed burns, *Geophysical Research Letters*, 39, 2012.
- Price, H. C., Mattsson, J., Zhang, Y., Bertram, A. K., Davies, J. F., Grayson, J. W., Martin, S. T., O’Sullivan, D., Reid, J. P., Rickards, A. M., 590 et al.: Water diffusion in atmospherically relevant α -pinene secondary organic material, *Chemical science*, 6, 4876–4883, 2015.
- Reid, J. P., Bertram, A. K., Topping, D. O., Laskin, A., Martin, S. T., Petters, M. D., Pope, F. D., and Rovelli, G.: The viscosity of atmospherically relevant organic particles, *Nature communications*, 9, 1–14, 2018.
- Reid, J. S., Koppmann, R., Eck, T. F., and Eleuterio, D. P.: A review of biomass burning emissions part II: intensive physical properties of biomass burning particles, *Atmospheric Chemistry and Physics*, 5, 799–825, <https://doi.org/10.5194/acp-5-799-2005>, <https://acp.copernicus.org/articles/5/799/2005/>, 2005.
- 595 Santín, C. and Doerr, S. H.: Fire effects on soils: the human dimension, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 371, 20150171, 2016.
- Schill, G. P., DeMott, P. J., Emerson, E. W., Rauker, A. M. C., Kodros, J. K., Suski, K. J., Hill, T. C. J., Levin, E. J. T., Pierce, J. R., Farmer, D. K., and Kreidenweis, S. M.: The contribution of black carbon to global ice nucleating particle concentrations relevant to mixed-phase clouds, *Proceedings of the National Academy of Sciences*, 117, 22705–22711, <https://doi.org/10.1073/pnas.2001674117>, <https://www.pnas.org/content/117/37/22705>, 2020.
- 600 Schoennagel, T., Balch, J. K., Brenkert-Smith, H., Dennison, P. E., Harvey, B. J., Krawchuk, M. A., Mietkiewicz, N., Morgan, P., Moritz, M. A., Rasker, R., et al.: Adapt to more wildfire in western North American forests as climate changes, *Proceedings of the National Academy of Sciences*, 114, 4582–4590, 2017.
- 605 Shiraiwa, M., Li, Y., Tsimpidi, A. P., Karydis, V. A., Berkemeier, T., Pandis, S. N., Lelieveld, J., Koop, T., and Pöschl, U.: Global distribution of particle phase state in atmospheric secondary organic aerosols, *Nature communications*, 8, 1–7, 2017.
- Torres, O., Bhartia, P. K., Taha, G., Jethva, H., Das, S., Colarco, P., Krotkov, N., Omar, A., and Ahn, C.: Stratospheric injection of massive smoke plume from Canadian boreal fires in 2017 as seen by DSCOVR-EPIC, CALIOP, and OMPS-LP observations, *Journal of Geophysical Research: Atmospheres*, 125, e2020JD032579, 2020.
- 610 Umo, N. S., Murray, B. J., Baeza-Romero, M. T., Jones, J. M., Lea-Langton, A. R., Malkin, T. L., O’Sullivan, D., Neve, L., Plane, J. M. C., and Williams, A.: Ice nucleation by combustion ash particles at conditions relevant to mixed-phase clouds, *Atmospheric Chemistry and Physics*, 15, 5195–5210, <https://doi.org/10.5194/acp-15-5195-2015>, <https://acp.copernicus.org/articles/15/5195/2015/>, 2015.
- Umo, N. S., Wagner, R., Ullrich, R., Kiselev, A., Saathoff, H., Weidler, P. G., Cziczo, D. J., Leisner, T., and Möhler, O.: Enhanced ice nucleation activity of coal fly ash aerosol particles initiated by ice-filled pores, *Atmospheric Chemistry and Physics*, 19, 8783–8800, 615 <https://doi.org/10.5194/acp-19-8783-2019>, <https://acp.copernicus.org/articles/19/8783/2019/>, 2019.
- Urbanski, S. P., Hao, W. M., and Baker, S.: Chemical composition of wildland fire emissions, *Developments in environmental science*, 8, 79–107, 2008.

- Vergara-Temprado, J., Holden, M. A., Orton, T. R., O'Sullivan, D., Umo, N. S., Browse, J., Reddington, C., Baeza-Romero, M. T., Jones, J. M., Lea-Langton, A., et al.: Is black carbon an unimportant ice-nucleating particle in mixed-phase clouds?, *Journal of Geophysical Research: Atmospheres*, 123, 4273–4283, 2018.
- 620 Veselovskii, I., Hu, Q., Goloub, P., Podvin, T., Korenskiy, M., Pujol, O., Dubovik, O., and Lopatin, A.: Combined use of Mie–Raman and fluorescence lidar observations for improving aerosol characterization: feasibility experiment, *Atmospheric Measurement Techniques*, 13, 6691–6701, 2020.
- Veselovskii, I., Hu, Q., Ansmann, A., Goloub, P., Podvin, T., and Korenskiy, M.: Fluorescence lidar observations of wildfire smoke inside cirrus: A contribution to smoke-cirrus – interaction research, *Atmospheric Chemistry and Physics Discussions*, 2021, 1–29, <https://doi.org/10.5194/acp-2021-1017>, <https://acp.copernicus.org/preprints/acp-2021-1017/>, 2021a.
- 625 Veselovskii, I., Hu, Q., Goloub, P., Podvin, T., Choël, M., Visez, N., and Korenskiy, M.: Mie–Raman–fluorescence lidar observations of aerosols during pollen season in the north of France, *Atmospheric Measurement Techniques*, 14, 4773–4786, <https://doi.org/10.5194/amt-14-4773-2021>, <https://amt.copernicus.org/articles/14/4773/2021/>, 2021b.
- 630 Wang, B., Lambe, A. T., Massoli, P., Onasch, T. B., Davidovits, P., Worsnop, D. R., and Knopf, D. A.: The deposition ice nucleation and immersion freezing potential of amorphous secondary organic aerosol: Pathways for ice and mixed-phase cloud formation, *Journal of Geophysical Research: Atmospheres*, 117, 2012.
- Wu, G., Wan, X., Gao, S., Fu, P., Yin, Y., Li, G., Zhang, G., Kang, S., Ram, K., and Cong, Z.: Humic-like substances (HULIS) in aerosols of central Tibetan Plateau (Nam Co, 4730 m asl): Abundance, light absorption properties, and sources, *Environmental science & technology*, 52, 7203–7211, 2018.
- 635 Yu, P., Toon, O. B., Bardeen, C. G., Zhu, Y., Rosenlof, K. H., Portmann, R. W., Thornberry, T. D., Gao, R.-S., Davis, S. M., Wolf, E. T., de Gouw, J., Peterson, D. A., Fromm, M. D., and Robock, A.: Black carbon lofts wildfire smoke high into the stratosphere to form a persistent plume, *Science*, 365, 587–590, <https://doi.org/10.1126/science.aax1748>, <https://doi.org/10.1126%2Fscience.aax1748>, 2019.
- Zhang, Y., Wang, L., Liu, P., Li, Y., Zhan, R., Huang, Z., and Lin, H.: Measurement and extrapolation modeling of PAH laser-induced fluorescence spectra at elevated temperatures, *Applied Physics B*, 125, 1–12, 2019.
- 640

Table 1. The averaging time intervals and altitude ranges of selected BBA layers detected by LILAS system for the analysis of statistics. P1, P2, ..., P9 represent different time intervals in Lille September 2020.

NO of period	Time interval [UTC]	Height [m]
P1	2020-09-11, 02:00:00 – 2020-09-11, 04:00:00	5500 – 7500
P2	2020-09-11, 22:00:00 – 2020-09-12, 02:03:00:00	5500 – 8500
P3	2020-09-12, 20:00:00 – 2020-09-12, 23:00:00	2500 – 3800
P4	2020-09-14, 00:00:00 – 2020-09-14, 03:00:00	5500 – 7500
P5	2020-09-14, 22:00:00 – 2020-09-15, 02:00:00	5500 – 6500
P6	2020-09-15, 02:00:00 – 2020-09-15, 04:00:00	4500 – 7200
P7	2020-09-17, 22:00:00 – 2020-09-18, 03:00:00	5400 – 8500
P8	2020-09-18, 20:00:00 – 2020-09-18, 22:00:00	4800 – 6500
P9	2020-09-20, 20:00:00 – 2020-09-21, 00:00:00	4400 – 6200




Figure 1. True color images from OMPS onboard Suomi NPP. (a) 06 September 2020, (b) 10 September 2020, (c) 12 September 2020. The two AERONET observation sites: NASA_Ames (37.420N, 122W) and PNNL (46N, 119W). ©Google Maps




Figure 2. The backscattering coefficients and PLDR ratio at 532 nm measured by CALIPSO. The smoke plumes in (a), (b) and (c) are attributed to Creek fires on 05-07 September 2020, and in (d), (e) and (f) are attributed to Oregon fires. The inserted maps display the granules of CALIPSO and the locations of the grey dots represent the region where the measurements are averaged. (a) Central US, 2020-09-08. (b) East of Quebec, 2020-09-09. (c) Eastern Europe, 2020-09-11. (d) Western coast of US, 2020-09-10. (e) Western coast of US, 2020-09-11. (f) Eastern US, 2020-09-14. The layer between the two red dashed lines is BBA. The blue dashed lines represent the location of the mixture of ice crystals and BBA particles.




Figure 3. AOD and AE (Level 1.5) measured by AERONET sun/photometers at two observation sites: (a) PNNL (46N, 119W) and (b) NASA_Ames (37.420N, 122W). NASA_Ames is within the influence of both the Creek fire and the Oregon fire. PNNL is mainly impacted by the smoke of Oregon fire. The smoke of Oregon fire arrived at NASA_Ames on about 9 September and at PNNL on 12 September 2020. The arrival of the Oregon smoke caused significant increase of AOD and decrease of the AE at both sites.

The backscattering coefficients and PLDR ratio at 532-nm measured by CALIPSO. The smoke plumes in (a), (b) and (c) are attributed to Creek fires on 05-07 September 2020, and in (d), (e) and (f) are attributed to Oregon fires. The inserted maps display the granules of CALIPSO and the locations of the grey dots represent the region where the measurements are averaged. (a) Central US, 2020-09-08. (b) East-of-Quebec, 2020-09-09. (c) Eastern Europe, 2020-09-11. (d) Western coast of US, 2020-09-10. (e) Western coast of US, 2020-09-11. (f) Eastern US, 2020-09-14.




Figure 4. Lidar and sun/sky photometer observations during the smoke episodes on 10-22 September 2020, LilleATOLL, France. (a) The range-corrected lidar signal at 1064 nm. (b) The volume depolarization ratio at 1064 nm. (c) The AOD and AE measured by the sun/sky photometer operated in Lilleat ATOLL. The layers with strong depolarization appearing in the boundary layer on 14-16 September are polluted dust from Saharan region. The smoke layers from Californian smoke are mostly distributed in the free troposphere.




Figure 5. Box plots of the variabilities of BBA intensive parameters during the smoke episode episodes in September 2020. P1, P2, ..., P9 refer to the time intervals defined in Table 1. Only nighttime observations are considered in this analysis. The measurements in the time intervals are averaged The lower and upper edges of the box represents respectively the first and third quartiles of data. The lower and upper range-bars represent respectively the minimum and maximum. The dot and bar in the box indicate the mean value and the middle-median value, respectively. (a) Lidar ratios, (b) PLDRs. The colors of the plots represent the different wavelengths (magenta: 355 nm, green: 532 nm, red: 1064 nm). (c) fluorescence capacity (in cyan), (d) EAE₃₅₅₋₅₃₂ (in blue) and BAE₃₅₅₋₅₃₂ (in orange).




Figure 6. HYSPLIT back trajectories overlaid on the UVAI measured by OMPS onboard SUOMI-NPP satellite. (a) 96-hour back trajectories ending at 00:00 UTC, 12 September 2020 overlaid with UVAI on 08 September. (b) 144-hour back trajectories ending at 00:00 UTC, 18 September 2020 overlaid with UVAI on 14 September. ©Google Earth.




Figure 7. Lidar observation observations on 11-12 September 2020. (a) Range corrected lidar signal at 1064 nm, (b) volume depolarization ratio at 532-1064 nm, (c) backscatter coefficient of fluorescence at calibrated signal ratio between 466 and 387 nm channels, i.e. $\frac{C_R P_{466}}{C_F P_{387}}$ in Equation (1). This quantity is used as intermediate parameter to represent the fluorescence signal. The calibration coefficient C_R/C_F represents the ratio of the electro-optical gain between the 2 channels. The white zone in the figures represents missing data due to the turnoff of the detectors (ex. to avoid daytime sunlight interferences) or intermediate operations during the measurements.




Figure 8. Aerosol vertical profiles from averaged measurements between 22:00, 11 September 2020 and 02:03:00, 12 September 2020, UTC. (a) Extinction coefficients, (b) backscattering coefficient, (c) lidar ratios. The colors of lines represent different wavelengths: 355 (magenta), 532 (green) and 1064 nm (red). (d) $EAE_{355-532}$ (orange), $BAE_{355-532}$ (blue) and temperature (cyan open circle), (e) PLDRs, (f) fluorescence backscattering coefficient (pink) and fluorescence capacity (cyan), (g) WVMR (pink solid circle), RH to liquid water (RH, pink solid circle) and ice (RH_{ice} , grey solid circle). The extinction and backscatter coefficients at 355 and 532 nm were calculated using Raman technique (Ansmann et al., 1992) and the backscatter coefficient at 1064 nm was calculated by Klett method with an assumption of $LR=50$ sr. The radio sounding station is in Beauvechain/Beauvechain, Belgium, about 100 km from the lidar observation site. The error bars in the figure represent the statistical errors. The method of error estimation is presented in the Appendix of (Hu et al., 2019). The relative humidity to ice is calculated using the improved Magnus formula in Alduchov and Eskridge (1996).




Figure 9. Lidar The same plot as Figure 7 but for lidar observation on 17-18 September 2020. (a) Range corrected lidar signal at 1064 nm, (b) volume depolarization ratio at 1064 nm, (c) backscatter coefficient of fluorescence at calibrated signal ratio between 466 and 387 nm channels.




Figure 10. Aerosol vertical profiles from averaged measurements The same as Figure 8 except for observations between 22:30, 17 September 2020 and 03:00, 18 September 2020, UTC. The calculation is done with the same method as in Figure 8. The radio sounding measurements are from Hertzmonceux, England, which is about 200 km from the observation site. Radio sounding at Beauvechain, Belgium is not available on this day. The error bars represent the statistical errors, the same as in Figure 8.




Figure 11. ~~Averaged~~ The profiles of averaged parameters in two consecutive periods: 20:30–22:30 UTC, 17 September 2020 (before the observation of ice crystals and in lighter colors) and 22:30–23:30 UTC, 17 September 2020 (during ice crystal observation and in darker colors). The dash-dotted line represents the position of altitude where the ice crystals appeared. (a) Backscattering coefficient at 532 nm, (b) fluorescence capacity at 466 nm, (c) fluorescence backscattering coefficient, (d) PLDR at 1064 nm.