

The semi-annual oscillation (SAO) in the upper troposphere and lower stratosphere (UTLS)

Ming Shangguan¹ and Wuke Wang²

¹School of Geography and Information Engineering, China University of Geosciences, Wuhan, China

²Department of Atmospheric Science, China University of Geosciences, Wuhan, China

Correspondence: Wuke Wang (wangwuke@cug.edu.cn)

Abstract. Both the scientific and operational communities are increasingly interested in subseasonal to seasonal variations of weather and climate. The Semi-Annual Oscillation (SAO) has been studied extensively at the surface as well as in the middle atmosphere (upper stratosphere and the lower mesosphere). However, the SAO in the upper troposphere and lower stratosphere (UTLS) has been less discussed. Here we find evident SAO of temperature in the UTLS (250-175 hPa) from the subtropics to mid-latitudes (22.5-42.5°) using high-quality satellite measurements, reanalysis data, and model simulations. We show the mechanism of its formation by an energy budget analysis. The temperature in the Northern Hemisphere (NH) UTLS shows the first peak in February according to the dynamical heating and shows the second peak in July due to the dynamical heating and moist processes. Similar to the NH, the austral winter time maximum of temperature in the SH is related to dynamical heating and the austral summer time maximum is related to both moist and dynamical heating in the UTLS. Model simulations indicate that the SAO in the UTLS is partly modified by the SAO of sea surface temperatures (SSTs).

1 Introduction

Subseasonal to seasonal predictions of weather and climate are increasingly important due to the urgent requirement from decision makers (Merryfield et al., 2020). As an important component of subseasonal to seasonal variations, the Semi-Annual Oscillation (SAO) has been well known at the surface (surface-SAO) (Meehl et al., 1998) as well as in the middle atmosphere (MA-SAO) (Garcia et al., 1997). It is well known that the annual cycle of temperature gradient and mean sea level pressure is dominated by a strong half-yearly oscillation between 50°S and 65°S, which exceeds the magnitude of the yearly wave in many locations in the SH near the surface (Walland and Simmonds, 1998; Simmonds and Jones, 1998). This surface-SAO is a coupled ocean-atmosphere phenomenon, and a change of the seasonal cycle of Sea Surface Temperatures (SSTs) at 50°S could alter the amplitude of the surface-SAO (Meehl et al., 1998). Bracegirdle (2011) showed that this surface SAO is coupled with the stratospheric circulation, and correlations are found over a large altitude range in the troposphere and stratosphere. Yang and Wu (2022) relate the semiannual surface air response to changes in the oceanic mixed layers. The SAO is also noticeable as the dominating mode of wind and temperature variations in the tropical middle atmosphere between the middle stratosphere and the upper mesosphere (Read, 1962; Garcia et al., 1997). This MA-SAO is a complex interplay of momentum advection,

planetary waves from the extratropics, and vertically propagating equatorial waves including both global-scale waves and
25 small-scale gravity waves (Hamilton and Mahlman, 1988; Richter and Garcia, 2006; Ern et al., 2015, 2021).

In comparison, only a few studies focus on the SAO in the upper troposphere and lower stratosphere (UTLS). The SAO in
temperature is reported by Loon (1967) over the middle troposphere in the SH with two maxima in March and September near
the equinoxes, which is related to varying heating/cooling rates in different latitude bands. Loon and Jenne (1969) described the
tropical-subtropical SAO of the zonal wind and temperature in the SH based on data of the 20 radiosonde stations in the upper
30 troposphere (100-300 hPa). They also found that the amplitude of SAO in temperature and wind are different along different
longitudes, i.e., with stronger signals in the eastern hemisphere (Loon and Jenne, 1969; Chen and Tsay, 2014). According to
Loon and Jenne (1969), the temperature oscillation is the result of an intensification of vertical motions from autumn to winter
and the zonal wind oscillation is associated with second harmonics of opposite phase in the temperature. Shea et al. (1995) made
a further study on the Tropical-Subtropical Semi-Annual Oscillation (TS-SAO) in the upper troposphere with more extensive
35 data coverage. Shea et al. (1995) found that the TS-SAO of temperature, which has maxima in the transitional seasons in the
tropics and peaks in the extreme seasons in the subtropics, resulted from a marked semi-annual variation of the winds in the
tropics with maxima in the inter-monsoonal months. The precise mechanism for this oscillation seems somewhat unclear, but
may be associated with thermal fluctuations connected with the movement of the intertropical convergence zone from one side
of the Equator to the other and associated modulations of the mean meridional circulation (Read and Castrejón-Pita, 2012).

40 While measurements in the UTLS are relatively sparse, reanalysis data are widely used to investigate temperature variabil-
ities (Broeke, 2000; Fueglistaler et al., 2009; Gettelman et al., 2011; Wang et al., 2016; Shangguan et al., 2019). Despite the
importance of the SAO in the mid-latitudes, there have been few articles focused on it since that of Loon (1967), and these
have been mostly based on radiosonde and reanalysis data. With the development of satellite techniques, many new remote
sensing data are available in the UTLS. According to many studies (Wickert et al., 2001, 2009; Ho et al., 2017), the Global
45 Navigation Satellite System Radio Occultation (GNSS RO) can provide highly accurate temperature profiles from the middle-
upper troposphere to lower stratosphere by measuring the time delay in occulted signals from one satellite to another. The first
RO mission (Challenging Minisatellite Payload (CHAMP)) was launched in 2001, and the Constellation Observing System for
Meteorology, Ionosphere and Climate (COSMIC), which is a constellation of six satellites since late 2006 is widely applied in
the climate research (Ho et al., 2014; Randel and Wu, 2015; Gao et al., 2017). Compared with radiosonde data, the GNSS RO
50 data have global coverage (both in continents and oceans) and are not affected by weather (Anthes et al., 2008).

In this study, we use the GNSS RO data and the latest ECMWF reanalysis ERA5 (Hersbach et al., 2019) and the NASA
MERRA2 reanalysis (GMAO, 2015a, b) to study the SAO in the UTLS (UTLS-SAO), and to discuss its origin. We expand the
oscillation using temperature data available from GNSS RO and reanalyses between 60°N to 60°S spanning the period 2001-
2017, and using Hadley Centre SST data set (Rayner et al., 2003) for a spatial coherence picture in the same period. The latest
55 ERA5 and MERRA2, which include more recent instrument observations and improved data assimilation methods, should
provide more reliable estimates than earlier studies. PSD (Power Spectrum Densities) analysis is used to analyze the time-scale
of temporal variability and a significance test is used to diagnose whether the SAO signal is significant and not an artifact
e.g., just a harmonic part of the seasonal cycle. The SAO variations investigated in this study are more sinus-like. The energy

budget, i.e., the heating rates according to dynamical, radiative, and moist processes, are analyzed to explain the formation of the SAO in temperature. To understand the relationship between SAO and SSTs, three model simulations with NCAR's Whole Atmosphere Community Climate Model, version 6 (WACCM6) are used. Details of the data and model simulations are described in section 2. In section 3, we show the analysis and results. Conclusions are summarized in section 4.

2 Data and Methods

2.1 GNSS RO temperature data

CHAMP provides ca. 150 occultation events globally per day from May 2001 to October 2008, and COSMIC began providing 1000-3000 occultation events per day since late 2006 (Wickert et al., 2001; Anthes et al., 2008). Many studies have demonstrated that GNSS RO temperature data have good quality in the range 8-30 km (Schmidt et al., 2005, 2010; Ho et al., 2009, 2012). In our study, we make use of monthly mean temperature data at 500-10 hPa. About 100 observations per month per 5° latitude grid can be provided by a single satellite CHAMP and more than 10 times the number of profiles are available since late 2006 due to the start of the COSMIC mission. WetPrf products are interpolated onto 100 m vertical resolution from 0.1 to 40 km based on the one-dimensional variational method (Kursinski et al., 2000; Wee and Kuo, 2015). According to Xu et al. (2017) biases could exist for altitudes below ~5-8 km in the wetPrf. We use the reprocessed and post-processed RO data, which are stable and accurate observations for climate studies. The CHAMP wetPf2 version is 2016.2430 and the COSMIC-1 wetPrf products are 2013.3520 and 2016.1120.

GNSS RO monthly zonal means of standard pressure level (400-10 hPa) were determined based on CHAMP and COSMIC-1 for the period 2001-2017, for this, 5° nonoverlapping latitude bands centered at 57.5°S–57.5°N were used. Data exceeding 3 times the standard deviation have been discarded at each level. For the COSMIC-1 data, we make additional monthly zonal means of standard pressure levels with a grid resolution 10° (latitude)x10° (longitude) for the period 2007-2017. The same averaging strategy is used.

Due to the missing data in the early part of the GNSS RO record, a spectral analysis of temperature data was performed using a Lomb-Scargle periodogram to identify the various periods in the time series (Lomb, 1976; Scargle, 1982; Horne and Baliunas, 1986; Press and Rybicki, 1989). The Lomb-Scargle periodogram can handle the periodic signals with missing data. For the SAO we choose the maximum power spectral density (PSD) between 5 and 7 months and use the probability 0.95 of detection for the significance test (Horne and Baliunas, 1986). The significance test is used to check whether the peak is a true signal peak and not the result of random fluctuations.

2.2 Reanalysis data

ERA5 is the latest ECMWF reanalysis (released in 2018) and various newly reprocessed data sets, recent instruments, improved data assimilation system are used in ERA5. Detailed information can be found in ERA5 data documentation <https://confluence.ecmwf.int/display/CKB/ERA5%3A+data+documentation>. The ERA5 monthly averaged data on pressure levels from 2001 to

90 2017 with 2.5° horizontal resolution is downloaded from Climate Data Store (CDS)(Hersbach et al., 2019). MERRA2 is the latest atmospheric reanalysis of NASA’s Global Modeling and Assimilation Office (GMAO) with data resolution 0.5°x0.625° (Gelaro et al., 2017). Both reanalyses contain the eastward wind (u), northward wind (v), vertical pressure velocity (ω) and temperature (T) on pressure levels. Both reanalyses have assimilated GNSS RO bending angles.

2.3 Model simulations

95 WACCM6 is one of the two available atmospheric components of the Community Earth System Model (CESM) from NCAR. It simulates atmospheric processes from the surface to about 140 km, which resolves the stratospheric dynamical and chemical processes well (Marsh et al., 2013; Gettelman et al., 2019). The standard version of WACCM has 70 vertical levels with a vertical resolution of about 1 km in the UTLS region (Gettelman et al., 2019; Wang et al., 2019). The horizontal resolution used here is 1.9°x2.5°.

100 The model is integrated into its atmosphere-only mode, with prescribed SSTs. We first employ a Control simulation, with SSTs prescribed to historically observed values over the period from 2001 to 2017. The SSTs are provided by Hadley Centre SST data set (Rayner et al., 2003). For comparison, we then employ two sensitivity simulations, which removed the SAO of SSTs (SST-SAO) globally (rmSAO run) and in the Tropics (rmSAO-TP run). The SST-SAO is removed by a band-pass filter (signals between 5 and 7 months are removed) using the Butterworth method (Butterworth, 1930). The differences between
 105 these two sensitivity simulations and the Control simulation, therefore, indicate the influences of the SST-SAO on the UTLS-SAO.

2.4 Thermal budget analysis

To explain the SAO of temperature in the UTLS, the thermal budget is analyzed using the MERRA2 reanalysis as well as model simulations. The thermal budget of the UTLS is a balance between the dynamical heating and the total diabatic (Gettelman and
 110 Birner, 2007). The thermodynamic balance formalism is expressed as (Andrews et al., 1987; Abalos et al., 2013):

$$\bar{T}_t = -\bar{v}^* \bar{T}_y - \bar{w}^* S - e^{\frac{z}{H}} [e^{-\frac{z}{H}} (\overline{v'T'} \frac{\bar{T}_y}{S} + \overline{w'T'})]_z + \bar{Q} \quad (1)$$

$$\bar{v}^* = \bar{v} - \rho_0^{-1} [\rho_0 \overline{v'\Theta'} / \bar{\Theta}_z]_z \quad (2)$$

$$\bar{w}^* = \bar{w} + (a \cos \phi)^{-1} [\cos \phi \overline{v'\Theta'} / \bar{\Theta}_z]_\phi \quad (3)$$

where H is 7 km, ρ_0 is the atmosphere density, ϕ is latitude, overbars in the equation denote zonal means, primes indicate
 115 deviations from it and subscripts denote partial derivatives. \bar{v}^* and \bar{w}^* are components of the residual circulations, which can be calculated by eq.2-3. \bar{w} is the zonal mean vertical velocity, which is converted from the vertical pressure velocity ω in the reanalysis. Θ denotes potential temperature. $S = H \frac{N^2}{R}$ with $R = 287 m^2 s^{-2} K^{-1}$ and N^2 is the Brunt-Väisälä frequency. The first three terms on the right side of the eq.1 indicate the dynamical heating, while Q is the diabatic heating. The dynamical heating includes the heating by the meridional component of the residual circulation (the first term on the right side of eq.1), the
 120 heating/cooling by downwelling/upwelling (the second term on the right side of eq.1), and heating related to eddys (the third

term on the right side of eq.1, mainly associated with the vertical eddy heat flux). In the UTLS region, Q is mainly determined by radiation and moisture condensation processes.

MERRA2 provides very detailed temperature tendencies related to dynamics, radiation, moist, friction, gravity wave drag and near-surface turbulence. In the UTLS region, the friction, gravity wave drag and near-surface turbulence terms are very small, then we only use the dynamics, radiation and moist terms in this study. In addition, to keep a balance of the thermal budget during the data assimilation, an extra term called the analysis tendencies (ANA) is used in MERRA2 reanalysis (Mapes and Bacmeister, 2012). ANA can be interpreted as the negative of model physical tendency error. Therefore, the ANA can be considered as residual (Mapes and Bacmeister, 2012). We further calculate the first two terms in eq.1 with the given equations and then separate the dynamical term of MERRA2 into the meridional component of the residual circulation, the heating by downwelling and eddy terms.

In the WACCM6 model, the temperature tendencies related to dynamics, radiation, moist and gravity wave drag processes are also diagnosed and have direct outputs (Gettelman and Birner, 2007). Here we also use the dynamics, radiation and moist terms in this study. The radiation term consists of long-wave and short-wave terms in the simulation. The dynamical term is also separated into the meridional component of the residual circulation, the heating by downwelling and eddy terms with the same method as MERRA2.

3 Analysis and Results

3.1 Spatial distribution of the SAO

According to the Lomb-Scargle PSD values for the temperature data, the annual cycle is strongest in most of the regions. Figure 1 shows the SAO PSD in the period 2001-2017 and the lapse rate tropopause determined from GNSS RO data. The most prominent SAO signal with strong and significant SAO PSD can be seen in the UTLS region (250-175 hPa) from the subtropics (22.5°) to mid-latitudes (42.5°) in both hemispheres. This is also observed in the ratio of semiannual/annual cycle PSD (Figure S1). The ratio is larger than 0.6 in the UTLS (300-175 hPa) from the subtropics to mid-latitudes ($20-45^\circ$) and middle-upper troposphere (500-200 hPa) from tropics ($10^\circ\text{S}-10^\circ\text{N}$). The most prominent SAO PSDs can also be seen in the tropics from 400 to 225 hPa (Figure 1 and Figure S1). This is consistent with the results from Loon and Jenne (1969), which shows clear SAO of temperature in the upper troposphere in the tropics and SH subtropics. Our results give a survey of the SAO and find that the significant SAO also persists in the lower stratosphere and the mid-latitudes, i.e., 70-30 hPa in SH ($32.5^\circ\text{S}-47.5^\circ\text{S}$) and NH ($37.5^\circ\text{N}-47.5^\circ\text{N}$) mid-latitudes. It is also noteworthy that the magnitude of SAO PSD in the tropics in the troposphere and lower stratosphere are much weaker than the signals in mid-latitudes. We then mainly focus on the SAO in the UTLS region (250-175 hPa) in mid-latitudes ($32.5^\circ-42.5^\circ$) hereafter in this study.

The reanalysis data and model simulations show good agreement with GNSS RO for the general pattern of SAO PSD. The ERA5 data show the best agreement with the GNSS RO, with a very similar spatial pattern and comparable magnitude as shown in Figure 1a-b. The SAO signals in MERRA2 are slightly weaker and less significant than those in GNSS RO and ERA5

(Figure 1c and Figure S1c). The model also shows a good representation of the SAO in the UTLS, except that the significant region of SAO in the tropical regions is not observed from 350 to 200 hPa (Figure 1d and Figure S1d).

155 Further information can be gained from Figure 2, which shows the latitude-longitude distributions of the SAO PSD at 200 hPa. The significant SAO region at 200 hPa in mid-latitudes is larger in the Eastern than in the Western Hemisphere, which is consistent with previous studies (Loon and Jenne, 1969). Note that due to data limitation, the PSD shown in Figure 2a is calculated for the period 2007-2017. Further analysis indicates that the different period of analysis does not influence the results. As shown in Figure S2, the PSD distribution 2007-2017 from the ERA5, MERRA2 and the Control simulation is very
160 similar to that shown in Figure 2, which is analyzed for the period 2001-2017. Large SAO PSD occurs around the Asian region (25°N-45°N, 20°E-100°E), the Australian region (45°S-30°S, 30°E-180°E), as well as the North American around Great Lakes. The strong SAO signals over that region are likely related to both moist and dynamical processes. As seen in Figures S3 and S4, there are strong upward motion over south Asia but downward motion from north Africa to central Asia. The strong upwelling brings a large number of water vapor from the surface to the upper troposphere, which condenses there and heats
165 the atmosphere at around 200 hPa. At the same time, the strong downwelling leads to dynamical heating over the regions from north Africa to central Asia. A further energy budget analysis indicates that the vertical transportation of water vapor and subsequent condensation in the upper troposphere over the Asian region is much stronger than in other regions and leads to a stronger peak of temperature in the summer season (Figure S5). In addition, the radiative cooling is relatively weaker than in other regions, which also contributes to the stronger temperature peak in the Asia region (Figure S5). Again, the ERA5 has the
170 best agreement with GNSS RO measurements with a consistent pattern and comparable magnitude (Figure 2b). The MERRA2 and the model simulation also show good agreement with the GNSS RO data (Figure 2c-d). It is also noteworthy that there are also significant SAO signals in the tropics except over the Pacific. However, the magnitude of the SAO PSD in the tropics is much weaker than that in mid-latitudes. According to previous studies, the SAO of the tropical stationary eddies is caused by the semi-annual east-west seesaw of the global divergent circulation between the areas of the Asian-Australian (AA) monsoon
175 (60°E-120°W) and the extra-AA monsoon (120°W-60°E) particularly in SH tropics (Chen and Wu, 1992; Chen et al., 1996).

3.2 Time evolution and mechanism of the SAO in the UTLS

To analyze the time evolution of SAO signals, Figure 3 shows the time series in the zonal mean temperature and corresponding annual cycle anomalies in the NH mid-latitudes (NHM) (32.5°N-42.5°N) and SH Mid-latitudes (SHM) (32.5°S-42.5°S) at 200 hPa. From the time series, the SAO of temperature is evident in both hemispheres, with two clear peaks of temperature in each
180 year. Such oscillation can be seen in all the observed and reanalysis data sets, including the GNSS RO measurements as well as the ERA5 and MERRA2 reanalysis. The model simulations can also simulate the SAO of temperature except that an offset is of about 1 K between the Control simulation and the observation/reanalysis. From the annual cycle anomalies, two peaks of temperature exist in February and July in the NHM and in February and August in the SHM. The peak in July/August is more intense than in February, especially in the NHM. For all the datasets, the peaks in January-February and July-August are quite
185 even in NHM. In SHM the peak at August of simulations is weaker than observation/reanalysis.

To further investigate the mechanism of the SAO in the UTLS, the thermal budget is analyzed using the MERRA2 data and model simulations. The annual cycle of the heating rates at 200 hPa averaged over the latitude bands from 32.5° to 42.5° in the two hemispheres is shown in Figure 4. To show the relationship between the heating rates and the temperature, the annual cycle of temperature from the MERRA2 reanalysis data is also shown in Figure 4. In the NHM, there are two peaks of total heating rates in December and June and two nadirs in March and September (Figure 4a). While the total heating rate is positive, the temperature will increase. The total heating rates are positive from mid-November to mid-January and from April to July, which leads to increases in temperature during the same periods. From mid-January and July, the total heating rate turns to be negative, and the temperature starts to decrease. Therefore, there are two peaks of temperature in February and July. At the same time, there are two nadirs of temperature in April and November. To clearly show the relative contribution of different terms to the total heating rate, the variation of each term at peaks and nadirs of total heating rates is selected and shown in Figure 5a. Overall, the dynamical processes contribute the most prominent positive heating rates through all terms, which is mostly offset by the radiative cooling. The radiative term can further be separated into short-wave warming and long-wave cooling (Figure S6a). The short-wave and long-wave terms are all strongest during summer and the radiation heating rates are negative due to larger negative value of long-wave terms. The moist and analysis terms are quite small. For the first temperature peak in January-February, it is mainly related to the dynamical heating, since the dynamical warming is slightly stronger than the radiative cooling and other terms are near zero in December. The second temperature peak in July is caused by a combination of moist and dynamical processes. There is a secondary peak of dynamical heating in June, although it is offset by radiative cooling. The moist processes delay the transition of total heating rates from positive to negative, which is in July but would be a month earlier otherwise. This positive peak in moist heating is related to the release of heat by the condensation of water vapor that is transported by deep convections in the upper troposphere. The negative total heating rates in March and September are mainly determined by the strong radiative cooling, which is stronger than the sum of other terms.

In the SHM (Figure 4b and Figure 5b), the total heating rates are positive from mid-November to mid-January and from May to August, which leads to a first peak of temperature in February and the second peak in August. For the SH austral summer, although the annual variation of the moist heating is relatively weak, it is stronger in DJF than in other seasons. In addition, the dynamical heating is also relatively strong in DJF. Therefore, both the dynamical and moist processes contribute to the positive total heating rates and lead to the temperature peak in SH austral summer. For the SH austral winter, the positive total heating rate and subsequent temperature maximum are mainly related to dynamical heating. Note that the ANA term in the SHM is large, which indicates a poor ability of the MERRA2 model to represent the temperature variations in SHM UTLS. The moist processes are lower in the SHM than in the NHM, which is perhaps due to overall weaker monsoon circulation. The energy budget analysis can also be confirmed by model simulations (Figure S7). Figure S7 is the same as Figure 4 but for the Control simulation. The model shows a very similar annual variation of dynamical, radiative, moist and total heating rates in both the NHM and SHM compared to the MERRA2 reanalysis.

In Figure 6 the dynamical term is further divided into the eddy, heating by downwelling ($-wS$) and heating by the meridional component of the residual circulation ($-vT_y$) terms according to Eq.1. The eddy term is mainly associated with the vertical eddy heat flux. As seen from Figure 6, the $-vT_y$ term is close to zero, which is less important in the UTLS region than other

terms as also indicated by previous studies (Abalos et al., 2013). The $-wS$ term is strongest during boreal winter (DJF) in both hemispheres, since the vertical component of the residual circulation (including the tropical upwelling and the extra-tropical downwelling) is most prominent in DJF. This is consistent with previous studies which shows a strongest dynamical cooling according to the upwelling in the tropics (Abalos et al., 2013). In the SH, there is a secondary peak of the $-wS$ term in JJA. The eddy term is non-negligible in the upper troposphere (at 200 hPa) and peaks from April to June. The first peak in dynamical heating during boreal winter (DJF) is dominated by the $-wS$ term while the secondary peak near June is significantly influenced by the eddy term.

3.3 Relationship between the UTLS-SAO and the SSTs

As introduced in the sect. 1, there is also a pronounced SAO signal at the surface (Meehl et al., 1998). While the surface is the main energy source of the atmosphere, it would be very interesting to investigate the relationship between the UTLS-SAO and the Surface-SAO. Since the surface-SAO is a coupled ocean-atmosphere phenomenon and is strongly modified by the seasonal cycle of SSTs (Meehl et al., 1998), we mainly focus on the relationship between the UTLS-SAO and the SAO in SSTs (SST-SAO) in this study. We checked the PSD of the SST-SAO (Figure S8), and there are significant SAO signals in the tropics from the Eastern Pacific (120°W) to the Atlantic Ocean, Indian Ocean and the Western Pacific (150°E), which is also found by earlier studies (Schott et al., 2009; Park and Lee, 2014; Yan et al., 2018). The SST-SAO in the Pacific and Atlantic oceans in the tropics is related to the SAO of solar radiation (Yashayaev and Zveryaev, 2001), while the SST-SAO in the tropical Indian Ocean is related to the Indian monsoon (Hu et al., 2005). The SAO signals over the Central Pacific (150°E-120°W) are not significant (Figure S8), which resembles the UTLS-SAO in the tropics as shown in Figure 2. This indicates the potential connection between the SST-SAO and the UTLS-SAO in the tropics.

To identify the relationship between the subtropical to mid-latitude UTLS-SAO and the SST-SAO, the correlation coefficients are calculated between the SST-SAO and ULTS SAO in the tropical (5°S-5°N) and the SH/NH mid-latitude (Figure S9). Firstly, we extracted ERA5 temperature SAO signals (5-7 months) using the 1-D wavelet Morlet transform (Lilly and Olhede, 2009, 2012). Secondly, we calculated the Pearson correlation coefficient between SAO signals and SST-SAOs. As shown in Figure S9, both the SH/NH mid-latitude and tropical UTLS-SAO are strongly and significantly correlated with SST variations. The UTLS-SAO shows a positive correlation with SST-SAO in the tropics (5°S-5°N), especially over the Indian Ocean and the Western Pacific (Figure S9c), whereas the SST-SAO are most pronounced (Figure S8). The SH/NH mid-latitudes show very similar correlation patterns with SST-SAO (Figure S9a-b) since the UTLS-SAO in the two hemispheres are almost in phase with each other (Figure 3). All correlations are mainly negative in Figure S9a-b, which indicate there are global phase variations.

We then further investigate the impacts of the SST-SAO on UTLS-SAO using our model simulations. As introduced in Section 2.3, we first employ a Control simulation using the WACCM6 model with SSTs prescribed to observed variations. As seen in Figures 1-3, the model represents the UTLS-SAO pretty well although the simulated temperature is ca. 1 K lower than observations. We then employ a sensitive simulation with the same configuration except that the SST-SAO has been removed globally (the rmSAO run). As discussed above, SST-SAO is mainly significant in the tropics (Figure S8), and the UTLS-SAO

255 is most correlated with tropical SSTs, therefore we employ a third simulation with the SST-SAO removed only in the tropics (the rmSAO-TP simulation).

Figure 7 shows the PSD of SAO in temperature from the rmSAO and rmSAO-TP runs, as well as their relative differences with respect to the Control simulation. Similar to the Control simulation, there are significant SAO signals in the rmSAO and rmSAO-TP simulations. While the SST-SAO is removed globally, the SAO in the tropical upper-middle troposphere (500-175
260 hPa) is not significant. However, if the SST-SAO is removed only in the tropics, the upper troposphere SAO in the tropics is still significant, which indicates that the upper-middle troposphere SAO in the tropics is significantly influenced by the SST-SAO in the extra-tropics. At the same time, the SAO in the lower stratosphere of mid-latitudes (70-30 hPa in SH 32.5°S-47.5°S and NH 37.5°N-47.5°N) is also significantly reduced in both the rmSAO and rmSAO-TP simulations.

Although the UTLS-SAO is still significant in the rmSAO and rmSAO-TP simulations, the magnitudes of the UTLS-SAO
265 in the two sensitive simulations are significantly reduced (Figures 7c-d). Compared with Control simulation, the magnitude of rmSAO simulation is reduced by 31% for NHM and 55% for SHM (rectangular box in Figure 7c). The averaged magnitude reduction of rmSAO-TP simulation is 14% in NHM and 41% in SHM (rectangular box in Figure 7d). Such reduction of SAO PSD caused by removing SST-SAO is more evident in the SHM and NHM than tropics. This clearly shows that the SST-SAO modifies the strength of the SAO in the UTLS.

This result can also be confirmed by the averaged seasonal cycle of the extracted UTLS-SAO signal as shown in Figure 8a. To compare the amplitude of SAO signals in model simulations, we extracted SAO signals (5-7 months) using the 1-D wavelet Morlet transform (Lilly and Olhede, 2009, 2012). As shown in Figure 8a, the averaged amplitude of the rmSAO simulation (blue lines) is smaller than the Control simulation (red lines) (11%), and the amplitude of the rmSAO-TP simulation (dash black lines) is slightly weaker than the Control simulation (1%). This demonstrates that the SST-SAO in extra-tropics for the
275 UTLS-SAO in NHM is more important. Reduction of the SAO amplitude related to removing SST-SAO is more evident in the SHM (Figure 8b) compared to that in NHM. Compared with the Control simulation, the averaged amplitude of the rmSAO and rmSAO-TP simulation is reduced by 33% and 29%, respectively. This might be related to the relatively large area of ocean in the SH compared to the NH. Anyway, a connection has been established between the surface and the UTLS SAO, and it is obvious that the SST-SAO influences the UTLS-SAO significantly.

To further find out through what processes of the SSTs modifies the UTLS-SAO, the thermal budget analysis is applied to the three model simulations. Figure 9 shows a comparison of the thermal budget in the UTLS between the two sensitivity simulations and the Control simulation. In the NH, the total heating rates are significantly reduced from May to July, which is mainly caused by the weaker dynamical and moist heating in the two sensitivity simulations. From August to September, the reduction of moist heating is also evident but is offset by less radiative cooling. In October, the negative values of the total
285 heating rates are also reduced in the two sensitivity simulations due to the reduced radiative cooling. During boreal winter, the impact of removing SST-SAO is not significant.

In the SH, the total heating rates are reduced overall and especially significantly reduced in austral summer (November-December). This is related to the weaker dynamical and moist heating, although it is partly offset by less radiative cooling. In winter, the changes in total heating rate are not significant either. In April, the negative values of the total heating rate are

290 also reduced due to the reduced radiative cooling. In summary, the UTLS-SAO is modified by the SST-SAO mainly through its modification to the summer time moist heating, dynamical heating and autumn radiative cooling.

4 Conclusions

Variability of zonal mean temperatures over 500-10 hPa is analyzed based on high-quality GNSS temperature measurements and reanalysis data (MERRA2 and ERA5) covering 2001-2017. All data sets show a good agreement and ERA5 has the best
295 agreement with GNSS RO data. The model simulations have also good agreement with other data sets except in the tropical region. Our results confirm the tropical and subtropical SAO of temperature in the upper troposphere as reported by previous studies (Loon and Jenne, 1969). In addition, we also find a significant SAO of temperature in the lower stratosphere (just above the tropopause and from 70 to 30 hPa) in the mid-latitudes of both hemispheres. From the PSD of the SAO of temperature, the SAO signal is most pronounced from the subtropics to mid-latitudes (22.5° - 42.5°) in the UTLS region (250-175 hPa).
300 The SAO of temperature with maxima in February and July/August in the NH/SH mid-latitudes in the UTLS is observed. Furthermore, the SAO signals are likely related to both moist and dynamical processes.

The explanation of the SAO of temperature in the UTLS is given by a thermal budget analysis. In the NH, the first temperature peak is in February is mainly related to the dynamical heating while the second peak in July is caused by a combination of moist and dynamical processes. The temperature peaks follow the transitions of total heating rates by definition, and the moist
305 processes delay the transition of total heating rate from positive to negative in July. In the SH, both the dynamical and moist processes contribute to the temperature peak in summer, while the temperature maximum in winter (August) is mainly related to dynamical heating. The further energy budget analysis indicates that the vertical transportation of water vapor and subsequent condensation in the upper troposphere is much stronger and leads to a stronger peak of temperature in the summer season. Through the thermal budget analysis, we also find that the UTLS-SAO in the SH can be well represented by the WACCM6
310 model. A significant part in the SH represented by MERRA2 model cannot be explained by ANA, dynamical and moist process, which indicates that the budget for the SH is not as reliable as the NH.

Based on a series of model simulations, we analyze the relationship between UTLS-SAO and the SST-SAO. The results indicate that the amplitude of the UTLS-SAO can be significantly reduced while the SST-SAO is removed in the tropics, and will be further reduced if the SST-SAO is removed globally. The UTLS-SAO of temperature is therefore partly modified by the
315 SSTs. A comparison of the thermal budget between the sensitivity simulations and the Control simulation further indicates that the SST-SAO mainly modifies the UTLS-SAO through its modification to the summer time moist heating, dynamical heating and autumn radiative cooling.

The annual and semi-annual variations are the most pronounced short-term climate signals. In the past decades, numerous attentions have been attracted to investigate various aspects of the semi-annual cycle in the surface and stratosphere. Few
320 attempts have been made to analyze the SAO in the UTLS especially in the mid-latitudes. The major effort of this study not only explores the structure of the SAO in the mid-latitudinal UTLS and its relationship with thermodynamic balance but also provides the relevance between the SAO in the UTLS and SSTs by climate model simulations. However, a full mechanistic

understanding of the SAO drivers in the UTLS, the predictability of the SAO in the UTLS as well as its potential impacts on surface weather and climate awaits further studies.

325 *Data availability.* The authors are greatly appreciative to the teams of the CDAAC for the use of the GNSS RO data sets (<http://cdaac-www.cosmic.ucar.edu/cdaac/index.html>), the NASA GSFC for MERRA2 data (<https://disc.gsfc.nasa.gov/datasets?project=MERRA-2>) and the Copernicus Climate Change Service and the ECWMF for the ERA5 data (<https://cds.climate.copernicus.eu>). The simulations can be provided to readers by contacting the corresponding author.

330 *Author contributions.* MS performed the computational implementation and the analysis, created the figures and wrote the first draft of the paper. WW made the model simulations, created the figures and provided advice on the analysis design and contributed to the text. All authors contributed to the study design.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. The research has been supported jointly by the National Natural Science Foundation of China (grant no. 41904023 and 42075055), and the Fundamental Research Funds for the Central Universities, China University of Geosciences (Wuhan) CUG2106357.

335 References

- Abalos, M., Randel, W. J., Kinnison, D. E., and Serrano, E.: Quantifying tracer transport in the tropical lower stratosphere using WACCM, *Atmospheric Chemistry and Physics*, 13, 10 591–10 607, <https://doi.org/10.5194/acp-13-10591-2013>, 2013.
- Andrews, D. G., Holton, J. R., and Leovy, C. B.: *Middle Atmosphere Dynamics*, Academic Press, Orlando, Florida, p. 489, 1987.
- Anthes, R. A., Bernhardt, P. A., Chen, Y., Cucurull, L., Dymond, K. F., Ector, D., Healy, S. B., Ho, S.-P., Hunt, D. C., and Kuo, Y.-H.: The
340 COSMIC/FORMOSAT-3 Mission: Early Results, *Bulletin of the American Meteorological Society*, 89, 313–333, 2008.
- Bracegirdle, T. J.: The seasonal cycle of stratosphere-troposphere coupling at southern high latitudes associated with the semi-annual oscillation in sea-level pressure, *Climate dynamics*, 37, 2323–2333, 2011.
- Broeke, M. R. V. D.: The semiannual oscillation and Antarctic climate, part 5: impact on the annual temperature cycle as derived from NCEP/NCAR re-analysis, *Climate Dyn.*, 16, 369–377, 2000.
- 345 Butterworth, S.: On the theory of filter amplifiers, *Experimental Wireless and the Wireless Engineer*, 7, 536–541, 1930.
- Chen, T.-C. and Tsay, J.-D.: Development and maintenance mechanism for the semiannual oscillation of the North Pacific upper-level circulation, *Journal of climate*, 27, 3767–3783, 2014.
- Chen, T.-C. and Wu, K.-D.: Semi-annual oscillation of the global divergent circulation, *Tellus A: Dynamic Meteorology and Oceanography*, 44, 357–365, <https://doi.org/10.3402/tellusa.v44i5.14967>, 1992.
- 350 Chen, T. C., Yen, M. C., and Van Loon, H.: An Observational Study of the Tropical-Subtropical Semiannual Oscillation, *Journal of Climate*, 9, 1993–2002, 1996.
- Ern, M., Preusse, P., and Riese, M.: Driving of the SAO by gravity waves as observed from satellite, *Annales Geophysicae*, 33, 483–504, <https://doi.org/10.5194/angeo-33-483-2015>, 2015.
- Ern, M., Diallo, M., Preusse, P., Mlynczak, M. G., Schwartz, M. J., Wu, Q., and Riese, M.: The semiannual oscillation (SAO) in the tropical
355 middle atmosphere and its gravity wave driving in reanalyses and satellite observations, *Atmospheric Chemistry and Physics*, 21, 13 763–13 795, <https://doi.org/10.5194/acp-21-13763-2021>, 2021.
- Fueglistaler, S., Dessler, A. E., Dunkerton, T. J., Folkins, I., Fu, Q., and Mote, P. W.: Tropical tropopause layer, *Reviews of Geophysics*, 47, 2009.
- Gao, P., Xu, X., and Zhang, X.: On the relationship between the QBO/ENSO and atmospheric temperature using COSMIC radio occultation
360 data, *Journal of Atmospheric and Solar-Terrestrial Physics*, 156, 103–110, 2017.
- Garcia, R. R., Dunkerton, T. J., Lieberman, R. S., and Vincent, R. A.: Climatology of the semiannual oscillation of the tropical middle atmosphere, *Journal of Geophysical Research: Atmospheres*, 102, 26 019–26 032, <https://doi.org/10.1029/97JD00207>, 1997.
- Gelaro, R., Mccarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., and Reichle, R.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *Journal of Climate*, 30, 5419–5454,
365 <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- Gottelman, A. and Birner, T.: Insights into Tropical Tropopause Layer processes using global models, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/10.1029/2007JD008945>, 2007.
- Gottelman, A., Hoor, P., Pan, L. L., Randel, W. J., Hegglin, M. I., and Birner, T.: The extratropical upper troposphere and lower stratosphere, *Reviews of Geophysics*, 49, <https://doi.org/10.1029/2011RG000355>, 2011.
- 370 Gottelman, A., Mills, M. J., Kinnison, D. E., Garcia, R. R., Smith, A. K., Marsh, D. R., Tilmes, S., Vitt, F., Bardeen, C. G., McInerney, J., Liu, H.-L., Solomon, S. C., Polvani, L. M., Emmons, L. K., Lamarque, J.-F., Richter, J. H., Glanville, A. S., Bacmeister, J. T., Phillips, A. S.,

- Neale, R. B., Simpson, I. R., DuVivier, A. K., Hodzic, A., and Randel, W. J.: The Whole Atmosphere Community Climate Model Version 6 (WACCM6), *Journal of Geophysical Research: Atmospheres*, 124, 12 380–12 403, <https://doi.org/10.1029/2019JD030943>, 2019.
- 375 GMAO: Global Modeling and Assimilation Office, instM_3d_asm_Np: MERRA2 3D IAU State, Meteorology monthly mean (p-coord, 0.625x0.5L42), version 5.12.4, Greenbelt, MD, USA: Goddard Space Flight Center Distributed Active Archive Center (GSFC DAAC) Accessed: January 2021, <https://doi.org/10.5067/2E096JV59PK7>, 2015a.
- GMAO: Global Modeling and Assimilation Office, tavgM_3d_tdt_Np: MERRA2 3D IAU State, Meteorology monthly mean (p-coord, 0.625x0.5L42), version 5.12.4, Greenbelt, MD, USA: Goddard Space Flight Center Distributed Active Archive Center (GSFC DAAC) Accessed: January 2021, <https://doi.org/10.5067/VILT59HI2MOY>, 2015b.
- 380 Hamilton, K. and Mahlman, J. D.: General Circulation Model Simulation of the Semiannual Oscillation of the Tropical Middle Atmosphere, *Journal of the Atmospheric Sciences*, 45, 3212–3235, 1988.
- Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 monthly averaged data on pressure levels from 1979 to present. Copernicus Climate Change Service (C3S) Climate Data Store (CDS). (Accessed on 24-01-2021), <https://doi.org/10.24381/cds.6860a573>, 2019.
- 385 Ho, S.-P., Kirchengast, G., Leroy, S., Wickert, J., Mannucci, A. J., Steiner, A., Hunt, D., Schreiner, W., Sokolovskiy, S., Ao, C., Borsche, M., Engeln, A. v., Foelsche, U., Heise, S., Iijima, B., Kuo, Y., Kursinski, R., Pirscher, B., Ringer, M., Rocken, C., and Schmidt, T.: Estimating the uncertainty of using GPS radio occultation data for climate monitoring: Intercomparison of CHAMP refractivity climate records from 2002 to 2006 from different data centers, *Journal of Geophysical Research Atmospheres*, 114, <https://doi.org/10.1029/2009JD011969>, 2009.
- 390 Ho, S.-P., Hunt, D., Steiner, A. K., Mannucci, b. J., Kirchengast, G., Gleisner, H., Heise, S., Engeln, A. v., Marquardt, C., Sokolovskiy, S., Schreiner, W., Scherllin-Pirscher, B., Ao, C., Wickert, J., Syndergaard, S., Lauritsen, K. B., Leroy, S., Kursinski, E. R., Kuo, Y. H., Foelsche, U., Schmidt, T., and Gorbunov, M.: Reproducibility of GPS radio occultation data for climate monitoring: Profile-to-profile inter-comparison of CHAMP climate records 2002 to 2008 from six data centers, *Journal of Geophysical Research Atmospheres*, 117, <https://doi.org/10.1029/2012JD017665>, 2012.
- 395 Ho, S.-P., Yue, X., Zeng, Z., Ao, C. O., Huang, C., and Kursinski, E. R.: Applications of COSMIC Radio Occultation Data from the Troposphere to Ionosphere and Potential Impacts of COSMIC-2 Data, *Bulletin of the American Meteorological Society*, 95, 18–22, 2014.
- Ho, S.-P., Peng, L., and Vömel, H.: Characterization of the long-term radiosonde temperature biases in the upper troposphere and lower stratosphere using COSMIC and Metop-A/GRAS data from 2006 to 2014, *Atmospheric Chemistry and Physics*, 17, 4493–4511, 2017.
- Horne, J. H. and Baliunas, S. L.: A prescription for period analysis of unevenly sampled time series, *Astrophysical Journal*, 302, 757–763, 400 1986.
- Hu, R., Liu, Q., Meng, X., and Godfrey, J. S.: On the Mechanism of the Seasonal Variability of SST in the Tropical Indian Ocean, *Adv. Atmos. Sci.*, 22, 12, <https://doi.org/10.1007/BF02918758>, 2005.
- Kursinski, E. R., Healy, S. B., and Romans, L. J.: Initial results of combining GPS occultations with ECMWF global analyses within a 1DVar framework, *Earth Planets Space*, 52, 885–892, 2000.
- 405 Lilly, J. M. and Olhede, S. C.: Higher-Order Properties of Analytic Wavelets, *IEEE Transactions on Signal Processing*, 57, 146–160, <https://doi.org/10.1109/TSP.2008.2007607>, 2009.
- Lilly, J. M. and Olhede, S. C.: Generalized Morse Wavelets as a Superfamily of Analytic Wavelets, *IEEE Transactions on Signal Processing*, 60, 6036–6041, <https://doi.org/10.1109/TSP.2012.2210890>, 2012.
- Lomb, N. R.: Least-squares frequency analysis of unequally spaced data, *Astrophysics and Space Science*, 39, 447–462, 1976.

- 410 Loon, H. V.: The Half-Yearly Oscillations in Middle and High Southern Latitudes and the Coreless Winter, *jas*, 24, 472–486, 1967.
- Loon, H. V. and Jenne, R. L.: The Half-Yearly Oscillations in the Tropics of the Southern Hemisphere, *J. Atmos. Sci.*, 26, 218–232, 1969.
- Mapes, B. E. and Bacmeister, J. T.: Diagnosis of Tropical Biases and the MJO from Patterns in the MERRA Analysis Tendency Fields, *Journal of Climate*, 25, 6202–6214, 2012.
- Marsh, D. R., Mills, M. J., Kinnison, D. E., Lamarque, J.-F., Calvo, N., and Polvani, L. M.: Climate change from 1850 to 2005 simulated in
415 CESM1 (WACCM), *J. Climate*, 26, 7372–7391, <https://doi.org/10.1175/JCLI-D-12-00558.1>, 2013.
- Meehl, G. A., Hurrell, J. W., and Loon, H. V.: A modulation of the mechanism of the semiannual oscillation in the Southern Hemisphere, *Tellus A: Dynamic Meteorology and Oceanography*, 50, 442–450, <https://doi.org/10.3402/tellusa.v50i4.14537>, 1998.
- Merryfield, W. J., Baehr, J., Batté, L., Becker, E. J., Butler, A. H., Coelho, C. A. S., Danabasoglu, G., Dirmeyer, P. A., Doblas-Reyes, F. J., Domeisen, D. I. V., Ferranti, L., Ilynia, T., Kumar, A., Müller, W. A., Rixen, M., Robertson, A. W., Smith, D. M., Takaya, Y., Tuma,
420 M., Vitart, F., White, C. J., Alvarez, M. S., Ardilouze, C., Attard, H., Baggett, C., Balmaseda, M. A., Beraki, A. F., Bhattacharjee, P. S., Bilbao, R., de Andrade, F. M., DeFlorio, M. J., Díaz, L. B., Ehsan, M. A., Fragkoulidis, G., Grainger, S., Green, B. W., Hell, M. C., Infanti, J. M., Isensee, K., Kataoka, T., Kirtman, B. P., Klingaman, N. P., Lee, J.-Y., Mayer, K., McKay, R., Mecking, J. V., Miller, D. E., Neddermann, N., Ng, C. H. J., Ossó, A., Pankatz, K., Peatman, S., Pegion, K., Perlwitz, J., Recalde-Coronel, G. C., Reintges, A., Renkl, C., Solaraju-Murali, B., Spring, A., Stan, C., Sun, Y. Q., Tozer, C. R., Vigaud, N., Woolnough, S., and Yeager, S.: Current
425 and Emerging Developments in Subseasonal to Decadal Prediction, *Bulletin of the American Meteorological Society*, 101, E869–E896, <https://doi.org/doi:10.1175/BAMS-D-19-0037.1>, 2020.
- Park, K.-A. and Lee, E.-Y.: Semi-annual cycle of sea-surface temperature in the East/Japan Sea and cooling process, *International Journal of Remote Sensing*, 35, 4287–4314, <https://doi.org/10.1080/01431161.2014.916437>, 2014.
- Press, W. H. and Rybicki, G. B.: Fast Algorithm for Spectral Analysis of Unevenly Sampled Data, *Astrophysical Journal*, 338, 277–280,
430 1989.
- Randel, W. J. and Wu, F.: Variability of zonal mean tropical temperatures derived from a decade of GPS radio occultation data, *J. Atmos. Sci.*, 72, 1261–1275, <https://doi.org/10.1175/JAS-D-14-0216.1>, 2015.
- Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., Rowell, D. P., Kent, E. C., and Kaplan, A.: Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2002JD002670>, 2003.
435
- Read, P. L. and Castrejón-Pita, A. A.: Phase synchronization between stratospheric and tropospheric quasi-biennial and semi-annual oscillations, *Quart. J. Roy. Meteor. Soc.*, 138, 1338–1349, 2012.
- Read, R.: Some features of the annual temperature regime in the tropical stratosphere, *Mon. Weather Rev.*, 90, 211–215, 1962.
- Richter, J. H. and Garcia, R. R.: On the forcing of the Mesospheric Semi-Annual Oscillation in the Whole Atmosphere Community Climate
440 Model, *Geophysical Research Letters*, 33, <https://doi.org/10.1029/2005GL024378>, 2006.
- Scargle, J. D.: Studies in astronomical time series analysis.II.Statistical aspects of spectral analysis of unevenly spaced data, *Astrophysical Journal*, 263, 835–835, 1982.
- Schmidt, T., Wickert, J., Beyerle, G., König, R., Galas, R., and Reigber, C.: The CHAMP Atmospheric Processing System for Radio Occultation Measurements, Springer Berlin Heidelberg, 2005.
- 445 Schmidt, T., Wickert, J., and Haser, A.: Variability of the upper troposphere and lower stratosphere observed with GPS radio occultation bending angles and temperatures, *Advances in Space Research*, 46, 150–161, 2010.

- Schott, F. A., Xie, S.-P., and McCreary Jr., J. P.: Indian Ocean circulation and climate variability, *Reviews of Geophysics*, 47, <https://doi.org/10.1029/2007RG000245>, 2009.
- Shangguan, M., Wang, W., and Jin, S.: Variability of temperature and ozone in the upper troposphere and lower stratosphere from multi-
450 satellite observations and reanalysis data, *Atmospheric Chemistry and Physics*, 19, 6659–6679, 2019.
- Shea, D. J., Loon, H. V., and Hurrell, J. W.: The tropical-subtropical semi-annual oscillation in the upper troposphere, *Int. J. Climatol.*, 15, 975–983, 1995.
- Simmonds, I. and Jones, D. A.: The mean structure and temporal variability of the semiannual oscillation in the southern extratropics, *Int. J. Climatol.*, 18, 473–504, 1998.
- 455 Walland, D. and Simmonds, I.: Baroclinicity, Meridional Temperature Gradients, and the Southern Semiannual Oscillation, *Journal of Climate*, 12, 3376–3382, 1998.
- Wang, W., Matthes, K., Omrani, N. E., and Mojib, M.: Decadal variability of tropical tropopause temperature and its relationship to the Pacific decadal oscillation, *Sci. Rep.*, 6, 29 537, <https://doi.org/10.1038/srep29537>, 2016.
- Wang, W., Shangguan, M., Tian, W., Schmidt, T., and Ding, A.: Large Uncertainties in Estimation of Tropical Tropopause Temperature
460 Variabilities Due to Model Vertical Resolution, *Geophysical Research Letters*, 46, 10 043–10 052, <https://doi.org/10.1029/2019GL084112>, 2019.
- Wee, T.-K. and Kuo, Y.-H.: Advanced stratospheric data processing of radio occultation with a variational combination for multifrequency GNSS signals, *Journal of Geophysical Research Atmospheres*, 119, 11,011–11,039, 2015.
- Wickert, J., Reigber, C., Beyerle, G., König, R., Marquardt, C., Schmidt, T., Grunwaldt, L., Galas, R., Meehan, T. K., and Melbourne, W. G.:
465 Atmosphere sounding by GPS radio occultation: First results from CHAMP, *Geophys. Res. Lett.*, 28, 3263–3266, 2001.
- Wickert, J., Michalak, G., Schmidt, T., Beyerle, G., Cheng, C., Healy, S. B., Heise, S., Huang, C., Jakowski, N., and Köhler, W.: GPS radio occultation: results from CHAMP, GRACE and FORMOSAT-3/COSMIC., *Terrestrial Atmospheric & Oceanic Sciences*, 20, 35–50, 2009.
- Xu, G., Yue, X., Zhang, W., and Wan, X.: Assessment of Atmospheric Wet Profiles Obtained from COSMIC Radio Occultation Observations over China, *Atmosphere*, 8, 208–208, 2017.
- 470 Yan, Y., Wang, G., Chen, C., and Ling, Z.: Annual and Semiannual Cycles of Diurnal Warming of Sea Surface Temperature in the South China Sea, *Journal of Geophysical Research: Oceans*, 123, 5797–5807, <https://doi.org/10.1029/2017JC013657>, 2018.
- Yang, F. and Wu, Z.: On the physical origin of the semiannual component of surface air temperature over oceans, *Climate Dynamics*, pp. 1–13, 2022.
- Yashayaev, I. M. and Zveryaev, I. I.: Climate of the seasonal cycle in the North Pacific and the North Atlantic oceans, *International Journal of Climatology*, 21, 401–417, <https://doi.org/10.1002/joc.585>, 2001.
475




Figure 1. The power spectrum densities (PSD) of SAO based on GNSS RO (a), ERA5 (b), MERRA2 (c) and model simulation (d) temperature for the period 2001-2017. The dots mark the significant area at 95% level. The dashed black lines mark the tropopause height calculated with GNSS RO data.




Figure 2. The PSD of SAO based on COSMIC-1 (2007-2017) (a), ERA5 (2001-2017) (b), MERRA2 (2001-2017) (c) and model simulation (2001-2017) (d) at 200 hPa. The dots mark the significant area at 95% level.




Figure 3. Time series of the zonal mean temperature (T) at 200 hPa averaged around the Northern Hemisphere mid-latitudes (NHM) 32.5°N-42.5°N (a) and their corresponding annual cycle anomalies (b). The Southern Hemisphere mid-latitudes (SHM) 32.5°S-42.5°S (c) and their corresponding annual cycle anomalies (d).




Figure 4. Annual cycle of the zonal mean temperature (T) and heating rates at 200 hPa averaged around the NHM 32.5°N-42.5°N (a) and the SHM 32.5°S-42.5°S (b) based on MERRA2 data. The red, blue, dashed blue and dashed red lines indicate the heating rates related to dynamics, radiation, condensation and analysis processes, respectively. The positive total heating rates are filled with light red color and the negative total heating rates are filled with light blue color. The total heating rates, which are the sum of analysis, dynamical, radiative and moist heating rates, have been 5 times enlarged to be more visible in the figures.




Figure 5. The variation of heating rates terms at 200 hPa at peaks and troughs of the total heating rates around the NHM 32.5°N-42.5°N (a) and the SHM 32.5°S-42.5°S (b) based on MERRA2 data. The blue, red, yellow, purple and green columns are total, analysis, dynamical, radiative and moist terms, respectively.




Figure 6. Annual cycle of dynamical heating rates at 200 hPa averaged around the NHM 32.5°N-42.5°N (a) and the SHM 32.5°S-42.5°S based on MERRA2 (a-b) and Control simulation (c-d). The cyan, red, green and black lines are the eddy, heating by downwelling ($-wS$), heating by the meridional component of the residual circulation ($-vT_y$) and dynamical term.




Figure 7. (a) The PSD of SAO analyzed for the period 2001-2017 based on model simulation with removed SST-SAO (rmSAO). (b) Same as (a), but for the simulation with removed SST-SAO in the tropics (rmSAO-TP). The dots mark the significant area at 95% level. (c) The difference of SAO PSD between the rmSAO and the Control simulations (rmSAO - Control) for the period 2001-2017. (d) Same as (c), but for the difference between the rmSAO-TP and the Control simulations (rmSAO-TP - Control). The black dots mark area with significant difference of the two time series at 95% level based on t-test. The dashed black lines mark the tropopause height calculated with GNSS RO data.




Figure 8. The average extracted SAO signals (5-7 months) based on 1-D wavelet Morlet transform for the Northern Hemisphere mid-latitudes (NHM) (a) and for the Southern Hemisphere mid-latitudes (SHM)(b). The red, blue and dashed black lines are results of the Control, rmSAO and rmSAO-TP runs, respectively.




Figure 9. Annual cycle of the heating rates at 200 hPa averaged around the Northern Hemisphere mid-latitudes 32.5°N - 42.5°N (a) and the Southern Hemisphere mid-latitudes 32.5°S - 42.5°S (b). The red, blue and yellow lines indicate the heating rates related to dynamics, radiation and condensation processes, respectively. The total heating rates, which are the sum of the dynamical, radiative and moist heating rates, are illustrated by black lines. The solid, dashed and dotted-dashed lines indicate data from the Control, rmSAO and rmSAO-TP simulations, respectively.