

Updated trends of the stratospheric ozone vertical distribution in the 60°S–60°N latitude range based on the LOTUS regression model

Sophie Godin-Beekmann¹, Niramson Azouz¹, Viktoria Sofieva², Daan Hubert³, Irina Petropavlovskikh⁴, Peter Effertz⁴, Gérard Ancellet¹, Doug. A. Degenstein⁵, Daniel Zawada⁵, Lucien Froidevaux⁶, Stacey Frith⁷, Jeannette Wild⁸, Sean Davis⁹, Wolfgang Steinbrecht¹⁰, Thierry Leblanc¹¹, Richard Quere¹², Kleareti Tourpali¹³, Robert Damadeo¹⁴, Eliane Maillard-Barras¹⁵, René Stübi¹⁵, Corinne Vigouroux³, Carlo Arosio¹⁶, Gerald Nedoluha¹⁷, Ian Boyd¹⁸, Roeland Van Malderen¹⁹, Emmanuel Mahieu²⁰, Dan Smale¹², Ralf Sussmann²¹

¹LATMOS Sorbonne Université, UVSQ, CNRS, France,

²Finnish Meteorological Institute, Finland,

³Royal Belgian Institute for Space Aeronomy (BIRA-IASB), Belgium,

⁴Cooperative Institute for Research in Environmental Sciences, U. of Colorado, Boulder, CO, USA,

⁵University of Saskatchewan, Canada,

⁶Jet Propulsion Laboratory, California Institute of Technology, Pasadena, USA,

⁷Science Systems and Applications, Inc & NASA Goddard Space Flight Center, USA,

⁸ESSIC/UMD & NOAA/NCEP/Climate Prediction Center, USA,

⁹NOAA Chemical Sciences Laboratory, USA

¹⁰Deutsche Wetterdienst, Germany,

¹¹Jet Propulsion Laboratory, California Institute of Technology, Wrightwood, USA,

¹²National Institute of Water and Atmospheric Research (NIWA), New Zealand,

¹³Aristotle University of Thessaloniki, Greece,

¹⁴NASA Langley, USA,

¹⁵Federal Office of Meteorology and Climatology, MeteoSwiss, Switzerland,

¹⁶Institute of Environmental Physics, Bremen Universität, Germany,

¹⁷Remote Sensing Division, Naval Research Laboratory, Washington, DC, USA,

¹⁸Bryan Scientific Consulting, Charlottesville, VA, USA,

¹⁹Royal Meteorological Institute, Belgium,

²⁰University of Liège, Institute of Astrophysics and Geophysics, Liège, Belgium,

²¹Karlsruhe Institute of Technology, IMK-IFU Garmisch-Partenkirchen, Germany

Correspondence to: sophie.godin-beekmann@latmos.ipsl.fr

Abstract.

This study presents an updated evaluation of stratospheric ozone profile trends in the 60°S – 60°N latitude range over the 2000 – 2020 period using an updated version of the Long-term Ozone Trends and Uncertainties in the Stratosphere (LOTUS) regression model that was used to evaluate such trends up to 2016 for the last WMO Ozone Assessment (2018). In addition to the derivation of detailed trends as a function of latitude and vertical coordinates, the regressions are performed with the data sets averaged over broad latitude bands, i.e. 60°S–35°S, 20°S–20°N and 35°N–60°N. The same methodology as in the last

Supprimé: v

Mis en forme : Anglais (G.B.)

Supprimé: f

Mis en forme : Exposit

Assessment is applied to combine trends in these broad latitude bands in order to compare the results with the previous studies. Longitudinally resolved merged satellite records are also considered in order to provide a better comparison with trends retrieved from ground-based records, e.g. lidar, ozone sondes, Umkehr, microwave and Fourier Transform Infrared (FTIR) spectrometers at selected stations where long-term time series are available. The study includes a comparison with trends derived from the REF-C2 simulations of the Chemistry Climate Model Initiative (CCMI-1). This work confirms past results showing an ozone increase in the upper stratosphere, which is now significant in the three broad latitude bands. The increase is largest in the northern and southern hemisphere midlatitudes, with -2.2 ± 0.7 %/decade at ~ 2.1 hPa, and -2.1 ± 0.6 %/decade at ~ 3.2 hPa respectively, compared to -1.6 ± 0.6 %/decade at ~ 2.6 hPa in the tropics. New trend signals have emerged from the records, such as a significant decrease of ozone in the tropics around 35 hPa and a non-significant increase of ozone in the southern mid-latitudes at about 20 hPa. Non-significant negative ozone trends are derived in the lowermost stratosphere, with the most pronounced trends in the tropics. While a very good agreement is obtained between trends from merged satellite records and the CCMI-1 REF-C2 simulation in the upper stratosphere, observed negative trends in the lower stratosphere are not reproduced by models at southern and, in particular, at northern midlatitudes, where models report an ozone increase. However, the lower stratospheric trend uncertainties are quite large, for both measured and modelled trends. Finally, 2000-2020 stratospheric ozone trends derived from the ground-based and longitudinally resolved satellite records are in reasonable agreement over the European Alpine and tropical regions, while at the Lauder station in the southern hemisphere mid-latitudes they show some differences.

Supprimé: latest

Supprimé: close

Supprimé: , especially

Supprimé: .

1. Introduction

The recovery of the ozone layer has been under scrutiny since the peak of ozone depleting substances (ODS) was reached in the stratosphere in the mid and end of the 1990s depending on the latitude region (e.g. Newman et al., 2007) in response to reduced ODS emissions imposed by the 1987 Montreal Protocol and its subsequent amendments. After first indications of a small ozone increase from various ground-based and satellite records in the upper stratosphere, i.e. above 35 km (WMO, 2010), clear evidence of the impact of decreasing ODS content on ozone levels in that altitude region was provided in WMO (2014) and references therein. Since then, an upper stratospheric ozone increase has been confirmed by various studies (e.g. Harris et al., 2015; Steinbrecht et al., 2017; Petropavlovskikh et al., 2019). In parallel, chemistry climate models (CCMs) have attributed half of this increase to decreased ODS concentrations and half to upper stratospheric cooling resulting from increased greenhouse gases (GHGs), which slows gas-phase ozone depleting reactions (e.g., chapter 5 of WMO, 2018). In contrast, an ozone increase in the lower stratosphere has not been detected to date, except for some emerging signs in the Antarctic polar region in spring (de Laat et al., 2015; Solomon et al., 2016; Pazmiño et al., 2018; WMO, 2018).

Supprimé: at the turn of the 21st century

Supprimé: signed in 1987

Supprimé: First

Supprimé: for ozone in the upper stratosphere

Supprimé: ,

Supprimé: ,

80 The issue of ozone evolution and recovery in the lower stratosphere has received a lot of attention in recent years. Using several long-term satellite combined records and derived trends based on the Dynamical Linear Modelling (DLM) method, Ball et al. (2018) found a decline of ozone in the lower stratosphere over the period 1998-2016. This result was challenged by Chipperfield et al. (2019), who argued that the ozone reduction was influenced by short-term dynamical variability at the end of the studied period. The cause for ozone decline or lack of recovery in the lower stratosphere was investigated by several model-based studies. Orbe et al. (2020) suggested that the observed decrease of ozone in the northern hemisphere could be explained by a poleward expansion of tropical upwelling and a reduced downwelling in the northern subtropical region. Other studies (Wargan et al., 2018; Ball et al., 2020) pointed to an enhanced meridional mixing between the tropics and the midlatitudes.

85 The present study is a follow-up of the ozone profile trend analysis performed within the LOTUS activity of the Stratosphere-Troposphere Processes And their Role in Climate (SPARC) programme (Petropavlovskikh et al., 2019) that contributed to chapter 3 of the last WMO/UNEP Ozone Assessment (WMO, 2018) and is referred to as LOTUS19 hereafter. In order to achieve a consistent interpretation of stratospheric ozone changes, multiple merged satellite and ground-based data records of ozone vertical distribution were collected to perform the same trend analyses. Previously published multiple linear regression (MLR) models were tested on a common ozone data set to evaluate the sensitivity of derived trends to the use of different models for the regression. This enabled the selection of the open source "LOTUS regression model" (https://arg.usask.ca/docs/LOTUS_regression), maintained by the University of Saskatchewan. The trends in the vertical distribution of stratospheric ozone profiles were assessed over the period 1985 – 2016. A new approach was established for combining the trend estimates from individual satellite-based records into a single best estimate of ozone profile trends representative of the three broad latitude bands: 35° – 60° at southern (SH) and northern hemisphere (NH) midlatitudes, and 20°N – 20°S in the tropics. Special attention was given to the evaluation of trend significance as a function of altitude. The LOTUS19 trend results were compared to those derived from previous studies (e.g., Harris et al., 2015, Steinbrecht et al., 2017). LOTUS19 found positive trends in the upper stratosphere in the post-ODS peak period (2000 – 2016) for both satellite and ground-based records. Results from merged satellite records showed statistically significant positive combined trends in the northern hemisphere midlatitudes of 2–3% per decade in the ~5–1 hPa pressure range and in the tropics of 1–1.5% per decade in the ~3–1 hPa pressure range. Combined trends were not statistically significant in the upper stratosphere at southern midlatitudes and no significant trends were obtained in the lower stratosphere. The LOTUS19-derived trends in broad latitude bands were also compared to trends from the CCM1-1 simulations. Both models and merged satellite records showed similar results in terms of trend values and significance, except at southern midlatitudes in the upper stratosphere where model trends were found to be significant.

Supprimé: at extrapolar latitudes was investigated by numerous studies ...

Supprimé: Ball et al. (2018) considered

Supprimé: using

Supprimé: . They

Supprimé: On the other hand, Wargan et al. (2018) used an idealized atmospheric tracer in a chemistry transport model constrained by MERRA-2 re-analyses in order to assess possible mechanisms responsible for the decrease of lower stratospheric ozone. Their analysis detected enhanced isentropic transport between the tropical (20°S – 20°N) and extratropical lower stratosphere in the past two decades, which could have influenced the downward trends. This result was confirmed by Ball et al. (2020), who investigated ozone, water vapor and temperature trends from both satellite records and CCM simulations performed in the frame of the REF-B2 CCM1, as well as changes in residual circulation upwelling and mixing efficiency from Japanese 55-year Reanalyses (JRA55) and European Centre for Medium-Range Weather Forecasts (ECMWF) ERA Interim reanalyses. These authors showed that observed ozone trends in the lower stratosphere can be explained by enhanced mixing between the tropics and extratropics and that these trends are not well reproduced by CCM simulations. ¶

Supprimé: unified

Supprimé: that is

Supprimé: of 2–3% per decade in the ~5–1 hPa pressure range and

Supprimé: respectively

Supprimé: model

Supprimé: -1 REF-C2 simulations

140 Since LOTUS19, other studies assessed global stratospheric ozone profile trends. Szelag et al. (2020) analyzed the seasonal dependence of stratospheric ozone trends from four merged satellite datasets over the 2000 – 2018 period using a two-step MLR model. They found positive trends in the upper stratosphere at middle and high latitudes, maximizing during the winter. This is consistent with findings that due to GHG concentration increases, the Brewer-Dobson, which is most effective in the winter season, should strengthen and accelerate the ozone recovery (e.g. Garcia et al., 2008). In the lower and middle stratosphere, negative trends were found in the tropics during all seasons, along with trends of varying sign depending on the season in the northern and southern midlatitudes. Another study by Sofieva et al. (2021) evaluated regional trends from the new merged gridded dataset MEGRIDOP that combines ozone profile data from six limb-viewing satellite instruments. Zonal trend estimates agreed with previously published results. Longitudinally resolved trends showed a zonal asymmetry in the upper stratosphere at high and middle latitudes in the northern hemisphere with larger trends over Scandinavia than over Siberia.

Supprimé: After

Supprimé: , which

Supprimé: photochemical control of ozone in that region.

150 In the present study, we compute trends from updated versions of the merged satellite records used in LOTUS19 and extended to the end of 2020, from the newly available MEGRIDOP merged time series and from updated versions of the ground-based data records selected from selected stations of the Network for the Detection of Atmospheric Composition Changes (NDACC), where observations of ozone profile as well as of other parameters are collected using a variety of ground-based techniques. Trends in the ozone vertical distribution derived from the various satellite and ground-based records as well as the CCMI-1 simulations considered in the study are evaluated using an updated version (version 0.8.0) of the LOTUS regression model.

Supprimé: the so-called supersite

Supprimé: ozone profile

Supprimé: over the post ODS peak period

155 The paper is organized as follows. Section 2 summarizes the satellite and ground-based records used in the study, while section 3 describes the updated version of the LOTUS regression model that was employed to retrieve trends from the various records. Section 4 displays the different trend results for the merged satellite data sets both as a function of latitude and vertical levels, and combined in broad latitude bands. Comparison between combined trends in broad latitude bands with corresponding LOTUS19 results and with trends computed from the updated CCMI-1 simulations are presented. In addition, trends from ground-based records at selected NDACC sites are compared to those from longitudinally resolved satellite data. Section 5 discusses improvements in trend retrievals with respect to LOTUS19, while conclusions of the study regarding long-term ozone profile changes are given in section 6.

Supprimé: super

2. Data

165 This section provides a brief description of the long-term ozone records used for trend retrieval. Readers can refer to Petropavlovskikh et al. (2019) for a more in-depth description of the various observational datasets (Chapter 1 of the report).

Supprimé: substantial

2.1 Merged satellite records

175 Seven merged satellite records that were extended to December 2020 are used for this study (see also Table 2.2 of LOTUS19). The Global Ozone Chemistry And Related trace gas Data records for the Stratosphere (GOZCARDS v.2.20) ozone monthly mean record includes HALogen Occultation Experiment (HALOE; v19), Aura Microwave Limb Sounder (MLS v4.2), SAGE I (version 5.9), SAGE II (v7) and covers the period 1979 – 2020. HALOE and Aura MLS measurements are adjusted with SAGE II data, which are used as a reference in the overlapping time periods (Froidevaux et al., 2015).

180 Data included in the Stratospheric Water and OzOne Satellite Homogenized (SWOOSH) merged record are Aura MLS v4.2, UARS MLS v5, UARS HALOE v19, SAGE II v7 and SAGE III v4. The merged records are homogenised to minimise artificial discontinuities and to account for inter-satellite biases in the record (Davis et al., 2016).

The Solar Backscatter Ultraviolet Merged total and profile Ozone Data (SBUV MOD) record includes data from the SBUV predominantly on the NOAA satellite series of instruments and the Ozone Mapping and Profiler Suite – Nadir profiler (OMPS-NP) on the Suomi National Polar-orbiting Partnership (SNPP). Data from all SBUV instruments except the NOAA-9 instrument and the morning portion of NOAA-14 and NOAA-16 are included, providing a continuous coverage of ozone profiles since 1978. For the merged data set no external calibration adjustments are applied, as in-instrument calibration adjustments have already been applied at the radiance level within the retrieval algorithm. Measurements are averaged during periods when more than one instrument is operational (Frith et al., 2017). Version 8.7 of SBUV MOD is used in this study, which includes refined in-instrument calibration adjustments (for NOAA-16 through OMPS NP, using NOAA-19 as a reference) and a diurnal correction to account for varying measurement times (Frith et al., 2020; Kramarova et al., 2022).

185 Another approach was adopted for the SBUV Cohesive (SBUV COH) merged dataset that uses much of the same SBUV and OMPS instruments as SBUV MOD, but retains use of Version 8.6 for SBUV processing as was used in LOTUS19. The COH approach identifies a representative satellite for each time period, and examines data for each overlapping period to improve the consistency of some satellite records with their neighbors. As in LOTUS19, the updated SBUV COH dataset used in this study adjusts NOAA-16, NOAA-17, NOAA-19 to the NOAA-18 SBUV record, and also extends the record to 2020 with OMPS-NP from SNPP (NOAA v3r2, from NOAA/NESDIS), also adjusted to NOAA-18. Early data are minimally adjusted. Nimbus-7 and NOAA-11 are not adjusted, and NOAA-9 is used for a minimal time period to fill a data gap and is adjusted to NOAA-11 (Wild et al., 2019). SBUV COH uses Version 8.6 for SBUV data and the NOAA v3r2 version of the OMPS-NP retrieval.

200 The merged SAGE-CCI-OMPS dataset was developed in the framework of the European Space Agency Climate Change Initiative for Ozone (Ozone-CCI). It includes data from SAGE II, several ozone measuring instruments on board the Environmental Satellite (EnviSat), OSIRIS on Odin, ACE-FTS on the SCience SATellite (SCISAT), and the OMPS - Limb

Mis en forme : Espace Avant : 6 pt

Supprimé: product

Supprimé: instrument

Supprimé:

Supprimé: though

Supprimé: 2r3

Supprimé: 2r3

Supprimé:

Profiler (OMPS-LP) (Sofieva et al., 2017). The merging method consists in merging long-term deseasonalized anomalies from the individual satellite ozone records. A similar methodology has been used for the merged SAGE-OSIRIS-OMPS time series that includes data from SAGE II, OSIRIS and OMPS-LP Usask 2D records (Bourassa et al., 2018; Zawada et al., 2018).

215 Finally, the SAGE-SCIAMACHY-OMPS record includes data from SAGE II, the SCanning Imaging Absorption spectroMeter for Atmospheric CHartography (SCIAMACHY) and OMPS-LP retrieved with U. Bremen code. The merging of SCIAMACHY and OMPS-LP records with SAGE II occultation observations is carried out from zonally averaged monthly anomalies (Arosio et al., 2019).

220 Compared to LOTUS19, the SAGE-MIPAS-OMPS record that was not extended to 2020 has been replaced by the SAGE-SCIAMACHY-OMPS record. Most other records were extended to 2020 with no substantial version change. SBUV MOD now includes OMPS-NP and uses the Version 8.7 retrieval algorithm for all data set used. SBUV-COH is the same dataset as used in LOTUS19 through 2010. The NOAA-19 component for 2011 to 2013 has been reprocessed since the LOTUS19 report with enhanced calibrations, but no algorithm change, and OMPS-NPP extends the data from 2014 to 2020. For better comparison with ground-based datasets, we also use the new Merged GRidded Dataset of Ozone Profiles (MEGRIDOP) record (Sofieva et al., 2021). This dataset has a resolved longitudinal structure, and is derived by merging of six limb and 225 occultation satellite data sets (GOMOS, SCIAMACHY and MIPAS on Envisat, OSIRIS, OMPS-LP, and Aura MLS), using a similar methodology as for SAGE-CCI-OMPS (Sofieva et al., 2021).

2.2 Ground-based records

230 Several NDACC stations were selected for trend comparison with merged satellite records. These stations provide multiple ground-based long-term ozone records using different techniques as mandated by the NDACC strategy (see also <http://ndaccdemo.org/stations>). Ground-based measurement techniques used for the comparison include balloon-borne ozonesondes, lidar (light detection and ranging), microwave radiometer, FTIR spectrometer, and Umkehr profile retrieval from Dobson sunrise and sunset measurements. Ozonesondes are small balloon-borne instruments attached to a standard radiosonde. Based on electrochemical sensing solution, they measure ozone in situ profile from the ground to about 30–35 km 235 altitude with a ~150 m vertical resolution linked to balloon ascent rate and ozone cell response time. There are several types of ozonesondes and two of them are used in this study: Electrochemical concentration cell (ECC) and Brewer-Mast (BM). Each ozonesonde is a unique instrument and biases have been found in ozonesonde records linked to the preparation method, the type of sonde or the sensing solution used, or even to the batch of instruments purchased from manufacturers. Since 2004, the WMO/GAW-sponsored Assessment of Standard Operating Procedures of Ozonesondes (ASOPOS) panel has evaluated 240 and intercompared ozonesonde measurements in the field or in laboratory chambers. The latest ASOPOS2 report (Smit, Thompson et al., 2021) provides recommendations on sonde preparation steps and measurement protocols, with the objective

Supprimé:

Supprimé: to

Supprimé: by available

Supprimé: with

Supprimé: from th

Supprimé: e

Supprimé: data

Supprimé: :

Supprimé: supersite

Supprimé: term "supersite"

Supprimé: is

Supprimé: linked to the

Supprimé: available at these stations

Supprimé: The selected NDACC stations are Mauna Loa (MLO) and Hilo (for ozonesondes) in the tropics, Lauder in the southern hemisphere midlatitudes and various stations located in the vicinity of the European Alps in the northern hemisphere midlatitudes. The so-called Alpine stations are Hohengeissenberg, Arosa, Payerne, Zugspitze, Jungfrauoch and Observatoire de Haute-Provence (OHP).

Supprimé: ¶

Supprimé:

Supprimé:

265 to achieve by adoption of these guidelines the 5% uncertainty level in tropospheric and stratospheric ozone requested by satellite and trends communities. Based on ASOPOS recommendations, ECC ozonesondes measurements records have been homogenized in multiple stations worldwide and the ECC ozonesonde data used in this study are from the Harmonization and Evaluation of Ground Based Instruments for Free Tropospheric Ozone Measurements (HEGIFTOM) prepared set of homogenized ozonesonde records.

270 Lidar is an active remote sensing technique. For the measurement of the ozone vertical distribution it uses the emission of two laser wavelengths with different ozone absorption cross-sections according to the so-called Differential Absorption Lidar (DIAL) technique. Pulsed lasers are used in order to obtain range resolved measurements (e.g. Godin-Beekmann et al., 2003, Leblanc et al., 2016). This study uses lidar ozone profile records extended to 2020.

275 Microwave ozone radiometers (MWR) detect emission spectra in the millimetre range produced by thermally excited rotational ozone transitions (e.g. Maillard-Barras et al., 2020). The ozone profile retrieval is based on the pressure broadening effect of the emitted line with the use of a priori profile and optimal estimation method (Rodgers et al., 2000, Bernet et al., 2019).

280 Umkehr ozone profiles are retrieved from the difference in zenith sky intensities selected from two spectral regions in the so-called C-pair at 311.5 and 332.4 nm of Dobson and Brewer spectrometer measurements. The long-term record of Umkehr measurements from four NOAA Dobson spectrophotometers located in the Boulder, OHP, MLO and Lauder stations has recently been reprocessed with the optimized homogenization technique (Petrovavlovskikh et al., 2022) and the three latter improved records are used here.

The FTIR ozone measurements are performed over the 600–4500cm⁻¹ spectral range, with high-resolution spectrometers, using the sun as source of light under clear sky conditions. On top of total ozone columns, low vertical resolution ozone profiles can be derived from the temperature and pressure dependence of the line shapes (Hase et al., 1999, Vigouroux et al., 2015).

285 The selected NDACC stations are Mauna Loa (MLO, lidar, microwave, Umkehr) and Hilo (for ozonesondes) in the tropics, Lauder in the southern hemisphere midlatitudes (lidar, ozonesondes, FTIR, Umkehr) and stations located in the vicinity of the European Alps in the northern hemisphere midlatitudes. These Alpine stations are Hohenpeissenberg (lidar, ozonesondes), Arosa (Umkehr), Payerne (microwave, ozonesondes), Zugspitze (FTIR), Jungfraujoch (FTIR) and Observatoire de Haute-Provence (OHP, lidar, ozonesondes, Umkehr). The location of these stations within a radius of less than 700 km corresponds to one grid cell of the longitudinally resolved satellite records used in this study, i.e. 10°lat x 20°long for MEGRIDOP and 10°lat x 30°long for SBUV-MOD and SWOOSH, which facilitates the satellite – ground-based trend comparison. Table 1 summarizes the ground-based measurements performed in the selected NDACC stations and the length of the record.

Supprimé: -

Supprimé: 2019

Mis en forme : Espace Avant : 0 pt

Supprimé: We have also compared trends from ground-based ozone records derived with the LOTUS model with those from FTIR measurements using another trend model (e.g. Vigouroux et al. 2015). ...

Mis en forme : Anglais (E.U.)

300 2.3 CCMI-1 Model data

In addition, we have used data from the chemistry–climate models (CCMs) participating in phase 1 of the Chemistry–Climate Model Initiative (CCMI-1; Eyring et al., 2013), which are able to capture the coupling between stratosphere and troposphere in terms of composition and physical climate processes more consistently than previous model generations. The REF-C2 simulation, which is a seamless simulation running from 1960 to 2100, was selected and the trend analysis was made over the period 1979–2020. REF-C2 experiments follow the WMO (2011) A1 scenario for ozone depleting substances and the RCP 6.0 for other greenhouse gases, tropospheric ozone precursors, and aerosol and aerosol precursor emissions. Ocean conditions are either modeled (from a separate climate model simulation), or internally generated (in the case of ocean-coupled models). The simulation includes state-of-knowledge historic forcings, with recommendations that the 11-year solar cycle and QBO forcings be either internally model-generated or nudged from the data set provided by Free University of Berlin. No volcanic forcing was used in this Reference simulation. For a detailed description of all forcings used in the reference simulations, see Eyring et al. (2013), Hegglin et al. (2016), and Morgenstern et al. (2017). For the CCMI-1 trend analysis, all necessary proxies were calculated directly from the pertaining individual model simulations. We calculated the appropriate QBO and ENSO proxies from the model data (zonal winds and SSTs), and used the external forcings (e.g. 11-year solar cycle) as provided to the modelling groups taking into account their implementation. We note that although the recommendation for the REF-C2 set of simulations did not include volcanic forcing, we found that some models did use it. Moreover, as volcanic effects could appear via different routes (e.g. SSTs or QBO), stratospheric AOD was used as a proxy in the trend analysis as forcing provided to the modelling groups, see section 4.5.2 of the LOTUS19 report for an in-depth explanation of CCMI-1 trends calculation.

Supprimé: W

Supprimé: We used t

Supprimé: that

Supprimé: with

Supprimé: our

Supprimé: c

Supprimé: ing

Supprimé: ,

Supprimé: were

3. The LOTUS regression model

An updated version of the LOTUS regression model (version 0.8.0) is used for the trend computation. It relies on the classical multiple linear regression method, which estimates the variability of time series from explanatory variables from the general least squares approach. The explanatory variables or proxies used in the LOTUS model are the quasi-biennial oscillation (QBO), the El Niño–Southern Oscillation (ENSO), the 11-year solar cycle, the stratospheric aerosol optical depth (sAOD) and a long-term trend. As in LOTUS19, we use independent linear trend (ILT) terms to evaluate long-term changes before and after the ODS peak, i.e., before January 1997 and after January 2000. The LOTUS model is applied to the ozone records without weight based on e.g. measurement uncertainty. Most data sets are provided as monthly mean time series and are deseasonalized within the LOTUS model using Fourier components representing annual and semi-annual variations. The fitting of the deseasonalized times series is based on the following equation:

Supprimé: -

Supprimé: the

Supprimé: e.g.,

Supprimé:

Supprimé: For the d

Supprimé: that are

Supprimé: not

Supprimé: , the model includes

Supprimé: model

Supprimé: following equation for deseasonalized time series:

$$y(z,t) = \beta_1(z,t) \cdot QBO_1(t) + \beta_2(z,t) \cdot QBO_2(t) + \beta_3(z,t) \cdot ENSO(t) + \beta_4(z,t) \cdot Solar(t) + \beta_5(z,t) \cdot sAOD(t) + (\beta_6(z,t) + \beta_7(z,t)(t - t_1)) \cdot L_{pre}(t) + (\beta_8(z,t) + \beta_9(z,t)(t - t_2)) \cdot L_{post}(t) + \beta_{10}(z,t) \cdot Gap(t) + \varepsilon(z,t) \quad (1)$$

$y(z,t)$ is the monthly mean ozone anomaly time series at altitude z , $\beta_{1-10}(z,t)$ are the fitted coefficients and $\varepsilon(z,t)$ represents the residual term. QBO₁ and QBO₂ are two orthogonal components of the QBO calculated with principal component analysis. No lag is applied to the ENSO, sAOD and Solar F10.7 proxies. Data sources of these proxies are provided in Table 2. Regarding the trend terms, $L_{pre}(t)$, $L_{post}(t)$ and $Gap(t)$ are written as follows:

$$L_{pre}(t) = \begin{cases} 1 & \text{if } t \leq t_1 \\ 0 & \text{if } t > t_1 \end{cases}$$

$$L_{post}(t) = \begin{cases} 0 & \text{if } t \leq t_2 \\ 1 & \text{if } t > t_2 \end{cases}$$

$$Gap(t) = \begin{cases} 0 & \text{if } t \leq t_1 \\ 1 & \text{if } t > t_1 \text{ and } t \leq t_2 \\ 0 & \text{if } t \geq t_2 \end{cases}$$

t_1 corresponds to January 1st 1997 and t_2 to January 1st 2000.

The Cochrane and Orcutt (1949) method is applied to correct for autocorrelation of residuals. Several improvements were made to the LOTUS model used in this work compared to the version used in LOTUS19. The new version (v0.8.0) of the model includes a seasonal variation of the regressed coefficient $\beta_k(z,t)$ for the various predictors, by adding Fourier components in the model as following:

$$\beta_k(z,t) = \beta_{k0}(z) + \sum_{i=1}^2 \beta_{k1i}(z) \sin\left(\frac{2\pi it}{12}\right) + \sum_{i=1}^2 \beta_{k2i}(z) \cos\left(\frac{2\pi it}{12}\right) \quad (2)$$

The new v0.8.0 LOTUS model includes also a new AOD predictor from the GloSSAC climatology instead of the NASA Goddard Institute for Space Science (GISS) AOD used before. For detailed information on the LOTUS regression model and its new features since LOTUS19, see http://argpages.usask.ca/docs/LOTUS_regression/index.html.

The improved LOTUS model with seasonal variation of fitted coefficients was applied to the merged satellite records included in the study over the 1985 – 2020 period for all latitude bins and altitude/pressure levels (depending on the native coordinates of the time series). It was also applied to each vertical level of the ground-based data used for comparison at the selected NDACC stations and to the gridded satellite data (e.g. MEGRIDOP, SWOOSH and SBUV MOD) in the vicinity of the stations. Trends from CCMI-1 model data were obtained from the updated v.0.8.0 LOTUS model in the same way.

Supprimé: ¶

Supprimé: +------(1)¶

Mis en forme : Police :Symbol

Supprimé: In equation (1), the coefficients A – J are the result of the regression and

Supprimé: represent

Supprimé:

Supprimé: from

Supprimé: .

Supprimé:

Supprimé: F10.7

Supprimé: , which are taken from <http://www.esrl.noaa.gov/psd/enso/mei/table.html> and http://www.spaceweather.ca/data-donnee/sol_flux/sx-5-mavg-eng.php respectively.

Mis en forme : Gauche

Mis en forme : Police :Non Italique

Mis en forme : Police :Non Italique

Mis en forme : Police :Italique

Mis en forme : Exposit

Mis en forme : Police :Italique

Mis en forme : Exposit

Supprimé: For the trend terms, linear components (e.g. $Linear_{pre}$ and $Linear_{post}$) and constant (i.e., C_1 and C_2) terms are applied for the “pre” (i.e., before Jan 1997) and “post” (Jan 2000 – Dec 2000) time periods, with zero values outside these periods, which makes it possible for the two trends to be treated independently of each other. A constant (C_3) term is added to fill the 3-year gap between the “pre” and “post” time periods. $\varepsilon(t)$ represents the residual term.

Mis en forme : Espace Avant : 0 pt

Supprimé: See http://argpages.usask.ca/docs/LOTUS_regression/index.html and LOTUS19 for more information on the LOTUS model. ¶ GloSSAC is taken from <https://asdc.larc.nasa.gov/project/GloSSAC>.

Supprimé: predictors (e.g. ENSO, Solar cycle and QBO)

Supprimé: which improves the characterization of ozone variability by the model (section 5) and

Supprimé: .

Supprimé: GloSSAC is taken from <https://asdc.larc.nasa.gov/project/GloSSAC>.¶

Supprimé: proxies

Supprimé: supersite

4. Results

4.1 Global trends as a function of altitude/pressure and latitude from merged satellite records

415 Figure 1 displays the trend results for the post-2000 period (i.e., from 01/2000 to 12/2020) retrieved from the merged satellite records for all latitude bands and vertical bins. The upper panel shows trend results for the merged satellite records provided in pressure levels (i.e., SBUV MOD, SBUV COH, GOZCARDS and SWOOSH), while trends from SAGE-CCI-OMPS, SAGE-SCIAMACHY-OMPS and SAGE-OSIRIS-OMPS provided in altitude levels are displayed the bottom panel. Dotted areas indicate trend values that are not significant at 2-sigma uncertainty. As in LOTUS19 (e.g., Figure 5.2 of the report), 420 positive and significant trend values are observed in the upper stratosphere for all datasets. Some discrepancies in the magnitude and latitude of the significant positive trends can be noticed among the records. In the upper panel, the SBUV MOD record show the largest positive trend values around 8 hPa and above 2hPa, while positive trends of the other records are observed above 7 – 5 hPa and are generally significant for all latitude bands. Both SBUV MOD and SBUV COH display non-significant trends in the tropical and subtropical latitudes (above 2 hPa in the SBUV COH case). SWOOSH and GOZCARDS, 425 which share similar individual satellite records, show similar trend patterns, with SWOOSH trend values slightly larger at 5 hPa in the midlatitudes. For records in the bottom panels, the various trends are also very similar in the upper stratosphere, with increasing trend values from left to right panels. It is also interesting to see significant positive trend values for most records except SBUV COH in the southern midlatitudes in the middle stratosphere (above ~25 km), while positive trends are not observed in this region in the northern hemisphere. In the lower stratosphere, trend values are generally negative but not significant, except in the lowermost stratosphere in the tropics, and especially in the bottom panel. These results are close to those of LOTUS19. The main feature of the zonally resolved trends is that most combined satellite records now show significant positive trends in vertical levels between ~ 5 – 2 hPa for all latitude bands. This was not the case in LOTUS19, 430 where trends from, e.g. GOZCARDS, SWOOSH, SAGE-CCI-OMPS for which the trends were not statistically significant in the tropics.

4.2 Trends over broad latitude bands

435 As in previous ozone profile trend studies, e.g. Harris et al. (2015), Steinbrecht et al. (2017) and LOTUS19, we also calculated trends over broad latitude bands, namely 60°S–35°S, 20°S–20°N, and 35°N–60°N. For GOZCARDS, SWOOSH, SBUV MOD, and SBUV COH, we first computed the deseasonalized monthly anomalies with respect to their own 1998–2008 climatology for each latitude and pressure bin, then averaged these anomalies over the broad latitude band with latitude weights, in a similar way as in LOTUS19. The SAGE-SCIA-OMPS, SAGE-CCI-OMPS and SAGE-OSIRIS-OMPS datasets were provided as deseasonalized records with the entire time period of the record used to compute the climatology. Ozone anomalies were averaged in a similar way as in the previous case. The LOTUS model was then applied to each broadband 440

Supprimé: (bottom)

Supprimé: (altitude)

Supprimé: respectively

Supprimé:

Supprimé: at

Supprimé: in the case of

Supprimé:

Supprimé: works

Supprimé: investigated

anomaly record. Figure 2 displays the results for the seven merged records, with each record plotted in its native vertical coordinate and with 2-sigma uncertainty. Results generally confirms the significant positive trends of ozone for all records in the upper stratosphere, i.e. between ~7 hPa and ~2 hPa in the three broad latitude bands, except for SBUV MOD in the tropics, where the trend is only slightly positive and not significant. Maximum positive trend for this record is seen at around 8 hPa in this region. Notwithstanding SBUV MOD behaviour in the tropics, the spread of trend values is more pronounced in the southern hemisphere where the lowest and largest positive trend values are obtained from SAGE-CCI-OMPS and SAGE-OSIRIS-OMPS respectively. At pressures larger than 10 hPa, trends are generally close to zero, except in the southern midlatitudes where some positive significant trend values are noticed, e.g. from SAGE-OSIRIS-OMPS, SAGE-SCIAMACHY-OMPS and SWOOSH. In the lowermost stratosphere, i.e. below 20 km, we see a hint of negative trends. This is most pronounced in the tropics but error bars are too large to conclude that there is a significant decrease. Note also the difference between the northern and southern hemisphere where negative trend values are larger in absolute value in the former albeit non-significant. Compared to LOTUS19 (e.g. Figure 5.6 of the report), the agreement between the records is much improved, especially in the upper stratosphere, e.g. due to the lower trend values of SBUV COH, which now agree quite well with the other records. Similarly, SBUV MOD trend values in the tropics, while still high near 8 hPa, are reduced relative to LOTUS19.

4.3 Combined trends

The various trend profiles in broad latitude bands were combined in order to facilitate comparison with LOTUS19 results and with CCMI-1 simulations. We adopted the same methodology as in LOTUS 19. The combined trend corresponds to the unweighted mean of the seven trend profiles shown in Figure 2. At pressures larger than 50 hPa level, the mean combines the results from five data records, since SBUV data in the lowermost stratosphere should not be considered. The average is calculated after converting the trends in altitude coordinate to pressure coordinate using a climatological ERA-INTERIM pressure – temperature profile over the period. For the combined trend uncertainty, we have to take into account the correlation between the individual trend estimates, which is due to the use of common underlying individual satellite records for some of them, e.g. SAGE II for many merged records, or OMPS-LP and the various SBUV time series in the case of SBUV MOD and SBUV COH records. The correlation also comes from similar atmospheric variability not characterized by the regression model (see LOTUS19 for more details). The variance of the mean is estimated as follows:

$$\sigma_{mean}^2 = \max \left(\frac{1}{N^2} \sum_{i,j} C_{i,j} \sigma_i \sigma_j, \frac{1}{n_{eff}} \sum_i \frac{(x_i - \bar{x})^2}{N-1} \right) \quad (3)$$

where σ_i is the uncertainty of individual trends x_i estimated from the fit, \bar{x} is the unweighted mean of the trends, N is the number of averaged records, $C_{i,j}$ are the correlation coefficients between the fit residuals of merged data records i and j , and n_{eff} is the effective number of independent values, evaluated as follows:

Supprimé: with respect to

Supprimé: It

Supprimé: e.g

Supprimé: (

Supprimé:)

Supprimé: Below

Supprimé: e

Supprimé: g

Supprimé: o

Supprimé: s

Supprimé:

Supprimé: For this process, w

Supprimé: Below

Supprimé: the

Supprimé: are

Supprimé: here

Supprimé: done

Supprimé: is also due to

Mis en forme : Normal, Espace Après : 0 pt, Interligne : simple

Supprimé: ¶

Supprimé: →→→(2)

Supprimé: s

Supprimé: for

$$n_{eff} = \frac{N^2}{\sum_{i,j=1}^N c_{ij}} \quad (4)$$

505 In equation 3, the first term on the right-hand side corresponds to the variance of the mean based on the classical propagation of errors for correlated variables (σ_{err}^2) and the second to the variance of the mean for n_{eff} independent estimates (σ_{obs}^2). The second term considers additional uncertainties in the trend average that are not identified in the first term, but might lead to different trend estimates, such as drifts in the individual time series. More information on this method is given in chapter 5 of LOTUS19 (section 5.3.1). Results of the combined trends from this study, called LOTUS22 hereafter, are displayed in Figure 3 (red curves) with comparison to the LOTUS19 combined trends (blue curve), see also Figure S1 for the comparison of LOTUS19 and LOTUS22 trends on both pre-1997 and post-2000 periods. In addition, Table S1 provides the correlation coefficients obtained for LOTUS22. The n_{eff} value for the seven merged ozone records is equal to 1.39, compared to 1.37 in LOTUS19, where six records were considered for the trend combination. Figure 3 shows that, compared to LOTUS19, the combined trend uncertainty is significantly smaller, especially in the upper stratosphere across all three broad latitude bands.

515 This confirms the previous findings that ozone is increasing in the upper stratosphere. The increase is somewhat larger in the NH with a maximum trend of ~2.2%/decade reached at ~2.2 hPa, compared to ~2.1%/decade at ~3.2 hPa in the SH and ~1.6%/decade at ~2.6 hPa in the tropics. Uncertainties of the combined trends are also smaller at pressures larger than 10 hPa, except in the southern hemisphere midlatitudes, where already mentioned positive trends retrieved from some of the records impact both the combined trend average and its uncertainty. It should be noted that slightly significant negative trends of

520 ~-1%/decade are retrieved in the tropics in the 30 – 40 hPa range. In the lowermost stratosphere, i.e. at pressures larger than 50 hPa, the LOTUS22 combined trends are negative and systematically larger in magnitude than the LOTUS19 derived trends. In the tropics, the trend uncertainty in our study increases, so that although the negative trends also increase in magnitude, they are not significant, as in LOTUS19. In the SH and NH lower stratosphere, the difference between both combined trends is very small, although LOTUS22 provides slightly more negative trends retrieved in this study with also smaller uncertainty. In all

525 cases and as noted previously, the uncertainties are large and preclude any definitive conclusion about ozone long-term changes in the lowermost stratosphere.

4.4 Trends over selected NDACC stations

530 The results of comparisons between ground-based and merged satellite ozone trends for the selected NDACC stations are displayed in Figure 4. The merged satellite records are MEGRIDOP, SWOOSH and SBUV MOD, for which longitudinally resolved data were provided. We use the satellite data in the grid cell closest to the stations for the trend computation. For the ground-based results in the Alpine stations, trends correspond to the average trend of the following records: (1) ozonesondes: Hohenpeissenberg, Payerne and OHP; (2) lidar: Hohenpeissenberg and OHP; (3) Umkehr: Arosa and OHP; (4) FTIR: Zugspitze and Jungfraujoch; (5) MWR: Payerne. For MWR trends, we used the Payerne record only, as some calibration

Supprimé: ¶
→

Mis en forme : Normal, Retrait : Gauche : 0 cm, Espace
Après : 0 pt, Interligne : simple

Supprimé: →(3)

Supprimé: 2

Supprimé: . We

Supprimé: below

Supprimé: -

Supprimé: e.g.,

Supprimé: below

Supprimé: with in the latter case

Supprimé: large

Supprimé: super-site

Supprimé: supersites

550 problems were found in the MWR Bern record. In LOTUS19, trends from ground-based instruments were compared to the combined trends from merged satellite records in broad latitude bands. This is different from the more direct comparison performed in the present study. Figure 4 shows a general good agreement between the ground-based and the gridded satellite trends for the NH Alpine and tropical stations, usually well within the respective uncertainties. Trend results differ more at the Lauder station. Results from the Alpine ground-based measurements reproduce quite well the trend patterns observed with the merged satellite records, e.g. an increase of about 2%/decade on average in the upper stratosphere, trend values around zero in the middle stratosphere, and mostly negative trends below 20 km, with large uncertainties. It is interesting to see that gridded satellite trend results, which rely on more individual data points, differ as much as the ground-based ones in the upper stratosphere.

560 At Mauna Loa/Hilo, similar patterns emerge also with (1) an ozone increase in most records in the upper stratosphere, except the MWR trend, (2) very small negative trends in the middle stratosphere that are most pronounced with the lidar record, and (3) larger negative trends in absolute values in the lower stratosphere below 20 km. For MEGRIDOP and the lidar, negative trends are significant in this altitude region. Compared to Figure 5.10 of LOTUS 19, ozonesondes and Umkehr results now show a better agreement with the other records. Regarding the MWR, it should be noted that there were major upgrades to the MLO MW instrument from 2015-2017, including the replacement of the filterbank spectrometer with an FFT spectrometer, which induced a gap in the dataset.

565 At Lauder, the trend profiles still show larger differences, which were not reduced compared to LOTUS19, despite the homogenization of the ozonesondes and Umkehr records. Trend results from MWR data are not shown as the record was not extended after 2016. Most records show positive trends in the upper stratosphere except the lidar and Umkehr records. Gridded satellite data trends are in between the ground-based ones in this altitude range. In the lowermost stratosphere, negative trends are obtained from most records except from Umkehr. Largest negative trends in absolute values are retrieved from the ozonesondes and lidar records.

570 In conclusion, trend comparison between ground-based and longitudinally resolved satellite records provide a similar picture as in the previous section, more specifically (1) positive and significant ozone trends in the upper stratosphere from most of the records, (2) negligible trends in the middle stratosphere and (3) negative trends below 20 km, which are statistically significant in lidar and ozonesondes records.

4.5 Comparison with trends derived from CCMI-1 simulations

575 The comparison of CCMI-1 trend results and merged satellite combined trends (LOTUS22) is displayed in Figure 5, which also includes results from LOTUS19. In the figure, multi-model mean trend estimates from the CCMI-1 REF-C2 simulations are represented by the black line, and the 2-sigma uncertainty of the multi-model mean trend estimates is given by the grey

Supprimé: trends

Supprimé: merged satellite combined trends, which makes direct

Supprimé: with

Supprimé: more difficult

Supprimé: are more scattered

Supprimé: by

Supprimé: instruments

Supprimé:

Mis en forme : Espace Avant : 6 pt

Supprimé: one,

Supprimé: a common behaviour of

Supprimé: is more difficult to detect due the discrepancies in the results, ...

Supprimé: It should be noted that MW trend results are similar to those displayed in LOTUS19 as the record was not extended after 2016. ...

Supprimé: with

Supprimé: also

595 envelope. Red (blue) solid curves show LOTUS22 (LOTUS19) combined trends, respectively, with corresponding 2-sigma
 uncertainties (Eq. 3) represented by the dashed lines. The individual model trends (from a total of 16 CCMI-1 models, as in
 LOTUS19) are estimated using the ILT regression method in the same way as for the satellite data and are then combined into
 a multi-model mean. Model simulations are updated to include 2020, and the necessary proxies are calculated directly from
 the pertaining individual model simulations where appropriate (e.g., QBO, ENSO), or taken from the external forcings
 600 provided to the modelling groups. Figure 5 shows very good agreement between the CCMI-1 and LOTUS22 trend estimates
 in the upper stratosphere, both regarding the average trend values and the uncertainties. In the northern hemisphere, the
 agreement is improved compared to the LOTUS19 results. In the middle stratosphere, some differences are observed, e.g., in
 the southern hemisphere, where the LOTUS22 trends are positive with large uncertainties, in contrast to CCMI-1 and
 LOTUS19 observed trends that are very close to zero. In the lowermost stratosphere, observed and simulated trend values
 605 diverge at midlatitudes of both hemispheres, with positive non-significant trends from the CCMI-1 simulations, and negative,
 but also non-significant trends from LOTUS 19 and LOTUS22. Agreement is better in the tropics, where CCMI and both
 LOTUS studies show small negative trends although non-significant. In conclusion, the LOTUS22 trend results confirm the
 findings of LOTUS19, and provide observed trends that are consistent in magnitude and uncertainty range with simulated
 ozone trends from the CCMI initiative.

610

5. Discussion

In this section, we discuss in more detail the differences in trend estimates between LOTUS19 and LOTUS22. Compared to
 Figure 5.6 of the LOTUS19 report, the agreement between the merged data sets is improved, resulting in smaller uncertainties
 in the combined trend results. This improvement is to a large extent driven by a better agreement of trend results between both
 615 SBUV merged records, e.g. SBUV MOD and SBUV COH, due to revised inter-calibration of the individual SBUV records
 and the addition of OMPS in both merged datasets. Trends from these datasets now agree better with other records' trends,
 especially in the upper stratosphere. We can have a better understanding of the improvement in trend uncertainties from Figure
 6, which displays the square root of both terms included in the variance of the combined trend (Eq. 3), i.e. the term linked to
 error propagation (left term on the right side in Eq. 3, σ_{err} , dashed line) and that linked to the standard error of the trend sample
 620 (right term of the right term in Eq. 3, σ_{obs} , dotted line) for LOTUS22 in red and LOTUS19 in blue. The three panels correspond
 to the three broad latitude bands. As indicated in Eq. 3, the reported uncertainty value is the maximum of both terms as a
 function of pressure. We can see from the figure that in both studies the uncertainty is dominated by the σ_{obs} term in the upper
 and middle stratosphere and by the σ_{err} term in the lowermost stratosphere. Reduction in the σ_{obs} term from LOTUS19 to
 LOTUS22 is clearly visible in the figure. It is most pronounced in the tropics around 5 hPa, then in the SH in the 5 – 1 hPa
 625 pressure range, and in the NH to a somewhat lower extent in the same pressure range. The LOTUS22 σ_{obs} term is reduced also

Supprimé: predictors

Supprimé: remarkable

Supprimé: T

Supprimé: average

Supprimé: in the northern hemisphere

Supprimé: larger

Supprimé: all results

Supprimé: and a better agreement with

Supprimé:

Supprimé: 2

Supprimé: 2

Supprimé: corresponds to

Supprimé: the

Supprimé: term

Supprimé: the

Supprimé: term

in the tropical middle stratosphere but it is increased with respect to LOTUS19 at southern midlatitudes at about the same pressure range. This is due to the already mentioned positive trends retrieved by most of the records in this latitude and pressure range. In the lowermost stratosphere, the dominance of the σ_{err} term is expected due to the large uncertainty retrieved for the trends of the majority of the records in this altitude range.

Another factor that allowed us to reduce the uncertainty of our trend retrieval is the improvement of the LOTUS trend model that includes now seasonal terms for the fitted coefficients (as described in Section 3). Thanks to this improvement, the model now better fits the ozone variability of the various records. This is shown in Figure 7, which displays adjusted R^2 values of the regression in broad latitude bands for the records with altitude as vertical coordinate on the left and those with pressure on the right. Adjusted R^2 provides an estimation of the amount of variance in the monthly data explained by the regression model. It is an indicator of the goodness of the fit. Displayed R^2 values correspond to the average of R^2 profiles for the three latitude bands considered in the study. Solid (dashed) lines show R^2 values retrieved from the LOTUS model with seasonal (non-seasonal) variation of the fitted coefficients. Larger R^2 values are systematically obtained with seasonal variation of the fitted coefficients.

Using both the improved merged satellite records for this study and the new version of the LOTUS regression model, we can further constrain ozone trends in the various altitude and latitude regions.

6. Conclusion

This study provides an updated evaluation of stratospheric ozone profile trends in the 60°S – 60°N latitude range from up-to-date merged satellite and ground-based records. Some satellite data series were improved with respect to those used in the previous assessments (WMO, 2018 and LOTUS19), e.g., SBUV MOD and SBUV COH, which resulted in a better agreement between trends from both records and with the other ones used in the study. Additional records that were absent from the LOTUS19 study are included, e.g., the SAGE-SCIAMACHY-OMPS and MEGRIDOP records. Regarding ground-based data, we use ECC homogenized ozonesonde and Umkehr data reprocessed with optimized homogenization technique that were not available previously. An updated version of the LOTUS regression trend model allows us to improve the fit of ozone variability for the various records. With these improvements, we can draw the following conclusions:

- The increase of ozone in the upper stratosphere at pressures lower than 5 hPa is confirmed, with a clearer recovery in the southern hemisphere compared to LOTUS19. In this altitude region, combined satellite trends are significant in the three broad latitude bands considered, i.e., southern and northern midlatitudes and tropics.
- In the middle stratosphere, i.e., between 50 and 10 hPa, we see the emergence of new signals that will need to be confirmed in the future: an increase of ozone in the southern hemisphere midlatitudes of about 1.2%/decade, though

Supprimé: was

Supprimé: predictors

Supprimé: , as

Supprimé: displaying

Supprimé: 3

Supprimé: predictors

Supprimé: proxies

Supprimé: thus

Supprimé: new

Supprimé:

Supprimé: e.g

Supprimé: .

685

non-significant, and an ozone decrease in the tropics that is (just) significant at around 35 hPa. In the northern hemisphere, ozone trends are close to zero in this altitude range.

690

- In the lower and **lowermost** stratosphere, negative ozone trends are obtained for all latitude bands, as in LOTUS19. Trends are negligible in the southern midlatitudes. The trends amount to about -2%/decade in the tropics and **are non-significant** due to the large uncertainties. Negative ozone trends are also obtained in the northern hemisphere, mainly **at pressures larger than** 70 hPa. They reach -2%/decade at 100 hPa but are also non-significant.
- Comparison of combined trends with those derived from updated CCMI-1 simulations in broad latitude bands show **very good** agreement in the upper stratosphere, both in trend magnitude and uncertainty. Larger differences are seen below 10 hPa, 40 hPa and 60 hPa in the tropics, SH and NH midlatitudes, respectively, with the CCMI-1 trends being generally **more positive** than the satellite ones. Differences are most pronounced in the NH midlatitudes, where average satellite trends are negative, while those of CCMI-1 are positive. **Due to the large uncertainty in both cases,** these differences are non-significant.

695

Compared to LOTUS19, we **find** a better agreement between trends from ground-based measurements and from satellite records, especially in the tropics and in the Alpine stations of the northern hemisphere, **for most of the records**. This can be due to **both** the use of longitudinally resolved satellite data and **to** improved ground-based and satellite records. Trends **values** are more scattered at the Lauder station. **The differences between trends from longitudinally resolved satellite records and some ground-based time series at the selected NDACC stations, warrant more detailed analyses in the future, focusing on the possible biases between the records.**

700

The ozone recovery signal that is mainly observed in the upper stratosphere has an influence on total ozone trends. A recent study indicates a total ozone recovery of 0.5 ± 0.2 %/decade (~ 1.5 DU/decade) since 1996 (Weber et al., 2022). However, total column ozone evolution is influenced even more by trends in the lower stratosphere and also by tropospheric ozone trends. The latter are estimated to be of the order of ~ 1.5 DU/decade (e.g. Gaudel et al., 2018; Ziemke et al., 2019) with larger changes found in the tropical regions. The precise impact of stratospheric and tropospheric partial ozone column trends on total column ozone trends needs thus further evaluation.

705

The consistency of ozone profile trends found in this study demonstrates that the global ozone observing system is still robust, thanks to the **continuing and improved** satellite and ground-based records. This allows us to quite accurately evaluate long-term ozone changes in the stratosphere. The cause of **some** larger discrepancies between combined satellite and CCMI trends in the lower stratosphere will have to be further investigated, as the new set of CCMI-2022 simulations become available (Plummer et al., 2021). More generally, the study of ozone trends in this region **may require a special focus with geophysically-based coordinate systems, based on e.g. tropopause level or equivalent latitude (Millan et al., 2021)** in order to better constrain ozone variability and provide a more accurate trend evaluation.

710

Supprimé: most

Supprimé: -

Supprimé: can still be considered

Supprimé: below

Supprimé: remarkable

Supprimé: larger

Supprimé: but d

Supprimé: observe

Supprimé: Examination of the

Supprimé: and merged satellite trends

Supprimé: this location and other

Supprimé:

Supprimé: should be the subject

Supprimé: of a more detail

Supprimé: ed study in the future.

Supprimé: is

Supprimé: also

Supprimé: continuous

Supprimé: improvement of

Supprimé: set of

Supprimé: s

Supprimé: e.g.

Supprimé: new

Supprimé: s as proposed by the SPARC Observed Composition Trends And Variability in the Upper Troposphere and Lower Stratosphere (OCTAV-UTLS) activity

Data availability

The satellite, ground-based and CCMI monthly mean data are available at the LOTUS FMI depository: ftp://ftp2.fmi.fi/Phase-2_2022-2019.

745

Authors contribution

NA computed the satellite and ground-based trends, VS computed the combined trends and provided Figure 4, DH provided Figures 3 and 6, KT provided the trends from CCMI-1 simulations, DD, DZ and RD provided and maintain the LOTUS regression model. Other co-authors contributed with satellite or ground-based data. The results of the study were discussed by all the co-authors. The manuscript was written by SGB with supporting comments by all co-authors.

750

Competing interests

Two co-authors (SGB and IP) are co-organizers of the special issue “Atmospheric ozone and related species in the early 2020s: latest results and trends (ACP/AMT inter-journal SI), 2021”.

755

Acknowledgements

The ground-based data used in this publication were obtained as part of the Network for the Detection of Atmospheric Composition Change (NDACC) and are available through the NDACC website www.ndacc.org. Optimized Umkehr data are available at <https://gml.noaa.gov/aftp/data/ozwv/Dobson/AC4/Umkehr/Optimized/>. Homogenized ozonesonde data were obtained through the HEGIFTOM Focus Working Group: <https://hegiftom.meteo.be/>. NA’s work was supported by a postdoctoral fellowship from Institut National des Sciences de l’Univers du Centre National de la Recherche Scientifique (INSU-CNRS). Work at the Jet Propulsion Laboratory, California Institute of Technology, was performed under contract with the National Aeronautics and Space Administration (80NM0018D004); we gratefully acknowledge the efforts of Ray Wang, John Anderson, and Ryan Fuller towards the initial GOZCARDS ozone data record, and its subsequent updates. IP and JW were supported by NOAA Climate Program Office’s Atmospheric Chemistry, Carbon Cycle, and Climate program, grant number NA19OAR4310169 (CU)/ NA19OAR4310171 (UMD). The SBUV Merged Ozone Data Set was constructed under the NASA MeaSURES Project and is maintained under NASA WBS 479717 (Long Term Measurement of Ozone). CA acknowledges the support of the University and State of Bremen, and the funding from DAAD and the Living Planet Fellowship SOLVE. The SAGE-CCI-OMPS and MEGRIDOP datasets are created in the framework of ESA Ozone-CCI + project. The work of VS was supported by European Space Agency (Ozone_cci+ project, contract 4000126562/19/I-NB), the EU Copernicus Climate Change Service for Atmospheric Composition ECVs (contract C3S_312b_Lot2_DLR_2018SC1), and the Academy of Finland (the Centre of Excellence of Inverse Modelling and Imaging (decision 336798). The multi-decadal

760

765

770

Supprimé: <https://igacproject.org/hegiftom-focus-working-group>

775 monitoring program of ULiège at the Jungfraujoch station has been primarily supported by the F.R.S.-FNRS and BELSPO (both in Brussels, Belgium) and by the GAW-CH programme of MeteoSwiss. The International Foundation High Altitude Research Stations Jungfraujoch and Gornergrat (HFSJG, Bern) supported the facilities needed to perform the FTIR observations.

780 **References**

- Arosio, C., Rozanov, A., Malinina, E., Weber, M., and Burrows, J. P.: Merging of ozone profiles from SCIAMACHY, OMPS and SAGE II observations to study stratospheric ozone changes, *Atmos. Meas. Tech.*, 12, 2423–2444, <https://doi.org/10.5194/amt-12-2423-2019>, 2019.
- 785 Ball, W. T., Alsing, J., Mortlock, D. J., Staehelin, J., Haigh, J. D., Peter, T., Tummon, F., Stübi, R., Stenke, A., Anderson, J., Bourassa, A., Davis, S. M., Degenstein, D., Frith, S., Froidevaux, L., Roth, C., Sofieva, V., Wang, R., Wild, J., Yu, P., Ziemke, J. R., and Rozanov, E. V.: Evidence for a continuous decline in lower stratospheric ozone offsetting ozone layer recovery, *Atmos. Chem. Phys.*, 18, 1379–1394, <https://doi.org/10.5194/acp-18-1379-2018>.
- Ball, W. T., Chiodo, G., Abalos, M., Alsing, J., and Stenke, A.: Inconsistencies between chemistry–climate models and observed lower stratospheric ozone trends since 1998, *Atmos. Chem. Phys.*, 20, 9737–9752, <https://doi.org/10.5194/acp-20-9737-2020>, 2020.
- 790 Bernet, L., von Clarmann, T., Godin-Beekmann, S., Ancellet, G., Maillard Barras, E., Stübi, R., Steinbrecht, W., Kämpfer, N., and Hocke, K.: Ground-based ozone profiles over central Europe: incorporating anomalous observations into the analysis of stratospheric ozone trends, *Atmos. Chem. Phys.*, 19, 4289–4309, <https://doi.org/10.5194/acp-19-4289-2019>, 2019.
- Bourassa, A. E., Roth, C. Z., Zawada, D. J., Rieger, L. A., McLinden, C. A., and Degenstein, D. A.: Drift-corrected Odin-OSIRIS ozone product: algorithm and updated stratospheric ozone trends, *Atmos. Meas. Tech.*, 11, 489–498, <https://doi.org/10.5194/amt-11-489-2018>, 2018.
- 795 Chipperfield, M. P., Dhomse, S., Hossaini, R., Feng, W., Santee, M. L., Weber, M., Burrows, J. P., Wild, J. D., Loyola, D., and Coldewey-Egbers, M.: On the Cause of Recent Variations in Lower Stratospheric Ozone, *Geophys. Res. Lett.*, 45, 5718–5726, <https://doi.org/10.1029/2018GL078071>
- 800 Cochran, D., and G. H Orcutt, Application of Least Squares Regression to Relationships Containing Auto-Correlated Error Term, *Journal of the American Statistical Association*, 1949, <https://doi.org/10.1080/01621459.1949.10483290>
- Davis, S. M., Rosenlof, K. H., Hassler, B., Hurst, D. F., Read, W. G., Vömel, H., Selkirk, H., Fujiwara, M., and Damadeo, R.: The Stratospheric Water and Ozone Satellite Homogenized (SWOOSH) database: A long-term database for climate studies, *Earth Syst. Sci. Data*, doi:10.5194/essd-8-461-2016, 2016.

- 805 de Laat, A. T. J., van der A, R. J., and van Weele, M.: Tracing the second stage of ozone recovery in the Antarctic ozone-hole
with a "big data" approach to multivariate regressions, *Atmos. Chem. Phys.*, 15, 79–97, [https://doi.org/10.5194/acp-15-79-](https://doi.org/10.5194/acp-15-79-2015)
2015, 2015.
- Eyring, V., Arblaster, J.M., Cionni, I., Sedláček, J., Perlwitz, J., Young, P.J., Bekki, S., Bergmann, S., Cameron-Smith, P.,
Collins, W.J., Faluvegi, G., Gottschaldt, K.-D., Horowitz, L.W., Kinnison, D.E., Lamarque, J.-F., Marsh, D.R., Saint-
810 Martin, D., Shindell, D.T., Sudo, K., Szopa, S., and Watanabe, S.: Long-term ozone changes and associated climate impacts
in CMIP5 simulations, *J. Geophys. Res. Atmos.*, 118, doi:10.1002/jgrd.50316, 2013.
- Frith, S. M., Stolarski, R. S., Kramarova, N. A., and McPeters, R. D.: Estimating uncertainties in the SBUV Version 8.6 merged
profile ozone data set, *Atmos. Chem. Phys.*, 17, 14695–14707, <https://doi.org/10.5194/acp-17-14695-2017>, 2017.
- Frith, S. M., Bhartia, P. K., Oman, L. D., Kramarova, N. A., McPeters, R. D., and Labow, G. J.: Model-based climatology of
815 diurnal variability in stratospheric ozone as a data analysis tool, *Atmos. Meas. Tech.*, 13, 2733–2749,
<https://doi.org/10.5194/amt-13-2733-2020>, 2020.
- Garcia, R. R. and Randel, W. J.: Acceleration of the Brewer–Dobson circulation due to increases in greenhouse gases, *J.*
Atmos. Sci., 65, 2731–2739, <https://doi.org/10.1175/2008JAS2712.1>, 2008.
- Godin-Beekmann, S., J. Porteneuve, A. Garnier, Systematic DIAL ozone measurements at Observatoire de Haute-Provence,
820 *J. Env. Monitoring*, 5, 57-67, 2003
- Hase, F., Blumenstock, T., and Paton-Walsh, C., 1999: Analysis of the instrumental line shape of high-resolution Fourier
transform IR spectrometers with gas cell measurements and new retrieval software, *Appl. Opt.*, 38, 3417-3422,
doi:10.1364/AO.38.003417.
- Harris, N. R. P., Hassler, B., Tummon, F., Bodeker, G. E., Hubert, D., Petropavlovskikh, I., Steinbrecht, W., Anderson, J.,
825 Bhartia, P. K., Boone, C. D., Bourassa, A., Davis, S. M., Degenstein, D., Delcloo, A., Frith, S. M., Froidevaux, L., Godin-
Beekmann, S., Jones, N., Kurylo, M. J., Kyrölä, E., Laine, M., Leblanc, S. T., Lambert, J.-C., Liley, B., Mahieu, E.,
Maycock, A., de Mazière, M., Parrish, A., Querel, R., Rosenlof, K. H., Roth, C., Sioris, C., Staehelin, J., Stolarski, R. S.,
Stübi, R., Tamminen, J., Vigouroux, C., Walker, K. A., Wang, H. J., Wild, J., and Zawodny, J. M.: Past changes in the
vertical distribution of ozone – Part 3: Analysis and interpretation of trends, *Atmos. Chem. Phys.*, 15, 9965–9982,
830 <https://doi.org/10.5194/acp-15-9965-2015>
- Hegglin, M. I., Lamarque, J.-F., Duncan, B., Eyring, V., Gettelman, A., Hess, P., Myhre, G., Nagashima, T., Plummer, D.,
Ryerson, T., Shepherd, T., and Waugh, D.: Report on the IGAC/SPARC Chemistry-Climate Model Initiative (CCMI) 2015
science workshop, SPARC Newsletter, 46, 37–42, 2016.
- Kramarova N., Bhartia P.K., Huang L-K., Ziemke J., Frith S. M., McPeters R., Labow G., Haffner D., Stolarski R., DeLand
835 M., A new approach to cross-calibrate satellite instruments”, *to be submitted* to *Atmospheric Measurement Techniques*,
2022.

Mis en forme : Français

Mis en forme : Français

- Leblanc, T., Robert J. Sica, Joanna A. E. van Gijsel, Sophie Godin-Beekmann, Alexander Haefele, Thomas Trickl, Guillaume Payen, and Gianluigi Liberti, Proposed standardized definitions for vertical resolution and uncertainty in the NDACC lidar ozone and temperature algorithms – Part 2: Ozone DIAL uncertainty budget. *Atmospheric Measurement Techniques*, European Geosciences Union, 2016, 9 (8), pp.4051-4078. <10.5194/amt-9-4051-2016>
- 840 Maillard Barras, E., Haefele, A., Nguyen, L., Tummon, F., Ball, W. T., Rozanov, E. V., Rüfenacht, R., Hocke, K., Bernet, L., Kämpfer, N., Nedoluha, G., and Boyd, I.: Study of the dependence of stratospheric ozone long-term trends on local solar time, *Atmos. Chem. Phys.* 20, 8453–8471, <https://doi.org/10.5194/acp-20-8453-2020>, 2020
- 845 Millan, L. F., G. Manney, and Z. D. Lawrence, Reanalysis intercomparison of potential vorticity and potential-vorticity-based diagnostics, *Atmos. Chem. Phys.*, 21, 5355–5376, 2021, <https://doi.org/10.5194/acp-21-5355-2021>
- Morgenstern, O., Hegglin, M.I., Rozanov, E., O'Connor, F.M., Abraham, N.L., Akiyoshi, H., Archibald, A.T., Bekki, S., Butchart, N., Chipperfield, M.P., Deushi, M., Dhomse, S.S., Garcia, R.R., Hardiman, S.C., Horowitz, L.W., Jöckel, P., Josse, B., Kinnison, D., Lin, M., Mancini, E., Manyin, M.E., Marchand, M., Marécal, V., Michou, M., Oman, L.D., Pitari, G., Plummer, D.A., Revell, L.E., Saint-Martin, D., Schofield, R., Stenke, A., Stone, K., Sudo, K., Tanaka, T.Y., Tilmes, S., Yamashita, Y., Yoshida, K., and Zeng, G.: Review of the global models used within phase 1 of the Chemistry–Climate Model Initiative (CCMI), *Geosci. Model Dev.*, 10, 639-671, <https://doi.org/10.5194/gmd-10-639-2017>, 2017.
- 850 Newman, P. A., Daniel, J. S., Waugh, D. W., and Nash, E. R.: A new formulation of equivalent effective stratospheric chlorine (EESC), *Atmos. Chem. Phys.*, 7, 4537–4552, <https://doi.org/10.5194/acp-7-4537-2007>, 2007.
- Orbe, C., Wargan, K., Pawson, S., & Oman, L. D.: Mechanisms linked to recent ozone decreases in the Northern Hemisphere lower stratosphere. *Journal of Geophysical Research: Atmospheres*, 125, e2019JD031631. <https://doi.org/10.1029/2019JD031631>, 2020.
- 855 Pazmiño, A., Godin-Beekmann, S., Hauchecorne, A., Claud, C., Khaykin, S., Goutail, F., Wolfram, E., Salvador, J., and Quel, E.: Multiple symptoms of total ozone recovery inside the Antarctic vortex during austral spring, *Atmos. Chem. Phys.*, 18, 7557–7572, <https://doi.org/10.5194/acp-18-7557-2018>.
- 860 Petropavlovskikh, I., Godin-Beekmann, S., Hubert, D., Damadeo, R., Hassler, B., and Sofieva, V.: SPARC/IO3C/GAW report on Long-term Ozone Trends and Uncertainties in the Stratosphere, SPARC/IO3C/GAW, SPARC Report No. 9, WCRP-17/2018, GAW Report No. 241, <https://doi.org/10.17874/f899e57a20b>, 2019.
- Petropavlovskikh, I., Miyagawa, K., McClure-Beegle, A., Johnson, B., Wild, J., Strahan, S., Wargan, K., Querel, R., Flynn, L., Beach, E., Ancellet, G., and Godin-Beekmann, S.: Optimized Umkehr profile algorithm for ozone trend analyses, *Atmos. Meas. Tech. Discuss.* [preprint], <https://doi.org/10.5194/amt-2021-203>, in review, 2021.
- 865 Plummer, D., et al., CCMI-2022: A new set of Chemistry–Climate Model Initiative (CCMI) Community Simulations to Update the Assessment of Models and Support Upcoming Ozone Assessment Activities, in SPARC newsletter, July 2021, https://www.sparc-climate.org/wp-content/uploads/sites/5/2021/07/SPARCnewsletter_Jul2021_web.pdf

Supprimé: Maillard-Barras Eliane, Dominique Ruffieux & Klemens Hocke (2009) Stratospheric ozone profiles over Switzerland measured by SOMORA, ozonesonde and MLS/AURA satellite, *International Journal of Remote Sensing*, 30:15-16, 4033-4041, DOI: 10.1080/01431160902821890

- Rodgers, C. D.: Inverse Methods for Atmospheric Sounding: Theory and Practice, World Scientific Publishing Co. Pte. Ltd, Singapore, <https://doi.org/10.1142/3171>, 2000.
- 875 Smit, H. G. J. & the Panel for the Assessment of Standard Operating Procedures for Ozonesondes (ASOPOS 2.0) (2021). Ozonesonde Measurement Principles and Best Operational Practices, World Meteorological Organization. GAW Report, 268. [Available at https://library.wmo.int/doc_num.php?explnum_id=10884]
- Sofieva, V. F., Kyrölä, E., Laine, M., Tamminen, J., Degenstein, D., Bourassa, A., Roth, C., Zawada, D., Weber, M., Rozanov, A., Rahpoe, N., Stiller, G., Laeng, A., von Clarmann, T., Walker, K. A., Sheese, P., Hubert, D., van Roozendael, M., Zehner, C., Damadeo, R., Zawodny, J., Kramarova, N., and Bhartia, P. K.: Merged SAGE II, Ozone_cci and OMPS ozone profile dataset and evaluation of ozone trends in the stratosphere, *Atmos. Chem. Phys.*, 17, 12533–12552, <https://doi.org/10.5194/acp-17-12533-2017>
- 880 Sofieva, V. F., Szelag, M., Tamminen, J., Kyrölä, E., Degenstein, D., Roth, C., Zawada, D., Rozanov, A., Arosio, C., Burrows, J. P., Weber, M., Laeng, A., Stiller, G. P., von Clarmann, T., Froidevaux, L., Livesey, N., van Roozendael, M., and Retscher, C.: Measurement report: regional trends of stratospheric ozone evaluated using the Merged GRidded Dataset of Ozone Profiles (MEGRIDOP), *Atmos. Chem. Phys.*, 21, 6707–6720, <https://doi.org/10.5194/acp-21-6707-2021>, 2021
- Solomon, S., Ivy, D. J., Kinnison, D., Mills, M. J., Neely, R. R., and Schmidt, A.: Emergence of healing in the Antarctic ozone layer, *Science*, 353, 269–274, <https://doi.org/10.1126/science.aae0061>
- 890 Steinbrecht, W., Froidevaux, L., Fuller, R., Wang, R., Anderson, J., Roth, C., Bourassa, A., Degenstein, D., Damadeo, R., Zawodny, J., Frith, S., McPeters, R., Bhartia, P., Wild, J., Long, C., Davis, S., Rosenlof, K., Sofieva, V., Walker, K., Rahpoe, N., Rozanov, A., Weber, M., Laeng, A., von Clarmann, T., Stiller, G., Kramarova, N., Godin-Beekmann, S., Leblanc, T., Querel, R., Swart, D., Boyd, I., Hocke, K., Kämpfer, N., Maillard Barras, E., Moreira, L., Nedoluha, G., Vigouroux, C., Blumenstock, T., Schneider, M., Garcia, O., Jones, N., Mahieu, E., Smale, D., Kotkamp, M., Robinson, J., Petropavlovskikh, I., Harris, N., Hassler, B., Hubert, D., and Tummon, F.: An update on ozone profile trends for the period 2000 to 2016, *Atmos. Chem. Phys.*, 17, 10675–10690, <https://doi.org/10.5194/acp-17-10675-2017>
- Szelag, M. E., Sofieva, V. F., Degenstein, D., Roth, C., Davis, S., and Froidevaux, L.: Seasonal stratospheric ozone trends over 2000–2018 derived from several merged data sets, *Atmos. Chem. Phys.*, 20, 7035–7047, <https://doi.org/10.5194/acp-20-7035-2020>, 2020.
- 900 Vigouroux, C., Blumenstock, T., Coffey, M., Errera, Q., Garcia, O., Jones, N. B., Hannigan, J. W., Hase, F., Liley, B., Mahieu, E., Mellqvist, J., Notholt, J., Palm, M., Persson, G., Schneider, M., Servais, C., Smale, D., Thölix, L., and De Mazière, M.: Trends of ozone total columns and vertical distribution from FTIR observations at eight NDACC stations around the globe, *Atmos. Chem. Phys.*, 15, 2915–2933, <https://doi.org/10.5194/acp-15-2915-2015>, 2015.

Wargan, K., Orbe, C., Pawson, S., Ziemke, J. R., Oman, L. D., Olsen, M. A., Coy, L., and Emma Knowland, K.: Recent
905 Decline in Extratropical Lower Stratospheric Ozone Attributed to Circulation Changes, *Geophys. Res. Lett.*, 45, 5166–
5176, <https://doi.org/10.1029/2018GL077406>

Wild, Jeannette, et al. "Ozone recovery as detected in NOAA Ground-Based and Satellite Ozone Measurements." AGU Fall
Meeting 2019. AGU, 2019.

WMO (World Meteorological Organization), Scientific Assessment of Ozone Depletion: 2010, Global Ozone Research and
910 Monitoring Project - Report No. 52, 516 pp., Geneva, Switzerland, 2011.

WMO: Scientific Assessment of Ozone Depletion: 2018, Global Ozone Research and Monitoring Project Report, World
Meteorological Organization, p. 588, Geneva, Switzerland, 2018.

Zawada, D. J., Rieger, L. A., Bourassa, A. E., and Degenstein, D. A.: Tomographic retrievals of ozone with the OMPS
LimbProfiler: algorithm description and preliminary results, *Atmos.Meas. Tech.*, 11, 2375–2393,
915 <https://doi.org/10.5194/amt-11-2375-2018>, 2018.

Code de champ modifié

Station		Latitude/Longitude	Ozone profile records	Record length
Alpine	Hohenpeissenberg	47.8°N/11.0°E	Ozonesonde Lidar	1966-2020 1987-2020
	Payerne	46.8°N/6.9°E	Ozonesonde Microwave	1968-2020 2000-2020
	Zugspitze	47.40N/11.0°E	FTIR	1995-2020
	Arosa	46.7°N/9.7°E	Umkehr	1956-2020
	Jungfraujoch	46.5°N/7.9°E	FTIR	1995-2020
	OHP	43.9°N/5.7°E	Umkehr Lidar Ozonesonde	1984-2020 1985-2020 1991-2020
Mauna Loa	19.5°N/155.6°W	Umkehr Lidar Microwave	1984-2020 1993-2020 1995-2020	
Hilo	19.7°N/155.1°W	Ozonesonde	1982-2020	
Lauder	45°S/169.7°E	Umkehr Lidar Ozonesonde Microwave FTIR	1987-2020 1994-2020 1986-2020 1992-2016 2001-2020	

Supprimé: w

920 **Table 1.** Long-term ground-based NDACC ozone profile records used in the study.

Variable	Proxy	Source
Solar(t)	10.7 cm Solar Flux	https://spaceweather.gc.ca/forecast-prevision/solar-solaire/solarflux/sx-5-mavg-en.php
QBO ₁ (t), QBO ₂ (t)	Orthogonal components of the QBO calculated with Principal Component Analysis	http://www.geo.fu-berlin.de/met/ag/strat/produkte/qbo/qbo.dat
sAOD(t)	Stratospheric aerosol optical depth from GloSSAC	https://asdc.larc.nasa.gov/project/GloSSAC/GloSSAC_2.0
ENSO(t)	Multivariate El Niño/Southern Oscillation (ENSO) index (MEI.v2)	https://psl.noaa.gov/enso/mei/

Table 2: Sources of explanatory variables / proxy timeseries used in the LOTUS regression model.





Figure 1. Ozone profile trends from merged satellite records in percent per decade for the post-2000 period (Jan 2000 – Dec 2020). Grey stippling denotes results that are not significantly different from zero at the 2-sigma level. Data are presented on their native latitudinal grid and vertical coordinate








Figure 2. Merged satellite ozone trends with their 2-sigma uncertainties for the post-2000 period as estimated by the LOTUS regression model for latitude bands 60°S–35°S (left), 20°S–20°N (center), and 35°N–60°N (right). Colored lines are the trend estimates from the individual merged data sets on their native vertical grid.

950



955

Figure 3. Combined post-2000 ozone profile trend estimates and uncertainties (2-sigma) from the seven merged satellite records (below the 50 hPa level: five records, see text). Red (blue) solid line and light red (blue) shaded areas indicate LOTUS22 (LOTUS19) trend values and uncertainties.







Figure 4. Ozone profile trends for the post-2000 periods from selected ground-based NDACC stations. Left panel: SH Lauder station, middle panel: Tropical Mauna Loa and Hilo (ozonesondes) stations, right panel: NH Alpine stations (see text)



Supprimé:



Supprimé:




Supprimé:

Supprimé:

Supprimé: supersites


Supprimé: Top panel: NH Alpine stations (see text), middle panel: Tropical Mauna Loa and Hilo (ozonesondes) stations, bottom panel: SH Lauder station



975

Figure 5. Black line: multi-model mean ozone profile trend estimates from the CCMi REF-C2 simulations over three broad latitude bands (left 60°S – 35°S, center 20°S-20°N, right 35°N- 60°N). Grey envelope: 2 σ uncertainty of the multi-model mean trend estimates. Red (blue) lines and light red (light blue) dashed lines represent LOTUS22 (LOTUS19) average and 2-sigma uncertainty respectively.

980



990 **Figure 6.** Decomposition of error terms for the combined trend estimates: propagation of errors from fit residuals (σ_{err} , dashed lines) and standard error of the trend sample (σ_{obs} , dotted lines) for LOTUS22 in red and LOTUS19 in blue for the three broad latitude bands. Combined trend uncertainty is shown by circle symbols that indicate the maximum of σ_{err} and σ_{obs} in each case.




Figure 7. Adjusted R^2 values of the LOTUS regression model using seasonal (solid lines) and non-seasonal (dashed lines) variation of fitted coefficients in the LOTUS regression model (see text).

1000

Supprimé: predictors