

Measurement Report: Black carbon properties and concentrations in Southern Sweden urban and rural air – The importance of long-range transport

Erik Ahlberg^{1*}, Stina Ausmeel^{1,a}, Lovisa Nilsson¹, Mårten Spanne², Julija Pauraite³, Jacob Klenø Nøjgaard^{4,b}, Michele Bertò⁵, Henrik Skov⁴, Pontus Roldin¹, Adam Kristensson¹, Erik Swietlicki¹, Axel Eriksson^{6*}

¹Division of Nuclear Physics, Lund University, Box 118, 221 00 Lund, Sweden

²Environment Department, City of Malmö, 208 50 Malmö, Sweden

³Department of Environmental Research, Center for Physical Sciences and Technology, Savanorių ave. 231, 02300 Vilnius, Lithuania

⁴Department of Environmental Science, iClimate, Aarhus University, Roskilde, Denmark

⁵Laboratory of Atmospheric Chemistry, Paul Scherrer Institute (PSI), 5232 Villigen PSI, Switzerland

⁶Ergonomics and Aerosol Technology, Lund University, Box 118, 221 00 Lund, Sweden

^aNow at: Swedish Environmental Protection Agency, 10648 Stockholm, Sweden

^bNow at: National Research Centre for the Working Environment, 2100 Copenhagen, Denmark

Correspondence to: Erik Ahlberg (erik.ahlberg@nuclear.lu.se) and Axel Eriksson (axel.eriksson@design.lth.se)

Abstract. Soot, or black carbon (BC), aerosol is a major climate forcer with severe health effects. The impacts depend strongly on particle number concentration, size and mixing state. This work reports on two field campaigns at nearby urban and rural sites, 65 km apart, in southern Sweden during late summer 2018. BC was measured using a single particle soot photometer (SP2) and Aethalometers (AE33). Differences in BC concentrations between the sites are driven primarily by local traffic emissions. Equivalent and refractory BC mass concentrations at the urban site were on average a factor 2.2 and 2.5, with peaks during rush hour up to a factor ~4, higher than the rural background levels. The number fraction of particles containing a soot core was significantly higher in the city. BC particles at the urban site were on average smaller by mass and had less coating owing to fresh traffic emissions. The organic components of the fresh traffic plumes were similar in mass spectral signature to “hydrocarbon-like organic aerosol” (HOA), commonly associated with traffic. Despite the intense local traffic (~30 000 vehicles passing per day), PM₁, including organic aerosol, was dominated by aged continental air masses even at the curbside site. The fraction of thickly coated particles at the urban site was highly correlated with the mass concentrations of all measured chemical species of PM₁, consistent with aged, internally mixed aerosol. Trajectory analysis for the whole year showed that air masses arriving at the rural site from eastern Europe contained approximately double the amount of BC compared to air masses from western Europe. Furthermore, BC from the largest region emissions in the Malmö/Copenhagen urban area transported to the rural site is discernible above background levels only when precipitation events are excluded. We show that continental Europe, and not the Malmö/Copenhagen region is the major contributor to the background BC mass concentrations in southern Sweden.

1 Introduction

35 Virtually everywhere in the world, a fraction of the ambient aerosol consists of soot. Soot is formed by incomplete combustion
of carbonaceous fuels at hot air-starved conditions, and commonly contains both highly absorbing graphitic-like black carbon
(BC) and organic carbon. It has severe effects on climate and human health (e.g. Bond et al., 2013; IPCC, 2013; WHO, 2021;
WHO, 2013; IARC, 2014; IPCC, 2021). Therefore, ambient measurements of soot concentrations and properties are of great
40 importance to constrain and model its effects. Soot measurements are complicated by the fact that the ambient aerosol is a
dynamic mixture, and that soot from different sources may not possess the same properties in terms of chemical content (and
nanostructure (Vander Wal et al., 2010; Malmberg et al., 2019), light absorption (Sandradewi et al. 2008), size (Schwarz et
al., 2008), and toxicity (Hakkarainen et al., 2022).

A soot particle survive in the atmosphere approximately 5-9 days (Textor et al., 2006). During its lifetime several processes
affects its properties. Soot is formed in a chain of steps, starting with the inception of the first condensed-phase particles from
45 gas-phase hydrocarbon soot precursor species (Michelsen et al 2020). These then grow rapidly by coagulation and gas-to-
particle conversion and become more graphitic, also making them more light-absorbing. These near-spherical 10-30 nm
primary particles coagulate to form agglomerates that are emitted from the source to the surrounding air, if not removed by
oxidation already in the combustion process or by exhaust after-treatment (e.g. particle filters). The properties of the emitted
soot particles depend on the combustion conditions and fuel used. For example, it is well known that the wavelength
50 dependence of light absorption is stronger for biomass burning BC than for traffic emissions (Sandradewi et al., 2008).

Once in the atmosphere, the freshly formed soot particles evolve from their initial agglomerated state, each consisting of large
numbers of primary soot particles, into compacted soot cores with significant coatings of inorganic and organic material
(Corbin et al., 2023). During such atmospheric ageing, the soot particles will not only increase their size and effective density,
but also their ability to absorb light will typically increase (Zhang et al., 2018). Further, the particles change from being nearly
55 hydrophobic into hygroscopic particles that can act as CCN (cloud condensation nuclei; Swietlicki et al., 2008) which increases
the likelihood for wet removal which is the main deposition process for BC in the atmosphere (Textor et al., 2006). This
transformation depends on atmospheric conditions and constituents but can be on the order of hours under favourable
conditions (Eriksson et al., 2017).

Already in 2012, the International Agency for Research on Cancer, which is part of the World Health Organization (WHO),
classified diesel engine exhaust as carcinogenic to. WHO further concluded that there is sufficient evidence for an association
60 between short-term BC levels and all-cause and cardiovascular mortality, and cardiopulmonary hospital admissions (WHO,
2013). Drawing similar conclusions as the WHO, the US EPA (2019) summarizes the associations between several health
effects and BC concentrations in their impact assessment (USEPA, 2019). In its latest update of the Air Quality Guidelines,
WHO did not yet recommend air quality guidelines for black carbon (WHO, 2021). Instead, they made a statement of good
65 practice recommending systematic measurements of black and elemental carbon, and production of BC/EC emission
inventories, exposure assessments and source apportionment.

Formatted: English (United States)

Formatted: English (United States)

A myriad of BC measurement techniques exist, and the terminology is based on the measured property (Petzold et al., 2013). From light absorption techniques, the equivalent black carbon (eBC) mass concentration is obtained from the ratio of measured light absorption coefficients to the corresponding mass absorption cross-section (MAC). More recently, techniques based on laser-induced incandescence (LII) have been developed for measuring the refractory black carbon (rBC) mass concentration. For instance, the single particle soot photometer (SP2) (Schwarz et al., 2006; Stephens et al., 2003) deploys LII to measure the single particle rBC mass. In addition, the SP2 can be used to retrieve the rBC core size, mass-size distribution and provide an estimation of the coating thickness of rBC particles (Moteki & Kondo, 2007).

Although BC mass concentrations are routinely estimated from optical methods, BC number concentrations, size distributions and mixing state are rarely measured. In a previous study it was shown that global aerosol microphysics models underestimate the BC particle size, by a factor of ~2-3, while overestimating the number concentrations, by more than a factor of 3, compared to airborne measurements using ~~single particle laser induced incandescence~~ the SP2 (Reddington et al., 2013). BC from different sources have different size distributions (Schwarz et al., 2008; Laborde et al., 2013; Saarikoski et al., 2021), that affects both transport (lifetime) and light interactions (Hinds, 2012), as well as deposited dose (Alfoldy et al., 2009; Rissler et al., 2012). ~~Upon ageing in the atmosphere, fresh BC particles will obtain a coating layer consisting of condensed materials changing both their hygroscopicity (Swietlicki et al., 2008) and light absorption properties (e.g. Zhang et al., 2018). This transformation depends on atmospheric conditions and constituents but can be on the order of hours under favourable conditions (Eriksson et al., 2017).~~ Recent studies have pointed to the importance of BC mixing state, governed by emission source and atmospheric ageing, in understanding the light absorbing properties, and hence climatic impacts, of BC containing particles (e.g. Liu et al., 2017; Liu et al., 2019; Fierce et al., 2020; Yuan et al., 2020). These properties of BC, that are crucial to understand both health and climate impacts, are not measured by the BC measurement techniques commonly used by monitoring networks.

The aim of this study was to compare BC particle properties in nearby urban and rural settings, and to investigate the influence of urban emissions on the rural background air. ~~BC was measured using single particle laser induced incandescence and, utilizing both~~ filter-based absorption measurements and single particle LII. We compare BC mass and number concentrations, size distributions and mixing state. Furthermore, the relation to total particle number concentrations and chemically resolved PM₁ are assessed. Trajectory analysis was used to assess the influence of long- and short-range transport.

2 Methods

2.1 Measurement sites

Consecutive aerosol measurement campaigns were performed at existing field sites in a rural and an urban setting in southern Sweden, during the period late July to early October 2018. The weather did not change dramatically between the two campaigns, hence seasonality is not expected to play a major role for the results. Figure S1 shows that the BC concentrations at the rural site during July-October did not change drastically between the times of the two campaigns. The urban campaign

Formatted: English (United States)

(September-October) took place at a curbside measurement site near a busy road junction surrounded by 4-7 story buildings (about 30 000 vehicles per day) in Malmö, which has about 300 000 inhabitants, situated about 30 km east of Copenhagen. Air was sampled at 3 meters above the pavement using ¼" stainless steel tubing and a PM_{2.5} cyclone. Total inlet length was 4-5 meters and the flow was 7-8 liters per minute. The validity of the curbside site measurements as an indicator of the BC concentrations in the city was assessed by comparing simultaneously collected data at a rooftop urban background site (20 m a.g., PM_{2.5} inlet). The rural measurements (July-August) were performed at the Hyltemossa Research Station, about 65 km northeast of Malmö/Copenhagen. This station is part of both the ACTRIS (Aerosol, Clouds, and Trace gases European Research Infrastructure) and ICOS (Integrated Carbon Observation System) European environmental research infrastructure networks. The field site is surrounded by a managed spruce forest. Aerosols were sampled at a height of 30 meters above ground through a PM₁₀ inlet and 25 mm stainless steel tube, at a flow rate of 16.7 liters per minute. The single particle soot measurements at the rural site were conducted through a ¼" tube at 20 liters per minute (turbulent flow), due to simultaneous Eddy Covariance flux measurements (not described here).

2.2 Instrumentation and Analysis

2.2.1 Overview

Instrumentation to measure aerosols and gases was similar at both sites. Figure 1 shows during which time the urban and rural campaigns took place and which data was used. BC properties and concentrations was determined using a Single Particle Soot Photometer (SP2) that was moved between the sites and Aethalometers (AE33) deployed permanently. Aethalometer data for the whole year was used from the rural site to assess source regions. The particle size distribution was measured using Scanning (urban) and Differential (rural) Mobility Particle Sizers. During the latter part of the urban campaign, chemically resolved particle constituents were measured simultaneously at both sites with Soot Particle Aerosol Mass Spectrometers (SP-AMS). An Aerosol Particle Mass Analyzer (APM) was at a later stage deployed at the urban site together with a Differential Mobility Analyzer (DMA) to measure the size resolved effective density and particle mass.




Figure 1. Overview of the data and instruments used. Not shown in the figure are DMA-APM measurements of effective density during spring 2019.

2.2.2 SP2

125 The SP2 (Droplet Measurement Technologies) measures the refractory black carbon (rBC) mass of individual particles using
laser-induced incandescence (LII) (Stephens et al., 2003; Schwarz et al., 2006). In the SP2, particles are passed through a
continuous 1064 nm Nd:YAG laser beam. Light absorbing particles (such as soot) are heated by the laser to temperatures
where they will start to incandesce, if they are large enough. The incandescent light, measured by optical detectors, is linearly
proportional to the BC core mass and independent of particle coating and fractal structure (Schwarz et al., 2006; Slowik et al.,
130 2007). The approximate range of the SP2 is 0.5 – 200 fg per particle, where the lower limit is due to energy dissipation, and
the particles in upper limit may either saturate the detectors or not absorb enough energy to incandesce. The overall relative
uncertainty in mass concentration by the SP2 was estimated by Sharma et al. (2017) to be within 25-38 %. To limit the data
collected, not all particle peak signals were saved (10 and 1-5 % was saved in the rural and urban campaigns respectively).

The SP2 incandescence channels were calibrated using monodisperse Aquadag (provided by Droplet Measurement
135 Technologies) before, during and after the field campaigns. Particle sizes were selected using a DMA and mass was calculated
using the effective density recommended by Gysel et al. (2011). Since the calibration during the rural campaign did not yield
satisfactory data, the calibration during the urban campaign was used to analyze both datasets. Although this is not optimal,
the incandescence detector calibration is relatively stable in time. The response of the broadband incandescence detector from
typical ambient BC particle masses varied with less than $\pm 15\%$ for calibrations performed before and after the rural campaign.

140 The SP2 sensitivity to ambient BC has been shown to resemble that of fullerene soot particles (Baumgardner et al., 2012).
Following Laborde et al. (2012b), the Aquadag calibration factors were therefore translated into a fullerene soot equivalent
calibration using a scaling factor of 1.34 at 8.9 fg and an intercept of 0.

The calibrations and campaign data were processed using the PSI SP2 toolkit version 4.111. BC particles between mass
equivalent diameters of 64-580 nm (using a density of 1800 kg m⁻³) were included in the result. The lower size was selected
145 from looking at the sharp decrease in the number-size distribution when including very low peak heights. The mass and number
concentrations presented are not corrected for particles outside this range. Pileci et al. (2021) found that the rBC mass outside
the range of the SP2 is on the order of 20 % for several European background stations, depending on the size distribution. No
particle detection efficiency calibration was performed. Lognormal fits to averages of 2 and 24h were used to calculate
geometric mean diameters.

150 BC particle coatings can be estimated by the SP2 using a separate scattering detector. The delay-time method was used to
estimate BC particle coatings in a qualitative way (Moteki and Kondo, 2007; Subramanian et al., 2010). This method separates
BC particles as being either thinly or thickly coated depending on the delay-time between the maximum scattering and
incandescence peak heights. Coated BC particles generate two scattering peaks, one from the evaporating coating and one
from the BC core itself. The scattering signal of the coating will be detected before the incandescence signal since BC particles
155 take some time to heat up to the point of incandescence, and absorbed energy will dissipate through evaporation of the coating

material. For thickly coated particles, the scattering of the coating will exceed the scattering of the BC core, while for thinly coated particles the opposite is true. Moteki and Kondo (2007) and Laborde et al. (2012a) showed that a coating volume fraction of 70 % is needed for a particle to be classified as thickly coated by the SP2.

2.2.3 Aethalometer

160 Aerosol light absorption was measured at both sites using a seven wavelength Aethalometer (model AE33, Magee Scientific)
(Drinovec et al., 2015). Equivalent black carbon (eBC) mass concentrations were calculated using absorption at 880 nm and
the default mass absorption cross section (MAC) value of $7.77 \text{ m}^2 \text{ g}^{-1}$. The absorption Ångström exponent (AAE), a measure
of the spectral dependence of light absorption and commonly used in source apportionment of BC (Kirchstetter et al., 2004;
Sandradewi et al., 2008), was calculated between the wavelengths 370 - 950 nm. The relative uncertainty of the absorption
165 coefficient measured by the aethalometer increases with lower BC mass concentrations, and has been shown to converge
around 30 % for an older version of the instrument (Backman et al., 2017). The uncertainty of eBC also includes the uncertainty
in the MAC value which was not estimated in the present study.

2.2.4 SP-AMS

The chemical composition of refractory and non-refractory particulate mass was measured in real-time with two soot particle
170 aerosol mass spectrometers (SP-AMS, Aerodyne Research Inc.) (Onasch et al., 2012), at the urban and rural measurement site,
respectively. The measurements were performed in parallel, and there are 15 days of overlap (Sept. 25 to Oct. 10). The SP-
AMS detects particles within the vacuum aerodynamic diameter size range of about 70 nm to 1 μm . The aerosol sample flow
is focused through an aerodynamic lens. The (non-refractory) components of particles in the air beam are vaporized on a heated
tungsten plate (600 °C), subjected to 70 eV electron ionization, and the ions produced are detected and categorized (organic,
175 nitrate, sulfate, ammonium or chloride) by a high-resolution time-of-flight mass spectrometer. In the SP-AMS set-up a laser
is used to also vaporize the refractory BC (rBC). We operated both SP-AMSs in the “dual vaporizer” configuration as further
discussed in the SI. Calibrations were performed in the field using 300 nm mobility diameter particles from nebulizing
ammonium nitrate as described in Onasch et al (2012). The absolute concentration of species at the urban site are not shown
due to uncertainties in the calibration and dissimilarities when comparing to other instruments. For the rural data, PM_{10} derived
180 from the DMPS (using a density of 1.5 g cm^{-3}) and a Palas Fidas 200 at a nearby site agreed well with that from the SP-AMS.
All rBC data presented in this work are derived from the SP2 measurements which are more accurate. Data analysis was
performed with IGOR Pro 7 (Wavemetrics, USA), and the AMS analysis software package including SQUIRREL 1.61F, and
PIKA 1.21F.

For the urban dataset, the fresh traffic aerosol was isolated through selection of distinct traffic plumes, i.e. short (seconds to
185 minutes) increase in concentration. Increases in rBC number concentration from the SP2 and the concentration of the HOA
marker ion C_4H_9^+ from the AMS were used as indicators of fresh traffic plumes, and 90 such plumes were selected. For each
plume, the current urban background aerosol was defined by selecting two windows, one before and one after the plume, of

about 2 minutes, i.e. a total of 4 minutes- (see Figure S2). The typical plume duration was about 30-60 seconds. Organic aerosol (OA) mass spectra from the AMS were calculated for the plume and for the background respectively, and the background spectra were subtracted from the plume spectra to obtain a net contribution to OA mass concentrations. The m/z tracer method (Ng et al., 2011) was used to estimate the different OA components.

2.2.5 DMPS/SMPS

Particle size distributions were measured between electrical mobility diameters of 3.2 - 900 nm at the rural site and 11.5 – 604 nm at the urban site. At the rural site the DMPS consisted of two Hauke type DMAs and two condensation particle counters (CPC, TSI 3772 and TSI UCPC 3025) (Wiedensohler et al., 2012). At the urban site the SMPS consisted of a TSI DMA 3082 and a TSI CPC 3772. At both sites, the aerosol was dried before sizing using Permapure driers with low pressure sheath air. Size distributions at both sites were averaged over 1 hour.

2.2.6 DMA-APM

Size dependent particle mass and effective density were measured with a DMA-APM. Pre-selection of particle size was done with a DMA (TSI 3082), which selects particles based on electrical mobility diameter. The quasi-monodisperse aerosol was then led through the APM (model 3600, Kanomax), which measures the relationship between electrical mobility and particle mass. In the APM, the aerosol passes a rotating cylinder with an applied voltage, and the particle mass is determined based on the balance equation between the centrifugal force and electrostatic force (McMurry et al., 2002). The mass selected particles were counted after the DMA-APM with a CPC (TSI 3772). The aerosol flow was 1 liter per minute. The effective density is calculated based on the electrical mobility diameter and the particle mass from the DMA-APM. For spherical particles, this is the bulk density, but for non-spherical particles, the effective density depends on the particle shape factor. Six different particle mobility diameters were tested, 50, 75, 100, 150, 250, and 350 nm. The APM was operated in a constant RPM mode, where the angular rotation speed was constant and the APM voltage was increased step-wise during a time period of 15 minutes per scan. The rotational speed and voltage were selected so that the effective density was measured in approximately the range 0.1-3.5 g cm⁻³. The APM was run during five days in spring 2019, after the intensive campaigns to verify the bimodality, in terms of effective density, of the city aerosol.

2.2.7 Particle inlet losses

Particle losses in the inlets were assessed using the Particle loss calculator (von der Weiden et al., 2009). The losses in the longest tube sections of each inlet that was used are shown in Fig. S2S3. Losses in additional tubing and the actual inlets are expected to be similar (and small) in all cases. In the laminar flow inlets at both sites, PM₁ losses are negligible, while the turbulent flow inlet has losses of 20-30 % for spherical particles with a diameter between 100 – 1000 nm (density 1600 kg m⁻³). Based on these calculations and the measured rBC size distribution (discussed below), the SP2 mass and number

concentrations measured at the rural site have been adjusted to correct for 25 and 30 % losses respectively since adjusting by size is not possible because the size of rBC cores including coating was not measured.

220 2.2.8 Trajectory analysis

Wind direction and speed data from the rural site at Hyltemossa were downloaded from the ICOS Carbon portal (ICOS 2019), and analyzed using the Igor pro tool Zefir (Petit et al., 2017). To investigate how the air mass origin influence the BC concentrations at the rural site, we used the Hybrid Single Particle Lagrangian Integrated Trajectory Model (HYSPLIT) with meteorological data from the Global Data Assimilation System (GDAS) (Stein et al., 2015; Rolph et al., 2017). Seven days


225 HYSPLIT backward trajectories, arriving with one hour intervals at 100 m a.g.l. at Hyltemossa, were calculated for year 2018.


The air masses were then classified into four different categories depending on which geographical region the air masses spent most time over. For this classification we only considered the air mass origin during the last 48 hours upwind Hyltemossa and the cumulative time which the transported air parcels were below 1000 m a.g.l.. Figure S3S4 in the supplementary material illustrate how the air mass origin were divided into North Westerly (NW), South Westerly (SW), South Easterly (SE) and

230 North Easterly (NE). In addition we separated the air mass origin depending on if they passed over the Malmö/Copenhagen region less than 12 hours before the air masses arrived at Hyltemossa.

The long-distance transported BC source contributions during the rural and urban measurement campaigns were estimated using the Lagrangian 1D-column chemistry transport model ADCHEM (Roldin et al., 2011; Roldin et al., 2019). ADCHEM was setup and run forward in time along pre-computed 14-days backward HYSPLIT air mass trajectories arriving 100 m a.g.l.

235 at Hyltemossa, one new trajectory every hour. Source specific anthropogenic BC emissions along the trajectories were taken from the CAMS-GLOB-ANT v4.2 global emission inventory, which has a spatial resolution of $0.1^\circ \times 0.1^\circ$ (Granier et al., 2019). BC emissions from wildfires were estimated using the GFED4 emission inventory (Randerson et al., 2018).





240 Figure 2. Diurnal cycles of rBC mass concentrations ($\pm 1\sigma$) from the urban and rural sites as measured by the SP2. Values are averages of data collected during 34 (rural) and 41 (urban) days of sampling.

3 Results and Discussion

3.1 BC concentrations

245 An overview of the BC measurement results is shown in Table 1. As expected, the curbside site has higher concentrations than the rural site, with campaign average factors of 2.5 for rBC mass, 2.2 for eBC mass, and 3.2 for rBC number concentration. The differences in concentrations are due to local emissions, clearly shown in the diurnal pattern (Fig. 2), which follows traffic intensity. The comparison between the urban street-level and urban background eBC levels are very similar in time-series but with a lower daily maximum for the roof-top measurements (Fig. S4S5). This suggests that the curbside measurements are indicative for the city. The average eBC mass concentration at the roof-top site during the campaign was 15% lower than at the curbside site. The largest difference between the urban and rural sites occur during morning rush hour when rBC mass concentrations are up to a factor 3.7 higher in the city. The mass concentrations are still higher in the city during nighttime, before the first rush hour peaks, but only by a factor of 1.2-1.3. The weekends have lower rBC mass concentration (M_{rBC} of $0.11 \pm 0.06 \mu\text{g m}^{-3}$) compared to weekdays in the city, but is still a factor of 1.8 higher than the average for the rural background station. The AAE is slightly higher at the urban site, which could be due to the differences in BC size distribution and coating (Virkkula, 2021).

250
255 Table 1. Averages $\pm 1\sigma$ (original averaging time in parenthesis) for the urban and rural campaigns. rBC mass (M) and number (N) concentrations are from SP2 measurements and eBC mass concentrations are from absorption at 880 nm using a MAC of $7.77 \text{ m}^2 \text{ g}^{-1}$. Mass-based geometric mean diameter (GMD_M) and geometric standard deviation (GSD_M) are from the rBC mass equivalent size distribution. rBC concentrations are not corrected for particles outside the range of the instrument. Rural rBC concentrations are corrected for inlet losses. Absorption Ångström exponents (AAE) are calculated between absorption at 370 and 950 nm.

	Rural	Urban
M_{rBC} (1h), $\mu\text{g m}^{-3}$	0.06 ± 0.05	0.15 ± 0.11
M_{eBC} (1h), $\mu\text{g m}^{-3}$	0.22 ± 0.16	0.49 ± 0.45
N_{rBC} (1h), cm^{-3}	31 ± 21	100 ± 80
GMD_M (24h), nm	168 ± 12	141 ± 11
GSD_M (24h), nm	1.74 ± 0.08	1.77 ± 0.04
$AAE_{370-950 \text{ nm}}$ (1h)	1.13 ± 0.12	1.24 ± 0.26

3.2 Estimation of BC particles number fractions

The number fraction of particles containing a BC core was estimated by comparing the daily average of BC number concentration from the SP2 to the number concentration of particles larger than 64 nm as measured by the mobility particle sizers. At the rural site this fraction was 2.7 ± 0.6 (1σ) %, while for the urban site the fraction was 13.4 ± 5.5 (1σ) %. These estimates are biased low because the BC size distribution (as measured by a mobility particle sizer) is shifted to larger sizes if shape factors and coating is added. Increasing the cut off size for total particle number measured by the SMPS to $D_p > 130$ nm (assuming a coating thickness of similar magnitude as the rBC core mass equivalent diameter) the fractions are increased to 6 % in the rural campaign and 45 % in the urban campaign. In a study outside Paris (Laborde et al., 2013), the number of BC particles counted by the SP2 (with no correction for particles outside the measurement range) was 0-15 % of the total particle number concentration (above 20 nm) measured by an SMPS. In airborne measurements over Europe (using an SP2 and a passive cavity aerosol spectrometer probe), Reddington et al. (2013) found that the number fraction of approximately 14 % of particles with dry diameters above 260 nm contained a BC core. In an older study, Rose et al. (2006), using a volatility tandem-DMA set-up, measured summertime BC particle number fractions of 2-7 % at a rural site and 20-60 % in a street canyon, depending on the selected size (30-150 nm in mobility diameter).

3.3 rBC vs. eBC

The difference between rBC and eBC is significant at both sites (Fig. S5S6). For the full campaign averages, eBC is more than 3 times higher than rBC at both sites, with median ratios (eBC/rBC) of 3.04 and 3.65 the urban and rural sites respectively. Several studies have found similar discrepancies between rBC and eBC (Holder et al., 2014; Raatikainen et al., 2015; Sharma et al., 2017; Tasoglou et al., 2018; Li et al., 2019). The rBC mass in this study is biased low because of the limited range of the SP2, but given the size distributions measured (example shown in Fig. S6S7), this mass bias should be on the order of 10-20 %, which still only would lower the ratio between eBC and rBC to 2.4-2.7. One reason for biased high eBC values is that the measured absorption can be amplified by non-absorbing coatings of BC cores (Kalbermatter et al., 2022). It is well known that site specific MAC can differ from the default values from the manufacturer of the AE33 (e.g. Cui et al., 2016). At a site nearby the rural site of this study, a MAC of $12.64 \text{ m}^2 \text{ g}^{-1}$ at 880 nm has previously been derived using aethalometer (AE33) absorption coefficients and elemental carbon (EC) measurements (Martinsson et al., 2017). This value is 1.6 times higher than the default and if used, the ratio between eBC and rBC of the rural campaign would be ~2.





Figure 3. Average mass-size distribution of BC cores for the urban (black) and rural (orange) campaigns. Filled areas show $\pm 1\sigma$.
 290 Insert shows the normalized distribution for comparison of the modes. Dip at 270 nm is due to erroneous stitching of high and low gain detector channels.

3.4 BC core size and coating

The average geometric mean diameter (GMD) of the rBC mass size distribution, from lognormal fits of 24 h data, was highest in the rural setting (Table 1). The average mass size distributions for the full campaigns are shown in Fig. 3. Mass equivalent diameters of 168 and 148 nm for the rural and urban sites respectively, correspond to BC particle masses of 4.5 and 3.1 fg. The number size distribution generally peaked close to the SP2 detection limit (64 nm, corresponding to 0.25 fg, for both campaigns), and the GMD of this distribution is therefore more uncertain, and not presented here. An example of 24 h number and mass size distributions are shown in Fig. S6S7. The difference in mass GMD is statistically significant ($P < 0.01$) and can be explained by different BC particle sources, air masses and coagulation. Figure S1 shows that while BC concentrations at the rural site were similar during the two campaigns, the source sectors changed slightly. Notably, more industrial emissions and less shipping emissions affected the concentrations during the urban campaign. Figure 4 shows the diurnal variation in GMD from mass size distributions averaged over 2 h. The urban site has a higher fraction of particles coming from local traffic, which have been shown to be smaller than particles from e.g. biomass burning (Schwarz et al., 2008; Laborde et al., 2013; Saarikoski et al., 2021). This shifts the size distribution during daytime to slightly lower sizes, although the measurement uncertainties, including calibrations, of the SP2 are close to the difference in average GMD between the sites.
 305



The BC particles at the rural site were characterized by a thicker coating than at the urban location, as measured by the SP2 delay time method (a typical example is shown in Fig. S7S8). The fraction of “thickly coated” BC particles with an incandescence peak height corresponding to a BC core diameter of mass 2.4 fg (~136 nm mass equivalent diameter) was on average $71 \pm 8\%$ (1σ) at the rural site and $38 \pm 7\%$ (1σ) at the urban site, with no big difference during weekends. The short-

310 term APM measurements verified the bimodal structure in effective density of the urban aerosol. The mass distribution of 150 nm particles is shown in Fig. 5. The two modes suggest that the aerosol is externally mixed, with more and less dense particles present. The two lognormal curve fittings have a mean mass of 0.86 ± 0.02 fg, and 2.49 ± 0.02 fg, respectively. These masses correspond to effective densities of 0.49 ± 0.01 g cm⁻³, and 1.41 ± 0.01 g cm⁻³, respectively, which is very similar to the effective densities presented by Rissler et al. (2014), who reported two effective densities, for 150 nm particles sampled in
315 downtown Copenhagen, of 0.53 ± 0.05 g cm⁻³, and 1.36 ± 0.15 g cm⁻³, respectively. For 50 and 75 nm particles, the average mass distribution was unimodal with a mean mass of 0.062 ± 0.001 fg and 0.26 ± 0.01 fg, respectively, corresponding to effective densities of 0.95 ± 0.01 g cm⁻³ and 1.19 ± 0.02 g cm⁻³. The average mass distribution of 100 nm particles was bimodal with mean masses of 0.33 ± 0.01 fg and 0.68 ± 0.01 fg, corresponding to effective densities of 0.63 ± 0.02 g cm⁻³ and 1.29 ± 0.01 g cm⁻³. These values are also similar to the densities in Rissler et al. (2014), but we did not observe an additional
320 mode at higher masses for 50 and 75 nm particles, but only the less dense, fresh soot mode. However, in single APM spectra, the mean mass mode was sometimes different from the average, and on a couple of occasions, a bimodal distribution was observed also for the smaller particles. No mass mode could be extracted for APM measurements of larger particle sizes (250 and 350 nm) due to low particle concentrations and consequently poor counting statistics.



325

Figure 4. Box-plot showing the daily pattern of rBC GMD (2 h averages) as measured by the SP2, at the urban and rural sites. Boxes show the median, 25th and 75th percentiles, with values more than 1.5 below or above those considered outliers (+).



330 Figure 5. Measured DMA-APM spectrum for urban particles with a diameter of 150 nm (black markers). The spectrum is ~~an~~the average number concentration (arbitrary units) of eleven samples from all sampling days. Two lognormal fits are shown in black.





Figure 6. Wind rose showing the frequency of different wind directions (30° resolution) during the rural campaign and corresponding concentrations of eBC.

335 3.6 Origins of BC in the rural background air

Figure 6 shows the wind direction probability together with eBC from the five weeks rural campaign. The Malmö/Copenhagen area is in the direction 200 – 230 degrees as seen from the rural site. Based on the HYSPLIT trajectory analysis (Sect. 2.2.8), some peaks in eBC, non-refractory PM1, and total particle number concentrations at the rural site coincide with air masses originating over Malmö/Copenhagen. These urban plumes were discernible when the air mass was relatively clean (typically of western or north-western origin). However, no conclusive results on the influence of the urban site on rural levels could be derived from analysis of the limited field campaign data. Also when analyzing the complete eBC dataset from 2018 there is no significant difference in the median eBC concentration for air masses with and without influence from Copenhagen and/or Malmö, according to the trajectory analysis. However, if we only consider air masses with insignificant precipitation within 48 hours upwind the rural site (i.e. less than <1 mm precipitation according the HYSPLIT model), the measured eBC median concentration is significantly higher in the air masses that have moved over Copenhagen/Malmö, 354.7 ng m⁻³ compared to 226.9 ng m⁻³ without influence from Copenhagen/Malmö. Similar results are also found when only the air masses originating from SW is considered (see Fig. 7 and Tables S2-S4). Most likely the contribution from the BC emissions from Copenhagen/Malmö is not apparent when analyzing the whole eBC dataset from 2018 because the SW air masses are generally influenced by more precipitation (more BC wet scavenging) compared to other air masses. The HYSPLIT trajectory simulations for the complete year show that 66 % of all air masses that passed over Malmö/Copenhagen were influenced by >1 mm precipitation within 48 hours upwind the rural site, compared to 39 % for all other air masses. The lowest median eBC concentration is found in air masses from NW (101 ng m⁻³) followed by NE air masses (149.9 ng m⁻³) and SW air masses (192.0 ng m⁻³). SE air masses clearly stand out from any other air masses with a median eBC concentration of 563 ng m⁻³. This can partly be explained by the low probability of precipitation in SE air masses. Only 23 % of these air masses are influenced by >1 mm precipitation within 48 hours upwind the rural site. However, also when considering the effect of precipitation upwind Hyltemossa the SE air masses have a median eBC concentration that is ~2 times larger than the SW air masses.



360 **Figure 7. Trajectory analysis of different air mass origins, with corresponding median eBC mass concentrations in the legend. Panel A shows histograms with the observed eBC from the Aethalometer at Hyltemossa from year 2018 for air masses originating from SE, NE, NW and SW as defined in Fig. S3S4. Panel B shows the eBC histogram from all SW air masses with or without influence from the Copenhagen (CPH) and Malmö region.**

365 Comparison of the 15 days of simultaneous urban and rural measurements are shown in Fig. 8. It is clear that non refractory-PM₁ concentration and composition are driven by long-range pollution at both sites, while black carbon is more influenced by local emissions. The organic aerosol at the urban site was clearly dominated by oxygenated organic aerosol (OOA, 78%) with only a minor contribution of HOA (9%), suggesting that aged long-range sources dominate even the organic aerosol, for which one could expect a larger difference between the sites owing to the high urban traffic density (Glasius et al., 2011). However, in the original time resolution of 20 seconds (not shown) the urban AMS data features transient plumes of traffic exhaust. Yet, as shown in Fig. 8, these plumes have little impact on average concentration. The OA mass spectrum from traffic plumes at the curbside site is shown in Fig. S8S9. The plume spectra are similar in mass spectral profiles to “hydrocarbon like organic aerosol” HOA, commonly associated with traffic emissions. This OA is likely present in thin coatings on the freshly emitted traffic BC, and possibly also externally mixed with the BC, and hence not detected by the SP2 time delay method.

370
375 With regards to eBC, Fig. 8 shows that approximately half of the mass at the urban site is due to long range transport, despite the high traffic intensity (about 30 000 vehicles per day). This can be deduced from the temporal variability and absolute concentrations at both sites. The result is corroborated by the rBC results shown in Fig. 2. Firstly, rBC concentrations are roughly doubled at the urban site compared to the rural site. Secondly, the diurnal pattern observed at the urban site (see Figure 2) shows that levels between 01.00 and 05.00 local time, during which local traffic density is low, are about half of the average concentration.




Figure 8. Concentrations of eBC and PM₁ chemical species at both sites during overlapping urban and rural measurement period. 1 hour averages for eBC, 20 minutes for PM₁ chemical species. Absolute concentrations of PM₁ at the urban site was not possible to extract due to uncertainties in the calibration.

Consistent with the high abundance of BC from long range transport at the urban site, on average 40 % of urban BC particles with a diameter close to the mass GMD were found to be thickly coated based on the SP2 data. We did not directly measure the chemical composition of the thick coatings. However, as shown in Fig. 9, non-refractory PM₁ concentration is highly correlated with thickly coated BC fraction ($R^2 = 0.87$). Indeed, all major chemical species in non refractory-PM₁ correlate well with the thickly coated BC fraction ($R^2 = 0.82 - 0.88$). This suggests that the BC coatings are similar in composition to non refractory-PM₁ (dominated by secondary material), despite the fact that only a minor fraction of the particles in PM₁ contain rBC as discussed above. As non refractory-PM₁ is more widely and accurately measured than BC coating composition such homogeneity simplifies assessment of BC properties and impacts, for example concerning cloud formation and lung deposition. However, direct measurements of BC coating composition, ideally on a single particle basis, are needed to support this conjecture.





Figure 9. Comparison of chemically resolved non-refractory PM_{10} concentration and the fraction of thickly coated BC cores, with a mass equivalent diameter corresponding to ~ 136 nm, in the urban setting. Left: time series. Right: scatterplots of thickly coated BC particle fraction vs non refractory- PM_{10} and subcategories thereof. 24 hour averages.

395 4 Summary and Conclusions

During 11 weeks in 2018, BC particles were sampled at two nearby sites in southern Sweden. Campaign average mass concentrations of BC, measured with two different methods, was a factor of 2.2-2.5 higher at the urban site, compared to the

rural background site. Hourly averages during rush hour peaks were up to a factor 4 higher in the city. Despite good correlation between rBC and eBC, a factor of 3 higher concentrations was consistently measured with the optical method. The number fraction of particles containing BC, compared to total particle concentration above 64 nm in mobility diameter, was 2.4 and 13.4 % at the rural and urban site respectively. However, these numbers depend strongly on the sizes that are integrated and would ideally be measured and reported in a size-segregated manner using pre-selection of particles. The BC particle size distributions were similar, with peak mode diameters slightly smaller in the city, which can be expected since a larger fraction of the particles are from fresh traffic emissions. This was also seen in the BC particle coating, which was assessed in a qualitative manner. The rural site had approximately double the amount of thickly coated particles compared to the street measurements.

Simultaneous measurements using SP-AMS at both sites showed similarities in both time-series and mass spectra of the organic aerosol, verifying the impact of long range aerosol and secondary aerosol production. Primary organic aerosol mass spectra were investigated based on transient plumes from local traffic at the urban site, but composed a small fraction of the total organic aerosol. The fraction of thickly coated particles at the urban site was highly correlated to all measured species of the AMS, again showing the importance of long range transport.

Plumes from the nearby urban site to the rural site were not clearly distinguishable during the field campaigns. Trajectory analysis of the full year of 2018 show that significant increases in eBC concentrations at the rural site for air masses passing over the Malmö/Copenhagen area are only clearly seen during days with low precipitation. This increase, however, is small in comparison with the influence of long-range transported BC. Transport of BC from continental, and especially eastern, Europe is what governs the BC concentrations in southern Sweden background air, when looking at eBC from the full year of 2018.

Data availability

Processed data for main results are available at <https://doi.org/10.5281/zenodo.6559236>. EBC and DMPS data from the rural site are available at <http://ebas.nilu.no>. Raw or specific data sets are available from the authors upon request.

Author contribution

EA: Conceptualization, Methodology, Validation, Formal analysis, Investigation, Data curation, Writing- original draft, Visualization, Project administration. SA: Validation, Formal analysis, Investigation, Data curation, Writing- original draft, Writing- Review & editing. LN: Formal analysis, Writing- review & editing, Visualization. MS: Investigation, Resources, Data curation, Writing- review & editing. JP: Formal analysis, Writing- review & editing, Visualization. JKN: Resources, Writing- review & editing. MB: Writing- review & editing. HS: Writing- review & editing. PR: Methodology, Software, Formal analysis, Resources, Data curation, Writing- Original draft, Writing- review & editing, Visualization, Funding acquisition. AK: Formal analysis, Data curation, Writing- review & editing. ES: Conceptualization, Methodology, Resources,

Writing- review & editing, Supervision, Project administration, Funding acquisition. AE: Conceptualization, Methodology, Validation, Formal analysis, Investigation, Data curation, Writing- Original draft, Writing- review & editing, Visualization.

435 **Competing interests**

The authors declare that they have no conflict of interest.

Acknowledgements

The authors would like to thank ICOS Sweden, especially Tobias Biermann, Michal Heliasz, Jutta Holst, and Thomas Holst, for meteorological data and Hyltemossa station management, Paul Hansson at Malmö Environment Department for help with
440 setting up the urban station measurements and Dalaplan station technical assistance. We also wish to acknowledge the support from the Swedish Research Council (Vetenskapsrådet) for ACTRIS Sweden under contract 2021-00177.

Financial Support

This work was supported by the Swedish research councils Formas (projects 2015-00994 and 2018-01745) and VR (projects 2019-05062 and 2019- 05006).

445 **References**

- Alfoldy, B., B. Giechaskiel, W. Hofmann and Y. Drossinos: Size-distribution dependent lung deposition of diesel exhaust particles, *Journal of Aerosol Science*, 40, 652-663, 2009.
- Backman, J., L. Schmeisser, A. Virkkula, J. A. Ogren, E. Asmi, S. Starkweather, S. Sharma, K. Eleftheriadis, T. Uttal and A. J. A. M. T. Jefferson: On Aethalometer measurement uncertainties and an instrument correction factor for the Arctic, 10, 5039-
450 5062, 2017.
- Baumgardner, D., O. Popovicheva, J. Allan, V. Bernardoni, J. Cao, F. Cavalli, J. Cozic, E. Diapouli, K. Eleftheriadis, P. J. Genberg, C. Gonzalez, M. Gysel, A. John, T. W. Kirchstetter, T. A. J. Kuhlbusch, M. Laborde, D. Lack, T. Muller, R. Niessner, A. Petzold, A. Piazzalunga, J. P. Putaud, J. Schwarz, P. Sheridan, R. Subramanian, E. Swietlicki, G. Valli, R. Vecchi and M. Viana: Soot reference materials for instrument calibration and intercomparisons: a workshop summary with recommendations, *Atmospheric Measurement Techniques*, 5, 1869-1887, 2012.
- 455 Bond, T. C., S. J. Doherty, D. W. Fahey, P. M. Forster, T. Berntsen, B. J. DeAngelo, M. G. Flanner, S. Ghan, B. Karcher, D. Koch, S. Kinne, Y. Kondo, P. K. Quinn, M. C. Sarofim, M. G. Schultz, M. Schulz, C. Venkataraman, H. Zhang, S. Zhang, N. Bellouin, S. K. Guttikunda, P. K. Hopke, M. Z. Jacobson, J. W. Kaiser, Z. Klimont, U. Lohmann, J. P. Schwarz, D. Shindell,

- T. Storelvmo, S. G. Warren and C. S. Zender: Bounding the role of black carbon in the climate system: A scientific assessment, 460 *Journal of Geophysical Research-Atmospheres*, 118, 5380-5552, 2013.
- [Corbin, J. C., R. L. Modini and M. Gysel-Beer: Mechanisms of soot-aggregate restructuring and compaction, *Aerosol Science and Technology*, 57, 89-111, 2023.](#)
- Cui, X. J., X. F. Wang, L. X. Yang, B. Chen, J. M. Chen, A. Andersson and O. Gustafsson: Radiative absorption enhancement from coatings on black carbon aerosols, *Science of the Total Environment*, 551, 51-56, 2016.
- 465 Drinovec, L., G. Mocnik, P. Zotter, A. S. H. Prevot, C. Ruckstuhl, E. Coz, M. Rupakheti, J. Sciare, T. Muller, A. Wiedensohler and A. D. A. Hansen: The "dual-spot" Aethalometer: an improved measurement of aerosol black carbon with real-time loading compensation, *Atmospheric Measurement Techniques*, 8, 1965-1979, 2015.
- Eriksson, A., C. Wittbom, P. Roldin, M. Sporre, E. Öström, P. Nilsson, J. Martinsson, J. Rissler, E. Nordin and B. J. S. r. Svenningsson: Diesel soot aging in urban plumes within hours under cold dark and humid conditions, 7, 1-10, 2017.
- 470 Fierce, L., T. B. Onasch, C. D. Cappa, C. Mazzoleni, S. China, J. Bhandari, P. Davidovits, D. A. Fischer, T. Helgestad, A. T. Lambe, A. J. Sedlacek, G. D. Smith and L. Wolff: Radiative absorption enhancements by black carbon controlled by particle-to-particle heterogeneity in composition, 201919723, 2020.
- Glasius, M., A. la Cour and C. Lohse: Fossil and nonfossil carbon in fine particulate matter: A study of five European cities, *Journal of geophysical research*, 116, D11302, 2011.
- 475 Granier, C., Darras, S., Denier van der Gon, H., Doubalova, J., Elguindi, N., Galle, B., Gauss, M., Guevara, M., Jalkanen, J.-P., Kuenen, J., Liousse, C., Quack, B., Simpson, D., Sindelarova, K., The Copernicus Atmosphere Monitoring Service global and regional emissions (April 2019 version), Copernicus Atmosphere Monitoring Service (CAMS) report, doi:10.24380/d0bn-kx16, 2019.
- Gysel, M., M. Laborde, J. S. Olfert, R. Subramanian and A. J. Grohn: Effective density of Aquadag and fullerene soot black carbon reference materials used for SP2 calibration, *Atmospheric Measurement Techniques*, 4, 2851-2858, 2011.
- 480 Hakkarainen, H., L. Salo, S. Mikkonen, S. Saarikoski, M. Aurela, K. Teinilä, M. Ihalainen, S. Martikainen, P. Marjanen and T. J. S. o. T. T. E. Lepistö: Black carbon toxicity dependence on particle coating: Measurements with a novel cell exposure method, 156543, 2022.
- Hinds, W. C. (2012). [Aerosol technology: properties, behavior, and measurement of airborne particles](#), John Wiley & Sons.
- 485 Holder, A. L., G. S. W. Hagler, T. L. B. Yelverton and M. D. Hays: On-road black carbon instrument intercomparison and aerosol characteristics by driving environment, *Atmospheric Environment*, 88, 183-191, 2014.
- IARC (2014). Diesel and gasoline engine exhausts and some nitroarenes. . [IARC monographs on the evaluation of carcinogenic risks to humans](#). **105**: 9.
- ICOS , R. I. (2019). ICOS ATC MTO Release, Hyltemossa (150.0 m), 2017-09-26–2019-04-30
- 490 https://hdl.handle.net/11676/DbbmB-ppi1ZsmQzHfuQ9y_oY.

IPCC (2013). Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge, United Kingdom and New York, NY, USA, Cambridge University Press.

IPCC (2021). Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change; Chapter Three: Human influence on the climate system.

- 495 Kalbermatter, D. M., G. Močnik, L. Drinovec, B. Visser, J. Röhrbein, M. Oscity, E. Weingartner, A. P. Hyvärinen and K. Vasilatou: Comparing black-carbon- and aerosol-absorption-measuring instruments – a new system using lab-generated soot coated with controlled amounts of secondary organic matter, *Atmos. Meas. Tech.*, 15, 561-572, 2022.
- Kirchstetter, T. W., T. Novakov and P. V. Hobbs: Evidence that the spectral dependence of light absorption by aerosols is affected by organic carbon, *Journal of Geophysical Research-Atmospheres*, 109, 2004.
- 500 Laborde, M., M. Crippa, T. Tritscher, Z. Juranyi, P. F. Decarlo, B. Temime-Roussel, N. Marchand, S. Eckhardt, A. Stohl, U. Baltensperger, A. S. H. Prevot, E. Weingartner and M. Gysel: Black carbon physical properties and mixing state in the European megacity Paris, *Atmospheric Chemistry and Physics*, 13, 5831-5856, 2013.
- Laborde, M., P. Mertes, P. Zieger, J. Dommen, U. Baltensperger and M. Gysel: Sensitivity of the Single Particle Soot Photometer to different black carbon types, *Atmospheric Measurement Techniques*, 5, 1031-1043, 2012a.
- 505 Laborde, M., M. Schnaiter, C. Linke, H. Saathoff, K. H. Naumann, O. Möhler, S. Berlenz, U. Wagner, J. W. Taylor, D. Liu, M. Flynn, J. D. Allan, H. Coe, K. Heimerl, F. Dahlkotter, B. Weinzierl, A. G. Wollny, M. Zanatta, J. Cozic, P. Laj, R. Hitzenberger, J. P. Schwarz and M. Gysel: Single Particle Soot Photometer intercomparison at the AIDA chamber, *Atmospheric Measurement Techniques*, 5, 3077-3097, 2012b.
- 510 Li, H. Y., K. D. Lamb, J. P. Schwarz, V. Selimovic, R. J. Yokelson, G. R. McMeeking and A. A. May: Inter-comparison of black carbon measurement methods for simulated open biomass burning emissions, *Atmospheric Environment*, 206, 156-169, 2019.
- Liu, D. T., J. Whitehead, M. R. Alfarra, E. Reyes-Villegas, D. V. Spracklen, C. L. Reddington, S. F. Kong, P. I. Williams, Y. C. Ting, S. Haslett, J. W. Taylor, M. J. Flynn, W. T. Morgan, G. McFiggans, H. Coe and J. D. Allan: Black-carbon absorption enhancement in the atmosphere determined by particle mixing state, *Nature Geoscience*, 10, 184-U132, 2017.
- 515 Liu, H., X. Pan, D. Liu, X. Liu, X. Chen, Y. Tian, Y. Sun, P. Fu and Z. Wang: Mixing characteristics of refractory black carbon aerosols determined by a tandem CPMA-SP2 system at an urban site in Beijing, *Atmos. Chem. Phys. Discuss.*, 2019, 1-25, 2019.
- Malmborg, V. B., A. C. Eriksson, S. Török, Y. Zhang, K. Kling, J. Martinsson, E. C. Fortner, L. Gren, S. Kook and T. B. J. C. Onasch: Relating aerosol mass spectra to composition and nanostructure of soot particles, 142, 535-546, 2019.
- 520 Martinsson, J., H. A. Azeem, M. K. Sporre, R. Bergstrom, E. Ahlberg, E. Ostrom, A. Kristensson, E. Swietlicki and K. E. Stenstrom: Carbonaceous aerosol source apportionment using the Aethalometer model - evaluation by radiocarbon and levoglucosan analysis at a rural background site in southern Sweden, *Atmospheric Chemistry and Physics*, 17, 4265-4281, 2017.

- 525 McMurry, P. H., X. Wang, K. Park and K. Ehara: The relationship between mass and mobility for atmospheric particles: A new technique for measuring particle density, *Aerosol Science and Technology*, 36, 227-238, 2002.
- [Michelsen, H. A., M. B. Colket, P. E. Bengtsson, A. D'Anna, P. Desgroux, B. S. Haynes, J. H. Miller, G. J. Nathan, H. Pitsch and H. Wang: A Review of Terminology Used to Describe Soot Formation and Evolution under Combustion and Pyrolytic Conditions, *Acs Nano*, 14, 12470-12490, 2020.](#)
- 530 Moteki, N. and Y. Kondo: Effects of mixing state on black carbon measurements by laser-induced incandescence, *Aerosol Science and Technology*, 41, 398-417, 2007.
- Ng, N.L., Canagaratna, M.R., Jimenez, J.L., Zhang, Q., Ulbrich, I.M. and Worsnop, D.R., 2011. Real-time methods for estimating organic component mass concentrations from aerosol mass spectrometer data. *Environmental science & technology*, 45(3), pp.910-916.
- 535 Onasch, T. B., A. Trimborn, E. C. Fortner, J. T. Jayne, G. L. Kok, L. R. Williams, P. Davidovits and D. R. Worsnop: Soot Particle Aerosol Mass Spectrometer: Development, Validation, and Initial Application, *Aerosol Science and Technology*, 46, 804-817, 2012.
- Petit, J.-E., O. Favez, A. Albinet and F. Canonaco: A user-friendly tool for comprehensive evaluation of the geographical origins of atmospheric pollution: Wind and trajectory analyses, *Environmental Modelling & Software*, 88, 183-187, 2017.
- 540 [Petzold, A., J. A. Ogren, M. Fiebig, P. Laj, S. M. Li, U. Baltensperger, T. Holzer-Popp, S. Kinne, G. Pappalardo, N. Sugimoto, C. Wehrli, A. Wiedensohler and X. Y. Zhang: Recommendations for reporting "black carbon" measurements, *Atmospheric Chemistry and Physics*, 13, 8365-8379, 2013.](#)
- Pileci, R. E., R. L. Modini, M. Berto, J. F. Yuan, J. C. Corbin, A. Marinoni, B. Henzing, M. M. Moerman, J. P. Putaud, G. Spindler, B. Wehner, T. Muller, T. Tuch, A. Trentini, M. Zanatta, U. Baltensperger and M. Gysel-Beer: Comparison of co-located refractory black carbon (rBC) and elemental carbon (EC) mass concentration measurements during field campaigns at several European sites, *Atmospheric Measurement Techniques*, 14, 1379-1403, 2021.
- 545 Raatikainen, T., D. Brus, A. P. Hyvarinen, J. Svensson, E. Asmi and H. Lihavainen: Black carbon concentrations and mixing state in the Finnish Arctic, *Atmospheric Chemistry and Physics*, 15, 10057-10070, 2015.
- Randerson, J.T., van der Werf, G.R., Giglio, L., Collatz, G.J., and Kasibhatl, P.S., Global Fire Emissions Database, Version 4.1 (GFEDv4). ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAC/1293>, 2018.
- 550 Reddington, C. L., G. McMeeking, G. W. Mann, H. Coe, M. G. Frontoso, D. Liu, M. Flynn, D. V. Spracklen and K. S. Carslaw: The mass and number size distributions of black carbon aerosol over Europe, *Atmospheric Chemistry and Physics*, 13, 4917-4939, 2013.
- Rissler, J., E. Swietlicki, A. Bengtsson, C. Boman, J. Pagels, T. Sandström, A. Blomberg and J. Löndahl: Experimental determination of deposition of diesel exhaust particles in the human respiratory tract, *Journal of Aerosol Science*, 48, 18-33, 2012.

- Rissler, J., E. Z. Nordin, A. C. Eriksson, P. T. Nilsson, M. Frosch, M. K. Sporre, A. Wierzbicka, B. Svenningsson, J. Londahl, M. E. Messing, S. Sjogren, J. G. Hemmingsen, S. Loft, J. H. Pagels and E. Swietlicki: Effective Density and Mixing State of Aerosol Particles in a Near-Traffic Urban Environment, *Environmental Science & Technology*, 48, 6300-6308, 2014.
- 560 Rolph, G., A. Stein and B. Stunder: Real-time environmental applications and display system: READY, *Environmental Modelling & Software*, 95, 210-228, 2017.
- Roldin, P., Swietlicki, E., Schurgers, G., Arneht, A., Lehtinen, K. E. J., Boy, M., and Kulmala, M.: Development and evaluation of the aerosol dynamics and gas phase chemistry model ADCHEM, *Atmos. Chem. Phys.*, 11, 5867-5896, <https://doi.org/10.5194/acp-11-5867-2011>, 2011.
- 565 Roldin, P., Ehn, M., Kurtén, T., Olenius, T., Rissanen, M. P., Sarnela, N., Elm, J., Rantala, P., Hao, L., Hyttinen, N., Heikkinen, L., Worsnop, D. R., Pichelstorfer, L., Xavier, C., Clusius, P., Öström, E., Petäjä, T., Kulmala, M., Vehkamäki, H., Virtanen, A., Riipinen, I., and Boy, M.: The role of highly oxygenated organic molecules in the Boreal aerosol-cloud-climate system, *Nat. Commun.*, 10, 4370, <https://doi.org/10.1038/s41467-019-12338-8>, 2019.
- Rose, D., B. Wehner, M. Ketzel, C. Engler, J. Voigtlander, T. Tuch and A. Wiedensohler: Atmospheric number size distributions of soot particles and estimation of emission factors, *Atmospheric Chemistry and Physics*, 6, 1021-1031, 2006.
- 570 Saarikoski, S., J. V. Niemi, M. Aurela, L. Pirjola, A. Kousa, T. Rönkkö and H. Timonen: Sources of black carbon at residential and traffic environments obtained by two source apportionment methods, *Atmos. Chem. Phys.*, 21, 14851-14869, 2021.
- Sandradewi, J., A. S. H. Prevot, S. Szidat, N. Perron, M. R. Alfarra, V. A. Lanz, E. Weingartner and U. Baltensperger: Using aerosol light absorption measurements for the quantitative determination of wood burning and traffic emission contributions to particulate matter, *Environmental Science & Technology*, 42, 3316-3323, 2008.
- 575 Schwarz, J. P., R. S. Gao, D. W. Fahey, D. S. Thomson, L. A. Watts, J. C. Wilson, J. M. Reeves, M. Darbeheshti, D. G. Baumgardner, G. L. Kok, S. H. Chung, M. Schulz, J. Hendricks, A. Lauer, B. Karcher, J. G. Slowik, K. H. Rosenlof, T. L. Thompson, A. O. Langford, M. Loewenstein and K. C. Aikin: Single-particle measurements of midlatitude black carbon and light-scattering aerosols from the boundary layer to the lower stratosphere, *Journal of Geophysical Research-Atmospheres*, 111, 2006.
- 580 Schwarz, J. P., R. S. Gao, J. R. Spackman, L. A. Watts, D. S. Thomson, D. W. Fahey, T. B. Ryerson, J. Peischl, J. S. Holloway, M. Trainer, G. J. Frost, T. Baynard, D. A. Lack, J. A. de Gouw, C. Warneke and L. A. Del Negro: Measurement of the mixing state, mass, and optical size of individual black carbon particles in urban and biomass burning emissions, *Geophysical Research Letters*, 35, 2008.
- 585 Sharma, S., W. R. Leitch, L. Huang, D. Veber, F. Kolonjari, W. Zhang, S. J. Hanna, A. K. Bertram and J. A. Ogren: An evaluation of three methods for measuring black carbon in Alert, Canada, *Atmospheric Chemistry and Physics*, 17, 15225-15243, 2017.
- Slowik, J. G., E. S. Cross, J. H. Han, P. Davidovits, T. B. Onasch, J. T. Jayne, L. R. Williams, M. R. Canagaratna, D. R. Worsnop, R. K. Chakrabarty, H. Moosmuller, W. P. Arnott, J. P. Schwarz, R. S. Gao, D. W. Fahey, G. L. Kok and A. Petzold:

- 590 An inter-comparison of instruments measuring black carbon content of soot particles, *Aerosol Science and Technology*, 41, 295-314, 2007.
- Stein, A., R. R. Draxler, G. D. Rolph, B. J. Stunder, M. Cohen and F. Ngan: NOAA's HYSPLIT atmospheric transport and dispersion modeling system, *Bulletin of the American Meteorological Society*, 96, 2059-2077, 2015.
- Stephens, M., N. Turner and J. Sandberg: Particle identification by laser-induced incandescence in a solid-state laser cavity, *Applied Optics*, 42, 3726-3736, 2003.
- 595 Subramanian, R., G. L. Kok, D. Baumgardner, A. Clarke, Y. Shinozuka, T. L. Campos, C. G. Heizer, B. B. Stephens, B. de Foy, P. B. Voss and R. A. Zaveri: Black carbon over Mexico: the effect of atmospheric transport on mixing state, mass absorption cross-section, and BC/CO ratios, *Atmospheric Chemistry and Physics*, 10, 219-237, 2010.
- Swietlicki, E., H.-C. Hansson, K. Hämeri, B. Svenningsson, A. Massling, G. McFiggans, P. McMurry, T. Petäjä, P. Tunved, M. J. T. B. C. Gysel and P. Meteorology: Hygroscopic properties of submicrometer atmospheric aerosol particles measured with H-TDMA instruments in various environments—a review, 60, 432-469, 2008.
- 600 Tasoglou, A., R. Subramanian and S. N. Pandis: An inter-comparison of black-carbon-related instruments in a laboratory study of biomass burning aerosol, *Aerosol Science and Technology*, 52, 1320-1331, 2018.
- [Textor, C., M. Schulz, S. Guibert, S. Kinne, Y. Balkanski, S. Bauer, T. Bernsten, T. Berglen, O. Boucher, M. Chin, F. Dentener, T. Diehl, R. Easter, H. Feichter, D. Fillmore, S. Ghan, P. Ginoux, S. Gong, J. E. Kristjansson, M. Krol, A. Lauer, J. F. Lamarque, X. Liu, V. Montanaro, G. Myhre, J. Penner, G. Pitari, S. Reddy, O. Seland, P. Stier, T. Takemura and X. Tie: Analysis and quantification of the diversities of aerosol life cycles within AeroCom. *Atmospheric Chemistry and Physics*, 6, 1777-1813, 2006.](#)
- 605 [US EPA \(2019\), Integrated Science Assessment for Particulate Matter, US Environmental Protection Agency, US EPA/600/R-19/188, Dec 2019.](#)
- 610 [Vander Wal, R. L., V. M. Bryg and M. D. Hays: Fingerprinting soot \(towards source identification\): Physical structure and chemical composition, *Journal of Aerosol Science*, 41, 108-117, 2010.](#)
- Virkkula, A.: Modeled source apportionment of black carbon particles coated with a light-scattering shell, *Atmos. Meas. Tech.*, 14, 3707-3719, 2021.
- 615 von der Weiden, S. L., F. Drewnick and S. Borrmann: Particle Loss Calculator - a new software tool for the assessment of the performance of aerosol inlet systems, *Atmospheric Measurement Techniques*, 2, 479-494, 2009.
- WHO (2013). Review of evidence on health aspects of air pollution—REVIHAAP project: final technical report. Copenhagen, WHO.
- WHO (2021). WHO global air quality guidelines. Particulate matter (PM_{2.5} and PM₁₀), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide. Geneva: World Health Organization.
- 620 Wiedensohler, A., W. Birmili, A. Nowak, A. Sonntag, K. Weinhold, M. Merkel, B. Wehner, T. Tuch, S. Pfeifer, M. Fiebig, A. M. Fjåraa, E. Asmi, K. Sellegri, R. Depuy, H. Venzac, P. Villani, P. Laj, P. Aalto, J. A. Ogren, E. Swietlicki, P. Williams, P. Roldin, P. Quincey, C. Hüglin, R. Fierz-Schmidhauser, M. Gysel, E. Weingartner, F. Riccobono, S. Santos, C. Grüning, K.

- Faloon, D. Beddows, R. Harrison, C. Monahan, S. G. Jennings, C. D. O'Dowd, A. Marinoni, H. G. Horn, L. Keck, J. Jiang, J. Scheckman, P. H. McMurry, Z. Deng, C. S. Zhao, M. Moerman, B. Henzing, G. de Leeuw, G. Löschau and S. Bastian: Mobility particle size spectrometers: harmonization of technical standards and data structure to facilitate high quality long-term observations of atmospheric particle number size distributions, *Atmos. Meas. Tech.*, 5, 657-685, 2012.
- 625 Yuan, J., R. L. Modini, M. Zanatta, A. B. Herber, T. Müller, B. Wehner, L. Poulain, T. Tuch, U. Baltensperger and M. Gysel-Beer: Variability in the mass absorption cross-section of black carbon (BC) aerosols is driven by BC internal mixing state at a central European background site (Melpitz, Germany) in winter, *Atmos. Chem. Phys. Discuss.*, 2020, 1-36, 2020.
- 630 Zhang, Y., O. Favez, F. Canonaco, D. Liu, G. Močnik, T. Amodeo, J. Sciare, A. S. Prévôt, V. Gros, A. J. n. C. Albinet and A. Science: Evidence of major secondary organic aerosol contribution to lensing effect black carbon absorption enhancement, 1, 1-8, 2018.