



Seasonal updraft speeds change cloud droplet number concentrations in low level clouds over the Western North Atlantic

Simon Kirschler^{1,2}, Christiane Voigt^{1,2}, Bruce Anderson³, Ramon Campos Braga⁴, Gao Chen³, Andrea F. Corral⁵, Ewan Crosbie³, Hossein Dadashazar⁵, Richard A. Ferrare³, Valerian Hahn^{1,2}, Johannes Hendricks¹, Stefan Kaufmann^{1,2}, Richard Moore³, Mira L. Pöhlker^{6,7}, Claire Robinson³, Amy J. Scarino³, Dominik Schollmayer^{1,2}, Michael A. Shook³, K. Lee Thornhill³, Edward Winstead³, Luke D. Ziemba³, and Armin Sorooshian^{5,8}

¹Institut für Physik der Atmosphäre, Deutsches Zentrum für Luft- und Raumfahrt (DLR), Oberpfaffenhofen, Germany

²Institut für Physik der Atmosphäre, Johannes Gutenberg-Universität, Mainz, Germany

³NASA Langley Research Center, Hampton, VA, USA

⁴Multiphase Chemistry Department, Max Planck Institute for Chemistry, Mainz, Germany

⁵Department of Chemical and Environmental Engineering, University of Arizona, Tucson, Arizona, USA

⁶Experimental Aerosol and Cloud Microphysics Department, Leibniz Institute for Tropospheric Research, Leipzig, Germany

⁷Faculty of Physics and Earth Sciences, Leipzig Institute for Meteorology, University of Leipzig, Leipzig, Germany

⁸Department of Hydrology and Atmospheric Sciences, University of Arizona, Tucson, Arizona, USA

Correspondence: Kirschler Simon (Simon.Kirschler@dlr.de)

Abstract. Low level clouds over the Western North Atlantic show a seasonal cycle in cloud properties which anticorrelates to aerosol concentrations. To determinate the impact of dynamic and aerosol processes within marine low clouds we examine the seasonal impact of updraft speed w and cloud condensation nuclei concentration at 0.43% supersaturation ($N_{CCN_{0.43\%}}$) on the cloud droplet number concentration (N_C) of low level clouds over the Western North Atlantic Ocean. Aerosol and cloud properties were measured with instruments on board the NASA LaRC Falcon HU-25 during the ACTIVATE (Aerosol Cloud meTeorology Interactions oVer the western ATLantic Experiment) mission in summer (August) and winter (February-March) 2020. The data are grouped in different $N_{CCN_{0.43\%}}$ loadings and the density functions of N_C and w near the cloud bases are compared. For low updrafts ($w < 1.3 \text{ m s}^{-1}$), N_C in winter are mainly limited by the updraft speed and in summer additionally by aerosols. At larger updrafts ($w > 3 \text{ m s}^{-1}$), N_C are impacted by the aerosol population, while at clean marine conditions cloud nucleation is aerosol limited and for high pollution it is influenced by aerosols and updraft. The aerosol size distribution in winter shows a bimodal distribution in clean marine environments, which transforms to a unimodal distribution in high pollution levels due to altering processes, whereas unimodal distributions prevail in summer with a significant difference in their aerosol concentration and composition. The increase in pollution level is accompanied with an increase of organic aerosol and sulfate compounds in both seasons. We demonstrate that N_C can be explained by cloud condensation nuclei activation through upwards processed air masses with varying fractions of activated aerosols. The activation highly depends on w and thus supersaturation between the different seasons, while the aerosol size distribution additionally affects N_C within a season. Our results quantify the seasonal influence of w and $N_{CCN_{0.43\%}}$ on N_C and can be used to improve the representation of low marine clouds in models.



20 1 Introduction

Understanding cloud formation processes and their influence on the Earth's climate system are fundamental to assess climate model forecast quality (Zelinka et al., 2014, 2017; Seinfeld et al., 2016; IPCC, 2021). The results of the model evaluation activities of the Coupled Model Intercomparison Project Phase 6 (CMIP6) show that improvements in cloud representation result in stronger shortwave cloud feedbacks and higher effective climate sensitivity to the global mean surface air temperature of the CMIP6 model ensemble (Bock et al., 2020). In particular, regions with large multi model mean biases in near-surface air temperature and their cloud feedback are of high interest (Andrews et al., 2015; Ceppi et al., 2017) and targeted by various field campaigns (e.g., Lu et al., 2007; Hersey et al., 2009; Wood et al., 2011; Russell et al., 2013; Knippertz et al., 2015; Wendisch et al., 2016; Flamant et al., 2018; Sorooshian et al., 2018; Formenti et al., 2019; Sorooshian et al., 2019).

Atmospheric aerosols can act as cloud condensation nuclei (CCN) and activate to cloud droplets in favourable conditions, determined by atmospheric ambient parameters such as supersaturation and aerosol size and chemical composition (Köhler, 1936; Twomey, 1959; Köhler et al., 2006; Reutter et al., 2009; Rosenfeld et al., 2014; Cecchini et al., 2017; Prabhakaran et al., 2020). This leads to an alteration of cloud droplet number concentration N_C (Twomey and Warner, 1967) and consequential cloud radiative effects (Twomey, 1977; Rosenfeld et al., 2019). Higher aerosol concentrations and additional CCN activation increases cloud lifetime and thickness by suppressing precipitation (Albrecht, 1989; Freud and Rosenfeld, 2012; Braga et al., 2017b). There are several approaches to quantify N_C with satellite measurements. A direct approach utilizes the cloud optical depth, the cloud droplet effective radius and cloud top temperature (Grosvenor et al., 2018). An indirect approach exploits the aerosol-cloud interaction and uses the aerosol optical depth (AOD) as a proxy for N_C (Quaas et al., 2008). (A list of symbols and abbreviations is given in Appendix A1.) Both approaches have high uncertainty, i.e. retrieving N_C with AOD from satellites remains a challenge (Gryspeerd et al., 2017; Painemal et al., 2020). Rosenfeld et al. (2016) have shown that based on a satellite methodology it is possible to retrieve cloud base N_C and supersaturation, which further yields the CCN concentration at a given supersaturation with an accuracy of $\pm 30\%$. However, satellites measure bulk properties which are limited in observing mechanisms on a microphysical scale (McComiskey et al., 2012). Consequently, in-situ measurements are needed to validate and enhance the understanding of the respective cloud processes.

This work focuses on the Western North Atlantic Ocean (WNAO) (Sorooshian et al., 2020) which provides ideal conditions for studying aerosol cloud interactions due to influence from the polluted East Coast of North America. Dadashazar et al. (2021b) find an anti-correlation in the seasonal cycle of AOD and N_C for this area, which is in contrast to findings in other regions (e.g., Penner et al., 2006, 2011; Quaas et al., 2008; Gryspeerd et al., 2016). Braga et al. (2017a) use a statistical approach (Haddad and Rosenfeld, 1997) to quantify the relationship of w to N_C at cloud bases of convective clouds over the Amazon basin. Braga et al. (2021) show good agreement of the derived relationship with an adiabatic parcel model. Our analysis focused on w and aerosol impact on N_C in marine boundary layer stratus and stratocumulus clouds near cloud base.



Global aerosol-climate simulations still suffer from large uncertainties in the representation of aerosol-cloud-radiation interactions (e.g., Myhre et al., 2013). Particularly large model uncertainties persist with regard to aerosol effects on marine clouds (e.g. McCoy et al., 2020, 2021). Simulating aerosol-cloud interactions in such models requires the application of micro-physical two-moment cloud schemes in combination with aerosol sub-models providing information about aerosol properties relevant for cloud formation (e.g., Lohmann et al., 2007; Lohmann and Hoose, 2009; Righi et al., 2020). Aerosol effects on N_C are described in these models on the base of dedicated parameterizations (e.g., Abdul-Razzak and Ghan, 2000; Ghan et al., 2011) which are driven by model information about the aerosol size distribution and composition as well as w . Comparisons with observational data are essential to evaluate the robustness of simulating these quantities as well as the resulting N_C . The present study provides consistent information about all of these quantities under marine conditions for different seasons. Hence it is a valuable contribution to the data base available for global aerosol-climate model evaluation and can, therefore, trigger important improvements of aerosol-climate simulations and the applied parameterizations of the cloud nucleation process.

In the following sections we show that the aerosol size distribution in combination with w determine N_C near cloud base of marine clouds regardless of thermodynamical conditions. Furthermore, we found that the aerosol size distribution indicates the availability of CCN from the aerosol population and w with the corresponding supersaturation signifying the fraction of activated CCN over the WNAO.

2 Methods

2.1 Region of study during the ACTIVATE campaign

The Aerosol Cloud meTerology Interactions oVer the western ATlantic Experiment (ACTIVATE) campaign (Sorooshian et al., 2019) is focused on probing clouds between $25^\circ - 50^\circ\text{N}$ and $60^\circ - 85^\circ\text{W}$. Clouds are characterized simultaneously by the low flying NASA Langley Falcon HU-25 and the King Air UC-12 flying above. The Falcon HU-25 provides detailed in-situ measurements of aerosol, cloud, gas and meteorological properties by sampling the marine boundary layer (MBL) at different altitudes down to 150 m above sea level, while the UC-12 probed clouds with remote sensing instruments flying at 8 – 10 km. The ACTIVATE mission plans for flights in three consecutive years (2020-2022) with 150 joint research flights (RF) and a total amount of 600 flight hours per aircraft, where 40 RF (35 joint, 5 alone with Falcon HU-25) with around 130 flight hours per plane were conducted in the first year 2020. Here we use measurements from the Falcon HU-25 RF of 2020 shown in Figure 1.

2.2 Instrumentation

2.2.1 Cloud measurements: The Fast Cloud Droplet Probe

The Fast Cloud Droplet Probe (FCDP) (O'Connor et al., 2008; Knop et al., 2021) manufactured by Stratton Park Engineering Company Incorporated (SPEC Inc.) is a forward-scattering probe, which counts single particles in the diameter size range of 1.5 – 50 μm . In this analysis we use only particles with diameters larger than 3 μm . The FCDP uses a laser beam at




Figure 1. The used subset of HU-25 flights track during the first year of ACTIVATE flights in 2020. Each line represents a research flight with the running number of deployment. Lines colored with shades of blue to green represent flights during winter (February–March) while dotted orange lines indicate flights during summer (August).

785 nm wavelength to collect light scattered by particles passing through the laser beam according to Mie theory in a $4^\circ - 12^\circ$ collection angle. A 70:30 beam splitter is used to split the collected light to a signal and qualifier detector. The signal detector has a $800 \mu\text{m}$ pinhole for coincidence reduction (Lance, 2012) and a rectangular slit aperture with $800 \mu\text{m}$ length and $200 \mu\text{m}$ width. Both detectors convert the incoming light intensity into corresponding voltages and amplify them over two stages. The beam diameter on the detectors depends on the distance of the measured particle from the focal plane of the collecting lens system. The ratio of the qualifier voltage to signal voltage is the so-called depth of field (DoF) criteria which can be used to limit the sample area of the probe, because the slit aperture width restricts the intensity on the qualifier detector depending on the magnification of the beam diameter. In this analysis we use a DoF criteria >0.6 , which is equivalent to a calibrated sample area of 0.248 mm^2 (Lance et al., 2010; Faber et al., 2018). With a sampling rate of 25 ns the FCDP additionally stores the transit time, inter-arrival time and waveform of each particle. These parameters are used for data corrections, see Baumgardner et al. (1985); SPEC inc (2012). Coincidence correction is applied by deriving a theoretical particle transit time, determined by particle air speed (PAS) and particle diameter, under consideration of a top hat intensity along the laser beam cross section. Measured particles with transit times larger than 125% of the theoretical transit time are deemed coincident and



95 are thus discarded. A shattering correction is done by using the adaptive method and a waveform symmetry filter is applied, both methods are described in SPEC inc (2012).

According to Baumgardner et al. (2017), light scattering probes have a propagated uncertainty in size due to Mie ambiguity, collection angles, coincidence, nonsphericity and shattering of 10 – 50% and a propagated uncertainty in N_C due to sample area uncertainty, coincidence and shattering of 10 – 30% (Kleine et al., 2018; Bräuer et al., 2021a, b). The FCDP with its fast
100 electronics, small pinhole feature for coincidence reduction and applicable filtering techniques can be classified among the lower end of both propagated uncertainties in size and N_C .

2.2.2 Cloud measurements: The Two-Dimensional Stereo Probe

The Two-Dimensional Stereo probe (2D-S) from SPEC Inc. is an optical array probe, which generates shadow images of particles with a linear array of 128 photodiodes (Lawson et al., 2006, 2019). It measures single particles in a size range of
105 $5.7 - 1465 \mu m$ with an effective pixel size of $11.4 \mu m$ for each photodiode channel. The 2D-S has two identical subsystems perpendicularly aligned with a combination of transmitting and receiving arm each. Both arms operate with a laser of $785 nm$ wavelength and traversing particles generate a diffraction pattern according to diffraction theory. Similar to the FCDP the 2D-S has an optical plane, determined by the focal points of the light collecting lens system. Each photodiode is triggered if light intensity falls below a threshold of 50%. All shadowed photodiodes are recorded at a fast succession, specified by the sampling
110 rate, while a particle passes the laser beam. A recorded ensemble of slices produces a two-dimensional image of the particle (Knollenberg, 1970).

The sample area of the 2D-S depends on the particle size. The diffraction pattern can be calculated analytically with angular spectrum theory or fresnel theory (Korolev et al., 1991) and depends on the particles distance to the optical plane. With increasing distance the diffracted light forms spots of destructive interference and the particle is magnified. These particles
115 are classified as out-of-focus. In the case of spheroidal liquid droplets the so called poisson spot forms at the center of the projected circle and a size correction can be applied by relating the radius of the poisson spot to that of the captured particle image (Korolev, 2007). The magnification results in a decrease of the shadowing on the photodiodes until it is below the 50% threshold and no photodiode is triggered. The size dependent optical depth of field was verified for optical array probes according to Korolev et al. (1998). The maximum optical depth of field equals half of the $63 mm$ distance of the 2D-S
120 transmitting arm to the receiving arm. The maximum is reached with a drop radius of $109 \mu m$ and the maximum sample area can be calculated by multiplying the arm distance with the array width. In this work the all-in method is used to determine the effective array width and thus the sample area. The all-in method rejects particles with occulted edge photodiodes and adjusts the sample area depending on size, because the possibility of large particles rejected is higher compared to small particles (Knollenberg, 1970).

125 The 2D-S measures with a constant sampling rate resulting in an artificially elongated/shortened particle image if the actual PAS deviates from the PAS for which the sampling rate was computed. The PAS was measured by a pitot tube attached to a Cloud Aerosol and Precipitation Spectrometer (Voigt et al., 2017, 2021), which was mounted on the opposite wing at the same



position as the 2D-S/FCDP Combination. With the PAS to sampling rate ratio the deformed images can be corrected (Weigel et al., 2016).

130 Sizing accuracy is affected by out of focus and shattered particles, the time response and discretization of the probe hardware and lies for imaging probes in a 10 – 100% range according to Baumgardner et al. (2017). While the uncertainty of out of focus spheroids is reduced with Korolev’s correction it remains for ice particles. The 2D-S has a relative fast response time of 41 *ns* and can be classified on the lower end of the uncertainty range for spheroids and in the middle for ice particles (Lawson and Baker, 2006; Baker and Lawson, 2006; Gurganus and Lawson, 2018). The concentration is affected by the size dependent
135 optical depth of field and shattering (Lawson, 2011). With shattering removal and an adjusted sample area the 2D-S here is similarly representative for the lower range of 10%-100% propagated uncertainty in N_C .

Measured size distributions of FCDP and 2D-S overlap in the size range of 16 – 51.3 μm . We perform the overlap calculation for the size range between the lower FCDP bin edge at 27 μm to the upper 2D-S bin edge of 39.9 μm . The particle distribution inside the overlap 2D-S bin is estimated with the next 2D-S bin by linear interpolation and attributed proportionally to the last
140 FCDP bin and a new 2D-S bin.

2.2.3 Vertical velocity

The winds on the HU-25 are measured by the NASA Langley TAMMS (turbulent air motion measurement system). The primary components include fast-response flow-angle and temperature sensors to determine the wind with respect to the aircraft along with an Applanix 650 inertial navigation system (Applanix Inc.) to provide the aircraft’s position, speed and altitude.
145 The data is recorded at 200 Hz on a UEIPAC-300 real-time controller (United Electronics Industries) and then averaged down to 20 Hz for processing, analysis, and data archiving. The flow-angle system includes five, flush-mounted pressure-ports installed in a cruciform pattern in the aircraft radome to provide angle of attack (vertically-aligned ports) and side-slip (horizontally aligned ports) measurements. Corresponding fast-response high-precision pressure transducers are placed as close as possible to the pressure ports in order to minimize delays and errors. Pitch and yaw maneuvers, speed variations and reverse
150 headings are performed periodically during deployments to verify system operation and calibration and validate derived mean horizontal-wind vectors. Three dimensional winds are computed from the full air motion equations (Lenschow, 1986). The aircraft platform velocity components are computed internally by the Applanix by combining the GPS and inertial data via a Kalman filtering technique. Ambient air temperature measurements needed to determine true air speed are made with a Rosemount Model 102 non-deiced total air temperature sensor with a fast response platinum sensing element (E102E4AL).
155 This setup has been used extensively for other campaigns on the NASA P-3 aircraft. Additional details on the instrumentation, calibration, and intercomparison results of the TAMMS when used on the NASA P-3 can be found at Thornhill et al. (2003). All wind measurements including horizontal and vertical winds have a 5% uncertainty.

2.2.4 Aerosol Measurements: Cloud Condensation Nuclei

The CCN number concentrations were measured with a CCN-100 counter manufactured by Droplet Measurement Technolo-
160 gies, which is based on the concept of Roberts and Nenes (2005) and characterized by Lance et al. (2006). The CCN-100 was



operated in two modes during ACTIVATE. The first is a continuous flow mode where ambient air enters a column shaped humidified chamber with a constant supersaturation of 0.43%. Aerosols are activated depending on their size and chemical properties. The droplets are measured afterwards by an optical particle counter. The second is a scanning flow mode where the flow rate in the chamber is changed while a constant temperature gradient is maintained (Moore and Nenes, 2009). Here an aerosol sample is exposed to a continuously changing supersaturation in the chamber and the concentration of activated aerosols N_{CCN} is measured depending on supersaturation. One scan is typically done in a 10 – 60 seconds time interval and in this analysis we use the mean of N_{CCN} in a supersaturation range of 0.40–0.46% to approximate $N_{CCN_{0.43\%}}$. The uncertainty in percent supersaturation is ± 0.04 and in $N_{CCN_{0.43\%}}$ $\pm 10\%$.

Since the instrument supersaturation is fixed in continuous flow mode and artificially generated in scanning flow mode we have to estimate the supersaturation in cloud base. The maximum supersaturation S_{max} is calculated according to Pinsky et al. (2012) with

$$S_{max} = Cw^{\frac{3}{4}}N_c^{-\frac{1}{2}} \quad (1)$$

where C is determined by cloud base temperature and pressure, and w and N_c are the updraft speed and cloud droplet number concentration, respectively, measured in cloud base.

2.2.5 Aerosol Measurements: Chemical Composition

Submicron non-refractory aerosol chemical composition was measured by a High Resolution Time-of-Flight Aerosol Mass Spectrometer (HR-ToF-AMS; Aerodyne Research Inc. DeCarlo et al., 2006; Hilario et al., 2021). Mass concentrations of sulfate, nitrate, chloride, ammonium, and organic matter were recorded at 1Hz and averaged to 30-s for all subsequent analyses. Measurements were made isokinetically using a forward-facing dual-diffuser aircraft inlet (model 1200, Brechtel Manufacturing Inc.) and were pressure-controlled at 500 torr. Mass concentrations were processed using default relative ionization efficiencies for each chemical component, with a collection efficiency of unity, and are reported at standard temperature and pressure (STP; 273.15 K and 1013.25 mb).

The particle-into-liquid sampler (PILS) obtained water-soluble aerosol composition data. Sampled aerosol particles were grown into droplets that were collected via inertial impaction and transported to vials on a rotating carousel. The liquid content of the vials was analyzed post-flight via ion chromatography for water-soluble ions (Sorooshian et al., 2006). This study reports on sea salt concentrations by attributing Na^+ , Cl^- and Mg^{2+} exclusively to sea salt and adding SO_4^{2-} (0.25), K^+ (0.036) and Ca^{2+} (0.039) fractions based on their ratio to Na^+ in sea water (Bowen, 1979; Farren et al., 2019; Ma et al., 2021). The PILS sample particles up to roughly 4 μm in diameter (McNaughton et al., 2007; Hilario et al., 2021).

2.2.6 Aerosol Measurements: Size Distribution

Aerosol size distributions were obtained from a combination of two particles sizers. A custom Scanning Mobility Particle Sizer (SMPS; Differential Mobility Analyzer model 3085 and Condensation Particle Counter model 3776, TSI, Inc., Moore




Figure 2. Typical flight pattern of one cloud ensemble of the HU-25 aerosol, cloud and vertical velocity data set. In this study we focus on the four legs in the beginning of each ensemble containing two pairs (yellow and orange shaded areas) consisting of a below cloud base leg followed by an above cloud base leg. The blue shading indicates the cloud layer.

et al., 2017) measured 3.2 – 89.1 nm diameter particle size distributions at approximately 60 s time response. A Laser Aerosol Spectrometer (LAS; model 3340, TSI, Inc.; Moore et al., 2021) measured 100 – 3162 nm diameter particles at 1 Hz time response. SMPS sizing is calibrated and frequently verified using NIST-traceable polystyrene latex spheres. LAS sizing is calibrated using lab-generated monodisperse ammonium sulfate (refractive index = 1.52). Each instrument sampled dried air from the same common inlet as the HR-ToF-AMS and data are reported at STP.

2.3 Methodology

In this study we select the data a priori into pairs of series of below cloud base (BCB) and above cloud base (ACB) legs resulting in two pairs per ensemble (ensemble is a collection of legs below, in, and above clouds) flown during ACTIVATE, shown in Figure 2. This flight design intends for measurements to reflect the same environment. Closely spaced aerosol and cloud measurements are ensured by taking the latest full N_{CCN} scan or 60 seconds of continuous $N_{CCN_{0.43\%}}$ measurements of the BCB leg and the last measurement of a cloud portion in the nearest ACB leg is restricted to never exceed a horizontal distance of 40 km to the aerosol measurement. Cloud periods are defined as seconds with a threshold of liquid water content $> 0.02 \text{ gm}^{-3}$ and $N_C > 20 \text{ cm}^{-3}$. We additionally excluded pairs with precipitation occurrences in the BCB leg by using the 2D-S size distribution and images, since the $N_{CCN_{0.43\%}}$ measurements are influenced by the large particles shattering on the aerosol inlet and precipitation indicates that the cloud is at a different point of its life cycle where agglomeration and coalescence altered N_C and aerosol removal occurred below cloud. Each flight leg pair consists of a $N_{CCN_{0.43\%}}$ distribution taken from the pair's BCB leg either in continuous flow or scanning flow mode, a mean aerosol loading derived from the



$N_{CCN_{0.43\%}}$ distribution, N_C and positive vertical velocity measurements (updraft speeds w) in cloud portions of the pair's ACB
210 leg.

For ensuring similar environmental conditions the pairs are classified with respect to their mean $N_{CCN_{0.43\%}}$ into a low pol-
luted (LP), medium polluted (MP) and high polluted (HP) group. For comparison both seasons share the boundaries separating
the groups and the bin boundaries are chosen by identifying modes in the distribution of all winter pair mean $N_{CCN_{0.43\%}}$ values.
The LP groups contains $N_{CCN_{0.43\%}}$ from the minimum measured to 372 cm^{-3} , the MP group extends from $> 372 - 769\text{ cm}^{-3}$
215 and the HP group is defined for $> 769\text{ cm}^{-3}$ to the maximum measured in the respective season. The Probability Matching
Method (PMM) is used on each group's set of N_C and w within a 2.5 to 97.5 percentile interval to quantify the impact of w for
the different pollution levels.

We use the effective updraft speed w_{eff} for approximating the updraft through the measured w density function in cloud base

$$w_{\text{eff}} = \frac{\int w^2}{\int w}. \quad (2)$$

220 With the help of the w to N_C relation from the PMM the corresponding N_C to w_{eff} can be derived and therefore a S_{max}
estimate, which is representative for the supersaturation in cloud base of the respective group. We use the variability and
magnitude of w with the related S_{max} estimates, the aerosol size distribution and chemical composition to quantify their
contribution to the activation of CCN in the winter and summer season 2020 for different pollution levels.

2.3.1 Probability Matching Method

225 The PMM was proposed by Calheiros and Zawadzki (1987) for a statistical comparison of radar reflectivity to rain rate. The
derived relationship is verified and performs significantly better than power law regression (Rosenfeld et al., 1994). Addi-
tional improvements by taking physical parameters into account for different rain type classification were done by Rosenfeld
et al. (1995). The PMM is mathematically justified with an error estimation by Haddad and Rosenfeld (1997), and Braga et al.
(2017a) showed that the PMM can be applied to get a reasonable relationship of w to N_C . The PMM is based on the assumption
230 that two related parameters taken in non-simultaneous measurements, sharing the same environment in terms of climatolog-
ical and physical means, are increasing monotonically with each other. The relationship can be computed by matching the
percentiles of the parameter's density functions, with more details on the mathematical background described in Haddad and
Rosenfeld (1997). Braga et al. (2021) showed good agreement between measurements of N_C at cloud bases of convective
clouds and estimations from an adiabatic parcel model.

235 3 Results and Discussion

3.1 Measurements of $N_{CCN_{0.43\%}}$ below and N_C , and w near Cloud Base

The measurements are grouped into pairs with consecutive BCB and ACB legs, illustrated in Figure 3a for a pair during RF02
on 15 February 2020. In Figure 3b the size distribution of the 2D-S/FCDP combination is shown. The FCDP measures a




Figure 3. a) Aerosol and cloud properties measured in an ensemble pair of BCB and ACB legs from RF02 on 15 February 2020. The Falcon altitude is given by the black line and the yellow and blue shading indicates the BCB and the ACB legs. N_C (red), $N_{CCN_{0.43\%}}$ (green) and w (orange) are shown as lines. The aerosol loading $N_{CCN_{0.43\%}}$ representative of the pair’s environment is $440(\pm 20) cm^{-3}$, calculated from the measurements mean within the BCB leg between 0.40–0.46% supersaturation of the CCN-100 scans nearest to the cloud measurement. The pair’s mean values with standard deviation for w and N_C are $1.85(\pm 0.82) ms^{-1}$ and $385(\pm 171) cm^{-3}$, respectively. b) Histogram showing the color-coded log-normalized number concentrations per bin on a 1-second basis of the 2D-S/FCDP combination with the diameter given in the ordinate.

constant particle background in a size range of 3–12 μm with concentrations between 0.1 to 1 per cubic centimeter. The background is visible in all BCB legs, but vanishes at flight levels above cloud top above the boundary layer (not shown). The relative humidity data show values mainly above 75% in the BCB legs for both seasons, suggesting the measured background concentrations being deliquescent sea salt particles.

240



Table 1. All pairs consisting of serial below cloud base (BCB) and above cloud base (ACB) legs during the February-March 2020 deployment. Mean values and standard deviation in parenthesis for w and N_C from ACB cloud portions, and $N_{CCN_{0.43\%}}$ from the BCB legs. D_{\max} is the maximal distance of cloud measurements to the aerosol measurements and h_{ACB} is the height above cloud base with standard deviation in parenthesis.

Flight	Date	t_{initial} [UTC]	CCN-100 Mode (Supersat [%])	D_{\max} [km]	in-cloud [s]	$N_{CCN_{0.43\%}}$ [cm^{-3}]	w [ms^{-1}]	N_C [cm^{-3}]	h_{ACB} [m]
RF01	14 Feb 2020	17:21:32	Scan (0.17-0.70)	37.2	19	647 (\pm 35)	0.83 (\pm 0.56)	298 (\pm 173)	127 (\pm 4)
RF01	14 Feb 2020	17:30:17	Scan* (0.17-0.70)	35.3	28	664 (\pm 50)	1.67 (\pm 0.70)	593 (\pm 492)	136 (\pm 14)
RF01	14 Feb 2020	17:58:43	Scan (0.16-0.69)	28.3	51	582 (\pm 46)	1.74 (\pm 1.21)	723 (\pm 344)	103 (\pm 5)
RF01	14 Feb 2020	18:05:17	Scan* (0.16-0.68)	35.6	44	582 (\pm 36)	2.07 (\pm 1.26)	570 (\pm 308)	111 (\pm 3)
RF02	15 Feb 2020	17:09:31	Scan (0.17-0.71)	22.9	59	436 (\pm 37)	0.62 (\pm 0.48)	389 (\pm 217)	82 (\pm 6)
RF02	15 Feb 2020	17:18:16	Scan (0.17-0.71)	19.8	58	630 (\pm 36)	0.63 (\pm 0.33)	648 (\pm 279)	73 (\pm 2)
RF02	15 Feb 2020	18:23:53	Scan (0.16-0.71)	29.4	34	489 (\pm 34)	0.87 (\pm 0.52)	297 (\pm 223)	147 (\pm 6)
RF02	15 Feb 2020	18:32:38	Scan (0.16-0.71)	38.4	130	440 (\pm 20)	1.85 (\pm 0.82)	385 (\pm 171)	200 (\pm 3)
RF03	17 Feb 2020	17:41:11	Scan* (0.17-0.71)	40.0	74	1564 (\pm 65)	0.25 (\pm 0.29)	930 (\pm 663)	93 (\pm 3)
RF09	27 Feb 2020	18:47:10	Scan* (0.16-0.72)	32.7	62	659 (\pm 39)	0.72 (\pm 0.53)	671 (\pm 357)	98 (\pm 5)
RF09	27 Feb 2020	18:55:55	Scan (0.17-0.72)	29.7	36	575 (\pm 46)	0.64 (\pm 0.53)	336 (\pm 218)	125 (\pm 6)
RF09	27 Feb 2020	19:28:43	Scan (0.16-0.71)	37.5	41	582 (\pm 29)	0.73 (\pm 0.54)	467 (\pm 250)	145 (\pm 5)
RF09	27 Feb 2020	19:39:39	Scan (0.17-0.71)	33.2	48	656 (\pm 42)	0.91 (\pm 0.77)	355 (\pm 224)	189 (\pm 19)
RF09	27 Feb 2020	20:10:17	Scan (0.16-0.71)	28.7	42	674 (\pm 29)	1.13 (\pm 0.94)	716 (\pm 377)	151 (\pm 4)
RF09	27 Feb 2020	20:19:02	Scan (0.16-0.71)	31.9	35	650 (\pm 35)	0.83 (\pm 0.68)	647 (\pm 292)	199 (\pm 4)
RF13	01 Mar 2020	14:10:32	Scan* (0.16-0.71)	28.2	96	1217 (\pm 93)	1.57 (\pm 1.28)	1020 (\pm 556)	113 (\pm 4)
RF13	01 Mar 2020	15:00:51	Scan (0.16-0.72)	37.2	74	361 (\pm 19)	1.54 (\pm 1.63)	372 (\pm 197)	169 (\pm 5)
RF13	01 Mar 2020	16:02:06	Scan* (0.17-0.71)	36.7	51	769 (\pm 41)	1.46 (\pm 1.30)	818 (\pm 721)	139 (\pm 3)
RF16	06 Mar 2020	19:34:26	Scan (0.17-0.71)	32.3	55	991 (\pm 46)	0.99 (\pm 0.73)	1367 (\pm 958)	208 (\pm 6)
RF16	06 Mar 2020	19:43:11	Scan (0.16-0.72)	28.3	36	1788 (\pm 109)	1.80 (\pm 1.06)	1157 (\pm 912)	100 (\pm 3)
RF16	06 Mar 2020	20:15:59	Scan (0.17-0.72)	29.9	49	1501 (\pm 71)	1.84 (\pm 1.06)	1014 (\pm 742)	130 (\pm 6)
RF16	06 Mar 2020	20:24:44	Scan* (0.17-0.72)	34.7	33	945 (\pm 53)	1.55 (\pm 1.27)	397 (\pm 358)	193 (\pm 5)
RF17	08 Mar 2020	14:34:49	Flow (0.43)	25.0	39	183 (\pm 28)	1.05 (\pm 0.88)	434 (\pm 228)	117 (\pm 3)
RF17	08 Mar 2020	14:44:29	Flow (0.43)	29.2	17	245 (\pm 31)	1.35 (\pm 0.68)	498 (\pm 214)	135 (\pm 3)
RF17	08 Mar 2020	15:11:45	Flow (0.43)	28.7	112	164 (\pm 26)	0.46 (\pm 0.45)	208 (\pm 93)	173 (\pm 4)
RF17	08 Mar 2020	15:23:22	Flow (0.43)	28.3	72	96 (\pm 18)	0.95 (\pm 0.98)	218 (\pm 101)	163 (\pm 4)
RF17	08 Mar 2020	15:52:58	Flow (0.43)	30.1	56	196 (\pm 27)	0.83 (\pm 0.85)	386 (\pm 212)	91 (\pm 3)
RF17	08 Mar 2020	16:02:17	Flow (0.43)	19.8	65	225 (\pm 33)	1.52 (\pm 1.34)	346 (\pm 149)	129 (\pm 4)
RF19	09 Mar 2020	17:27:27	Flow (0.43)	28.6	26	291 (\pm 34)	0.63 (\pm 0.48)	208 (\pm 146)	125 (\pm 6)
RF19	09 Mar 2020	17:57:47	Flow (0.43)	23.7	20	299 (\pm 44)	0.61 (\pm 0.44)	247 (\pm 125)	121 (\pm 4)



245

Flight	Date	t _{initial} [UTC]	CCN-100 Mode (Supersat [%])	D _{max} [km]	in-cloud [s]	N _{CCN_{0.43%}} [cm ⁻³]	w [ms ⁻¹]	N _C [cm ⁻³]	h _{ACB} [m]
RF19	09 Mar 2020	18:41:47	Flow (0.43)	17.5	18	335 (± 46)	0.43 (± 0.28)	215 (± 114)	224 (± 2)
RF19	09 Mar 2020	18:50:13	Flow (0.43)	37.4	24	307 (± 36)	0.64 (± 0.56)	285 (± 171)	196 (± 11)
RF20	11 Mar 2020	13:46:55	Flow (0.43)	25.3	22	875 (± 101)	0.45 (± 0.46)	780 (± 430)	62 (± 2)
RF20	11 Mar 2020	14:26:13	Flow (0.43)	23.3	10	986 (± 134)	0.26 (± 0.21)	320 (± 221)	42 (± 3)
RF21	12 Mar 2020	14:43:10	Flow (0.43)	25.2	19	586 (± 84)	1.64 (± 1.07)	675 (± 383)	141 (± 4)
RF21	12 Mar 2020	14:51:22	Flow (0.43)	27.4	30	500 (± 91)	0.77 (± 0.70)	458 (± 275)	140 (± 4)
RF21	12 Mar 2020	15:19:57	Flow (0.43)	21.2	42	587 (± 102)	0.78 (± 0.72)	654 (± 418)	71 (± 2)
RF21	12 Mar 2020	16:06:00	Flow (0.43)	22.3	34	494 (± 58)	0.68 (± 0.48)	559 (± 255)	116 (± 3)
RF21	12 Mar 2020	16:14:01	Flow (0.43)	38.8	25	455 (± 72)	0.69 (± 0.57)	584 (± 261)	124 (± 46)
All RF	Average			29.3	45	612	1.02	535	130

* Only one scan in BCB leg.

All data ensemble pairs used in this study from the ACTIVATE winter February-March 2020 deployment are given in Table 1. We use selected pairs with a minimum in-cloud time above or equal 10 seconds for sufficient statistics. N_C is predicted to reach its maximum at a height above cloud base depending on *w* and subsequent S_{max} estimates in an adiabatic parcel model (Braga et al., 2021). That the activation of CCN into cloud droplets had sufficient time is ensured by taking only pairs into account with cloud measurements at a height above cloud base h_{ACB} greater than 35 m in this analysis. The h_{ACB} is gauged by calculating the middle of the difference between leg-mean values at BCB and ACB altitudes. In total we use 39 pairs from 10 RF, where all needed data are available for the PMM application, with a combined duration of 1786 seconds in cloud. The aerosol loading mean N_{CCN_{0.43%}} values range from 96 cm⁻³ in clean conditions up to 1788 cm⁻³ in high polluted environments. The mean of N_C is between 208 – 1367 cm⁻³. The measured *w* distributional mean ranges from 0.25 up to 2.07 ms⁻¹. During RF02 15 February 2020 flight a distinct shift of N_{CCN_{0.43%}} was measured between 17:42 to 17:57 UTC which can be attributed to a plume crossing event and affected pairs were excluded from the analysis, because the link between aerosol environment and measured N_C through cloud formation is questionable. The horizontal distance between aerosol measurements below cloud and cloud measurements in cloud base is mainly below 30 km and never exceeds 40 km. Results derived from the PMM are more robust with a choice of narrow a priori boundaries for classifying similar environmental conditions.

The same procedure was applied to the flights of the ACTIVATE August 2020 deployment resulting in the pairs listed in Table 2. We use a total of 16 pairs from 5 RF with a combined duration of 360 seconds in cloud. The full data set of the ACTIVATE August-September 2020 deployment including CCN measurements is only available for the August period limiting available pairs. The reduced fraction of time in cloud is in line with the observed lower cloud fraction and horizontal dimension of clouds during summer. In addition to excluding pairs affected by precipitation the pairs in RF28 were not used in the analysis because of a smoke layer possibly altering the cloud formation process. The aerosol loading mean N_{CCN_{0.43%}} values range from 122 cm⁻³ in clean conditions up to 1995 cm⁻³ in high polluted environments. The pairs in summer exhibit



Table 2. All pairs consisting of serial below cloud base (BCB) and above cloud base (ACB) legs during the August 2020 deployment. Mean values and standard deviation in parenthesis for w and N_C from ACB cloud portions, and $N_{CCN_{0.43\%}}$ from the BCB legs. D_{\max} is the maximal distance of cloud measurements to the aerosol measurements and h_{ACB} is the height above cloud base with standard deviation in parenthesis.

Flight	Date	t_{initial} [UTC]	CCN-100 Mode (Supersat [%])	D_{\max} [km]	in-cloud [s]	$N_{CCN_{0.43\%}}$ [cm^{-3}]	w [ms^{-1}]	N_C [cm^{-3}]	h_{ACB} [m]
RF23	13 Aug 2020	14:48:15	Scan (0.16-0.71)	19.4	31	225 (± 22)	0.55 (± 0.32)	169 (± 71)	129 (± 13)
RF23	13 Aug 2020	16:59:29	Scan (0.17-0.71)	25.1	55	267 (± 30)	0.39 (± 0.27)	145 (± 68)	164 (± 2)
RF24	17 Aug 2020	14:54:38	Scan (0.16-0.72)	22.7	26	304 (± 30)	0.68 (± 0.37)	208 (± 87)	152 (± 12)
RF24	17 Aug 2020	15:01:12	Scan* (0.17-0.71)	36.7	18	372 (± 22)	0.64 (± 0.42)	163 (± 105)	101 (± 10)
RF24	17 Aug 2020	15:34:01	Scan* (0.16-0.71)	27.0	31	122 (± 10)	0.87 (± 0.70)	103 (± 62)	71 (± 16)
RF24	17 Aug 2020	16:57:07	Scan* (0.17-0.72)	36.5	15	204 (± 16)	0.82 (± 0.54)	173 (± 66)	127 (± 6)
RF25	20 Aug 2020	14:42:26	Scan (0.18-0.71)	20.0	18	1744 (± 110)	0.89 (± 0.69)	649 (± 510)	94 (± 3)
RF25	20 Aug 2020	14:49:00	Scan* (0.17-0.71)	30.1	11	1586 (± 82)	0.95 (± 1.16)	658 (± 605)	79 (± 2)
RF25	20 Aug 2020	15:13:03	Scan* (0.17-0.71)	35.0	20	1291 (± 54)	0.78 (± 0.58)	484 (± 462)	123 (± 4)
RF25	20 Aug 2020	15:54:36	Scan* (0.17-0.72)	31.7	24	1113 (± 76)	0.71 (± 0.54)	557 (± 449)	89 (± 2)
RF25	20 Aug 2020	16:03:21	Scan (0.16-0.71)	26.2	24	1266 (± 47)	0.72 (± 0.48)	739 (± 537)	61 (± 3)
RF26	21 Aug 2020	15:35:56	Scan (0.16-0.71)	32.2	20	1261 (± 78)	0.35 (± 0.28)	458 (± 373)	57 (± 9)
RF27	25 Aug 2020	15:10:29	Scan (0.16-0.72)	28.8	15	1627 (± 101)	0.84 (± 0.66)	338 (± 232)	129 (± 4)
RF27	25 Aug 2020	15:19:14	Scan (0.16-0.71)	16.1	10	1529 (± 91)	0.70 (± 0.65)	440 (± 324)	136 (± 6)
RF27	25 Aug 2020	16:18:18	Scan* (0.16-0.71)	33.4	19	1794 (± 89)	0.59 (± 0.50)	536 (± 434)	82 (± 3)
RF27	25 Aug 2020	16:27:02	Scan (0.16-0.72)	35.3	23	1995 (± 186)	0.35 (± 0.27)	575 (± 455)	105 (± 3)
All RF	Average			28.5	23	1044	0.68	400	106

* Only one scan in BCB leg.

a bimodal distribution of either very clear or high polluted conditions and are similar within a day while the high polluted conditions occur within a higher frequency. The N_C mean ranges from 103 up to 739 cm^{-3} which is 25% lower in terms of all pairs average of 400 cm^{-3} compared to the wintertime 535 cm^{-3} average and in good agreement with the findings of Dadashazar et al. (2021b). A similar trend is observed in the w measurements where the mean w is from 0.35 to 0.95 ms^{-1} and thus 33% lower in terms of all pairs average 0.68 ms^{-1} in comparison with the wintertime 1.02 ms^{-1} average. Also, less variability of updraft speed was measured with an average of 0.53 ms^{-1} (76%) compared to 0.76 ms^{-1} (78%) during wintertime indicating a higher dynamical influence during winter, i.e., with a high intraday variability.




Figure 4. Probability distributions of wintertime $N_{CCN_{0.43\%}}$ of the low polluted (LP/blue), medium polluted (MP/orange) and high polluted (HP/red) group with their mean values (a). All values are binned with a bin width of 30 cm^{-3} and the abscissa gives the probability of occurrence in the group. The same is done in b) for the summertime $N_{CCN_{0.43\%}}$ distribution of the low polluted and high polluted group. No summertime pair is attributable to the MP group and thus not shown.

275 3.2 Seasonal Aerosol Distribution and Composition below Cloud Base

The pair measurements indicate a correlation between N_C and w which can be quantified by the PMM. For this purpose, all pairs of a season are sorted according to their mean aerosol loading into three groups where each has a new set of $N_{CCN_{0.43\%}}$, N_C and w respectively. Figure 4a depicts the $N_{CCN_{0.43\%}}$ distribution of each group observed during winter. The LP group ranges from $115 - 451 \text{ cm}^{-3}$ with a mean of 251 cm^{-3} for $N_{CCN_{0.43\%}}$. The MP(HP) group have values from $337(678) - 941(1903) \text{ cm}^{-3}$ with a mean $N_{CCN_{0.43\%}}$ of $542(1036) \text{ cm}^{-3}$ respectively. There are overlap regions between the groups due to the usage of single values of $N_{CCN_{0.43\%}}$ in Figure 4, but the separation is sufficient for the applicability of the PMM.

The same is shown for the summer period in Figure 4b, where only pairs corresponding to the LP and HP group were measured. For the summertime the LP(HP) group's $N_{CCN_{0.43\%}}$ values range from $181(971) - 403(2275) \text{ cm}^{-3}$ with a mean of $273(1513) \text{ cm}^{-3}$. The minimum value of the summertime HP group is noticeably higher compared to the wintertime HP group, does not have a distinct peak and is more equally distributed. The summer HP group's mean $N_{CCN_{0.43\%}}$ is 46% higher than the winter HP mean value while the LP groups are quite similar in shape. Interestingly, no pair in the MP group was measured during summer and $N_{CCN_{0.43\%}}$ is either in clean or high polluted conditions within a research flight.

$N_{CCN_{0.43\%}}$ are a subsample of the available aerosol population. The aerosol size distributions during wintertime are shown in Figure 5a. The clean marine environment (LP) has a distinct bimodal distribution consisting of an Aitken mode ($10 - 100 \text{ nm}$) peaking at around 20 nm and an accumulation mode ($100 - 1000 \text{ nm}$) at 100 nm . In contrast the HP group has a unimodal distribution with a flat peak at $40 - 100 \text{ nm}$ at similar $dN_{\log D_p}$ concentrations to the LP and MP group and exhibits a plateau below 20 nm which hints to an overlapping ultra fine particle mode. The integrated number concentration for particles greater than 85 nm N_{gt85} in the BCB leg, depicted in Figure 5c, shows that the steady increase from 472 cm^{-3} (LP) over




Figure 5. Wintertime aerosol size distributions from the SMPS/LAS instrument combination of the low polluted (LP) (blue), medium polluted (MP) (orange) and high polluted (HP) (red) group with their mean distribution and shaded distributions for a single pair (a). The same is done in b) for the summertime aerosol size distribution of the low polluted and high polluted group. Whisker plots of N_{gt85} with the mean marked by a cross for wintertime (c) and summertime (d). No summertime pair is attributable to the MP group and thus not shown in b) and d).

658 cm^{-3} (MP) to 1108 cm^{-3} (HP) is significant for LP to HP and probably stems from condensational growth and coagulation processes for both Aitken and accumulation modes. The unimodal HP size distribution originates from an overlapping Aitken and accumulation mode where the aerosol properties can differ. The MP group has a bimodal distribution with the maximum of the Aitken mode at around 15 nm and accumulation mode peaks similarly at 100 nm .

295



Table 3. All pair's aerosol composition below cloud base (BCB) during the February-March 2020 deployment. Mean values and standard deviation in parenthesis for organic aerosol (OA), SO_4^{2-} , NO_3^- , NH_4^+ from AMS measurements and sea salt from PILS measurements. All measurements are given at standard temperature and pressure.

Group	Flight	date	t_{initial} [UTC]	SeaSalt [μgm^{-3}]	OA [μgm^{-3}]	SO_4^{2-} [μgm^{-3}]	NO_3^- [μgm^{-3}]	NH_4^+ [μgm^{-3}]
LP	RF13	01 Mar 2020	14:53:22	1.97(±0.35)	0.57(±0.14)	0.93(±0.05)	0.06(±0.04)	0.41(±0.14)
LP	RF17	08 Mar 2020	14:32:31	2.39(±0.00)	0.35(±0.20)	0.30(±0.03)	0.05(±0.02)	0.18(±0.07)
LP	RF17	08 Mar 2020	14:41:44	4.23(±2.96)	0.32(±0.13)	0.35(±0.05)	0.07(±0.04)	0.06(±0.04)
LP	RF17	08 Mar 2020	15:09:24	3.69(±0.00)	0.15(±0.10)	0.36(±0.05)	0.02(±0.03)	0.03(±0.11)
LP	RF17	08 Mar 2020	15:50:27	4.87(±0.00)	0.42(±0.08)	0.39(±0.02)	0.07(±0.03)	0.26(±0.13)
LP	RF17	08 Mar 2020	15:59:48	4.42(±0.49)	0.43(±0.12)	0.42(±0.03)	0.04(±0.05)	0.04(±0.16)
LP	RF19	09 Mar 2020	17:25:14	3.14(±0.00)	0.16(±0.08)	0.30(±0.04)	0.04(±0.03)	0.07(±0.12)
LP	RF19	09 Mar 2020	17:55:32	3.80(±0.00)	0.08(±0.16)	0.34(±0.02)	0.03(±0.04)	0.03(±0.09)
LP	RF19	09 Mar 2020	18:39:22	3.80(±0.13)	0.27(±0.19)	0.36(±0.03)	0.02(±0.02)	0.07(±0.11)
LP	Average			3.59	0.31	0.42	0.04	0.13
MP	RF01	14 Feb 2020	17:19:21	0.98(±0.65)	1.20(±0.31)	1.24(±0.05)	2.07(±0.28)	1.36(±0.10)
MP	RF01	14 Feb 2020	17:28:38	2.16(±0.17)	1.26(±0.19)	1.33(±0.16)	1.94(±0.23)	1.39(±0.25)
MP	RF01	14 Feb 2020	17:56:27	4.19(±0.00)	0.92(±0.12)	0.89(±0.06)	0.58(±0.06)	0.54(±0.16)
MP	RF01	14 Feb 2020	18:04:21	5.84(±0.00)	1.08(±0.09)	0.96(±0.07)	0.64(±0.06)	0.72(±0.13)
MP	RF02	15 Feb 2020	17:07:07	1.89(±0.93)	0.80(±0.16)	0.66(±0.06)	0.14(±0.05)	0.29(±0.09)
MP	RF02	15 Feb 2020	17:15:06	3.22(±0.25)	0.99(±0.12)	0.66(±0.05)	0.18(±0.04)	0.37(±0.16)
MP	RF02	15 Feb 2020	18:17:32	4.20(±1.28)	0.83(±0.14)	0.82(±0.05)	0.08(±0.04)	0.37(±0.20)
MP	RF02	15 Feb 2020	18:30:07	5.51(±0.84)	0.64(±0.14)	0.82(±0.05)	0.10(±0.02)	0.16(±0.11)
MP	RF09	27 Feb 2020	18:46:24	3.65(±0.37)	1.56(±0.10)	1.16(±0.07)	0.17(±0.04)	0.35(±0.12)
MP	RF09	27 Feb 2020	18:53:48	3.29(±0.76)	1.41(±0.12)	1.07(±0.02)	0.15(±0.05)	0.32(±0.17)
MP	RF09	27 Feb 2020	19:26:17	2.51(±0.00)	1.44(±0.10)	1.05(±0.07)	0.14(±0.05)	0.30(±0.20)
MP	RF09	27 Feb 2020	19:36:09	2.42(±0.14)	1.40(±0.17)	0.71(±0.06)	0.10(±0.03)	0.35(±0.10)
MP	RF09	27 Feb 2020	20:08:09	2.03(±0.00)	1.70(±0.07)	1.04(±0.05)	0.13(±0.04)	0.35(±0.15)
MP	RF09	27 Feb 2020	20:16:39	2.80(±0.00)	1.54(±0.09)	1.01(±0.04)	0.10(±0.02)	0.48(±0.15)
MP	RF13	01 Mar 2020	16:00:14	1.48(±0.12)	0.81(±0.12)	0.73(±0.04)	0.60(±0.05)	0.58(±0.12)
MP	RF21	12 Mar 2020	14:41:07	1.94(±0.00)	2.09(±0.21)	0.81(±0.04)	0.14(±0.05)	0.37(±0.12)
MP	RF21	12 Mar 2020	14:48:49	2.39(±0.00)	1.79(±0.23)	0.87(±0.05)	0.16(±0.04)	0.36(±0.13)
MP	RF21	12 Mar 2020	15:17:07	1.90(±0.00)	1.66(±0.26)	0.68(±0.04)	0.13(±0.03)	0.25(±0.10)
MP	RF21	12 Mar 2020	16:03:43	1.00(±0.03)	1.75(±0.11)	0.78(±0.05)	0.12(±0.04)	0.31(±0.06)
MP	RF21	12 Mar 2020	16:11:45	0.99(±0.04)	1.73(±0.17)	0.84(±0.04)	0.10(±0.02)	0.40(±0.05)
MP	Average			2.72	1.33	0.91	0.39	0.48



Group	Flight	date	t _{initial} [UTC]	SeaSalt [μgm^{-3}]	OA [μgm^{-3}]	SO ₄ ²⁻ [μgm^{-3}]	NO ₃ ⁻ [μgm^{-3}]	NH ₄ ⁺ [μgm^{-3}]
HP	RF03	17 Feb 2020	17:39:28	1.76(±0.59)	2.94(±0.18)	1.04(±0.10)	0.49(±0.04)	0.66(±0.18)
HP	RF13	01 Mar 2020	14:08:38	1.27(±0.00)	0.99(±0.21)	0.66(±0.09)	0.96(±0.19)	0.62(±0.17)
HP	RF16	06 Mar 2020	19:32:13	3.65(±0.00)	0.93(±0.16)	0.64(±0.06)	0.05(±0.03)	0.07(±0.06)
HP	RF16	06 Mar 2020	19:40:24	6.72(±0.00)	2.19(±0.18)	0.91(±0.04)	0.23(±0.04)	0.45(±0.08)
HP	RF16	06 Mar 2020	20:13:19	4.95(±1.45)	1.93(±0.16)	0.96(±0.07)	0.13(±0.04)	0.44(±0.13)
HP	RF16	06 Mar 2020	20:22:54	7.26(±0.00)	0.81(±0.10)	0.56(±0.05)	0.03(±0.04)	0.20(±0.08)
HP	RF20	11 Mar 2020	13:44:36	3.25(±0.25)	1.25(±0.12)	0.29(±0.02)	0.18(±0.03)	0.07(±0.11)
HP	RF20	11 Mar 2020	14:23:48	3.70(±0.00)	1.70(±0.15)	0.38(±0.03)	0.18(±0.05)	0.08(±0.08)
HP	Average			4.07	1.59	0.68	0.28	0.32

300

305

310

315

320

Since the groups are categorized by their mean $N_{CCN_{0.43\%}}$ and the group's N_{gt85} are constantly higher than their $N_{CCN_{0.43\%}}$, the activation radii of the size distribution at 0.43% supersaturation is probably between 85 – 93 nm for the MP and HP group and around 106 nm for the LP group. The winter groups differ for particles smaller than 40 nm, which contributes a high fraction to the available aerosol population for the LP and MP group. We consider particles smaller than 40 nm as irrelevant for the cloud formation process itself, but as a critical reservoir for the accumulation mode through altering processes, which can be seen in the HP group's distribution. The MP group with its high fraction of particles below 20 nm could hint to the process of new particle formation (Zheng et al., 2021). However, the aerosol size distributions display that for a critical activation radii down to 40 nm the HP group has the highest amount of particles being possible CCN, followed by the MP group and finally the LP group.

During summertime the aerosol size distribution of the LP and HP group are comparable by adhering to a unimodal distribution, but differ significantly in the dN_{dlogD_p} concentrations between 10 nm and 400 nm (see Figure 5b). This difference is reflected in N_{gt85} in Figure 5d with mean values of 241 cm^{-3} (LP) compared to 1418 cm^{-3} (HP). The summer group's mean N_{gt85} is smaller than their mean $N_{CCN_{0.43\%}}$, suggesting a critical activation radius of the size distribution below 85 nm at 0.43% supersaturation for both groups. Here the altering processes of Aitken and accumulation modes are negligible and the difference in the HP group suggest another source of pollution during summer. The WNAO is directly located in the Northern hemisphere west wind band in winter, but during summertime the anticyclonic circulation driven by the Bermuda-Azores High influences the study region with a south west wind component (Sorooshian et al., 2020; Painemal et al., 2021; Dadashazar et al., 2021a). Therefore the sources of pollution can change between the seasons.

The wintertime aerosol mass concentrations in the BCB legs are given in Table 3. Sea salt is the dominant species with respect to mass throughout the season and has a high variability day to day and within a research flight. The highest concentrations were measured during RF16 on 6 March 2020, which can be attributed to the HP group and thus yield high $N_{CCN_{0.43\%}}$. However, there is no observable trend of sea salt mass concentration between the groups. On the other hand OA shows a significant increase from the LP to the MP/HP group. It can be deduced that the MP and HP group are influenced by pollution sources like



Table 4. All pair's aerosol composition below cloud base (BCB) during the August 2020 deployment. Mean values and standard deviation in parenthesis for organic aerosol (OA), SO_4^{2-} , NO_3^- , NH_4^+ from AMS measurements and sea salt from PILS measurements. All measurements are given at standard temperature and pressure.

Group	Flight	date	t_{initial} [UTC]	SeaSalt [μgm^{-3}]	OA [μgm^{-3}]	SO_4^{2-} [μgm^{-3}]	NO_3^- [μgm^{-3}]	NH_4^+ [μgm^{-3}]
LP	RF23	13 Aug 2020	14:43:44	-	0.25(± 1.00)	1.32(± 0.11)	0.04(± 0.11)	0.26(± 0.21)
LP	RF23	13 Aug 2020	16:56:11	-	0.78(± 0.62)	1.19(± 0.06)	0.07(± 0.04)	-0.13(± 0.63)
LP	RF24	17 Aug 2020	14:51:54	1.62(± 0.00)	2.18(± 0.25)	1.31(± 0.14)	0.16(± 0.06)	0.48(± 0.37)
LP	RF24	17 Aug 2020	15:00:00	4.32(± 0.00)	1.81(± 0.37)	1.39(± 0.14)	0.14(± 0.09)	-0.12(± 0.52)
LP	RF24	17 Aug 2020	16:55:16	0.24(± 0.00)	1.22(± 0.21)	0.67(± 0.08)	0.08(± 0.04)	-0.27(± 0.28)
LP	Average			2.06	1.25	1.17	0.10	0.04
HP	RF25	20 Aug 2020	14:40:06	0.75(± 0.20)	10.38(± 0.28)	3.24(± 0.15)	0.64(± 0.12)	1.29(± 0.43)
HP	RF25	20 Aug 2020	14:47:13	0.44(± 0.24)	9.88(± 0.48)	3.50(± 0.13)	0.40(± 0.07)	1.31(± 0.33)
HP	RF25	20 Aug 2020	15:12:04	0.60(± 0.00)	7.40(± 0.29)	2.53(± 0.08)	0.26(± 0.08)	0.78(± 0.31)
HP	RF25	20 Aug 2020	15:53:07	1.71(± 0.00)	6.18(± 0.52)	2.28(± 0.10)	0.23(± 0.10)	1.08(± 0.22)
HP	RF25	20 Aug 2020	16:01:02	6.25(± 0.00)	7.10(± 0.43)	2.55(± 0.07)	0.24(± 0.05)	1.05(± 0.21)
HP	RF26	21 Aug 2020	15:32:54	0.41(± 0.00)	5.33(± 0.55)	2.45(± 0.08)	0.22(± 0.07)	1.05(± 0.33)
HP	RF27	25 Aug 2020	15:07:52	3.53(± 0.00)	8.91(± 0.55)	1.84(± 0.08)	0.47(± 0.07)	0.77(± 0.48)
HP	RF27	25 Aug 2020	15:16:54	4.25(± 0.00)	9.03(± 0.32)	2.10(± 0.11)	0.44(± 0.09)	0.40(± 0.31)
HP	RF27	25 Aug 2020	16:16:47	4.04(± 0.88)	9.66(± 0.41)	1.66(± 0.08)	0.48(± 0.05)	0.40(± 0.32)
HP	RF27	25 Aug 2020	16:24:52	3.21(± 0.06)	9.67(± 0.58)	1.71(± 0.09)	0.47(± 0.11)	0.72(± 0.21)
HP	Average			2.52	8.35	2.39	0.39	0.89

the North East Coast American outflow, while the LP group represents natural marine conditions. The SO_4^{2-} , NO_3^- and NH_4^+ mass concentrations have a slight increase from the clean marine condition (LP) to polluted conditions (MP/HP) and i.e. RF01 on 14 February 2020 is an outlier and has the highest values, which decreases farther offshore during the flight.

In Table 4 is the BCB aerosol mass concentration below cloud depicted for the August 2020 summertime period. The sea salt mass concentration is highly variable like wintertime with low statistics in the LP group. Negative values for NH_4^+ mean that the mass concentration is lower than the calibrated background concentration and thus real. A significant increase from the LP to the HP group is measured for all species expect sea salt and suggest more pollution in the summer season. The difference in mass concentration is not equally distributed with the smallest rate of a doubling for SO_4^{2-} , followed by a factor of 4 for NO_3^- and a factor of over 6(20) for OA(NH_4^+). The chemical composition of the aerosol population alters N_C (Hoose and Möhler, 2012), i.e. the organic carbon species have variable influences depending on solubility, molecular weight and surface tension (Ervens et al., 2005).




Figure 6. a) The lines represent the w to N_C relation derived with the PMM for the low polluted (LP/blue), medium polluted (MP/orange) and high polluted (HP/red) group with their boundaries of mean $N_{CCN_{0.43\%}}$ in parenthesis. The dark shaded areas represent the measurement uncertainty of 20% in addition with the relative error calculated according to Haddad and Rosenfeld (1997) with the assumption that the standard deviation of N_C in each group represents the ratio of the noisy variation in the N_C measurements to the true variation in N_C . c) The S_{max} estimate of each group is given in 'x' markers for the same w spectrum with the error as shallow lined shaded area. The vertical lines are the w_{eff} with associated S_{max} . The same PMM and S_{max} analysis in b) and d) for the summertime LP and HP group.

335 3.3 Seasonal impact of w and $N_{CCN_{0.43\%}}$ on N_C

Figure 6a shows the application of the PMM to all groups of the winter season. The w to N_C relations shows the fraction of activated aerosol from the aerosol size distribution for a given updraft of supersaturation, respectively. The LP group, which has a mean N_C of $315(\pm 165) \text{ cm}^{-3}$, shows the highest impact of w to N_C for $w < 1.4 \text{ ms}^{-1}$ and reaches saturation for higher w values. The MP group exhibits a similar trend with a mean of $518(\pm 304) \text{ cm}^{-3}$, but the impact of w is decreasing slower compared to the LP group for higher w . The HP group shows the strongest impact for $w < 1.6 \text{ ms}^{-1}$ and as a mean N_C of $930(\pm 630) \text{ cm}^{-3}$. In addition, it has a second mode with a strong increase in N_C for $w > 3 \text{ ms}^{-1}$.

The two domains of w in the HP group could represent the activation of smaller aerosol particles from the aerosol population. Since the critical diameter of aerosol activation depends on the supersaturation and is shifted towards smaller diameters for higher supersaturation, the positive correlation of w and supersaturation results in smaller aerosols getting activated for higher w (Köhler, 1936; Dusek et al., 2006; Schulze et al., 2020). N_C are slightly smaller than the respective group's $N_{CCN_{0.43\%}}$



leading to a mean supersaturation below 0.43% in winter. The LP group exhibits some characteristics of an aerosol-limited regime with N_C highly depending on the available aerosol population, while the HP group shows the characteristics of an updraft-limited regime with N_C being directly proportional to w (Reutter et al., 2009). The MP group is between both regimes and tend to the characteristics of an updraft-limited regime, since N_C does not reach saturation for high w .

350 The S_{\max} estimate for each group's w_{eff} in Figure 6c is decreasing with increasing pollution level and are 0.33%(LP), 0.26%(MP) and 0.18%(HP), respectively. Since the variability of updraft speed is higher with larger w , the local supersaturation can deviate from the derived S_{\max} estimates. The reduction of S_{\max} for increasing pollution levels demonstrate the water vapor competition of more activated CCN and thus function as a buffer for preventing higher supersaturation. The LP group's mean N_C is above its mean $N_{\text{CCN}_{0.43\%}}$ although S_{\max} is near, and below 0.43%, which could be explained by a contribution of the
355 soluble Aitken mode particles in the bimodal aerosol size distribution (Pöhlker et al., 2021). However, the winter groups exhibit mean N_C near $N_{\text{CCN}_{0.43\%}}$ with a trend of a reduced fraction of activated aerosol with increasing pollution level.

In Figure 6b the PMM is applied to the summer season in the same way. The impact of w on N_C has a similar trend in summer and winter for the LP group up to the maximal measured w of 1.3 m s^{-1} during summer and has a mean N_C of $196(\pm 55) \text{ cm}^{-3}$. The HP group has a nearly constant impact for the full range of w up to 2.1 m s^{-1} and a mean N_C of
360 $642(\pm 389) \text{ cm}^{-3}$. The w to N_C relation coincides with the wintertime equivalent for w below 1.7 m s^{-1} . The S_{\max} estimate for each group's w_{eff} in Figure 6d is analogously reduced from the LP to the HP group in summer as in winter, while between the seasons a halving of the S_{\max} takes place.

N_{gt85} of the summer LP group is significantly lower than its winter counterpart, thus less aerosol for cloud formation is available in clean conditions during summer compared to winter. On the other hand N_{gt85} of the HP summer group is
365 substantially higher than during winter. Another key feature is the lower mean N_{gt85} in comparison to the mean $N_{\text{CCN}_{0.43\%}}$, showing a higher fraction of activated CCN in summer for a given supersaturation of 0.43%, which hints to a lower mean critical supersaturation needed for activation of the summer aerosol composition. Table 3 and Table 4 show an increased mass concentration of OA and SO_4^{2-} between the respective groups. The high hygroscopicity of SO_4^{2-} is most likely accountable for the observed lower mean N_{gt85} than mean $N_{\text{CCN}_{0.43\%}}$, because the raised OA mass concentrations from the LP to HP group is
370 not reflected. Lower supersaturation in summer due to the smaller updrafts results in less activated CCN. The bisection of w_{eff} in Figure 6b propagates through derived S_{\max} estimates to N_C .

4 Conclusions

In this study we examine the seasonal impact of w and $N_{\text{CCN}_{0.43\%}}$ on N_C over the WNAO from an in-situ perspective during the ACTIVATE campaign. The impact is determined by a statistical approach with the PMM where pairs of flight legs below
375 and in cloud base are used to categorize in-situ measurements into similar environmental conditions and $N_{\text{CCN}_{0.43\%}}$. We also give detailed information on the aerosol size distribution and composition below cloud base. Key findings are summarized and related to 2020 winter (February-March) and summer (August) conditions as follows:



- 380
- N_C in low clouds over the WNAO show a positive correlation with w and $N_{CCN_{0.43\%}}$. Updrafts smaller than 1.3 m s^{-1} have the highest impact on N_C in both seasons. Polluted environments exhibit a stronger w impact over the full w distribution in a season, while in clean marine environments the available N_{CCN} limit N_C for higher w .
 - The WNAO exhibits an anti-correlated seasonal cycle of N_C and $N_{CCN_{0.43\%}}$ at cloud base with 25% less N_C and 71% more $N_{CCN_{0.43\%}}$ in their overall observed mean values in summer compared to winter. The seasonal cycle is consistent with the anti-correlated AOD and N_C cycle measured by remote sensing and satellite instruments (Dadashazar et al., 2021b).
 - 385 – The mean values of w at cloud bases are 33% lower in summer compared to winter. Simultaneously the variability of updraft speeds is reduced by 31% in summer. Both indicate a higher dynamical influence during winter. A correlation of N_C and w is observed in the seasonal cycle and suggest that the difference between the seasons is driven by dynamics.
 - The winter $N_{CCN_{0.43\%}}$ directly below cloud shows a broad distribution due to different aerosol sources and pollution levels, while only clear sky or high polluted conditions were measured in summer. For high polluted environments,
390 summer exhibits a 46% increased mean $N_{CCN_{0.43\%}}$.
 - The aerosol size distribution during winter exhibits a bimodal distribution in clean marine and medium polluted condition, which transforms into a unimodal distribution for higher pollution levels. The Aitken mode acts as reservoir for the accumulation mode, since N_{gt85} increases while the aerosol number concentrations do not differ significantly. In contrast to the winter period, the summer period is characterized by unimodal distributions and a clear difference between the
395 aerosol concentrations of the pollution levels.
 - The aerosol composition shows a constant proportion of sea salt in each season, with an increased aerosol mass concentration measured in winter, which could be related to the increased surface wind speeds resulting in more efficient wind-driven sea salt emissions (Painemal et al., 2021). With the increase in pollution levels, a concomitant increase in OA , SO_4^{2-} , NO_3^- and NH_4^+ mass concentrations is measured in summer. In winter, the increase is comparatively
400 moderate.
 - w and related S_{max} determine the range of activated CCN and S_{max} is reduced at increasing pollution levels. As shown, w dominantly affects the activation of CCN and determines the fraction of activated aerosol and thus explains generally higher N_C values during winter compared to summer.

405 The observational data presented in this study includes key parameters which are used in state-of-the-art aerosol-climate models to describe aerosol-induced cloud modifications. Consistent observations of the aerosol number concentration, size distribution and composition, w as well as N_C are provided for a wide range of conditions in the winter and summer seasons. Hence the data could serve as a valuable basis for evaluating and further improving the representation of aerosol-cloud interactions in future climate simulations.



Data availability. The ACTIVATE data are available at <http://doi.org/10.5067/SUBORBITAL/ACTIVATE/DATA001>

410 Appendix A

A1 List of symbols and abbreviations

D_{\max}	maximum distance of cloud measurements to aerosol measurements
h_{ACB}	height above cloud base
N_{C}	cloud droplet number concentration
$N_{\text{CCN}_{0.43\%}}$	cloud condensation nuclei concentration at 0.43% supersaturation
S_{\max}	maximum supersaturation in cloud base
w	updraft speed
w_{eff}	effective updraft speed
2D-S	Two-Dimensional Stereo probe
ACB	above cloud base
ACTIVATE	Aerosol Cloud meTerology Interactions oVer the western ATlantic Experiment
AMS	Aerosol Mass Spectrometer
AOD	aerosol optical depth
BCB	below cloud base
CCN	cloud condensation nuclei
CMIP6	Coupled Model Intercomparison Project Phase 6
DoF	depth of field
FCDP	Fast Cloud Droplet Probe
LAS	Laser Aerosol Spectrometer
MBL	marine boundary layer
OA	organic aerosol
PAS	particle air speed
PILS	Particle-Into-Liquid Sampler
PMM	Probability Matching Method
RF	research flights
SMPS	Scanning Mobility Particle Sizer
SPEC Inc.	Stratton Park Engineering Company Incorporated
TAMMS	turbulent air motion measurement system
WNAO	Western North Atlantic Ocean



Author contributions. S.K. conducted the analysis and wrote the manuscript. C.V. advised the study and provided intensive feedback on the manuscript. K.T. wrote section 2.2.3. A.S. and L.Z. wrote section 2.2.5. L.Z. wrote section 2.2.6. J.H. contributed to section 1 and 4. E.C, 415 E.W and L.Z. performed the flight measurements. S.K., V.H., St.K., C.R. and D.S. participated in instrument calibration. B.A., R.M., L.Z. and A.S. participated in mission planning. R.F., A.J.S. and M.S. conducted the weather forecast. E.C, R.F., A.J.S., R.M., M.S. and A.S. participated in strategic flight planning. S.K., R.M, K.T., C.R, E.W., L.Z., M.S. and A.S. participated in mission operation. G.C. and M.S. conducted the data management. All authors commented on the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

420 *Acknowledgements.* The work was funded by NASA grant 80NSSC19K0442 in support of ACTIVATE, a NASA Earth Venture Suborbital-3 (EVS-3) investigation funded by NASA's Earth Science Division and managed through the Earth System Science Pathfinder Program Office. C.V, S.K. and St.K. were funded by the Helmholtz excellence programme (grant number W2/W3-060, by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – TRR 301 – Project-ID 428312742 and the SPP 1294 HALO under contract VO 1504/7-1.



References

- 425 Abdul-Razzak, H. and Ghan, S. J.: A Parameterization of Aerosol Activation: 2. Multiple Aerosol Types, *Journal of Geophysical Research: Atmospheres*, 105, 6837–6844, <https://doi.org/10.1029/1999JD901161>, 2000.
- Albrecht, B. A.: Aerosols, Cloud Microphysics, and Fractional Cloudiness, *Science*, 245, 1227–1230, <https://doi.org/10.1126/science.245.4923.1227>, 1989.
- Andrews, T., Gregory, J. M., and Webb, M. J.: The Dependence of Radiative Forcing and Feedback on Evolving Patterns of Surface Temperature Change in Climate Models, *Journal of Climate*, 28, 1630–1648, <https://doi.org/10.1175/JCLI-D-14-00545.1>, 2015.
- 430 Baker, B. and Lawson, R. P.: Improvement in Determination of Ice Water Content from Two-Dimensional Particle Imagery. Part I: Image-to-Mass Relationships, *Journal of Applied Meteorology and Climatology*, 45, 1282–1290, <https://doi.org/10.1175/JAM2398.1>, 2006.
- Baumgardner, D., Strapp, W., and Dye, J. E.: Evaluation of the Forward Scattering Spectrometer Probe. Part II: Corrections for Coincidence and Dead-Time Losses, *Journal of Atmospheric and Oceanic Technology*, 2, 626–632, [https://doi.org/10.1175/1520-0426\(1985\)002<0626:EOTFSS>2.0.CO;2](https://doi.org/10.1175/1520-0426(1985)002<0626:EOTFSS>2.0.CO;2), 1985.
- 435 Baumgardner, D., Abel, S. J., Axisa, D., Cotton, R., Crosier, J., Field, P., Gurganus, C., Heymsfield, A., Korolev, A., and Kraemer, M.: Cloud Ice Properties: In Situ Measurement Challenges, *Meteorological monographs*, 58, 9–1, 2017.
- Bock, L., Lauer, A., Schlund, M., Barreiro, M., Bellouin, N., Jones, C., Meehl, G. A., Predoi, V., Roberts, M. J., and Eyring, V.: Quantifying Progress Across Different CMIP Phases With the ESMValTool, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD032321, <https://doi.org/10.1029/2019JD032321>, 2020.
- 440 Bowen, H. J. M.: *Environmental Chemistry of the Elements*, Academic press, 1979.
- Braga, R. C., Rosenfeld, D., Weigel, R., Jurkat, T., Andreae, M. O., Wendisch, M., Pöhlker, M. L., Klimach, T., Pöschl, U., Pöhlker, C., Voigt, C., Mahnke, C., Borrmann, S., Albrecht, R. I., Molleker, S., Vila, D. A., Machado, L. A. T., and Artaxo, P.: Comparing Parameterized versus Measured Microphysical Properties of Tropical Convective Cloud Bases during the ACRIDICON–CHUVA Campaign, *Atmospheric Chemistry and Physics*, 17, 7365–7386, <https://doi.org/10.5194/acp-17-7365-2017>, 2017a.
- 445 Braga, R. C., Rosenfeld, D., Weigel, R., Jurkat, T., Andreae, M. O., Wendisch, M., Pöschl, U., Voigt, C., Mahnke, C., Borrmann, S., Albrecht, R. I., Molleker, S., Vila, D. A., Machado, L. A. T., and Grulich, L.: Further Evidence for CCN Aerosol Concentrations Determining the Height of Warm Rain and Ice Initiation in Convective Clouds over the Amazon Basin, *Atmospheric Chemistry and Physics*, 17, 14433–14456, <https://doi.org/10.5194/acp-17-14433-2017>, 2017b.
- 450 Braga, R. C., Ervens, B., Rosenfeld, D., Andreae, M. O., Förster, J.-D., Fütterer, D., Hernández Pardo, L., Holanda, B. A., Jurkat-Witschas, T., Krüger, O. O., Lauer, O., Machado, L. A. T., Pöhlker, C., Sauer, D., Voigt, C., Walser, A., Wendisch, M., Pöschl, U., and Pöhlker, M. L.: Cloud Droplet Formation at the Base of Tropical Convective Clouds: Closure between Modeling and Measurement Results of ACRIDICON–CHUVA, *Atmospheric Chemistry and Physics*, 21, 17513–17528, <https://doi.org/10.5194/acp-21-17513-2021>, 2021.
- Bräuer, T., Voigt, C., Sauer, D., Kaufmann, S., Hahn, V., Scheibe, M., Schlager, H., Diskin, G. S., Nowak, J. B., DiGangi, J. P., Huber, F., Moore, R. H., and Anderson, B. E.: Airborne Measurements of Contrail Ice Properties—Dependence on Temperature and Humidity, *Geophysical Research Letters*, 48, e2020GL092166, <https://doi.org/10.1029/2020GL092166>, 2021a.
- 455 Bräuer, T., Voigt, C., Sauer, D., Kaufmann, S., Hahn, V., Scheibe, M., Schlager, H., Huber, F., Le Clercq, P., Moore, R., and Anderson, B.: Reduced Ice Number Concentrations in Contrails from Low Aromatic Biofuel Blends, *Atmospheric Chemistry and Physics Discussions*, pp. 1–14, <https://doi.org/10.5194/acp-2021-582>, 2021b.



- 460 Calheiros, R. V. and Zawadzki, I.: Reflectivity-Rain Rate Relationships for Radar Hydrology in Brazil, *Journal of Applied Meteorology and Climatology*, 26, 118–132, [https://doi.org/10.1175/1520-0450\(1987\)026<0118:RRRRFR>2.0.CO;2](https://doi.org/10.1175/1520-0450(1987)026<0118:RRRRFR>2.0.CO;2), 1987.
- Cecchini, M. A., Machado, L. A. T., Andreae, M. O., Martin, S. T., Albrecht, R. I., Artaxo, P., Barbosa, H. M. J., Borrmann, S., Fütterer, D., Jurkat, T., Mahnke, C., Minikin, A., Molleker, S., Pöhlker, M. L., Pöschl, U., Rosenfeld, D., Voigt, C., Weinzierl, B., and Wendisch, M.: Sensitivities of Amazonian Clouds to Aerosols and Updraft Speed, *Atmospheric Chemistry and Physics*, 17, 10037–
465 10050, <https://doi.org/10.5194/acp-17-10037-2017>, 2017.
- Ceppi, P., Brient, F., Zelinka, M. D., and Hartmann, D. L.: Cloud Feedback Mechanisms and Their Representation in Global Climate Models, *WIREs Climate Change*, 8, e465, <https://doi.org/10.1002/wcc.465>, 2017.
- Dadashazar, H., Alipanah, M., Hilario, M. R. A., Crosbie, E., Kirschler, S., Liu, H., Moore, R. H., Peters, A. J., Scarino, A. J., Shook, M., Thornhill, K. L., Voigt, C., Wang, H., Winstead, E., Zhang, B., Ziemba, L., and Sorooshian, A.: Aerosol Responses to
470 Precipitation along North American Air Trajectories Arriving at Bermuda, *Atmospheric Chemistry and Physics*, 21, 16 121–16 141, <https://doi.org/10.5194/acp-21-16121-2021>, 2021a.
- Dadashazar, H., Painemal, D., Alipanah, M., Brunke, M., Chellappan, S., Corral, A. F., Crosbie, E., Kirschler, S., Liu, H., Moore, R., Robinson, C., Scarino, A. J., Shook, M., Sinclair, K., Thornhill, K. L., Voigt, C., Wang, H., Winstead, E., Zeng, X., Ziemba, L., Zuidema, P., and Sorooshian, A.: Cloud Drop Number Concentrations over the Western North Atlantic Ocean: Seasonal Cycle, Aerosol Interrelationships,
475 and Other Influential Factors, *Atmospheric Chemistry and Physics Discussions*, pp. 1–36, <https://doi.org/10.5194/acp-2021-153>, 2021b.
- DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, M. J., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K., Horvath, T., Docherty, K. S., Worsnop, D. R., and Jimenez, J. L.: Field-Deployable, High-Resolution, Time-of-Flight Aerosol Mass Spectrometer, *Analytical Chemistry*, 78, 8281–8289, <https://doi.org/10.1021/ac061249n>, 2006.
- Dusek, U., Frank, G. P., Hildebrandt, L., Curtius, J., Schneider, J., Walter, S., Chand, D., Drewnick, F., Hings, S., Jung, D., Borrmann, S.,
480 and Andreae, M. O.: Size Matters More Than Chemistry for Cloud-Nucleating Ability of Aerosol Particles, *Science*, 312, 1375–1378, <https://doi.org/10.1126/science.1125261>, 2006.
- Ervens, B., Feingold, G., and Kreidenweis, S. M.: Influence of Water-Soluble Organic Carbon on Cloud Drop Number Concentration, *Journal of Geophysical Research: Atmospheres*, 110, <https://doi.org/10.1029/2004JD005634>, 2005.
- Faber, S., French, J. R., and Jackson, R.: Laboratory and In-Flight Evaluation of Measurement Uncertainties from a Commercial Cloud
485 Droplet Probe (CDP), *Atmospheric Measurement Techniques*, 11, 3645–3659, <https://doi.org/10.5194/amt-11-3645-2018>, 2018.
- Farren, N. J., Dunmore, R. E., Mead, M. I., Mohd Nadzir, M. S., Samah, A. A., Phang, S.-M., Bandy, B. J., Sturges, W. T., and Hamilton, J. F.: Chemical Characterisation of Water-Soluble Ions in Atmospheric Particulate Matter on the East Coast of Peninsular Malaysia, *Atmospheric Chemistry and Physics*, 19, 1537–1553, <https://doi.org/10.5194/acp-19-1537-2019>, 2019.
- Flamant, C., Knippertz, P., Fink, A. H., Akpo, A., Brooks, B., Chiu, C. J., Coe, H., Danuor, S., Evans, M., Jegede, O., Kalthoff, N., Konaré,
490 A., Lioussé, C., Lohou, F., Mari, C., Schlager, H., Schwarzenboeck, A., Adler, B., Amekudzi, L., Aryee, J., Ayoola, M., Batenburg, A. M., Bessardon, G., Borrmann, S., Brito, J., Bower, K., Burnet, F., Catoire, V., Colomb, A., Denjean, C., Fosu-Amankwah, K., Hill, P. G., Lee, J., Lathon, M., Maranan, M., Marsham, J., Meynadier, R., Ngamini, J.-B., Rosenberg, P., Sauer, D., Smith, V., Stratmann, G., Taylor, J. W., Voigt, C., and Yoboué, V.: The Dynamics–Aerosol–Chemistry–Cloud Interactions in West Africa Field Campaign: Overview and Research Highlights, *Bulletin of the American Meteorological Society*, 99, 83–104, <https://doi.org/10.1175/BAMS-D-16-0256.1>, 2018.
- 495 Formenti, P., D’Anna, B., Flamant, C., Mallet, M., Piketh, S. J., Schepanski, K., Waquet, F., Auriol, F., Brogniez, G., Burnet, F., Chaboureaud, J.-P., Chauvigné, A., Chazette, P., Denjean, C., Desboeufs, K., Doussin, J.-F., Elguindi, N., Feuerstein, S., Gaetani, M., Giorio, C., Klopper, D., Mallet, M. D., Nabat, P., Monod, A., Solmon, F., Namwoonde, A., Chikwililwa, C., Mushi, R., Welton, E. J., and Holben, B.: The



- Aerosols, Radiation and Clouds in Southern Africa Field Campaign in Namibia: Overview, Illustrative Observations, and Way Forward, *Bulletin of the American Meteorological Society*, 100, 1277–1298, <https://doi.org/10.1175/BAMS-D-17-0278.1>, 2019.
- 500 Freud, E. and Rosenfeld, D.: Linear Relation between Convective Cloud Drop Number Concentration and Depth for Rain Initiation, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/10.1029/2011JD016457>, 2012.
- Ghan, S. J., Abdul-Razzak, H., Nenes, A., Ming, Y., Liu, X., Ovchinnikov, M., Shipway, B., Meskhidze, N., Xu, J., and Shi, X.: Droplet Nucleation: Physically-based Parameterizations and Comparative Evaluation, *Journal of Advances in Modeling Earth Systems*, 3, <https://doi.org/10.1029/2011MS000074>, 2011.
- 505 Grosvenor, D. P., Sourdeval, O., Zuidema, P., Ackerman, A., Alexandrov, M. D., Bennartz, R., Boers, R., Cairns, B., Chiu, J. C., and Christensen, M.: Remote Sensing of Droplet Number Concentration in Warm Clouds: A Review of the Current State of Knowledge and Perspectives, *Reviews of Geophysics*, 56, 409–453, 2018.
- Gryspeerdt, E., Quaas, J., and Bellouin, N.: Constraining the Aerosol Influence on Cloud Fraction, *Journal of Geophysical Research: Atmospheres*, 121, 3566–3583, <https://doi.org/10.1002/2015JD023744>, 2016.
- 510 Gryspeerdt, E., Quaas, J., Ferrachat, S., Gettelman, A., Ghan, S., Lohmann, U., Morrison, H., Neubauer, D., Partridge, D. G., and Stier, P.: Constraining the Instantaneous Aerosol Influence on Cloud Albedo, *Proceedings of the National Academy of Sciences*, 114, 4899–4904, 2017.
- Gurganus, C. and Lawson, P.: Laboratory and Flight Tests of 2D Imaging Probes: Toward a Better Understanding of Instrument Performance and the Impact on Archived Data, *Journal of Atmospheric and Oceanic Technology*, 35, 1533–1553, <https://doi.org/10.1175/JTECH-D-17-0202.1>, 2018.
- 515 Haddad, Z. S. and Rosenfeld, D.: Optimality of Empirical Z-R Relations, *Quarterly Journal of the Royal Meteorological Society*, 123, 1283–1293, <https://doi.org/10.1002/qj.49712354107>, 1997.
- Hersey, S. P., Sorooshian, A., Murphy, S. M., Flagan, R. C., and Seinfeld, J. H.: Aerosol Hygroscopicity in the Marine Atmosphere: A Closure Study Using High-Time-Resolution, Multiple-RH DASH-SP and Size-Resolved C-ToF-AMS Data, *Atmospheric Chemistry and Physics*, 9, 2543–2554, <https://doi.org/10.5194/acp-9-2543-2009>, 2009.
- 520 Hilario, M. R. A., Crosbie, E., Shook, M., Reid, J. S., Cambaliza, M. O. L., Simpas, J. B. B., Ziemba, L., DiGangi, J. P., Diskin, G. S., Nguyen, P., Turk, F. J., Winstead, E., Robinson, C. E., Wang, J., Zhang, J., Wang, Y., Yoon, S., Flynn, J., Alvarez, S. L., Behrangi, A., and Sorooshian, A.: Measurement Report: Long-range Transport Patterns into the Tropical Northwest Pacific during the CAMP²Ex Aircraft Campaign: Chemical Composition, Size Distributions, and the Impact of Convection, *Atmospheric Chemistry and Physics*, 21, 3777–3802, <https://doi.org/10.5194/acp-21-3777-2021>, 2021.
- 525 Hoose, C. and Möhler, O.: Heterogeneous Ice Nucleation on Atmospheric Aerosols: A Review of Results from Laboratory Experiments, *Atmospheric Chemistry and Physics*, 12, 9817–9854, <https://doi.org/10.5194/acp-12-9817-2012>, 2012.
- IPCC: Climate Change 2021 - the Physical Science Basis, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, v., p. Zhai, a. Pirani, s. L. Connors, c. Péan, s. Berger, n. Caud, y. Chen, l. Goldfarb, m. I. Gomis, m. Huang, k. Leitzell, e. Lonnoy, j. B. R. Matthews, t. K. Maycock, t. Waterfield, o. Yelekçi, r. Yu and b. Zhou (Eds.)], p. 1535, Cambridge University Press, cambridge edn., 2021.
- 530 Kleine, J., Voigt, C., Sauer, D., Schlager, H., Scheibe, M., Jurkat-Witschas, T., Kaufmann, S., Kärcher, B., and Anderson, B. E.: In Situ Observations of Ice Particle Losses in a Young Persistent Contrail, *Geophysical Research Letters*, 45, 13,553–13,561, <https://doi.org/10.1029/2018GL079390>, 2018.



- 535 Knippertz, P., Coe, H., Chiu, J. C., Evans, M. J., Fink, A. H., Kalthoff, N., Liousse, C., Mari, C., Allan, R. P., Brooks, B., Danour, S., Flamant, C., Jegede, O. O., Lohou, F., and Marsham, J. H.: The DACCIWA Project: Dynamics–Aerosol–Chemistry–Cloud Interactions in West Africa, *Bulletin of the American Meteorological Society*, 96, 1451–1460, <https://doi.org/10.1175/BAMS-D-14-00108.1>, 2015.
- Knollenberg, R. G.: The Optical Array: An Alternative to Extinction and Scattering for Particle Size Measurements, *J. Appl. Meteor*, 9, 86–103, 1970.
- 540 Knop, I., Bansmer, S. E., Hahn, V., and Voigt, C.: Comparison of Different Droplet Measurement Techniques in the Braunschweig Icing Wind Tunnel, *Atmospheric Measurement Techniques*, 14, 1761–1781, <https://doi.org/10.5194/amt-14-1761-2021>, 2021.
- Koehler, K. A., Kreidenweis, S. M., DeMott, P. J., Prenni, A. J., Carrico, C. M., Ervens, B., and Feingold, G.: Water Activity and Activation Diameters from Hygroscopicity Data - Part II: Application to Organic Species, *Atmospheric Chemistry and Physics*, 6, 795–809, <https://doi.org/10.5194/acp-6-795-2006>, 2006.
- 545 Köhler, H.: The Nucleus in and the Growth of Hygroscopic Droplets, *Transactions of the Faraday Society*, 32, 1152–1161, 1936.
- Korolev, A.: Reconstruction of the Sizes of Spherical Particles from Their Shadow Images. Part I: Theoretical Considerations, *Journal of Atmospheric and Oceanic Technology*, 24, 376–389, <https://doi.org/10.1175/JTECH1980.1>, 2007.
- Korolev, A. V., Kuznetsov, S. V., Makarov, Y. E., and Novikov, V. S.: Evaluation of Measurements of Particle Size and Sample Area from Optical Array Probes, *Journal of Atmospheric and Oceanic Technology*, 8, 514–522, 1991.
- 550 Korolev, A. V., Strapp, J. W., and Isaac, G. A.: Evaluation of the Accuracy of PMS Optical Array Probes, *Journal of Atmospheric and Oceanic Technology*, 15, 708–720, 1998.
- Lance, S.: Coincidence Errors in a Cloud Droplet Probe (CDP) and a Cloud and Aerosol Spectrometer (CAS), and the Improved Performance of a Modified CDP, *Journal of Atmospheric and Oceanic Technology*, 29, 1532–1541, 2012.
- Lance, S., Nenes, A., Medina, J., and Smith, J. N.: Mapping the Operation of the DMT Continuous Flow CCN Counter, *Aerosol Science and Technology*, 40, 242–254, <https://doi.org/10.1080/02786820500543290>, 2006.
- 555 Lance, S., Brock, C. A., Rogers, D., and Gordon, J. A.: Water Droplet Calibration of the Cloud Droplet Probe (CDP) and in-Flight Performance in Liquid, Ice and Mixed-Phase Clouds during ARCPAC, *Atmospheric Measurement Techniques*, 3, 1683–1706, <https://doi.org/10.5194/amt-3-1683-2010>, 2010.
- Lawson, R. P.: Effects of Ice Particles Shattering on the 2D-S Probe, *Atmospheric Measurement Techniques*, 4, 1361–1381, <https://doi.org/10.5194/amt-4-1361-2011>, 2011.
- 560 Lawson, R. P. and Baker, B. A.: Improvement in Determination of Ice Water Content from Two-Dimensional Particle Imagery. Part II: Applications to Collected Data, *Journal of Applied Meteorology and Climatology*, 45, 1291–1303, <https://doi.org/10.1175/JAM2399.1>, 2006.
- Lawson, R. P., O'Connor, D., Zmarzly, P., Weaver, K., Baker, B., Mo, Q., and Jonsson, H.: The 2D-S (Stereo) Probe: Design and Preliminary Tests of a New Airborne, High-Speed, High-Resolution Particle Imaging Probe, *Journal of Atmospheric and Oceanic Technology*, 23, 1462–1477, <https://doi.org/10.1175/JTECH1927.1>, 2006.
- 565 Lawson, R. P., Woods, S., Jensen, E., Erfani, E., Gurganus, C., Gallagher, M., Connolly, P., Whiteway, J., Baran, A. J., May, P., Heymsfield, A., Schmitt, C. G., McFarquhar, G., Um, J., Protat, A., Bailey, M., Lance, S., Muehlbauer, A., Stith, J., Korolev, A., Toon, O. B., and Krämer, M.: A Review of Ice Particle Shapes in Cirrus Formed In Situ and in Anvils, *Journal of Geophysical Research: Atmospheres*, 124, 10 049–10 090, <https://doi.org/10.1029/2018JD030122>, 2019.
- 570 Lenschow, D. H.: Aircraft Measurements in the Boundary Layer, in: *Probing the Atmospheric Boundary Layer*, edited by Lenschow, D. H., pp. 39–55, American Meteorological Society, Boston, MA, https://doi.org/10.1007/978-1-944970-14-7_5, 1986.



- Lohmann, U. and Hoose, C.: Sensitivity Studies of Different Aerosol Indirect Effects in Mixed-Phase Clouds, *Atmospheric Chemistry and Physics*, 9, 8917–8934, <https://doi.org/10.5194/acp-9-8917-2009>, 2009.
- 575 Lohmann, U., Stier, P., Hoose, C., Ferrachat, S., Kloster, S., Roeckner, E., and Zhang, J.: Cloud Microphysics and Aerosol Indirect Effects in the Global Climate Model ECHAM5-HAM, *Atmospheric Chemistry and Physics*, 7, 3425–3446, <https://doi.org/10.5194/acp-7-3425-2007>, 2007.
- Lu, M.-L., Conant, W. C., Jonsson, H. H., Varutbangkul, V., Flagan, R. C., and Seinfeld, J. H.: The Marine Stratus/Stratocumulus Experiment (MASE): Aerosol-cloud Relationships in Marine Stratocumulus, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/10.1029/2006JD007985>, 2007.
- 580 Ma, L., Dadashazar, H., Hilario, M. R. A., Cambaliza, M. O., Lorenzo, G. R., Simpas, J. B., Nguyen, P., and Sorooshian, A.: Contrasting Wet Deposition Composition between Three Diverse Islands and Coastal North American Sites, *Atmospheric Environment*, 244, 117919, <https://doi.org/10.1016/j.atmosenv.2020.117919>, 2021.
- McComiskey, A., Feingold, G., and Quaas, J.: The Scale Problem in Quantifying Aerosol Indirect Effects., *Atmospheric Chemistry & Physics*, 12, <https://doi.org/10.5194/acp-12-1031-2012>, 2012.
- 585 McCoy, I. L., McCoy, D. T., Wood, R., Regayre, L., Watson-Parris, D., Grosvenor, D. P., Mulcahy, J. P., Hu, Y., Bender, F. A.-M., Field, P. R., Carslaw, K. S., and Gordon, H.: The Hemispheric Contrast in Cloud Microphysical Properties Constrains Aerosol Forcing, *Proceedings of the National Academy of Sciences*, 117, 18 998–19 006, <https://doi.org/10.1073/pnas.1922502117>, 2020.
- McCoy, I. L., Bretherton, C. S., Wood, R., Twohy, C. H., Gettelman, A., Bardeen, C. G., and Toohey, D. W.: Influences of Recent Particle Formation on Southern Ocean Aerosol Variability and Low Cloud Properties, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033 529, <https://doi.org/10.1029/2020JD033529>, 2021.
- 590 McNaughton, C. S., Clarke, A. D., Howell, S. G., Pinkerton, M., Anderson, B., Thornhill, L., Hudgins, C., Winstead, E., Dibb, J. E., Scheuer, E., and Maring, H.: Results from the DC-8 Inlet Characterization Experiment (DICE): Airborne Versus Surface Sampling of Mineral Dust and Sea Salt Aerosols, *Aerosol Science and Technology*, 41, 136–159, <https://doi.org/10.1080/02786820601118406>, 2007.
- 595 Moore, R. H. and Nenes, A.: Scanning Flow CCN Analysis—A Method for Fast Measurements of CCN Spectra, *Aerosol Science and Technology*, 43, 1192–1207, <https://doi.org/10.1080/02786820903289780>, 2009.
- Moore, R. H., Thornhill, K. L., Weinzierl, B., Sauer, D., D’Ascoli, E., Kim, J., Lichtenstern, M., Scheibe, M., Beaton, B., Beyersdorf, A. J., Barrick, J., Bulzan, D., Corr, C. A., Crosbie, E., Jurkat, T., Martin, R., Riddick, D., Shook, M., Slover, G., Voigt, C., White, R., Winstead, E., Yasky, R., Ziemba, L. D., Brown, A., Schlager, H., and Anderson, B. E.: Biofuel Blending Reduces Particle Emissions from Aircraft Engines at Cruise Conditions, *Nature*, 543, 411–415, <https://doi.org/10.1038/nature21420>, 2017.
- 600 Moore, R. H., Wiggins, E. B., Ahern, A. T., Zimmerman, S., Montgomery, L., Campuzano Jost, P., Robinson, C. E., Ziemba, L. D., Winstead, E. L., Anderson, B. E., Brock, C. A., Brown, M. D., Chen, G., Crosbie, E. C., Guo, H., Jimenez, J. L., Jordan, C. E., Lyu, M., Nault, B. A., Rothfuss, N. E., Sanchez, K. J., Schueneman, M., Shingler, T. J., Shook, M. A., Thornhill, K. L., Wagner, N. L., and Wang, J.: Sizing Response of the Ultra-High Sensitivity Aerosol Spectrometer (UHSAS) and Laser Aerosol Spectrometer (LAS) to Changes in Submicron Aerosol Composition and Refractive Index, *Atmospheric Measurement Techniques*, 14, 4517–4542, <https://doi.org/10.5194/amt-14-4517-2021>, 2021.
- 605 Myhre, G., Shindell, D., Bréon, F.-M., Collins, W., Fuglestedt, J., Huang, J., Koch, D., Lamarque, J.-F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T., and Zhang, H.: Anthropogenic and Natural Radiative Forcing, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Stocker, T., Qin, D., Plattner, G.-K., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V.,



- and Midgley, P., book Section 8, pp. 659–740, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/CBO9781107415324.018>, 2013.
- O'Connor, D., Baker, B., and Lawson, R. P.: Upgrades to the FSSP-100 Electronics, in: Proc. 15th Int. Conf. on Clouds and Precipitation, 2008.
- 615 Painemal, D., Chang, F.-L., Ferrare, R., Burton, S., Li, Z., Smith Jr., W. L., Minnis, P., Feng, Y., and Clayton, M.: Reducing Uncertainties in Satellite Estimates of Aerosol–Cloud Interactions over the Subtropical Ocean by Integrating Vertically Resolved Aerosol Observations, *Atmospheric Chemistry and Physics*, 20, 7167–7177, <https://doi.org/10.5194/acp-20-7167-2020>, 2020.
- Painemal, D., Corral, A. F., Sorooshian, A., Brunke, M. A., Chellappan, S., Gorooh, V. A., Ham, S.-H., O'Neill, L., Smith, W. L., Tselioudis, G., Wang, H., Zeng, X., and Zuidema, P.: An Overview of Atmospheric Features Over the Western North Atlantic Ocean and North American East Coast—Part 2: Circulation, Boundary Layer, and Clouds, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033423, <https://doi.org/10.1029/2020JD033423>, 2021.
- 620 Penner, J. E., Quaas, J., Storelvmo, T., Takemura, T., Boucher, O., Guo, H., Kirkevåg, A., Kristjánsson, J. E., and Seland, Ø.: Model Intercomparison of Indirect Aerosol Effects, *Atmospheric Chemistry and Physics*, 6, 3391–3405, <https://doi.org/10.5194/acp-6-3391-2006>, 2006.
- 625 Penner, J. E., Xu, L., and Wang, M.: Satellite Methods Underestimate Indirect Climate Forcing by Aerosols, *Proceedings of the National Academy of Sciences*, 108, 13404–13408, <https://doi.org/10.1073/pnas.1018526108>, 2011.
- Pinsky, M., Khain, A., Mazin, I., and Korolev, A.: Analytical Estimation of Droplet Concentration at Cloud Base, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/10.1029/2012JD017753>, 2012.
- Pöhlker, M. L., Zhang, M., Campos Braga, R., Krüger, O. O., Pöschl, U., and Ervens, B.: Aitken Mode Particles as CCN in
630 Aerosol- and Updraft-Sensitive Regimes of Cloud Droplet Formation, *Atmospheric Chemistry and Physics*, 21, 11723–11740, <https://doi.org/10.5194/acp-21-11723-2021>, 2021.
- Prabhakaran, P., Shawon, A. S. M., Kinney, G., Thomas, S., Cantrell, W., and Shaw, R. A.: The Role of Turbulent Fluctuations in Aerosol Activation and Cloud Formation, *Proceedings of the National Academy of Sciences*, 117, 16831–16838, <https://doi.org/10.1073/pnas.2006426117>, 2020.
- 635 Quaas, J., Boucher, O., Bellouin, N., and Kinne, S.: Satellite-Based Estimate of the Direct and Indirect Aerosol Climate Forcing, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2007JD008962>, 2008.
- Reutter, P., Su, H., Trentmann, J., Simmel, M., Rose, D., Gunthe, S. S., Wernli, H., Andreae, M. O., and Pöschl, U.: Aerosol- and Updraft-Limited Regimes of Cloud Droplet Formation: Influence of Particle Number, Size and Hygroscopicity on the Activation of Cloud Condensation Nuclei (CCN), *Atmospheric Chemistry and Physics*, 9, 7067–7080, <https://doi.org/10.5194/acp-9-7067-2009>, 2009.
- 640 Righi, M., Hendricks, J., Lohmann, U., Beer, C. G., Hahn, V., Heinold, B., Heller, R., Krämer, M., Ponater, M., Rolf, C., Tegen, I., and Voigt, C.: Coupling Aerosols to (Cirrus) Clouds in the Global EMAC-MADE3 Aerosol–Climate Model, *Geoscientific Model Development*, 13, 1635–1661, <https://doi.org/10.5194/gmd-13-1635-2020>, 2020.
- Roberts, G. C. and Nenes, A.: A Continuous-Flow Streamwise Thermal-Gradient CCN Chamber for Atmospheric Measurements, *Aerosol Science and Technology*, 39, 206–221, <https://doi.org/10.1080/027868290913988>, 2005.
- 645 Rosenfeld, D., Wolff, D. B., and Amitai, E.: The Window Probability Matching Method for Rainfall Measurements with Radar, *Journal of Applied Meteorology and Climatology*, 33, 682–693, [https://doi.org/10.1175/1520-0450\(1994\)033<0682:TWPMMF>2.0.CO;2](https://doi.org/10.1175/1520-0450(1994)033<0682:TWPMMF>2.0.CO;2), 1994.
- Rosenfeld, D., Amitai, E., and Wolff, D. B.: Improved Accuracy of Radar WPMM Estimated Rainfall upon Application of Objective Classification Criteria, *Journal of Applied Meteorology and Climatology*, 34, 212–223, <https://doi.org/10.1175/1520-0450-34.1.212>, 1995.



- Rosenfeld, D., Fischman, B., Zheng, Y., Goren, T., and Giguzin, D.: Combined Satellite and Radar Retrievals of Drop Concentration and
650 CCN at Convective Cloud Base, *Geophysical Research Letters*, 41, 3259–3265, <https://doi.org/10.1002/2014GL059453>, 2014.
- Rosenfeld, D., Zheng, Y., Hashimshoni, E., Pöhlker, M. L., Jefferson, A., Pöhlker, C., Yu, X., Zhu, Y., Liu, G., Yue, Z., Fischman, B.,
Li, Z., Giguzin, D., Goren, T., Artaxo, P., Barbosa, H. M. J., Pöschl, U., and Andreae, M. O.: Satellite Retrieval of Cloud Condensa-
tion Nuclei Concentrations by Using Clouds as CCN Chambers, *Proceedings of the National Academy of Sciences*, 113, 5828–5834,
<https://doi.org/10.1073/pnas.1514044113>, 2016.
- 655 Rosenfeld, D., Zhu, Y., Wang, M., Zheng, Y., Goren, T., and Yu, S.: Aerosol-Driven Droplet Concentrations Dominate Coverage and Water
of Oceanic Low-Level Clouds, *Science*, 363, <https://doi.org/10.1126/science.aav0566>, 2019.
- Russell, L. M., Sorooshian, A., Seinfeld, J. H., Albrecht, B. A., Nenes, A., Ahlm, L., Chen, Y.-C., Coggon, M., Craven, J. S., Flagan, R. C.,
Frossard, A. A., Jonsson, H., Jung, E., Lin, J. J., Metcalf, A. R., Modini, R., Mülmenstädt, J., Roberts, G., Shingler, T., Song, S., Wang, Z.,
and Wonaschütz, A.: Eastern Pacific Emitted Aerosol Cloud Experiment, *Bulletin of the American Meteorological Society*, 94, 709–729,
660 <https://doi.org/10.1175/BAMS-D-12-00015.1>, 2013.
- Schulze, B. C., Charan, S. M., Kenseth, C. M., Kong, W., Bates, K. H., Williams, W., Metcalf, A. R., Jonsson, H. H., Woods, R.,
Sorooshian, A., Flagan, R. C., and Seinfeld, J. H.: Characterization of Aerosol Hygroscopicity Over the Northeast Pacific Ocean: Im-
pacts on Prediction of CCN and Stratocumulus Cloud Droplet Number Concentrations, *Earth and Space Science*, 7, e2020EA001 098,
<https://doi.org/10.1029/2020EA001098>, 2020.
- 665 Seinfeld, J. H., Bretherton, C., Carslaw, K. S., Coe, H., DeMott, P. J., Dunlea, E. J., Feingold, G., Ghan, S., Guenther, A. B., and Kahn,
R.: Improving Our Fundamental Understanding of the Role of Aerosol- Cloud Interactions in the Climate System, *Proceedings of the
National Academy of Sciences*, 113, 5781–5790, 2016.
- Sorooshian, A., Brechtel, F. J., Ma, Y., Weber, R. J., Corless, A., Flagan, R. C., and Seinfeld, J. H.: Modeling and Characterization of a
Particle-into-Liquid Sampler (PILS), *Aerosol Science and Technology*, 40, 396–409, <https://doi.org/10.1080/02786820600632282>, 2006.
- 670 Sorooshian, A., MacDonald, A. B., Dadashazar, H., Bates, K. H., Coggon, M. M., Craven, J. S., Crosbie, E., Hersey, S. P., Hodas, N., and
Lin, J. J.: A Multi-Year Data Set on Aerosol-Cloud-Precipitation-Meteorology Interactions for Marine Stratocumulus Clouds, *Scientific
data*, 5, 180 026, 2018.
- Sorooshian, A., Anderson, B., Bauer, S. E., Braun, R. A., Cairns, B., Crosbie, E., Dadashazar, H., Diskin, G., Ferrare, R., Flagan, R. C., Hair,
J., Hostetler, C., Jonsson, H. H., Kleb, M. M., Liu, H., MacDonald, A. B., McComiskey, A., Moore, R., Painemal, D., Russell, L. M.,
675 Seinfeld, J. H., Shook, M., Smith, W. L., Thornhill, K., Tselioudis, G., Wang, H., Zeng, X., Zhang, B., Ziemba, L., and Zuidema, P.:
Aerosol-Cloud-Meteorology Interaction Airborne Field Investigations: Using Lessons Learned from the U.S. West Coast in the Design
of ACTIVATE off the U.S. East Coast, *Bulletin of the American Meteorological Society*, 100, 1511–1528, [https://doi.org/10.1175/BAMS-
D-18-0100.1](https://doi.org/10.1175/BAMS-
D-18-0100.1), 2019.
- Sorooshian, A., Corral, A. F., Braun, R. A., Cairns, B., Crosbie, E., Ferrare, R., Hair, J., Kleb, M. M., Mardi, A. H., Maring, H., McComiskey,
680 A., Moore, R., Painemal, D., Scarino, A. J., Schlosser, J., Shingler, T., Shook, M., Wang, H., Zeng, X., Ziemba, L., and Zuidema, P.:
Atmospheric Research Over the Western North Atlantic Ocean Region and North American East Coast: A Review of Past Work and
Challenges Ahead, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD031 626, <https://doi.org/10.1029/2019JD031626>, 2020.
- SPEC inc: FFSSP and FCDP Data Processing Manual, 10/27/2012, Available at: [http://www.specinc.com/sites/default/files/Soft-
ware_and_manuals/FCDP_PostProcessingSoftwareManual_Rev2.6_20121027.Pdf](http://www.specinc.com/sites/default/files/Software_and_manuals/FCDP_PostProcessingSoftwareManual_Rev2.6_20121027.Pdf) (Last Access: 7 June 2021), 2012.



- 685 Thornhill, K. L., Anderson, B. E., Barrick, J. D. W., Bagwell, D. R., Friesen, R., and Lenschow, D. H.: Air Motion Intercomparison Flights during Transport and Chemical Evolution in the Pacific (TRACE-P)/ACE-ASIA, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2002JD003108>, 2003.
- Twomey, S.: The Nuclei of Natural Cloud Formation Part II: The Supersaturation in Natural Clouds and the Variation of Cloud Droplet Concentration, *Geofisica pura e applicata*, 43, 243–249, <https://doi.org/10.1007/BF01993560>, 1959.
- 690 Twomey, S.: The Influence of Pollution on the Shortwave Albedo of Clouds, *Journal of the Atmospheric Sciences*, 34, 1149–1152, [https://doi.org/10.1175/1520-0469\(1977\)034<1149:TIOPTO>2.0.CO;2](https://doi.org/10.1175/1520-0469(1977)034<1149:TIOPTO>2.0.CO;2), 1977.
- Twomey, S. and Warner, J.: Comparison of Measurements of Cloud Droplets and Cloud Nuclei, *Journal of the Atmospheric Sciences*, 24, 702–703, 1967.
- Voigt, C., Schumann, U., Minikin, A., Abdelmonem, A., Afchine, A., Borrmann, S., Boettcher, M., Buchholz, B., Bugliaro, L., Costa, A., Curtius, J., Dollner, M., Dörnbrack, A., Dreiling, V., Ebert, V., Ehrlich, A., Fix, A., Forster, L., Frank, F., Fütterer, D., Giez, A., Graf, K., Groß, J.-U., Groß, S., Heimerl, K., Heinold, B., Hüneke, T., Järvinen, E., Jurkat, T., Kaufmann, S., Kenntner, M., Klingebiel, M., Klimach, T., Kohl, R., Krämer, M., Krisna, T. C., Luebke, A., Mayer, B., Mertes, S., Molleker, S., Petzold, A., Pfeilsticker, K., Port, M., Rapp, M., Reutter, P., Rolf, C., Rose, D., Sauer, D., Schäfler, A., Schlage, R., Schnaiter, M., Schneider, J., Spelten, N., Spichtinger, P., Stock, P., Walser, A., Weigel, R., Weinzierl, B., Wendisch, M., Werner, F., Wernli, H., Wirth, M., Zahn, A., Ziereis, H., and Zöger, M.:
- 700 ML-CIRRUS: The Airborne Experiment on Natural Cirrus and Contrail Cirrus with the High-Altitude Long-Range Research Aircraft HALO, *Bulletin of the American Meteorological Society*, 98, 271–288, <https://doi.org/10.1175/BAMS-D-15-00213.1>, 2017.
- Voigt, C., Kleine, J., Sauer, D., Moore, R. H., Bräuer, T., Le Clercq, P., Kaufmann, S., Scheibe, M., Jurkat-Witschas, T., Aigner, M., Bauder, U., Boose, Y., Borrmann, S., Crosbie, E., Diskin, G. S., DiGangi, J., Hahn, V., Heckl, C., Huber, F., Nowak, J. B., Rapp, M., Rauch, B., Robinson, C., Schripp, T., Shook, M., Winstead, E., Ziemba, L., Schlager, H., and Anderson, B. E.: Cleaner Burning Aviation Fuels Can
- 705 Reduce Contrail Cloudiness, *Communications Earth & Environment*, 2, 1–10, <https://doi.org/10.1038/s43247-021-00174-y>, 2021.
- Weigel, R., Spichtinger, P., Mahnke, C., Klingebiel, M., Afchine, A., Petzold, A., Krämer, M., Costa, A., Molleker, S., and Jurkat, T.: Thermodynamic Correction of Particle Concentrations Measured by Underwing Probes on Fast Flying Aircraft, *Atmospheric Measurement Techniques*, 9, 5135–5162, 2016.
- Wendisch, M., Pöschl, U., Andreae, M. O., Machado, L. A. T., Albrecht, R., Schlager, H., Rosenfeld, D., Martin, S. T., Abdelmonem, A., Afchine, A., Araújo, A. C., Artaxo, P., Aufmhoff, H., Barbosa, H. M. J., Borrmann, S., Braga, R., Buchholz, B., Cecchini, M. A., Costa, A., Curtius, J., Dollner, M., Dorf, M., Dreiling, V., Ebert, V., Ehrlich, A., Ewald, F., Fisch, G., Fix, A., Frank, F., Fütterer, D., Heckl, C., Heidelberg, F., Hüneke, T., Jäkel, E., Järvinen, E., Jurkat, T., Kanter, S., Kästner, U., Kenntner, M., Kesselmeier, J., Klimach, T., Knecht, M., Kohl, R., Kölling, T., Krämer, M., Krüger, M., Krisna, T. C., Lavric, J. V., Longo, K., Mahnke, C., Manzi, A. O., Mayer, B., Mertes, S., Minikin, A., Molleker, S., Münch, S., Nillius, B., Pfeilsticker, K., Pöhlker, C., Roiger, A., Rose, D., Rosenow, D., Sauer, D., Schnaiter, M., Schneider, J., Schulz, C., de Souza, R. A. F., Spanu, A., Stock, P., Vila, D., Voigt, C., Walser, A., Walter, D., Weigel, R., Weinzierl, B., Werner, F., Yamasoe, M. A., Ziereis, H., Zinner, T., and Zöger, M.: ACRIDICON-CHUVA Campaign: Studying Tropical Deep Convective Clouds and Precipitation over Amazonia Using the New German Research Aircraft HALO, *Bulletin of the American Meteorological Society*, 97, 1885–1908, <https://doi.org/10.1175/BAMS-D-14-00255.1>, 2016.
- 715 Wood, R., Mechoso, C. R., Bretherton, C. S., Weller, R. A., Huebert, B., Straneo, F., Albrecht, B. A., Coe, H., Allen, G., Vaughan, G., Daum, P., Fairall, C., Chand, D., Gallardo Klenner, L., Garreaud, R., Grados, C., Covert, D. S., Bates, T. S., Krejci, R., Russell, L. M., de Szoeki, S., Brewer, A., Yuter, S. E., Springston, S. R., Chaigneau, A., Tonizzo, T., Minnis, P., Palikonda, R., Abel, S. J., Brown, W. O. J., Williams, S., Fochesatto, J., Brioude, J., and Bower, K. N.: The VAMOS Ocean-Cloud-Atmosphere-Land Study Regional Experiment



- (VOCALS-REx): Goals, Platforms, and Field Operations, *Atmospheric Chemistry and Physics*, 11, 627–654, <https://doi.org/10.5194/acp-11-627-2011>, 2011.
- 725 Zelinka, M. D., Andrews, T., Forster, P. M., and Taylor, K. E.: Quantifying Components of Aerosol-Cloud-Radiation Interactions in Climate Models, *Journal of Geophysical Research: Atmospheres*, 119, 7599–7615, <https://doi.org/10.1002/2014JD021710>, 2014.
- Zelinka, M. D., Randall, D. A., Webb, M. J., and Klein, S. A.: Clearing Clouds of Uncertainty, *Nature Climate Change*, 7, 674–678, <https://doi.org/10.1038/nclimate3402>, 2017.
- Zheng, G., Wang, Y., Wood, R., Jensen, M. P., Kuang, C., McCoy, I. L., Matthews, A., Mei, F., Tomlinson, J. M., Shilling, J. E., Zawadowicz,
730 M. A., Crosbie, E., Moore, R., Ziemba, L., Andreae, M. O., and Wang, J.: New Particle Formation in the Remote Marine Boundary Layer, *Nature Communications*, 12, 527, <https://doi.org/10.1038/s41467-020-20773-1>, 2021.