

Seasonal Variation of Oxygenated Organic Molecules in Urban Beijing and their Contribution to Secondary Organic Aerosol

Yishuo Guo¹, Chao Yan^{1,2,3*}, Yuliang Liu², Xiaohui Qiao⁴, Feixue Zheng¹, Ying Zhang¹, Ying Zhou¹, Chang Li¹, Xiaolong Fan¹, Zhuohui Lin¹, Zemin Feng¹, Yusheng Zhang¹, Penggang Zheng^{5,6}, Linhui Tian⁷, Wei Nie², Zhe Wang⁵, Dandan Huang⁸, Kaspar R. Daellenbach^{3,9}, Lei Yao^{1,3}, Lubna Dada^{3,9}, Federico Bianchi³, Jingkun Jiang⁴, Yongchun Liu¹, Veli-Matti Kerminen³, Markku Kulmala^{1,3}

Affiliations:

¹ Aerosol and Haze Laboratory, Beijing Advanced Innovation Center for Soft Matter Science and Engineering, Beijing University of Chemical Technology, Beijing, China

² Joint International Research Laboratory of Atmospheric and Earth System Research, School of Atmospheric Sciences, Nanjing University, Nanjing, China

³ Institute for Atmospheric and Earth System Research / Physics, Faculty of Science, University of Helsinki, Finland

⁴ State Key Joint Laboratory of Environment Simulation and Pollution Control, State Environmental Protection Key Laboratory of Sources and Control of Air Pollution Complex, School of Environment, Tsinghua University, Beijing, China

⁵ Division of Environment and Sustainability, The Hong Kong University of Science and Technology, Hong Kong SAR, China

⁶ Department of Civil and Environmental Engineering, The Hong Kong Polytechnic University, Hong Kong SAR

⁷ Department of Civil and Environmental Engineering, Faculty of Science and Technology, University of Macau, Taipa, Macau, China

⁸ State Environmental Protection Key Laboratory of Formation and Prevention of Urban Air Pollution Complex, Shanghai Academy of Environmental Sciences, Shanghai, China

⁹ Laboratory of Atmospheric Chemistry, Paul Scherrer Institute, Villigen, Switzerland.

*Correspondence to: Chao Yan (chaoyan@nju.edu.cn)

Abstract Oxygenated organic molecules (OOMs) are crucial for atmospheric new particle formation and secondary organic aerosol (SOA) growth. Therefore, understanding their chemical composition, temporal behavior, and sources is of great importance. Previous studies on OOMs mainly focus on environments where biogenic sources are predominant, yet studies on sites with dominant anthropogenic emissions, such as megacities, have been lacking. Here, we conducted long-term measurements of OOMs covering four seasons of the year 2019 in urban Beijing. The OOM concentration was found to be the highest in summer ($1.6 \times 10^8 \text{ cm}^{-3}$), followed by autumn ($7.9 \times 10^7 \text{ cm}^{-3}$), spring ($5.7 \times 10^7 \text{ cm}^{-3}$) and winter ($2.3 \times 10^7 \text{ cm}^{-3}$), suggesting that enhanced photo-oxidation together with the rise of temperature promote the formation of OOMs. Most OOMs contained 5 to 10 carbon atoms and 3 to 7 effective oxygen atoms ($\text{nO}_{\text{eff}} = \text{nO} - 2 \times \text{nN}$). The average nO_{eff} increased with increasing atmospheric photo-oxidation capacity, which was the highest in summer and the lowest in winter and autumn. By performing a newly developed workflow, OOMs were classified into four types: aromatic OOMs, aliphatic OOMs, isoprene OOMs, and monoterpene OOMs. Among them, aromatic OOMs (29-41 %) and aliphatic OOMs (26-41 %) were the main contributors in all seasons, indicating that OOMs in Beijing were dominated by anthropogenic sources. The contribution of isoprene OOMs increased significantly in summer (33 %), which is much higher than those in other three seasons (8-10 %).

38 Concentrations of isoprene ($0.2\text{--}5.3\times 10^7\text{ cm}^{-3}$) and monoterpene ($1.1\text{--}8.4\times 10^6\text{ cm}^{-3}$) OOMs in Beijing were lower
39 than those reported at other sites, and they possessed lower oxygen and higher nitrogen contents due to high NO_x
40 levels ($9.5\text{--}38.3\text{ ppbv}$) in Beijing. With regard to the nitrogen content of the two anthropogenic OOMs, aromatic
41 OOMs were mainly composed of CHO and CHON species, while aliphatic OOMs were dominated by CHON and
42 CHON_2 ones. Such prominent differences suggest varying formation pathways between these two OOMs. By
43 combining the measurements and an aerosol dynamic model, we estimated that the SOA growth rate through OOM
44 condensation could reach $0.64\text{ }\mu\text{g}\cdot\text{m}^{-3}\cdot\text{h}^{-1}$, $0.61\text{ }\mu\text{g}\cdot\text{m}^{-3}\cdot\text{h}^{-1}$, $0.41\text{ }\mu\text{g}\cdot\text{m}^{-3}\cdot\text{h}^{-1}$, and $0.30\text{ }\mu\text{g}\cdot\text{m}^{-3}\cdot\text{h}^{-1}$ in autumn, summer,
45 spring, and winter, respectively. Despite the similar concentrations of aromatic and aliphatic OOMs, the former had
46 lower volatilities and, therefore, showed higher contributions (46-62%) to SOA than the latter (14-32%). By contrast,
47 monoterpene OOMs and isoprene OOMs, limited by low abundances or high volatilities, had low contributions of
48 8-12% and 3-5%, respectively. Overall, our results improve the understanding of the concentration, chemical
49 composition, seasonal variation and potential atmospheric impacts of OOMs, which can help formulate refined
50 restriction policy specific to SOA control in urban areas.

51

1. INTRODUCTION

Atmospheric aerosols affect global climate both directly and indirectly (Stocker, 2014) and are known to have a detrimental influence on human health (Lelieveld et al., 2015). Modeling studies have suggested that new particle formation (NPF) dominates the number concentration of particles and is an important contributor to cloud condensation nuclei (CCN) in the global atmosphere (Merikanto et al., 2009; Gordon et al., 2017). In terms of aerosol mass, it has been shown that a significant fraction is composed of secondary organic aerosol (SOA) (Zhang et al., 2007; Jimenez et al., 2009; Hallquist et al., 2009). In both NPF and SOA formation processes, oxygenated organic molecules (OOMs, see the definition in Sect. S2 the work of Nie et al. (Nie et al., 2022)) have been acknowledged as an important contributor, and thus advanced understanding of OOMs is crucial.

The role of OOMs in NPF was first suggested in 2002, but they could not be identified or quantified at that time (O'Dowd et al., 2002). Then, the emergence of Atmospheric Pressure interface Time-of-Flight (API-ToF) mass spectrometer (Junninen et al., 2010; Ehn et al., 2010; Ehn et al., 2012) and Chemical Ionization-API-ToF (CI-API-ToF) mass spectrometer (Jokinen et al., 2012; Ehn et al., 2014) provided the first direct measurement of highly oxygenated organic molecules (HOMs), a subgroup of OOMs with the most oxygen content, which inspired later studies on their role in NPF and SOA growth (Kulmala et al., 2013; Schobesberger et al., 2013; Riccobono et al., 2014; Ehn et al., 2014; Kirkby et al., 2016; Tröstl et al., 2016; Bianchi et al., 2016; Lehtipalo et al., 2018; Rose et al., 2018; Stolzenburg et al., 2018; Mohr et al., 2019; Heinritzi et al., 2020; Yan et al., 2020; Caudillo et al., 2021). These studies found that the functionality and volatility of OOMs are the key factors in determining whether OOM species can participate in NPF (Donahue et al., 2013). More specifically, ultra-low-volatility organic compounds (ULVOCs, whose mass saturation concentrations, C^* , are smaller than $3 \times 10^{-9} \mu\text{g}\cdot\text{m}^{-3}$, or number saturation concentrations, N^* , are smaller than 6 cm^{-3} by assuming an average molar mass of 300 Da) are the main participator of the initial nucleation at certain conditions (Schervish and Donahue, 2020), and extremely-low-volatility organic compounds (ELVOCs, $3 \times 10^{-9} < C^* < 3 \times 10^{-4} \mu\text{g}\cdot\text{m}^{-3}$, $6 < N^* < 6 \times 10^5 \text{ cm}^{-3}$) and low-volatility organic compounds (LVOCs, $3 \times 10^{-4} < C^* < 0.3 \mu\text{g}\cdot\text{m}^{-3}$, $6 \times 10^5 < C^* < 6 \times 10^8 \text{ cm}^{-3}$) can have a dominant contribution to the growth of newly formed particles.

Owing to the significance of OOMs in atmospheric aerosols formation and growth, its reliable measurement is of high importance. Up till now, the majority of reported sites in the lower troposphere with OOM measurement are non-urban areas, such as forest, agricultural pasture and countryside, where the most abundant OOM species are oxidized products from monoterpenes and isoprene. In the boreal forest of southern Finland, the reported OOM concentration was the highest in summer ($4.6 \times 10^8 \text{ cm}^{-3}$) (Huang et al., 2020), followed by autumn ($8.0 \times 10^7 \text{ cm}^{-3}$) (Zha et al., 2018) and spring ($\sim 4.0 \times 10^7 \text{ cm}^{-3}$) (Yan et al., 2016; Roldin et al., 2019; Bianchi et al., 2017). The level of OOMs also varied significantly at different sites. In Melpitz agricultural-forest of central Europe, OOM concentration ($2.5 \times 10^8 \text{ cm}^{-3}$ in summer) (Mutzel et al., 2015) was comparable with that in Hyytiälä, while in Alabama forest of the United States, OOM concentration was much higher ($4.8 \times 10^9 \text{ cm}^{-3}$ in summer) (Massoli et

al., 2018;Krechmer et al., 2015), possibly due to higher UVB and temperature. Besides, monoterpene OOMs at agricultural-rural mixed Vielbrunn were also detected ($3.6 \times 10^6 \text{ cm}^{-3}$ in spring), and results showed that many other unidentified species also took a large fraction, especially at night (Kürten et al., 2016). All of these studies highlight the importance of OOM measurement worldwide. Several urban observations were also reported (Brean et al., 2019;Ye et al., 2020). Although they showed that OOMs in Chinese urban cities contain a significant fraction of compounds with 6 to 9 carbons and that many contain nitrogen, they either reported concentrations of a few chosen species or just spectral signals. Moreover, due to the limitation of short measurement periods, they were incapable of exploring the seasonal behavior of OOM concentration and detailed composition, which are crucial for fully evaluating their potential contribution to the growth of SOA.

In this work, we studied the OOMs measured by a CI-API-ToF mass spectrometer using nitrate (NO_3^-) as reagent ions. The dataset covers four seasons of Year 2019. We performed detailed molecular analyses within the mass to charge ratio between 200-400 Th and identified around 1000 OOMs for each season. The seasonal variations of their concentration, molecular composition, volatility distribution, and potential SOA contribution were systematically investigated for the first time. Furthermore, with a newly developed workflow, we traced their potential sources, including aromatics, aliphatics, monoterpenes, and isoprene. Finally, we evaluated the relative contribution of anthropogenic and biogenic sources in different seasons.

2. MEASUREMENTS AND METHODS

2.1. Measurements

The measurement was conducted at the west campus of Beijing University of Chemical Technology (39.95° N , 116.31° E) on the fifth floor of the teaching building, which is about 15 m above the ground level. This station is a representative urban site, and a detailed description can be found elsewhere (Liu et al., 2020;Yan et al., 2021;Guo et al., 2021).

The concentration of OOMs was measured by a nitrate (NO_3^-)-CI-API-ToF mass spectrometer (abbreviated as nitrate CIMS) (Aerodyne Research, Inc.). The basic working principle of this instrument can be found elsewhere (Jokinen et al., 2012), and the detailed sampling configuration is the same as that reported by Yan et al. (Yan et al., 2021). Two steps were included in the quantification of OOM concentration. First, a mass-dependent transmission experiment was conducted according to a previous study (Heinritzi et al., 2016), and the transmission curve was obtained by comparing the decrease of primary ion signals and the increase of added perfluorinated acid signals. Second, the calibration factor of sulfuric acid was applied to estimate OOM concentration. Some studies have shown that OOM molecules with less oxygen number are not ionized as efficiently as sulfuric acid by NO_3^- (Hyytinen et al., 2015;Hyytinen et al., 2018;Riva et al., 2019). Therefore, the reported OOM concentration in this study should be regarded as the lower limit. The concentration of each OOM molecule can be calculated as follows:

$$[OOM] = \frac{\sum_{i=0}^1 (OOM)(HNO_3)_i NO_3^- + (OOM-H)^- (HNO_3)_i}{\sum_{i=0}^2 (HNO_3)_i NO_3^-} \times C \div T_{OOM}$$

The numerator on the right-hand side is the sum of detected signal of that OOM in the unit of counts per second (cps), either as a cluster of a neutral molecule combined with NO_3^- , $(OOM)(HNO_3)_i NO_3^-$, or as a deprotonated ion, $(OOM-H)^-$. It should be pointed out that in the nitrate CIMS, most OOMs were detected as the cluster form of $(OOM)NO_3^-$, and detailed information can be found in Sect. S1. The denominator is the sum of all reagent ion signals in cps. C is the calibration factor of sulfuric acid, ranging from 6.07×10^9 to $7.47 \times 10^9 \text{ cm}^{-3} / (\text{normalized cps})$ during the whole year. Such a narrow range of the calibration factor also indicated that our instrument had a stable performance during the measurement period. T_{OOM} is the relative transmission efficiency of a specific OOM molecule in comparison with that of the reagent ions.

The number concentration of aerosol particles from 6 to 840 nm was measured by a Differential Mobility Particle Sizer (DMPS) (Aalto et al., 2001). The mass concentration of $PM_{2.5}$ was measured with a Tapered Element Oscillating Microbalance Dichotomous Ambient Particulate Monitor (TEOM 1405-DF, Thermo Fisher Scientific Inc, USA). The chemical composition of the $PM_{2.5}$ was obtained from an Aerosol Chemical Speciation Monitor (ACSM) (Jayne et al., 2000; Drewnick et al., 2005), and PMF analysis was further performed to separate secondary organic aerosols from primary ones. Meteorological parameters were measured with a weather station (AWS310, Vaisala Inc.) located on the rooftop of the building. Concentrations of trace gases, including carbon monoxide (CO), sulfur dioxide (SO_2), nitrogen oxides (NO_x), and ozone (O_3), were monitored using Thermo Environmental Instruments (models 48i, 43i-TLE, 42i, 49i, respectively).

The measurement period covers four seasons of the year 2019, including 135 days in total. Winter, spring, summer, and autumn periods range from 5th Jan. to 14th Feb., 15th Mar. to 14th Apr., 10th Jul. to 9th Aug., and 19th Oct. to 18th Nov., respectively.

2.2. A revised workflow for the classification of OOM Sources

A recently developed workflow, based on the molecular composition as well as the up-to-date knowledge of atmospheric OOM formation chemistry, was used for retrieving their possible sources (Xu et al., 2021; Nie et al., 2022). In this approach, mass spectral binning combined with positive matrix factorization (binPMF) (Zhang et al., 2019) needs to be performed first to extract the factor of monoterpene OOMs. However, as performing binPMF is time-consuming and not suitable for large data sets as used in this study, we replaced the binPMF step by the criteria of $nC=10$, $nO_{eff} \geq 4$, and $2 \leq DBE \leq 4$ (Fig. 1) for the selection of monoterpene OOMs. Such standards were set based on their reported composition (Ehn et al., 2012; Yan et al., 2016; Jokinen et al., 2014; Boyd et al., 2015; Berndt et al., 2016; Berndt et al., 2018). Here, nC is the carbon number. $nO_{eff} (= nO - 2 \times nN)$ is the effective oxygen number, which subtracts the number of oxygen bonded to nitrogen by assuming that all nitrogen atoms are in the form of nitrate groups ($-ONO_2$) or peroxyxynitrate nitrate group ($-OONO_2$). To our best knowledge, this is the common case

for all nitrogen-containing compounds formed through the reaction between RO_2 and NO_x (Orlando and Tyndall, 2012; Seinfeld and Pandis, 2016). Exceptions are those nitrophenols (Yan et al., 2016; Wang et al., 2019; Song et al., 2021) that were classified separately (Nie et al., 2022). The reason for choosing $n\text{O}_{\text{eff}}$ rather than $n\text{O}$ is that it better reflects the oxidation state of closed-shell molecules as well as their parent RO_2 radicals. For example, $\text{C}_7\text{H}_9\text{O}_5$ peroxy radical can produce both $\text{C}_7\text{H}_{10}\text{O}_4$ and $\text{C}_7\text{H}_9\text{O}_6\text{N}$ when reacting with NO , and all of them have the same $n\text{O}_{\text{eff}}$ of four. In addition, $n\text{O}_{\text{eff}}$ considers the influence of nitrogen and represents volatility more directly (Yan et al., 2020), and thus makes it easier for the volatility comparison among OOMs with different nitrogen atoms.

DBE denotes the double bond equivalence and is calculated as $(2n\text{C}+2-n\text{H}-n\text{N})/2$, which is the same as the term degree of unsaturation. The DBE of one OOM molecule is influenced by both its precursor and the oxidation processes. For example, aromatic VOCs have DBE values no smaller than 4. For their oxidation products, a previous study has shown that under OH exposures equivalent to approximately 10 h to 15 days in typical atmospheric conditions, they possess DBE values no smaller than 2 (Garmash et al., 2020). However, reported monoterpene OOMs also have DBE values the same as aromatic OOMs, which makes them difficult to distinguish. According to laboratory studies, the majority of monomer products from monoterpene oxidation are C_{10} compounds (Yan et al., 2020). Measurement results also showed that the concentrations of C_{10} aromatic VOCs are very low (Zhang et al., 2017) compared with other $\text{C}_6\text{-C}_9$ ones. Therefore, those C_{10} OOMs with DBE values of 2 to 4 are likely monoterpene OOMs. For OOMs with DBE values smaller than 2, neither aromatics nor monoterpenes oxidation could explain their formation. Hence, the precursors of those OOMs should be the ones without aromatic rings and have smaller DBE values, such as alkanes, alkenes, and some unsaturated oxygen-containing VOCs (OVOCs). OOMs with DBE values of 2 are rather complex. Their precursors could be aromatics, aliphatics, or other unknown sources, and a detailed discussion of the classification criteria can be found in Nie et al. (Nie et al., 2022).

Due to the complexity of atmosphere, there indeed remain some uncertainties in the workflow. For example, a recent study found that some aliphatic VOCs were able to produce OOMs with DBE values of 2 – 3 and $n\text{O}_{\text{eff}}$ no smaller than 6 (Wang et al., 2021), which however, are classified as aromatic OOMs in the workflow. Nie et al. (Nie et al., 2022) has tested the performance of the workflow on OOMs from Wang et al. 2021. Results show that the accuracy is almost 100 % for OOMs from *n*-decane and ~ 75 % for OOMs from cyclohexane (Nie et al., 2022). Thus, the workflow is generally reliable in classifying aliphatic OOMs. In addition, the VOC to NO_x ratios in the experiments of Wang et al. (Table S17) is much higher than in urban Beijing (Table S18). This high-level NO_x might suppress the autooxidation as the $\text{RO}_2 + \text{NO}$ termination reaction likely dominates the fate of RO_2 radicals, leading to comparatively lower $n\text{O}_{\text{eff}}$. In this way, aliphatic OOMs is difficult to reach $n\text{O}_{\text{eff}}$ of 6. The revised workflow finally divided OOMs into five groups: isoprene (IP) OOMs, monoterpene (MT) OOMs, aromatic OOMs, aliphatic OOMs, and a small amount of undistinguished OOMs (6-9 %) that cannot be classified into those four types.




Figure 1. Workflow for retrieving OOM sources. “IP OOMs” represents isoprene-derived OOMs. nO_{eff} and nN are the numbers of effective oxygen and nitrogen in each OOM molecule, respectively. “Y” and “N” denote “Yes” and “No”, respectively.

3. RESULTS AND DISCUSSIONS

3.1. Seasonal Variation of OOM Concentration and Composition

The concentration and molecular composition are the most fundamental characteristics of OOMs. We summarized the OOM concentrations in Beijing and other lower tropospheric sites in Table 1 and Fig. 2 for better comparison. Generally, a clear seasonal trend of OOM concentration in Beijing can be observed, where total OOM concentration is highest in summer ($1.6 \times 10^8 \text{ cm}^{-3}$), followed by autumn ($7.9 \times 10^7 \text{ cm}^{-3}$) and spring ($5.7 \times 10^7 \text{ cm}^{-3}$), and the lowest in winter ($2.3 \times 10^7 \text{ cm}^{-3}$). This apparent increase of OOM concentrations with an increased temperature and theoretical global radiation indicates that elevated solar radiation along with higher temperature favors the generation of OOMs. In comparison to other locations, the level of OOMs in urban Beijing varied within the ranges of previously reported ones (Yan et al., 2016; Roldin et al., 2019; Bianchi et al., 2017; Zha et al., 2018; Huang et al., 2020; Mutzel et al., 2015; Massoli et al., 2018; Nie et al., 2022). Interestingly, the above clear correlation between global radiation (or temperature) and OOM concentration can also be seen in other locations, yet OOMs in forest environments are in general higher than in urban or suburban areas. On the one hand, this observation suggests that OOM formation at a specific environment is prevalently influenced by the strength of atmospheric photochemistry; on the other hand, forest environment appears to have more abundant OOMs than urban environments, possibly because the OOM yield of biogenic VOCs is higher than that of anthropogenic VOCs (Berndt et al., 2016; Teng et al., 2017; Garmash et al., 2020; Molteni et al., 2018). Yet, a quantitative explanation of the OOM variation between seasons and locations requires comprehensive measurements as well as analyses on both the production and loss of OOMs.

Table 1. Mean, standard deviation (Std), median, 25 and 75 percentiles (25th and 75th) of measured OOM concentrations at various lower tropospheric sites.

Measurement Site	Period	Mean (cm ⁻³)	Std (cm ⁻³)	Median (cm ⁻³)	25 th (cm ⁻³)	75 th (cm ⁻³)	Reference
Beijing, China	2019 Jan.-Feb.	2.7×10^7	1.7×10^7	2.3×10^7	1.3×10^7	3.6×10^7	This study
Beijing, China	2019 Mar.-Apr.	6.9×10^7	5.1×10^7	5.7×10^7	3.1×10^7	8.9×10^7	This study
Beijing, China	2019 Jul.-Aug.	1.6×10^8	7.5×10^7	1.6×10^8	1.1×10^8	2.2×10^8	This study
Beijing, China	2019 Oct.-Nov.	8.3×10^7	5.2×10^7	7.9×10^7	4.0×10^7	1.2×10^8	This study
Hong Kong, China	2018 Nov.	2.3×10^8	1.1×10^8	2.1×10^8	1.5×10^8	2.9×10^8	2022, Nie et al.
Shanghai, China	2018 Nov.	7.8×10^7	6.3×10^7	6.1×10^7	2.6×10^7	1.2×10^8	2022, Nie et al.
Nanjing, China	2018 Nov.	7.7×10^7	5.4×10^7	7.2×10^7	3.1×10^7	1.1×10^8	2022, Nie et al.
Hyytiälä forest, Finland	2012 Apr.-May	7.5×10^7	6.1×10^7	5.7×10^7	3.6×10^7	9.2×10^7	2016, Yan et al.
Hyytiälä forest, Finland	2013 May	1.4×10^7	7.9×10^6	1.2×10^7	8.1×10^6	1.8×10^7	2019, Roldin et al.
Hyytiälä forest, Finland	2013 Apr.-Jun.	4.5×10^7	1.2×10^7	4.9×10^7	3.2×10^7	5.5×10^7	2017, Bianchi et al.
Hyytiälä forest, Finland	2016 Sep.	1.2×10^8	1.0×10^8	8.0×10^7	3.8×10^7	1.7×10^8	2018, Zha et al.
Melpitz, Germany	2013 Jul.	2.7×10^8	1.7×10^8	2.5×10^8	1.4×10^8	3.5×10^8	2015, Mutzel et al.
Alabama forest, USA	2013 Jun.-Jul.	4.7×10^9	1.5×10^9	4.8×10^9	3.7×10^9	5.3×10^9	2018, Massoli et al.




Figure 2. OOM concentration vs. temperature at various lower tropospheric sites during daytime (07:00 – 17:00). Data points are colored by theoretical global radiation (GlobRad). Square, circle, and triangle markers represent urban, suburban, and forest areas, respectively. The gray error bars show standard deviations (1 σ). Nanjing, Shanghai and Hong Kong datasets are from Nie et al. (Nie et al., 2022), Melpitz data is from Mutzel (Mutzel et al., 2015), Alabama data is from Massoli et al. (Massoli et al., 2018), and Hyytiälä datasets are from Yan et al. (Yan et al., 2016) and Zha et al. (Zha et al., 2018).

To further demonstrate the seasonal influence of solar radiation and precursor VOCs on OOM concentration, we classified OOMs of each season into four groups based on the brightness parameter (Dada et al., 2017) and PM_{2.5} level, respectively. As shown in Fig. S6, in seasons other than summer, OOM concentration under polluted conditions is much higher than that under clean conditions, which likely results from the elevation of precursors coming along with polluted air masses and the accumulation during the pollution. Besides, the concentration of

total OOMs on sunny days is higher than that on cloudy days, implying that photochemical oxidation plays a key role in the production of OOM molecules. There is one exception that OOM concentration is not significantly different between sunny and cloudy days under clean condition in autumn, and the cause cannot be concluded in this study without a complete VOC measurement.

For OOM composition, the two-dimensional H/C-O_{eff}/C (ratio of hydrogen number to carbon number vs. ratio of effective oxygen number to carbon number) diagrams are plotted to show its characteristics (Fig. S7). And main CHO, CHON, and CHON₂ OOM species are also summarized in Table 2. Generally, the composition of OOM molecules exhibits high similarity among different seasons, suggesting no significant changes in OOM formation in general. However, two seasonal characteristics can be found. First, the most oxygenated OOM molecules, such as C_nH_{2n-2}O_{6,7}, C_nH_{2n-4}O_{7,8}, C_nH_{2n+1}O₈N and C_nH_{2n-1}O₉N OOMs, are mainly observed in summer, and meanwhile, the least oxygenated ones, e.g., C_nH_{2n-7}O₂N and C_nH_{2n-9}O_{4,5}N are mostly detected in winter. These observations indicate that, in addition to the enhanced OOM concentration, strong photochemistry also leads to a high oxidation state of OOM. And these summer-specific OOMs can be classified as highly oxygenated organic molecules (HOMs). Second, C₅H₁₀O₈N₂ is exceedingly high in summer. A previous study suggested that C₅H₁₀O₈N₂ is one dominant oxidation product from isoprene (Xu et al., 2021), and therefore, the high concentration of C₅H₁₀O₈N₂ is a clear indication of intensive isoprene oxidation in summer, which will be discussed in detail in Sect. 3.2.

Table 2. Main CHO, CHON and CHON₂ OOM species measured in this study.

DBE	CHO OOMs	CHON OOMs	CHON ₂ OOMs
0	C _n H _{2n+2} O ₆	C _n H _{2n+1} O ₃₋₈ N	C _n H _{2n} O ₄₋₁₁ N ₂
1	C _n H _{2n} O ₂₋₈	C _n H _{2n-1} O ₃₋₉ N	C _n H _{2n-2} O ₄₋₁₀ N ₂
2	C _n H _{2n-2} O ₃₋₇	C _n H _{2n-3} O ₃₋₉ N	C _n H _{2n-4} O ₅₋₁₁ N ₂
3	C _n H _{2n-4} O ₂₋₈	C _n H _{2n-5} O ₃₋₁₀ N	C _n H _{2n-6} O ₈₋₁₁ N ₂
4	C _n H _{2n-6} O ₃₋₉	C _n H _{2n-7} O ₃₋₉ N	C _n H _{2n-8} O _{7,8} N ₂
5	C _n H _{2n-8} O ₃₋₈	C _n H _{2n-9} O ₄₋₁₀ N	C _n H _{2n-10} O ₆₋₁₀ N ₂

For a better understanding of OOM composition variation among seasons, the distributions of nC, nO_{eff}, nN, and DBE, as well as their seasonal variations, are further analyzed. It should be pointed out that the concentration (5.3×10⁷ cm⁻³) and the fraction (33 %) of IP OOMs in summer is much higher than those in other three seasons, and therefore they are plotted in bars with diagonal lines individually (Fig. 3). In terms of carbon content, the majority of OOMs contain 5 to 10 carbon atoms. For OOMs with 6 to 10 carbon atoms, in seasons other than summer, C₆ are the most abundant, and a decreasing trend can be seen along with an increasing nC, while in summer, an opposite trend is observed, i.e., the relative contribution increases with an increasing nC. The causes behind the different trends in summer and other seasons are complex but might include changes in the precursor VOC distribution, varying reactivity responses of VOCs to temperature, and the volatilities of OOMs that influence their atmospheric lifetime. Further analysis on this topic will be made in the future. For C₅ OOMs, the contribution from isoprene varies from less than half in winter and spring to ~70 % in summer. The high contribution of IP OOMs in summer

is in line with the strong isoprene emission coupled with the enhanced photo-oxidation (Cheng et al., 2018;Zhang et al., 2020).

Concerning the oxygen content, most OOMs contain 3 to 7 effective oxygen atoms, accounting for 86-95 % of total OOMs in all seasons. With the increase of effective oxygen number, the contribution of corresponding OOMs first increases and then decreases, with $nO_{eff}=4$ OOMs having the highest fraction. Fig. S8 shows that the concentration-weighted average nO_{eff} is the highest in summer and lowest in winter and autumn, which is consistent with the observation of individual molecules where the most highly oxygenated ones are usually more abundant in summer. The enhanced multi-step oxidation (Garmash et al., 2020;Wu et al., 2021) and favored auto-oxidation (Molteni et al., 2018;Wang et al., 2017;Wang et al., 2018b;Bianchi et al., 2019) at high temperatures in warmer seasons are most likely the causes. Furthermore, when taking IP OOMs into account, $nO_{eff}=4$ becomes even more prominent in summer, in which $C_5H_{10}O_8N_2$ takes the largest portion and accounts for 77 % of $nO_{eff}=4$ IP OOMs. In winter and autumn, however, $nO_{eff}=3$ has a much higher fraction than those in the other two seasons, and those OOMs are mainly composed of low-DBE compounds, such as $C_nH_{2n}O_7N_2$, $C_nH_{2n-2}O_7N_2$, and $C_nH_{2n-1}O_5N$ species.

As for nitrogen content, the vast majority (98-99 %) of OOMs contain 0 to 2 nitrogen atoms, in which CHON OOMs take the largest fraction, varying from 42 % to 51 % among seasons. It should be noted that, although the mixing ratios of NO_x in different seasons change significantly, the nitrogen distributions of non-isoprene OOMs are similar, which is probably due to the fact that NO (0.6-10.0 ppbv) and NO_2 (8.9-28.3 ppbv) concentrations in urban Beijing are always high throughout the year. Those nitrogen atoms could come from either NO_3 radical oxidation or NO_x termination. During the day, nitrogen is likely added mainly through NO_x termination as NO_3 radical should be photolyzed or titrated by NO. However, in the absence of NO_3 photolysis at night, when NO concentration is low, the concentration of NO_3 radical could reach up to ~ 10 pptv ($2.7 \times 10^8 \text{ cm}^{-3}$) in Beijing (Wang et al., 2018a). Under such levels, the NO_3 radical could even dominate the oxidation of biogenic VOCs (i.e., isoprene and monoterpenes) and some aliphatic VOCs, yet the oxidation of aromatic VOCs are driven by OH radicals (Table S4 and Table S5). Therefore, nitrogen is possibly added through both processes at night.

In the case of DBE distribution, most OOMs comprise 0 to 6 DBE values and there is not too much difference among seasons for non-isoprene OOMs. Generally, with the increase of DBE, the fraction of corresponding non-isoprene OOMs first increases and then decreases, with DBE=2 OOMs having the highest contribution (20-29 %). And this is possibly caused by the fact that almost all precursor VOCs, such as aromatics, aliphatics and monoterpenes, can form oxidation products with DBE value of 2. OOMs with DBE larger than 4 and nC no smaller than 10 are likely derived from polycyclic aromatic hydrocarbons (PAHs, $DBE \geq 7$), and their fraction varies from 5 % in summer to 7-8% in the other three seasons. This demonstrates that PAHs may also have a non-negligible contribution to total OOMs. For IP OOMs, most of them possess 0 or 1 DBE, and only a small fraction (2-16 %) of them retains DBE of precursor isoprene. This indicates that hydroxyl or hydroperoxyl are the major functional groups of these products, whereas products with higher DBE values are not prominent. It could mean that

compounds containing carbonyl or epoxide groups are either not efficiently formed under the urban environment, or fast lost via heterogeneous reactions (Riedel et al., 2015; Zhang et al., 2022). Meanwhile, the instrumental detection bias, i.e., nitrate CIMS is less sensitive to carbonyl and epoxide groups, cannot be excluded. Therefore, further combination of CIMS instruments with different chemical ionization methods are highly desirable in the future.




Figure 3. Number of carbon (nC), effective oxygen (nO_{eff}), nitrogen (nN), and double bond equivalence (DBE) distribution of OOMs for four seasons. The abbreviations “IP” and “non-IP” represent IP OOMs and other non-isoprene OOMs, respectively. The bars with diagonal lines and filled colors represent IP OOMs and non-isoprene OOMs, respectively.

3.2. Characteristics of Source-classified OOMs

With the workflow described in Sect.2.2, total OOMs were classified into four types: IP OOMs, MT OOMs, aromatic OOMs, and aliphatic OOMs. As shown in Fig. 4, the seasonal concentrations of OOMs from different sources vary with the same trend, highest in summer, followed by autumn, spring, and winter. During the whole year, aromatic OOMs (29-41 %) and aliphatic OOMs (26-41 %) are the most abundant categories, demonstrating that OOMs in Beijing are dominantly from anthropogenic sources. This is also consistent with the observation of SOA composition in previous studies (Le Breton et al., 2018; Mehra et al., 2021). In terms of OOMs from biogenic sources, IP OOMs show a prominent contribution in summer (33 %), which is much higher than those in other seasons (8-10 %). Although it is recently suggested that isoprene can have both biogenic and anthropogenic sources (Wagner and Kuttler, 2014; Panopoulou et al., 2020), the much higher enhancement of IP OOMs in summer can only be explained by the large additional biogenic emission (Cheng et al., 2018; Mo et al., 2018). For MT OOMs,

301 however, the fractional contribution does not show a seasonal variation as clear as that of IP OOMs – it only varies
 302 between 5 % and 6 %.




Figure 4. Concentration (left panel) and fraction (right panel) of source-classified OOMs in four seasons. The abbreviations “IP” “MT” “Aro” “Ali” and “Un” stand for IP OOMs, MT OOMs, aromatic OOMs, aliphatic OOMs and undistinguished OOMs respectively. “Win” “Spr” “Sum” and “Aut” represent winter, spring, summer and autumn separately.

3.2.1 Characteristics of biogenic OOMs

The spectral profiles and the fractions of IP OOMs with different nitrogen numbers in four seasons are shown in Fig. 5. Prominent IP OOM species include C₄H₆O₄, C₅H₈O₄, C₄H₇O_{6,7}N, C₅H₉O₅₋₇N, C₅H₁₀O_{7,8}N₂ and C₅H₉O₁₀N₃. CHON₃ OOM (C₅H₉O₁₀N₃) is detected in all four seasons, suggesting that multi-generation oxidation is involved throughout the year. Besides, the composition of IP OOMs exhibits clear seasonal variation. First, compared with other three seasons, C₅H₁₀O₈N₂ and C₅H₉O₁₀N₃ have much higher contributions in summer, indicating that NO_x may be involved more efficiently in the oxidation process of isoprene despite its lower concentration (Fig. S10 and Table S1). In addition, nighttime NO₃ radicals produced efficiently during summer nights (Wang et al., 2018a) should also promote their formation. Second, despite the overall lowest concentrations of IP OOMs in winter, C₄H₈O₇ exhibits a maximum concentration (Fig. S9) and the highest fraction, implying that it may have additional sources other than isoprene oxidation in winter. Third, due to the influence of C₅H₁₀O₈N₂ and C₅H₉O₁₀N₃, CHON₂ and CHON₃ IP OOMs take extremely large proportion (~ 67 %) in summer. And interestingly, the seasonal trend of nitrate IP OOM fraction (from largest to smallest is summer, autumn, spring and winter) did not follow the variation of NO_x concentration (from highest to lowest is autumn, winter, spring and summer, Fig. S5 and Table S1), which suggests that the formation of nitrate IP OOMs probably has a non-linear response to NO_x.

The concentrations of these prominent IP OOM molecules during summertime in Beijing, Nanjing (32.12° N, 118.95° E) (Liu et al., 2021) and Alabama mixed-forest (32.90° N, 87.25° W) (Krechmer et al., 2015; Massoli et al., 2018) are further compared (Fig. 6). Please note that these molecules plotted are not all the IP OOMs but rather selected abundant ones reported in the literature and in this study. As shown in Fig. 6 (A), IP OOMs exhibit the

highest concentration in Alabama forest and the lowest one in Beijing, and the level of IP OOMs in Beijing and Nanjing are comparable. This concentration difference is likely caused by the variation of biogenic isoprene emissions, since Alabama measurement was conducted in a forest, Nanjing site is a suburban area with large vegetation coverage nearby, and Beijing site is located in urban downtown. Besides, the overall varying patterns of IP OOM species in Beijing and Nanjing are very similar, indicating that isoprene in those two urban sites undergo similar oxidation pathways. In terms of oxygen distribution, Beijing and Nanjing are rather similar in that $nO_{eff}=4$ OOMs contribute the most, whereas in Alabama $nO_{eff}=5$ ones are the most abundant (Fig. 6 (B)). This lower oxygen content in urban cities is probably caused by the high NO_x levels (11.1 ppbv, 8.5 ppbv and 0.5 ppbv for Beijing, Nanjing and Alabama respectively, Fig. S10), since NO_x efficiently suppresses the oxygen addition of RO_2 radicals (Zhao et al., 2018). Furthermore, the different NO_x levels among the three sites also influence the nitrogen content that Beijing is the highest and Alabama is the lowest (Fig. 6 (C)).

From the perspective of diurnal variation, CHO IP OOMs in Beijing possess one daytime peak, while CHON OOMs mainly contain day-night-dual-peak or nocturnal-peak-only types (Fig. S11), which is similar to that reported in Alabama forest (Massoli et al., 2018). But it should be noted that the diurnal variations of some CHON IP OOMs with same molecular composition in this study and Massoli et al. (2018) are not identical, suggesting that their formation pathways are different under various atmospheric conditions.




Figure 5. Fractional profiles of each IP OOM molecule in (A) winter, (B) spring, (C) summer and (D) autumn. The mass to charge ratio (m/z) denotes OOM clustered with NO_3^- or as deprotonated ones. The fraction of each compound is calculated as the ratio of its concentration to the total concentration of IP OOMs. The red, green, blue and orange bars are for CHO, CHON, $CHON_2$ and $CHON_3$ OOMs respectively. Please note that $CHON_3$ OOMs only include $C_5H_9O_{10}N_3$.





Figure 6. (A) Concentration comparison of specific fingerprint IP OOM molecules between our study and previously reported ones (Krechmer et al., 2015; Massoli et al., 2018; Liu et al., 2021). The markers are median concentration values, and the bottom and up ranges of the error bars denote 25th and 75th percentiles respectively. Distribution of (B) effective oxygen number, nO_{eff}, and (C) nitrogen number, nN, of IP OOMs plotted in figure (A).

Different from IP OOMs, the overall composition distributions of MT OOMs in the four seasons are quite similar and vary with identical oxygen addition patterns (Fig. 7). Predominant MT OOM molecules are C₁₀H₁₄O₄₋₈, C₁₀H₁₆O₄₋₆, C₁₀H₁₈O_{4,5}, C₁₀H_{13,15,17}O₆₋₉N and C₁₀H_{14,16}O₈₋₁₀N₂. Besides, most MT OOMs belong to CHON category (54-59 %), and the CHO (18-21 %) and CHON₂ (19-27 %) ones, with a comparable contribution during the year.

Then, for a better understanding of MT OOM characteristics under different atmospheric environments, representative MT OOM molecules in summer Beijing, spring Hyytiälä forest (61.8° N) (Yan et al., 2016) and summer Alabama mixed-forest (Massoli et al., 2018) are further compared. The following differences can be identified. First, MT OOM concentrations are the highest in Alabama and the lowest in urban Beijing (Fig. 8 (A)), which should result from the synergetic influence of UVB, temperature and precursor monoterpenes. Second, the levels of the two MT radicals that have high concentrations in the Hyytiälä forest, C₁₀H₁₅O₈[•] and C₁₀H₁₅O₁₀[•], are not detected in Beijing. This is possibly caused by both low monoterpene abundance (Cheng et al., 2018) and high NO_x concentration in Beijing (11.06 ppbv, Fig. S10), which lead to a low production rate and high loss rate of RO₂ radicals. Third, most MT OOMs in urban Beijing possess 5 or 6 effective oxygen, whereas in the forest environment a large fraction of them can hold 7 to 10 effective oxygen (Fig. 8 (B)). This again suggests that high NO_x in Beijing effectively inhibits the oxygen addition processes (Zhao et al., 2018; Orlando and Tyndall, 2012). Additionally, high NO_x in Beijing also leads to high nitrogen content (Fig. 8 (C)) by promoting the termination reaction between RO₂

368 and NO_x (Orlando and Tyndall, 2012) and facilitating the formation of NO₃ radical (Wang et al., 2018a), which
 369 further leads to the formation of nitrate MT OOMs (Boyd et al., 2015; Nah et al., 2015).



370 **Figure 7.** Fractional profiles of each MT OOM molecule in (A) winter, (B) spring, (C) summer and (D) autumn. The mass to
 371 charge ratio (m/z) denotes OOM clustered with NO₃⁻ or as deprotonated ones. The fraction of each compound is calculated as
 372 the ratio of its concentration to the total concentration of MT OOMs. The red, green, blue and orange bars are for CHO, CHON,
 373 CHON₂ and CHON₃ OOMs respectively.
 374




Figure 8. (A) Concentration comparison of specific fingerprint MT OOM molecules between our study and previously reported ones (Massoli et al., 2018; Yan et al., 2016). The markers are median concentration values, and the bottom and up ranges of the error bars denote 25th and 75th percentiles respectively. Please note that only the summer Beijing data was plotted as the overall pattern of MT OOMs in Beijing summer and spring are very similar (Fig. 7). Distribution of (B) effective oxygen number, nO_{eff}, and (C) nitrogen number, nN, of MT OOMs plotted in figure (A).

3.2.2 Characteristics of anthropogenic OOMs

Although the oxidation pathways and product composition of a few aromatic VOCs have been studied previously, the reported products are of much less diversity compared to the complex real atmosphere. Therefore, we rely on the workflow (see Sect. 2.2 and (Nie et al., 2022)) to find out possible aromatic OOMs in our measurement. Among the deduced aromatic OOMs, almost all C₆-C₉ CHO and C₆ CHON compounds have been detected in previous benzene, toluene, xylene, ethylbenzene and mesitylene experiments (Molteni et al., 2018; Garmash et al., 2020) (see detail in Table S9), which demonstrates the reliability of our workflow.

As shown in Fig. 9 (A) to (D), predominant aromatic species in different seasons possess high similarity and they could be classified into C₆₋₉H₈₋₁₄O₄, C₆₋₈H₈₋₁₂O₅, C_{7,8}H_{8,10}O₅, C₆₋₁₀H₇₋₁₅O₆N, C₇₋₉H₉₋₁₃O₇N, C₈₋₁₀H₁₁₋₁₅O₈N and C_{8,9}H₁₂₋₁₄O₁₀N₂ categories, among which a prominent CH₂ spacing is seen. Such patterns are most likely due to the co-existence of homologous precursor VOCs, although fragmentation processes during the oxidation could also play a role (Pan and Wang, 2014; Zaytsev et al., 2019; Xu et al., 2020). Besides, the distribution of CHON aromatic OOMs in winter and autumn are very similar, with C₈H₁₁O₆N being the highest; in comparison, the overall distribution moves to higher oxygen content in summer, e.g., C₈H₁₁O₇N becomes the largest one. This suggests that enhancement of radiation, which leads to strong photochemistry and high temperature, and the reduction of NO_x in

summer benefit the formation of highly oxygenated organic molecules (Garmash et al., 2020; Orlando and Tyndall, 2012). There are also fingerprint molecules for different seasons, such as $C_{10}H_8O_6$ in spring, $C_9H_{10}O_{11}N_2$ in summer, and $C_{10}H_6O_4$ in summer and autumn. Due to the complexity of real atmosphere, the reason for their seasonal variation is unclear, and further analysis is warranted. In terms of nitrogen content, aromatic OOMs contain large fraction of CHO (42-52 %) and CHON (39-51 %) species. The contribution of $CHON_2$ OOMs reaches the highest in summer, which again indicates that the involvement of NO_x is enhanced under the influence of elevated UVB and temperature. The carbon distribution among seasons are very similar (Fig. S13), where C4 to C9 aromatic OOMs, probably derived from monocyclic aromatic hydrocarbons, make up 68-76 %, and other $C \geq 10$ ones, of which 59-68 % are likely the oxidation products from PAHs ($DBE \geq 5$) (Table S10), take up 24-32 %. This implies that the relative abundance of emitted aromatic precursors with different carbon atoms is quite stable during the year.

Major aliphatic OOM molecules in different seasons are highly similar, and they possess more evident homologous patterns than aromatic OOMs (Fig. 9 (E) to (H) and Table S11). The dominant species of aliphatic OOMs are $C_{6-9}H_{10-16}O_4$, $C_{6-10}H_{11-19}O_6N$, $C_{5-10}H_{7-17}O_6N$, $C_{5-10}H_{8-18}O_8N_2$ and $C_{6-9}H_{12-18}O_7N_2$, and some less oxygenated compounds, e.g., $C_{6-8}H_{11-15}O_5N$, also have considerable contributions in winter and autumn. For nitrogen content, CHO aliphatic OOMs only take a small fraction of 8-11 %, implying that NO_x termination may dominate the formation of aliphatic close-shell molecules. Meanwhile, this CHO OOM fraction of aliphatic OOMs is much smaller than that of aromatic OOMs (42-52 %), suggesting that the branching ratio of aliphatic $RO_2 - NO_x$ reaction forming $CHON_x$ species is higher than that of the aromatic one. Besides, unlike aromatic OOMs, aliphatic $CHON_2$ OOMs have a bigger contribution in winter than in the other three seasons. This is because a major sequence of $CHON_2$ OOMs, $C_{6-14}H_{12-28}O_7N_2$, is found to be coincided with $PM_{2.5}$ (Fig. S14), which is frequently high in winter. In winter Beijing, the regional transportation of air masses from the south and southeast areas is a large source of $PM_{2.5}$ (Wang et al., 2015; Chen et al., 2021; Chen et al., 2022; Tan et al., 2022), therefore, the good correlation between $CHON_2$ species and $PM_{2.5}$ suggests that those compounds might also have regional sources. Table S12 shows that $C_{6-10}H_{12-20}O_7N_2$ are relatively volatile and are classified as semi-volatile organic compounds, thus they could originate from long-distance transport. It is very likely that they are equilibrated in larger gas-phase concentrations as SOA also increases with the elevation of $PM_{2.5}$ (Fig. S15). Whereas for $C_{11-14}H_{22-28}O_7N_2$, they are less volatile and lie in the range of low-volatility organic compounds, hence, they can hardly be transported and are more likely produced during the air mass transportation. Those pollution-related OOMs take the largest (14 %) and smallest (2 %) fraction in winter and summer respectively (Table S11). In terms of carbon distribution, there is not too much difference among seasons, in which the relatively short C4 to C9 aliphatic OOMs make up 83-90 % and the longer ones take up 10-17 % (Fig. S13).




Figure 9. Fractional profiles of each aromatic OOM molecule in (A) winter, (B) spring, (C) summer and (D) autumn, and of each aliphatic OOM molecule in (E) winter, (F) spring, (G) summer and (H) autumn. The mass to charge ratio (m/z) denotes OOM clustered with NO_3^- or as deprotonated ones. The fraction of each compound is calculated as the ratio of its concentration to the total concentration of aromatic OOMs ((A) to (D)) or aliphatic OOMs ((E) to (H)). The red, green, blue and orange bars are for CHO, CHON, CHON₂ and CHON₃ OOMs respectively. Species marked with grey dashed lines are primary ones during the year, and species in grey background are special ones for specific seasons.

Up till now, field measurements of anthropogenic OOMs are rare (Nie et al., 2022;Liu et al., 2021). In general, the concentrations of aromatic and aliphatic OOMs in Beijing are comparable with those in other Chinese megacities (Table S13), and fingerprint aromatic and aliphatic OOM molecules in Beijing and Nanjing are also identical (Table S14). This suggests that the OOM production, including both the precursor emissions and oxidation mechanisms, may share high similarities in megacities. Yet, a more systematic comparison can only be made when measurements at more locations are available in the future.

3.3. Atmospheric implication: OOM contribution to SOA through condensation

The volatility of organic compound determines its partitioning between gas and particle phases, and thus influences its atmospheric lifetime, gas-phase concentration, and contribution to SOA. Therefore, we estimate the volatility of source-classified OOMs (detailed method can be found in Sect. S3) and summarize the results in Fig. 10. The seasonal variations of OOMs classified as ELVOCs (extremely low-volatility organic compounds), LVOCs (low-volatility organic compounds) and SVOCs (semi-volatile organic compounds) follow the same trend as that of total OOMs, with the highest concentrations in summer ($1.3 \times 10^7 \text{ cm}^{-3}$, $4.0 \times 10^7 \text{ cm}^{-3}$ and $8.4 \times 10^7 \text{ cm}^{-3}$ for ELVOCs, LVOCs, and SVOCs, respectively) and the lowest ones in winter ($4.4 \times 10^6 \text{ cm}^{-3}$, $9.4 \times 10^6 \text{ cm}^{-3}$ and $5.3 \times 10^6 \text{ cm}^{-3}$ for ELVOCs, LVOCs, and SVOCs, respectively). Here, we focus particularly on OOMs with relatively low volatility with high potential contributing to the formation of SOA.

Due to the concentration variation of four source-classified OOMs and their temperature-dependent volatility distribution (Table S16), their fractions within different volatility ranges have distinct seasonal characteristics (Fig. 10 (B)). Among ELVOCs, aromatic OOMs take the largest fractions, ranging from 72 to 94 % throughout the year. For LVOCs, aromatic (34-51 %) and aliphatic OOMs (17-42 %) are the two that have the largest proportions. And MT OOMs, favored by its low volatility (Table S16) (Tröstl et al., 2016;Yan et al., 2020), also take up ~ 14 % of LVOCs in seasons other than winter. IP OOMs, however, due to its high volatility (Table S16) (Krechmer et al., 2015;Xu et al., 2021), do not have an appreciable contribution to ELVOCs and LVOCs even in summer when its concentration is exceedingly high. Consequently, it is likely that the pure condensation of IP OOMs has a minor contribution to SOA growth regardless of the season.




Figure 10. (A) Concentration and (B) fraction of source-classified OOMs in ELVOCs (extremely low volatile organic compounds), LVOCs (low volatile organic compounds) and SVOCs (semi-volatile organic compounds) in four seasons. Please also note that the fractions smaller than 4% are not marked.

The rate of OOM condensation onto particles, referred to as condensation flux hereafter, was calculated based on the particle dynamic model proposed by Trostl et al. (Tröstl et al., 2016) (see details in Sect. S5). In terms of seasonal variation (Fig. 11 (A)), OOM condensation flux exhibits the highest level in autumn ($0.64 \mu\text{g}\cdot\text{m}^{-3}\cdot\text{h}^{-1}$), followed by summer ($0.61 \mu\text{g}\cdot\text{m}^{-3}\cdot\text{h}^{-1}$), spring ($0.41 \mu\text{g}\cdot\text{m}^{-3}\cdot\text{h}^{-1}$), and decreases to the lowest in winter ($0.30 \mu\text{g}\cdot\text{m}^{-3}\cdot\text{h}^{-1}$). For seasonal comparison of SOA formation rate caused by OOM condensation, the characteristic accumulation time (AccTime), defined as SOA divided by OOM condensation flux, is calculated as an indicator (see details in Sect. S4). As shown in Fig. 11 (B), a characteristic time of 24 hours is enough to explain the observed SOA concentration by OOM condensation in winter, and it is reduced to 7 hours, 10 hours, and 7 hours for spring, summer, and autumn, respectively. It should be noted that this should not be interpreted as the entire SOA being formed via OOM condensation during this characteristic time, but rather that OOM condensation is efficient and can have a significant contribution to SOA formation. A recent study (Nie et al., 2022) suggested that OOM condensation can account for about 40% of the SOA formation in wintertime Beijing. Our analysis on seasonal variation indicates that the condensation of OOMs could have a larger contribution to SOA formation in seasons other than winter.

For OOMs from different sources, aromatic OOMs contributes the most during the year, varying from 46 % to 62 %, followed by aliphatic OOMs (14-32 %). In comparison, the two biogenic ones, MT OOMs (8-12 %) and IP OOMs (3-5 %), have smaller contribution in all the four seasons. This indicates that the formation of SOA through condensation in urban Beijing is dominated by anthropogenic sources, which is in line with the previously reported SOA composition (Le Breton et al., 2018; Mehra et al., 2021). Overall, our results suggest that in order to control the formation of SOA, the emission of anthropogenic VOCs, especially aromatics, should be restricted with a high priority.




Figure 11. (A) Condensation flux of OOMs calculated by the particle dynamic model by Trostl et al. (Tröstl et al., 2016) in four seasons. (B) Characteristic accumulation time of SOA (AccTime), calculated as SOA divided by OOM condensation flux, in four seasons. This parameter is used as indicator for the relative accumulation rate of SOA caused by OOM condensation in different seasons. The values in each box of (A) and (B) are the median values of corresponding parameters. (C) Estimated condensation flux contribution of four source-classified OOMs in four seasons. The abbreviations “IP” “MT” “Aro” “Ali” and “Un” stand for IP OOMs, MT OOMs, aromatic OOMs, aliphatic OOMs and undistinguished OOMs respectively.

4. Summary and Conclusions

A long-term measurement of OOMs based on nitrate CIMS was conducted in urban Beijing. OOMs in the mass range of 200-400 Th were systematically investigated. Total OOM concentration in Beijing shows a clear dependence on UVB and temperature, suggesting the importance of photo-oxidation and temperature on OOM formation. In comparison to other atmospheric sites, total OOM concentration in Beijing (2.3×10^7 - 1.6×10^8 cm⁻³) is generally comparable to urban and suburban areas, and is clearly lower than those measured in forested areas. In the case of composition, most OOMs have 5 to 10 carbon atoms, 3 to 7 effective oxygen atoms, 0 to 2 nitrogen atoms and 0 to 6 DBE values. The seasonal variation of average effective oxygen atom follows the same trend as the overall atmospheric oxidation capacity, being the highest in summer and the lowest in winter and autumn. While for nitrogen and DBE distribution, there are not too much difference among seasons disregarding isoprene OOMs, indicating that the dominant formation pathways of each source-classified OOMs stay constant during the year.

With a revised workflow, we further separate OOMs into isoprene, monoterpene, aromatic, and aliphatic OOMs. For relative abundance, aromatic (29-41 %) and aliphatic OOMs (26-41 %) are major contributors throughout the year, suggesting that OOMs in urban atmospheric environment are controlled by anthropogenic activities. In addition, isoprene OOMs play an important role in summer and their fraction reaches to 33 %, indicating that biogenic sources are also large contributors to total OOMs in warmer seasons. The concentration of isoprene OOMs

($0.2\text{--}5.3\times 10^7\text{ cm}^{-3}$) and monoterpene OOMs ($1.1\text{--}8.4\times 10^6\text{ cm}^{-3}$) are smaller than those in forest areas, and they exhibit higher nitrogen and lower oxygen content compared with other cleaner sites. One recent study (Nie et al., 2022) reported that the composition of wintertime OOMs among four Chinese megacities, including Beijing, were similar. Our study further demonstrates that the composition of summertime OOMs between Beijing and Nanjing also have strong resemblance. Consequently, the seasonal characteristics of Beijing OOMs in this study could be representative of OOMs in other Chinese metropolises.

In terms of volatility, monoterpene OOMs are the most condensable, isoprene OOMs are the most volatile, and aromatic OOMs are more condensable than aliphatic ones. Based on the volatility and concentration characteristics of the four source-classified OOMs, an aerosol growth model was utilized to calculate their contribution to SOA growth. Results show that the condensation flux of total OOMs ($0.30\text{--}0.64\text{ }\mu\text{g}\cdot\text{m}^{-3}\cdot\text{h}^{-1}$) are high enough to produce a considerable amount of SOA within a day, and that aromatic (46–62 %) and aliphatic (14–32 %) OOMs are found to be dominant contributors regardless of seasons. This suggests that the formation of SOA in urban cities are likely driven by OOMs from anthropogenic sources, and highlights the importance of reducing anthropogenic emissions, especially aromatics, for pollution mitigation.

Data and materials availability: Data and materials are available upon contacting the first author and corresponding author.

Author contributions: CY and YG designed the study and wrote the manuscript. YG, YL, FZ, Ying Zhang, Ying Zhou, CL, XF, ZL, ZF, Yusheng Zhang, PZ and LT conducted the measurement and collected the data. CY, WN, ZW, DH, XQ, YL, YG, PZ and LT built the workflow and contributed to the aerosol dynamic model. JJ and VMK modified the manuscript. And all co-authors have read and commented on the manuscript.

Competing interests: The authors declare no competing interest.

Acknowledgements: This work was supported by the National Natural Science Foundation of China (NSFC) project (41875175, 42075101 and 22188102) and Samsung PM_{2.5} SRP. Kaspar R. Daellenbach acknowledges support by the Swiss National Science Foundation Ambizione grant PZPGP2_201992. Heikki Junninen is acknowledged for providing the tofTool package used for processing LTOF-CIMS data.

539 References

- 540 Aalto, P., Hämeri, K., Becker, E., Weber, R., Salm, J., Mäkelä, J. M., Hoell, C., O'dowd, C. D., Hansson, H.-C.,
 541 Väkevä, M., Koponen, I. K., Buzorius, G., and Kulmala, M.: Physical characterization of aerosol particles during
 542 nucleation events, *Tellus B: Chemical and Physical Meteorology*, 53, 344-358, 10.3402/tellusb.v53i4.17127, 2001.
- 543 Berndt, T., Richters, S., Jokinen, T., Hyttinen, N., Kurtén, T., Otkjær, R. V., Kjaergaard, H. G., Stratmann, F.,
 544 Herrmann, H., Sipilä, M., Kulmala, M., and Ehn, M.: Hydroxyl radical-induced formation of highly oxidized
 545 organic compounds, *Nature Communications*, 7, 13677, 10.1038/ncomms13677, 2016.
- 546 Berndt, T., Mentler, B., Scholz, W., Fischer, L., Herrmann, H., Kulmala, M., and Hansel, A.: Accretion Product
 547 Formation from Ozonolysis and OH Radical Reaction of α -Pinene: Mechanistic Insight and the Influence of
 548 Isoprene and Ethylene, *Environmental Science & Technology*, 52, 10.1021/acs.est.8b02210, 2018.
- 549 Bianchi, F., Tröstl, J., Junninen, H., Frege, C., Henne, S., Hoyle, C. R., Molteni, U., Herrmann, E., Adamov, A.,
 550 Bukowiecki, N., Chen, X., Duplissy, J., Gysel, M., Hutterli, M., Kangasluoma, J., Kontkanen, J., Kürten, A.,
 551 Manninen, H. E., Münch, S., Peräkylä, O., Petäjä, T., Rondo, L., Williamson, C., Weingartner, E., Curtius, J.,
 552 Worsnop, D. R., Kulmala, M., Dommen, J., and Baltensperger, U.: New particle formation in the free troposphere:
 553 A question of chemistry and timing, *Science*, 352, 1109-1112, 10.1126/science.aad5456, 2016.
- 554 Bianchi, F., Garmash, O., He, X., Yan, C., Iyer, S., Rosendahl, I., Xu, Z., Rissanen, M. P., Riva, M., Taipale, R.,
 555 Sarnela, N., Petäjä, T., Worsnop, D. R., Kulmala, M., Ehn, M., and Junninen, H.: The role of highly oxygenated
 556 molecules (HOMs) in determining the composition of ambient ions in the boreal forest, *Atmos. Chem. Phys.*, 17,
 557 13819-13831, 10.5194/acp-17-13819-2017, 2017.
- 558 Bianchi, F., Kurten, T., Riva, M., Mohr, C., Rissanen, M. P., Roldin, P., Berndt, T., Crounse, J. D., Wennberg, P.
 559 O., Mentel, T. F., Wildt, J., Junninen, H., Jokinen, T., Kulmala, M., Worsnop, D. R., Thornton, J. A., Donahue, N.,
 560 Kjaergaard, H. G., and Ehn, M.: Highly Oxygenated Organic Molecules (HOM) from Gas-Phase Autoxidation
 561 Involving Peroxy Radicals: A Key Contributor to Atmospheric Aerosol, *Chemical Reviews*, 119, 3472-3509,
 562 10.1021/acs.chemrev.8b00395, 2019.
- 563 Boyd, C. M., Sanchez, J., Xu, L., Eugene, A. J., Nah, T., Tuet, W. Y., Guzman, M. I., and Ng, N. L.: Secondary
 564 organic aerosol formation from the β -pinene+NO₃ system: effect of humidity and peroxy radical fate, *Atmos. Chem.*
 565 *Phys.*, 15, 7497-7522, 10.5194/acp-15-7497-2015, 2015.
- 566 Brean, J., Harrison, R. M., Shi, Z., Beddows, D. C. S., Acton, W. J. F., Hewitt, C. N., Squires, F. A., and Lee, J.:
 567 Observations of highly oxidized molecules and particle nucleation in the atmosphere of Beijing, *Atmos. Chem.*
 568 *Phys.*, 19, 14933-14947, 10.5194/acp-19-14933-2019, 2019.
- 569 Caudillo, L., Rörup, B., Heinritzi, M., Marie, G., Simon, M., Wagner, A. C., Müller, T., Granzin, M., Amorim, A.,
 570 Ataei, F., Baalbaki, R., Bertozzi, B., Brasseur, Z., Chiu, R., Chu, B., Dada, L., Duplissy, J., Finkenzeller, H.,
 571 Gonzalez Carracedo, L., He, X. C., Hofbauer, V., Kong, W., Lamkaddam, H., Lee, C. P., Lopez, B., Mahfouz, N.
 572 G. A., Makhmutov, V., Manninen, H. E., Marten, R., Massabò, D., Mauldin, R. L., Mentler, B., Molteni, U., Onnela,
 573 A., Pfeifer, J., Philippov, M., Piedehierro, A. A., Schervish, M., Scholz, W., Schulze, B., Shen, J., Stolzenburg, D.,
 574 Stozhkov, Y., Surdu, M., Tauber, C., Tham, Y. J., Tian, P., Tomé, A., Vogt, S., Wang, M., Wang, D. S., Weber, S.
 575 K., Welti, A., Yonghong, W., Yusheng, W., Zauner-Wieczorek, M., Baltensperger, U., El Haddad, I., Flagan, R. C.,
 576 Hansel, A., Höhler, K., Kirkby, J., Kulmala, M., Lehtipalo, K., Möhler, O., Saathoff, H., Volkamer, R., Winkler, P.
 577 M., Donahue, N. M., Kürten, A., and Curtius, J.: Chemical composition of nanoparticles from α -pinene nucleation
 578 and the influence of isoprene and relative humidity at low temperature, *Atmos. Chem. Phys. Discuss.*, 2021, 1-26,
 579 10.5194/acp-2021-512, 2021.

Chen, D., Xia, L., Guo, X., Lang, J., Zhou, Y., Wei, L., and Fu, X.: Impact of inter-annual meteorological variation from 2001 to 2015 on the contribution of regional transport to PM_{2.5} in Beijing, China, *Atmospheric Environment*, 260, 118545, <https://doi.org/10.1016/j.atmosenv.2021.118545>, 2021.

Chen, D., Jin, X., Fu, X., Xia, L., Guo, X., Lang, J., Zhou, Y., and Wei, W.: Impact of Inter-Annual Variation in Meteorology from 2010 to 2019 on the Inter-City Transport of PM_{2.5} in the Beijing–Tianjin–Hebei Region, *Sustainability*, 14, 10.3390/su14106210, 2022.

Cheng, X., Li, H., Zhang, Y., Li, Y., Zhang, W., Wang, X., Bi, F., Zhang, H., Gao, J., Chai, F., Lun, X., Chen, Y., Gao, J., and Lv, J.: Atmospheric isoprene and monoterpenes in a typical urban area of Beijing: Pollution characterization, chemical reactivity and source identification, *Journal of Environmental Sciences*, 71, 150-167, <https://doi.org/10.1016/j.jes.2017.12.017>, 2018.

Dada, L., Paasonen, P., Nieminen, T., Buenrostro Mazon, S., Kontkanen, J., Peräkylä, O., Lehtipalo, K., Hussein, T., Petäjä, T., Kerminen, V. M., Bäck, J., and Kulmala, M.: Long-term analysis of clear-sky new particle formation events and nonevents in Hyytiälä, *Atmos. Chem. Phys.*, 17, 6227-6241, 10.5194/acp-17-6227-2017, 2017.

Donahue, N. M., Ortega, I. K., Chuang, W., Riipinen, I., Riccobono, F., Schobesberger, S., Dommen, J., Baltensperger, U., Kulmala, M., Worsnop, D. R., and Vehkamäki, H.: How do organic vapors contribute to new-particle formation?, *Faraday Discussions*, 165, 91-104, 10.1039/C3FD00046J, 2013.

Drewnick, F., Hings, S. S., DeCarlo, P., Jayne, J. T., Gonin, M., Fuhrer, K., Weimer, S., Jimenez, J. L., Demerjian, K. L., Borrmann, S., and Worsnop, D. R.: A New Time-of-Flight Aerosol Mass Spectrometer (TOF-AMS)—Instrument Description and First Field Deployment, *Aerosol Science and Technology*, 39, 637-658, 10.1080/02786820500182040, 2005.

Ehn, M., Junninen, H., Petäjä, T., Kurtén, T., Kerminen, V. M., Schobesberger, S., Manninen, H. E., Ortega, I. K., Vehkamäki, H., Kulmala, M., and Worsnop, D. R.: Composition and temporal behavior of ambient ions in the boreal forest, *Atmos. Chem. Phys.*, 10, 8513-8530, 10.5194/acp-10-8513-2010, 2010.

Ehn, M., Kleist, E., Junninen, H., Petäjä, T., Lönn, G., Schobesberger, S., Dal Maso, M., Trimborn, A., Kulmala, M., Worsnop, D. R., Wahner, A., Wildt, J., and Mentel, T. F.: Gas phase formation of extremely oxidized pinene reaction products in chamber and ambient air, *Atmos. Chem. Phys.*, 12, 5113-5127, 10.5194/acp-12-5113-2012, 2012.

Ehn, M., Thornton, J. A., Kleist, E., Sipilä, M., Junninen, H., Pullinen, I., Springer, M., Rubach, F., Tillmann, R., Lee, B., Lopez-Hilfiker, F., Andres, S., Acir, I.-H., Rissanen, M., Jokinen, T., Schobesberger, S., Kangasluoma, J., Kontkanen, J., Nieminen, T., Kurtén, T., Nielsen, L. B., Jørgensen, S., Kjaergaard, H. G., Canagaratna, M., Maso, M. D., Berndt, T., Petäjä, T., Wahner, A., Kerminen, V.-M., Kulmala, M., Worsnop, D. R., Wildt, J., and Mentel, T. F.: A large source of low-volatility secondary organic aerosol, *Nature*, 506, 476-479, 10.1038/nature13032, 2014.

Garmash, O., Rissanen, M. P., Pullinen, I., Schmitt, S., Kausiala, O., Tillmann, R., Zhao, D., Percival, C., Bannan, T. J., Priestley, M., Hallquist, A. M., Kleist, E., Kiendler-Scharr, A., Hallquist, M., Berndt, T., McFiggans, G., Wildt, J., Mentel, T., and Ehn, M.: Multi-generation OH oxidation as a source for highly oxygenated organic molecules from aromatics, *Atmospheric Chemistry and Physics*, 20, 515-537, 10.5194/acp-20-515-2020, 2020.

Gordon, H., Kirkby, J., Baltensperger, U., Bianchi, F., Breitenlechner, M., Curtius, J., Dias, A., Dommen, J., Donahue, N. M., Dunne, E. M., Duplissy, J., Ehrhart, S., Flagan, R. C., Frege, C., Fuchs, C., Hansel, A., Hoyle, C. R., Kulmala, M., Kürten, A., Lehtipalo, K., Makhmutov, V., Molteni, U., Rissanen, M. P., Stozhkov, Y., Tröstl, J., Tsagkogeorgas, G., Wagner, R., Williamson, C., Wimmer, D., Winkler, P. M., Yan, C., and Carslaw, K. S.: Causes and importance of new particle formation in the present-day and preindustrial atmospheres, *Journal of Geophysical Research: Atmospheres*, 122, 8739-8760, <https://doi.org/10.1002/2017JD026844>, 2017.

622 Guo, Y., Yan, C., Li, C., Ma, W., Feng, Z., Zhou, Y., Lin, Z., Dada, L., Stolzenburg, D., Yin, R., Kontkanen, J.,
 623 Daellenbach, K. R., Kangasluoma, J., Yao, L., Chu, B., Wang, Y., Cai, R., Bianchi, F., Liu, Y., and Kulmala, M.:
 624 Formation of nighttime sulfuric acid from the ozonolysis of alkenes in Beijing, *Atmos. Chem. Phys.*, 21, 5499-5511,
 625 10.5194/acp-21-5499-2021, 2021.

626 Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N.
 627 M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M.
 628 E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, T. F., Monod, A., Prevot, A. S. H.,
 629 Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary
 630 organic aerosol: current and emerging issues, *Atmospheric Chemistry and Physics*, 9, 5155-5236, 10.5194/acp-9-
 631 5155-2009, 2009.

632 Heinritzi, M., Simon, M., Steiner, G., Wagner, A. C., Kürten, A., Hansel, A., and Curtius, J.: Characterization of
 633 the mass-dependent transmission efficiency of a CIMS, *Atmos. Meas. Tech.*, 9, 1449-1460, 10.5194/amt-9-1449-
 634 2016, 2016.

635 Heinritzi, M., Dada, L., Simon, M., Stolzenburg, D., Wagner, A. C., Fischer, L., Ahonen, L. R., Amanatidis, S.,
 636 Baalbaki, R., Baccarini, A., Bauer, P. S., Baumgartner, B., Bianchi, F., Brilke, S., Chen, D., Chiu, R., Dias, A.,
 637 Dommen, J., Duplissy, J., Finkenzeller, H., Frege, C., Fuchs, C., Garmash, O., Gordon, H., Granzin, M., El Haddad,
 638 I., He, X., Helm, J., Hofbauer, V., Hoyle, C. R., Kangasluoma, J., Keber, T., Kim, C., Kürten, A., Lamkaddam, H.,
 639 Laurila, T. M., Lampilahti, J., Lee, C. P., Lehtipalo, K., Leiminger, M., Mai, H., Makhmutov, V., Manninen, H. E.,
 640 Marten, R., Mathot, S., Mauldin, R. L., Mentler, B., Molteni, U., Müller, T., Nie, W., Nieminen, T., Onnela, A.,
 641 Partoll, E., Passananti, M., Petäjä, T., Pfeifer, J., Pospisilova, V., Quéléver, L. L. J., Rissanen, M. P., Rose, C.,
 642 Schobesberger, S., Scholz, W., Scholze, K., Sipilä, M., Steiner, G., Stozhkov, Y., Tauber, C., Tham, Y. J., Vazquez-
 643 Pufleau, M., Virtanen, A., Vogel, A. L., Volkamer, R., Wagner, R., Wang, M., Weitz, L., Wimmer, D., Xiao, M.,
 644 Yan, C., Ye, P., Zha, Q., Zhou, X., Amorim, A., Baltensperger, U., Hansel, A., Kulmala, M., Tomé, A., Winkler,
 645 P. M., Worsnop, D. R., Donahue, N. M., Kirkby, J., and Curtius, J.: Molecular understanding of the suppression of
 646 new-particle formation by isoprene, *Atmos. Chem. Phys.*, 20, 11809-11821, 10.5194/acp-20-11809-2020, 2020.

647 Huang, W., Li, H., Sarnela, N., Heikkinen, L., Tham, Y. J., Mikkilä, J., Thomas, S. J., Donahue, N. M., Kulmala,
 648 M., and Bianchi, F.: Measurement report: Molecular composition and volatility of gaseous organic compounds in
 649 a boreal forest: from volatile organic compounds to highly oxygenated organic molecules, *Atmos. Chem. Phys.*
 650 *Discuss.*, 2020, 1-27, 10.5194/acp-2020-1257, 2020.

651 Hyttinen, N., Kupiainen-Määttä, O., Rissanen, M. P., Muuronen, M., Ehn, M., and Kurtén, T.: Modeling the
 652 Charging of Highly Oxidized Cyclohexene Ozonolysis Products Using Nitrate-Based Chemical Ionization, *The*
 653 *Journal of Physical Chemistry A*, 119, 6339-6345, 10.1021/acs.jpca.5b01818, 2015.

654 Hyttinen, N., Otkjær, R. V., Iyer, S., Kjaergaard, H. G., Rissanen, M. P., Wennberg, P. O., and Kurtén, T.:
 655 Computational Comparison of Different Reagent Ions in the Chemical Ionization of Oxidized Multifunctional
 656 Compounds, *The Journal of Physical Chemistry A*, 122, 269-279, 10.1021/acs.jpca.7b10015, 2018.

657 Jayne, J. T., Leard, D. C., Zhang, X., Davidovits, P., Smith, K. A., Kolb, C. E., and Worsnop, D. R.: Development
 658 of an Aerosol Mass Spectrometer for Size and Composition Analysis of Submicron Particles, *Aerosol Science and*
 659 *Technology*, 33, 49-70, 10.1080/027868200410840, 2000.

660 Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan,
 661 J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy,
 662 J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T.,
 663 Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., Dunlea, J.,

Huffman, J. A., Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimojo, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of Organic Aerosols in the Atmosphere, *Science*, 326, 1525, 10.1126/science.1180353, 2009.

Jokinen, T., Sipilä, M., Junninen, H., Ehn, M., Lönn, G., Hakala, J., Petäjä, T., Mauldin III, R. L., Kulmala, M., and Worsnop, D. R.: Atmospheric sulphuric acid and neutral cluster measurements using CI-API-TOF, *Atmos. Chem. Phys.*, 12, 4117-4125, 10.5194/acp-12-4117-2012, 2012.

Jokinen, T., Sipilä, M., Richters, S., Kerminen, V.-M., Paasonen, P., Stratmann, F., Worsnop, D., Kulmala, M., Ehn, M., Herrmann, H., and Berndt, T.: Rapid Autoxidation Forms Highly Oxidized RO₂ Radicals in the Atmosphere, *Angewandte Chemie International Edition*, 53, 14596-14600, <https://doi.org/10.1002/anie.201408566>, 2014.

Junninen, H., Ehn, M., Petäjä, T., Luosujärvi, L., Kotiaho, T., Kostianen, R., Rohner, U., Gonin, M., Fuhrer, K., Kulmala, M., and Worsnop, D. R.: A high-resolution mass spectrometer to measure atmospheric ion composition, *Atmos. Meas. Tech.*, 3, 1039-1053, 10.5194/amt-3-1039-2010, 2010.

Kirkby, J., Duplissy, J., Sengupta, K., Frege, C., Gordon, H., Williamson, C., Heinritzi, M., Simon, M., Yan, C., Almeida, J., Tröstl, J., Nieminen, T., Ortega, I. K., Wagner, R., Adamov, A., Amorim, A., Bernhammer, A.-K., Bianchi, F., Breitenlechner, M., Brilke, S., Chen, X., Craven, J., Dias, A., Ehrhart, S., Flagan, R. C., Franchin, A., Fuchs, C., Guida, R., Hakala, J., Hoyle, C. R., Jokinen, T., Junninen, H., Kangasluoma, J., Kim, J., Krapf, M., Kürten, A., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Mathot, S., Molteni, U., Onnela, A., Peräkylä, O., Piel, F., Petäjä, T., Praplan, A. P., Pringle, K., Rap, A., Richards, N. A. D., Riipinen, I., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Scott, C. E., Seinfeld, J. H., Sipilä, M., Steiner, G., Stozhkov, Y., Stratmann, F., Tomé, A., Virtanen, A., Vogel, A. L., Wagner, A. C., Wagner, P. E., Weingartner, E., Wimmer, D., Winkler, P. M., Ye, P., Zhang, X., Hansel, A., Dommen, J., Donahue, N. M., Worsnop, D. R., Baltensperger, U., Kulmala, M., Carslaw, K. S., and Curtius, J.: Ion-induced nucleation of pure biogenic particles, *Nature*, 533, 521-526, 10.1038/nature17953, 2016.

Krechmer, J. E., Coggon, M. M., Massoli, P., Nguyen, T. B., Crounse, J. D., Hu, W., Day, D. A., Tyndall, G. S., Henze, D. K., Rivera-Rios, J. C., Nowak, J. B., Kimmel, J. R., Mauldin, R. L., Stark, H., Jayne, J. T., Sipilä, M., Junninen, H., St. Clair, J. M., Zhang, X., Feiner, P. A., Zhang, L., Miller, D. O., Brune, W. H., Keutsch, F. N., Wennberg, P. O., Seinfeld, J. H., Worsnop, D. R., Jimenez, J. L., and Canagaratna, M. R.: Formation of Low Volatility Organic Compounds and Secondary Organic Aerosol from Isoprene Hydroxyhydroperoxide Low-NO Oxidation, *Environmental Science & Technology*, 49, 10330-10339, 10.1021/acs.est.5b02031, 2015.

Kulmala, M., Kontkanen, J., Junninen, H., Lehtipalo, K., Manninen, H. E., Nieminen, T., Petäjä, T., Sipilä, M., Schobesberger, S., Rantala, P., Franchin, A., Jokinen, T., Järvinen, E., Äijälä, M., Kangasluoma, J., Hakala, J., Aalto, P. P., Paasonen, P., Mikkilä, J., Vanhanen, J., Aalto, J., Hakola, H., Makkonen, U., Ruuskanen, T., Mauldin, R. L., 3rd, Duplissy, J., Vehkamäki, H., Bäck, J., Kortelainen, A., Riipinen, I., Kurtén, T., Johnston, M. V., Smith, J. N., Ehn, M., Mentel, T. F., Lehtinen, K. E., Laaksonen, A., Kerminen, V. M., and Worsnop, D. R.: Direct observations of atmospheric aerosol nucleation, *Science*, 339, 943-946, 10.1126/science.1227385, 2013.

Kürten, A., Bergen, A., Heinritzi, M., Leiminger, M., Lorenz, V., Piel, F., Simon, M., Sitals, R., Wagner, A. C., and Curtius, J.: Observation of new particle formation and measurement of sulfuric acid, ammonia, amines and highly oxidized organic molecules at a rural site in central Germany, *Atmos. Chem. Phys.*, 16, 12793-12813, 10.5194/acp-16-12793-2016, 2016.

706 Le Breton, M., Wang, Y., Hallquist, Å. M., Pathak, R. K., Zheng, J., Yang, Y., Shang, D., Glasius, M., Bannan, T.
 707 J., Liu, Q., Chan, C. K., Percival, C. J., Zhu, W., Lou, S., Topping, D., Wang, Y., Yu, J., Lu, K., Guo, S., Hu, M.,
 708 and Hallquist, M.: Online gas- and particle-phase measurements of organosulfates, organosulfonates and nitrooxy
 709 organosulfates in Beijing utilizing a FIGAERO ToF-CIMS, *Atmos. Chem. Phys.*, 18, 10355-10371, 10.5194/acp-
 710 18-10355-2018, 2018.

711 Lehtipalo, K., Yan, C., Dada, L., Bianchi, F., Xiao, M., Wagner, R., Stolzenburg, D., Ahonen, L., Amorim, A.,
 712 Baccarini, A., Bauer, P., Baumgartner, B., Bergen, A., Bernhammer, A.-K., Breitenlechner, M., Brilke, S., Buchholz,
 713 A., Mazon, S., Chen, D., and Worsnop, D.: Multicomponent new particle formation from sulfuric acid, ammonia,
 714 and biogenic vapors, *Science Advances*, 4, eaau5363, 10.1126/sciadv.aau5363, 2018.

715 Lelieveld, J., Evans, J. S., Fnais, M., Giannadaki, D., and Pozzer, A.: The contribution of outdoor air pollution
 716 sources to premature mortality on a global scale, *Nature*, 525, 367-371, 10.1038/nature15371, 2015.

717 Liu, Y., Yan, C., Feng, Z., Zheng, F., Fan, X., Zhang, Y., Li, C., Zhou, Y., Lin, Z., Guo, Y., Zhang, Y., Ma, L.,
 718 Zhou, W., Liu, Z., Dada, L., Dällenbach, K., Kontkanen, J., Cai, R., Chan, T., Chu, B., Du, W., Yao, L., Wang, Y.,
 719 Cai, J., Kangasluoma, J., Kokkonen, T., Kujansuu, J., Rusanen, A., Deng, C., Fu, Y., Yin, R., Li, X., Lu, Y., Liu,
 720 Y., Lian, C., Yang, D., Wang, W., Ge, M., Wang, Y., Worsnop, D. R., Junninen, H., He, H., Kerminen, V.-M.,
 721 Zheng, J., Wang, L., Jiang, J., Petäjä, T., Bianchi, F., and Kulmala, M.: Continuous and comprehensive atmospheric
 722 observations in Beijing: a station to understand the complex urban atmospheric environment, *Big Earth Data*, 4,
 723 295-321, 10.1080/20964471.2020.1798707, 2020.

724 Liu, Y., Nie, W., Li, Y., Ge, D., Liu, C., Xu, Z., Chen, L., Wang, T., Wang, L., Sun, P., Qi, X., Wang, J., Xu, Z.,
 725 Yuan, J., Yan, C., Zhang, Y., Huang, D., Wang, Z., Donahue, N. M., Worsnop, D., Chi, X., Ehn, M., and Ding, A.:
 726 Formation of condensable organic vapors from anthropogenic and biogenic volatile organic compounds (VOCs) is
 727 strongly perturbed by NO_x in eastern China, *Atmos. Chem. Phys.*, 21, 14789-14814, 10.5194/acp-21-14789-2021,
 728 2021.

729 Massoli, P., Stark, H., Canagaratna, M. R., Krechmer, J. E., Xu, L., Ng, N. L., Mauldin, R. L., Yan, C., Kimmel, J.,
 730 Misztal, P. K., Jimenez, J. L., Jayne, J. T., and Worsnop, D. R.: Ambient Measurements of Highly Oxidized Gas-
 731 Phase Molecules during the Southern Oxidant and Aerosol Study (SOAS) 2013, *ACS Earth and Space Chemistry*,
 732 2, 653-672, 10.1021/acsearthspacechem.8b00028, 2018.

733 Mehra, A., Canagaratna, M., Bannan, T. J., Worrall, S. D., Bacak, A., Priestley, M., Liu, D., Zhao, J., Xu, W., Sun,
 734 Y., Hamilton, J. F., Squires, F. A., Lee, J., Bryant, D. J., Hopkins, J. R., Elzein, A., Budisulistiorini, S. H., Cheng,
 735 X., Chen, Q., Wang, Y., Wang, L., Stark, H., Krechmer, J. E., Brean, J., Slater, E., Whalley, L., Heard, D., Ouyang,
 736 B., Acton, W. J. F., Hewitt, C. N., Wang, X., Fu, P., Jayne, J., Worsnop, D., Allan, J., Percival, C., and Coe, H.:
 737 Using highly time-resolved online mass spectrometry to examine biogenic and anthropogenic contributions to
 738 organic aerosol in Beijing, *Faraday Discussions*, 10.1039/D0FD00080A, 2021.

739 Merikanto, J., Spracklen, D. V., Mann, G. W., Pickering, S. J., and Carslaw, K. S.: Impact of nucleation on global
 740 CCN, *Atmos. Chem. Phys.*, 9, 8601-8616, 10.5194/acp-9-8601-2009, 2009.

741 Mo, Z., Shao, M., Wang, W., Liu, Y., Wang, M., and Lu, S.: Evaluation of biogenic isoprene emissions and their
 742 contribution to ozone formation by ground-based measurements in Beijing, China, *Science of The Total
 743 Environment*, 627, 1485-1494, <https://doi.org/10.1016/j.scitotenv.2018.01.336>, 2018.

744 Mohr, C., Thornton, J. A., Heitto, A., Lopez-Hilfiker, F. D., Lutz, A., Riipinen, I., Hong, J., Donahue, N. M.,
 745 Hallquist, M., Petäjä, T., Kulmala, M., and Yli-Juuti, T.: Molecular identification of organic vapors driving
 746 atmospheric nanoparticle growth, *Nature Communications*, 10, 4442, 10.1038/s41467-019-12473-2, 2019.

747 Molteni, U., Bianchi, F., Klein, F., El Haddad, I., Frege, C., Rossi, M. J., Dommen, J., and Baltensperger, U.:
 748 Formation of highly oxygenated organic molecules from aromatic compounds, *Atmos. Chem. Phys.*, 18, 1909-1921,
 749 10.5194/acp-18-1909-2018, 2018.
 750 Mutzel, A., Poulain, L., Berndt, T., Iinuma, Y., Rodigast, M., Böge, O., Richters, S., Spindler, G., Sipilä, M.,
 751 Jokinen, T., Kulmala, M., and Herrmann, H.: Highly Oxidized Multifunctional Organic Compounds Observed in
 752 Tropospheric Particles: A Field and Laboratory Study, *Environmental Science & Technology*, 49, 7754-7761,
 753 10.1021/acs.est.5b00885, 2015.
 754 Nah, T., Sanchez, J., Boyd, C., and Ng, N.: Photochemical Aging of α -pinene and β -pinene Secondary Organic
 755 Aerosol formed from Nitrate Radical Oxidation, *Environmental Science & Technology*, 50,
 756 10.1021/acs.est.5b04594, 2015.
 757 Nie, W., Yan, C., Huang, D. D., Wang, Z., Liu, Y., Qiao, X., Guo, Y., Tian, L., Zheng, P., Xu, Z., Li, Y., Xu, Z.,
 758 Qi, X., Sun, P., Wang, J., Zheng, F., Li, X., Yin, R., Dallenbach, K. R., Bianchi, F., Petäjä, T., Zhang, Y., Wang,
 759 M., Schervish, M., Wang, S., Qiao, L., Wang, Q., Zhou, M., Wang, H., Yu, C., Yao, D., Guo, H., Ye, P., Lee, S.,
 760 Li, Y. J., Liu, Y., Chi, X., Kerminen, V.-M., Ehn, M., Donahue, N. M., Wang, T., Huang, C., Kulmala, M., Worsnop,
 761 D., Jiang, J., and Ding, A.: Secondary organic aerosol formed by condensing anthropogenic vapours over China's
 762 megacities, *Nature Geoscience*, 15, 255-261, 10.1038/s41561-022-00922-5, 2022.
 763 O'Dowd, C. D., Aalto, P., Hmeri, K., Kulmala, M., and Hoffmann, T.: Atmospheric particles from organic vapours,
 764 *Nature*, 416, 497-498, 10.1038/416497a, 2002.
 765 Orlando, J. J., and Tyndall, G. S.: Laboratory studies of organic peroxy radical chemistry: an overview with
 766 emphasis on recent issues of atmospheric significance, *Chemical Society Reviews*, 41, 6294-6317,
 767 10.1039/C2CS35166H, 2012.
 768 Pan, S., and Wang, L.: Atmospheric Oxidation Mechanism of m-Xylene Initiated by OH Radical, *The Journal of*
 769 *Physical Chemistry A*, 118, 10778-10787, 10.1021/jp506815v, 2014.
 770 Panopoulou, A., Liakakou, E., Sauvage, S., Gros, V., Locoge, N., Stavroulas, I., Bonsang, B., Gerasopoulos, E.,
 771 and Mihalopoulos, N.: Yearlong measurements of monoterpenes and isoprene in a Mediterranean city (Athens):
 772 Natural vs anthropogenic origin, *Atmospheric Environment*, 243, 117803,
 773 <https://doi.org/10.1016/j.atmosenv.2020.117803>, 2020.
 774 Riccobono, F., Schobesberger, S., Scott, C., Dommen, J., Ortega, I., Rondo, L., Almeida, J., Amorim, A., Bianchi,
 775 F., Breitenlechner, M., David, A., Downard, A., Dunne, E., Duplissy, J., Ehrhart, S., Flagan, R., Franchin, A.,
 776 Hansel, A., Junninen, H., and Baltensperger, U.: Oxidation Products of Biogenic Emissions Contribute to
 777 Nucleation of Atmospheric Particles, *Science (New York, N.Y.)*, 344, 717-721, 10.1126/science.1243527, 2014.
 778 Riedel, T. P., Lin, Y.-H., Budisulistiorini, S. H., Gaston, C. J., Thornton, J. A., Zhang, Z., Vizuete, W., Gold, A.,
 779 and Surratt, J. D.: Heterogeneous Reactions of Isoprene-Derived Epoxides: Reaction Probabilities and Molar
 780 Secondary Organic Aerosol Yield Estimates, *Environmental Science & Technology Letters*, 2, 38-42,
 781 10.1021/ez500406f, 2015.
 782 Riva, M., Rantala, P., Krechmer, J. E., Peräkylä, O., Zhang, Y., Heikkinen, L., Garmash, O., Yan, C., Kulmala, M.,
 783 Worsnop, D., and Ehn, M.: Evaluating the performance of five different chemical ionization techniques for
 784 detecting gaseous oxygenated organic species, *Atmos. Meas. Tech.*, 12, 2403-2421, 10.5194/amt-12-2403-2019,
 785 2019.
 786 Roldin, P., Ehn, M., Kurtén, T., Olenius, T., Rissanen, M. P., Sarnela, N., Elm, J., Rantala, P., Hao, L., Hyttinen,
 787 N., Heikkinen, L., Worsnop, D. R., Pichelstorfer, L., Xavier, C., Clusius, P., Öström, E., Petäjä, T., Kulmala, M.,

788 Vehkamäki, H., Virtanen, A., Riipinen, I., and Boy, M.: The role of highly oxygenated organic molecules in the
 789 Boreal aerosol-cloud-climate system, *Nature Communications*, 10, 4370, 10.1038/s41467-019-12338-8, 2019.
 790 Rose, C., Zha, Q., Dada, L., Yan, C., Lehtipalo, K., Junninen, H., Mazon, S. B., Jokinen, T., Sarnela, N., Sipilä, M.,
 791 Petäjä, T., Kerminen, V.-M., Bianchi, F., and Kulmala, M.: Observations of biogenic ion-induced cluster formation
 792 in the atmosphere, *Science Advances*, 4, eaar5218, 10.1126/sciadv.aar5218, 2018.
 793 Schervish, M., and Donahue, N. M.: Peroxy radical chemistry and the volatility basis set, *Atmos. Chem. Phys.*, 20,
 794 1183-1199, 10.5194/acp-20-1183-2020, 2020.
 795 Schobesberger, S., Junninen, H., Bianchi, F., Lönn, G., Ehn, M., Lehtipalo, K., Dommen, J., Ehrhart, S., Ortega, I.
 796 K., Franchin, A., Nieminen, T., Riccobono, F., Hutterli, M., Duplissy, J., Almeida, J., Amorim, A., Breitenlechner,
 797 M., Downard, A. J., Dunne, E. M., Flagan, R. C., Kajos, M., Keskinen, H., Kirkby, J., Kupc, A., Kürten, A., Kurtén,
 798 T., Laaksonen, A., Mathot, S., Onnela, A., Praplan, A. P., Rondo, L., Santos, F. D., Schallhart, S., Schnitzhofer, R.,
 799 Sipilä, M., Tomé, A., Tsagkogeorgas, G., Vehkamäki, H., Wimmer, D., Baltensperger, U., Carslaw, K. S., Curtius,
 800 J., Hansel, A., Petäjä, T., Kulmala, M., Donahue, N. M., and Worsnop, D. R.: Molecular understanding of
 801 atmospheric particle formation from sulfuric acid and large oxidized organic molecules, *Proceedings of the National*
 802 *Academy of Sciences*, 110, 17223, 10.1073/pnas.1306973110, 2013.
 803 Seinfeld, J. H., and Pandis, S. N.: *Atmospheric chemistry and physics: from air pollution to climate change*, John
 804 Wiley & Sons, 2016.
 805 Song, K., Guo, S., Wang, H., Yu, Y., Wang, H., Tang, R., Xia, S., Gong, Y., Wan, Z., Lv, D., Tan, R., Zhu, W.,
 806 Shen, R., Li, X., Yu, X., Chen, S., Zeng, L., and Huang, X.: Measurement Report: Online Measurement of Gas-
 807 Phase Nitrated Phenols Utilizing CI-LToF-MS: Primary Sources and Secondary Formation, *Atmos. Chem. Phys.*
 808 *Discuss.*, 2021, 1-28, 10.5194/acp-2020-1294, 2021.
 809 Stocker, T.: *Climate change 2013: the physical science basis: Working Group I contribution to the Fifth assessment*
 810 *report of the Intergovernmental Panel on Climate Change*, 2014.
 811 Stolzenburg, D., Fischer, L., Vogel, A. L., Heinritzi, M., Schervish, M., Simon, M., Wagner, A. C., Dada, L.,
 812 Ahonen, L. R., Amorim, A., Baccarini, A., Bauer, P. S., Baumgartner, B., Bergen, A., Bianchi, F., Breitenlechner,
 813 M., Brilke, S., Buenrostro Mazon, S., Chen, D., Dias, A., Draper, D. C., Duplissy, J., El Haddad, I., Finkenzeller,
 814 H., Frege, C., Fuchs, C., Garmash, O., Gordon, H., He, X., Helm, J., Hofbauer, V., Hoyle, C. R., Kim, C., Kirkby,
 815 J., Kontkanen, J., Kürten, A., Lampilahti, J., Lawler, M., Lehtipalo, K., Leiminger, M., Mai, H., Mathot, S., Mentler,
 816 B., Molteni, U., Nie, W., Nieminen, T., Nowak, J. B., Ojdanic, A., Onnela, A., Passananti, M., Petäjä, T., Quéléver,
 817 L. L. J., Rissanen, M. P., Sarnela, N., Schallhart, S., Tauber, C., Tomé, A., Wagner, R., Wang, M., Weitz, L.,
 818 Wimmer, D., Xiao, M., Yan, C., Ye, P., Zha, Q., Baltensperger, U., Curtius, J., Dommen, J., Flagan, R. C., Kulmala,
 819 M., Smith, J. N., Worsnop, D. R., Hansel, A., Donahue, N. M., and Winkler, P. M.: Rapid growth of organic aerosol
 820 nanoparticles over a wide tropospheric temperature range, *Proceedings of the National Academy of Sciences*, 115,
 821 9122, 10.1073/pnas.1807604115, 2018.
 822 Tan, Q., Ge, B., Xu, X., Gan, L., Yang, W., Chen, X., Pan, X., Wang, W., Li, J., and Wang, Z.: Increasing impacts
 823 of the relative contributions of regional transport on air pollution in Beijing: Observational evidence, *Environmental*
 824 *Pollution*, 292, 118407, <https://doi.org/10.1016/j.envpol.2021.118407>, 2022.
 825 Teng, A. P., Crounse, J. D., and Wennberg, P. O.: Isoprene Peroxy Radical Dynamics, *Journal of the American*
 826 *Chemical Society*, 139, 5367-5377, 10.1021/jacs.6b12838, 2017.
 827 Tröstl, J., Chuang, W. K., Gordon, H., Heinritzi, M., Yan, C., Molteni, U., Ahlm, L., Frege, C., Bianchi, F., Wagner,
 828 R., Simon, M., Lehtipalo, K., Williamson, C., Craven, J. S., Duplissy, J., Adamov, A., Almeida, J., Bernhammer,
 829 A.-K., Breitenlechner, M., Brilke, S., Dias, A., Ehrhart, S., Flagan, R. C., Franchin, A., Fuchs, C., Guida, R., Gysel,

830 M., Hansel, A., Hoyle, C. R., Jokinen, T., Junninen, H., Kangasluoma, J., Keskinen, H., Kim, J., Krapf, M., Kürten,
831 A., Laaksonen, A., Lawler, M., Leiminger, M., Mathot, S., Möhler, O., Nieminen, T., Onnela, A., Petäjä, T., Piel,
832 F. M., Miettinen, P., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Sengupta, K., Sipilä, M., Smith, J.
833 N., Steiner, G., Tomè, A., Virtanen, A., Wagner, A. C., Weingartner, E., Wimmer, D., Winkler, P. M., Ye, P.,
834 Carslaw, K. S., Curtius, J., Dommen, J., Kirkby, J., Kulmala, M., Riipinen, I., Worsnop, D. R., Donahue, N. M.,
835 and Baltensperger, U.: The role of low-volatility organic compounds in initial particle growth in the atmosphere,
836 *Nature*, 533, 527-531, 10.1038/nature18271, 2016.

837 Wagner, P., and Kuttler, W.: Biogenic and anthropogenic isoprene in the near-surface urban atmosphere — A case
838 study in Essen, Germany, *Science of The Total Environment*, 475, 104-115,
839 <https://doi.org/10.1016/j.scitotenv.2013.12.026>, 2014.

840 Wang, H., Lu, K., Guo, S., Wu, Z., Shang, D., Tan, Z., Wang, Y., Le Breton, M., Lou, S., Tang, M., Wu, Y., Zhu,
841 W., Zheng, J., Zeng, L., Hallquist, M., Hu, M., and Zhang, Y.: Efficient N₂O₅ uptake and NO₃ oxidation in the
842 outflow of urban Beijing, *Atmos. Chem. Phys.*, 18, 9705-9721, 10.5194/acp-18-9705-2018, 2018a.

843 Wang, L., Liu, Z., Sun, Y., Ji, D., and Wang, Y.: Long-range transport and regional sources of PM_{2.5} in Beijing
844 based on long-term observations from 2005 to 2010, *Atmospheric Research*, 157, 37-48,
845 <https://doi.org/10.1016/j.atmosres.2014.12.003>, 2015.

846 Wang, S., Wu, R., Berndt, T., Ehn, M., and Wang, L.: Formation of Highly Oxidized Radicals and Multifunctional
847 Products from the Atmospheric Oxidation of Alkylbenzenes, *Environmental Science & Technology*, 51, 8442-8449,
848 10.1021/acs.est.7b02374, 2017.

849 Wang, S., Riva, M., Yan, C., Ehn, M., and Wang, L.: Primary Formation of Highly Oxidized Multifunctional
850 Products in the OH-Initiated Oxidation of Isoprene: A Combined Theoretical and Experimental Study,
851 *Environmental Science & Technology*, 52, 12255-12264, 10.1021/acs.est.8b02783, 2018b.

852 Wang, Y., Hu, M., Wang, Y., Zheng, J., Shang, D., Yang, Y., Liu, Y., Li, X., Tang, R., Zhu, W., Du, Z., Wu, Y.,
853 Guo, S., Wu, Z., Lou, S., Hallquist, M., and Yu, J. Z.: The formation of nitro-aromatic compounds under high NO_x
854 and anthropogenic VOC conditions in urban Beijing, China, *Atmos. Chem. Phys.*, 19, 7649-7665, 10.5194/acp-19-
855 7649-2019, 2019.

856 Wang, Z., Ehn, M., Rissanen, M. P., Garmash, O., Quéléver, L., Xing, L., Monge-Palacios, M., Rantala, P.,
857 Donahue, N. M., Berndt, T., and Sarathy, S. M.: Efficient alkane oxidation under combustion engine and
858 atmospheric conditions, *Communications Chemistry*, 4, 18, 10.1038/s42004-020-00445-3, 2021.

859 Wu, R., Vereecken, L., Tsiligiannis, E., Kang, S., Albrecht, S. R., Hantschke, L., Zhao, D., Novelli, A., Fuchs, H.,
860 Tillmann, R., Hohaus, T., Carlsson, P. T. M., Shenolikar, J., Bernard, F., Crowley, J. N., Fry, J. L., Brownwood, B.,
861 Thornton, J. A., Brown, S. S., Kiendler-Scharr, A., Wahner, A., Hallquist, M., and Mentel, T. F.: Molecular
862 composition and volatility of multi-generation products formed from isoprene oxidation by nitrate radical, *Atmos.*
863 *Chem. Phys.*, 21, 10799-10824, 10.5194/acp-21-10799-2021, 2021.

864 Xu, L., Møller, K. H., Crounse, J. D., Kjaergaard, H. G., and Wennberg, P. O.: New Insights into the Radical
865 Chemistry and Product Distribution in the OH-Initiated Oxidation of Benzene, *Environmental Science &*
866 *Technology*, 54, 13467-13477, 10.1021/acs.est.0c04780, 2020.

867 Xu, Z. N., Nie, W., Liu, Y. L., Sun, P., Huang, D. D., Yan, C., Krechmer, J., Ye, P. L., Xu, Z., Qi, X. M., Zhu, C.
868 J., Li, Y. Y., Wang, T. Y., Wang, L., Huang, X., Tang, R. Z., Guo, S., Xiu, G. L., Fu, Q. Y., Worsnop, D., Chi, X.
869 G., and Ding, A. J.: Multifunctional Products of Isoprene Oxidation in Polluted Atmosphere and Their Contribution
870 to SOA, *Geophysical Research Letters*, 48, e2020GL089276, <https://doi.org/10.1029/2020GL089276>, 2021.

871 Yan, C., Nie, W., Äijälä, M., Rissanen, M. P., Canagaratna, M. R., Massoli, P., Junninen, H., Jokinen, T., Sarnela,
 872 N., Häme, S. A. K., Schobesberger, S., Canonaco, F., Yao, L., Prévôt, A. S. H., Petäjä, T., Kulmala, M., Sipilä, M.,
 873 Worsnop, D. R., and Ehn, M.: Source characterization of highly oxidized multifunctional compounds in a boreal
 874 forest environment using positive matrix factorization, *Atmos. Chem. Phys.*, 16, 12715-12731, 10.5194/acp-16-
 875 12715-2016, 2016.

876 Yan, C., Nie, W., Vogel, A. L., Dada, L., Lehtipalo, K., Stolzenburg, D., Wagner, R., Rissanen, M. P., Xiao, M.,
 877 Ahonen, L., Fischer, L., Rose, C., Bianchi, F., Gordon, H., Simon, M., Heinritzi, M., Garmash, O., Roldin, P., Dias,
 878 A., Ye, P., Hofbauer, V., Amorim, A., Bauer, P. S., Bergen, A., Bernhammer, A. K., Breitenlechner, M., Brilke, S.,
 879 Buchholz, A., Mazon, S. B., Canagaratna, M. R., Chen, X., Ding, A., Dommen, J., Draper, D. C., Duplissy, J.,
 880 Frege, C., Heyn, C., Guida, R., Hakala, J., Heikkinen, L., Hoyle, C. R., Jokinen, T., Kangasluoma, J., Kirkby, J.,
 881 Kontkanen, J., Kürten, A., Lawler, M. J., Mai, H., Mathot, S., Mauldin, R. L., Molteni, U., Nichman, L., Nieminen,
 882 T., Nowak, J., Ojdanic, A., Onnela, A., Pajunoja, A., Petäjä, T., Piel, F., Quéléver, L. L. J., Sarnela, N., Schallhart,
 883 S., Sengupta, K., Sipilä, M., Tomé, A., Tröstl, J., Väisänen, O., Wagner, A. C., Ylisirniö, A., Zha, Q., Baltensperger,
 884 U., Carslaw, K. S., Curtius, J., Flagan, R. C., Hansel, A., Riipinen, I., Smith, J. N., Virtanen, A., Winkler, P. M.,
 885 Donahue, N. M., Kerminen, V. M., Kulmala, M., Ehn, M., and Worsnop, D. R.: Size-dependent influence of NO_x
 886 on the growth rates of organic aerosol particles, *Science Advances*, 6, eaay4945, 10.1126/sciadv.aay4945, 2020.

887 Yan, C., Yin, R., Lu, Y., Dada, L., Yang, D., Fu, Y., Kontkanen, J., Deng, C., Garmash, O., Ruan, J., Baalbaki, R.,
 888 Schervish, M., Cai, R., Bloss, M., Chan, T., Chen, T., Chen, Q., Chen, X., Chen, Y., Chu, B., Dällenbach, K.,
 889 Foreback, B., He, X., Heikkinen, L., Jokinen, T., Junninen, H., Kangasluoma, J., Kokkonen, T., Kurppa, M.,
 890 Lehtipalo, K., Li, H., Li, H., Li, X., Liu, Y., Ma, Q., Paasonen, P., Rantala, P., Pileci, R. E., Rusanen, A., Sarnela,
 891 N., Simonen, P., Wang, S., Wang, W., Wang, Y., Xue, M., Yang, G., Yao, L., Zhou, Y., Kujansuu, J., Petäjä, T.,
 892 Nie, W., Ma, Y., Ge, M., He, H., Donahue, N. M., Worsnop, D. R., Veli-Matti, K., Wang, L., Liu, Y., Zheng, J.,
 893 Kulmala, M., Jiang, J., and Bianchi, F.: The Synergistic Role of Sulfuric Acid, Bases, and Oxidized Organics
 894 Governing New-Particle Formation in Beijing, *Geophysical Research Letters*, 48, e2020GL091944,
 895 <https://doi.org/10.1029/2020GL091944>, 2021.

896 Ye, C., Yuan, B., Lin, Y., Wang, Z., Hu, W., Li, T., Chen, W., Wu, C., Wang, C., Huang, S., Qi, J., Wang, B.,
 897 Wang, C., Song, W., Wang, X., Zheng, E., Krechmer, J. E., Ye, P., Zhang, Z., Wang, X., Worsnop, D. R., and Shao,
 898 M.: Chemical characterization of oxygenated organic compounds in gas-phase and particle-phase using iodide-
 899 CIMS with FIGAERO in urban air, *Atmos. Chem. Phys. Discuss.*, 2020, 1-62, 10.5194/acp-2020-1187, 2020.

900 Zaytsev, A., Koss, A. R., Breitenlechner, M., Krechmer, J. E., Nihill, K. J., Lim, C. Y., Rowe, J. C., Cox, J. L.,
 901 Moss, J., Roscioli, J. R., Canagaratna, M. R., Worsnop, D. R., Kroll, J. H., and Keutsch, F. N.: Mechanistic study
 902 of the formation of ring-retaining and ring-opening products from the oxidation of aromatic compounds under urban
 903 atmospheric conditions, *Atmos. Chem. Phys.*, 19, 15117-15129, 10.5194/acp-19-15117-2019, 2019.

904 Zha, Q., Yan, C., Junninen, H., Riva, M., Sarnela, N., Aalto, J., Quéléver, L., Schallhart, S., Dada, L., Heikkinen,
 905 L., Peräkylä, O., Zou, J., Rose, C., Wang, Y., Mammarella, I., Katul, G., Vesala, T., Worsnop, D. R., Kulmala, M.,
 906 Petäjä, T., Bianchi, F., and Ehn, M.: Vertical characterization of highly oxygenated molecules (HOMs) below and
 907 above a boreal forest canopy, *Atmos. Chem. Phys.*, 18, 17437-17450, 10.5194/acp-18-17437-2018, 2018.

908 Zhang, H., Li, H., Zhang, Q., Zhang, Y., Zhang, W., Wang, X., Bi, F., Chai, F., Gao, J., Meng, L., Yang, T., Chen,
 909 Y., Cheng, Q., and Xia, F.: Atmospheric Volatile Organic Compounds in a Typical Urban Area of Beijing: Pollution
 910 Characterization, Health Risk Assessment and Source Apportionment, *Atmosphere*, 8, 10.3390/atmos8030061,
 911 2017.

912 Zhang, H., Zhang, Y., Huang, Z., Acton, W. J. F., Wang, Z., Nemitz, E., Langford, B., Mullinger, N., Davison, B.,
 913 Shi, Z., Liu, D., Song, W., Yang, W., Zeng, J., Wu, Z., Fu, P., Zhang, Q., and Wang, X.: Vertical profiles of biogenic
 914 volatile organic compounds as observed online at a tower in Beijing, *Journal of Environmental Sciences*, 95, 33-42,
 915 <https://doi.org/10.1016/j.jes.2020.03.032>, 2020.
 916 Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami, A.,
 917 Middlebrook, A. M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P. F., Salcedo, D., Onasch, T.,
 918 Jayne, J. T., Miyoshi, T., Shimojo, A., Hatakeyama, S., Takegawa, N., Kondo, Y., Schneider, J., Drewnick, F.,
 919 Borrmann, S., Weimer, S., Demerjian, K., Williams, P., Bower, K., Bahreini, R., Cottrell, L., Griffin, R. J.,
 920 Rautiainen, J., Sun, J. Y., Zhang, Y. M., and Worsnop, D. R.: Ubiquity and dominance of oxygenated species in
 921 organic aerosols in anthropogenically-influenced Northern Hemisphere midlatitudes, *Geophysical Research Letters*,
 922 34, <https://doi.org/10.1029/2007GL029979>, 2007.
 923 Zhang, Y.-Q., Ding, X., He, Q.-F., Wen, T.-X., Wang, J.-Q., Yang, K., Jiang, H., Cheng, Q., Liu, P., Wang, Z.-R.,
 924 He, Y.-F., Hu, W.-W., Wang, Q.-Y., Xin, J.-Y., Wang, Y.-S., and Wang, X.-M.: Observational Insights into
 925 Isoprene Secondary Organic Aerosol Formation through the Epoxide Pathway at Three Urban Sites from Northern
 926 to Southern China, *Environmental Science & Technology*, 56, 4795-4805, 10.1021/acs.est.1c06974, 2022.
 927 Zhang, Y., Peräkylä, O., Yan, C., Heikkinen, L., Äijälä, M., Daellenbach, K. R., Zha, Q., Riva, M., Garmash, O.,
 928 Junninen, H., Paatero, P., Worsnop, D., and Ehn, M.: A novel approach for simple statistical analysis of high-
 929 resolution mass spectra, *Atmos. Meas. Tech.*, 12, 3761-3776, 10.5194/amt-12-3761-2019, 2019.
 930 Zhao, Y., Thornton, J. A., and Pye, H. O. T.: Quantitative constraints on autoxidation and dimer formation from
 931 direct probing of monoterpene-derived peroxy radical chemistry, *Proceedings of the National Academy of Sciences*,
 932 115, 12142, 10.1073/pnas.1812147115, 2018.

933