

Long-term Variability in Immersion-mode Marine Ice Nucleating Particles from Climate Model Simulations and Observations

Aishwarya Raman¹, Thomas Hill², Paul J. DeMott², Balwinder Singh¹, Kai Zhang¹, Po-Lun Ma¹, Mingxuan Wu¹, Hailong Wang¹, Simon P. Alexander^{3,4}, and Susannah M. Burrows¹

¹Pacific Northwest National Laboratory, 902 Battelle Blvd, Richland, WA 99354

²Department of Atmospheric Science, Colorado State University, Fort Collins, CO 80523-1371, USA

³Australian Antarctic Division, Kingston, Tasmania, Australia

⁴Australian Antarctic Program Partnership, Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania, Australia

Correspondence: Aishwarya Raman(aishwarya.raman@pnnl.gov)

Table S1. Descriptions of aerosol aware INP parameterizations

INP parameterization	Aerosol property	Study area	Sample type	Limitations
D15	Number concentrations of dust particles with diameters greater than 500 nm [m^{-3}]	Saharan and Asian deserts	Mineral dust particles larger than $0.5 \mu\text{m}$	Does not distinguish between different dust mineralogies.
W15	Total organic carbon mass concentration [g m^{-3}]	Arctic (July–August 2013)	Marine Organic Aerosol (MOA)	Does not include the differences in the emission and atmospheric chemistry of INPs
M18	Sea spray aerosol surface area concentration [$\text{m}^{-2} \text{m}^{-3}$]	Mace Head station (August 2015)	Sea salt aerosol	between the airborne sea-spray (SSA) and sea surface microlayer water samples ; Entrainment of multiple INP species into a single SSA particle is not considered.
M18+D15	Sea spray aerosol surface area concentration [$\text{m}^{-2} \text{m}^{-3}$] and number concentrations of dust particles [m^{-3}] with diameters greater than 500 nm	Mace Head (August 2015) and South ern Ocean (March-April, 2016)	Dust and Sea salt	Derived for clean marine conditions. Only background sea-spray INPs are captured. Represents only background INPs. Potential variations in marine INPs due to ocean biology or dust mineralogy are not captured.




Figure S1. Climatology of dust concentrations from ground stations in the Northern Hemisphere compared against E3SM aerosol climatology. For the model, monthly average dust concentrations are shown for the period 2016–2018. Error bars in the model represent standard deviation of aerosol mass concentrations for 2016–2018. Error bars in the observations for each ground station represent standard deviation of measurements. Both CTL and EXP are shown for comparison with observations.




Figure S2. Climatology of sea salt concentrations from ground stations in the Northern Hemisphere compared against E3SM aerosol climatology. For the model, monthly average sea salt concentrations are shown for the period 2016–2018. Error bars in the model represent standard deviation of aerosol mass concentrations for 2016–2018. Error bars in the observations for each ground station represent standard deviation of measurements. Both CTL and EXP are shown for comparison with observations.




Figure S3. Differences in ratios of CNT and D15 INP parameterizations between -20°C and -28°C




Figure S4. Comparison of INP dependence on temperature in CNT and D15 INP parameterizations from E3SM simulations. Panel (a) shows INPs simulated by CNT dust for temperatures -28.5°C (black), -28°C (magenta), -20.5°C (green), -20°C (blue), -16.5°C (grey), -16°C (red). For a given temperature, different data points shown here correspond to the days during the MICRE campaign. Panel (b) shows the INP temperature spectra for D15.



Figure S5. (Top) Aerial photograph of Macquarie Island isthmus, with the location of MICRE and RSV Aurora Australis (on which MARCUS was located) indicated. (Bottom) MICRE campaign filter on Macquarie Island. Waves, kelp and tussock upwind (toward the ocean) over which air must pass before reaching the filter are also visible in the photo. Photo credits: Andrew Klekociuk, Australian Antarctic Division.




Figure S6. Observed INP concentrations (L^{-1}) at Macquarie Island from MICRE and simulated INPs using the updated dry deposition parameterization in E3SMv1. (a) and (b) : D15, (c) and (d) : M18, (e) and (f) : M18+D15, (g) and (h) : W15. INP concentrations are colored by activation temperatures used for measurements. Solid line in each panel represents 1:1 comparison, while dashed lines represent a factor of 2 and 10 from the observations. Error metrics in each panel include fractional gross error (FGE), modified normalized mean bias (MNMB), spearman correlation (R), percentage of model INPs within a factor of two from observations (2x), and percentage of model INPs within a factor of 10 from observations (10x). Scatter plots for austral summer (October-February) are shown in the left column and those for austral winter (March - September) are shown in the right column.