

Significant formation of sulfate aerosols contributed by the heterogeneous drivers of dust surface

Tao Wang¹, Yangyang Liu¹, Hanyun Cheng¹, Zhenzhen Wang¹, Hongbo Fu¹, Jianmin Chen¹, Liwu Zhang^{1,2*}

5 1 Shanghai Key Laboratory of Atmospheric Particle Pollution and Prevention, National Observations and Research Station for Wetland Ecosystems of the Yangtze Estuary, IRDR international Center of Excellence on Risk Interconnectivity and Governance on Weather, Department of Environmental Science & Engineering, Fudan University, Shanghai, 200433, Peoples' Republic of China

2 Shanghai Institute of Pollution Control and Ecological Security, Shanghai, 200092, Peoples' Republic of China

10 *Correspondence to:* Liwu Zhang (zhanglw@fudan.edu.cn)

Abstract. The importance of dust heterogeneous oxidation in the removal of atmospheric SO₂ and formation of sulfate aerosols is not adequately understood. In this study, the Fe, Ti, Al-bearing components, Na⁺, Cl⁻, K⁺, and Ca²⁺ of the dust surface were discovered to be closely associated with the heterogeneous formation of sulfate. Regression models were then developed to make a reliable prediction for the heterogeneous reactivity based on the particle chemical compositions. Further, the recognized gas-phase, aqueous-phase and heterogeneous oxidation routes were quantitatively assessed and kinetically compared by combining the laboratory work with modeling study. In the presence of 55 μg m⁻³ airborne dust, heterogeneous oxidation accounts for approximately 28.6% of the secondary sulfate aerosols during nighttime, while the proportion decreases to 13.1% in the presence of solar irradiation. On the dust surface, heterogeneous drivers (e.g. transition metal constituents, water-soluble ions) are more efficient than surface adsorbed oxidants (e.g. H₂O₂, NO₂, O₃) in the conversion of SO₂, particularly during nighttime. Dust heterogeneous oxidation offers an opportunity to explain the missing sulfate source during severe haze pollution events, and its contribution proportion in the complex atmospheric environments could be even higher than the current calculation results. Overall, the dust surface drivers are responsible for the significant formation of sulfate aerosols and have profound impacts on the atmospheric sulfur cycling.

15
20

1 Introduction

25 As an important component of atmospheric particulate matters, sulfate exerts profound impacts on the Earth's climate system, air quality, and public health (Seinfeld and Pandis, 2016; Wang et al., 2021a). The rapid formation of sulfate was proven as largely responsible for London Fog and Beijing Haze (Wang et al., 2016; Cheng et al., 2016). Secondary sulfate aerosols originate predominately from the conversion of sulfur dioxide (SO₂) via the gas-phase oxidation in gaseous environments, aqueous-phase oxidation in liquid media, and heterogeneous oxidation on aerosol surfaces (Ravishankara, 1997; Mauldin III et al., 2012; Su et al., 2020; Liu et al., 2021a). In recent years, the newly found sulfate formation pathways were kinetically compared with the documented ones to evaluate the relative importance of them (Cheng et al., 2016; Gen et al., 2019; Liu et al., 2020a; Wang et al., 2020a). In addition, the reported oxidation channels were compared with each other by aerosol observations or modeling investigations (Berglen et al., 2004; Sarwar et al., 2013; He et al., 2018; Ye et al., 2018; Fan et al., 2020; Tao et al., 2020; Zheng et al., 2020; Song et al., 2021; Tilgner et al., 2021; Liu et al., 2021b; Gao et al., 2022; Wang et al., 2022a; Ye et al., 2022). In general, these studies emphasized the importance of certain newly discovered aqueous-phase process or compared the contributions from the documented gas- and aqueous-phase pathways. Nevertheless, heterogeneous reaction was scarcely involved in discussion, thus hindering the deeper understanding of the atmospheric relevance of aerosol surfaces.

Heterogeneous reaction alters the concentrations of gas-phase SO₂ and particle-phase sulfate, and its atmospheric influences were considered by observation and modeling works (Fairlie et al., 2010; Alexander et al., 2012; Chen et al., 2017; Wang et al., 2021b). As summarized by Table S1, when simulating the sulfate burst events, researchers observed the positive feedbacks after implementing heterogeneous mechanism into the WRF-Chem (Li et al., 2017), GEOS-Chem (Shao et al., 2019), CAMx (Huang et al., 2019) and CMAQ (Zhang et al., 2019a) models. However, these improved models highlighted the heterogeneous oxidation motivated by the surface adsorbed oxidants rather than the heterogeneous drivers of aerosol surface. To address the knowledge gap, the revised GEOS-Chem (Wang et al., 2014) and WRF-CMAQ (Zheng et al., 2015) models considered the heterogeneous oxidation driven by aerosol surfaces, and successfully reproduced the rapid sulfate formation. Xue et al. (2016) moved forward to develop an observation-based model and found that heterogeneous pathway contributed up to one third of the secondary sulfate in a typical haze-fog event. However, the "aerosol surface" mentioned in the previous works was not distinguished by its physical or chemical properties (Wang et al., 2014; Zheng et al., 2015; Xue et al., 2016), thereby making it difficult to compare the atmospheric importance of the diverse airborne surfaces. As discussed above, it is of great importance to investigate the heterogeneous drivers of one selected aerosol surface, and further evaluate the atmospheric significance of the relevant heterogeneous pathway.

The resurgence of sandstorms in North China makes the air pollution situation more complex than ever before. The concentration of PM₁₀ (particulate matter with an aerodynamic diameter of less than ten micrometers) in Beijing reached up to 3600 µg m⁻³, largely beyond the standards of World Health Organization and Chinese government (Li et al., 2021a). As the

most abundant primary aerosol in the troposphere (Textor et al., 2006; Tang et al., 2016), dust particles could transport more than one full circuit around the globe within ~ 2 weeks (Uno et al., 2009) and concurrently participate into an array of atmospheric reactions. Heterogeneous reactions over dust surface consume and produce various trace gases, thereby affecting the dust property and tropospheric oxidation capacity (Tang et al., 2016). One of the most extensive concerns is that the numerous surface sites of windblown dust provide opportunities for a variety of atmospheric reactions to occur, e.g. oxidation of SO_2 and formation of sulfate (Usher et al., 2003). In the past decades, plenty of laboratory works have been performed to explore the heterogeneous behaviors of SO_2 on dust surfaces.

When discussing the heterogeneous oxidation on dust proxy, environmental factors like humidity, temperature, irradiance were frequently concerned. Adsorbed water not only helps to accumulate surface $\text{S}_{(\text{IV})}$ species, but also competes with SO_2 for surface sites (Rubasinghege and Grassian, 2013). The reversible adsorption of SO_2 is believed to be exothermic (Clegg and Abbatt, 2001a). In contrast, there were positive temperature dependences observed for CaCO_3 below 250 K over the entire reaction (Wu et al., 2011) and for Fe_2O_3 within 284-318 K during the initial reaction stage (Wang et al., 2018a). Light irradiation normally accelerates the transformation of (bi)sulfites to (bi)sulfates (Li et al., 2010; Nanayakkara et al., 2012; Han et al., 2021), while iron oxides may undergo photoreactive dissolution and thus exhibit weaker reactivity under sunlight (Fu et al., 2009). The reactivity can be enhanced by H_2O_2 (Huang et al., 2016), or O_3 (Li et al., 2006; Li et al., 2007; Zhang et al., 2018), or NO_x (Ma et al., 2008; Liu et al., 2012; He et al., 2014; Yu et al., 2018; Zhao et al., 2018; Yang et al., 2018a; Wang et al., 2020b), or NH_3 (Yang et al., 2016; Yang et al., 2018b; Yang et al., 2019), or Cl_2 (Huang et al., 2017). By contrast, organic compounds, like CH_3CHO (Zhao et al., 2015), HCOOH (Wu et al., 2013), CH_3COOH (Yang et al., 2020; Wang et al., 2022b) and C_3H_6 (Chu et al., 2019), were found to suppress the interactions due to the accumulation or production of particle-phase organic acids. Moreover, CO_2 hinders the heterogeneous oxidation under dark condition (Liu et al., 2020b), whereas presents positive impacts on sulfate formation under solar irradiation (Liu et al., 2022). The particle physical properties, including size (Baltrusaitis et al., 2010; Zhang et al., 2016), morphology (Li et al., 2019) and crystal structure (Yang et al., 2017), display varied impacts on heterogeneous reaction. When considering the particle chemical properties, Fe_2O_3 is more active than Al_2O_3 and SiO_2 in the heterogeneous uptake of SO_2 and formation of sulfate (Chughtai et al., 1993; Zhang et al., 2006; He et al., 2014). Furthermore, the presence of moderate nitrate (Kong et al., 2014; Du et al., 2019), or Al_2O_3 (Wang et al., 2018b), or surfactant (Zhanzakova et al., 2019), or oxalate (Li et al., 2021b), or (bi)carbonate (Liu et al., 2022) on dust surfaces could favor the heterogeneous kinetics under specific conditions. Relative to the environmental factors, the heterogeneous effects relevant to particle property are still under debate.

The usage of simple mineral oxide as a substitution for natural dust may be problematic as such approach could undermine the atmospheric importance of more complex mineralogy. Authentic dusts were utilized in laboratory works (Table S2). Some studies concerned single samples like Saharan dust (Ullerstam et al., 2002; Ullerstam et al., 2003; Adams et al., 2005; Harris et al., 2012), Arizona test dust (Park and Jang, 2016; Zhang et al., 2019a; Zhang et al., 2019b), China loess (Usher et al., 2002) and Asian dust (Ma et al., 2012). The comparison of diverse samples has been attracting increasing attention. Zhou et al. (2014) observed the positive temperature effect on Xinjiang siozem, in contrast to the negative temperature dependence for Inner

90 Mongolia desert dust. Huang et al. (2015) discovered the accelerated oxidation of SO₂ by H₂O₂, and attributed the different moisture effects to the dusts' varying components. Maters et al. (2017) explored the heterogeneous uptake on volcanic ash and glass samples, and related the reactivity differences to the varying abundances of surface basic and reducing sites. Park et al. (2017) compared the heterogeneous reactions on Gobi desert dust and Arizona test dust, and linked the sulfate formation to the quantity of semi-conductive metals. Wang et al. (2019) discovered the enhanced uptake of SO₂ on clay minerals after the simulated cloud processing, and explained this evolution by the modification of iron speciation. Urupina et al. (2019) discussed the kinetics of diverse volcanic dusts, and further experimentally proved that neither one selected pristine oxide nor a mixture of them can adequately typify the behavior of natural dust (Urupina et al., 2021). Recently, based on the measurement method for sulfite and sulfate (Urupina et al., 2020), they determined the associations between sulfate production and dust chemical properties like (Fe+Al)/Si and Na abundance (Urupina et al., 2022). The aforementioned works broaden the horizons of the heterogeneous drivers on dust surface. Up to now, the dominate dust surface drivers remain controversial due to the limited statistical linkages between the chemical composition of dust and the production rate of sulfate.

While the atmospheric relevance of the oxidation of SO₂ on dust has been widely recognized (Wu et al., 2020; Xu et al., 2020), the contribution of dust heterogeneous oxidation to secondary sulfate aerosols has not been quantitatively determined. By means of the improved WRF-CMAQ model, Wang et al. (2012) attributed a 27% decrement of SO₂ concentration and a 12% increment of sulfate concentration to the heterogeneous processes during an Asian dust storm. Moreover, Tian et al. (2021) recently simulated the heterogeneous formation of dust sulfate by the revised GEOS-Chem model and found that, during the dust episodes in North China, up to 30% of the secondary sulfate resulted from the heterogeneous processes on dust surface. However, the photocatalytic reactivity of dust was not considered by the advanced models. Yu et al. (2017) developed the atmospheric mineral aerosol reaction (AMAR) model based on the laboratory works, and suggested that the heterogeneous photocatalysis of mineral dust surface contributed more than half of the secondary sulfate. However, the heterogeneous reactivity was measured by quantifying all the adsorbed SO₂ rather than calculating the yield of particle-phase sulfate. Moreover, limited gas- and aqueous-phase pathways were included in these models, and the heterogeneous reactivities of surface adsorbed oxidants and dust surface drivers have not been distinguished yet. Therefore, it is highly desirable to comprehensively compare the dust heterogeneous pathways with other documented sulfate formation pathways.

115 Hereby, upon understanding the driving factors and driving force of the airborne dust surface, this work compared dust heterogeneous pathways with the gas- and aqueous-phase ones with respect to the formation rate of sulfate and atmospheric lifetime of SO₂. In order to characterize the sensitivity of heterogeneous reaction to dust loading, the scenarios with different dust concentrations were also considered. The joint influences of ionic strength and aerosol liquid water content on the aqueous-phase oxidation of SO₂ were further discussed to prove the significance of heterogeneous kinetics under diverse atmospheric conditions. The recently reported microdroplet interfacial oxidations were additionally compared with the dust-related heterogeneous pathways to emphasize the atmospheric relevance of dust surface drivers. This study attempted to verify the significant formation of sulfate aerosols contributed by the heterogeneous drivers of dust surface.

2 Methods

2.1 Methodology overview

125 This study attempted to investigate whether the heterogeneous oxidation of SO₂ on dust surface, particularly that induced by
the dust surface drivers, makes great impacts on the loss of gaseous SO₂ and formation of sulfate aerosols (Text S1, Fig. S1).
The gas- and aqueous-phase pathways were assessed by the documented methodologies and parameterizations (Cheng et al.,
2016; Seinfeld and Pandis, 2016; Shao et al., 2019; Song et al., 2021), as briefly introduced by Sect. 2.2 (more details in
Supporting Information). The heterogeneous conversion of SO₂ comprises dust-mediated and dust-driven modes, emphasizing
130 the oxidants co-adsorbed with SO₂ and the drivers of dust surface, respectively. In dust-mediated mode, dust surface functions
as a reaction medium that supports the interaction between adsorbed oxidants and SO₂. In dust-driven mode, the oxidation of
adsorbed SO₂ is initiated by the active components of dust surface. The former mode was assessed by the particle properties
based on the reported methodology, while the later mode was quantitatively characterized by the laboratory works.

2.2 Gas- and aqueous-phase oxidation pathways

135 The gas-phase oxidation of SO₂ is initiated by hydroxyl radical (OH), stabilized Criegee intermediates (CIs) and nitrate radical
(NO₃). The former two oxidants promote the sulfate formation during daytime, while the latter one works mainly during
nighttime. More details can be found in Text S2.

The aqueous-phase sulfate formation is pH-dependent and can be quantified based on the published documents and the
references cited therein (Cheng et al., 2016; Su et al., 2020; Liu et al., 2021a). Herein, aqueous-phase pathway refers to the
140 liquid SO₂ conversion by the transition-metal ion-catalyzed oxygen (TMI-O₂), ozone (O₃), hydrogen peroxide (H₂O₂), nitrogen
oxide (NO₂), methyl hydroperoxide (CH₃OOH), peroxyacetic acid (CH₃COOOH), hypochlorous acid (HOCl), hypobromous
acid (HOBr), dissolved nitrous acid (HONO), photosensitization (T*) and nitrate photolysis (P_{NO₃⁻}). The processes relevant to
T* and P_{NO₃⁻} are only considered for the daytime scenario. More details can be found in Text S3 and Tables S3-S4.

The oxidant concentration data were derived from the atmospheric observation campaigns performed in Beijing, North
145 China. The measurements conducted in warm seasons were considered in priority to correspond the experimental temperature.
The relevant temperature range is 293-303 K in representative of warm season. Considering the relatively high irradiance used
in the laboratory experiments, the oxidant parameters for daytime discussion were selected from the observations performed
at noon time. The influences of dust loading on the oxidant concentrations, nitrate photolysis kinetics and TMI abundances
were considered to reflect the linkages between different reactions. More details can be found in Text S4 and Table S5.

150 2.3 Heterogeneous oxidation-particle characterizations

Five airborne clay minerals, including Nontronite (NAu), Chlorite (CCa), Montmorillonite (SWy), Kaolin (KGa) and Illite

(IMt), were obtained from the Source Clay Minerals Repository (Purdue University, West Lafayette, Indiana, USA). The purchased clays were sent for the following measurements.

155 The clay minerals were analyzed by X-ray fluorescence spectrometer (Axios Advanced, PANalytical, Netherlands) for element distributions (Table S6). The Brunauer-Emmett-Teller (BET) specific surface areas (S_{BET}) of NAu, CCa, SWy, KGa and IMt were measured by a Quantachrome Nova 1200 BET apparatus to be 19.76, 5.67, 22.64, 18.77 and 20.05 $\text{m}^2 \text{g}^{-1}$, respectively.

160 Particles were suspended in Milli-Q water (18.2 $\text{M}\Omega\cdot\text{cm}$ at 25 °C) before the size measurement (ViewSizerTM 3000, MANTA Instruments, USA). The particle diameters range mostly from 50 to 1000 nm and are averaged to be 399 nm for NAu, 272 nm for CCa, 438 nm for SWy, 396 nm for KGa, and 366 nm for IMt (Fig. S2).

165 Prepared particles were ultrasonically extracted in Milli-Q water, followed by the filtration through a polytetrafluoroethylene membrane filter. The obtained solution was analyzed by an ion chromatography (883 Basic, Metrohm, Switzerland) for the concentrations of anions (HCOO^- , Cl^- , NO_3^- , SO_4^{2-}) and cations (Na^+ , NH_4^+ , K^+ , Mg^{2+} , Ca^{2+}) (Fig. S3) by the reported methods (Wang et al., 2020c). The water-soluble ions account for 4.6‰, 0.1‰, 8.5‰, 0.3‰ and 0.8‰ of the mass contents of NAu, CCa, SWy, KGa and IMt, respectively.

170 A mixture, denoted as natural dust hereafter, was prepared by mechanically mixing the studied clay minerals by their atmospheric abundances. Because the clays in Kaolinite group (NAu, KGa), Montmorillonite group (SWy), Illite group (IMt) and Chlorite group (CCa) occupy respectively 6.6%, 4.0%, 53.8% and 4.3% mass fractions of the airborne dust (Usher et al., 2003), the prepared natural dust sample comprises 8.8 wt% of NAu, 8.8 wt% of KGa, 5.3 wt% of CCa, 71.4 wt% of IMt, and 5.7 wt% of CCa.

2.4 Heterogeneous oxidation-DRIFTS measurements

175 The *in-situ* DRIFTS (diffuse reflectance infrared Fourier transform spectroscopy) spectra were collected using a FTIR spectrometer (Tracer-100, Shimadzu, Japan) equipped with a mercury-cadmium-telluride detector cooled by liquid nitrogen. A gas supply system was constructed by linking the experimental units through Teflon tubes. Mass flow controllers (D07-19, Severstar, China) adjusted the reactant gases to the expected reactant concentration and relative humidity (RH). Gas cylinders: high-pure air (79% N_2 and 21% O_2), SO_2 (2.46×10^{15} molecules cm^{-3} diluted by N_2).

180 Before each experiment, a ceramic cup holding particles was placed into the reaction chamber. The particles were treated in a stream of dry air (300 ml min^{-1}) for 30 min to minimize the surface water and impurities (Wang et al., 2018a, 2018c). After the pretreatment, the sample was exposed to humidified air (RH = 50%) to reach moisture saturation, followed by the collection of background spectrum and then the introduction of reactant gas for 240 min. The SO_2 concentration was 3.69×10^{13} molecules cm^{-3} in a total flow rate of 100 ml min^{-1} .

The simulated solar irradiation with an actinic flux of 6.51×10^{15} photons $\text{cm}^{-2} \text{ s}^{-1}$ was provided by an Xenon lamp (TCX-250, Ceaulight, China). The reaction temperature (296.8 K) was controlled by a heater attached to a recirculating cooling water system and determined by a calibration curve introduced previously (Wang et al., 2018a). The kinetic parameters involved in

185 the gas- and aqueous-phase oxidation pathways (listed by Sect. 2.2) were corrected by the experimental temperature as much as possible. All exposures were performed in triplicates. The experimental setups are displayed by Fig. S4, and the recorded spectra are shown in Fig. S5 and S6.

2.5 Heterogeneous oxidation-acidity and kinetics

190 The recorded spectra were analyzed by referring to the previous literatures and the references cited therein (Persson and Vgren, 1996; Peak et al., 1999; Goodman et al., 2001; Zhang et al., 2006; Wu et al., 2011; Liu et al., 2012; Nanayakkara et al., 2012; Nanayakkara et al., 2014; Huang et al., 2016; Ma et al., 2017; Yang et al., 2017). Totally six sulfur-containing species can be identified: hydrated SO₂ (SO₂·H₂O), bisulfite, sulfite, solvated sulfate, coordinated sulfate, bisulfate, and the assignments are summarized by Table S7. The overlapping bands were further analyzed by Gaussian/Lorentzian deconvolution to obtain the product distributions (Fig. S7). The consistent deconvolution procedure was performed for the infrared spectra derived from
195 the repeated experiments.

Because part of the measured ions exist in the surface coordinated forms or crystalline states, the particle acidity (pH) may not be accurately characterized by the typical proxy methods (Hennigan et al., 2015). Herein, the ionization equilibrium of dissolved SO₂ in the water layers of particle surface is considered to be associated with particle acidity. As reported, the dissolved SO₂ would transform from SO₂·H₂O to HSO₃⁻, and then to SO₃²⁻ as the medium evolves from the extremely acidic
200 to the nearly alkaline (Haynes, 2014; Zhang et al., 2015). The relative abundance of SO₂·H₂O and SO₃²⁻, as assumed to be equivalent to the relative integral area of their characteristic peaks, is utilized to calculate the particle acidity (more details in Text S3-2 of Supporting Information). The relative abundance of HSO₄⁻ and SO₄²⁻, which can be used to calculate the acidity (Rindelaub et al., 2016; Ault, 2020), was not considered by the present study because the characteristic signals of bisulfate cannot be observed in some of the recorded spectra. As noted by Keene et al. (2004) and Hennigan et al. (2015), estimating
205 the pH of airborne aerosol by ion balance may be difficult because it is unable to distinguish between free and undissociated H⁺ (e.g. protons associated with HSO₄⁻ and HSO₃⁻). Herein, the relative abundance of S_(IV) species is derived from the different infrared absorption signals relevant to atomic sulfur. Thus, the current method is recommended for the pH determination of humidified gas-solid interface.

Heterogeneous kinetics can be assessed by the reactive uptake coefficient (γ) by assuming a first-order loss of SO₂. The γ
210 for dust-driven heterogeneous reaction can be calculated by:

$$\gamma = \frac{d[\text{SO}_4^{2-}]/dt}{Z} \quad (1)$$

$$Z = \frac{1}{4} \times A_s \times [\text{SO}_2] \times v_{\text{SO}_2} \quad (2)$$

$$v_{\text{SO}_2} = \sqrt{\frac{8RT}{\pi M_{\text{SO}_2}}} \quad (3)$$

where $d[\text{SO}_4^{2-}]/dt$ is the rate of sulfate production on dust surface (ion s⁻¹), A_s is the reactive surface area (m²), $[\text{SO}_2]$ is the

215 experimental concentration of SO₂ (molecules m⁻³), v_{SO₂} is the molecular velocity of SO₂ (m s⁻¹), R is gas constant (J mol⁻¹ K⁻¹), T is the experimental temperature (K), M_{SO₂} is the molecular weight of SO₂ (kg mol⁻¹). Because the infrared intensity is proportional to the amount of surface product, [SO₄²⁻] can be translated by the integral area of the sulfate characteristic peaks: [SO₄²⁻] = f × (integral area) (4)

220 where f is the conversion factor and represents the number of SO₄²⁻ corresponding to per unit integral area. The sulfate production rate of dust surface can be translated by the calibration curves by mixing weighed Na₂SO₄ with the target particle sample to a set of concentrations (Martin et al., 1987; Li et al., 2006; Wu et al., 2011; Wang et al., 2020d). The conversion factors are calculated to be 1.32 × 10¹⁸, 4.62 × 10¹⁷, 6.97 × 10¹⁷, 8.20 × 10¹⁷, 9.44 × 10¹⁷, and 9.25 × 10¹⁷ for NAu, CCa, SWy, KGa, IMt, and natural dust, respectively. Because Na₂SO₄ was thoroughly mixed with the particles, S_{BET} was used to calculate γ.

225 All samples except SWy presented steady sulfate production potentials over the entire experiment, while the products on SWy increased by reaction time in the beginning and then gradually remained unchanged (Fig. S8). Accordingly, the process on SWy was assessed by the experimental data of the first 30 min of reaction, and the kinetics of other samples were calculated by the spectra recorded throughout the experiment.

230 Besides the dust surface drivers, the heterogeneous conversion of SO₂ can be additionally initiated by the gaseous oxidants (O₃, H₂O₂, NO₂, HOCl, HOBr, CH₃OOH, CH₃COOOH, HONO) co-adsorbed with SO₂. Considering that the adsorption of oxidant onto dust surfaces would produce gas- and particle-phase species (Usher et al., 2003; Tang et al., 2017), the dust-mediated heterogeneous oxidation can be assumed to primarily proceed in the surface water layers. Such assumption can be experimentally proved as the acceleration of SO₂ oxidation induced by the co-oxidant (e.g. H₂O₂, NO₂, O₃) becomes more significant under higher humidity (Huang et al., 2015; Park et al., 2017; Zhang et al., 2018). Based on the literatures (Hanson et al., 1994; Jacob, 2000; Seinfeld and Pandis, 2016; Shao et al., 2019), the γ for dust-mediated heterogeneous reaction can be calculated by:

$$\gamma = \left[\frac{1}{\alpha} + \frac{v}{4H^*RT\sqrt{D_a k_{\text{chem}}}} \times \frac{1}{f_r} \right]^{-1} \quad (5)$$

$$k_{\text{chem}} = \frac{R_a}{\sum S_{(IV)}} \quad (6)$$

$$f_r = \coth \frac{r_p}{l} - \frac{l}{r_p} \quad (7)$$

240
$$l = \sqrt{\frac{D_a}{k_{\text{chem}}}} \quad (8)$$

where k_{chem} is the pseudo first-order reaction rate constant of the studied S_(IV) specie(s) (s⁻¹) that produce in the rate of R_a (M s⁻¹), f_r is the diffusive correction term comparing the radius of natural dust r_p (m) with the diffuse-reactive length l (m), α is the mass accommodation coefficient of SO₂ (dimensionless), D_a is the aqueous-phase diffusion coefficient of SO₂ (1.78 × 10⁻⁵ m²

245 s^{-1} at 296.8 K) (Himmelblau, 1964; Haynes, 2014), H^* is the effective Henry's law constant for the studied $S_{(IV)}$ specie(s) ($M \text{ atm}^{-1}$), and is jointly determined by the gas-liquid equilibrium of SO_2 and the ionization equilibriums of dissolved sulfur species, and R is gas constant ($L \text{ atm mol}^{-1} \text{ K}^{-1}$). The terms on the right-hand side (RHS) of Eq. (5) show the two contributions to the overall resistance to dust-mediated heterogeneous uptake: mass accommodation at surface, diffusion and reaction in surface liquid water layers. The $S_{(IV)}$ species include $\text{SO}_2 \cdot \text{H}_2\text{O}$, HSO_3^- and SO_3^{2-} .

The sulfate formation rate in the atmosphere can be calculated by the following equations (Jacob, 2000; Li et al., 2020a).

$$250 \quad \frac{d[\text{SO}_4^{2-}]}{dt} = \left(\frac{r_p}{D_g} + \frac{4}{v\gamma} \right)^{-1} S_p [\text{SO}_2] \quad (9)$$

$$S_p = C \times F \times S_{\text{BET}} \quad (10)$$

where $d[\text{SO}_4^{2-}]/dt$ is the atmospheric sulfate formation rate ($\mu\text{g m}^{-3} \text{ h}^{-1}$), r_p is the particle radius (m), D_g is the gas-phase diffusion coefficient of SO_2 ($\text{m}^2 \text{ h}^{-1}$), v is the molecular velocity of SO_2 (m s^{-1}), γ is the reactive uptake coefficient (dimensionless), S_p is the particle surface area density ($\text{m}^2 \text{ m}^{-3}$), $[\text{SO}_2]$ is the atmospheric SO_2 concentration ($\mu\text{g m}^{-3}$), C is the dust concentration of $55 \mu\text{g m}^{-3}$ in representative of the common atmospheric condition of China (Zhang et al., 2012), F is the mass fraction of clay mineral in the natural dust community (%), S_{BET} is the BET specific surface area ($\text{m}^2 \text{ g}^{-1}$).

2.6 Atmospheric lifetime of SO_2

For the gas- and aqueous-phase pathways, the lifetime of SO_2 (τ) can be calculated by Eq. (11) (Jacob, 2000).

$$\tau = \frac{1}{k_{\text{chem}}} \quad (11)$$

260 where k_{chem} is the assumed first-order reaction rate constant of the studied $S_{(IV)}$ specie(s) (s^{-1}).

For the heterogeneous pathways, the τ can be calculated by Eq. (12) (Clegg and Abbatt, 2001b).

$$\tau = \frac{4}{\gamma v S_p} \quad (12)$$

where γ is the reactive uptake coefficient (dimensionless), v is the molecular velocity of SO_2 (m s^{-1}), S_p is the particle surface area density ($\text{m}^2 \text{ m}^{-3}$), as described by Eq. (10).

265 The atmospheric lifetime caused by the multiple pathways (τ_{total}) can be estimated by Eq. (13) (Seinfeld and Pandis, 2016).

$$\tau_{\text{total}} = \left(\sum \frac{1}{\tau} \right)^{-1} \quad (13)$$

3 Results and discussion

3.1 Driving factors of dust surface

270 Correlation analysis is performed to identify the dust surface drivers (Fig. 1), based on which the reaction mechanism of dust heterogeneous pathway can be better understood (Scheme 1).

The mineral drivers are investigated at first (Fig. 1a). Under dark condition, the sulfate production rate correlates positively with Fe, while presents negative dependence against Al. As documented, airborne sulfate was highly associated with Fe-rich dusts, and the heterogeneous reaction of SO₂ can be regarded as a possible explanation (Sullivan et al., 2007). Typical Fe-bearing minerals accelerate the oxidation of S_(IV) species by either the surface active oxygen (O[•]) derived from the adsorbed O₂ in oxygen vacancy or the iron redox cycling initiated by Fe³⁺ from surface acidic media, or both (Baltrusaitis et al., 2007; Fu et al., 2007; Yang et al., 2016). Conversely, Al₂O₃ presents weaker heterogeneous reactivity than Fe₂O₃ (Zhang et al., 2006; Yang et al., 2018b; Xu et al., 2021), and may hinder heterogeneous reaction by blocking the active sites of other mineral constituents (Wang et al., 2018b). Generally, Fe-bearing component plays a crucial role in the heterogeneous reaction of SO₂, while the presence of Al-bearing component directly weakens the dust's reactivity or indirectly decreases the proportion of other active mineral constituents.

280 The sulfate yield enhanced by solar irradiation associates positively with the abundance of Ti or Al. Transition metal oxides in dust act as photocatalyst that yields electron-hole pairs, followed by the formation of reactive oxygen species (ROS) such as hydroxyl radical ([•]OH), superoxide (O₂^{•-}), hydroperoxyl radical (HO₂[•]) and dissociated active oxygen species (O*) (Chen et al., 2012; Abou-Ghanem et al., 2020; Wang et al., 2020c; Sakata et al., 2021). Apart from the site-blocking effect under dark condition, irradiated Al-bearing constituents additionally facilitate the sulfate formation. Physically, Al-bearing components disperse other efficient mineralogical constituents in case of agglomeration (Darif et al., 2016). Chemically, sunlight may alter the electronic configuration of α-Al₂O₃, which presented photoactivity in the reported heterogeneous process (Guan et al., 2014). Generally, the kinetic discrepancy between dark reaction and photoreaction relates primarily to the existence of Ti- and Al-bearing minerals.

290 No correlation can be observed between the abundance of element and the sulfate production rate of photoreaction. The dust with higher proportion of elemental Ti was reported to exhibit greater reactivity toward SO₂ (Park et al., 2017), or NO₂ (Ndour et al., 2009), or O₃ (Abou-Ghanem et al., 2020) in the photochemical processes. However, these results were derived from qualitative comparisons rather than the quantitative analysis performed herein. Analogous to the Al₂O₃ discussed above, semi-conductive metal exhibits dual behaviors in dark and light reactions. For instance, the iron oxide in dark reaction owns great reactivity toward SO₂, whereas that under solar irradiation may present impaired heterogeneous kinetics due to dissolution (Fu et al., 2009). In addition, the heterogeneous reaction on the surface of TiO₂ or Ti-bearing mineral is thermodynamically favored under dark condition but depends largely on photocatalysis under solar irradiation (Chen et al., 2012). Due to the complexity of photoreaction that involves both dark and light reaction mechanisms, the sulfate formation rate of photoreaction may not be

directly linked to particle chemical compositions.

300 The driving effects of water-soluble ions are further studied (Fig. 1b). Herein, Na⁺ and Cl⁻ are observed to present positive impacts on the sulfate formation under dark condition. The presence of halite (NaCl) has positive implications for the dust's hygroscopic growth (Wang et al., 2014; Tang et al., 2019), and the moderate surface water provides an efficient medium for the adsorption of gas-phase SO₂ and the ionization of dissolved sulfur species (Rubasinghege and Grassian, 2013). Moreover, there are negative associations between the photoinduced sulfate enhancement and the abundances of Na⁺, K⁺, Ca²⁺. These
305 cations can be hydrolyzed by adsorbed water to produce H⁺, thereby elevating the particle acidity. In the surface aqueous medium, increased acidity retards the hydrolysis and dissociation of SO₂ (Park et al., 2008; Huang et al., 2015), and inhibits the production of OH by irradiated dust surface (Zheng et al., 1997; Yang et al., 2008; Liu et al., 2017a). Urupina et al. (2022) reported the positive correlations between secondary dust sulfate and the amount of elemental Na. The results here provide additional insights into the heterogeneous effects of the water-soluble constituents on dust surface.

310 Apart from the dust surface drivers and inhibitors, sulfate radical (SO₄^{•-}) may also participate into the heterogeneous event. In the absence of solar irradiation, laboratory studies discovered the uncatalyzed SO₂ autoxidation by SO₄^{•-} at the acidic droplet interface (Hung and Hoffmann, 2015; Hung et al., 2018; Chen et al., 2022). With an appropriate humidity and sufficient saline components, there could be an air-liquid interface occurring over the solid dust surface via water uptake (Gaston et al., 2017; Tang et al., 2019; Wu et al., 2020), then producing SO₄^{•-}. Moreover, on the surface of iron oxide, the •OH derived from Fenton
315 reaction reacts with SO₄²⁻ to form SO₄^{•-} (Kim et al., 2019; Li et al., 2020b). The presence of solar irradiation would accelerate the formation of SO₄^{•-} due to the abundant •OH produced via photocatalysis (Antonioni et al., 2018). Generally, the dust surface with higher proportion of saline components and transition metals may cause the more significant oxidation by SO₄^{•-}, especially under solar irradiation with higher ambient humidity and particle acidity.

Based on the correlation analysis, regression analysis can be further performed to predict the surface kinetics by the chemical
320 compositions of dust. Referring to the publication of Zhang et al. (2020), we derived an exponential parameterization for the production rate of particle-phase sulfate:

$$SF_{\text{dark}} = M[A]^a[B]^b[C]^c[D]^d[E]^e[F]^f + N \quad (14)$$

where SF_{dark} is the yield of SO₄²⁻ on the particle sample of per unit mass within per unit time (ions g⁻¹ s⁻¹); [A], [B], [C], [D], [E] and [F] are the mass fractions of element and ion (%); M and N are the constant parameters (dimensionless).

325 After the statistical procedures by SPSS (version 22.0), we obtained the regression equation for dark reaction:

$$SF_{\text{dark}} (\times 10^{15}) = 22.858 [Al]^{-0.001} [Fe]^{0.111} [Cl^-]^{0.001} [Na^+]^{0.001} - 13.404 \quad (15)$$

Analogously, the discrepancy between the sulfate production rates under dark and illuminated conditions, denoted as R_{light/dark}, can be described by:

$$R_{\text{light/dark}} = SF_{\text{light}}/SF_{\text{dark}} = 14.539 [Al]^{0.073} [Ti]^{0.167} [Cl^-]^{-0.001} [Na^+]^{-0.001} [K^+]^{-0.001} [Mg^{2+}]^{-0.001} - 3.186 \quad (16)$$

330 Then,

$$SF_{\text{light}} = SF_{\text{dark}} \times R_{\text{light/dark}} \quad (17)$$

Because the abundance variation of water-soluble ion presents significantly less influence on the prediction result relative to the elemental transition metal, the complete regression models can be simplified by merely considering the mineral element abundances:

$$335 \quad SF_{\text{dark}} (\times 10^{15}) = 30.880 [\text{Fe}]^{0.077} - 21.679 \quad (18)$$

$$R_{\text{light/dark}} = 15.581[\text{Al}]^{0.062}[\text{Ti}]^{0.141} - 4.277 \quad (19)$$

$$SF_{\text{light}} = SF_{\text{dark}} \times R_{\text{light/dark}} \quad (20)$$

The complete and simplified regression models both accurately simulate the experimental data with all linear slopes approaching 1.0 and all R^2 values larger than 0.995 (Fig. 2). The simplified regression model can be considered as the preferred
340 recommendation due to the fewer parameters needed and its greater performance in the photoreaction prediction. Shang et al. (2010) found that the heterogeneous sulfate production on pristine TiO_2 (Degussa P25) can be accelerated by 8.4 times by the presence of ultraviolet light (365 nm, $350 \mu\text{W cm}^{-2}$). Such photoinduced enhancement is comparable with the prediction result (10.2 times) derived from the simplified regression model developed by this study. Additionally, because the chemical
345 composition of bulk sample may not fully explain the sulfate formation over dust surface, further study to discuss the model uncertainty is warranted.





Figure 1. Correlation analysis on the sulfate production rate of dust surface.

Spearman correlation coefficient for the relationship between the rate of sulfate production and the abundance of (a) mineral element or (b) water-soluble ion. The significant correlations with p values lower than 0.05 or 0.01 are highlighted by the labels in white.




350

Scheme 1. Mechanism illustration on the dust-driven heterogeneous conversion of SO₂.

The heterogeneous reaction of airborne SO₂ on natural dust surface includes the initial adsorption of gas-phase SO₂ and the later conversion of S_(IV) species to S_(VI) products. The formed S_(IV) species include both adsorbed and dissolved SO₂, bisulfite and sulfite (dashed box in blue). The generation of S_(VI) products can be attributed to the surface active oxygen (O⁻), Fe(III)-Fe(II) redox cycling, reactive oxygen species (ROS), and sulfate radical (SO₄^{·-}) (dotted box in deep yellow). Heterogeneous oxidation is largely influenced by the dust surface drivers and inhibitors, which promote and hinder the heterogeneous procedures, respectively (solid box in black). Dust surface drivers are determined as (a) the Fe-bearing constituents and saline ions (Na⁺, Cl⁻) under dark condition, and (b) the Ti- and Al-bearing constituents under solar irradiation. Dust surface inhibitors are identified as (c) the Al-bearing constituents under dark condition, and (d) the metal cations (Na⁺, K⁺, Ca²⁺) under solar irradiation.

355



360

Figure 2. Regression analysis on the sulfate production rate of dust surface.

Linear relationships between the experimental SF_{dark} , SF_{light} , $R_{\text{light/dark}}$ and those simulated by the (a) complete and (b) simplified regression models. The unit of SF (ions g^{-1} s^{-1}) indicates the number of SO_4^{2-} formed on the particle sample of per unit mass within per unit time, and $R_{\text{light/dark}}$ indicates the ratio of SF values in presence and absence of simulated solar irradiation.

365 3.2 Driving force of dust surface

The driving force of dust surface can be characterized by reactive uptake coefficient and particle acidity (Fig. 3, Table S8). Under dark condition, the γ values are highest for SWy and NAu, followed by IMt and CCa, with that of KGa being the lowest. The presence of simulated solar irradiation causes a different rank: IMt>SWy>CCa>NAu>KGa. The γ values of CCa, SWy, KGa, IMt under dark condition are respectively 1.52, 1.01, 2.94, 2.30 times greater than those for photoreactions, reflecting
370 the distinct photocatalytic performances of the clay minerals. Conversely, NAu presents the decreased heterogeneous uptake capacity when exposed to solar irradiation due to its rich abundance of Fe, whose oxides may occur photoreductive dissolution in acidic media (Fu et al., 2010; Shi et al., 2012). Comparing the studies performed by the same experimental approach and assessment procedures, these clay minerals are more efficient than the previously concerned mineral dust proxies, including CaCO₃ (Li et al., 2006; Wu et al., 2011; Zhang et al., 2018), Al₂O₃ (Liu et al., 2017b), TiO₂ (He and Zhang, 2019) and manganese oxides (Wang et al., 2020d), in the heterogeneous production of sulfate under the parallel conditions.

The γ values for the natural dust-driven heterogeneous reactions are calculated to be 6.08×10^{-6} under dark condition and 1.14×10^{-5} under illuminated condition (Fig. 3a). Within the data set of authentic dust, the γ for dark reaction is lower than the reported uptake coefficients of China loess (3.0×10^{-5}), Inner Mongolia desert dust (2.41×10^{-5}), Xinjiang sioezem (8.34×10^{-5}), Saharan dust (6.6×10^{-5}), Asian mineral dust (2.54×10^{-5}), Tengger desert dust (4.48×10^{-5}) and ATD (1.92×10^{-5}) under
380 the similar experimental conditions (Usher et al., 2002; Adams et al., 2005; Zhou et al., 2014; Huang et al., 2015). It should be noted that, the previous studies measured net uptake coefficient that quantifies all the heterogeneously adsorbed SO₂, and some of them assessed the oxidations accelerated by surface adsorbed oxidants (e.g. NO₂, O₃, H₂O₂). In order to quantify the driving force of dust surface in the contribution of particle-phase sulfate, S_(IV) species were not involved in the current kinetics calculation. Quantitatively, the dust surface drivers are responsible for the atmospheric sulfate formation rates of 0.195 and
385 0.365 $\mu\text{g m}^{-3} \text{h}^{-1}$ during nighttime and daytime, respectively.

Particle acidity is further calculated to discuss the driving force of dust surface for sulfate formation. After the SO₂ exposure, CCa is the most acidic, followed by IMt, KGa and NAu, leaving SWy being more neutral. Because no significant correlation can be found between the γ for dust-driven sulfate formation and the pH of reacted clay mineral, the absolute acidity level depends largely on the basic nature of dust. The lowest pH assigned to CCa can be explained by its highest content of elemental
390 Mg relative to the other clays (see Table S6). The Mg-bearing constituents dissolve to be Mg²⁺ that can be further hydrolyzed by water to produce H⁺, thus accelerating the acidification of particle (Park et al., 2008; Huang et al., 2015). For one clay mineral, its different acidities after dark and light reactions reflect the distinct heterogeneous kinetics. All the studied clays, with the exception of NAu, become more acidic after the photoreaction than the dark reaction, which can be explained by the photoinduced SO₂ adsorption and sulfate production on these samples. The opposite situation of NAu coincides to the
395 decreased heterogeneous kinetics over its surface by the presence of solar irradiation. Generally, the natural dust presents the particle acidity (pH) of 4.18 after dark reaction and 4.41 after photoreaction (Fig. 3b). Such results locate within the acidity ranges of dust seeds (3.0-7.0) (Ault, 2020; Pye et al., 2020) and haze aerosols (3.5-4.8) (Ding et al., 2019; Song et al., 2019),

suggesting that the dust-driven heterogeneous pathway could affect aerosol acidity by a certain extent.

Figure 4 presents the pH-dependent γ for natural dust-mediated heterogeneous pathway, along with the experimental data of natural dust-driven heterogeneous pathway. The dust-mediated γ is orders of magnitude lower than the α of SO_2 (~ 0.14 under the experimental temperature of 296.8 K). By definition, α is the probability that a SO_2 molecule striking a liquid surface finally enters into the liquid phase, whereas γ is the sulfate formation rate normalized by the total surface collision rate of SO_2 (Jacob, 2000; Davidovits et al., 2006). That is, γ involves all the uptake processes, including the mass accommodation at surface (Seinfeld and Pandis, 2016). Moreover, unlike the pH-independent α , γ varies with pH, somewhat coinciding to the evolution of dissolved sulfur species that can be quantitatively described by the H^* of SO_2 . The γ values of H_2O_2 , CH_3OOH and CH_3COOOH are pH-independent under specific conditions, in relation to the acid-catalyzed rate-limiting steps relevant to these peroxides (Lind et al., 1987; Liu et al., 2021a). Generally, the dust-mediated heterogeneous γ is largely determined by the diffusion and reaction processes in the water layers of dust surface, as characterized by the second item on the RHS of Eq. (5).

The dust-mediated heterogeneous sulfate formation is primarily contributed by the surface adsorbed H_2O_2 below the nocturnal pH of 5.50 or the diurnal pH of 5.27. When the acidity exceeds the thresholds, the dust-mediated pathway would be kinetically dominated by NO_2 and O_3 during nighttime, or HOBr and HOCl during daytime. Comparing the γ values with the same acidity, the dust-driven pathway appears to be more efficient than the dust-mediated one, thus having the possibility to account for more secondary sulfate aerosols. In the following sections, the oxidation of SO_2 mediated or driven by the natural dust would be set as the typical dust-mediated and dust-driven heterogeneous reactions to compare with the widely documented gas- and aqueous-phase pathways.




Figure 3. Analysis results of the *in-situ* infrared spectra recorded for the heterogeneous reaction of SO₂ on clay minerals and natural dust.

(a) Reactive uptake coefficients (γ) for the heterogeneous formation of sulfate. (b) Particle acidity (pH) of the reacted particle samples. The dark (grey square) and light (yellow circle) conditions were both considered. Dots represent the results of clay minerals, and those of natural dust are shown by the lines. All error bars represent 1 SD.

420




Figure 4. Particle acidity-dependent reactive uptake coefficients (γ) for the dust-mediated and dust-driven heterogeneous pathways.

The dust-mediated pathway can be induced by the surface oxidants co-adsorbed with SO_2 , including hydrogen peroxide (H_2O_2), nitrogen oxide (NO_2), ozone (O_3), hypochlorous acid (HOCl), hypobromous acid (HOBr), methyl hydroperoxide (CH_3OOH), peroxyacetic acid (CH_3COOOH) and dissolved nitrous acid (HONO). The dust-driven pathway can be induced by the heterogeneous drivers (transition-metal-bearing components and water-soluble ions) on natural dust and clay minerals [Nontronite (NAu), Chlorite (CCa), Montmorillonite (SWy), Kaolin (KGa), Illite (IMt)].

425

3.3 Comparison of atmospheric oxidation pathways

430 Figure 5 compares the sulfate formation rates of diverse atmospheric oxidation pathways by a newly developed comparison model. Based on the parameterization scheme in model and the experimental results discussed above, the total sulfate formation rates are summed to be $0.795 \mu\text{g m}^{-3} \text{h}^{-1}$ during nighttime and $5.179 \mu\text{g m}^{-3} \text{h}^{-1}$ during daytime, under the acidity of natural dust (Fig. 5a and e). The higher sulfate flux during daytime can be explained by the photo-increased oxidation channels, oxidant concentrations, heterogeneous reactivity, and the elevated particle acidity that facilitates the TMI catalysis. It is worthwhile to mention that, the nocturnal sulfate concentration was reported to approach or exceed the subsequent diurnal level, as explained by the higher nocturnal humidity facilitating the liquid oxidations or the lower boundary layer at night causing the adverse diffusion conditions (Liu et al., 2017c; Tutsak and Koçak, 2019; Li et al., 2020c). Meteorological factors like humidity are not considered by the current model, which emphasizes the comparison of diverse pathways through kinetic regime. The estimated sulfate fluxes are generally lower than some published data derived from the same concentration of SO_2 (40 ppb) because the gas- and aqueous-phase parameters here are corrected by the experimental temperature (296.8 K) rather than the much lower level (271 K) used previously (Cheng et al., 2016; Gen et al., 2019; Liu et al., 2020a; Su et al., 2020; Wang et al., 2020a; Liu et al., 2021a). Considering that some kinetic parameters were experimentally determined under room temperature with the lack of temperature dependence, uncertainties may exist in the discussion on cold environments. The current sulfate formation is assessed near room temperature that is pertinent to the nature of sandstorm occurring during late spring and early summer in East Asia (Wu et al., 2020; Ren et al., 2021).

During nighttime, the gas-phase, aqueous-phase and heterogeneous pathways explain 31.6, 39.8 and 28.6% of secondary sulfate, respectively (Fig. 5b). The diurnal contribution proportions of gas-phase (45.5%) and aqueous-phase pathways (41.4%) exceed their nocturnal levels, thus lowering the heterogeneous proportion to 13.1% during daytime (Fig. 5f). Although the heterogeneous sulfate yield during nighttime is lower than that during daytime (see Sect. 3.2), nocturnal heterogeneous process accounts for the higher proportion of secondary sulfate. The diurnal sulfate formation rates of gas-phase, aqueous-phase and heterogeneous pathways are respectively 9.4, 6.8 and 3.0 times greater than the nocturnal levels, indicating that the oxidations in gaseous and liquid media could be more kinetically susceptible to the occurrence of sunlight than those relevant to the humidified gas-solid interface. In gas-phase oxidation, OH is the predominant oxidant, followed by CIs, while NO_3 contributes little. In aqueous-phase oxidation, TMI- O_2 and H_2O_2 play crucial roles in the conversion of SO_2 , coinciding to the reported results (Fan et al., 2020; Song et al., 2021). Besides, there are lower contributions from the NO_2 during nighttime and the HOBr, T^* , $\text{P}_{\text{NO}_3^-}$ during daytime. The heterogeneously formed sulfate is mostly ascribed to the dust-driven pathway rather than the dust-mediated one. In comparison, the sulfate contribution proportions of the studied dust heterogeneous oxidation are comparable with those obtained by the OBM model (30.6% for nighttime and 19.4% for daytime) (Xue et al., 2016), and the revised GEOS-Chem model (20-30%) (Tian et al., 2021) and WRF-CMAQ model (up to 12%) (Wang et al., 2012). While the uptake coefficients used in the previous studies are generally greater than the present experimental results, more pathways are implemented in the current comparison model, thereby causing the parallel comparison results. By contrast, the AMAR

model highlighted the dust's photocatalytic surface that contributes remarkable secondary sulfate (>50%) under the constraint simulation conditions (Yu et al., 2017).


465 In dust-mediated heterogeneous oxidation, H_2O_2 is the most efficient oxidant, followed by the nocturnal NO_2 and diurnal hypohalous acids (HOBr, HOCl) responsible for less secondary sulfate (Fig. 5c, g). The dust-driven heterogeneous sulfate formation is mainly attributed to IMt that owns the largest proportion in dust community, followed by NAu and SWy with relatively significant contributions by heterogeneous sulfate formation (Fig. 5d and h). The nocturnal and diurnal sulfate fluxes of natural dust-driven heterogeneous pathway are respectively 5.8 and 1.2 times greater than those of the natural dust-mediated one. Hence, the heterogeneous sulfate formation is primarily ascribed to the dust surface drivers rather than the surface adsorbed oxidants, as experimentally proved by the laboratory studies concerning the heterogeneous SO_2 oxidation on authentic dusts accelerated by the presence of NO_2 (Park et al., 2017), or O_3 (Park et al., 2017), or H_2O_2 (Huang et al., 2015).
470 The kinetic discrepancy between the dust-mediated and dust-driven heterogeneous pathways is more significant during nighttime than daytime. In the estimation, H_2O_2 is the predominant dust-mediated oxidant and its concentration becomes lower under weaker sunlight. The dust-mediated contribution during nighttime is thus relatively small by the presence of the relatively low H_2O_2 concentration. In general, when investigating the heterogeneous process on dust particles, particle variables could be more important than gas variables in elucidating the reaction characteristics and atmospheric implications.

Figure 6 summaries the particle acidity-dependent sulfate formation rates of gas-phase, aqueous-phase and heterogeneous pathways. Relative to the pH-independent gas-phase process, the aqueous-phase oxidation becomes relatively productive under the extremely acidic and near-neutral situations, while fails to support the rapid sulfate formation within the pH range of 4-5 (minimums at 4.85 for nighttime and 4.55 for daytime), which overlaps the acidity range of the weak dust-mediated heterogeneous pathway. The significant sulfate formation in acidic medium is primarily contributed by the TMI- O_2 pathway where the concentrations of TMIs and H^+ keep increasing as the aerosol water becomes more acidic. When $\text{pH}>5.0$, the sulfate formation rate increases with elevated pH, in step with the evolution of the concentration of dissolved sulfur species. Noticeably, the pH range of 4-5 involves the acidity of the aged natural dust and overlaps that of the common haze aerosols, implicating that the dust surface drivers have profound impacts on the secondary sulfate burst in highly polluted environment (Ding et al., 2019; Song et al., 2019). The important role of dust in sulfate formation was confirmed by atmospheric observation research. For example, secondary sulfate was observed to accumulate on the dust-dominant super-micron particles collected in the North China Plain, and the mass fraction of coarse-mode sulfate dramatically increased during the evolutionary stages of haze episode (Xu et al., 2020). Within the entire pH range, the aqueous-phase pathway is more active than the dust-mediated heterogeneous one in contributing to secondary sulfate.
480
485
490

Figure 7 compares the lifetimes of SO_2 influenced by the diverse atmospheric oxidation pathways. Calculations of lifetimes can be useful in estimating how long the SO_2 is likely to remain airborne before it is removed from the atmosphere (Seinfeld and Pandis, 2016). Theoretically, the lifetime of SO_2 determined by heterogeneous reaction is negatively correlated with γ and S_p , and S_p is positively associated with the concentration and S_{BET} of dust, as respectively described by Eq. (12) and (10). As a result, the dust with greater heterogeneous reactivity, or higher atmospheric loading, or larger S_{BET} is prone to cause the
495

shorter lifetime. During nighttime, IMt causes the shortest SO₂ lifetime (46.90 days) due to its highest concentration relative to the other clay minerals. The relatively large heterogeneous uptake capacity of NAu and SWy link to the second and third shortest SO₂ lifetimes (221.30 and 260.12 days, respectively). On the other hand, the weakest heterogeneous reactivity of KGa leads to the second longest lifetime (1758.65 days), and the longest result caused by CCa (2256.25 days) can be interpreted by its lowest S_{BET} that causes the lowest Sp. The presence of solar irradiation alters the lifetime ranking: IMt (20.39 days) < SWy (256.96 days) < NAu (437.85 days) < KGa (597.54 days) < CCa (1488.46 days), as influenced by the different photoactivities of the clay minerals. The heterogeneous drivers of natural dust surface are comparable with TMI-O₂ in the loss of SO₂, and both of them function as the most important lifespan influencers in the absence of sunlight (Fig. S9a). During daytime, the natural dust-driven pathway is only next to the oxidations induced by OH, TMI-O₂ and H₂O₂ (Fig. S9b). The comparison results of lifetime agree well with those of sulfate formation rate (Fig. 5), illustrating that the heterogeneous drivers of dust surface are somewhat responsible for altering the concentrations of gas-phase and particulate sulfur species.

Gas- and aqueous-phase pathways serve as the most significant influencers of the diurnal lifespan of SO₂, followed by dust-driven heterogeneous pathway, while during nighttime these three present closer impacts (Fig. S10). The atmospheric lifetimes of SO₂ induced by the considered atmospheric oxidation pathways are calculated to be 6.17 days during nighttime and 0.99 days during daytime. Neglecting dust heterogeneous pathways may lengthen the SO₂ lifespans to 10.27 and 1.12 days in the absence and presence of solar irradiation, respectively. Analogously, scientists obtained the shorter lifetime of SO₂ from climate model after considering the heterogeneous reaction occurring on volcanic ash particles (Zhu et al., 2020). Clay minerals are more concentrated in the troposphere than volcanic ash and thus have more significant impacts on the removal of atmospheric SO₂.



515

Figure 5. Contributions of diverse atmospheric oxidation pathways to secondary sulfate aerosols.

Gas-phase oxidation can be induced by hydroxyl radical (OH), stabilized Criegee intermediates (CIs), as well as the nitrate radical (NO_3) only for nighttime. Aqueous-phase oxidation can be induced by hydrogen peroxide (H_2O_2), nitrogen oxide (NO_2), ozone (O_3), hypochlorous acid (HOCl), hypobromous acid (HOBr), methyl hydroperoxide (CH_3OOH), peroxyacetic acid (CH_3COOOH), dissolved nitrous acid (HONO), transition-metal ion-catalyzed oxygen (TMI- O_2), and the photosensitization (T^*) and nitrate photolysis (P_{NO_3}) only for daytime. Dust-mediated heterogeneous oxidation can be initiated by the surface oxidants (H_2O_2 , NO_2 , O_3 , HOCl, HOBr, CH_3OOH , CH_3COOOH , HONO) co-adsorbed with SO_2 . Dust-driven heterogeneous oxidation can be ascribed to the heterogeneous drivers (transition-metal-bearing components and water-soluble ions) on the surfaces of natural dust and clay minerals [Nontronite (NAu), Chlorite (CCa), Montmorillonite (SWy), Kaolin (KGa), Illite (IMt)]. The (a-d) nighttime and (e-h) daytime conditions were distinguished by the different parameterizations. (a, e) Particle acidity-dependent sulfate formation rates of the diverse atmospheric oxidation pathways. The elements' colors and shapes are characterized by the legends in solid boxes. (b, f) Quantified sulfate contribution proportions of the studied gas-phase (red), aqueous-phase (blue), and heterogeneous (yellow) reaction pathways. Sulfate formation rates of the dust-mediated and dust-driven pathways during (c-d) nighttime and (g-h) daytime. The effects of ionic strength on the aqueous-phase oxidation were not taken into account. The dust concentration was set to be $55 \mu\text{g m}^{-3}$, in representative of the common atmospheric condition of North China (Zhang et al., 2012). The panels in dashed boxes right to the legends illustrate the primary physical-chemical processes of atmospheric sulfate formation. More parameterization and methodology details can be found in the Texts S2-S4 of Supporting Information and Sect. 2.1-2.5 of the main content.

525

530




Figure 6. Formation rate of sulfate attributed to the gas-phase, aqueous-phase and dust heterogeneous pathways as a function of particle acidity (pH).

535 Grey areas indicate the pH ranges of the polluted particulate matters, with darker ones being more common (Ding et al., 2019).




Figure 7. Atmospheric lifetimes of SO₂ induced by the diverse sulfate formation pathways.

Both nighttime (grey square) and daytime (yellow circle) conditions were considered. The lifetimes of SO₂ caused by gas-phase, aqueous-phase and dust heterogeneous pathways are displayed by the dashed lines. The atmospheric lifetimes of SO₂ induced by all the studied oxidation pathways are presented by the solid lines. The effects of ionic strength on the aqueous-phase SO₂ oxidation were not taken into account. The dust concentration was set to be 55 μg m⁻³ to reflect the common atmospheric condition of North China. More methodology details can be found in Sect. 2.6 of the main content.

540

3.4 Sensitivity analysis

The aforementioned calculations set the concentration of natural dust to be $55 \mu\text{g m}^{-3}$. In contrast to the common atmospheric loading, the burst of sandstorm was normally accompanied by the quickly elevated dust concentration up to thousands of $\mu\text{g m}^{-3}$ (Li et al., 2021a; Yin et al., 2021; Filonchik, 2022). It would be meaningful to estimate the heterogeneous contributions in the dust-rich environments. Theoretically, dust concentration within the ranges of $72\text{--}770 \mu\text{g m}^{-3}$ during nighttime and $24\text{--}260 \mu\text{g m}^{-3}$ during daytime could cause the extra sulfate formation of $0.3\text{--}3.0 \mu\text{g m}^{-3} \text{h}^{-1}$ (Fig. 8a), in line with the acknowledged range of missing sulfate formation rate (Cheng et al., 2016; Liu et al., 2020a), which was found to be positively correlated with PM concentration as well (Cheng et al., 2016). Therefore, the heterogeneous reaction of SO_2 on dust surface is a considerable sulfate formation pathway and may evolve into the missing sulfate source in the atmosphere.

The occurrence of sandstorm, particularly during nighttime, aggravates the sulfate pollution in coarse aerosol mode (Fig. 8b). For instance, the dust concentration of $200 \mu\text{g m}^{-3}$, which approaches the PM_{10} level in North China on March 2021 (Yin et al., 2021), could heterogeneously explain 44.9% of the secondary sulfate during nighttime, as well as 29.6% during daytime. It is worthwhile to note that, the heterogeneous contribution proportion is susceptible to the evolution of relatively low dust concentration, further increase of dust loading will not significantly elevate the heterogeneous proportion, resulting in a plateau (nighttime) or decrease (daytime) under severe dust pollution. In fact, the increased dust concentration facilitates the aqueous-phase TMI- O_2 pathway and dust-mediated and dust-driven heterogeneous reactions, whereas presents negative impacts on the others by the removal of gaseous oxidants over dust surface. The unsusceptible response to dust concentration, as shown by Fig. 8b, is related to the increased sulfate contributions from TMI- O_2 . Unlike the TMI-derived oxidation and dust-driven pathway receiving constant positive feedbacks from the increased dust concentration, the dust-mediated pathway is somewhat affected by the decreased oxidants adsorbed on dust surface. Since the importance of dust-mediated pathway is more significant during daytime than nighttime, there is a negative correlation between the high dust concentration and heterogeneous contribution proportion in the presence of solar irradiation.

The increased dust concentration, in fact, not only facilitates the heterogeneous process by providing more reactive surfaces, but also affects the gas- and aqueous-phase reactions by altering the atmospheric abundances of reactive species, as explained below. The evolution of dust pollution from slight to heavy conditions would cause the loss of various gaseous oxidants by heterogeneous uptake, and therefore the gas- and aqueous-phase sulfate fluxes, except that induced by TMI- O_2 , decrease against dust concentration (Bian and Zender, 2003; Tang et al., 2017). Furthermore, the dissolution of mineral constituents produces TMIs in aerosol liquid media (Alexander et al., 2009; Shao et al., 2019), and the irradiated mineral dust was reported to emit gaseous ROS by surface photocatalysis (Dupart et al., 2012; Chen et al., 2021). Herein, the studied oxidation pathways are distinguished by their contribution proportions (Fig. 8c and d). The contribution proportions of OH and H_2O_2 decrease against dust loading, whereas the importance of TMI-derived oxidation and dust-driven pathway become more significant as the dust concentrates. The contribution proportion assigned to dust-mediated process increases with dust concentration at first and then decreases. While the increased dust loading provides more physical space for the dust-mediated reactions, the

simultaneously decreased gas-phase oxidants restrain the accumulation of particle-phase oxidants.

580 Atmospheric lifetime of SO₂ is also affected by the concentration of dust. As shown by Fig. 9a, the lifespan of airborne SO₂ during nighttime is higher than that during daytime, and both of them decrease against dust concentration. Analogous to the heterogeneous contribution proportion (Fig. 8b), the lifespan of SO₂ is more susceptible to the variation of dust concentration in clean and slightly polluted environments than that under heavily polluted conditions. The mild dust pollution, especially its level variations, should be paid more attention. The heterogeneous loss of SO₂ by dust surface was normally evaluated against the gas-phase loss by OH (Ullerstam et al., 2003; Adams et al., 2005; Li et al., 2006; Huang et al., 2015; Ma et al., 2018). Current estimation indicates that the airborne dust with the concentration of 45 μg m⁻³ during nighttime, or 91 μg m⁻³ during daytime, can be regarded as comparable with OH in controlling the removal of SO₂ (Fig. 9b). Such dust concentrations are 585 sometimes common in the troposphere, especially during the dust storm periods (Li et al., 2021a; Yin et al., 2021; Filonchyk, 2022) or near the dust source regions (Ke et al., 2022). Therefore, the heterogeneous loss of SO₂ by airborne dust surface may have a similar magnitude as the main gas-phase loss process and can be taken as an important sink for SO₂.





Figure 8. Sensitivity tests of sulfate formation rate to dust concentration.

590

(a) Sulfate formation rate of diverse dust-related heterogeneous pathways as a function of dust concentration. The grey area suggests the missing sulfate formation rate ranging from 0.3 to 3 $\mu\text{g m}^{-3} \text{h}^{-1}$ as a reference (Cheng et al., 2016; Liu et al., 2021a). (b) Heterogeneous sulfate proportion varying with dust concentration. Secondary sulfate contributions attributed to the studied oxidation pathways during (c) nighttime and (d) daytime.



595

Figure 9. Sensitivity tests of SO₂ lifetime to dust concentration.

(a) Atmospheric lifetime of SO₂ induced by all the studied pathways varying with dust concentration. (b) Comparison between the SO₂ lifetimes ascribed to dust heterogeneous pathway and OH-initiated gas-phase oxidation.

3.5 Uncertainty analysis

600 The contribution proportion of dust heterogeneous oxidation could be over- or underestimated if considering the uncertainty factors. Herein, the joint impacts of ionic strength (I) and aerosol liquid water content (ALWC) on the aqueous-phase oxidation and the kinetic comparison between microdroplet interfacial oxidation and dust-driven heterogeneous oxidation are discussed to further understand the atmospheric relevance of the heterogeneous drivers on dust surface under the complex atmospheric conditions.


605 The aqueous-phase oxidation of SO₂ by H₂O₂, O₃, NO₂ and TMI-O₂ was quantified under different I-ALWC settings. At first, the sulfate formation rate was calculated as a function of ionic strength under the studied ALWC of 300 μg m⁻³ (Fig. S11). TMI-O₂ and H₂O₂ dominate the liquid oxidation, while the impacts of NO₂ and O₃ only slightly peak at ~ 1.0 M during nighttime. Specifically, during nighttime, TMI-O₂ dominates the sulfate formation under the relatively low ionic strength, while the contribution of H₂O₂ exceeds that of TMI-O₂ over the ionic strength of 0.028 M. During daytime, H₂O₂ is the predominate oxidant within the studied ionic strength range. Relative to the ionic strength-free calculations, the aqueous oxidation could be weakened by the ionic strength lower than 17.8 M during nighttime or 14.3 M during daytime. Such values can be taken as criteria to distinguish the over- or underestimation of liquid kinetics under the ALWC of 300 μg cm⁻³, which was widely used to characterize the haze events in North China (Cheng et al., 2016).

610 The joint influences of I and ALWC were further considered (Fig. 10). At each ionic strength, sulfate formation rate associates positively with ALWC. At each ALWC, the increase of ionic strength hinders the aqueous oxidation at first and then facilitates this process, as a consequence resulting in a threshold line distinguishing the negative or positive effects of ionic strength, as presented by Fig. 10a and b. Furthermore, the I-ALWC relationships of California, USA (Stelson and Seinfeld, 1981); Beijing, China (Song et al., 2021); Mexico City, Mexico (Volkamer et al., 2007; Hennigan et al., 2015); and the nine cities of Germany (Scheinhardt et al., 2013) were found to locate left to the thresholds, indicating the negative effect of ionic strength under the investigated scenarios (Fig. 10c). Calculated by the reported I-ALWC relationships, the dust-mediated pathway contributes 4.3-20.1% of the secondary sulfate during nighttime and 6.8-22.0% during daytime, and the dust-driven pathway accounts for respectively 29.1-41.6% and 9.9-12.4% of the sulfate formation in the absence and presence of sunlight (Fig. 10d). Therefore, the heterogeneous contribution proportions in the complex atmospheric environments can be even higher than those estimated by this study.

625 Besides the humidified gas-solid interface, gas-liquid interface of microdroplet is another type of medium that supports the heterogeneous formation of sulfate. The oxidation of SO₂ was found to proceed at the interfacial layer of a droplet with higher kinetics than the bulk process (Jayne and Davidovits, 1990). Recently, the interfacial roles of O₂ (Hung and Hoffmann, 2015; Hung et al., 2018; Chen et al., 2022), NO₂ (Liu and Abbatt, 2021; Yu, 2021) and Mn²⁺ (Zhang et al., 2021; Wang et al., 2021a; Wang et al., 2022c) have been quantitatively described. Figure 11 compares dust-driven heterogeneous pathway with the documented gas-liquid interfacial oxidations. As reported by Hung et al. (2015, 2018), acidic droplet interface favors the noncatalyzed oxidation by the presence of sufficient SO₃⁻ and SO₄⁻ when pH<4.0, which can be further explained by the

structure differences of water at the interface and in the bulk phase. Recently, Chen et al. (2022) reevaluated the sulfate formation by interfacial O₂ within the pH range of 3.5-4.5 and discovered the positive dependence toward ionic strength. The oxidation of SO₂ by interfacial NO₂ displays the similar pH dependence as that occurs in bulk solution. The secondary sulfate contributed by interfacial Mn²⁺ is associated positively with particle acidity. Generally, the O₂ at acidic interface dominates the oxidation when pH<4.0, whereas at pH>4.0 the interfacial oxidation is primarily controlled by Mn²⁺. Comparing the sulfate formation rates under the same aerosol acidity, the dust surface drivers generally present greater reactivity than the interfacial NO₂ and O₂. Under the ionic strength of 40 M, the reactivity of interfacial O₂ can be regarded as equivalent to that of the dust surface drivers during nighttime. Interfacial Mn²⁺ exceeds the dust surface drivers through kinetic regime. For instance, the Mn²⁺-catalyzed oxidation characterized by Wang et al. (2021a) is 6.5 or 1.7 times more efficient than the dust-driven pathway in sulfate formation during nighttime or daytime, respectively.

The dust-driven heterogeneous pathway here was investigated by the infrared technique focusing on the bulk particle sample rather than air-suspended particles. Up to now, the micro-scale effects of dust particle have not been systematically studied. Suspended ATD was concerned by a smog chamber research and its heterogeneous reactivity toward SO₂ was characterized by the net uptake coefficient of 1.71×10^{-6} under dark condition (Park and Jang, 2016). Such kinetic constant is approximately one order of magnitude higher than those obtained from the film-based flow tube tests ($\sim 1.75 \times 10^{-7}$) under the parallel experimental conditions (Zhang et al., 2019b). Therefore, in the atmosphere, dust surface drivers may kinetically approach the microdroplet interface in the removal of SO₂ and formation of sulfate.



650 **Figure 10.** Joint influences of ionic strength (I) and aerosol liquid water content (ALWC) on the aqueous-phase oxidation of SO₂.
 Aqueous-phase sulfate formation rate varying with I and ALWC during (a) nighttime and (b) daytime. The dash-dotted lines indicate the
 thresholds distinguishing the negative and positive effects of ionic strength. (c) Reported I-ALWC relationships versus the nocturnal and
 diurnal thresholds. The field-observed data were collected from the measurements performed in California, USA (Stelson and Seinfeld,
 1981); Beijing, China (Song et al., 2021); Mexico City, Mexico (Volkamer et al., 2007; Hennigan et al., 2015); and the nine cities of Germany
 655 (Scheinhardt et al., 2013). (d) Heterogeneous contribution proportions calculated by the reported I-ALWC relationships and parameterization
 of this study (ionic strength-free settings and an ALWC of 300 $\mu\text{g m}^{-3}$ for aqueous-phase oxidation; dust concentration of 55 $\mu\text{g m}^{-3}$ for
 heterogeneous oxidation).




Figure 11. Kinetic comparison between microdroplet interfacial oxidation and dust-driven heterogeneous pathway.

660

The interfacial oxidation of SO_2 can be induced by the O_2 , or NO_2 , or Mn^{2+} at aerosol particle interfaces. The O_2 -dominated oxidation can be assessed by the method concluded by Hung et al. (2018). Chen et al. (2022) refreshed the assessment method relevant to interfacial O_2 , and the result here was obtained under the ionic strength of 40 M within the pH range of 3.5-4.5. The NO_2 -dominated oxidation was assessed based on the work of Liu and Abbatt (2021). The Mn^{2+} -dominated oxidation can be assessed by the methods from Wang et al. (2021a) and Zhang et al. (2021). More parameterization and methodology details can be found in the Text S5 of Supporting Information.

665 4 Conclusions and implications

This study attempted to deeply understand the importance of heterogeneous oxidation, particularly that induced by dust surface drivers, in the loss of airborne SO₂ and formation of sulfate aerosols. Laboratory research was performed at first to investigate the heterogeneous driving factors and driving force. Based on the correlation and regression analysis, transition metal elements, particularly Fe for dark condition, and Al and Ti for photoreaction, were determined to dominate the heterogeneous oxidation, while water-soluble ions present minor influences. A series of empirical equations were developed to kinetically predict the dust-driven process. The Al, Fe, Ti-bearing mineralogical components, as well as their mixtures, are thus recommended as appropriate proxies for the following laboratory research. The dust-driven heterogeneous γ values were calculated to be 6.08×10^{-6} for dark reaction and 1.14×10^{-5} for photoreaction, corresponding respectively to the atmospheric sulfate formation rates of 0.195 and 0.365 $\mu\text{g m}^{-3} \text{h}^{-1}$ in the presence of 55 $\mu\text{g m}^{-3}$ natural dust.

670 A comparison model was further developed to reveal the atmospheric relevance of the heterogeneous drivers on dust surface. Dust heterogeneous oxidation is suggested to explain 28.6% of the secondary sulfate aerosols during nighttime and 13.1% during daytime, and the dust surface drivers act as the dominate contributors. Furthermore, dust heterogeneous oxidation affects the atmospheric lifetime of SO₂. The increased dust concentration may aggravate the secondary sulfate pollution and has the potential to explain the acknowledged missing sulfate formation rate (0.3-3.0 $\mu\text{g m}^{-3} \text{h}^{-1}$). The joint effects of I and ALWC on liquid kinetics influence the importance of heterogeneous pathway. The heterogeneous contribution proportions estimated by the reported I-ALWC relationships are generally greater than those calculated by the parameterization in current comparison model. Additionally, dust heterogeneous pathway in the atmosphere is believed to be kinetically comparable with droplet interfacial oxidation.

685 Overall, this study suggests that the implementation of heterogeneous processes into atmospheric models shall vastly improve the agreement between the modeled and observed sulfate concentrations. Dust heterogeneous pathway should be treated as a significant contributor of secondary sulfate. More necessarily, dust surface drivers are needed to be viewed as an important research focus. Zheng et al. (2015) revised the CMAQ model by adding heterogeneous mechanism and observed the accurate simulation run upon determining the uptake coefficient to be 2.0×10^{-5} , which is higher than the γ values of dust heterogeneous pathway herein (7.04×10^{-6} during nighttime and 1.55×10^{-5} during daytime). Hence, other solid aerosols, such as sea salts (Laskin et al., 2003; Rossi, 2003) and carbonaceous particles (He and He, 2020; Zhang et al., 2020), would better be taken into the following heterogeneous discussions together with the dust-related processes we studied.

695 Meteorological factors impact the atmospheric relevance of dust heterogeneous pathway. Taking humidity as an example, on dust surface, the heterogeneous reaction of SO₂ is humidity-dependent, and the exact dependence varies with the type of dust and the condition of reaction (Huang et al., 2015; Park et al., 2017; Urupina et al., 2022). The uptake of gas-phase oxidants over dust surface is also influenced by humidity (Kumar et al., 2014). For aerosol droplet, increased humidity elevates its liquid volume and radius (Wu et al., 2018; Ding et al., 2019). Aerosol liquid water serves as an efficient medium for multiphase

700 reactions (Wu et al., 2018; Yue et al., 2019), whereas radius is negatively associated with the sulfate formation at droplet interface (Hung et al., 2018; Wang et al., 2021a; Chen et al., 2022). Furthermore, droplet acidity decreases as the liquid volume increases, thereby increasing the ionization of dissolved sulfur species, followed by the increased sulfate formation rate (Yue et al., 2019; Jin et al., 2020; Gao et al., 2022). Overall, how meteorological factors influence the relative importance of gas-phase, aqueous-phase and heterogeneous pathways warrants further research.

705 Heterogeneous laboratory results were scarcely discussed together with other SO₂ conversion routes. This study attempted to set an example for kinetically comparing the heterogeneous oxidation characterized by laboratory work with the documented gas- and aqueous-phase data. Relative to the three-dimensional numerical models, the developed comparison model involves more oxidation pathways. Since the aqueous-phase oxidation here is relevant to aerosol droplet rather than cloud/fog droplet, the current parameterization could be more appropriate for simulating the fine particulate matters that were frequently collected by atmospheric observations and compared with modeling data. Furthermore, the heterogeneous oxidation was classified into dust-mediated and dust-driven modes to better distinguish the key surface impactors. Therefore, this comparison model has advantages over the traditional atmospheric chemistry models, and is recommended for the following heterogeneous laboratory research to systematically compare the experimental data with the acknowledged gas-phase/aqueous-phase/heterogeneous oxidation pathways.

710 This work broadens the application of infrared technique in atmospheric laboratory research. Apart from the uptake coefficient calculations normally concerned, this study moved forward to bridge the relationship between particle acidity and sulfate formation rate by analyzing the shape and intensity of infrared spectrum, and further compared heterogeneous oxidation with other atmospheric pathways. This research not only provides a promising methodology for the future heterogeneous research utilizing the classic *in-situ* DRIFTS approach, but also helps to take good advantages of the infrared technique in the laboratory studies in relation to atmospheric heterogeneous oxidation.

Data availability. A dataset for this paper can be accessed at <https://data.mendeley.com/datasets/hyvdz7khs6/1>.

720 **Author Contributions.** T.W. designed the experiments and processed the data, and wrote the paper together with Y.L; H.C. and Z.W. measured the physical and chemical properties of the particle samples; H.F., J.C., and L.Z. provided guidance in the data analysis and paper writing.

Competing interests. The authors declare no competing financial interest.

725 **Financial support.** The research was financially supported by National Natural Science Foundation of China (22176036, 21976030, 22006020, 42205099), Natural Science Foundation of Shanghai (19ZR1471200) and China Postdoctoral Science Foundation (2021M700792).

References

- 730 Abou-Ghanem, M., Oliynyk, A. O., Chen, Z., Matchett, L. C., McGrath, D. T., Katz, M. J., Locock, A. J., and Styler, S. A.: Significant Variability in the Photocatalytic Activity of Natural Titanium-Containing Minerals: Implications for Understanding and Predicting Atmospheric Mineral Dust Photochemistry, *Environ. Sci. Technol.*, 54, 13509-13516, <https://doi.org/10.1021/acs.est.0c05861>, 2020.
- Adams, J. W., Rodriguez, D., and Cox, R. A.: The uptake of SO₂ on Saharan dust: a flow tube study, *Atmos. Chem. Phys.*, 5, 2679-2689, <https://doi.org/10.5194/acpd-5-2643-2005>, 2005.
- 735 Alexander, B., Allman, D. J., Amos, H. M., Fairlie, T. D., Dachs, J., Hegg, D. A., and Sletten, R. S.: Isotopic constraints on the formation pathways of sulfate aerosol in the marine boundary layer of the subtropical northeast Atlantic Ocean, *J. Geophys. Res.: Atmos.*, 117, D6304, <https://doi.org/10.1029/2011JD016773>, 2012.
- Alexander, B., Park, R. J., Jacob, D. J., and Gong, S.: Transition metal-catalyzed oxidation of atmospheric sulfur: Global implications for the sulfur budget, *J. Geophys. Res.*, 114, D2309, <https://doi.org/10.1029/2008JD010486>, 2009.
- 740 Antoniou, M. G., Boraei, I., Solakidou, M., Deligiannakis, Y., Abhishek, M., Lawton, L. A., and Edwards, C.: Enhancing photocatalytic degradation of the cyanotoxin microcystin-LR with the addition of sulfate-radical generating oxidants, *J. Hazard. Mater.*, 360, 461-470, <https://doi.org/10.1016/j.jhazmat.2018.07.111>, 2018.
- Ault, A. P.: Aerosol Acidity: Novel Measurements and Implications for Atmospheric Chemistry, *Accounts Chem. Res.*, 53, 1703-1714, <https://doi.org/10.1021/acs.accounts.0c00303>, 2020.
- 745 Baltrusaitis, J., Cwiertny, D. M., and Grassian, V. H.: Adsorption of sulfur dioxide on hematite and goethite particle surfaces, *Phys. Chem. Chem. Phys.*, 9, 5542-5554, <https://doi.org/10.1039/b709167b>, 2007.
- Baltrusaitis, J., Jayaweera, P. M., and Grassian, V. H.: Sulfur Dioxide Adsorption on TiO₂ Nanoparticles: Influence of Particle Size, Coadsorbates, Sample Pretreatment, and Light on Surface Speciation and Surface Coverage, *J. Phys. Chem. C*, 115, 492-500, <https://doi.org/10.1021/jp108759b>, 2010.
- 750 Berglen, T. F., Berntsen, T. K., Isaksen, I. S. A., and Sundet, J. K.: A global model of the coupled sulfur/oxidant chemistry in the troposphere: The sulfur cycle, *J. Geophys. Res.*, 109, D19310, <https://doi.org/10.1029/2003JD003948>, 2004.
- Bian, H., and Zender, C. S.: Mineral dust and global tropospheric chemistry: Relative roles of photolysis and heterogeneous uptake, *J. Geophys. Res.: Atmos.*, 108, 4672, <https://doi.org/10.1029/2002JD003143>, 2003.
- 755 Chen, H., Nanayakkara, C. E., and Grassian, V. H.: Titanium dioxide photocatalysis in atmospheric chemistry, *Chem. Rev.*, 112, 5919-5948, <https://doi.org/10.1021/cr3002092>, 2012.
- Chen, Q., Schmidt, J. A., Shah, V., Jaeglé, L., Sherwen, T., and Alexander, B.: Sulfate production by reactive bromine: Implications for the global sulfur and reactive bromine budgets, *Geophys. Res. Lett.*, 44, 7069-7078, <https://doi.org/10.1002/2017GL073812>, 2017.
- 760 Chen, Y., Tong, S., Li, W., Liu, Y., Tan, F., Ge, M., Xie, X., and Sun, J.: Photocatalytic Oxidation of SO₂ by TiO₂: Aerosol Formation and the Key Role of Gaseous Reactive Oxygen Species, *Environ. Sci. Technol.*, 55, 9784-9793, <https://doi.org/10.1021/acs.est.1c01608>, 2021.
- Chen, Z., Liu, P., Wang, W., Cao, X., Liu, Y., Zhang, Y., and Ge, M.: Rapid Sulfate Formation via Uncatalyzed Autoxidation of Sulfur Dioxide in Aerosol Microdroplets, *Environ. Sci. Technol.*, 56, 7637-7646, <https://doi.org/10.1021/acs.est.2c00112>, 2022.
- 765 Cheng, Y., Zheng, G., Wei, C., Mu, Q., Zheng, B., Wang, Z., Gao, M., Zhang, Q., He, K., Carmichael, G., Pöschl, U., and Su, H.: Reactive nitrogen chemistry in aerosol water as a source of sulfate during haze events in China, *Sci. Adv.*, 2, e1601530, <https://doi.org/10.1126/sciadv.1601530>, 2016.
- 770 Chu, B., Wang, Y., Yang, W., Ma, J., Ma, Q., Zhang, P., Liu, Y., and He, H.: Effects of NO₂ and C₃H₆ on the heterogeneous oxidation of SO₂ on TiO₂ in the presence or absence of UV-Vis irradiation, *Atmos. Chem. Phys.*, 19, 14777-14790, <https://doi.org/10.5194/acp-19-14777-2019>, 2019.
- Chughtai, A. R., Brooks, M. E., and Smith, D. M.: Effect of metal oxides and black carbon (soot) on SO₂/O₂/H₂O reaction systems, *Aerosol Sci. Tech.*, 19, 121-132, <https://doi.org/10.1080/02786829308959626>, 1993.
- Clegg, M., and Abbatt, D.: Uptake of Gas-Phase SO₂ and H₂O₂ by Ice Surfaces: Dependence on Partial Pressure, Temperature,

- and Surface Acidity, *J. Phys. Chem. A*, 105, 6630-6636, <https://doi.org/10.1021/jp010062r>, 2001a.
- 775 Clegg, S. M., and Abbatt, J. P. D.: Oxidation of SO₂ by H₂O₂ on ice surfaces at 228 K: a sink for SO₂ in ice clouds, *Atmos. Chem. Phys.*, 1, 73-78, <https://doi.org/10.5194/acp-1-73-2001>, 2001b.
- Darif, B., Ojala, S., Pirault-Roy, L., Bensitel, M., Brahmi, R., and Keiski, R. L.: Study on the catalytic oxidation of DMDS over Pt-Cu catalysts supported on Al₂O₃, AlSi₂₀ and SiO₂, *Appl. Catal., B*, 181, 24-33, <https://doi.org/10.1016/j.apcatb.2015.07.050>, 2016.
- 780 Davidovits, P., Kolb, C. E., Williams, L. R., Jayne, J. T., and Worsnop, D. R.: Mass Accommodation and Chemical Reactions at Gas-Liquid Interfaces, *Chem. Rev.*, 106, 1323-1354, <https://doi.org/10.1021/cr040366k>, 2006.
- Ding, J., Zhao, P., Su, J., Dong, Q., Du, X., and Zhang, Y.: Aerosol pH and its driving factors in Beijing, *Atmos. Chem. Phys.*, 19, 7939-7954, <https://doi.org/10.5194/acp-19-7939-2019>, 2019.
- 785 Du, C., Kong, L., Zhanzakova, A., Tong, S., Yang, X., Wang, L., Fu, H., Cheng, T., Chen, J., and Zhang, S.: Impact of adsorbed nitrate on the heterogeneous conversion of SO₂ on α -Fe₂O₃ in the absence and presence of simulated solar irradiation, *Sci. Total Environ.*, 649, 1393-1402, <https://doi.org/10.1016/j.scitotenv.2018.08.295>, 2019.
- Dupart, Y., King, S. M., Nekat, B., Nowak, A., Wiedensohler, A., Herrmann, H., David, G., Thomas, B., Miffre, A., Rairoux, P., Anna, B. D., and George, C.: Mineral dust photochemistry induces nucleation events in the presence of SO₂, *Proc. Natl. Acad. Sci. U. S. A.*, 109, 20842-20847, <https://doi.org/10.1073/pnas.1212297109>, 2012.
- 790 Fairlie, T. D., Jacob, D. J., Dibb, J. E., Alexander, B., Avery, M. A., van Donkelaar, A., and Zhang, L.: Impact of mineral dust on nitrate, sulfate, and ozone in transpacific Asian pollution plumes, *Atmos. Chem. Phys.*, 10, 3999-4012, <https://doi.org/10.5194/acp-10-3999-2010>, 2010.
- Fan, M., Zhang, Y., Lin, Y., Li, J., Cheng, H., An, N., Sun, Y., Qiu, Y., Cao, F., and Fu, P.: Roles of Sulfur Oxidation Pathways in the Variability in Stable Sulfur Isotopic Composition of Sulfate Aerosols at an Urban Site in Beijing, China, *Environ. Sci. Technol. Lett.*, 7, 883-888, <https://doi.org/10.1021/acs.estlett.0c00623>, 2020.
- 795 Filonchik, M.: Characteristics of the severe March 2021 Gobi Desert dust storm and its impact on air pollution in China, *Chemosphere*, 287, 132219, <https://doi.org/10.1016/j.chemosphere.2021.132219>, 2022.
- Fu, H., Cwiertny, D. M., Carmichael, G. R., Scherer, M. M., and Grassian, V. H.: Photoreductive dissolution of Fe-containing mineral dust particles in acidic media, *J. Geophys. Res.*, 115, D11304, <https://doi.org/10.1029/2009JD012702>, 2010.
- 800 Fu, H., Wang, X., Wu, H., Yin, Y., and Chen, J.: Heterogeneous uptake and oxidation of SO₂ on iron oxides, *J. Phys. Chem. C*, 111, 6077-6085, <https://doi.org/10.1021/jp070087b>, 2007.
- Fu, H., Xu, T., Yang, S., Zhang, S., and Chen, J.: Photoinduced formation of Fe(III)-sulfato complexes on the surface of α -Fe₂O₃ and their photochemical performance, *J. Phys. Chem. C*, 113, 11316-11322, <https://doi.org/10.1021/jp8088275>, 2009.
- 805 Gao, J., Shi, G., Zhang, Z., Wei, Y., Tian, X., Feng, Y., Russell, A. G., and Nenes, A.: Targeting Atmospheric Oxidants Can Better Reduce Sulfate Aerosol in China: H₂O₂ Aqueous Oxidation Pathway Dominates Sulfate Formation in Haze, *Environ. Sci. Technol.*, <https://doi.org/10.1021/acs.est.2c01739>, 2022.
- Gaston, C. J., Pratt, K. A., Suski, K. J., May, N. W., Gill, T. E., and Prather, K. A.: Laboratory Studies of the Cloud Droplet Activation Properties and Corresponding Chemistry of Saline Playa Dust, *Environ. Sci. Technol.*, 51, 1348-1356, <https://doi.org/10.1021/acs.est.6b04487>, 2017.
- 810 Gen, M., Zhang, R., Huang, D., Li, Y., and Chan, C. K.: Heterogeneous Oxidation of SO₂ in Sulfate Production during Nitrate Photolysis at 300 nm: Effect of pH, Relative Humidity, Irradiation Intensity, and the Presence of Organic Compounds, *Environ. Sci. Technol.*, 53, 8757-8766, <https://doi.org/10.1021/acs.est.9b01623>, 2019.
- Goodman, A. L., Li, P., Usher, C. R., and Grassian, V. H.: Heterogeneous uptake of sulfur dioxide on aluminum and magnesium oxide particles, *J. Phys. Chem. A*, 105, 6109-6120, <https://doi.org/10.1021/jp004423z>, 2001.
- 815 Guan, C., Li, X., Luo, Y., and Huang, Z.: Heterogeneous Reaction of NO₂ on α -Al₂O₃ in the Dark and Simulated Sunlight, *J. Phys. Chem. A*, 118, 6999-7006, <https://doi.org/10.1021/jp503017k>, 2014.
- Han, L., Liu, X., Chen, Y., Xiang, X., Cheng, S., and Wang, H.: Key factors influencing the formation of sulfate aerosol on the surface of mineral aerosols: Insights from laboratory simulations and ACSM measurements, *Atmos. Environ.*, 253, 118341, <https://doi.org/10.1016/j.atmosenv.2021.118341>, 2021.
- 820 Hanson, D. R., Ravishankara, A. R., and Solomon, S.: Heterogeneous reactions in sulfuric acid aerosols: A framework for model calculations, *J. Geophys. Res.: Atmos.*, 99, 3615-3629, <https://doi.org/10.1029/93JD02932>, 1994.
- Harris, E., Sinha, B., Foley, S., Crowley, J. N., Borrmann, S., and Hoppe, P.: Sulfur isotope fractionation during heterogeneous oxidation of SO₂ on mineral dust, *Atmos. Chem. Phys.*, 12, 4867-4884, <https://doi.org/10.5194/acp-12-4867-2012>, 2012.

- Haynes, W. M.: Handbook of Chemistry and Physics, CRC Press, New York, 2014.
- 825 He, G., and He, H.: Water Promotes the Oxidation of SO₂ by O₂ over Carbonaceous Aerosols, *Environ. Sci. Technol.*, 54, 7070-7077, <https://doi.org/10.1021/acs.est.0c00021>, 2020.
- He, H., Wang, Y., Ma, Q., Ma, J., Chu, B., Ji, D., Tang, G., Liu, C., Zhang, H., and Hao, J.: Mineral dust and NO_x promote the conversion of SO₂ to sulfate in heavy pollution days, *Sci. Rep.-UK*, 4, 4172, <https://doi.org/10.1038/srep04172>, 2014.
- 830 He, P., Alexander, B., Geng, L., Chi, X., Fan, S., Zhan, H., Kang, H., Zheng, G., Cheng, Y., Su, H., Liu, C., and Xie, Z.: Isotopic constraints on heterogeneous sulfate production in Beijing haze, *Atmos. Chem. Phys.*, 18, 5515-5528, <https://doi.org/10.5194/acp-18-5515-2018>, 2018.
- He, X., and Zhang, Y.: Influence of relative humidity on SO₂ oxidation by O₃ and NO₂ on the surface of TiO₂ particles: Potential for formation of secondary sulfate aerosol, *Spectrochim. Acta, Part A*, 219, 121-128, <https://doi.org/10.1016/j.saa.2019.04.046>, 2019.
- 835 Hennigan, C. J., Izumi, J., Sullivan, A. P., Weber, R. J., and Nenes, A.: A critical evaluation of proxy methods used to estimate the acidity of atmospheric particles, *Atmos. Chem. Phys.*, 15, 2775-2790, <https://doi.org/10.5194/acp-15-2775-2015>, 2015.
- Himmelblau, D. M.: Diffusion of dissolved gases in liquids, *Chem. Rev.*, 64, 527-550, <https://doi.org/10.1021/cr60231a002>, 1964.
- 840 Huang, L., An, J., Koo, B., Yarwood, G., Yan, R., Wang, Y., Huang, C., and Li, L.: Enhanced sulfate formation through SO₂+NO₂ heterogeneous reactions during heavy winter haze in the Yangtze River Delta region, China, *Atmos. Chem. Phys.*, 19, 14311-14328, <https://doi.org/10.5194/acp-2019-292>, 2019.
- Huang, L., Zhao, Y., Li, H., and Chen, Z.: Kinetics of heterogeneous reaction of sulfur dioxide on authentic mineral dust: effects of relative humidity and hydrogen peroxide, *Environ. Sci. Technol.*, 49, 10797-10805, <https://doi.org/10.1021/acs.est.5b03930>, 2015.
- 845 Huang, L., Zhao, Y., Li, H., and Chen, Z.: Hydrogen peroxide maintains the heterogeneous reaction of sulfur dioxide on mineral dust proxy particles, *Atmos. Environ.*, 141, 552-559, <https://doi.org/10.1016/j.atmosenv.2016.07.035>, 2016.
- Huang, Z., Zhang, Z., Kong, W., Feng, S., Qiu, Y., Tang, S., Xia, C., Ma, L., Luo, M., and Xu, D.: Synergistic effect among Cl₂, SO₂ and NO₂ in their heterogeneous reactions on gamma-alumina, *Atmos. Environ.*, 166, 403-411, <https://doi.org/10.1016/j.atmosenv.2017.06.041>, 2017.
- 850 Hung, H., Hsu, M., and Hoffmann, M. R.: Quantification of SO₂ Oxidation on Interfacial Surfaces of Acidic Micro-Droplets: Implication for Ambient Sulfate Formation, *Environ. Sci. Technol.*, 52, 9079-9086, <https://doi.org/10.1021/acs.est.8b01391>, 2018.
- Hung, H., and Hoffmann, M. R.: Oxidation of Gas-Phase SO₂ on the Surfaces of Acidic Microdroplets: Implications for Sulfate and Sulfate Radical Anion Formation in the Atmospheric Liquid Phase, *Environ. Sci. Technol.*, 49, 13768-13776, <https://doi.org/10.1021/acs.est.5b01658>, 2015.
- 855 Jacob, D. J.: Heterogeneous chemistry and tropospheric ozone, *Atmos. Environ.*, 34, 2131-2159, [https://doi.org/10.1016/S1352-2310\(99\)00462-8](https://doi.org/10.1016/S1352-2310(99)00462-8), 2000.
- Jayne, J. T., and Davidovits, P.: Uptake of SO₂(g) by Aqueous Surfaces as a Function of pH: The Effect of Chemical Reaction at the Interface, *J. Phys. Chem.*, 94, 6041-6048, <https://doi.org/10.1021/j100378a076>, 1990.
- 860 Jin, X., Wang, Y., Li, Z., Zhang, F., Xu, W., Sun, Y., Fan, X., Chen, G., Wu, H., Ren, J., Wang, Q., and Cribb, M.: Significant contribution of organics to aerosol liquid water content in winter in Beijing, China, *Atmos. Chem. Phys.*, 20, 901-914, <https://doi.org/10.5194/acp-20-901-2020>, 2020.
- Ke, Z., Liu, X., Wu, M., Shan, Y., and Shi, Y.: Improved Dust Representation and Impacts on Dust Transport and Radiative Effect in CAM5, *J. Adv. Model. Earth Sy.*, 14, <https://doi.org/10.1029/2021MS002845>, 2022.
- 865 Keene, W. C., Pszenny, A. A. P., Maben, J. R., Stevenson, E., and Wall, A.: Closure evaluation of size-resolved aerosol pH in the New England coastal atmosphere during summer, *J. Geophys. Res.: Atmos.*, 109, D23307, <https://doi.org/10.1029/2004JD004801>, 2004.
- Kim, J., Choe, Y. J., Kim, S. H., and Jeong, K.: Enhancing the decomposition of refractory contaminants on SO₄²⁻-functionalized iron oxide to accommodate surface SO₄⁻-generated via radical transfer from ·OH, *Appl. Catal., B*, 252, 62-76, <https://doi.org/10.1016/j.apcatb.2019.04.015>, 2019.
- 870 Kong, L. D., Zhao, X., Sun, Z. Y., Yang, Y. W., Fu, H. B., Zhang, S. C., Cheng, T. T., Yang, X., Wang, L., and Chen, J. M.: The effect of nitrate on the heterogeneous uptake of sulfur dioxide on hematite, *Atmos. Chem. Phys.*, 14, 9451-9467, <https://doi.org/10.5194/acp-14-9451-2014>, 2014.

- 875 Kumar, R., Barth, M. C., Madronich, S., Naja, M., Carmichael, G. R., Pfister, G. G., Knote, C., Brasseur, G. P., Ojha, N., and Sarangi, T.: Effects of dust aerosols on tropospheric chemistry during a typical pre-monsoon season dust storm in northern India, *Atmos. Chem. Phys.*, 14, 6813-6834, <https://doi.org/10.5194/acp-14-6813-2014>, 2014.
- Laskin, A., Gaspar, D. J., Wang, W., Hunt, S. W., Cowin, J. P., Colson, S. D., and Finlayson-Pitts, B. J.: Reactions at Interfaces as a Source of Sulfate Formation in Sea-Salt Particles, *Science*, 301, 340-344, <https://doi.org/10.1126/science.1085374>, 2003.
- 880 Li, G., Bei, N., Cao, J., Huang, R., and Wu, J.: A possible pathway for rapid growth of sulfate during haze days in China, *Atmos. Chem. Phys.*, 17, 3301-3316, <https://doi.org/10.5194/acp-17-3301-2017>, 2017.
- Li, G., Lu, D., Yang, X., Zhang, H., Guo, Y., Qu, G., Wang, P., Chen, L., Ruan, T., Hou, X., Jin, X., Zhang, R., Tan, Q., Zhai, S., Ma, Y., Yang, R., Fu, J., Shi, J., Liu, G., Wang, Q., Liang, Y., Zhang, Q., Liu, Q., and Jiang, G.: Resurgence of Sandstorms Complicates China's Air Pollution Situation, *Environ. Sci. Technol.*, 55, 11467-11469, <https://doi.org/10.1021/acs.est.1c03724>, 2021a.
- 885 Li, J., Shang, J., and Zhu, T.: Heterogeneous reactions of SO₂ on ZnO particle surfaces, *Sci. China: Chem.*, 54, 161-166, <https://doi.org/10.1007/s11426-010-4167-9>, 2010.
- Li, J., Zhang, Y., Cao, F., Zhang, W., Fan, M., Lee, X., and Michalski, G.: Stable Sulfur Isotopes Revealed a Major Role of Transition-Metal Ion-Catalyzed SO₂ Oxidation in Haze Episodes, *Environ. Sci. Technol.*, 54, 2626-2634, <https://doi.org/10.1021/acs.est.9b07150>, 2020a.
- 890 Li, K., Fang, X., Wang, T., Gong, K., Ali Tahir, M., Wang, W., Han, J., Cheng, H., Xu, G., and Zhang, L.: Atmospheric organic complexation enhanced sulfate formation and iron dissolution on nano α -Fe₂O₃, *Environ. Sci.: Nano*, 8, 698-710, <https://doi.org/10.1039/D0EN01220C>, 2021b.
- Li, K., Kong, L., Zhanzakova, A., Tong, S., Shen, J., Wang, T., Chen, L., Li, Q., Fu, H., and Zhang, L.: Heterogeneous Conversion of SO₂ on Nano α -Fe₂O₃: the Effect of Morphology, Light Illumination and Relative Humidity, *Environ. Sci.: Nano*, 6, 1838-1851, <https://doi.org/10.1039/C9EN00097F>, 2019.
- 895 Li, L., Chen, Z. M., Zhang, Y. H., Zhu, T., Li, J. L., and Ding, J.: Kinetics and mechanism of heterogeneous oxidation of sulfur dioxide by ozone on surface of calcium carbonate, *Atmos. Chem. Phys.*, 6, 2453-2464, <https://doi.org/10.5194/acp-6-2453-2006>, 2006.
- 900 Li, L., Chen, Z. M., Zhang, Y. H., Zhu, T., Li, S., Li, H. J., Zhu, L. H., and Xu, B. Y.: Heterogeneous oxidation of sulfur dioxide by ozone on the surface of sodium chloride and its mixtures with other components, *J. Geophys. Res.*, 112, D18301, <https://doi.org/10.1029/2006JD008207>, 2007.
- Li, T., Wang, X., Chen, Y., Liang, J., and Zhou, L.: Producing 'OH, SO₄⁻ and 'O₂' in heterogeneous Fenton reaction induced by Fe₃O₄-modified schwertmannite, *Chem. Eng. J.*, 393, 124735, <https://doi.org/10.1016/j.ccej.2020.124735>, 2020b.
- 905 Li, S., Zhang, F., Jin, X., Sun, Y., Wu, H., Xie, C., Chen, L., Liu, J., Wu, T., Jiang, S., Cribb, M., and Li, Z.: Characterizing the ratio of nitrate to sulfate in ambient fine particles of urban Beijing during 2018-2019, *Atmos. Environ.*, 237, 117662, <https://doi.org/10.1016/j.atmosenv.2020.117662>, 2020c.
- Lind, J. A., Lazrus, A. L., and Kok, G. L.: Aqueous phase oxidation of sulfur(IV) by hydrogen peroxide, methylhydroperoxide, and peroxyacetic acid, *Journal of Geophysical Research*, 92, 4171-4177, <https://doi.org/10.1029/JD092iD04p04171>, 1987.
- 910 Liu, C., Ma, Q., Liu, Y., Ma, J., and He, H.: Synergistic reaction between SO₂ and NO₂ on mineral oxides: a potential formation pathway of sulfate aerosol, *Phys. Chem. Chem. Phys.*, 14, 1668-1676, <https://doi.org/10.1039/C1CP22217A>, 2012.
- Liu, J., Wu, D., Fan, S., Mao, X., and Chen, H.: A one-year, on-line, multi-site observational study on water-soluble inorganic ions in PM_{2.5} over the Pearl River Delta region, China, *Sci. Total Environ.*, 601-602, 1720-1732, <https://doi.org/10.1016/j.scitotenv.2017.06.039>, 2017c.
- 915 Liu, T., Chan, A. W. H., and Abbatt, J. P. D.: Multiphase Oxidation of Sulfur Dioxide in Aerosol Particles: Implications for Sulfate Formation in Polluted Environments, *Environ. Sci. Technol.*, 55, 4227-4242, <https://doi.org/10.1021/acs.est.0c06496>, 2021a.
- Liu, T., Clegg, S. L., and Abbatt, J. P. D.: Fast oxidation of sulfur dioxide by hydrogen peroxide in deliquesced aerosol particles, *Proc. Natl. Acad. Sci. U. S. A.*, 117, 1354-1359, <https://doi.org/10.1073/pnas.1916401117>, 2020a.
- 920 Liu, Y., Wang, T., Fang, X., Deng, Y., Cheng, H., Fu, H., and Zhang, L.: Impact of greenhouse gas CO₂ on the heterogeneous reaction of SO₂ on alpha-Al₂O₃, *Chinese Chem. Lett.*, 31, 2712-2716, <https://doi.org/10.1016/j.ccl.2020.04.037>, 2020b.
- Liu, T., and Abbatt, J. P. D.: Oxidation of sulfur dioxide by nitrogen dioxide accelerated at the interface of deliquesced aerosol particles, *Nat. Chem.*, 13, 1173-1177, <https://doi.org/10.1038/s41557-021-00777-0>, 2021.

- 925 Liu, W., He, X., Pang, S., and Zhang, Y.: Effect of relative humidity on O₃ and NO₂ oxidation of SO₂ on α-Al₂O₃ particles, *Atmos. Environ.*, 167, 245-253, <https://doi.org/10.1016/j.atmosenv.2017.08.028>, 2017b.
- Liu, X., Chen, W., and Jiang, H.: Facile synthesis of Ag/Ag₃PO₄/AMB composite with improved photocatalytic performance, *Chem. Eng. J.*, 308, 889-896, <https://doi.org/10.1016/j.cej.2016.09.125>, 2017a.
- 930 Liu, Y., Deng, Y., Liu, J., Fang, X., Wang, T., Li, K., Gong, K., Bacha, A. U., Nabi, I., Ge, Q., Zhang, X., George, C., and Zhang, L.: A novel pathway of atmospheric sulfate formation through carbonate radicals, *Atmos. Chem. Phys.*, 22, 9175-9197, <https://doi.org/10.5194/acp-22-9175-2022>, 2022.
- Liu, Y., Feng, Z., Zheng, F., Bao, X., Liu, P., Ge, Y., Zhao, Y., Jiang, T., Liao, Y., Zhang, Y., Fan, X., Yan, C., Chu, B., Wang, Y., Du, W., Cai, J., Bianchi, F., Petäjä, T., Mu, Y., He, H., and Kulmala, M.: Ammonium nitrate promotes sulfate formation through uptake kinetic regime, *Atmos. Chem. Phys.*, 21, 13269-13286, <https://doi.org/10.5194/acp-21-13269-2021>, 2021b.
- 935 Ma, Q., Liu, Y., Liu, C., Ma, J., and He, H.: A case study of Asian dust storm particles: Chemical composition, reactivity to SO₂ and hygroscopic properties, *J. Environ. Sci.*, 24, 62-71, [https://doi.org/10.1016/S1001-0742\(11\)60729-8](https://doi.org/10.1016/S1001-0742(11)60729-8), 2012.
- Ma, Q., Liu, Y., and He, H.: Synergistic effect between NO₂ and SO₂ in their adsorption and reaction on γ-Alumina, *J. Phys. Chem. A*, 112, 6630-6635, <https://doi.org/10.1021/jp802025z>, 2008.
- Ma, Q., Wang, L., Chu, B., Ma, J., and He, H.: Contrary Role of H₂O and O₂ in the Kinetics of Heterogeneous Photochemical Reactions of SO₂ on TiO₂, *J. Phys. Chem. A*, 123, 1311-1318, <https://doi.org/10.1021/acs.jpca.8b11433>, 2018.
- 940 Ma, Q., Wang, T., Liu, C., He, H., Wang, Z., Wang, W., and Liang, Y.: SO₂ initiates the efficient conversion of NO₂ to HONO on MgO surface, *Environ. Sci. Technol.*, 51, 3767-3775, <https://doi.org/10.1021/acs.est.6b05724>, 2017.
- Martin, M. A., Childers, J. W., and Palmer, R. A.: Fourier transform infrared photoacoustic spectroscopy characterization of sulfur-oxygen species resulting from the reaction of SO₂ with CaO and CaCO₃, *Appl. Spectrosc.*, 41, 120-126, <https://doi.org/10.1366/0003702874868151>, 1987.
- 945 Maters, E. C., Delmelle, P., Rossi, M. J., and Ayris, P. M.: Reactive Uptake of Sulfur Dioxide and Ozone on Volcanic Glass and Ash at Ambient Temperature, *J. Geophys. Res.: Atmos.*, 122, 10077-10088, <https://doi.org/10.1002/2017JD026993>, 2017.
- Mauldin III, R. L., Berndt, T., Sipilä, M., Paasonen, P., Petäjä, T., Kim, S., Kurtén, T., Stratmann, F., Kerminen, V. M., and Kulmala, M.: A new atmospherically relevant oxidant of sulphur dioxide, *Nature*, 488, 193-196, <https://doi.org/10.1038/nature11278>, 2012.
- 950 Nanayakkara, C. E., Larish, W. A., and Grassian, V. H.: Titanium dioxide nanoparticle surface reactivity with atmospheric gases, CO₂, SO₂, and NO₂: roles of surface hydroxyl groups and adsorbed water in the formation and stability of adsorbed products, *J. Phys. Chem. C*, 118, 23011-23021, <https://doi.org/10.1021/jp504402z>, 2014.
- Nanayakkara, C. E., Pettibone, J., and Grassian, V. H.: Sulfur dioxide adsorption and photooxidation on isotopically-labeled titanium dioxide nanoparticle surfaces: roles of surface hydroxyl groups and adsorbed water in the formation and stability of adsorbed sulfite and sulfate, *Phys. Chem. Chem. Phys.*, 14, 6957-6966, <https://doi.org/10.1039/c2cp23684b>, 2012.
- 955 Ndour, M., Nicolas, M., D Anna, B., Ka, O., and George, C.: Photoreactivity of NO₂ on mineral dusts originating from different locations of the Sahara desert, *Phys. Chem. Chem. Phys.*, 11, 1312-1319, <https://doi.org/10.1039/b806441e>, 2009.
- 960 Park, J., Ivanov, A. V., and Molina, M. J.: Effect of Relative Humidity on OH Uptake by Surfaces of Atmospheric Importance, *J. Phys. Chem. A*, 112, 6968-6977, <https://doi.org/10.1021/jp8012317>, 2008.
- Park, J., Jang, M., and Yu, Z.: Heterogeneous Photo-oxidation of SO₂ in the Presence of Two Different Mineral Dust Particles: Gobi and Arizona Dust, *Environ. Sci. Technol.*, 51, 9605-9613, <https://doi.org/10.1021/acs.est.7b00588>, 2017.
- Park, J., and Jang, M.: Heterogeneous photooxidation of sulfur dioxide in the presence of airborne mineral dust particles, *RSC Adv.*, 6, 58617-58627, <https://doi.org/10.1039/C6RA09601H>, 2016.
- 965 Peak, D., Ford, R. G., and Sparks, D. L.: An in situ ATR-FTIR investigation of sulfate bonding mechanisms on Goethite., *J. Colloid Interf. Sci.*, 218, 289-299, <https://doi.org/10.1006/jcis.1999.6405>, 1999.
- Persson, P., and Vgren, L. L.: Potentiometric and spectroscopic studies of sulfate complexation at the goethite-water interface, *Geochim. Cosmochim. Ac.*, 60, 2789-2799, [https://doi.org/10.1016/0016-7037\(96\)00124-X](https://doi.org/10.1016/0016-7037(96)00124-X), 1996.
- 970 Pye, H. O. T., Nenes, A., Alexander, B., Ault, A. P., Barth, M. C., Clegg, S. L., Collett Jr., J. L., Fahey, K. M., Hennigan, C. J., Herrmann, H., Kanakidou, M., Kelly, J. T., Ku, I., McNeill, V. F., Riemer, N., Schaefer, T., Shi, G., Tilgner, A., Walker, J. T., Wang, T., Weber, R., Xing, J., Zaveri, R. A., and Zuend, A.: The acidity of atmospheric particles and clouds, *Atmos. Chem. Phys.*, 20, 4809-4888, <https://doi.org/10.5194/acp-20-4809-2020>, 2020.
- Ravishankara, A. R.: Heterogeneous and Multiphase Chemistry in the Troposphere, *Science*, 276, 1058-1064,

<https://doi.org/10.1126/science.276.5315.1058>, 1997.

- 975 Ren, Y., Wei, J., Wu, Z., Ji, Y., Bi, F., Gao, R., Wang, X., Wang, G., and Li, H.: Chemical components and source identification of PM_{2.5} in non-heating season in Beijing: The influences of biomass burning and dust, *Atmos. Res.*, 251, 105412, <https://doi.org/10.1016/j.atmosres.2020.105412>, 2021.
- Rindelaub, J. D., Craig, R. L., Nandy, L., Bondy, A. L., Dutcher, C. S., Shepson, P. B., and Ault, A. P.: Direct Measurement of pH in Individual Particles via Raman Microspectroscopy and Variation in Acidity with Relative Humidity, *J. Phys. Chem. A*, 120, 911-917, <https://doi.org/10.1021/acs.jpca.5b12699>, 2016.
- 980 Rossi, M. J.: Heterogeneous Reactions on Salts, *Chem. Rev.*, 103, 4823-4882, <https://doi.org/10.1021/cr020507n>, 2003.
- Rubasinghege, G., and Grassian, V. H.: Role(s) of adsorbed water in the surface chemistry of environmental interfaces, *Chem. Commun.*, 49, 3071-3094, <https://doi.org/10.1039/C3CC38872G>, 2013.
- 985 Sakata, K., Takahashi, Y., Takano, S., Matsuki, A., Sakaguchi, A., and Tanimoto, H.: First X-ray Spectroscopic Observations of Atmospheric Titanium Species: Size Dependence and the Emission Source, *Environ. Sci. Technol.*, 55, 10975-10986, <https://doi.org/10.1021/acs.est.1c02000>, 2021.
- Sarwar, G., Fahey, K., Kwok, R., Gilliam, R. C., Roselle, S. J., Mathur, R., Xue, J., Yu, J., and Carter, W. P. L.: Potential impacts of two SO₂ oxidation pathways on regional sulfate concentrations: Aqueous-phase oxidation by NO₂ and gas-phase oxidation by Stabilized Criegee Intermediates, *Atmos. Environ.*, 68, 186-197, <https://doi.org/10.1016/j.atmosenv.2012.11.036>, 2013.
- 990 Scheinhardt, S., Müller, K., Spindler, G., and Herrmann, H.: Complexation of trace metals in size-segregated aerosol particles at nine sites in Germany, *Atmos. Environ.*, 74, 102-109, <https://doi.org/10.1016/j.atmosenv.2013.03.023>, 2013.
- Seinfeld, J. and Pandis, S.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, 3rd edn., Wiley, ISBN 978-1-118-94740-1, 2016.
- 995 Shang, J., Li, J., and Zhu, T.: Heterogeneous reaction of SO₂ on TiO₂ particles, *Science China Chemistry*, 53, 2637-2643, <https://doi.org/10.1007/s11426-010-4160-3>, 2010.
- Shao, J., Chen, Q., Wang, Y., Lu, X., He, P., Sun, Y., Shah, V., Martin, R. V., Philip, S., Song, S., Zhao, Y., Xie, Z., Zhang, L., and Alexander, B.: Heterogeneous sulfate aerosol formation mechanisms during wintertime Chinese haze events: air quality model assessment using observations of sulfate oxygen isotopes in Beijing, *Atmos. Chem. Phys.*, 19, 6107-6123, <https://doi.org/10.5194/acp-19-6107-2019>, 2019.
- 1000 Shi, Z., Krom, M. D., Jickells, T. D., Bonneville, S., Carslaw, K. S., Mihalopoulos, N., Baker, A. R., and Benning, L. G.: Impacts on iron solubility in the mineral dust by processes in the source region and the atmosphere: A review, *Aeolian Res.*, 5, 21-42, <https://doi.org/10.1016/j.aeolia.2012.03.001>, 2012.
- Song, H., Lu, K., Ye, C., Dong, H., Li, S., Chen, S., Wu, Z., Zheng, M., Zeng, L., Hu, M., and Zhang, Y.: A comprehensive observation-based multiphase chemical model analysis of sulfur dioxide oxidations in both summer and winter, *Atmos. Chem. Phys.*, 21, 13713-13727, <https://doi.org/10.5194/acp-21-13713-2021>, 2021.
- 1005 Song, S., Nenes, A., Gao, M., Zhang, Y., Liu, P., Shao, J., Ye, D., Xu, W., Lei, L., Sun, Y., Liu, B., Wang, S., and McElroy, M. B.: Thermodynamic Modeling Suggests Declines in Water Uptake and Acidity of Inorganic Aerosols in Beijing Winter Haze Events during 2014/2015–2018/2019, *Environ. Sci. Technol. Lett.*, 6, 752-760, <https://doi.org/10.1021/acs.estlett.9b00621>, 2019.
- 1010 Stelson, A. W., and Seinfeld, J. H.: Chemical Mass Accounting of Urban Aerosol, *Environ. Sci. Technol.*, 15, 671-679, <https://doi.org/10.1021/es00088a005>, 1981.
- Su, H., Cheng, Y., and Pöschl, U.: New Multiphase Chemical Processes Influencing Atmospheric Aerosols, Air Quality, and Climate in the Anthropocene, *Accounts Chem. Res.*, 53, 2034-2043, <https://doi.org/10.1021/acs.accounts.0c00246>, 2020.
- 1015 Sullivan, R. C., Guazzotti, S. A., Sodeman, D. A., and Prather, K. A.: Direct observations of the atmospheric processing of Asian mineral dust, *Atmos. Chem. Phys.*, 7, 1213-1236, <https://doi.org/10.5194/acp-7-1213-2007>, 2007.
- Tang, M., Cziczo, D. J., and Grassian, V. H.: Interactions of water with mineral dust aerosol: water adsorption, hygroscopicity, cloud condensation, and ice nucleation, *Chem. Rev.*, 116, 4205-4259, <https://doi.org/10.1021/acs.chemrev.5b00529>, 2016.
- Tang, M., Huang, X., Lu, K., Ge, M., Li, Y., Cheng, P., Zhu, T., Ding, A., Zhang, Y., Gligorovski, S., Song, W., Ding, X., Bi, X., and Wang, X.: Heterogeneous reactions of mineral dust aerosol: implications for tropospheric oxidation capacity, *Atmos. Chem. Phys.*, 17, 11727-11777, <https://doi.org/10.5194/acp-17-11727-2017>, 2017.
- 1020 Tang, M., Zhang, H., Gu, W., Gao, J., Jian, X., Shi, G., Zhu, B., Xie, L., Guo, L., Gao, X., Wang, Z., Zhang, G., and Wang, X.: Hygroscopic Properties of Saline Mineral Dust From Different Regions in China: Geographical Variations,

- Compositional Dependence, and Atmospheric Implications, *J. Geophys. Res.: Atmos.*, 124, 10844-10857, <https://doi.org/10.1029/2019JD031128>, 2019.
- 1025 Tao, W., Su, H., Zheng, G., Wang, J., Wei, C., Liu, L., Ma, N., Li, M., Zhang, Q., Pöschl, U., and Cheng, Y.: Aerosol pH and chemical regimes of sulfate formation in aerosol water during winter haze in the North China Plain, *Atmos. Chem. Phys.*, 20, 11729-11746, <https://doi.org/10.5194/acp-20-11729-2020>, 2020.
- 1030 Textor, C., Schulz, M., Guibert, S., Kinne, S., Balkanski, Y., Bauer, S., Berntsen, T., Berglen, T., Boucher, O., Chin, M., Dentener, F., Diehl, T., Easter, R., Feichter, H., Fillmore, D., Ghan, S., Ginoux, P., Gong, S., Grini, A., Hendricks, J., Horowitz, L., Huang, P., Isaksen, I., Iversen, T., Kloster, S., Koch, D., A. Kirkevåg, Kristjansson, J. E., Krol, M., Lauer, A., Lamarque, J. F., Liu, X., Montanaro, V., Myhre, G., Penner, J., Pitari, G., Reddy, S., Seland, Ø., Stier, P., Takemura, T., and Tie, X.: Analysis and quantification of the diversities of aerosol life cycles within AeroCom, *Atmos. Chem. Phys.*, 6, 1777-1813, <https://doi.org/10.5194/acp-6-1777-2006>, 2006.
- 1035 Tian, R., Ma, X., Sha, T., Pan, X., and Wang, Z.: Exploring dust heterogeneous chemistry over China: Insights from field observation and GEOS-Chem simulation, *Sci. Total Environ.*, 798, 149307, <https://doi.org/10.1016/j.scitotenv.2021.149307>, 2021.
- 1040 Tilgner, A., Schaefer, T., Alexander, B., Barth, M., Collett Jr., J. L., Fahey, K. M., Nenes, A., Pye, H. O. T., Herrmann, H., and McNeill, V. F.: Acidity and the multiphase chemistry of atmospheric aqueous particles and clouds, *Atmos. Chem. Phys.*, 21, 13483-13536, <https://doi.org/10.5194/acp-21-13483-2021>, 2021.
- Tutsak, E., and Koçak, M.: High time-resolved measurements of water-soluble sulfate, nitrate and ammonium in PM_{2.5} and their precursor gases over the Eastern Mediterranean, *Sci. Total Environ.*, 672, 212-226, <https://doi.org/10.1016/j.scitotenv.2019.03.451>, 2019.
- 1045 Ullerstam, M., Johnson, M. S., Vogt, R., and Ljungstrom, E.: DRIFTS and Knudsen cell study of the heterogeneous reactivity of SO₂ and NO₂ on mineral dust, *Atmos. Chem. Phys.*, 3, 2043-2051, <https://doi.org/10.5194/acp-3-2043-2003>, 2003.
- Ullerstam, M., Vogt, R., Langer, S., and Ljungström, E.: The kinetics and mechanism of SO₂ oxidation by O₃ on mineral dust, *Phys. Chem. Chem. Phys.*, 4, 4694-4699, <https://doi.org/10.1039/B203529B>, 2002.
- Uno, I., Eguchi, K., Yumimoto, K., Takemura, T., Shimizu, A., Uematsu, M., Liu, Z., Wang, Z., Hara, Y., and Sugimoto, N.: Asian dust transported one full circuit around the globe, *Nat. Geosci.*, 2, 557-560, <https://doi.org/10.1038/ngeo583>, 2009.
- 1050 Urupina, D., Gaudion, V., Romanias, M. N., Verrielle, M., and Thevenet, F.: Method development and validation for the determination of sulfites and sulfates on the surface of mineral atmospheric samples using reverse-phase liquid chromatography, *Talanta*, 219, 121318, <https://doi.org/10.1016/j.talanta.2020.121318>, 2020.
- Urupina, D., Gaudion, V., Romanias, M. N., and Thevenet, F.: Surface Distribution of Sulfites and Sulfates on Natural Volcanic and Desert Dusts: Impact of Humidity and Chemical Composition, *ACS Earth Space Chem.*, 6, 642-655, <https://doi.org/10.1021/acsearthspacechem.1c00321>, 2022.
- 1055 Urupina, D., Lasne, J., Romanias, M. N., Thiery, V., Dagsson-Waldhauserova, P., and Thevenet, F.: Uptake and surface chemistry of SO₂ on natural volcanic dusts, *Atmos. Environ.*, 217, 116942, <https://doi.org/10.1016/j.atmosenv.2019.116942>, 2019.
- Urupina, D., Romanias, M. N., and Thevenet, F.: How Relevant Is It to Use Mineral Proxies to Mimic the Atmospheric Reactivity of Natural Dust Samples? A Reactivity Study Using SO₂ as Probe Molecule, *Minerals*, 11, 282, <https://doi.org/10.3390/min11030282>, 2021.
- 1060 Usher, C. R., Al-Hosney, H., Carlos-Cuellar, S., and Grassian, V. H.: A laboratory study of the heterogeneous uptake and oxidation of sulfur dioxide on mineral dust particles, *J. Geophys. Res.: Atmos.*, 107, 4713, <https://doi.org/10.1029/2002JD002051>, 2002.
- 1065 Usher, C. R., Michel, A. E., and Grassian, V. H.: Reactions on mineral dust, *Chem. Rev.*, 103, 4883-4940, <https://doi.org/10.1021/cr020657y>, 2003.
- Volkamer, R., San Martini, F., Molina, L. T., Salcedo, D., Jimenez, J. L., and Molina, M. J.: A missing sink for gas-phase glyoxal in Mexico City: Formation of secondary organic aerosol, *Geophys. Res. Lett.*, 34, <https://doi.org/10.1029/2007GL030752>, 2007.
- 1070 Wang, G., Zhang, R., Gomez, M. E., Yang, L., Levy Zamora, M., Hu, M., Lin, Y., Peng, J., Guo, S., Meng, J., Li, J., Cheng, C., Hu, T., Ren, Y., Wang, Y., Gao, J., Cao, J., An, Z., Zhou, W., Li, G., Wang, J., Tian, P., Marrero-Ortiz, W., Secrest, J., Du, Z., Zheng, J., Shang, D., Zeng, L., Shao, M., Wang, W., Huang, Y., Wang, Y., Zhu, Y., Li, Y., Hu, J., Pan, B., Cai, L., Cheng, Y., Ji, Y., Zhang, F., Rosenfeld, D., Liss, P. S., Duce, R. A., Kolb, C. E., and Molina, M. J.: Persistent sulfate

- 1075 formation from London Fog to Chinese haze, *Proc. Natl. Acad. Sci. U. S. A.*, 113, 13630-13635, <https://doi.org/10.1073/pnas.1616540113>, 2016.
- Wang, H., Zhong, C., Ma, Q., Ma, J., and He, H.: The adsorption and oxidation of SO₂ on MgO surface: experimental and DFT calculation studies, *Environ. Sci.: Nano*, 7, 1092-1101, <https://doi.org/10.1039/C9EN01474H>, 2020b.
- Wang, K., Hattori, S., Lin, M., Ishino, S., Alexander, B., Kamezaki, K., Yoshida, N., and Kang, S.: Isotopic constraints on atmospheric sulfate formation pathways in the Mt. Everest region, southern Tibetan Plateau, *Atmos. Chem. Phys.*, 21, 8357-8376, <https://doi.org/10.5194/acp-21-8357-2021>, 2021b.
- 1080 Wang, K., Zhang, Y., Nenes, A., and Fountoukis, C.: Implementation of dust emission and chemistry into the Community Multiscale Air Quality modeling system and initial application to an Asian dust storm episode, *Atmos. Chem. Phys.*, 12, 10209-10237, <https://doi.org/10.5194/acp-12-10209-2012>, 2012.
- Wang, R., Yang, N., Li, J., Xu, L., Tsona, N. T., Du, L., and Wang, W.: Heterogeneous reaction of SO₂ on CaCO₃ particles: Different impacts of NO₂ and acetic acid on the sulfite and sulfate formation, *J. Environ. Sci.*, <https://doi.org/10.1016/j.jes.2021.08.017>, 2022b.
- 1085 Wang, S., Wang, L., Fan, X., Wang, N., Ma, S., and Zhang, R.: Formation pathway of secondary inorganic aerosol and its influencing factors in Northern China: Comparison between urban and rural sites, *Sci. Total Environ.*, 840, 156404, <https://doi.org/10.1016/j.scitotenv.2022.156404>, 2022a.
- 1090 Wang, T.: Date for “Significant formation of sulfate aerosols contributed by the heterogeneous drivers of dust surface”, Mendeley Data [data set], <https://doi.org/10.17632/hyvdz7khs6.1>, 2022.
- Wang, T., Liu, Y., Deng, Y., Cheng, H., Fang, X., and Zhang, L.: Heterogeneous Formation of Sulfur Species on Manganese Oxides: Effects of Particle Type and Moisture Condition, *J. Phys. Chem. A*, 124, 7300-7312, <https://doi.org/10.1021/acs.jpca.0c04483>, 2020d.
- 1095 Wang, T., Liu, Y., Deng, Y., Cheng, H., Yang, Y., Feng, Y., Zhang, L., Fu, H., and Chen, J.: Photochemical Oxidation of Water-Soluble Organic Carbon (WSOC) on Mineral Dust and Enhanced Organic Ammonium Formation, *Environ. Sci. Technol.*, 54, 15631-15642, <https://doi.org/10.1021/acs.est.0c04616>, 2020c.
- Wang, T., Liu, Y., Deng, Y., Fu, H., Zhang, L., and Chen, J.: The influence of temperature on the heterogeneous uptake of SO₂ on hematite particles, *Sci. Total Environ.*, 644, 1493-1502, <https://doi.org/10.1016/j.scitotenv.2018.07.046>, 2018a.
- 1100 Wang, T., Liu, Y., Deng, Y., Fu, H., Zhang, L., and Chen, J.: Emerging investigator series: heterogeneous reactions of sulfur dioxide on mineral dust nanoparticles: from single component to mixed components, *Environ. Sci.: Nano*, 5, 1821-1833, <https://doi.org/10.1039/C8EN00376A>, 2018b.
- Wang, T., Liu, Y., Deng, Y., Fu, H., Zhang, L., and Chen, J.: Adsorption of SO₂ on mineral dust particles influenced by atmospheric moisture, *Atmos. Environ.*, 191, 153-161, <https://doi.org/10.1016/j.atmosenv.2018.08.008>, 2018c.
- 1105 Wang, T., Liu, M., Liu, M., Song, Y., Xu, Z., Shang, F., Huang, X., Liao, W., Wang, W., Ge, M., Cao, J., Hu, J., Tang, G., Pan, Y., Hu, M., and Zhu, T.: Sulfate Formation Apportionment during Winter Haze Events in North China, *Environ. Sci. Technol.*, 56, 7771-7778, <https://doi.org/10.1021/acs.est.2c02533>, 2022c.
- Wang, W., Liu, M., Wang, T., Song, Y., Zhou, L., Cao, J., Hu, J., Tang, G., Chen, Z., Li, Z., Xu, Z., Peng, C., Lian, C., Chen, Y., Pan, Y., Zhang, Y., Sun, Y., Li, W., Zhu, T., Tian, H., and Ge, M.: Sulfate formation is dominated by manganese-catalyzed oxidation of SO₂ on aerosol surfaces during haze events, *Nat. Commun.*, 12, 1993, <https://doi.org/10.1038/s41467-021-22091-6>, 2021a.
- 1110 Wang, X., Gemayel, R., Hayeck, N., Perrier, S., Charbonnel, N., Xu, C., Chen, H., Zhu, C., Zhang, L., Wang, L., Nizkorodov, S. A., Wang, X., Wang, Z., Wang, T., Mellouki, A., Riva, M., Chen, J., and George, C.: Atmospheric Photosensitization: A New Pathway for Sulfate Formation, *Environ. Sci. Technol.*, 54, 3114-3120, <https://doi.org/10.1021/acs.est.9b06347>, 2020a.
- 1115 Wang, Y., Zhang, Q., Jiang, J., Zhou, W., Wang, B., He, K., Duan, F., Zhang, Q., Philip, S., and Xie, Y.: Enhanced sulfate formation during China's severe winter haze episode in January 2013 missing from current models, *J. Geophys. Res.*, 119, 10425-10440, <https://doi.org/10.1002/2013jd021426>, 2014.
- Wang, Z., Wang, T., Fu, H., Zhang, L., Tang, M., George, C., Grassian, V. H., and Chen, J.: Enhanced heterogeneous uptake of sulfur dioxide on mineral particles through modification of iron speciation during simulated cloud processing, *Atmos. Chem. Phys.*, 19, 12569-12585, <https://doi.org/10.5194/acp-19-12569-2019>, 2019.
- 1120 Wu, C., Zhang, S., Wang, G., Lv, S., Li, D., Liu, L., Li, J., Liu, S., Du, W., Meng, J., Qiao, L., Zhou, M., Huang, C., and Wang, H.: Efficient Heterogeneous Formation of Ammonium Nitrate on the Saline Mineral Particle Surface in the Atmosphere of

East Asia during Dust Storm Periods, *Environ. Sci. Technol.*, 54, 15622-15630, <https://doi.org/10.1021/acs.est.0c04544>, 2020.

1125 Wu, L. Y., Tong, S. R., Wang, W. G., and Ge, M. F.: Effects of temperature on the heterogeneous oxidation of sulfur dioxide by ozone on calcium carbonate, *Atmos. Chem. Phys.*, 11, 6593-6605, <https://doi.org/10.5194/acp-11-6593-2011>, 2011.

Wu, L., Tong, S., Zhou, L., Wang, W., and Ge, M.: Synergistic effects between SO₂ and HCOOH on α -Fe₂O₃, *J. Phys. Chem. A*, 117, 3972-3979, <https://doi.org/10.1021/jp400195f>, 2013.

1130 Wu, Z., Wang, Y., Tan, T., Zhu, Y., Li, M., Shang, D., Wang, H., Lu, K., Guo, S., Zeng, L., and Zhang, Y.: Aerosol Liquid Water Driven by Anthropogenic Inorganic Salts: Implying Its Key Role in Haze Formation over the North China Plain, *Environ. Sci. Technol. Lett.*, 5, 160-166, <https://doi.org/10.1021/acs.estlett.8b00021>, 2018.

Xu, M., Qiu, P., He, Y., Guo, S., Bai, Y., Zhang, H., Zhao, S., Shen, X., Zhu, B., Guo, Q., and Guo, Z.: Sulfur isotope composition during heterogeneous oxidation of SO₂ on mineral dust: The effect of temperature, relative humidity, and light intensity, *Atmos. Res.*, 254, 105513, <https://doi.org/10.1016/j.atmosres.2021.105513>, 2021.

1135 Xu, W., Kuang, Y., Liang, L., He, Y., Cheng, H., Bian, Y., Tao, J., Zhang, G., Zhao, P., Ma, N., Zhao, H., Zhou, G., Su, H., Cheng, Y., Xu, X., Shao, M., and Sun, Y.: Dust-Dominated Coarse Particles as a Medium for Rapid Secondary Organic and Inorganic Aerosol Formation in Highly Polluted Air, *Environ. Sci. Technol.*, 54, 15710-15721, <https://doi.org/10.1021/acs.est.0c07243>, 2020.

1140 Xue, J., Yuan, Z., Griffith, S. M., Yu, X., Lau, A. K. H., and Yu, J. Z.: Sulfate formation enhanced by a cocktail of high NO_x, SO₂, particulate matter, and droplet pH during haze-fog events in megacities in China: an observation-based modeling investigation, *Environ. Sci. Technol.*, 50, 7325-7334, <https://doi.org/10.1021/acs.est.6b00768>, 2016.

Yang, L., Yu, L. E., and Ray, M. B.: Degradation of paracetamol in aqueous solutions by TiO₂ photocatalysis, *Water Res.*, 42, 3480-3488, <https://doi.org/10.1016/j.watres.2008.04.023>, 2008.

1145 Yang, N., Tsona, N. T., Cheng, S., Li, S., Xu, L., Wang, Y., Wu, L., and Du, L.: Competitive reactions of SO₂ and acetic acid on α -Al₂O₃ and CaCO₃ particles, *Sci. Total Environ.*, 699, 134362, <https://doi.org/10.1016/j.scitotenv.2019.134362>, 2020.

Yang, W., Chen, M., Xiao, W., Guo, Y., Ding, J., Zhang, L., and He, H.: Molecular Insights into NO-Promoted Sulfate Formation on Model TiO₂ Nanoparticles with Different Exposed Facets, *Environ. Sci. Technol.*, 52, 14110-14118, <https://doi.org/10.1021/acs.est.8b02688>, 2018a.

1150 Yang, W., He, H., Ma, Q., Ma, J., Liu, Y., Liu, P., and Mu, Y.: Synergistic formation of sulfate and ammonium resulting from reaction between SO₂ and NH₃ on typical mineral dust, *Phys. Chem. Chem. Phys.*, 18, 956-964, <https://doi.org/10.1039/C5CP06144J>, 2016.

Yang, W., Ma, Q., Liu, Y., Ma, J., Chu, B., Wang, L., and He, H.: Role of NH₃ in the Heterogeneous Formation of Secondary Inorganic Aerosols on Mineral Oxides, *J. Phys. Chem. A*, 122, 6311-6320, <https://doi.org/10.1021/acs.jpca.8b05130>, 2018b.

1155 Yang, W., Ma, Q., Liu, Y., Ma, J., Chu, B., and He, H.: The effect of water on the heterogeneous reactions of SO₂ and NH₃ on the surfaces of α -Fe₂O₃ and γ -Al₂O₃, *Environ. Sci.: Nano*, 6, 2749-2758, <https://doi.org/10.1039/C9EN00574A>, 2019.

Yang, W., Zhang, J., Ma, Q., Zhao, Y., Liu, Y., and He, H.: Heterogeneous reaction of SO₂ on manganese oxides: the effect of crystal structure and relative humidity, *Sci. Rep.-UK*, 7, 4550, <https://doi.org/10.1038/s41598-017-04551-6>, 2017.

1160 Ye, C., Liu, P., Ma, Z., Xue, C., Zhang, C., Zhang, Y., Liu, J., Liu, C., Sun, X., and Mu, Y.: High H₂O₂ Concentrations Observed during Haze Periods during the Winter in Beijing: Importance of H₂O₂ Oxidation in Sulfate Formation, *Environ. Sci. Technol. Lett.*, 5, 757-763, <https://doi.org/10.1021/acs.estlett.8b00579>, 2018.

Ye, C., Lu, K., Song, H., Mu, Y., Chen, J., and Zhang, Y.: A critical review of sulfate aerosol formation mechanisms during winter polluted periods, *J. Environ. Sci.*, <https://doi.org/10.1016/j.jes.2022.07.011>, 2022.

1165 Yin, Z., Wan, Y., Zhang, Y., and Wang, H.: Why super sandstorm 2021 in North China, *Natl. Sci. Rev.*, <https://doi.org/10.1093/nsr/nwab165>, 2021.

Yu, J.: An interfacial role for NO₂, *Nat. Chem.*, 13, 1158-1160, <https://doi.org/10.1038/s41557-021-00845-5>, 2021.

Yu, T., Zhao, D., Song, X., and Zhu, T.: NO₂-initiated multiphase oxidation of SO₂ by O₂ on CaCO₃ particles, *Atmos. Chem. Phys.*, 18, 6679-6689, <https://doi.org/10.5194/acp-18-6679-2018>, 2018.

1170 Yu, Z., Jang, M., and Park, J.: Modeling atmospheric mineral aerosol chemistry to predict heterogeneous photooxidation of SO₂, *Atmos. Chem. Phys.*, 17, 10001-10017, <https://doi.org/10.5194/acp-17-10001-2017>, 2017.

Yue, F., Xie, Z., Zhang, P., Song, S., He, P., Liu, C., Wang, L., Yu, X., and Kang, H.: The role of sulfate and its corresponding S(IV)+NO₂ formation pathway during the evolution of haze in Beijing, *Sci. Total Environ.*, 687, 741-751, <https://doi.org/10.1016/j.scitotenv.2019.06.096>, 2019.

- 1175 Zhang, F., Wang, Y., Peng, J., Chen, L., Sun, Y., Duan, L., Ge, X., Li, Y., Zhao, J., Liu, C., Zhang, X., Zhang, G., Pan, Y., Wang, Y., Zhang, A. L., Ji, Y., Wang, G., Hu, M., Molina, M. J., and Zhang, R.: An unexpected catalyst dominates formation and radiative forcing of regional haze, *Proc. Natl. Acad. Sci. U. S. A.*, 117, 3960-3966, <https://doi.org/10.1073/pnas.1919343117>, 2020.
- Zhang, H., Xu, Y., and Jia, L.: A chamber study of catalytic oxidation of SO₂ by Mn²⁺/Fe³⁺ in aerosol water, *Atmos. Environ.*, 245, 118019, <https://doi.org/10.1016/j.atmosenv.2020.118019>, 2021.
- 1180 Zhang, R., Wang, G., Guo, S., Zamora, M. L., Ying, Q., Lin, Y., Wang, W., Hu, M., and Wang, Y.: Formation of urban fine particulate matter, *Chem. Rev.*, 115, 3803-3855, <https://doi.org/10.1021/acs.chemrev.5b00067>, 2015.
- Zhang, S., Xing, J., Sarwar, G., Ge, Y., He, H., Duan, F., Zhao, Y., He, K., Zhu, L., and Chu, B.: Parameterization of heterogeneous reaction of SO₂ to sulfate on dust with coexistence of NH₃ and NO₂ under different humidity conditions, *Atmos. Environ.*, 208, 133-140, <https://doi.org/10.1016/j.atmosenv.2019.04.004>, 2019a.
- 1185 Zhang, X. Y., Wang, Y. Q., Niu, T., Zhang, X. C., Gong, S. L., Zhang, Y. M., and Sun, J. Y.: Atmospheric aerosol compositions in China: spatial/temporal variability, chemical signature, regional haze distribution and comparisons with global aerosols, *Atmos. Chem. Phys.*, 12, 779-799, <https://doi.org/10.5194/acp-12-779-2012>, 2012.
- Zhang, X., Zhuang, G., Chen, J., Wang, Y., Wang, X., An, Z., and Zhang, P.: Heterogeneous reaction of sulfur dioxide on typical mineral particles, *J. Phys. Chem. B*, 110, 12588-12596, <https://doi.org/10.1021/jp0617773>, 2006.
- 1190 Zhang, Y., Bao, F., Li, M., Chen, C., and Zhao, J.: Nitrate-Enhanced Oxidation of SO₂ on Mineral Dust: A Vital Role of a Proton, *Environ. Sci. Technol.*, 53, 10139-10145, <https://doi.org/10.1021/acs.est.9b01921>, 2019b.
- Zhang, Y., Tong, S. R., and Ge, M. F.: A study about the influence of the size of CaCO₃ on the heterogeneous oxidation of sulfur dioxide by ozone, *Spectrosc. Spect. Anal.*, 36, 126-127, 2016.
- Zhang, Y., Tong, S., Ge, M., Jing, B., Hou, S., Tan, F., Chen, Y., Guo, Y., and Wu, L.: The influence of relative humidity on the heterogeneous oxidation of sulfur dioxide by ozone on calcium carbonate particles, *Sci. Total Environ.*, 633, 1253-1262, <https://doi.org/10.1016/j.scitotenv.2018.03.288>, 2018.
- 1195 Zhanzakova, A., Tong, S., Yang, K., Chen, L., Li, K., Fu, H., Wang, L., and Kong, L.: The effects of surfactants on the heterogeneous uptake of sulfur dioxide on hematite, *Atmos. Environ.*, 213, 548-557, <https://doi.org/10.1016/j.atmosenv.2019.06.050>, 2019.
- 1200 Zhao, D., Song, X., Zhu, T., Zhang, Z., and Liu, Y.: Multiphase oxidation of SO₂ by NO₂ on CaCO₃ particles, *Atmos. Chem. Phys.*, 18, 2481-2493, <https://doi.org/10.5194/acp-2017-610>, 2018.
- Zhao, X., Kong, L., Sun, Z., Ding, X., Cheng, T., Yang, X., and Chen, J.: Interactions between heterogeneous uptake and adsorption of sulfur dioxide and acetaldehyde on hematite, *J. Phys. Chem. A*, 119, 4001-4008, <https://doi.org/10.1021/acs.jpca.5b01359>, 2015.
- 1205 Zheng, B., Zhang, Q., Zhang, Y., He, K. B., Wang, K., Zheng, G. J., Duan, F. K., Ma, Y. L., and Kimoto, T.: Heterogeneous chemistry: a mechanism missing in current models to explain secondary inorganic aerosol formation during the January 2013 haze episode in North China, *Atmos. Chem. Phys.*, 15, 2031-2049, <https://doi.org/10.5194/acp-15-2031-2015>, 2015.
- Zheng, H., Song, S., Sarwar, G., Gen, M., Wang, S., Ding, D., Chang, X., Zhang, S., Xing, J., Sun, Y., Ji, D., Chan, C. K., Gao, J., and McElroy, M. B.: Contribution of Particulate Nitrate Photolysis to Heterogeneous Sulfate Formation for Winter Haze in China, *Environ. Sci. Technol. Lett.*, <https://doi.org/10.1021/acs.estlett.0c00368>, 2020.
- 1210 Zheng, S., Huang, Q., Zhou, J., and Wang, B.: A study on dye photoremoval in TiO₂ suspension solution, *Journal of photochemistry and photobiology. A, Chemistry.*, 108, 235-238, [https://doi.org/10.1016/S1010-6030\(97\)00014-2](https://doi.org/10.1016/S1010-6030(97)00014-2), 1997.
- Zhou, L., Wang, W., Gai, Y., and Ge, M.: Knudsen cell and smog chamber study of the heterogeneous uptake of sulfur dioxide on Chinese mineral dust, *J. Environ. Sci.*, 26, 2423-2433, <https://doi.org/10.1016/j.jes.2014.04.005>, 2014.
- 1215 Zhu, Y., Toon, O. B., Jensen, E. J., Bardeen, C. G., Mills, M. J., Tolbert, M. A., Yu, P., and Woods, S.: Persisting volcanic ash particles impact stratospheric SO₂ lifetime and aerosol optical properties, *Nat. Commun.*, 11, 4526, <https://doi.org/10.1038/s41467-020-18352-5>, 2020.