



Understanding greenhouse gas (GHG) column concentrations in Munich using WRF

Xinxu Zhao¹, Jia Chen¹, Julia Marshall², Michal Galkowski^{3,4}, Stephan Hachinger⁵, Florian Dietrich¹, Ankit Shekhar⁶, Johannes Gensheimer^{*1}, Adrian Wenzel¹, and Christoph Gerbig³

¹Electrical and Computer Engineering, Technische Universität München, 80333 Munich, Germany

²Deutsches Zentrum für Luft- und Raumfahrt (DLR), Institut für Physik der Atmosphäre, Oberpfaffenhofen, Germany

³Max Planck Institute for Biogeochemistry, 07745 Jena, Germany

⁴AGH University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland

⁵Leibniz Supercomputing Center (Leibniz-Rechenzentrum, LRZ) of Bavarian Academy of Sciences and Humanities, Boltzmannstr. 1, 85748 Garching, Germany

⁶Institute for Agricultural Sciences (IAS), ETH Zurich, Universitätstrasse 2, 8092 Zürich, Switzerland

Correspondence: Jia Chen (jia.chen@tum.de)

Abstract. To address ambitious goals of carbon neutrality set at national and city scales, a number of atmospheric networks have been deployed to monitor greenhouse gas (GHG) concentrations in and around cities. To convert these measurements into estimates of emissions from cities, atmospheric models are used to simulate the transport of various tracer gases and help interpret these measurements. We set up a modelling framework using the Weather Research and Forecasting (WRF) model applied at a high spatial resolution (up to 400 m) to simulate the atmospheric transport of GHGs and interpret the observations provided by the Munich Urban Carbon Column Network (MUCCnet). Building on previous analyses using similar measurements performed within a campaign for the city of Berlin and its surroundings (Zhao et al., 2019), our modelling framework has been improved regarding the initialization of tagged tracers, model settings, and input data. To assess the model performance, we validate the modelled output against two local weather stations and observed column GHG concentrations provided by MUCCnet from 1 to 30 August, 2018. The modelled wind matches well with the measurements from the weather stations, with wind speeds slightly overestimated. The measured slant column concentrations of GHGs and their variability are generally reproduced by the model, with a bias in CO₂ of around 3.7 ppm that can be attributed to the initial and boundary conditions used. The differential column method (DCM) has been applied to cancel out the influence from the background concentrations of CO₂. We optimize its application by selecting suitable days on which the assumption of the DCM holds true: a relatively uniform air mass travels over the city, passing from an upwind site to a downwind site. In particular, the Stochastic Time-Inverted Lagrangian Transport model (STILT) is used here and driven by our WRF modelled meteorological fields to obtain footprints (i.e., the potential areas of influence for signals observed at specific points), further used for interpreting measurement results. Combining these footprints with knowledge of local emission sources, we find evidence of CH₄ sources near Munich that are missing or underestimated in the emission inventory used. This demonstrates the potential of this data-model framework to constrain local sources and improve emission inventories.

^{*}Now at Max Planck Institute for Biogeochemistry, 07745 Jena, Germany



1 Introduction

Human activities have resulted in an increase of the global average temperature relative to pre-industrial levels of approx. 1.1 °C, a number which is expected to reach around 1.5 °C between 2030 and 2052 (Masson-Delmotte et al., 2018; Allen et al., 2018). To achieve the long-term goal of the Paris Agreement to limit temperature increase to 2 °C relative to pre-industrial levels, effective and efficient adaptation at national, regional and local levels is needed, leading to deep reductions in atmospheric emissions of greenhouse gases (GHGs) over the coming decades (Masson-Delmotte et al., 2021). More than half of the world's population resides in urban areas, which are directly responsible for over 30% of the global GHG emissions (Masson-Delmotte et al., 2018) and approx. 65% of global energy use (IRENA, 2016). Thus, cities play a vital role in addressing the challenge of carbon mitigation. The development of science-based methods to estimate carbon emissions from urban areas is crucial for developing effective and coherent adaptation actions, and monitoring their success.

As the continent with the highest population density, Europe plays a major role in future mitigation efforts. In recognition of this fact, the European Commission aims to make Europe climate neutral by 2050 (EU Commission, 2018). Furthermore, member countries of the European Union (EU) have also adopted individual strategies consistent with that goal. For example, the German government plans to reduce the national GHG emissions by more than 65% compared to 1990 by 2030, achieving climate neutrality by 2045. Local-scale initiatives have also been put in place. Munich, currently the third largest city in Germany with over 1.5 million inhabitants, has set an even more ambitious goal, aiming to be climate neutral by 2030.

To confront the challenge of carbon mitigation in cities and reach the goals set by individual municipalities, a multitude of urban atmospheric networks have been built worldwide to optimize urban emissions (DeCola et al., 2018). Using the measurements they provide, GHG concentrations can thus be monitored in and around cities, and more accurate emission estimates can be derived interpreting these measurements with atmospheric transport modelling and statistical techniques (Lauvaux et al., 2016; Staufer et al., 2016). Based on these quantitative assessments, more reliable scientific guidance can be provided to policymakers in order to plan local emission reductions effectively and monitor mitigation efforts.

Two standard approaches are widely used for estimating emission fluxes: the bottom-up approach and the top-down approach. Using the bottom-up approach, the total fluxes are estimated on the basis of statistical activity data from individual sectors (e.g., power plants, traffic) and the corresponding emission factors. This approach is widely used for generating global and national sector-by-sector emission inventories (e.g., the Emissions Database for Global Atmospheric Research, 'EDGAR' Janssens-Maenhout et al. (2019)). The spatially and temporally disaggregated emissions produced using this technique are often quite uncertain, owing to missing or simplified knowledge of emission processes, their considerable heterogeneity in space and time (Klausner et al., 2020), or both. By using the top-down approach, estimated emission fluxes can be refined using additional information provided by measurements of atmospheric concentrations. Prior emission estimates, usually estimated by the bottom-up approach, are used as inputs for an atmospheric transport model and the resultant concentrations are compared to the atmospheric composition measurements (Shekhar et al., 2020). The emission estimates can then be optimized using either a mass-balance approach (Heimbürger et al., 2017) or other inverse techniques (Jones et al., 2021). In terms of GHG emission estimation for cities or an area of interest, the top-down inversion approach has frequently been applied in



55 modelling studies accompanied by urban measurement networks, e.g., in California (Turner et al., 2016), Paris (Stauffer et al., 2016), Boston (Sargent et al., 2018), Berlin (Klausner et al., 2020) and Indianapolis (Jones et al., 2021). Inversion models still show considerable potential for improvement, owing to limited knowledge about the characteristics and spatial distributions of emission sources (e.g., missing stationary point sources and inner-city traffic), uncertainties in background concentrations, and the difficulty of modelling transport in complex urban environments.

60 To aid in reaching the goal of climate neutrality and track emissions in Munich, our group has established a novel automated urban sensor network (MUCCnet: Munich Urban Carbon Column Network, accessible via <http://atmosphere.ei.tum.de/>, Dietrich et al. (2021)) for continuous, long-term monitoring of GHGs in and around Munich. In brief, MUCCnet takes measurements simultaneously at five locations to capture concentration signals in and around the city, with the goal of estimating city emissions using continuous measurements accompanied by atmospheric models. MUCCnet is designed with the differential column methodology (DCM, Chen et al. (2016)) in mind, developed to quantify the emissions within a certain area, e.g.,
65 a single city. This is done by capturing concentration enhancements between downwind and upwind sites, so that a signal can be attributed to the emissions from the area in between. Several studies have used this approach combined with atmospheric models to study urban and local emissions, e.g., in Berlin (Hase et al., 2015; Zhao et al., 2019), Paris (Vogel et al., 2019), Munich (Toja-Silva et al., 2017) and Chino, California (Chen et al., 2016; Viatte et al., 2017). Detailed descriptions of MUCCnet
70 and its measurement principle are presented in Sect. 4.1 and Dietrich et al. (2021).

On the basis of MUCCnet measurements and our previous study that interpreted data from a measurement campaign around Berlin (Hase et al., 2015; Zhao et al., 2019), we have set up a modelling framework for Munich. This is based on the Weather Research and Forecasting model (WRF) enhanced with a biospheric flux module (WRF-GHG; Beck et al. (2012)), which runs
75 at a horizontal resolution of up to 400 m over the city. WRF is a mesoscale model commonly used for weather and atmospheric studies, and WRF-Chem has been extended with additional modules for tracer transport and chemistry (now including the GHG modules; Skamarock et al. (2008)). With WRF-Chem (Peckham et al., 2017), modelled meteorological fields are used to drive simulations of the atmospheric transport of trace gases, e.g., GHGs (Zhao et al., 2019) and air pollutants (Georgiou et al., 2018). Our model for Munich aims to reproduce observations from the five measurement locations in MUCCnet so as to aid in their interpretation and better understand the processes driving the emission and uptake of GHGs around the city. Apart
80 from generating concentration fields for different emission tracers, the output from the modelling system will be further used as input for other studies. For instance, highly-resolved meteorological fields can drive particle transport models (Fasoli et al., 2018) and the simulated biogenic signals have been used in a Bayesian inversion model (Jones et al., 2021) to infer emission values.

In this paper, we describe our model framework in detail and apply it in the interpretation of the observations collected by
85 MUCCnet from 1 to 30 August, 2018. Compared to the previous study in Berlin (Zhao et al., 2019), the model has been updated and several aspects related to model settings and initialization processes have been improved. As an example, more precise anthropogenic fluxes have been used for the tagged emission tracers. All these model-related aspects are described in Sect. 2. In Sect. 3, we assess the performance of our model by comparing its output to the measurements from two local weather stations. A model-measurement comparison of GHG column concentrations is presented in Sect. 4. In Sect. 5, we optimize the



90 application of DCM, and use it to assess the model performance and further track missing or underestimated emission sources around MUCCnet, through combining footprints generated by the particle transport model STILT (Fasoli et al., 2018) and the knowledge of local sources.

2 Modelling Framework Description

We use WRF-Chem Version 3.9.1.1 (Skamarock et al., 2008; Peckham et al., 2017) with an updated GHG module (Beck et al., 95 2012) in order to simulate the fluxes and transport processes of atmospheric GHGs in and around Munich at a horizontal resolution of up to 400 m. The main component of the setup is the WRF-ARW model, based on the fully compressible non-hydrostatic Euler equations (Skamarock et al., 2008). We take realistic meteorological driving data from the ERA5 reanalysis (Hersbach et al., 2020), extracted at approx. 31 km horizontal resolution and with 137 vertical levels from ground level to 0.01 hPa. These data provide the initial and boundary conditions for the meteorological fields.

100 In our model, biogenic fluxes of GHGs are simulated online, driven by meteorological parameters at native model resolution, in addition to other input. Furthermore, fluxes from external emission inventories are included as input and these surface fluxes are transported as passive tracers in WRF (Beck et al., 2012). Compared to the originally developed WRF-GHG, in which the GHG modules had to be explicitly integrated with WRF Version 3.2 (cf. Zhao et al., 2019), these modules have been added to the official WRF-Chem repository since WRF-Chem Version 3.4. It is worth noting that the GHG module does not take 105 into account atmospheric chemical reactions (as it treats GHGs as passive tracers). This is, however, not expected to produce significant biases, owing to the long lifetimes of GHGs compared to the relatively short residence time of tracers in the regional domain (Super et al., 2016; Dekker et al., 2017).

The model is configured in a three-domain nested configuration, with horizontal resolutions of 10 km for the outermost domain (D01), 2 km for the intermediate domain (D02) and 400 m for the innermost domain (D03), as illustrated in Fig. 1. The 110 spatial grids are assigned using the Lambert Conformal Conic (LCC) projection. The simulations are carried out with model integration time steps of 30 seconds, 6 seconds 1.2 seconds for each domain, with model outputs saved at time intervals of 3 hours, 1 hour, and 15 minutes, respectively. We define 46 vertical levels from the surface up to 50 hPa, 21 of which are in the lowest 1 km of the atmosphere. All five total column measurement sites from MUCCnet are located inside D03.

To better capture the urban landscape features and improve the urban model performance (Brousse et al., 2016; Ching et al., 115 2018; Mughal, 2020), we provide extra morphological urban parameters for the innermost domain (D03, area of Munich), which enables us to use the urban canopy multi-layer scheme in WRF (Brousse et al., 2016). This is done by re-classifying the land-cover categories for urban areas, while keeping the other land-cover categories unmodified. The re-categorized land-cover types are derived from the European Local Climate Zones (LCZ) map (Demuzere et al., 2019), extracted for our high resolution domain (Fig. 1). More information regarding this procedure can be found in the supplement.

120 The initial and lateral boundary conditions in the simulated background concentration fields of CO₂ and CH₄ are taken from the Integrated Forecasting System (IFS) Cycle 45r1, implemented by ECMWF, at a horizontal resolution of approx. 40 km (Rémy et al., 2019; Browne et al., 2019).




Figure 1. Topography map for the entire domain area (left panel). The right panel shows the land use classification in D03, including the 21 IGBP-modified MODIS land-cover types (from 1 to 21, as illustrated in the color bar and labels), and 10 LCZ land-cover categories defined for the urban areas of D03 (from 30 to 40, as illustrated in the color bar and labels). The five measurement sites in our MUCCnet campaign are marked as black dots on the right panel. The national boundaries and coastlines in the left panel from National Center for Atmospheric Research (NCAR) Graphics Version 4.1 are depicted (© UCAR/NCAR).

Biogenic CO₂ fluxes are implemented online utilizing the Vegetation Photosynthesis and Respiration Model (VPRM; Mahadevan et al. (2008); Chen et al. (2020)), a simple diagnostic light-use-efficiency (LUE) model coupled to WRF-Chem. VPRM does not reproduce the physiological processes of vegetation, but rather calculates Gross Primary Production (GPP) using the input of meteorological variables and vegetation indices derived from remote sensing. Ecosystem respiration (RES) is estimated using a simple linear model related to the air temperature and vegetation-specific parameters. Finally, the hourly CO₂ Net Ecosystem Exchange (NEE) is the difference between GPP and RES. In detail, the entire calculation is based on satellite-derived indices, short wave radiation and surface temperature at 2 m above the ground level as simulated by WRF



130 (Beck et al., 2012). The indices (i.e., Enhanced Vegetation Index (EVI) and Land Surface Water Index (LSWI)) here are derived from reflectance data measured by MODIS, specifically the product MOD09A1 Version 6. MODIS has resolutions of 0.5-1 km depending on the wavelength band and a temporal resolution of eight days (Vermote, 2015). The MODIS reflectance data are aggregated and interpolated onto the LCC projection, and the vegetation is classified following the SynerCover Product (SYNMAP) data with a resolution of 1 km (Jung et al., 2006).

135 Because the 1-km resolution of the SYNMAP dataset is unable to resolve vegetation within cities, urban areas are essentially masked out and VPRM does not produce any fluxes within cities (see the middle panel in Fig. S1 of the supplement). Thus, we extend and refine the vegetation classification using the Dynamic Land Cover map of the Copernicus Global Land Service at a resolution of 100 m (CGLS-LC100). This refined classification is used for our innermost domain to better capture the urban biogenic signals of CO₂. Details of this reclassification are described in Table S1 of the supplement. A comparative
140 visualisation of the vegetation classification maps with and without the refinement around Munich is given in Fig. S1 of the supplement.

CH₄ fluxes from wetlands are estimated using the Kaplan model (Kaplan et al., 2006), which is run online within WRF-Chem. This model calculates CH₄ emissions from anaerobic microbial production in wetlands as a fraction of heterotrophic respiration (Beck et al., 2013). The flux estimates depend on the modelled soil moisture, soil temperature and the carbon pool
145 from the Lund-Potsdam-Jena model, which is used for classifying the wetland fractions in the domain (Beck et al., 2012).

The first version of the GHG and co-emitted species emission database produced by the Netherlands Organisation for Applied Scientific Research (TNO; dataset TNO_GHGco_v1.1; Super et al. (2020)) is used to initialize anthropogenic fluxes as tagged tracers. This dataset provides annual gridded anthropogenic emissions for 2015 at a horizontal resolution of 0.05° x 0.1° (latitude x longitude, approx. 6 km x 6 km), covering most of Europe and part of North Africa. In addition to the
150 gridded emissions, point emission sources (e.g., power plants) are reported separately with geographical coordinates. These emissions are classified into fourteen sectors, following the Gridding Nomenclature for Reporting (GNFR) emission categories (A to L), i.e., industry, public power, road transport and other anthropogenic sources. Furthermore, a high-resolution version of TNO_GHGco_v1.1, at a resolution of 1/120° x 1/60° (latitude x longitude, approx. 1 km x 1 km) is available for central Europe (i.e., all of Germany and parts of France, Poland and the Netherlands, etc.). This version of TNO_GHGco_v1.1 was prepared
155 to support model studies at the local scale in the CO₂ Human Emissions project (CHE; van der Gon et al. (2019)). Since our outermost domain is not fully covered by the high-resolution version of TNO_GHGco_v1.1, we use the lower resolution (6 km) emissions for the outermost domain (D01), and the high-resolution version for the other two domains (D02 and D03 in Fig.1). To prepare the input for WRF-Chem, the required temporal disaggregation of the annual emissions was performed based on time-dependent scaling factors for monthly, weekly, and diurnal variations (Zhao et al., 2019; Super et al., 2020). In addition,
160 we release the emission fluxes from the point sources in TNO_GHGco_v1.1 from different heights above the ground, using the vertical profiles provided in Table 2 of Brunner et al. (2019). The re-allocation of point sources to vertical levels in our domain is illustrated in Fig. S2 of the supplement.



3 Model-measurement comparison for wind fields

To assess the performance of our model framework and evaluate the modelled meteorological variables used for transporting
165 the fluxes, some key meteorological parameters are compared to measured values provided by two local weather stations.

The first station is located at the Meteorological Institute of the Ludwig Maximilian University of Munich (LMU; latitude:
48.15°, longitude: 11.57°, altitude: 561 m), close to the center of Munich. This station can provide time series of meteorological
variables second by second. We compare the model to five meteorological variables measured at LMU: the temperatures at
heights of 2 m and 30 m above the ground (T2 & T30), the precipitation, and the wind speed and direction at 30 m above
170 the ground (WS30 & WD30). The other station we consider is operated by the German Meteorological Service (Deutscher
Wetterdienst in German, DWD). This automatic weather station is located at the Munich airport and has station ID 01262
(latitude: 48.35°, longitude: 11.81°, altitude: 446 m). We use the following variables measured at the airport for comparison
with our model: T2, precipitation, relative humidity, air pressure and the wind speed and direction at 10 m above the ground
(WS10 & WD10 respectively).

175 As one of the key drivers for the transport of trace gases in the model, the simulated wind field directly impacts the transport
patterns of the tracers. Thus, it is particularly important to assess the model performance with regards to the wind field. Here,
we employ the measured WS30 & WD30 at the LMU station, and WS10 & WD10 at the DWD station. The LMU station
measures the winds every second, while the wind data given by DWD are recorded as 10-minute means. We apply a cut-off
wind speed threshold (0.5 m/s in our case) to the values shown in Fig. 2, owing to large uncertainties in wind directions during
180 low wind speed periods (Zhao et al., 2019).

A comparison between the modelled and measured winds at the LMU station is shown in Fig. 2. Prevailing wind directions
both in the simulations and measurements are either easterly or westerly during the daytime, while the prevailing winds at night
are generally from the southwest. The measurements (panels (a.1), (a.3) and (a.5)) show larger scatter in the wind direction
over August compared to the simulations. Along the time series, the simulated (Fig. 2(b): red crosses) and measured (blue
185 dots) wind speeds show similar variability, but the model generally overestimates wind speeds with a root mean squared error
(RMSE) of 2.0 m/s and a mean absolute error (MAE) of 1.2 m/s. Regarding a comparison of the wind directions between the
model and measurements (see Fig. 2(c)), the model mostly follows the measured fluctuations of wind directions but with some
difference over time (RMSE=64.1 degrees & MAE=1.8 degrees). The model performance is reduced in some periods, e.g.,
between 24 and 26 August, when the variability of the wind direction is remarkably lower in the model.

190 In addition to this model-measurement comparison for WS30 & WD30, similar comparisons regarding the other meteorological
variables have also been performed and are presented in Sect. S5 of the supplement. These comparisons indicate that
our model has the capability to provide reasonable simulated meteorological fields for driving the transport of trace gases.

4 Model-measurement comparison for concentration fields and model tracer analysis

This part of our study is dedicated to a comparison of the measurements from MUCCnet to column concentrations for different
195 tagged tracers extracted from the simulation output fields. Section 4.1 contains relevant information on the measurements used




Figure 2. Wind roses (a) and time series of simulated and observed wind speeds (b) and wind directions (c) at a height of 30 m above the ground at LMU. Wind roses for the measurements over August 2018 are plotted in Panels (a.1): 24 hours, (a.3): daytime from 6 am to 5 pm only and (a.5): nighttime only, while Panels (a.2), (a.4) and (a.6) represent the modelled values. Each wind rose indicates WS30, WD30 and the frequency (% scale) of wind coming from a particular direction during the targeted period. The blue dots in Panels (b) & (c) represent the measured values from the LMU station and the red crosses represent the simulation. The grey shaded areas mark the measurement periods used for the model-measurement comparison of column concentrations in Sect. 4.

for interpreting the model, and the model-measurement comparison is discussed in Sect. 4.2. The final section (Sect. 4.3) characterizes changes in concentrations caused by individual anthropogenic emission processes, as represented by our model.

4.1 Description of MUCCnet and the measurement campaign

Our WRF model framework for Munich is designed to study GHG concentrations in connection with the Munich Urban
 200 Carbon Column network (MUCCnet; Dietrich et al. (2021)). In MUCCnet, five compact Fourier-transform infrared (FTIR) Spectrometers (EM27/SUN by Bruker Optics) have been deployed, four of which were located around Munich at a radius of 20 km for the 2018 period analysed in this study (this was later changed to 10 km in 2019). The fifth instrument has been set up close to the center of Munich, at the TUM campus (see Fig. 1). By using the sun as a light source, the EM27/SUN measures near-infrared solar spectra (Hedelius et al., 2016). In MUCCnet, the recorded interferograms are automatically transformed



205 to spectra, converted to abundances of CO₂ and CH₄ in a column between the instrument and the end of the atmosphere in the direction towards the sun, and further uploaded to the official website of MUCCnet. All five instruments are automatically operated and controlled using our universal enclosure systems and two software programs (Heinle and Chen, 2018; Dietrich et al., 2021). Detailed information on the EM27/SUN instrument can be found, e.g., in Gisi et al. (2012), Hedelius et al. (2016), Chen et al. (2016), Hase et al. (2016) and Frey et al. (2019).

210 In this study, we compare simulations to measurements collected during a campaign that was carried out from 1 to 30 August, 2018. Table S4 in the supplement shows relevant parameters for assessing the measurement performance during that period, including the number of observations per day for each site and the ground-level wind information for each day, i.e., the daily mean of WS30 & the approximate change in WD30 during the day, provided by the LMU station.

The EM27/SUN measures column-averaged dry air mole fractions of CO₂ and CH₄, hereafter referred to as XCO₂ and
215 XCH₄. Over the entire campaign period in 2018, the mean of the measured XCO₂ for all five sites is 404.4 ppm with a standard deviation of 1.2 ppm, ranging from 400.8 ppm to 408.1 ppm. For XCH₄, the measurements range from 1840.5 ppb to 1896.0 ppb, with a mean of 1865.5 ppb and a standard deviation of 9.1 ppb. Since the operation of the instruments is strongly influenced by weather conditions, such as sunlight (Vogel et al., 2019), the spatial and temporal measurement coverage for some days (e.g., 1-3 August) is limited (see Table S4). By assessing the measurement performance of the campaign, we in total
220 selected fifteen days with good measurement conditions (with a quality level better than '++', cf. Table S4 and Sect. S6 of the supplement) to make the model-measurement comparison: 4-6, 9, 11, 16-22 and 27-29 August, 2018. Details of the campaign and side-by-side calibrations are discussed in Sect. 4.1 and 5 of Dietrich et al. (2021).

4.2 Model-measurement comparison of XCO₂ and XCH₄

In our modelling framework, the anthropogenic emission fluxes from an external inventory (namely TNO-GHGco, see Sect. 2
225 for details) are used as fluxes for tagged tracers, each representing a source category group available in the inventory (Super et al., 2020). These tracers are transported passively throughout the model domains using internal WRF-Chem transport schemes. Then, the total concentrations for a trace gas are derived by summing up the contributions from individual emission processes (i.e., the different tagged tracers) and the background concentrations for this gas, provided by the Copernicus Monitoring Service (CAM5) and advected from the model boundaries as a separate tracer.

230 4.2.1 Calculation of smoothed slant column concentrations

The modelled vertical concentration profiles are converted to pressure-weighted column-averaged concentrations. That is to say, for the trace gas G , the simulated column-concentration at a specific location and time ($XG(x, y, t)$) can be calculated as follows:

$$XG(x, y, t) = \sum_{l=1}^{L_{\text{ver}}} [w_l(x, y, t) \times G_l(x, y, t)] \quad (1)$$



235 where G_l stands for the simulated mole fraction at the location (x, y) and time t at the l^{th} vertical layer of WRF. L_{ver} is the total number of the vertical layers (i.e., 45 in our study) and w_l is the weight of the l^{th} vertical layer which can be obtained as:

$$w_l(x, y, t) = \frac{\Delta P_l(x, y, t)}{P_{sf}(x, y, t) - P_{tp}} \quad (2)$$

where P_{tp} is the hydrostatic pressure at the top of the model (i.e., 50 hPa) and P_{sf} is the surface pressure. ΔP_l denotes the pressure difference between the top and the bottom of the l^{th} vertical layer.

240 However, when comparing our modeled values to the measurements here, we need to consider the characteristics of the instruments in a more accurate way. The EM27/SUN records the spectra along a slant column from the sun to the ground, instead of a vertical column perpendicular to the ground. Simulated concentration fields of CO_2 and CH_4 used for the model-measurement comparison in this study must therefore be aggregated along the slant columns from the ground to the sun.

In addition, when reconstructing the vertical structure of the atmosphere during the retrieval process of ground-based remote
245 sensing instruments (e.g., Vogel et al. (2019); Zhao et al. (2019)) and satellites (e.g., Ohyama et al. (2020)), an averaging kernel (AK) is used to represent the altitude-dependent column sensitivity (Borsdorff et al., 2014). For solar-viewing instruments, its shape is strongly dependent on the solar zenith angle (SZA). The retrieved quantity then typically depends considerably on the AK as well as on the a-priori profile used in the retrieval. Accordingly, when the modelled values are compared to such measurements, they also need to be smoothed using the AK and the a-priori profile in order for the comparison to be valid.
250 Since the EM27/SUN has a spectral resolution of only 0.5 cm^{-1} , we can use a fitted AK matrix, which is obtained by applying a simple least-squares fit to a given a-priori AK profile. The details related to applying the AK to the model data were previously described in detail in Sect. 3.3 of Zhao et al. (2019).

Thus, the simulations are mostly interpreted by comparing the aggregated and AK-smoothed modelled values to column measurements. The smoothed slant column concentration for a target gas XG_{sla}^S is calculated following Eq. 3 (cf. Vogel et al.
255 (2019); Zhao et al. (2019)),

$$XG_{sla}^S(x, y, t) = \sum_{l=1}^{L_{ver}} w_{sla,l}(x, y, t) \times [AK_{G,l}(t) \times G_{sla,l}(x, y, t) + (1 - AK_{G,l}(t)) \times G_{pri,l}] \quad (3)$$

where $G_{sla,l}$ are modelled concentrations for trace gas G at the l^{th} vertical layer following the slant column along the line of the sun, $AK_{G,l}$ is the fitted AK of the gas G at the l^{th} vertical layer and time t , $G_{pri,l}$ stands for the mixing ratio of the a-priori profile for trace gas G at the l^{th} vertical layer, and $w_{sla,l}$ is the weight of the l^{th} vertical layer along the slant column. We have
260 used the a-priori profile from the Whole Atmosphere Community Climate Model (WACCM) Version 6 for 2018.

It should also be noted here that the measured samples are filtered during the autonomous retrieval process in MUCCnet (Dietrich et al., 2021). Specifically, to reduce uncertainties caused by high air masses, measurements are discarded when they are observed at SZA larger than 75 degrees (Tu et al., 2020). Thus, the measurement period each day ranges from around 6:00 UTC to 17:00 UTC and lasts for approx. 11 hours in summer. In the following figures, the grey shaded areas are used to mark
265 these measurement periods used for the model-measurement comparisons.




Figure 3. Time series and scatter plots for XCO_2 (a & b) and XCH_4 (c & d). In panels (a & c), the dashed lines represent the daily mean modelled $XCO_{2,sla}^S$ and $XCH_{4,sla}^S$, while the solid lines denote the measurements. Colors in panels (a & c) mark the different measurement sites. In scatter plots (panels (b & d)), colors represent the values for different measurement days, as marked in the color bar. The error bar represents the standard deviation of the measured and simulated values at each site.

4.2.2 Comparison of daily mean concentrations and estimation of CO_2 mean bias

The daily mean measured and modelled values ($XCO_{2,sla}^S$ and $XCH_{4,sla}^S$) for the 15 studied days and their scatter plots are shown in Fig. 3. When producing time-averaged modelled values, we have considered the limited measurement period, as mentioned above. For CO_2 , the simulated smoothed column concentrations ($XCO_{2,sla}^S$, dotted lines in Fig. 3(a)) are overall
 270 overestimated compared to the measurements, with a mean bias (MB) of $3.7 \text{ ppm} \pm 0.9 \text{ ppm}$, the latter value giving the standard deviation of the MB over all measurement days. This bias for $XCO_{2,sla}^S$ is mainly attributed to the initial and boundary conditions of the concentration fields in the model as provided by CAMS, which has also been seen in other studies. Gałkowski et al. (2021) found a similar bias between the CAMS product and airborne measurements in the free troposphere over Europe, with a MB of $3.7 \pm 1.5 \text{ ppm}$. Moreover, Tu et al. (2020) also reported a bias when comparing CAMS to their column
 275 measurements from the Collaborative Carbon Column Observing Network (COCCON) site at Kiruna, Sweden, with a MB of $3.7 \pm 1.8 \text{ ppm}$.



As can be seen in the scatter plot of $XCO_{2,sla}^S$ (Fig. 3(b)), the measurements generally exhibit more scatter (seen in the magnitude of the error bars in the x-direction) compared to the simulation (the error bars in the y-direction), and the slope of the linear regression is only 0.45. The smaller standard deviations in the model represent weaker fluctuations over the daily mean. This will be discussed further in Sect. 4.2.3 when looking into the model-measurement comparison at higher temporal resolutions.

Figure. 3(c & d) shows that the daily mean modelled values of $XCH_{4,sla}^S$ (solid lines in Fig. 3(c)) agree well with the daily mean measurements. The model is able to capture most of the variations in the daily mean values, while in general the observed values are slightly higher, with a linear regression slope of 0.73 and a negative MB (-1.8 ± 4.0 ppb). This small bias could be caused by the initial and lateral boundary conditions from CAMS, or due to unknown or underestimated emissions. Comparing CH_4 in the CAMS product with in-situ observations in the troposphere, Gałkowski et al. (2021) also reported a negligible MB, but a relatively large standard deviation (0 ± 14 ppb) in their setup.

4.2.3 Comparison between model and measurements: intra-day concentrations

In order to obtain a more detailed view on how the model behaves at higher temporal resolution, the daily curves of $XCO_{2,sla}^S$ and $XCH_{4,sla}^S$ at five sites for 7 consecutive days from 16 to 22 August are plotted in Figs. 4 & 5 against the corresponding modeled values with hourly temporal resolution. Model-measurement comparisons for the rest of the selected days are shown in Figs. S5 & S6 of the supplement.

As mentioned in Sect. 4.2.1, the modelled and observed slant column concentrations used for the model-measurement comparisons are smoothed using the SZA-dependent AK based on Eq. 3. Due to the restriction of SZAs and the corresponding availability of measured values provided by the MUCCnet (cf. Sect. 4.1), these model-measurement comparisons of the total column concentrations for GHGs can only be made during the daytime, approx. from 6:00 UTC to 17:00 UTC (cf. Fig. 4). However, Figures 4 & 5 also illustrate the contributions to the total column concentrations of CO_2 & CH_4 from different tracers in the model at nighttime, using pressure-weighted column concentrations (Eq. 1) as a proxy. This can be used to interpret the model and the measurements, e.g., the contribution of nighttime vegetation respiration to the changes in total column concentrations of CO_2 .

Figure 4 shows the modelled CO_2 column concentrations from all tracers. In general, there is little difference in the column background concentrations among the five sites (black lines). Variations of the modelled total CO_2 corrected by the MB (solid red lines) are mostly dominated by biogenic activity (dashed blue lines), with only a minor influence predicted from anthropogenic emissions (dashed yellow lines).

After smoothing, the modelled, bias-corrected $XCO_{2,sla}^S$ (green '+') is slightly higher than the corresponding XCO_2 modelled values (solid red lines), especially during the morning and afternoon, with a RMSE of 0.37 ppm and a MB with its standard deviation of 0.34 ± 0.13 ppm. This is caused by the steeper shape of the vertical profile of the AK under larger SZAs. During the daytime from around 6:00 UTC to 17:00 UTC, there is no obvious difference between the modelled values with and without smoothing.




Figure 4. Modelled $XCO_{2,sla}^S$, attribution of variations to different tracers, and the measurements at the five MUCNet sites from 16 to 22 August, 2018. The purple circles represent the column measurements from MUCNet and the green '+' stands for the modelled $XCO_{2,sla}^S$ after subtracting the MB. The other lines in the plot give pressure-weighted modelled column concentrations along the full time series. These are calculated following Eq. 1 (i.e., without smoothing using the SZA-dependent AK) and are corrected by MB. The black curve represents the modelled background (XCO_{2_BCK}), and the red curve shows the modelled XCO_2 . The dashed yellow and blue curves highlight the concentration changes caused by human activities (XCO_{2_ANT}) and biogenic activities (XCO_{2_BIO}). The grey shaded areas mark the measurement periods used for comparing observations to model results.

310 The modelled $XCO_{2,sla}^S$ (green '+') reproduces the variability in the measurements (purple 'o') reasonably well (RMSE: 1.33 ppm; MB \pm std: -0.79 ± 0.14 ppm), as it turns out. However, the measurements often show a steep decrease in concentration during the morning, while the model only shows slight declines. This difference could be the result of an underestimation of the modelled biogenic respiration (RES) from VPRM. During the growing season (June-September), VPRM (Mahadevan et al., 2008) has been found to underestimate RES, especially at nighttime and overestimate GPP during the daytime (Hu et al., 2021; 315 Gourdji et al., 2021). This causes an overestimation of the magnitude of NEE (i.e., the difference between RES and GPP), which could explain the difference between the modelled column concentrations and what is observed by MUCNet. In our case, the observations suggested higher RES fluxes at nighttime than what was simulated by the model. This led to much lower modelled column concentrations in the early morning, which was also seen in Hu et al. (2021).



Compared to the modelled values for the other six dates, a slight rise was seen (approx. 1.5 ppm on average) in the morning of 21 August. This was induced by a combined effect of elevated background and biogenic tracer mole fractions. Closer analysis of this case (see the animation of XCO_2_BCK from 19 to 21 August in the supplement) has shown that the background enhancement entered from the northwest of the outermost boundary in the morning of 19 August and was transported into the innermost domain by the late afternoon of 20 August. This enhanced background signal contributes around two thirds of the modelled 1.5-ppm rise. The rest is the result of air masses with a strong biospheric CO_2 signature coming from the northwest of our outermost domain. Animations of XCO_2_BIO and XCO_2_BCK from 19 to 21 August are attached in the supplement to further illustrate this as well.




Figure 5. Modelled $XCH_4^S_{4,sla}$, attribution of variations to different tracers and the measurements at five sites of the MUCCnet from 16 to 22 August, 2018. The purple 'o' represents the column measurements from MUCCnet and the green '+' stands for the modelled $XCH_4^S_{4,sla}$. The other curves in the plot show the modelled column concentrations calculated following Eq. 1, i.e., without smoothing using the SZA-dependent AK: black curve – modelled background (XCH_4_BCK); red curve – modelled XCH_4 . The dashed yellow and blue curves highlight the concentration changes caused by human activities (XCH_4_ANT) and induced by biogenic activities (XCH_4_BIO), including the background. The grey shaded areas mark the measurement periods used for comparing to the simulations.

On 22 August, compared to the simulations for the other days, the modelled $XCO_2^S_{2,sla}$ shows a larger deficit with respect to the measurement in Fig. 4. The model can capture the variation during the day quite well, but produces too low XCO_2 values when bias corrected through subtraction of the MB. This more extreme mismatch is probably due to the advection of



330 air masses heavily impacted by biogenic activities (and thus with less CO₂) in the model, coming into the domain from e.g., Italy, Slovenia and Croatia. As can be seen in the map of the modelled NEE from VPRM in Fig. S7 of the supplement, much stronger biogenic fluxes are found in the south of the outermost domain, compared to the other areas. More CO₂ was taken up and the affected air masses (i.e., with comparatively less CO₂) are able to reach Munich when the wind is strong enough to drive them past the Alps. A constant MB over all days, as was applied here, may not always be realistic, as we see here. An
335 evaluation of the signals within MUCNet using the DCM is still possible, however, as will be discussed in Sect. 5.

In terms of CH₄, we conduct the same visual analysis (cf. Fig. 5). The variations of the modelled total XCH₄ (solid red) are dominated by anthropogenic activities (XCH₄_ANT, dashed yellow) and these two variations mostly overlap, since no significant signal induced by wetland emissions (XCH₄_BIO, dashed blue) is predicted by the model. The time series of the modelled XCH_{4,sla}^S (green '+') shows general agreement with the measurements (purple 'o'). The measurements seem to
340 capture stronger emission signals (e.g., on 22 August), perhaps due to gaps in our knowledge of the spatio-temporal distribution of CH₄ emissions. The modelled values show little diurnal variability at all sites compared to the measurements (RMSE: 6.7 ppb and MB ± std: -3.3 ± 5.9 ppb). Regarding the strong enhancements observed by the instruments during the daytime, especially on 22 August, these might be the result of sources which are missing from inventories, or are underestimated in their magnitude. In Sect. 5, we attempt to detect such unknown or underestimated emission sources over the domain using DCM.

345 4.3 Tracer analysis related to human activities

Beyond the major contributors to the concentration enhancements above the background as discussed above, we also analyse the contributions from individual anthropogenic emission processes to understand how these processes impact concentrations quantitatively. To be specific, we use the GNFR emission categories from TNO-GHGco for separately advected tagged tracers. For CO₂, the categories are: "A. Power Plants", "B. Industry", "C. Other Stationary Combustion", "D. Road Transport" and "E.
350 Other". For CH₄, the emission processes are: "A. Power Plants", "B. Industry", "C. Agriculture", "D. Waste Management", "E. Fugitives and solvents", and "E. Other".

The changes in concentrations induced by different human activities along the full time series are plotted in Fig. 6. For CO₂, the emissions from road transport (red) contribute the largest portion (around 37.4 %) of the total concentration enhancements caused by anthropogenic activities in August, 2018. This is consistent with the finding that over 30 % of the total GHG
355 emissions are induced by on-road transportation for around one-third of 167 cities worldwide (Wei et al., 2021). This finding may also be due to the seasonal features of emissions in Munich, as there is no heating and less electricity generation in summer. For other contributors, power plants account for around 7 %, other stationary combustion for 13.1 %, and both industry and other contributors for approx. 21 %, respectively. Figure 6(b) shows that for CH₄ the emissions are dominated by agriculture (brown) and waste management (purple), which are estimated to contribute approx. 50 % and 25 % of the total human-related
360 concentration enhancements.




Figure 6. Average modelled concentration contributions by individual anthropogenic emission processes of (a) CO_2 (XCO_2_ANT) and (b) CH_4 (XCH_4_ANT) from 1 to 30 August, 2018 in and around Munich (D03). The grey shaded areas mark the measurement periods used for comparing to the simulations.

5 DCM-based evaluation of models, detection and tracing of additional emission sources

In order to analyse the differences between the measurements and the model, described in Sect. 4.2, we apply DCM. This method allows for a cleaner interpretation, eliminating biases such as the offset discussed for CO_2 in Sect. 4.2.2. Combining this approach with particle transport modelling using STILT (Fasoli et al., 2018), we explore how we can trace unexpectedly high measured CH_4 signals (cf. Sect. 4.2.3, Fig. 5) to potential additional sources that are not included in the emission inventory.

In general, DCM is an approach used to assess the emissions over a certain area through the concentration differences (gradients) between down- and upwind GHG measurement sites (Chen et al., 2016; Dietrich et al., 2021). In Zhao et al. (2019), DCM was shown to be a useful post-processing approach in model analysis and model-measurement comparisons, due to its ability to cancel out biases related to initial and boundary conditions. The aim here is to compare our measurements to the model using DCM for selected days from 16 to 22 August, 2018, before we attempt to track CH_4 emission sources based on this analysis with the help of STILT.

To begin with, we select up- and downwind sites for our analysis based on the wind data presented in Fig. 2 (Sect. 5.1). We have significantly improved the application of DCM compared to our previous work: we make use of the transport model STILT



(Sect. 5.2) at this point, together with wind information along the whole column, to select dates for which our methodology is applicable (Sect. 5.3). This accounts for differences in wind shear. After comparing the modelled concentration gradients to the observations (Sect. 5.4), we further track the potential location(s) of unknown or underestimated CH₄ sources by using STILT footprint contours. These footprints mark the area of upstream fluxes that influence air masses which arrive at the measurement sites (Sect. 5.5).

5.1 Selection of up- and downwind sites

In DCM, we start by dividing our measurement sites into down- and upwind sites day by day, based on measured wind directions.

As seen in Fig. 3, the measured concentration values at the center site of MUCCnet (TUM) are found to always be higher than the values observed at the other sites, owing to the dense distribution of emission sources close to the city center, combined with the higher sensitivity to fluxes in the near-field of the observation location. To better understand the concentration gradients between the down- and upwind sites, we choose to exclude the center site.

The four remaining sites are grouped according to the wind directions observed at LMU. As shown in the wind rose of the measurements (see Fig. 2(a.3)), the prevailing wind directions during the daytime for our simulation period (WD30) are usually northeasterly or northwesterly. Table 1 shows the list of the down- and upwind sites for different prevailing wind conditions, which are used to calculate the concentration gradients.

Doing this, we assume that the surface winds measured in the city center are representative of the regional wind conditions over our domain during the day, and that they are sufficiently stationary for application of DCM. When rapid, regional-scale horizontal and vertical wind shifts occur, as during a summer cold front passage around the Alps, these assumptions might fail.

Therefore, we set up (Sect. 5.2) and use (Sect. 5.3) the STILT model (Fasoli et al., 2018) to assess transport patterns. For our discussion of CO₂ and CH₄ measurements vs. simulations, and most importantly for our attempt to locate unknown CH₄ emissions (Sect. 5.5), we have thus been able to select days where the assumptions for DCM are likely to be met. In particular, our method ensures the validity of DCM for the interpretation of the CH₄ concentration peaks.

Table 1. Table of up- and downwind sites depending on wind directions

Wind Direction	Upwind Sites	Downwind Sites
Northeasterly/Easterly (NE/N)	Markt Schwaben (East) Garching (North)	Weßling (West)
Northwesterly/Westerly (NW/W)	Weßling (West)	Garching (North) Markt Schwaben (East) Höhenkirchen (South)



5.2 STILT Model Setup

Before applying DCM to any of the days from 16 to 22 August, we assessed its applicability for each day (see Sect. 5.3) by tracking the origin of air masses at different measurement sites with the transport model STILT (cf. Fasoli et al. (2018)), and
400 by assessing the modelled vertical wind profiles used for generating footprints (see Sect. 5.3.2). Here, we briefly describe the STILT setup.

In our study, STILT with R code base (Version 2, as available via [https://uataq.github.io/stilt/index.html/](https://uataq.github.io/stilt/index.html#/), last access: 11, Jan 2022) was implemented using around 168 core hours provided by the high performance computer center LRZ. To assure transport consistency with previously presented results, STILT was driven by the WRF meteorological fields generated for our
405 second domain (D02) at a horizontal resolution of 2 km. In order to trace back the origin of air masses at a given spatio-temporal receptor point (corresponding to the time at which an instrument performs a given measurement), STILT uses ensembles of tracer particles which are propagated backwards in time. Specifically, the model provides us with the sensitivity of the analysed slant columns to surface fluxes. These sensitivities are calculated by considering the residence time of released particles when they traverse the lower planetary boundary layer before reaching the measurement location, and are further aggregated over the
410 STILT simulation time to produce footprint maps. In our configuration of STILT, we have released 500 particles at 13 altitudes along the slant column, namely at 20, 180, 350, 520, 700, 880, 1060, 1250, 1440, 1620, 1920, 2020 and 2220 m above the ground level for each simulation.

STILT then yields so-called footprint contours (i.e., contours enclosing a certain percentage, e.g., 90 %, of the accumulated surface sensitivity) for each altitude layer. Details of the percentile footprint contours are described in Sect. S11 of the
415 supplement. In order to yield an estimated effective footprint independent of emission height (depending on source and local conditions), we have aggregated these altitude-dependent footprint data for the different layers using the pressure differences between layers as weights (Jones et al., 2021).

5.3 Preparation of DCM: Date selection with footprints from STILT and vertical wind data

Leveraging the footprint contours from STILT, the differences in the origin and path of air masses arriving at up- and downwind
420 sites can be determined. Understanding these differences is a key prerequisite for determining the location of potential unknown GHG sources based on noteworthy signals in the downwind-upwind concentration gradients. Whenever air masses reaching up- and downwind sites have very different areas of influence, the upwind site cannot be used as a relative background site when calculating concentration gradients. When the footprints fully overlap, however, such that air passes over the upwind to the downwind site, and additional GHG contributions are from in between, DCM can be used. But even when this condition
425 is not strictly met, if differences in the footprint areas are small, the small non-overlapping parts are potential locations for unknown GHG emitters and sinks to be pinned down, rendering DCM effective. Signals coming from the overlapping area of the footprints, in contrast, will be visible at multiple measurement sites with a characteristic time delay. Clearly, for all this to hold, we need to check whether our footprints are realistic. The main prerequisite for this is the accuracy of the WRF wind fields, driving STILT, at different heights. We check this criterion at the end of this section using radiosonde data.



430 5.3.1 Assessment of air mass origin (footprints) at different sites on different days

The approach outlined above can be applied to understand up-/downwind differences and obtain information about GHG sources and transport in the target area. Here, we adopt the following strategy: for each measurement site, we compute the footprint contours with the receptor time in STILT set to the time of the daily XCH₄ peak value². Then, we accept only days for our study where the overlap of these footprints is large. The peak times of the stations are usually different by only a few
435 hours.

Our strategy results in three outcomes: (*i*) days with unstable wind conditions in time or with large variations of wind directions from one site to another are excluded (unless the two effects compensate quite exactly, which is unlikely); (*ii*) the large overlap and small differences in footprint contours allow for a clear localisation of potential origins of differences, as discussed above; and (*iii*) we can understand whether the peak is of the same origin at all sites. We chose to analyse
440 CH₄ signals at the end of this study (Sect. 5.5), as this offers a realistic possibility of tracking human emission sources (cf. Sect. 4.2.3) in this exploratory work. In contrast, the current state of the art makes it more difficult to trace anthropogenic effects in CO₂ signals, where biogenic activity plays a much larger role.

We applied our strategy to all measurement days, and finally selected 16, 20 and 21 August 2018 as days suitable for further analysis (Sects. 5.4 and 5.5). Figure 7 and the supplement (in particular Fig. S11) lay out the reasons for our decision and
445 show all the footprint contours. In the following paragraphs, we discuss the examples from Fig. 7 (16 and 22 August) further as typical days deemed appropriate (16 August) or inappropriate (22 August) for further analysis using DCM. Besides the footprints, Figure 7 also shows the peak times used as receptor times for STILT.

On August 16, with easterly wind prevailing, the instruments deployed upwind (Garching/Markt Schwaben) captured peak CH₄ signals in the first half of the day and a similar signal was then seen at TUM about an hour later (blue dots in Fig. 7(a)).
450 However, the sensor at the downwind site (Weßling) did not detect a major peak. Using the knowledge of station locations and the observed peak times, we would have predicted a peak at Weßling (Fig. 7(a), black dot) as follows: the upwind and central sites captured the peaks at around 11:00 and 12:00 UTC, respectively. These air masses would then possibly have reached the downwind site after approximately two hours (estimated from wind speeds as given in the supplement, Table S4), corresponding to the distance between TUM and Weßling, i.e., at 14:00 UTC (Fig. 7(a), black dot). Note that the receptor times
455 for all up- and downwind sites and dates, determined from observed peaks or estimates in this manner, are listed in Table S5.

On 22 August, the origins of air masses at most of our sites differ significantly (Fig. 7(b)). The footprints (with receptor times corresponding to the blue dots in Fig. 7(b), top panel) do not overlap as cleanly, and any 'tracing experiment' would be poorly controlled. The peaks seen at the Markt Schwaben, TUM, and Garching sites may well be of different origin, given the different footprint shapes. Thus, we do not study this day in further detail.




Figure 7. Observed XCH₄ over time (upper panel) and the 90th percentile contours of column footprints (lower panel) on (a/left) 16 and (b/right) 22 August at up- and downwind sites with different colors: red for Weßling (West), green for Markt Schwaben (East), purple for Garching (North) and pink for TUM (Center). The peaks in the observations are marked by dots (blue: clear peaks, black: inferred peaks, see main content/footnote). Two black squares in (b) mark the measurement sites of IGRA (cf. Sect. 5.3.2). The background maps use tiles from Stamen Design (<https://maps.stamen.com/>, under CC BY 3.0, with data by OpenStreetMap, under ODbL, from 2021 Dec. 21). The map colouring reflects the emissions from the initial inventory (i.e., TNO-MACCco).

460 5.3.2 Assessment of vertical wind profiles for calculating realistic footprints

In order to assess whether our WRF meteorological fields allow STILT to produce realistic footprints, we have further evaluated our modelled height-dependent wind fields using observations. This assesses their accuracy in the vertical dimension more deeply than the wind-field comparison shown in Sect. 3.

465 Modelled and measured wind speed and direction profiles are shown in Fig. 8 for 16 and 22 August. The comparison uses radiosonde data from two sites (Altenstadt and Hohenpeißenberg, marked as black squares in Fig. 7(b)), provided by Integrated Global Radiosonde Archive (IGRA). This archive collects radiosonde and pilot balloon observations along significant vertical levels historically and in near-real-time from around 800 distributed stations worldwide (Durre et al., 2018). On both days at Altenstadt (Fig. 8(a, b)), the model reproduces the observed winds well, especially at noon, while outliers in wind direction

²In a few cases where the peak times have been hardly detectable, they have been inferred using the peak times at nearby sites and considering time delays derived from the daily surface wind speeds from LMU.




Figure 8. Wind profile comparison between the model (orange dashed line) and the measurements (blue solid line) from two radiosonde locations, i.e., (a) & (b) Altenstadt on 16 and 22 August, and (c) Hohenpeißenberg on 22 August.

always exist at the lowest level in the morning and some mismatches appear at 18:00 UTC. Regarding the wind profiles
 470 measured at Hohenpeißenberg, the modelled wind direction at higher altitudes agree quite well with the observations, with mismatches close to the ground level. It is worth noting that our domain is close to the Alps, with complex topography making meteorological modelling more challenging. The results need to be interpreted in this context.

5.4 Model-measurement comparison of concentration gradients

After checking the prerequisites in the previous sections, we focus our analysis of the differential concentrations (gradients)
 475 for the selected dates of 16, 20 and 21 August.

Figure 9 shows $\Delta XCO_{2,sla}^S$ for the days selected based on the prevailing wind directions. The modelled concentration
 480 gradients of $XCO_{2,sla}^S$ between the down- and the upwind sites (blue solid lines in Fig. 9) are driven by both biogenic activities (light green) and human activities (grey). The biogenic part can be attributed to the special spatial distribution of biogenic sinks in Munich and its surroundings (see also Fig. S8 of the supplement): The southeastern and southwestern parts around Munich are more biologically active and have greater carbon sinks, compared to the other areas. This is an interesting difference to Berlin, where our previous study (Zhao et al., 2019) showed no such signal, corresponding to a relatively even distribution of biogenic fluxes over Berlin. In the model-measurement comparison of $\Delta XCO_{2,sla}^S$, the model (blue) was able to reproduce




Figure 9. Time series of $\Delta XCO_{2,sla}^S$ for three targeted days and their scatter plot: 16 and 21 August with NE/E winds in the upper two rows, and 20 August with W/NW winds in the bottom row. The column concentration differences between the down- and the upwind sites are plotted as red solid lines for measurements, blue solid lines for the modelled full signal values, light green dashed lines for the simulated biogenic signal values and grey for the modelled contributions related to anthropogenic activities.

the general variations when comparing to the measurements (red) shown in Fig. 9, with a Pearson correlation coefficient of 0.74 and RMSE of 0.37 ppm. However, the modelled and measured concentration gradients show some differences, e.g.,
 485 the modelled concentration gradients before 10:00 UTC were underestimated compared to the observations. The differences between the modelling results and the measurements could be potentially caused by underestimated concentration gradients from biogenic fluxes in and around the city (light green) in the model. In particular, underestimating RES during nighttime (details in Sect. 4.2.3) could result in the underestimation of concentration gradients of CO₂ in the early morning.

As seen in the scatter plot of $\Delta XCH_{4,sla}^S$ in Fig. 10(a), the modelled values are mostly positive during the day, while the
 490 measured concentration gradients between the down- and the upwind sites are mainly negative. That is, the instruments always measure strong signals at the upwind sites compared to the downwind sites, which cannot be reproduced by the model. As a large methane sink over the city is not expected, the most likely cause for this phenomenon is missing or underestimated emission sources located upstream of the upwind sites, located somewhere to the northeast or east of the Garching and Markt Schwaben stations in the case with NE/E winds. Likewise, with W/NW winds, the negative measured concentration gradients
 495 between the three down- and one upwind sites are found with -1.89 ppb in daily means and the model fails to reproduce




Figure 10. (a) Scatter plot of modelled and measured $\Delta XCH_{4,sla}^S$ for three targeted days and distribution of known emission sources located in the (b) eastern and (c) western area of Munich and the footprint contours for down- and upwind sites on (b) 16 and (c) 20 August: green for Markt Schwaben (East), red for Weßling (West), purple for Garching (North) and yellow for Höhenkirchen (South). The white solid line in (b) shows the location of the Burghausen-Fising-Amerdingen high pressure natural gas pipeline. The background maps use tiles from Stamen Design (<https://maps.stamen.com/>, under CC BY 3.0, with data by OpenStreetMap, under ODbL, from 2021 Dec. 21). The map colouring reflects the emissions from the initial inventory (i.e., TNO-MACCCo), with the same color bar as in Fig. 7.

these signals. Again, the measured column concentrations at the upwind sites (i.e., Weßling) are generally higher than at the downwind sites. Especially in the morning of 20 August, a clear strong increase was captured at the upwind side (see Fig. 5). However, none of these features could be replicated by the model. We postulate the presence of an unknown or underestimated source of emissions located upstream of Weßling as the most likely explanation.

500 5.5 Localizing unknown emission sources

To further localize the underestimated or unknown emissions at the upstream areas of both upwind sides (i.e., the western area of Weßling and the eastern area of Markt Schwaben), the footprint contours are used to interpret the transport of air masses. Figure 10 shows the footprint contours of the up- and the downwind sites with two different wind conditions (b) for 16 and (c) for 20 August and their receptor times are listed in Table S11 of the supplement.

505 As mentioned in Sect. 4.3, the major contributors of CH_4 related to human activities are waste management, agriculture and industries (USEPA, 2019). A number of sources from these sectors in and around the areas covered by the footprint contours are identified based on the European industrial emissions portal (available as <https://industry.eea.europa.eu/>, last access: 14 December, 2021), the initial emission inventory and local knowledge, which are marked in Fig. 10(b & c). The waste management here refers to landfill and wastewater treatment facilities, the emissions induced by agriculture come from
510 livestock and the industry emissions are from heat and gas production, manufacturing, etc. Another potential source of the



observed signals that was not represented in our emission inventory was a high-pressure natural gas pipeline that passes through the eastern section of our domain. This pipeline was reportedly under construction and tested in 2018 (constructed by Bayernets GmbH; Macht (2017); Bayernets).

Clearly, we cannot yet determine the exact cause of the observed peaks that were not reproduced in the model. However, with
515 a longer observation record and refinements to the modelling approach, we see potential to track down strong emitters of GHGs, which could provide information for supporting mitigation strategies. Here, the year-round measurements from MUCCnet, which cover a wide range of weather conditions and complete seasonal cycles, will help to complement and improve general inventories.

6 Conclusion

520 We have developed a WRF modelling framework for Munich to accompany MUCCnet and provide regularly updated concentration maps. Compared to a previous study for the city of Berlin (Zhao et al., 2019), we have introduced additional tagged tracers and improved model inputs. Measurements from MUCCnet and meteorological stations have been used to validate the model. Simulated slant column concentrations, extracted from the model with a smoothing consistent with the instrument characteristics, show encouraging general agreement with observations. We have then focused on comparing modelled column
525 concentrations to measurements and identified the flux categories responsible for the observed signals. The diurnal cycle of CO₂ was not well captured, with the model showing lower concentration enhancements in the morning, while agreeing well in the afternoon. We suspect that this is related to underestimated nighttime RES fluxes in VPRM, which has been reported in another recent study. Our study concludes with a refined application of DCM, aided by air-mass transport tracing with the STILT model. In an exploratory application using CH₄, this enabled us to identify unexpected signals in the measurements and
530 to roughly estimate the potential source regions.

This study focused on a one-month period of measurements to refine the modelling approach, but the continuous measurements of MUCCnet are ongoing. This increasingly long and complete data set will enable emission sources to be monitored on a regular basis. The evaluation of model output and measurements for longer time periods, with various wind conditions, will allow for a better localisation of sources, and for an improvement of localised emission inventories. We are looking forward
535 to conducting further studies in this direction, in particular as Munich is a main pilot city in the EU H2020 project ICOS Cities. An application to the period of the COVID pandemic with varying lockdown conditions is also foreseen. Inclusion of measurements from aircraft (e.g., In-service Aircraft for a Global Observing System (IAGOS) project – www.iagos.org) and satellites (e.g., the Orbiting Carbon Observatory-2/3 (OCO-2/3) operated by NASA) will help to further validate and make use of our models. Our data-model framework will bring us closer to the aim of effectively tracing GHG emissions and improving
540 emission inventories.

Author contributions.



The modelling framework designed for Munich is mainly built by XZ with joint effort from JM, MG, CG, SH and JC. Regarding the initialization, JM, MG and CG retrieved the CAMS fields, and preprocessed the vegetation indices and the necessary variables for the Kaplan model. JC, DF and AW supplied the measurement. CG, JG, JM, AS and XZ put the effort
545 in the improvement of VPRM. SH gave guidance on running the WRF model on the Linux Cluster of Leibniz Supercomputing Centre (LRZ, Garching b. M., Germany). XZ, JC and SH designed the computational framework. XZ and JC performed the analysis of the results with support of JM, CG and MG. With the input from all authors, XZ wrote the manuscript. All authors provided critical feedback and helped shape the research, analysis and manuscript.

Data availability.

550 The model data used to support the results described in this paper are available upon request to the first and corresponding authors.

Competing interests.

The authors declare that they have no conflict of interest.

Acknowledgements. TUM authors as well as MUCCnet are supported by the German Research Foundation (DFG, grant nos. CH 1792/2-1,
555 INST 95/1544).. WRF and STILT workflows were run on the CoolMUC-2 High-Performance Computing (HPC) system, part of the Linux Cluster at LRZ, where also two nodes of ESM are housed.

Ankit Shekhar acknowledges funding by ETH Zürich project FEVER ETH-27 19-1.



References

- Allen, M., Babiker, M., Chen, Y., de Coninck, H., Connors, S., van Diemen, R., Dube, O., Ebi, K., Engelbrecht, F., Ferrat, M., et al.: Summary
560 for policymakers Global Warming of 1.5 °C: an IPCC Special Report on the Impacts of Global Warming of 1.5 °C Above Pre-Industrial
Levels and Related Global Greenhouse Gas Emissions Pathways, in the Context of Strengthening the Global Response to the Threat of
Climate Change, World Meteorological Organization, pp. 1–24, 2018.
- Bayernets: Erdgasfernleitung Monaco Von Burghausen Nach Finsing, [https://www.bayernets.de/fileadmin/Dokumente/Broschueren/
bayernets_Informationsbroschuere_MONACO.pdf](https://www.bayernets.de/fileadmin/Dokumente/Broschueren/bayernets_Informationsbroschuere_MONACO.pdf), bayernets GmbH, last access: 11, April 2022.
- 565 Beck, V., Koch, T., Kretschmer, R., Marshall, J. and Ahmadov, R., Gerbig, C., Pillai, D., and Heimann, M.: The WRF Greenhouse Gas Model
(WRF-GHG). Technical Report No. 25, Max Planck Institute for Biogeochemistry, Jena, Germany., 2012.
- Beck, V., Gerbig, C., Koch, T., Bela, M., Longo, K., Freitas, S., Kaplan, J., Prigent, C., Bergamaschi, P., and Heimann, M.: WRF-Chem
simulations in the Amazon region during wet and dry season transitions: evaluation of methane models and wetland inundation maps,
Atmospheric Chemistry and Physics, 13, 7961–7982, <https://doi.org/10.5194/acp-13-7961-2013>, 2013.
- 570 Borsdorff, T., Hasekamp, O., Wassmann, A., and Landgraf, J.: Insights into Tikhonov regularization: application to trace gas col-
umn retrieval and the efficient calculation of total column averaging kernels, Atmospheric Measurement Techniques, 7, 523–535,
<https://doi.org/10.5194/amt-7-523-2014>, 2014.
- Brousse, O., Martilli, A., Foley, M., Mills, G., and Bechtel, B.: WUDAPT, an efficient land use producing data tool for mesoscale models?
Integration of urban LCZ in WRF over Madrid, Urban Climate, 17, 116–134, <https://doi.org/10.1016/j.uclim.2016.04.001>, 2016.
- 575 Browne, P. A., de Rosnay, P., Zuo, H., Bennett, A., and Dawson, A.: Weakly coupled ocean–atmosphere data assimilation in the ECMWF
NWP system, Remote Sensing, 11, 234, <https://doi.org/10.3390/rs11030234>, 2019.
- Brunner, D., Kuhlmann, G., Marshall, J., Clément, V., Fuhrer, O., Broquet, G., Löscher, A., and Meijer, Y.: Accounting for the vertical dis-
tribution of emissions in atmospheric CO₂ simulations, Atmospheric Chemistry and Physics, 19, 4541–4559, [https://doi.org/10.5194/acp-
19-4541-2019](https://doi.org/10.5194/acp-
19-4541-2019), 2019.
- 580 Chen, J., Viatte, C., Hedelius, J. K., Jones, T., Franklin, J. E., Parker, H., Gottlieb, E. W., Wennberg, P. O., Dubey, M. K., and Wofsy, S. C.:
Differential column measurements using compact solar-tracking spectrometers, Atmospheric Chemistry and Physics, 16, 8479–8498,
<https://doi.org/10.5194/acp-16-8479-2016>, 2016.
- Chen, J., Gerbig, C., Marshall, J., and Totsche, K. U.: Short-term forecasting of regional biospheric CO₂ fluxes in Europe using a light-use-
efficiency model (VPRM, MPI-BGC version 1.2), Geoscientific Model Development, 13, 4091–4106, [https://doi.org/10.5194/gmd-13-
4091-2020](https://doi.org/10.5194/gmd-13-
4091-2020), 2020.
- 585 Ching, J., Mills, G., Bechtel, B., See, L., Feddema, J., Wang, X., Ren, C., Brousse, O., Martilli, A., Neophytou, M., et al.: WUDAPT: An
urban weather, climate, and environmental modeling infrastructure for the anthropocene, Bulletin of the American Meteorological Society,
99, 1907–1924, <https://doi.org/10.1175/BAMS-D-16-0236.1>, 2018.
- DeCola, P., Tarasova, O., Brunner, D., Maksyutov, S., Manning, A., Vogel, F., Gurney, K., Turnbull, J., Zavala-Araiza, D., Kort, E., Robinson,
590 R., Canadell, P., Ciais, P., Vladu, F., Houweling, S., Lauvaux, T., and Mueller, K.: An Integrated Global Greenhouse Gas Information
System (IG3IS) Science Implementation Plan Approved by EC-70, World Meteorological Organization (WMO), 2018.
- Dekker, I. N., Houweling, S., Aben, I., Röckmann, T., Krol, M., Martínez-Alonso, S., Deeter, M. N., and Worden, H. M.: Quantification of
CO emissions from the city of Madrid using MOPITT satellite retrievals and WRF simulations, Atmospheric chemistry and physics, 17,
14 675–14 694, <https://doi.org/10.5194/acp-2017-418>, 2017.



- 595 Demuzere, M., Bechtel, B., Middel, A., and Mills, G.: Mapping Europe into local climate zones, *PloS one*, 14, e0214474, <https://doi.org/10.1371/journal.pone.0214474>, 2019.
- Dietrich, F., Chen, J., Voggenreiter, B., Aigner, P., Nachtigall, N., and Reger, B.: MUCCnet: Munich Urban Carbon Column network, *Atmospheric Measurement Techniques*, 14, 1111–1126, <https://doi.org/10.5194/amt-14-1111-2021>, 2021.
- Durre, I., Yin, X., Vose, R. S., Applequist, S., and Arnfield, J.: Enhancing the data coverage in the Integrated Global Radiosonde Archive, *Journal of Atmospheric and Oceanic Technology*, 35, 1753–1770, <https://doi.org/10.1175/JTECH-D-17-0223.1>, 2018.
- 600 EU Commission, .: A Clean Planet for all-A European strategic long-term vision for a prosperous, modern, competitive and climate neutral economy, Tech. rep., Technical report, European Commission, Brussels. Federal Ministry of Interior Building and Community (IBC)-Sustainable Building Assessment System (BNB), 2018.
- Fasoli, B., Lin, J. C., Bowling, D. R., Mitchell, L., and Mendoza, D.: Simulating atmospheric tracer concentrations for spatially distributed
605 receptors: updates to the Stochastic Time-Inverted Lagrangian Transport model's R interface (STILT-R version 2), *Geoscientific Model Development*, 11, 2813–2824, <https://doi.org/10.5194/gmd-11-2813-2018>, 2018.
- Frey, M., Sha, M. K., Hase, F., Kiel, M., Blumenstock, T., Harig, R., Surawicz, G., Deutscher, N. M., Shiomi, K., Franklin, J. E., et al.: Building the COllaborative Carbon Column Observing Network (COCCON): long-term stability and ensemble performance of the EM27/SUN Fourier transform spectrometer, *Atmospheric Measurement Techniques*, 12, 1513–1530, <https://doi.org/10.5194/amt-12-1513-2019>, 2019.
- 610 Gałkowski, M., Jordan, A., Rothe, M., Marshall, J., Koch, F.-T., Chen, J., Agusti-Panareda, A., Fix, A., and Gerbig, C.: In situ observations of greenhouse gases over Europe during the CoMet 1.0 campaign aboard the HALO aircraft, *Atmospheric Measurement Techniques*, 14, 1525–1544, <https://doi.org/10.5194/amt-14-1525-2021>, 2021.
- Georgiou, G. K., Christoudias, T., Proestos, Y., Kushta, J., Hadjinicolaou, P., and Lelieveld, J.: Air quality modelling in the summer over the eastern Mediterranean using WRF-Chem: chemistry and aerosol mechanism intercomparison, *Atmospheric Chemistry and Physics*, 18,
615 1555–1571, <https://doi.org/10.5194/acp-18-1555-2018>, 2018.
- Gisi, M., Hase, F., Dohe, S., Blumenstock, T., Simon, A., and Keens, A.: XCO₂-measurements with a tabletop FTS using solar absorption spectroscopy, *Atmospheric Measurement Techniques*, 5, 2969–2980, <https://doi.org/10.5194/amt-5-2969-2012>, 2012.
- Gourdji, S., Karion, A., Lopez-Coto, I., Ghosh, S., Mueller, K. L., Zhou, Y., Williams, C. A., Baker, I. T., Haynes, K., and Whetstone, J.: A modified Vegetation Photosynthesis and Respiration Model (VPRM) for the eastern USA and Canada, evaluated with comparison to atmospheric observations and other biospheric models, *Earth and Space Science Open Archive*, p. 50,
620 <https://doi.org/10.1002/essoar.10506768.1>, 2021.
- Hase, F., Frey, M., Blumenstock, T., Groß, J., Kiel, M., Kohlhepp, R., Mengistu Tsidu, G., Schäfer, K., Sha, M., and Orphal, J.: Application of portable FTIR spectrometers for detecting greenhouse gas emissions of the major city Berlin, *Atmospheric Measurement Techniques*, 8, 3059–3068, <https://doi.org/10.5194/amt-8-3059-2015>, 2015.
- 625 Hase, F., Frey, M., Kiel, M., Blumenstock, T., Harig, R., Keens, A., and Orphal, J.: Addition of a channel for XCO observations to a portable FTIR spectrometer for greenhouse gas measurements, *Atmospheric Measurement Techniques*, 9, 2303–2313, <https://doi.org/10.5194/amt-9-2303-2016>, 2016.
- Hedelius, J. K., Viatte, C., Wunch, D., Roehl, C. M., Toon, G. C., Chen, J., Jones, T., Wofsy, S. C., Franklin, J. E., Parker, H., et al.: Assessment of errors and biases in retrievals of XCO₂, XCH₄, XCO, and XN₂O from a 0.5 cm⁻¹ resolution solar-viewing spectrometer,
630 *Atmospheric Measurement Techniques*, 9, 3527–3546, <https://doi.org/10.5194/amt-9-3527-2016>, 2016.



- Heimburger, A. M., Harvey, R. M., Shepson, P. B., Stirn, B. H., Gore, C., Turnbull, J., Cambaliza, M. O., Salmon, O. E., Kerlo, A.-E. M., Lavoie, T. N., et al.: Assessing the optimized precision of the aircraft mass balance method for measurement of urban greenhouse gas emission rates through averaging, *Elementa: Science of the Anthropocene*, 5, <https://doi.org/10.1525/elementa.134>, 2017.
- Heinle, L. and Chen, J.: Automated enclosure and protection system for compact solar-tracking spectrometers, *Atmospheric Measurement Techniques*, 11, 2173–2185, <https://doi.org/10.5194/amt-11-2173-2018>, 2018.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., et al.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- Hu, X.-M., Gourdj, S. M., Davis, K. J., Wang, Q., Zhang, Y., Xue, M., Feng, S., Moore, B., and Crowell, S. M.: Implementation of improved parameterization of terrestrial flux in WRF-VPRM improves the simulation of nighttime CO₂ peaks and a daytime CO₂ band ahead of a cold front, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD034362, <https://doi.org/10.1029/2020JD034362>, 2021.
- IRENA, I.: Renewable energy in cities, International Renewable Agency: Abu Dhabi, UAE, 2016.
- Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Dentener, F., Bergamaschi, P., Pagliari, V., Olivier, J. G., Peters, J. A., et al.: EDGAR v4.3.2 Global Atlas of the three major greenhouse gas emissions for the period 1970–2012, *Earth System Science Data*, 11, 959–1002, <https://doi.org/10.5194/essd-11-959-2019>, 2019.
- Jones, T. S., Franklin, J. E., Chen, J., Dietrich, F., Hajny, K. D., Paetzold, J. C., Wenzel, A., Gately, C., Gottlieb, E., Parker, H., et al.: Assessing Urban Methane Emissions using Column Observing Portable FTIR Spectrometers and a Novel Bayesian Inversion Framework, *Atmospheric Chemistry and Physics Discussions*, pp. 1–30, <https://doi.org/10.5194/acp-2020-1262>, 2021.
- Jung, M., Henkel, K., Herold, M., and Churkina, G.: Exploiting synergies of global land cover products for carbon cycle modeling, *Remote Sensing of Environment*, 101, 534–553, <https://doi.org/10.1016/j.rse.2006.01.020>, 2006.
- Kaplan, J., Folberth, G., and Hauglustaine, D.: Role of methane and biogenic volatile organic compound sources in late glacial and Holocene fluctuations of atmospheric methane concentrations, *Global Biogeochemical Cycles*, 20, <https://doi.org/10.1029/2005GB002590>, 2006.
- Klausner, T., Mertens, M., Huntrieser, H., Galkowski, M., Kuhlmann, G., Baumann, R., Fiehn, A., Jöckel, P., Pühl, M., Roiger, A., et al.: Urban greenhouse gas emissions from the Berlin area: A case study using airborne CO₂ and CH₄ in situ observations in summer 2018, *Elementa: Science of the Anthropocene*, 8, 2020.
- Lauvaux, T., Miles, N. L., Deng, A., Richardson, S. J., Cambaliza, M. O., Davis, K. J., Gaudet, B., Gurney, K. R., Huang, J., O’Keefe, D., et al.: High-resolution atmospheric inversion of urban CO₂ emissions during the dormant season of the Indianapolis Flux Experiment (INFLUX), *Journal of Geophysical Research: Atmospheres*, 121, 5213–5236, <https://doi.org/10.1002/2015JD024473>, 2016.
- Macht, V.: Die Pipeline ist im Anmarsch, <https://www.merkur.de/lokales/erding/isen-ort86592/erdgasfernleitung-monaco-pipeline-ist-im-anmarsch-8372.html>, merkur.de, last access: 11, April 2022, 2017.
- Mahadevan, P., Wofsy, S. C., Matross, D. M., Xiao, X., Dunn, A. L., Lin, J. C., Gerbig, C., Munger, J. W., Chow, V. Y., and Gottlieb, E. W.: A satellite-based biosphere parameterization for net ecosystem CO₂ exchange: Vegetation Photosynthesis and Respiration Model (VPRM), *Global Biogeochemical Cycles*, 22, <https://doi.org/10.1029/2006GB002735>, 2008.
- Masson-Delmotte, V., Zhai, P., Pörtner, H.-O., Roberts, D., Skea, J., Shukla, P., Pirani, A., Moufouma-Okia, W., Péan, C., Pidcock, R., Connors, S., Matthews, J., Chen, Y., Zhou, X., Gomis, M., Lonnoy, E., Maycock, T., and Tignor, M. and Waterfield, T.: IPCC, 2018: Summary for policymakers. In: Global warming of 1.5 °C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty, World Meteorological Organization, Geneva, Switzerland, 10, 2018.



- Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M.,
670 Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B.: Summary for Policymakers. In:
Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovern-
mental Panel on Climate Change, Cambridge University Press, pp. 1–24, 2021.
- Mughal, M. O.: Modelling the Urban Heat Island in Singapore-state of the art WRF model technical details, Technical Report Cooling
Singapore, 500, <https://doi.org/10.3929/ethz-b-000412358>, 2020.
- 675 Ohyama, H., Morino, I., Velazco, V. A., Klausner, T., Bagtasa, G., Kiel, M., Frey, M., Hori, A., Uchino, O., Matsunaga, T., et al.: Validation of
XCO₂ and XCH₄ retrieved from a portable Fourier transform spectrometer with those from in situ profiles from aircraft-borne instruments,
Atmospheric Measurement Techniques, 13, 5149–5163, <https://doi.org/10.5194/amt-13-5149-2020>, 2020.
- Peckham, S. E., Grell, G. A., McKeen, S. A., Ahmadov, R., Barth, M., Pfister, G., Wiedinmyer, C., Fast, J. D., Gustafson, W. I., Ghan, S. J.,
Zaveri, R., Easter, R. C., Barnard, J., Chapman, E., Hewson, M., Schmitz, R., Salzmann, M., and Freitas, S. R.: WRF-Chem Version
680 3.9.1.1 User's Guide, Tech. rep., National Center for Atmospheric Research, https://ruc.noaa.gov/wrf/wrf-chem/Users_guide.pdf, 2017.
- Rémy, S., Kipling, Z., Flemming, J., Boucher, O., Nabat, P., Michou, M., Bozzo, A., Ades, M., Huijnen, V., Benedetti, A., et al.: Description
and evaluation of the troposphere aerosol scheme in the European Centre for Medium-Range Weather Forecasts (ECMWF) Integrated
Forecasting System (IFS-AER, cycle 45R1), Geoscientific Model Development, 12, 4627–4659, [https://doi.org/10.5194/gmd-12-4627-](https://doi.org/10.5194/gmd-12-4627-2019)
2019, 2019.
- 685 Sargent, M., Barrera, Y., Nehrkorn, T., Hutyra, L. R., Gately, C. K., Jones, T., McKain, K., Sweeney, C., Hegarty, J., Hardiman, B., et al.:
Anthropogenic and biogenic CO₂ fluxes in the Boston urban region, Proceedings of the National Academy of Sciences, 115, 7491–7496,
<https://doi.org/10.1073/pnas.1803715115>, 2018.
- Shekhar, A., Chen, J., Paetzold, J. C., Dietrich, F., Zhao, X., Bhattacharjee, S., Ruisinger, V., and Wofsy, S. C.: Anthropogenic CO₂
emissions assessment of Nile Delta using XCO₂ and SIF data from OCO-2 satellite, Environmental Research Letters, 15, 095 010,
690 <https://doi.org/10.1088/1748-9326/ab9cfe>, 2020.
- Skamarock, William C. and Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Duda, M. G., Huang, X.-Y., Wang, W., and Pow-
ers, J. G.: Description of the Advanced Research WRF Version 3, Tech. rep., National Center for Atmospheric Research, <https://pdfs.semanticscholar.org/ace5/4d4d1d6c9914997ad8f4e410044fdeb95b9d.pdf>, 2008.
- Staufer, J., Broquet, G., Bréon, F.-M., Puygrenier, V., Chevallier, F., Xueref-Rémy, I., Dieudonné, E., Lopez, M., Schmidt, M., Ramonet, M.,
695 et al.: The first 1-year-long estimate of the Paris region fossil fuel CO₂ emissions based on atmospheric inversion, Atmospheric Chemistry
and Physics, 16, 14 703–14 726, <https://doi.org/10.5194/acp-16-14703-2016>, 2016.
- Super, I., Denier van der Gon, H. A., van der Molen, M. K., Sterk, H. A., Hensen, A., and Peters, W.: A multi-model approach
to monitor emissions of CO₂ and CO in an urban-industrial complex, Atmospheric Chemistry and Physics Discussions, pp. 1–31,
<https://doi.org/10.5194/acp-2016-807>, 2016.
- 700 Super, I., Dellaert, S. N., Visschedijk, A. J., and Denier van der Gon, H. A.: Uncertainty analysis of a European high-resolution emis-
sion inventory of CO₂ and CO to support inverse modelling and network design, Atmospheric Chemistry and Physics, 20, 1795–1816,
<https://doi.org/10.5194/acp-20-1795-2020>, 2020.
- Toja-Silva, F., Chen, J., Hachinger, S., and Hase, F.: CFD simulation of CO₂ dispersion from urban thermal power plant: Analysis of
turbulent Schmidt number and comparison with Gaussian plume model and measurements, Journal of Wind Engineering and Industrial
705 Aerodynamics, 169, 177–193, <https://doi.org/10.1016/j.jweia.2017.07.015>, 2017.



- Tu, Q., Hase, F., Blumenstock, T., Kivi, R., Heikkinen, P., Sha, M. K., Raffalski, U., Landgraf, J., Lorente, A., Borsdorff, T., Chen, H., Dietrich, F., and Chen, J.: Atmospheric CO₂ and CH₄ abundances on regional scales in boreal areas using Copernicus Atmosphere Monitoring Service (CAMS) analysis, Collaborative Carbon Column Observing Network (COCCON) spectrometers, and Sentinel-5 Precursor satellite observations, *Atmospheric Measurement Techniques*, 13, 4751–4771, <https://doi.org/amt-13-4751-2020>, 2020.
- 710 Turner, A. J., Shusterman, A. A., McDonald, B. C., Teige, V., Harley, R. A., and Cohen, R. C.: Network design for quantifying urban CO₂ emissions: assessing trade-offs between precision and network density, *Atmospheric Chemistry and Physics*, 16, 13465–13475, <https://doi.org/10.5194/acp-16-13465-2016>, 2016.
- USEPA: Global Non-CO₂ Greenhouse Gas Emission Projections Mitigation Potential: 2015-2050, Tech. rep., US Environmental Protection Agency Washington, DC, <https://www.epa.gov/globalmitigation-non-co2-greenhouse-gases>, ePA-430-R-19-010, 2019.
- 715 van der Gon, H. D., Kuenen, J., Boleti, E., Muntean, M., Greet, M., Marshall, J., and Haussaire, J.-M.: Emissions and natural fluxes Dataset, 2019.
- Vermote, E.: MOD09A1 MODIS Surface Reflectance 8-Day L3 Global 500m SIN Grid V006, <https://modis.gsfc.nasa.gov/data/dataproduct/mod09.php>, <https://doi.org/10.5067/MODIS/MOD09A1.006>, 2015.
- Viatte, C., Lauvaux, T., Hedelius, J. K., Parker, H., Chen, J., Jones, T., Franklin, J. E., Deng, A. J., Gaudet, B., Verhulst, K., Duren, R.,
720 Wunch, D., Roehl, C., Dubey, M., Wofsy, S., and Wennberg, P.: Methane emissions from dairies in the Los Angeles Basin, *Atmospheric Chemistry and Physics*, 17, 7509–7528, <https://doi.org/10.5194/acp-17-7509-2017>, 2017.
- Vogel, F. R., Frey, M., Staufer, J., Hase, F., Broquet, G., Xueref-Remy, I., Chevallier, F., Ciais, P., Sha, M. K., Chelin, P., et al.: XCO₂ in an emission hot-spot region: the COCCON Paris campaign 2015, *Atmospheric Chemistry and Physics*, 19, 3271–3285, <https://doi.org/10.5194/acp-19-3271-2019>, 2019.
- 725 Wei, T., Wu, J., and Chen, S.: Keeping track of greenhouse gas emission reduction progress and targets in 167 cities worldwide, *Frontiers in Sustainable Cities*, p. 64, <https://doi.org/10.3389/frsc.2021.696381>, 2021.
- Zhao, X., Marshall, J., Hachinger, S., Gerbig, C., Frey, M., Hase, F., and Chen, J.: Analysis of total column CO₂ and CH₄ measurements in Berlin with WRF-GHG, *Atmospheric Chemistry and Physics*, 19, 11279–11302, <https://doi.org/10.5194/acp-19-11279-2019>, 2019.