



Parameterizing the aerodynamic effect of trees in street canyons for the street-network model MUNICH using the CFD model Code_Saturne

Alice Maison^{1,2}, Cédric Flageul³, Bertrand Carissimo¹, Andrée Tuzet², Yunyi Wang¹, and Karine Sartelet¹

¹CEREA, École des Ponts, EDF R&D, Marne-la-Vallée, France

²Université Paris-Saclay, INRAE, AgroParisTech, UMR EcoSys, 78850 Thiverval-Grignon, France

³PPRIME Institute, Curiosity Group, Université de Poitiers, CNRS, ISAE-ENSMA, France

Correspondence: Alice Maison (alice.maison@enpc.fr), Karine Sartelet (karine.sartelet@enpc.fr)

Abstract. Trees provide many ecosystem services in cities such as urban heat island reduction, water runoff limitation and carbon storage. However, the presence of trees in street canyons reduces the wind velocity in the street and limits pollutant dispersion. Thus, to get accurate simulations of pollutant concentrations, the aerodynamic effect of trees should be taken into account in air-quality models at the street level.

5 The Model of Urban Network of Intersecting Canyons and Highways (MUNICH) simulates the pollutant concentrations in a street-network, considering dispersion and physico-chemical processes. It can be coupled to a regional-scale chemical-transport model to simulate air quality over districts or cities. The aerodynamic effect of tree crown is parameterized here, through their impacts on the average wind velocity in the street direction and the vertical transfer coefficient associated with the dispersion of a tracer. The parameterization is built using local-scale simulations performed with the Computational Fluid Dynamics
10 (CFD) code Code_Saturne. Two-dimensional CFD simulations in an infinite street canyon are used to quantify the effect of trees depending on the tree characteristics (Leaf Area Index, crown volume fraction and tree height to street height ratio) using a drag-porosity approach. The tree crown slows down the flow and produces turbulent kinetic energy in the street, thus impacting the tracer dispersion. This effect increases with the Leaf Area Index and the crown volume fraction of the trees: the average horizontal velocity in the street is reduced up to 68 % and the vertical transfer coefficient up to 23 % in the simulations
15 performed here.

A parameterization of these effects on horizontal and vertical transfers for the street model MUNICH is proposed. Existing parameterizations in MUNICH are modified based on Code_Saturne simulations to account for both building and tree effects on vertical and horizontal transfers. The parameterization is built to get similar tree effects (quantified by a relative deviation between the cases without and with trees) between Code_Saturne and MUNICH. The vertical wind profile and mixing length
20 depend on Leaf Area Index, crown radius and tree height to street height ratio. The interaction between the trees and the street aspect ratio is also considered.



1 Introduction

Cities are, by definition, areas with high density of people, infrastructures and activities, and this urbanization causes many issues. First, air quality is poor because of the numerous air pollutants emitted by anthropic activities such as traffic, industries or residential activities, and the reduction of air flow by high buildings limits the dispersion of these pollutants (Faiz, 1993; Akimoto, 2003; Yuan et al., 2014; Zhang et al., 2020). In addition to air pollution issues, radiative and water budgets are strongly modified in cities compared to countryside (Bozonnet et al., 2015). Temperatures are, in average, higher than in the countryside because of the urban heat island created by additional anthropogenic energy released, storage of radiative energy in dark materials, radiation multireflection and lack of vegetation and associated evapotranspiration (Oke, 1982; Pigeon et al., 2007; Stewart, 2011; Hebbert and Jankovic, 2013). Impervious soils also decrease water infiltration and intensify runoff (Leopold, 1968). In addition, growing urbanisation and increasing extreme events (due to climate change) such as pollution peaks, heat waves and floods have negative consequences on environment and human health (Robine et al., 2007; Angel et al., 2011; Pascal et al., 2013; West et al., 2016; IPCC, 2021).

One nature-based solution is to green the city by planting vegetation as lawns, trees in streets or in parks, green walls and roofs (Livesley et al., 2016; Revelli and Porporato, 2018). Vegetation and especially trees contribute to improve human thermal comfort by creating favorable micro-climate with lower air temperature (through solar radiation interception and creation of shade) and higher evaporation (through ground and vegetation evapotranspiration) (Taha et al., 1991; Bowler et al., 2010; Gillner et al., 2015; Klemm et al., 2015; Lobaccaro and Acero, 2015; Gunawardena et al., 2017). This positive effect of trees is significant in particular during heat wave episodes, which will be more frequent in the future due to climate change (IPCC, 2021). Trees and vegetated areas also favorize infiltration in soils that contributes to offset water runoff induced by soil artificialization (Armson et al., 2013; Berland et al., 2017). Besides, vegetation is known to store carbon (Nowak and Crane, 2002; Svirejeva-Hopkins et al., 2004) and to enhance human well-being (van Dillen et al., 2012; Bertram and Rehdanz, 2015; Krekel et al., 2015). For all these ecosystem services in urban areas, city greening is often promoted and for example the city of Paris has about 205,000 trees of which 52 % are roadside trees (Municipality of Paris, 2021).

Many studies have tried to figure out the impact of trees on air pollution in a street canyon and they have shown that trees are an important parameter to take into account if we want to understand and simulate accurately the pollutant concentrations in the streets (Beckett et al., 1998; Nowak et al., 1998, 2006; Jayasooriya et al., 2017). Vegetation and especially trees represent surfaces available for pollutant dry deposition, and hence can contribute to reduce air pollutants concentrations. However, pollutant removal and its impact on air quality vary greatly depending on tree characteristics, tree species and pollutant type (Hwang et al., 2011; Selmi et al., 2016; Xue and Li, 2017; Ozdemir, 2019). Trees may alter air quality by influencing aerodynamic processes and limiting the pollutant dispersion (Buccolieri et al., 2011; Wania et al., 2012; Vos et al., 2013; Gromke and Ruck, 2007, 2009, 2012; Gromke and Blocken, 2015). The aerodynamic effect of trees is defined as the drag force resulting from the friction between the air and the leaves. Since the tree crown can be seen as a porous medium, as air passes through it but is slowed down and the drag force increases with the leaf surface.

For a wind perpendicular to a street, the air recirculate inside the street canyon (Harman et al., 2004). Pollutants emitted at the



bottom of the street (by traffic) accumulate on the leeward side of the street inducing higher local concentrations (Vardoulakis et al., 2003; Cai et al., 2008; Huang et al., 2019). Obstacles in the street, such as trees, can strongly impact the air flow and intensify the pollutant accumulation (Vos et al., 2013). This effect has been studied using Computational Fluid Dynamics (CFD) models where the aerodynamic influence of trees on the flow is represented by a porosity model (Buccolieri et al., 2009; Zaïdi et al., 2013; Wei et al., 2016; Jeanjean et al., 2017; Santiago et al., 2017). Vos et al. (2013) showed that pollutant concentrations may increase by 20 % in a street, because of the presence of two rows of trees in the street. The effects of vegetation depend on the height, width and density (Leaf Area Index) of trees (Vos et al., 2013; Janhäll, 2015), as well as the height to width ratio of streets (Wania et al., 2012). Finally, it is necessary to assess accurately the effect of trees on pollutant dispersion in order to find what configurations are more effective in reducing air pollution in street canyon and to guide urban development policy (Janhäll, 2015).

The effect of trees on aerodynamic processes should also be considered at the street level in air quality models. As discussed previously, CFD models including trees are used to study wind fields and pollutant transport in street canyons (Li et al., 2006). However, as the street is discretized with a fine mesh, the computational cost is high and simulations at the city scale are too expensive today. Fast-running codes such as simplified street-network or street-in-grid models are developed to simulate street pollutant concentrations over neighbourhood or cities, but they do not take into account the effect of trees in the streets. The objective of this study is to parameterize the effect of trees on air flow in the Model of Urban Network of Intersecting Canyons and Highways (MUNICH) (Kim et al. (2018); Lugon et al. (2019); <http://cerea.enpc.fr/munich/>, last access on 17 December 2021). To build this parameterization, simulations in street canyons are performed with Code_Saturne (Archambeau et al. (2004); <https://www.code-saturne.org/>, last access on 17 December 2021), a CFD code, which can represent the tree aerodynamic effect with a drag porosity approach (Katul et al., 2004) and has previously been compared with field measurements (Zaïdi et al., 2013). MUNICH parameterizations have already been compared with Code_Saturne results in a treeless canyon and a new parameterization for horizontal and vertical transfers have been developed in MUNICH based on Wang (2012, 2014) and Code_Saturne simulations (Maison et al., 2022). In the present study, the tree aerodynamic effect is added to this parameterization and Code_Saturne (version 6.0) is used as a reference to parameterize the aerodynamic effect of trees in the street network model MUNICH. CFD simulations are performed in three streets of aspect ratios varying from 0.3 to 1.0 and a large range of tree leaf area index, crown radius and heights is tested. The tree aerodynamic effect quantified with Code_Saturne is analysed depending on these street and tree characteristics.

The approach to model the dispersion of pollutants in MUNICH and Code_Saturne is fundamentally different due to their physical modeling and discretization in space and time. MUNICH is a street-network model that simulates air pollutant concentration in a urban canopy. Street dimensions and pollutant concentrations are assumed to be homogeneous in each street segment. Air flow is divided into horizontal flux from one street to another, and vertical flux between the street and the background (Maison et al., 2022). Background concentrations above the street can be computed by 3D Chemistry-transport models (CTMs), such as Polair3D (Sartelet et al., 2018; Lugon et al., 2019, 2021). To build the tree parameterization, the CFD simulation setup is adapted for the comparison with MUNICH and several simulations are performed with Code_Saturne considering a range of street and tree characteristics.



The structure of the paper is as follows. The MUNICH and Code_Saturne models are presented in Sect. 2. Then, the tree effect on horizontal wind speed and vertical transfer coefficient is quantified with Code_Saturne simulations in Sect. 3 and parameterized in MUNICH in Sect. 4. Conclusions are presented in Sect. 5.

2 Materials and methods

95 2.1 Description of MUNICH

In MUNICH, each street segment is assumed to be homogeneous, i.e. with uniform building height H and street width W , and of length L (m). The street is characterized by its height-to-width ratio called aspect ratio, $a_r = H/W$ (-). Only the average pollutant concentrations over the street are considered. Pollutants are transported by the horizontal wind speed (advection) in the street network and by a vertical transfer coefficient between the streets and the background. Several parameterizations of the horizontal wind speed and of the vertical transfer coefficient exist in MUNICH. The ones recently developed in Maison et al. (2022) and based on Code_Saturne simulations are used and detailed here. The vertical profile of the wind speed in the street direction is calculated as an attenuation of the wind speed in the street direction and at the roof level, $U_{H,\varphi}$ ($\text{m}\cdot\text{s}^{-1}$) as (Maison et al., 2022):

$$U(z) = U_{H,\varphi} [C_1 I_0(g(z)) + C_2 K_0(g(z))] \quad \text{with } U_{H,\varphi} = U_H |\cos(\varphi)|, \quad (1)$$

105 where C_1 and C_2 are integration coefficients, I_0 and K_0 are the first and second type modified Bessel functions of order 0. Besides, the wind speed at the roof level U_H has to be multiplied by $|\cos(\varphi)|$ to select the component of the wind speed in the street direction. This vertical wind profile is then integrated between the soil roughness z_{0s} (m) and H to compute the average horizontal wind speed in the street direction. The function g is calculated as (Wang, 2012, 2014):

$$g(z) = 2\sqrt{\alpha \frac{z}{H}}, \quad (2)$$

110 and:

$$C_1 = \frac{1}{I_0(g(H)) - I_0(g(z_{0s}))} \quad \text{and} \quad C_2 = -\frac{C_1 I_0(g(z_{0s}))}{K_0(g(z_{0s}))}, \quad (3)$$

where α is a dimensionless coefficient expressing the effects of wind angle on wind attenuation in the street, and it is computed as:

$$\alpha = \frac{C_B a_r}{\kappa s_H} \quad \text{with } C_B = 0.31 [1 - \exp(-1.6 a_r)] f_\varphi \quad (4)$$

$$115 \quad \text{and } f_\varphi = \begin{cases} |\cos(2\varphi)|^3 & \text{if } \varphi \in [0, 45^\circ] \cup [135, 225^\circ] \cup [315, 360^\circ] \\ 0 & \text{if } \varphi \in]45, 135^\circ[\cup]225, 315^\circ[\end{cases} \quad (5)$$

where $s_H = s(z = H)$ is a dimensionless factor describing the effect of canopy on the mixing length l_m (m) (Wang, 2012, 2014).

The mixing length is calculated as:

$$\frac{1}{l_m} = \frac{1}{\kappa z} + \frac{1}{l_{cb}} \implies l_m = \kappa z \frac{l_{cb}}{l_{cb} + \kappa z} = \kappa z s(z), \quad (6)$$



where κz corresponds to the mixing length over a rough bare soil (without canopy), and l_{cb} is the characteristic length (m) in
120 the street canyon, corresponding to the mixing length of the urban canopy alone ($l_{cb} = 0.5W$).

The vertical transfer coefficient that drives pollutant exchange between the street and the background zone is calculated at
 $z = H$ as:

$$q_{vert} = \sigma_W l_m = \sigma_W \kappa H s_H, \quad (7)$$

where σ_W (in $\text{m}\cdot\text{s}^{-1}$) is a velocity scale equal to the standard deviation of the vertical wind speed. It depends both on the
125 friction velocity above the urban canopy, u_* , and on atmospheric stability (Soulhac et al., 2011).

In section 4, the tree effect is parameterized by taking into account the characteristic length of the trees in the mixing length
 l_m (Eq. (19), which leads to modifications in the coefficient α Eq. (23)).

2.2 Description of Code_Saturne

2.2.1 Street and tree modeling setup

130 In the present study, the tree effect is studied in three street canyons of different street aspect ratios: a wide street canyon (WC),
an intermediate canyon (IC) and a narrow canyon (NC). Their characteristics are presented in Table 1:

Table 1. Characteristics of the three canyons studied.

Canyon	Building height H (m)	Street width W (m)	Street aspect ratio a_r (-)	Maximum height of the domain (m)
WC	8.5	27.5	0.3	25.5
IC	14.0	27.5	0.5	42.0
NC	27.5	27.5	1.0	82.5

A color code is introduced to simplify the reading of the figures: red symbols for the wide canyon (WC), purple ones for the
intermediate canyon (IC) and blue ones for the narrow canyon (NC). Abbreviations, parameters and variables used are listed
in Appendix A.

135 The $k - \varepsilon$ linear production turbulence model is used in Code_Saturne. Stationary simulations are performed with a thermally
neutral atmosphere. A 2D infinite street canyon is modeled with periodic condition in Y axis but the flow and the wind speed
vector are 3D. The mesh is composed of hexahedral cells of 1 m in Y axis direction and 0.5 m in X and Z axis. The vertical
profiles of U , k and ε are set in inlet (top left border of the domain). The complete description of Code_Saturne simulation
setup in treeless canyons can be found in Maison et al. (2022).

140 The tree geometry and its representation in Code_Saturne is shown in Fig. 1. Here, two rows of trees ($n = 2$) are considered,
one on each side of the street, whose positions on the X axis are $X = 34.5$ m and $X = 48.0$ m (the wall positions are $X = 27.5$
and $X = 55.0$ m). r is the tree radius (m), h , h_{min} and h_{max} correspond respectively to the middle, minimum and maximum
heights of the tree crown (m). Note that all the simulations verify $h_{max} \leq H$, i.e. the top of the trees do not exceed the top of
the street.




Figure 1. 3D scheme of the street canyon with cylindrical tree crown dimensions.

145 In the cells containing the trees, an additional drag term is added to the Navier-Stokes equations (Katul et al., 2004; Zaïdi et al., 2013):

$$S_{u,i} = -\rho LAD C_{D_t} |U| \vec{U}_i, \quad (8)$$

where $|U|$ is the modulus of U , U_i is the velocity in the i -direction, LAD is the Leaf area density in $\text{m}_{\text{leaves}}^2 \cdot \text{m}_{\text{tree crown}}^{-3}$, ρ is the air density and C_{D_t} is the tree drag coefficient set to 0.2, a representative value for trees (Katul and Albertson, 1998). The

150 Leaf Area Index (LAI) is the one-sided green leaf area per unit ground surface area and the Leaf Area Density (LAD) is the one-sided green leaf area per unit volume. They are calculated as:

$$LAI = \frac{\text{surface of leaves}}{\text{soil projected surface area}} \left(\text{m}_{\text{leaf}}^2 \cdot \text{m}_{\text{soil}}^{-2} \right) \quad \text{and} \quad LAD = \frac{\text{surface of leaves}}{\text{volume of tree crown}} \left(\text{m}_{\text{leaf}}^2 \cdot \text{m}_{\text{crown}}^{-3} \right). \quad (9)$$

Source terms for trees are also implemented in the $k - \varepsilon$ equations:

$$S_k = \rho LAD C_{D_t} (\beta_p |U|^3 - \beta_d |U| k) \quad (10)$$

155 $S_\varepsilon = \rho LAD C_{D_t} \left(C_{4\varepsilon} \beta_p |U|^3 \frac{\varepsilon}{k} - C_{5\varepsilon} \beta_d |U| \varepsilon \right), \quad (11)$

where $C_{4\varepsilon} = C_{5\varepsilon} = 0.9$, $\beta_p = 1.0$ and $\beta_d = 5.03$ are constants of the model (Zaïdi et al., 2013). Note that this type of tree aerodynamic effect modeling is commonly used and evaluated by comparison with experimental results (Buccolieri et al., 2018).

Only the impact of the tree leaves is considered and the impact of the tree trunk and branches on the flow is not modeled. As
 160 the streets are modeled in 2D and an infinite length in the Y direction is assumed (see section 2), the LAI considered here is the equivalent cylindrical LAI_{2D} . Usually, studies consider the tree LAI , which is well defined for a 3D tree. The relation between the tree LAI_{3D} and the equivalent cylindrical LAI_{2D} is given in Appendix B. To check that this effect of trees is



well modeled using this simplified 2D setup, 3D simulations were performed and compared to some of the 2D simulations presented here, showing very good agreement between the 2D and 3D simulations in terms of the influence of the trees on the flow (Appendix B).
165

To get a complete vision of the effect of trees, 45 simulations are performed per canyon, and the impact of three independent tree parameters are studied:

- the 2D equivalent LAI: $LAI_{2D} = 0.5, 1, 2, 3$ and 4 (see Appendix B for the conversion between LAI_{3D} and equivalent LAI_{2D}).
- 170 – The Crown Volume Fraction (CVF) calculated as the tree (2D cylindrical) crown volume divided by the street volume: $CVF = \frac{n\pi r^2}{HW}$. Ratio already used in studies quantifying the effect of trees on street pollution such as Gromke and Blocken (2015), (see Appendix B for the relation between 2D and 3D CVF). Note that apart from the Appendix B, all CVF mentioned in the figures and in the text refer to the 2D CVF . Three CVF ranges are simulated: $CVF \approx 5, 10$ and 25 %.
- 175 – The tree to street height ratio calculated as the crown middle height divided by the building height (h/H). Three h/H ranges are simulated: $h/H \approx 1/3, 1/2$ and $2/3$.

Note that these three tree characteristics are normalized by street characteristics, and similar values are chosen in the three canyons to be able to compare them and to quantify if the street aspect ratio influences the tree effect (i.e. if there is an interaction between tree and building effects). However, the normalized tree characteristics are not exactly equal in the three canyons due to the street sizes and the 0.5 m mesh cells that limit the possible range of tree sizes. The detailed list of the tree characteristics and the different values used for the tree parameters are presented in Table A4.
180

2.2.2 Calculation of vertical and horizontal transfers in Code_Saturne for comparison to MUNICH

To evaluate the vertical transfer between the street and the background zone in Code_Saturne, a passive tracer is emitted in each mesh cell of the street with an arbitrary stationary emission rate $e = 1000 \mu\text{g}\cdot\text{s}^{-1}$ for a street canyon of length $L = 1$ m. For the tracer to be dispersed only by vertical transfers and not by horizontal winds within the street, simulations are performed with wind perpendicular to the street at the inlet ($\varphi = 90^\circ$). The initial street and background tracer concentrations are zero. At the end of the simulation (stationary state reached), the tracer concentration is averaged in the street and in the background zone (noted C_{street} and C_{bg} in $\mu\text{g}\cdot\text{m}^{-3}$) to reproduce MUNICH homogeneous street assumption. The background zone corresponds to the area of the same volume as the street but located just above the street (see Maison et al. (2022) for more details). In absence of other processes (horizontal transport, deposition and chemical reactions), the tracer mass balance in the street yields to:
190



$$Q_{vert} = e \quad (12)$$

$$\implies q_{vert} WL \frac{C_{street} - C_{bg}}{H} = e \quad (13)$$

$$\implies q_{vert} = \frac{eH}{WL (C_{street} - C_{bg})}, \quad (14)$$

195 where Q_{vert} is the vertical flux of pollutant at the roof level for the whole street ($\mu\text{g}\cdot\text{s}^{-1}$), q_{vert} is the vertical transfer coefficient ($\text{m}^2\cdot\text{s}^{-1}$) and WL is the exchange surface (m^2). Thus, the vertical transfer coefficient can be compared between both models: in Code_Saturne, q_{vert} is calculated from the emission rate e and the concentration gradient $\left(\frac{C_{street}-C_{bg}}{H}\right)$ following Eq. (14), and in MUNICH it is calculated from Eq. (7).

In MUNICH, the horizontal transfer velocity is equal to the street average horizontal wind speed in the street direction U_{street} , which is calculated by integrating Eq. (1) between $z = z_{0s}$ and $z = H$. In Code_Saturne, U_{street} is estimated from the wind speed in the Y direction averaged over the street mesh cells. In addition to U_{street} , the MUNICH vertical profile of wind speed (Eq. (1)) can also be compared to Code_Saturne by averaging the wind speed in the Y direction over the street width (X axis). As the objective of the study is to parameterize the aerodynamic effect of trees, chemistry and deposition on built and vegetated surfaces are not considered here. Further details on the simulations setup of MUNICH and Code_Saturne and on the comparison of vertical and horizontal transfers in a treeless canyon are presented in Maison et al. (2022).

3 Quantification of the tree crown effect on horizontal and vertical transfers by Code_Saturne simulations

To quantify tree effect on the horizontal wind speed along the street and on the vertical transfer coefficient, Code_Saturne simulations are performed for a large range of tree characteristics (LAI_{2D} , CVF and height ratio h/H), as summarized in Table A4. The tree effect is expressed as a relative deviation between the simulations without and with trees.

210 3.1 Tree effect on horizontal transfer

To quantify and compare the effect of tree crowns in Code_Saturne and MUNICH, and thus to overcome eventual differences between the two models observed in a treeless canyon for the horizontal velocity U_{street} , a relative deviation (RD) of U_{street} , between the simulations with and without tree, is computed as:

$$RD_{U_{street}} = \frac{U_{street} - U_{street_0}}{U_{street_0}}, \quad (15)$$

215 where U_{street_0} stands for the average wind velocity in a treeless street and U_{street} is the average wind velocity in a street with trees, as computed with Code_Saturne. This RD between the Code_Saturne simulations with and without tree is shown in Fig. 2 (a) for WC, (b) for IC and (c) for NC. It shows that $RD_{U_{street}}$ becomes increasingly negative meaning that U_{street} decreases as the LAI_{2D} and the CVF increase. On the opposite, the tree height has either no impact or a small impact compared to LAI_{2D} and CVF . When the effect of tree height is noticeable, an increase of the tree height induces a decrease of RD.



220 In the range of the tree characteristics studied, U_{street} is attenuated from 7.3 to 62.3 %. The tree effect on U_{street} can be compared between the three canyons. The tree effect on U_{street} increases as the canyon is deeper, highlighting the complex interaction between the street dimensions and the tree effect on the velocity. This observation is consistent with the study of Wania et al. (2012).




Figure 2. Relative deviation RD of U_{street} computed from Code_Saturne simulations for different tree LAI_{2D} , CVF and height ratio for WC (a), IC (b) and NC (c). The graphic is divided into three columns corresponding to the three height ratios, and higher CVF correspond to darker colors. $|RD_{U_{street}}|$ values are specified with data labels.



3.2 Tree effect on vertical transfer

225 The relative deviation of the vertical transfer coefficient ($RD_{q_{vert}}$) between simulations with and without tree is introduced to quantify the tree effect on the vertical transfer coefficient. Similarly to the relative deviation of U_{street} (Eq. (15)), the relative deviation $RD_{q_{vert}}$ is expressed as:

$$RD_{q_{vert}} = \frac{q_{vert} - q_{vert0}}{q_{vert0}}, \quad (16)$$

where q_{vert0} stands for the vertical transfer coefficient in a treeless street and q_{vert} is the vertical transfer coefficient in a street
230 with trees. Code_Saturne $RD_{q_{vert}}$ is plotted for different tree parameters LAI_{2D} , CVF and height ratio h/H in Fig. 3 (a) for WC, 3 (b) for IC and 3 (c) for NC.

For WC, $RD_{q_{vert}}$ increases with tree LAI_{2D} , CVF and height ratio (Fig. 3(a)). For IC, $RD_{q_{vert}}$ also increases with CVF
and height ratio but $RD_{q_{vert}}$ tends to slightly decrease as the LAI increase when the LAI is high and the ratio h/H is small
(differences between $LAI = 3$ and 4 for $h/H = 0.36$ and 0.50) (Fig. 3(b)). For NC, a small increase of $RD_{q_{vert}}$ with LAI_{2D}
235 and CVF is observed for $h/H = 0.65$. However, for the two other smaller height ratios the tree effect is very low ($-1.1 \leq RD_{q_{vert}} \leq 0.5$) (Fig. 3(c)).

For vertical transfer, a wind perpendicular to the street ($\varphi = 90^\circ$) is used to focus on the effects on vertical transfers only. The
complex air flow occurring in street canyons for a perpendicular wind leads to heterogeneous tracer concentration in the street.
In general for the three canyons, the presence of the two tree crowns tends to increase the tracer concentration on the leeward
240 side of the street and to decrease it on the windward side (Buccolieri et al., 2009; Gromke and Ruck, 2012). Depending on
street aspect ratio and therefore on the air flow regime (Oke, 1988; Harman et al., 2004) as well as on the tree characteristics,
the tree crown effect on local tracer concentration (leeward versus windward side) is more or less important. For example, for
NC, the flow regime is skimming, and in average, the variation of C_{street} due to the presence of trees compensates between
the two sides of the street explaining why $RD_{q_{vert}}$ is very low for NC.

245 Note that unlike U_{street} (Sect. 3.1), the tree effect on q_{vert} decreases when the canyon gets deeper. For example, the tree effect,
as quantified by $RD_{q_{vert}}$, ranges between -1.0 and -20.3% for WC, -1.0 and -18.7% for IC and 0.5 and -2.7% for NC. The
averaged tree effect is relatively less strong when the canyon is deeper. In other words, regarding vertical transfers the street
effect dominates over the tree effect.

This section demonstrated that the tree effect parameterizations should depend on tree characteristics and also on building
250 characteristics to account for the building-tree interactions. The next section aims to parameterize in MUNICH the tree effect
on U_{street} and q_{vert} observed in Code_Saturne simulations.




Figure 3. Relative deviation RD of q_{vert} computed from Code_Saturne simulations for different tree LAI_{2D} , CVF and height ratio for WC (a), IC (b) and NC (c). The graphic is divided into three columns corresponding to the three height ratios, and higher CVF correspond to darker colors. $|RD_{q_{vert}}|$ values are specified with data labels.

4 Parameterization of the aerodynamic effect of tree crowns in MUNICH

4.1 Model description

The MUNICH parameterizations of horizontal and vertical transfers detailed in Maison et al. (2022) are modified to take into account the tree effects. These parameterizations are based on Wang (2012, 2014) equations, which were originally developed for homogeneous vegetated cover. To remain consistent with this hypothesis, the parameterization will depend on the homo-



geneous Leaf Area Index in the street noted LAI_{street} . As for the conversion from LAI_{3D} to LAI_{2D} , LAI_{street} is estimated from LAI_{2D} conserving the leaf surface. LAI_{street} is calculated by spreading the tree crown cylindrical LAI_{2D} over the whole street width:

$$260 \quad LAI_{street} \times S_{street} = LAI_{2D} \times S_{2D} \quad (17)$$

$$\Rightarrow LAI_{street} = LAI_{2D} \times \frac{S_{2D}}{S_{street}} = \frac{2rLn LAI_{2D}}{WL} = \frac{2rn LAI_{2D}}{W}, \quad (18)$$

where S_{street} is the soil projected area of the street homogeneous tree crown and S_{2D} is the soil projected area of the Code_Saturne 2D cylindrical tree crown (in m^2). Note that regardless of the tree crown geometry, LAI_{street} is always equal to the street total leaf surface divided by the street ground area (WL).

265 To account for the tree effect on the mixing length, the characteristic length of the trees (l_{ct} in m) is added into the equation describing the mixing length l_m :

$$\frac{1}{l_m} = \frac{1}{\kappa H} + \frac{1}{l_{cb}} + \frac{1}{l_{ct} f_{b \times t}}, \quad (19)$$

$$\text{with } l_{ct} = \frac{E_t H}{C_{D_t} \frac{1}{2} LAI_{street}}, \quad (20)$$

270 where $\frac{1}{2} LAI_{street}$ corresponds to the leaf frontal area density assuming a random leaf orientation distribution, E_t is a proportionality constant taken equal to 0.054 for vegetated cover as suggested by Wang (2014), and $f_{b \times t}$ is a function parameterized based on Code_Saturne simulations in Sect. 4.2 and representing the interaction between buildings, trees and the tree crown height.

The dimensionless factor s_H expressing the effects of the tree canopy on the mixing length l_m is calculated using its definition
 275 in Eq. (6) and the expression of l_m in Eq. (19):

$$l_m = \kappa H s_H \text{ at roof level} \quad (21)$$

$$\text{with } s_H = \begin{cases} \frac{l_{cb}}{l_{cb} + \kappa H} & \text{without tree} \\ \frac{l_{cb} l_{ct} f_{b \times t}}{\kappa H (l_{cb} + l_{ct} f_{b \times t}) + l_{cb} l_{ct} f_{b \times t}} & \text{with trees.} \end{cases} \quad (22)$$

The simulations with and without trees have to be distinguished to avoid any convergence issue since l_{ct} tends to $+\infty$ when LAI_{street} tends to 0. This s_H factor includes now rough soil, building and tree effects on mixing length and is then used in the
 280 calculation of the attenuation coefficient α (Eq. (2)). In the numerator, the expressions for buildings and trees are added:

$$\alpha = \frac{C_B a_r + C_{D_t} C_u \frac{1}{2} LAI_{street}}{\kappa s_H}, \quad (23)$$

where C_{D_t} is the tree drag coefficient (dimensionless) taken equal to the one used in Code_Saturne ($C_{D_t} = 0.2$). Wang (2014) presents C_u as a dimensionless coefficient homogeneous on the vertical axis, but that can depend on canopy features. This coefficient has to be determined based on observation data and will be parameterized in Sect. 4.2.



285 4.2 Parameter determination based on Code_Saturne simulations

Two parameters introduced in l_m and α equations (Eq. (19) and (23)) have to be determined based on Code_Saturne: $RD_{q_{vert}}$ and $RD_{U_{street}}$. The function $f_{b\times t}$ depends on a_r to account for the interaction between trees and buildings observed in Fig. 3 and also on h_{max} to account for the effect of the tree height. Note that since tree crowns are assumed to be homogeneous within the canopy in the original Wang formulation, the tree crown height h_{max} is not taken into account (see Eq. (20)), and therefore
290 it needs to be included in the function $f_{b\times t}$. The $f_{b\times t}$ expression is determined by maximizing the fit between Code_Saturne $RD_{q_{vert}}$ and MUNICH $RD_{q_{vert}}$:

$$f_{b\times t} = \frac{a_0 + a_1 \exp(a_2 a_r)}{(h_{max}/H)^2} \quad \text{with } a_0 = 3.26, a_1 = 0.0256 \text{ and } a_2 = 6.70. \quad (24)$$

A comparison of Code_Saturne $RD_{q_{vert}}$ and the $RD_{q_{vert}}$ parameterized with Eqs. (7), (19), (20) and (24) is presented for the three canyons in Fig. 4 (a,c,e). Normalized Mean Absolute Error (NMAE) and Bias (NMB) are calculated to compare
295 Code_Saturne and MUNICH results (see Appendix C for the definition of the statistical indicators). Figures 4 (a,c,e) show a good agreement between Code_Saturne $RD_{q_{vert}}$ and the parameterized $RD_{q_{vert}}$ because the $f_{b\times t}$ function was determined to minimize bias. The NMAE can be high for $RD_{q_{vert}}$ (up to 63 % in NC) because the parameterization does not reproduce the slight decrease of $RD_{q_{vert}}$ when LAI_{2D} increases from $LAI_{2D} = 3$ to 4 when $h/H \approx 1/3$ and $1/2$ (see section 3.2, Fig. 3 (b) and (c)). But this is not an issue for NC since the q_{vert} values are low.

300 For U_{street} , the parameter C_u also needs to be determined. A constant value of $C_u = 6.7$ is sufficient to get a good fit between Code_Saturne $RD_{U_{street}}$ and MUNICH $RD_{U_{street}}$. No dependency of C_u on building or tree features is needed, as shown in Fig. 4 (b,d,f), which compares Code_Saturne $RD_{U_{street}}$ and the parameterized $RD_{U_{street}}$. Note that the parameterizations of l_{ct} and $f_{b\times t}$ impact not only the vertical transfers but also the horizontal wind speed, because they are involved in the calculation of the s_H factor and hence of the α coefficient. The good comparisons between Code_Saturne $RD_{U_{street}}$ and
305 MUNICH $RD_{U_{street}}$ also shows that the parameterized s_H reproduces well the horizontal wind speed, and therefore the interactions between trees and building effects and the influence of h_{max} .




Figure 4. Comparison of (a,c,e) RD_{qvert} and (b,d,f) $RD_{Ustreet}$ computed from Code_Saturne simulations and parameterized in MUNICH for different tree LAI_{2D} , CVF and height ratio and for (a,b) WC, (c,d) IC and (e,f) NC.



4.3 Comparison of q_{vert} , U_{street} and wind profiles

Figure 5 (a) and (b) presents respectively a comparison of Code_Saturne and parameterized q_{vert} and U_{street} . The statistical indicators are presented in Table 2. As MUNICH was parameterized to reproduce well the tree effect observed in Code_Saturne, and as the two models agree well in a treeless canyon (Maison et al., 2022), the vertical transfer coefficient and wind speed with trees are close between the two models as expected (Fig. 5 and Table 2). The parameterized vertical transfer coefficients with trees agree well with Code_Saturne ones with normalized mean absolute errors ranging from 2.2 to 4.1 % and bias from -4.1 to 2.8 %. The parameterized average wind speed with trees agree well with Code_Saturne ones with normalized mean absolute errors ranging from 3.4 to 6.8 % and a normalized mean bias from -2.8 to -1.1 %.




Figure 5. Normalized U_{street} and q_{vert} computed from Code_Saturne simulations for different tree LAI_{2D} , CVF and height ratios.

Table 2. Statistical indicators (Normalized Mean Absolute Error (NMAE) and Bias (NMB) in %) for the comparison of Code_Saturne and MUNICH q_{vert} and U_{street} .

Canyon	q_{vert}		U_{street}	
	NMAE	NMB	NMAE	NMB
WC	2.2	-2.1	3.4	-1.1
IC	3.5	2.8	4.4	-1.7
NC	4.1	-4.1	6.8	-2.8

315 For each street canyon, Fig. 6 compares the Code_Saturne and parameterized vertical wind profiles for fixed CVF and h/H ratio but for five different LAI_{2D} . It shows that an increase of the LAI_{2D} induces a decrease of the wind velocity. Concerning the vertical profile shape, trees induce a lower wind velocity on the entire street and not only in the tree crown. They even



slightly impact the velocity just above the street. The maximum of attenuation of the wind is located in the middle of the tree crown and the wind velocity is re-accelerated under the tree crown. In the middle of the tree crown, the parameterized wind speed is close to the one of Code_Saturne. In the lower part and under the tree crown, the parameterized wind speed is underestimated. In fact, the re-acceleration under the tree crown is complex to consider in parameterized models. Besides, in real streets the trees height is not homogeneous so this re-acceleration under the tree crown might be unrealistic in Code_Saturne simulations. In the parameterized wind profile with trees, the reduction of the wind speed just above the street due to the presence of trees was neglected. So the parameterized wind speed above and in the upper part of the tree crown is overestimated compared to Code_Saturne. Note that Kent et al. (2017) proposed a method to include vegetation effect in aerodynamic roughness parameters and so to account for vegetation in the above urban canopy wind profile of Macdonald et al. (1998). This method considers vegetation at the city scale and is not applicable here, since we work at the street scale. For now, as MUNICH assumes an homogeneous street canyon, the profile is averaged to compute U_{street} and the shape of the profile is not used.




Figure 6. Comparison of Code_Saturne (solid lines) and parameterized (dotted lines) vertical wind profiles. The profiles in a treeless canyon are in black and darker colors correspond to increasing LAI_{2D} . For the three canyons, $h/H \approx 0.5$ and $CVF \approx 25\%$.

5 Conclusions

Although the discretizations and the physical modeling used in MUNICH and Code_Saturne are fundamentally different, the setup of the CFD simulations was adapted to compare the two models through the average horizontal wind speed along the street and the calculation of the vertical transfer coefficient at the roof level by averaging a passive tracer concentration in the street and the background (above the streets). To build a parameterization suited to the street network MUNICH, the CFD simulations were simplified with several hypothesis such as an infinite street canyon with 2D CFD simulations and homogeneous emissions over the street. Furthermore, quantities of interest were averaged over the street in the CFD simulations.

The aerodynamic effect of trees in street canyons is quantified with Code_Saturne simulations. Simulations were performed



with 2 rows of trees and a large range of tree Leaf Area Index, crown radius and heights. In the range of the simulations performed, the average horizontal velocity in the street is reduced by -7.3 to -62.3 % and the vertical transfer coefficient by +0.5 to -20.3 %. This highlights the necessity of adding the tree aerodynamic effects into street models, such as MUNICH.

340 Maison et al. (2022) proposed a parameterization of horizontal and vertical transfers in a treeless canyon and the present study adds the tree effect to the building effect in this parameterization. The differences of wind speed and vertical transfer coefficient in a treeless canyon are very low between the two models (Maison et al., 2022). However, to overcome any model-specific difference and to build a parameterization for MUNICH of the tree effect on the aerodynamic parameters, relative deviation ratios between cases without and with trees are defined.

345 First, the Maison et al. (2022) vertical transfer coefficient parameterization is modified by adding a term representing the tree effect and the interaction between the trees and the buildings in the mixing-length expression. This term is function of two dimensionless parameters to characterize the trees: street Leaf Area Index (LAI_{street}) and tree to street height ratio (h_{max}/H) as well as building height (H) and street width (W). A parameterization of the tree mixing length was defined from Code_Saturne simulations to obtain a tree effect as close as possible in the two models. The parameterized vertical transfer coefficients with

350 trees agree well with those of Code_Saturne with normalized mean absolute error ranging from 2.2 to 4.1 % and bias from -4.1 to 2.8 %.

Second, this new mixing length expression is also used to compute the vertical wind profile and average wind speed along the street. Only one constant is fixed in the wind attenuation coefficient to maximize the fit between Code_Saturne and MUNICH tree effect on average wind speed. The comparison of the average wind speed gives a normalized mean absolute error ranging

355 from 3.4 to 6.8 % and a normalized mean bias from -2.8 to -1.1 %.

MUNICH now includes a relatively simple parameterization of tree effect on both horizontal and vertical aerodynamic processes based on commonly used tree parameters, that can be easily computed from urban databases and can reproduce the main effects obtained in much more detailed and costly CFD simulations. This parameterization can also be used in urban climate models to compute water and heat transfers in tree-lined streets. The perspectives of this study are to quantify the effect of street

360 trees on air quality from the street level to the scale of the city of Paris and to compare biogenic and anthropic contributions to air pollution over an entire city with the fast running code MUNICH.

Code availability. The last version of MUNICH source code is available online at <https://doi.org/10.5281/zenodo.4168984> and <https://github.com/cerea-lab/munich> (last access on 5 January 2022).

For the three 2D canyons considered in the present study, the mesh, the source code and the XML setup file allowing reproduction of

365 the CFD results using Code_Saturne version 6.0 are available online at https://gitlab.enpc.fr/alice.maison/tree_parametrization and at <http://dx.doi.org/10.17632/fzfrjsz3mv.2> under the GNU GPL2.0 licence (last access on 3 December 2021).



Appendix A: Lists of abbreviations, variables parameters and tree dimensions

Table A1. List of abbreviations

Acronym	Definition
WC	Wide canyon
IC	Intermediate canyon
NC	Narrow canyon
CTM	Chemistry-transport model
CFD	Computational fluid dynamics
RD	Relative deviation

Table A2. List of parameters

Symbol	Definition	Value	Unit
κ	Von Kàrmàn constant	0.42	-
PBLH	Planetary Boundary Layer Height	1000	m
z_{0s}	Code_Saturne inside street walls roughness length	0.10	m
u_*	Friction velocity	0.727	m.s ⁻¹
E	Parameter in modified Wang (2014) parameterization	0.5	-
n	Number of tree rows	2	-
C_{Dt}	Tree crown drag coefficient (Katul and Albertson, 1998)	0.2	-



Table A3. List of variables

Group of variables	Symbol	Definition	Unit
Street characteristics	H	Buildings height	m
	W	Street width	m
	L	Street length	m
	a_r	Street aspect ratio	-
	l_{cb}	Characteristic length in the street	m
Tree characteristics	LAD	Tree Leaf Area Density	$\text{m}_{\text{leaf}}^2 \cdot \text{m}_{\text{tree crown}}^{-3}$
	LAI	Tree Leaf Area Index	$\text{m}_{\text{leaf}}^2 \cdot \text{m}_{\text{soil}}^{-2}$
	LAI_{3D}	3D Leaf Area Index of the spherical tree crown	$\text{m}_{\text{leaf}}^2 \cdot \text{m}_{\text{soil}}^{-2}$
	LAI_{2D}	2D equivalent Leaf Area Index of the cylindrical tree crown	$\text{m}_{\text{leaf}}^2 \cdot \text{m}_{\text{soil}}^{-2}$
	LAI_{street}	Leaf Area Index of the homogeneous street tree crown	$\text{m}_{\text{leaf}}^2 \cdot \text{m}_{\text{soil}}^{-2}$
	S_{3D}	Soil projected area of the 3D spherical tree crown	m^2
	S_{2D}	Soil projected area of the 2D cylindrical tree crown	m^2
	S_{street}	Soil projected area of the street homogeneous tree crown	m^2
	r	Tree radius	m
	CVF	Crown Volume Fraction	- or %
	h	Middle crown height	m
	h_{max}	Maximum tree crown height	m
	h_{min}	Minimum tree crown height	m
	δ	Spacing between two trees within a row	m
l_{ct}	Tree characteristic length	m	
Horizontal wind speed	U_{street}	Average street horizontal wind speed	$\text{m} \cdot \text{s}^{-1}$
	U	Norm of the horizontal wind speed	$\text{m} \cdot \text{s}^{-1}$
	U_X	Horizontal wind speed in the X direction	$\text{m} \cdot \text{s}^{-1}$
	U_Y	Horizontal wind speed in the Y direction	$\text{m} \cdot \text{s}^{-1}$
	U_H	Average horizontal wind speed at the roof level	$\text{m} \cdot \text{s}^{-1}$
	φ	Angle of the wind direction	rad or °
	α	Wind attenuation coefficient	-
	s_H	Characteristic length factor	-
	C_u	Empiric coefficient in α equation	-
C_B	Function of a_r and φ	-	
Vertical transfer	Q_{vert}	Vertical flux of pollutant	$\mu\text{g} \cdot \text{s}^{-1}$
	q_{vert}	Vertical transfer coefficient	$\text{m}^2 \cdot \text{s}^{-1}$
	σ_W	Standard deviation of the vertical wind speed at $z = H$ (Soulhac et al., 2011)	$\text{m} \cdot \text{s}^{-1}$
	l_m	Mixing length in the street	m
	e	Passive tracer emission rate for the street	$\mu\text{g} \cdot \text{s}^{-1}$
	C_{street}	Street concentration	$\mu\text{g} \cdot \text{m}^{-3}$
C_{bg}	Background concentration	$\mu\text{g} \cdot \text{m}^{-3}$	



Table A4. List of tree dimensions simulated with Code_Saturne (r : radius, h : middle crown height, h_{min} : crown bottom height h_{max} : crown top height) and calculated tree parameters (CVF : crown volume fraction and h/H : height ratio).

Canyon	r (m)	h (m)	h_{min} (m)	h_{max} (m)	CVF (%)	h/H (-)
WC	1.5	3.5	2.0	5.0	6.0	0.41
	1.5	4.5	3.0	6.0	6.0	0.53
	1.5	5.5	4.0	7.0	6.0	0.65
	2.0	3.5	1.5	5.5	10.8	0.41
	2.0	4.5	2.5	6.5	10.8	0.53
	2.0	5.5	3.5	7.5	10.8	0.65
	3.0	3.5	0.5	6.5	24.2	0.41
	3.0	4.5	1.5	7.5	24.2	0.53
	3.0	5.5	2.5	8.5	24.2	0.65
IC	2.0	5.0	3.0	7.0	6.5	0.36
	2.0	7.0	5.0	9.0	6.5	0.50
	2.0	9.0	7.0	11.0	6.5	0.64
	2.5	5.0	2.5	7.5	10.2	0.36
	2.5	7.0	4.5	9.5	10.2	0.50
	2.5	9.0	6.5	11.5	10.2	0.64
	4.0	5.0	1.0	9.0	26.1	0.36
	4.0	7.0	3.0	10.0	26.1	0.50
	4.0	9.0	5.0	13.0	26.1	0.64
NC	2.5	9.0	6.5	11.5	5.2	0.33
	2.5	14.0	11.5	16.5	5.2	0.51
	2.5	18.0	15.5	20.5	5.2	0.65
	3.5	9.0	5.5	12.5	10.2	0.33
	3.5	14.0	10.5	17.5	10.2	0.51
	3.5	18.0	14.5	21.5	10.2	0.65
	5.5	9.0	3.5	14.5	25.1	0.33
	5.5	14.0	8.5	19.5	25.1	0.51
	5.5	18.0	12.5	23.5	25.1	0.65

Each line of the Table A4 corresponds to the parameters of a simulation. For each case, simulations with different LAI_{2D} were performed: 0.5, 1, 2, 3 and 4.



370 Appendix B: Comparison of 2D and 3D Code_Saturne simulations with trees

To save computing time, Code_Saturne simulations were performed in a periodic 2D canyon of length $L = 1$ m, and the street was therefore considered infinite due to this periodicity. In this case, the tree crown is represented as an infinite cylinder of radius r . To take into account the fact that in reality most of the tree crowns are spherical and are spaced from each other, an equivalent cylindrical LAI_{2D} is calculated in the 2D simulations. For the tree effect to be similar, the surface of leaves is kept
 375 constant between the 2D and 3D simulations:

$$LAI_{3D} \times S_{3D} = LAI_{2D} \times S_{2D} \quad (B1)$$

$$LAI_{2D} = LAI_{3D} \times \frac{S_{3D}}{S_{2D}} = LAI_{3D} \times \frac{\pi r^2}{2r(2r + \delta)}, \quad (B2)$$

where δ is the spacing between two tree crowns within the row and $2r + \delta$ is the length of the street section for one spherical tree (m). S_{3D} and S_{2D} are respectively the soil projected area of the 3D and 2D tree crowns (m^2). In addition to the leaf
 380 surface and tree height, the tree radius is also conserved between 2D and 3D simulations, however the value of the Crown Volume Fraction (CVF) varies:

$$CVF_{2D} = \frac{V_{2D}}{V_{street}} = \frac{n\pi r^2 L}{HWL} \quad \text{and} \quad CVF_{3D} = \frac{V_{3D}}{V_{street}} = \frac{n\frac{4}{3}\pi r^3}{HW(2r + \delta)} \quad (B3)$$

V_{street} is the street volume (m^3), V_{3D} and V_{2D} are respectively the 3D and 2D tree crown volumes (m^3). The relation between 2D and 3D CVF is therefore:

$$385 \quad CVF_{2D} = \frac{3}{4} \times \frac{2r + \delta}{r} \times CVF_{3D} \quad (B4)$$

2D and 3D simulations with different LAI are performed to check the validity of the 2D infinite street canyon hypothesis. 3D simulations are realized with a periodic street of 10 m-length and cell meshes of $0.5 \times 0.5 \times 0.5$ m in the X , Y and Z directions. In both 2D and 3D simulations, C_{street} and C_{bg} are computed with a normal wind to the street ($\varphi = 90^\circ$) and U_{street} with $\varphi = 45^\circ$. The results are presented in Fig. B1 (a-d) for WC, (e-h) for IC and (i-l) for NC.

390 For each canyon, the cases without tree ($LAI = 0$) are equivalent: C_{street} , C_{bg} , q_{vert} and U_{street} are similar between the 2D and 3D cases with a maximum relative difference of 0.1 % for U_{street} . For the cases with trees, concerning C_{street} , C_{bg} and q_{vert} , the differences between the 2D and 3D cases are relatively low over the range of simulations performed with a maximum of 3.3 % for C_{street} and C_{bg} in WC. For U_{street} , slightly larger differences are observed (up to -5.4 % in WC for $LAI = 4(2D)/12.7(3D)$). These differences can be explained by the distribution of the turbulence around the crown. For each
 395 variable and canyon, the relative deviation between the 2D and 3D cases increases with the tree LAI . Given the low differences observed between the 2D and 3D simulations, the hypothesis of a 2D canyon with a cylindrical tree crown associated with an equivalent LAI_{2D} to represent a spherical tree crown spaced of δ m in a street of length L is reasonable.




Figure B1. Comparison of C_{street} (a,e,i), C_{bg} (b,f,j), q_{vert} (c,g,k) and U_{street} (d,h,l) for 2D and 3D simulations in the three canyons without and with trees of $LAI_{2D} = 1$ and 4. $CVF_{2D} \approx 10\%$ and $h/H \approx 0.5$ are fixed. The percentages written in the middle of the histogram bars correspond to the RD between the 2D and 3D cases.



Appendix C: Definition of the statistical indicators

Code_Saturne and MUNICH simulations are noted cs_i and m_i , respectively. In this section, n is the total number of simulations
400 which is equal to 45 per street canyon.

- Normalized Mean Absolute Error (%):

$$NMAE = 100 \times \frac{\sum_{i=1}^n |m_i - cs_i|}{\left| \sum_{i=1}^n cs_i \right|} \quad (C1)$$

- Normalized Mean Bias (%):

$$NMB = 100 \times \frac{\sum_{i=1}^n (m_i - cs_i)}{\sum_{i=1}^n cs_i} \quad (C2)$$

405 *Author contributions.* KS, CF, BC and AM were responsible for conceptualization. AM developed the MUNICH model code and performed the Code_Saturne simulations. AM, KS, CF and BC performed the formal analysis. AM conducted the visualization. AM, KS and CF were responsible for writing the original draft and BC and AT reviewed it. CF, BC and YW provided support in Code_Saturne computing resources. KS and AT were responsible for funding acquisition.

Competing interests. The authors declare that they have no conflict of interest.

410 *Acknowledgements.* This work was partially funded by the sTREEt ANR project (ANR-19-CE22-0012). The authors thank Youngseob KIM and Lya LUGON for their support in the development of the MUNICH model and Martin FERRAND for his support in the understanding of the Code_Saturne model.



References

- Akimoto, H.: Global Air Quality and Pollution, *Science*, 302, 1716–1719, <https://doi.org/10.1126/science.1092666>, 2003.
- 415 Angel, S., Parent, J., Civco, D. L., Blei, A., and Potere, D.: The dimensions of global urban expansion: Estimates and projections for all countries, 2000–2050, *Progress in Planning*, 75, 53–107, <https://doi.org/10.1016/j.progress.2011.04.001>, 2011.
- Archambeau, F., Méchitoua, N., and Sakiz, M.: Code Saturne: A finite volume code for the computation of turbulent incompressible flows—Industrial applications, *International Journal on Finite Volumes*, 1, 1–62, 2004.
- Armson, D., Stringer, P., and Ennos, A.: The effect of street trees and amenity grass on urban surface water runoff in Manchester, UK, *Urban For. Urban Gree.*, 12, 282–286, <https://doi.org/10.1016/j.ufug.2013.04.001>, 2013.
- 420 Beckett, K., Freer-Smith, P., and Taylor, G.: Urban woodlands: their role in reducing the effects of particulate pollution, *Environ. Pollut.*, 99, 347–360, [https://doi.org/10.1016/S0269-7491\(98\)00016-5](https://doi.org/10.1016/S0269-7491(98)00016-5), 1998.
- Berland, A., Shiflett, S. A., Shuster, W. D., Garmestani, A. S., Goddard, H. C., Herrmann, D. L., and Hopton, M. E.: The role of trees in urban stormwater management, *Landscape Urban Plan.*, 162, 167–177, <https://doi.org/10.1016/j.landurbplan.2017.02.017>, 2017.
- 425 Bertram, C. and Rehdanz, K.: The role of urban green space for human well-being, *Ecological Economics*, 120, 139–152, <https://doi.org/10.1016/j.ecolecon.2015.10.013>, 2015.
- Bowler, D. E., Buyung-Ali, L., Knight, T. M., and Pullin, A. S.: Urban greening to cool towns and cities: A systematic review of the empirical evidence, *Landscape Urban Plan.*, 97, 147–155, <https://doi.org/10.1016/j.landurbplan.2010.05.006>, 2010.
- Bozonnet, E., Musy, M., Calmet, I., and Rodriguez, F.: Modeling methods to assess urban fluxes and heat island mitigation measures from street to city scale, *International Journal of Low-Carbon Technologies*, 10, 62–77, <https://doi.org/10.1093/ijlct/ctt049>, 2015.
- 430 Buccolieri, R., Gromke, C., Di Sabatino, S., and Ruck, B.: Aerodynamic effects of trees on pollutant concentration in street canyons, *Science of The Total Environment*, 407, 5247–5256, <https://doi.org/10.1016/j.scitotenv.2009.06.016>, 2009.
- Buccolieri, R., Salim, M., Leo, L. S., Di Sabatino, S., Chan, A., Ielpo, P., de Gennaro, G., and Gromke, C.: Analysis of local scale tree–atmosphere interaction on pollutant concentration in idealized street canyons and application to a real urban junction, *Atmos. Environ.*, 45, 1702–1713, <https://doi.org/10.1016/j.atmosenv.2010.12.058>, 2011.
- 435 Buccolieri, R., Santiago, J.-L., Rivas, E., and Sanchez, B.: Review on urban tree modelling in CFD simulations: Aerodynamic, deposition and thermal effects, *Urban For. Urban Gree.*, 31, 212–220, <https://doi.org/10.1016/j.ufug.2018.03.003>, 2018.
- Cai, X.-M., Barlow, J., and Belcher, S.: Dispersion and transfer of passive scalars in and above street canyons—Large-eddy simulations, *Atmos. Environ.*, 42, 5885–5895, <https://doi.org/10.1016/j.atmosenv.2008.03.040>, 2008.
- 440 Faiz, A.: Automotive emissions in developing countries—relative implications for global warming, acidification and urban air quality, *Transportation Research Part A: Policy and Practice*, 27, 167–186, [https://doi.org/10.1016/0965-8564\(93\)90057-R](https://doi.org/10.1016/0965-8564(93)90057-R), 1993.
- Gillner, S., Vogt, J., Tharang, A., Dettmann, S., and Roloff, A.: Role of street trees in mitigating effects of heat and drought at highly sealed urban sites, *Landscape Urban Plan.*, 143, 33–42, <https://doi.org/10.1016/j.landurbplan.2015.06.005>, 2015.
- Gromke, C. and Blocken, B.: Influence of avenue-trees on air quality at the urban neighborhood scale. Part II: Traffic pollutant concentrations at pedestrian level, *Environ. Pollut.*, 196, 176–184, <https://doi.org/10.1016/j.envpol.2014.10.015>, 2015.
- 445 Gromke, C. and Ruck, B.: Influence of trees on the dispersion of pollutants in an urban street canyon—Experimental investigation of the flow and concentration field., *Atmos. Environ.*, 41, 3287–3302, <https://doi.org/10.1016/j.atmosenv.2006.12.043>, 2007.
- Gromke, C. and Ruck, B.: On the Impact of Trees on Dispersion Processes of Traffic Emissions in Street Canyons, *Bound.-Lay. Meteorol.*, 131, 19–34, <https://doi.org/10.1007/s10546-008-9301-2>, 2009.



- 450 Gromke, C. and Ruck, B.: Pollutant Concentrations in Street Canyons of Different Aspect Ratio with Avenues of Trees for Various Wind Directions, *Bound.-Lay. Meteorol.*, 144, 41–64, <https://doi.org/10.1007/s10546-012-9703-z>, 2012.
- Gunawardena, K., Wells, M., and Kershaw, T.: Utilising green and bluespace to mitigate urban heat island intensity, *Science of The Total Environment*, 584–585, 1040–1055, <https://doi.org/10.1016/j.scitotenv.2017.01.158>, 2017.
- Harman, I. N., Barlow, J. F., and Belcher, S. E.: Scalar fluxes from urban street canyons. Part II Model, *Bound.-Lay. Meteorol.*, 113, 387–409, <https://doi.org/10.1007/s10546-004-6205-7>, 2004.
- 455 Hebbert, M. and Jankovic, V.: Cities and Climate Change: The Precedents and Why They Matter, *Urban Studies*, 50, 1332–1347, <https://doi.org/10.1177/0042098013480970>, 2013.
- Huang, Y.-D., Hou, R.-W., Liu, Z.-Y., Song, Y., Cui, P.-Y., and Kim, C.-N.: Effects of Wind Direction on the Airflow and Pollutant Dispersion inside a Long Street Canyon, *Aerosol and Air Quality Research*, 19, 1152–1171, <https://doi.org/10.4209/aaqr.2018.09.0344>, 2019.
- 460 Hwang, H.-J., Yook, S.-J., and Ahn, K.-H.: Experimental investigation of submicron and ultrafine soot particle removal by tree leaves, *Atmospheric Environment*, 45, 6987–6994, <https://doi.org/10.1016/j.atmosenv.2011.09.019>, 2011.
- IPCC: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Report, Intergovernmental Panel on Climate Change, United Nations, [Masson-Delmotte V., Zhai P., Pirani A., Connors S.L., Péan C., Berger S., Caud N., Chen Y., Goldfarb L., Gomis M.I., Huang M., Leitzell K., Lonnoy E., Matthews
- 465 J.B.R., Maycock T.K., Waterfield T., Yelekçi O., Yu R., and Zhou B. (eds.)]. Cambridge University Press. In Press., 2021.
- Janhäll, S.: Review on urban vegetation and particle air pollution – Deposition and dispersion, *Atmos. Environ.*, 105, 130–137, <https://doi.org/10.1016/j.atmosenv.2015.01.052>, 2015.
- Jayasooriya, V., Ng, A., Muthukumaran, S., and Perera, B.: Green infrastructure practices for improvement of urban air quality, *Urban For. Urban Gree.*, 21, 34–47, <https://doi.org/10.1016/j.ufug.2016.11.007>, 2017.
- 470 Jeanjean, A., Buccolierib, R., Eddy, J., Monks, P., and Leigh, R.: Air quality affected by trees in real street canyons: The case of Marylebone neighbourhood in central London., *Urban For. Urban Gree.*, 22, 41–43, <https://doi.org/10.1016/j.ufug.2017.01.009>, 2017.
- Katul, G. and Albertson, J.: An Investigation of Higher-Order Closure Models for a Forested Canopy, *Bound.-Lay. Meteorol.*, 89, 47–74, <https://doi.org/10.1023/A:1001509106381>, 1998.
- Katul, G. G., Mahrt, L., Poggi, D., and Sanz, C.: ONE- and TWO-Equation Models for Canopy Turbulence, *Bound.-Lay. Meteorol.*, 113, <https://doi.org/10.1023/B:BOUN.0000037333.48760.e5>, 2004.
- 475 Kent, C. W., Grimmond, S., and Gatey, D.: Aerodynamic roughness parameters in cities: Inclusion of vegetation, *Journal of Wind Engineering and Industrial Aerodynamics*, 169, 168–176, <https://doi.org/10.1016/j.jweia.2017.07.016>, 2017.
- Kim, Y., Wu, Y., Seigneur, C., and Roustan, Y.: Multi-scale modeling of urban air pollution: development and application of a Street-in-Grid model (v1.0) by coupling MUNICH (v1.0) and Polair3D (v1.8.1), *Geosci. Model Dev.*, 11, 611–629, [https://doi.org/10.5194/gmd-11-611-](https://doi.org/10.5194/gmd-11-611-2018)
- 480 2018, 2018.
- Klemm, W., Heusinkveld, B. G., Lenzholzer, S., and van Hove, B.: Street greenery and its physical and psychological impact on thermal comfort, *Landscape Urban Plan.*, 138, 87–98, <https://doi.org/10.1016/j.landurbplan.2015.02.009>, 2015.
- Krekel, C., Kolbe, J., and Wüstemann, H.: The Greener, The Happier? The Effects of Urban Green and Abandoned Areas on Residential Well-Being, *The German Socio-Economic Panel study at DIW Berlin*, 728, 65, 2015.
- 485 Leopold, L. B.: Hydrology for urban land planning - A guidebook on the hydrologic effects of urban land use, *US. Geological Survey*, 554, 18, 1968.



- Li, X., Liu, C., Leung, D., and Lam, K.: Recent progress in CFD modelling of wind field and pollutant transport in street canyons, *Atmos. Environ.*, 40, 5640–5658, <https://doi.org/10.1016/j.atmosenv.2006.04.055>, 2006.
- Livesley, S. J., McPherson, E. G., and Calfapietra, C.: The Urban Forest and Ecosystem Services: Impacts on Urban Water, Heat, and Pollution Cycles at the Tree, Street, and City Scale, *Journal of Environmental Quality*, 45, 119–124, <https://doi.org/10.2134/jeq2015.11.0567>, 2016.
- Lobaccaro, G. and Acero, J.: Comparative analysis of green actions to improve outdoor thermal comfort inside typical urban street canyons, *Urban Climate*, 14, 251–267, <https://doi.org/10.1016/j.uclim.2015.10.002>, 2015.
- Lugon, L., Sartelet, K., Kim, Y., Vigneron, J., and Chrétien, O.: Street-in-Grid modeling of gas-phase pollutants in Paris city, *Atmos. Chem. Phys. Discuss.*, accepted for publication in ACP, <https://doi.org/10.5194/acp-2019-1087>, 2019.
- Lugon, L., Sartelet, K., Kim, Y., Vigneron, J., and Chrétien, O.: Simulation of primary and secondary particles in the streets of Paris using MUNICH, *Faraday Discuss.*, <https://doi.org/10.1039/D0FD00092B>, 2021.
- Macdonald, R., Griffiths, R., and Hall, D.: An improved method for the estimation of surface roughness of obstacle arrays, *Atmos. Environ.*, 32, 1857–1864, [https://doi.org/10.1016/S1352-2310\(97\)00403-2](https://doi.org/10.1016/S1352-2310(97)00403-2), 1998.
- Maison, A., Flageul, C., Carissimo, B., Tuzet, A., and Sartelet, K.: Parametrization of Horizontal and Vertical Transfers for the Street-Network Model MUNICH Using the CFD Model Code_Saturne, *Atmosphere*, 13, 527, <https://doi.org/10.3390/atmos13040527>, 2022.
- Municipality of Paris: Paris Data : Les arbres, <https://opendata.paris.fr/explore/dataset/les-arbres/>, accessed: 2021-12-17, 2021.
- Nowak, D., McHale, P., Ibarra, M., Crane, D., Stevens, J., and Luley, C.: Modeling the Effects of Urban Vegetation on Air Pollution, *Air Pollution Modeling and Its Application XII*, pp. 399–407, https://doi.org/10.1007/978-1-4757-9128-0_41, 1998.
- Nowak, D. J. and Crane, D. E.: Carbon storage and sequestration by urban trees in the USA, *Environ. Pollut.*, 116, 381–389, [https://doi.org/10.1016/S0269-7491\(01\)00214-7](https://doi.org/10.1016/S0269-7491(01)00214-7), 2002.
- Nowak, D. J., Crane, D. E., and Stevens, J. C.: Air pollution removal by urban trees and shrubs in the United States, *Urban For. Urban Gree.*, 4, 115–123, <https://doi.org/10.1016/j.ufug.2006.01.007>, 2006.
- Oke, T.: Street design and urban canopy layer climate, *Energy and Buildings*, 11, 103–113, [https://doi.org/10.1016/0378-7788\(88\)90026-6](https://doi.org/10.1016/0378-7788(88)90026-6), 1988.
- Oke, T. R.: The energetic basis of the urban heat island, *Quarterly Journal of the Royal Meteorological Society*, 108, 1–24, <https://doi.org/10.1002/qj.49710845502>, 1982.
- Ozdemir, H.: Mitigation impact of roadside trees on fine particle pollution, *Science of The Total Environment*, 659, 1176–1185, <https://doi.org/10.1016/j.scitotenv.2018.12.262>, 2019.
- Pascal, M., Corso, M., Chanel, O., Declercq, C., Badaloni, C., Cesaroni, G., Henschel, S., Meister, K., Haluza, D., Martin-Olmedo, P., and Medina, S.: Assessing the public health impacts of urban air pollution in 25 European cities: Results of the Aphekom project, *Science of The Total Environment*, 449, 390–400, <https://doi.org/10.1016/j.scitotenv.2013.01.077>, 2013.
- Pigeon, G., Legain, D., Durand, P., and Masson, V.: Anthropogenic heat release in an old European agglomeration (Toulouse, France), *Int. J. Climatol.*, 27, 1969–1981, <https://doi.org/10.1002/joc.1530>, 2007.
- Revelli, R. and Porporato, A.: Ecohydrological model for the quantification of ecosystem services provided by urban street trees, *Urban Ecosystems*, 21, 489–504, <https://doi.org/10.1007/s11252-018-0741-2>, 2018.
- Robine, J., Cheung, S., and Roy, S. L.: Report on excess mortality in Europe during summer 2003, *Tech. rep.*, EU Community Action Programme for Public Health, 2007.
- Santiago, J.-L., Rivas, E., Sanchez, B., Buccolieri, R., and Martin, F.: The Impact of Planting Trees on NO_x Concentrations: The Case of the Plaza de la Cruz Neighborhood in Pamplona (Spain), *Atmosphere*, 8, <https://doi.org/10.3390/atmos8070131>, 2017.



- 525 Sartelet, K., Zhu, S., Moukhtar, S., André, M., André, J., Gros, V., Favez, O., Brasseur, A., and Redaelli, M.: Emission of intermediate, semi and low volatile organic compounds from traffic and their impact on secondary organic aerosol concentrations over Greater Paris, *Atmos. Environ.*, 180, 126–137, 2018.
- Selmi, W., Weber, C., Rivière, E., Blond, N., Mehdi, L., and Nowak, D.: Air pollution removal by trees in public green spaces in Strasbourg city, France, *Urban For. Urban Gree.*, 17, 192–201, <https://doi.org/10.1016/j.ufug.2016.04.010>, 2016.
- 530 Souhac, L., Salizzoni, P., Cierco, F.-X., and Perkins, R.: The model SIRANE for atmospheric urban pollutant dispersion; part I, presentation of the model, *Atmos. Environ.*, 45, 7379–7395, <https://doi.org/10.1016/j.atmosenv.2011.07.008>, 2011.
- Stewart, I.: A systematic review and scientific critique of methodology in modern urban heat island literature, *Int. J. Climatol.*, 31, 200–217, <https://doi.org/10.1002/joc.2141>, 2011.
- Svirejeva-Hopkins, A., Schellnhuber, H., and Pomaz, V.: Urbanised territories as a specific component of the Global Carbon Cycle, *Ecological Modelling*, 173, 295–312, <https://doi.org/10.1016/j.ecolmodel.2003.09.022>, 2004.
- 535 Taha, H., Akbari, H., and Rosenfeld, A.: Heat island and oasis effects of vegetative canopies: Micro-meteorological field-measurements, *Theor. Appl. Climatol.*, 44, 123–138, <https://doi.org/10.1007/BF00867999>, 1991.
- van Dillen, S. M. E., de Vries, S., Groenewegen, P. P., and Spreeuwenberg, P.: Greenspace in urban neighbourhoods and residents' health: adding quality to quantity, *Journal of Epidemiology and Community Health*, 66, e8–e8, <https://doi.org/10.1136/jech.2009.104695>, 2012.
- 540 Vardoulakis, S., Fisher, B. E., Pericleous, K., and Gonzalez-Flesca, N.: Modelling air quality in street canyons: a review, *Atmos. Environ.*, 37, 155–182, [https://doi.org/10.1016/S1352-2310\(02\)00857-9](https://doi.org/10.1016/S1352-2310(02)00857-9), 2003.
- Vos, P. E., Maiheu, B., Vankerkom, J., and Janssen, S.: Improving local air quality in cities: To tree or not to tree?, *Environ. Pollut.*, 183, 113–122, <https://doi.org/10.1016/j.envpol.2012.10.021>, 2013.
- Wang, W.: An Analytical Model for Mean Wind Profiles in Sparse Canopies, *Bound.-Lay. Meteorol.*, 142, 383–399, <https://doi.org/10.1007/s10546-011-9687-0>, 2012.
- 545 Wang, W.: Analytically Modelling Mean Wind and Stress Profiles in Canopies, *Bound.-Lay. Meteorol.*, 151, 239–256, <https://doi.org/10.1007/s10546-013-9899-6>, 2014.
- Wania, A., Bruse, M., Blond, N., and Weber, C.: Analysing the influence of different street vegetation on traffic-induced particle dispersion using microscale simulations., *J. Env. Manag.*, pp. 91–101, <https://doi.org/10.1016/j.jenvman.2011.06.036>, 2012.
- 550 Wei, X., Dupont, E., Gilbert, E., Musson-Genon, L., and Carissimo, B.: Experimental and Numerical Study of Wind and Turbulence in a Near-Field Dispersion Campaign at an Inhomogeneous Site., *Bound.-Lay. Meteorol.*, pp. 475–499, <https://doi.org/10.1007/s10546-016-0148-7>, 2016.
- West, J. J., Cohen, A., Dentener, F., Brunekreef, B., Zhu, T., Armstrong, B., Bell, M. L., Brauer, M., Carmichael, G., Costa, D. L., Dockery, D. W., Kleeman, M., Krzyzanowski, M., Künzli, N., Lioussé, C., Lung, S.-C. C., Martin, R. V., Pöschl, U., Pope, C. A., Roberts, J. M., Russell, A. G., and Wiedinmyer, C.: What We Breathe Impacts Our Health: Improving Understanding of the Link between Air Pollution and Health, *Environ. Sci. Technol.*, 50, 4895–4904, <https://doi.org/10.1021/acs.est.5b03827>, 2016.
- 555 Xue, F. and Li, X.: The impact of roadside trees on traffic released PM 10 in urban street canyon: Aerodynamic and deposition effects, *Sustainable Cities and Society*, 30, 195–204, <https://doi.org/10.1016/j.scs.2017.02.001>, 2017.
- Yuan, C., Ng, E., and Norford, L. K.: Improving air quality in high-density cities by understanding the relationship between air pollutant dispersion and urban morphologies, *Building and Environment*, 71, 245–258, <https://doi.org/10.1016/j.buildenv.2013.10.008>, 2014.
- 560 Zaïdi, H., Dupont, E., Milliez, M., Musson-Genon, L., and Carissimo, B.: Numerical Simulations of the Microscale Heterogeneities of Turbulence Observed on a Complex Site, *Bound.-Lay. Meteorol.*, 147, 237–259, <https://doi.org/10.1007/s10546-012-9783-9>, 2013.

<https://doi.org/10.5194/acp-2022-287>
Preprint. Discussion started: 29 April 2022
© Author(s) 2022. CC BY 4.0 License.



Zhang, Y., Gu, Z., and Yu, C. W.: Impact Factors on Airflow and Pollutant Dispersion in Urban Street Canyons and Comprehensive Simulations: a Review, *Current Pollution Report*, 6, 425–439, <https://doi.org/10.1007/s40726-020-00166-0>, 2020.