

Robust evidence for reversal in the aerosol effective climate forcing trend

Johannes Quaas¹, Hailing Jia¹, Chris Smith^{2,3}, Anna Lea Albright⁴, Wenche Aas⁵, Nicolas Bellouin^{6,7}, Olivier Boucher⁶, Marie Doutriaux-Boucher⁸, Piers M. Forster², Daniel Grosvenor², Stuart Jenkins⁹, Zbigniew Klimont³, Norman G. Loeb¹⁰, Xioyan Ma¹¹, Vaishali Naik¹², Fabien Paulot¹², Philip Stier⁹, Martin Wild¹³, Gunnar Myhre¹⁴, and Michael Schulz¹⁵

¹Universität Leipzig, Institute for Meteorology, Leipzig, Germany

²University of Leeds, School of Earth and Environment, Leeds, U.K.

³International Institute for Applied Systems Analysis, Laxenburg, Austria

⁴Laboratoire de Météorologie Dynamique, Institut Pierre Simon Laplace, Sorbonne Université, Paris, France

⁵Norwegian Institute for Air Research, Kjeller, Norway

⁶Institut Pierre-Simon Laplace, Sorbonne Université / CNRS, Paris, France

⁷University of Reading, UK

⁸EUMETSAT, Darmstadt, Germany

⁹University of Oxford, Atmospheric, Oceanic and Planetary Physics, Oxford, U.K.

¹⁰NASA Langley Research Center, Hampton, USA

¹¹Nanjing University of Information Science & Technology, School of Atmospheric Physics, Nanjing, China

¹²Geophysical Fluid Dynamics Laboratory, Princeton, USA

¹³ETH Zürich, Department of Environmental Systems Science, Zürich, Switzerland

¹⁴CICERO, Oslo, Norway

¹⁵Norwegian Meteorological Institute, Oslo, Norway

Correspondence: Johannes Quaas (johannes.quaas@uni-leipzig.de)

Abstract. Anthropogenic aerosols exert a cooling influence that offsets part of the greenhouse gas warming. Due to their short tropospheric lifetime of only up to several days, the aerosol forcing responds quickly to emissions. Here we present and discuss the evolution of the aerosol forcing since 2000. There are multiple lines of evidence that allow us to robustly conclude that the anthropogenic aerosol effective radiative forcing – both aerosol-radiation and aerosol-cloud interactions – has become globally less negative, i.e. that the trend in aerosol effective radiative forcing changed sign from negative to positive. Bottom-up inventories show that anthropogenic primary aerosol and aerosol precursor emissions declined in most regions of the world; observations related to aerosol burden show declining trends, in particular of the fine-mode particles that make up most of the anthropogenic aerosols; satellite retrievals of cloud droplet numbers show trends in regions with aerosol declines that are consistent with these in sign, as do observations of top-of-atmosphere radiation. Climate model results, including a revised set that is constrained by observations of the ocean heat content evolution show a consistent sign and magnitude for a positive forcing relative to 2000 due to reduced aerosol effects. This reduction leads to an acceleration of the forcing of climate change, i.e. an increase in forcing by 0.1 to 0.3 W m⁻², up to 12% of the total climate forcing in 2019 compared to 1750 according to IPCC.

1 Introduction

15 Anthropogenic pollution particles, aerosols, exert an effective radiative forcing on climate due to aerosol-radiation interactions (ERF_{ari}, also known as "aerosol direct effect", combined with the "semi-direct effect") and aerosol-cloud interactions (ERF_{aci}, "aerosol indirect effect") (Chýlek and Coakley, 1974; Boucher et al., 2013; Forster et al., 2021; Szopa et al., 2021). ERF_{ari} occurs through the scattering and absorption of sunlight by aerosols while for ERF_{aci} aerosols act as cloud condensation nuclei (Twomey, 1974). Both entail rapid adjustments that tend to enhance the radiative forcing. A recent assessment provided
20 an estimated total ERF due to aerosols (ERF_{ae}) in the range of -2.0 to -0.35 W m^{-2} (5 to 95% confidence interval; 2005 to 2015 compared to 1850, Bellouin et al., 2020b). The latest assessment report by the Intergovernmental Panel on Climate Change (IPCC) concluded that the 2019 vs. 1750 ERF_{ae} has a best estimate of -1.1 W m^{-2} and 5 to 95% confidence interval of -1.7 to -0.4 W m^{-2} (Forster et al., 2021). This negative forcing offsets a sizeable fraction of the current CO₂ ERF. Throughout this manuscript, we consider ERF with 1750 as baseline, or changes in ERF over certain periods (most often from 2000 to 2019).
25 Forster et al. (2021) quantify a temperature increase in 2019 relative to 1750 of $+1.01^\circ\text{C}$ due to the ERF by CO₂ ($+1.81^\circ\text{C}$ considering all anthropogenic greenhouse gases), and a temperature change by -0.50°C due to aerosols in that period. This implies that without the cooling effect of aerosols, the world would already have reached the 1.5°C temperature threshold of "dangerous" climate change as set out by the Paris agreement.

A fundamental difference between radiative forcing by aerosols and long-lived greenhouse gases is tied to their atmospheric
30 lifetimes: greenhouse gases have lifetimes of decades to millennia (Solomon et al., 2009), while the lifetime of tropospheric aerosols is only up to several days. Climate thus responds to long-lived greenhouse gases such as CO₂ largely in terms of their cumulative emissions, but to aerosols in direct link to its current emissions rate. Shorter lived greenhouse gases such as methane have an intermediate effect, whereby deep reductions in emissions can have substantial effects on temperature within a few decades (Shindell and Smith, 2019; Smith et al., 2021b; Allen et al., 2022). A further reduction in aerosol emissions – required
35 due to their environmental and health impacts (Lelieveld et al., 2015; Cohen et al., 2017) – thus takes out the negative aerosol forcing and leads to a warming relative to the period prior to emission reduction (Brasseur and Roeckner, 2005; Dufresne et al., 2005), an effect also known as climate penalty of air quality improvements (Ekman et al., 2020; Hong et al., 2020). Also, the importance of the aerosol forcing relative to the CO₂ forcing was largest in the early industrial period (Stevens, 2015). It will continue to decrease, since anthropogenic aerosol emissions will likely decrease at the global level (Myhre et al., 2015; Szopa
40 et al., 2021).

At what point did the aerosol forcing became substantially less negative at a scale relevant for global forcing? There are suggestions that decreasing started in the last decades in different regions, and for several regions this trend reversal has been documented (e.g., Cermak et al., 2010). The decrease stems in particular from reductions of SO₂ emissions from coal use in the residential sector, power plants, and industry. For other regions, evidence is lacking or more anecdotal. However, for
45 understanding of global climate change, it is relevant to ask to which extent the aerosol climate forcing has become less negative at the global scale.

Here we propose that aerosol trends and their effects can be best investigated in the satellite era since the turn of the century. We analyse multiple observational and model datasets to demonstrate that both ERF_{ari} and ERF_{aci} show reduced trends since 2000 in regions which demonstrate a robust and substantial aerosol ERF trend in models.

50 2 Changes in aerosol emissions




Figure 1. Linear trends (2000 to 2019) of (a) anthropogenic emissions in sulfur dioxide (SO₂) from the Community Emissions Data System (CEDS v_2021_04_21; Hoesly et al., 2018); (b) and (c) as (a), but for anthropogenic emissions in organic carbon (OC) and black carbon (BC), respectively. Regions with small absolute trends (less than $7 \mu\text{g m}^{-2} \text{day}^{-1} \text{yr}^{-1}$) are masked by grey shading. Isolines enclose regions with trends in clear-sky solar ERF (see later, Fig. 4) larger than $0.05 \text{ W m}^{-2} \text{yr}^{-1}$ in absolute terms. The average values in these regions are shown in Fig. 6, listed in Table 1 and discussed in Section 7. A figure that combines the panels of Fig. 1 to 4 is provided as Supplementary material. A figure that shows the trends in absolute units is also provided as Supplementary material.

Despite substantial differences in their absolute magnitude especially at the regional level (Elguindi et al., 2020), the different emission inventories agree in general on the sign of the historical trends at regional and global levels (Granier et al., 2011; Klimont et al., 2017; Hoesly et al., 2018; Aas et al., 2019; Elguindi et al., 2020), especially over Europe and North America (Elguindi et al., 2020). A number of clear conclusions have thus been drawn in the literature for aerosol emissions in specific regions. Aerosol emissions have seen a steep increase since the beginning of the industrial period (e.g., Szopa et al., 2021). In several regions, declines after a peak are documented. An example is Europe, where since the 1980s, aerosol emissions declined strongly following air quality policies (Krüger and Graßl, 2002; Vestreng et al., 2007; Tørseth et al., 2012; Cherian et al., 2014; Crippa et al., 2016; Costa-Surós et al., 2019). A similar behaviour is documented for North America (e.g., Streets et al., 2009; Aas et al., 2019; Elguindi et al., 2020). Sulphur and nitrogen deposition over the USA, reflecting anthropogenic emissions, have been declining by between 1 and 3% yr⁻¹ in the period 1989–2010 (Sickles II and Shadwick, 2015). In contrast, anthropogenic aerosol emissions over China have been increasing until around 2010, and decreasing thereafter (Klimont et al., 2017; Zheng et al., 2018; Aas et al., 2019; Wang et al., 2021). The exact temporal evolution of aerosol emissions over the past 20 years especially over China was erroneously represented (a too weak decline since 2010) in some emission datasets, leading to some incompatibility of aerosols in the 6th Coupled Model Intercomparison Project (CMIP6; Eyring et al., 2016; Hoesly et al., 2018; Elguindi et al., 2020) in comparison to observations (Paulot et al., 2018; Wang et al., 2021). Aerosol emissions over

India continued to rise throughout the period 2000 to 2019 (Klimont et al., 2013; Wang et al., 2021). Over remote oceanic regions, ship emissions played a substantial, increasing role in the first part of the time period of interest (Smith et al., 2011). Since 2010, they have declined first in emission control areas (IMO, 2008) and since 2020 over much of the global oceans (IMO, 2019). This declining signal is visible also in cloud properties (Gryspeerdt et al., 2019).

70 Here we consider more specifically emissions from the the newest version of the Community Emissions Data System (CEDS; O'Rourke et al., 2021). The trends are shown in Fig. 1. A previous version of this dataset was used in CMIP6 and described by Hoesly et al. (2018). Sulfur emissions were mostly declining since 2000, in particular there were substantial declines over North America and Europe, continuing decreasing trends that started in the last decades of the 20th century. Also over East Asia, due to reductions after 2010, the overall trend is negative, despite the fact that in the first decade of the 21st century, emissions did
75 still increase. Over Southeast Asia, including India, and also over parts of Africa, sulfate precursor emissions showed increasing trends. Some shipping routes over ocean show increasing trends in this period. OC and BC emissions broadly show the same, but with more widespread increases especially over more regions in East Asia, Africa, and also South America. All considered aerosol species show increasing trends in emissions for high latitudes of both hemispheres. The updates of CEDS emissions (Elguindi et al., 2020) show that more recent evidence points to even stronger decline in SO₂ emissions in the second part of
80 the last decade and the BC and OC trends are showing a decline rather than increase, especially in China (see also Kanaya et al., 2020). These are further discussed in Elguindi et al. (2020).

3 Changes in aerosol abundance

The emission trends are reflected in observations of aerosol abundance. Due to their short lifetime, it is expected that regional trends in emissions are also reflected by regional trends in concentrations that are somewhat smoothed out spatially, in case
85 of typically prevailing wind directions mostly leewards. Trends in surface concentrations from in situ observations were found to show the expected trends in a global compilation (Collaud Coen et al., 2020) for sulfate and PM_{2.5}, and specifically so for the declining trends over Europe (Stjern et al., 2011; Aas et al., 2019) and North America (Jongeward et al., 2016; Aas et al., 2019), and the first increasing, then decreasing behaviour over China (Zheng et al., 2018; Aas et al., 2019).

The analysis of trends from remote sensing, especially from satellite remote sensing, is challenging, because datasets may not
90 be homogeneous over the lifetime of a satellite instrument, due to changing instrument response and satellite orbit. However, for NASA's Earth Observation Satellites Terra and Aqua, care has been taken to avoid many of the issues that hamper satellite trend analysis such as orbital drift (Levy et al., 2013). Studies show that trends from various satellites are, at least qualitatively, consistent (Wei et al., 2019). Declining trends in aerosols in certain regions, such as over Europe (Stjern et al., 2011; Cherian et al., 2014; Li et al., 2014; Georgoulias et al., 2016; Cherian and Quaas, 2020) and over the USA (Li et al., 2014; Jongeward et al., 2016; Cherian and Quaas, 2020), as seen from satellite analysis, have been documented earlier. The changes in aerosols
95 over East Asia, especially China, are not monotonic over the period of interest. Rather, the trends reversed from positive (2000–2010) to negative (since 2010), and this is seen in satellite observations of aerosols (Paulot et al., 2018; Sogacheva et al., 2018; Filonchyk et al., 2019; Ma et al., 2019; Samset et al., 2019). In contrast, over Southeast Asia, especially India, aerosol




Figure 2. Linear trends (2000 to 2019) of (a) aerosol optical depth (AOD) as retrieved from the Multi-angle Imaging Spectroradiometer (MISR; Garay et al., 2017) on board the Terra satellite, where the coloured circles show the AOD trends from the AERONET ground-based sunphotometer network (Holben et al., 1998; Giles et al., 2019) where data since 2000 are available; (b) as (a) but for the fine-mode AOD, i.e. the AOD due to aerosols with radius smaller than $1 \mu\text{m}$. (c) and (d) as for (a) and (b), but retrievals from the MODerate Resolution Imaging Spectroradiometer (MODIS; Levy et al., 2013) (fine-mode AOD unavailable over land) from the Terra satellite averaged (starting 2002) with MODIS retrievals from the Aqua satellite; (e) PMAp aerosol optical depth as retrieved from the Global Ozone Monitoring Experiment–2 (GOME-2) instrument on-board EUMETSAT’s Metop-A satellite that is available only for 2008 to 2017. Isolines as in Fig. 1. A figure that shows the trends in absolute units is provided as Supplementary material.

retrievals from satellites show continuing increases throughout the period (Li et al., 2014; Zhao et al., 2017; Dahutia et al., 2018; Hammer et al., 2018; Cherian and Quaas, 2020). Model-data synergy allowed to attribute these satellite-derived trends to the specific emission changes (Bauer et al., 2020; Yu et al., 2020), and to quantify the changes at between -3.1 and -1.2% yr^{-1} for the different regions affected by the declines in anthropogenic aerosol emissions from 2000 to 2014 (Mortier et al., 2020). Mortier et al. (2020) further documented that climate models were able to reproduce these trends quantitatively.

We report AOD trends from various satellite datasets on a common scale in Fig 2. It specifically shows aerosol optical depth (AOD) and fine-mode AOD (AODf) from the MODerate Resolution Imaging Spectroradiometer (MODIS, Levy et al., 2013) instrument on board the EOS Terra and Aqua satellites, and the Multi-Angle Imaging Spectro-Radiometer (MISR, Garay et al., 2017) instrument on board the EOS Terra satellite. Also the – presumably more stable – ground-based retrievals from the AERONET network (Holben et al., 1998, 2001; Giles et al., 2019) are analysed for the stations for which the time series since 2000 are available. The regression coefficients are reported, and it is to be kept in mind that none of the quantities really follows a straight line. Rather, the overall tendency of the noisy time change is referred to. The trends in both AOD and AODf from the different satellite instruments in the Southern hemisphere oceanic, and also in the Northern hemisphere high-latitude oceanic

regions differ – MODIS shows increases and MISR, decreases or scattered results in both quantities. As a third estimate, the EUMETSAT Polar Multi-sensor Aerosol optical properties product (PMAp, Grzegorski et al., 2021) climate data record derived using the Global Ozone Monitoring Experiment–2 (GOME-2) instrument on board EUMETSAT’s Metop-A satellite are used. These are available for a shorter, 10-year period, for 2008 to 2017. In the Southern hemisphere oceanic regions, Metop-A shows very small trends for this shorter period; in the Northern hemisphere high latitude oceans, it tends to confirm the decreases shown by MISR. The increase in AOD retrieved by MODIS has been reported in previous studies. Bai et al. (2020) report an increasing trend over the 2003 to 2017 period, they propose that there might have been an increase in sea salt consistent with increasing wind speed in reanalysis. In a different conclusion, Fan et al. (2018) demonstrate that while trends in AOD from MODIS are consistent with those derived from Aeronet over land in the Northern hemisphere, the trends over Australia and Southern America are inconsistent. Also in the study by Wei et al. (2019), the MODIS trends in the Southern hemisphere were reported to show stronger positive trends than the six other satellite products they examined for the 2003 to 2010 period in the oceanic regions of the Southern hemisphere they investigated (South Atlantic and Indian oceans). However, in the regions discussed above with pronounced trends in anthropogenic aerosol (precursor) emissions, the satellite trends show the same behaviour qualitatively in all three datasets. These trends are largely consistent with those from AERONET data (circles in Fig. 2). The decreasing trends over North America, Europe, and East Asia are clearly seen and at many grid points statistically significant (at 5% significance level according to a t-test with correction as in Santer et al., 2000), as are the increasing trends over India. It is particularly interesting to note that the trends in AODf are more consistent still in spatial extent to the changes in sulfate (precursor) emissions. These smaller particles, with radii $< 1 \mu\text{m}$, contain the bulk of the anthropogenic contribution to the aerosol (Bellouin et al., 2005; Kaufman et al., 2005; Kinne, 2019).

4 Changes in cloud properties




Figure 3. Linear trends (2000 to 2019) in cloud properties retrieved for liquid-water clouds from MODIS (Platnick et al., 2017) where cloud droplet number concentration (CDNC, panel a) and cloud liquid water path (LWP, panel b) are computed assuming adiabatic clouds (Quaas et al., 2006; Grosvenor et al., 2018); (c) liquid cloud fraction. Isolines as in Fig. 1. A figure that shows the trends in absolute units is provided as Supplementary material.

Clouds are a key determinant for variability and trends of the Earth's energy budget. Due to their large spatio-temporal variability it is not easy to distinguish long-term signals from weather noise. Clouds respond not only to aerosols, but also to global warming and interannual as well as decadal internal climate variability (Forster et al., 2021). Overall, satellite analysis documented changes in clouds that are consistent with several of the hypotheses relevant for cloud-climate feedbacks (Norris et al., 2016), but little evidence for patterns of cloud cover or cloud-top altitude trends that would be expected due to aerosol-cloud interactions (Norris et al., 2016). The most immediate impact of aerosols is on cloud droplet number concentration (Bellouin et al., 2020b; Quaas et al., 2020). For this microphysical quantity, some clear and significant trends were identified in satellite observations for the outflow region east of East Asia (Bennartz et al., 2011), albeit the declining trends in cloud water path and cover are not necessarily what is expected in relation to aerosol-cloud interactions (Benas et al., 2020).

The trends in satellite-derived cloud droplet number concentrations were reported previously to be consistent with the aerosol trends in several regions (McCoy et al., 2018; Cherian and Quaas, 2020). Trends in cloudiness and cloud radiative properties are, however, less conclusive possibly due to their large variability beyond the variability driven by aerosols (Norris et al., 2016; Cherian and Quaas, 2020), and also since their response to aerosols depends on cloud regime (Mülmenstädt and Feingold, 2018; Zhang et al., 2022).

The trends in MODIS retrievals of cloud properties (Platnick et al., 2017) are shown in Fig. 3. MODIS Terra (10.30 h overpass time) is combined with MODIS Aqua (13.30 h overpass time) from 2002 onwards. Cloud droplet number concentration is derived from the MODIS retrievals as discussed in Grosvenor et al. (2018). For all three cloud quantities presented, only liquid-water clouds, as determined by the retrieval algorithm, are selected. The results confirm the qualitative consistency between droplet number and aerosol trends. This consistency is inferred from the similarity in sign across the regions in which aerosols show spatially contingent trends attributable to the anthropogenic aerosol emissions as evident from Fig. 1 in comparison to Fig. 2, and in particular as summarized later in Table 1. Cloud droplet concentrations show declines especially over the oceans of the Northern hemisphere mid-latitudes, in particular downwind of the regions where aerosol emissions declined. The signal is much weaker over the continents, though (as also discussed by Ma et al., 2018). Cloud liquid water path (related to cloud thickness; defined in the satellite retrievals as cloudy-sky rather than all-sky) does not show trend patterns that would be strongly related to the pattern of trends in droplet concentration. It was documented earlier that the adjustment of liquid water path to cloud droplet concentration perturbation appears to be weak in comparison to natural variability (Malavelle et al., 2017; Toll et al., 2019; Haghghatnasab et al., 2022; Chen et al., 2022). In contrast, the change in cloud fraction is broadly consistent in pattern and sign with the trends in droplet concentration. This was also suggested by satellite correlation studies (Gryspeerd et al., 2016; Rosenfeld et al., 2019; Christensen et al., 2020) and analysis of the response of clouds to volcanic eruptions (Chen et al., 2022). It is to be noted that cloud properties, especially outside the regions with strong aerosol changes, also respond to global warming (in particular, sea surface temperature trends under stratocumulus regions such as the Eastern Pacific) and natural variability. Thus, only the averaging over the larger contingent regions with consistent aerosol changes may allow to reduce this "noise" to infer a possible signal.

Nevertheless, the conclusion of this review of trends in cloud quantities is that cloud droplet concentrations show trends that are spatially consistent with the expectation of declining anthropogenic aerosol emissions, and that apparently also liquid-cloud

fraction trends show pattern consistent with the aerosol declines. Since these retrievals are independent of the aerosol retrievals discussed earlier, this is a strong corroboration of the earlier conclusion that satellites show a declining trend in aerosols in regions of anthropogenic emissions.

170 5 Changes in radiation




Figure 4. Linear trends (2000 to 2019) in (a) net broadband solar flux for clear-sky situations as retrieved from the Clouds and the Earth’s Radiant Energy System (CERES) Energy Balanced and Filled (EBAF) product (Loeb et al., 2018a) from Terra, averaged for years after 2002 with retrievals also from Aqua; (b) and (c) as (a) but for all-sky solar radiation fluxes and all-sky solar plus terrestrial net fluxes, respectively; (d – f) trends in effective radiative forcings due to aerosols as computed from the dedicated RFMIP (Pincus et al., 2016) / AerChemMIP (Collins et al., 2017) for which output was available from the CanESM5 (Swart et al., 2019), GISS-E2-1-G (Kelley et al., 2020), HadGEM3-GC31-LL (Andrews et al., 2019), IPSL-CM6A-LR (Boucher et al., 2020), MIROC6 (Hajima et al., 2020), NOAA-GFDL (Held et al., 2019), and NorESM2-LM (Seland et al., 2020) of which the ensemble average is shown for (d) clear sky solar, (e) all-sky solar and (f) all sky net (solar plus terrestrial) spectra. Signs are inverted for consistency with the CERES results (negative trends in ERF_{aer} mean decreases in absolute magnitude). Isolines enclose regions with trends in clear-sky solar ERF (panel d) larger than $0.05 \text{ W m}^{-2} \text{ yr}^{-1}$ in absolute terms. The average values in these regions as listed in Table 1 and discussed in Section 7. Net fluxes, defined as positive if downward, are plotted, consistently in the simulations and the satellite retrievals.

Changes in net top-of-atmosphere radiation fluxes in a period correspond to the changes in ERF in that period, but also include the signal of natural variability and of feedbacks to changing climate. Previous analysis of model simulations suggested that between 2000 and 2015, ERF_{aer} was reduced in absolute magnitude, i.e. increased (became less negative), by about $0.003 \text{ W m}^{-2} \text{ yr}^{-1}$ at a global scale (Myhre et al., 2017), mainly over the Northern hemisphere mid-latitudes, especially

175 over North America, the North Atlantic ocean, Europe and adjacent Asia. In Fig. 4, the trends in ERFaer as simulated by models contributing to CMIP6 are analysed. This makes use of the dedicated simulations of the Radiative Forcing Model Intercomparison Project (RFMIP, Pincus et al., 2016) that trace the aerosol ERF over time (the piClim-histaer simulations). For seven Earth System Models, the relevant diagnostics for these simulations were submitted, namely for the Canadian Centre for Climate Modelling and Analysis (CCMA, Swart et al., 2019), for the US National Aeronautics and Space Administration
180 Goddard Institute for Space Studies Earth system model (GISS-E2-1-G, Kelley et al., 2020), for the UK Hadley Centre Global Environment Model (HadGEM3-GC31-LL, Andrews et al., 2019), for the Institut Pierre Simon Laplace Climate Model (IPSL-CM6A-LR, Boucher et al., 2020), for the Model for Interdisciplinary Research on Climate (MIROC6, Hajima et al., 2020), for the National Oceanic and Atmospheric Administration / Geophysical Fluid Dynamics Laboratory Climate Model (CM4, Held et al., 2019), and for the Norwegian Earth System Model (NorESM2-LM, Seland et al., 2020). The ensemble average of these
185 models is considered. The results show that the pattern in the clear sky solar ERFaer trends is closely related to the pattern in the trends in sulfate precursors. It reflects the strong declines in the main source areas over North America, Europe, and East Asia, along with the increases over India and surrounding areas. The patterns in all-sky ERFaer, i.e. including the cloud effects, both in solar and terrestrial spectra, are noisier but also show trends that are consistent with the pattern seen for the clear-sky, solar ERFaer and in aerosols and droplet concentrations also in the observations. It is interesting to assess the relative
190 importance of aerosols to other agents that are examined in the RFMIP piClim-histall simulations (Supplementary Fig. S5). In the solar spectrum, as expected, clearly the aerosol signal dominates, whereas in the terrestrial spectrum, the additional signal by the increase in greenhouse gases is seen. Supplementary Fig. S6 to S8 examine the differences across models in the simulated aerosol ERF trends. Despite differences in particular in the absolute magnitude, the pattern identified in the multi-model mean is qualitatively similar in all seven individual models. Similarly, for individual ensemble members of a selected ESM
195 (Supplementary Fig. S9 to S11) the pattern is robustly simulated.

In the multi-model mean, global mean changes show a decline of the clear-sky, solar ERFaer by $0.0117 \text{ W m}^{-2} \text{ yr}^{-1}$, of the all-sky solar ERFaer a decline by $0.0172 \text{ W m}^{-2} \text{ yr}^{-1}$ and of the all-sky terrestrial ERFaer a (compensating) increase by $0.0013 \text{ W m}^{-2} \text{ yr}^{-1}$. The integral, net decline over the 20-year period according to these models thus was 0.32 W m^{-2} .

This result can be compared to the assessment by IPCC AR6 (Forster et al., 2021). Their assessment is based on multiple lines
200 of evidence that are incorporated in an emulator ensemble simulation. The time series of the diagnosed ERFaer is available via the IPCC web site and at <https://doi.org/10.5281/zenodo.5705391>. Computing the linear trend between 2000 and 2019 on the basis of the emulator ensemble yields an increase by $+0.0145 \text{ W m}^{-2} \text{ yr}^{-1}$ between 2000 and 2019 (5 to 95% confidence interval of $+0.0068$ to $+0.0253$), i.e. by $+0.29$ ($+0.14$ to $+0.51$) W m^{-2} over the full period (Gulev et al., 2021; Forster et al., 2021).

205 The ERFaer may be inferred from the Earth radiation budget which is measurable at the top of the atmosphere. Several studies have investigated the retrievals of this quantity from the Clouds and the Earth's Radiant Energy System (CERES) Energy instrument that is also on the EOS Terra and Aqua satellites. CERES shows patterns for clear-sky broadband radiation that are consistent with the aerosol spatio-temporal changes (Loeb et al., 2018b; Paulot et al., 2018; Loeb et al., 2021b). Further, Loeb et al. (2021a) document an increase in the Earth's energy imbalance, seen in both Earth radiation budget satellite observations

210 and ocean heat content. They find this to be due to a strongly decreasing trend in reflected solar radiation, which they attribute to decreased reflection by clouds and sea ice, and a declining trend in emitted terrestrial radiation due to increases in greenhouse gases and water vapour. Using partial radiative perturbation analysis, Loeb et al. (2021a) attribute the trend in solar radiation mostly to changes in clouds, with a very small contribution only due to the effect by aerosol-radiation interactions. This is also a result of a new study by Jenkins et al. (2022). CERES observations were also analysed by Raghuraman et al. (2021). They
215 find for the period 2001 to 2020 an increasing trend by $0.038 \pm 0.024 \text{ W m}^{-2} \text{ yr}^{-1}$. They attribute about one third of this trend to the reduction in aerosol ERF.

Kramer et al. (2021) disentangle the trends in satellite-retrieved radiation fluxes using radiative kernels, notably isolating the impact of radiative forcings. They quantify the change in absorbed solar radiation over the 2003 to 2018 period at $0.044 \pm 0.02 \text{ W m}^{-2} \text{ yr}^{-1}$. Singling out the instantaneous radiative forcing in the solar spectrum, they obtain a change of
220 $0.006 \pm 0.003 \text{ W m}^{-2} \text{ yr}^{-1}$ which they largely attribute to aerosol changes. Paulot et al. (2018) constrained the radiative forcing due to aerosol-radiation interactions (aerosol direct effect) in the GFDL climate model and obtained an almost negligible trend in ERFaer of $0.0002 \text{ W m}^{-2} \text{ yr}^{-1}$. Their study, however, considered the period from 2001 to 2015 only and thus a time when then increasing emissions over China were much more relevant.

Bellouin et al. (2020a) used the Copernicus reanalysis of atmospheric composition, which assimilates MODIS AODs, to
225 estimate RFaer and found statistically significant decreasing (less negative) trends over North and South America, Europe, and China, and an increasing (more negative) trend over India for the period 2003 to 2017. Their globally averaged trend in RFaer is $0.00 \text{ W m}^{-2} \text{ yr}^{-1}$, but limitations in their estimate may imply that the real trend is positive.

Surface measurements of radiation also show increasing trends over large regions (Wild, 2009, 2012; Cherian et al., 2014; Hatzianastassiou et al., 2020). Trends in aerosol effects become particularly apparent in surface radiation records under cloud-
230 free conditions. Such records indicate in Europe increasing clear-sky surface solar radiation and thus decreasing aerosol effects throughout the 2000s with some tendency for saturation (leveling off) after 2010 (Manara et al., 2016; Wild et al., 2021). Surface radiation records in China suggest a trend reversal in clear sky surface solar radiation from decrease to increase in the late 2000s (Yang et al., 2019), in line with anthropogenic aerosol emission trends (Section 2). It has been shown for Europe (Pfeifroth et al., 2018) and China (Wang et al., 2019) that solar radiation consistently increases in both surface and satellite
235 observations.

The CERES data are also shown in Fig. 4. The clear-sky solar radiation changes in the areas where the models show decreases in the clear-sky solar ERFaer, with a pattern consistent in sign and magnitude with the model results. For all-sky radiation, the data show larger trends and also much more noise in the patterns. However, the sign of the changes in the regions where an aerosol signal is expected is consistent between the models and the data. These results are consistent with what was documented
240 in the literature before (see above). A quantitative comparison is provided later in Section 7.

6 Ocean heat uptake and surface temperatures as constraints for the simulated ERFaer evolution

The temporal evolution of observed climate change – specifically, surface temperature changes and their pattern – has been proposed as a constraint on the magnitude of the aerosol effective radiative forcing (Ekman, 2014; Rotstayn et al., 2015; Stevens, 2015; Kretzschmar et al., 2017; Aas et al., 2019; Albright et al., 2021; Smith and Forster, 2021). Increasingly it is now
245 recognized that the ocean heat uptake is of overwhelming interest for monitoring the Earth energy imbalance (von Schuckmann et al., 2016; Palmer, 2017; Allison et al., 2020; Forster et al., 2021), since it is a non-volatile indicator of climate change.

Based on this, Smith et al. (2021a) constrained the aerosol ERF from the CMIP6 models by considering the ocean heat uptake from observations between 1971 and 2018, in addition to observations of surface temperature. In Smith et al. (2021a) a 100,000 member ensemble of time series of historical aerosol forcing was generated from emissions of BC, OC and SO₂ using
250 simple formulas calibrated to CMIP6 models, considering ERFari and ERFaci separately, with 1850–2010 ERFari and ERFaci constrained to the distribution of Bellouin et al. (2020b) in the prior. Weights were assigned to each ensemble member based on how closely historical surface temperature and ocean heat content change were simulated compared to observations when the aerosol forcing was combined with other historical forcings in a two-layer energy balance model (Geoffroy et al., 2013) to generate a posterior distribution of historical aerosol forcing. This study assessed the ERFaer between 1750 and 2019 at
255 -0.9 W m^{-2} and suggested a slightly positive trend between 1980 and 2014 of $0.0025 \text{ W m}^{-2} \text{ yr}^{-1}$. The method of Smith et al. (2021a) is applied to assess the trend in aerosol ERF between 2000 and 2019 (Fig. 5) focusing on the emissions trends during this period. This yields a constrained trend of $0.0114 (-0.003 \text{ to } 0.0274) \text{ W m}^{-1} \text{ yr}^{-1}$, much stronger than the one considering the longer period (5 to 95% confidence interval in brackets). The integral change in the 2000 – 2019 period in ERFaer is thus 0.23 W m^{-2} in the best estimate, very close to the suggestion by the analysed models.

260 Albright et al. (2021) explore bounds on ERFaer using a Bayesian model of aerosol forcing and Earth’s multi-timescale temperature response to radiative forcing, finding a best estimate present-day lower bound of -1.3 W m^{-2} . In Fig. 5, their method is applied to the period investigated here, from 2000 to 2019. Their baseline estimate yields a mean trend of $0.0047 \text{ W m}^{-2} \text{ yr}^{-1}$ (5 to 95% confidence interval of $-0.000912 \text{ to } 0.0106 \text{ W m}^{-2} \text{ yr}^{-1}$). The best estimate of the change in ERFaer for the 20-year-period is thus 0.094 W m^{-2} (5 to 95% confidence interval of $-0.018 \text{ to } 0.21 \text{ W m}^{-2}$). Prescribing internal climate variability
265 that is a factor of five larger than the CMIP6 mean and assuming large, correlated errors in global temperature observations yields a fifth-percentile ERFaer lower bound of -1.8 W m^{-2} and a mean estimate of the change in ERFaer for the 20-year-period of 0.16 W m^{-2} (5 to 95% confidence interval of $0.04 \text{ to } 0.32 \text{ W m}^{-2}$, see ‘increased variance’ in Fig. 5b).

Albright et al. (2021) caution that ocean heat content data do not, at present, offer robust additional constraints on ERFaer. Biases in coupled climate models towards too positive radiative feedbacks since about 1980, compared to radiative feedbacks
270 in models forced by historical surface temperatures (e.g., Zhou et al., 2016), suggest that emulators trained on coupled models yield inferences of ERFaer that are biased low when attempting to fit to recent planetary heat uptake (see Sec. 4d in Albright et al., 2021)). That is, the discrepancy between the observed planetary heat uptake and net top-of-atmosphere radiative imbalance in coupled models can be thought of as a ‘ghost forcing’ that can either be attributed to more negative radiative feedbacks or more negative radiative forcing (e.g., more negative ERFaer). They conclude that constraining ERFaer with ocean heat

275 content data requires independent observational constraints on the true radiative damping and a better understanding whether recent SST patterns that cause more negative feedbacks are forced or unforced (Andrews et al., 2018; Sherwood et al., 2020).

Jenkins et al. (2022) also suggests that 0.2 W m^{-2} is a plausible best-estimate ERF change for the considered period, based on analysis of satellite observations and energy balance of global temperatures. However, they also conclude that large variability signals implies that a very weak trend change cannot be ruled out either.




Figure 5. Assessment of the linear trend in ERFaer between 2000 and 2019. (a) as in Smith et al. (2021a, their Fig. 7; abbreviated as S21 in the labels), and (b) as in Albright et al. (2021, labelled A21). The constraint is as in the cited studies, but applied to the period 2000 to 2019. Increased variance in Albright et al. (2021) corresponds to a scenario prescribing internal climate variability that is a factor of five larger than the CMIP6 mean and assuming large, correlated errors in global temperature observations, yielding a fifth-percentile ERFaer lower bound of -1.8 W m^{-2} . The labels along the x-axis correspond to 5 and 95% percentiles, as well as mean, of the distribution of the curves in the corresponding colour.

280 7 Discussion and conclusions

The trends of aerosols, clouds and radiation in the observations are subject not only to changes in anthropogenic aerosol emissions, but also to other influencing factors. These include changes in natural aerosol emissions, which remain poorly constrained and contribute a substantial fraction of total observed AOD, interannual variability, and responses to greenhouse-




Figure 6. Mean values derived from the maps in Fig. 1 to 4, averaged over (green/downward hatching) the regions with substantial negative trends (defined as larger than $0.05 \text{ W m}^{-2} \text{ yr}^{-1}$ in absolute terms; isolines in Fig. 4) in ERF clear sky, solar from RFMIP (Fig. 4d) and (purple/upward hatching) substantial positive trends. The grey bar (crosses) is the global mean between 60°S and 60°N . The regions with negative trends cover 7.3% of the Earth surface, the ones with positive trends, 1.1%. AODFM is retrieved only over oceans, and the PMAp time series only spans ten years from 2008 to 2017. From left to right: Trends in emissions of SO_2 , OC and BC, in AOD and AODFM from MISR, MODIS and Metop-A, in cloud droplet concentration N_d , LWP and cloud fraction from MODIS. All these are provided in units of $\% \text{ yr}^{-1}$ and refer to the left axis. Trends in CERES retrievals of solar clear-sky, solar all-sky and net all-sky radiation, as well as in model-derived ERF for solar clear-sky, solar all-sky, and net all-sky are shown in units of $\text{W m}^{-2} \text{ yr}^{-1}$ (right axis) and are inverted in sign: negative trends here are for reduction in magnitude (less negative fluxes / effective forcings).

gas-induced global warming; aerosol-cloud interactions may also be altered in a changing climate (Zhou et al., 2021; Murray-
285 Watson and Gryspeerdt, 2022).

(a) SO ₂ emissions (% yr ⁻¹)			(b) OC emissions (% yr ⁻¹)			(c) BC emissions (% yr ⁻¹)		
Decreases	Increases	Global	Decreases	Increases	Global	Decreases	Increases	Global
-5.34	+3.80	-0.61	-0.71	+1.03	+0.88	-2.50	+1.37	+0.78
(d) MISR AOD (% yr ⁻¹)			(e) MISR AODFM (% yr ⁻¹)					
-1.15	+1.35	-0.41	-1.66	+1.34	-0.92			
(f) MODIS AOD (% yr ⁻¹)			(g) MODIS AODFM (% yr ⁻¹)			(h) PMAp AOD (% yr ⁻¹)		
-0.85	+1.74	+0.16	-0.88	+2.36	+0.53	-1.79	+0.11	-0.87
(i) N _d (% yr ⁻¹)			(j) LWP (% yr ⁻¹)			(k) Cloud fraction (% yr ⁻¹)		
-0.43	+0.07	-0.15	-0.16	-0.11	-0.04	-0.16	+0.11	-0.10
(l) CERES rsutcs (W m ⁻² yr ⁻¹)			(m) CERES rsut (W m ⁻² yr ⁻¹)			(n) CERES net (W m ⁻² yr ⁻¹)		
-0.104	+0.041	-0.029	-0.186	+0.040	-0.059	-0.226	+0.030	-0.042
(o) ERF SW clr (W m ⁻² yr ⁻¹)			(p) ERF SW (W m ⁻² yr ⁻¹)			(q) ERF net (W m ⁻² yr ⁻¹)		
-0.087	+0.102	-0.012	-0.096	+0.087	-0.017	-0.083	+0.074	-0.016

Table 1. Values corresponding to Fig. 6. The trends in absolute units are reported in Supplementary Table S1.

IPCC AR6	+0.29 (+0.14 to +0.51) W m ⁻²
Method of Smith et al. (2021), constraint from ocean heat uptake	+0.23 (-0.05 to 0.55) W m ⁻²
Method of Albright et al. (2021), constraint from surface temperature changes	+0.094 (-0.02 to 0.21) W m ⁻²
RFMIP models (Fig. 4)	+0.32 W m ⁻²
Kramer et al. (2021)	+0.12 W m ⁻²
Raghuraman et al. (2021)	+0.24±0.20 W m ⁻²

Table 2. Estimates of ERFaer change between 2000 and 2019. The 5 to 95% uncertainty ranges are provided. Kramer et al. (2021) assess RF due to aerosols and use the period 2003 to 2018; Raghuraman et al. (2021) the period 2001 to 2019.

Natural aerosol emissions, especially of dust, are highly variable and impact the distribution of AOD in specific regions (Chin et al., 2014). Natural aerosol emissions may respond to increasing temperatures (Yli-Juuti et al., 2021). Also volcanic aerosol emissions, both from eruptions and from degassing, are an important contribution in particular to atmospheric sulfate aerosol. However, in satellite retrievals for the 2005 to 2016 period, no strong trends across volcanoes have been observed (Carn et al., 2017). Fires may emit large amounts of aerosols. This was in particular the case for the Australian bush fires in 2020 (Boer et al., 2020; Heinold et al., 2021). Whether there were substantial trends in fire aerosol emissions in recent decades is not quite clear (Doerr et al., 2016), even if burned area decreased in many regions (Andela et al., 2017). However, global warming increases the risk of fire (van Oldenborgh et al., 2021). More broadly, biomass burning aerosols are typically considered separately from the anthropogenic emissions presented in Fig. 1. In particular in the northern hemisphere high

295 latitudes, increasing emissions in biomass burning occurred in the period of interest (van et al., 2017) and likely explain the increases in aerosol abundance (Fig. 2) seen in these regions.

Sea salt aerosols, that are a function of near-ocean-surface wind speed, are subject to variability, both forced and unforced, albeit to a lesser extent than dust (Stier et al., 2006). To the extent that the MODIS, rather than MISR, AOD and AODFM trends above the southern hemisphere oceans are right, such variability in sea salt aerosol may cause the increasing trends (Struthers et al., 2013). Trends in long-range transport of aerosols could be another reason for such increases. However, the satellite retrievals are particularly uncertain in this region, due to the large zenith angles and large cloud cover that both hamper aerosol retrievals.

Towards the end of the time series investigated here, there were specific effects due to the COVID-19 pandemic in 2020 (Forster et al., 2020; Gettelman et al., 2021; Fiedler et al., 2021). For this reason, and due to the particularly large fire activity, the end year of the data analysed here was chosen as 2019.

The different quantities investigated here are not independent. The climate models are driven by the emissions. In turn, the emissions inventories consider satellite retrievals for some of their aspects such as fires and shipping. The satellite retrievals of cloud properties and radiation are not linked to the other quantities but are more noisy in their results. Cloud properties respond to variability in climate dynamics, both forced and unforced, beyond the impact of anthropogenic aerosols. Norris et al. (2016) document global patterns of changes in cloud coverage and cloud albedo that show a reduction in cloud cover and albedo in the mid-latitudes and an increase in the Tropics from the 1980s to the first decade of the 21st century, and that are consistent with the expectations due to cloud responses to global warming.

We now turn to discussing quantifying the changes in aerosols, clouds, and radiation. For this, the regions with clear trends in aerosols are identified by subjectively choosing the regions in which the ERF simulated by the CMIP6 models (Fig. 4) exceeds $\pm 0.05 \text{ W m}^{-2} \text{ yr}^{-1}$ for the solar, clear sky component. Regions with increasing and decreasing ERF are distinguished. Table 1 summarizes all quantities analysed in Fig. 1 to 4. In the regions with declining clear-sky solar ERF, in particular SO_2 emissions decreased strongly, but also OC and BC emissions decrease according to the inventory of Hoesly et al. (2018). In regions with increasing clear-sky solar ERF, emissions of all three species increased. Both MODIS and MISR show corresponding declining trends in column-aerosol metrics for the regions with aerosol emission reductions, and increasing trends where aerosols increased. The numbers are much larger for MISR than for MODIS for the declining-trend regions (almost a factor of 2 larger in case of AODFM). Also in the global average, the MODIS-derived AOD and AODFM trends are positive, in consistent with the result from MISR and Metop-A, and also inconsistent with the trends in clouds and radiation. This is due to the aerosol increases in MODIS over much of the oceans, especially in the southern hemisphere. The reasons are debated in the literature (see above). In contrast, MODIS-retrieved cloud droplet number, liquid water path and cloud fraction in these regions all increase (decrease) where aerosol emissions increase (decrease). Globally, all three quantities decrease. Droplet number concentrations change by a rate that is a factor of 2 (compared to MODIS AOD) to 4 (compared to MISR AODFM) less than for aerosol optical depth, highlighting that there is not a 1:1 relationship of droplet number and aerosol measured as AOD (e.g., Quaas et al., 2020; Jia et al., 2022). Since the regions with increasing aerosol are more limited in extent, the absolute numbers are expected to be more uncertain. There is only a small LWP response that is inconsistent in

330 sign between regions with increasing and decreasing aerosol emissions. LWP does not just respond to N_d perturbations, but
also to global warming (with an expected increase in LWP on average; e.g., Norris et al., 2016). However, the fact that there
is little LWP trend where N_d trends are substantial is consistent with other observations-based assessments (Malavelle et al.,
2017; Toll et al., 2019). Cloud fraction, in turn, appears to show decreasing trends in the regions with anthropogenic aerosol
decreases, and increasing trends where aerosol increases. Although it is also a function of other drivers, this could be a hint at
335 a systematic (positive, i.e. negative in terms of forcing) aerosol effect on cloud fraction as also documented in earlier statistical
studies (Gryspeerd et al., 2016; Rosenfeld et al., 2019; Christensen et al., 2020; Chen et al., 2022). It is to be noted that
the spatial consistency of the trends in cloudiness are not a proof of causality. Consistent with these results, CERES shows
decreasing trends in top-of-atmosphere radiation budget. The changes in net radiation retrieved by CERES are expected to
reflect the trends in ERF, but also natural variability and feedbacks to climate change. The numbers are stronger for all-sky
340 than for clear-sky, indicating a comparatively strong contribution by aerosol-cloud interactions (Forster et al., 2021; Loeb
et al., 2021a). The numbers are consistent in sign with what the CMIP6 models suggest for changes in ERFaer, although more
negative (less positive where aerosols increase) in particular when the cloud effects are included. The global average values
similarly show stronger declines than CMIP6 but are also consistent in sign.

In conclusion, there are clear, robust and consistent signals for net declining anthropogenic aerosol influence on climate in the
345 period since 2000, i.e. the period for which high-quality satellite retrievals of all relevant quantities are available. The regions in
which aerosol emissions declined (in particular North America, Europe and East Asia) dominate over regions with increasing
trends. The summary of the results in terms of aerosol effective climate forcings are listed in Table 2. This demonstrates
consistency of the findings of this study to previous ones. The overall climate-relevant signal is a decline in negative aerosol
effective radiative forcing by about 0.1 to 0.3 W m^{-2} , i.e. between 15 and 50% of the 0.6 W m^{-2} increase in CO_2 ERF (Forster
350 et al., 2021) in the same time period. This signal will very likely continue in the future, increasing the urgency for strong
measures on reducing greenhouse gas emissions (McKenna et al., 2021).

Data availability. The MODIS cloud products MYD08_D3 from Aqua and MOD08_D3 from Terra were used in this study from the Atmo-
sphere Archive and Distribu- tion System (LAADS) Distributed Active Archive Center (DAAC), <https://ladsweb.nascom.nasa.gov/>. MISR
data were obtained from the NASA Langley Research Center Atmospheric Science Data Center (<https://opendap.larc.nasa.gov/opendap/>
355 MISR/MIL3YAEN.004). CERES data were obtained from <https://ceres.larc.nasa.gov/data/>. The Metop-A data are available as PMAp Cli-
mate Data Record (CDR) at https://doi.org/10.15770/EUM_SEC_CLM_0053. AERONET data were used from https://aeronet.gsfc.nasa.gov/data_push/AOT_Level2_Monthly.tar.gz. RFMIP model output is available from the Earth System Grid Federation (ESGF).

Author contributions. The fundamentals stem from active discussions with all authors. J.Q. coordinated the study and led the writing of the
manuscript with significant contributions from all authors. H.J. created Fig. 1 to 4, including processing the data, and Table 1. with substantial
360 help from C.S. for the RFMIP models and M.D.-B. for the Metop-A data. A.-L.A. and C.S. prepared Fig. 5 and the data processed for it.

Competing interests. The authors declare there are no competing interests.

8 Acknowledgments

This work stems from discussions and work in the EU Horizon2020 projects FORCES (GA no. 821205) and CONSTRAIN (GA no. 820829). H.J. and J.Q. further acknowledge support by the German Research Foundation (Joint call between National
365 Science Foundation of China and Deutsche Forschungsgemeinschaft, DFG, GZ QU 311/28-1, project "CloudTrend"). PS acknowledges funding from the European Research Council (ERC) project RECAP under the European Union's Horizon 2020 research and innovation program with grant agreement 724602.

We would like to thank Annica Ekman (Stockholm University), Bjorn Stevens (Max Planck Institute for Meteorology, Hamburg), and Songmiao Fan (GFDL Princeton) for insightful comments.

370 We thank all data producers to make their data available. We acknowledge the World Climate Research Programme, which, through its Working Group on Coupled Modelling, coordinated and promoted CMIP6. We thank the climate modeling groups for producing and making available their model output, and the multiple funding agencies who support CMIP6 and ESGF. We would also like to thank Michael Diamond and two anonymous reviewers for constructive and helpful comments. We further acknowledge the Open Access Publishing Fund of Leipzig University supported by the German Research Foundation within
375 the program Open Access Publication Funding.

References

- Aas, W., Mortier, A., Bowersox, V., Cherian, R., Faluvegi, G., Fagerli, H., Hand, J., Klimont, Z., Galy-Lacaux, C., Lehmann, C. M. B., Lund Myhre, C., Myhre, G., Olivié, D., Sato, K., Quaas, J., Rao, P. S. P., Schulz, M., Shindell, D., Skeie, R. B., Stein, A., Takemura, T., Tsyro, S., Vet, R., and Xu, X.: Global and regional trends of atmospheric sulfur, *Sci. Rep.*, 9, 953, <https://doi.org/10.1038/s41598-018-37304-0>, 2019.
- Albright, A. L., Proistosescu, C., and Huybers, P.: Origins of a Relatively Tight Lower Bound on Anthropogenic Aerosol Radiative Forcing from Bayesian Analysis of Historical Observations, *J. Climate*, 34, 8777–8792, <https://doi.org/10.1175/JCLI-D-21-0167.1>, 2021.
- Allen, M., Peters, G., Shine, K., Azar, C., Balcombe, P., Boucher, O., Cain, M., Ciais, P., Collins, W., Forster, P. M., Frame, D. J., Friedlingstein, P., Fyson, C., Gasser, T., Hare, B., Jenkins, S., Hamburg, S. P., Johansson, D. J. A., Lynch, J., Macey, A., Morfeldt, J., Nauels, A., Ocko, I., Oppenheimer, M., Pacala, S. W., Pierrehumbert, R., Rogelj, J., Schaeffer, M., Schleussner, C. F., Shindell, D., Skeie, R. B., Smith, S. M., and Tanaka, K.: Indicate separate contributions of long-lived and short-lived greenhouse gases in emission targets, *npj Clim. Atmos. Sci.*, 5, 5, <https://doi.org/10.1038/s41612-021-00226-2>, 2022.
- Allison, L. C., Palmer, M. D., Allan, R. P., Hermanson, L., Liu, C., and Smith, D. M.: Observations of planetary heating since the 1980s from multiple independent datasets, *Environ. Res. Comm.*, 2, 101 001, <https://doi.org/10.1088/2515-7620/abb39>, 2020.
- Andela, N., Morton, D. C., Giglio, L., Chen, Y., van der Werf, G. R., Kasibhatla, P. S., DeFries, R. S., Collatz, G. J., Hantson, S., Kloster, S., Bachelet, D., Forrest, M., Lasslop, G., Li, F., Mangeon, S., Melton, J. R., Yue, C., and Randerson, J. T.: A human-driven decline in global burned area, *Science*, 356, 1356–1362, <https://doi.org/10.1126/science.aal4108>, 2017.
- Andrews, T., Gregory, J. M., Paynter, D., Silvers, L. G., Zhou, C., Mauritsen, T., Webb, M. J., Armour, K. C., Forster, P. M., and Titchner, H.: Accounting for changing temperature patterns increases historical estimates of climate sensitivity, *Geophys. Res. Lett.*, 45, 8490–8499, 2018.
- Andrews, T., Andrews, M. B., Bodas-Salcedo, A., Jones, G. S., Kuhlbrodt, T., Manners, J., Menary, M. B., Ridley, J., Ringer, M. A., Sellar, A. A., Senior, C. A., and Tang, Y.: Forcings, Feedbacks, and Climate Sensitivity in HadGEM3-GC3.1 and UKESM1, *J. Adv. Model. Earth Syst.*, 11, 4377–4394, <https://doi.org/https://doi.org/10.1029/2019MS001866>, 2019.
- Bai, H., Wang, M., Zhang, Z., and Liu, Y.: Synergetic satellite trend analysis of aerosol and warm cloud properties over ocean and its implication for aerosol-cloud interactions, *J. Geophys. Res. Atmos.*, 125, e2019JD031 598, <https://doi.org/10.1029/2019JD031598>, 2020.
- Bauer, S. E., Tsigaridis, K., Faluvegi, G., Kelley, M., Lo, K. K., Miller, R. L., Nazarenko, L., Schmidt, G. A., and Wu, J.: Historical (1850–2014) Aerosol Evolution and Role on Climate Forcing Using the GISS ModelE2.1 Contribution to CMIP6, *J. Adv. Model. Earth Syst.*, 12, e2019MS001 978, <https://doi.org/10.1029/2019MS001978>, 2020.
- Bellouin, N., Boucher, O., Haywood, J., and Reddy, M. S.: Global estimate of aerosol direct radiative forcing from satellite measurements, *Nature*, 438, 1138–1141, <https://doi.org/10.1038/nature04348>, 2005.
- Bellouin, N., Davies, W. H., Shine, K. P., Quaas, J., Mülmenstädt, J., Forster, P. M., Smith, C., Lee, L., Regayre, L., Brasseur, G., Sudarchikova, N., Bouarar, I., Boucher, O., and Myhre, G.: Radiative forcing of climate change from the Copernicus reanalysis of atmospheric composition, *Earth Syst. Sci. Data*, 12, 1649–1677, <https://doi.org/10.5194/essd-12-1649-2020>, 2020a.
- Bellouin, N., Quaas, J., Gryspeerdt, E., Kinne, S., Stier, P., Watson-Parris, D., Boucher, O., Carslaw, K., Christensen, M., Daniau, A.-L., Dufresne, J.-L., Feingold, G., Fiedler, S., Forster, P., Gettelman, A., Haywood, J. M., Malavelle, F., Lohmann, U., Mauritsen, T., McCoy, D., Myhre, G., Mülmenstädt, J., Neubauer, D., Possner, A., Rugenstein, M., Sato, Y., Schulz, M., Schwartz, S. E., Sourdeval,

- O., Storelvmo, T., Toll, V., Winker, D., and Stevens, B.: Bounding global aerosol radiative forcing of climate change, *Rev. Geophys.*, 58, e2019RG000660, <https://doi.org/10.1029/2019RG000660>, 2020b.
- 415 Benas, N., Meirink, J. F., Karlsson, K.-G., Stengel, M., and Stammes, P.: Satellite observations of aerosols and clouds over southern China from 2006 to 2015: analysis of changes and possible interaction mechanisms, *Atmos. Chem. Phys.*, 20, 457–474, <https://doi.org/10.5194/acp-20-457-2020>, 2020.
- Bennartz, R., Fan, J., Rausch, J., Leung, L. R., and Heidinger, A. K.: Pollution from China increases cloud droplet number, suppresses rain over the East China Sea, *Geophys. Res. Lett.*, 38, L09704, <https://doi.org/10.1029/2011GL047235>, 2011.
- 420 Boer, M. M., Resco de Dios, V., and Bradstock, R. A.: Unprecedented burn area of Australian mega forest fires, *Nat. Clim. Chang.*, 10, 171–172, <https://doi.org/10.1038/s41558-020-0716>, 2020.
- Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S., Sherwood, S., Stevens, B., and Zhang, X.: Clouds and Aerosols, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Stocker, T., Qin, D., Plattner, G.-K., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P., chap. 7, pp. 571–658, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/CBO9781107415324.016>, 2013.
- 425 Boucher, O., Servonnat, J., Albright, A. L., Aumont, O., Balkanski, Y., Bastrikov, V., Bekki, S., Bonnet, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Caubel, A., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., D’Andrea, F., Davini, P., de Lavergne, C., Denvil, S., Deshayes, J., Devilliers, M., Ducharne, A., Dufresne, J.-L., Dupont, E., Éthé, C., Fairhead, L., Falletti, L., Flavoni, S., Foujols, M.-A., Gardoll, S., Gastineau, G., Ghattas, J., Grandpeix, J.-Y., Guenet, B., Guez, Lionel, E., Guilyardi, E., Guimberteau, M., Hauglustaine, D., Hourdin, F., Idelkadi, A., Joussaume, S., Kageyama, M., Khodri, M., Krinner, G., Lebas, N., Levvasseur, G., Lévy, C., Li, L., Lott, F., Lurton, T., Luysaert, S., Madec, G., Madeleine, J.-B., Maignan, F., Marchand, M., Marti, O., Mellul, L., Meurdesoif, Y., Mignot, J., Musat, I., Ottlé, C., Peylin, P., Planton, Y., Polcher, J., Rio, C., Rochetin, N., Rousset, C., Sepulchre, P., Sima, A., Swingedouw, D., Thiéblemont, R., Traore, A. K., Vancoppenolle, M., Vial, J., Vialard, J., Viovy, N., and Vuichard, N.: Presentation and Evaluation of the IPSL-CM6A-LR Climate Model, *J. Adv. Model. Earth Syst.*, 12, e2019MS002010, <https://doi.org/https://doi.org/10.1029/2019MS002010>, 2020.
- 430 Brasseur, G. P. and Roeckner, E.: Impact of improved air quality on the future evolution of climate, *Geophys. Res. Lett.*, 32, L23704, <https://doi.org/10.1029/2005GL023902>, 2005.
- Carn, S., Fioletov, V., McLinden, C. A., Li, C., and Krotkov, N. A.: A decade of global volcanic SO₂ emissions measured from space, *Sci. Rep.*, 7, 44095, <https://doi.org/10.1038/srep44095>, 2017.
- Cermak, J., Wild, M., Knutti, R., Mishchenko, M., and Heidinger, K.: Consistency between global satellite-derived aerosol and cloud data sets on recent brightening trends, *Geophys. Res. Lett.*, 37, L21704, <https://doi.org/10.1029/2010GL044632>, 2010.
- 440 Chen, Y., Haywood, J., Wang, Y., Malavelle, F., Jordan, G., Partridge, D., Fieldsend, J., Johannes De L., Schmidt, A., Cho, N., Oreopoulos, L., Platnick, S., Grosvenor, D., Field, P., and Lohmann, U.: Machine learning reveals climate forcing from aerosols is dominated by increased cloud cover, *Nat. Geosci.*, 15, 609–614, <https://doi.org/10.1038/s41561-022-00991-6>, 2022.
- Cherian, R. and Quaas, J.: Trends in AOD, clouds and cloud radiative effects in satellite data and CMIP5 and CMIP6 model simulations over aerosol source regions, *Geophys. Res. Lett.*, 47, e2020GL087132, <https://doi.org/10.1029/2020GL087132>, 2020.
- 445 Cherian, R., Quaas, J., Salzmann, M., and Wild, M.: Pollution trends over Europe constrain global aerosol forcing as simulated by climate models, *Geophys. Res. Lett.*, 41, 2176–2181, <https://doi.org/10.1002/2013GL058715>, 2014.
- Chin, M., Diehl, T., Tan, Q., Prospero, J. M., Kahn, R. A., Remer, L. A., Yu, H., Sayer, A. M., Bian, H., Geogdzhayev, I. V., Holben, B. N., Howell, S. G., Huebert, B. J., Hsu, N. C., Kim, D., Kucsera, T. L., Levy, R. C., Mishchenko, M. I., Pan, X., Quinn, P. K., Schuster,

- 450 G. L., Streets, D. G., Strode, S. A., Torres, O., and Zhao, X.-P.: Multi-decadal aerosol variations from 1980 to 2009: a perspective from observations and a global model, *Atmos. Chem. Phys.*, 14, 3657–3690, <https://doi.org/10.5194/acp-14-3657-2014>, 2014.
- Christensen, M. W., Jones, W. K., and Stier, P.: Aerosols enhance cloud lifetime and brightness along the stratus-to-cumulus transition, *Proc. Nat. Acad. Sci. USA*, 117, 17 591–17 598, <https://doi.org/10.1073/pnas.1921231117>, 2020.
- Chýlek, P. and Coakley, J. A.: Aerosols and Climate, *Science*, 183, 75–77, <https://doi.org/10.1126/science.183.4120.75>, 1974.
- 455 Cohen, A., Brauer, M., Burnett, R., Anderson, H. R., Frostad, J., Estep, K., Balakrishnan, K., Brunekreef, B., Dandona, L., Dandona, R., Feigin, V., Freedman, G., Hubbell, B., Jobling, A., Kan, H., Knibbs, L., Liu, Y., Martin, R., Morawska, L., Pope III, C. A., Shin, H., Straif, K., Shaddick, G., Thomas, M., van Dingenen, R., van Donkelaar, A., Vos, T., Murray, C. J. L., and Forouzanfar, M. H.: Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015, *The Lancet*, 389, 1907–1918, [https://doi.org/10.1016/S0140-6736\(17\)30505-6](https://doi.org/10.1016/S0140-6736(17)30505-6), 2017.
- 460 Collaud Coen, M. C., Andrews, E., Alastuey, A., Arsov, T., Backman, J., Brem, B., Bukowiecki, N., Couret, C., Eleftheriadis, K., Flentje, H., Fiebig, M., Gysel-Beer, M., Hand, J., Hoffer, A., Hooda, R., Hueglin, C., Joubert, W., Keywood, M., Kim, J., Kim, S.-W., Labuschagne, C., Lin, N.-H., Lin, Y., Lund Myhre, C., Luoma, K., Lyamani, H., Marinoni, A., Mayol-Bracero, O. L., Mihalopoulos, N., Pandolfi, M., Prats, N., Prenni, A. J., Putaud, J.-P., Ries, L., Reisen, F., Sellegri, K., Sharma, S., Sheridan, P., Sherman, J. P., Sun, J., Titos, G., Torres, E., Tuch, T., Weller, R., Wiedensohler, A., Zieger, P., and Laj, P.: Multidecadal trend analysis of aerosol radiative properties at a global
465 scale, *Atmos. Chem. Phys.*, 20, 8867–8908, <https://doi.org/10.5194/acp-20-8867-2020>, 2020.
- Collins, W. J., Lamarque, J.-F., Schulz, M., Boucher, O., Eyring, V., Hegglin, M. I., Maycock, A., Myhre, G., Prather, M., Shindell, D., and Smith, S. J.: AerChemMIP: quantifying the effects of chemistry and aerosols in CMIP6, *Geosci. Model Dev.*, 10, 585–607, <https://doi.org/10.5194/gmd-10-585-2017>, 2017.
- Costa-Surós, M., Sourdeval, O., Acquistapace, C., Baars, H., Carbajal Henken, C., Genz, C., Hesemann, J., Jimenez, C., König, M.,
470 Kretzschmar, J., Madenach, N., Meyer, C. I., Schrödner, R., Seifert, P., Senf, F., Brueck, M., Cioni, G., Engels, J. F., Fieg, K., Gorges, K., Heinze, R., Siligam, P. K., Burkhardt, U., Crewell, S., Hoose, C., Seifert, A., Tegen, I., and Quaas, J.: Detection and attribution of aerosol-cloud interactions in large-domain large-eddy simulations with ICON, *Atmos. Chem. Phys. Discuss*, in revision, <https://doi.org/10.5194/acp-2019-850>, 2019.
- Crippa, M., Janssens-Maenhout, G., Dentener, F., Guizzardi, D., Sindelarova, K., Muntean, M., Van Dingenen, R., and Granier, C.: Forty years
475 of improvements in European air quality: regional policy-industry interactions with global impacts, *Atmos. Chem. Phys.*, 16, 3825–3841, <https://doi.org/10.5194/acp-16-3825-2016>, 2016.
- Dahutia, P., Pathak, B., and Bhuyan, P. K.: Aerosols characteristics, trends and their climatic implications over Northeast India and adjoining South Asia, *Int. J. Climatol.*, 38, 1234–1256, <https://doi.org/10.1002/joc.5240>, 2018.
- Doerr, S., Santín, C., and Santín Nuno, C.: Global trends in wildfire and its impacts: perceptions versus realities in a changing world, *Phil. Transact. Royal Soc. B: Biol. Sci.*, 371, 20150 345, <https://doi.org/10.1098/rstb.2015.0345>, 2016.
- 480 Dufresne, J.-L., Quaas, J., Boucher, O., Denvil, S., and Fairhead, L.: Contrast of the climate effects of anthropogenic sulfate aerosols between the 20th and 21st century, *Geophys. Res. Lett.*, 32, L21 703, <https://doi.org/10.1029/2005GL023619>, 2005.
- Ekman, A., Kjellström, E., Hansson, H.-C., Riipinen, I., Salter, M., Pandis, S., Klimont, Z., Martins, H., Cordeiro, A., Berntsen, T. K., and Jakobsson, I.: Is there a conflict between the clean air goals of the European Green Deal and climate neutrality?, <https://forces-project.eu/publications/policy-brief/>, 2020.
485
- Ekman, A. M. L.: Do sophisticated parameterizations of aerosol-cloud interactions in CMIP5 models improve the representation of recent observed temperature trends?, *J. Geophys. Res. Atmos.*, 119, 817–832, <https://doi.org/10.1002/2013JD020511>, 2014.

- Elguindi, N., Granier, C., Stavrakou, T., Darras, S., Bauwens, M., Cao, H., Chen, C., Denier van der Gon, H. A. C., Dubovik, O., Fu, T. M., Henze, D. K., Jiang, Z., Keita, S., Kuenen, J. J. P., Kurokawa, J., Liousse, C., Miyazaki, K., Müller, J.-F., Qu, Z., Solmon, F., and Zheng, B.: Intercomparison of Magnitudes and Trends in Anthropogenic Surface Emissions From Bottom-Up Inventories, Top-Down Estimates, and Emission Scenarios, *Earth's Future*, 8, e2020EF001 520, <https://doi.org/https://doi.org/10.1029/2020EF001520>, 2020.
- 490 Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geosci. Model Dev.*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- 495 Fan, X., Xia, X., and Chen, H.: Can MODIS detect trends in aerosol optical depth over land?, *Adv. Atmos. Sci.*, 35, 135–145, <https://doi.org/10.1007/s00376-017-7017-2>, 2018.
- Fiedler, S., Wyser, K., Rogelj, J., and van Noije, T.: Radiative effects of reduced aerosol emissions during the COVID-19 pandemic and the future recovery, *Atmos. Res.*, 264, 105 866, <https://doi.org/10.1016/j.atmosres.2021>, 2021.
- Filonchyk, M., Yan, H., Zhang, Z., Yang, S., Li, W., and Li, Y.: Combined use of satellite and surface observations to study aerosol optical depth in different regions of China, *Sci. Rep.*, 9, 6174, <https://doi.org/10.1038/s41598-019-42466-6>, 2019.
- 500 Forster, P., Forster, H., Evans, M., Gidden, M. J., Jones, C. D., Keller, C. A., Lamboll, R. D., Le Quéré, C., Rogelj, J., Rosen, D., Schleussner, C.-F., Richardson, T. B., Smith, C. J., and Turnock, S. T.: Current and future global climate impacts resulting from COVID-19, *Nat. Clim. Chang.*, 10, 913–919, <https://doi.org/10.1038/s41558-020-0883-0>, 2020.
- Forster, P., Storelvmo, T., Armour, K., Collins, W., Dufresne, J.-L., Frame, D., Lunt, D., Mauritsen, T., Palmer, M., Watanabe, M., Wild, M., and Zhang, H.: The Earth's Energy Budget, Climate Feedbacks, and Climate Sensitivity, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., chap. 7, pp. 923–1054, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896.009>, 2021.
- 505 Garay, M. J., Kalashnikova, O. V., and Bull, M. A.: Development and assessment of a higher-spatial-resolution (4.4 km) MISR aerosol optical depth product using AERONET-DRAGON data, *Atmos. Chem. Phys.*, 17, 5095–5106, <https://doi.org/10.5194/acp-17-5095-2017>, 2017.
- Geoffroy, O., Saint-Martin, D., Bellon, G., Voldoire, A., Olivié, D. J. L., and Tytéca, S.: Transient Climate Response in a Two-Layer Energy-Balance Model. Part II: Representation of the Efficacy of Deep-Ocean Heat Uptake and Validation for CMIP5 AOGCMs, *J. Climate*, 26, 1859 – 1876, <https://doi.org/10.1175/JCLI-D-12-00196.1>, 2013.
- 515 Georgoulas, A. K., Alexandri, G., Kourtidis, K. A., Lelieveld, J., Zanis, P., Pöschl, U., Levy, R., Amiridis, V., Marinou, E., and Tsikerdekis, A.: Spatiotemporal variability and contribution of different aerosol types to the aerosol optical depth over the Eastern Mediterranean, *Atmos. Chem. Phys.*, 16, 13 853–13 884, <https://doi.org/10.5194/acp-16-13853-2016>, 2016.
- Gettelman, A., Lamboll, R., Bardeen, C. G., Forster, P. M., and Watson-Parris, D.: Climate impacts of COVID-19 induced emission changes, *Geophys. Res. Lett.*, 48, e2020GL091 805, <https://doi.org/10.1029/2020GL091805>, 2021.
- 520 Giles, D. M., Sinyuk, A., Sorokin, M. G., Schafer, J. S., Smirnov, A., Slutsker, I., Eck, T. F., Holben, B. N., Lewis, J. R., Campbell, J. R., Welton, E. J., Korkin, S. V., and Lyapustin, A. I.: Advancements in the Aerosol Robotic Network (AERONET) Version 3 database – automated near-real-time quality control algorithm with improved cloud screening for Sun photometer aerosol optical depth (AOD) measurements, *Atmos. Meas. Tech.*, 12, 169–209, <https://doi.org/10.5194/amt-12-169-2019>, 2019.
- Granier, C., Bessagnet, B., Bond, T., D'Angiola, A., van der Gon, H. D., Frost, G. J., Heil, A., Kaiser, J. W., Kinne, S., Klimont, Z., Kloster, S., Lamarque, J.-F., Liousse, C., Masui, T., Meleux, F., Mieville, A., Ohara, T., Raut, J.-C., Riahi, K., Schultz, M. G., Smith, S. J.,
- 525

- Thompson, A., van Aardenne, J., van der Werf, G. R., and van Vuuren, D. P.: Evolution of anthropogenic and biomass burning emissions of air pollutants at global and regional scales during the 1980–2010 period, *Climatic Change*, 109, 163, <https://doi.org/10.1007/s10584-011-0154-1>, 2011.
- Grosvenor, D. P., Sourdeval, O., Zuidema, P., Ackerman, A., Alexandrov, M. D., Bennartz, R., Boers, R., Cairns, B., Chiu, C., Christensen, M., Deneke, H., Diamond, M., Feingold, G., Fridlind, A., Hünerbein, A., Knist, C., Kollias, P., Marshak, A., McCoy, D., Merk, D., Painemal, D., Rausch, J., Rosenfeld, D., Russchenberg, H., Seifert, P., Sinclair, K., Stier, P., B. van D., Wendisch, M., Werner, F., Wood, R., Zhang, Z., and Quaas, J.: Remote sensing of cloud droplet number concentration in warm clouds: A review of the current state of knowledge and perspectives, *Rev. Geophys.*, 56, 409–453, <https://doi.org/10.1029/2017RG000593>, 2018.
- Gryspeerdt, E., Quaas, J., and Bellouin, N.: Constraining the aerosol influence on cloud fraction, *J. Geophys. Res.*, 121, 3566–3583, <https://doi.org/10.1002/2015JD023744>, <http://onlinelibrary.wiley.com/doi/10.1002/2015JD023744/abstract>, ulei, 2016.
- Gryspeerdt, E., Smith, T. W. P., O’Keeffe, E., Christensen, M. W., and Goldsworth, F. W.: The impact of ship emission controls recorded by cloud properties, *Geophys. Res. Lett.*, 46, 12 547–12 555, <https://doi.org/10.1029/2019GL084700>, 2019.
- Grzegorski, M., Poli, G., Cacciari, A., Jafariserajehlou, S., Holdak, A., Lang, R., Vazquez-Navarro, M., Munro, R., and Fougnie, B.: Multi-Sensor Retrieval of Aerosol Optical Properties for Near-Real-Time Applications Using the Metop Series of Satellites: Concept, Detailed Description, and First Validation, *Remote Sensing*, 14, 85, <https://doi.org/10.3390/rs14010085>, 2021.
- Gulev, S., Thorne, P., Ahn, J., Dentener, F., Domingues, C., Gerland, S., Gong, D., Kaufman, D., Nnamchi, H., Quaas, J., Rivera, J., Sathyendranath, S., Smith, S., Trewin, B., von Schuckmann, K., and Vose, R.: Changing State of the Climate System, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., chap. 2, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2021.
- Haghighatnasab, M., Kretschmar, J., Block, K., and Quaas, J.: Impact of Holuhraun volcano aerosols on clouds in cloud-system resolving simulations, *Atmos. Chem. Phys. Discuss.*, in review, <https://doi.org/10.5194/acp-2022-38>, <https://acp.copernicus.org/preprints/acp-2022-38/>, ulei, 2022.
- Hajima, T., Watanabe, M., Yamamoto, A., Tatebe, H., Noguchi, M. A., Abe, M., Ohgaito, R., Ito, A., Yamazaki, D., Okajima, H., Ito, A., Takata, K., Ogochi, K., Watanabe, S., and Kawamiya, M.: Development of the MIROC-ES2L Earth system model and the evaluation of biogeochemical processes and feedbacks, *Geosci. Model Dev.*, 13, 2197–2244, <https://doi.org/10.5194/gmd-13-2197-2020>, 2020.
- Hammer, M. S., Martin, R. V., Li, C., Torres, O., Manning, M., and Boys, B. L.: Insight into global trends in aerosol composition from 2005 to 2015 inferred from the OMI Ultraviolet Aerosol Index, *Atmos. Chem. Phys.*, 18, 8097–8112, <https://doi.org/10.5194/acp-18-8097-2018>, 2018.
- Hatzianastassiou, N., Ioannidis, E., Korras-Carraca, M.-B., Gavrouzou, M., Papadimas, C. D., Matsoukas, C., Benas, N., Fotiadi, A., Wild, M., and Vardavas, I.: Global Dimming and Brightening Features during the First Decade of the 21st Century, *Atmosphere*, 11, 308, <https://doi.org/10.3390/atmos11030308>, 2020.
- Heinold, B., Baars, H., Barja, B., Christensen, M., Kubin, A., Ohneiser, K., Schepanski, K., Schutgens, N., Senf, F., Schrödner, R., Villanueva, D., and Tegen, I.: Important role of stratospheric injection height for the distribution and radiative forcing of smoke aerosol from the 2019/2020 Australian wildfires, *Atmos. Chem. Phys. Discuss.*, in review, <https://doi.org/10.5194/acp-2021-862>, 2021.
- Held, I. M., Guo, H., Adcroft, A., Dunne, J. P., Horowitz, L. W., Krasting, J., Shevliakova, E., Winton, M., Zhao, M., Bushuk, M., Wittenberg, A. T., Wyman, B., Xiang, B., Zhang, R., Anderson, W., Balaji, V., Donner, L., Dunne, K., Durachta, J., Gauthier, P. P. G., Ginoux, P., Golaz,

- J.-C., Griffies, S. M., Hallberg, R., Harris, L., Harrison, M., Hurlin, W., John, J., Lin, P., Lin, S.-J., Malyshev, S., Menzel, R., Milly, P.
565 C. D., Ming, Y., Naik, V., Paynter, D., Paulot, F., Rammawamy, V., Reichl, B., Robinson, T., Rosati, A., Seman, C., Silvers, L. G.,
Underwood, S., and Zadeh, N.: Structure and Performance of GFDL's CM4.0 Climate Model, *J. Adv. Model. Earth Syst.*, 11, 3691–3727,
<https://doi.org/10.1029/2019MS001829>, 2019.
- Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M.,
Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J.-I., Li, M., Liu, L., Lu, Z., Moura, M. C. P., O'Rourke, P. R., and Zhang, Q.:
570 Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS),
Geosci. Model Dev., 11, 369–408, <https://doi.org/10.5194/gmd-11-369-2018>, 2018.
- Holben, B., Eck, T., Slutsker, I., Tanré, D., Buis, J., Setzer, A., Vermote, E., Reagan, J., Kaufman, Y., Nakajima, T., Lavenu, F., Jankowiak, I.,
and Smirnov, A.: AERONET—A Federated Instrument Network and Data Archive for Aerosol Characterization, *Remote Sens. Environ.*,
66, 1–16, [https://doi.org/10.1016/S0034-4257\(98\)00031-5](https://doi.org/10.1016/S0034-4257(98)00031-5), 1998.
- 575 Holben, B. N., Tanré, D., Smirnov, A., Eck, T. F., Slutsker, I., Abuhassan, N., Newcomb, W. W., Schafer, J. S., Chatenet, B., Lavenu, F.,
Kaufman, Y. J., Castle, J. V., Setzer, A., Markham, B., Clark, D., Frouin, R., Halthore, R., Karneli, A., O'Neill, N. T., Pietras, C., Pinker,
R. T., Voss, K., and Zibordi, G.: An emerging ground-based aerosol climatology: Aerosol optical depth from AERONET, *J. Geophys.
Res.*, 106, 12 067—12 097, <https://doi.org/10.1029/2001JD900014>, 2001.
- Hong, C., Zhang, Q., Zhang, Y., Davis, S. J., Zhang, X., Tong, D., Guan, D., Liu, Z., and He, K.: Weakening aerosol direct radiative effects
580 mitigate climate penalty on Chinese air quality, *Nat. Clim. Chang.*, 10, 845–850, <https://doi.org/10.1038/s41558-020-0840-y>, 2020.
- IMO: MEPC.176(58) Amendments to the Annex of the Protocol of 1997 to amend the International Convention for the Prevention of
Pollution from Ships, 1973, as modified by the Protocol of 1978 relating thereto (Revised MARPOL Annex VI), [https://wwwcdn.imo.org/
localresources/en/OurWork/Environment/Documents/176\(58\).pdf](https://wwwcdn.imo.org/localresources/en/OurWork/Environment/Documents/176(58).pdf), 2008.
- IMO: Annex 15, Resolution MEPC.321(74) 2019 Guidelines for Port State Control Under MARPOL Annex VI Chapter 3, [https://wwwcdn.
585 imo.org/localresources/en/OurWork/Environment/Documents/MEPC.321%2874%29.pdf](https://wwwcdn.imo.org/localresources/en/OurWork/Environment/Documents/MEPC.321%2874%29.pdf), 2019.
- Jenkins, S., Grainger, R., Povey, A., Gettelman, A., Stier, P., and Allen, M.: Is Anthropogenic Global Warming Accelerating?, *J. Climate*, in
press, 2022.
- Jia, H., Quaas, J., Gryspeerdt, E., Böhm, C., and Sourdeval, O.: Addressing the difficulties in quantifying the Twomey effect for marine warm
clouds from multi-sensor satellite observations and reanalysis, *Atmos. Chem. Phys. Discuss.*, in review, [https://doi.org/10.5194/acp-2021-
590 999](https://doi.org/10.5194/acp-2021-999), 2022.
- Jongeward, A. R., Li, Z., He, H., and Xiong, X.: Natural and Anthropogenic Aerosol Trends from Satellite and Surface Observations and
Model Simulations over the North Atlantic Ocean from 2002 to 2012, *J. Atmos. Sci.*, 73, 4469–4485, [https://doi.org/10.1175/JAS-D-15-
0308.1](https://doi.org/10.1175/JAS-D-15-0308.1), 2016.
- Kanaya, Y., Yamaji, K., Miyakawa, T., Taketani, F., Zhu, C., Choi, Y., Komazaki, Y., Ikeda, K., Kondo, Y., and Klimont, Z.: Rapid reduction in
595 black carbon emissions from China: evidence from 2009–2019 observations on Fukue Island, Japan, *Atmos. Chem. Phys.*, 20, 6339–6356,
<https://doi.org/10.5194/acp-20-6339-2020>, 2020.
- Kaufman, Y. J., Boucher, O., Tanré, D., Chin, M., Remer, L. A., and Takemura, T.: Aerosol anthropogenic component estimated from satellite
data, *Geophys. Res. Lett.*, 32, L17 804, <https://doi.org/10.1029/2005GL023125>, 2005.
- Kelley, M., Schmidt, G. A., Nazarenko, L. S., Bauer, S. E., Ruedy, R., Russell, G. L., Ackerman, A. S., Aleinov, I., Bauer, M., Bleck, R.,
600 Canuto, V., Cesana, G., Cheng, Y., Clune, T. L., Cook, B. I., Cruz, C. A., Del Genio, A. D., Elsaesser, G. S., Faluvegi, G., Kiang, N. Y.,
Kim, D., Lacis, A. A., Leboissetier, A., LeGrande, A. N., Lo, K. K., Marshall, J., Matthews, E. E., McDermid, S., Mezuman, K., Miller,

- R. L., Murray, L. T., Oinas, V., Orbe, C., García-Pando, C. P., Perlwitz, J. P., Puma, M. J., Rind, D., Romanou, A., Shindell, D. T., Sun, S., Tausnev, N., Tsigaridis, K., Tselioudis, G., Weng, E., Wu, J., and Yao, M.-S.: GISS-E2.1: Configurations and Climatology, *J. Adv. Model. Earth Syst.*, 12, e2019MS002 025, <https://doi.org/https://doi.org/10.1029/2019MS002025>, 2020.
- 605 Kinne, S.: Aerosol radiative effects with MACv2, *Atmos. Chem. Phys.*, 19, 10 919–10 959, <https://doi.org/10.5194/acp-19-10919-2019>, 2019.
- Klimont, Z., Smith, S. J., and Cofala, J.: The last decade of global anthropogenic sulfur dioxide: 2000–2011 emissions, *Environ. Res. Lett.*, 8, 014 003, <https://doi.org/10.1088/1748-9326/8/1/014003>, 2013.
- Klimont, Z., Kupiainen, K., Heyes, C., Purohit, P., Cofala, J., Rafaj, P., Borcken-Kleefeld, J., and Schöpp, W.: Global anthropogenic emissions of particulate matter including black carbon, *Atmos. Chem. Phys.*, 17, 8681–8723, <https://doi.org/10.5194/acp-17-8681-2017>, <https://acp.copernicus.org/articles/17/8681/2017/>, 2017.
- 610 Kramer, R. J., He, H., Soden, B. J., Oreopoulos, L., Myhre, G., Forster, P. M., and Smith, C. J.: Observational evidence of increasing global radiative forcing, *Geophys. Res. Lett.*, 48, e2020GL091 585, <https://doi.org/10.1029/2020GL091585>, 2021.
- Kretzschmar, J., Salzmann, M., Mülmenstädt, J., Boucher, O., and Quaas, J.: Comment on “Rethinking the lower bound on aerosol radiative forcing”, *J. Clim.*, 30, 6579–6584, <https://doi.org/10.1175/JCLI-D-16-0668.1>, 2017.
- 615 Krüger, O. and Graßl, H.: The indirect aerosol effect over Europe, *Geophys. Res. Lett.*, 29, 1925, <https://doi.org/10.1029/2001GL014081>, 2002.
- Lelieveld, J., Evans, J. S., Fnais, M., Giannadaki, D., and Pozzer, A.: The contribution of outdoor air pollution sources to premature mortality on a global scale, *Nature*, 525, 367–371, <https://doi.org/10.1038/nature15371>, 2015.
- Levy, R. C., Mattoo, S., Munchak, L. A., Remer, L. A., Sayer, A. M., Patadia, F., and Hsu, N. C.: The Collection 6 MODIS aerosol products over land and ocean, *Atmos. Meas. Tech.*, 6, 2989–3034, <https://doi.org/10.5194/amt-6-2989-2013>, <https://www.atmos-meas-tech.net/6/2989/2013/>, 2013.
- 620 Li, J., Carlson, B. E., Dubovik, O., and Lacis, A. A.: Recent trends in aerosol optical properties derived from AERONET measurements, *Atmos. Chem. Phys.*, 14, 12 271–12 289, <https://doi.org/10.5194/acp-14-12271-2014>, 2014.
- Loeb, N. G., Doelling, D. R., Wang, H., Su, W., Nguyen, C., Corbett, J. G., Liang, L., Mitrescu, C., Rose, F. G., and Kato, S.: Clouds and the Earth’s Radiant Energy System (CERES) Energy Balanced and Filled (EBAF) Top-of-Atmosphere (TOA) Edition-4.0 Data Product, *J. Climate*, 31, 895–918, <https://doi.org/10.1175/JCLI-D-17-0208.1>, 2018a.
- 625 Loeb, N. G., Thorsen, T. J., Norris, J. R., Wang, H., and Su, W.: Changes in Earth’s energy budget during and after the “Pause” in global warming: An observational perspective, *Climate*, 6, <https://doi.org/10.3390/cli6030062>, 2018b.
- Loeb, N. G., Johnson, G. C., Thorsen, T. J., Lyman, J. M., Rose, F. G., and Kato, S.: Satellite and ocean data reveal marked increase in Earth’s heating rate, *Geophys. Res. Lett.*, 48, e2021GL093 047, <https://doi.org/10.1029/2021GL093047>, 2021a.
- 630 Loeb, N. G., Su, W., Bellouin, N., and Ming, Y.: Changes in Clear-Sky Shortwave Aerosol Direct Radiative Effects Since 2002, *J. Geophys. Res.*, 126, e2020JD034 090, <https://doi.org/10.1029/2020JD034090>, 2021b.
- Ma, X., Jia, H., Yu, F., and Quaas, J.: Opposite aerosol index-cloud droplet effective radius correlations over major industrial regions and their adjacent oceans, *Geophys. Res. Lett.*, 45, 5771–5778, <https://doi.org/10.1029/2018GL077562>, 2018.
- 635 Ma, Z., Liu, R., Liu, Y., and Bi, J.: Effects of air pollution control policies on PM2.5 pollution improvement in China from 2005 to 2017: a satellite-based perspective, *Atmos. Chem. Phys.*, 19, 6861–6877, <https://doi.org/10.5194/acp-19-6861-2019>, 2019.
- Malavelle, F. F., Haywood, J. M., Jones, A., Gettelman, A., Clarisse, L., Bauduin, S., Allan, R. P., Karset, I. H. H., Kristjánsson, J. E., Oreopoulos, L., Cho, N., Lee, D., Bellouin, N., Boucher, O., Grosvenor, D. P., Carslaw, K. S., Dhomse, S., Mann, G. W., Schmidt, A., Coe, H., Hartley, M. E., Dalvi, M., Hill, A. A., Johnson, B. T., Johnson, C. E., Knight, J. R., O’Connor, F. M., Partridge, D. G., Stier, P.,

- 640 Myhre, G., Platnick, S., Stephens, G. L., Takahashi, H., and Thordarson, T.: Strong constraints on aerosol-cloud interactions from volcanic eruptions, *Nature*, 546, 485–491, <https://doi.org/10.1038/nature22974>, 2017.
- Manara, V., Brunetti, M., Celozzi, A., Maugeri, M., Sanchez-Lorenzo, A., and Wild, M.: Detection of dimming/brightening in Italy from homogenized all-sky and clear-sky surface solar radiation records and underlying causes (1959–2013), 16, 11 145–11 161, <https://doi.org/10.5194/acp-16-11145-2016>, 2016.
- 645 McCoy, D. T., Bender, F. A.-M., Grosvenor, D. P., Mohrmann, J. K., Hartmann, D. L., Wood, R., and Field, P. R.: Predicting decadal trends in cloud droplet number concentration using reanalysis and satellite data, *Atmos. Chem. Phys.*, 18, 2035–2047, <https://doi.org/10.5194/acp-18-2035-2018>, 2018.
- McKenna, C., Maycock, A., Forster, P., Smith, C. J., and Tokarska, K. B.: Stringent mitigation substantially reduces risk of unprecedented near-term warming rates, *Nat. Clim. Chang.*, 11, 126–131, <https://doi.org/10.1038/s41558-020-00957-9>, 2021.
- 650 Mortier, A., Gliß, J., Schulz, M., Aas, W., Andrews, E., Bian, H., Chin, M., Ginoux, P., Hand, J., Holben, B., Zhang, H., Kipling, Z., Kirkevåg, A., Laj, P., Lurton, T., Myhre, G., Neubauer, D., Olivíé, D., von Salzen, K., Skeie, R. B., Takemura, T., and Tilmes, S.: Evaluation of climate model aerosol trends with ground-based observations over the last 2 decades – an AeroCom and CMIP6 analysis, *Atmos. Chem. Phys.*, 20, 13 355–13 378, <https://doi.org/10.5194/acp-20-13355-2020>, 2020.
- Mülmenstädt, J. and Feingold, G.: The Radiative Forcing of Aerosol–Cloud Interactions in Liquid Clouds: Wrestling and Embracing Uncer-
655 tainty, *Curr. Clim. Change Rep.*, 4, 23–40, <https://doi.org/10.1007/s40641-018-0089-y>, 2018.
- Murray-Watson, R. J. and Gryspeerdt, E.: Stability dependent increases in liquid water with droplet number in the Arctic, *Atmos. Chem. Phys. Discuss.*, in review, <https://doi.org/10.5194/acp-2021-861>, 2022.
- Myhre, G., Boucher, O., Bréon, F. M., Forster, P., and Shindell, D.: Declining uncertainty in transient climate response as CO₂ forcing dominates future climate change, *Nature Geosci.*, 8, 181–185, <https://doi.org/10.1038/ngeo2371>, 2015.
- 660 Myhre, G., Aas, W., Cherian, R., Collins, W., Faluvegi, G., Flanner, M., Forster, P., Hodnebrog, Ø., Klimont, Z., Lund, M. T., Mülmenstädt, J., Lund Myhre, C., Olivíé, D., Prather, M., Quaas, J., Samset, B. H., Schnell, J. L., Schulz, M., Shindell, D., Skeie, R., Takemura, T., and Tsyro, S.: Multi-model simulations of aerosol and ozone radiative forcing due to anthropogenic emission changes during the period 1990–2015, *Atmos. Chem. Phys.*, 17, 2709–2720, <https://doi.org/10.5194/acp-17-2709-2017>, 2017.
- Norris, J. R., Allen, R. J., Evan, A. T., Zelinka, M. D., O’Dell, C. W., and Klein, S. A.: Evidence for climate change in the satellite cloud
665 record, *Nature*, 536, 72–75, <https://doi.org/10.1038/nature18273>, 2016.
- O’Rourke, P., Smith, S. J., Mott, A. R., Ahsan, H., McDuffie, E. E., Crippa, M., Klimont, Z., McDonald, B., Wang, S., Nicholson, M. B., Hoesly, R. M., and Feng, L.: CEDS v_2021_04_21 Gridded emissions data, <https://doi.org/10.25584/PNNLDataHub/1779095>, 2021.
- Palmer, M.: Reconciling Estimates of Ocean Heating and Earth’s Radiation Budget, *Curr. Clim. Change Rep.*, 3, 78–86, <https://doi.org/10.1007/s40641-016-0053-7>, 2017.
- 670 Paulot, F., Paynter, D., Ginoux, P., Naik, V., and Horowitz, L. W.: Changes in the aerosol direct radiative forcing from 2001 to 2015: observational constraints and regional mechanisms, *Atmos. Chem. Phys.*, 18, 13 265–13 281, <https://doi.org/10.5194/acp-18-13265-2018>, 2018.
- Pfeyfroth, U., Sanchez-Lorenzo, A., Manara, V., Trentmann, J., and Hollmann, R.: Trends and variability of surface solar radiation in Europe based on surface- and satellite-based data records, *J. Geophys. Res.*, 123, 1735–1754, <https://doi.org/10.1002/2017JD027418>, 2018.
- 675 Pincus, R., Forster, P. M., and Stevens, B.: The Radiative Forcing Model Intercomparison Project (RFMIP): experimental protocol for CMIP6, *Geosci. Model Devel.*, 9, 3447–3460, <https://doi.org/10.5194/gmd-9-3447-2016>, 2016.

- Platnick, S., Meyer, K. G., King, M. D., Wind, G., Amarasinghe, N., Marchant, B., Arnold, G. T., Zhang, Z., Hubanks, P. A., Holz, R. E., Yang, P., Ridgway, W. L., and Riedi, J.: The MODIS Cloud Optical and Microphysical Products: Collection 6 Updates and Examples From Terra and Aqua, *IEEE Transact. Geosci. Remote Sens.*, 55, 502–525, <https://doi.org/10.1109/TGRS.2016.2610522>, 2017.
- 680 Quaas, J., Boucher, O., and Lohmann, U.: Constraining the total aerosol indirect effect in the LMDZ and ECHAM4 GCMs using MODIS satellite data, *Atmos. Chem. Phys.*, 6, 947–955, <https://doi.org/10.5194/acp-6-947-2006>, 2006.
- Quaas, J., Arola, A., Cairns, B., Christensen, M., Deneke, H., Ekman, A. M. L., Feingold, G., Fridlind, A., Gryspeerdt, E., Hasekamp, O., Li, Z., Lipponen, A., Ma, P.-L., Mülmenstädt, J., Nenes, A., Penner, J., Rosenfeld, D., Schrödner, R., Sinclair, K., Sourdeval, O., Stier, P., Tesche, M., van Diedenhoven, B., and Wendisch, M.: Constraining the Twomey effect from satellite observations: Issues and perspectives, *Atmos. Chem. Phys.*, 20, 15 079–15 099, <https://doi.org/10.5194/acp-20-15079-2020>, <https://acp.copernicus.org/articles/20/15079/2020/>, ulei, 2020.
- Raghuraman, S., Paynter, D., and Ramaswamy, V.: Anthropogenic forcing and response yield observed positive trend in Earth’s energy imbalance, *Nat. Commun.*, 12, 4577, <https://doi.org/10.1038/s41467-021-24544-4>, 2021.
- Rosenfeld, D., Zhu, Y., Wang, M., Zheng, Y., Goren, T., and Yu, S.: Aerosol-driven droplet concentrations dominate coverage and water of oceanic low-level clouds, *Science*, 364, <https://doi.org/10.1126/science.aay4194>, 2019.
- 690 Rotstajn, L. D., Collier, M. A., Shindell, D. T., and Boucher, O.: Why does aerosol forcing control historical global-mean surface temperature change in CMIP5 models?, *J. Climate*, 28, 6608–6625, <https://doi.org/10.1175/JCLI-D-14-00712.1>, 2015.
- Samset, B. H., Lund, M. T., Bollasina, M., Myhre, G., and Wilcox, L.: Emerging Asian aerosol patterns, *Nature Geoscience*, 12, 582–584, <https://doi.org/10.1038/s41561-019-0424-5>, 2019.
- 695 Seland, Ø., Bentsen, M., Olivić, D., Toniazzo, T., Gjermundsen, A., Graff, L. S., Debernard, J. B., Gupta, A. K., He, Y.-C., Kirkevåg, A., Schwinger, J., Tjiputra, J., Aas, K. S., Bethke, I., Fan, Y., Griesfeller, J., Grini, A., Guo, C., Ilicak, M., Karset, I. H. H., Landgren, O., Liakka, J., Moseid, K. O., Nummelin, A., Spensberger, C., Tang, H., Zhang, Z., Heinze, C., Iversen, T., and Schulz, M.: Overview of the Norwegian Earth System Model (NorESM2) and key climate response of CMIP6 DECK, historical, and scenario simulations, *Geosci. Model Devel.*, 13, 6165–6200, <https://doi.org/10.5194/gmd-13-6165-2020>, 2020.
- 700 Sherwood, S., Webb, M. J., Annan, J. D., Armour, K., Forster, P. M., Hargreaves, J. C., Hegerl, G., Klein, S. A., Marvel, K. D., Rohling, E. J., et al.: An assessment of Earth’s climate sensitivity using multiple lines of evidence, *Rev. Geophys.*, p. e2019RG000678, 2020.
- Shindell, D. and Smith, C.: Climate and air-quality benefits of a realistic phase-out of fossil fuels, *Nature*, 573, 408–411, <https://doi.org/10.1038/s41586-019-1554-z>, 2019.
- Sickles II, J. E. and Shadwick, D. S.: Air quality and atmospheric deposition in the eastern US: 20 years of change, *Atmos. Chem. Phys.*, 15, 705 173–197, <https://doi.org/10.5194/acp-15-173-2015>, 2015.
- Smith, C. J. and Forster, P. M.: Suppressed late-20th Century warming in CMIP6 models explained by forcing and feedbacks, *Geophys. Res. Lett.*, 48, e2021GL094 948, <https://doi.org/10.1029/2021GL094948>, 2021.
- Smith, C. J., Harris, G. R., Palmer, M. D., Bellouin, N., Collins, W., Myhre, G., Schulz, M., Golaz, J.-C., Ringer, M., Storelvmo, T., and Forster, P. M.: Energy budget constraints on the time history of aerosol forcing and climate sensitivity, *J. Geophys. Res. Atmos.*, p. e2020JD033622, <https://doi.org/10.1029/2020JD033622>, 2021a.
- 710 Smith, M., Cain, M., and Allen, M.: Further improvement of warming-equivalent emissions calculation, *npj Clim. Atmos. Sci.*, 4, 19, <https://doi.org/10.1038/s41612-021-00169-8>, 2021b.
- Smith, S. J., van Aardenne, J., Klimont, Z., Andres, R. J., Volke, A., and Delgado Arias, S.: Anthropogenic sulfur dioxide emissions: 1850–2005, *Atmos. Chem. Phys.*, 11, 1101–1116, <https://doi.org/10.5194/acp-11-1101-2011>, 2011.

- 715 Sogacheva, L., de Leeuw, G., Rodriguez, E., Kolmonen, P., Georgoulias, A. K., Alexandri, G., Kourtidis, K., Proestakis, E., Marinou, E., Amiridis, V., Xue, Y., and van der A, R. J.: Spatial and seasonal variations of aerosols over China from two decades of multi-satellite observations – Part 1: ATSR (1995–2011) and MODIS C6.1, *Atmos. Chem. Phys.*, 18, 11 389–11 407, <https://doi.org/10.5194/acp-18-11389-2018>, 2018.
- Solomon, S., Plattner, G.-K., Knutti, R., and Friedlingstein, P.: Irreversible climate change due to carbon dioxide emissions, *Proc. Nat. Acad. Sci. USA*, 106, 1704–1709, <https://doi.org/10.1073/pnas.0812721106>, 2009.
- 720 Stevens, B.: Rethinking the Lower Bound on Aerosol Radiative Forcing, *J. Climate*, 28, 4794–4819, <https://doi.org/10.1175/JCLI-D-14-00656.1>, 2015.
- Stier, P., Feichter, J., Roeckner, E., Kloster, S., and Esch, M.: The evolution of the global aerosol system in a transient climate simulation from 1860 to 2100, *Atmos. Chem. Phys.*, 6, 3059–3076, <https://doi.org/10.5194/acp-6-3059-2006>, 2006.
- 725 Stjern, C. W., Stohl, A., and Kristjánsson, J. E.: Have aerosols affected trends in visibility and precipitation in Europe?, *J. Geophys. Res.*, 116, D02 212, <https://doi.org/10.1029/2010JD014603>, 2011.
- Streets, D. G., Yan, F., Chin, M., Diehl, T., Mahowald, N., Schultz, M., Wild, M., Wu, Y., and Yu, C.: Anthropogenic and natural contributions to regional trends in aerosol optical depth, 1980–2006, *J. Geophys. Res.*, 114, D00D18, <https://doi.org/10.1029/2008JD011624>, 2009.
- Struthers, H., Ekman, A. M. L., Glantz, P., Iversen, T., Kirkevåg, A., Seland, O., Martensson, E. M., Noone, K., and Nilsson, E. D.: Climate-induced changes in sea salt aerosol number emissions: 1870 to 2100, *J. Geophys. Res.*, 118, 670–682, <https://doi.org/https://doi.org/10.1002/jgrd.50129>, 2013.
- 730 Swart, N. C., Cole, J. N. S., Kharin, V. V., Lazare, M., Scinocca, J. F., Gillett, N. P., Anstey, J., Arora, V., Christian, J. R., Hanna, S., Jiao, Y., Lee, W. G., Majaess, F., Saenko, O. A., Seiler, C., Seinen, C., Shao, A., Sigmund, M., Solheim, L., von Salzen, K., Yang, D., and Winter, B.: The Canadian Earth System Model version 5 (CanESM5.0.3), *Geosci. Model Dev.*, 12, 4823–4873, <https://doi.org/10.5194/gmd-12-4823-2019>, 2019.
- 735 Szopa, S., Naik, V., Adhikary, B., Artaxo, P., Bernsten, T., Collins, W., Fuzzi, S., Gallardo, L., Scharr, A. K., Klimont, Z., Liao, H., Unger, N., and Zanis, P.: Short-Lived Climate Forcers, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., chap. 6, pp. 817–922, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896.008>, 2021.
- 740 Toll, V., Christensen, M., Quaas, J., and Bellouin, N.: Weak average liquid-cloud-water response to anthropogenic aerosols, *Nature*, 572, 51–55, <https://doi.org/10.1038/s41586-019-1423-9>, 2019.
- Tørseth, K., Aas, W., Breivik, K., Fjæraa, A. M., Fiebig, M., Hjellbrekke, A. G., Lund Myhre, C., Solberg, S., and Yttri, K. E.: Introduction to the European Monitoring and Evaluation Programme (EMEP) and observed atmospheric composition change during 1972–2009, *Atmos. Chem. Phys.*, 12, 5447–5481, <https://doi.org/10.5194/acp-12-5447-2012>, 2012.
- 745 Twomey, S.: Pollution and the planetary albedo, *Atmos. Environ.*, 8, 1251–1256, [https://doi.org/10.1016/0004-6981\(74\)90004-3](https://doi.org/10.1016/0004-6981(74)90004-3), 1974.
- van Marle, M. J. E., Kloster, S., Magi, B. I., Marlon, J. R., Daniiau, A.-L., Field, R. D., Arneeth, A., Forrest, M., Hantson, S., Kehrwald, N. M., Knorr, W., Lasslop, G., Li, F., Mangeon, S., Yue, C., Kaiser, J. W., and van der Werf, G. R.: Historic global biomass burning emissions for CMIP6 (BB4CMIP) based on merging satellite observations with proxies and fire models (1750–2015), *Geosci. Model Dev.*, 10, 3329–3357, <https://doi.org/10.5194/gmd-10-3329-2017>, 2017.
- 750

- van Oldenborgh, G. J., Krikken, F., Lewis, S., Leach, N. J., Lehner, F., Saunders, K. R., van Weele, M., Haustein, K., Li, S., Wallom, D., Sparrow, S., Arrighi, J., Singh, R. K., van Aalst, M. K., Philip, S. Y., Vautard, R., and Otto, F. E. L.: Attribution of the Australian bushfire risk to anthropogenic climate change, *Nat. Hazards Earth Syst. Sci.*, 21, 941–960, <https://doi.org/10.5194/nhess-21-941-2021>, 2021.
- 755 Vestreng, V., Myhre, G., Fagerli, H., Reis, S., and Tarrasón, L.: Twenty-five years of continuous sulphur dioxide emission reduction in Europe, *Atmos. Chem. Phys.*, 7, 3663–3681, <https://doi.org/10.5194/acp-7-3663-2007>, 2007.
- von Schuckmann, K., Palmer, M., Trenberth, K., Cazenave, A., Chambers, D., Champollion, N., Hansen, J., Josey, S. A., Loeb, N., Mathieu, P.-P., Meyssignac, B., and Wild, M.: An imperative to monitor Earth’s energy imbalance, *Nature Clim. Change*, 6, 138–144, <https://doi.org/10.1038/nclimate2876>, 2016.
- 760 Wang, Y., Trentmann, J., Pfeifroth, U., Yuan, W., and Wild, M.: Improvement of Air Pollution in China Inferred from Changes between Satellite-Based and Measured Surface Solar Radiation, *Remote Sens.*, 11, 2910, <https://doi.org/10.3390/rs11242910>, 2019.
- Wang, Z., Lin, L., Xu, Y., Che, H., Zhang, X., Zhang, H., Dong, W., Wang, C., Gui, K., and Xie, B.: Incorrect Asian aerosols affecting the attribution and projection of regional climate change in CMIP6 models, *npj Clim. Atmos. Sci.*, 4, <https://doi.org/10.1038/s41612-020-00159-2>, 2021.
- 765 Wei, J., Peng, Y., Mahmood, R., Sun, L., and Guo, J.: Intercomparison in spatial distributions and temporal trends derived from multi-source satellite aerosol products, *Atmos. Chem. Phys.*, 19, 7183–7207, <https://doi.org/10.5194/acp-19-7183-2019>, 2019.
- Wild, M.: Global dimming and brightening: A review, *J. Geophys. Res.*, 114, D00D16, <https://doi.org/10.1029/2008JD011470>, 2009.
- Wild, M.: Enlightening Global Dimming and Brightening, *Bull. Amer. Meteor. Soc.*, 93, 27–37, 2012.
- Wild, M., Wacker, S., Yang, S., and Sanchez-Lorenzo, A.: Evidence for clear-sky dimming and brightening in central Europe, *Geophys. Res. Lett.*, p. e2020GL092216, <https://doi.org/10.1029/2020GL092216>, 2021.
- 770 Yang, S., Wang, X., and Wild, M.: Causes of Dimming and Brightening in China Inferred from Homogenized Daily Clear-Sky and All-Sky in situ Surface Solar Radiation Records (1958–2016), *J. Climate*, 32, 5901–5913, <https://doi.org/10.1175/JCLI-D-18-0666.1>, 2019.
- Yli-Juuti, T., Mielonen, T., Heikkinen, L., Arola, A., Ehn, M., Isokääntä, S., Keskinen, H.-M., Kulmala, M., Laakso, A., Lipponen, A., Luoma, K., Mikkonen, S., Nieminen, T., Paasonen, P., Petäjä, T., Romakkaniemi, S., Tonttila, J., Kokkola, H., and Virtanen, A.: Significance of the organic aerosol driven climate feedback in the boreal area, *Nat. Commun.*, 12, 5637, <https://doi.org/10.1038/s41467-021-25850-7>, 2021.
- 775 Yu, H., Yang, Y., Wang, H., Tan, Q., Chin, M., Levy, R. C., Remer, L. A., Smith, S. J., Yuan, T., and Shi, Y.: Interannual variability and trends of combustion aerosol and dust in major continental outflows revealed by MODIS retrievals and CAM5 simulations during 2003–2017, *Atmos. Chem. Phys.*, 20, 139–161, <https://doi.org/10.5194/acp-20-139-2020>, 2020.
- Zhang, J., Zhou, X., Goren, T., and Feingold, G.: Albedo susceptibility of northeastern Pacific stratocumulus: the role of covarying meteorological conditions, *Atmos. Chem. Phys.*, 22, 861–880, <https://doi.org/10.5194/acp-22-861-2022>, 2022.
- 780 Zhao, B., Jiang, J. H., Gu, Y., Diner, D., Worden, J., Liou, K.-N., Su, H., Xing, J., Garay, M., and Huang, L.: Decadal-scale trends in regional aerosol particle properties and their linkage to emission changes, *Environ. Res. Lett.*, 12, 054 021, <https://doi.org/10.1088/1748-9326/aa6cb2>, 2017.
- Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., Qi, J., Yan, L., Zhang, Y., Zhao, H., Zheng, Y., He, K., and Zhang, Q.: Trends in China’s anthropogenic emissions since 2010 as the consequence of clean air actions, *Atmos. Chem. Phys.*, 18, 14 095–14 111, <https://doi.org/10.5194/acp-18-14095-2018>, 2018.
- 785 Zhou, C., Zelinka, M. D., and Klein, S. A.: Impact of decadal cloud variations on the Earth’s energy budget, *Nature Geosci.*, 9, 871–874, 2016.

790 Zhou, X., Zhang, J., and Feingold, G.: On the Importance of Sea Surface Temperature for Aerosol-Induced Brightening of Marine Clouds and Implications for Cloud Feedback in a Future Warmer Climate, *Geophys. Res. Lett.*, 48, e2021GL095896, <https://doi.org/https://doi.org/10.1029/2021GL095896>, 2021.