

Observation-Based Constraints on Modeled Aerosol Surface Area: Implications for Heterogeneous Chemistry

Rachel A. Bergin¹, Monica Harkey^{2,3}, Alicia Hoffman^{2,3}, Richard H. Moore⁴, Bruce Anderson⁴, Andreas Beyersdorf^{4,*}, Luke Ziemba⁴, Lee Thornhill^{4,5}, Edward Winstead^{4,5}, Tracey Holloway^{2,3}, and Timothy H. Bertram¹

¹Department of Chemistry, University of Wisconsin - Madison, Madison, WI 53703, USA

²Nelson Institute Center for Sustainability and the Global Environment, University of Wisconsin–Madison, Madison, WI 53703, USA

³Department of Atmospheric and Oceanic Sciences, University of Wisconsin–Madison, Madison, WI 53703, USA

⁴NASA Langley Research Center, Hampton, VA 23666, USA

⁵Science Systems and Applications, Inc., Hampton, VA 23666, USA

*Now at California State University, San Bernardino, San Bernardino, CA 92407, USA

Correspondence to: T.H. Bertram (timothy.bertram@wisc.edu)

Abstract. Heterogeneous reactions occurring at the surface of atmospheric aerosol particles regulate the production and lifetime of a wide array of atmospheric gases. Aerosol surface area plays a critical role in setting the rate of heterogeneous reactions in the atmosphere. Despite the central role for aerosol surface area, there are few assessments of the accuracy of aerosol surface area concentrations in regional and global models. In this study, we compare aerosol surface area concentrations in the EPA’s Community Multiscale Air Quality (CMAQ) model with commensurate observations from the 2011 NASA flight-based DISCOVER-AQ (Deriving Information on Surface Conditions from COlumn and VERtically Resolved Observations Relevant to Air Quality) campaign. The study region includes the Baltimore and Washington, DC metropolitan area. Dry aerosol surface area was measured aboard the NASA P-3B aircraft using an Ultra-High Sensitivity Aerosol Spectrometer (UHSAS). We show that modeled and measured dry aerosol surface area, $S_{a,mod}$ and $S_{a,meas}$ respectively, are modestly correlated ($r^2 = 0.52$) and on average agree to within a factor of two ($S_{a,mod}/S_{a,meas} = 0.44$) over the course of the 13 research flights. We show that $S_{a,mod}/S_{a,meas}$ does not depend strongly on photochemical age or the concentration of secondary biogenic aerosol, suggesting that the condensation of low-volatility gas-phase compounds does not strongly affect model-measurement agreement. In comparison, there is strong agreement between measured and modeled aerosol number concentration ($N_{mod}/N_{meas} = 0.87$, $r^2 = 0.63$). The persistent underestimate of S_a in the model, combined with strong agreement in modeled and measured aerosol number concentrations, suggests that model representation of the size distribution of primary emissions or secondary aerosol formed at the early stages of oxidation may contribute to the observed differences.

For reactions occurring on small particles, the rate of heterogeneous reactions is a linear function of both S_a and the reactive uptake coefficient (γ). To assess the importance of uncertainty in modeled S_a for the representation of heterogeneous reactions in models, we compare both the mean and the variance in $S_{a,mod}/S_{a,meas}$ to that in $\gamma(N_2O_5)_{mod}/\gamma(N_2O_5)_{meas}$. We find that the

35 uncertainty in model representation of heterogeneous reactions is primarily driven by uncertainty in the parametrization of reactive uptake coefficients, although the discrepancy between $S_{a,mod}$ and $S_{a,meas}$ is not insignificant. Our analysis suggests that model improvements to aerosol surface area concentrations, in addition to more accurate parameterizations of heterogeneous kinetics, will advance the representation of heterogeneous chemistry in regional models.

1 Introduction

40 1.1 The Role of Aerosol Surface Area in Heterogeneous Reaction Kinetics

Reactions occurring at atmospheric interfaces, such as suspended aerosol particles, catalyze the production and loss of key gas-phase compounds in Earth's atmosphere with important implications for regional air quality (Chang et al., 2011). The rate of heterogeneous reactions occurring at the surface of aerosol particles is a function of the gas-aerosol collision frequency and the per collision reaction probability. Variability in gas-aerosol collision frequency is determined by the aerosol surface area concentration. The probability of reaction, or the net reactive uptake coefficient (γ), is reaction specific and dependent on chemical kinetics, gas accommodation at the surface, and near surface diffusion (Abbatt et al., 2012). Collectively, the first-order removal rate of a gas-phase species (A) from the atmosphere can be written as:

$$\frac{d[A]}{dt} = -k_{het}[A] \quad \text{E1}$$

where the heterogeneous reaction rate constant (k_{het}), in the absence of gas-phase diffusion limitations, can be written as:

$$50 \quad k_{het} = \frac{\gamma_{species}\omega_{species}S_a}{4} \quad \text{E2}$$

where ω is the mean molecular speed of the gas-phase molecule (m s^{-1}) and S_a is the surface area concentration of aerosol particles ($\text{m}^2 \text{m}^{-3}$).

To date, most evaluations of the role of heterogeneous chemistry on gas-phase composition have focused on uncertainty in parameterizations of reactive uptake coefficients, such as the reactive uptake of dinitrogen pentoxide (N_2O_5) due to its role as a NO_x sink (Brown et al., 2009; Evans and Jacob, 2005; MacIntyre and Evans, 2010; McDuffie et al., 2018). In comparison, there has been less focus on model representation of aerosol surface area concentrations, despite the fact that k_{het} is linearly dependent on S_a . An accurate representation of aerosol surface area in regional and global chemical transport models is challenging, as S_a is a complex function of size-dependent aerosol particle emissions, chemical transformations, and removal

60 processes. Here, we directly compare aerosol surface area concentrations in a regional chemical transport model with commensurate aircraft measurements to assess the representation of S_a in regional air quality models.

1.2 Calculation of Aerosol Surface Area in Regional Air Quality Models

The total aerosol particle surface area concentration has been calculated in air quality models using a variety of approaches, including the discrete representation of the particle size distribution in defined size ranges, known as the sectional method (Adams and Seinfeld, 2002; Gelbard et al., 1980; Jacobson, 2001; Lee et al., 2009; Lee and Adams, 2012; Luo and Yu, 2011; Spracklen et al., 2006; Trivitanurak et al., 2008; Yu and Luo, 2009), and a continuous modal representation of the particle size distribution (Kleeman et al., 1997; Mann et al., 2010; Meng, 1998; Pringle et al., 2010; Sartelet et al., 2006; Stier et al., 2005; Vignati et al., 2004; Zhang et al., 2010a). Here, we review the modal representation of particle size distributions implemented in the Community Multiscale Air Quality (CMAQ) model and the calculation of both wet and dry total aerosol surface area (S_a). Aerosol particle size distributions in CMAQ follow the method developed for the Regional Particulate Model, an extension of the Regional Acid Deposition Model (Binkowski, 1999; Binkowski and Roselle, 2003) where the total particle size distribution is treated as the superposition of three separate lognormal distributions (or modes); Aitken, accumulation, and coarse modes (Binkowski, 1999; Whitby, 1978). The lognormal particle size distribution for each mode is defined as:

75

$$n(\ln D) = \frac{N}{\sqrt{2\pi} \ln \sigma_g} \exp \left[-0.5 \left(\frac{\ln \frac{D}{D_g}}{\ln \sigma_g} \right)^2 \right] \quad \text{E3}$$

where N is the total number concentration, D is the particle diameter, and D_g and σ_g are the geometric mean diameter and geometric standard deviation. Under this definition, the Aitken mode describes aerosol particles of diameter smaller than approximately $0.1 \mu\text{m}$ with a median diameter of $0.03 \mu\text{m}$, while the accumulation mode encompass the diameter range of 0.1 to $2.5 \mu\text{m}$ with a median diameter of $0.3 \mu\text{m}$ (Binkowski, 1999). The coarse mode describes particles of diameter 0.3 to about $10 \mu\text{m}$ with a median diameter of $6 \mu\text{m}$. It should be noted that there is uncertainty in the exact size distributions within CMAQ dependent on emissions parameters, so the median diameters within modes are approximate (Elleman and Covert, 2010). Particle nucleation and growth result in changes to the mode diameter and can result in the transfer of particle number, surface

area, and mass to the next larger size (e.g. Aitken to accumulation). Outside of particle growth and nucleation, the approximate
85 median diameters are unchanged. The size ranges for each mode are based on Whitby 1978, and geometric standard deviation
is also based on Whitby 1978 but has been updated to those from Elleman and Covert 2010.

Three integral properties of the aerosol size distribution are calculated in CMAQ: the zeroth (M_0), second (M_2), and third
moments (M_3), where the k^{th} moment of the size distribution is calculated as:

$$90 \quad M_k = \int_{-\infty}^{\infty} D^k (\ln D) d(\ln D) = ND_g^k \exp\left[\frac{k^2}{2} \ln^2 \sigma_g\right] \quad E4$$

In this representation, $N = M_0$, $S_a = \pi M_2$, and $V = \left(\frac{\pi}{6}\right) M_3$, where V is the total aerosol volume (Binkowski, 1999; Binkowski
and Roselle, 2003). Though M_2 is utilized in CMAQ's aerosol subroutines, it is multiplied by π prior to use in main CMAQ
routines, such that it is identified as a modal surface area (Binkowski and Roselle, 2003).

95 The time rate of change of each moment is calculated for each grid box and time interval as:

$$\frac{\partial M_k}{\partial t} = P_k - L_k M_k \quad E5$$

where P and L represent the production and loss of M_k in each aerosol mode. With respect to S_a ($S_a = \pi M_2$), neglecting
transport terms, P_2 includes new particle formation (Aitken mode only), condensational growth, and primary emissions and L_2
includes intramodal coagulation, dry and wet deposition. Primary aerosol emission rates are sourced from the 2011 EPA
100 National Emissions Inventory, which characterizes emissions based on source type and location. Within CMAQ, all primary
aerosol emissions, independent of source type, are parameterized with modal size distributions per Elleman and Covert 2010
(see SI). In the version of CMAQ used here, new particle formation is based on classical, binary homogeneous nucleation
(Kulmala et al., 1998). Particle growth is described by Binkowski and Roselle (2003), and secondary organic aerosol schemes
in Carlton et al.(2010).

105

In the interpretation of model S_a , the following model specific details should be considered: 1) Fine particles (Aitken and
accumulation modes) do not coagulate with coarse mode particles and coarse mode particles do not coagulate with each other
(Binkowski and Roselle, 2003). 2) The size distribution for primary PM_{2.5} emissions are assumed to have a geometric mean

($D_g = 0.3\mu\text{m}$) and geometric standard deviation ($\sigma_g = 2$) and > 99% of $\text{PM}_{2.5}$ emission are assigned to the accumulation mode
110 (Binkowski and Roselle, 2003), which may have consequent effects on the aerosol surface area distribution. 3) Particles are
assumed to be spherical.

The condensation of water is accounted for in the chemical evolution of M_2 , thus M_2 is inherently the wet second moment
(M_2^w) which is used in the calculation of heterogeneous chemical reactions. In addition to M_2^w , a dry second moment (M_2^d) is
115 calculated as a function of the third moment (M_3) as:

$$M_2^d = M_2^w \left(\frac{M_3^d}{M_3^w} \right)^{\frac{2}{3}} \quad \text{E6}$$

In the following analyses, we concentrate on the comparison of modeled and measured S_a to evaluate the relative uncertainty
associated with model descriptions of heterogeneous kinetic mechanisms (i.e. reactive uptake coefficients, γ) and aerosol
120 particle size distributions (i.e. aerosol surface area, S_a) that combined dictate the fate of reactive gas-phase molecules.

1.3 Previous Model-Measurement Comparisons of Aerosol Surface Area

Evaluation of regional air quality models has largely focused on criteria air pollutants such as ozone (O_3) and particle mass
(e.g. $\text{PM}_{2.5}$) (Appel et al., 2021). Previous model evaluation of particle mass has focused on an array of metrics including mass
125 concentration (Gantt et al., 2012; Spak and Holloway, 2009; Wang et al., 2009), number concentration (Park et al., 2006;
Ranjithkumar et al., 2021; Wang et al., 2009; Zhang et al., 2010b), size distribution (Kelly et al., 2011; Nolte et al., 2015; Park
et al., 2006; Zhang et al., 2010b), composition (Knote et al., 2011; Nolte et al., 2015; Prank et al., 2016), and aerosol optical
depth (Ghan et al., 2001; Knote et al., 2011), among others. There has been a very long and detailed history of CMAQ
evaluation of $\text{PM}_{2.5}$, including both ground-based (Baker et al., 2018; Fan et al., 2005; Ghim et al., 2017; Hogrefe et al., 2009,
130 2015; Liu and Zhang, 2011; Prank et al., 2016; Smyth et al., 2006; Wang et al., 2021; Yu et al., 2012, 2008b, 2008a; Zhang et
al., 2019, 2006, 2010c), ship-based (Yu et al., 2012), and aircraft-based measurements (Baker et al., 2018; Chen et al., 2020;
Yu et al., 2012). For fifteen studies comparing ground-based measurements of $\text{PM}_{2.5}$ to CMAQ outputs between 1999 and

2018, ten saw an underestimation of $PM_{2.5}$ by the model ranging between 6-75% (Ghim et al., 2017; Liu and Zhang, 2011; Prank et al., 2016; Wang et al., 2021; Yu et al., 2008a, 2012, 2008b; Zhang et al., 2019, 2006, 2010c), dependent on pollution
135 events and rural versus urban location, while four found that CMAQ predicted observations well (Baker et al., 2018; Fan et al., 2005; Hogrefe et al., 2009; Smyth et al., 2006), matching general trends in the observational data, and one saw an overestimation of observational data (Hogrefe et al., 2015). Of the three aircraft studies, two saw significant underestimation of $PM_{2.5}$ aloft (Baker et al., 2018; Chen et al., 2020), while one saw overestimation in some $PM_{2.5}$ compositional components and underestimation in others (Yu et al., 2012).

140

Particle surface area specifically is not regulated as a criteria air pollutant as standards of measurement and air quality controls are determined on a mass per unit volume basis. However, particle surface area indirectly affects the concentration of $PM_{2.5}$ and O_3 as it can serve to regulate the lifetime of nitrogen oxides (Chang et al., 2011; Geyer and Stutz, 2004; Portmann et al., 1996; Stadler et al., 2018) and hydrogen oxides (George et al., 2013; Lakey et al., 2015; Martin et al., 2003; Thornton et al.,
145 2008; Thornton and Abbatt, 2005), the production rate of secondary organic aerosol (Gaston et al., 2014), and new particle formation and growth rates, as preexisting aerosol surface area serves as a condensation sink for low volatility gas-phase compounds (Donahue et al., 2014; Trump et al., 2014). There are few reports of model-measurement comparisons of particle surface area, and those that have been reported in the literature have focused on comparisons of heavily spatially and temporally averaged concentrations (e.g., field campaign averages). For example, Simon et al. compared ground-based aerosol surface
150 area concentrations calculated in the CAMx model to measurements made aboard the *R.V. Ronald H. Brown* in the Gulf of Mexico and the Houston Ship Channel with two differential mobility particle sizers and an aerodynamic particle sizer (Bates et al., 2008; Simon et al., 2010). The results of these studies are given in Table 1. Model prediction of median S_a in the Gulf of Mexico was similar to the measurement data ($S_{a,mod}/S_{a,meas} = 0.96$), with median values and ranges again shown in Table 1. In comparison, model prediction of median S_a in the Houston Ship Channel, where there is large spatial and temporal
155 fluctuation in S_a relative to the smaller interquartile range seen in the Gulf of Mexico, yielded $S_{a,mod}/S_{a,meas} = 1.6$. Modeled S_a was also compared to measured S_a aloft on two research flights from the TexAQS II/GoMACCS field study in September and October of 2006. The range of measured S_a values ($<600 \mu m^2/cm^3$) matched well with the average values predicted in the

CAMx model, though the maximum modeled values were much larger than those measured ($4000-8000\mu\text{m}^2/\text{cm}^3$ compared to $<600\mu\text{m}^2/\text{cm}^3$), consistent with the *R.V. Ronald H. Brown* comparisons. Overall, it should be noted that on a regional scale, modeled values agree well with measurements of aerosol S_a , however maximum modeled values were larger than those measured both for ground-based measurements and those aloft.

Region		Median ($\mu\text{m}^2/\text{cm}^3$)	Q1, Q3 ($\mu\text{m}^2/\text{cm}^3$)	Reference
Gulf of Mexico	Measured	361	277, 398	Simon et al. 2010
	CAMx Model	347	298, 394	
Houston Ship Channel	Measured	592	513, 800	Simon et al. 2010
	CAMx Model	949	667, 1760	
Northeast U.S. (surface - 0.1km)	Measured	135	90.6, 223	Jaeglé et al., 2018
	GEOS-Chem Model	169	-	
Northeast U.S. (3.5 - 4.5 km)	Measured	4.4	2.49, 17.1	Jaeglé et al., 2018
	GEOS-Chem Model	3.8	-	

165 **Table 1: Comparison of model and measured aerosol surface area concentrations. Simon et al. (2010) compared CAMx model results with ground-based measurements in the Gulf of Mexico and the Houston Ship Channel. Jaeglé et al. (2018) compared measurements from the WINTER campaign and GEOS-Chem modeled data at the surface and aloft between 3.5 and 4.5km.**

More recently, modeled dry surface area concentrations were assessed over the northeast US during the 2015 Wintertime
 170 INvestigation of Transport, Emissions, and Reactivity (WINTER) aircraft campaign (Jaeglé et al., 2018). While quantitative assessment of aerosol surface area was not the focus of this study, dry aerosol S_a was calculated by combining dry aerosol size distribution observations from a Passive Cavity Aerosol Spectrometer Probe and Ultra-High Sensitivity Aerosol Spectrometer

and comparing to the GEOS-Chem chemical transport model. Two versions of the model, a reference and improved model were compared to the observations within 13 altitude bins, ranging from surface to 4.5km, here we focus on the improved
175 model results. The GEOS-Chem model medians were encompassed by the observed interquartile ranges in each altitude bin. The improved model showed excellent agreement with measurements when compiled over large spatial and temporal scales, where $S_{a,mod}/S_{a,meas}$ was 1.25 and 0.68 for the surface and 4.5km comparisons, respectively.

Given the importance of accurate model representation of aerosol surface area to multiple atmospheric processes, and the
180 limited number of prior studies conducted in urban environments, we revisit this comparison using a regional air quality model aerosol and commensurate aircraft observations conducted in an urban environment.

2 Methods and Models

2.1 Aerosol Evaluation in the CMAQ Model

CMAQ simulations were performed as described by Abel et al. 2018, Abel et al. 2019, and Harkey et al. 2021, with carbon
185 bond 5 chemistry, anthropogenic emissions from the 2011 National Emissions Inventory, and input meteorology from the Weather Research and Forecasting (WRF) version 3.2.1 (Skamarock et al., 2008), constrained to the North American Regional Reanalysis (NARR; Messinger et al., 2006; Abel et al., 2018, 2019; Harkey et al., 2021). The CMAQ simulation utilized here employed CMAQ version 5.2.1 (Byun and Schere, 2006; Nolte et al., 2015), and was run with 25 vertical layers from the surface to 100 hPa, a 12×12 km grid, and hourly temporal resolution. Anthropogenic emissions and emissions from fires (both
190 prescribed and not) were based on the 2011 National Emissions Inventory version 2, with in-line estimates of NO and NO₂ produced by lightning, boundary conditions from the Model for Ozone and Related Chemical Tracers version 4 (MOZART; Emmons et al., 2010), and biogenic emissions from WRF output in the Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN; Guenther et al., 2012). The model was run from 20 May through 31 August 2011, to include 11 days of spin-up (Harkey et al., 2021). The data set utilized in this analysis is only a subset of the model data set originally run at
195 UW-Madison.

CMAQ was also run for the time period of the 2015 WINTER field campaign for comparison of modelled and measured N_2O_5 uptake coefficients. This CMAQ simulation also employs input meteorology constrained to NARR, calculated using WRF version 3.8.1 (Skamarock et al., 2008). Anthropogenic emissions were taken from the 2016 National Emissions Inventory Collaborative, version 1 (NEIC, 2019). Emissions from fires and boundary conditions were taken from the EPA Air Quality Time Series (EQUATES) project (<https://www.epa.gov/cmaq/equates>). Biogenic emissions and lightning NO_x emissions were both calculated in-line. This simulation employed CMAQ version 5.3.2 (Appel et al., 2021), with carbon bond 6 chemistry (Emery et al., 2015; Luecken et al., 2019). The WINTER-period simulation was run from 21 January through 16 March 2015, to include 11 days of spin-up, with hourly output on a 12 x 12 km grid and on 35 vertical levels from the surface to 100 hPa.

205

The CMAQ simulation for the 2011 DISCOVER-AQ period employed the “AERO6” aerosol module (Binkowski and Roselle, 2003; Carlton et al., 2010; Foley et al., 2010; Sonntag et al., 2014), where primary and secondary aerosols are characterized by bimodal lognormal size distributions, and the total size distribution is the sum of three aerosol size modes: Aitken, accumulation, and coarse modes with median diameters of 0.03, 0.3, and $6\mu m$ respectively. The CMAQ simulation for the 2015 WINTER period employed the “AERO7” aerosol module, which builds on the AERO6 module, with updates to aerosols formed by monoterpene oxidation, anthropogenic volatile organic compounds, and to aerosol liquid water (Pye et al., 2015; Pye et al., 2017; Qin et al., 2021; Xu et al., 2018). The CMAQ simulation for the DISCOVER-AQ period employed the default heterogeneous N_2O_5 uptake (Davis et al., 2008), while the simulation covering the WINTER period employed a N_2O_5 uptake modified per Bertram and Thornton (2009).

215

Due to the modality of the CMAQ representation of aerosols, we calculate each parameter relating to the aerosol data set separately for each mode, and these are then combined to result in a total value that can be directly compared to the DISCOVER-AQ observational data. The total CMAQ dry surface area is computed as the sum of the modal dry surface areas. The variable SRF is an output of CMAQ but is defined as $SRF = \pi M_2^d$. SRF is a modal variable like each moment, such that total surface area = $SRF_{ATKN} + SRF_{ACC} + SRF_{COR}$ (dry surface area in the Aitken, accumulation, and coarse modes, respectively).

2.2 DISCOVER-AQ 2011 Campaign

Research flights conducted during the NASA DISCOVER-AQ (Deriving Information on Surface Conditions from Column
225 and VERTically Resolved Observations Relevant to Air Quality) campaigns were designed to measure the vertical and spatial
distribution of key air pollutants in urban environments, with a focus on connecting surface measurements with vertically
integrating satellite observations. The first DISCOVER-AQ campaign, conducted aboard the NASA P-3B aircraft during July
2011, comprised of 14 science flights in the Baltimore and Washington, DC area (Crawford et al., 2014; Crawford and
Pickering, 2014; NASA, 2012). The 2011 DISCOVER-AQ campaign was the first of a series of flights with an objective of
230 narrowing the gap of satellite and observational data and air quality utilizing near-surface air pollution measurements. Science
flights concentrated on high time resolution measurements of atmospheric composition in the convective boundary layer. Here,
we focus on measurements of dry aerosol surface area concentration (S_a), determined from high time resolution (1Hz) size
distributions made using an Ultra-High Sensitivity Aerosol Spectrometer (Droplet Measurement Technologies, UHSAS)
integrating between $60 < d_p < 1000$ nm, which captures the peak of the surface area distribution, shown in Figure 1 below. The
235 UHSAS measures the particle size from optical light scattering, which was calibrated during DISCOVER-AQ using NIST-
traceable polystyrene latex spheres whose refractive index may differ slightly from that of real-world, aerosols that may result
in a slight under-sizing bias (Moore et al., 2021). Ambient air was sampled through an isokinetic inlet, allowing the aerosol
temperature to quickly equilibrate to that of the cabin of the P3-B aircraft. Along with additional ram pressure heating in the
inlet during flow deceleration, particles reach an RH of below 40-50%, meaning the aerosol is considered to be dry. The term
240 dry here distinguishes that the particle hydration state is greatly reduced compared to that of the unperturbed ambient air. It is
important to note that aerosol water is not accounted for in the following model-measurement comparison, though there would
be some aerosol water present at the higher end of the 40-50% RH threshold, which may impact the comparison of the measured
aerosol to dry aerosol in the model. The particle mobility size from 10-310 nm diameters was measured with a TSI Scanning
Mobility Particle Sizer (SMPS) with 45-second time resolution and the particle aerodynamic size from 500-4000 nm diameters
245 was measured with a TSI Aerodynamic Particle Sizer (APS) at 1 Hz. On the representative day shown in Figure 1, SMPS data
showed that approximately 5.8% of the surface area fell below the 60nm threshold of the UHSAS measurement for an average

surface area distribution for the average of all altitudes, while the APS indicated that supermicron particles did not contribute to the particle surface area. The distribution comparison between the model and measurement differs in shape at different altitudes. Figure 1b shows a comparison of the near surface data, below 1km, which gives a very similar distribution to that of the average of all altitudes. However, figure 1c, or that above 3km altitude shows a different shape, and one that is more comparable between model and measurement. It should be noted that the axis scales in figure 1b and 1c do not match, showing a discrepancy in distribution calculation between CMAQ and DISCOVER-AQ that will be discussed in later sections. For simplicity, we choose to focus exclusively on the UHSAS size distribution data given their high frequency and wide size range of particle diameter, but it is important to note that not all of the particle surface area is captured by the UHSAS instrument.

UHSAS data is available to the public at the NASA Langley Atmospheric Science's Data Center and Distributed Active Archive Center (<http://doi.org/10.5067/Aircraft/DISCOVER-AQ/Aerosol-TraceGas>). In the following analysis we utilize observations of nitric oxide (NO) and nitrogen dioxide (NO₂) measured with the NCAR four channel chemiluminescence instrument (Ridley and Grahek, 1990) and carbon monoxide (CO) measured via Differential Absorption CO Measurement (DACOM) (Sachse et al., 1987) to assess differences in modeled and measured aerosol surface area.

260




Figure 1: Modeled (blue) and measured (orange) dry aerosol surface area distributions averaged over the DISCOVER-AQ sampling domain of Baltimore and Washington, DC at a) all altitudes, b) altitudes below 1km, and c) altitudes above 3km for 1 July 2011.

2.3 Model-Measurement Comparison

265 To compare measured and modeled S_a , we sample the hourly 12km \times 12 km CMAQ output at the time and location of each DISCOVER-AQ sampling point. The spatial resolution of CMAQ, relative the DISCOVER-AQ flight, is shown in Figure 2 for the 28 July 2011 research flight, where the color corresponds to the modeled surface-level dry S_a (in $\mu\text{m}^2 \text{cm}^{-3}$) at noon EST. Indexing and analysis of the two aforementioned data sets was completed in MATLAB. Each 1s data point from the DISCOVER-AQ 2011 campaign was mapped to the nearest (as described below) 4D index (time of day, latitude, longitude, 270 and altitude) in the lower time and spatial resolution CMAQ model for direct comparison. The nearest 1hr averaged CMAQ time point was selected based on the time window that encompassed the aircraft flight time, i.e. a flight time of 09:16:00 was mapped to CMAQ time period of 9:00 – 9:59. The nearest 12km \times 12km CMAQ grid box was also selected based on the grid box that encompassed the aircraft location at the time of sampling. The nearest CMAQ altitude (or layer) was identified by locating the aircraft height within one of the 25 indexes in which the altitude was encompassed. With all four CMAQ indexes 275 assigned to each data point, the 1s DISCOVER-AQ data set could be fully mapped and compared to that from the CMAQ model. It should be noted that there are far more data points in the observed data than in the model due to resolution constraints, and thus the model is being oversampled. Each of the four indexes were concatenated together in the order defined in CMAQ, namely latitude, longitude, layer (altitude), and time. This process was utilized for each data point for an entire flight of DISCOVER-AQ and was then replicated for each subsequent flight. The result of this approach is shown in Figure 3, for the 280 comparison of modeled and measured carbon monoxide (CO). The coefficient of determination (r^2) for the linear regression of modeled vs measured CO concentration ($\text{CO}_{\text{mod}} / \text{CO}_{\text{meas}}$) was 0.44 with a slope of 1.0499 ± 0.0007 . The large variance highlights the spatial and temporal mismatch of model sampling and measurement, while the near unit correlation coefficient indicates that on average modeled and measured CO agree. This agreement implies that the model-measurement comparison of many well-understood parameters should be accurate and that there is not a fundamental issue in comparing modeled and 285 measured data between the two datasets.




Figure 2: Map of the Baltimore-Washington DC area representative of the July 2011 DISCOVER-AQ flights with an example flight path from 28 July 2011. The flight path is overlaid on the CMAQ model grid, showing the modeled surface area in each 12km \times 12km grid box at noon EST (UTC 16).




Figure 3: a) Linear regression of modeled vs measured carbon monoxide (CO) mixing ratio in ppbv and b) histogram of the modeled-measured CO for the entire DISCOVER-AQ campaign.


295 3 Results

3.1 Campaign Averaged Comparison of Aerosol Surface Area Concentrations

First, we assess general agreement between campaign averaged modeled and measured surface area concentrations. The observational data is from the 13 DISCOVER-AQ flights during July 2011 (1-29 July). The final research flight consisted of a dual highway leg conducted south along the Baltimore-Washington Parkway and north along I-95 at low altitude to compare
 300 the two roadways. Given the proximity to a large point source, this research flight was not included in the following analysis. The observational data from the UHSAS included number, surface area, and volume measurements, though the focus of this study is primarily surface area (S_a).

In this analysis, we compare dry surface area concentrations as the DISCOVER-AQ measurements were made at an RH below
 305 40-50% and are thus considered dry, and we do not have direct measurements of particle growth factors for comparison of wet S_a . However, it is important to note that S_a used in E2 is the surface area concentration at ambient humidity and any uncertainty in modeled aerosol hygroscopicity will propagate to the aerosol surface area concentration used in E2. Figure 4 shows the campaign-averaged vertical profile of both the measured dry UHSAS surface area ($S_{a,meas}$) as well as the modeled dry CMAQ

surface area ($S_{a,mod}$), along with the interquartile ranges separated into 1km altitude bins. Since the UHSAS measurement
 310 frequency is 1 Hz and the CMAQ modeled data is at one-hour time resolution, and the model samples a full domain in 12km
 grid boxes compared to the smaller domain sampled by aircraft, there are many more measurement data points ($N = 330204$)
 than comparable modeled data points ($N = 5196$) over the course of the flight campaign. In Figure 4, the UHSAS measurements
 have been averaged to the spatial and temporal resolution of the model, such that the number of observational points is the
 same as the number of model points. The light gray error bars shown in Figure 4a reflect the standard deviation of the data
 315 from the mean at that point in time and space. It should be noted that the error bars on this dataset are large, due to the spatial
 and temporal mismatch between model and measurement in a highly heterogeneous sampling domain. For both model and
 measurement, the surface area increases towards the surface, as is to be expected, and decreases with altitude. The vertical
 profile is well captured by the CMAQ model, however, there is a larger range of measured surface area concentrations than is
 seen in the corresponding model altitude bin.



320

Figure 4: Average vertical profile of a) measured DISCOVER-AQ aerosol surface area concentration ($S_{a,meas}$) and b) CMAQ aerosol surface area concentration ($S_{a,mod}$) over the entirety of the DISCOVER-AQ campaign. Each measured point is an average of the points included in that 4D index corresponding to CMAQ. The overlaid boxplots show the median (red line within blue box), and interquartile ranges (blue box with the 25th percentile at the left end and 75th percentile at the right end) in 1km altitude bins. The labels on the altitude axis lie at the midpoint of the 1km altitude bin, and red crosses indicate outliers from the majority of the dataset at that altitude portion.

325

As shown in Figure 4, measured S_a is on average larger than modeled S_a , particularly at low altitude, where the ratio of the median, modeled S_a to measured S_a near the surface ($z < 1$ km) is 0.47. This contrasts with what has been reported previously in the literature. For example, both Jaeglé et al and Simon et al. found that the median modeled near-surface S_a was consistently
 330 larger than measured S_a ($S_{a,mod}/S_{a,meas} = 1.04 - 1.6$) (Simon et al., 2010).

The comparison between modeled and measured S_a is also shown with histograms in Figure 5 for all altitudes (5a,b) and for the surface level measurements (0-1km; 5c,d). While the number of points is not consistent between the model and measurement datasets due to the 12km grid box constraint and time frequency in CMAQ, differences in the range in surface
 335 area concentrations are observed. Measured surface area concentrations range between 0 - $1.87 \times 10^3 \mu\text{m}^2/\text{cm}^3$ with the vast majority of data below $420 \mu\text{m}^2/\text{cm}^3$, while modeled S_a ranges between 0- $300\mu\text{m}^2/\text{cm}^3$.




Figure 5: Histograms of a) measured aerosol surface area concentration (DISCOVER-AQ) at all altitudes, b) modeled aerosol surface area concentration (CMAQ) at all altitudes, c) measured aerosol surface area in the 0 to 1km altitude bin, and d) modeled aerosol surface area in the 0 to 1km altitude bin over the entirety of the DISCOVER-AQ campaign.
 340

3.2 Direct Model-Measurement Comparison

A linear regression of CMAQ modeled dry aerosol surface area concentration and measured aerosol surface area concentration is shown in Figure 6. The measured data has been averaged to the space and time domain of CMAQ (latitude, longitude, altitude, and time). The coefficient of determination (r^2) for the linear regression of modeled and measured S_a was 0.52 with a slope of 0.437 ± 0.004 , indicating that the measured S_a is on average twice that of the model value.





Figure 6: Comparison of $S_{a, \text{mod}}$ and $S_{a, \text{meas}}$ over the full DISCOVER-AQ campaign with measurement data averaged to the corresponding model latitude, longitude, altitude, and time point.

The histogram of the surface area ratio ($S_{a, \text{mod}}/S_{a, \text{meas}}$) throughout the campaign in Figure 7 shows that the model underpredicts the measured surface area ratio in 81% of the comparison points. The model underpredicts S_a by a factor of two 44% of the time. In the following section, we explore potential causes for model-measurement disagreement, including model-

measurement spatial and temporal differences, the spatial distribution of primary emissions, and/or treatment of secondary aerosol formation.



355

Figure 7: Histogram of the ratio of model to measured aerosol surface area concentration ($S_{a,mod}/S_{a,meas}$) over the full DISCOVER-AQ campaign.


4 Discussion

In the following section, we explore the source of model-measurement discrepancy in S_a discussed in section 3. We begin by
360 investigating the dependence of $S_{a,mod}/S_{a,meas}$ on altitude and proximity to primary aerosol sources. We then investigate the
role of temporal and spatial resolution as CMAQ has a much coarser resolution, both spatially and temporally, than the
measured data. Finally, we investigate the possibility of impacts on $S_{a,mod}/S_{a,meas}$ from anthropogenic and biogenic indicators
as they are tied to aerosol emissions. It is also important to acknowledge that some of the model-measurement disagreement
could be due to processes not considered in the model such as phase separation, viscosity changes of aerosols, and direct
365 modeling of clouds impacting cloud processing of aerosols, though the impacts of these processes are not investigated further
in this work. The lack of a fourth mode below the Aitken mode for nucleation of particles and growth to the Aitken mode also

impacts the accuracy of the size distribution within CMAQ and may explain a portion of the model-measurement disagreement, though it is known that improving the default parameterization does not reduce all errors to the size distribution (Elleman and Covert, 2009b).”

370 4.1 Dependence of $S_{a,mod}/S_{a,meas}$ on Altitude

Given the strong dependence of S_a on altitude as shown in Figure 4, we first explore if part of the variance in $S_{a,mod}/S_{a,meas}$ shown in Figure 6 can be explained by altitude. In Figure 8, we show $S_{a,mod}/S_{a,meas}$ as a function of altitude. As shown, there is an altitude dependence in $S_{a,mod}/S_{a,meas}$, where the mean, median, and interquartile range (25th to 75th percentile) are given in Table 2 for the 1km altitude bins from 0-5km. Model-measurement discrepancy in S_a is largest at low altitude, where particle number concentrations are highest, proximity to particle sources is close, and heterogeneity in particle number concentrations are largest.



380 **Figure 8: Ratio of modeled to measured aerosol surface area concentration ($S_{a,mod}/S_{a,meas}$) including median (red line within blue box), and interquartile ranges (blue box with the 25th percentile at the left end and 75th percentile at the right end) in 1km altitude bins. The red crosses outside of the bounds of the plot note outliers. The labels on the altitude axis lie at the midpoint of the 1km altitude bin.**

0-1km 1-2km 2-3km 3-4km 4-5km

$S_{a,mod}/S_{a,meas}$ (mean)	0.56	0.97	0.89	1.05	1.42
$S_{a,mod}/S_{a,meas}$ (median)	0.47	0.51	0.63	0.82	0.75
$S_{a,mod}/S_{a,meas}$ (interquartile range)	0.33-0.65	0.33-0.80	0.37-0.97	0.56-1.31	0.53-2.14

Table 2: Mean, median, and interquartile range (range of 25th to 75th percentile) surface area ratio ($S_{a,mod}/S_{a,meas}$) for each 1km altitude bin from 0-5km.

385 4.2 Dependence of $S_{a,mod}/S_{a,meas}$ on Spatial and Temporal Resolution

The 12 km \times 12 km spatial resolution and 1 hour temporal resolution of CMAQ is significantly larger and longer than the spatial and temporal resolution of the aircraft data, resulting in an inherent contrast in resolution between model and measurement that may play a role in the variance in $S_{a,mod}/S_{a,meas}$. Within any individual 12 km \times 12 km model pixel, in the Baltimore-Washington sampling area, there is heterogeneity in S_a as shown in Figure 9. Sub grid scale variability in S_a would
 390 lead to increased variance in $S_{a,mod}/S_{a,meas}$, but likely with a mean and median close to 1 if the domain sampling was not biased, comparable to what is observed in the CO comparison (Figure 3), where the histogram of the CO data showcases a clear center around 1, with very few data points beyond a CO_{mod}/CO_{meas} value of 2.




Figure 9: Flight path from 28 July with overlaid UHSAS S_a data within the 10th layer of CMAQ altitude (~850-1000m) and gridded CMAQ S_a from layer 10 in the background. The CMAQ data is specifically at noon EST (UTC 16).

To investigate the discrepancy more quantitatively, we compare the probability density functions (PDF) of the model-to-measured CO, NO_x, particle number concentration and particle surface area concentrations. We use the PDF to characterize the population of data based on the standard deviation and mean, which provides a quantitative and comparable assessment of the variability in the comparison. Assuming that the research flights sampled the CMAQ model domain in an unbiased way (i.e. flights did not target or avoid point sources) we would expect that the PDFs of the model-to-measured ratio in CO, NO_x, number concentration, and surface area concentration would all center at 1 (or $\log_{10}(1) = 0$ as shown in Figure 10) and the standard deviation of the distribution (σ) would reflect heterogeneity in the scalar concentration at scales smaller than the model spatial or temporal domain. The histogram, PDF, and cumulative distribution function (CDF) for $\log_{10}(S_{a, \text{mod.}}/S_{a, \text{meas.}})$ is shown in Figure 10. The PDF of the histogram of $\log_{10}(S_{a, \text{mod.}}/S_{a, \text{meas.}})$ has a mean (μ) of -0.26 and standard deviation (σ) of 0.34.





Figure 10: Normalized histogram, probability density function (PDF), and cumulative distribution function (CDF) for the model-to-measured aerosol surface area ($S_{a, \text{mod.}}/S_{a, \text{meas.}}$). The histogram and PDF serve to indicate the median and spread of the dataset, while the CDF indicates the percentage of data encompassed at a certain data threshold.

Comparison of the peak and width of the PDF of the model-to-measured ratios of CO, particle number concentration (N), and NO_x provides an objective measure for assessing the impact of spatial and temporal resolution on the comparison. As shown in Figure 11 and Table 3, the mean of each PDF is -0.0029, 0.047, and -0.14 for CO, N , and NO_x. Each of these values is

415 significantly closer to 0 than that measured for S_a (-0.26), suggesting that the methodology for assessing model-measurement agreement should not be significantly impacted by model resolution, especially given the large range in atmospheric lifetimes for CO, N , and NO_x . However, without utilizing a smaller model resolution to directly test for impacts of the grid size, resolution issues cannot be fully ruled out. Interestingly, the mean $N_{\text{mod}}/N_{\text{meas}}$ is close to 1 ($10^{0.047} = 1.11$), where a value closer to 1 indicates agreement between model and measurement and that closer to zero indicates a large discrepancy between
 420 datasets. The mean $N_{\text{mod}}/N_{\text{meas}}$ is significantly different than that observed for $S_{a, \text{mod}}/S_{a, \text{meas}}$ ($10^{-0.26} = 0.55$), perhaps suggesting that the model-measurement disagreement is related to the shape of the size distribution either due to the *a priori* emissions size distribution or secondary aerosol processes.



425 **Figure 11: Normalized probability density functions for the \log_{10} of the model to measurement ratio in particle number concentration (yellow), carbon monoxide (red), NO_x (purple), and particle surface area concentration (blue).**

	mean (μ)	standard deviation (σ)
$\log_{10}(S_{a, \text{mod}}/S_{a, \text{meas}})$	-0.26	0.34
$\log_{10}(N_{\text{mod}}/N_{\text{meas}})$	0.047	0.27
$\log_{10}([\text{CO}]_{\text{mod}}/[\text{CO}]_{\text{meas}})$	-0.0029	0.15

$$\log_{10}([\text{NO}_x]_{\text{mod}}/[\text{NO}_x]_{\text{meas}})$$

$$-0.14$$

$$0.34$$

Table 3: Probability Density Function (PDF) fit parameters for the distributions shown in Figure 11.

430 Also shown in Figure 11 and Table 3 the standard deviation (σ) of the PDF for the CO, N , and NO_x model-to-measurement ratios, is 0.15, 0.27, and 0.34 respectively. The standard deviation of the PDF of $[\text{CO}]_{\text{mod}}/[\text{CO}]_{\text{meas}}$ is the narrowest, likely reflecting the longer lifetime of CO and a damping of sub grid scale variability of CO in each pixel. The width of the N and S_a ratio distributions are comparable, again highlighting that the deviation of $S_{a,\text{mod}}/S_{a,\text{meas}}$ from 1 may reflect differences in the model-measured aerosol size distribution (as shown in Figure 1). Collectively, this analysis suggests that there is not a
435 significant bias on average in the methodology based on model resolution and that the apparent differences in the number and surface area model-measurement ratios are most likely driven by the shape of the underlying aerosol size distribution.

It is interesting to note that the model-measurement agreement in particle number concentrations is significantly better than that of particle surface area concentrations, implying that the differences in S_a may be related to the shape of the aerosol size
440 distribution. There have been numerous analyses of model-measurement comparison of the aerosol number concentration and size distributions specific to CMAQ (Elleman and Covert, 2009, 2010; Kelly et al., 2011; Zhang et al., 2010b). Elleman and Covert compared the 4km CMAQ v4.4 model's size distributions to measurement data from the 2001 Pacific Northwest and Pacific field campaigns. The Pacific Northwest field campaign (PNW2001) was conducted in August 2001 with both airborne and ground-based measurements of pollution in the Puget Sound urban area around Seattle, Washington, and included
445 northwest Oregon, western Washington, and southwest British Columbia. PNW2001 was conducted to complement that of the Pacific 2001 field campaign, which was a major regional air pollution study in the Lower Fraser Valley of metropolitan Vancouver, British Columbia focusing on ground-based observations, conducted from 10 August to 2 September 2001. Analyses of these two campaigns and model predictions found that CMAQ underpredicted airborne particle number concentrations by a factor of 10-100 and was least accurate in the smallest size mode: the Aitken mode (Elleman and Covert,
450 2009). The underprediction was consistent between measurement studies and did not depend on time and location. Zhang et al compared CMAQ v4.4 to the 1999 Southern Oxidants Study and corroborated the findings of Elleman and Covert, that the


Aitken mode was significantly underpredicted in total number concentration (varying by up to 3 orders of magnitude), yielding an overall underprediction of $PM_{2.5}$ in Atlanta (Zhang et al., 2010b).

455 In a follow-up analysis, Elleman and Covert used updated emissions size distributions to compare a summer 2001 case study comprising data from a period of August 2001 with airborne and surface measurements from Pacific 2001 and PNW2001, as was used in the original base case to CMAQ (Elleman and Covert, 2010). CMAQ still underpredicted the observable aerosol number concentrations by about one order of magnitude with updated emission size distributions, which was an improvement from the 1-2 orders of magnitude previously but pointed to issues within the model's prediction of aerosol number. Kelly et al. then utilized the updated emissions size distributions from Elleman and Covert as well as the original distributions with in
460 CMAQ to compare to the 1998 California Regional $PM_{10}/PM_{2.5}$ Air Quality Study (CRPAQS) (Kelly et al., 2011). It was noted that the simulated number size distributions from the improved emission simulation were about 20% lower than the observations, while the standard-emission simulation was about a factor of 5 lower than the observations, confirming that the updated emissions improved model-measurement agreement. The observed shape of the distributions also better matched the
465 updated emissions simulations. The improvement in model-measurement agreement showcases the necessity for accurate size distributions and emissions within CMAQ and the impact on S_a data.

4.3 Dependence of $S_{a,mod}/S_{a,meas}$ on Secondary Aerosol Production

Two potential reasons for the discrepancy between mean $S_{a,mod}/S_{a,meas}$ (0.55) and mean N_{mod}/N_{meas} (1.11) are: 1) uncertainty
470 in the size distribution of primary aerosol particles, and 2) uncertainty in secondary aerosol production (i.e. the condensation of low volatility material to existing aerosol particles). To investigate these two potential sources, we investigate the response of $S_{a,mod}/S_{a,meas}$ to photochemical age. We start by looking at the response of $S_{a,mod}/S_{a,meas}$ to the NO_x/HNO_3 ratio (Figure 12), where high NO_x/HNO_3 in this sampling region is indicative of air masses near an anthropogenic source, similar to that of a NO_x/NO_y clock (Kleinman et al., 2008; Pan et al., 2015; Tie et al., 2009). If the aerosol surface area of primary emissions is
475 underestimated in the model, we would expect $S_{a,mod}/S_{a,meas}$ to be biased low at high NO_x/HNO_3 . If the condensation rate of low volatility anthropogenic species is underestimated in the model, we would expect $S_{a,mod}/S_{a,meas}$ to decrease with a

decreasing NO_x/HNO_3 ratio as the air mass ages. As shown in Figure 12, $S_{a,\text{mod.}}/S_{a,\text{meas}}$ is remarkably constant over a wide span of NO_x/HNO_3 ratios (0.5-10), before tending to larger values at low NO_x/HNO_3 . This trend is also seen in the dependence of $N_{\text{mod.}}/N_{\text{meas}}$ on NO_x/HNO_3 , suggesting a potential discrepancy in the model-measured lifetime of aerosol or treatment of background aerosol particles in the region. This trend suggests that an underestimate in the condensation of low-volatility gas-phase compounds of anthropogenic origin are not a significant driver of model-measurement discrepancy in S_a . Rather, the persistent underestimate of S_a in the model at high NO_x/HNO_3 points to uncertainty in the size distribution of primary emissions or secondary aerosol formed at the early stages of oxidation.



485 **Figure 12: Interquartile ranges in $S_{a,\text{mod.}}/S_{a,\text{meas}}$ as a function of the modeled (CMAQ) NO_x/HNO_3 concentration ratio.**

To address secondary aerosol formation more generally, we also assessed the response of $S_{a,\text{mod.}}/S_{a,\text{meas}}$ to temperature as equilibrium partitioning in the gas-phase based on temperature and RH is a primary driver of secondary aerosol formation. No statistically significant trend in $S_{a,\text{mod.}}/S_{a,\text{meas}}$ was observed over the range of temperatures observed during DISCOVER-AQ.

490

To further investigate secondary aerosol formation as a factor in driving the discrepancy in modeled S_a , we assess the response of $S_{a,\text{mod.}}/S_{a,\text{meas}}$ to isoprene oxidation products in the aerosol phase as an example of biogenic VOC oxidation. As shown in

Figure 13, there does not appear to be a trend with concentration of isoprene SOA. Though we cannot test for all biogenic oxidation products, the lack of a trend with isoprene SOA in the aerosol phase may mean that the discrepancy in $S_{a,mod.}/S_{a,meas}$ is not biogenic in nature. There is also no trend with the total SOA concentration parameterized in CMAQ, shown in figure 13b.





Figure 13: Interquartile ranges in $S_{a,mod.}/S_{a,meas}$ as a function of the modeled (CMAQ) a) isoprene SOA concentration and b) total SOA concentration.

500 5 Implications for the Treatment of Heterogeneous Reactions in Air Quality Models

As shown in E2, the rate constant for the heterogeneous loss of gas-phase compounds to aerosol (k_{het}) is linearly dependent on both aerosol surface area concentration (S_a) and the reactive uptake coefficient (γ). In section 3, we showed that the average $S_{a,mod.}/S_{a,meas}$, determined from the regression of the average model and measurement S_a was 0.437, which would result in approximately a factor of 2 underestimate in k_{het} . A similar underestimation has been seen previously in select ground (Ghim et al., 2017; Liu and Zhang, 2011; Prank et al., 2016; Wang et al., 2021; Yu et al., 2008a, 2012, 2008b; Zhang et al., 2019, 2006, 2010c) and aircraft-based (Baker et al., 2018; Chen et al., 2020) studies of CMAQ prediction of $PM_{2.5}$, which may point

to a larger issue in model representation of particle mass. For some heterogeneous reactions, where the reactive uptake coefficients are well parameterized in model (e.g., extremely low volatility species) uncertainty in S_a likely determines uncertainty in k_{het} . To assess the dominant source of uncertainty in model derived k_{het} , we focus on the N_2O_5 system as an example. Recently, McDuffie et al. (2018) assessed the accuracy of model parameterizations of $\gamma(N_2O_5)$ using ambient observations from the WINTER campaign. In Figure 14a, we show the histogram and PDF of the ratio of $\gamma(N_2O_5)_{mod}$ calculated in CMAQ using the Bertram and Thornton parameterization for the WINTER campaign, compared with $\gamma(N_2O_5)_{meas}$, which was determined in McDuffie et al. from an observationally constrained analysis of flight data via a box model solved along the flight path to determine the uptake coefficient (McDuffie et al., 2018). The PDF of the directly compared model-measurement ratio is centered above zero ($\mu = 0.22$ or $\gamma(N_2O_5)_{mod}/\gamma(N_2O_5)_{meas} = 1.65$) (Figure 14, Table 3). Interestingly, since $k_{het}(N_2O_5)$ is proportional to the product of S_a and $\gamma(N_2O_5)$, the underestimate in model S_a is compensated by an overestimate in $\gamma(N_2O_5)$ in the mean state if the underestimate in model S_a is consistent for WINTER, though that is not necessarily the case. While the width of the PDF of $\log_{10}(\gamma(N_2O_5)_{mod}/\gamma(N_2O_5)_{meas})$ for WINTER is similar to that seen for S_a for DISCOVER-AQ, it should be noted that neither the histogram of the $\gamma(N_2O_5)$ ratio or the S_a ratio is easily fit to a gaussian peak shape. As shown in Figure 14, the histogram of the $\log_{10}(\gamma(N_2O_5)_{mod}/\gamma(N_2O_5)_{meas})$ for WINTER has a broader range of values than that of S_a in this study. Collectively, this analysis highlights that while model uncertainty in $k_{het}(N_2O_5)$ is largely a function of quality of the $\gamma(N_2O_5)$ parameterization, future improvements in modeled surface area concentrations, particularly in urban environments, will also result in more accurate representations of heterogeneous chemical reactions.



525 **Figure 14: Normalized probability density functions for the \log_{10} of the model to measurement ratio in (a) $\gamma(\text{N}_2\text{O}_5)$ from the WINTER campaign and (b) particle surface area concentration, S_a from this study.**

6. Summary

This study examined the ability of the CMAQ model to accurately predict aerosol surface area as it directly affects
 530 heterogeneous chemistry within the model. The CMAQ data was compared to dry measured aerosol surface area data from the 2011 DISCOVER-AQ campaign utilizing a UHSAS. A discrepancy between modeled and measured dry aerosol surface area, $S_{a,\text{mod.}}$ and $S_{a,\text{meas}}$ respectively, are modestly correlated ($r^2 = 0.52$) and on average agree to within a factor of two ($S_{a,\text{mod.}}/S_{a,\text{meas}} = 0.44$) over the course of the 13 research flights. However, there was a strong correlation between measured and modeled number concentration ($N_{\text{mod.}}/N_{\text{meas}} = 0.87$, $r^2 = 0.63$). When looking into possible sources of the discrepancy, there was not a
 535 strong dependence on photochemical age or secondary biogenic aerosol concentration. The strong agreement in aerosol number concentration may indicate that the modeled size distribution contributes to the observed discrepancy, though the exact source of discrepancy is outside of the scope of this study.

The discrepancy in aerosol surface area was also compared to that of the reactive uptake coefficient of N₂O₅ during the 2015
540 WINTER campaign due to the fact that the uptake coefficient also directly impacts heterogeneous reaction rates. The
uncertainty in the modeled heterogeneous chemistry remains primarily driven by that of the uptake coefficient, as the
uncertainty in those values is larger than that seen by *S_a*. Model improvements to aerosol surface area concentrations along
with improvements to the parameterization of reactive uptake coefficients will greatly impact the accuracy of heterogeneous
chemistry within regional models.

545 **Data Availability**

Data from the 2011 DISCOVER-AQ Campaign is publicly available at <http://doi.org/10.5067/Aircraft/DISCOVER-AQ/Aerosol-TraceGas> CMAQ output data will be made available on the University of Wisconsin – Madison MINDS database at <https://minds.wisconsin.edu/handle/1793/76304>.

Author Contributions

550 Monica Harkey, Alicia Hoffman, and Tracey Holloway ran the CMAQ model and assisted with analysis of model outputs.
Rachel Bergin analyzed the model and measurement data. Rachel Bergin and Timothy Bertram wrote the paper. All authors
reviewed and edited the paper.

Competing Interests

At least one of the (co-)authors is a member of the editorial board of Atmospheric Chemistry and Physics. The authors have
555 no other competing interests to declare.

Acknowledgements

We would like to thank the DISCOVER-AQ NASA Langley Aerosol Research Group Experiment (LARGE) research team,
including Richard Moore, Bruce Anderson, Andreas Beyersdorf, Luke Ziemba, Lee Thornhill, and Edward Winstead for the

use of their data in this work. We would also like to thank Heather Simon, Lyatt Jaeglé, for their data contributions, and Erin
560 McDuffie and Steve Brown specifically for the use of the WINTER data in this work.

Financial Support

This research has been supported by EPA project number R840006 and the NOAA Climate Program Office's Atmospheric
Chemistry, Carbon Cycle, and Climate program (NA18OAR4310109). The NASA LARGE research team was supported by
funding from the NASA Earth Venture Suborbital-1 Program.

565 References

- Abbatt, J. P. D., Lee, A. K. Y. and Thornton, J. A.: Quantifying trace gas uptake to tropospheric aerosol : recent advances and
remaining challenges w, *Chem. Soc. Rev.*, 41, 6555–6581, doi:10.1039/c2cs35052a, 2012.
- Abel, D. W., Holloway, T., Harkey, M., Meier, P., Ahl, D., Limaye, V. S. and Patz, J. A.: Air-quality-related health impacts
from climate change and from adaptation of cooling demand for buildings in the eastern United States: An interdisciplinary
570 modeling study, *PLoS Med.*, 15(7), 1–27, doi:10.1371/journal.pmed.1002599, 2018.
- Abel, D. W., Holloway, T., Martínez-Santos, J., Harkey, M., Tao, M., Kubes, C. and Hayes, S.: Air Quality-Related Health
Benefits of Energy Efficiency in the United States, *Environ. Sci. Technol.*, 53(7), 3987–3998, doi:10.1021/acs.est.8b06417,
2019.
- Adams, P. J. and Seinfeld, J. H.: Predicting global aerosol size distributions in general circulation models, *J. Geophys. Res.*
575 *Atmos.*, 107(19), AAC 4-1-AAC 4-23, doi:10.1029/2001JD001010, 2002.
- Appel, K. W., Bash, J., Fahey, K., Foley, K., Gilliam, R., Hogrefe, C., Hutzell, W., Kang, D., Mathur, R., Murphy, B.,
Napelenok, S., Nolte, C., Pleim, J., Pouliot, G., Pye, H., Ran, L., Roselle, S., Sarwar, G., Schwede, D., Sidi, F., Spero, T. and
Wong, D.: The Community Multiscale Air Quality (CMAQ) Model Versions 5.3 and 5.3.1: System Updates and Evaluation,
Geosci. Model Dev. Discuss., 1–41, doi:10.5194/gmd-2020-345, 2021.
- 580 Baker, K. R., Woody, M. C., Valin, L., Szykman, J., Yates, E. L., Iraci, L. T., Choi, H. D., Soja, A. J., Koplitz, S. N., Zhou,
L., Campuzano-Jost, P., Jimenez, J. L. and Hair, J. W.: Photochemical model evaluation of 2013 California wild fire air quality
impacts using surface, aircraft, and satellite data, *Sci. Total Environ.*, 637–638, 1137–1149,
doi:10.1016/j.scitotenv.2018.05.048, 2018.
- Bates, T. S., Quinn, P. K., Coffman, D., Schulz, K., Covert, D. S., Johnson, J. E., Williams, E. J., Lerner, B. M., Angevine, W.
585 M., Tucker, S. C., Brewer, W. A. and Stohl, A.: Boundary layer aerosol chemistry during TexAQS/GoMACCS 2006: Insights
into aerosol sources and transformation processes, *J. Geophys. Res.*, 113, 1–18, doi:10.1029/2008jd010023, 2008.

- Bertram, T. H., and Thornton, J. A.: Toward a general parameterization of N₂O₅ reactivity on aqueous particles: the competing effects of particle liquid water, nitrate, and chloride, *Atmos. Chem. Phys.*, 9(21), 8351–8363, doi:10.5194/acp-9-8351-2009, 2009.
- 590 Binkowski, F. S.: Chapter 10 - AEROSOLS IN MODELS-3 CMAQ, Washington, DC., 1999.
- Binkowski, F. S. and Roselle, S. J.: Models-3 Community Multiscale Air Quality (CMAQ) model aerosol component 1. Model description, *J. Geophys. Res. Atmos.*, 108(6), doi:10.1029/2001jd001409, 2003.
- Brown, S. S., Dubé, W. P., Fuchs, H., Ryerson, T. B., Wollny, A. G., Brock, C. A., Bahreini, R., Middlebrook, A. M., Neuman, T. A., Atlas, E., Roberts, J. M., Osthoff, H. D., Trainer, M., Fehsenfeld, F. C. and Ravishankara, A. R.: Reactive uptake
595 coefficients for N₂O₅ determined from aircraft measurements during the Second Texas Air Quality Study: Comparison to current model parameterizations, *J. Geophys. Res. Atmos.*, 114(11), 1–16, doi:10.1029/2008JD011679, 2009.
- Byun, D., and Schere, K. L.: Review of the governing equations, computational algorithms, and other components of the Models-3 Community Multiscale Air Quality (CMAQ) modeling, *Appl. Mech. Rev.*, 59(2), 51–77, doi:10.1115/1.2128636, 2006.
- 600 Carlton, A. G., Bhave, P. V., Napelenok, S. L., Edney, E. O., Sarwar, G., Pinder, R. W., Pouliot, G. A. and Houyoux, M.: Model representation of secondary organic aerosol in CMAQv4.7, *Environ. Sci. Technol.*, 44(22), 8553–8560, doi:10.1021/es100636q, 2010.
- Chang, W. L., Bhave, P. V., Brown, S. S., Riemer, N., Stutz, J. and Dabdub, D.: Heterogeneous atmospheric chemistry, ambient measurements, and model calculations of N₂O₅: A review, *Aerosol Sci. Technol.*, 45(6), 665–695,
605 doi:10.1080/02786826.2010.551672, 2011.
- Chen, J., Yin, D., Zhao, Z., Kaduwela, A. P., Avise, J. C., DaMassa, J. A., Beyersdorf, A., Burton, S., Ferrare, R., Herman, J. R., Kim, H., Neuman, A., Nowak, J. B., Parworth, C., Scarino, A. J., Wisthaler, A., Young, D. E. and Zhang, Q.: Modeling air quality in the San Joaquin valley of California during the 2013 Discover-AQ field campaign, *Atmos. Environ. X*, 5, 100067, doi:10.1016/j.aeaoa.2020.100067, 2020.
- 610 Crawford, J. H. and Pickering, K. E.: DISCOVER-AQ: Advancing strategies for air quality observations in the next decade, *EM Air Waste Manag. Assoc. Mag. Environ. Manag.*, (September), 4–7, 2014.
- Crawford, J. H., Dickerson, R. R. and Hains, J. C.: DISCOVER-AQ: Observations and early results, *EM Air Waste Manag. Assoc. Mag. Environ. Manag.*, (September), 8–15, 2014.
- Davis, J. M., Bhave, P. V., and Foley, K. M.: Parameterization of N₂O₅ reaction probabilities on the surface of particles
615 containing ammonium, sulfate, and nitrate, *Atmos. Chem. Phys.*, 8(17), 5295–5311, doi:10.5194/acp-8-5295-2008, 2008.
- Donahue, N. M., Robinson, A. L., Trump, E. R., Riipinen, I. and Kroll, J. H.: Volatility and Aging of Atmospheric Organic Aerosol, in *Atmospheric and Aerosol Chemistry*, edited by V. F. McNeill and P. A. Ariya, pp. 97–143, Springer Berlin Heidelberg, Berlin, Heidelberg., 2014.
- Elleman, R. A. and Covert, D. S.: Aerosol size distribution modeling with the Community Multiscale Air Quality modeling
620 system in the Pacific Northwest: 1. Model comparison to observations, *J. Geophys. Res. Atmos.*, 114(11),

- doi:10.1029/2008JD010791, 2009.
- Elleman, R. A., and Covert, D. S.: Aerosol size distribution modeling with the Community Multiscale Air Quality modeling system in the Pacific Northwest: 2. Parameterizations for ternary nucleation and nucleation mode processes, *J. Geophys. Res.*, 114, D11207, doi:10.1029/2009JD012187, 2009b.
- 625 Elleman, R. A. and Covert, D. S.: Aerosol size distribution modeling with the Community Multiscale Air Quality modeling system in the Pacific Northwest: 3. Size distribution of particles emitted into a mesoscale model, *J. Geophys. Res. Atmos.*, 115(3), 1–14, doi:10.1029/2009JD012401, 2010.
- Emery, C., Jung, J., Koo, B., and Yarwood, G.: Improvements to CAMx snow cover treatments and carbon bond chemical mechanism for winter ozone, *Ramboll Environ*, 43pp., Report Number UDAQ PO 480 52000000001, 2015.
- 630 Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J. F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L. and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geosci. Model Dev.*, 3(1), 43–67, doi:10.5194/gmd-3-43-2010, 2010.
- Evans, M. J. and Jacob, D. J.: Impact of new laboratory studies of N₂O₅ hydrolysis on global model budgets of tropospheric
635 nitrogen oxides, ozone, and OH, *Geophys. Res. Lett.*, 32(9), 1–4, doi:10.1029/2005GL022469, 2005.
- Fan, J., Zhang, R., Li, G., Nielsen-Gammon, J. and Li, Z.: Simulations of fine particulate matter (PM_{2.5}) in Houston, Texas, *J. Geophys. Res. D Atmos.*, 110(16), 1–9, doi:10.1029/2005JD005805, 2005.
- Foley, K. M., Roselle, S. J., Appel, K. W., Bhave, P. V., Pleim, J. E., Otte, T. L., Mathur, R., Sarwar, G., Young, J. O., Gilliam, R. C., Nolte, C. G., Kelly, J. T., Gilliland, A. B., and Bash, J. O.: Incremental testing of the Community Multiscale Air Quality
640 (CMAQ) modeling system version 4.7, *Geosci. Model Dev.*, 3(1), 205–226, doi:10.5194/gmd-3-205-2010, 2010.
- Gantt, B., Johnson, M. S., Meskhidze, N., Sciare, J., Ovadnevaite, J., Ceburnis, D. and O’Dowd, C. D.: Model evaluation of marine primary organic aerosol emission schemes, *Atmos. Chem. Phys.*, 12(18), 8553–8566, doi:10.5194/acp-12-8553-2012, 2012.
- Gaston, C. J., Riedel, T. P., Zhang, Z., Gold, A., Surratt, J. D. and Thornton, J. A.: Reactive uptake of an isoprene-derived
645 epoxydiol to submicron aerosol particles, *Environ. Sci. Technol.*, 48(19), 11178–11186, doi:10.1021/es5034266, 2014.
- Gelbard, F., Tambour, Y. and Seinfeld, J. H.: Sectional representations for simulating aerosol dynamics, *J. Colloid Interface Sci.*, 76(2), 541–556, doi:10.1016/0021-9797(80)90394-X, 1980.
- George, I. J., Matthews, P. S. J., Whalley, L. K., Brooks, B., Goddard, A., Baeza-Romero, M. T. and Heard, D. E.:
650 Measurements of uptake coefficients for heterogeneous loss of HO₂ onto submicron inorganic salt aerosols, *Phys. Chem. Chem. Phys.*, 15, 12829–12845, doi:10.1039/c3cp51831k, 2013.
- Geyer, A. and Stutz, J.: Vertical profiles of NO₃, N₂O₅, O₃, and NO_x in the nocturnal boundary layer: 2. Model studies on the altitude dependence of composition and chemistry, *J. Geophys. Res. D Atmos.*, 109(12), 1–18, doi:10.1029/2003JD004211, 2004.
- Ghan, S., Laulainen, N., Easter, R., Wagener, R., Nemesure, S., Chapman, E., Zhang, Y. and Leung, R.: Evaluation of aerosol

- 655 direct radiative forcing in MIRAGE, *J. Geophys. Res.*, 106(D6), 5295–5316, 2001.
- Ghim, Y. S., Choi, Y., Kim, S., Bae, C. H., Park, J. and Shin, H. J.: Evaluation of model performance for forecasting fine particle concentrations in Korea, *Aerosol Air Qual. Res.*, 17(7), 1856–1864, doi:10.4209/aaqr.2016.10.0446, 2017.
- Guenther, A.B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling
660 biogenic emissions, *Geosci. Model Dev.*, 5, 1471–1492, doi:10.5194/gmd-5-1471-2012, 2012.
- Harkey, M., Holloway, T., Kim, E. J., Baker, K. R. and Henderson, B.: Satellite Formaldehyde to Support Model Evaluation, *J. Geophys. Res. Atmos.*, 126(4), 1–18, doi:10.1029/2020JD032881, 2021.
- Hogrefe, C., Lynn, B., Goldberg, R., Rosenzweig, C., Zalewsky, E., Hao, W., Doraiswamy, P., Civerolo, K., Ku, J. Y., Sistla, G. and Kinney, P. L.: A combined model-observation approach to estimate historic gridded fields of PM_{2.5} mass and species
665 concentrations, *Atmos. Environ.*, 43(16), 2561–2570, doi:10.1016/j.atmosenv.2009.02.031, 2009.
- Hogrefe, C., Pouliot, G., Wong, D., Torian, A., Roselle, S., Pleim, J. and Mathur, R.: Annual application and evaluation of the online coupled WRF-CMAQ system over North America under AQMEII phase 2, *Atmos. Environ.*, 115, 683–694, doi:10.1016/j.atmosenv.2014.12.034, 2015.
- Jacobson, M. Z.: GATOR-GCMM: A global- through urban-scale air pollution and weather forecast model 1. Model design
670 and treatment of subgrid soil, vegetation, roads, rooftops, water, sea ice, and snow, *J. Geophys. Res. Atmos.*, 106(D6), 5385–5401, doi:10.1029/2000JD900560, 2001.
- Jaeglé, L., Shah, V., Thornton, J. A., Lopez-Hilfiker, F. D., Lee, B. H., McDuffie, E. E., Fibiger, D., Brown, S. S., Veres, P., Sparks, T. L., Ebben, C. J., Wooldridge, P. J., Kenagy, H. S., Cohen, R. C., Weinheimer, A. J., Campos, T. L., Montzka, D. D., Digangi, J. P., Wolfe, G. M., Hanisco, T., Schroder, J. C., Campuzano-Jost, P., Day, D. A., Jimenez, J. L., Sullivan, A. P.,
675 Guo, H. and Weber, R. J.: Nitrogen Oxides Emissions, Chemistry, Deposition, and Export Over the Northeast United States During the WINTER Aircraft Campaign, *J. Geophys. Res. Atmos.*, 123(21), 12,368–12,393, doi:10.1029/2018JD029133, 2018.
- Kelly, J. T., Avise, J., Cai, C. and Kaduwela, A. P.: Simulating particle size distributions over California and impact on lung deposition fraction, *Aerosol Sci. Technol.*, 45(2), 148–162, doi:10.1080/02786826.2010.528078, 2011.
- 680 Kleeman, M. J., Cass, G. R. and Eldering, A.: Modeling the airborne particle complex as a source-oriented external mixture, *J. Geophys. Res. Atmos.*, 102(17), 21355–21372, doi:10.1029/97jd01261, 1997.
- Kleinman, L. I., Springston, S. R., Daum, P. H., Lee, Y. N., Nunnermacker, L. J., Senum, G. I., Wang, J., Weinstein-Lloyd, J., Alexander, M. L., Hubbe, J., Ortega, J., Canagaratna, M. R. and Jayne, J.: The time evolution of aerosol composition over the Mexico City plateau, *Atmos. Chem. Phys.*, 8(6), 1559–1575, doi:10.5194/acp-8-1559-2008, 2008.
- 685 Knote, C., Brunner, D., Vogel, H., Allan, J., Asmi, A., Äijälä, M., Carbone, S., van der Gon, H. D., Jimenez, J. L., Kiendler-Scharr, A., Mohr, C., Poulain, L., Prévôt, A. S. H., Swietlicki, E. and Vogel, B.: Towards an online-coupled chemistry-climate model: evaluation of trace gases and aerosols in COSMO-ART, *Geosci. Model Dev.*, 4(4), 1077–1102, doi:10.5194/gmd-4-1077-2011, 2011.

- Kulmala, M., Laaksonen, A. and Pirjola, L.: Parameterizations for sulfuric acid/water nucleation rates, *J. Geophys. Res. Atmos.*, 103(D7), 8301–8307, doi:10.1029/97JD03718, 1998.
- 690 Lakey, P. S. J., George, I. J., Whalley, L. K., Baeza-Romero, M. T. and Heard, D. E.: Measurements of the HO₂ Uptake Coefficients onto Single Component Organic Aerosols, *Environ. Sci. Technol.*, 49, 4878–4885, doi:10.1021/acs.est.5b00948, 2015.
- Lee, Y. H. and Adams, P. J.: A fast and efficient version of the Two-Moment Aerosol Sectional (TOMAS) global aerosol microphysics model, *Aerosol Sci. Technol.*, 46(6), 678–689, doi:10.1080/02786826.2011.643259, 2012.
- 695 Lee, Y. H., Chen, K. and Adams, P. J.: Development of a global model of mineral dust aerosol microphysics, *Atmos. Chem. Phys.*, 9(7), 2441–2458, doi:10.5194/acp-9-2441-2009, 2009.
- Liu, P. and Zhang, Y.: Use of a process analysis tool for diagnostic study on fine particulate matter predictions in the U.S. - Part I: Model evaluation, *Atmos. Pollut. Res.*, 2(1), 49–60, doi:10.5094/APR.2011.007, 2011.
- 700 Luecken, D. J., Yarwood, G., and Hutzell, W. T.: Multipollutant modeling of ozone, reactive nitrogen and HAPs across the continental US with CMAQ-CB6, *Atmos. Environ.*, 201, 62-72, doi:10.1016/j.atmosenv.2018.11.060, 2019.
- Luo, G. and Yu, P.: Simulation of particle formation and number concentration over the Eastern United States with the WRF-Chem + APM model, *Atmos. Chem. Phys.*, 11(22), 11521-11533, doi:10.5194/acp-11-11521-2011, 2011.
- MacIntyre, H. L. and Evans, M. J.: Sensitivity of a global model to the uptake of N₂O₅ by tropospheric aerosol, *Atmos. Chem. Phys.*, 10(15), 7409–7414, doi:10.5194/acp-10-7409-2010, 2010.
- 705 Mann, G. W., Carslaw, K. S., Spracklen, D. V., Ridley, D. A., Manktelow, P. T., Chipperfield, M. P., Pickering, S. J. and Johnson, C. E.: Description and evaluation of GLOMAP-mode: A modal global aerosol microphysics model for the UKCA composition-climate model, *Geosci. Model Dev.*, 3(2), 519–551, doi:10.5194/gmd-3-519-2010, 2010.
- Martin, R. V., Jacob, D. J. and Yantosca, R. M.: Global and regional decreases in tropospheric oxidants from photochemical effects of aerosols, *J. Geophys. Res.*, 108, doi:10.1029/2002JD002622, 2003.
- 710 McDuffie, E. E., Fibiger, D. L., Dubé, W. P., Lopez-Hilfiker, F., Lee, B. H., Thornton, J. A., Shah, V., Jaeglé, L., Guo, H., Weber, R. J., Michael Reeves, J., Weinheimer, A. J., Schroder, J. C., Campuzano-Jost, P., Jimenez, J. L., Dibb, J. E., Veres, P., Ebben, C., Sparks, T. L., Wooldridge, P. J., Cohen, R. C., Hornbrook, R. S., Apel, E. C., Campos, T., Hall, S. R., Ullmann, K. and Brown, S. S.: Heterogeneous N₂O₅ Uptake During Winter: Aircraft Measurements During the 2015 WINTER Campaign and Critical Evaluation of Current Parameterizations, *J. Geophys. Res. Atmos.*, 123(8), 4345–4372, doi:10.1002/2018JD028336, 2018.
- Meng, Z.: of Atmospheric Aerosol Dynamics, *Environ. Eng.*, 103, 3419–3435, 1998.
- Messinger, F., DiMego, G., Kalnay, E., Mitchell, K., Shafran, P. C., Ebisuzaki, W., Jovic, D., Woollen, J., Rogers, E., Berbery, E. H., Ek, M. B., Fan, Y., Grumbine, R., Higgins, W., Li, H., Lin, Y., Manikin, G., Parrish, D., and Shi, W.: North American Regional Reanalysis, *Bull. Am. Meteorol.*, 343-360, doi:10.1175/BAMS-87-3-343, 2006.
- 720 Moore, R. H., Wiggins, E. B., Ahern, A. T., Zimmerman, S., Montgomery, L., Campuzano Jost, P., Robinson, C. E., Ziemba, L. D., Winstead, E. L., Anderson, B. E., Brock, C. A., Brown, M. D., Chen, G., Crosbie, E. C., Guo, H., Jimenez, J. L., Jordan,

- C. E., Lyu, M., Nault, B. A., Rothfuss, N. E., Sanchez, K. J., Schueneman, M., Shingler, T. J., Shook, M. A., Thornhill, K. L., Wagner, N. L. and Wang, J.: Sizing response of the Ultra-High Sensitivity Aerosol Spectrometer (UHSAS) and Laser Aerosol Spectrometer (LAS) to changes in submicron aerosol composition and refractive index, *Atmos. Meas. Tech.*, 14(6), 4517–4542, doi:10.5194/amt-14-4517-2021, 2021.
- NASA: DISCOVER-AQ: Diagnosing the Air We Breathe, [online] Available from: <http://www.nasa.gov/discover-aq>, 2012.
- NEIC: National Emissions Inventory Collaborative, 2016v1 Emiss. Model. Platf. [online] Available from: <http://views.cira.colostate.edu/wiki/wiki/10202>, 2019.
- 725 Nolte, C. G., Appel, K. W., Kelly, J. T., Bhawe, P. V., Fahey, K. M., Collett, J. L., Zhang, L. and Young, J. O.: Evaluation of the Community Multiscale Air Quality (CMAQ) model v5.0 against size-resolved measurements of inorganic particle composition across sites in North America, *Geosci. Model Dev.*, 8(9), 2877–2892, doi:10.5194/gmd-8-2877-2015, 2015.
- Pan, X., Kanaya, Y., Tanimoto, H., Inomata, S., Wang, Z., Kudo, S. and Uno, I.: Examining the major contributors of ozone pollution in a rural area of the Yangtze River Delta region during harvest season, *Atmos. Chem. Phys.*, 15(11), 6101–6111, doi:10.5194/acp-15-6101-2015, 2015.
- 735 Park, S. K., Marmur, A., Kim, S. B., Tian, D., Hu, Y., McMurry, P. H. and Russell, A. G.: Evaluation of fine particle number concentrations in CMAQ, *Aerosol Sci. Technol.*, 40(11), 985–996, doi:10.1080/02786820600907353, 2006.
- Portmann, R. W., Solomon, S., Garcia, R. R., Thomason, L. W., Poole, L. R. and McCormick, M. P.: Role of aerosol variations in anthropogenic ozone depletion in the polar regions, *J. Geophys. Res. Atmos.*, 101(17), 22991–23006, doi:10.1029/96jd02608, 1996.
- 740 Prank, M., Sofiev, M., Tsyro, S., Hendriks, C., Semeena, V., Francis, X. V., Butler, T., Van Der Gon, H. D., Friedrich, R., Hendricks, J., Kong, X., Lawrence, M., Righi, M., Samaras, Z., Sausen, R., Kukkonen, J. and Sokhi, R.: Evaluation of the performance of four chemical transport models in predicting the aerosol chemical composition in Europe in 2005, *Atmos. Chem. Phys.*, 16(10), 6041–6070, doi:10.5194/acp-16-6041-2016, 2016.
- 745 Pringle, K. J., Tost, H., Message, S., Steil, B., Giannadaki, D., Nenes, A., Fountoukis, C., Stier, P., Vignati, E. and Lelieveld, J.: Description and evaluation of GMXe: A new aerosol submodel for global simulations (v1), *Geosci. Model Dev.*, 3(2), 391–412, doi:10.5194/gmd-3-391-2010, 2010.
- Pye, H. O. T., Luecken, D. J., Xu, L., Boyd, C. M., Ng, N. L., Baker, K. R., Ayres, B. R., Bash, J. O., Baumann, K., Carter, W. P. L., Edgerton, E., Fry, J. L., Hutzell, W. T., Schwede, D. B., and Shepson, P. B.: Modeling the current and future roles of particulate organic nitrates in the southeastern United States, *Environ. Sci. Technol.*, 49(24), 14195–14203, doi:10.1021/acs.est.5b03738, 2015.
- 750 Pye, H. O. T., Murphy, B. N., Xu, L., Ng, N. L., Carlton, A. G., Guo, H., Weber, R., Vasilakos, P., Appel, K. W., Budisulistiorini, S. H., Surratt, J. D., Nenes, A., Hu, W., Jimenez, J. L., Isaacman-VanWertz, G., Misztal, P. K., and Goldstein, A. H.: On the implications of aerosol liquid water and phase separation for organic aerosol mass, *Atmos. Chem. Phys.*, 17(1), 343–369, doi:10.5194/acp-17-343-2017, 2017.
- 755

- Qin, M., Murphy, B. N., Isaacs, K. K., McDonald, B. C., Lu, Q., McKeen, S. A., Koval, L., Robinson, A. L., Efstathiou, C., Allen, C., and Pye, H. O. T.: Criteria pollutant impacts of volatile chemical products informed by near-field modeling, *Nat. Sustain.*, 4, 129-137, doi:10.1038/s41893-020-00614-1, 2021.
- 760 Ranjithkumar, A., Gordon, H., Williamson, C., Rollins, A., Pringle, K., Kupc, A., Luke Abraham, N., Brock, C. and Carslaw, K.: Constraints on global aerosol number concentration, SO₂ and condensation sink in UKESM1 using ATom measurements, *Atmos. Chem. Phys.*, 21(6), 4979–5014, doi:10.5194/acp-21-4979-2021, 2021.
- Ridley, B. A. and Grahek, F. E.: A small, low-flow, high-sensitivity reaction vessel for NO chemiluminescence detectors, *J. Atmos. Ocean. Technol.*, 7(2), 307–311, doi:10.1175/1520-0426(1990)007<0307:aslfhs>2.0.co;2, 1990.
- 765 Sachse, G., Hill, G., Wade, L. and Perry, M.: Fast-response, high-precision carbon monoxide sensor using a tunable diode laser absorption technique, *J. Geophys. Res. Atmos.*, 92(D2), 2071–2081, doi:10.1029/JD092iD02p02071, 1987.
- Sartelet, K. N., Hayami, H., Albriet, B. and Sportisse, B.: Development and preliminary validation of a modal aerosol model for tropospheric chemistry: MAM, *Aerosol Sci. Technol.*, 40(2), 118–127, doi:10.1080/02786820500485948, 2006.
- 770 Simon, H., Kimura, Y., McGaughey, G., Allen, D. T., Brown, S. S., Coffman, D., Dibb, J., Osthoff, H. D., Quinn, P., Roberts, J. M., Yarwood, G., Kemball-Cook, S., Byun, D. and Lee, D.: Modeling heterogeneous ClNO₂ formation, chloride availability, and chlorine cycling in Southeast Texas, *Atmos. Environ.*, 44(40), 5476–5488, doi:10.1016/j.atmosenv.2009.09.006, 2010.
- Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Duda, M. G., Huang, X.-Y., Wang, W., and Powers, J. G.: A description of the Advanced Research WRF Version 3, 113 pp., NCAR Tech. Note NCAR/TN-475+STR, doi:10.5065/D68S4MVH, 2008.
- 775 Smyth, S. C., Jiang, W., Yin, D., Roth, H. and Giroux, É.: Evaluation of CMAQ O₃ and PM_{2.5} performance using Pacific 2001 measurement data, *Atmos. Environ.*, 40(15), 2735–2749, doi:10.1016/j.atmosenv.2005.10.068, 2006.
- Sonntag, D. B., Baldauf, R. W., Yanca, C. A., and Fulper, C. R.: Particulate matter speciation profiles for light-duty gasoline vehicles in the United States, *J. Air Waste Manage. Assoc.*, 64(5), 529-545, doi:10.1080/10962247.2013.870096, 2014.
- Spak, S. N. and Holloway, T.: Seasonality of speciated aerosol transport over the Great Lakes region, *J. Geophys. Res. Atmos.*, 114(8), 1–18, doi:10.1029/2008JD010598, 2009.
- 780 Spracklen, D. V., Carslaw, K. S., Kulmala, M., Kerminen, V. M., Mann, G. W. and Sihto, S. L.: The contribution of boundary layer nucleation events to total particle concentrations on regional and global scales, *Atmos. Chem. Phys.*, 6(12), 5631–5648, doi:10.5194/acp-6-5631-2006, 2006.
- Stadtler, S., Simpson, D., Schröder, S., Taraborrelli, D., Bott, A. and Schultz, M.: Ozone impacts of gas-aerosol uptake in global chemistry transport models, *Atmos. Chem. Phys.*, 18(5), 3147–3171, doi:10.5194/acp-18-3147-2018, 2018.
- 785 Stier, P., Feichter, J., Kinne, S., Kloster, S., Vignati, E., Wilson, J., Ganzeveld, L., Tegen, I. and Werner, M.: The aerosol-climate model ECHAM5-HAM, *Atmos. Chem. Phys.*, 5, 1125–1156, 2005.
- Thornton, J. and Abbatt, J. P. D.: Measurements of HO₂ uptake to aqueous aerosol: Mass accommodation coefficients and net reactive loss, *J. Geophys. Res.*, 110, 1–12, doi:10.1029/2004JD005402, 2005.
- Thornton, J. A., Jaegle, L. and McNeill, V. F.: Assessing known pathways for HO₂ loss in aqueous atmospheric aerosols:

- 790 Regional and global impacts on tropospheric oxidants, *J. Geophys. Res.*, 113, 1–16, doi:10.1029/2007JD009236, 2008.
- Tie, X., Madronich, S., Li, G., Ying, Z., Weinheimer, A., Apel, E. and Campos, T.: Simulation of Mexico City plumes during the MIRAGE-Mex field campaign using the WRF-Chem model, *Atmos. Chem. Phys.*, 9(14), 4621–4638, doi:10.5194/acp-9-4621-2009, 2009.
- Trivitayanurak, W., Adams, P. J., Spracklen, D. V. and Carslaw, K. S.: Tropospheric aerosol microphysics simulation with
795 assimilated meteorology: Model description and intermodel comparison, *Atmos. Chem. Phys.*, 8(12), 3149–3168, doi:10.5194/acp-8-3149-2008, 2008.
- Trump, E. R., Riipinen, I. and Donahue, N. M.: Interactions between atmospheric ultrafine particles and secondary organic aerosol mass: A model study, *Boreal Environ. Res.*, 19(5–6), 352–362, 2014.
- Vignati, E., Wilson, J. and Stier, P.: M7: An efficient size-resolved aerosol microphysics module for large-scale aerosol
800 transport models, *J. Geophys. Res. D Atmos.*, 109(22), 1–17, doi:10.1029/2003JD004485, 2004.
- Wang, M., Penner, J. E. and Liu, X.: Coupled IMPACT aerosol and NCAR CAM3 model: Evaluation of predicted aerosol number and size distribution, *J. Geophys. Res. Atmos.*, 114(6), 1–30, doi:10.1029/2008JD010459, 2009.
- Wang, X., Li, L., Gong, K., Mao, J., Hu, J., Li, J., Liu, Z., Liao, H., Qiu, W., Yu, Y., Dong, H., Guo, S., Hu, M., Zeng, L. and Zhang, Y.: Modelling air quality during the EXPLORE-YRD campaign – Part I. Model performance evaluation and impacts
805 of meteorological inputs and grid resolutions, *Atmos. Environ.*, 246(December 2020), 118131, doi:10.1016/j.atmosenv.2020.118131, 2021.
- Whitby, K. T.: The physical characteristics of sulfur aerosols, *Atmos. Environ.*, 12, 135–159, doi:10.1016/j.atmosenv.2007.10.057, 1978.
- Xu, L., Pye, H. O. T., He, J., Chen, Y., Murphy, B. N., and Ng, N. L.: Experimental and model estimates of the contributions
810 from biogenic monoterpenes and sesquiterpenes to secondary organic aerosol in the southeastern United States, *Atmos. Chem. Phys.*, 18(17), 12613–12637, doi:10.5194/acp-18-12613-2018, 2018.
- Yu, F. and Luo, G.: Simulation of particle size distribution with a global aerosol model: contribution of nucleation to aerosol and CCN number concentrations, *Atmos. Chem. Phys.*, 9(20), 7691–7710, doi:10.5194/acp-9-7691-2009, 2009.
- Yu, S., Mathur, R., Schere, K., Kang, D., Pleim, J., Young, J., Tong, D., Pouliot, G., McKeen, S. A. and Rao, S. T.: Evaluation
815 of real-time PM_{2.5} forecasts and process analysis for PM_{2.5} formation over the eastern United States using the Eta-CMAQ forecast model during the 2004 ICARTT study, *J. Geophys. Res. Atmos.*, 113(6), doi:10.1029/2007JD009226, 2008a.
- Yu, S., Mathur, R., Pleim, J., Pouliot, G., Wong, D., Eder, B., Schere, K., Gilliam, R. and Rao, S. T.: Comparative evaluation of the impact of WRF/NMM and WRF/ARW meteorology on CMAQ simulations for PM_{2.5} and its related precursors during the 2006 TexAQS/GoMACCS study, *Atmos. Chem. Phys.*, 12(9), 4091–4106, doi:10.5194/acp-12-4091-2012, 2012.
- 820 Yu, Y., Sokhi, R. S., Kitwiroon, N., Middleton, D. R. and Fisher, B.: Performance characteristics of MM5-SMOKE-CMAQ for a summer photochemical episode in southeast England, United Kingdom, *Atmos. Environ.*, 42(20), 4870–4883, doi:10.1016/j.atmosenv.2008.02.051, 2008b.
- Zhang, K., Wan, H., Wang, B., Zhang, M., Feichter, J. and Liu, X.: Tropospheric aerosol size distributions simulated by three

- online global aerosol models using the M7 microphysics module, *Atmos. Chem. Phys.*, 10(13), 6409–6434, doi:10.5194/acp-
825 10-6409-2010, 2010a.
- Zhang, Q., Xue, D., Liu, X., Gong, X. and Gao, H.: Process analysis of PM 2.5 pollution events in a coastal city of China using CMAQ, *J. Environ. Sci. (China)*, 79, 225–238, doi:10.1016/j.jes.2018.09.007, 2019.
- Zhang, Y., Liu, P., Queen, A., Misenis, C., Pun, B., Seigneur, C. and Wu, S. Y.: A comprehensive performance evaluation of MM5-CMAQ for the Summer 1999 Southern Oxidants Study episode-Part II: Gas and aerosol predictions, *Atmos. Environ.*,
830 40(26), 4839–4855, doi:10.1016/j.atmosenv.2005.12.048, 2006.
- Zhang, Y., Liu, P., Liu, X. H., Jacobson, M. Z., McMurry, P. H., Yu, F., Yu, S. and Schere, K. L.: A comparative study of nucleation parameterizations: 2. Three-dimensional model application and evaluation, *J. Geophys. Res. Atmos.*, 115(20), doi:10.1029/2010JD014151, 2010b.
- Zhang, Y., Liu, P., Liu, X. H., Pun, B., Seigneur, C., Jacobson, M. Z. and Wang, W. X.: Fine scale modeling of wintertime
835 aerosol mass, number, and size distributions in central California, *J. Geophys. Res. Atmos.*, 115(15), doi:10.1029/2009JD012950, 2010c.