



Measurement Report: Increasing trend of atmospheric ion concentrations in the boreal forest

Juha Sulo¹, Janne Lampilahti¹, Xuemeng Chen², Jenni Kontkanen^{1,3}, Tuomo Nieminen^{1,4}, Veli-Matti Kerminen¹, Tuukka Petäjä¹, Markku Kulmala^{1,5}, and Katrianne Lehtipalo^{1,6}

5

¹Institute for Atmospheric and Earth System Research/Physics, University of Helsinki, 00014 Helsinki, Finland

²Laboratory of Environmental Physics, University of Tartu, Tartu, Estonia

³CSC - IT Center for Science Ltd., Espoo, Finland

10 ⁴Institute for Atmospheric and Earth System Research / Forest Sciences, Faculty of Agriculture and Forestry, University of Helsinki, 00014 Helsinki, Finland

⁵Aerosol and Haze Laboratory, Beijing Advanced Innovation Center for Soft Matter Science and Engineering, Beijing University of Chemical Technology, Beijing, China

⁶Finnish Meteorological Institute, 00560 Helsinki, Finland

Correspondence to: Juha Sulo (juha.sulo@helsinki.fi)

15 **Abstract.** The concentration of atmospheric ions affects the total aerosol particle number concentrations in the atmosphere as well as atmospheric new particle formation via ion-induced nucleation, ion-ion recombination, and effects on condensational growth. In this study, we investigate the concentrations and long-term trends of atmospheric ions in a boreal forest environment using 16 years of cluster ion (0.8 – 2 nm) and intermediate ion (2 – 7 nm) measurements and characterize the most important factors that explain those trends. We found that the median concentration of cluster ions in a boreal forest
20 was 710 cm⁻³, the median concentration of 2 – 4 intermediate ions was 14 cm⁻³ and the median concentration of 4 – 7 nm intermediate ions was 9 cm⁻³. The concentrations of both cluster and intermediate ions have been increasing over the 16-year measurement period, with cluster ion concentrations increasing by about 1% per year and intermediate ion concentrations increasing 1.7 – 3.9 % a year. The increase in cluster-ion concentrations can be best explained by the decrease in the coagulation sink caused by larger aerosol particles. Meanwhile, the dependence of intermediate ion concentrations on
25 meteorological conditions is evident, but ionization sources and the coagulation sink do not seem to explain the increasing trend. This is likely because the dynamics of intermediate ions are more complicated, so that ionization sources and coagulation sink alone cannot directly explain the variation. Season-specific analysis of the ion concentrations suggests that while coagulation sink is the limiting factor for the ion concentrations in spring and summer, the dynamics are different in autumn and winter. Based on our findings, we recommend that a more comprehensive analysis is needed to determine if the
30 increase in ambient ion concentrations, increasing temperature and changing abundance of condensable vapors makes ion-mediated and ion-induced nucleation pathways in the boreal forest more relevant in the future.



1 Introduction

35 Atmospheric ions are produced via the ionization of air molecules (Rutherford, 1897), typically the ionization of nitrogen and oxygen due to their abundance in the atmosphere (Israël, 1970). The three main atmospheric ionization sources are radon decay, terrestrial gamma radiation and cosmic rays (Eisenbud and Gesell, 1997; Mironova et al., 2015; Bazilevskaya et al., 2008). Anthropogenic sources of ionizing radiation come mostly from radiation leaks from nuclear power plants and nuclear weapon detonations (WHO, 1968). The charge can then be neutralized by ion-ion recombination, lost to a foreign surface or transferred to other charged or neutral molecules, clusters or aerosol particles (Seinfeld and Pandis, 1998; Tammet et al., 40 2006). Of the charged particles, cluster ions (0.8 – 2 nm) exist in the atmosphere all the time (Venzac et al., 2007, 2008; Mirme et al., 2010). Intermediate ions (2 – 7 nm) are typically formed by the propagation of charge onto initially neutral particles of the same size (Tammet et al., 2013) or through ion-mediated nucleation (Hirsikko et al., 2011). Ions of this size are typically connected to atmospheric new particle formation (NPF), snow fall or rain (Manninen et al., 2010; Kerminen et al., 2018; Leino et al. 2016).

45 Atmospheric ions are the carriers of electric charge in the atmosphere. The ability for air to conduct electricity was first discovered by Richmann (1751) and Coulomb (1785), and Faraday (1834) proposed an explanation to this by suggesting that neutral molecules are ionized. Early results focused on understanding the charging of air molecules and aerosols in specific charge events, such as thunderstorms and electrification of clouds (Canton, 1753; Franklin, 1751). It was not until after the discovery of radioactivity by Wilhelm Röntgen, Henri Becquerel and Marie and Pierre Curie that the ionization of neutral air 50 ions in fair weather conditions was successfully explained (Carlson and De Angelis, 2011; De Angelis, 2014; Wilson, 1895, 1899). Air ions were then subsequently used to measure radioactivity as well as air quality (Misaki et al., 1972a,b, 1975; Tuomi, 1989; Israelsson and Knudsen, 1986; Retalis and Pitta, 1989).

The phenomenon of atmospheric ions was historically studied in the field of atmospheric electricity (Israël, 1970), because measuring atmospheric ions and their flow allows for the measurement of the conduction current in the atmosphere 55 (Harrison and Carslaw, 2003; Wilson, 1921). However, in the past 30 years, the relevance of atmospheric ions and aerosols has been recognized and the role of charged aerosols in both cloud processes and NPF has been debated (Dickinson, 1975; Arnold, 1980; Nadytko et al., 2003, Kulmala and Tammet, 2007; Enghoff and Svensmark, 2008). The current view is that ions can contribute to both the formation (Kirkby et al. 2011; 2016) and growth of small particles (Lehtipalo et al. 2016; Leppä et al. 2011; Stolzenburg et al. 2020; Svensmark et al. 2017), but the overall significance of ion-induced nucleation in 60 the present-day atmosphere is rather limited, at least in the continental boundary layer (Manninen et al. 2010; Kulmala et al. 2013; Dunne et al. 2016).

The concentration of atmospheric cluster ions can be expressed as a balance equation:

$$\frac{dn}{dt} = q - CoagS \cdot n - \alpha n^2,$$

(1)




65 in which n is the cluster ion concentration, q is the rate of air molecule ionization, $CoagS$ is the coagulation sink coefficient
(Kulmala et al., 2001) and α is the ion-ion recombination coefficient. The coagulation sink coefficient describes the rate at
which cluster ions are lost via coagulation to larger particles, and this rate can be up to 2 times higher than the corresponding
rate of neutral clusters of the same size (Mahfouz and Donahue, 2021). The dynamics of the intermediate ions are more
70 complicated, as their concentration dependent largely on the growth of smaller aerosols, which in turn is dependent on other
factors such as meteorological conditions and the abundance of condensable vapors (Kerminen et al. 2018).

Despite the increased interest in atmospheric ion concentration in the past decades, there has been relatively little research on
how atmospheric ion concentrations have changed over time. Hirsikko et al. (2011) compared ion concentration
measurements around the world in different environments. Their study included measurements from several different time
periods, but no analysis of the long-term changes in concentrations were made. Other studies have investigated the diurnal
75 and seasonal cycles of atmospheric ions based on a few years of data at best (Chen et al., 2016, Manninen et al., 2009), or
have been focused on characterizing the charged aerosol size distribution (Hörrak, 2001). Long-term trends of atmospheric
ions and their characteristic properties have not been properly assessed.

In this study, we investigate the long-term trends of atmospheric cluster (0.8 – 2 nm) and intermediate ions (2 – 7 nm) in the
boreal forest and attempt to characterize the main drivers behind these trends. We use 16 years of cluster and intermediate
80 ion concentration data from the SMEAR II station in Hyytiälä, Finland, as well as ionization rates calculated from gamma
and radon measurements and cosmic ray models.

2 Materials and Methods

The data used in this study is from the SMEAR II station (Hari and Kulmala, 2005) in Hyytiälä, Finland (61°5 00 00 N,
24°1700 E; 181 m a.s.l.). The station is considered a rural background station, with the largest urban settlement, Tampere,
85 located about 60 kilometers to the south-west. Scots pine forest surrounds the station. The station hosts a comprehensive set
of long-term measurements, including aerosol, radiation and meteorological measurements used in this study. The location
of the ion measurements and the radiation measurements used to calculate the ionization rates are shown in Figure 1. The
measurement locations are between 45 and 125 meters apart from each other at the site. The Balanced Scanning Mobility
Analyzer (BSMA; Tammet, 2006) is at location A. The radiation measurements are currently located at site C but were at
90 site B prior to 2019. Figure 2 and Table 1 show the data availability of all the variables used in this study, which includes the
ion concentrations as well as their most important sources and sinks. Temperature and relative humidity are included as the
most relevant meteorological variables.



95 **Figure 1: The locations of the ion size distribution measurement at site A (BSMA) and the radiation measurements at site C (at site B prior to 2019), and their distances from one another. (© Google Maps 2022)**




Figure 2: Time periods when data is available for variables we used in our analysis. I_{CR} is the cosmic ray ionization rate, I_g is the gamma ionization rate, I_R is the radon ionization rate, RH is relative humidity, T is temperature and CS is the condensation sink. BSMA is the ion spectrometer used in the study.

100



2.1. Measurements

2.1.1. Balanced Scanning Mobility Analyzer

The Balanced Scanning Mobility Analyzer (BSMA; Tammet, 2006) is a differential aspiration condenser with two parallel aspiration condensers connected as a balanced capacitance bridge. Because this eliminates the electrostatically induced current, the BSMA can continuously scan through a mobility range of $0.023 - 3.2 \text{ cm}^2 \text{ V}^{-1} \text{ s}^{-1}$, which corresponds to a mobility diameter range of 0.82 to 8.3 nm in Millikan diameters. The inlet flow rate is set to 2400 liters per minute (lpm) to reduce ion losses in the inlet. The time resolution of the instrument is around 10 minutes, which is the time it takes to complete one full scan of the mobility range.

2.1.2. Ion sources and sinks

We have calculated the ionization rates from gamma radiation and radon decay using the method prescribed by Chen et al. (2016) and used the modeled cosmic ray ionization rate from (<http://cosmicrays oulu.fi/CR1/CR11.html>) and calculated the relevant ionization rate at the atmospheric depth relevant to our station by interpolation (Kulmala et al., 2010). Other sources such as traffic emissions, power lines, rain and lightning strikes can produce cluster and intermediate ions (Eisele 1989a,b; Haverkamp et al., 2004; Tammet et al., 2009), but we estimate that their effects are minor in the long-term at our site because the durations of these events are in general quite short and the measurement site is not in the vicinity of major power lines or traffic.

The main sink for cluster and intermediate ions is their coagulation with pre-existing particles. For sub-10 nm particles, condensation sink (CS) can be used as a good estimate for the changes in coagulation sink (CoagS), if not necessarily its order of magnitude (Dal Maso et al., 2002). It is not necessary to know the absolute value of the coagulation sink, because we are only attempting to gauge how the coagulation sink changes over time. As Lehtinen et al. (2007) shows, the coagulation sink and CS are directly linked and it can therefore be used to evaluate changes in the coagulation sink. The CS was calculated in this study by integrating over the aerosol size distribution as presented in Kulmala et al. (2001) from a combined particle number size distribution between 3 nm and 20 μm derived from the measurement with a Differential Mobility Particle Size (DMPS) and an Aerodynamic Particle Sizer (APS).

2.1.3. Meteorological variables

Temperature and relative humidity (RH) data used in this study were acquired from the smartSMEAR portal (<https://smear.avaa.csc.fi/>; Junninen et al., 2009) with a one-second time resolution averaged to monthly or annual resolution, depending on our needs. Temperature and RH are typical indicators of the overall conditions in the ambient air, as changes in RH are linked to changes in cloud cover (Dada et al., 2017) and precipitation. BSMA as an instrument is also sensitive to changes in humidity (Tammet, 2006), and therefore it bears monitoring in this study. Temperature is linked to



the growing season and biological activity in the surrounding forest (Nieminen et al., 2014), which affects the atmospheric composition and availability of vapors for particle formation and growth, but it can also have an direct effect on the ion-enhanced particle formation processes (Laakso et al. (2002); Curtius et al. (2006); Yu (2010); Kürten et al., 2016).

135

Table 1: The variables used to explain the variance in cluster and intermediate ion concentrations. Coverage is defined as the number of data points available in the time period given.

Variable	Time period	Coverage	Measurement device	If calculated, definition in
Ion size distribution (0.82 – 8.3 nm)	2005 – 2021	90 %	BSMA	-
Condensation Sink	2005 – 2021	96%	DMPS	Kulmala et al., 2001
Temperature	2004 – 2021	97%	Pt100 inside custom shield	-
Relative Humidity	2004 – 2021	98%	Rotronic MP102H RH sensor	-
Radon ionisation rate	2004 – 2021	77%	Description in Chen et al., 2016	Chen et al., 2016
Gamma ionisation rate	2004 – 2015	100%	Description in Chen et al., 2016	Chen et al., 2016
Cosmic ray ionisation rate	2004 – 2019	100%	Modeled	Usoskin and Kovaltsov, 2006

2.2. Ion data verification

Prior to trends analysis, the ion concentration time series need to be verified for consistency and breakpoints. Instrument upgrades, relocations, inlet changes and such can cause breakpoints in which the time series has a discontinuity due to changes in instrument calibration, detection limit, activation probability or sizing. Therefore, it is important to detect these breakpoints and if possible, homogenize the time series before attempting trend analysis. The most common way to determine whether a time series has a breakpoint is visually (Collaud Cohen et al., 2020), but care must be taken when looking for breakpoints in ion concentrations that include multichannel instruments and/or are interpolated to a specific size range from the initial inverted number size distribution. This is because the summing or interpolation can hide possible breakpoints. We investigated our measured ion number size distributions for the breakpoints in two ways, visually and by making sure the signal variance of the instruments remained steady over the measurement period.

The visual inspection of the BSMA time series revealed no notable breakpoints in the time series. This was further supported by the channel-by-channel signal of the time series, in which the variance for each channel remained steady over the entire



150 measurement period (See Appendix, Figure A1). The variance increased slightly over time due to the instrument getting older, but this was not a notable change.

2.3. Size selection

Ions are traditionally split according to their size into small, intermediate and large ions (Hirsikko et al., 2011). Cluster ions, also called small ions, are typically smaller than 2 nm, while intermediate ions are between 2 and 7 nm and large ions are
155 larger than that (Hirsikko et al., 2011; Hörrak et al., 2000,2003) We chose to use three size ranges in our analysis. One for the cluster ions (0.8 – 2 nm) and two for the intermediate ions (2 – 4 nm and 4 – 7 nm). We split the intermediate ions into two size ranges, because earlier studied have shown that the 2 – 4 nm ion concentration is typically the most sensitive to NPF (Leino et al., 2016), while depending on growth rate, it can take several hours for the newly formed particle to reach 4 nm.

160 2.4. Trend analysis and multiple linear regression analysis

When investigating trends from atmospheric time series, it is important to account for autocorrelation which exists in most atmospheric variables. We used the 3PW algorithm described in Collaud Cohen et al. (2020) to remove the autocorrelation before applying the Mann-Kendall (MK) test on the prewhitened time series. The yearly trend was then calculated by using the Sen's slope and presented as an annual change.

165 Finally, we tested the effect of various factors on the concentration of cluster and intermediate ions in the atmosphere by using n-way analysis of variance (ANOVAn) from the MATLAB Statistics and Machine Learning toolbox.

3 Results

3.1 Cluster and intermediate ion concentrations

In this section we present and analyze the time series of cluster and intermediate ion concentrations in the boreal forest.

170 The median concentrations and variability of ions measured with BSMA are presented in Table 2, and as violin plots in Figure 3. Negative and positive ion concentrations are very similar to one another, with the exception that the 4 – 7 nm negative ion concentrations are more spread out with higher concentrations being more common (Figure 3). This might suggest that more negative particles are created by ion-induced nucleation than positive particles or the negative ions grow faster to the 4-7 nm size bin before being neutralized or lost.

175 Our measured concentrations agree well with concentrations reported by previous studies. Komppula et al. (2007) reported mean cluster ion concentrations in Hyytiälä to be typically around 840 cm⁻³ for positive ions and 770 cm⁻³ for negative ions when measured with the Air Ion Spectrometer (AIS) (Mirme et al., 2007) and Hirsikko et al. (2005) reported the monthly mean values to be between 600 and 800 cm⁻³ for both polarities, measured with the BSMA. Leino et al. (2016) reported intermediate ion concentrations at Hyytiälä to be between 1 and 25 cm⁻³ for negative ions measured with a NAIS and



180 Manninen et al. (2009) reported median 1.8 – 3 nm ion concentrations measured with the same instrument to be between 1 and 10 cm⁻³ with the negative ion concentration being slightly higher. Hirsikko et al. (2005) reported the intermediate ions concentrations measured with the BSMA to be between 0 and 25 cm⁻³. The concentrations measured by the BSMA in this study are very similar when averaged over the whole time period of 2004 – 2021.




Figure 3: Measured ion concentrations. Negative polarity is on the left in blue and positive polarity is on the right in orange. The cluster ion (0.8 – 2 nm) concentrations are on the top row, and intermediate ion concentrations are in the middle and bottom, 2 – 4 nm and 4 – 7 nm, respectively. The white dot is the median concentration and the whiskers mark the location of the 95th and 5th percentile data points.

190

Table 2: Descriptive statistics of the ion measurements dataset between 2005 and 2021.

Ion	Concentration (cm ⁻³)	5th percentile	25th percentile	Median	75th percentile	95th percentile
0.8 – 2 nm (-)		350	560	710	860	1100
2 – 4 nm (-)		1.5	6.7	14	27	97
4 – 7 nm (-)		0.0	3.4	9.2	21	110
0.8 – 2 nm (+)		380	580	720	850	1100
2 – 4 nm (+)		2.0	7.4	14	26	72
4 – 7 nm (+)		0.0	3.0	8.0	18	89




3.2 Trend analysis

3.2.1 Long-term trends

The long-term ion concentration trends are shown in Figure 4 and Table 3. We observe a slight positive trend of 1% annual change for the cluster ion concentration in both polarizations, corresponding to a median increase of $7.1 \text{ cm}^{-3}/\text{year}$. The trends are very similar in both polarities and within the confidence limits for each other. For the intermediate ions, the BSMA time series exhibits a positive trend of around 3.5 – 3.9 % per year for 2 – 4 nm ions in both polarities (median increase of $0.55 \text{ cm}^{-3}/\text{year}$) and a positive trend of around 1.7 – 1.9 % per year for 4 – 7 nm ions in both polarities (median increase of $0.17 \text{ cm}^{-3}/\text{year}$).

Table 3: The relative change in ion concentration for BSMA for 2004 – 2020. The relative changes per year have been calculated from Sen’s slopes.

(% a year)	2004 – 2020
0.8 – 2 nm	1.1 (0.6 – 1.7)
(-)	
2 – 4 nm (-)	3.9 (1.9 – 6.1)
(-)	
4 – 7 nm	1.9 (0.1 – 3.5)
(-)	
0.8 – 2 nm	0.9 (0.4 – 1.3)
(+)	
2 – 4 nm	3.4 (0.9 – 5.9)
(+)	
4 – 7 nm	1.7 (-0.3 – 3.5)
(+)	
CS	-1.9 (-3.3 – -0.6)
T	-0.8 (-3.7 – 2.2)
RH	-0.4 (-4.9 – 4.3)
I_{Radon}	-
I_{gamma}	-1.0 (-1.8 – -0.4)
I_{CR}	-



205 **Figure 4: The time series of the cluster and intermediate ion concentrations. The relative changes per year have been calculated from Sen's slopes.**

Out of the three main ionization sources (radon, gamma and cosmic ray ionization), radon and gamma ionization rates exhibit generally decreasing trends before 2015 (Figure 5, Table 3), which would suggest that cluster ion concentrations should be decreasing during this period. However, our observations do not follow this. The main sink for cluster and intermediate ions, the sink from larger aerosol particles, which is here described by the CS (see section 2.1.3.), also exhibits a decreasing trend. Therefore, the competing influence of the decreasing trends will be further analyzed in section 3.2.3. to identify the most important factor to the ion concentrations in the boreal forest. Unfortunately, there is a large fraction of missing ionization rate data in the later part of the analyzed time period. Radon and cosmic ray ionization rates exhibit multi-year sinusoidal behavior and we do not list a trend for them here. This is expected for the cosmic ray ionization rate, but the reason for the changes in radon ionization rate are unclear. Temperature and relative humidity both exhibit no significant trends (Figure 5).

215





Figure 5: The time series of main sources and sinks as well as temperature and relative humidity. The relative changes per year have been calculated from Sen's slopes.


3.2.2 Seasonal cycle

220 We investigated the annual variability in the ion concentrations by calculating the median seasonal cycle of the ions in both polarities (Figure 6) and ion sources and sinks as well as T and RH (Figure 7). Cluster ions (0.8 – 2 nm) in both polarities have an annual maximum in the late autumn in October and a local maximum in spring in May. Chen et al (2016) and Hirsikko et al (2005) observed similar maxima for cluster ions in a boreal forest. The 2 – 4 nm intermediate ion concentration has an annual maximum in spring in May in both polarities which coincides with the known annual maximum for NPF event frequency (Nieminen et al., 2014). The negative 2 – 4 nm ion concentration has a second, smaller maximum in September, coinciding with the second annual maximum in NPF event frequency (Nieminen et al., 2014). This is in good agreement with the result that 2 – 4 nm ions in particular are good indicators for NPF events (Leino et al., 2016). The fall maximum is not clearly observed in the positive 2 – 4 nm ion concentration. This may suggest that there is different chemistry involved in autumn NPF than in spring, but further analysis of this is outside the scope of the manuscript. The 4 – 230 7 nm ion concentration in both polarities has an annual maximum in March and a second local maximum in September, which matches the maxima observed by Leino et al (2016) for 2 – 7 nm ions.



235 **Figure 6:** The annual variance of ion concentrations in the size bins, 0.8 – 2 nm, 2 – 4 nm and 4 – 7 nm. The left-hand side boxplots are in the negative polarity and the right-hand side are in the positive polarity. The numbers on the x-axis are month numbers (Jan-Dec). The red line is the median concentration, and the blue box contains 50% (25th to 75th percentile) of all data points. The whiskers mark the location of the 95th and 5th percentile data points.

CS has an annual maximum in the summer (Figure 7a), which has been suggested to suppress NPF during the summer (Buenrostro Mazon et al., 2016; Nieminen et al., 2014). The annual minimum is in December, with a local minimum in
240 March. This roughly follows the seasonal behavior of CS reported previously (Lyubovtseva et al. (2005); Vana et al., 2016). Temperature has a maximum in the summer (Figure 7b) as expected, while RH has a minimum in the summer (Figure 7c; Lyubovtseva et al. 2005; Hirsikko et al., 2005). The radon ionization rate has an annual maximum in February and a second local maximum in September (Figure 7d). This has been connected to changes in the mixing layer depth in Chen et al. (2016). The gamma ionization on the other hand has a local minimum in February and March (Figure 7e), which coincides
245 with the typical maximum snow depth at the station, which suppresses gamma radiation from the soil (Chen et al., 2016). Finally, the cosmic radiation ionization rate remains practically constant (Figure 7f), as its changes are known to occur in an 11-year-period and a median annual value should appear fairly constant.



250 **Figure 7: The annual variation of CS, T, RH, I_r , I_g , I_{CR} . The red line is the median concentration, and the blue box contains 50% (25th to 75th percentile) of all data points. The whiskers mark the location of the 95th and 5th percentile data points. The numbers on the x-axis are month numbers.**


3.2.3 Analysis of variance

We investigated the reasons behind the variations in ion concentrations by performing an analysis of variance (ANOVA) for a simple multiple linear regression model. Our assumption was that cluster and intermediate ion concentration variation is caused by the interplay of their sources and sinks. Additionally, concentrations can be modulated by the prevailing meteorological conditions, which are connected to atmospheric air composition and ion processes. The degrees of freedom in the ANOVA were adjusted to reflect the number of explanatory variables included. The explanatory variables used are listed in Table 1.

260 The percentage of variability explained by each explanatory variable is presented in Figure 8. 50% of the cluster ion concentration variability is explained by changes in CS, which corroborates the assumption in the previous chapter, that the decrease in CS could be the main reason for the observed negative trend in ion concentrations. The second most important factor was RH. Intermediate ion concentrations, both 2 – 4 and 4 – 7 nm, are best explained by a combination of sources and meteorological variability. However, the combination is different for negative and positive ions. The chemical composition
265 of the negative and positive ions is known to be different (Luts et al., 2011, Ehn et al., 2011), and it is possible that humidity



as well as chemical differences in condensable vapors can cause differences in the dynamics of the negative and positive ions.



270 **Figure 8: The amount of variation explained by our chosen explanatory variables for each size range and polarity for BSMA. The symbol (-) denotes negative polarity, (+) a positive polarity and (+-) the sum of both polarities. The bolded red values are statistically significant ($p < 0.05$).**

275 However, because of strong seasonal effects in ion processes, the variation of ion concentrations could be dominated by different factors in different seasons. For this reason, we also investigated how these factors explained the variation in ion concentrations in different seasons. In Figures A2 – A5 (See Appendix) we show the amount of variability explained by our selected variables when isolating the different seasons.


280 The different seasons each have some characteristics that are not directly visible in the variation of the entire time series. In spring and summer, the variation in cluster ion concentrations is most strongly explained by the radon ionization rate, about 40%. CS explains a further 23 – 41% of the variance in spring and summer, with meteorological variables being more important in summer. In autumn, temperature explains 27 – 37% of the variation in cluster ion concentrations while ion sources and sinks contribute less to the variation. Radon and gamma ionization rates contribute roughly 15% each to the cluster ion concentration variance. Our chosen variables fail to explain the changes in cluster ion concentrations in winter. Gamma ionization rate explains about 10% of the variation cluster ion concentrations in winter. The fraction of variance explained in autumn and winter suggests that our chosen variables are not ideally suited to explaining the variance in those seasons. Other factors such as the abundance of relevant chemical vapors to grow the ions into the size ranges considered
 285 here may be more important during these seasons.



The intermediate ion variation in spring is in general explained best by the gamma ionization rate and cosmic ray ionization rate. The 2 – 4 nm concentration of negative ions, however, is largely explained by the CS, notably different from the positive polarity. Meteorological parameters have the least effect in spring. Summer and autumn intermediate ion concentration variation is explained by a combination of ion sources and meteorology. Radon ionization rate explains 25 – 290 42 % of the variance in intermediate ion concentration in the summer, while in the autumn all three ion sources explain 10 – 20 % each, but these numbers are not statistically significant. In winter, the variation in intermediate ion concentration is explained by a combination of radon ionization rate and temperature, likely linked to varying snow depth and its suppressive effect on the ionization rates (Chen et al., 2016).

295 We also analyzed the seasonal variance by using median concentrations calculated for each month. (Figure 9). The variance in cluster ion concentration in the negative polarity is explained by a combination of CS (19%) and temperature (22 %). Meanwhile, in the positive polarity the variation is best explained by CS (15 %) and I_g (14 %). However, our chosen variables do not explain the seasonal variation in cluster ion concentrations very well. The variance in intermediate ion concentrations is explained in both polarities by a combination of I_g and the meteorological variables T and RH. 2 – 4 nm 300 intermediate ion concentrations are particularly dependent on the variance in RH. This can be explained by the tendency of NPF events to occur during clear sky, and thus low RH days (Dada et al., 2017). The dependency is clearer in the seasonal variation because seasonal effects are modulated by the long-term trends.

All in all, the effect of ionization rates on the trends and variability of ion concentrations were surprisingly low. It could be that although the ionization rates control the production of primary ions, the measured concentrations, even in the smallest 305 size bin studied here, depend largely on the factors affecting ion dynamics and growth to large enough sizes to be detected by our instruments, which in turn depend on multiple other factors such as air composition (Chen et al., 2016).



310 **Figure 9: The amount of seasonal variation explained by our chosen explanatory variables for each size range and polarity. The symbol (-) denotes negative polarity, (+) a positive polarity and (+-) the sum of both polarities. The bolded red values are statistically significant ($p < 0.05$).**

4 Discussion & Conclusions

In this work we have presented a 16-year long time series of ion concentrations at SMEAR II station in southern Finland. Using this unique data set, we investigated long-term trends of cluster and intermediate ion concentrations in the atmosphere at a boreal forest. A connection between the steadily increasing ambient cluster ion concentrations and decreasing CoagS, estimated in this analysis with CS, appears clear. The decreasing CoagS is likely due to overall decreasing level of anthropogenic aerosols in boreal forest (Luoma et al. 2019), but possible changes in the air mass origins cannot be easily dismissed and were not investigated here. Although some trends were evident in the ambient ionization sources as well, they did not explain the overall trend of the ambient cluster ion concentrations.

320 The reasons for the rise of ambient intermediate ion concentration are more complex. This is to be expected, as the dynamics of 2 – 7 nm aerosols are much more complicated, not only dependent on formation and loss, but ion-ion recombination, ion-aerosol attachment and ion growth (Kulmala et al., 2013). And because growth is dependent on many factors, such as the availability of condensable vapors and favorable meteorological conditions, the dependencies will not be as simple (Kerminen et al., 2018). The connection to meteorological conditions is evident in our analysis as well, whereas the season-specific analysis of variation suggests that while our chosen variables appear to be the limiting factors during the summer and spring months, other factors may be more important to the intermediate ion concentration in autumn and especially

325



winter. These factors include boundary layer dynamics and the abundance of condensable vapors (Hao et al., 2018; Hao et al., 2021; Kirkby et al., 2011; Almeida et al., 2013; Vakkari et al., 2015).

Our analysis has several sources of uncertainty worth noting. As mentioned previously, because the BSMA data with RH above 80 % were excluded, our analysis does not include any possible dynamics related to precipitation, which also is
330 known to produce intermediate ions. Therefore, NPF events are likely overrepresented in the data. Additionally, we are missing three years of radon ionization rate data and five years of gamma ionization data. This makes comparing trends over the entire time period difficult and introduces uncertainty in our interpretation of the results.

Increase in ion concentrations can mostly be attributed to a decreasing CS. The changes in the ionization rates explain the observed trend to a lesser degree. It is possible that this is simply because the ion sink is the dominant limiting factor to the
335 formation of cluster and intermediate ions. However, several open questions remain. Although studies indicate that the neutral pathway dominates NPF in the boreal forest, the role of ions is not totally clear (Wagner et al., 2017). Ion-induced nucleation can be significant in certain conditions, e.g., when the sulfuric acid concentration is low (Rose et al., 2018; Yan et al., 2018; Gagné et al., 2010). Additionally, ion-ion recombination can create neutral aerosols. It is possible that ions are especially important in the initial stages of NPF, which is difficult to measure with current instrumentation. It will have to be
340 determined in the future whether the number of aerosols produced by ion-ion recombination (Kontkanen et al. 2013; Franchin et al., 2015) is affected by the changes in ion concentrations. It can be speculated that the increasing ion concentrations, decreasing concentrations of sulfuric acid in boreal forest (Nieminen et al., 2014) and increasing temperatures will make the ion-induced or ion-mediated NPF pathways more important in future. A more comprehensive analysis of this is needed.

345

Data availability. The meteorological data from the SMEAR II station can be accessed from the smartSMEAR website: <https://smear.avaa.csc.fi/> (last access: February 28th, 2022) (Junninen et al., 2009). The ion concentration data, the ionization rates and condensation sink data are available at <https://doi.org/10.5281/zenodo.6597201> (Sulo et al., 2022). The data are licensed under a Creative Commons 4.0 Attribution (CC BY) license.

350

Author contributions. JS performed the data analysis and wrote the paper. JL was responsible for the ion measurements, JL and JS handled the ion data inversion, XC calculated the radon and gamma ionization rates, TN calculated the cosmic ray ionization rate. JK, MK and KL designed the study and provided supervision. All authors contributed to the discussion of the results and provided comments on the paper.

355

Competing interests. Author VMK is a member of the editorial board for Atmospheric Chemistry and Physics.

Acknowledgements. Hanna E. Manninen, Stephany Buenrostro Mazon, Jussi Paatero, the SMEAR II technical and scientific staff and everyone else contributing to the ion and radiation measurements are gratefully acknowledged. This work was



360 supported by ACCC Flagship funded by the Academy of Finland grant number 337549, Academy professorship funded by the Academy of Finland (grant no. 302958), Academy of Finland projects no. 1325656, 316114 and 325647; “Quantifying carbon sink, CarbonSink+ and their interaction with air quality” INAR project funded by Jane and Aatos Erkko Foundation and European Research Council (ERC) project ATM-GTP Contract No. 742206.

References

- 365 Almeida, J., Schobesberger, S., Kürten, A., Ortega, I., Kupiainen-Määttä, O., Praplan, A., Adamov, A., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Donahue, N., Downard, A., Dunne, E., Duplissy, J., Ehrhart, S., Flagan, R., Franchin, A., Guida, R., Hakala, J., Hansel, A., Heinritzi, M., Henschel, H., Jokinen, T., Junninen, H., Kajos, M., Kangasluoma, J., Keskinen, H., Kupc, A., Kurtén, T., Kvashin, A., Laaksonen, A., Lehtipalo, K., Leiminger, M., Leppä, J., Loukonen, V., Makhmutov, V., Mathot, S., McGrath, M., Nieminen, T., Olenius, T., Onnela, A., Petäjä, T., Riccobono, F.,
- 370 Riipinen, I., Rissanen, M., Rondo, L., Ruuskanen, T., Santos, F., Sarnela, N., Schallhart, S., Schnitzhofer, R., Seinfeld, J., Simon, M., Sipilä, M., Stozhkov, Y., Stratmann, F., Tomé, A., Tröstl, J., Tsagkogeorgas, G., Vaattovaara, P., Viisanen, Y., Virtanen, A., Vrtala, A., Wagner, P., Weingartner, E., Wex, H., Williamson, C., Wimmer, D., Ye, P., Yli-Juuti, T., Carslaw, K., Kulmala, M., Curtius, J., Baltensperger, U., Worsnop, D., Vehkamäki, H. and Kirkby, J.: Molecular understanding of sulphuric acid–amine particle nucleation in the atmosphere, *Nature*, 502(7471), 359–363, doi:10.1038/nature12663,
- 375 2013. Arnold, F.: Multi-ion complexes in the stratosphere-implications for trace gases and aerosol, *Nature*, 284, 610–611, 1980.
- Arnold, F.: Atmospheric Ions and Aerosol Formation, *Space Sci. Rev.*, 137, 225–239, 2008.
- Bazilevskaya, G. A., Usoskin, I. G., Fluckiger, E. O., Harrison, R. G., Desorgher, L., Butikofer, R., Krainev, M. B., Makhmutov, V. S., Stozhkov, Y. I., Svirzhevskaya, A. K., Svirzhevsky, N. S., and Kovaltsov, G. A.: Cosmic Ray Induced
- 380 Ion Production in the Atmosphere, *Space Sci. Rev.*, 137, 149–173, 2008.
- Buenrostro Mazon S., Kontkanen J., Manninen H.E., Nieminen T., Kerminen V.-K. & Kulmala M: A long-term comparison of nighttime cluster events and daytime ion formation in a boreal forest. *Boreal Env. Res.* 21: 242–261, 2016.
- Canton, J.: Electrical Experiments, with an Attempt to Account for Their Several Phaenomena; Together with Some Observations on Thunder-Clouds, *Phil. Trans.*, 48, 350–358, 1753.
- 385 Carlson, P. and De Angelis, A.: Nationalism and internationalism in science: the case of the discovery of cosmic rays, *Eur. Phys. J. H*, 35, 309–329, doi:10.1140/epjh/e2011-10033-6, 2011.
- Chen, X., Kerminen, V.-M., Paatero, J., Paasonen, P., Manninen, H. E., Nieminen, T., Petäjä, T., and Kulmala, M.: How do air ions reflect variations in ionising radiation in the lower atmosphere in a boreal forest?, *Atmos. Chem. Phys.*, 16, 14297–14315, https://doi.org/10.5194/acp-16-14297-2016, 2016.



- 390 Collaud Coen, M., Andrews, E., Bigi, A., Martucci, G., Romanens, G., Vogt, F. P. A., and Vuilleumier, L.: Effects of the prewhitening method, the time granularity, and the time segmentation on the Mann–Kendall trend detection and the associated Sen's slope, *Atmos. Meas. Tech.*, 13, 6945–6964, <https://doi.org/10.5194/amt-13-6945-2020>, 2020.
- Coulomb, C. A.: Troisième Mémoire sur l'Electricité et le Magnétisme, *Histoire de l'Académie Royale des Sciences*, l'Académie Royale des Sciences Paris, 612–638, 1785.
- 395 Curtius, J., Lovejoy, E. R., and Froyd, K. D.: Atmospheric ion-induced aerosol nucleation, *Space Sci. Rev.*, 125, 159–167, 2006
- Dada, L., Paasonen, P., Nieminen, T., Buenrostro Mazon, S., Kontkanen, J., Peräkylä, O., Lehtipalo, K., Hussein, T., Petäjä, T., Kerminen, V.-M., Bäck, J., and Kulmala, M.: Long-term analysis of clear-sky new particle formation events and nonevents in Hyytiälä, *Atmos. Chem. Phys.*, 17, 6227–6241, <https://doi.org/10.5194/acp-17-6227-2017>, 2017.
- 400 Dal Maso, M., Kulmala, M., Lehtinen, K. E. J., Mäkelä, J. M., Aalto, P., and O'Dowd, C. D.: Condensation and coagulation sinks and formation of nucleation mode particles in coastal and boreal forest boundary layers, *J. Geophys. Res.*, 107(D19), doi:10.1029/2001JD001053, 2002.
- De Angelis, A.: Atmospheric ionization and cosmic rays: studies and measurements before 1912, *Astropart. Phys.*, 53, 19–26, doi:10.1016/j.astropartphys.2013.05.010, 2014.
- 405 Dickinson, R.: Solar Variability and the Lower Atmosphere, *Bulletin of the American Meteorological Society*, 56(12), 1240–1248, doi:10.1175/1520-0477(1975)056<1240:svatla>2.0.co;2, 1975.
- Ehn, M., Junninen, H., Schobesberger, S., Manninen, H., Franchin, A., Sipilä, M., Petäjä, T., Kerminen, V., Tammet, H., Mirme, A., Mirme, S., Hörrak, U., Kulmala, M. and Worsnop, D.: An Instrumental Comparison of Mobility and Mass Measurements of Atmospheric Small Ions, *Aerosol Science and Technology*, 45(4), 522–532, doi:10.1080/02786826.2010.547890, 2011.
- 410 Eisele, F. L.: Natural and atmospheric negative ions in the troposphere, *J. Geophys. Res.*, 94, 2183–2196, 1989a.
- Eisele, F. L.: Natural and transmission line produced positive ions, *J. Geophys. Res.*, 94, 6309–6318, 1989b.
- Eisenbud, M., & Gesell, T.: *Environmental Radioactivity: From Natural, Industrial, and Military Sources*. 4th ed. San Diego: San Diego Academic Press, 1997.
- 415 Enghoff, M. B. and Svensmark, H.: The role of atmospheric ions in aerosol nucleation – a review, *Atmos. Chem. Phys.*, 8, 4911–4923, <https://doi.org/10.5194/acp-8-4911-2008>, 2008.
- Faraday, M.: Experimental researches on electricity, 7th series, *Phil. Trans. R. Soc. (Lond.)*, 124, 77–122, 1834.
- Franklin, B.: A Letter of Benjamin Franklin, Esq; to Mr. Peter Collinson, F. R. S. concerning an Electrical Kite, *Phil. Trans.*, 47, 565–567, 1751.
- 420 Franchin, A., Ehrhart, S., Leppä, J., Nieminen, T., Gagné, S., Schobesberger, S., Wimmer, D., Duplissy, J., Riccobono, F., Dunne, E. M., Rondo, L., Downard, A., Bianchi, F., Kupc, A., Tsagkogeorgas, G., Lehtipalo, K., Manninen, H. E., Almeida, J., Amorim, A., Wagner, P. E., Hansel, A., Kirkby, J., Kürten, A., Donahue, N. M., Makhmutov, V., Mathot, S., Metzger, A., Petäjä, T., Schnitzhofer, R., Sipilä, M., Stozhkov, Y., Tomé, A., Kerminen, V.-M., Carslaw, K., Curtius, J., Baltensperger,



- U., and Kulmala, M.: Experimental investigation of ion–ion recombination under atmospheric conditions, *Atmos. Chem. Phys.*, 15, 7203–7216, <https://doi.org/10.5194/acp-15-7203-2015>, 2015.
- 425 Gagné, S., Nieminen, T., Kurtén, T., Manninen, H. E., Petäjä, T., Laakso, L., Kerminen, V.-M., Boy, M., and Kulmala, M.: Factors influencing the contribution of ion-induced nucleation in a boreal forest, Finland, *Atmos. Chem. Phys.*, 10, 3743–3757, <https://doi.org/10.5194/acp-10-3743-2010>, 2010.
- Hao, L., Garmash, O., Ehn, M., Miettinen, P., Massoli, P., Mikkonen, S., Jokinen, T., Roldin, P., Aalto, P., Yli-Juuti, T.,
430 Joutsensaari, J., Petäjä, T., Kulmala, M., Lehtinen, K. E. J., Worsnop, D. R., and Virtanen, A.: Combined effects of boundary layer dynamics and atmospheric chemistry on aerosol composition during new particle formation periods, *Atmos. Chem. Phys.*, 18, 17705–17716, <https://doi.org/10.5194/acp-18-17705-2018>, 2018.
- Hari, P. and Kulmala, M.: Station for measuring ecosystematmosphere relations (SMEAR II), *Boreal Environ. Res.*, 10, 315–322, 2005.
- 435 Harrison, R. G. and Carslaw, K. S.: Ion-aerosol-cloud processes in the lower atmosphere, *Rev. Geophys.*, 41(3), 1012, doi:10.1029/2002RG000114, 2003.
- Haverkamp, H., Wilhelm, S., Sorokin, A., and Arnold, F.: Positive and negative ion measurements in jet aircraft engine exhaust: concentrations, sizes and implications for aerosol formation, *Atmos. Environ.*, 38, 2879–2884, 2004.
- Hirsikko, A., Laakso, L., Horrak, U., Aalto, P. P., Kerminen, V.-M., and Kulmala, M.: Annual and size dependent variation
440 of growth rates and ion concentrations in boreal forest, *Boreal Environ. Res.*, 10, 357–369, 2005.
- Hirsikko, A., Nieminen, T., Gagné, S., Lehtipalo, K., Manninen, H. E., Ehn, M., Hörrak, U., Kerminen, V.-M., Laakso, L., McMurry, P. H., Mirme, A., Mirme, S., Petäjä, T., Tammet, H., Vakkari, V., Vana, M., and Kulmala, M.: Atmospheric ions and nucleation: a review of observations, *Atmos. Chem. Phys.*, 11, 767–798, <https://doi.org/10.5194/acp-11-767-2011>, 2011.
- Hörrak, U.: Air ion mobility spectrum at a rural area, PhD, Institute of Environmental Physics, University of Tartu, Tartu,
445 Estonia, 81 pp., 2001.
- Israel, H.: Atmospheric Electricity, Israel Program for Scientific Translations, Jerusalem, 1, 317 pp., 1970.
- Israelsson, S. and Knudsen, E.: Effects of radioactive fallout from a nuclear power plant accident on electrical parameters, *J. Geophys. Res.*, 91(D11), 11909–11910, 1986.
- Junninen, H., Lauri, A., Keronen, P., Aalto, P., Hiltunen, V., Hari, P., and Kulmala, M.: Smart-SMEAR: on-line data
450 exploration and visualization tool for SMEAR stations, *Boreal Environ. Res.*, 14, 447–457, 2009.
- Kirkby, J., Curtius, J., Almeida, J., Dunne, E., Duplissy, J., Ehrhart, S., Franchin, A., Gagné, S., Ickes, L., Kürten, A., Kupc, A., Metzger, A., Riccobono, F., Rondo, L., Schobesberger, S., Tsagkogeorgas, G., Wimmer, D., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Downard, A., Ehn, M., Flagan, R., Haider, S., Hansel, A., Hauser, D., Jud, W., Junninen, H., Kreissl, F., Kvashin, A., Laaksonen, A., Lehtipalo, K., Lima, J., Lovejoy, E., Makhmutov, V., Mathot, S.,
455 Mikkilä, J., Minginette, P., Mogo, S., Nieminen, T., Onnela, A., Pereira, P., Petäjä, T., Schnitzhofer, R., Seinfeld, J., Sipilä, M., Stozhkov, Y., Stratmann, F., Tomé, A., Vanhanen, J., Viisanen, Y., Vrtala, A., Wagner, P., Walther, H., Weingartner, E.,



- Wex, H., Winkler, P., Carslaw, K., Worsnop, D., Baltensperger, U. and Kulmala, M.: Role of sulphuric acid, ammonia and galactic cosmic rays in atmospheric aerosol nucleation, *Nature*, 476(7361), 429-433, doi:10.1038/nature10343, 2011.
- 460 Kerminen, V., Chen, X., Vakkari, V., Petäjä, T., Kulmala, M. and Bianchi, F.: Atmospheric new particle formation and growth: review of field observations, *Environmental Research Letters*, 13(10), 103003, doi:10.1088/1748-9326/aadf3c, 2018.
- Komppula, M., Vana, M., Kerminen, V.-M., Lihavainen, H., Viisanen, Y., Hörrak, U., Komsaare, K., Tamm, E., Hirsikko, A., Laakso, L. & Kulmala, M.: Size distributions of atmospheric ions in the Baltic Sea region. *Boreal Env. Res.* 12: 323–336, 2007.
- 465 Kontkanen, J., Lehtinen, K. E. J., Nieminen, T., Manninen, H. E., Lehtipalo, K., Kerminen, V.-M., and Kulmala, M.: Estimating the contribution of ion–ion recombination to sub-2 nm cluster concentrations from atmospheric measurements, *Atmos. Chem. Phys.*, 13, 11391–11401, https://doi.org/10.5194/acp-13-11391-2013, 2013.
- Kulmala, M., Dal Maso, M., Mäkelä, J.M., Pirjola, L., Väkevä, M., Aalto, P., Miikkulainen, P., Hameri, K. and O’Dowd, C.: On the formation, growth and composition of nucleation mode particles, *Tellus B*, 53, 479–490, 2001.
- 470 Kulmala, M., and H. Tammet.: Finnish-Estonian air ion and aerosol workshops, *Boreal Environ. Res.*, 12, 237 – 245, 2007.
- Kulmala, M., Kontkanen, J., Junninen, H., Lehtipalo, K., Manninen, H., Nieminen, T., Petäjä, T., Sipilä, M., Schobesberger, S., Rantala, P., Franchin, A., Jokinen, T., Järvinen, E., Äijälä, M., Kangasluoma, J., Hakala, J., Aalto, P., Paasonen, P., Mikkilä, J., Vanhanen, J., Aalto, J., Hakola, H., Makkonen, U., Ruuskanen, T., Mauldin, R., Duplissy, J., Vehkamäki, H., Bäck, J., Kortelainen, A., Riipinen, I., Kurtén, T., Johnston, M., Smith, J., Ehn, M., Mentel, T., Lehtinen, K., Laaksonen, A.,
- 475 Kerminen, V. and Worsnop, D.: Direct Observations of Atmospheric Aerosol Nucleation, *Science*, 339(6122), 943-946, doi:10.1126/science.1227385, 2013.
- Laakso, L., Mäkelä, J. M., Pirjola, L., and Kulmala, M.: Model studies on ion-induced nucleation in the atmosphere, *J. Geophys. Res.*, 107(D20), 4427, doi:10.1029/2002JD002140, 2002.
- Leino K., Nieminen T., Manninen H.E., Petäjä T., Kerminen V.-M. & Kulmala M.: Intermediate ions as a strong indicator
- 480 for new particle formation bursts in a boreal forest. *Boreal Env. Res.* 21: 274–286, 2016.
- Leppä, J., Anttila, T., Kerminen, V.-M., Kulmala, M., and Lehtinen, K. E. J.: Atmospheric new particle formation: real and apparent growth of neutral and charged particles, *Atmos. Chem. Phys.*, 11, 4939–4955, https://doi.org/10.5194/acp-11-4939-2011, 2011.
- Luoma, K., Virkkula, A., Aalto, P., Petäjä, T., and Kulmala, M.: Over a 10-year record of aerosol optical properties at
- 485 SMEAR II, *Atmos. Chem. Phys.*, 19, 11363–11382, https://doi.org/10.5194/acp-19-11363-2019, 2019.
- Luts, A., Parts, T., Hörrak, U., Junninen, H. and Kulmala, M.: Composition of negative air ions as a function of ion age and selected trace gases: Mass- and mobility distribution, *Journal of Aerosol Science*, 42(11), 820-838, doi:10.1016/j.jaerosci.2011.07.007, 2011.
- Lyubovtseva, Y. S., Sogacheva, L., Dal Maso, M., Bonn, B., Keronen, P. & Kulmala, M.: Seasonal variation of trace gases,
- 490 meteorological parameters, and formation of aerosols in boreal forests. *Boreal Env. Res.* 10: 493-510, 2005.



- Mahfouz, N. G. A. and Donahue, N. M.: Technical note: The enhancement limit of coagulation scavenging of small charged particles, *Atmos. Chem. Phys.*, 21, 3827–3832, <https://doi.org/10.5194/acp-21-3827-2021>, 2021.
- Manninen, H. E., Petäjä, T., Asmi, E., Riipinen, I., Nieminen, T., Mikkilä, J., Hörrak, U., Mirme, A., Mirme, S., Laakso, L., Kerminen, V.-M., and Kulmala, M.: Long-time field measurements of charged and neutral clusters using Neutral cluster and
495 Air Ion Spectrometer (NAIS), *Boreal Environ. Res.*, 14, 591–605, 2009.
- Manninen, H. E., Nieminen, T., Asmi, E., Gagne, S., Häkkinen, S., Lehtipalo, K., Aalto, P., Vana, M., Mirme, A., Mirme, S., Horrak, U., Plass-Dulmer, C., Stange, G., Kiss, G., Hoffer, A., Toro, N., Moerman, M., Henzing, B., de Leeuw, G., Brinkenberg, M., Kouvarakis, G. N., Bougiatioti, A., Mihalopoulos, N., O’Dowd, C., Ceburnis, D., Arneth, A., Svenningsson, B., Swietlicki, E., Tarozzi, L., Decesari, S., Facchini, M.C., Birmili, W., Sonntag, A., Wiedensohler, A.,
500 Boulon, J., Sellegri, K., Laj, P., Gysel, M., Bukowiecki, N., Weingartner, E., Wehrle, G., Laaksonen, A., Hamed, A., Joutsensaari, J., Petäjä, T., Kerminen, V.-M., and Kulmala, M.: EUCAARI ion spectrometer measurements at 12 European sites – analysis of new-particle formation events, *Atmos. Chem. Phys.*, 10, 7907–7927, doi:10.5194/acp-10-7907-2010, 2010.
- Mirme, A., Tamm, E., Mordas, G., Vana, M., Uin, J., Mirme, S., Bernotas, T., Laakso, L., Hirsikko, A., and Kulmala, M.: A
505 widerange multi-channel Air Ion Spectrometer, *Boreal Environ. Res.*, 12, 247–264, 2007.
- Mirme, S., Mirme, A., Minikin, A., Petzold, A., Horrak, U., Kerminen, V.-M., and Kulmala, M.: Atmospheric sub-3nm particles at high altitudes, *Atmos. Chem. Phys.*, 10, 437–451, doi:10.5194/acp-10-437-2010, 2010.
- Mirme, S. and Mirme, A.: The mathematical principles and design of the NAIS – a spectrometer for the measurement of cluster ion and nanometer aerosol size distributions, *Atmos. Meas. Tech.*, 6, 1061–1071, [https://doi.org/10.5194/amt-6-1061-](https://doi.org/10.5194/amt-6-1061-2013)
510 2013, 2013.
- Mironova, I. A., Aplin, K. L., Arnold, F., Bazilevskaya, G. A., Harrison, R. G., Krivolutsky, A. A., & Usoskin, I. G.: Energetic particle influence on the Earth’s atmosphere. *Space Science Reviews*, 194, 1–96. <https://doi.org/10.1007/s11214-015-0185-4>, 2016.
- Misaki, M., Ohtagaki, M., and Kanazawa, I.: Mobility spectrometry of the atmospheric pollution, *Pure Appl. Geophys.*, 100,
515 133–145, 1972a.
- Misaki, M., Ikegami, M., and Kanazawa, I.: Atmospheric electrical conductivity measurement in the Pacific Ocean, exploring the background level of global pollution, *J. Meteorol. Soc. Japan*, 50, 497–500, 1972b.
- Misaki, M., Ikegami, M., and Kanazawa, I.: Deformation of the size distribution of aerosol particles dispersing from land to ocean, *J. Meteorol. Soc. Japan*, 53, 111–120, 1975.
- 520 Nadykto, A. B. and Yu, F.: Uptake of neutral polar vapor molecules by charged clusters/particles: Enhancement due to dipole-charge interaction, *J. Geophys. Res.*, 108, 4717, doi:10.1029/2003JD003664, 2003.
- Nieminen, T., Asmi, A., Dal Maso, M., Aalto, P., Keronen, P., Petäjä, T., Kulmala, M., and Kerminen, V.-M.: Trends in atmospheric new-particle formation: 16 years of observations in a boreal-forest environment, *Boreal Environ. Res.*, 19, 191–214, 2014.



- 525 Retalis, D. and Pitta, A.: Effects of electrical parameters at Athens Greece by radioactive fallout from a nuclear power plant accident, *J. Geophys. Res.*, 94(D11), 13093–13097, 1989.
- Richmann, G. W.: De electricitate in corporibus producenda nova tentamina, *Commentarii Acad. Sci. Imper. Petropolitanae.*, 14, 299–326, 1744-1746, 1751.
- Rose, C., Zha, Q., Dada, L., Yan, C., Lehtipalo, K., Junninen, H., Mazon, S. B., Jokinen, T., Sarnela, N., Sipilä, M., Petäjä, T., Kerminen, V.-M., Bianchi, F. and Kulmala, M.: Observations of biogenic ion-induced cluster formation in the atmosphere, *Science Advances*, 4(4), doi:10.1126/sciadv.aar5218, 2018.
- 530 Rutherford, E.: The velocity and rate of recombination of the ions of gases exposed to Rontgen radiation, *Philos. Mag.*, 44, 422–440, 1897.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric chemistry and physics: From air pollution to climate change*, John Wiley & Sons, Inc., New York, USA, 1998
- 535 Stolzenburg, D., Simon, M., Ranjithkumar, A., Kürten, A., Lehtipalo, K., Gordon, H., Ehrhart, S., Finkenzeller, H., Pichelstorfer, L., Nieminen, T., He, X.-C., Brilke, S., Xiao, M., Amorim, A., Baalbaki, R., Baccarini, A., Beck, L., Bräkling, S., Caudillo Murillo, L., Chen, D., Chu, B., Dada, L., Dias, A., Dommen, J., Duplissy, J., El Haddad, I., Fischer, L., Gonzalez Carracedo, L., Heinritzi, M., Kim, C., Koenig, T. K., Kong, W., Lamkaddam, H., Lee, C. P., Leiminger, M., Li, Z.,
- 540 Makhmutov, V., Manninen, H. E., Marie, G., Marten, R., Müller, T., Nie, W., Partoll, E., Petäjä, T., Pfeifer, J., Philippov, M., Rissanen, M. P., Rörup, B., Schobesberger, S., Schuchmann, S., Shen, J., Sipilä, M., Steiner, G., Stozhkov, Y., Tauber, C., Tham, Y. J., Tomé, A., Vazquez-Pufleau, M., Wagner, A. C., Wang, M., Wang, Y., Weber, S. K., Wimmer, D., Wlasits, P. J., Wu, Y., Ye, Q., Zauner-Wieczorek, M., Baltensperger, U., Carslaw, K. S., Curtius, J., Donahue, N. M., Flagan, R. C., Hansel, A., Kulmala, M., Lelieveld, J., Volkamer, R., Kirkby, J., and Winkler, P. M.: Enhanced growth rate of atmospheric particles from sulfuric acid, *Atmos. Chem. Phys.*, 20, 7359–7372, <https://doi.org/10.5194/acp-20-7359-2020>, 2020.
- 545 Sulo, J., Sarnela, N., Kontkanen, J., Ahonen, L., Paasonen, P., Laurila, T., Jokinen, T., Kangasluoma, J., Junninen, H., Sipilä, M., Petäjä, T., Kulmala, M., and Lehtipalo, K.: Long-term measurement of sub-3 nm particles and their precursor gases in the boreal forest, *Atmos. Chem. Phys.*, 21, 695–715, <https://doi.org/10.5194/acp-21-695-2021>, 2021.
- Sulo, Juha, Lampilahti, Janne, Chen, Xuemeng, Kontkanen, Jenni, Nieminen, Tuomo, Kerminen, Veli-Matti, Petäjä, Tuukka,
- 550 Kulmala, Markku, & Lehtipalo, Katrianne.: Dataset from measurement report: Increasing trend of atmospheric ion concentrations in the boreal forest [Data set]. Zenodo. <https://doi.org/10.5281/zenodo.6597201>, 2022.
- Svensmark, H., Enghoff, M., Shaviv, N. and Svensmark, J.: Increased ionization supports growth of aerosols into cloud condensation nuclei, *Nature Communications*, 8(1), doi:10.1038/s41467-017-02082-2, 2017.
- Tammet, H.: Continuous scanning of the mobility and size distribution of charged clusters and nanometer particles in atmospheric air and the Balanced Scanning Mobility Analyzer BSMA, *Atmos. Res.*, 82, 523–535, doi:10.1016/j.atmosres.2006.02.009, 2006.
- 555 Tammet, H.: A joint dataset of fair-weather atmospheric electricity, *Atmos. Res.*, 91, 194–200, <http://dx.doi.org/doi:10.1016/j.atmosres.2008.01.012>, 2009.




- Tammet, H.: Symmetric Inclined Grid Mobility Analyzer for the Measurement of Charged Clusters and Fine Nanoparticles
560 in *Atmospheric Air, Aerosol Science and Technology*, 45(4), 468-479, doi:10.1080/02786826.2010.546818, 2011.
- Tammet, H. & Komsaare, K. & Hörrak, U.: Intermediate ions in the atmosphere. *Atmospheric Research*. 135-136.
10.1016/j.atmosres.2012.09.009, 2013.
- Tuomi, T. J.: Ten year summary 1977–1986 of atmospheric electricity measured at Helsinki-Vantaa airport, Finland,
Geophysica, 25, 1–20, 1989.
- 565 Vakkari, V., O'Connor, E. J., Nisantzi, A., Mamouri, R. E., and Hadjimitsis, D. G.: Low-level mixing height detection in
coastal locations with a scanning Doppler lidar, *Atmos. Meas. Tech.*, 8, 1875–1885, [https://doi.org/10.5194/amt-8-1875-](https://doi.org/10.5194/amt-8-1875-2015)
2015, 2015.
- Vana M., Komsaare K., Hörrak U., Mirme S., Nieminen T., Kontkanen J., Manninen H.E., Petäjä T., Noe S.M. & Kulmala
M.: Characteristics of new-particle formation at three SMEAR stations. *Boreal Env. Res.* 21: 345–362, 2016.
- 570 Venzac, H., Sellegri, K., and Laj, P.: Nucleation events detected at the high altitude site of the Puy de Dome Research
Station, ^ France, *Boreal Environ. Res.*, 12, 345–359, 2007.
- Venzac, H., Sellegri, K., Laj, P., Villani, P., Bonasoni, P., Marinoni, A., Cristofanelli, P., Calzolari, F., Fuzzi, S., Decesari,
S., Facchini, M.-C., Vuillermoz, E., and Verza, G. P.: High frequency new particle formation in the Himalayas, *P. Natl.*
Acad. Sci. USA, 105, 15666–15671, 2008.
- 575 Wagner, R., Yan, C., Lehtipalo, K., Duplissy, J., Nieminen, T., Kangasluoma, J., Ahonen, L. R., Dada, L., Kontkanen, J.,
Manninen, H. E., Dias, A., Amorim, A., Bauer, P. S., Bergen, A., Bernhammer, A.-K., Bianchi, F., Brilke, S., Mazon, S. B.,
Chen, X., Draper, D. C., Fischer, L., Frege, C., Fuchs, C., Garmash, O., Gordon, H., Hakala, J., Heikkinen, L., Heinritzi, M.,
Hofbauer, V., Hoyle, C. R., Kirkby, J., Kürten, A., Kvashnin, A. N., Laurila, T., Lawler, M. J., Mai, H., Makhmutov, V.,
Mauldin III, R. L., Molteni, U., Nichman, L., Nie, W., Ojdanic, A., Onnela, A., Piel, F., Quéléver, L. L. J., Rissanen, M. P.,
580 Sarnela, N., Schallhart, S., Sengupta, K., Simon, M., Stolzenburg, D., Stozhkov, Y., Tröstl, J., Viisanen, Y., Vogel, A. L.,
Wagner, A. C., Xiao, M., Ye, P., Baltensperger, U., Curtius, J., Donahue, N. M., Flagan, R. C., Gallagher, M., Hansel, A.,
Smith, J. N., Tomé, A., Winkler, P. M., Worsnop, D., Ehn, M., Sipilä, M., Kerminen, V.-M., Petäjä, T., and Kulmala, M.:
The role of ions in new particle formation in the CLOUD chamber, *Atmos. Chem. Phys.*, 17, 15181–15197,
<https://doi.org/10.5194/acp-17-15181-2017>, 2017.
- 585 Wilson, C. T. R.: The effect of rontgen's rays on cloudy condensation, *P. R. Soc. Lond.*, 59, 338–339, 1895.
- Wilson, C. T. R.: On the condensation nuclei produced in gases by the action of rontgen rays, uranium rays, ultra-violet
light, and other agents, *Philos. T. R. Soc. Lond.*, 192, 403–453, 1899.
- Wilson, C. T. R.: Investigations on lightning discharges and on the electric field of thunderstorms, *Philos. T. R. Soc. Lond.*,
221, 73– 115, 1921.
- 590 WHO: Routine surveillance for radionuclides in air and water, pp. 64, 1968.
- Wu, H., Li, Z., Li, H., Luo, K., Wang, Y., Yan, P., Hu, F., Zhang, F., Sun, Y., Shang, D., Liang, C., Zhang, D., Wei, J., Wu,
T., Jin, X., Fan, X., Cribb, M., Fischer, M., Kulmala, M. and Petäjä, T.: The impact of the atmospheric turbulence-

development tendency on new particle formation: a common finding on three continents, *National Science Review*, 8(3), doi:10.1093/nsr/nwaa157, 2020.


595 Yan, C., Dada, L., Rose, C., Jokinen, T., Nie, W., Schobesberger, S., Junninen, H., Lehtipalo, K., Sarnela, N., Makkonen, U., Garmash, O., Wang, Y., Zha, Q., Paasonen, P., Bianchi, F., Sipilä, M., Ehn, M., Petäjä, T., Kerminen, V.-M., Worsnop, D. R., and Kulmala, M.: The role of H₂SO₄-NH₃ anion clusters in ion-induced aerosol nucleation mechanisms in the boreal forest, *Atmos. Chem. Phys.*, 18, 13231–13243, <https://doi.org/10.5194/acp-18-13231-2018>, 2018.

600 Yu, F., Luo, G., Bates, T. S., Andersson, B., Clarke, A., Kapustin, V., Yantosca, R. M., Wang, Y., and Wu, S.: Spatial distributions of particle number concentrations in the global troposphere: Simulations, observations, and implications for nucleation mechanism, *J. Geophys. Res.*, 115, D17205, doi:10.1029/2009JD013473, 2010.

Appendix




605 **A1: Variance of the concentration time series of the BSMA size channels.**



A2: The amount of variation explained by our chosen explanatory variables for each size range and polarity. The data points selected are only between March and May and represent changes in springtime concentrations. The symbol (-) denotes negative polarity, (+) a positive polarity and (+-) the sum of both polarities. The bolded red values are statistically significant ($p < 0.05$).


610



A3: The amount of variation explained by our chosen explanatory variables for each size range and polarity. The data points selected are only between June and August and represent changes in summertime concentrations. The symbol (-) denotes negative polarity, (+) a positive polarity and (+-) the sum of both polarities. The bolded red values are statistically significant ($p < 0.05$).




615



A4: The amount of variation explained by our chosen explanatory variables for each size range and polarity. The data points selected are only between September and November and represent changes in autumntime concentrations. The symbol (-) denotes negative polarity, (+) a positive polarity and (+-) the sum of both polarities. The bolded red values are statistically significant ($p < 0.05$).

620



A5: The amount of variation explained by our chosen explanatory variables for each size range and polarity. The data points selected are only between December and February and represent changes in wintertime concentrations. The symbol (-) denotes negative polarity, (+) a positive polarity and (+-) the sum of both polarities. The bolded red values are statistically significant ($p < 0.05$).

625