

# Turbulent structure of the Arctic boundary layer in early summer driven by stability, wind shear and cloud top radiative cooling: ALOUD airborne observations

Dmitry G. Chechin<sup>1,4</sup>, Christof Lüpkes<sup>2</sup>, Jörg Hartmann<sup>2</sup>, André Ehrlich<sup>3</sup>, and Manfred Wendisch<sup>3</sup>

<sup>1</sup>A.M. Obukhov Institute of Atmospheric Physics of the Russian Academy of Sciences, Moscow, Russia

<sup>2</sup>Alfred-Wegener-Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany

<sup>3</sup>Leipzig Institute for Meteorology, Leipzig University, Leipzig, Germany

<sup>4</sup>Moscow Center for Fundamental and Applied Mathematics, Moscow, Russia

**Correspondence:** Christof Lüpkes (christof.luepkes@awi.de)

## Abstract.

Clouds are assumed to play an important role for the Arctic amplification process. This motivated a detailed investigation of cloud processes including radiative and turbulent fluxes. Data from the aircraft campaign ALOUD were analyzed with a focus on the mean and turbulent structure of the cloudy boundary layer over the Fram Strait marginal sea ice zone in late spring/early summer 2017. Vertical profiles of turbulence moments are presented from contrasting atmospheric boundary layers (ABLs) from four days. They differ by the magnitude of wind speed, boundary-layer height, stability, by the strength of the cloud-top radiative cooling and by the number of cloud layers. Turbulence statistics up to third order moments are presented, which were obtained from horizontal level flights and from slanted profiles. It is shown that both of these flight patterns complement each other and form a data set that resolves the vertical structure of the ABL turbulence well. The comparison of the four days shows that especially during weak wind, even in shallow Arctic ABLs with mixing ratios below  $3 \text{ g kg}^{-1}$  cloud-top cooling can serve as a main source of turbulent kinetic energy (TKE). Well-mixed ABLs are generated where TKE is increased and vertical velocity variance shows pronounced maxima in the cloud layer. Negative vertical velocity skewness points then to upside-down convection. Turbulent heat fluxes are directed upward in the cloud layer as a result of cold downdrafts. In two cases with single-layer stratocumulus, turbulent transport of heat flux and of temperature variance are both negative in the cloud layer suggesting an important role of large eddies. In contrast, in a case with weak cloud-top cooling, these quantities are positive in the ABL due to the heating from the surface.

Based on observations and results of a mixed-layer model it is shown that the maxima of turbulent fluxes are, however, smaller than the jump of the net terrestrial radiation flux across the upper part of a cloud due to the i) shallowness of the mixed-layer and ii) the presence of a downward entrainment heat flux. The mixed-layer model also shows that the production of TKE is substantially smaller in stratocumulus over Arctic sea ice compared to subtropics due to a smaller surface moisture flux and smaller decrease in specific humidity (or even humidity inversions) right above cloud top.

In a strong wind case, wind shear is shaping the ABL turbulent structure, especially over rough sea ice, despite the presence of a strong cloud-top cooling. In the presence of mid-level clouds, cloud-top radiative cooling and thus also TKE in the

lowermost cloud layer are strongly reduced and the ABL turbulent structure becomes governed by stability, i.e., by the surface-air temperature difference and wind speed. A comparison of slightly unstable and weakly stable cases shows a strong reduction of TKE due to increased stability even though the absolute value of wind speed was similar. In summary, the presented study documents vertical profiles of the ABL turbulence with a high resolution in a wide range of conditions. It can serve as a basis for turbulence closure evaluation and process studies in Arctic clouds.

## 1 Introduction

Within the past two decades an extraordinary climate change has been observed in the Arctic. Numerous processes and feedback mechanisms have been proposed and discussed as possible reasons for the currently ongoing changes resulting in an enhanced Arctic warming compared to mid-latitudes, which is generally called the Arctic amplification. The effect of the processes and feedbacks has well been documented by a large number of modelling and observational studies (e.g., Serreze and Francis, 2006; Graversen et al., 2008; Overland et al., 2011; Wendisch et al., 2017; Osborne et al., 2018). While the changes are obvious, the chain of interlinked processes and complex air-sea ice-ocean interactions leading, e.g., to reduced sea ice cover is not yet fully explored and quantified. One of the key factors in the framework of Arctic amplification feedbacks are clouds and their various effects, which are in the center of this work. In early studies (e.g., Curry et al., 1996; Morrison et al., 2011; Shupe et al., 2011) it was found that low-level stratocumulus clouds frequently occur over the Arctic and influence the surface heat budget and consequently the sea ice mass budget. In more recent investigations, clouds have been recognized to be involved in different feedback mechanisms contributing to the observed warming (e.g., Serreze and Francis, 2006; Pithan and Mauritsen, 2014; Goosse et al., 2018; Stapf et al., 2020). However, despite the progress, there are still gaps in knowledge related to the quantification of the role of specific processes affecting cloud properties over sea ice.

One of the difficulties is the complex interaction of processes of different physical nature and scale, such as radiation, turbulence, and cloud microphysics, as well as their interplay with large-scale synoptic forcing and variable surface conditions (e.g., Curry et al., 1996). In particular, interactions with turbulence, as well as the turbulent exchange with the surface and overlying air have a strong impact on clouds (Morrison et al., 2011). Low-level clouds often reside in the atmospheric boundary layer (ABL) and thus affect its properties and the energy exchange with sea ice. However, cloud layers are also frequently decoupled from the surface-based atmospheric boundary layer (e.g., Curry, 1986). Both the treatment of clouds and ABL over Arctic sea ice remain to be weak points of modern atmospheric models (Tjernström et al., 2005, 2008; Pithan et al., 2014). Therefore, further research focused on a combined consideration of cloud, radiation and turbulence properties is needed.

Although the crucial effects of Arctic clouds on the ABL energetics and turbulence structure are widely accepted, a quantification, especially over sea ice covered regions is difficult since data are available from a limited number of campaigns only. Mixed-phase clouds represent one of the most complex challenges (e.g., Morrison et al., 2011). They are occurring most frequently in the cold seasons but also during early summer, which is considered in this work. The first analyses of the impact of summer Arctic boundary-layer clouds on the turbulent structure and on ABL energy fluxes using in situ aircraft observations were presented by Curry (1986) and Curry et al. (1988) based on a data set obtained more than four decades ago. The detailed

analysis offers insight into the combined effect of radiation and turbulence but findings are based on only four cases with very limited vertical resolution for turbulence measurements. Only three more cases were documented later, namely by Finger and Wendling (1990), Brümmer et al. (1994) and Inoue et al. (2005), providing additional yet still insufficient data for process understanding and model evaluation.

5 These studies have shown that the main effect of clouds is associated with radiative cooling at the cloud top. In particular, the latter leads to convective overturning and generation of turbulence due to buoyancy. This process is well-known for stratocumulus in lower latitudes. As stressed by Curry et al. (1988) Arctic clouds occur frequently in multiple layers. In such cases, the strongest radiative cooling as well as the buoyancy production of turbulence occurs in the uppermost cloud layer, which is far from the surface and thus decoupled. In the lowest, surface-based layer, which is often stably stratified, wind shear becomes the  
10 main mechanism of turbulence generation (Inoue et al., 2005). These findings were confirmed using a large dataset obtained from surface-based cloud radar during the Arctic Summer Cloud Ocean Study (ASCOS, Shupe et al., 2013; Tjernström et al., 2014; Sedlar and Shupe, 2014; Sotiropoulou et al., 2014).

However, due to the very limited number of in situ observations, gaps remain in a large region of the parameter space. Namely, except from the ASCOS cases, the mentioned studies presented cases with a rather strong wind speed ( $10 \text{ m s}^{-1}$  and  
15 stronger). Thus, weak to moderate wind speed conditions still remain uncovered by observations. Also, the multi-layer cases, studied earlier, were characterized exclusively by negative heat fluxes in the surface-based boundary layer. To conclude, the effect of a variable wind shear and stability on the turbulent structure of a cloud-topped ABL and on the cloud properties cannot be well assessed based on these studies alone.

One of the goals of this paper is therefore to extend the earlier data sets by turbulence data obtained for another four cases  
20 from the airborne measurements during polar Day (ACLOUD) campaign (Wendisch et al., 2019). After that, the available data base is still small but it is another step forward, which helps to reach the second goal, namely to gain a better understanding of processes affecting the turbulent structure of the Arctic ABL in the presence of clouds. The data will be used to describe and analyze the mean and turbulent structure of the cloudy boundary layer in terms of turbulent and radiative energy quantities over sea ice found in early summer. The ACLOUD campaign was carried out in May-June 2017 over the Fram Strait marginal  
25 sea ice zone (MIZ) to the north-west from Svalbard. Unlike in most previous studies, we consider not only the more qualitative turbulence structure (as possible by surface-based radar measurements) but quantify also the fluxes of momentum, heat and radiative energy as well as further turbulence quantities such as turbulent kinetic energy, variances and higher-order turbulence moments. Compared with Finger and Wendling (1990), a new aspect is to consider the variability due to the impact of external factors such as the mean wind field and multi-layer clouds and to obtain a better vertical resolution. The latter is also an  
30 advantage compared with the studies of Curry (1986) and Curry et al. (1988).

Our strategy is - as to some extent also in the study of Curry (1986) - to investigate several contrasting cases with low-level clouds highlighting the forcing mechanisms of turbulence and their variability. One of the important forcing parameters is the strength of mean wind, which varies from case to case due to a different synoptic situation. Second, we will show that the intensity of the cloud-top radiative cooling and thus of turbulence is strongly modulated by the presence of multi-layer clouds.  
35 This extends the above mentioned study of Inoue et al. (2005) on the effect of a 2-layer cloud system. Third, we document

how surface roughness and wind direction modulate stratification and turbulence parameters in a cloudy boundary layer and, in turn the ABL height and cloud thickness. We will discuss similarities and differences to earlier findings using the new data set, which will help to better understand the most important features in Arctic boundary layer clouds.

The paper is structured as follows. We introduce the campaign and describe the measurement equipment in Section 2. Results of different flights are presented in Section 3. A final discussion and conclusions follow in Sections 4 and 5, respectively. Appendix A contains information concerning the accuracy of the measurements, Appendix B presents vertical profiles of net terrestrial radiative flux densities and associated cooling rates and Appendix C describes a diagnostic mixed-layer model addressed in Section 4.

## 2 ACLOUD campaign

During the ACLOUD campaign the two research aircraft Polar 5 and Polar 6 (Wesche et al., 2016) of the German Alfred Wegener Institute were used for collocated measurements in the cloudy and cloud free atmospheric boundary layer (ABL) over the open ocean and the marginal sea ice zone to the North and West of Svalbard. The general strategy was to use Polar 5 as a remote-sensing platform for cloud observations from above and Polar 6 for in situ measurements in clouds and below them (Wendisch et al., 2019). During some flights both aircraft were used for in situ observations.

ACLOUD took place during May/June 2017 covering the transition from late spring to early summer meteorological conditions at the end of May (Knudsen et al., 2018). During the first two weeks of June 2017 several episodes with warm air advection took place so that the air temperature increased over sea ice but it remained still below zero. The majority of flights above, in and below low-level stratocumulus clouds over sea ice and open ocean were performed during this period.

During the ACLOUD period the research vessel (RV) Polarstern (PS) of Alfred Wegener Institute was drifting in the sea ice north of Svalbard (Wendisch et al., 2019). Some flights were carried out next to the ship but, e.g. safety rules did not allow parallel measurements in clouds with the tethered balloon (Egerer et al., 2019), which was operated next to the ship.

### 2.1 Aircraft instrumentation and data processing

The instrumentation of both aircraft including radiation and turbulence equipment is overviewed in Wendisch et al. (2019) and in Ehrlich et al. (2019). Here, we only summarize the relevant information about the instrumentation for turbulence and boundary-layer observations. For measurement uncertainty we refer to Hartmann et al. (2018) and to further aspects described in Appendix A.

Both aircraft were equipped with identical meteorological instrumentation mounted at the nosebooms. It includes the Rosemount 858 five-hole probes, and two fast-response open-wire Pt100 temperature sensors in deiced and non-deiced Rosemount housings. Each aircraft was also instrumented with slow-response Vaisala temperature and humidity sensors (HMT-333) and also with static pressure sensors. Downward and upward shortwave and terrestrial longwave radiative energy fluxes were measured using Kipp and Zonen radiometers (CMP22 and CGR-4) installed at both aircraft.

The five-hole probe measures dynamic and static pressures and two differential pressures in orthogonal directions. Those data are used for the calculations of the true air speed, of the angle of attack  $\alpha$  and the sideslip angle  $\beta$  of the aircraft. Both five-hole probes were equipped with effective deicing and purging systems for liquid water entering the pressure tubes in cloudy air. If not purged, the liquid water can cause serious problems, especially when it freezes in the tubes.

- 5 The measurements of position, movement, and attitude of the aircraft are based on an inertial navigation system (INS) and are merged with the GPS signal. The data provide the aircraft roll, pitch, and yaw angles, ground speed and heading and vertical speed. These measurements yield the aircraft velocity vector relative to the Earth's fixed coordinate system. From the latter and the airflow vector provided by the 5-hole probe, the three components of the wind vector in geographic coordinates are calculated (e.g., Lenschow et al., 1986).
- 10 The majority of the ABL clouds observed during the campaign was dominated by liquid particles (Wendisch et al., 2019). Thus, only the liquid water content (LWC) is considered here as measured by a Nevzorov probe (Korolev et al., 1998) installed on Polar 6.

## 2.2 Flight patterns

Usually, horizontal flight sections are considered when turbulence moments are derived from high frequency aircraft measurements of meteorological variables. Turbulence statistics are calculated then by application of the eddy-covariance method. Different requirements exist for the length of horizontal flight sections. First of all, they should be long enough to keep the statistical sampling error small (Lenschow and Stankov, 1986; Lenschow et al., 1994). Practically, the necessary flight length depends on the size of the relevant eddies, which in turn is a function, e.g. of the stratification and ABL height. The optimal length must consider also horizontal homogeneity and stationarity to separate mesoscale circulations along the track from turbulence. We found (see Appendix A) that the length of flight sections between 8 km and 18 km was sufficient to gain reliable vertical profiles of turbulence statistics. This result is based on both the statistical accuracy and the physically explainable structure of the obtained profiles of turbulence moments. Also the comparability of results of repeated flights with one or two aircraft at the same location was considered.

25 The difficulty of measuring the vertical flux profiles is that due to the range limitation of an aircraft only few flight levels are possible. Our strategy was to fly about 5-10 horizontal sections in staggered altitude levels (partly exactly upon each other, partly as a double-triangle pattern (Figures 1 and 2) as described also in Ehrlich et al. (2019). The aircraft range allowed us to sample profiles at 2-4 locations during one measurement flight.

30 The drawback of these horizontal patterns is that the vertical resolution of a related flux profile is limited. For this reason we used also slanted profiles (see Figure 1) with low ascending and descending rates of about  $0.5 \text{ m s}^{-1}$  to  $1.5 \text{ m s}^{-1}$ . This method has been used earlier by Mahrt (1985), Lenschow et al. (1988), Tjernström (1993), Aliabadi et al. (2016). Turbulence statistics were calculated using a moving window of about 100 s width corresponding to a layer of 100 m thickness and to a flight distance of about 5-10 km. Within each moving window a polynomial trend was removed. One can interpret the obtained profiles in a way that every point represents an approximation of the true values averaged over the corresponding height interval. Thus, this method provides continuous vertical profiles but suffers from two drawbacks: 1) relatively short time spent

at a certain height resulting in a higher statistical uncertainty; 2) smoothing of the vertical profiles. It should be stressed that the obtained statistics from slanted profiles do not strictly approximate the values from horizontal legs, except the cases when turbulence statistics change with height very slowly. It is important that the flight segments crossing the inversion at the ABL top are excluded from the analysis of slanted profiles so that the considered profiles start below the ABL top and steep jumps of quantities like temperature and wind components in the capping inversion are not misinterpreted as turbulence effects.

As in Tetzlaff et al. (2015), who considered measurements over sea ice with sometimes also low values of heat fluxes, we show in Appendix A that the accuracy of the used turbulence probe was in horizontal legs sufficiently high to measure heat fluxes in the range of at least  $5 \text{ W m}^{-2}$ . Also in Appendix A, specific uncertainties related with measurements in clouds are addressed. Comparison of the results of the two different aircraft and plausibility of obtained profiles point to even higher accuracy (see Figure 13 and its description).

### 3 Results


The main goal of our analysis is to discuss the variability of the cloud impact and to explain the differences between the observed cases. This will be done on the basis of mean variables but also of turbulent moments such as the vertical fluxes of sensible heat  $H$ , and components of momentum fluxes  $M_u$  and  $M_v$ . Here, the wind vector was rotated so that the u-component is pointing into the mean wind direction averaged over the boundary layer and the v-component points to the orthogonal direction. Furthermore, we consider the turbulent kinetic energy (TKE), variances of velocity components (also after rotation)  $\sigma_u^2$ ,  $\sigma_v^2$ , and  $\sigma_w^2$  and of potential temperature  $\sigma_\theta^2$ , the skewness of vertical velocity  $S_w = \overline{w'^3} / \overline{w'^2}^{3/2}$ , the vertical turbulent transport of potential temperature flux  $T_{SH} = \overline{w'\theta'}$  and the vertical transport of potential temperature variance  $T_{TV} = \overline{w'\theta'^2}$ .

The description of cases does not follow a chronological order but starts with two single-layer low cloud cases and ends with multi-layer cloud cases. A weak-wind single-layer case is described first in detail because it serves as a reference case where the cloud effect on the ABL structure was clearly dominant throughout the ABL.


#### 3.1 A single-layer cloud case with weak wind (5 June 2017)

On 5 June, a single-layer stratocumulus deck was present over the sea ice to the north of Svalbard. On the MODIS image one can see the contours of sea ice floes (Figure 2) through the cloud layer, however, the clouds found during the flight, two hours after the satellite overpass, appeared opaque (Fig. 3f). The region of observations was over sea ice only on the periphery of an atmospheric high-pressure ridge with weak horizontal pressure gradients causing a weak south-westerly flow along the flight track in the lowest 500 m over the surface.

On the way north to PS and back to south, Polar 6 was performing slow ascents and descents (saw-tooth profiling), as shown by colored segments in Figure 2. Over PS, Polar 6 performed a double-triangle pattern with horizontal legs in different altitudes. Clearly, all profiles were obtained over sea ice but patches of open water (lead-like structures) were also observed. In the following, results of horizontal flight legs over PS and of slanted profiles along the track from T1 to the ship and back are discussed.



**Figure 1.** Flight pattern in clouds showing staggered horizontal legs and slanted profiles. Turns between legs were not included in the data analyses.



**Figure 2.** Polar 6 track on 5 June 2017 overlaid over the MODIS satellite image. T1-T5 represent descends and ascents of the aircraft and PS is the position of RV Polarstern. Wind bars indicate the ABL averaged wind. The bars are plotted for wind stronger than 2.5 knots, while weaker wind is indicated by an open circle. Isobars represent the mean sea level pressure field based on the ERA5 reanalysis (Hersbach et al., 2020).



**Table 1.** The characteristics of the observed profiles on 5 June 2017.

Profile	$h_{top}$ (m)	$h_{cb}$ (m)	$\Delta\theta_{top}$ (K)	$\Delta q_v$ (g/kg)	$\Delta LW_{net}$ ( $Wm^{-2}$ )	$LWP$ ( $gm^{-2}$ )	$w_*$ ( $ms^{-1}$ )	$\theta_*$ (K)
T1	420	90	2.8	0.4	75	72	0.41	0.017
T2	540	320	5.4	0.6	60	20	0.59	0.023
T3	525	300	4.9	0.5	58	21	0.45	0.014
T4	480	240	4.8	0.5	56	44	0.42	0.014
T5	500	300	4.8	-0.17	61	–	0.47	0.016
PS	420	170	4.5	0.45	63	30	0.50	0.022

The values of  $w_*$  and  $\theta_*$  are calculated using Equations C14-C16 with the observed heat flux profiles.


We first consider Figure 3 showing vertical profiles of dynamic and thermodynamic parameters measured during profile patterns introduced in Figure 1. The profiles document similarities at the different positions but also certain variability. Main characteristics of the shallow boundary layer are summarized in Table 1.

We find the following most important features. All profiles of potential temperature (Fig. 3a) indicate that the ABL was well mixed and capped by a sharp temperature inversion at cloud top around 400 to 500 m height. The latter is visible from LWC shown in Fig 3f. The temperature inversions went along with strong jumps in relative humidity at cloud top (Fig. 3d). Also the mixing ratio decreased above the cloud, although less pronounced and with some variability allowing small maxima (Fig. 3e) right above the cloud top and some of them still within the capping inversion. This phenomenon is best seen at T1, but also at T5 where another maximum occurs also 200 m above cloud top. We point to this because such maxima and their possible origin have been described, e.g., by Egerer et al. (2021) who found a stronger maximum at the PS location on the same day using a tethered balloon. However, the aircraft measurements point to the spatial variability of the phenomenon.

The ABL wind was very weak with a horizontal speed of  $1\text{ m s}^{-1}$  to  $2\text{ m s}^{-1}$ . In the southern profiles (profiles T1 and T5 in Fig. 3b), a low-level jet was present above the mixed-layer. Vertical profiles of wind direction (Fig. 3c) show that wind appeared rather uniform in the cloud layer (visible from Fig 3 f), while a step change in direction below the cloud base is clearly visible in several profiles, especially at T4 and T5. This might hint to a decoupling between the well-mixed cloud layer and the surface-based boundary layer in some profiles but might also represent random variability. At the same time, the potential temperature and specific humidity profiles show no signs of decoupling.

The ABL variability along the considered track was most obvious in the LWC profiles (Fig. 3f), showing LWC-maxima between  $0.2$  and  $0.35\text{ g m}^{-3}$ . Cloud bases were between 100 and 300 m and thus were more variable than cloud tops (see above). As it is typical for a well-mixed stratocumulus layer, the LWC was increasing almost linearly with height with the same slope in all profiles. This was confirmed also by the results of the averaged horizontal flight legs.

In the case investigated here a strong cloud-top cooling was present in all profiles (see radiation profiles in Appendix B). The jump of the net terrestrial energy flux  $LW_{net}$  across the cloud-top layer was in the range of  $50$  to  $70\text{ W m}^{-2}$  (Figure B1). We



**Figure 3.** Vertical profiles of the mean potential temperature, wind speed and direction, absolute and specific humidity and the liquid water content obtained during slant ascents and descents of Polar 6 on its way to PS and back on 5 June 2017. The locations of the ascents and descents are shown in Figure 2.

assume here that  $LW_{net}$  is positive, when the net flux is directed upward (as in, e.g., Nicholls and Leighton (1986), their Fig. 7). The impact of the radiative cloud top cooling is obvious in the profiles of turbulence moments discussed in the following.

To allow a comparison of the fluxes measured at positions T1 to T5 and at PS we introduce the scaling

$$z' = \frac{z - c_b}{h - c_b} \quad \text{if } z \geq c_b \quad \text{and} \quad z' = \frac{z - c_b}{c_b} \quad \text{if } z < c_b \quad (1)$$

5 of the vertical axis where  $z$  is height,  $h$  is cloud top and  $c_b$  is cloud base.

In Fig. 4 we show the results of the slanted profiles and of the horizontal sections. For their comparison one should keep in mind their horizontal distance from each other. This may explain some of the differences between both types of results. However, main characteristics are similar. For example, values of turbulent heat flux  $H$  were clearly upward in the cloud and reached maxima of about  $10 \text{ W m}^{-2}$ . This was slightly larger than in the layer below the cloud where at PS the surface-generated convective conditions led to clearly upward fluxes. At the positions of the slanted profiles  $H$  was near zero in the subcloud layer (Fig. 4a). Thus, the profiles of  $H$  deviated from a shape typical for quasi-stationary convective ABLs

developing over a heated surface where  $H$  decreases linearly with height. Obviously, positive heat flux was generated by the cold downdrafts (due to cloud-top cooling) and warm updrafts. At the cloud top, one would expect a substantial negative heat flux due to entrainment considering the presence of a temperature jump across the inversion. But only at PS the value of  $H$  was negative near the ABL top (about  $-3 \text{ W m}^{-2}$  to  $-4 \text{ W m}^{-2}$ ) and thus downward. A possible reason is that capturing the downward fluxes is difficult especially for the slanted profile flights due to the very shallow inversion layer but besides that, the maximum in the spread of  $H$  near cloud base possibly indicates also that the heat flux was carried by eddies of varying size, some which did and some which did not penetrate below the cloud base.


Values of both components of the momentum flux were similar, which is due to the very low wind speed so that mainly convective mixing rather than wind shear was responsible. Nevertheless, directional shear was present at the cloud top and below the cloud base and might also have influenced momentum flux profiles. It can be seen that both components were increasing with height in the cloud with the largest negative values at the cloud base. Below the cloud, both components decreased with height and they were near zero in the layer reached by the aircraft below the cloud (Fig. 4b). Negative and positive values were found close to the cloud top. The scatter in that region is most probably associated with the variability of mixing in the entrainment zone (intermittency) but also with the measurement uncertainty (see below and Section 4.)

The positive heat flux within the cloud layer as a consequence of the radiative cooling at cloud top was responsible for the  $TKE$  generation in the cloud as well as for the maximum in  $\sigma_w^2$  (Fig. 4d,f) in the upper portion of the mixed-layer. The latter is typical for upside-down convection and differs from Lenschow et al. (1980), who find the maximum of  $\sigma_w^2$  in the lower portion of a strong convective, surface forced ABL.  $TKE$  was slightly elevated in the cloud layer below  $z' = 0.9$  (Fig. 4f). Both  $\sigma_u$  and  $\sigma_v$  were nearly constant with height and of same size (Fig. 4d,e) apart from the region below the ABL top. Such a turbulence structure is typical for convective turbulence in a stratocumulus-topped ABL also in lower latitudes (Nicholls, 1984, 1989). But  $TKE$  and  $\sigma_u$  have some large values in the results of horizontal sections close to cloud top. These maxima could be related to the observed directional wind shear across the inversion, as discussed in Section 4.

Another evidence of the upside-down convection was the negative skewness  $S_w$  of vertical velocity throughout the cloud layer (Fig. 4g). Close to the cloud top,  $S_w$  was close to zero. Such a structure of  $S_w$  in the Arctic stratocumulus was reported earlier by Sedlar and Shupe (2014), as well as by Hogan et al. (2009) based on radar and lidar data. In the subcloud layer, a sharp transition of  $S_w$  to zero or small positive values is observed. This might represent a transition from the upside-down convection in the cloud layer to the turbulence generated by shear or slight heating from the surface.

It is also important to consider the turbulent transport of heat flux  $T_{SH}$  (Fig. 4h). In nonlocal turbulence closure schemes this term is seen as the origin of the nonlocal, sometimes also called “countergradient”, transport by large eddies (e.g. Zilitinkevich et al., 1999). We see here again a structure pointing to upside-down forcing, namely downward transport in the cloud layer.

Another third moment, namely, the transport of temperature variance  $T_{TV}$  (Fig. 4i), is also associated with nonlocal transport by large eddies. In the considered case,  $T_{TV}$  had a similar vertical structure as  $T_{SH}$  with negative values in the cloud layer. It should be noted that only two horizontal flight segments show substantial negative values of  $T_{SH}$  and  $T_{TV}$  in the cloud layer. However, vertical profiles of  $T_{SH}$  and  $T_{TV}$  based on slanted profiles provide a clear evidence of negative  $T_{SH}$  and  $T_{TV}$  in the cloud.



**Figure 4.** Vertical profiles of turbulence statistics obtained from horizontal legs over PS (black crosses) and slanted profiles on 5 June 2017. Red lines represent mean values based on four slanted profiles, while red shading indicates  $\pm$  one standard deviation. The cloud layer as observed over PS is shown with grey shading.

Based on the considered vertical profiles of turbulence statistics, as well as on the profile of mean wind direction (Fig. 3c) we can conclude that the fully mixed layer associated with the cloud was perhaps weakly coupled to the surface-based mixed layer. As discussed above, this is most obvious in the vertical profile of  $S_w$ . The absolute values of other statistics such as  $H$ ,  $\sigma_u$ ,  $\sigma_v$ ,  $TKE$ ,  $T_{SH}$ , and  $T_{TV}$  also show a slight, yet a clear decrease in the subcloud layer. It is important to note that based on horizontal legs alone it would be harder to observe this. This shows the advantage of considering slanted profiles together with level flights.


Figure 4c also shows an increased temperature variance close to the ABL top. This is typical for convective and cloud-topped ABLs and is related to the entrainment of air from the inversion layer with strong temperature gradient (Nicholls, 1984, 1989). Young (1988) found that one of the reasons for the increasing variance of  $\theta$  at the top of a strong convective ABL is the forcing by the vertical gradient of temperature rather than turbulent transport of the temperature variance through the capping inversion. This is confirmed here since at the cloud top  $T_{TV}$  is near zero. But the visually observed undulating cloud top, as well as possible gravity waves in the stably stratified inversion layer might have contributed also to an increased temperature variance near the capping inversion.

### 3.2 A single-layer cloud case with strong wind (2 June 2017)


Another single-layer stratocumulus case observed during ACLOUD on 2 June represents a flow directed almost parallel to the MIZ but with a slight on-ice flow component. It differs from the 5 June case by a stronger wind and stronger horizontal gradients of the ABL height and temperature. Another difference is that the flight was performed not only over sea ice as the flight on 5 June, but the first part was over open water and thus, as discussed further, strong horizontal inhomogeneity in the cloud field was observed along the flight track from South to North. On that day, Polar 6 performed saw-tooth profiling over open water and sea ice on the way north and a double-triangle pattern in the cloud-topped boundary layer over sea ice at the northern end of the track (Fig. 5).

Figure 6 shows vertical profiles of mean variables as observed during the saw-tooth ascents and descents. Clearly, a thick mixed layer was present over the open water up to about 750-800 m height (profiles T1, T2 and T3). The ABL height gradually decreased along the south-north flight track and reached about 350-400 m in the northernmost profiles T6 and T7. The cloud layer thickness also decreased from 500 m in the southern profiles to 250-300 m in the northern profiles (Fig. 6f). A strong capping inversion was observed over both open water and sea ice. The magnitude of the temperature jump at the cloud top reached 5 K to 7 K (Fig. 6a). Drier air was lying over the ABL and the jump-like decrease of the mixing ratio was about 2  $\text{g kg}^{-1}$  across the cloud top (Fig. 6e).

All wind speed profiles show a decrease of wind speed towards the mixed layer top, either starting from the lowest measuring level (profiles over water) or from a maximum in a low-level jet as in the profiles over sea ice (Fig. 6b). There, wind speed was highest (profiles T5 and T6) and reached about  $14 \text{ m s}^{-1}$ . Such a low-level jet could have been produced by the sloping inversion. As shown by Chechin and Lüpkes (2017), the increase of the ABL height towards South in the presence of a capping inversion would result in an increase of the easterly wind component in the ABL.



**Figure 5.** Polar 6 flight track on 2 June 2017 overlaid on the MODIS satellite image. T1-T7 represent descends and ascents of the aircraft. The wind barbs are showing the wind speed and direction averaged over the profile segments within the ABL. Isobars represent the mean sea level pressure field based on the ERA5 reanalysis.



**Figure 6.** Vertical profiles of the mean potential temperature, wind speed and direction, absolute and specific humidity and the liquid water content obtained during slant ascents and descents of Polar 6 on its to Polarstern and back mageon 2 June 2017. The locations of the ascents and descents are shown in Figure 5.

The main reason for such a South-North variability of the ABL height and temperature might have been that the ABL wind was not directed along the flight track, but from southwest (220 degree). Thus air parcels that arrived at positions T6 and T7 were influenced by sea ice in the MIZ while T1-T3 rather represent open ocean conditions.

For the presentation of turbulence statistics we use in the following  $z/h$  as the vertical coordinate and not the more complicated  $z'$  as for 5 June. This is done although again several profiles are considered (slanted profiles T5, T6 and horizontal legs over Polarstern) at various positions and with different values of the ABL height  $h$ . However, the ratio  $c_b/h$  where  $c_b$  is the cloud base height, is about the same for these profiles. Thus, using  $z'$  would not result in any difference as compared to using simply  $z/h$ .

Similar to the 5 June case, a strong cloud-top terrestrial radiative cooling was present on 2 June both over open water and sea ice. The cooling rates were similar with about  $-9 \text{ K h}^{-1}$  maximum cooling for all profiles independent if over sea ice or open water (Fig. B1). The cooling explains some of the observed characteristics of the turbulent structure. Over sea ice, these

characteristics differ strongly from those of the 5 June case. This becomes clear from a comparison of the results from slanted profiles shown in Figure 4 with those in Figure 7 (Profiles T5 and T6). The main difference is the occurrence of a pronounced  $TKE$  maximum in the lower portion of the ABL on 2 June, which does not exist on 5 June. It is clearly related to the maxima of  $\sigma_u^2$  and  $\sigma_v^2$  and not to the maximum of  $\sigma_w^2$ . Near the surface, the absolute values of  $\sigma_u^2$  and  $\sigma_v^2$  reach 1.0 and 0.5  $\text{m}^2\text{s}^{-2}$  (during slanted profiles), respectively, which is two to three times larger than the largest value of  $\sigma_w$ . On 5 June,  $\sigma_u^2$  and  $\sigma_v^2$  were much smaller and  $\sigma_w^2$  had a pronounced maximum in the upper portion of the ABL and thus in the cloud. Although the vertical resolution obtained by the horizontal legs is not high enough on 2 June to resolve the lower maximum, these measurements do not contradict the slanted profiles (Figure 7). In the cloud layer, some values of  $\sigma_w$  are somewhat elevated relative to the value obtained by the lowest horizontal leg on 2 June. This hints to two sources of turbulence: one in the cloud layer and the second one close to the ground.

The downward increase of  $TKE$  towards maxima in the lower part of the ABL, which is most pronounced at positions T5 and T6, hints to the dominant role of the  $TKE$  production by wind shear. One can conclude furthermore that the overall buoyancy production of  $TKE$  was relatively small because only small positive values of sensible heat flux occurred in the cloud layer. In the lower part of the ABL we detected negative heat flux. The latter was due to the warm air advection over colder sea ice. Thus, the ABL was being cooled both at its top and bottom. The cooling at the ABL top produced negative vertical velocity skewness  $S_w$  close to the top. In the middle of the ABL, we observed a pronounced shift to positive values of  $S_w$  indicating surface generated turbulence. Another evidence of the influence of the surface is positive  $T_{TV}$  in slanted profiles, which corresponds to the upward transfer of temperature variance.


We conclude that the surface was shaping the turbulence structure but slight cloud-generated turbulence was still present, although it did not govern the entire ABL turbulent structure.

### 3.3 Multi-layer cloud cases


#### 3.3.1 A two-layer cloud case with a slightly unstable boundary layer (14 June 2017)

We consider results from 14 June obtained north of Svalbard where the estimated sea ice fraction was about 90-95 % based on visual observations (Figure 8). The air advected from North was only slightly colder than the average surface temperature of the mixture of ice and water in leads and small polynyas. But this temperature difference was high enough for the development of a slightly unstable ABL in the lowermost 400 m. In contrast to the profiles shown in the previous sections, a mid-level cloud layer existed with base at about 1900 m and top at 2400 m (see LWC in Figure 9f). This cloud layer produced strong cloud top radiative cooling as can be concluded from  $LW_{net}$  indicating a strong vertical divergence at cloud top (Fig. B1e,f). There, the maximal cloud top cooling rate amounted to about  $-11 \text{ K h}^{-1}$ . In contrast, the lower cloud layer could not produce a pronounced cloud top cooling because the loss of terrestrial radiation was compensated by the emission of the cloud base of the overlying cloud layer. This is obvious in the low vertical divergence of  $LW_{net}$ , which was much less pronounced at the top of the lower cloud layer at about 400 m and resulted in about  $-1 \text{ K h}^{-1}$  cooling rate.






**Figure 7.** Vertical profiles of turbulent sensible heat flux  $H$ , vertical velocity variance  $\sigma_w^2$  and skewness  $S_w$ , horizontal velocity components variances  $\sigma_u^2$  and  $\sigma_v^2$  and turbulent kinetic energy  $TKE$  as observed over sea ice on 02 June 2017 based on horizontal flight sections (black crosses) and slanted profiles T5 and T6. The colors of T5 and T6 are the same as in Figures 6 and 5.



**Figure 8.** Tracks of Polar 5 (blue) and Polar 6 (red) on 14 June 2017 overlaid over MODIS satellite images. The wind barbs are showing the wind speed and direction averaged over the profile segments within the ABL. Only the northernmost cross wind leg of Polar 6 and the two northernmost two cross wind legs of Polar 5 are analyzed. The yellow line marks the descent of Polar 6 used for the analysis.

Concerning results of the horizontal flight sections, we concentrate here only on the three northernmost cross-wind staggered legs (Figure 10). The reason is that in this region the surface and cloud conditions showed less variability between and along the different horizontal sections. Also, there were icing problems along the southern legs.


Apparently, a well-mixed layer had developed below cloud top. In the ABL, heat fluxes were directed upward and show a more or less linear decrease with height from values of about  $10 \text{ W m}^{-2}$  at  $z = 0.2h$  to  $0 \text{ W m}^{-2}$  at cloud top ( $z = h$ ). This led to negative values of the bulk Richardson numbers  $Ri_b$  where gradients of temperatures and wind entering  $Ri_b$  refer to the two lowermost legs. Values are  $Ri_b = -0.07$ ,  $Ri_b = -0.05$ , and  $Ri_b = -0.20$  from South to North (or  $z/L = -0.34$ ,  $-0.96$  and  $-0.81$  respectively where  $z$  is the altitude of the lowest horizontal flight leg and  $L$  is the Obukhov length). Based on these values the regime in the observed case can be classified as convection with shear (e.g. Fedorovich and Conzemius, 2008).




**Figure 9.** Results for 14 June 2017 obtained from a descent of Polar 6 in between the locations of the northernmost horizontal legs and the middle ones shown in Figure 8.

Since the external conditions (temperature, wind, ABL top, cloud thickness) did not change substantially between the locations of the flight legs, it is expected that the turbulent quantities varied also very little. And indeed, despite the small signals, the accuracy of both aircraft data is obviously high enough to show very similar results at the different locations (Figure 10).

It is interesting to compare this case with the cases observed on 2 June and 5 June (see Figures 4 and 7). While 5 June was purely cloud driven, 2 June was both cloud driven in the upper part of the ABL and surface driven in the lowest part. Both had a similar heat flux  $O(10) \text{ Wm}^{-2}$  at the cloud top, caused by cloud-top cooling, and which was obviously absent on 14 June. Thus on both days 2 and 5 June the heat flux profile with the maximum near the cloud top was reversed relative to the profile on 14 June with a maximum near the surface. Further differences are: On 14 June, the absolute values of  $\sigma_w^2$  and of  $TKE$  were only half as large as on 5 June despite stronger wind and positive surface heat flux on 14 June.  $\sigma_w^2$  and  $TKE$  were larger on 2 June than on both other days, obviously, due to the much larger wind speed. Another difference to 2 and 5 June is that on 14 June the vertical transport of heat flux  $T_{SH}$ , and of temperature variance  $T_{TV}$  and also  $S_w$  remained positive in the entire ABL (except near the capping inversion). This indicates once more that the processes were not cloud-top driven but surface driven.



**Figure 10.** Turbulence statistics based on horizontal flight legs of 14 June 2017 obtained from Polar 5 (closed symbols) and Polar 6 (open symbols). Results of Polar 5 refer to the two northernmost dashed blue legs shown in Figure 8. The Polar 6 result refers to its northernmost flight leg (red in Figure 8). Altitude is normalized by the cloud top height  $h = 400$  m. The bright symbols (lightblue and orange) refer to  $M_v$ ,  $\sigma_v^2$  while the dark (red and blue) symbols in the corresponding panels refer to  $M_u$  and  $\sigma_u^2$ .




**Figure 11.** Tracks of Polar 5 (blue) and Polar 6 (red) on 20 June 2017 overlaid over the corresponding MODIS image. The yellow line refers to the Polar 5 descent shown in Figure 12.

The reason for this different behavior is clearly the existence of the mid-level clouds, which led to a reduction of the cloud top cooling and thus suppressed mixing at the cloud top.

### 3.3.2 A three-layer cloud case with a stable boundary layer (20 June 2017)


On 20 June, again a multi-layer cloud system was observed in westerly wind conditions and over a region with an ice fraction of about 95 %. Figure 12 shows vertical profiles obtained during a descent of Polar 5 in the northern part of the observational area. No Nevzorov probe was installed on Polar 5, but visual observations and the vertical profile of relative humidity (Fig. 12b) allowed us to detect even three cloud layers with cloud tops at about 350 m, 2100 m, and 3000 m. The strongest cooling occurred at the uppermost cloud top where the maximal cooling rate reached  $-10 \text{ K h}^{-1}$ . Below that cloud top, also the strongest turbulence and positive turbulent heat flux were observed during the slanted descent (not shown here).



**Figure 12.** Results for 20 June 2017 obtained from a descent of Polar 5 at the location of the northernmost horizontal legs (see Figure 11).

Figure 13 shows results for the lowest cloud layer and indicates a fundamental difference between the ABLs observed on this day and on 14 June. On 20 June, the westerly winds led to a near-surface stable stratification. This caused downward but very small absolute values of fluxes of sensible heat. The uncertainty of these values is in the order of the statistical error, nevertheless, the good qualitative agreement of all turbulent quantities obtained from both aircraft shows that the results are reliable enough to see the differences between the ABL structures of 5 June, 14 June and 20 June. Turbulence decreased to almost zero at about  $z = h$ . This is roughly where the capping inversion started. Although no Nevzorov data was available for this flight, visual observations suggested that the cloud layer reached far into this capping inversion. This phenomenon was much more pronounced than on 14 March, when probably surface- and cloud-forced convection led to a well-mixed ABL.

To understand the differences between the results from both flight days with multi-layer clouds, it is important to note that the absolute wind speed did not differ much on these two days. Thus differences between the days are most probably caused by the different density stratifications. The stability observed on 20 June caused strongly reduced values of  $\sigma_w^2$  (also of  $\sigma_u^2$  and  $\sigma_v^2$ , not shown for 20 June) relative to the data on 14 June and consequently, also the  $TKE$  was much lower on 20 June. Variances and  $TKE$  decreased to zero at cloud top, which is not really the case on 14 June. Nevertheless, despite the reduced values



**Figure 13.** Turbulence statistics for the lowermost cloud layer based on horizontal flight legs of 20 June 2017 obtained from Polar 5 (blue symbols) and Polar 6 (red symbols). The results refer to the two blue and red northernmost flight legs of the aircraft (see Figure 11). Closed (open) symbols refer to the northern (southern) positions. Altitude is normalized by the cloud top height, for Polar 5  $h = 190$  m (North) and  $h = 225$  m (South) and for Polar 6  $h = 250$  m (both positions). The bright symbols (lightblue and orange) refer to  $M_v$ ,  $\sigma_v^2$  while the dark (red and blue) symbols in the corresponding panels refer to  $M_u$  and  $\sigma_u^2$ .

of  $TKE$ , a continuously turbulent ABL was observed on 20 June. This is in agreement with the observed values of stability parameters ( $Ri_b = 0.16$  and  $Ri_b = 0.27$  or  $z/L = 0.29$  and  $z/L = 0.56$ , respectively for the Polar 5 legs at the lowest heights), according to which the ABL state can be classified as slightly stable (e.g. Golder, 1972; Mohan and Siddiqui, 1998).

This comparison shows for both flight days with multiple cloud layers an impressive agreement between the measurements of both nosebooms installed at two aircraft. This holds especially for the results of 20 June, when the signals from both nosebooms were very low, but nevertheless, very close to each other. Obviously, the situation was horizontally also very homogeneous, note that the maximum distance between legs was almost 150 km.

#### 4 Discussion

It is important to compare the presented results with previous turbulence datasets obtained in liquid-water clouds over Arctic sea ice. The largest dataset was presented by Curry (1986) and Curry et al. (1988), who considered vertical profiles of mean variables and of turbulence statistics for six cases of summertime clouds in the Arctic.

First of all, we can confirm their conclusion that one of the differences of the Arctic stratocumulus clouds as compared to their marine counterparts in lower latitudes is the frequent occurrence of multiple cloud layers. Curry et al. (1988) concludes that in such situations, strong turbulent mixing occurs in the uppermost cloud layer, which is decoupled from the surface. The lowermost cloud layer can occur as a kind of fog in a stably stratified boundary layer. Exactly such a scenario was observed also by us on 20 June and thus it represents a further hint that such cases might be typical for the summertime Arctic.

However, the dataset presented by Curry et al. (1988) does not contain a multi-layer case with an unstably stratified boundary layer, as we observed on 14 June. The comparison of the two multi-layer cases observed on 14 and 20 June reveals a strong difference in the magnitude of variances and of TKE. This highlights that also in the presence of shallow boundary-layer clouds with low cloud base surface fluxes may strongly affect the lower ABL turbulent structure.

Curry (1986) compares the magnitude of the radiative flux divergence at cloud top with the values of turbulent heat flux in the cloud and finds that only a small fraction of radiative cooling is compensated by turbulent heat transport. She further concludes that only “a small portion of cloud-top cooling results in the production of the mixed-layer convection”. Such a conclusion is based on the values of turbulent heat flux measured below the upper part of the cloud where the strongest longwave cooling is observed (see Table 5 in Curry (1986)). Indeed, the reported values of the turbulent heat flux amount to about  $3 \text{ W m}^{-2}$ , which is lower than the  $10\text{-}15 \text{ W m}^{-2}$  observed by us on 5 June. Our values are, however still much lower than the jump of the net longwave radiative flux  $\Delta LW$  across the cloud top. According to our observations  $\Delta LW$  is about  $50\text{-}70 \text{ W m}^{-2}$  and thus similar to the values reported by Curry (1986). Therefore, the above-mentioned conclusions of Curry (1986) seem to be applicable also to our data, while the reasons behind it remain unclear.

To consider this matter in more detail a simple mixed-layer model is utilized in the following. In particular, it is used to i) understand why the observed turbulent heat flux is much smaller than  $\Delta LW$  and ii) to estimate which portion of  $\Delta LW$  is forcing the mixed-layer convection. Mixed-layer models of different complexity have been frequently used to describe stratocumulus-topped mixed layers (e.g. Lilly, 1968; Nicholls, 1984; Turton and Nicholls, 1987; Stevens, 2002) and the model



used here is similar to the most simple ones formulated earlier (see details in Appendix C). The idea is that we use the observed profile of net terrestrial radiation as well as turbulent heat fluxes observed at the bottom of the mixed layer to diagnose the heat flux profile within the mixed layer. Furthermore, the model allows a calculation of the buoyancy forcing produced by longwave cooling and an estimation of the resulting vertical velocity variance. For the latter we use the Lenschow et al. (1980) parameterization as given by Equation C17.


First, the results of the mixed-layer model are compared with the observed profiles of  $H$  and  $\sigma_w^2$ . The following scenario is considered for 5 June: the bottom of the mixed layer is assumed to be at 50 m height where both sensible and latent heat fluxes are set to zero; the value of the liquid-water potential temperature jump across the mixed-layer is  $\Delta\theta_l = 6$  K, while the jump of total specific humidity is set to  $\Delta q_t = -0.5$  g kg<sup>-1</sup>. Cloud base is assumed to be at 170 m. Entrainment fluxes of  $\theta_l$  and  $q_t$  are parameterized using Equations C11 and C13.

Figure 14a shows the diagnosed sensible heat flux in comparison with the observed one. The mixed-layer model is producing positive heat flux across the mixed-layer with a maximum of about 20 W m<sup>-2</sup> located in the upper part of the cloud. This agrees qualitatively with observations, however, the model overestimates the flux by about a factor of 2.5. The diagnosed entrainment flux of sensible heat is about -4 W m<sup>-2</sup> which is very close to the observed one. Due to the overestimation of the sensible heat flux, the diagnosed value  $w_* = 0.66$  m s<sup>-1</sup> exceeds the values 0.45-0.5 m s<sup>-1</sup> based on observations (see Table 1).

The low values of the observed heat flux in comparison to the diagnosed one might indirectly hint to a more negative entrainment heat flux than measured. In Finger and Wendling (1990), the entrainment heat flux amounted to about -20 W m<sup>-2</sup> which is about 5 times larger than in our case. Prescribing the value -20 W m<sup>-2</sup> at the top of the ABL in the mixed-layer model results in a decrease of the diagnosed heat flux in the cloud layer and in a good agreement with observations (not shown here). We should once more stress the difficulty to interpret observations near the capping inversion due to several factors such as undulating flight height, large vertical gradients of heat flux, and undulations of cloud top due to non-turbulent processes.

The parameterized profiles of  $\sigma_w^2$  using  $w_* = 0.5$  m s<sup>-1</sup> (based on the observed heat flux profile) and  $w_* = 0.66$  m s<sup>-1</sup> (based on the mixed-layer model heat flux profile) are shown in Figure 14b. For both values, the shape of the profile is apparently well described by the Lenschow et al. (1980) parameterization (Equation C17) including the location of the maximum. Despite the agreement of the shape, the smaller (and thus observed) value ( $w_* = 0.5$  m s<sup>-1</sup>) leads, however, to an underestimation of  $\sigma_w^2$  in the whole ABL when compared to the values from horizontal legs. The larger value 0.66 m s<sup>-1</sup> shifts the parameterized  $\sigma_w^2$  profile closer to the observations. An even better fit to observations can be obtained with  $w_* = 0.5$  m s<sup>-1</sup> but only when the constants  $c_1 = 3.6$  and  $c_2 = 0.75$  are used in the Lenschow parameterization (dashed curve in Figure 14b).

The question is why the observed  $\sigma_w^2$  is larger than  $\sigma_w^2$  calculated with  $w_*$  based on the observed heat flux profile. It is helpful to consider the ratio  $\sigma_{w,max}/w_*$  where  $\sigma_{w,max}$  is the maximum value of  $\sigma_w^2$ . In particular, for  $w_* = 0.5$  m s<sup>-1</sup> we obtain  $\sigma_{w,max}/w_* \approx 1$ , which is larger than the values obtained earlier. Previous studies report  $\sigma_{w,max}/w_* \approx 0.6 - 0.65$  for stratocumulus in lower latitudes (Nicholls, 1984; Caughey et al., 1982; Deardorff, 1980) and in a convective boundary layer heated from below (Young, 1988). A possible explanation for this discrepancy is given by Smedman and Hoegstroem (1983) who showed that the ratio  $\sigma_{w,max}/w_*$  is increased when wind shear contributes to the production of turbulence in addition to buoyancy. In the presence of shear, Finger and Wendling (1990) observed similar values of  $\sigma_w^2$  and heat flux as we did on 5



**Figure 14.** Observed and diagnosed turbulent heat flux profiles for 5 June (a) and vertical velocity variance (b) obtained using the Lenschow et al. (1980) parameterization (Equation C17) with  $w_* = 0.5 \text{ m s}^{-1}$  and  $w_* = 0.66 \text{ m s}^{-1}$  and with modified constants  $c_1$  and  $c_2$ .

June, which would also result in  $\sigma_{w,max}/w_* \approx 1$  in the cloud layer. In their case, the ABL wind speed was, however, much stronger and amounted to about  $10 \text{ m s}^{-1}$ . Although the wind speed was weak on 5 June, a directional wind shear was present at the cloud top and base (Fig. 3c), which might have contributed to TKE and  $\sigma_w^2$  production and would explain the large value of  $\sigma_{w,max}/w_*$ .

- 5 Thus, the mixed-layer diagnostics shows several things. First, we can conclude that the heat fluxes observed by Curry (1986) in single-layer stratocumulus are much smaller than one would expect. Our observations are much closer to the mixed-layer model and are also in agreement with Finger and Wendling (1990) who observed turbulent heat fluxes of  $10\text{-}15 \text{ W m}^{-2}$  in the cloud layer. Second, the mixed-layer model shows that the turbulent heat flux can indeed be several times smaller than  $\Delta LW$  and yet it is enough to redistribute the longwave cooling uniformly throughout the mixed layer, as explained in more detail
- 10 further.

To understand why turbulent heat flux can be much smaller than  $\Delta LW$  it is important to consider the following property of the mixed layer. Namely, the total flux of the conservative variable  $\theta_l$  has to change linearly with height as expressed in Equation C3. There, the total flux is the sum of  $LW_{net}$  and  $\overline{w'\theta_l}$ . Due to the fact that usually  $LW_{net}$  changes nonlinearly with height (Figure B1), some amount of turbulent heat flux is needed to compensate for this nonlinearity. The needed amount of


15 turbulent heat flux is proportional to how far from linear the  $LW_{net}$  profile is. Obviously, for a very thin layer (e.g. a layer between  $z/h = 0.8$  and  $z/h = 1$  in Figure B1)  $LW_{net}$  alone is already close to linear so that there the cloud layer would cool almost uniformly with height even with a small amount of turbulent mixing. This is not the case for optically and geometrically thick mixed layers, where the strong longwave cooling in the upper part of the cloud layer generates a strongly nonlinear profile of  $LW_{net}$ . To obtain well-mixed profiles of the cloud liquid water potential temperature and cloud water mixing ratio the loss

of heat at cloud top has to be redistributed then throughout the mixed layer by strong turbulent motions. In such thick clouds, the maximum of the turbulent heat flux would be closer to  $\Delta LW$ . Apart from this, the presence of a negative entrainment flux also leads to a decrease of the turbulent heat flux maximum in the cloud because the entrainment is compensating a part of the longwave cooling. Thus, we can conclude that in thin mixed layers, indeed, turbulent heat flux can be substantially smaller than  
5 the longwave cooling expressed by  $\Delta LW$ . This explains why not the whole amount of  $\Delta LW$  is used to force the mixed-layer convection.

In connection with the above discussion, one should also keep in mind the uncertainty of the observed turbulence statistics. The magnitude of the uncertainty for heat and momentum fluxes obtained from horizontal legs is discussed in Appendix A. Similar to earlier studies (e.g. Curry et al., 1988; Finger and Wendling, 1990; Inoue et al., 2005), the uncertainty is estimated  
10 empirically by comparing results from shorter segments of a long horizontal leg. Theoretical estimates of sampling errors due to an insufficient length of flight legs also exist (Lenschow and Stankov, 1986; Lenschow et al., 1994). Based on large Eddy simulation (LES) Petty (2021) showed that there is a good agreement between the empirical and theoretical estimates. However, neither theoretical estimates nor LES account for the horizontal non-turbulent inhomogeneities that exist in nature and complicate the analysis of the observations. It is even harder to obtain theoretical estimates for slanted profiles as statistical  
15 properties of turbulence are functions of height within the ABL. Thus, we rely on empirical estimates of the uncertainty also for slanted profiles. So, we can consider the range of the slanted profiles (red hatched regions in Figures 4 and 7) obtained at different positions as a measure of accuracy obtained from these profiles. This is not more than a rough estimate and we expect that especially near the inversion (above  $z/h = 0.8$  the measurement errors are large for the slanted profiles but also for the horizontal legs. This hold also for previous aircraft campaigns (Lenschow and Stankov, 1986). In the future, flights should be  
20 performed repeating the same profiles and including horizontal flight legs at the same positions to determine the accuracy.

Finally, it should be noted that theoretically the uncertainty for third moments is higher than for the second ones. It can be reduced by considering longer averaging intervals, but the drawback of long horizontal legs is that horizontal inhomogeneity starts to affect the results. In case of slanted profiles, it is simply not possible to increase the leg length, as it will either cover a very large height interval or require a very slow ascent or descent, so that the horizontal distance flown during ascent/descent  
25 would become too large.

The mixed-layer model also highlights an important difference between the Arctic and subtropic stratocumulus. For subtropics, a substantial latent heat flux at the bottom of the mixed layer is typical, while at the top of the mixed layer dry air is entrained. In the mixed-layer model, this leads to an increased buoyancy and heat flux in the cloud layer and consequently to a stronger turbulence. The reason for that is an increase of condensation and latent heat release in updrafts and of evaporation and associated cooling in downdrafts. This is illustrated by the sensitivity experiments with the mixed-layer model (Fig. 15). The  
30 parameters of the experiments are the same as for the 5 June experiment, apart that we prescribe a gradual increase of the latent heat flux at the bottom of the mixed layer as well as an increase of the total humidity jump at the top of the mixed layer. The typical values for subtropical stratocumulus are  $\rho L_e (\overline{w'q'_t})_0 = 115 \text{ W m}^{-2}$  and  $\Delta q_t = -7.5 \text{ g kg}^{-1}$  (Duynderke et al., 2004; Stevens et al., 2005). In contrast, in the Arctic the surface latent heat flux is small. During the ASCOS campaign, latent heat  
35 flux in the surface layer did not exceed  $5 \text{ W m}^{-2}$  (Brooks et al., 2017). Furthermore, a humidity inversion often occurs at the



**Figure 15.** Sensitivity of the sensible heat flux profile (a) and vertical velocity variance (b) obtained from the mixed-layer model to the value of latent heat flux at the bottom of the mixed layer  $\rho L_e \overline{(w'q'_t)}_0$  and to the value of the jump of total specific humidity at the top of the mixed layer  $\Delta q_t$ . Vertical velocity variance is obtained using the parameterization of Lenschow et al. (1980) (Equation C17) with the original values of  $c_1$  and  $c_2$ .

top of the Arctic mixed layer, which leads to the entrainment of moist air. As a result, an increase of buoyancy is much smaller in the Arctic stratocumulus.

Table 2 summarizes the investigated cases together with observations from earlier studies. It shows that, to our knowledge, our study presents for the first time i) a case with a strong cloud-top cooling and almost zero wind speed, as well as ii) two multi-  
5 layer cases with contrasting stability in the boundary layer. This broadens the parameter space with available observations, which is important for process understanding and model verification and testing.

## 5 Conclusions

We presented and analyzed airborne measurements of turbulence and radiation obtained during the campaign ACLOUD over the marginal sea ice zone to the north and northwest of Svalbard. By combining two types of flight patterns, namely staggered  
10 horizontal legs and slanted profiles, we obtained vertical profiles of turbulent quantities using eddy covariance method. A combination of both flight patterns is shown to be clearly beneficial for the analysis of turbulence profiles.

The presented analysis attempts to distinguish the effect of clouds on turbulence statistics in comparison to the role of other mechanisms of turbulence generation. This was possible because one of the cases, namely, 5 June 2017, was characterized by a near-zero wind speed in the ABL. Thus, the terrestrial radiative cooling at the cloud top was the only relevant mechanism of  
15 turbulence generation apart from a possible influence of directional wind shear, which was of secondary importance. To our

**Table 2.** Summary of the observed cases during ACLOUD in comparison with earlier turbulence observations in stratocumulus-topped ABL over the Arctic sea ice in summer.

Case	Campaign	Cloud layers	$h_{ABL}$ (m)	$\Delta LW_{net}$ or cooling rate	$LWC_{max}$ ( $g\ m^{-3}$ )	$H_{min,max}$ ( $W\ m^{-2}$ )	$ U $ ( $m\ s^{-1}$ )	stratification
2.06.2017	ACLOUD	1	350-450	$50-70\ W\ m^{-2}$	0.3	-10	11-14	stable/mixed
5.06.2017	ACLOUD	1	400-500	$60-80\ W\ m^{-2}$	0.3	10	0-2	mixed
14.06.2017	ACLOUD	2	400	$15-20\ W\ m^{-2}$	0.2	10	6-7	unstable
20.06.2017	ACLOUD	3	190-225	$0\ W\ m^{-2}$	–	-5	6-8	stable
20.06.1980	Arctic Stratus Experiment	1	240	$45\ W\ m^{-2}$	0.4	3	10	mixed
28.06.1980	—”—	2	250	–	0.1	-3	10	stable
26.06.1984	MIZEX	1	400	$5\ K\ hour^{-1}$	0.35-0.4	15	10	mixed
8-9.05.1988	ARKTIS 1988	2	200-400	–	0.15	-27	15-20	stable/mixed
29.07.1998	SHEBA	2	500	$10\ K\ day^{-1}$	0.3	-15	14	stable/mixed

Cases are described in: Arctic Stratus Experiment (Curry et al., 1988), MIZEX (Finger and Wendling, 1990), ARKTIS 1988 (Brümmer et al., 1994), SHEBA (Inoue et al., 2005)

knowledge, this is the first time that such a case of summertime stratocumulus over Arctic sea ice is presented. We showed that the cloud-top cooling is forcing upside-down convection, which shapes the ABL turbulent structure, especially in its upper part and during weak synoptic wind. In situations with only one cloud layer, the cloud impact is identified by local maxima of the vertical velocity variance and heat flux in the cloud layer, as well as by a negative vertical velocity skewness. In this respect, an Arctic stratocumulus-topped ABL is qualitatively similar to other cloud-topped ABLs in lower latitudes despite its shallowness and lower humidity.

We showed that strong cloud-top cooling causes the third moments  $T_{SH} = \overline{w'^2\theta'}$  and  $T_{TV} = \overline{w'\theta'^2}$  to be negative in the cloud layer hinting at the possible importance of nonlocal transport associated with large eddies. However, this is only a hint and quantifying the role of the third-order transport has to be a subject of a separate study. Here, we did not analyze the second-order moment budgets and thus cannot conclude on the relative role of various processes, as was done for a cloud-topped ABL in the LES-based study by Heinze et al. (2015).

Based on results of a mixed-layer model we showed that just a part of the net longwave cooling is forcing mixed-layer convection. In particular, the amount of sensible heat flux generated by the longwave cooling depends on how far from linear is the profile of the net longwave flux. Thus, the maximum of sensible heat flux in a boundary layer cloud is expected to be small in a shallow mixed layer and large in a thick mixed layer. In particular, this explains why the observed heat flux maximum is

much smaller than the jump of the net longwave flux across the upper part of the cloud. Thus, the amount of buoyancy forcing for the mixed-layer turbulence depends on the mixed-layer thickness.

In conditions with strong wind and a very shallow boundary layer shear-generated turbulence played a more important role for the ABL turbulent structure than the cloud impact. This statement holds true even in the presence of strong cloud-top cooling. Namely, the strong-wind case that has been investigated was characterized by a pronounced maximum of the variance of all three velocity components and TKE located at low altitude in the subcloud layer.

When mid-level clouds were present, the cloud-top cooling in the lower cloud layer located in the ABL became essentially weaker. This resulted in a much weaker turbulence in the ABL even despite stronger wind. In such conditions, the ABL stability associated with surface-air temperature difference and wind speed apparently becomes an important factor influencing the ABL turbulence. This concerns both turbulence magnitude and the ABL turbulent structure.

To conclude, our results suggest that cloud-top cooling is one of the major sources of turbulence in the spring-summer Arctic ABL even despite its typical shallowness and relative dryness. We found that the amount of such cooling strongly depends on the presence of mid-level clouds, as concluded earlier also by other studies based on radar observations. The importance for the ABL structure depends among other factors mainly on wind speed, ABL height, and stability.


Finally, based on mixed-layer model diagnostics, we showed that despite qualitative similarities of the stratocumulus-topped boundary layers in the Arctic and lower latitudes, there are also substantial differences. In particular, low latent heat flux at the surface and small total specific humidity jump at the ABL top (or even a humidity inversion) result in a much smaller buoyancy forcing as compared to subtropical stratocumulus. In addition, as already stressed above, in shallow mixed layers a smaller amount of turbulent heat transport is needed to redistribute the effect of cloud-top cooling across the mixed layer. This further decreases buoyancy heat flux in the mixed-layer. As a result, smaller magnitudes of turbulent heat flux maxima and TKE are expected in clouds over Arctic sea ice as compared to lower latitudes. The situation might be different, however, for clouds over open water, especially in case of large air-sea temperature differences.

Thus for an adequate representation of the ABL turbulence in coarse-resolution models a tight coupling between the radiative, microphysical, and turbulence parameterizations is needed. The presented cases may serve as a reference for further studies focusing on the evaluation of such parameterizations.


We stress that the number of cases, which we considered, was limited and the results should motivate for future research. However, we have shown by the few cases that there is large complexity due to the many forcing parameters but a detailed consideration helps to understand the mechanisms. Moreover, an understanding of these processes is relevant for the polar climate due to the strong impact on the energy fluxes. Future measurements would be helpful to further study the described phenomena.

## **Appendix A: Accuracy**

Before ACLOUD the Polar 5 and 6 turbulence equipment was never used inside clouds and especially not in clouds above sea ice. Such conditions are challenging because, e.g. heat fluxes are mostly much smaller than in strong convective conditions



**Figure A1.** Frequency weighted power spectra ( $S_u$ ,  $S_v$ ,  $S_w$  for wind components,  $S_t$  for temperature, and  $S_{wT}$  for temperature flux) and ogive  $O_{w,\theta}$  as measured on 5 June 2017 in the cloud at 254 m height.



**Figure A2.** Left panel: Fluxes of sensible heat and of absolute values of momentum as measured on 5 June 2017. Whiskers in both panels show the sampling error  $\epsilon$  determined following Friehe et al. (1991)

over the open ocean in a surface-forced convective ABL during cold air advection (see airborne measurements described by Gryanik and Hartmann (2002)) or over leads described in earlier applications of the used turbulence probe by Tetzlaff et al. (2015). Thus the accuracy of the measurements needs to be reconsidered. Moreover, as in earlier studies (Curry, 1986; Finger and Wendling, 1990) it is important to quantify empirically the uncertainty of the derived turbulence statistics. Apart from  
 5 instrumentation characteristics, other factors are influencing the accuracy as described in Section 2.2. The accuracy of the turbulence measurements in clouds will be considered in the following for the typical case of 5 June 2017.

On this day, eight horizontal flight legs have been flown, one clearly below the cloud layer, five legs within the cloud and another two sections above the cloud. Cloud base and cloud top have been observed at 200 m and 400 m, respectively. Frequency weighted power spectra for three wind components, potential temperature and heat flux are shown in Figure A1 as  
 10 measured in the cloud. Spectra of  $u$ ,  $v$ , and  $w$  show the characteristic slope of  $-2/3$  (corresponding to  $-5/3$  without weighting) in the inertial subrange over more than two orders of magnitude. Similarly, heat flux multiplied with the frequency shows the required  $-4/3$  slope. Temperature variations at high frequencies (above 5 Hz) become very small and reach the detection limit. Thus the temperature variance spectra begin to exhibit some white noise beyond 5 Hz. However, this has no serious impact on the measured heat flux, which becomes obvious by considering ogives  $O_{w,\theta}$  representing the cumulative integration of the  
 15 cospectrum  $C_{o_{w,\theta}}$  between the highest and lowest measured frequency (Friehe et al., 1991). Based on the ogives shown in Figure A1 one can conclude that at least 95 % of the fluxes are caused by contributions below 1 Hz. Maxima for  $w$  and  $\theta$  are



between 0.1 Hz and 0.2 Hz dependent on the aircraft height. This corresponds to wave lengths of about 1.5-3.5 km, when we assume an average aircraft speed of  $55 \text{ ms}^{-1}$ . The ogive shows some deviation from an ideal S-shape at the low frequency (large wave length) end of the spectrum. This might point to a too short leg length, however, a prolongation would have led to other difficulties since usually, homogeneity of turbulence is not guaranteed over larger distances than the 10-18 km during our  
5 measurements because of inhomogeneities in the cloud structures.


Also in horizontal flight sections an aircraft cannot fly always exactly in one altitude. Especially in the uppermost part of the cloud with the capping inversion the remaining fluctuation of aircraft height of about  $\pm 10 \text{ m}$  is therefore correlated with changes of potential temperature caused by its vertical gradient. We corrected this impact on the temperature series based on the mean measured vertical temperature gradient along the flight leg. The impact of this correction is not large (in the range of  
10 the sampling error  $\epsilon$ ).  $\epsilon$  was determined as in Friehe et al. (1991) dependent on the leg length and flight altitude above ground. Figure A2 also shows the momentum flux  $M$  with the corresponding  $\epsilon$ .

We need to address, however, also some uncertainty of air temperature observations in clouds caused by evaporation of liquid from the sensing element (Lenschow and Pennell, 1974) and of droplets during the adiabatic heating of air in the housing of the sensor (Lemone, 1980; Albrecht et al., 1979). We have to leave an experimental determination of this error for future  
15 research for the used Rosemount temperature sensor and can only speculate that the related error for turbulence is probably not too large in the cold Arctic conditions with relatively low absolute humidity. With respect to the accuracy of the temperature measurements it is a good sign at least that results of both aircraft agree well when flight legs have been flown close to each other (see Ehrlich et al. (2019)) within a thin cloud at a distance of about 200 m. At such a distance one can expect that both  
20 temperature sensors have not been exposed to the same humidity conditions so that a possible error due to evaporation effects might have caused differences in the temperatures.


Also, in Sections 3 and 4 we show that all vertical profiles can be physically well explained. This concerns both the profiles derived from horizontal flight legs and those from the slanted profile flights. Concerning the latter, we need to assume that the statistical error is probably larger as compared with the horizontal legs, so that especially those results need to be considered with some caution. Nevertheless, results are at least reasonable. It is also important to see that profiles of different turbu-  
25 lence moments do not show contrasting results. This gives additional confidence in the data that is beyond the pure statistical evaluation and should also be kept in mind when the accuracy is considered.

## Appendix B: Terrestrial radiation flux density and cooling rates

Figure B1 shows the net terrestrial radiative flux (left panels) as well as the radiative cooling rates in the atmospheric boundary layer for the considered cases. Note that on 14 and 20 June the strongest radiative cooling was observed in the upper cloud  
30 layers at heights about 2500 and 3000 m (not shown here).



**Figure B1.** Vertical profiles of the net terrestrial radiative energy flux  $LW_{net}$  and of the cooling rate associated with the vertical divergence of  $LW_{net}$ . The altitude is normalized by the cloud top height  $h$  so that individual profiles can be compared.



**Figure C1.** Schematic representation of the vertical profiles of the conserved variables  $\theta_l$  and  $q_t$  in the mixed layer.

### Appendix C: Diagnostic mixed-layer model

We choose the following variables that are approximately conservative during the moist-adiabatic processes in the mixed-layer: the liquid-water potential temperature  $\theta_l$  and the total liquid water mixing ratio  $q_t$ . The latter two are defined as

$$\theta_l = \theta - \left(\frac{\theta}{T}\right) \frac{L}{c_p} q_c, \quad (\text{C1})$$

$$5 \quad q_t = q_v + q_c, \quad (\text{C2})$$

where  $L$  is the heat of vaporization,  $c_p$  is specific heat at constant pressure,  $q_c$  is the liquid water content and  $q_v$  is specific humidity. Below the cloud base, i.e. for  $z < h_{cb}$  we have simply  $\theta_l = \theta$  and  $q_t = q_v$  as shown in Fig. C1.

As follows from the mixed-layer assumption, the total vertical fluxes of the conserved variables are linear functions of  $z$  for  $h_b \geq z \geq h_t$ , where  $h_{b,t}$  are the bottom- and top heights of the mixed-layer. Namely, one obtains

$$10 \quad \overline{w'\theta'_l} + LW_{net} = \frac{z - h_b}{h_t - h_b} (\overline{w'\theta'_l} + LW_{net})|_{z=h_t} + \frac{h_t - z}{h_t - h_b} (\overline{w'\theta'_l} + LW_{net})|_{z=h_b}, \quad (\text{C3})$$

$$\overline{w'q'_t} = \frac{z - h_b}{h_t - h_b} (\overline{w'q'_t})|_{z=h_t} + \frac{h_t - z}{h_t - h_b} (\overline{w'q'_t})|_{z=h_b}. \quad (\text{C4})$$

The above equations allow one to diagnose the profiles of  $\overline{w'\theta'_l}$  from the observed profiles of  $LW_{net}$ . However, we are more interested in diagnosing the vertical profiles of  $\overline{w'\theta'}$  which can then be directly compared to the observed ones. Such diagnostic relations for  $\overline{w'\theta'}$  are given further following Deardorff (1976).

Following the definition of  $\theta_l$ , in the cloud layer, the turbulent flux  $\overline{w'\theta'}$  is

$$5 \quad \overline{w'\theta'} = \overline{w'\theta'_l} + \frac{\theta}{T} \frac{L}{c_p} \overline{w'q'_c} . \quad (C5)$$

In saturated air

$$\overline{w'q'_v} = \overline{w'q'_s} = \frac{L}{R_v T \theta} q_s \overline{w'\theta'} , \quad (C6)$$

where  $q_s$  is saturation specific humidity and  $R_v$  is the gas constant for water vapor. Since  $\overline{w'q'_c} = \overline{w'q'_t} - \overline{w'q'_v}$  we can rewrite Equation C5 as

$$10 \quad \overline{w'\theta'} = \left[ \overline{w'\theta'_l} + \left( \frac{\theta}{T} \frac{L}{c_p} \overline{w'q'_t} \right) \right] \left[ 1 + \frac{L^2}{c_p R_v T^2} q_s \right]^{-1} . \quad (C7)$$

Below the cloud base we simply have  $\overline{w'\theta'} = \overline{w'\theta'_l}$ .

It is also useful to obtain a diagnostic relation for the virtual potential temperature flux  $\overline{w'\theta'_v}$  which is associated with the buoyancy flux. If we neglect the small contribution of the liquid cloud water  $q_c$  to air density, the virtual potential flux is

$$\overline{w'\theta'_v} = \overline{w'\theta'} + \theta (1 + 0.61 \overline{w'q'_v}) . \quad (C8)$$

15 In the cloud layer we can use C6 in C8 to obtain

$$\overline{w'\theta'_v} = \overline{w'\theta'} \left[ 1 + 0.61 q_s \left( 1 + \frac{L}{R_v T} \right) \right] , \quad (C9)$$

while below the cloud layer

$$\overline{w'\theta'_v} = \overline{w'\theta'_l} + \theta (1 + 0.61 \overline{w'q'_t}) . \quad (C10)$$

The turbulent entrainment fluxes  $\overline{w'\theta'_l}|_{z=h_t}$  and  $\overline{w'q'_t}|_{z=h_t}$  across the boundary-layer top are parameterized as

$$20 \quad \overline{w'\theta'_l}|_{z=h_t} = -w_e \Delta \theta_l , \quad (C11)$$

$$\overline{w'q'_t}|_{z=h_t} = -w_e \Delta q_t , \quad (C12)$$

where  $\Delta \theta_l = \Delta \theta - L/c_p \Delta q_c$  and  $\Delta q_t = \Delta q_v + \Delta q_c$  are the jumps of the conservative variables across the mixed-layer top. For the entrainment velocity, we use one of the most simple parameterizations, namely

$$w_e = A \frac{\theta_0}{g} \frac{w_*^3}{\Delta \theta_v \Delta h} , \quad (C13)$$

25 where  $A = 0.2$  is the typically used value of the proportionality constant. In Equation C13, we assume  $\Delta \theta_v \approx \Delta \theta$ .

In the used parameterization of entrainment, similar to Randall (1980) we assume that all of the observed radiative flux divergence occurs in the turbulent cloudy layer.

Finally, following Deardorff (1980); Nicholls (1989) we can use the diagnosed profile of the buoyancy flux to obtain the vertical velocity and temperature scales  $w_*$  and  $\theta_*$ , respectively, as

$$5 \quad w_* = (2.5I)^{1/3}, \quad (\text{C14})$$

$$\theta_* = w_*^2 \theta_v / gh, \quad (\text{C15})$$

where

$$I = (g/\theta_0) \int_{h_{bot}}^{h_{top}} \overline{w'\theta'_v} dz, \quad (\text{C16})$$

Furthermore, Equation C14 can be used to estimate the expected values of  $\sigma_w$  in the mixed-layer. Namely, Lenschow et al.

10 (1980) suggested the following parameterization for  $\sigma_w/w_*$  as function of  $z$ , namely

$$\frac{\sigma_w^2}{w_*^2} = c_1 \left[ \frac{z_{top} - z}{z_{top} - z_{bot}} \right]^{2/3} \left[ 1 - c_2 \frac{z_{top} - z}{z_{top} - z_{bot}} \right]^2, \quad (\text{C17})$$

where  $z_{top}$  and  $z_{bot}$  are the mixed-layer top and bottom heights, respectively;  $c_1 = 1.8$  and  $c_2 = 0.8$  are the empirical constants. Equation C17 is valid for  $z_{bot} \leq z < z_{top}$ .

*Author contributions.* DC, CL, AE and MW conceptualized the study and worked out the methodology. All of the authors took part in the  
 15 ACLOUD campaign in 2017 to obtain the observational data used in the study. JH carried out the initial data processing and the five-hole probe calibration. DC and CL processed the data to obtain turbulence statistics and carried out the analysis of the obtained results. DC formulated and applied the mixed-layer model.

*Competing interests.* None of the authors have any competing interests

*Data availability.* The 1Hz-averaged noseboom data is available in Hartmann et al. (2019a). The full-resolution noseboom data at 100Hz is  
 20 available in Hartmann et al. (2019b). The 1Hz-averaged liquid and total water content from the Nevzorov data is available in Chechin (2019). The vertical profiles of the net terrestrial radiation are available in Stapf et al. (2019). MODIS images are available at <https://ladsweb.modaps.eosdis.nasa.gov/>. ERA5 reanalysis data is available at <https://climate.copernicus.eu/>.

*Acknowledgements.* We gratefully acknowledge the funding of the German Research Foundation DFG (Deutsche Forschungsgemeinschaft) of the Transregional Collaborative Research Center SFB/TRR 172 (Project-ID 268020496). Part of the analysis of the turbulence statistics by

D. Chechin was funded by the Russian Science Foundation grant 18-77-10072-P and data processing by D. Chechin was funded by Russian Ministry of Science and Higher Education, agreement No. 075-15-2019-1621. DC thanks Johannes Stapf for processing observations of radiative fluxes, for the initial script for plotting MODIS images as well as for fruitful discussions.

## References

- Albrecht, B. A., Cox, S. K., and Schubert, W. H.: Radiometric measurements of in-cloud temperature fluctuations, *Journal of Applied Meteorology and Climatology*, 18, 1066–1071, 1979.
- Aliabadi, A. A., Staebler, R. M., Liu, M., and Herber, A.: Characterization and Parametrization of Reynolds Stress and Turbulent Heat Flux in the Stably-Stratified Lower Arctic Troposphere Using Aircraft Measurements, *Boundary-Layer Meteorology*, 161, 99–126, <https://doi.org/10.1007/s10546-016-0164-7>, <https://doi.org/10.1007/s10546-016-0164-7>, 2016.
- Brooks, I. M., Tjernström, M., Persson, P. O. G., Shupe, M. D., Atkinson, R. A., Canut, G., Birch, C. E., Mauritsen, T., Sedlar, J., and Brooks, B. J.: The Turbulent Structure of the Arctic Summer Boundary Layer During The Arctic Summer Cloud-Ocean Study, *Journal of Geophysical Research: Atmospheres*, 122, 9685–9704, <https://doi.org/https://doi.org/10.1002/2017JD027234>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2017JD027234>, 2017.
- Brümmer, B., Busack, B., Hoerber, H., and Kruspe, G.: Boundary-layer observations over water and Arctic sea-ice during on-ice air flow, *Boundary-layer meteorology*, 68, 75–108, 1994.
- Caughey, S. J., Crease, B. A., and Roach, W. T.: A field study of nocturnal stratocumulus II Turbulence structure and entrainment, *Quarterly Journal of the Royal Meteorological Society*, 108, 125–144, <https://doi.org/https://doi.org/10.1002/qj.49710845508>, <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.49710845508>, 1982.
- Chechin, D.: Liquid water content measured by the Nevzorov probe during the aircraft ALOUD campaign in the Arctic, <https://doi.org/10.1594/PANGAEA.906658>, <https://doi.org/10.1594/PANGAEA.906658>, 2019.
- Chechin, D. G. and Lüpkes, C.: Boundary-Layer Development and Low-level Baroclinicity during High-Latitude Cold-Air Outbreaks: A Simple Model, *Boundary-Layer Meteorology*, 162, 91–116, <https://doi.org/10.1007/s10546-016-0193-2>, <https://doi.org/10.1007/s10546-016-0193-2>, 2017.
- Curry, J. A.: Interactions among turbulence, radiation and microphysics in Arctic stratus clouds, *Journal of the atmospheric sciences*, 43, 90–106, 1986.
- Curry, J. A., Ebert, E. E., and Herman, G. F.: Mean and turbulence structure of the summertime Arctic cloudy boundary layer, *Quarterly Journal of the Royal Meteorological Society*, 114, 715–746, <https://doi.org/https://doi.org/10.1002/qj.49711448109>, <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.49711448109>, 1988.
- Curry, J. A., Schramm, J. L., Rossow, W. B., and Randall, D.: Overview of Arctic Cloud and Radiation Characteristics, *Journal of Climate*, 9, 1731–1764, [https://doi.org/10.1175/1520-0442\(1996\)009<1731:OOACAR>2.0.CO;2](https://doi.org/10.1175/1520-0442(1996)009<1731:OOACAR>2.0.CO;2), [https://doi.org/10.1175/1520-0442\(1996\)009<1731:OOACAR>2.0.CO;2](https://doi.org/10.1175/1520-0442(1996)009<1731:OOACAR>2.0.CO;2), 1996.
- Deardorff, J. W.: Usefulness of Liquid-Water Potential Temperature in a Shallow-Cloud Model, *Journal of Applied Meteorology*, 15, 98–102, [https://doi.org/10.1175/1520-0450\(1976\)015<0098:UOLWPT>2.0.CO;2](https://doi.org/10.1175/1520-0450(1976)015<0098:UOLWPT>2.0.CO;2), [https://doi.org/10.1175/1520-0450\(1976\)015<0098:UOLWPT>2.0.CO;2](https://doi.org/10.1175/1520-0450(1976)015<0098:UOLWPT>2.0.CO;2), 1976.
- Deardorff, J. W.: Stratocumulus-capped mixed layers derived from a three-dimensional model, *Boundary-Layer Meteorology*, 18, 495–527, 1980.
- Duynkerke, P. G., de Roode, S. R., van Zanten, M. C., Calvo, J., Cuxart, J., Cheinet, S., Chlond, A., Grenier, H., Jonker, P. J., Köhler, M., et al.: Observations and numerical simulations of the diurnal cycle of the EUROCS stratocumulus case, *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 130, 3269–3296, 2004.

- Egerer, U., Gottschalk, M., Siebert, H., Ehrlich, A., and Wendisch, M.: The new BELUGA setup for collocated turbulence and radiation measurements using a tethered balloon: first applications in the cloudy Arctic boundary layer, *Atmospheric Measurement Techniques*, 12, 4019–4038, <https://doi.org/10.5194/amt-12-4019-2019>, <https://www.atmos-meas-tech.net/12/4019/2019/>, 2019.
- Egerer, U., Ehrlich, A., Gottschalk, M., Griesche, H., Neggers, R. A., Siebert, H., and Wendisch, M.: Case study of a humidity layer above Arctic stratocumulus and potential turbulent coupling with the cloud top, *Atmospheric Chemistry and Physics*, 21, 6347–6364, 2021.
- Ehrlich, A., Wendisch, M., Lüpkes, C., Buschmann, M., Bozem, H., Chechin, D., Clemen, H.-C., Dupuy, R., Eppers, O., Hartmann, J., et al.: A comprehensive in situ and remote sensing data set from the Arctic CLOUD Observations Using airborne measurements during polar Day (ACLOUD) campaign, *Earth System Science Data*, 11, 1853–1881, 2019.
- Fedorovich, E. and Conzemius, R.: Effects of wind shear on the atmospheric convective boundary layer structure and evolution, *Acta Geophysica*, 56, 114, 2008.
- Finger, J. E. and Wendling, P.: Turbulence structure of Arctic stratus clouds derived from measurements and calculations, *Journal of Atmospheric Sciences*, 47, 1351–1373, 1990.
- Friehe, C., Shaw, W., Rogers, D., Davidson, K., Large, W., Stage, S., Crescenti, G., Khalsa, S., Greenhut, G., and Li, F.: Air-sea fluxes and surface layer turbulence around a sea surface temperature front, *Journal of Geophysical Research: Oceans*, 96, 8593–8609, 1991.
- Golder, D.: Relations among stability parameters in the surface layer, *Boundary-Layer Meteorology*, 3, 47–58, 1972.
- Goosse, H., Kay, J. E., Armour, K. C., Bodas-Salcedo, A., Chepfer, H., Docquier, D., Jonko, A., Kushner, P. J., Lecomte, O., Massonnet, F., et al.: Quantifying climate feedbacks in polar regions, *Nature communications*, 9, 1–13, 2018.
- Graversen, R. G., Mauritsen, T., Tjernström, M., Källén, E., and Svensson, G.: Vertical structure of recent Arctic warming, *Nature*, 451, 53–56, 2008.
- Gryanik, V. M. and Hartmann, J.: A Turbulence Closure for the Convective Boundary Layer Based on a Two-Scale Mass-Flux Approach, *Journal of the Atmospheric Sciences*, 59, 2729 – 2744, [https://doi.org/10.1175/1520-0469\(2002\)059<2729:ATCFTC>2.0.CO;2](https://doi.org/10.1175/1520-0469(2002)059<2729:ATCFTC>2.0.CO;2), [https://journals.ametsoc.org/view/journals/atsc/59/18/1520-0469\\_2002\\_059\\_2729\\_atcftc\\_2.0.co\\_2.xml](https://journals.ametsoc.org/view/journals/atsc/59/18/1520-0469_2002_059_2729_atcftc_2.0.co_2.xml), 2002.
- Hartmann, J., Gehrman, M., Kohnert, K., Metzger, S., and Sachs, T.: New calibration procedures for airborne turbulence measurements and accuracy of the methane fluxes during the AirMeth campaigns, *Atmospheric Measurement Techniques*, 11, 4567–4581, <https://doi.org/10.5194/amt-11-4567-2018>, <https://www.atmos-meas-tech.net/11/4567/2018/>, 2018.
- Hartmann, J., Lüpkes, C., and Chechin, D.: 1Hz resolution aircraft measurements of wind and temperature during the ACLOUD campaign in 2017, <https://doi.org/10.1594/PANGAEA.902849>, <https://doi.org/10.1594/PANGAEA.902849>, 2019a.
- Hartmann, J., Lüpkes, C., and Chechin, D.: High resolution aircraft measurements of wind and temperature during the ACLOUD campaign in 2017, <https://doi.org/10.1594/PANGAEA.900880>, <https://doi.org/10.1594/PANGAEA.900880>, 2019b.
- Heinze, R., Mironov, D., and Raasch, S.: Second-moment budgets in cloud topped boundary layers: A large-eddy simulation study, *Journal of Advances in Modeling Earth Systems*, 7, 510–536, 2015.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., et al.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, 2020.
- Hogan, R. J., Grant, A. L., Illingworth, A. J., Pearson, G. N., and O’Connor, E. J.: Vertical velocity variance and skewness in clear and cloud-topped boundary layers as revealed by Doppler lidar, *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 135, 635–643, 2009.
- Inoue, J., Kosović, B., and Curry, J. A.: Evolution of a storm-driven cloudy boundary layer in the Arctic, *Boundary-layer meteorology*, 117, 213–230, 2005.



- Knudsen, E. M., Heinold, B., Dahlke, S., Bozem, H., Crewell, S., Gorodetskaya, I. V., Heygster, G., Kunkel, D., Maturilli, M., Mech, M., Viceto, C., Rinke, A., Schmithüsen, H., Ehrlich, A., Macke, A., Lüpkes, C., and Wendisch, M.: Meteorological conditions during the ACLOUD/PASCAL field campaign near Svalbard in early summer 2017, *Atmospheric Chemistry and Physics*, 18, 17995–18022, <https://doi.org/10.5194/acp-18-17995-2018>, <https://www.atmos-chem-phys.net/18/17995/2018/>, 2018.
- 5 Korolev, A. V., Strapp, J. W., Isaac, G. A., and Nevzorov, A. N.: The Nevzorov Airborne Hot-Wire LWC–TWC Probe: Principle of Operation and Performance Characteristics, *Journal of Atmospheric and Oceanic Technology*, 15, 1495–1510, [https://doi.org/10.1175/1520-0426\(1998\)015<1495:TNAHWL>2.0.CO;2](https://doi.org/10.1175/1520-0426(1998)015<1495:TNAHWL>2.0.CO;2), [https://doi.org/10.1175/1520-0426\(1998\)015<1495:TNAHWL>2.0.CO;2](https://doi.org/10.1175/1520-0426(1998)015<1495:TNAHWL>2.0.CO;2), 1998.
- Lemone, M. A.: On the difficulty of measuring temperature and humidity in cloud: Comments on "Shallow convection on day 261 of GATE: Mesoscale arcs", *Monthly Weather Review*, 108, 1702–1705, 1980.
- 10 Lenschow, D., Wyngaard, J. C., and Pennell, W. T.: Mean-field and second-moment budgets in a baroclinic, convective boundary layer, *Journal of Atmospheric Sciences*, 37, 1313–1326, 1980.
- Lenschow, D., Society, A. M., and for Atmospheric Research (U.S.), N. C.: Probing the atmospheric boundary layer, American Meteorological Society, <https://books.google.de/books?id=EC9RAAAAMAAJ>, 1986.
- Lenschow, D. H. and Pennell, W. T.: On the Measurement of In-Cloud and Wet-Bulb Temperatures from an Aircraft, *Monthly Weather Review*, 102, 447 – 454, [https://doi.org/10.1175/1520-0493\(1974\)102<0447:OTMOIC>2.0.CO;2](https://doi.org/10.1175/1520-0493(1974)102<0447:OTMOIC>2.0.CO;2), [https://journals.ametsoc.org/view/journals/mwre/102/6/1520-0493\\_1974\\_102\\_0447\\_otmoic\\_2\\_0\\_co\\_2.xml](https://journals.ametsoc.org/view/journals/mwre/102/6/1520-0493_1974_102_0447_otmoic_2_0_co_2.xml), 1974.
- 15 Lenschow, D. H. and Stankov, B. B.: Length scales in the convective boundary layer, *Journal of Atmospheric Sciences*, 43, 1198–1209, 1986.
- Lenschow, D. H., Li, X. S., Zhu, C. J., and Stankov, B. B.: The stably stratified boundary layer over the great plains, *Boundary-Layer Meteorology*, 42, 95–121, <https://doi.org/10.1007/BF00119877>, <https://doi.org/10.1007/BF00119877>, 1988.
- 20 Lenschow, D. H., Mann, J., and Kristensen, L.: How Long Is Long Enough When Measuring Fluxes and Other Turbulence Statistics?, *Journal of Atmospheric and Oceanic Technology*, 11, 661 – 673, [https://doi.org/10.1175/1520-0426\(1994\)011<0661:HLILEW>2.0.CO;2](https://doi.org/10.1175/1520-0426(1994)011<0661:HLILEW>2.0.CO;2), [https://journals.ametsoc.org/view/journals/atot/11/3/1520-0426\\_1994\\_011\\_0661\\_hlilew\\_2\\_0\\_co\\_2.xml](https://journals.ametsoc.org/view/journals/atot/11/3/1520-0426_1994_011_0661_hlilew_2_0_co_2.xml), 1994.
- Lilly, D. K.: Models of cloud-topped mixed layers under a strong inversion, *Quarterly Journal of the Royal Meteorological Society*, 94, 292–309, 1968.
- 25 Mahrt, L.: Vertical Structure and Turbulence in the Very Stable Boundary Layer, *Journal of Atmospheric Sciences*, 42, 2333 – 2349, [https://doi.org/10.1175/1520-0469\(1985\)042<2333:VSATIT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1985)042<2333:VSATIT>2.0.CO;2), [https://journals.ametsoc.org/view/journals/atoc/42/22/1520-0469\\_1985\\_042\\_2333\\_vsatit\\_2\\_0\\_co\\_2.xml](https://journals.ametsoc.org/view/journals/atoc/42/22/1520-0469_1985_042_2333_vsatit_2_0_co_2.xml), 1985.
- Mohan, M. and Siddiqui, T.: Analysis of various schemes for the estimation of atmospheric stability classification, *Atmospheric Environment*, 32, 3775–3781, 1998.
- 30 Morrison, H., de Boer, G., Feingold, G., Harrington, J., Shupe, M. D., and Sulia, K.: Resilience of persistent Arctic mixed-phase clouds, *Nature Geoscience*, 5, 11, <https://doi.org/10.1038/ngeo1332>, 2011.
- Nicholls, S.: The dynamics of stratocumulus: Aircraft observations and comparisons with a mixed layer model, *Quarterly Journal of the Royal Meteorological Society*, 110, 783–820, 1984.
- Nicholls, S.: The structure of radiatively driven convection in stratocumulus, *Quarterly Journal of the Royal Meteorological Society*, 115, 487–511, 1989.
- 35 Nicholls, S. and Leighton, J.: An observational study of the structure of stratiform cloud sheets: Part I. Structure, *Quarterly Journal of the Royal Meteorological Society*, 112, 431–460, 1986.
- Osborne, E., Richter-Menge, J., and Jeffries, M.: Arctic report card 2018, Washington, DC: National Park Service, 2018.

- Overland, J. E., Wood, K. R., and Wang, M.: Warm Arctic<sup>2014</sup>cold continents: climate impacts of the newly open Arctic Sea, *Polar Research*, 30, 15 787, 2011.
- Petty, G. W.: Sampling error in aircraft flux measurements based on a high-resolution large eddy simulation of the marine boundary layer, *Atmospheric Measurement Techniques*, 14, 1959–1976, 2021.
- 5 Pithan, F. and Mauritsen, T.: Arctic amplification dominated by temperature feedbacks in contemporary climate models, *Nature Geoscience*, 7, 181–184, <https://doi.org/10.1038/ngeo2071>, <https://doi.org/10.1038/ngeo2071>, 2014.
- Pithan, F., Medeiros, B., and Mauritsen, T.: Mixed-phase clouds cause climate model biases in Arctic wintertime temperature inversions, *Climate Dynamics*, 43, 289–303, 2014.
- Randall, D. A.: Entrainment into a Stratocumulus Layer with Distributed Radiative Cooling, *Journal of Atmospheric Sciences*, 37, 148 – 159, [https://doi.org/10.1175/1520-0469\(1980\)037<0148:EIASLW>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<0148:EIASLW>2.0.CO;2), [https://journals.ametsoc.org/view/journals/atsc/37/1/1520-0469\\_1980\\_037\\_0148\\_eiaslw\\_2\\_0\\_co\\_2.xml](https://journals.ametsoc.org/view/journals/atsc/37/1/1520-0469_1980_037_0148_eiaslw_2_0_co_2.xml), 1980.
- 10 Sedlar, J. and Shupe, M. D.: Characteristic nature of vertical motions observed in Arctic mixed-phase stratocumulus, *Atmospheric Chemistry and Physics*, 14, 3461–3478, <https://doi.org/10.5194/acp-14-3461-2014>, <https://www.atmos-chem-phys.net/14/3461/2014/>, 2014.
- Serreze, M. C. and Francis, J. A.: The Arctic amplification debate, *Climatic change*, 76, 241–264, 2006.
- 15 Shupe, M. D., Walden, V. P., Eloranta, E., Uttal, T., Campbell, J. R., Starkweather, S. M., and Shiobara, M.: Clouds at Arctic Atmospheric Observatories. Part I: Occurrence and Macrophysical Properties, *Journal of Applied Meteorology and Climatology*, 50, 626–644, <https://doi.org/10.1175/2010JAMC2467.1>, <https://doi.org/10.1175/2010JAMC2467.1>, 2011.
- Shupe, M. D., Persson, P. O. G., Brooks, I. M., Tjernström, M., Sedlar, J., Mauritsen, T., Sjogren, S., and Leck, C.: Cloud and boundary layer interactions over the Arctic sea ice in late summer, *Atmospheric Chemistry and Physics*, 13, 9379–9399, <https://doi.org/10.5194/acp-13-9379-2013>, <https://www.atmos-chem-phys.net/13/9379/2013/>, 2013.
- 20 Smedman, A.-S. and Hoegstroem, U.: Turbulent characteristics of a shallow convective internal boundary layer, *Boundary-Layer Meteorology*, 25, 271–287, 1983.
- Sotiropoulou, G., Sedlar, J., Tjernström, M., Shupe, M. D., Brooks, I. M., and Persson, P. O. G.: The thermodynamic structure of summer Arctic stratocumulus and the dynamic coupling to the surface, *Atmospheric Chemistry and Physics*, 14, 12 573–12 592, <https://doi.org/10.5194/acp-14-12573-2014>, <https://www.atmos-chem-phys.net/14/12573/2014/>, 2014.
- 25 Stapf, J., Ehrlich, A., Jäkel, E., and Wendisch, M.: Cloud radiative forcing, LWP and cloud-free albedo derived from airborne broadband irradiance observations during the ACLOUD campaign, <https://doi.org/10.1594/PANGAEA.909289>, <https://doi.org/10.1594/PANGAEA.909289>, 2019.
- Stapf, J., Ehrlich, A., Jäkel, E., Lüpkes, C., and Wendisch, M.: Reassessment of shortwave surface cloud radiative forcing in the Arctic: consideration of surface-albedo–cloud interactions, *Atmospheric Chemistry and Physics*, 20, 9895–9914, 2020.
- Stevens, B.: Entrainment in stratocumulus-topped mixed layers, *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 128, 2663–2690, 2002.
- Stevens, B., Moeng, C.-H., Ackerman, A. S., Bretherton, C. S., Chlond, A., de Roode, S., Edwards, J., Golaz, J.-C., Jiang, H., Khairoutdinov, M., et al.: Evaluation of large-eddy simulations via observations of nocturnal marine stratocumulus, *Monthly weather review*, 133, 1443–1462, 2005.
- 35 Tetzlaff, A., Lüpkes, C., and Hartmann, J.: Aircraft-based observations of atmospheric boundary-layer modification over Arctic leads, *Quarterly Journal of the Royal Meteorological Society*, 141, 2839–2856, <https://doi.org/https://doi.org/10.1002/qj.2568>, <https://rmetsonlineibrary.wiley.com/doi/abs/10.1002/qj.2568>, 2015.

- Tjernström, M.: Turbulence Length Scales in Stably Stratified Free Shear Flow Analyzed from Slant Aircraft Profiles, *Journal of Applied Meteorology and Climatology*, 32, 948 – 963, [https://doi.org/10.1175/1520-0450\(1993\)032<0948:TLSISS>2.0.CO;2](https://doi.org/10.1175/1520-0450(1993)032<0948:TLSISS>2.0.CO;2), [https://journals.ametsoc.org/view/journals/apme/32/5/1520-0450\\_1993\\_032\\_0948\\_tlsiss\\_2\\_0\\_co\\_2.xml](https://journals.ametsoc.org/view/journals/apme/32/5/1520-0450_1993_032_0948_tlsiss_2_0_co_2.xml), 1993.
- 5 Tjernström, M., Žagar, M., Svensson, G., Cassano, J. J., Pfeifer, S., Rinke, A., Wyser, K., Dethloff, K., Jones, C., Semmler, T., and Shaw, M.: ‘Modelling the Arctic Boundary Layer: An Evaluation of Six Arcmip Regional-Scale Models using Data from the Sheba Project’, *Boundary-Layer Meteorology*, 117, 337–381, <https://doi.org/10.1007/s10546-004-7954-z>, <https://doi.org/10.1007/s10546-004-7954-z>, 2005.
- Tjernström, M., Sedlar, J., and Shupe, M. D.: How well do regional climate models reproduce radiation and clouds in the Arctic? An evaluation of ARCMIP simulations, *Journal of Applied Meteorology and Climatology*, 47, 2405–2422, 2008.
- 10 Tjernström, M., Leck, C., Birch, C. E., Bottenheim, J. W., Brooks, B. J., Brooks, I. M., Bäcklin, L., Chang, R. Y.-W., de Leeuw, G., Di Liberto, L., de la Rosa, S., Granath, E., Graus, M., Hansel, A., Heintzenberg, J., Held, A., Hind, A., Johnston, P., Knulst, J., Martin, M., Matrai, P. A., Mauritsen, T., Müller, M., Norris, S. J., Orellana, M. V., Orsini, D. A., Paatero, J., Persson, P. O. G., Gao, Q., Rauschenberg, C., Ristovski, Z., Sedlar, J., Shupe, M. D., Sierau, B., Sirevaag, A., Sjogren, S., Stetzer, O., Swietlicki, E., Szczodrak, M., Vaattovaara, P., Wahlberg, N., Westberg, M., and Wheeler, C. R.: The Arctic Summer Cloud Ocean Study (ASCOS): overview and experimental design, *Atmospheric Chemistry and Physics*, 14, 2823–2869, <https://doi.org/10.5194/acp-14-2823-2014>, <https://www.atmos-chem-phys.net/14/2823/2014/>, 2014.
- 15 Turton, J. and Nicholls, S.: A study of the diurnal variation of stratocumulus using a multiple mixed layer model, *Quarterly Journal of the Royal Meteorological Society*, 113, 969–1009, 1987.
- Wendisch, M., Brückner, M., Burrows, J., Crewell, S., Dethloff, K., Ebell, K., Lüpkes, C., Macke, A., Notholt, J., Quaas, J., et al.: Arctic Amplification: climate relevant atmospheric and Surface processes, and feedback mechanisms:(AC) 3, *Eos, Trans. Amer. Geophys. Union*, 98, 2017.
- 20 Wendisch, M., Macke, A., Ehrlich, A., Lüpkes, C., Mech, M., Chechin, D., Dethloff, K., Velasco, C. B., Bozem, H., Brückner, M., Clemen, H.-C., Crewell, S., Donth, T., Dupuy, R., Ebell, K., Egerer, U., Engelmann, R., Engler, C., Eppers, O., Gehrman, M., Gong, X., Gottschalk, M., Gourbeyre, C., Griesche, H., Hartmann, J., Hartmann, M., Heinold, B., Herber, A., Herrmann, H., Heygster, G., Hoor, P., Jafariserajehlou, S., Jäkel, E., Järvinen, E., Jourdan, O., Kästner, U., Kecorius, S., Knudsen, E. M., Köllner, F., Kretzschmar, J., Lelli, L., Leroy, D., Maturilli, M., Mei, L., Mertes, S., Mioche, G., Neuber, R., Nicolaus, M., Nomokonova, T., Notholt, J., Palm, M., van Pinxteren, M., Quaas, J., Richter, P., Ruiz-Donoso, E., Schäfer, M., Schmieder, K., Schnaiter, M., Schneider, J., Schwarzenböck, A., Seifert, P., Shupe, M. D., Siebert, H., Spreen, G., Stapf, J., Stratmann, F., Vogl, T., Welti, A., Wex, H., Wiedensohler, A., Zanatta, M., and Zeppenfeld, S.: The Arctic Cloud Puzzle: Using ACLOUD/PASCAL Multiplatform Observations to Unravel the Role of Clouds and Aerosol Particles in Arctic Amplification, *Bulletin of the American Meteorological Society*, 100, 841–871, <https://doi.org/10.1175/BAMS-D-18-0072.1>, <https://doi.org/10.1175/BAMS-D-18-0072.1>, 2019.
- 30 Wesche, C., Steinhage, D., and Nixdorf, U.: Polar aircraft Polar 5 and POLar6 operated by the Alfred Wegener Institute. *J. Large-Scale Res. Facil.*, 2, A87, 2016.
- Young, G. S.: Turbulence structure of the convective boundary layer. Part I. Variability of normalized turbulence statistics, *Journal of Atmospheric Sciences*, 45, 719–726, 1988.
- 35 Zilitinkevich, S., Gryanik, V. M., Lykossov, V. N., and Mironov, D. V.: Third-Order Transport and Nonlocal Turbulence Closures for Convective Boundary Layers, *Journal of the Atmospheric Sciences*, 56, 3463 – 3477, <https://doi.org/10.1175/1520->

0469(1999)056<3463:TOTANT>2.0.CO;2, [https://journals.ametsoc.org/view/journals/atasc/56/19/1520-0469\\_1999\\_056\\_3463\\_totant\\_2.0.co\\_2.xml](https://journals.ametsoc.org/view/journals/atasc/56/19/1520-0469_1999_056_3463_totant_2.0.co_2.xml), 1999.