

# Composition and Reactivity of Volatile Organic Compounds in the South Coast Air Basin and San Joaquin Valley of California

Shang Liu<sup>1</sup>, Barbara Barletta<sup>2</sup>, Rebecca S. Hornbrook<sup>3</sup>, Alan Fried<sup>4</sup>, Jeff Peischl<sup>5,6</sup>, Simone Meinardi<sup>2</sup>, Matthew Coggon<sup>5,6</sup>, Aaron Lamplugh<sup>5,6</sup>, Jessica B. Gilman<sup>6</sup>, Georgios I. Gkatzelis<sup>5,6,7</sup>,  
5 Carsten Warneke<sup>6</sup>, Eric C. Apel<sup>3</sup>, Alan J. Hills<sup>3</sup>, Ilann Bourgeois<sup>5,6</sup>, James Walega<sup>4</sup>, Petter Weibring<sup>4</sup>,  
Dirk Richter<sup>4</sup>, Toshihiro Kuwayama<sup>1</sup>, Michael FitzGibbon<sup>1</sup>, Donald Blake<sup>2</sup>

<sup>1</sup>Research Division, California Air Resources Board, Sacramento, 95814, USA

<sup>2</sup>Department of Chemistry, University of California, Irvine, 92697, USA

10 <sup>3</sup>Atmospheric Chemistry Observations & Modeling Laboratory, National Center for Atmospheric Research, Boulder, 80301, USA

<sup>4</sup>Institute of Arctic & Alpine Research, University of Colorado, Boulder, 80303 USA

<sup>5</sup>Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder 80309, USA

<sup>6</sup>NOAA Chemical Science Laboratory, Boulder, 80305, USA

15 <sup>7</sup>Now at: Institute of Energy and Climate Research, IEK-8: Troposphere, Forschungszentrum Jülich GmbH, Jülich, 52428, Germany

*Correspondence to:* Shang Liu (shang.liu@arb.ca.gov)

**Abstract.** Comprehensive aircraft measurements of volatile organic compounds (VOCs) covering the South Coast Air Basin (SoCAB) and San Joaquin Valley (SJV) of California were obtained in the summer of 2019. Combined with the CO, CH<sub>4</sub>,  
20 and NO<sub>x</sub> data, the total calculated gas-phase hydroxyl radical reactivity (cOHR<sub>TOTAL</sub>) was quantified to be 6.1 s<sup>-1</sup> and 4.6 s<sup>-1</sup>  
for the SoCAB and SJV, respectively. VOCs accounted for ~60%–70% of the cOHR<sub>TOTAL</sub> in both basins. In particular, oxygenated VOCs (OVOCs) contributed > 60% of the cOHR of total VOCs (cOHR<sub>VOC</sub>) as well as the total observed VOC mixing ratio. Primary biogenic VOCs (BVOCs) represented a minor fraction (< 2%) of the total VOC mixing ratio but accounted for 21% and 6% of the cOHR<sub>VOC</sub> in the SoCAB and SJV, respectively. Furthermore, the contribution of BVOCs  
25 to the cOHR<sub>VOC</sub> increased with increasing cOHR<sub>VOC</sub> in the SoCAB, suggesting that BVOCs were important ozone precursors during high ozone episodes. Spatially, the trace gases were heterogeneously distributed in the SoCAB with their mixing ratios and cOHR significantly greater over the inland regions than the coast, while their levels were more evenly distributed in the SJV. The results highlight that a better grasp of the emission rates and sources of OVOCs and BVOCs is essential for a predictive understanding of the ozone abundance and distribution in California.

## 30 1 Introduction

Ambient ozone (O<sub>3</sub>) is a criteria pollutant that forms from complex photochemical reactions of volatile organic compounds (VOCs) and oxides of nitrogen (NO<sub>x</sub> = NO + NO<sub>2</sub>) in the presence of sunlight. Exposure to O<sub>3</sub> can lead to adverse health problems such as airway inflammation and decreased lung function, especially in susceptible populations such as children

(Nuvolone et al., 2017; Kampa and Castanas, 2008). Persistent NO<sub>x</sub> emission control measures in California have resulted in the substantially reduced ambient O<sub>3</sub> levels, e.g., the ozone design value (ODV) decreased by over 60% in California's South Coast Air Basin (SoCAB) from 273 parts per billion (ppb) in 1980 to 102 ppb in 2015 (Parrish et al., 2017). However, the SoCAB and San Joaquin Valley (SJV) continue to exceed the health-based, 8-hour National Ambient Air Quality Standard (NAAQS) of 70 ppb particularly during the warm summer months (Faloona et al., 2020; Parrish et al., 2017). The plateauing of O<sub>3</sub> levels since 2010 (Wu et al., 2022) poses additional challenges for California's air pollution reduction pathways.

40

Further reducing ambient O<sub>3</sub> may be more difficult than expected due to the nonlinear O<sub>3</sub> production rate in response to decreases of NO<sub>x</sub> and VOCs and the relatively high O<sub>3</sub> background. For example, Parrish et al. (2017) showed that the observational-based background ODV for southern California was 62±1.9 ppb, which was comparable to the modeled background ODV of 45–65 ppb with an outlier of ~92 ppb. This background ODV has decreased very slowly at ~1 ppb decade<sup>-1</sup> since the mid-2000s (Parrish et al., 2021). Compared to NO<sub>x</sub>, VOCs are more complex in their emission sources, composition, and reactivity. VOCs are emitted to the atmosphere from a variety of anthropogenic, pyrogenic, and biogenic sources (Goldstein and Galbally, 2007). After emission, VOCs undergo complex photochemical reactions resulting in the production of O<sub>3</sub>. VOCs are comprised of hundreds of molecules with different functionalities. The composition of VOCs in California has been rapidly changing over the past years (Warneke et al., 2012), due in large part to the aggressive emission control measures implemented for on-road mobile sources. This emission source of VOCs is expected to decrease in California over the next decades, driven by the Governor's Executive Order (N-79-20) with a goal of 100% Zero Emission Vehicle sales of new passenger cars and light-duty trucks by 2035. In contrast, the relative contributions of industrial activities and consumer products to ambient VOCs are growing (McDonald et al., 2018; Kim et al., 2022). With a clear path for NO<sub>x</sub> reduction in the state, it is critical that the current sources of VOCs and their potential for O<sub>3</sub> formation are better characterized to guide future policies that effectively reduce ambient O<sub>3</sub>.

55

In this study, we analyzed an extensive set of VOCs and other key trace gases measured from the National Aeronautics and Space Administration (NASA) DC-8 research aircraft that flew over the SoCAB and SJV in the summer of 2019. We characterized the composition and the source signatures of the VOCs in these two basins. By examining the calculated hydroxyl radical (OH) reactivity of VOCs (cOHR<sub>VOC</sub>), we further determined the key VOC species that are responsible for O<sub>3</sub> production. The spatial distributions of the cOHR<sub>VOC</sub> in the SoCAB and SJV are also investigated and compared.

60

## 2 Methods and data processing


The study was conducted in partnership with the Fire Influence on Regional to Global Environments and Air Quality (FIREX-AQ) campaign in the summer of 2019 (Warneke et al., in preparation). We used data from the two California flights (Fig. S1), which were designed to study the spatial distribution of VOCs from non-fire emissions over the SoCAB and SJV in California. The measurements were carried out during ~11:00–18:00 (local time) on board the NASA DC-8 research

65

aircraft on 22 July and 5 September 2019, both of which were weekdays. The flights covered the SJV by flying in a raster pattern. In the SoCAB, the measurements were made over the northern and inland regions of the SoCAB, where the O<sub>3</sub> peak events usually occur (Cai et al., 2019). The flights had fewer measurements in the coastal areas and no measurements over central Los Angeles due to flight restrictions (Fig. 1).

To demonstrate the representativeness of the two flights, we synthesized data from the surface monitoring network ([https://aqs.epa.gov/aqsweb/airdata/download\\_files.html](https://aqs.epa.gov/aqsweb/airdata/download_files.html)) in the SoCAB and SJV for the summer of 2019. We then compared the diurnal profiles of the air pollutants and meteorological parameters between the two measurement days with the days during July–September, 2019. Examples are shown for Los Angeles County in the SoCAB and Kern County in the SJV (Fig. S2). The O<sub>3</sub> and air temperature profiles from the two measurement days overlapped with those from July–September. These comparisons show that the measurement days are representative of typical summer days in the SoCAB and SJV. It is expected that the summer period features high O<sub>3</sub> pollution as well as high BVOC emissions and secondary OVOC productions.

80



85 **Figure 1:** Spatial distribution of (a) O<sub>3</sub>, (b) NO<sub>x</sub>, (c) VOCs, and (d) cOHR<sub>voc</sub>. The flight tracks and sample locations are represented by colored lines and points, respectively. The points are sized by flight altitude and colored by the mixing ratios of O<sub>3</sub>, NO<sub>x</sub>, VOCs, and cOHR<sub>voc</sub> values in (a), (b), (c), and (d), respectively, with the scales shown in the legend. The background map (grey) shows the boundaries of the California air basins, with black indicating the SoCAB and SJV.

**Table 1.** Summary of measurements.

#	Technique	Species Measured	Sample Duration/ Frequency	Institution	Investigator	Reference
1	Chemiluminescence (CL)	NO, NO <sub>2</sub> , O <sub>3</sub> , NO <sub>y</sub>	1 s	NOAA CSL	Thomas Ryerson	Bourgeois et al. (2022); (Bourgeois et al., 2021)
2	Laser Absorption Spectroscopy (LAS)	CO, CH <sub>4</sub>	1 s	NASA LaRC	Glenn Diskin	Sachse et al. (1987)
3	Whole Air Sampling (WAS)	C <sub>2</sub> -C <sub>10</sub> Alkanes, C <sub>2</sub> -C <sub>4</sub> Alkenes, C <sub>6</sub> -C <sub>9</sub> Aromatics, C <sub>1</sub> -C <sub>5</sub> Alkyl nitrates, etc.	20–100 s, average 40 s, variable frequency	UC Irvine	Donald Blake	Simpson et al. (2020)
4	HR-ToF-GC/MS	C <sub>3</sub> -C <sub>10</sub> hydrocarbons, C <sub>1</sub> -C <sub>7</sub> OVOCs, HCN, CH <sub>3</sub> CN, C <sub>1</sub> -C <sub>2</sub> halocarbons, etc.	33 s every 1.75 min	NCAR ACOM	Eric Apel	Apel et al. (2015)
5	Whole Air Sampling (WAS)	C <sub>2</sub> -C <sub>10</sub> Alkanes, C <sub>2</sub> -C <sub>4</sub> Alkenes, C <sub>6</sub> -C <sub>9</sub> Aromatics, C <sub>1</sub> -C <sub>5</sub> Alkyl nitrates, etc.	5 s sample collection every 10-1600 seconds	NOAA CSL	Jessica Gilman	Lerner et al. (2017)
6	Laser Absorption Spectroscopy (LAS)	HCHO	1 s	CU Boulder	Alan Fried	Richter et al. (2015), Fried et al. (2020)
7	H <sub>3</sub> O <sup>+</sup> ToF-CIMS	Speciated hydrocarbons and OVOCs	1 s	NOAA CSL	Carsten Warneke	Yuan et al. (2017a)
8	CIMS	PAN, PPN, other PANs	1 s	Georgia Tech	Gregory Huey	Zheng et al. (2011)
9	CIMS	Phenol	1 s	CalTech	Paul Wennberg	Crouse et al. (2006)
10	Iodide ToF-CIMS	Formic acid	1 s	NOAA CSL	Patrick Veres	Veres et al. (2020)

- 90 Here we present analyses of the VOCs, CO, NO<sub>x</sub>, and O<sub>3</sub> measured by eight research groups operating instrumentation on board the DC-8 (Table 1). The technical details of the instruments are described in the Supplementary Materials. Since the diverse techniques have different sampling frequencies, the data were merged to the sampling time (~40 s in duration) of the whole air samples collected by the University of California Irvine (UCI-WAS). The merged data set was created using the data merge tool provided in the FIREX-AQ data archive (<https://www-air.larc.nasa.gov/cgi-bin/ArcView/firexaq>).
- 95 Specifically, the measurements from data set #1 to data set #10 in Table 1 were sequentially merged. The duplicated measurements were removed during each merging step, i.e., if the measurement of a compound existed in the merged data set and was above detection limit, additional measurements of that compound was not added to the merged data set during the following merging steps.

100 Fig. S3 shows the comparison of the study-average mixing ratios and standard deviations calculated from the merged data set (a subset of the original data sets) and the original data sets for the species merged to the UCI-WAS measurements. Each data set was pre-processed such that only the measurements covering the SoCAB and SJV were used for the comparison. The agreement (slope of 1.07 and  $r^2$  of 1.00) between the merged data set and the original data set suggests that the merged data set can represent the original measurements.

105

To better represent the near-surface mixing ratios, the samples collected at altitudes higher than 4,000 ft (1.2 km) above ground level (a.g.l) were removed from the merged data set. The samples collected over the ocean were also excluded. After screening, the final data set for this study has 69 and 95 samples for the SoCAB and SJV, respectively. These samples are shown in Fig. 1. A total of 168 gas-phase compounds were measured, of which 137 had detectable mixing ratio (Table 2).

110 Given that the aircraft speed was about  $125 \text{ m s}^{-1}$  and the sample collection time was about 40 s, the measurements represented approximately 5-km integrated space along the flight track. The average altitude ( $\pm$  standard deviation) of the samples was  $625 (\pm 237) \text{ m a.g.l.}$  and  $360 (\pm 107) \text{ m a.g.l.}$  in the SoCAB and SJV, respectively. Observations have shown that the daily maximum planetary boundary layer (PBL) height was approximately 1–1.5 km in the SoCAB and 0.5–1 km in the SJV (Cui et al., 2019; Bianco et al., 2011). Therefore the samples were most likely collected within the PBL. Since the

115 samples were collected at least a few hundred meters away from emission sources, the reactive VOCs likely have undergone photochemical processing. Using the  $\text{NO}_x$  and  $\text{NO}_y$  measurements and assuming that the ambient  $[\text{OH}]$  was  $4 \times 10^6 \text{ molecules cm}^{-3}$ , the photochemical age of the samples was estimated to be approximately 12 hours on average (Cappa et al., 2012). This means a species with an OH rate coefficient of  $10^{-11} \text{ cm}^3 \text{ molecules}^{-1} \text{ s}^{-1}$  had lost ~80% of its mass before they were sampled by the aircraft. Compounds with higher rate coefficients (e.g., the BVOCs) had more loss. The statistics of the

120 observed mixing ratios of the gas-phase species and their OH rate coefficients are summarized in Table 2.

### 3 Results and discussion

#### 3.1 Mixing ratio and spatial distribution of $\text{O}_3$ , VOCs, and $\text{NO}_x$

The study-average aircraft-measured  $\text{O}_3$  ( $\pm 1 \sigma$ ) mixing ratio was  $70.2 \pm 26.7 \text{ ppb}$  in the SoCAB and  $73.5 \pm 9.5 \text{ ppb}$  in the SJV. The maximum observed  $\text{O}_3$  mixing ratio was  $122.0 \text{ ppb}$  and  $101.0 \text{ ppb}$  in the SoCAB and SJV, respectively. The average

125 non- $\text{CH}_4$  VOC mixing ratio ( $\pm 1 \sigma$ ) in the SoCAB and SJV was  $36.9.0 \pm 7.3 \text{ ppb}$  and  $45.8 \pm 5.6 \text{ ppb}$ , respectively (for simplicity, we use VOC to represent non- $\text{CH}_4$  VOC hereafter). The higher VOC levels in the SJV were mainly driven by methanol. The average observed methanol was  $17.1 \text{ ppb}$  in the SJV, which doubled the average  $7.7 \text{ ppb}$  observed in the SoCAB. The average aircraft-measured  $\text{NO}_x$  ( $\pm 1 \sigma$ ) was  $3.8 \pm 4.8 \text{ ppb}$  in the SoCAB, which was 3.5 times the  $\text{NO}_x$  ( $\pm 1 \sigma$ ) of  $1.1 \pm 0.6 \text{ ppb}$  in the SJV. The high  $\text{NO}_x$  levels in the SoCAB resulted in a relatively lower VOC-to- $\text{NO}_x$  ratio of  $9.7 \pm 12.4$

130 compared to  $41.6 \pm 23.3$  in the SJV.

The spatial distributions of O<sub>3</sub>, VOCs, and NO<sub>x</sub> are shown in Fig. 1 (a-c). In the SoCAB, the mixing ratios of O<sub>3</sub>, VOCs, and NO<sub>x</sub> generally increased from coastal to inland regions, with the highest mixing ratios occurring near the northern boundary of the SoCAB. The coast-inland gradient likely resulted from the accumulation and chemical aging of the air pollutants as they were transported eastwards. In the SJV, the mixing ratios of O<sub>3</sub>, VOCs, and NO<sub>x</sub> were more homogeneously distributed, i.e., the occurrence frequency of the high mixing ratios was smaller compared to that in the SoCAB (Figs. S4-S6). Since the very high mixing ratios determine the design values (Parrish et al., 2017), the result suggests that O<sub>3</sub> in the SoCAB is more prone to exceed the NAAQS. This is consistent with the higher O<sub>3</sub> design values in the SoCAB compared to the SJV resulted from analyses using the monitoring network data (Parrish et al., 2017).

140

**Table 2.** Statistics of the mixing ratios (median, average, 1- $\sigma$  standard deviation, and maximum values in ppt unless otherwise specified) and OH reaction rate coefficients of the measured VOCs in the SoCAB and the SJV.

Compound <sup>a</sup>	SoCAB				SJV				$k_{OH}^c$	Meas. Tech. <sup>d</sup>
	Med. <sup>b</sup>	Avg. <sup>b</sup>	S.D. <sup>b</sup>	Max. <sup>b</sup>	Med. <sup>b</sup>	Avg. <sup>b</sup>	S.D. <sup>b</sup>	Max. <sup>b</sup>		
Alkanes (excluding CH <sub>4</sub> )										
Ethane	2408	2675	2041	7678	1025	1282	586	4052	0.25	WAS
Propane	1041	1458	1323	4927	421	620	429	2096	1.1	WAS
<i>n</i> -Butane	395	415	321	1247	72	124	130	777	2.36	WAS
Isobutane	259	259	189	719	45	78	80	378	2.1	WAS
<i>n</i> -Pentane	248	220	167	603	30	54	68	554	6.911	WAS
Isopentane	530	508	409	1565	65	103	113	811	3.6	WAS
Cyclopentane	26	22	14	56	6	9	8	37	5.0	WAS
<i>n</i> -Hexane	73	74	58	242	12	18	15	73	5.2	WAS
2-Methylpentane	118	110	80	292	15	23	19	93	5.2	WAS
3-Methylpentane	67	64	50	182	8	13	12	61	5.2	WAS
2,2-Dimethylbutane	29	28	19	75	5	7	4	23	2.2	WAS
2,3-Dimethylbutane	40	39	29	116	7	9	6	28	5.8	WAS
Methylcyclopentane	69	62	47	189	12	16	13	64	5.7	WAS
Cyclohexane	22	25	17	72	6	9	7	32	7.0	WAS
<i>n</i> -Heptane	27	30	22	83	7	11	8	37	6.8	WAS
2-Methylhexane	32	33	24	95	6	8	5	26	6.9	WAS
3-Methylhexane	34	36	27	104	7	9	6	27	7.2	WAS
2,3-Dimethylpentane	43	44	32	124	6	7	4	19	6.5	WAS
2,4-Dimethylpentane	47	46	27	103	6	8	6	20	4.8	WAS (NOAA)
Methylcyclohexane	19	21	14	58	4	8	7	32	9.6	WAS
<i>n</i> -Octane	13	15	10	46	6	6	2	11	8.1	WAS
2,2,4-Trimethylpentane	55	65	50	196	6	8	6	39	3.3	WAS
2,3,4-Trimethylpentane	21	21	15	61	6	7	4	17	6.6	WAS

<i>n</i> -Nonane	9	10	6	34	4	5	1	9	9.7	WAS
<i>n</i> -Decane	7	9	6	29	4	5	1	7	11	WAS
<i>n</i> -Undecane	4	6	3	16	4	4	1	8	12	WAS
Alkenes										
Ethene	266	312	303	1181	47	63	14	248	8.5	WAS
Propene	38	45	44	181	12	14	8	59	26	WAS
Propadiene	6	8	4	16	NA	NA	NA	NA	9.8	WAS
1-Butene	6	8	6	25	4	4	1	7	31	WAS
Isobutene	9	11	7	23	9	10	4	15	51	WAS
1,3-Butadiene	5	6	2	8	7	7	1	7	67	WAS
<i>trans</i> -2-Butene	1	1	1	5	1	2	4	15	64	WAS (NOAA)
1-Pentene	4	5	2	8	3	4	1	6	31	WAS
<i>trans</i> -2-Pentene	4	5	2	8	NA	NA	NA	NA	67	WAS
<i>Cis</i> -2-Pentene	1	1	1	2	0	1	1	2	65	WAS (NOAA)
2-Methyl-1-butene	5	5	2	8	5	6	3	10	61	WAS
3-Methyl-1-butene	4	4	1	5	NA	NA	NA	NA	32	WAS
2-Methyl-2-butene	4	4	1	4	3	3	0	3	87	WAS
1,3-Pentadiene	NA	NA	NA	NA	4	4	1	4	101	WAS
1-Hexene	5	6	1	8	6	9	8	23	37	WAS
1-Heptene	10	11	8	32	5	7	4	14	40	WAS
1-Octene	NA	NA	NA	NA	7	7	6	11	30	WAS
1-Nonene	3	3	0	4	NA	NA	NA	NA	42	WAS
Aromatics										
Benzene	74	82	65	222	21	25	13	76	1.2	WAS
Toluene	137	160	141	551	17	24	21	131	5.6	WAS
Ethylbenzene	25	26	18	65	5	7	4	17	7.0	WAS
m/p-Xylene	42	49	47	173	7	9	7	33	19	WAS
o-Xylene	22	24	19	79	5	6	3	15	14	WAS
Styrene	6	8	7	29	4	4	1	4	58	WAS
Isopropylbenzene	4	4	1	6	NA	NA	NA	NA	6.3	WAS
<i>n</i> -Propylbenzene	5	6	2	11	3	4	1	5	5.8	WAS
2-Ethyltoluene	4	6	3	13	4	4	1	5	12	WAS
3-Ethyltoluene	6	8	7	24	3	4	1	5	19	WAS
4-Ethyltoluene	5	6	4	17	4	4	1	7	12	WAS
1,3,5-Trimethylbenzene	4	4	1	5	NA	NA	NA	NA	57	WAS
1,2,4-Trimethylbenzene	6	9	8	27	5	5	2	8	33	WAS
Naphthalene	10	10	6	30	9	9	3	17	23	H <sub>3</sub> O <sup>+</sup> ToF- CIMS
BVOCs and Related Oxidation Products										



Methacrolein (MAC)	126	196	193	692	31	41	33	156	29	WAS
Methyl vinyl ketone (MVK)	166	303	317	1292	120	134	88	474	20	WAS
Isoprene	156	178	178	651	14	36	55	298	100	WAS
$\alpha$ -Pinene	6	8	5	20	6	6	3	11	53	WAS
$\beta$ -Pinene	5	4	1	6	NA	NA	NA	NA	74	WAS
OVOCs										
Methanol	7734	7703	3858	15980	16349	17062	4350	29719	0.94	HR-ToF-GC/MS
Formaldehyde	4042	4475	3021	9503	3703	4239	1507	10067	9.4	LAS (CU Boulder)
Formic acid	5444	4344	2861	8506	9837	9702	1872	13211	0.45	CIMS (NOAA)
Carbonyl sulfide	616	611	38	678	611	615	26	702	0.0020	WAS
Nitromethane	40	61	63	360	81	105	76	422	0.016	WAS
Methyl nitrate	9	10	4	17	10	10	1	14	0.023	WAS
Ethyl nitrate	4	4	2	9	3	4	1	6	0.18	WAS
Isocyanic acid	238	354	269	1151	210	337	290	1104	0.0012	CIMS (NOAA)
Ethanol	793	1804	2500	10031	2590	2933	2020	11019	3.2	WAS
Acetaldehyde	1182	1307	914	3229	1229	1516	797	4410	15	HR-ToF-GC/MS
Methyl formate	151	148	52	226	172	177	24	209	0.22	HR-ToF-GC/MS
PAN	974	914	812	3220	546	668	393	2463	0.042	CIMS (Georgia Tech)
Isopropanol	104	153	187	854	55	60	28	187	5.1	WAS
Propanal	144	157	105	434	122	138	54	317	20	HR-ToF-GC/MS
Acrolein	44	58	61	434	50	53	31	186	20	WAS
Acetone	4285	4415	2655	9953	2896	3176	760	5649	0.17	HR-ToF-GC/MS
<i>n</i> -Propyl nitrate	1	2	1	5	1	1	0	2	0.58	WAS
Isopropyl nitrate	11	12	9	35	7	8	3	17	0.29	WAS
PPN	100	96	95	475	49	62	40	215	0.46	CIMS (Georgia Tech)
APAN	17	28	31	131	20	21	10	58	16	CIMS (Georgia Tech)
Carbon suboxide	6	6	2	9	5	5	1	7	2.6	HR-ToF-GC/MS
Methyl acetate	68	95	84	404	74	78	22	139	0.26	WAS
Butanal	37	40	25	145	36	42	21	120	24	WAS
Isobutanal	18	21	12	58	18	19	8	58	26	WAS

Crotonaldehyde	13	15	9	34	15	15	2	21	36	HR-ToF-GC/MS
Methyl ethyl ketone (MEK)	185	250	199	898	140	186	116	545	1.1	WAS
2,3-Butanedione	16	16	9	44	24	24	6	44	0.25	HR-ToF-GC/MS
Furan	NA	NA	NA	NA	8	8	7	13	40	WAS
2-Furanone	39	36	20	74	75	70	25	111	45	PTR-MS
Tetrahydrofuran	5	6	4	19	1	2	1	3	18	HR-ToF-GC/MS
Ethyl acetate	23	25	19	85	20	33	37	235	1.63	HR-ToF-GC/MS
Methyl propionate	2	2	1	3	2	2	1	5	0.88	HR-ToF-GC/MS
Maleic anhydride	40	37	24	82	18	23	10	50	1.5	H <sub>3</sub> O <sup>+</sup> ToF-CIMS
Isobutyl nitrate	19	21	11	40	13	12	5	22	1.5	WAS and HR-ToF-GC/MS
2-Butyl nitrate	10	12	10	38	5	6	3	22	0.86	WAS
PBN	25	28	23	103	11	14	10	72	4.7	CIMS (Georgia Tech)
2-Pentyl nitrate	3	4	3	12	1	2	1	7	1.7	WAS
3-Pentyl nitrate	2	2	2	7	1	1	1	4	1.0	WAS
3-Methyl-2-butyl nitrate	4	4	4	17	1	2	1	6	1.7	WAS
Phenol	6	6	4	16	4	7	6	21	27	CIMS (CalTech)
Catechol/5-Methylfurfural	45	38	23	68	37	37	12	60	78	H <sub>3</sub> O <sup>+</sup> ToF-CIMS
Guaiacol	14	15	11	32	19	19	5	25	54	H <sub>3</sub> O <sup>+</sup> ToF-CIMS
Benzaldehyde	30	29	21	72	27	30	17	80	12	H <sub>3</sub> O <sup>+</sup> ToF-CIMS
Creosol	12	11	7	25	10	11	3	17	75	H <sub>3</sub> O <sup>+</sup> ToF-CIMS
Syringol	4	5	2	9	4	4	2	9	81	H <sub>3</sub> O <sup>+</sup> ToF-CIMS
Other VOCs										
Chloromethane	572	567	36	651	561	569	33	679	0.036	WAS
Dichloromethane	80	99	73	450	69	68	5	80	0.12	WAS
Chloroform	21	24	13	57	18	19	3	29	0.11	WAS
Tetrachloromethane	78	78	1	81	79	79	1	82	0.00	WAS
Bromomethane	7	8	2	15	11	12	9	89	0.029	WAS
Dibromomethane	1	1	0	1	1	1	0	1	0.11	WAS
Bromoform	2	2	1	5	1	1	0	1	0.15	WAS

Dibromochloromethane	3	4	3	12	1	1	0	2	0.22	WAS
Bromodichloromethane	2	2	2	6	1	1	0	2	1.2	WAS
Iodomethane	2	1	1	4	1	1	0	2	0.10	WAS
Methanethiol	9	9	6	21	11	10	3	18	33	HR-ToF-GC/MS
Hydrogen cyanide	209	246	85	457	195	212	42	321	0.03	CIMS (CalTech)
Ethyne	284	342	308	1185	87	104	47	318	0.75	WAS
Chloroethane	3	3	1	6	3	3	1	6	0.40	WAS
1,2-Dichloroethane	16	16	6	30	17	17	1	20	0.23	WAS
Methyl chloroform	2	2	0	3	2	2	0	2	0.0095	WAS
Trichloroethene	2	4	4	17	0	1	0	3	1.9	WAS
Tetrachloroethene	5	11	10	37	2	3	4	39	0.16	WAS
Dimethyl sulfide	4	5	3	16	5	8	10	63	7.0	WAS
Carbon disulfide	5	5	3	12	5	5	1	8	1.2	HR-ToF-GC/MS
Acetonitrile	169	201	76	438	160	166	23	240	0.022	HR-ToF-GC/MS
Propyne	13	14	7	35	5	5	1	8	0.94	WAS
Acrylonitrile	12	14	6	22	NA	NA	NA	NA	4.0	WAS
Propanenitrile	13	15	8	36	14	15	5	29	0.14	WAS
D5	21	18	13	40	5	5	3	12	2.1	HR-ToF-GC/MS
NO, NO <sub>2</sub> , CO, and CH <sub>4</sub>										
NO	229	747	1276	5921	138	154	75	426	10	CL
NO <sub>2</sub>	1425	2997	3661	17891	879	976	514	2691	12	CL
CO (ppb)	196	193	92	354	130	137	18	206	0.24	LAS (NASA)
CH <sub>4</sub> (ppb)	1946	1957	71	2091	2070	2158	239	3437	0.0064	LAS (NASA)

145 <sup>a</sup>The following species were measured but not reported as their mixing ratios were below detection limit: *cis*-2-butene, 1,2-butadiene, 1-buten-3-yne, 1,3-butadiyne, 1-butyne, 2-butyne, *cis*-2-pentene, 3-methyl-1-pentene/4-methyl-1-pentene, 1-decene, cyclopentene, ethynylbenzene, chlorobenzene, tricyclene, camphene, myrcene, limonene,  $\Delta$ 3-carene, 2-methylfuran, 3-methylfuran, benzofuran, 1,3-butadiyne, chloriodomethane, 2-methyl-3-buten-2-ol, methylacrylonitrile, pyrrole, 2-ethylfuran, dimethylfurans, vinylfuran, 3-furaldehyde, and 1,3-pentadiene.

The mixing ratios of carbon suboxide, 2,3-butanedione, tetrahydrofuran, ethyl acetate, methyl propionate, and methanethiol from the HR-ToF-GC/MS measurement was estimated using relative sensitivities. The sum of these species accounted for less than 0.2% of the total VOC mixing ratio in the SoCAB and SJV.

150 The following abbreviations are used: D5 represents decamethylcyclopentasiloxane, PAN represents peroxyacetyl nitrate, PPN represents peroxypropionyl nitrate, APAN represents peroxyacryloyl nitrate, and PBN represents peroxybutyryl nitrate.

<sup>b</sup>NA represents not applicable because the VOCs are below detection limit.

<sup>c</sup>The unit for  $k_{OH}$  is  $10^{-12}$  cm<sup>3</sup> molecule<sup>-1</sup> s<sup>-1</sup>.  $k_{OH}$  values are given at 298 K. References for the  $k_{OH}$  values are provided in the text.

<sup>d</sup>WAS refers to the UC Irvine WAS unless otherwise stated.


155 The H<sub>3</sub>O<sup>+</sup> ToF-CIMS measurements and the formic acid mixing ratio (from iodide ToF-CIMS) were only available during the flight on 5 September 2019.

Hydrogen cyanide was measured by the CalTech CIMS and HR-ToF-GC/MS. The high time resolution (1 Hz) CalTech CIMS measurement was used.

The mixing ratio of isobutyl nitrate was derived by the difference between the sum of isobutyl nitrate and 2-butyl nitrate (measured by the HR-ToF-GC/MS) and 2-butyl nitrate (measured by the UC Irvine WAS).

### 3.2 Composition and source signatures of VOCs

160 The measured VOCs are grouped into chemical families based on their characteristic functional groups or emission sources (for biogenic VOCs and their oxidation products). The average VOC composition from the two research flights was similar, so the study-average composition (i.e., average of all samples collected during the two flights) is presented (Fig. 2). The mean, median, and maximum mixing ratios for each VOC measured in this study are listed in Table 2. Oxygenated VOCs (OVOCs) and alkanes dominated the measured VOC mixing ratios, accounting for 91%–96% of the total measured VOC  
165 mixing ratio in both the SoCAB and SJV. This is comparable to the measurements conducted at Pasadena in 2010 during the California Research at the Nexus of Air Quality and Climate Change (CalNex) field study, where OVOCs and alkanes comprised ~80% of the measured reactive organic carbon (ROC) mass (Heald et al., 2020). They are also the two most abundant chemical families in other regions of the world, e.g., in Seoul, South Korea (Simpson et al., 2020; Kim et al., 2018) and Hong Kong (Ling et al., 2014). Specifically, OVOCs accounted for 74% and 91% of the total measured VOC mixing  
170 ratio in the SoCAB and SJV, respectively, and the contributions of alkanes were 17% and 5% in the SoCAB and SJV, respectively. In contrast, primary biogenic VOCs (BVOCs) are a minor fraction (< 2%) of the total measured VOCs mixing ratio in both basins.



**Figure 2:** The campaign-average VOC composition for the SoCAB and SJV.

Methanol was the most abundant OVOC in both basins, which was responsible for 28% and 41% of the OVOC mixing ratio in the SoCAB and SJV, respectively. Methanol in the SoCAB (7.7 ppb) was about half of the methanol levels in the SJV (17.1 ppb). Very high methanol (> 25 ppb) was observed over the southeast (e.g., Delano, Tulare, Visalia) regions of Fresno, where dense, confined animal farming operations (CAFOs) are located, suggesting direct methanol emissions from dairy operations. Similar to methanol, formic acid in the SoCAB (4.3 ppb) was about half of the formic acid in the SJV (9.7 ppb). Formic acid has been identified as a major fatty acid from dairy operations (Page et al., 2014; Mårtensson et al., 1999). Formic acid also increased with methanol ( $r^2 = 0.7$ ), suggesting that dairies may be a major source of formic acid in the SJV. The region with high methanol mixing ratio also features high ethanol levels, which likely resulted from dairy emissions as well. Studies have shown that methanol and ethanol are the dominant VOCs emitted from dairy cows and their wastes (Gentner et al., 2014; Shaw et al., 2007; Sun et al., 2008). In addition, ethanol is likely the most abundant VOC species emitted from dairy silages in the SJV (Malkina et al., 2011; Yuan et al., 2017b), while consumer product use has a sizeable contribution to its ambient mixing ratios in urban cores (McDonald et al., 2018; Coggon et al., 2021). Vehicle operations also emit ethanol into the atmosphere due to the increasing use of the E10 and E85 ethanol-gasoline blends (De Gouw et al., 2012; Gkatzelis et al., 2021). More detailed surface-level source apportionment methods must be employed to determine the specificity of its origin. The high ethanol levels in the SJV may lead to high mixing ratios of its oxidation product, e.g., acetaldehyde, which was 16% higher in the SJV than that in the SoCAB.

Formaldehyde (HCHO) was the second and third most abundant OVOC, with comparable mixing ratio (~4.5 ppb) in the SoCAB and SJV, respectively. HCHO contributed to 16% and 10% to the OVOC mixing ratio in the SoCAB and SJV, respectively. Primary sources of HCHO include automotive exhaust (Anderson et al., 1996), industrial emissions (Salthammer et al., 2010), biomass burning (Holzinger et al., 1999), and off-gassing from building materials (Gilbert et al., 2008; Liu et al., 2016). HCHO can also form from the photooxidation of VOCs, such as alkenes and BVOCs (Choi et al., 2010; Parrish et al., 2012). Given the contrast between anthropogenic emission sources in the SoCAB and SJV (e.g., the CO emissions in the SoCAB are 1.4 times the CO emissions in the SJV, estimated from California Air Resources Board's emission inventory), the similarity of HCHO mixing ratio between the two regions during the two flight days suggests that photooxidation of non-anthropogenic emissions (e.g., BVOCs) may be the dominant regional driver of HCHO, which is consistent with the findings from previous studies in North America (Lee et al., 1998; Palmer et al., 2003). The high methanol, ethanol, and formic acid mixing ratios in the SJV made the OVOC level in the SJV 52% higher compared to the OVOC level in the SoCAB. Acetone also had sizable contributions to the OVOC mixing ratio (Fig. 2).

The composition of alkanes in the SoCAB and SJV was similar, but the total alkane mixing ratio was 2.6 times higher in the SoCAB. Ethane and propane were the dominant species. Specifically, ethane accounted for ~40%–50% of the total alkane mixing ratio, followed by propane (~25%) in both regions. The linear regressions of propane versus ethane for the SoCAB

and SJV fell on the same line, with a Pearson's  $r$ -value of 0.96 and a slope of 0.6. Peischl et al. (2013) summarized the propane versus ethane slope values from different sources. The slope obtained from this study is consistent with the aircraft observation over the SoCAB in 2010 and fell within the slope range for local oil and gas well emissions (Peischl et al., 2013). In contrast, the propane-to-ethane ratios for pipeline quality natural gas are  $\sim 0.15$  as heavier alkanes are removed during processing (Wennberg et al., 2012; Peischl et al., 2013). On-road emission ratios of propane-to-ethane are also very low, but relatively minor as gasoline and diesel do not contain large amounts of these alkanes (Fraser et al., 1998). This result suggests that oil and gas production activities are likely the major source of propane and ethane in the SoCAB and SJV. Isopentane and  $n$ -pentane were tightly correlated in both basins but with different slopes. The isopentane/ $n$ -pentane ratio was 2.4 in the SoCAB. This value is consistent with ratios of 2.3–3.8 observed for regions highly impacted by vehicular emissions (Gilman et al., 2013). In contrast, the isopentane/ $n$ -pentane ratio was 1.8 for SJV, suggesting that the air mass was mixed with non-urban emissions.

The BVOCs accounted for  $< 2\%$  of the total VOC mixing ratio in both basins. In the BVOC category, isoprene is assumed to be only from biogenic emissions and anthropogenically-driven isoprene emissions are negligible in summer (Guenther et al., 2012; Reimann et al., 2000). The BVOCs were dominated by isoprene and its oxidation products methacrolein (MAC) and methyl vinyl ketone (MVK) (Wennberg et al., 2018), the sum of which accounted for  $\sim 95\%$  of the total observed BVOC mixing ratio. MVK was the most abundant BVOC species that contributed  $\sim 50\%$  of BVOC mixing ratio, followed by nearly equal contributions of isoprene and MAC. The monoterpenes (such as  $\alpha$ - and  $\beta$ -pinene) were minor, with their average mixing ratios smaller than 10 ppt.


### 3.3 Total OH reactivity

Quantifying the OH reactivity (cOHR) of reactive gases provides estimates of the potential roles of individual  $O_3$  precursors (Ling et al., 2014; Gilman et al., 2009; McDonald et al., 2018). The cOHR of a compound is calculated from the compound concentration multiplied by its reaction rate coefficient with the OH radical. The total cOHR is the sum of the cOHR of all reactants in the atmosphere, which is given by the following equation:

$$\begin{aligned} \text{cOHR}_{\text{TOTAL}} &= \text{cOHR}_{\text{CH}_4} + \text{cOHR}_{\text{CO}} + \text{cOHR}_{\text{NO}_x} + \text{cOHR}_{\text{VOC}} \\ &= k_{\text{OH}+\text{CH}_4}[\text{CH}_4] + k_{\text{OH}+\text{CO}}[\text{CO}] + k_{\text{OH}+\text{NO}_x}[\text{NO}_x] + \sum(k_{\text{OH}+\text{VOC}}[\text{VOC}]) \quad (1) \end{aligned}$$

where the subscript of cOHR indicates the species  $X$  used to calculate the cOHR,  $[X]$  is the mixing ratio of  $X$ , and  $k_{\text{OH}+X}$  is the reaction rate coefficient of  $X$  with the OH radical. The  $k_{\text{OH}+X}$  values were obtained from literature or kinetic databases (Atkinson and Arey, 2003; Atkinson et al., 2004; Atkinson et al., 2008; Kwok and Atkinson, 1995; Burkholder et al., 2015; Alton and Browne, 2020; Atkinson, 1986; Atkinson et al., 2006; Atkinson et al., 1997; Bierbach et al., 1995; Bierbach et al.,

1994; Borduas et al., 2016; Lauraguais et al., 2015; Roberts et al., 2003; Semadeni et al., 1995) (National Institute of Standards and Technology Chemical Kinetics Database) and are tabulated in Table 2. We used the rate constants at 298 K. This is reasonable as the average ambient air temperature ( $\pm$  standard deviation) of the samples was  $26.7(\pm 2.1)$  °C and  $31.0(\pm 1.6)$  °C in the SoCAB and SJV, respectively. There is likely missing cOHR from unmeasured VOCs due to the limitation of the quantification methods. Previous studies show that the fraction of the missing cOHR ranges from 5% to 80%, depending on the sampling location, time, and the number of measured VOC species (Yang et al., 2016; Hansen et al., 2021). With the extensive VOC measurements in this study, it is expected that the fraction of the missing cOHR is close to the lower end of the 5%–80% range. Nevertheless, the  $\text{cOHR}_{\text{VOC}}$  reported in this study should be interpreted as a minimum for the cOHR of the total VOCs.




**Figure 3:** The total measured cOHR for the SoCAB and SJV.

The  $\text{cOHR}_{\text{TOTAL}}$  was  $6.1 \text{ s}^{-1}$  and  $4.6 \text{ s}^{-1}$  for the SoCAB and SJV, respectively (Fig. 3). The  $\text{O}_3$  mixing ratio generally increased nonlinearly with increasing  $\text{cOHR}_{\text{VOC}}$ ,  $\text{cOHR}_{\text{CO}}$ ,  $\text{cOHR}_{\text{CH}_4}$ , and  $\text{cOHR}_{\text{NO}_x}$  in both basins, especially for  $\text{O}_3 > 40$  ppb (Fig. S7). The  $\text{cOHR}_{\text{TOTAL}}$  values observed in this study are generally less than half of the  $\text{cOHR}_{\text{TOTAL}}$  reported in Pasadena and Central California 10–20 years ago (Table S1). However, we note that the comparison may be biased as the previous studies are from ground measurements. The percentage contributions of  $\text{CH}_4$ , CO,  $\text{NO}_x$ , and VOCs to  $\text{cOHR}_{\text{TOTAL}}$  were similar between the SoCAB and SJV, with the VOCs accounting for  $\sim 60\%$  of the  $\text{cOHR}_{\text{TOTAL}}$  (Fig. 3). For reference, the global annual mean contribution of VOCs to  $\text{cOHR}_{\text{TOTAL}}$  is  $\sim 50\%$  from a recent review study (Heald and Kroll, 2020).

The large contribution of VOCs to  $\text{cOHR}_{\text{TOTAL}}$  is consistent with previous measurements in California (Table S1), suggesting that VOCs remain a high potential for  $\text{O}_3$  formation in the SoCAB and SJV. CO was the second most important species following VOCs, and was responsible for  $\sim 18\%$  of the  $\text{cOHR}_{\text{TOTAL}}$  in both basins. The contribution of  $\text{NO}_x$  to  $\text{cOHR}_{\text{TOTAL}}$  was 17% in the SoCAB and 6% in the SJV.

### 3.4 OH reactivity of individual VOCs

The  $\text{cOHR}$  of the individual species for the SoCAB and SJV generally follows similar order. Linear regression of the  $\text{cOHR}_{\text{SoCAB}}$  versus the  $\text{cOHR}_{\text{SJV}}$  for the individual species shows a slope of 1.3 and  $r^2$  of 0.8 (Fig. 4), confirming that the overall  $\text{cOHR}$  in the SoCAB was 30% higher than that in the SJV. The top five VOCs that have the largest  $\text{cOHR}$  in the SoCAB were HCHO, acetaldehyde, isoprene, methanol, and MVK in decreasing order. In the SJV, the top five most abundant VOCs in terms of  $\text{cOHR}$  were formaldehyde, acetaldehyde, methanol, ethanol, and formic acid. The sum of these species accounted for 63% and 73% of the calculated  $\text{cOHR}_{\text{VOC}}$  in the SoCAB and SJV, respectively. These results suggest that (i) aldehydes are the key compounds contributing to  $\text{cOHR}$  in both basins, (ii) biogenic VOCs are important for  $\text{cOHR}$  in the SoCAB, and (iii) dairy emissions are likely important for  $\text{cOHR}$  in the SJV. Compared to the  $\text{cOHR}$  of the VOCs measured in Pasadena in 2010 during the CalNex campaign (57 species measured in common), the majority of the species measured in the SoCAB (52 out of 57) in 2019 had smaller  $\text{cOHR}$  values, with a mean percentage decrease of 53%.



275


**Figure 4:** Scatter plot of  $\text{cOHR}_{\text{SoCAB}}$  versus  $\text{cOHR}_{\text{SJV}}$  for the individual species in (a) linear scale and (b) logarithmic scale. The red line represents linear regression using the orthogonal distance regression method.

The OVOCs category was the largest contributor to the  $\text{cOHR}_{\text{VOC}}$  among all the categories (Fig. 5; Table S2). The  $\text{cOHR}_{\text{OVOC}}$  was  $3.6 \text{ s}^{-1}$  in the SoCAB and  $3.1 \text{ s}^{-1}$  in the SJV, which was responsible for 59% and 68% of the  $\text{cOHR}_{\text{VOC}}$  in the



280 SoCAB and SJV, respectively. The contribution of the OVOCs to the  $cOHR_{VOC}$  was among the highest compared to the  
 $cOHR_{OVOC}$  measurements in other parts of the world (Table S2). For example, the percentage of OVOC contribution was  
substantially higher than that in mainland China (calculated from emission inventories), Seoul (South Korea, surface  
measurement), and Mexico City (Mexico, surface measurement) (Table S2). (We note that part of the difference may be due  
to the different study approaches, e.g., ground vs flight measurements.) HCHO was the most important OVOC in terms of  
285  $cOHR$ , which was responsible for 45% and 36% of the  $cOHR_{OVOC}$  in the SoCAB and SJV, respectively. This percentage  
translates to HCHO accounting for ~30% of the  $cOHR_{VOC}$  in both basins. Acetaldehyde contributed 21% of the  $cOHR_{OVOC}$   
and ~15% of the  $cOHR_{VOC}$  in both basins. Since ambient measurements of HCHO and acetaldehyde are scant compared to  
many other VOCs (e.g., aromatics), much remains to be learned on the sources and fate of these species. Methanol  
contributed 8% and 15% of the  $cOHR_{OVOC}$  in the SoCAB and SJV, respectively, while the ethanol contribution to the  
290  $cOHR_{OVOC}$  was ~7% in both basins. The sum of HCHO, acetaldehyde, methanol, and ethanol accounted for 51% and 69% of  
the  $cOHR_{VOC}$  in the SoCAB and SJV, respectively. Our measurements are consistent with the modeling results showing that  
30%–50% of  $cOHR_{VOC}$  in California is due to aldehydes and other oxygenated species from over a decade prior (Steiner et  
al., 2008).

295 Although BVOCs were only responsible for < 2% of the total measured VOC mixing ratio, they contributed 21% and 6% of  
the  $cOHR_{VOC}$  in the SoCAB and SJV, respectively. Therefore it is important that the reactivity of BVOCs is well represented  
in photochemical models, especially in the SoCAB. The relatively high  $cOHR_{BVOC}$  is due to the high chemical reactivity of  
the BVOCs, i.e., the reaction rate constants of BVOCs with OH radicals are generally 100 times greater than those of alkanes  
(Table 2). Among the BVOCs, isoprene accounted for 59% and 44% of the  $cOHR_{BVOC}$  in the SoCAB and SJV, respectively.  
300 The isoprene oxidation products MAC and MVK, together, accounted for ~45% of  $cOHR_{BVOC}$  in both regions. The  
contribution of monoterpenes to  $cOHR_{BVOC}$  was < 10% in both basins. This should be considered as a lower limit as a large  
fraction of monoterpenes may have reacted away by the time they were measured from the aircraft. The dominance of  
isoprene and its oxidation products in  $cOHR_{BVOC}$  is consistent with the high isoprene emissions from the oak woodlands  
throughout the foothills of the Sierra Nevada mountains and near the northern boundaries of the SoCAB (Fig. S8a) (Arey et  
305 al., 1995; Benjamin et al., 1997; Misztal et al., 2014).




**Figure 5:** The measured cOHR of VOCs in the SoCAB and SJV.

In contrast, the non-CH<sub>4</sub> alkanes only contributed 7% and 3% of the cOHR<sub>VOC</sub> in the SoCAB and SJV, respectively, despite  
 310 their greater atmospheric abundances. The sum of cOHR<sub>ethane</sub> and cOHR<sub>propane</sub> accounted for 21% and 31% of the cOHR<sub>alkanes</sub>  
 in the SoCAB and SJV, respectively. The substantially smaller contribution of alkanes to cOHR<sub>VOC</sub> compared to their  
 contribution to the total VOC mixing ratio is due to their low chemical reactivity. On the other hand, the longer-chain  
 alkanes dominated cOHR<sub>alkanes</sub>. Alkenes, aromatics, and other uncategorized VOCs (the “Other VOCs” in Table 2), together,  
 accounted for < 10% of the cOHR<sub>VOC</sub> in both basins.

### 315 3.5 Spatial variability of cOHR<sub>VOC</sub>

The cOHR<sub>VOC</sub> showed distinct frequency distributions in the SoCAB and SJV (Fig. 6). The cOHR<sub>VOC</sub> in the SoCAB spanned  
 from 0 to 8 s<sup>-1</sup>, with the highest occurrence frequency at 0–1 s<sup>-1</sup> and lowest at 7–8 s<sup>-1</sup>. Higher cOHR<sub>VOC</sub> occurred in more  
 inland regions (Fig. 1), which was similar to the spatial distribution of O<sub>3</sub>. This is likely due to the accumulation of air  
 pollutants in downwind locations as they were transported from west to east during the day in the SoCAB (Carreras-  
 320 Sospedra et al., 2006). In contrast, the cOHR<sub>VOC</sub> in the SJV was narrowly distributed and centered at 1–3 s<sup>-1</sup>. In addition, the

BVOCs in the SoCAB showed increasing contribution to the  $cOHR_{VOC}$  as the  $cOHR_{VOC}$  increased (the pie charts in Fig. 6a). The BVOCs contribution reached up to 20-30% when the  $cOHR_{VOC}$  was above  $4 \text{ s}^{-1}$  near the northern boundary of the SoCAB (Fig. 1, Fig. S8b). This result suggests that BVOCs are likely key  $O_3$  precursors during high  $O_3$  episodes in the SoCAB. A similar result was not observed in the SJV, i.e., the OVOCs dominated the  $cOHR_{VOC}$  across the  $cOHR_{VOC}$  range. The minor role of the BVOCs in  $O_3$  formation in the SJV was likely because of the daytime up-valley winds that prevented the entrainment of the BVOCs emitted over the Sierra Nevada mountains into the valley (Zhong et al., 2004).



**Figure 6:** Frequency distribution of  $cOHR_{VOC}$  in the (a) SoCAB and (b) SJV. Also shown above each  $1\text{-s}^{-1}$  binned  $cOHR_{VOC}$  range is a pie chart presenting the average contribution to the  $cOHR_{VOC}$  by VOC categories (shown in the legend).

330

#### 4 Conclusions and implications

In this work, we have updated the mixing ratios and  $cOHR$  of a wide range of speciated VOCs in the SoCAB and SJV of California. The limited number of flights is compensated by the wide range of speciated VOCs analyzed. The spatially resolved data set and analyses can be used to evaluate the performance of photochemical models. Our measurements suggest

335 that OVOCs are an important chemical class contributing to the cOHR in California. Specifically, HCHO, acetaldehyde,  
methanol, and ethanol are the key OVOC species, the sum of which accounted for ~50%–70% of the cOHR<sub>VOC</sub> in the  
SoCAB and SJV. The observation is consistent with previous field and modeling studies showing that OVOCs play critical  
roles in atmospheric chemistry (Lou et al., 2010; Wu et al., 2020). OVOCs have also been suggested to be responsible for a  
major fraction of the “missing OH reactivity” (Dolgorouky et al., 2012; Karl et al., 2009; Lou et al., 2010), i.e., the  
340 difference between directly-measured OHR<sub>TOTAL</sub> and the sum of the cOHR from individually-measured species. However,  
quantifying OVOCs is still challenging and measurements of their ambient mixing ratios are still lacking in many regions.  
Our work reinforces that elucidating the chemical composition of OVOCs and their sources will advance our capability to  
predict O<sub>3</sub> abundance and develop O<sub>3</sub> reduction strategies.

345 Biogenic emissions represent a significant source of VOCs for O<sub>3</sub> formation, especially in the SoCAB. On average 21% and  
6% of the cOHR<sub>VOC</sub> are contributed by primary BVOCs in the SoCAB and SJV, respectively, assuming that isoprene is  
solely from biogenic sources. The contribution of BVOCs to cOHR<sub>VOC</sub> was greatest near the northern edge of the SoCAB.  
This is consistent with the measurements in Pasadena, California, during the CalNex campaign, where isoprene was the  
single largest contributor to the cOHR<sub>VOC</sub> (Heald et al., 2020). The importance of BVOCs as a contributor to cOHR has also  
350 been reported for other urban environments, such as London, United Kingdom (Whalley et al., 2016), Beijing, China (Mo et  
al., 2018), and Seoul, South Korea (Kim et al., 2018). The cOHR<sub>BVOC</sub> reported in this study should be considered as an  
underestimation, since a fraction of HCHO may be produced from the oxidation of BVOCs (Choi et al., 2010). Source  
apportionment of HCHO on the regional scale is thus critical to improving our future understanding of the contribution of  
biogenic emissions to O<sub>3</sub> production. Besides reacting with the OH radical, HCHO is also a source of HO<sub>x</sub> radicals (HO<sub>x</sub> =  
355 HO + HO<sub>2</sub> + RO<sub>2</sub>) via photolysis. Reducing HCHO has the added benefit of reducing the production rate of HO<sub>x</sub>, thereby  
lowering the production rate of O<sub>3</sub> (Pusede and Cohen, 2012).

As policies continue to curb anthropogenic emissions, BVOCs will continue to become increasingly important in  
atmospheric chemistry and O<sub>3</sub> formation (Gu et al., 2021). The temperature rise due to a warming climate will likely further  
360 enhance the emissions of BVOCs in the immediate future. Since BVOCs are highly reactive, a small increase in BVOCs will  
disproportionately enhance their contribution to the total OHR. This effect makes BVOCs even more critical during peak O<sub>3</sub>  
events that may lead to O<sub>3</sub> exceedance. Our current knowledge of the role of BVOCs in the O<sub>3</sub> formation in the South Coast  
region is still highly uncertain, e.g., a recent photochemical modeling study suggests a significant underestimation of  
biogenic emissions in the SoCAB (Cai et al., 2019). As the contribution of BVOCs to O<sub>3</sub> production may represent a  
365 substantial fraction of the background O<sub>3</sub> that determines how stringent the O<sub>3</sub> standards will be, a more precise  
understanding of the magnitude and timing of BVOCs in O<sub>3</sub> formation is warranted. Such evaluation must be carefully  
paired with our evolving understanding of the biospheric feedback in response to changing climate to improve mitigation  
and adaptation actions as well as future air quality management planning.

## Data Availability

370 The data used in this study can be downloaded from the NASA FIREX-AQ data archive <https://www-air.larc.nasa.gov/cgi-bin/ArcView/firexaq>.

## Author contributions

BB, RSH, AF, JP, SM, MS, AL, JBG, GIG, CW, ECA, AJH, IB, JW, PW, DR, and DB conducted the aircraft measurements. SL analyzed the data and wrote the manuscript. BB, RSH, AF, JP, MC, JBG, GIG, CW, TK, and MF  
375 provided constructive comments and suggestions.

## Competing interests

The authors declare that they have no conflict of interest.

## Acknowledgements

The authors acknowledge the funding from California Air Resources Board (contract #RD19014) to support the aircraft  
380 measurements. The HR-ToF-GC/MS measurement was supported by the National Center for Atmospheric Research, which is a major facility sponsored by the National Science Foundation under Cooperative Agreement No. 1852977. E.C.A., R.S.H., and A.J.H. were also funded in part by NASA Award No. 80NSSC18K0633. We thank the FIREX-AQ team, the coordinators, NASA, and NOAA for making this data collection possible with the DC-8 research aircraft. We thank Thomas Ryerson for providing the O<sub>3</sub> and NO<sub>x</sub> data, Glenn Diskin for providing the CO and CH<sub>4</sub> data, Paul Wennberg, Lu Xu,  
385 Krystal Vasquez, Hannah Allen, and John Crouse for providing the phenol and hydrogen cyanide data, Patrick R. Veres and J. Andrew Neuman for providing the NOAA CIMS data, and Gregory Huey for providing the Georgia Tech CIMS data. We are grateful to Jeremy Avise and Chenxia Cai for their technical review and logistical coordination.

## Disclaimer

The statements and conclusions in this manuscript are those of the authors and do not represent the official views of the  
390 California Air Resources Board.

## References

Alton, M. W. and Browne, E. C.: Atmospheric chemistry of volatile methyl siloxanes: Kinetics and products of oxidation by OH radicals and Cl atoms, *Environmental Science & Technology*, 54, 5992-5999, 2020.

- 395 Anderson, L. G., Lanning, J. A., Barrell, R., Miyagishima, J., Jones, R. H., and Wolfe, P.: Sources and sinks of formaldehyde and acetaldehyde: An analysis of Denver's ambient concentration data, *Atmospheric Environment*, 30, 2113-2123, 1996.
- Apel, E., Hornbrook, R., Hills, A., Blake, N., Barth, M., Weinheimer, A., Cantrell, C., Rutledge, S., Basarab, B., and Crawford, J.: Upper tropospheric ozone production from lightning NO<sub>x</sub>-impacted convection: Smoke ingestion case study from the DC3 campaign, *Journal of Geophysical Research: Atmospheres*, 120, 2505-2523, 2015.
- 400 Arey, J., Crowley, D. E., Crowley, M., Resketo, M., and Lester, J.: Hydrocarbon emissions from natural vegetation in California's South Coast Air Basin, *Atmospheric Environment*, 29, 2977-2988, 1995.
- Atkinson, R.: Kinetics and mechanisms of the gas-phase reactions of the hydroxyl radical with organic compounds under atmospheric conditions, *Chemical Reviews*, 86, 69-201, 1986.
- 405 Atkinson, R. and Arey, J.: Atmospheric degradation of volatile organic compounds, *Chemical reviews*, 103, 4605-4638, 2003.
- Atkinson, R., Baulch, D., Cox, R., Hampson Jr, R., Kerr, J., Rossi, M., and Troe, J.: Evaluated kinetic, photochemical and heterogeneous data for atmospheric chemistry: Supplement V. IUPAC Subcommittee on Gas Kinetic Data Evaluation for Atmospheric Chemistry, *Journal of Physical and Chemical Reference Data*, 26, 521-1011, 1997.
- 410 Atkinson, R., Baulch, D., Cox, R., Crowley, J., Hampson, R., Hynes, R., Jenkin, M., Rossi, M., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume I-gas phase reactions of O<sub>x</sub>, HO<sub>x</sub>, NO<sub>x</sub> and SO<sub>x</sub> species, *Atmospheric chemistry and physics*, 4, 1461-1738, 2004.
- Atkinson, R., Baulch, D., Cox, R., Crowley, J., Hampson, R., Hynes, R., Jenkin, M., Rossi, M., Troe, J., and Subcommittee, I.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume II-gas phase reactions of organic species, *Atmospheric chemistry and physics*, 6, 3625-4055, 2006.
- 415 Atkinson, R., Baulch, D., Cox, R., Crowley, J., Hampson, R., Hynes, R., Jenkin, M., Rossi, M., Troe, J., and Wallington, T.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume IV-gas phase reactions of organic halogen species, *Atmospheric Chemistry and Physics*, 8, 4141-4496, 2008.
- Benjamin, M. T., Sudol, M., Vorsatz, D., and Winer, A. M.: A spatially and temporally resolved biogenic hydrocarbon emissions inventory for the California South Coast Air Basin, *Atmospheric Environment*, 31, 3087-3100, 1997.
- 420 Bianco, L., Djalalova, I., King, C., and Wilczak, J.: Diurnal evolution and annual variability of boundary-layer height and its correlation to other meteorological variables in California's Central Valley, *Boundary-layer meteorology*, 140, 491-511, 2011.
- Bierbach, A., Barnes, I., and Becker, K.: Product and kinetic study of the OH-initiated gas-phase oxidation of furan, 2-methylfuran and furanaldehydes at ≈300 K, *Atmospheric Environment*, 29, 2651-2660, 1995.
- 425 Bierbach, A., Barnes, I., Becker, K. H., and Wiesen, E.: Atmospheric chemistry of unsaturated carbonyls: Butenedial, 4-oxo-2-pentenal, 3-hexene-2, 5-dione, maleic anhydride, 3H-furan-2-one, and 5-methyl-3H-furan-2-one, *Environmental science & technology*, 28, 715-729, 1994.
- Borduas, N., Abbatt, J. P., Murphy, J. G., So, S., and da Silva, G.: Gas-phase mechanisms of the reactions of reduced organic nitrogen compounds with OH radicals, *Environmental science & technology*, 50, 11723-11734, 2016.

- 430 Bourgeois, I., Peischl, J., Neuman, J. A., Brown, S. S., Allen, H. M., Campuzano-Jost, P., Coggon, M. M., DiGangi, J. P., Diskin, G. S., and Gilman, J. B.: Comparison of airborne measurements of NO, NO<sub>2</sub>, HONO, NO<sub>y</sub> and CO during FIREX-AQ, *Atmospheric Measurement Techniques Discussions*, 1-47, 2022.
- Bourgeois, I., Peischl, J., Neuman, J. A., Brown, S. S., Thompson, C. R., Aikin, K. C., Allen, H. M., Angot, H., Apel, E. C., and Baublitz, C. B.: Large contribution of biomass burning emissions to ozone throughout the global remote troposphere, 435 *Proceedings of the National Academy of Sciences*, 118, 2021.
- Burkholder, J. B., Sander, S. P., Abbatt, J. P. D., Barker, J. R., Huie, R. E., Kolb, C. E., Kurylo, M. J., Orkin, V. L., Wilmouth, D. M., and Wine, P. H.: Chemical kinetics and photochemical data for use in atmospheric studies, *Evaluation Number 18*, JPL Publication, 15-10, 2015.
- 440 Cai, C., Avise, J., Kaduwela, A., DaMassa, J., Warneke, C., Gilman, J. B., Kuster, W., Gouw, J., Volkamer, R., Stevens, P., Lefer, B., Holloway, J. S., Pollack, I. B., Ryerson, T., Atlas, E., Blake, D., Rappenglueck, B., Brown, S. S., and Dube, W. P.: Simulating the weekly cycle of NO<sub>x</sub>-VOC-HO<sub>x</sub>-O<sub>3</sub> photochemical system in the South Coast of California during CalNex-2010 campaign, *Journal of Geophysical Research: Atmospheres*, 124, 3532-3555, 10.1029/2018jd029859, 2019.
- 445 Cappa, C. D., Onasch, T. B., Massoli, P., Worsnop, D. R., Bates, T. S., Cross, E. S., Davidovits, P., Hakala, J., Hayden, K. L., and Jobson, B. T.: Radiative absorption enhancements due to the mixing state of atmospheric black carbon, *Science*, 337, 1078-1081, 2012.
- Carreras-Sospedra, M., Dabdub, D., Rodriguez, M., and Brouwer, J.: Air quality modeling in the South Coast Air Basin of California: What do the numbers really mean?, *J. Air & Waste Manage. Assoc.*, 56, 1184-1195, 2006.
- 450 Choi, W., Faloon, I., Bouvier-Brown, N., McKay, M., Goldstein, A., Mao, J., Brune, W., LaFranchi, B., Cohen, R., and Wolfe, G.: Observations of elevated formaldehyde over a forest canopy suggest missing sources from rapid oxidation of arboreal hydrocarbons, *Atmospheric Chemistry and Physics*, 10, 8761-8781, 2010.
- Coggon, M. M., Gkatzelis, G. I., McDonald, B. C., Gilman, J. B., Schwantes, R. H., Abuhassan, N., Aikin, K. C., Arend, M. F., Berkoff, T. A., and Brown, S. S.: Volatile chemical product emissions enhance ozone and modulate urban chemistry, *Proceedings of the National Academy of Sciences*, 118, 2021.
- 455 Crounse, J. D., McKinney, K. A., Kwan, A. J., and Wennberg, P. O.: Measurement of gas-phase hydroperoxides by chemical ionization mass spectrometry, *Analytical chemistry*, 78, 6726-6732, 2006.
- Cui, Y. Y., Vijayan, A., Falk, M., Hsu, Y.-K., Yin, D., Chen, X. M., Zhao, Z., Avise, J., Chen, Y., and Verhulst, K.: A multiplatform inversion estimation of statewide and regional methane emissions in California during 2014–2016, *Environmental science & technology*, 53, 9636-9645, 2019.
- 460 De Gouw, J., Gilman, J., Borbon, A., Warneke, C., Kuster, W., Goldan, P., Holloway, J., Peischl, J., Ryerson, T., and Parrish, D.: Increasing atmospheric burden of ethanol in the United States, *Geophysical Research Letters*, 39, 2012.
- Dolgorouky, C., Gros, V., Sarda-Esteve, R., Sinha, V., Williams, J., Marchand, N., Sauvage, S., Poulain, L., Sciare, J., and Bonsang, B.: Total OH reactivity measurements in Paris during the 2010 MEGAPOLI winter campaign, *Atmospheric Chemistry and Physics*, 12, 9593-9612, 10.5194/acp-12-9593-2012, 2012.
- 465 Faloon, I. C., Chiao, S., Eiserloh, A. J., Alvarez, R. J., Kirgis, G., Langford, A. O., Senff, C. J., Caputi, D., Hu, A., Iraci, L. T., Yates, E. L., Marrero, J. E., Ryoo, J.-M., Conley, S., Tanrikulu, S., Xu, J., and Kuwayama, T.: The California baseline ozone transport study (CABOTS), *Bulletin of the American Meteorological Society*, 101, E427-E445, 10.1175/bams-d-18-0302.1, 2020.

- Fraser, M. P., Cass, G. R., and Simoneit, B. R.: Gas-phase and particle-phase organic compounds emitted from motor vehicle traffic in a Los Angeles roadway tunnel, *Environmental science & technology*, 32, 2051-2060, 1998.
- 470 Fried, A., Walega, J., Weibring, P., Richter, D., Simpson, I. J., Blake, D. R., Blake, N. J., Meinardi, S., Barletta, B., and Hughes, S. C.: Airborne formaldehyde and volatile organic compound measurements over the Daesan petrochemical complex on Korea's northwest coast during the Korea-United States Air Quality study: Estimation of emission fluxes and effects on air quality, *Elementa: Science of the Anthropocene*, 8, 2020.
- 475 Gentner, D. R., Ford, T. B., Guha, A., Boulanger, K., Brioude, J., Angevine, W. M., de Gouw, J. A., Warneke, C., Gilman, J. B., Ryerson, T. B., Peischl, J., Meinardi, S., Blake, D. R., Atlas, E., Lonneman, W. A., Kleindienst, T. E., Beaver, M. R., Clair, J. M. S., Wennberg, P. O., VandenBoer, T. C., Markovic, M. Z., Murphy, J. G., Harley, R. A., and Goldstein, A. H.: Emissions of organic carbon and methane from petroleum and dairy operations in California's San Joaquin Valley, *Atmospheric Chemistry and Physics*, 14, 4955-4978, 10.5194/acp-14-4955-2014, 2014.
- 480 Gilbert, N. L., Guay, M., Gauvin, D., Dietz, R. N., Chan, C. C., and Lévesque, B.: Air change rate and concentration of formaldehyde in residential indoor air, *Atmospheric Environment*, 42, 2424-2428, 10.1016/j.atmosenv.2007.12.017, 2008.
- Gilman, J. B., Lerner, B. M., Kuster, W. C., and De Gouw, J. A.: Source signature of volatile organic compounds from oil and natural gas operations in northeastern Colorado, *Environmental science & technology*, 47, 1297-1305, 2013.
- 485 Gilman, J. B., Kuster, W. C., Goldan, P. D., Herndon, S. C., Zahniser, M. S., Tucker, S. C., Brewer, W. A., Lerner, B. M., Williams, E. J., Harley, R. A., Fehsenfeld, F. C., Warneke, C., and de Gouw, J. A.: Measurements of volatile organic compounds during the 2006 TexAQS/GoMACCS campaign: Industrial influences, regional characteristics, and diurnal dependencies of the OH reactivity, *Journal of Geophysical Research*, 114, 10.1029/2008jd011525, 2009.
- Gkatzelis, G. I., Coggon, M. M., McDonald, B. C., Peischl, J., Gilman, J. B., Aikin, K. C., Robinson, M. A., Canonaco, F., Prevot, A. S., and Trainer, M.: Observations confirm that volatile chemical products are a major source of petrochemical emissions in US cities, *Environmental science & technology*, 55, 4332-4343, 2021.
- 490 Goldstein, A. H. and Galbally, I. E.: Known and unexplored organic constituents in the earth's atmosphere, *Environmental science & technology*, 41, 1514-1521, 2007.
- Gu, S., Guenther, A., and Faiola, C.: Effects of anthropogenic and biogenic volatile organic compounds on Los Angeles air quality, *Environmental Science & Technology*, 55, 12191-12201, 2021.
- 495 Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The model of emissions of gases and aerosols from nature version 2.1 (MEGAN2.1): An extended and updated framework for modeling biogenic emissions, *Geoscientific Model Development*, 5, 1471-1492, 10.5194/gmd-5-1471-2012, 2012.
- 500 Hansen, R. F., Griffith, S. M., Dusanter, S., Gilman, J. B., Graus, M., Kuster, W. C., Veres, P. R., de Gouw, J. A., Warneke, C., Washenfelder, R. A., Young, C. J., Brown, S. S., Alvarez, S. L., Flynn, J. H., Grossberg, N. E., Lefter, B., Rappenglueck, B., and Stevens, P. S.: Measurements of total OH reactivity during CalNex-LA, *Journal of Geophysical Research: Atmospheres*, 126, 10.1029/2020jd032988, 2021.
- Heald, C. L. and Kroll, J. H.: The fuel of atmospheric chemistry: Toward a complete description of reactive organic carbon, *Sci Adv*, 6, eaay8967, 10.1126/sciadv.aay8967, 2020.
- Heald, C. L., Gouw, J., Goldstein, A. H., Guenther, A. B., Hayes, P. L., Hu, W., Isaacman-VanWertz, G., Jimenez, J. L., Keutsch, F. N., Koss, A. R., Misztal, P. K., Rappenglueck, B., Roberts, J. M., Stevens, P. S., Washenfelder, R. A., Warneke,



- 505 C., and Young, C. J.: Contrasting reactive organic carbon observations in the southeast United States (SOAS) and southern California (CalNex), *Environ Sci Technol*, 54, 14923-14935, 10.1021/acs.est.0c05027, 2020.
- Holzinger, R., Warneke, C., Hansel, A., Jordan, A., Lindinger, W., Scharffe, D. H., Schade, G., and Crutzen, P. J.: Biomass burning as a source of formaldehyde, acetaldehyde, methanol, acetone, acetonitrile, and hydrogen cyanide, *Geophysical Research Letters*, 26, 1161-1164, 10.1029/1999gl1900156, 1999.
- 510 Kampa, M. and Castanas, E.: Human health effects of air pollution, *Environ Pollut*, 151, 362-367, 10.1016/j.envpol.2007.06.012, 2008.
- Karl, T., Guenther, A., Turnipseed, A., Tyndall, G., Artaxo, P., and Martin, S.: Rapid formation of isoprene photo-oxidation products observed in Amazonia, *Atmospheric Chemistry and Physics*, 9, 7753-7767, 2009.
- 515 Kim, S., Jeong, D., Sanchez, D., Wang, M., Seco, R., Blake, D., Meinardi, S., Barletta, B., Hughes, S., Jung, J., Kim, D., Lee, G., Lee, M., Ahn, J., Lee, S.-D., Cho, G., Sung, M.-Y., Lee, Y.-H., and Park, R.: The controlling factors of photochemical ozone production in Seoul, South Korea, *Aerosol and Air Quality Research*, 18, 2253-2261, 10.4209/aaqr.2017.11.0452, 2018.
- Kim, S. W., McDonald, B. C., Seo, S., Kim, K. M., and Trainer, M.: Understanding the paths of surface ozone abatement in the Los Angeles Basin, *Journal of Geophysical Research: Atmospheres*, 127, e2021JD035606, 2022.
- 520 Kwok, E. S. and Atkinson, R.: Estimation of hydroxyl radical reaction rate constants for gas-phase organic compounds using a structure-reactivity relationship: An update, *Atmospheric Environment*, 29, 1685-1695, 1995.
- Lauraguais, A., Bejan, I., Barnes, I., Wiesen, P., and Coeur, C.: Rate coefficients for the gas-phase reactions of hydroxyl radicals with a series of methoxylated aromatic compounds, *The Journal of Physical Chemistry A*, 119, 6179-6187, 2015.
- 525 Lee, Y. N., Zhou, X., Kleinman, L., Nunnermacker, L., Springston, S., Daum, P., Newman, L., Keigley, W., Holdren, M., and Spicer, C.: Atmospheric chemistry and distribution of formaldehyde and several multioxygenated carbonyl compounds during the 1995 Nashville/Middle Tennessee Ozone Study, *Journal of Geophysical Research: Atmospheres*, 103, 22449-22462, 1998.
- Lerner, B. M., Gilman, J. B., Aikin, K. C., Atlas, E. L., Goldan, P. D., Graus, M., Hendershot, R., Isaacman-VanWertz, G. A., Koss, A., and Kuster, W. C.: An improved, automated whole air sampler and gas chromatography mass spectrometry analysis system for volatile organic compounds in the atmosphere, *Atmospheric Measurement Techniques*, 10, 291-313, 2017.
- 530 Ling, Z. H., Guo, H., Lam, S. H. M., Saunders, S. M., and Wang, T.: Atmospheric photochemical reactivity and ozone production at two sites in Hong Kong: Application of a Master Chemical Mechanism-photochemical box model, *Journal of Geophysical Research: Atmospheres*, 119, 10567-10582, 10.1002/2014jd021794, 2014.
- 535 Liu, S., Li, R., Wild, R., Warneke, C., De Gouw, J., Brown, S., Miller, S., Luongo, J., Jimenez, J., and Ziemann, P.: Contribution of human-related sources to indoor volatile organic compounds in a university classroom, *Indoor Air*, 26, 925-938, 2016.
- 540 Lou, S., Holland, F., Rohrer, F., Lu, K., Bohn, B., Brauers, T., Chang, C., Fuchs, H., Häsel, R., and Kita, K.: Atmospheric OH reactivities in the Pearl River Delta–China in summer 2006: measurement and model results, *Atmospheric Chemistry and Physics*, 10, 11243-11260, 2010.

- Malkina, I. L., Kumar, A., Green, P. G., and Mitloehner, F. M.: Identification and quantitation of volatile organic compounds emitted from dairy silages and other feedstuffs, *J Environ Qual*, 40, 28-36, 10.2134/jeq2010.0302, 2011.
- Mårtensson, L., Magnusson, M., Shen, Y., and Jönsson, J. Å.: Air concentrations of volatile organic acids in confined animal buildings—determination with ion chromatography, *Agriculture, ecosystems & environment*, 75, 101-108, 1999.
- 545 McDonald, B. C., De Gouw, J. A., Gilman, J. B., Jathar, S. H., Akherati, A., Cappa, C. D., Jimenez, J. L., Lee-Taylor, J., Hayes, P. L., and McKeen, S. A.: Volatile chemical products emerging as largest petrochemical source of urban organic emissions, *Science*, 359, 760-764, 2018.
- Misztal, P., Karl, T., Weber, R., Jonsson, H., Guenther, A. B., and Goldstein, A. H.: Airborne flux measurements of biogenic isoprene over California, *Atmospheric Chemistry and Physics*, 14, 10631-10647, 2014.
- 550 Mo, Z., Shao, M., Wang, W., Liu, Y., Wang, M., and Lu, S.: Evaluation of biogenic isoprene emissions and their contribution to ozone formation by ground-based measurements in Beijing, China, *Sci Total Environ*, 627, 1485-1494, 10.1016/j.scitotenv.2018.01.336, 2018.
- Nuvolone, D., Petri, D., and Voller, F.: The effects of ozone on human health, *Environ Sci Pollut Res Int*, 25, 8074-8088, 10.1007/s11356-017-9239-3, 2017.
- 555 Page, L. H., Ni, J.-Q., Heber, A. J., Mosier, N. S., Liu, X., Joo, H.-S., Ndegwa, P. M., and Harrison, J. H.: Characteristics of volatile fatty acids in stored dairy manure before and after anaerobic digestion, *Biosystems Engineering*, 118, 16-28, 2014.
- Palmer, P. I., Jacob, D. J., Fiore, A. M., Martin, R. V., Chance, K., and Kurosu, T. P.: Mapping isoprene emissions over North America using formaldehyde column observations from space, *Journal of Geophysical Research: Atmospheres*, 108, 2003.
- 560 Parrish, D., Ryerson, T., Mellqvist, J., Johansson, J., Fried, A., Richter, D., Walega, J., Washenfelder, R. d., De Gouw, J., and Peischl, J.: Primary and secondary sources of formaldehyde in urban atmospheres: Houston Texas region, *Atmospheric Chemistry and Physics*, 12, 3273-3288, 2012.
- Parrish, D. D., Derwent, R. G., and Faloon, I. C.: Long-term baseline ozone changes in the Western US: A Synthesis of Analyses, *Journal of the Air & Waste Management Association*, 71, 1397-1406, 2021.
- 565 Parrish, D. D., Young, L. M., Newman, M. H., Aikin, K. C., and Ryerson, T. B.: Ozone design values in southern California's air basins: Temporal evolution and U.S. background contribution, *Journal of Geophysical Research: Atmospheres*, 122, 11,166-111,182, 10.1002/2016jd026329, 2017.
- Peischl, J., Ryerson, T., Brioude, J., Aikin, K., Andrews, A., Atlas, E., Blake, D., Daube, B., De Gouw, J., and Dlugokencky, E.: Quantifying sources of methane using light alkanes in the Los Angeles basin, California, *Journal of Geophysical Research: Atmospheres*, 118, 4974-4990, 2013.
- 570 Pusede, S. E. and Cohen, R. C.: On the observed response of ozone to NO<sub>x</sub> and VOC reactivity reductions in San Joaquin Valley California 1995–present, *Atmospheric Chemistry and Physics*, 12, 8323-8339, 10.5194/acp-12-8323-2012, 2012.
- Reimann, S., Calanca, P., and Hofer, P.: The anthropogenic contribution to isoprene concentrations in a rural atmosphere, *Atmospheric Environment*, 34, 109-115, 2000.
- 575 Richter, D., Weibring, P., Walega, J. G., Fried, A., Spuler, S. M., and Taubman, M. S.: Compact highly sensitive multi-species airborne mid-IR spectrometer, *Applied Physics B*, 119, 119-131, 2015.

- 580 Roberts, J. M., Jobson, B. T., Kuster, W., Goldan, P., Murphy, P., Williams, E., Frost, G., Riemer, D., Apel, E., and Stroud, C.: An examination of the chemistry of peroxy-carboxylic nitric anhydrides and related volatile organic compounds during Texas Air Quality Study 2000 using ground-based measurements, *Journal of Geophysical Research: Atmospheres*, 108, 2003.
- Sachse, G. W., Hill, G. F., Wade, L. O., and Perry, M. G.: Fast-response, high-precision carbon monoxide sensor using a tunable diode laser absorption technique, *Journal of Geophysical Research: Atmospheres*, 92, 2071-2081, 1987.
- Salthammer, T., Mentese, S., and Marutzky, R.: Formaldehyde in the indoor environment, *Chemical reviews*, 110, 2536-2572, 2010.
- 585 Semadeni, M., Stocker, D. W., and Kerr, J. A.: The temperature dependence of the OH radical reactions with some aromatic compounds under simulated tropospheric conditions, *International journal of chemical kinetics*, 27, 287-304, 1995.
- Shaw, S. L., Mitloehner, F. M., Jackson, W., DePeters, E. J., Fadel, J. G., Robinson, P. H., Holzinger, R., and Goldstein, A. H.: Volatile organic compound emissions from dairy cows and their waste as measured by proton-transfer-reaction mass spectrometry, *Environmental science & technology*, 41, 1310-1316, 2007.
- 590 Simpson, I. J., Blake, D. R., Blake, N. J., Meinardi, S., Barletta, B., Hughes, S. C., Fleming, L. T., Crawford, J. H., Diskin, G. S., and Emmons, L. K.: Characterization, sources and reactivity of volatile organic compounds (VOCs) in Seoul and surrounding regions during KORUS-AQ, *Elementa: Science of the Anthropocene*, 8, 2020.
- Steiner, A., Cohen, R., Harley, R., Tonse, S., Millet, D., Schade, G., and Goldstein, A.: VOC reactivity in central California: comparing an air quality model to ground-based measurements, *Atmospheric Chemistry and Physics*, 8, 351-368, 2008.
- 595 Sun, H., Trabue, S. L., Scoggin, K., Jackson, W. A., Pan, Y., Zhao, Y., Malkina, I. L., Koziel, J. A., and Mitloehner, F. M.: Alcohol, volatile fatty acid, phenol, and methane emissions from dairy cows and fresh manure, *Journal of environmental quality*, 37, 615-622, 2008.
- Veres, P. R., Neuman, J. A., Bertram, T. H., Assaf, E., Wolfe, G. M., Williamson, C. J., Weinzierl, B., Tilmes, S., Thompson, C. R., and Thames, A. B.: Global airborne sampling reveals a previously unobserved dimethyl sulfide oxidation mechanism in the marine atmosphere, *Proceedings of the National Academy of Sciences*, 117, 4505-4510, 2020.
- 600 Warneke, C., De Gouw, J. A., Holloway, J. S., Peischl, J., Ryerson, T. B., Atlas, E., Blake, D., Trainer, M., and Parrish, D. D.: Multiyear trends in volatile organic compounds in Los Angeles, California: Five decades of decreasing emissions, *Journal of Geophysical Research: Atmospheres*, 117, 2012.
- Warneke, C., Schwarz, J. P., Crawford, J., Dibb, J., Kalashnikova, O., Frost, G., Seidel, F., Al-Saadi, J., Brown, S., Washenfelder, R., Brewer, A., Moore, R., Anderson, B., Yacovitch, T., Herndon, S., Liu, S., Jaffe, D., Johnston, N., Selimovic, V., Yokelson, B., Giles, D., Holben, B., Goloub, P., Popovici, I., Trainer, M., Pierce, B., Fahey, D., Roberts, J., Soja, A., Peterson, D., Saide, P. E., Holmes, C., Wang, S., Coggon, M., Decker, Z., Ye, X., Stockwell, C., Xu, L., Gkatzelis, G. I., Lefer, B., and Crawford, J.: Fire influence on regional to global environments and air quality (FIREX-AQ), in preparation.
- 610 Wennberg, P. O., Mui, W., Wunch, D., Kort, E. A., Blake, D. R., Atlas, E. L., Santoni, G. W., Wofsy, S. C., Diskin, G. S., Jeong, S., and Fischer, M. L.: On the sources of methane to the Los Angeles atmosphere, *Environ Sci Technol*, 46, 9282-9289, 10.1021/es301138y, 2012.

- 615 Wennberg, P. O., Bates, K. H., Crouse, J. D., Dodson, L. G., McVay, R. C., Mertens, L. A., Nguyen, T. B., Praske, E., Schwantes, R. H., Smarte, M. D., St Clair, J. M., Teng, A. P., Zhang, X., and Seinfeld, J. H.: Gas-Phase reactions of isoprene and its major oxidation products, *Chem Rev*, 118, 3337-3390, 10.1021/acs.chemrev.7b00439, 2018.
- Whalley, L. K., Stone, D., Bandy, B., Dunmore, R., Hamilton, J. F., Hopkins, J., Lee, J. D., Lewis, A. C., and Heard, D. E.: Atmospheric OH reactivity in central London: observations, model predictions and estimates of in situ ozone production, *Atmospheric Chemistry and Physics*, 16, 2109-2122, 2016.
- 620 Wu, C., Wang, C., Wang, S., Wang, W., Yuan, B., Qi, J., Wang, B., Wang, H., Wang, C., and Song, W.: Measurement report: Important contributions of oxygenated compounds to emissions and chemistry of volatile organic compounds in urban air, *Atmospheric Chemistry and Physics*, 20, 14769-14785, 2020.
- Wu, S., Lee, H. J., Rohrbacher, A., Liu, S., Kuwayama, T., Seinfeld, J. H., and Kleeman, M. J.: Direct measurements of ozone response to emissions perturbations in California, *Atmos Chem Phys*, 22, 4929-4949, 2022.
- 625 Yang, Y., Shao, M., Wang, X., Nölscher, A. C., Kessel, S., Guenther, A., and Williams, J.: Towards a quantitative understanding of total OH reactivity: A review, *Atmospheric Environment*, 134, 147-161, 10.1016/j.atmosenv.2016.03.010, 2016.
- Yuan, B., Koss, A. R., Warneke, C., Coggon, M., Sekimoto, K., and de Gouw, J. A.: Proton-transfer-reaction mass spectrometry: Applications in atmospheric sciences, *Chemical Reviews*, 117, 13187-13229, 2017a.
- 630 Yuan, B., Coggon, M. M., Koss, A. R., Warneke, C., Eilerman, S., Peischl, J., Aikin, K. C., Ryerson, T. B., and de Gouw, J. A.: Emissions of volatile organic compounds (VOCs) from concentrated animal feeding operations (CAFOs): chemical compositions and separation of sources, *Atmospheric Chemistry and Physics*, 17, 4945-4956, 2017b.
- Zheng, W., Flocke, F., Tyndall, G., Swanson, A., Orlando, J., Roberts, J., Huey, L., and Tanner, D.: Characterization of a thermal decomposition chemical ionization mass spectrometer for the measurement of peroxy acyl nitrates (PANs) in the atmosphere, *Atmospheric Chemistry and Physics*, 11, 6529-6547, 2011.
- 635 Zhong, S., Whiteman, C. D., and Bian, X.: Diurnal evolution of three-dimensional wind and temperature structure in California's Central Valley, *Journal of Applied Meteorology*, 43, 1679-1699, 2004.