

Quantifying hard coal mines CH₄ emissions from TROPOMI and IASI observations using ~~high-resolution CAMS forecast data and~~ the wind-assigned anomaly method

Qiansi Tu^{1,2}, Matthias Schneider², Frank Hase², Farahnaz Khosrawi², Benjamin Ertl^{1,3}, Jaroslaw Necki⁴,
5 Darko Dubravica², Christopher J. Diekmann^{2,5}, Thomas Blumenstock², Dianjun Fang^{1,5}

¹School of Mechanical Engineering, Tongji University, Shanghai, China

² Karlsruhe Institute of Technology (KIT), Institute of Meteorology and Climate Research (IMK-ASF), Karlsruhe, Germany
a now at: Telespazio Germany GmbH

³ Karlsruhe Institute of Technology, Steinbuch Centre for Computing (SCC), Karlsruhe, Germany

10 ⁴AGH – University of Science and Technology, Krakow, Poland

⁵Qingdao Sino-German Institute of Intelligent Technologies, Qingdao, China

Correspondence to: Qiansi Tu (qiansi.tu@kit.edu), Matthias Schneider (matthias.schneider@kit.edu)

Abstract. Intensive coal mining activities in the Upper Silesian Coal Basin (USCB) in southern Poland are resulting in large amounts of methane (CH₄) emissions. Annual CH₄ emission reached 448 kt according to the European Pollutant Release and Transfer Register (E-PRTR, 2017). As a CH₄ emission hot spot in Europe, it is of importance to investigate its emission sources and make accurate emission estimates.

In this study, we use satellite-based total column-averaged dry-air mole fraction of CH₄ (XCH₄) from the TROPOspheric Monitoring Instrument (TROPOMI) and tropospheric XCH₄ (TXCH₄) from the Infrared Atmospheric Sounding Interferometer (IASI). In addition, the high-resolution model forecast XCH₄ and TXCH₄ from the Copernicus Atmosphere Monitoring Service (CAMS) are used to estimate the CH₄ emission rate averaged over three years (November 2017 to December 2020) in the

20 USCB region (49.3° - 50.8° N and 18° - 20° E). The wind-assigned anomaly method is first validated using CAMS forecast data (XCH₄ and TXCH₄), showing a good agreement to the CAMS-GLOB-ANT inventory. It indicates that the wind-assigned method works well. Using the CAMS inventory (CAMS-GLOB-ANT) as the a priori knowledge (location and the proportion

25 of the emission rate for each source in the total emissions) of the sources, together with ERA5 wind at 330 m, the wind-assigned XCH₄ anomalies for two opposite wind directions are calculated, which yields an estimated CH₄ emission of 815 ± 13 kt/year (9.7E26 ± 1.5E25 molec./s) for CAMS XCH₄ and 798 ± 11 kt/year (9.5E26 ± 1.3E25 molec./s) for CAMS TXCH₄. These values are very close to the total emission of the CAMS-GLOB-ANT (815 kt/year). Very good agreements between CAMS and the wind-assigned model results ($R^2 = 0.85$ for XCH₄ and TXCH₄) indicate that our wind-assigned method is quite

30 robust. The similar estimates of XCH₄ and TXCH₄ also imply that for a strong source, the dynamically induced variations of the CH₄ mixing ratio in the upper troposphere and lower stratosphere region are of secondary importance. This wind-assigned method is further applied to the TROPOMI XCH₄ and TROPOMI+IASI TXCH₄ by using the Carbon dioxide and Methane (CoMet) inventory derived for the year 2018. The calculated averaged total CH₄ emission over the USCB region is about 496

kt/year (5.9E26 molec./s) for TROPOMI XCH₄ and 437 kt/year (5.2E26 molec./s) for TROPOMI+IASI TXCH₄. These values are very close to the ones given in the E-PRTR inventory (448 kt/year) and the ones in the CoMet inventory (555 kt/year) and are thus in agreement with these inventories. The similar estimates of XCH₄ and TXCH₄ also imply that for a strong source, the dynamically induced variations of the CH₄ mixing ratio in the upper troposphere and lower stratosphere region are of secondary importance. Uncertainties from different error sources (background removal and noise in the data, vertical wind shear, wind field segmentation, and angle of the emission cone) are approximate 14.8% for TROPOMI XCH₄ and 11.4% for TROPOMI+IASI TXCH₄. These results suggest that our wind-assigned method is quite robust and might also serve as a simple method to estimate CH₄ or CO₂ emissions for other regions.

~~Since the wind speed is increasing with altitude, sensitivity tests show that higher CH₄ emission strengths are yielded with increasing altitude and vice versa. About 23% lower and 13% higher emission estimates are obtained when instead of the wind information at 330 m, the wind information at a lower altitude (10 m) and at a higher altitude (500 m) is used, respectively. When using different wind coverage and different wind segmentation, the estimated emission strengths change by 13.4% and 5.2%, respectively. These results suggest that our wind assigned method is quite robust and might also serve as a simple method to estimate CH₄ or CO₂ emissions for other regions.~~

1 Introduction

Atmospheric methane (CH₄) is the second most important anthropogenic greenhouse gas (GHG) with a larger global warming potential than carbon dioxide (CO₂) (IPCC, 2014). The globally averaged amount of atmospheric CH₄ has increased by 260% to 1877 ± 2 ppb from the preindustrial era until 2019 (World Meteorological Organization, 2020). Methane sources induced by anthropogenic activities include fossil fuel production and use (e.g., coal mining, gas/oil extraction), waste disposal, and agriculture, which in total accounts for about 60% of the total CH₄ emissions (Saunois et al., 2020). Although most sources and sinks of CH₄ have been characterized, their spatial-temporal variations and relative contributions to the atmospheric CH₄ level are still highly uncertain (Kirschke et al., 2013; Saunois et al., 2020).

Approximately 33% of the CH₄ emissions from coal mining (42,000 kt/year) are estimated to come from the total fossil-fuel-related emissions during 2008-2017 (Saunois et al., 2020). CH₄ is released primarily to the atmosphere via ventilation shafts located at the surface during the production and processing of the coal (Saunois et al., 2020; Andersen et al. 2021). The largest contribution of CH₄ emission related with the coal mining activities in Europe is from southern Poland—the Upper Silesian Coal Basin (USCB) (Luther et al., 2019; Krautwurst et al., 2021). The UCSB is in the Silesian Upland, which is a plateau between 200 and 300 m above sea level with a predominant south-west wind. The UCSB within Poland covers an area of over 5800 km², and to its south is the Tatra Mountain ridge with elevations larger than 2000 m a.s.l. The European Pollutant Release and Transfer Register (E-PRTR, 2017; <https://prtr.eea.europa.eu/>, last access: 25 October 2021) reports that the total CH₄ emissions from the UCSB region amount to 448 kt/year. Most of these emissions are from mining activities and heavy industry (Kostinek et al., 2021), which makes this region a hot spot of CH₄ emission in Europe.

To investigate the CH₄ emission from this hot spot, the Carbon Dioxide and Methane (CoMet) campaign was performed, covering roughly 3 weeks from May to June 2018. A variety of state-of-art instruments, including in situ and remote sensing instruments on the ground and aboard five research aircraft, were deployed in order to provide independent observations of GHG emissions on local to regional scale and provide data for satellite validation (more details can be found in Luther et al., 70 Fiehn et al., 2020; [Galkowski et al., 2021](#); Kostinek et al., 2021; Krautwurst et al., 2021; [Wolff et al., 2021](#)). For example, [Galkowski et al. \(2021\) present results of in situ GHG measurements obtained over nine research flights of the German research Aircraft HALO \(High Altitude and LOnge Range Research Aircraft\) acting as the airborne flagship of the CoMet campaign, together with simultaneous flasks measurements for isotopic composition of CH₄. A new lidar CHARM-F \(CO₂ and CH₄ Atmospheric Remote Monitoring Flugzeug\) was also onboard HALO and its measurements were investigated to determine](#) 75 [CO₂ emission rates from the power plant \(Wolff et al., 2021\)](#). Many studies present similar CH₄ emission estimates for the region based on different instruments and methods. Luther et al. (2019) estimated CH₄ emissions ranging from 6 ± 1 kt/year for a single shaft to up to 109 ± 33 kt/year for a subregion of the USCB covering several shafts, by using several portable Fourier Transform Infrared (FTIR) spectrometers (Bruker EM27/SUN). [Active AirCore system aboard an unmanned aerial vehicle \(UAV\) was used to measure CH₄ downwind of a single ventilation shaft and emission rates ranging from 0.5 to 14.5](#) 80 [kt/year based on a mass balance approach and ranging from 1.1 to 9.0 kt/year based on an inverse Gaussian method were estimated \(Andersen et al., 2021\)](#). Fiehn et al (2020) analyzed aircraft- and ground-based in situ observations and reported an emission estimate of 436 ± 115 kt/year and 477 ± 101 kt/year from two selected flights. An advanced model approach was introduced by Kostinek et al. (2021) to investigate two research flights in the morning and afternoon, resulting in estimated 85 CH₄ emissions of 451 ± 77 kt/year and 423 ± 79 kt/year, respectively. Another emission estimate based on the observations from the nadir-looking passive remote sensing Methane Airborne MAPper (MAMAP) instrument accounted for 8.8 kt/year to 78.8 kt/year for a sub-clusters of ventilation shafts (Krautwurst et al., 2021). A recent study ([Luther et al., 2021](#)[Luther et al., 2022](#)) displays a larger emission rate of 414 – 790 kt/year based on a network of four portable FTS instruments (EM27/SUN) 90 during the CoMet campaign.

Launched in October 2017, the TROPOspheric Monitoring Instrument (TROPOMI) on board the Sentinel-5 Precursor satellite provides an unprecedent high spatial resolution (5.5×7 km²) of the CH₄ total column-averaged dry-air mole fraction (XCH₄) (Veefkind et al., 2012; Lorente et al., 2021). An a posteriori method has been developed by Schneider et al. (2021) to obtain tropospheric XCH₄ (TXCH₄) by combining observations from TROPOMI and the Infrared Atmospheric Sounding Interferometer (IASI). This synergetic product is not influenced by the changing tropopause height, and it offers improved sensitivity to the tropospheric variations than the total column XCH₄ data from either sensor. The improved real-time forecast 95 data with high resolution ($0.1^\circ \times 0.1^\circ \sim 9$ km \times 9km) are produced by the Copernicus Atmosphere Monitoring Service (CAMS) (Agustí-Panareda et al., 2019; Barré et al., 2021). All data sets provide a large spatial coverage and long-term XCH₄/TXCH₄ observations, which help to better estimate CH₄ emission in the USCB region.

In Sect. 2 we present the data sets and methodology used in this study to derive estimated CH₄ emissions. The results and discussions are presented in Sect. 3. We present a novel wind-assigned method introduced by Tu et al., 2022, which is firstly

100 verified by the CAMS model forecasts and then applied to the TROPOMI XCH₄ and TROPOMI+IASI TXCH₄ data to estimate
the CH₄ emissions in the USCB region for the time period from November 2017 to December 2020, together with an
uncertainty analysis. Finally, the summary and conclusions are given in Sect. 4.

2 Data sets and method

105 There are over 50 active ventilation shafts in the USCB region (49.3° - 50.8° N and 18° - 20° E), Poland, whose emission rates
range between 0.17 kt/year and 41.02 kt/year (Gałkowski et al., 2021) (Figure 4b). Most of them are located near Katowice
and further west and southwest of Katowice.


2.1 CAMS CH₄ forecast and emission inventories

110 The Integrated Forecasting System (IFS, <https://www.ecmwf.int/en/publications/ifs-documentation>, last access: 27 October,
2021) from the European Centre for Medium-Range Weather Forecasts (ECMWF) is used in the CAMS atmospheric
composition analysis and forecasts system to simulate five-day CO₂ and CH₄ forecasts (Agustí-Panareda et al., 2019, Barré et
al., 2021), as well as other chemical species and aerosols (Flemming et al., 2015; Morcrette et al., 2009). This model is also
used in the operational Numerical Weather Prediction (NWP) system, but with additional modules (Agustí-Panareda et al.,
2019). The forecast data used in this study is the same suit as the one used in Barré et al. (2021), where the Cycle 45r1 IFS
model cycle was implemented. The CAMS GHGs operational dataset includes analysis and forecast data at medium and high
115 resolution with 137 model levels from the surface to 0.01 hPa (Barré et al., 2021). In this study we will focus on using the
high-resolution CH₄ forecasts, which have a spatial resolution of 0.1° × 0.1° and a temporal resolution of 3h, starting from
00:00 UTC. Here we use the daily averaged CAMS forecasts during 9:00 UTC - 18:00 UTC at each resolution grid point. The
corresponding standard deviation (STD) is considered as the noise/error.


120 The anthropogenic methane emissions used in the global CAMS forecasts are from the CAMS global anthropogenic
emission inventory (CAMS-GLOB-ANT, Granier et al., 2019; <https://permalink.aeris-data.fr/CAMS-GLOB-ANT>, last access:
27 October, 2021). The CAMS-GLOB-ANT inventory is based on the emissions provided by the EDGARv4.3.2 inventory for
the time period 2000-2012 (Crippa et al., 2018) and linearly extrapolated to 2020 using the trends from the CEDA global
inventory for the time period 2011-2014 (Hoesly et al., 2018). The latest version (CAMS-GLOB-ANT v4.2) was released in
125 March 2020, using the same set-up as v4.1 except for adding the emissions in 2020. The anthropogenic sources in the standard
v4.2 are divided into 12 sectors and the agriculture sections are split into three sectors, including livestock, soils and waste
burning (<https://eccad3.sedoo.fr/>, last access: 27 October, 2021). The inventory is provided as monthly mean with the same
spatial resolution (0.1° × 0.1°) as the CAMS forecast data (Granier et al., 2019).

130 The monthly averages of the CAMS global anthropogenic emissions for different sectors in the study area of USCB are
presented in Figure 1. The emissions from the sectors “agriculture soils” and “solvents” are zeros. The CH₄ emitted from ships
has 19 orders of magnitude, which are much lower than the other sectors. Thus, these three sectors are not shown here. The

sources from agriculture/livestock ($1.7\text{E}25 \pm 4.0\text{E}25$ molec./s) amount only 4% of the total emissions in this region. The dominant CH₄ sources in this region are fugitive sources from energy production and distribution (e.g. fuel use). With a mean value of $7.9\text{E}26$ molec./s and a standard deviation of $2.2\text{E}25$ molec./s, they account for 82% of the anthropogenic CH₄ emissions in the CAMS-GLOB-ANT inventory ($9.7\text{E}26$ molec./s in total). This becomes particularly visible in the spatially overlapping distribution within the USCB (see Figure 2). The seasonal emission variations of the fugitive sector are minor and can be ignored. Therefore, we apply the three-year mean of total emissions at grids with significant emissions without considering seasonal variations in the simple cone plume model (see Sect. 2.3). The total emissions amount to $9.7\text{E}26$ molec./s over this study area.



140 **Figure 1: Stacked area plot for different sectors of the monthly averaged CAMS global anthropogenic emissions (>1E20 molec./s) in the USCB region for 2017-2020 (<https://permalink.aeris-data.fr/CAMS-GLOB-ANT>, last access: 22 December 2021. Granier et al., 2019).**



145 **Figure 2: Spatial distribution of (a) the CAMS global anthropogenic emissions from all sectors and (b) percentage share of the fugitive emissions compared to the overall anthropogenic emissions over the USCB region on a $0.1^\circ \times 0.1^\circ$ latitude/longitude grid. The fugitives are the dominant CH₄ sources.**

2.2 TROPOMI and IASI data sets

The TROPOMI instrument is a nadir-viewing, imaging spectrometer, which uses passive remote sensing techniques to perform measurements of the solar radiation reflected by and radiated from the earth in the ultraviolet, the visible, the near-infrared and the shortwave infrared spectral bands (Veefkind et al., 2012). The instrument crosses the equator at about 13:30 local solar time at each orbit with a repeat cycle of 17 days. It observes a full swath (2600 km) per second with an orbit duration of 100 min. The algorithm for CH₄ column retrieval is called RemoTeC algorithm and it has been extensively used to derive CO₂ and CH₄ retrievals from the Greenhouse Gases Observing Satellite (GOSAT) and Orbiting Carbon Observatory-2 (OCO-2; Boesch et al., 2011; Butz et al., 2009, 2011; Hasekamp and Butz, 2008; Schepers et al., 2012). An updated retrieval algorithm has been implemented by Lorente et al. (2021) to obtain a data suit with less scatter and a higher resolution surface altitude database. This updated TROPOMI XCH₄ dataset has been validated with the Total Carbon Column Observing Network (TCCON) (-3.4 \pm 5.6 ppb) and GOSAT (-10.3 \pm 16.8 ppb), showing very good agreements. In this study the TROPOMI XCH₄ during November 2017 and December 2020 within the study area over the USCB region is investigated. The data provided by Lorente et al. (2021) includes an additional quality filter parameter (quality value, qa). TROPOMI XCH₄ with qa=1.0 represents the data under clear-sky and low-cloud atmospheric conditions and the problematic data points are removed as well. This quality filter has been applied in this study and about 16,000 data are derived over the three-year time period considered in this study.

The IASI instrument is a nadir viewing Fourier-transform spectrometer that measures the infrared part of the electromagnetic spectrum. IASI measurements are performed with a horizontal resolution of 12 km and a full swath width of about 2200 km on the ground. It is the key payload element of the polar-orbiting Metop-A -B and -C satellites. These satellites overpass the equator at 09:30 in the morning and 21:30 local time in the evening with about 14 orbits per day. It provides unprecedented accurate vertical information of atmospheric temperature and humidity, which helps to improve numerical weather prediction (NWP) (Collard, 2007; Coopmann et al., 2020). The thermal infrared nadir spectra of IASI have been successfully used in retrieving different atmospheric trace gas profiles and these retrievals are especially sensitive between the middle troposphere and the stratosphere (García et al., 2018; Diekmann et al., 2021; Schneider et al., 2021, 2022). By combining the IASI CH₄ profiles and the TROPOMI CH₄ total column, which has a higher sensitivity near ground, we are able to detect the tropospheric XCH₄ (TXCH₄) independently from CH₄ at higher altitudes. The combined product cannot be obtained by either the TROPOMI or IASI product independently. The combined product shows a weak positive bias of about 1 % with respect to the reference data (Schneider et al., 2021). We refer to this product in the following as the TROPOMI+IASI TXCH₄ and it comprises about 12,000 data points for the time period considered in this study.

175 2.3 Simple cone plume model and wind-assigned anomaly method

The averaged distribution of emitted CH₄ over a long-term period can be modeled simply as an evenly-distributed cone-shape dispersion based on the wind and source strength. Since CH₄ is a long-lived gas, its decay is negligible for short periods and not considered in the model. This model is referred to as simple cone plume model (see Figure A- 1). This model is easy to apply, and the estimated emission strengths are reasonable compared with the ones from other studies (Tu et al., 2022).

180 Based on the simple cone plume model, the enhanced CH₄ column (ΔCH_4) at the downwind side of the location (x_i, y_i) is computed as:

$$\Delta\text{CH}_4(x_i, y_i) = \frac{\varepsilon}{v \cdot d(x_i, y_i) \cdot \alpha} \quad \text{Eq. 1}$$

where the emission strength ε is the a priori knowledge from the CAMS-GLOB-ANT data set or from the coal mine ventilation shafts in this study (see Sect. 3.2). Their emission rates are assumed to be constant with time from 2017 to 2020. α is the angle of the emission cone and has an empirical value of 60°, which has been derived from TROPOMI NO₂ measurements (Tu et al., 2022). v is the wind speed from ERA5, which is the fifth generation ECMWF reanalysis product using 4D-Var data assimilation and model forecasts in Cycle 41R2 of the ECMWF IFS model (Copernicus Climate Change Service, C3S, 2017, Hersbach et al., 2020). ERA5 provides hourly estimates on 137 pressure levels in the vertical covering the atmosphere from the surface up to 0.01 hPa, with a spatial resolution of 0.25°×0.25° (Hersbach et al., 2020). d is the distance between the downwind location and the CH₄ emission source. Each individual source either from the CAMS-GLOB-ANT inventory or from the knowledge of the ventilation shafts is considered as an individual point source. The daily plume from each point source (location at (i,j)) is averaged over daytime (8:00 UTC - 18:00 UTC):

$$\overline{\text{XCH}_4}_{(i,j)} = \frac{1}{11} \sum_{t=1}^{11} \text{XCH}_4_{(i,j),t} \quad \text{Eq. 2}$$

these daily plumes are super-positioned over all point sources to obtain a daily plume ($\overline{\text{XCH}_4}_{\text{daily}}$):

$$\overline{\text{XCH}_4}_{\text{daily}} = \sum_{s=1}^{N_s} \overline{\text{XCH}_4}_{(i,j),s} \quad \text{Eq. 3}$$

where N_s represents the number of the sources.

195 The wind distributions at different height levels (10 m, ~330 m, ~500 m) over the USCB region are presented in Figure 3. The wind speed increases with increasing altitude (see **Table 1**). The ERA5 wind is divided into two opposite wind regimes based on directions (e.g., 135°-315° for SW and the rest for NE). For each wind regime **sector**, an averaged plume is computed:

$$\overline{\text{XCH}_4}_{\text{SW/NE}} = \frac{1}{N_d} \sum_{d=1}^{N_d} \overline{\text{XCH}_4}_{\text{daily},d} \quad \text{Eq. 4}$$

where N_d is the number of the days with SW wind or NE wind.

The difference of the two plumes is therefore the wind-assigned anomaly:

$$\text{wind-assigned anomaly} = \overline{\text{XCH}_4}_{\text{NE}} - \overline{\text{XCH}_4}_{\text{SW}} \quad \text{Eq. 5}$$

200 The estimated emission strengths can be calculated by fitting the modelled anomalies to the known anomalies from e.g. CAMS XCH₄/TXCH₄, and TROPOMI and TROPOMI+IASI observations. Note that CH₄ has a lifetime of around 12 years, which results in a high background concentration compared to the newly emitted CH₄. Thus, the contributions from the

background should be removed for correctly estimating emissions (Liu et al., 2021). The background is considered to consist of a constant value, a linear increase with time, a seasonal cycle, a daily anomaly and a horizontal anomaly. For more details, see the Appendix in Tu et al., 2022. [The uncertainties \(\$\pm\$ values\) in the emission estimate are determined by considering the deficits of the background model due to the imperfect elimination of the background and the noise in the data set.](#)

205

This method was firstly used to estimate the CH₄ emission from landfills in Madrid, Spain based on nearly three-year space-borne XCH₄ data, and different opening angles were investigated to obtain an empirical value (60°) (Tu et al., 2022). The CH₄ emission strengths derived from satellite products have the same orders of magnitude as the ones from single-day observations by ground-based instruments, showing that this method works properly.

210

Table 1: Number of days and the averaged wind speed (\pm standard deviation) per specific wind [regime](#) during daytime (08:00 UTC – 18:00 UTC) at different vertical levels from November 2017 to December 2020 over the USCB region. The days for the three-year average coincide with the TROPOMI overpass days.

	NE / >315° or <135°		SW / 135° – 315°	
	Number of days in total (%)	Averaged wind speed \pm standard deviation (m s ⁻¹)	Number of days in total (%)	Averaged wind speed \pm standard deviation (m s ⁻¹)
10 m	39.1	3.2 \pm 1.5	56. 9	3.4 \pm 1.6
~330 m (975 hPa)	38.7	4.1 \pm 2.2	56.9	4.3 \pm 2.3
~500 m (950 hPa)	38.7	5.0 \pm 2.7	57.3	5.9 \pm 3.5

215




Figure 3: Windrose plots for daytime (08:00 UTC – 18:00 UTC) from November 2017 to December 2020 for the ERA5 model wind at different vertical levels (10 m, ~330 m and ~500 m). The days for the three-year average coincide with the TROPOMI overpass days.

3. Results and discussion

220 3.1 Estimated emissions derived from CAMS forecasts (evaluation of the method)

The CAMS forecast XCH₄ data from November 2017 to December 2020 within the study area are illustrated in Figure 4 left. The areas with high XCH₄ amounts fit well with the CAMS anthropogenic CH₄ emissions (square symbols). Similar to the CoMet inventory, high sources in the CAMS-GLOB-ANT inventory are centered in this region, but there are other weaker sources outside. The total emission rate of the CoMet inventory is 555 kt/year (6.6E26 molec./s), which is slightly less than the CAMS-GLOB-ANT emissions (815 kt/year). This is probably because the CAMS-GLOB-ANT includes more CH₄ emission sources, e.g., wastes, and combustion from residential, commercial, which account to about 20%.




Figure 4: Averaged (a) CAMS forecast XCH₄, (b) TROPOMI XCH₄, and (c) TROPOMI+IASI TXCH₄ in the USCB region on a $0.1^\circ \times 0.1^\circ$ latitude/longitude grid during November 2017–December 2020. The square and triangle symbols represent the locations of CAMS-GLOB-ANT sources (for better viewing, only the emission strengths larger than $1E24$ molec./s are shown here) and the active coal mine shafts from the CoMet inventory (Galkowski et al., 2021), respectively. Different colors denote the amount of emission rates. The white grids represent no data from TROPOMI or the number of the points in the grid less than 5. A zoom version of panel (b) is shown in the appendix (Figure A- 2). Note, a different colorbar has been used in panel (c).

Based on the CAMS emissions, the wind-assigned method is applied to CAMS XCH₄. The XCH₄ enhancement (raw-background) and the wind-assigned anomalies are presented in Figure 5a and b, respectively. [The example plumes of the enhancements for wind coming from NE and SW are presented in Figure A- 3](#). Note, that the CAMS XCH₄ is coincided with TROPOMI XCH₄ for better comparison. Some data are thus missing here mostly due to the quality filter of TROPOMI observations. After removing the XCH₄ background, the XCH₄ anomalies well represent the CAMS sources. The highest CH₄ sources from the CAMS-GLOB-ANT inventory are also obviously visible in the 2D anomalies. In addition, the spatial distributions of the three XCH₄ data products show different patterns (Figure 4), whereas the anomalies (after removing background) patterns are similar (Figure 5(a) and (d), Figure 7(a) and (d)). This indicates that the background removal is of importance for XCH₄ and our method works well.

The wind-assigned anomalies for CAMS and [simple](#) cone plume model show a very good agreement with a slope of 1.11 and a R^2 of 0.85 (Figure 5c). Our results are derived from the CAMS emission information, and they agree very good with the CAMS model data. The estimated emission rate is about 815 ± 17 kt/year ($9.7E26 \pm 2.0E25$ molec./s) when using the ERA5 wind at 975 hPa (~330 m) and this value is quite close to the CAMS-GLOB-ANT (estimated emission rate at other levels are


presented in Sec. 3.2, see Figure 8 as well). Therefore, we use ERA5 wind at this level in the following. Note that the points whose distances to the nearest dominant sources are less than 10 km, are removed here, because they are very close to the significant sources and small changes in wind (either speed or direction) can result in high uncertainties.

250 XCH₄ is affected by local surface emissions and a varying stratospheric contribution due to changes in the tropopause altitude (Liu et al., 2021; Schneider et al., 2021). This stratospheric contribution has to be taken into account, in order to be able to use XCH₄ for a reliable investigation of local surface CH₄ sources and sinks (Pandey et al., 2016). Our background removal method effectively accounts for the stratospheric contribution. To show this we apply the approach to CAMS forecasts of XCH₄ (which has a significant stratospheric contribution) and TXCH₄ (calculated from the CAMS forecast as the CH₄ averaged from surface to 6 km, which should have a very limited stratospheric contribution). The results are presented in

255 Figure 5d-f. The CAMS TXCH₄ anomalies have similar distribution as CAMS XCH₄ anomalies, suggesting that our background removal approach reliably removes the stratospheric contribution. The retrieved CH₄ from satellite observations are based on total columns and therefore, these are strongly affected by the stratospheric abundance, i.e., by the changing tropopause altitude (Liu et al., 2021; Schneider et al., 2021). The model simulation uncertainties in representing XCH₄ in the stratosphere might introduce biases in investigating CH₄ sources and sinks (Pandey et al., 2016). To remove this influence, we calculate the tropospheric CAMS forecast CH₄ (TXCH₄) from the surface up to 7 km. The results are presented in Figure 5d-f. The CAMS TXCH₄ anomalies have similar distribution as CAMS XCH₄, showing that removing the background also works for the tropospheric CH₄. The wind-assigned plume and the correlation between CAMS and the wind-assigned model results are very similar between XCH₄ and TXCH₄. The estimated CH₄ emission strength derived from CAMS TXCH₄ is 798 ± 15

260 kt/year (9.5E26 ± 1.8E25 molec./s).

265 kt/year (9.5E26 ± 1.8E25 molec./s).






Figure 5: (a)-(c): CAMS XCH₄ enhancement (XCH₄-background), the wind-assigned anomalies (NE-SW), and correlation plot of the wind-assigned anomalies between CAMS and the [simple](#) cone plume model with using the CAMS-GLOB-ANT inventory (9.7E26 molec./s in total) and ERA5 wind at 330 m during November 2017–December 2020 over the USCB region. (d)-(f): the same as for the upper panel but for CAMS TXCH₄ (colorbars in (d) and (e) are different from that for XCH₄). The square symbols represent the locations of the CAMS-GLOB-ANT (>1E24 molec./s) inventory and different colors denote the amount of emission rates. The hatched areas in (a)-(b) and (d)-(e) represent no data in these grids. The uncertainties in (c) and (f) represent the mean error bars, i.e., error propagation of the background uncertainty and the CAMS standard deviation.

275 3.2 Estimated emissions derived from satellite observations

The high-resolution TROPOMI XCH₄ provides the ability to detect and quantify the CH₄ emissions (e.g., oil and gas sector, coal mining) on fine and large scales (Pandey et al., 2019; Varon et al., 2019; Gouw et al., 2020; Schneising et al., 2020). Figure 6 illustrates the enhanced XCH₄ (raw XCH₄-background in the upwind) distribution over the USCB region on an example day (6 June 2018), in which the wind mostly came from northeast. As expected, obvious XCH₄ enhancements were 280 observed by TROPOMI along the downwind direction (southwest of Katowice where most ventilation shafts are located), as well as simulated by the CAMS forecast. The downwind-enhanced XCH₄ modeled by our simple [cone](#) plume model and based on the CAMS-GLOB-ANT inventory also shows a similar shape of plume. This enhancement was also observed by portable FTIR instruments (COCCON) employed during the CoMet campaign (Figure 4 in Luther et al., 2019). The observations support the statement that TROPOMI is able to detect the CH₄ emission signals. In addition, the spatial pattern of the downwind 285 plume is similar to that of the cone-shaped plume, which implies our cone-shape assumption is reasonable.




Figure 6: ΔXCH_4 together with the ERA5 wind at 12:00 UTC from (a): TROPOMI observations at 11:34 UTC, (b): CAMS forecast at 12:00 UTC, and (c): from the simple [cone](#) plume model (averaged over the daytime) based on the CAMS-GLOB-ANT inventory

290 over the USCB region on an example day (6 June 2018). The “bg” in the title of (a) and (b) represents the average background, derived from the mean XCH₄ in the upwind region (50.3°-50.8° N, 19.5° -20.0° E). [Note, a different colorbar has been used in panel \(c\) for improved recognizability.](#)

295 The three-year averaged TROPOMI XCH₄ observations presented in Figure 4b shows scattered high XCH₄ amounts, whereas CAMS XCH₄ is more concentrated on the center of the study area, and those agree well with its anthropogenic emission sources (CAMS-GLOB-ANT inventory). This might be because TROPOMI detects other real CH₄ sources that are not included in the CAMS forecast model data.

300 For better comparison with other studies discussing the coal mine emissions in the USCB region, we apply the CoMet inventory as the a priori known sources in the wind-assigned method to estimate the CH₄ emissions. The results are illustrated in Figure 7. The TROPOMI XCH₄ and TROPOMI+IASI TXCH₄ anomalies show high concentrations around the areas where the ventilation shafts are located and the region in the northeast of Katowice. Although the anomalies of the satellite observations are lower than the CAMS results (Figure 5a), their spatial distributions are similar. Positive and negative plumes can be clearly seen in Figure 7b and e. The correlation of the wind-assigned anomalies between the TROPOMI and [simple](#) cone plume model has a very good agreement with an R² value of 0.76. Similar results are also derived from TROPOMI+IASI TXCH₄ with a R² value of 0.62. Compared to CAMS data, higher scatter is expected, because satellite observations suffer from observational errors and might contain more CH₄ sources (e.g., landfills, gas distribution network). Although none of 305 these sources have the same orders of magnitude of coal mining emission, they might still bring some errors.

310 The estimated CH₄ emission strengths are 496 ± 17 kt/year ($5.9\text{E}26 \pm 2.1\text{E}25$ molec./s) for XCH₄ and 437 ± 27 kt/year ($5.2\text{E}26 \pm 3.2\text{E}25$ molec./s) for TXCH₄, and both are close to the E-PRTR inventory (448 kt/year). The TROPOMI+IASI result has a higher uncertainty than the TROPOMI result, because (1) the vertical distribution of CH₄ is in general much more difficult to measure than the total column of CH₄ and (2) the vertical distribution is derived by considering two independent measurements, each with its own noise error. This might change for a larger number of data points (e.g., by using data from more years or by applying the method to IASI and TROPOMI successors on the upcoming METOP-SG satellite, which offers much more collocated observations).

315 However, in our study using TXCH₄ data in addition to XCH₄ data nicely documents the robustness of the method. Important for a correct estimation of the emission is the correct removal of the methane background signal. For XCH₄ the stratospheric and the tropospheric backgrounds have to be removed, whereas only the tropospheric background has to be removed for TXCH₄. Despite this difference, we estimate very similar emission rates from both data sets and the emission rate uncertainties of using XCH₄ or TXCH₄ are small compared to the estimated emission rates.

320 Figure 8 summarizes the estimated emission strengths derived from different products based on different a priori knowledge of inventories and wind information at different altitudes (for specific values see Table A- 1). Different a priori inventories result in 16%-32% changes in strength at different altitudes, which is generally smaller than the 47% difference in the total amount of inventories (9.7E26 for CAMS-GLOB-ANT and 6.6E26 molec./s for CoMet inventory). This is probably due to the different locations of sources and different proportions of each emission source in the total strengths in the two inventories.

When using the CAMS-GLOB-ANT inventory, CH₄ emission rates derived from CAMS XCH₄ and TXCH₄ are ~37% and ~56% higher than those derived from TROPOMI XCH₄ and IASI+TROPOMI TXCH₄, respectively. This difference is mainly due to the difference between the CAMS forecast and satellite products. The strength increases with respect to the increasing wind speed at higher altitude. Whereas the increment is not always proportional to the wind speed, i.e., less increase in the strength with respect to the wind speed at higher altitude (see Sect. 3.3.1).





Figure 7: Similar to Figure 5, but for (a-c) TROPOMI XCH₄ and (d-f) TROPOMI+IASI TXCH₄. The a priori knowledge of sources are based on the CoMet inventory (6.6E26 molec./s in total, Galkowski et al., 2021). The triangle symbols represent the locations of the active coal mine shafts and different colors denote the amount of emission rates.



335 **Figure 8: Estimated CH₄ emission rates derived from the CAMS forecasts (XCH₄ and TXCH₄), TROPOMI XCH₄, and TROPOMI+IASI TXCH₄ data based on different a priori knowledge of emission sources (CAMS-GLOB-ANT and CoMet inventories) and on ERA5 model winds at different altitudes (10 m, 330 m, 500 m). Square symbols represent the a priori emission sources from the CAMS-GLOB-ANT inventory and triangle symbols represent the a priori emission sources from the CoMet inventory. The two horizontal lines represent the number of total emissions for the CAMS-GLOB-ANT inventory (lavender color) and for the CoMet inventory (orange color), respectively. Note, error bars represent the uncertainties from background removal and noise in the data, which are much smaller than the results and they are not visible here. For specific values see Table A- 1.**

340

3.3 Uncertainty analysis

CH₄ signal is weak compared to the background concentration which shows an increasing trend with obvious seasonality and strong day-to-day signals. It is necessary to remove the background signals before estimating the emission strengths. However, the imperfect elimination of the background introduces uncertainties, which can be determined by considering the deficits of the background model and the noise in the background (Tu et al., 2022). In this study, the uncertainties of the estimated strengths include the background uncertainties.

Winds, particularly near the surface, are significantly altered by topography, which yields uncertainties in knowing the transport pathway from emission sources to the measurement location (Chen et al., 2016; Babenhauserheide et al., 2020). Thus, wind is one of the most important factors in correctly estimating the emission rates. Here, we investigate the wind uncertainties based on the CAMS XCH₄ and the CAMS-GLOB-ANT inventory. The wind used in Sect. 3.3.2 and 3.3.3 are from ERA5 at 10 m.

3.3.1 Vertical wind shear

Compared to the wind at 330 m, the distributions of wind directions are similar at lower or higher altitudes (10 m and 500 m) but the wind speed increases with higher altitude (Figure 3). The wind speed at 10 m is 20% weaker than that at 330 m (Table 1), which yields a corresponding lower emission estimate of 613 ± 13 kt/year ($7.3E26 \pm 1.5E25$ molec./s, -25%) based on the CAMS XCH₄ and CAMS emission inventory (Figure A- 4a).

Assuming that the height of the Planetary Boundary Layer (PBL) is typically less than a kilometer, we use the ERA5 wind at 500 m above the ground (Figure 3c) for describing the transport of methane released in the study region. The wind speed at 500 m increases by 22% and 37% for NE and SW regimes, respectively, i.e., 32% on average, compared to the wind at 330 m. The share of SW directed winds is slightly larger at the 500 m level. These differences result in an increase of 13% of the estimated emission rate (924 ± 19 kt/year, $1.1E27 \pm 2.3E25$ molec./s). The wind speed is linear in the calculation of ϵ (Eq. 2), but the wind speeds do not all linearly change for each grid and for each time at different levels. This results in unequal changes between the wind speed and the enhanced columns, and later unequal changes in the estimated emission strength. In addition, the simple cone plume model introduces biases, i.e., the enhanced column in the downwind is set to zero when its location is out of the cone angle (60°). Slight changes in the wind directions might result in a huge difference in the enhanced columns.

3.3.2 Use of narrowed angular wind regimes

The long-term wind comes from all directions (0° - 360°) (Figure 3). To define the uncertainty of wind regimes' coverage, the wind is separated into two groups with narrow coverage fields: $NE_{1/4}$ (0° - 90°) – $SW_{1/4}$ (180° - 270°) and $NW_{1/4}$ (270° - 360°) – $SE_{1/4}$ (90° - 180°). The final estimated emission strength is weighted by the number of days on which, on average, the wind blew in the respective wind regime (i.e., 115 days for $NE_{1/4}$ – $SW_{1/4}$ and 71 days for $NW_{1/4}$ – $SE_{1/4}$, respectively). The XCH₄ anomalies and the plume for the $NE_{1/4}$ – $SW_{1/4}$ regime are quite similar to those with using wider-coverage NE and SW fields (Figure 9a-c). The wind-assigned anomalies derived from CAMS and the [simple](#) cone plume model show very good agreement as well. Slightly less data points are found here because of the choice of narrower wind fields, especially for $NW_{1/4}$ – $SE_{1/4}$ wind fields. The estimated emission rate is about 773 ± 19 kt/year ($9.2E26 \pm 2.3E25$ molec./s) for the $NE_{1/4}$ – $SW_{1/4}$ field. This indicates that the effect of the segment in the wind field coverage is negligible when there are enough measurements. The use of $NW_{1/4}$ – $SE_{1/4}$ wind fields yields an emission strength of 1176 ± 134 kt/year ($1.4E27 \pm 1.6E26$ molec./s). The higher uncertainty is probably due to less measurements in these wind fields. The weighted rate is therefore about 927 kt/year ($1.1E27$ molec./s), 13.4% higher than based on the wider NE-SW wind regime (Sec. 3.1).




Figure 9: Similar figures to Figure 5b-c. Results are derived from CAMS XCH₄, CAMS emission inventory and ERA5 wind at 330 m for (a)-(b) narrow wind coverage ($NE_{1/4}$ and $SW_{1/4}$), and (c)-(d) narrow wind coverage ($NW_{1/4}$ and $SE_{1/4}$).

3.3.3 Investigation of different choices for wind field segmentation

The wind category here is based on the predominant wind fields over the USCB region and is divided into two opposite regimes (SW and NE). To investigate the effect of the segmentation on the uncertainty in the emission rate estimation, we additionally apply another kind of segmentation: N ($<90^\circ$ or $>270^\circ$) and S ($90^\circ - 270^\circ$) categories. Similar results are found and are shown in Figure 10. Though the 2D distribution of the plume changes due to the new wind category, an obvious plume can be seen. The estimated emission rate is 773 ± 19 kt/year ($9.2E26 \pm 2.3E25$ molec./s), which is only 5.2% less than that using NE and SW wind categories. The correlation of the wind-assigned anomalies derived from the CAMS and the simple cone plume model shows a very good agreement as well, with a similar R^2 value of 0.85 to that in the NE-SW wind category. This result demonstrates that our method is not significantly influenced by the wind regime division.




Figure 10: Similar figures to Figure 5b-c. Results are derived from CAMS XCH₄, CAMS emission inventory and ERA5 wind at 330 m but using a new wind category (N and S).

3.3.4 Investigation of different choices for angle of the emission cone

The angle ($\alpha = 60^\circ$) used in the simple cone plume model is an empirical value which affects the deduced emission strengths. Figure 11 shows the results when α is decreased or increased by 10° . Changes in the spatial distributions of wind-assigned anomalies and in the correlations derived from CAMS and the simple cone plume model are nearly negligible when using different angles in the model. The estimated emissions are 789 ± 16 kt/a ($9.5E26 \pm 1.9E25$ molec./s) for $\alpha = 50^\circ$ and 832 ± 17 kt/a ($9.9E26 \pm 2.0E25$ molec./s) for $\alpha = 70^\circ$, which are 3% lower and 2% higher than that with using the empirical angle ($\alpha = 60^\circ$).




Figure 11: Similar figures to Figure 5b-c. Results are derived from CAMS XCH₄, CAMS emission inventory and ERA5 wind at 330 m for (a)-(b) $\alpha = 50^\circ$, and (c)-(d) $\alpha = 70^\circ$.

The changes in the estimated emission rates for different products due to different error sources are summarized in Table A- 2. Based on the error propagation, the total uncertainty in the estimated emission rates from the different error sources (background removal and noise in the data, vertical wind shear at 500 m, wind field segmentation, and opening angle $\alpha = 70^\circ$) is approximately 14.7% for CAMS XCH₄, 14.8% for TROPOMI XCH₄ and 11.4% for TROPOMI+IASI TXCH₄. Note that, the use of narrowed angular wind regimes is not a preferable way due to few amounts of data in narrowed wind regimes and thus, is not considered an error source. In addition, the 500 m wind shear was used as a contribution to the budget, as the 10 m wind is not expected to be representative of the PBL.

4. Conclusion

Intensive mining activities are the dominant CH₄ emission sources in the USCB region, Poland, where one of the largest coal mining areas in Europe is located. It is thus of importance to quantify the CH₄ emissions from this area. In this study we use the combination of a simple cone plume model and a novel wind-assigned model to estimate CH₄ emission rates from high-resolution CAMS forecast XCH₄ and TXCH₄, along with satellite data (TROPOMI XCH₄ and TROPOMI+IASI TXCH₄) over the USCB region (49.3°N-50.8°N and 18°E-20°E) from November 2017 to December 2020.

Based on the CAMS-GLOB-ANT inventory, the dominant CH₄ source is emitted from energy production and distribution, 420 and the significant sources are spread around the city of Katowice and its southwest region. We firstly apply the wind-assigned method to the CAMS forecasts based on the a priori knowledge of CAMS-GLOB-ANT inventory (815 kt/year, 9.7E26 molec./s in total) and ERA5 wind at ~330 m. We use the wind-assigned anomalies of XCH₄/TXCH₄ to represent the difference of XCH₄/TXCH₄ between the conditions of two opposite wind fields (NE and SW). The wind-assigned anomalies derived from CAMS XCH₄/TXCH₄ show very good agreements with the output of the [simple](#) cone plume model with an R² value of 0.85 425 for CAMS XCH₄ and CAMS TXCH₄. This nice correlation indicates that our background removal works well. In addition, similar estimates are derived from CAMS XCH₄ (815 kt/year, [i.e.](#), 9.7E26 E25 molec./s) and TXCH₄ (798 kt/year, [i.e.](#), 9.5E26 molec./s).

To investigate the CH₄ emissions from this hot spot, the CoMet campaign was performed in 2018. Locations and emission 430 rates of the ventilation shafts of the coal mine used in this study are based on this campaign. Based on this knowledge, the emissions are estimated as 496 kt/year (5.9E26 molec./s) and 437 kt/year (5.2E26 molec./s) from the TROPOMI XCH₄ and combined TROPOMI+IASI TXCH₄, respectively. These results are 40% less than that derived from the CAMS model and 435 CAMS-GLOB-ANT inventory. It is probably because the CAMS-GLOB-ANT includes many sectors of anthropogenic sources, like wastes, and combustion from residential and commercial, which account for about 20%. Nevertheless, our results derived from satellite observations are close to the E-PRTR inventory of 448 kt/year and reasonably compared to the CoMet inventory (555 kt/year), and to previous studies over the USCB region (ranging from 9 kt/year to 79 kt/year for a sub-cluster 440 of shafts (Krautwurst et al., 2021) up to 477 kt/year derived from one flight (Fiehn et al. (2021)).

Similar 2D anomalies and plumes are also observed for TROPOMI XCH₄ and TROPOMI+IASI TXCH₄. This nicely documents the robustness of the method. The TROPOMI+IASI result has a higher uncertainty than the TROPOMI result, 445 because (1) the vertical distribution of CH₄ is in general much more difficult to measure than the total column of CH₄ and (2) the vertical distribution is derived by considering two independent measurements, each with its own noise error. This might change for a larger number of data points (e.g., by using data from more years or by applying the method to IASI and TROPOMI successors on the upcoming METOP-SG satellite, which offers much more collocated observations). Nonetheless, the uncertainties are insignificant compared to the estimated emission rates.

Wind contains uncertainties in knowing the transport pathway from emission sources to the measurement location and thus, 450 we analyze the effects in selecting wind at lower and higher [altitudes](#) (10 m and 500 m), wind field [coverage](#), and wind category. Wind distributions at higher levels are similar to the ones at 330 m. However, their speeds decrease by 20% at 10 m and increase by 32% at 500 m, which results in changes in the emission rates by -25% and 13 % [for CAMS XCH₄](#), respectively. Narrower wind field coverage (0°-90° for NE regime and 180°-270° for SW regime) and different wind segmentation (<90° or >270° for N regime and 90°-270° for S regime) introduce uncertainties of +13.4% and -5.2% [for CAMS XCH₄](#), respectively. The agreements for these sensitivity tests of the wind-assigned anomalies derived from the CAMS and from the [simple](#) cone plume model are as good as that using previous NE and SW wind fields. [The impact of a suboptimal choice for the angle \(60°\) used in the simple cone plume model is also discussed. The estimation is decreased by 3% for \$\alpha = 50^\circ\$ and increased by 2%](#)

455 for $\alpha = 70^\circ$ for CAMS XCH₄. This small change supports the empirical choice for α . Based on the error propagation, the total uncertainty in the estimated emission rates from the different error sources (background removal and noise in the data, vertical wind shear at 500 m, wind field segmentation, and opening angle $\alpha = 70^\circ$) is approximately 14.7% for CAMS XCH₄, 14.8% for TROPOMI XCH₄ and 11.4% for TROPOMI+IASI TXCH₄. These results suggest that the wind-assigned method is robust and is also suitable for estimating CH₄ and CO₂ emissions in other regions.

460 *Data availability.* The data are accessible by contacting the corresponding author (qiansi.tu@kit.edu). The SRON S5P-RemoTeC scientific TROPOMI CH₄ dataset from this study is available for download at <https://doi.org/10.5281/zenodo.4447228> (Lorente et al., 2021, last access: 08 November 2021). The TROPOMI data set is publicly available from <https://scihub.copernicus.eu/> (last access: 08 November 2021; ESA, 2020). The access and use of any Copernicus Sentinel data available through the Copernicus Open Access Hub are governed by the legal notice on the use of Copernicus Sentinel Data and Service Information, which is given here: 465 https://sentinels.copernicus.eu/documents/247904/690755/Sentinel_Data_Legal_Note (last access: 08 November 2021; European Commission, 2020). The MUSICA IASI data set is available for download via <https://doi.org/10.35097/408> (Schneider et al. 2021).

470 *Author contributions.* Qiansi Tu, Frank Hase and Matthias Schneider developed the research question. Qiansi Tu wrote the manuscript and performed the data analysis with input from Frank Hase, Matthias Schneider and Farahnaz Khosrawi. Matthias Schneider, Benjamin Ertl and Christopher J. Diekmann provided the combined (MUSICA IASI + TROPOMI) data and supported technically for the analysis of these data. Jarosław Necki supported in consultation of the local situation and CoMet inventory. All authors discussed the results and contributed to the final manuscript.

475 *Competing interests.* The authors declare that they have no conflict of interest.

480 *Acknowledgements.* The CAMS results were generated using Copernicus Atmosphere Monitoring Service (2017–2020) information. Neither the European Commission nor ECMWF is responsible for any use that may be made of the Copernicus information or data it contains. We also thank Michela Giusti and Kevin Marsh in the Data Support Team at ECMWF for retrieving and providing comments about the CAMS data. This research has largely benefit from funds of the Deutsche Forschungsgemeinschaft (provided for the two projects MOTIV and TEDDY with IDs/290612604 and 416767181, respectively). Important part of this work was performed on the supercomputer ForHLR funded by the Ministry of Science, Research and the Arts Baden-Württemberg and by the German Federal Ministry of Education and Research. We acknowledge Emissions of atmospheric Compounds and Compilation of Ancillary Data (ECCAD) for providing CAMS-GLOB-ANT 485 inventory data. We also give thanks to Claire Granier from Laboratoire d'Aérologie, Toulouse, France for providing information about the uncertainties of the CAMS-GLOB-ANT inventory.

We acknowledge the support by the Deutsche Forschungsgemeinschaft and the Open Access Publishing Fund of the Karlsruhe Institute of Technology.

490

Appendix




Figure A- 1: Sketch of the simple **cone** plume model used to explain the CH_4 emission estimation method. The methane at the point source is distributed along the wind direction (wind speed: v) in the cone-shaped area with an opening angle of α . The point source emits the methane at an emission rate of ϵ . We assumed the methane molecules are evenly distributed in the dotted area A, and the distance from area A to the point source is d . Therefore, the emitted methane in dt time period equals to the amount of methane in the area A. It yields the equation $\epsilon \times dt \approx \Delta \text{column} \times \frac{\alpha}{\pi} \times \pi \times d \times v \times dt$. This figure is adopted from Tu et al. (2022).




Figure A- 2: A zoomed figure of Figure 4(b).




Figure A- 3: The enhancements for wind coming from NE and SW.




Figure A- 4: Similar to Figure 5 but using ERA5 wind at 10 m.

505

Table A- 1: Estimated CH₄ emission rates derived from CAMS forecasts (XCH₄ and TXCH₄), TROPOMI XCH₄, and IASI&TROPOMI TXCH₄ data based on different a priori knowledge of emission sources (CAMS-GLOB-ANT and CoMet campaign inventories) and ERA5 model winds at different altitudes (10 m, 100 m and ~500 m).

	ERA5 wind at 10 m		ERA5 wind at 330 m (975 hPa)		ERA5 wind at 500 m (950 hPa)	
prior emission sources	CAMS-GLOB-ANT (total = 9.7E26 molec./s)	CoMet inventory (total = 6.6E26 molec./s)	CAMS-GLOB-ANT (total = 9.7E26 molec./s)	CoMet inventory (total = 6.6E26 molec./s)	CAMS-GLOB-ANT (total = 9.7E26 molec./s)	CoMet inventory (total = 6.6E26 molec./s)
CAMS XCH ₄	7.3E26 ± 1.5E25	6.3E26 ± 1.3E25	9.7E26 ± 2.0E25	8.3E26 ± 1.8E25	1.1E27 ± 2.3E25	9.2E26 ± 2.1E25
CAMS TXCH ₄	7.2E26 ± 1.3E25	6.1E26 ± 1.2E25	9.5E26 ± 1.8E25	8.1E26 ± 1.6E25	1.0E27 ± 2.1E25	8.9E26 ± 1.9E25
TROPOMI XCH ₄	5.4E26 ± 1.8E25	4.5E26 ± 1.5E25	7.1E26 ± 2.4E25	5.9E26 ± 2.1E25	8.1E26 ± 2.8E25	6.3E26 ± 2.3E25

IASI&TROPOMI TXCH ₄	4.7E26 ± 2.7 E25	4.0E26 ± 2.3 E25	6.2E24 ± 3.7 E25	5.2E26 ± 3.2 E25	6.8E24 ± 4.3 E25	5.5E26 ± 3.6 E25
-----------------------------------	-------------------------------------	-------------------------------------	-------------------------------------	-------------------------------------	-------------------------------------	-------------------------------------

510 [Table A- 2: Changes in the estimated emission rates for different products when using different input data or under different situations compared to their results using the default setting \(wind at 330 m, NE-SW wind segmentation, \$\alpha =60^\circ\$, CAMS-GLOB-ANT for CAMS XCH₄ and CoMet inventory for TROPOMI XCH₄ and TROPOMI+IASI TXCH₄\).](#)

	CAMS XCH₄	TROPOMI XCH₄	TROPOMI+IASI TXCH₄
Background removal & noise in the data	2.1%	3.6%	6.1%
Vertical wind shear (500 m)	13.4%	6.8%	5.8%
wind field segmentation (N-S)	-5.2%	12.7%	7.7%
angle of the emission cone ($\alpha=70^\circ$)	2.1%	0.07%	-0.02%
Total:	14.7%	14.8%	11.4%

Agustí-Panareda, A., Diamantakis, M., Massart, S., Chevallier, F., Muñoz-Sabater, J., Barré, J., Curcoll, R., Engelen, R., Langerock, B., Law, R. M., Loh, Z., Morguí, J. A., Parrington, M., Peuch, V.-H., Ramonet, M., Roehl, C., Vermeulen, A. T., Warneke, T., and Wunch, D.: Modelling CO₂ weather – why horizontal resolution matters, *Atmos. Chem. Phys.*, 19, 7347–7376, <https://doi.org/10.5194/acp-19-7347-2019>, 2019.

520 Andersen, T., Vinkovic, K., de Vries, M., Kers, B., Necki, J., Swolkien, J., Roiger, A., Peters, W. and Chen, H.: Quantifying methane emissions from coal mining ventilation shafts using an unmanned aerial vehicle (UAV)-based active AirCore system, *Atmos. Environ. X*, 12, 100135, doi:<https://doi.org/10.1016/j.aeaoa.2021.100135>, 2021.

Babenhauserheide, A., Hase, F., and Morino, I.: Net CO₂ fossil fuel emissions of Tokyo estimated directly from measurements of the Tsukuba TCCON site and radiosondes, *Atmos. Meas. Tech.*, 13, 2697–2710, <https://doi.org/10.5194/amt-13-2697-2020>, 2020.

525 Barré, J., Aben, I., Agustí-Panareda, A., Balsamo, G., Bouserez, N., Dueben, P., Engelen, R., Inness, A., Lorente, A., McNorton, J., Peuch, V.-H., Radnoti, G., and Ribas, R.: Systematic detection of local CH₄ anomalies by combining satellite measurements with high-resolution forecasts, *Atmos. Chem. Phys.*, 21, 5117–5136, <https://doi.org/10.5194/acp-21-5117-2021>, 2021.

530 Boesch, H., Baker, D., Connor, B., Crisp, D. and Miller, C.: Global Characterization of CO₂ Column Retrievals from Shortwave-Infrared Satellite Observations of the Orbiting Carbon Observatory-2 Mission, *Remote Sens.*, 3(2), doi:10.3390/rs3020270, 2011.

Butz, A., Hasekamp, O. P., Frankenberg, C., and Aben, I.: Retrievals of atmospheric CO₂ from simulated space-borne measurements of backscattered near-infrared sunlight: accounting for aerosol effects, *Appl. Opt.* 48, 3322–3336, 535 2009.

Butz, A., Hasekamp, O. P., Frankenberg, C., Vidot, J. and Aben, I.: CH₄ retrievals from space-based solar backscatter measurements: Performance evaluation against simulated aerosol and cirrus loaded scenes, *J. Geophys. Res. Atmos.*, 115(D24), doi:<https://doi.org/10.1029/2010JD014514>, 2010.

540 Butz, A., Guerlet, S., Hasekamp, O., Schepers, D., Galli, A., Aben, I., Frankenberg, C., Hartmann, J.-M., Tran, H., Kuze, A., Keppel-Aleks, G., Toon, G., Wunch, D., Wennberg, P., Deutscher, N., Griffith, D., Macatangay, R., Messerschmidt, J., Notholt, J., and Warneke, T.: Toward accurate CO₂ and CH₄ observations from GOSAT, *Geophysical Research Letters*, 38, <https://doi.org/10.1029/2011GL047888>, 2011. <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2011GL047888>.

545 Chen, J., Viatte, C., Hedenius, J. K., Jones, T., Franklin, J. E., Parker, H., Gottlieb, E. W., Wennberg, P. O., Dubey, M. K., and Wofsy, S. C.: Differential column measurements using compact solar-tracking spectrometers, *Atmos. Chem. Phys.*, 16, 8479–8498, <https://doi.org/10.5194/acp-16-8479-2016>, 2016.

Collard, A. D.: Selection of IASI Channels for Use in Numerical Weather Prediction, ECMWF, <https://www.ecmwf.int/node/8760>, 2007.

Coopmann, O., Guidard, V., Fourrié, N., Josse, B., and Marécal, V.: Update of Infrared Atmospheric Sounding Interferometer (IASI) channel selection with correlated observation errors for numerical weather prediction (NWP), *Atmos. Meas. Tech.*, 13, 2659–2680, <https://doi.org/10.5194/amt-13-2659-2020>, 2020.

Copernicus Climate Change Service (C3S): ERA5: Fifth generation of ECMWF atmospheric reanalyses of the global climate. Copernicus Climate Change Service Climate Data Store (CDS), date of access. <https://cds.climate.copernicus.eu/cdsapp#!/home>, 2017.

Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Dentener, F., van Aardenne, J. A., Monni, S., Doering, U., Olivier, J. G. J., Pagliari, V., and Janssens-Maenhout, G.: Gridded emissions of air pollutants for the period 1970–2012 within EDGAR v4.3.2, *Earth Syst. Sci. Data*, 10, 1987–2013, <https://doi.org/10.5194/essd-10-1987-2018>, 2018.

Diekmann, C. J., Schneider, M., Ertl, B., Hase, F., García, O., Khosrawi, F., Sepúlveda, E., Knippertz, P., and Braesicke, P.: The global and multi-annual MUSICA IASI {H₂O, δD} pair dataset, *Earth Syst. Sci. Data*, 13, 5273–5292, <https://doi.org/10.5194/essd-13-5273-2021>, 2021.

Fiehn, A., Kostinek, J., Eckl, M., Klausner, T., Gałkowski, M., Chen, J., Gerbig, C., Röckmann, T., Maazallahi, H., Schmidt, M., Korbeń, P., Nećki, J., Jagoda, P., Wildmann, N., Mallaun, C., Bun, R., Nickl, A.-L., Jöckel, P., Fix, A., and Roiger, A.: Estimating CH₄, CO₂ and CO emissions from coal mining and industrial activities in the Upper Silesian Coal Basin using an aircraft-based mass balance approach, *Atmos. Chem. Phys.*, 20, 12675–12695, <https://doi.org/10.5194/acp-20-12675-2020>, 2020.

Flemming, J., Huijnen, V., Arteta, J., Bechtold, P., Beljaars, A., Blechschmidt, A.-M., Diamantakis, M., Engelen, R. J., Gaudel, A., Inness, A., Jones, L., Josse, B., Katragkou, E., Marecal, V., Peuch, V.-H., Richter, A., Schultz, M. G., Stein, O., and Tsikerdekis, A.: Tropospheric chemistry in the Integrated Forecasting System of ECMWF, *Geosci. Model Dev.*, 8, 975–1003, <https://doi.org/10.5194/gmd-8-975-2015>, 2015.

De Gouw, J. A., Veefkind, J. P., Roosenbrand, E., Dix, B., Lin, J. C., Landgraf, J., and Levelt, P. F.: Daily Satellite Observations of Methane from Oil and Gas Production Regions in the United States, *Scientific Reports*, 10, 1379, <https://doi.org/10.1038/s41598-020-57678-4>, 2020.

Gałkowski, M., Fiehn, A., Swolkien, J., Stanisavljevic, M., Korben, P., Menoud, M., Necki, J., Roiger, A., Röckmann, T., Gerbig, C., & Fix, A.: Emissions of CH₄ and CO₂ over the Upper Silesian Coal Basin (Poland) and its vicinity (4.01) [Data set]. ICOS ERIC - Carbon Portal. <https://doi.org/10.18160/3K6Z-4H73>, 2021.

[Gałkowski, M., Jordan, A., Rothe, M., Marshall, J., Koch, F.-T., Chen, J., Agusti-Panareda, A., Fix, A., and Gerbig, C.: In situ observations of greenhouse gases over Europe during the CoMet 1.0 campaign aboard the HALO aircraft, *Atmos. Meas. Tech.*, 14, 1525–1544, <https://doi.org/10.5194/amt-14-1525-2021>, 2021.](#)

García, O. E., Schneider, M., Ertl, B., Sepúlveda, E., Borger, C., Diekmann, C., Wiegele, A., Hase, F., Barthlott, S., Blumenstock, T., Raffalski, U., Gómez-Peláez, A., Steinbacher, M., Ries, L., and de Frutos, A. M.: The MUSICA

IASI CH4 and N2O products and their comparison to HIPPO, GAW and NDACC FTIR references, *Atmos. Meas. Tech.*, 11, 4171–4215, <https://doi.org/10.5194/amt-11-4171-2018>, 2018.

585 Granier, C., S. Darras, H. Denier van der Gon, J. Doubalova, N. Elguindi, B. Galle, M. Gauss, M. Guevara, J.-P. Jalkanen, J. Kuenen, C. Lioussse, B. Quack, D. Simpson, K. Sindelarova, The Copernicus Atmosphere Monitoring Service global and regional emissions (April 2019 version), Copernicus Atmosphere Monitoring Service (CAMS) report, doi:10.24380/d0bn-kx16, 2019.

Hasekamp, O. P., and Butz, A.: Efficient calculation of intensity and polarization spectra in vertically inhomogeneous scattering and absorbing atmospheres, *J. Geophys. Res.*, 113, D20309, doi:10.1029/2008JD010379, 2008.

590 Hasekamp, O., Lorente, A., Hu, H., Butz, A., aan de Brugh, J., and Landgraf, J.: Algorithm Theoretical Baseline Document for Sentinel-5 Precursor methane retrieval, <http://www.tropomi.eu/documents/atbd/>, 2019.

Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellán, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., 595 Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S. and Thépaut, J.-N.: The ERA5 global reanalysis, *Q. J. R. Meteorol. Soc.*, 146(730), 1999–2049, doi:<https://doi.org/10.1002/qj.3803>, 2020.

Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M., Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J.-I., Li, M., Liu, L., Lu, Z., Moura, M. C. P., O'Rourke, P. R., and Zhang, Q.: Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from 600 the Community Emissions Data System (CEDS), *Geosci. Model Dev.*, 11, 369–408, <https://doi.org/10.5194/gmd-11-369-2018>, 2018.

Kirschke, S., Bousquet, P., Ciais, P., Saunois, M., Canadell, J. G., Dlugokencky, E. J., Bergamaschi, P., Bergmann, D., Blake, D. R., Bruhwiler, L., Cameron-Smith, P., Castaldi, S., Chevallier, F., Feng, L., Fraser, A., Heimann, M., Hodson, E., Houweling, S., Josse, B., Fraser, P. J., Krummel, P. B., Lamarque, J.-F., Langenfelds, R. L., Le Quéré, C., Naik, V., O'Doherty, S., Palmer, P. I., Pison, I., Plummer, D., Poulter, B., Prinn, R. G., Rigby, M., Ringeval, B., Santini, M., Schmidt, M., Shindell, D. T., Simpson, I. J., Spahni, R., Steele, L. P., Strode, S. A., Sudo, K., Szopa, S., van der Werf, G. R., Voulgarakis, A., van Weele, M., Weiss, R. F., Williams, J. E. and Zeng, G.: Three decades of global 605 methane sources and sinks, *Nat. Geosci.*, 6(10), 813–823, doi:10.1038/ngeo1955, 2013.

Kostinek, J., Roiger, A., Eckl, M., Fiehn, A., Luther, A., Wildmann, N., Klausner, T., Fix, A., Knote, C., Stohl, A., and Butz, 610 A.: Estimating Upper Silesian coal mine methane emissions from airborne in situ observations and dispersion modeling, *Atmos. Chem. Phys.*, 21, 8791–8807, <https://doi.org/10.5194/acp-21-8791-2021>, 2021.

Krautwurst, S., Gerilowski, K., Borchardt, J., Wildmann, N., Gałkowski, M., Swolkień, J., Marshall, J., Fiehn, A., Roiger, A., Ruhtz, T., Gerbig, C., Necki, J., Burrows, J. P., Fix, A., and Bovensmann, H.: Quantification of CH₄ coal mining emissions in Upper Silesia by passive airborne remote sensing observations with the Methane Airborne MAPper

(MAMAP) instrument during the CO₂ and Methane (CoMet) campaign, *Atmos. Chem. Phys.*, 21, 17345–17371, <https://doi.org/10.5194/acp-21-17345-2021>, 2021.

Landgraf, J., Butz, A., Hasekamp, O., Hu, H., and aan de Brugh, J.: Sentinel 5 L2 Prototype Processors, Algorithm Theoretical Baseline Document: Methane Retrieval, 2019.

620 Liu, M., van der A, R., van Weele, M., Eskes, H., Lu, X., Veefkind, P., de Laat, J., Kong, H., Wang, J., Sun, J., Ding, J., Zhao, Y. and Weng, H.: A New Divergence Method to Quantify Methane Emissions Using Observations of Sentinel-5P TROPOMI, *Geophys. Res. Lett.*, 48(18), e2021GL094151, doi:<https://doi.org/10.1029/2021GL094151>, 2021.

625 Lorente, A., Borsdorff, T., Butz, A., Hasekamp, O., aan de Brugh, J., Schneider, A., Wu, L., Hase, F., Kivi, R., Wunch, D., Pollard, D. F., Shiomi, K., Deutscher, N. M., Velazco, V. A., Roehl, C. M., Wennberg, P. O., Warneke, T., and Landgraf, J.: Methane retrieved from TROPOMI: improvement of the data product and validation of the first 2 years of measurements, *Atmos. Meas. Tech.*, 14, 665–684, <https://doi.org/10.5194/amt-14-665-2021>, 2021.

630 Luther, A., Kleinschek, R., Scheidweiler, L., Defratyka, S., Stanisavljevic, M., Forstmaier, A., Dandoci, A., Wolff, S., Dubravica, D., Wildmann, N., Kostinek, J., Jöckel, P., Nickl, A.-L., Klausner, T., Hase, F., Frey, M., Chen, J., Dietrich, F., Nečki, J., Swolkień, J., Fix, A., Roiger, A., and Butz, A.: Quantifying C₄ emissions from hard coal mines using mobile sun-viewing Fourier transform spectrometry, *Atmos. Meas. Tech.*, 12, 5217–5230, <https://doi.org/10.5194/amt-12-5217-2019>, 2019.

635 Luther, A., Kostinek, J., Kleinschek, R., Defratyka, S., Stanisavljevic, M., Forstmaier, A., Dandoci, A., Scheidweiler, L., Dubravica, D., Wildmann, N., Hase, F., Frey, M. M., Chen, J., Dietrich, F., Nečki, J., Swolkien, J., Knote, C., Vardag, S. N., Roiger, A., and Butz, A.: Observational constraints on methane emissions from Polish coal mines using a ground-based remote sensing network, *Atmos. Chem. Phys. Discuss. [preprint]*, <https://doi.org/10.5194/acp-2021-978>, in review, 2021.

640 Luther, A., Kostinek, J., Kleinschek, R., Defratyka, S., Stanisavljević, M., Forstmaier, A., Dandoci, A., Scheidweiler, L., Dubravica, D., Wildmann, N., Hase, F., Frey, M. M., Chen, J., Dietrich, F., Nečki, J., Swolkień, J., Knote, C., Vardag, S. N., Roiger, A., and Butz, A.: Observational constraints on methane emissions from Polish coal mines using a ground-based remote sensing network, *Atmos. Chem. Phys.*, 22, 5859–5876, <https://doi.org/10.5194/acp-22-5859-2022>, 2022.

Pandey, S., Houweling, S., Krol, M., Aben, I., Chevallier, F., Dlugokencky, E. J., Gatti, L. V., Gloor, E., Miller, J. B., Detmers, R., Machida, T., and Röckmann, T.: Inverse modeling of GOSAT-retrieved ratios of total column CH₄ and CO₂ for 2009 and 2010, *Atmos. Chem. Phys.*, 16, 5043–5062, <https://doi.org/10.5194/acp-16-5043-2016>, 2016.

645 Pandey, S., Gautam, R., Houweling, S., van der Gon, H. D., Sadavarte, P., Borsdorff, T., Hasekamp, O., Landgraf, J., Tol, P., van Kempen, T., Hoogeveen, R., van Hees, R., Hamburg, S. P., Maasakkers, J. D., and Aben, I.: Satellite observations reveal extreme methane leakage from a natural gas well blowout, *Proc Natl Acad Sci USA*, 116, 26376, <https://doi.org/10.1073/pnas.1908712116>, 2019.

Prather, M. J., Holmes, C. D. and Hsu, J.: Reactive greenhouse gas scenarios: Systematic exploration of uncertainties and the role of atmospheric chemistry, *Geophys. Res. Lett.*, 39(9), doi:<https://doi.org/10.1029/2012GL051440>, 2012.

650 IPCC, 2014: Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]. IPCC, Geneva, Switzerland, 151 pp.

Saunois, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A., Dlugokencky, E. J., Houweling, S., Patra, P. K., Ciais, P., Arora, V. K., Bastviken, D., Bergamaschi, P., Blake, D. R., Brailsford, G., Bruhwiler, L., Carlson, K. M., Carroll, M., Castaldi, S., Chandra, N., Crevoisier, C., Crill, P. M., Covey, K., Curry, C. L., Etiope, G., Frankenberg, C., Gedney, N., Hegglin, M. I., Höglund-Isaksson, L., Hugelius, G., Ishizawa, M., Ito, A., Janssens-Maenhout, G., Jensen, K. M., Joos, F., Kleinen, T., Krummel, P. B., Langenfelds, R. L., Laruelle, G. G., Liu, L., Machida, T., Maksyutov, S., McDonald, K. C., McNorton, J., Miller, P. A., Melton, J. R., Morino, I., Müller, J., Murguia-Flores, F., Naik, V., Niwa, Y., Noce, S., O'Doherty, S., Parker, R. J., Peng, C., Peng, S., Peters, G. P., Prigent, C., Prinn, R., Ramonet, M., Regnier, P., Riley, W. J., Rosentreter, J. A., Segers, A., Simpson, I. J., Shi, H., Smith, S. J., Steele, L. P., Thornton, B. F., Tian, H., Tohjima, Y., Tubiello, F. N., Tsuruta, A., Viovy, N., Voulgarakis, A., Weber, T. S., van Weele, M., van der Werf, G. R., Weiss, R. F., Worthy, D., Wunch, D., Yin, Y., Yoshida, Y., Zhang, W., Zhang, Z., Zhao, Y., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: The Global Methane Budget 2000–2017, *Earth Syst. Sci. Data*, 12, 1561–1623, <https://doi.org/10.5194/essd-12-1561-2020>, 2020.

660 Schneider, M., Ertl, B., Diekmann, C. J., Khosrawi, F., Röhling, A. N., Hase, F., Dubravica, D., García, O. E., Sepúlveda, E., Borsdorff, T., Landgraf, J., Lorente, A., Chen, H., Kivi, R., Laemmel, T., Ramonet, M., Crevoisier, C., Pernin, J., Steinbacher, M., Meinhardt, F., Deutscher, N. M., Griffith, D. W. T., Velazco, V. A., and Pollard, D. F.: Synergetic use of IASI and TROPOMI space borne sensors for generating a tropospheric methane profile product, *Atmos. Meas. Tech. Discuss.* [preprint], <https://doi.org/10.5194/amt-2021-31>, in review, 2021.

670 Schneider, M., Ertl, B., Diekmann, C. J., Khosrawi, F., Weber, A., Hase, F., Höpfner, M., García, O. E., Sepúlveda, E., and Kinnison, D.: Design and description of the MUSICA IASI full retrieval product, *Earth Syst. Sci. Data*, 14, 709–742, <https://doi.org/10.5194/essd-14-709-2022>, 2022.

Schepers, D., Guerlet, S., Butz, A., Landgraf, J., Frankenberg, C., Hasekamp, O., Blavier, J.-F., Deutscher, N. M., Griffith, D. W. T., Hase, F., Kyro, E., Morino, I., Sherlock, V., Sussmann, R. and Aben, I.: Methane retrievals from Greenhouse Gases Observing Satellite (GOSAT) shortwave infrared measurements: Performance comparison of proxy and physics retrieval algorithms, *J. Geophys. Res. Atmos.*, 117(D10), doi:<https://doi.org/10.1029/2012JD017549>, 2012.

675 Schneising, O., Buchwitz, M., Reuter, M., Vanselow, S., Bovensmann, H., and Burrows, J. P.: Remote sensing of methane leakage from natural gas and petroleum systems revisited, *Atmos. Chem. Phys.*, 20, 9169–9182, <https://doi.org/10.5194/acp-20-9169-2020>, 2020.

680 Tu, Q., Hase, F., Schneider, M., García, O., Blumenstock, T., Borsdorff, T., Frey, M., Khosrawi, F., Lorente, A., Alberti, C., Bustos, J. J., Butz, A., Carreño, V., Cuevas, E., Curcoll, R., Diekmann, C. J., Dubravica, D., Ertl, B., Estruch, C.,

León-Luis, S. F., Marrero, C., Morgui, J.-A., Ramos, R., Scharun, C., Schneider, C., Sepúlveda, E., Toledano, C., and Torres, C.: Quantification of CH₄ emissions from waste disposal sites near the city of Madrid using ground- and space-based observations of COCCON, TROPOMI and IASI, *Atmos. Chem. Phys.*, 22, 295–317, <https://doi.org/10.5194/acp-22-295-2022>, 2022.

685 Varon, D. J., McKeever, J., Jervis, D., Maasakkers, J. D., Pandey, S., Houweling, S., Aben, I., Scarpelli, T. and Jacob, D. J. :
Satellite discovery of anomalously large methane point sources from oil/gas production, *Geophysical Research Letters*, 46, 13507– 13516. <https://doi.org/10.1029/2019GL083798>, 2019.

690 Veefkind, J. P., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., Claas, J., Eskes, H. J., de Haan, J. F., Kleipool, Q.,
van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B.,
Vink, R., Visser, H., and Levelt, P. F.: TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global
observations of the atmospheric composition for climate, air quality and ozone layer applications, *Remote Sens. Environ.*, 120, 70–83, <https://doi.org/10.1016/j.rse.2011.09.027>, 2012.

695 [Wolff, S., Ehret, G., Kiemle, C., Amediek, A., Quatrevaelet, M., Wirth, M., and Fix, A.: Determination of the emission rates of](#)
[CO₂ point sources with airborne lidar, Atmos. Meas. Tech., 14, 2717–2736, https://doi.org/10.5194/amt-14-2717-2021, 2021.](#)

World Meteorological Organisation, WMO greenhouse gas bulletin No. 16.23 Novembet 2020 [Online]. Available at:
<https://public.wmo.int/en/resources/library/wmo-greenhouse-gas-bulletin>, 2020.

700