

Measurement report: High Arctic aerosol hygroscopicity at sub- and supersaturated conditions during spring and summer

Andreas Massling¹, Robert Lange^{1,2}, Jakob Boyd Pernov^{1,3}, Ulrich Gosewinkel¹, Lise-Lotte Sørensen¹, and Henrik Skov¹

5 ¹Department of Environmental Science, iClimate, Aarhus University, 4000 Roskilde, Denmark

²ROCKWOOL Group, 2640 Hedehusene, Denmark

³Extreme Environments Research Laboratory, École Polytechnique Fédérale de Lausanne, 1951 Sion, Switzerland

Correspondence to: Andreas Massling (anma@envs.au.dk)

10

Abstract. Aerosol hygroscopic growth and cloud droplet formation influence the radiation transfer budget of the atmosphere and thereby the climate. In the Arctic, these aerosol properties may have a more pronounced effect on the climate compared to the mid-latitudes. Hygroscopic growth and cloud condensation nuclei (CCN) concentrations of High Arctic aerosols were measured during two field studies in the spring and summer of 2016. The study site was the Villum Research Station (Villum) at Station Nord in the northeastern region of Greenland. Aerosol hygroscopic growth was measured with a hygroscopic tandem differential mobility analyzer (HTDMA) over a total of 23 days, and CCN concentrations were measured over a period of 95 days. Continuous particle number size distributions were recorded, facilitating calculations of aerosol CCN activation diameters and aerosol κ -values. In spring, average CCN concentrations, at supersaturations (SS) of 0.1 to 0.3 %, ranged from 53.7 to 85.3 cm⁻³, with critical activation diameters ranging from 130.2 to 80.2 nm, and κ_{CCN} ranging from 0.28-0.35. In summer, average CCN concentrations were 20.8 to 47.6 cm⁻³, while critical activation diameters and κ_{CCN} were from 137.1 to 76.7 nm and 0.23-0.35, respectively. Mean particle hygroscopic growth factors ranged from 1.60 to 1.75 at 90% relative humidity in spring, while values between 1.47 and 1.67 were observed in summer depending on the initial dry size. Although the summer aerosol number size distributions were characterized by frequent new particle formation events, the CCN population at cloud-relevant supersaturations was determined by accumulation mode aerosols.

15
20
25

1 Introduction:

In the Arctic region, the average temperature is rising about three times as fast as the global average (Lenssen et al., 2019). Aerosols influence the radiation balance of the atmosphere and the surface and thus play a key role in the climatic changes observed in the Arctic (Bellouin et al., 2011; Yang et al., 2014; Chylek et al., 2016). During sunlit seasons, the climatic effects of Arctic aerosols are diverse. Through aerosol-radiation interactions, aerosols scatter incoming shortwave radiation and absorb outgoing longwave radiation (Twomey, 1974). Arctic aerosols also impact clouds by altering cloud lifetime, precipitation rate, and cloud albedo (Zhao and Garrett, 2015). These aerosol-cloud interactions also influence the radiation balance. It has been shown that the Arctic atmosphere is limited in cloud condensation nuclei (CCN) (Mauritsen et al., 2011). This means that a relatively small increase in the concentration of aerosol particles that can function as CCN, can have a comparably large effect on clouds and thus the radiation balance. The radiative forcing (RF) of aerosol-radiation interactions in the Arctic is not well quantified but appears to be a net cooling of the surface (Quinn et al., 2008; Breider et al., 2017).

Aerosol light scattering is highly dependent on particle size and refractive index (Moffet and Prather, 2009; Raut et al., 2009). Particles interact with atmospheric water vapor and potentially grow due to hygroscopicity, which is the ability to uptake water (Pilat and Charlson, 1966; Carrico et al., 2000; Fierz-Schmidhauser et al., 2010). This water uptake at subsaturated conditions changes their light scattering behavior (Liu et al., 2013; Titos et al., 2016). Burgos et al. (2019) found by direct measurements of the aerosol enhancement factor that the effect of water uptake on light scattering is higher for Arctic aerosols compared to other atmospheric reservoirs, which is based on the special interplay between size and hygroscopicity in the Arctic (Zieger et al., 2010). One parameter to deduct from hygroscopicity measurements of aerosols is the κ -value, which provides a theoretical framework for deriving the hygroscopicity for bulk aerosols, assuming an internal mixture (Petters and Kreidenweis, 2007). Knowledge about the hygroscopic growth of Arctic aerosols is crucial for determining the effects of their aerosol-radiation and aerosol-cloud interactions.

Ambient Arctic aerosol size, chemical composition, and number concentrations undergo a strong seasonal cycle (Tunved et al., 2013; Nguyen et al., 2016; Freud et al., 2017). Very few anthropogenic aerosol sources exist within the Arctic. During summer, the Arctic lower troposphere is a largely isolated system (Iversen and Joranger, 1985; Stohl, 2006). Anthropogenic influence from long-range transport is mostly absent, although intrusion from high latitude regions, introducing anthropogenic aerosols, does occur. In summer, the ambient aerosol is mainly of biogenic origin, consisting mostly of biogenic sulfate and organic compounds (Willis et al., 2017). The aerosol is characterized by low particle number concentrations (50-200 cm^{-3}), interrupted by relatively frequent new particle formation (NPF) events which strongly increase the particle number concentration on timescales of hours to days (Nguyen et al., 2016; Freud et al., 2017; Collins et al., 2017; Dall'Osto et al., 2017). During autumn months, the occurrence of NPF decreases, and particle number concentrations reach a minimum. During winter, the polar front moves southward, extending the Arctic lower tropospheric system to include mid-latitude regions with anthropogenic emissions (Shaw et al., 2010). During the winter, inefficient wet deposition coupled with a stably stratified atmosphere allows for pollution to build up forming an increasingly prominent accumulation mode fraction of aerosols. This accumulation mode reaches its maximum number concentration in late spring (March-April) (Matsui et al., 2011; Freud et al., 2017; Lange et al., 2018), a phenomenon known as Arctic haze (Barrie et al., 1981; Shaw, 1995; Quinn et al., 2007). During the Arctic winter and early spring, when sunlight is absent or scarce, it has been demonstrated that especially anthropogenic aerosols have a warming effect due to resulting increases in cloud long-wave emissivity (Garrett et al., 2004; Garrett and Zhao, 2006). During this period, sulfate is the dominating particle component (Heidam et al., 1999; Heidam et al., 2004; Nielsen et al. 2019). Recent findings indicate that sulfate, sea spray aerosol and organic carbon might be internally mixed also in the submicron size range (Kirpes et al., 2018). When temperatures increase around May, the polar dome recedes and wet scavenging becomes an important aerosol sink, consequently, removing the Arctic haze (Engvall et al., 2008; Browse et al., 2012; Croft et al., 2016). This leads to the reestablishment of clean summer conditions.

Concentrations and temporal development of CCN in the Arctic are characterized in several previous studies. Jung et al. (2018), evaluated a multi-year CCN dataset from Zeppelin Mountain, Svalbard, and found CCN concentrations at 0.2 % SS of 70–133 cm^{-3} for the late spring/early summer and 16–57 cm^{-3} for the late summer/early autumn period. Silvergren et al. (2014) evaluated a full CCN seasonal cycle (2007-2008) on Zeppelin Mountain, Svalbard, from re-aerosolized filter

75 sampled aerosol. For supersaturations (SS) between 0.2–0.5 %, they found low CCN concentrations of $< 40 \text{ cm}^{-3}$ for
months September–October, $\sim 40\text{--}80 \text{ cm}^{-3}$ for November–February, $80\text{--}160 \text{ cm}^{-3}$ for March–June and $40\text{--}110 \text{ cm}^{-3}$ for
July–August, with the highest variation between supersaturations in the latter period. In contrast, Zabori et al. (2015)
directly determined κ -values from size-resolved CCN measurements on Svalbard to be between 0.4 and 0.3 at 0.4 % SS
80 on separate occasions in June and August 2008, respectively. A review of long-term studies from around the world,
including Utqiaġvik/Barrow, Alaska from 2007 to 2008, was published by Schmale et al. (2018). CCN concentrations at
0.2 % SS were between 20 and 200 cm^{-3} and CCN particle critical activation diameters ($D_{p,\text{crit}}$) showed a bimodal
distribution, with a smaller mode at $70\text{--}100 \text{ nm}$ and a larger one at $100\text{--}140 \text{ nm}$. The location showed the highest CCN
number concentrations and lowest $D_{p,\text{crit}}$ from February to June. In a study conducted in Northwestern Canada in May
2014, Herenz et al. (2018) found CCN concentrations between 10 and 200 cm^{-3} at $0.1\text{--}0.78 \%$ SS and found a κ -value of
85 0.19 at 0.1% SS and κ -values of $0.21\text{--}0.28$ at $0.2\text{--}0.7 \%$ SS. They detected both well-aged Arctic haze aerosols, as well
as more recently formed aerosols, characteristic of the Arctic summer period. It should be mentioned here, that these
studies concentrated on remote locations in the high Arctic. Properties of Arctic aerosols can be quite diverse depending
on the exact location and time of the year (Schmale et al., 2021; Schmale & Baccharini, 2021; Schmale et al., 2022). Certain
90 parts of the Arctic, e.g., Siberian and North American Arctic, are exposed to high levels of anthropogenic/industrial
activities, while the high/central Arctic often shows different seasonality of aerosol properties compared to the lower
Arctic. The exact location and seasonality also have consequences for the precursor gases of new particle formation
(Schmale and Baccharini, 2021) and their subsequent growth to CCN-relevant sizes.

From late spring to summer, the Arctic aerosol organic fraction changes from having a strong anthropogenic influence to
being mainly biogenic (Nielsen et al., 2019; Willis et al., 2017). Likewise, the total organic fraction increases during
95 summer as well (Chang et al., 2011; Breider et al., 2014). The origin and influence of these organic components on aerosol
CCN ability (Leck et al., 2002) and hygroscopic behavior are currently not well understood and are still the subject of
intense research (Myriokefalitakis et al., 2010; Fu et al., 2013; Willis et al., 2016). An organic marine source has been
proposed to be an important source for primary aerosols, NPF, and secondary growth (Leck and Bigg, 2005; Burkart et
al., 2017; Willis et al., 2017), which contributes significantly to the summertime aerosol mass load. Methanesulfonic acid
100 (MSA) (Nielsen et al. 2019; Dall'Osto et al., 2018b) and organic acids (Mungall et al., 2017), high molecular weight
compounds like proteins (Fu et al., 2015), and polymeric gels (Orellana et al., 2011) likewise play a role. Further,
inorganic non-sea salt sulfate from the oxidation of dimethyl sulfide (DMS) contributes importantly to aerosol mass
during summer and autumn (Rempillo et al., 2011; Leaitch et al., 2013). Surface active organics potentially increase
aerosol CCN activity by lowering droplet surface tension (Lohmann and Leck, 2005). However, studies on non-Arctic
105 marine aerosols at Mace Head have suggested a complex behavior, where biogenic marine particles with low hygroscopic
growth featured high CCN activity (Ovadnevaite et al., 2011). This behavior can potentially be explained by liquid phase
separation, where less-hygroscopic but surface-active components are located in an outer shell limiting hygroscopic
growth at subsaturated conditions. During CCN activation, the phase separation breaks down and the surface-active
compounds enhance CCN activity (Ovadnevaite et al., 2017). This process could also be relevant for Arctic aerosols.
110 However, Lange et al. (2018) found indications that surface activity is not necessary to explain κ -values of Arctic
accumulation mode aerosols.

Longer field studies reporting hygroscopic growth measured by HTDMA instruments in the Arctic are scarce. Zhou et al.
(2001) conducted a shipboard campaign in the Arctic Ocean up to $87^\circ 30' \text{ N}$ during the summer of 1996. They reported
hygroscopic growth factors (HGF) of $1.4\text{--}1.9$ at 90% relative humidity (RH) to be most prominent, however, cases of
115 $\text{HGF} > 1.9$ and $\text{HGF} < 1.4$ were also observed. The higher HGFs probably result from sea-salt aerosol, while the lower
HGF could originate from an organic sea surface source. Using the same measurement series, Silvergren et al. (2014)
determined HGF-derived κ_{HTDMA} -values to be $0.35\text{--}0.5$ in March–August and $0.6\text{--}0.9$ in September–January. During a
ship cruise in July–August 2013, Allan et al. (2015) observed that a possible contribution from iodine oxides lowered the
HGF of 50 nm particles at $90\% \text{ RH}$ to 1.34 , whereas HGF otherwise was consistently > 1.5 .

120 The present study evaluates measurements from two field studies in 2016, where CCN concentrations and HGFs were
measured at the High Arctic site, Villum Research Station (Villum). We present the measurement results and evaluate
which aerosol sources could be responsible for the observed findings at subsaturated and supersaturated conditions. Our

results can directly be used in models of Arctic aerosol–cloud–climate interactions, while simultaneously adding to the understanding of which aerosol sources are important for Arctic CCN concentrations.

125


2 Methods

2.1 Sampling site

All measurements presented in this study were recorded at the Villum Research Station (81° 36' N, 16° 40' W) in northeast Greenland. Its location is shown in Fig. 1. Villum is located at the Danish military station of Station Nord on the Princess Ingeborg Peninsula, about 750 m from the coast of Denmark Fjord, close to the Wandel Sea. This coastal site is within the polar dome year-round and is in close proximity to the Arctic pack ice during both spring and summer (Fig. 1). Currently, the entrance from the Wandel Sea into the Denmark Fjord is covered by sea ice year-round, although in recent years, the waters near the station are unfrozen from August to September. The sampling took place 2 km southwest of the station's main facilities in the "Air Observatory" which provides controlled laboratory conditions for instrument operation. The location is upwind of the main station premises >95 % of the time, pollution influx from the station itself is therefore minimal. Polar sunrise occurs on February 25th, polar day prevails from April 5th to September 3rd and polar sunset occurs on October 16th. From October 16th to February 25th, the station experiences polar night. Further description of the station is found elsewhere (Nguyen et al., 2016; Lange et al., 2018).

130

135



140 **Figure 1. Location of Villum Research Station at Station Nord.** Villum is shown with a red star. Average sea-ice concentration is included for the spring campaign (left) and summer campaign (right). Sea-ice concentrations were taken from the National Sea & Ice Data Center (https://nsidc.org/data/seaice_index/archives/).

The field studies described in this work were conducted in 2016 over two campaigns, one in spring and one in summer. The exact periods when the field studies were carried out are listed in Table 1 below. Basic meteorological parameters (temperature, wind speed and direction, atmospheric pressure, RH and solar radiation) are monitored at Villum.

145

Table 1. Measurement periods during spring and summer field study.

Instruments	Spring 2016	Summer 2016
-------------	-------------	-------------

CCN	20 Apr – 6 June	15 Aug – 2 Oct
HTDMA	22 Apr – 2 May	15 Aug – 28 Aug
SMPS	Full coverage	Full coverage

150 Although the duration of the spring field study arguably extends into the meteorological summer, and the summer field study extends into the meteorological autumn, the naming is kept for convenience.

2.2 Aerosol number size distributions

155 A scanning mobility particle sizer (SMPS) has been operated continuously at Villum since 2010. Studies of aerosol physical properties at Villum, including the measurement setup, have been published elsewhere (Nguyen et al., 2016; Freud et al., 2017; Lange et al., 2018, 2019). In brief, the SMPS system measures size distributions of particles with electrical mobility diameters in the range from 9 to 915 nm, with 5 min time resolution. It consists of a custom-built electrostatic classifier with a Vienna type medium DMA column, operating with a 5:1 sheath-to-sample flow ratio (Wiedensohler et al., 2012), either with a TSI 3010 condensation particle counter (CPC) or TSI 3772 CPC for particle
160 detection. Ambient aerosol is sampled through a specially designed heated inlet for total suspended particles, with laminar flow to minimize losses. There is no additional drying of the sample air prior to entering the analytical instruments, as the transition from the cold outside temperatures to the >20 °C inside of the “Air Observatory” ensures sufficient particle drying. The aerosol sample flow RH is below 35% more than 99% of the time. Raw SMPS measurements are inverted offline by an algorithm as described in Pfeifer et al. (2014).

165 All measurements recorded by the SMPS were quality-controlled by inspection of instrument flows, temperatures, and RHs as well as by visual inspection of size distribution plots. Measurements influenced by local pollution from e.g., nearby vehicles, or when the instrument was malfunctioning, were removed from further consideration.

2.3 Cloud condensation nuclei properties

170 Concentrations of CCN were measured by a Droplet Measurement Technologies CCN-100 CCN counter, cycling ten supersaturations of 0.1–1.0 % SS. The highest reachable supersaturation (typically ~1.8 % SS) was also included in the measurement protocol. CCN concentrations at the highest SS (CCN_{max}) were used as a reference against the number concentration of particles larger than 25 nm (N_{25}) inferred from SMPS measurements. For all settings of SS, the temperature gradient in the CCN column was allowed to stabilize for 5 min, before acquiring data for 5 min. An exception
175 to this procedure was the return from ~1.8 % SS to 0.1 % SS, where a 15 min stabilization time was allowed. The resulting cycling time for all 11 supersaturations was 120 min. Measurements at 0.25, 0.35 and 0.40 % SS showed to be largely redundant with measurements at other SSs and are therefore not presented further here. The SS setting of the instrument was calibrated with monodisperse ammonium sulfate (AS) aerosol, produced by an atomizer and a scanning electrical mobility spectrometer (SEMS), both from Brechtel. By varying AS aerosol particle diameter (D_p), $D_{p,crit}$ was determined
180 for 0.1–0.47 % SS at the sampling site four times during the field studies. Subsequently, calibrations were made at 0.1–1.0 % SS in a more rigorous procedure in our laboratory at the Department of Environmental Science at Aarhus University, Risø Campus, Roskilde, Denmark. The combined experimentally obtained values for $D_{p,crit}$ were compared with the corresponding calculated values obtained using the Köhler equation (Köhler, 1936) and AS bulk solution properties derived from the E-AIM model (Clegg et al., 1992; Wexler and Clegg, 2002). Hereby, a linear relationship
185 between the SS set on the CCN counter (SS_{set}) and the real SS value (SS_{real}) based on theory was established in the full range of 0.1–1.0 % SS (Least squares fit: $SS_{real} = 0.9019 \cdot SS_{set} + 0.0629$, $SD_{SS} = 0.031$ %, $r^2 = 0.995$), applying a method similar to that used by Kristensen et al. (2012), and Nakao et al. (2014). The linear calibration curve is shown in supplementary material 1 (Fig. S1). For convenience, the reported SS values in this paper are the SS_{set} values. Quality control of CCN data was based on the achievement and stability of the CCN column temperature gradient, and the stability

190 of the instrument temperatures and flows. Additionally, CCN measurements recorded simultaneously with SMPS measurements that were discarded during quality control were removed from further consideration.

Aerosol Dp_{crit} was determined by sequential integration of simultaneously measured particle number size distributions and CCN concentrations at a given SS, decreasing Dp_{crit} until the following relationship was satisfied.

$$\int_{Dp_{crit}}^{Dp_{max}} n_N(Dp) dDp = CCN_{SS} \quad (1)$$

195 Here $n_N(Dp)$ is the particle number size distribution and Dp_{max} its upper limit, CCN_{SS} is the measured CCN concentration at a given SS. Particle losses inside the CCN counter were accounted for by applying a size-dependent transmission function (Rose et al., 2010) to particle number size distributions. With the obtained Dp_{crit} , the κ hygroscopicity parameter (κ_{CCN} -value) was calculated according to Petters and Kreidenweis, 2007:

$$\kappa_{CCN} = \frac{4A^3}{27Dp_{crit}^3 \ln(S)^2} \quad (2)$$

200 Where S is the saturation ratio, related to SS as $S = SS/100+1$. The above-described linear SS calibration curve was applied hereto. The constant A is defined as:

$$A = \frac{4\sigma_{a/w} M_w}{RT\rho_w} \quad (3)$$

Where $\sigma_{a/w} = 0.072 J m^{-2}$ is the surface tension of the air/water interface, $T = 298.15$ °C is the standard temperature and ρ_w is the density of water at temperature T, R the gas constant and M_w is the molecular weight of water.

205 We found a discrepancy between the total particle number concentration measured by the CCN counter and the SMPS during the first ten days of the spring study, that we could not explain. We found that CCN_{max} was significantly higher than the simultaneous number concentration of particles larger than 25 nm (N_{25}) measured with the SMPS. This could not be explained by either non-isokinetic sampling conditions in the duct from where the CCN sample air was drawn (Supplementary material 2), or by decreased SMPS sample flow or other irregularities. We suspect the discrepancy to be
 210 caused by conditions in the sampling duct, possibly turbulence, as a change of the position of the CCN sampling port in the duct remedied the situation. The discrepancy was observed during 20–30 April. As the SMPS serves as a long-term monitoring device, we choose to accept it as our reference instrument. The ratio CCN_{max}/N_{25} serves as a tool for assessing the behavior of the CCN counter against the SMPS measurements. The ratio was on average 1.614 in the period 20–30 April, whereas it was 0.939 for the rest of the CCN measurement period. We observed a strong linear correlation between
 215 CCN_{max} and N_{25} for the period after 30 April. By plotting N_{25} as a function of CCN_{max} we observed that CCN_{max} increased by a constant factor during the period of discrepancy, and that there seemed to be no patterns in the offset. Thus, we have chosen to include this period of discrepancy, as it contains highly valuable measurements. We corrected all measured CCN concentrations within this period with a constant factor so that CCN_{max}/N_{25} matches that of the rest of the measurement period. A plot showing N_{25} as a function of CCN_{max} for both the uncorrected and corrected measurements
 220 before 30 April, and the remaining measurements after 30 April is included in Supplementary material 3 (Fig. S2).

2.4 Particle hygroscopic growth factor

Particle hygroscopic growth factor was measured with a Humidified Tandem Differential Mobility Analyzer (HTDMA) Model 3100 from Brechtel (Lopez-Yglesias et al., 2014). The instrument measured HGF of particles with 30, 60, 120 and
 225 240 nm dry diameter (Dp_{dry}) at 85 and 90 % RH. The HTDMA has a built-in aerosol dryer in the upstream classification unit. Due to the low ambient particle number concentrations in the high Arctic environment, the humidified size scans at each Dp_{dry} were conducted over 10 min, also scans with increasing and decreasing DMA voltage were averaged. This procedure ensured sufficient particle counts in most instances. Data inversion was done by an Igor® algorithm based on the principles in Stolzenburg and McMurry (2008) provided by Brechtel. As in the case of the CCN measurements, HGF
 230 measurements coinciding with polluted or faulty SMPS measurements were discarded. Moreover, maximum temperature

changes of 1 °C h⁻¹ during a humidified scan and minimum total particle concentrations of 1 cm⁻¹ in the classified and humidified aerosol size spectra were used as quality control criteria.

235 The HTDMA system was fitted with an automatic atomizer setup, allowing for regular HGF control measurements of AS aerosol. Control measurements of 100 nm AS particles at 85 and 90 % RH were conducted twice daily. Also, the consistency of the two particle sizing stages of the HTDMA was verified with size-certified polystyrene latex (PSL) spheres against a TSI 3080 SMPS system. We found that the set RH was reached within <2 % RH (absolute value) accuracy (Supplementary material 4). We used κ -Köhler theory to correct for these deviations (Petters and Kreidenweis, 2007). The κ_{HTDMA} -value can be inferred from HTDMA measurements by:

$$\frac{RH}{\exp\left(\frac{A}{Dp_{dry} \cdot GF}\right)} = \frac{GF^3 - 1}{GF^3 - (1 - \kappa_{\text{HTDMA}})} \quad (4)$$

240 In this procedure, the actual operating RH, determined from AS control measurements, was used to calculate the κ_{HTDMA} -value. The HGF at 85 or 90 % RH was found by applying this κ_{HTDMA} -value to the formula and determining HGF at the specific RH by an iterative approach described in Supplementary material 5. Ideally, the κ -value of a multispecies aerosol can be determined by the κ -mixing rule:

$$\kappa = \sum \varepsilon_i \kappa_i \quad (5)$$

245 where ε_i is the respective particle volume fraction of component i, and κ_i is the κ -value for that component. This relationship can be used both for κ -values determined by CCN- or HGF measurements.

2.5 Determination of uncertainties

250 An iterative Monte Carlo approach was used to determine Dp_{crit} , κ_{HTDMA} and the associated uncertainty of these parameters. The method is similar in concept to that used in Kristensen et al. (2016) and Herenz et al. (2018).

255 The counting process of the CCN counter was assumed to be Poisson. Hence, the standard deviation (SD) of the CCN concentration within the 5 min of data acquisition is related to the square root of total counts within that period. Even though the CCN concentration was generally low, the long data acquisition time resulted in counting SD rarely exceeding 1%. The uncertainty of SMPS sizing and concentration was assumed to be normally distributed. When determining Dp_{crit} , the SD of the particle diameter was estimated to be 2.5%, while a SD of 5% was estimated on the particle number concentration. The value of Dp_{crit} was calculated according to Eq. (1) where the parameter values were drawn randomly from normal distributions centered at the given value with the above-stated standard deviations. Each calculation was iterated 1000 times, and the final Dp_{crit} and its associated SD were determined by an automatic fit of a normal distribution to the obtained set of results. An example histogram from this procedure is given in supplemental material 6 (Fig. S5).

260 To determine κ_{HTDMA} a similar approach as for Dp_{crit} was applied. The SD of RH found during the monitoring of the HTDMA RH was used, together with the same estimate of particle sizing uncertainty as above 2.5%. Again, fitting normal distributions to the obtained set of results yielded values of κ_{HTDMA} with associated SD. The HGF at exactly 85 and 90% was then recalculated in accordance with Eq. (4), along with low and high estimates by subtracting from or adding the SD to κ_{HTDMA} .

265 Meaningful uncertainty intervals could not be determined for κ_{CCN} as Eq. (2) is highly sensitive to deviations in Dp_{crit} , resulting in skewed result distributions and unrealistic uncertainty intervals. Values of κ_{CCN} are therefore reported without an uncertainty interval, but it is noted that these quantities inherently are associated with large uncertainties.

2.6 Sampling set-up


270 The SMPS sampled aerosols through a total air inlet designed and manufactured by TROPOS (Leipzig, Germany). The inlet consisted of stainless steel which sampled aerosols about 7 m above ground level and was heated to +5 °C to smoothly condition the aerosol to room air temperature where the aerosol instrumentation was placed. The total air inlet had an inner diameter of about 75 mm and was connected to an isokinetic flow splitter where the instruments were connected with ¼ inch conductive tubing. The SMPS sampled from this inlet for both campaigns, while the HTDMA and
275 the CCN counter sampled from a stainless-steel inlet that was designed for gases but fulfilled the requirements for particle sampling during the spring field study, and for the summer field study they sampled from a second total air inlet as described above. The SMPS was operated without a pre-impactor. Losses for the ¼-inch tubing downstream of the flow splitter and in the SMPS were accounted for in the inversion routine from TROPOS (Leipzig, Germany). The SMPS has been running in parallel to a total CPC continuously for the spring field study where the measurements originating from
280 the SMPS (integration) and total CPC agreed reasonably well (mean ratio of CPC to SMPS was 0.85). CPC measurements were unavailable for the summer field study. Meteorological parameters were measured at a 10 m mast next to the measurement hut and originated from an average of two sensors in each height (9 m height and 3 m height).

3 Results

285 3.1 Meteorology

Figures 2 and 3 show the evolution of meteorological parameters during the two field studies. We use UVA radiation as a proxy for solar radiation and gas-phase photo-oxidation of organic compounds, and thus new particle formation and growth. Precipitation measurements were not available for 4 – 27 May. A power outage caused measurements of temperature, RH, radiation, wind speed, and pressure to fail from 25 August to 28 September. During this period, backup
290 measurements of temperature, RH, wind speed, and pressure were available, however, a noticeable offset is seen for the pressure measurements.

During the first weeks of the spring study (Fig. 2a) temperatures were low, about -30 to -10 °C, with occasional snowfall. Later in the study, the temperature increased to just below the freezing point, and the composition of the concurrent precipitation was probably a snow-rain mixture. RH ranged mostly between 70 and 90 % with a short decrease on 26
295 April and occasional peaks later in the study. Time-lapse photography revealed that these peaks were associated with fog, and concurrent removal of accumulation mode particles was also observed. The entire spring study occurred during polar day (Fig. 2b). Correspondingly, UVA radiation was present, with an increasing trend during the study. Atmospheric pressure was high, especially during the first period until 13 May, but occasions with air pressure drops and high wind speeds appeared several times in the latter half of this first study period.



300

Figure 2. Overview of meteorological conditions during the spring field study. a) Temperature, RH and precipitation. The dashed line indicates 0 °C. b) UVA radiation, wind speed and atmospheric pressure.

305

During the summer field study, temperatures were mostly above freezing in August (Fig. 3a). Thereafter they decreased to -10 °C towards the end of the summer study period. RH was more variable in this period compared to spring, ranging from 50 to 90% RH. The beginning of the summer study was still conducted during polar day, however, the UVA radiation was decreasing and almost disappeared at the end of this study period. This is expected, as polar sunset occurred about two weeks after the conclusion of the study. Atmospheric pressure was lower in summer compared to spring, with several occasions of strong decreases of pressure and high wind speeds. Especially the period of 28 - 31 August, which featured higher wind speeds.

310




Figure 3. Overview of meteorological conditions during the summer field study. Lighter shaded lines are backup measurements. a) Temperature, RH and precipitation. The dashed line indicates 0 °C. b) UVA radiation, wind speed and atmospheric pressure.


315

3.2 Ambient particle number size distributions

The average ambient particle number size distributions during the two measurement periods shown in Fig. 4 are typical for Arctic conditions during spring and summer. During the first part of the spring period (20 April – 8 May), a distinct accumulation mode is visible at approximately 100-300 nm. This is the characteristic Arctic haze mode, which predominantly consists of anthropogenic aerosols that are built up during winter and spring. In the second part of the spring period (9 May – 2 June), the haze mode largely disappeared and in turn, a nucleation mode appeared.

In the first part of the summer period (15 August – 28 August), the nucleation mode has grown compared to the latter part of the spring study. An Aitken mode at around 20-30 nm can also be identified. This signifies that secondarily formed particles succeeded to grow into the Aitken mode. In the latter part of the summer period, the number of freshly formed particles decreased, and as an accumulation mode has not yet significantly developed, the total particle number concentration becomes low. The large uncertainty bars on the graphs show that deviations from the average distributions are quite large. For example, new particle formation still happened in the latter part of September, but on fewer occasions.


325



330 **Figure 4. Mean particle number size distributions** during a) the spring measurement period and b) the summer measurement period. Error bars show 25th/75th percentiles in each SMPS size bin over the respective period.

3.3 Cloud condensation nuclei

335 The time series of CCN concentration, Dp_{crit} and κ_{CCN} for 0.1–1.0 % SS are shown for the spring period in Fig. 5, and the summer period in Fig. 6. In less than 0.1 % of all measurements at 1.0 % SS, a $Dp_{\text{crit}} < 25$ nm can be identified. The average Dp_{crit} at ~1.8 % SS was 33.7 nm, while the average activation fraction ($\text{CCN}_{\text{max}}/N_{25}$), as mentioned in Sect. 2.3, was 0.939, which gives credibility to our CCN measurements.



340 **Figure 5. CCN properties at various SSs during the spring measurement period.** a) particle and CCN number concentration, b) critical activation diameter ($D_{p,crit}$), and c) κ_{CCN} -value. Black line: total particle number concentration inferred from SMPS measurements (N_{tot}), gray line: number concentration of particles >25 nm (N_{25}), colored symbols: CCN measurements at 0.1–1.0 % SS.

345 Up to 8 May, CCN concentrations at varying SSs follow a similar pattern and differences between the values are relatively small (Fig. 5). From then and ongoing, differences for measured CCN concentrations are much larger across all SSs. This indicates that before May 8 a quite homogenous aerosol was observed, where even the activation of particles with smaller diameters at higher SS did not appreciably change the total number of activated particles, because accumulation mode particles dominated during this period. In contrast, during the latter period (May 9–June 2), applied SS played a more critical role with respect to the total number of activated particles. While there was little difference between $CCN_{0.1}$ and $CCN_{1.0}$ in the beginning of the spring field study, the difference between these two parameters was larger towards its end (Fig. 5). A subsequent increase in SS led to more activated particles in the Aitken mode range, identifying the importance of Aitken mode aerosol during this period. The days around 8 May seem to mark a transition from the typical Arctic haze period in late winter/early spring to the late spring/summer regime where transport of air masses from mid-latitudes to the high Arctic is minimized. This is supported by a significant decrease in non-sea salt sulfate concentrations measured


350

355

at Villum during the same period (see Skov et al., 2017 for further details and filter pack data from EBAS database, <https://ebas.nilu.no/>).

360

It is interesting to note, that while $D_{p,crit}$ varies at the higher supersaturations, it appears almost constant at 0.1% and 0.15% SS (Figure 5b). Correspondingly, $D_{p,crit}$ and κ_{CCN} -values at 0.1 and 0.15% SS show the smallest temporal variation throughout the spring field study, from all the listed supersaturations (Table 2). This behavior could be explained by a relatively unchanged chemical composition of the larger accumulation mode particles that activated at these lower SSs. A general trend is that the spread of CCN concentrations seen at different SSs increased during the second part of the spring field study (Figure 5a). Further CCN property statistics are stated in Tables 2 and 3.



365

Figure 6. CCN properties at various SSs during the summer measurement period. a) Particle and CCN number concentration, b) critical activation diameter ($D_{p,crit}$), and c) κ_{CCN} value. Black line: total particle number concentration inferred from SMPS measurements (N_{tot}), gray line: number concentration of particles >25 nm (N_{25}), colored symbols: CCN measurements at 0.1-1.0% SS.

370

375 The summer field study is characterized by relatively small variations of the observed critical diameters and corresponding κ -values. This finding indicates a relatively stable and homogeneous chemical composition over the whole summer period of observations, which is not surprising as the high Arctic summer aerosol is dominated by local and regional emissions. Simultaneously, relatively high variations are observed in the number of total activated particles, with all SSs showing similar patterns. An explanation for this finding might be relatively stable emission sources and processing pathways, with the emission source strength varying, which results in a stable shape of the particle number size distribution with varying total particle number concentration. As observed during the second period of the spring field study, differences in activated particle numbers at different applied SSs are still relatively large. This indicates a dominating Aitken mode where higher SSs activate larger fractions (lower $D_{p_{crit}}$) of this aerosol mode, which predominantly originates from local and regional emissions.

380

Table 2. Statistics of CCN measurements for the spring field study. The uncertainty values stated with CCN mean, CCN median, $D_{p_{crit}}$ mean, and $D_{p_{crit}}$ median are the respective statistics of the associated uncertainty, found in the fitting procedure. The values of SD are the standard deviations of the parameter distributions over the temporal evolution of the study. *signifies a significant difference between spring and summer field studies (Wilcoxon rank sum test, $p < 0.05$). CCN measurements for the spring field study were obtained from 20 April to 6 June.

385

Spring	0.1% SS	0.15% SS	0.2% SS	0.3% SS	0.5% SS	0.7% SS	1.0% SS
CCN (uncert) mean [cm^{-3}]	53.7* (± 0.5)	69.3* (± 0.5)	76.5* (± 0.5)	85.3* (± 0.6)	98.4* (± 0.6)	112.7* (± 0.6)	133.8* (± 0.7)
CCN (uncert) median [cm^{-3}]	53.3* (± 0.5)	71.3* (± 0.5)	78.6* (± 0.6)	85.1* (± 0.6)	94.9* (± 0.6)	108.3* (± 0.7)	116.6* (± 0.7)
CCN SD [cm^{-3}]	20.2	25.6	28.0	32.7	41.7	53.9	75.3
$D_{p_{crit}}$ (uncert) mean [nm]	130.2* (± 21.4)	102.9* (± 25.3)	92.2 (± 26.8)	80.2 (± 25.8)	64.2 (± 22.0)	54.4* (± 19.2)	44.8* (± 16.6)
$D_{p_{crit}}$ (uncert) median [nm]	129.2* (± 19.2)	100.7* (± 25.0)	88.9 (± 25.9)	75.9 (± 25.3)	58.7 (± 21.4)	46.7* (± 18.0)	37.3* (± 16.0)
$D_{p_{crit}}$ SD [nm]	9.5	10.9	13.1	14.1	16.8	19.1	19.2
κ_{CCN} mean	0.28*	0.35*	0.33	0.29	0.27	0.27*	0.30*
κ_{CCN} median	0.28*	0.35*	0.34	0.29	0.27	0.29*	0.30*
κ_{CCN} SD	0.06	0.10	0.11	0.12	0.15	0.16	0.19

390 **Table 3. Statistics of CCN measurements for the summer field study.** The uncertainty values stated with CCN mean, CCN median, $D_{p_{crit}}$ mean, and $D_{p_{crit}}$ median are the respective statistics of the associated uncertainty, found in the fitting procedure. The values of SD are the standard deviations of the parameter distributions over the temporal evolution of the study. *signifies a significant difference between spring and summer field studies (Wilcoxon rank-sum test, $p < 0.05$). CCN measurements for the summer field study were obtained from 15 August to 2 October.

Summer	0.1% SS	0.15% SS	0.2% SS	0.3% SS	0.5% SS	0.7% SS	1.0% SS
CCN (uncert) mean [cm^{-3}]	20.8* (± 0.3)	33.2* (± 0.4)	39.6* (± 0.4)	47.6* (± 0.4)	56.6* (± 0.5)	65.6* (± 0.5)	79.5* (± 0.5)
CCN (uncert) median [cm^{-3}]	19.4* (± 0.3)	32.5* (± 0.4)	39.2* (± 0.4)	46.2* (± 0.4)	53.0* (± 0.5)	59.8* (± 0.5)	73.9* (± 0.5)
CCN SD [cm^{-3}]	6.2	12.4	15.7	19.4	24.9	31.1	40.0
$D_{p_{crit}}$ (uncert) mean [nm]	137.1* (± 10.0)	103.3* (± 19.1)	89.7 (± 20.7)	76.7 (± 22.0)	61.6 (± 20.7)	51.3* (± 18.6)	40.5* (± 16.3)
$D_{p_{crit}}$ (uncert) median [nm]	138.7* (± 6.4)	103.6* (± 15.8)	90.0 (± 18.0)	76.6 (± 20.2)	60.7 (± 19.8)	49.7* (± 16.9)	39.3* (± 14.2)
$D_{p_{crit}}$ SD [nm]	7.78	6.51	6.7	8.2	9.15	9.12	5.55
κ_{CCN} mean	0.23*	0.34*	0.35	0.30	0.25	0.24*	0.27*
κ_{CCN} median	0.22*	0.32*	0.33	0.28	0.24	0.23*	0.27*
κ_{CCN} SD	0.04	0.13	0.23	0.15	0.11	0.12	0.10

395

The difference between the mean and median values in Tables 2 and 3 illustrates to which extent extreme values influence the respective parameters. Symmetric distributions have equal values of mean and median, hence differences are caused by skewed distributions. The mean CCN concentration was influenced by low concentrations during the spring study for 0.15 - 0.2 % SS, while higher concentrations elevated the mean above the median for 0.5 - 1.0 % SS. Elevated mean CCN concentrations were observed for all supersaturations during the summer field study, although mean and median values were within the uncertainty range for 0.15–0.3 % SS. Given the low aerosol concentrations during the summer, any perturbation in the CCN burden will elevate the mean. The stated standard deviation for CCN, $D_{p_{crit}}$ and κ_{CCN} signifies the temporal variation of these properties. For $D_{p_{crit}}$, the standard deviation was smaller for the summer study compared to spring. Meanwhile, $D_{p_{crit}}$ median was consistently larger during summer compared to spring. Median and mean κ_{CCN} was consistently larger in spring compared to summer. The standard deviation of κ_{CCN} was larger at 0.15 - 0.3 % SS during summer, but larger at 0.5 - 1.0 % SS during spring. This indicates that particles in the accumulation mode range, with $D_{p_{crit}}$ of about 70 - 100 nm, had a more variable chemical composition during summer, while Aitken mode particles, with $D_{p_{crit}} \sim 35 - 60$ nm, had a more variable chemical composition during the spring measurement period. Thus, it can be concluded that sources of accumulation mode particles were more stable and consistent in spring, while they were more variable and diverse in summer. Conversely, sources of Aitken mode particles were more variable in spring, and more stable and confined in summer.

Except for the total CCN measured at different supersaturations, significant differences for the retrieved hygroscopicity parameters $D_{p_{crit}}$ and κ_{CCN} (mean and median) were observed only for the two lowest and the two highest supersaturations. As low supersaturations - based on our cumulative approach to retrieve $D_{p_{crit}}$ and therewith κ_{CCN} - are mostly affected by the larger particles, we conclude that there are substantial differences in the chemical composition of the larger particles between the two seasons. As high supersaturations are mostly affected by the whole size distribution including the very

415


small particles, we can also conclude that there are differences in the chemical composition of the smallest particles between the two seasons. This is supported by the chemical composition of nucleating vapors at Villum (Beck et al., 2021), which showed that during the spring iodic acid is the main precursor vapor while in the summer sulfuric acid and ammonium become the dominant precursor species. These differences between the seasons for different sizes are somewhat expected as the spring particle number size distribution is mostly affected by processed and larger particles compared to the summer where freshly and locally formed particles affect the particle number size distribution.

We deliberately choose to report the total mean and median values of the entire set of calculated uncertainties, rather than combining the uncertainties by dividing the standard deviation by the square root of the number of samples. This is because the individual sample points cannot be assumed to sample the same quantity. As the aerosol changed with time, the CCN measurement process sampled a changing variable. Hence, the individual measurements did not sample identical probability distribution functions, and the traditional approach for finding the error of the mean is not valid.

During late summer, on 31 Aug, an increased κ_{CCN} of 0.79 at 0.3% occurred with high wind speeds from the direction of the unfrozen fjord, linking this measurement with hygroscopic sea spray aerosols (Martensson et al., 2003; de Leeuw et al., 2011). This is further supported by the concurrent ambient number size distributions (Fig. S6), which showed a bimodal appearance that is indicative of sea spray aerosol (Quinn et al., 2017).

3.4 Particle hygroscopic growth factor

To compensate for RH deviations inside the HTDMA, we determined the κ_{HTDMA} according to the method described in Sect. 2.4 and recalculated the HGF at exact values of 85 and 90 % RH. The time series of these recalculated particle hygroscopic growth factors are presented for the spring study in Fig. 7 and the summer study in Fig. 8. In only very rare cases did we observe bimodal HGF distributions (2.4 % in spring, 0.2 % in summer), signifying a predominantly internally mixed aerosol for both study periods. These instances are neglected, and an internal mixture is assumed at all times.



440


Figure 7. Particle hygroscopic growth factors during the spring study. a) Growth factors at 30 nm $D_{p,dry}$, b) at 60 nm $D_{p,dry}$, c) at 120 nm $D_{p,dry}$ and, d) at 240 nm $D_{p,dry}$. Dots represent values at 85% RH and triangles at 90% RH. The color

indicates the raw particle number concentration measured at that certain $D_{p,dry}$, downstream of the first and upstream of the second DMA.

445

Figure 7a contains fewer data points than the other panels. This was caused by the often too low particle concentrations observed at 30 nm $D_{p,dry}$ during the spring study period. The HGFs at 85 and 90 % followed similar trends for all selected dry particle diameters. Generally, hygroscopic growth factors did not vary greatly during the spring study but it appears that variations were slightly stronger at 90% compared to 85 % RH as expected. There was a general increase of HGF with increasing $D_{p,dry}$, which is further shown in Table 4 below. In general, relatively high HGFs were found for the spring campaign assuming ammonium sulfate or ammonium bisulfate being a major compound of the observed aerosol. While κ_{HTDMA} mean values range between 0.54 and 0.60 at RH = 85 % and between 0.46 and 0.51 at RH = 90 % (Table 4), for comparison a theoretical value for ammonium sulfate is calculated as 0.49 at RH = 89.7 % and 0.53 at RH 84.8 % using the Extended AIM Aerosol Thermodynamics Model (Clegg et al., 1998). Calculated values using the same model for ammonium bisulfate are 0.56 at RH = 89.8 % and 0.61 at RH = 84.9 %. The occurrence of partly unneutralized aerosols has been found earlier by Nielsen et al. (2019) during a spring study in 2015 using a Soot Particle - Aerosol Mass Spectrometer (SP-AMS), which is in accordance with our findings here. The overall HGF timeline shows a relatively stable aerosol, especially for accumulation mode particles with respect to subsaturated hygroscopic growth which might be related to a transport period including predominantly haze aerosol with only little variation. This finding can be confirmed by the small dependence of total CCN on different SSs implying that the majority of the particles are linked to an accumulation mode. The overall HGF timeline shows a relatively stable aerosol, especially for accumulation mode particles with respect to subsaturated hygroscopic growth which might be related to a transport period including predominantly haze aerosol with only little variation. This finding can be confirmed by the small dependence of total CCN on different SSs implying that the majority of the particles are linked to an accumulation mode.

460
465 However, fluctuations of individual values in HGF around the mean values can well be explained by a variation in the degree of neutralization of the aerosol. The distribution of HGF over the size regime as well as the corresponding calculated κ_{HTDMA} imply only little changes of the chemical composition over the size range of the Arctic aerosol investigated at Villum during this spring study.



470

Figure 8. Particle hygroscopic growth factors during the summer study. a) Growth factor at 30 nm $D_{p,dry}$, b) at 60 nm $D_{p,dry}$, and c) at 120 nm $D_{p,dry}$. Dots represent values at 85% RH and triangles at 90% RH. The color indicates the raw particle number concentration measured at that certain $D_{p,dry}$, downstream of the first and upstream of the second DMA.

475

Stronger variations of HGF were observed during the summer study period compared to the spring study. Also, a larger number of measurements were carried out at 30 nm $D_{p,dry}$, but no valid measurements could be obtained at 240 nm. In general, the pattern of HGF at 60 and 120 nm appear to covary, while measurements at 30nm $D_{p,dry}$ are inconsistent with those at larger sizes indicating a different chemical composition for the smallest particle size range. As an overall finding, HGF during the summer study was lower compared to the spring study. These differences are statistically significant for the mean and median values measured at RH = 85% for all sizes and at RH = 90 % for 30 nm and 60 nm $D_{p,dry}$. Correspondingly, κ_{HTDMA} mean values ranging from 0.37 to 0.45 (RH = 85 %) and from 0.36 to 0.45 (RH = 90 %), which no longer reflect a pure chemical inorganic composition of ammonium sulfate or ammonium bisulfate, but rather imply that the observed aerosol might be composed, to a substantial extent, of organic compounds mixed with inorganics species. This finding reflects the overall theory that the summer Arctic aerosol is predominantly of local and regional origin with microbial activity in marginal ice zones and phytoplankton sea surface emissions delivering the inorganic as well as the organic ingredients for particle growth. Organic aerosol precursors, which partly condense onto existing aerosol, might be an explanation for this finding by lowering the overall hygroscopicity of the observed particles leading to lower κ_{HTDMA} mean values. Low-level transport of anthropogenic pollution is to a large extent very limited during this time of the year. Statistics of measured particle hygroscopic growth factors and associated κ_{HTDMA} -values for both study periods are listed in Tables 4 and 5.

480

485

490

Table 4. Statistics of HTDMA measurements during the spring study. The stated uncertainties are the respective statistics of the associated uncertainty retrieved from the fitting procedure as described in Sect. 2.5. The values of SD are the standard deviations of the parameter distributions over the temporal evolution of the studies. *signifies a significant difference between spring and summer field studies (Wilcoxon rank-sum test, $p < 0.05$). HTDMA measurements for the spring field study were obtained from 22 April to 2 May.

Spring	30 nm	60 nm	120 nm	240 nm
HGF 85% RH (low-upper) mean	1.53* (1.48-1.58)	1.55* (1.49-1.61)	1.56* (1.50-1.62)	1.62 (1.55-1.68)
HGF 85% RH (low-upper) median	1.55* (1.50-1.60)	1.54* (1.49-1.60)	1.56* (1.50-1.62)	1.62 (1.55-1.68)
HGF 85% RH SD	0.05	0.04	0.03	0.03
HGF 90% RH (low-upper) mean	1.60* (1.51-1.68)	1.69* (1.58-1.79)	1.68 (1.56-1.79)	1.75 (1.62-1.87)
HGF 90% RH (low-upper) median	1.61* (1.52-1.69)	1.66* (1.55-1.76)	1.69 (1.57-1.79)	1.76 (1.62-1.87)
HGF 90% RH SD	0.06	0.09	0.04	0.04
κ_{HTDMA} 85% RH (uncert) mean	0.60* (± 0.08)	0.56* (± 0.08)	0.54* (± 0.08)	0.59 (± 0.09)
κ_{HTDMA} 85% RH (uncert) median	0.63* (± 0.08)	0.55* (± 0.08)	0.53* (± 0.08)	0.59 (± 0.09)
κ_{HTDMA} 85% RH SD	0.07	0.06	0.04	0.02
κ_{HTDMA} 90% RH (uncert) mean	0.50* (± 0.10)	0.52* (± 0.12)	0.46 (± 0.11)	0.51 (± 0.13)
κ_{HTDMA} 90% RH (uncert) median	0.51* (± 0.10)	0.48* (± 0.11)	0.47 (± 0.11)	0.52 (± 0.13)
κ_{HTDMA} 90% RH SD	0.07	0.11	0.04	0.04

505

Table 5. Statistics of HTDMA measurements during the summer field study. The stated uncertainties are the respective statistics of the associated uncertainty retrieved from the fitting procedure as described in Sect. 2.5. The values of SD are the standard deviations of the parameter distributions over the temporal evolution of the studies. *signifies a significant difference between spring and summer field studies (Wilcoxon rank-sum test, $p < 0.05$), there are no measurements at 240 nm $D_{p, \text{dry}}$ in the summer field study. HTDMA measurements for the summer field study were obtained from 15 August to 28 August.

Summer	30 nm	60 nm	120 nm	240 nm
HGF 85% RH (low-upper) mean	1.36* (1.33-1.39)	1.43* (1.40-1.46)	1.50* (1.46-1.53)	n.a.
HGF 85% RH (low-upper) median	1.34* (1.31-1.37)	1.45* (1.42-1.47)	1.51* (1.46-1.55)	n.a.
HGF 85% RH SD	0.12	0.09	0.05	n.a.
HGF 90% RH (low-upper) mean	1.47* (1.41-1.52)	1.58* (1.50-1.66)	1.67 (1.58-1.75)	n.a.
HGF 90% RH (low-upper) median	1.45* (1.39-1.51)	1.61* (1.52-1.68)	1.69 (1.59-1.77)	n.a.
HGF 90% RH SD	0.09	0.11	0.10	n.a.
κ_{HTDMA} 85% RH (uncert) mean	0.37* (± 0.04)	0.40* (± 0.04)	0.45* (± 0.05)	n.a.
κ_{HTDMA} 85% RH (uncert) median	0.34* (± 0.04)	0.42* (± 0.05)	0.46* (± 0.05)	n.a.
κ_{HTDMA} 85% RH SD	0.20	0.10	0.07	n.a.
κ_{HTDMA} 90% RH (uncert) mean	0.36* (± 0.06)	0.41* (± 0.08)	0.45 (± 0.09)	n.a.
κ_{HTDMA} 90% RH (uncert) median	0.34* (± 0.06)	0.43* (± 0.08)	0.47 (± 0.09)	n.a.
κ_{HTDMA} 90% RH SD	0.10	0.10	0.08	n.a.

510

The HGF low and upper values in Tables 4 and 5 were derived by either subtracting or adding the uncertainty of κ_{HTDMA} in the recalculation procedure, as described in Sect. 2.5. The uncertainty of κ_{HTDMA} , in turn, is the standard deviation of the normal distribution fit applied to the iteratively obtained set of κ_{HTDMA} -values. As noted for the CCN dataset, it cannot be assumed that the HTDMA measurements probe identical probability density functions, thus the combination of the standard deviation to a standard error is not a valid procedure. Again, here we instead state the mean and median values of the κ_{HTDMA} and the corresponding uncertainties.

515

In summary, we can see that for all smaller diameters (30 nm and 60 nm $D_{p, \text{dry}}$) the HGF values (mean and median) at RH 85 and RH 90 % are significantly different between the two field studies. This is also valid for the retrieved κ_{HTDMA} values. The two periods showed no significant difference at 120 nm $D_{p, \text{dry}}$ at 90 % RH. Similarly, κ_{HTDMA} had a larger temporal variability in the summer period. Especially κ_{HTDMA} at 30 nm had a comparably high temporal variability in summer. This indicated that as the hygroscopicity of 30 nm particles was on average lower during summer compared to spring, these

520 particles had a more variable hygroscopicity, and correspondingly a more variable chemical composition. As these particles are expected to originate from nucleation processes in the region, this indicates that various processes and thus different chemical compounds were responsible for the aerosol formation and/or further growth processes of these 30 nm particles during the summer study period (Beck et al., 2021). It has to be noted that there is also evidence for Aitken mode particles of primary origin. Several hypotheses exist which propose different production mechanisms for these particles.
525 This could be, for example, the breakup of larger particles or the collapse of marine gel particles in droplets to a nanoparticle state (Lawler et al., 2021). It was also found that sea spray aerosols can contribute to the Aitken mode population down to sizes of 35 nm (Xu et al., 2022). Contributions from such sources to the Aitken mode aerosol at Villum cannot be fully excluded at least during summertime, when open waters are closer to the station compared to wintertime.


530

3.5 Comparison of particle hygroscopicity derived based on CCN and HTDMA measurements

The comparison of κ -values determined simultaneously (within 2 h) by HTDMA and CCN measurements, revealed differing patterns between the two study periods. The differing pattern was observed at all SSs, but here results from 0.3% SS and both 85% and 90% RH are shown, as this illustration presents the clearest picture. Spring and summer κ_{HTDMA} versus κ_{CCN} datasets are displayed in Fig. 9. In general, only weak correlations were found between κ_{HTDMA} and κ_{CCN} .
535 Earlier studies also found different values between the two techniques (e.g. Petters and Kreidenweis, 2007; Rastak et al., 2017; Rosati et al., 2020).

Particles composed of partly neutralized sulfates or sulfuric acid are expected to exhibit similar hygroscopic behavior compared to that observed for the spring measurements (Petters and Kreidenweis, 2007; Biskos et al., 2009), i.e.,
540 comparably higher hygroscopic growth at subsaturated conditions and lower CCN activity at supersaturation. This corresponds to other findings from Villum, where the haze aerosol was found to be acidic, containing a large fraction of non-neutralized sulfates (Nielsen et al., 2019). However, as the κ -values of un-neutralized sulfates are quite high (e.g. $\kappa_{\text{HTDMA}}(\text{H}_2\text{SO}_4) = 1.19$, $\kappa_{\text{CCN}}(\text{H}_2\text{SO}_4) = 0.90$), additional less hygroscopic species, e.g. biogenic or anthropogenic organics, other inorganic components, and/or black carbon, need to be a significant fraction of the observed internally mixed aerosol
545 as well to explain our observations.

During the summer period, it appears that the aerosol was less acidic. Ammonium sulfate exhibits lower κ -values compared to sulfuric acid and gives a better agreement between the two techniques ($\kappa_{\text{HTDMA}}(\text{AS}) = 0.53$, $\kappa_{\text{CCN}}(\text{AS}) = 0.61$). Basically, a ratio larger than 1 was found between median κ_{HTDMA} and κ_{CCN} for both seasons in our study. Reasons why an increase for median κ_{CCN} was found from spring to summer compared to a decrease for median κ_{HTDMA} for the same period in our study remain unresolved and need further investigation, likely requiring a longer dataset. A possible
550 explanation is stated in the following paragraph.



555 **Figure 9. Correlation of κ_{HTDMA} and κ_{CCN} .** The κ_{HTDMA} -values include measurements at both 85 and 90% RH, of 30, 60, 120 and 240 nm $D_{\text{p,dry}}$. κ_{CCN} was recorded at 0.3% SS. Blue circles are spring measurements, red crosses are summer measurements. The black circle (cross) indicates the median for κ_{HTDMA} and κ_{CCN} during the summer (spring) field study. This Figure is adopted from the supplement material in Lange et al., 2019.

560 The full-year study from Svalbard by Silvergren et al. 2014 shows an increase in κ_{CCN} from April to August, from about 0.39 to 0.54, and an increase in κ_{HTDMA} from about 0.38 to 0.52 (both ranges originate from Figure 9 in the referred study). While we observe a corresponding increase in κ_{CCN} , we also observe a general decrease in κ_{HTDMA} over the corresponding period. This difference might be caused by a smaller relative contribution of sea salt components to the aerosol composition at Villum, compared to Svalbard. Also, a smaller contribution from DMS-derived sulfate aerosol is expected at Villum compared to Svalbard, based on the larger distance to open waters at Villum during summer. Marine microgel aerosols have been described to feature efficient CCN properties while showing low hygroscopic growth at subsaturated conditions (Ovadnevaite et al., 2011; Leck and Svensson, 2015; Hodas et al., 2016). As the coastline in the immediate proximity of Villum was ice-free during the summer study period, and the Arctic pack ice extent was close to a minimum

565

during August, areas of open water and open pack ice were available as potential sources for marine aerosols and gels from the sea surface microlayer. However, we only see a very limited indication of these properties in our data.

570

4 Discussion and outlook

Our measurements show that aerosols of the Arctic atmosphere demonstrate relatively high CCN activity and hygroscopic growth at subsaturated conditions. This is noteworthy as organic compounds make up a significant mass fraction (Leck et al., 2002; Chang et al., 2011; Lange et al., 2018; Nielsen et al. 2019) in Arctic aerosols. Correspondingly, the organic mass fraction must be assumed to contribute significantly to the hygroscopicity of the observed aerosol at Villum. For the spring and the summer measurement period, we determined average κ_{CCN} -values of 0.27–0.35 and 0.23–0.35, respectively. These values are mainly the result of the combined hygroscopicity of the main aerosol sulfate- and organic components. Nielsen et al. (2019) determined the organic mass component in PM_{10} to be on average 24% over the period February–May at Villum. However, the organic component of the aerosol mass in the Arctic is highly size-dependent in the particle size range probed both by the CCN counter and HTDMA in this study (Croft et al., 2019). Croft et al. (2019) performed model calculations for Arctic aerosols during summertime showing that about 50% of the mass fraction around 100 nm in diameter can be of organic origin. Also, the atmospheric aging of Arctic aerosols, which during summer to a large extent originated from local or regional sources, supports that biogenic precursors can largely contribute to the organic mass fraction of Arctic aerosols that were investigated by HTDMA and CCN counter techniques in this study. Tremblay et al. (2019) found large organic mass fractions compared to sulfate during particle formation events in the submicrometer and even ultrafine size range at Eureka station using AMS techniques. Despite the relatively high κ_{CCN} -values, average CCN concentrations were quite low in our study, especially during summer. For supersaturations $\leq 0.3\%$, the spring period featured average CCN concentrations of 53.7–85.3 cm^{-3} while average concentrations of 20.8–47.6 cm^{-3} were observed during the summer period. For SS $\geq 0.5\%$, CCN concentrations in spring were 98.4–133.8 cm^{-3} while CCN concentrations of 56.6–79.5 cm^{-3} were observed in summer. This shows that even when frequent NPF events taking place in summer, the atmosphere contains only few potential CCN, and increases in anthropogenic emissions, either by industrial activities or increased ship traffic may have a correspondingly large impact.

The CCN properties determined in this study compare, to a varying extent, with previous findings from other parts of the Arctic. Our mean CCN concentration at 0.7 % SS of 112.7 and 65.6 cm^{-3} in spring and summer, respectively, are somewhat higher than those of 14–47 cm^{-3} determined in the central Arctic Ocean by Martin et al. (2011). Oppositely, the summer median CCN concentration at 0.55 % SS of 247 cm^{-3} determined by aircraft in Arctic air masses by Latham et al. (2013) is significantly higher than the median CCN concentration at 0.50 % SS of 53.0–95.0 cm^{-3} that was found during both our field studies. Martin et al. (2011) and Latham et al. (2013) determined κ_{CCN} (mean \pm SD) to be 0.33 ± 0.13 and 0.32 ± 0.21 , respectively, which compares well to our study even if they are slightly higher than our total field study (mean \pm SD) κ_{CCN} of 0.30 ± 0.21 . A study by Jung et al. (2018), evaluated a several-year CCN dataset from Zeppelin Mountain, Svalbard. They found CCN concentrations at 0.2 % SS of 70–133 cm^{-3} in April–June and 16–57 cm^{-3} in August–September. This compares well to the CCN concentrations at 0.2 % SS of $76.5 \pm 28.0 \text{ cm}^{-3}$ and $39.6 \pm 15.7 \text{ cm}^{-3}$ we determined in spring and summer, respectively. Overall, the aerosol at Villum appears to be slightly less hygroscopic compared to that of the Central Arctic Ocean and the Canadian Arctic. However, the measured CCN concentrations at Villum are higher than those measured in the Arctic Ocean by Martin et al. (2011) but lower than those found over the Canadian Arctic (Latham et al., 2013) and at Zeppelin Mountain (Jung et al., 2018). This highlights that within the Arctic region, aerosol CCN activity, and therewith probably the aerosol sources, are somewhat heterogeneous in both time and space. Additionally, the aircraft measurements, as well as Zeppelin mountain measurements, originate in the free troposphere while Villum measurements represent boundary layer sampling making the local origin of observed aerosol more likely.

During summer, we observed the above-described heterogeneity of aerosol hygroscopicity. This was likely to be caused by the numerous and diverse active aerosol sources during the course of the summer measurement period. These sources are mostly of local and regional origin, including open waters, sea-ice edges or snow- and ice-covered surfaces.

Dall'Osto et al. (2017) demonstrated the influence of partially open sea-ice on NPF frequency, with a quantitative analysis combining air mass history and sea-ice coverage. A similar approach, including the time spent over different types of Arctic surface areas combined with meteorological parameters and potential biological activity, could potentially

615

determine individual sources of CCN-relevant aerosols in case a substantially long dataset of CCN measurements is available.

620 The actual supersaturation that develops during the atmospheric updraft of an air parcel is critically important for determining which fraction of the ambient aerosol acts as CCN. Yet little is known about this process in the Arctic atmosphere. Peak supersaturations (SS_{peak}) were calculated by Earle et al. (2011), based on aircraft measurements of cloud properties and characterization of aerosols below cloud, over the Beaufort Sea in April. For a bimodal aerosol, which is also well representative of our data, and under unpolluted conditions, they determined SS_{peak} of 0.33–0.44 % for updraft velocities of 30–40 cm s^{-1} and reported calculated κ -values of 0.2–0.3. Lower κ -values delay the onset of liquid water condensation during updraft, resulting in the aerosol population being exposed to higher SS_{peak} , which partly offsets the importance of κ -values. Oppositely, polluted aerosols, characterized by higher number concentrations, deplete available water vapor faster during condensation, which lowers the effective SS_{peak} . Values of 0.12–0.18 % SS are presented in the above-mentioned study for polluted cases. At the subarctic station Pallas in northern Finland SS_{peak} of 0.27–0.63 % was determined in Sep–Oct (Anttila et al., 2012), though they observed generally lower κ -values compared to our studies at 630 Villum, which could be explained by their proximity to anthropogenic emission sources.

Based on these peak supersaturations, the ambient $D_{\text{p,crit}}$ at Villum would be approximately 68–80 nm, yielding actual ambient CCN concentrations of about 85–92 cm^{-3} and 48–53 cm^{-3} , during the spring and summer period, respectively. These critical activation diameters are between the Aitken and accumulation mode (Fig. 2), indicating that even if newly formed particles grow into the lower Aitken mode, which is an established phenomenon in the Arctic atmosphere (Nguyen et al., 2016; Willis et al., 2016; Burkart et al., 2017; Dall'Osto et al., 2018a, 2018b), significant condensational growth will be required for these particles to reach cloud relevant sizes. Consequently, sources of accumulation mode aerosols could be more relevant for the CCN population compared to sources of new particle formation. This underlines the relevance of biogenic primary marine sources because these could emit CCN-relevant aerosols directly during summer when Arctic haze accumulation mode aerosol is absent. Marine gels and water-insoluble organic material from the sea surface microlayer have been shown to be directly emitted as particles in the CCN relevant size ranges (Facchini et al., 2008; Orellana et al., 2011; Karl et al., 2013). Hamacher-Barth et al. (2016) found gel particles at D_{p} of predominantly >45 nm within a morphological analysis. A chemical analysis of gel polysaccharide monomers showed that pentoses and hexoses, associated with cellular material of phytoplankton, and deoxysugars from microbial exudates were present in the Aitken and accumulation mode ranges of Arctic marine aerosols (Leck et al., 2013). A similar targeted chemical and morphological analysis of aerosol filter samples from Villum could therefore clarify the role of marine gels for late spring and summertime CCN and is proposed here for further studies. 640 645

The average RH during the spring period at Villum was 80.3 ± 7.9 %, while it was 77.3 ± 11.6 % during the summer period. For our HTDMA measurements, we found average HGFs of 1.56 and 1.62 at 85 % RH for 120 and 240 nm particles in spring, respectively, and 1.50 at 85 % RH for 120 nm particles during summer (no valid measurements of HGFs for 240 nm were obtained in summer). The ambient RH was similar to the RH measured at Ny-Ålesund, Svalbard, reported by Rastak et al. (2014). They used calculated HGFs for 200 nm particles at 85% RH to determine the direct radiative forcing exerted by hygroscopic aerosols. For the months of April and May, they used HGFs with an average of about 1.56, and for August, they used HGFs of about 1.59. They determined an average annual aerosol scattering enhancement factor of 4.30 ± 2.26 resulting in an aerosol-radiation RF of -0.83 to -2.60 W m^{-2} . Similarly, we expect particle hygroscopic growth at Villum at the ambient RH to yield a negative impact on radiative transfer as well. As we observed both higher RH and HGFs in spring compared to summer, the direct scattering effect should be more pronounced in spring and early summer compared to periods later in the year. Hygroscopic growth appears to be more pronounced in spring, whereas CCN processes seem to be relatively more important during summer. In spring, aerosols of anthropogenic origin, dominated by acidic sulfate particles and carbonaceous aerosols, exhibit high hygroscopic growth and aerosol-radiation effects potentially are the dominating RF mechanisms during this time of year. In summer, accumulation mode aerosol concentrations are smaller, and aerosol hygroscopic growth is generally lower. In this period, natural CCN active marine aerosols could be the dominating aerosol climate forcing agent through aerosol-cloud effects. Detailed modeling studies of the climatic influence of Arctic aerosols through aerosol-radiation and aerosol-cloud interactions could clarify this hypothesis. It must be noted that the Arctic cloud regime is very susceptible to small changes in CCN concentrations (Mauritsen et al., 2011). During the summertime, particle number concentrations in Arctic environments can undergo values below 10 cm^{-3} (Freud et al., 2017). This is why detailed measurements of subsaturated hygroscopicity and CCN ability are needed to understand the role of Arctic aerosols in aerosol-cloud-climate interactions. 650 655 660 665

Ongoing nephelometer measurements at Villum will provide a basis for further investigations of the impacts of aerosol hygroscopic growth on aerosol-radiation interactions in the nearer future. Furthermore, measurement series of hygroscopic properties of longer duration at Villum would be highly valuable, as a full yearly cycle of CCN and HGF measurements would be a leap forward in the understanding of Arctic climate-relevant aerosol properties.

5 Conclusions

Arctic aerosols, and their radiation- and cloud interactions, are key factors in the changing Arctic climate. Still though, many details about their sources, hygroscopic properties, and cloud interactions are not well understood. In this work, we provide results from two field studies conducted at the High Arctic site, Villum Research Station. One field study was carried out in the spring to early summer of 2016, while the other study was conducted from summer to autumn of the same year. During these field studies, particle hygroscopic growth factors and CCN properties were measured. Based on other studies, we expect peak supersaturations of about 0.3-0.4% to occur in Arctic clouds. This yields actual ambient average CCN concentrations of 85-92 cm^{-3} and 48-53 cm^{-3} during the spring and summer measurement period, respectively. The corresponding $D_{p,\text{crit}}$ at 0.3-0.4% SS was approximately 68-80 nm, showing that accumulation mode aerosols are more important for the CCN population compared to nucleation- and Aitken mode aerosols. These results can be used directly in model parameterisation. During spring, accumulation mode aerosols originated mostly from long-range transport and Arctic haze. During summer, the concentration of accumulation mode aerosols was generally low, and we cannot present a clear indication of its origin. Sea salt influences seem to be low, and only weak indications of primary marine organic particles were observed in the combined analysis of HGF and CCN measurements.

The ambient aerosol appears to be influenced by a multitude of sources during summer. Based on other studies of the impact of hygroscopic properties on radiative forcing in the High Arctic, the hygroscopic properties at Villum very likely result in direct aerosol-radiation interactions that lead to atmospheric cooling, especially in the spring period. Future studies, combining several additional methods, could further clarify the role of marine gels from the sea surface microlayer on CCN concentration and aerosol hygroscopic growth on the radiation balance in the High Arctic atmosphere. Targeted chemical and morphological analysis for marine gels on aerosol filter samples, aerosol scattering, or on-line aerosol composition measurements of the submicrometer aerosol could be relevant tools to further clarify open questions about the climate-relevant role of Arctic aerosols.

695

6 Acknowledgements

RL and UG were funded by the Independent Research Fund Denmark, grant DFF-4005-00482B-FTP “NUMEN”. The overall study was financially supported by the Danish Environmental Protection Agency with means from the MIKA/DANCEA funds for Environmental Support to the Arctic Region, which is part of the Danish contribution to “Arctic Monitoring and Assessment Program” (AMAP) and the Danish research project “Short lived Climate Forcers” (SLCF). We greatly acknowledge the Danish Center for the Environment for funding a grant to write this article. In addition, we want to thank our technician Bjarne Jensen and IT-expert Keld Mortensen for their great support within this study. The military staff at Station Nord is highly acknowledged for their excellent logistical help as well as Villum Foundation for providing the station infrastructure and corresponding applied instrumentation.

705

7 Data Availability

All data used in this manuscript can be found at <https://doi.org/10.5281/zenodo.7307293>. Sea-ice concentrations were taken from the National Sea & Ice Data Center (https://nsidc.org/data/seaice_index/archives/).

710

References

- Allan, J. D., Williams, P. I., Najera, J., Whitehead, J. D., Flynn, M. J., Taylor, J. W., Liu, D., Darbyshire, E., Carpenter, L. J., Chance, R., Andrews, S. J., Hackenberg, S. C., and McFiggans, G.: Iodine observed in new particle formation events in the Arctic atmosphere during ACCACIA, *Atmos Chem Phys*, 15, 5599-5609, <https://doi.org/10.5194/acp-15-5599-2015>, 2015.
- Anttila, T., Brus, D., Jaatinen, A., Hyvarinen, A. P., Kivekas, N., Romakkaniemi, S., Komppula, M., and Lihavainen, H.: Relationships between particles, cloud condensation nuclei and cloud droplet activation during the third Pallas Cloud Experiment, *Atmos Chem Phys*, 12, 11435-11450, <https://doi.org/10.5194/acp-12-11435-2012>, 2012.
- Barrie, L. A., Hoff, R. M., and Daggupaty, S. M.: The Influence of Mid-Latitudinal Pollution Sources on Haze in the Canadian Arctic, *Atmos Environ*, 15, 1407-1419, [https://doi.org/10.1016/0004-6981\(81\)90347-4](https://doi.org/10.1016/0004-6981(81)90347-4), 1981.
- Beck, L. J., Sarnela, N., Junninen, H., Hoppe, C. J. M., Garmash, O., Bianchi, F., Riva, M., Rose, C., Peräkylä, O., Wimmer, D., Kausiala, O., Jokinen, T., Ahonen, L., Mikkilä, J., Hakala, J., He, X.-C., Kontkanen, J., Wolf, K. K. E., Cappelletti, D., Mazzola, M., Traversi, R., Petroselli, C., Viola, A. P., Vitale, V., Lange, R., Massling, A., Nøjgaard, J. K., Krejci, R., Karlsson, L., Zieger, P., Jang, S., Lee, K., Vakkari, V., Lampilahti, J., Thakur, R. C., Leino, K., Kangasluoma, J., Duplissy, E.-M., Siivola, E., Marbouti, M., Tham, Y. J., Saiz-Lopez, A., Petäjä, T., Ehn, M., Worsnop, D. R., Skov, H., Kulmala, M., Kerminen, V.-M., and Sipilä, M.: Differing Mechanisms of New Particle Formation at Two Arctic Sites, *Geophys Res Lett*, 48, e2020GL091334, <https://doi.org/10.1029/2020GL091334>, 2021.
- Bellouin, N., Rae, J., Jones, A., Johnson, C., Haywood, J., and Boucher, O.: Aerosol forcing in the Climate Model Intercomparison Project (CMIP5) simulations by HadGEM2-ES and the role of ammonium nitrate, *J Geophys Res-Atmos*, 116, D20206, <https://doi.org/10.1029/2011jd016074>, 2011.
- Biskos, G., Buseck, P. R., and Martin, S. T.: Hygroscopic growth of nucleation-mode acidic sulfate particles, *J Aerosol Sci*, 40, 338-347, <https://doi.org/10.1016/j.jaerosci.2008.12.003>, 2009.
- Breider, T. J., Mickley, L. J., Jacob, D. J., Wang, Q. Q., Fisher, J. A., Chang, R. Y. W., and Alexander, B.: Annual distributions and sources of Arctic aerosol components, aerosol optical depth, and aerosol absorption, *J Geophys Res-Atmos*, 119, 4107-4124, <https://doi.org/10.1002/2013jd020996>, 2014.
- Breider, T. J., Mickley, L. J., Jacob, D. J., Ge, C., Wang, J., Sulprizio, M. P., Croft, B., Ridley, D. A., McConnell, J. R., Sharma, S., Husain, L., Dutkiewicz, V. A., Eleftheriadis, K., Skov, H., and Hopke, P. K.: Multidecadal trends in aerosol radiative forcing over the Arctic: Contribution of changes in anthropogenic aerosol to Arctic warming since 1980, *J Geophys Res-Atmos*, 122, 3573-3594, <https://doi.org/10.1002/2016jd025321>, 2017.
- Browse, J., Carslaw, K. S., Arnold, S. R., Pringle, K., and Boucher, O.: The scavenging processes controlling the seasonal cycle in Arctic sulphate and black carbon aerosol, *Atmos Chem Phys*, 12, 6775-6798, <https://doi.org/10.5194/acp-12-6775-2012>, 2012.
- Burgos, M., Andrews, E., Titos, G., Alados-Arboledas, L., Baltensperger, U., Day, D., Jefferson, A., Kalivitis, N., Mihalopoulos, N., Sherman, J., Sun, J., Weingartner, E., and Zieger, P.: A global view on the effect of water uptake on aerosol particle light scattering, *Scientific Data*, 6, 157, <https://doi.org/10.1038/s41597-019-0158-7>, 2019.
- Burkart, J., Hodshire, A. L., Mungall, E. L., Pierce, J. R., Collins, D. B., Ladino, L. A., Lee, A. K. Y., Irish, V., Wentzell, J. J. B., Liggio, J., Papakyriakou, T., Murphy, J., and Abbatt, J.: Organic Condensation and Particle Growth to CCN Sizes in the Summertime Marine Arctic is Driven by Materials More Semivolatile Than at Continental Sites, *Geophys Res Lett*, 44, 10725-10734, <https://doi.org/10.1002/2017gl075671>, 2017.
- Carrico, C. M., Rood, M. J., Ogren, J. A., Neususs, C., Wiedensohler, A., and Heintzenberg, J.: Aerosol optical properties at Sagres, Portugal during ACE-2, *Tellus B*, 52, 694-715, <https://onlinelibrary.wiley.com/doi/abs/10.1034/j.1600-0889.2000.00049.x>, 2000.
- Chang, R. Y. W., Leck, C., Graus, M., Muller, M., Paatero, J., Burkhart, J. F., Stohl, A., Orr, L. H., Hayden, K., Li, S. M., Hansel, A., Tjernstrom, M., Leaitch, W. R., and Abbatt, J. P. D.: Aerosol composition and sources in the central Arctic Ocean during ASCOS, *Atmos Chem Phys*, 11, 10619-10636, <https://doi.org/10.5194/acp-11-10619-2011>, 2011.
- Chylek, P., Vogelsang, T. J., Klett, J. D., Hengartner, N., Higdon, D., Lesins, G., and Dubey, M. K.: Indirect Aerosol Effect Increases CMIP5 Models' Projected Arctic Warming, *J Climate*, 29, 1417-1428, <https://doi.org/10.1175/Jcli-D-15-0362.1>, 2016.
- Clegg, S. L., Pitzer, K. S., and Brimblecombe, P.: Thermodynamics of Multicomponent, Miscible, Ionic-Solutions. 2. Mixtures Including Unsymmetrical Electrolytes, *J Phys Chem-Us*, 96, 9470-9479, <https://doi.org/10.1021/j100202a074>, 1992.
- Clegg, S. L., Brimblecombe, P., Wexler, A. S.: A thermodynamic model of the system $H^+ - NH_4^+ - Na^+ - SO_4^{2-} - NO_3^- - Cl^- - H_2O$ at 298.15 K, *J Phys Chem A*, 102, 2155-2171, <https://doi.org/10.1021/jp973043j>, 1998.
- Collins, D. B., Burkart, J., Chang, R. Y. W., Lizotte, M., Boivin-Rioux, A., Blais, M., Mungall, E. L., Boyer, M., Irish, V. E., Masse, G., Kunkel, D., Tremblay, J. E., Papakyriakou, T., Bertram, A. K., Bozem, H., Gosselin, M.,

- Levasseur, M., and Abbatt, J. P. D.: Frequent ultrafine particle formation and growth in Canadian Arctic marine and coastal environments, *Atmos Chem Phys*, 17, 13119-13138, <https://doi.org/10.5194/acp-17-13119-2017>, 2017.
- 770 Croft, B., Martin, R. V., Leaitch, W. R., Tunved, P., Breider, T. J., D'Andrea, S. D., and Pierce, J. R.: Processes controlling the annual cycle of Arctic aerosol number and size distributions, *Atmos Chem Phys*, 16, 3665-3682, <https://doi.org/10.5194/acp-16-3665-2016>, 2016.
- 775 Croft, B., Martin, R. V., Leaitch, W. R., Burkart, J., Chang, R. Y.-W., Collins, D. B., Hayes, P. L., Hodshire, A. L., Huang, L., Kodros, J. K., Moravek, A., Mungall, E. L., Murphy, J. G., Sharma, S., Tremblay, S., Wentworth, G. R., Willis, M. D., Abbatt, J. P. D., and Pierce, J. R.: Arctic marine secondary organic aerosol contributes significantly to summertime particle size distributions in the Canadian Arctic Archipelago, *Atmos Chem Phys*, 19, 2787-2812, <https://doi.org/10.5194/acp-19-2787-2019>, 2019.
- 780 Dall'Osto, M., Beddows, D. C. S., Tunved, P., Krejci, R., Strom, J., Hansson, H. C., Yoon, Y. J., Park, K. T., Becagli, S., Udisti, R., Onasch, T., O'Dowd, C. D., Simo, R., and Harrison, R. M.: Arctic sea ice melt leads to atmospheric new particle formation, *Sci Rep-Uk*, 7, 3318, <https://doi.org/10.1038/s41598-017-03328-1>, 2017.
- Dall'Osto, M., Geels, C., Beddows, D. C. S., Boertmann, D., Lange, R., Nøjgaard, J. K., Harrison, R. M., Simo, R., Skov, H., and Massling, A.: Regions of open water and melting sea ice drive new particle formation in North East Greenland, *Sci Rep-Uk*, 8, 6109, <https://doi.org/10.1038/s41598-018-24426-8>, 2018a.
- 785 Dall'Osto, M., Simo, R., Harrison, R. M., Beddows, D. C. S., Saiz-Lopez, A., Lange, R., Skov, H., Nøjgaard, J. K., Nielsen, I. E., and Massling, A.: Abiotic and biotic sources influencing spring new particle formation in North East Greenland, *Atmos Environ*, 190, 126-134, <https://doi.org/10.1016/j.atmosenv.2018.07.019>, 2018b.
- de Leeuw, G., Andreas, E. L., Anguelova, M. D., Fairall, C. W., Lewis, E. R., O'Dowd, C., Schulz, M., and Schwartz, S. E.: Production Flux of Sea Spray Aerosol, *Rev Geophys*, 49, RG2001, <https://doi.org/10.1029/2010rg000349>, 2011.
- 790 Earle, M. E., Liu, P. S. K., Strapp, J. W., Zelenyuk, A., Imre, D., McFarquhar, G. M., Shantz, N. C., and Leaitch, W. R.: Factors influencing the microphysics and radiative properties of liquid-dominated Arctic clouds: Insight from observations of aerosol and clouds during ISDAC, *J Geophys Res-Atmos*, 116, D00T09, <https://doi.org/10.1029/2011jd015887>, 2011.
- 795 Engvall, A. C., Krejci, R., Strom, J., Treffeisen, R., Scheele, R., Hermansen, O., and Paatero, J.: Changes in aerosol properties during spring-summer period in the Arctic troposphere, *Atmos Chem Phys*, 8, 445-462, <https://doi.org/10.5194/acp-8-445-2008>, 2008.
- Fachini, M. C., Rinaldi, M., Decesari, S., Carbone, C., Finessi, E., Mircea, M., Fuzzi, S., Ceburnis, D., Flanagan, R., Nilsson, E. D., de Leeuw, G., Martino, M., Woeltjen, J., and O'Dowd, C. D.: Primary submicron marine aerosol dominated by insoluble organic colloids and aggregates, *Geophys Res Lett*, 35, L17814, <https://doi.org/10.1029/2008gl034210>, 2008.
- 800 Fierz-Schmidhauser, R., Zieger, P., Wehrle, G., Jefferson, A., Ogren, J. A., Baltensperger, U., and Weingartner, E.: Measurement of relative humidity dependent light scattering of aerosols, *Atmos Meas Tech*, 3, 39-50, <https://doi.org/10.5194/amt-3-39-2010>, 2010.
- 805 Freud, E., Krejci, R., Tunved, P., Leaitch, R., Nguyen, Q. T., Massling, A., Skov, H., and Barrie, L.: Pan-Arctic aerosol number size distributions: seasonality and transport patterns, *Atmos Chem Phys*, 17, 8101-8128, <https://doi.org/10.5194/acp-17-8101-2017>, 2017.
- Fu, P. Q., Kawamura, K., Chen, J., Charriere, B., and Sempere, R.: Organic molecular composition of marine aerosols over the Arctic Ocean in summer: contributions of primary emission and secondary aerosol formation, *Biogeosciences*, 10, 653-667, <https://doi.org/10.5194/bg-10-653-2013>, 2013.
- 810 Fu, P. Q., Kawamura, K., Chen, J., Qin, M. Y., Ren, L. J., Sun, Y. L., Wang, Z. F., Barrie, L. A., Tachibana, E., Ding, A. J., and Yamashita, Y.: Fluorescent water-soluble organic aerosols in the High Arctic atmosphere, *Sci Rep-Uk*, 5, 9845, <https://doi.org/10.1038/srep09845>, 2015.
- Garrett, T. J., Zhao, C., Dong, X., Mace, G. G., and Hobbs, P. V.: Effects of varying aerosol regimes on low-level Arctic stratus, *Geophys Res Lett*, 31, L17105, <https://doi.org/10.1029/2004gl019928>, 2004.
- 815 Garrett, T. J. and Zhao, C. F.: Increased Arctic cloud longwave emissivity associated with pollution from mid-latitudes, *Nature*, 440, 787-789, <https://doi.org/10.1038/nature04636>, 2006.
- Hamacher-Barth, E., Leck, C., and Jansson, K.: Size-resolved morphological properties of the high Arctic summer aerosol during ASCOS-2008, *Atmos Chem Phys*, 16, 6577-6593, <https://doi.org/10.5194/acp-16-6577-2016>, 2016.
- 820 Heidam, N. Z., Wählin, P., and Christensen, J. H.: Tropospheric gases and aerosols in northeast Greenland, *J Atmos Sci*, 56, 261-278, [https://doi.org/10.1175/1520-0469\(1999\)056<0261:Tgaain>2.0.Co;2](https://doi.org/10.1175/1520-0469(1999)056<0261:Tgaain>2.0.Co;2), 1999.
- Heidam, N. Z., Christensen, J., Wählin, P., and Skov, H.: Arctic atmospheric contaminants in NE Greenland: levels, variations, origins, transport, transformations and trends 1990-2001, *Sci Total Environ*, 331, 5-28, <https://doi.org/10.1016/j.scitotenv.2004.03.033>, 2004.
- 825 Herenz, P., Wex, H., Henning, S., Kristensen, T. B., Rubach, F., Roth, A., Borrmann, S., Bozem, H., Schulz, H., and Stratmann, F.: Measurements of aerosol and CCN properties in the Mackenzie River delta (Canadian Arctic) during

- spring-summer transition in May 2014, *Atmos Chem Phys*, 18, 4477-4496, <https://doi.org/10.5194/acp-18-4477-2018>, 2018.
- 830 Hodas, N., Zuend, A., Schilling, K., Berkemeier, T., Shiraiwa, M., Flagan, R. C., and Seinfeld, J. H.: Discontinuities in hygroscopic growth below and above water saturation for laboratory surrogates of oligomers in organic atmospheric aerosols, *Atmos Chem Phys*, 16, 12767-12792, <https://doi.org/10.5194/acp-16-12767-2016>, 2016.
- Iversen, T. and Joranger, E.: Arctic Air-Pollution and Large-Scale Atmospheric Flows, *Atmos Environ*, 19, 2099-2108, [https://doi.org/10.1016/0004-6981\(85\)90117-9](https://doi.org/10.1016/0004-6981(85)90117-9), 1985.
- 835 Jung, C. H., Yoon, Y. J., Kang, H. J., Gim, Y., Lee, B. Y., Ström, J., Krejci, R., Tunved, P.: The seasonal characteristics of cloud condensation nuclei (CCN) in the arctic lower troposphere, *Tellus B: Chemical and Physical Meteorology*, 70, 1-13, <https://doi.org/10.1080/16000889.2018.1513291>, 2018.
- Karl, M., Leck, C., Coz, E., and Heintzenberg, J.: Marine nanogels as a source of atmospheric nanoparticles in the high Arctic, *Geophys Res Lett*, 40, 3738-3743, <https://doi.org/10.1002/grl.50661>, 2013.
- 840 Kirpes, R. M., Bondy, A. L., Bonanno, D., Moffet, R. C., Wang, B. B., Laskin, A., Ault, A. P., and Pratt, K. A.: Secondary sulfate is internally mixed with sea spray aerosol and organic aerosol in the winter Arctic, *Atmos Chem Phys*, 18, 3937-3949, <https://doi.org/10.5194/acp-18-3937-2018>, 2018.
- Köhler, H.: The nucleus in and the growth of hygroscopic droplets., *T Faraday Soc*, 32, 1152-1161, <https://doi.org/10.1039/tf9363201152>, 1936.
- 845 Kristensen, T. B., Wex, H., Nekat, B., Nojgaard, J. K., van Pinxteren, D., Lowenthal, D. H., Mazzoleni, L. R., Dieckmann, K., Koch, C. B., Mentel, T. F., Herrmann, H., Hallar, A. G., Stratmann, F., and Bilde, M.: Hygroscopic growth and CCN activity of HULIS from different environments, *J Geophys Res-Atmos*, 117, D22203, <https://doi.org/10.1029/2012jd018249>, 2012.
- Kristensen, T. B., Muller, T., Kandler, K., Benker, N., Hartmann, M., Prospero, J. M., Wiedensohler, A., and Stratmann, F.: Properties of cloud condensation nuclei (CCN) in the trade wind marine boundary layer of the western North Atlantic, *Atmos Chem Phys*, 16, 2675-2688, <https://doi.org/10.5194/acp-16-2675-2016>, 2016.
- 850 Lange, R., Dall'Osto, M., Skov, H., Nojgaard, J. K., Nielsen, I. E., Beddowse, D. C. S., Simob, R., Harrison, R. M., and Massling, A.: Characterization of distinct Arctic aerosol accumulation modes and their sources, *Atmos Environ*, 183, 1-10, <https://doi.org/10.1016/j.atmosenv.2018.03.060>, 2018.
- Lange, R., Dall'Osto, M., Skov, H., and Massling, A.: Large summer contribution of organic biogenic aerosols to Arctic cloud condensation nuclei, *Geophys Res Lett*, 46, 11500-11509, <https://doi.org/10.1029/2019GL084142>, 2019.
- 855 Latham, T. L., Beyersdorf, A. J., Thornhill, K. L., Winstead, E. L., Cubison, M. J., Hecobian, A., Jimenez, J. L., Weber, R. J., Anderson, B. E., and Nenes, A.: Analysis of CCN activity of Arctic aerosol and Canadian biomass burning during summer 2008, *Atmos Chem Phys*, 13, 2735-2756, <https://doi.org/10.5194/acp-13-2735-2013>, 2013.
- Lawler, M. J., Saltzman, E. S., Karlsson, L., Zieger, P., Salter, M., Baccarini, A., Schmale, J., Leck, C.: New insights into the composition and origins of ultrafine aerosol in the summertime high Arctic. *Geophysical Research Letters*, 48, 1-11, <https://doi.org/10.1029/2021GL094395>, 2021.
- 860 Leaitch, W. R., Sharma, S., Huang, L., Toom-Sauntry, D., Chivulescu, A., Macdonald, A. M., von Salzen, K., Pierce, J. R., Bertram, A. K., Schroder, J. C., Shantz, N. C., Chang, R. Y. W., and Norman, A. L.: Dimethyl sulfide control of the clean summertime Arctic aerosol and cloud, *Elem Sci Anth* 1, 000017, <https://doi.org/10.12952/journal.elementa.000017>, 2013.
- 865 Leck, C., Norman, M., Bigg, E. K., and Hillamo, R.: Chemical composition and sources of the high Arctic aerosol relevant for cloud formation, *J Geophys Res-Atmos*, 107, 4135, <https://doi.org/10.1029/2001jd001463>, 2002.
- Leck, C. and Bigg, E. K.: Biogenic particles on the surface microlayer and overlaying atmosphere in the central Arctic Ocean during summer, *Tellus B*, 57B, 305-316, <https://doi.org/10.1111/j.1600-0889.2005.00148.x>, 2005.
- 870 Leck, C., Gao, Q., Rad, F. M., and Nilsson, U.: Size-resolved atmospheric particulate polysaccharides in the high summer Arctic, *Atmos Chem Phys*, 13, 12573-12588, <https://doi.org/10.5194/acp-13-12573-2013>, 2013.
- Leck, C. and Svensson, E.: Importance of aerosol composition and mixing state for cloud droplet activation over the Arctic pack ice in summer, *Atmos Chem Phys*, 15, 2545-2568, <https://doi.org/10.5194/acp-15-2545-2015>, 2015.
- 875 Lenssen, N. J. L., Schmidt, G. A., Hansen, J. E., Menne, M. J., Persin, A., Ruedy, R., Zyss, D.: Improvements in the GISTEMP Uncertainty Model, *J Geophys Res-Atmos*, 124, 12, 6307-6326, <https://doi.org/10.1029/2018JD029522>, 2019.
- Liu, X. G., Gu, J. W., Li, Y. P., Cheng, Y. F., Qu, Y., Han, T. T., Wang, J. L., Tian, H. Z., Chen, J., and Zhang, Y. H.: Increase of aerosol scattering by hygroscopic growth: Observation, modeling, and implications on visibility, *Atmos Res*, 132, 91-101, <https://doi.org/10.1016/j.atmosres.2013.04.007>, 2013.
- 880 Lohmann, U. and Leck, C.: Importance of submicron surface-active organic aerosols for pristine Arctic clouds, *Tellus B*, 57, 261-268, <https://doi.org/10.1111/j.1600-0889.2005.00144.x>, 2005.
- Lopez-Yglesias, X. F., Yeung, M. C., Dey, S. E., Brechtel, F. J., and Chan, C. K.: Performance Evaluation of the Brechtel Mfg. Humidified Tandem Differential Mobility Analyzer (BMI HTDMA) for Studying Hygroscopic

- Properties of Aerosol Particles, *Aerosol Sci Tech*, 48, 969-980, <https://doi.org/10.1080/02786826.2014.952366>, 2014.
- 885 Martensson, E. M., Nilsson, E. D., de Leeuw, G., Cohen, L. H., and Hansson, H. C.: Laboratory simulations and parameterization of the primary marine aerosol production, *J Geophys Res-Atmos*, 108, 4297, <https://doi.org/10.1029/2002jd002263>, 2003.
- Martin, M., Chang, R. Y. W., Sierau, B., Sjogren, S., Swietlicki, E., Abbatt, J. P. D., Leck, C., and Lohmann, U.: Cloud condensation nuclei closure study on summer arctic aerosol, *Atmos Chem Phys*, 11, 11335-11350, 890 <https://doi.org/10.5194/acp-11-11335-2011>, 2011.
- Matsui, H., Kondo, Y., Moteki, N., Takegawa, N., Sahu, L. K., Koike, M., Zhao, Y., Fuelberg, H. E., Sessions, W. R., Diskin, G., Anderson, B. E., Blake, D. R., Wisthaler, A., Cubison, M. J., and Jimenez, J. L.: Accumulation-mode aerosol number concentrations in the Arctic during the ARCTAS aircraft campaign: Long-range transport of polluted and clean air from the Asian continent, *J Geophys Res-Atmos*, 116, D20217, 895 <https://doi.org/10.1029/2011jd016189>, 2011.
- Mauritsen, T., Sedlar, J., Tjernstrom, M., Leck, C., Martin, M., Shupe, M., Sjogren, S., Sierau, B., Persson, P. O. G., Brooks, I. M., and Swietlicki, E.: An Arctic CCN-limited cloud-aerosol regime, *Atmos Chem Phys*, 11, 165-173, <https://doi.org/10.5194/acp-11-165-2011>, 2011.
- Moffet, R. C. and Prather, K. A.: In-situ measurements of the mixing state and optical properties of soot with implications for radiative forcing estimates, *P Natl Acad Sci USA*, 106, 11872-11877, 900 <https://doi.org/10.1073/pnas.0900040106>, 2009.
- Mungall, E. L., Abbatt, J. P. D., Wentzell, J. J. B., Lee, A. K. Y., Thomas, J. L., Blais, M., Gosselin, M., Miller, L. A., Papakyriakou, T., Willis, M. D., and Liggio, J.: Microlayer source of oxygenated volatile organic compounds in the summertime marine Arctic boundary layer, *P Natl Acad Sci USA*, 114, 6203-6208, 905 <https://doi.org/10.1073/pnas.1620571114>, 2017.
- Myriokefalitakis, S., Vignati, E., Tsigaridis, K., Papadimas, C., Sciare, J., Mihalopoulos, N., Facchini, M. C., Rinaldi, M., Dentener, F. J., Ceburnis, D., Hatzianastasiou, N., O'Dowd, C. D., van Weele, M., and Kanakidou, M.: Global Modeling of the Oceanic Source of Organic Aerosols, *Adv Meteorol*, 10, 939171, <https://doi.org/10.1155/2010/939171>, 2010.
- 910 Nakao, S., Suda, S. R., Camp, M., Petters, M. D., and Kreidenweis, S. M.: Droplet activation of wet particles: development of the Wet CCN approach, *Atmos Meas Tech*, 7, 2227-2241, <https://doi.org/10.5194/amt-7-2227-2014>, 2014.
- Nguyen, Q. T., Glasius, M., Sorensen, L. L., Jensen, B., Skov, H., Birmili, W., Wiedensohler, A., Kristensson, A., Nojgaard, J. K., and Massling, A.: Seasonal variation of atmospheric particle number concentrations, new particle formation and atmospheric oxidation capacity at the high Arctic site Villum Research Station, Station Nord, *Atmos Chem Phys*, 16, 11319-11336, <https://doi.org/10.5194/acp-16-11319-2016>, 2016.
- 915 Nielsen, I. E., Skov, H., Massling, A., Eriksson, A. C., Dall'Osto, M., Junninen, H., Sarnela, N., Lange, R., Collier, S., Zhang, Q., Cappa, C. D., and Nojgaard, J. K.: Biogenic and anthropogenic sources of aerosols at the High Arctic site Villum Research Station, *Atmos Chem Phys*, 19, 10239-10256, <https://doi.org/10.5194/acp-19-10239-2019>, 2019.
- 920 Orellana, M. V., Matrai, P. A., Leck, C., Rauschenberg, C. D., Lee, A. M., and Coz, E.: Marine microgels as a source of cloud condensation nuclei in the high Arctic, *P Natl Acad Sci USA*, 108, 13612-13617, <https://doi.org/10.1073/pnas.1102457108>, 2011.
- Ovadnevaite, J., Ceburnis, D., Martucci, G., Bialek, J., Monahan, C., Rinaldi, M., Facchini, M. C., Berresheim, H., Worsnop, D. R., and O'Dowd, C.: Primary marine organic aerosol: A dichotomy of low hygroscopicity and high CCN activity, *Geophys Res Lett*, 38, L21806, <https://doi.org/10.1029/2011gl048869>, 2011.
- 925 Ovadnevaite, J., Zuend, A., Laaksonen, A., Sanchez, K. J., Roberts, G., Ceburnis, D., Decesari, S., Rinaldi, M., Hodas, N., Facchini, M. C., Seinfeld, J. H., and Dowd, C. O.: Surface tension prevails over solute effect in organic-influenced cloud droplet activation, *Nature*, 546, 637-641, <https://doi.org/10.1038/nature22806>, 2017.
- Petters, M. D. and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmos Chem Phys*, 7, 1961-1971, <https://doi.org/10.5194/acp-7-1961-2007>, 2007.
- 930 Pfeifer, S., Birmili, W., Schladitz, A., Muller, T., Nowak, A., and Wiedensohler, A.: A fast and easy-to-implement inversion algorithm for mobility particle size spectrometers considering particle number size distribution information outside of the detection range, *Atmos Meas Tech*, 7, 95-105, <https://doi.org/10.5194/amt-7-95-2014>, 2014.
- 935 Pilat, J. and Charlson, J.: Theoretical and optical studies of humidity effects on the size distribution of a hygroscopic aerosol, *J Rech Atmosphér*, 1, 165-170, 1966.
- Quinn, P. K., Shaw, G., Andrews, E., Dutton, E. G., Ruoho-Airola, T., and Gong, S. L.: Arctic haze: current trends and knowledge gaps, *Tellus B*, 59, 99-114, <https://onlinelibrary.wiley.com/doi/full/10.1111/j.1600-0889.2006.00238.x>, 2007

- 940 Quinn, P. K., Bates, T. S., Baum, E., Doubleday, N., Fiore, A. M., Flanner, M., Fridlind, A., Garrett, T. J., Koch, D.,
Menon, S., Shindell, D., Stohl, A., and Warren, S. G.: Short-lived pollutants in the Arctic: their climate impact and
possible mitigation strategies, *Atmos Chem Phys*, 8, 1723-1735, <https://doi.org/10.5194/acp-8-1723-2008>, 2008.
- 945 Quinn, P. K., Coffman, D. J., Johnson, J. E., Upchurch, L. M., and Bates, T. S.: Small fraction of marine cloud
condensation nuclei made up of sea spray aerosol, *Nat Geosci*, 10, 674-679, <https://doi.org/10.1038/Ngeo3003>,
2017.
- Rastak, N., Silvergren, S., Zieger, P., Wideqvist, U., Strom, J., Svenningsson, B., Maturilli, M., Tesche, M., Ekman, A.
M. L., Tunved, P., and Riipinen, I.: Seasonal variation of aerosol water uptake and its impact on the direct radiative
effect at Ny-Ålesund, Svalbard, *Atmos Chem Phys*, 14, 7445-7460, <https://doi.org/10.5194/acp-14-7445-2014>,
2014.
- 950 Rastak, N., Pajunoja, A., Acosta Navarro, J. C., Ma, J., Song, M., Partridge, D. G., Kirkevåg, A., Leong, Y., Hu, W. W.,
Taylor, N. F., Lambe, A., Cerully, K., Bougiatioti, A., Liu, P., Krejci, R., Petäjä, T., Percival, C., Davidovits, P.,
Worsnop, D. R., Ekman, A. M. L., Nenes, A., Martin, S., Jimenez, J. L., Collins, D. R., Topping, D. O., Bertram, A.
K., Zuend, A., Virtanen, A., and Riipinen, I.: Microphysical explanation of the RH-dependent water affinity of
biogenic organic aerosol and its importance for climate, *Geophys Res Lett*, 44, 5167-5177,
955 <https://doi.org/10.1002/2017GL073056>, 2017.
- Raut, J. C., Chazette, P., and Fortain, A.: Link between aerosol optical, microphysical and chemical measurements in an
underground railway station in Paris, *Atmos Environ*, 43, 860-868, <https://doi.org/10.1016/j.atmosenv.2008.10.038>,
2009.
- Rosati, B., Paul, A., Iversen, E. M., Massling, A., and Bilde, M.: Reconciling atmospheric water uptake by hydrate
forming salts, *Environ. Sci.: Processes Impacts*, 22, 1759-1767, <https://doi.org/10.1039/D0EM00179A>,
960 <http://dx.doi.org/10.1039/D0EM00179A>, 2020.
- Rose, D., Nowak, A., Achtert, P., Wiedensohler, A., Hu, M., Shao, M., Zhang, Y., Andreae, M. O., and Pöschl, U.:
Cloud condensation nuclei in polluted air and biomass burning smoke near the mega-city Guangzhou, China - Part
1: size-resolved measurements and implications for the modeling of aerosol particle hygroscopicity and CCN
965 activity, *Atmos. Chem. Phys.*, 10, 3365-3383, <https://doi.org/10.5194/acp-10-3365-2010>, 2010.
- Rempillo, O., Seguin, A. M., Norman, A. L., Scarratt, M., Michaud, S., Chang, R., Sjøstedt, S., Abbatt, J., Else, B.,
Papakyriakou, T., Sharma, S., Grasby, S., and Lévassieur, M.: Dimethyl sulfide air-sea fluxes and biogenic sulfur as
a source of new aerosols in the Arctic fall, *J Geophys Res-Atmos*, 116, D00S04,
970 <https://doi.org/10.1029/2011jd016336>, 2011.
- Schmale, J., Henning, S., Decesari, S., Henzing, B., Keskinen, H., Sellegri, K., Ovadnevaite, J., Pohlker, M. L., Brito,
J., Bougiatioti, A., Kristensson, A., Kalivitis, N., Stavroulas, I., Carbone, S., Jefferson, A., Park, M., Schlag, P.,
Iwamoto, Y., Aalto, P., Aijala, M., Bukowiecki, N., Ehn, M., Frank, G., Frohlich, R., Frumau, A., Herrmann, E.,
Herrmann, H., Holzinger, R., Kos, G., Kulmala, M., Mihalopoulos, N., Nenes, A., O'Dowd, C., Petaja, T., Picard,
975 D., Pohlker, C., Poschl, U., Poulain, L., Prevot, A. S. H., Swietlicki, E., Andreae, M. O., Artaxo, P., Wiedensohler,
A., Ogren, J., Matsuki, A., Yum, S. S., Stratmann, F., Baltensperger, U., and Gysel, M.: Long-term cloud
condensation nuclei number concentration, particle number size distribution and chemical composition
measurements at regionally representative observatories, *Atmos Chem Phys*, 18, 2853-2881,
980 <https://doi.org/10.5194/acp-18-2853-2018>, 2018.
- Schmale, J., Zieger, P., and Ekman, A. M. L.: Aerosols in current and future Arctic climate, *Nature Climate Change*, 11,
95-105. <https://doi.org/10.1038/s41558-020-00969-5>, 2021.
- Schmale, J., and Baccarini, A.: Progress in unraveling atmospheric new particle formation and growth across the Arctic,
Geophysical Research Letters, 48, e2021GL094198. <https://doi.org/10.1029/2021GL094198>, 2021.
- 985 Schmale, J., Sharma, S., Decesari, S., Pernov, J., Massling, A., Hansson, H.-C., von Salzen, K., Skov, H., Andrews, E.
B., Quinn, T., Upchurch, L., Eleftheriadis, K., and Traversi, R.: Pan-Arctic seasonal cycles and long-term trends of
aerosol properties from 10 observatories, *Atmos. Chem. and Phys.*, 22 (5), 3067-3096, 2022.
- Shaw, G. E.: The arctic haze phenomenon, *B Am Meteorol Soc*, 76, 2403-2413, [https://doi.org/10.1175/1520-0477\(1995\)076<2403:TAHP>2.0.CO;2](https://doi.org/10.1175/1520-0477(1995)076<2403:TAHP>2.0.CO;2), 1995.
- Shaw, P. M., Russell, L. M., Jefferson, A., and Quinn, P. K.: Arctic organic aerosol measurements show particles from
mixed combustion in spring haze and from frost flowers in winter, *Geophys Res Lett*, 37, L10803,
990 <https://doi.org/10.1029/2010gl042831>, 2010.
- Silvergren, S., Wideqvist, U., Strom, J., Sjogren, S., and Svenningsson, B.: Hygroscopic growth and cloud forming
potential of Arctic aerosol based on observed chemical and physical characteristics (a 1 year study 2007-2008), *J
Geophys Res-Atmos*, 119, 14080-14097, <https://doi.org/10.1002/2014JD021657>, 2014.
- 995 Skov, H. Massling, A. Nielsen, I.E. Nordstrom, C. Bossi, R. Katrin Vorkamp, K. Christensen, J. Larsen, M.M. Hansen,
K.M. Liisberg, J.B. and Poulsen, M.B. 2017. AMAP CORE – ATMOSPHERIC PART. Results from Villum
Research Station. Aarhus University, Technical Report from DCE – Danish Centre for Environment and Energy No.
101, 77 pp., <http://dce2.au.dk/pub/TR101.pdf>, 2017.

- Stohl, A.: Characteristics of atmospheric transport into the Arctic troposphere, *J Geophys Res-Atmos*, 111, D11306, <https://doi.org/10.1029/2005jd006888>, 2006.
- 1000 Stolzenburg, M. R. and McMurry, P. H.: Equations governing single and tandem DMA configurations and a new lognormal approximation to the transfer function, *Aerosol Sci Tech*, 42, 421-432, <https://doi.org/10.1080/02786820802157823>, 2008.
- Titos, G., Cazorla, A., Zieger, P., Andrews, E., Lyamani, H., Granados-Munoz, M. J., Olmo, F. J., and Alados-Arboledas, L.: Effect of hygroscopic growth on the aerosol light-scattering coefficient: A review of measurements, techniques and error sources, *Atmos Environ*, 141, 494-507, <https://doi.org/10.1016/j.atmosenv.2016.07.021>, 2016.
- 1005 Tremblay, S., Picard, J. C., Bachelder, J. O., Lutsch, E., Strong, K., Fogal, P., Leaitch, W.R., Sharma, S., Kolonjari, F., Cox, C.J., Chang, R.Y.W., and Hayes, P.L.: Characterization of aerosol growth events over Ellesmere Island during the summers of 2015 and 2016, *Atmos Chem Phys*, 19, 5589-5604, <https://doi.org/10.5194/acp-19-5589-2019>, 2019.
- 1010 Tunved, P., Strom, J., and Krejci, R.: Arctic aerosol life cycle: linking aerosol size distributions observed between 2000 and 2010 with air mass transport and precipitation at Zeppelin station, Ny-Ålesund, Svalbard, *Atmos Chem Phys*, 13, 3643-3660, <https://doi.org/10.5194/acp-13-3643-2013>, 2013.
- Twomey, S.: Pollution and the Planetary Albedo, *Atmos Environ*, 8, 1251-1256, [https://doi.org/10.1016/0004-6981\(74\)90004-3](https://doi.org/10.1016/0004-6981(74)90004-3), 1974.
- 1015 Wexler, A. S. and Clegg, S. L.: Atmospheric aerosol models for systems including the ions H^+ , NH_4^+ , Na^+ , SO_4^{2-} , NO_3^- , Cl^- , Br^- , and H_2O , *J Geophys Res-Atmos*, 107, ACH 14-1-ACH 14-14, <https://doi.org/10.1029/2001jd000451>, 2002.
- Wiedensohler, A., Birmili, W., Nowak, A., Sonntag, A., Weinhold, K., Merkel, M., Wehner, B., Tuch, T., Pfeifer, S., Fiebig, M., Fjaraa, A. M., Asmi, E., Sellegri, K., Depuy, R., Venzac, H., Villani, P., Laj, P., Aalto, P., Ogren, J. A., Swietlicki, E., Williams, P., Roldin, P., Quincey, P., Hüglin, C., Fierz-Schmidhauser, R., Gysel, M., Weingartner, E., Riccobono, F., Santos, S., Gruning, C., Faloon, K., Beddows, D., Harrison, R. M., Monahan, C., Jennings, S. G., O'Dowd, C. D., Marinoni, A., Horn, H. G., Keck, L., Jiang, J., Scheckman, J., McMurry, P. H., Deng, Z., Zhao, C. S., Moerman, M., Henzing, B., de Leeuw, G., Loschau, G., and Bastian, S.: Mobility particle size spectrometers: harmonization of technical standards and data structure to facilitate high quality long-term observations of atmospheric particle number size distributions, *Atmos Meas Tech*, 5, 657-685, <https://doi.org/10.5194/amt-5-657-2012>, 2012.
- 1025 Willis, M. D., Burkart, J., Thomas, J. L., Kollner, F., Schneider, J., Bozem, H., Hoor, P. M., Aliabadi, A. A., Schulz, H., Herber, A. B., Leaitch, W. R., and Abbatt, J. P. D.: Growth of nucleation mode particles in the summertime Arctic: a case study, *Atmos Chem Phys*, 16, 7663-7679, <https://doi.org/10.5194/acp-16-7663-2016>, 2016.
- 1030 Willis, M. D., Köllner, F., Burkart, J., Bozem, H., Thomas, J. L., Schneider, J., Aliabadi, A. A., Hoor, P. M., Schulz, H., Herber, A. B., Leaitch, W. R., and Abbatt, J. P. D.: Evidence for marine biogenic influence on summertime Arctic aerosol, *Geophys Res Lett*, 44, 6460-6470, <https://doi.org/10.1002/2017gl073359>, 2017.
- Xu, W., Ovadnevaite, J., Fossom, K.N., Lin, C., Huang, R.-J., Ceburnis, D., and O'Dowd, C.: Sea spray as an obscured source for marine cloud nuclei, *Nat. Geosci.* 15, 282-286, <https://doi.org/10.1038/s41561-022-00917-2>, 2022.
- 1035 Yang, Q., Bitz, C. M., and Doherty, S. J.: Offsetting effects of aerosols on Arctic and global climate in the late 20th century, *Atmos Chem Phys*, 14, 3969-3975, <https://doi.org/10.5194/acp-14-3969-2014>, 2014.
- Zabori, J., Rastak, N., Yoon, Y. J., Riipinen, I., and Strom, J.: Size-resolved cloud condensation nuclei concentration measurements in the Arctic: two case studies from the summer of 2008, *Atmos Chem Phys*, 15, 13803-13817, <https://doi.org/10.5194/acp-15-13803-2015>, 2015.
- 1040 Zhao, C. F. and Garrett, T. J.: Effects of Arctic haze on surface cloud radiative forcing, *Geophys Res Lett*, 42, 557-564, <https://doi.org/10.1002/2014gl062015>, 2015.
- Zhou, J. C., Swietlicki, E., Berg, O. H., Aalto, P. P., Hameri, K., Nilsson, E. D., and Leck, C.: Hygroscopic properties of aerosol particles over the central Arctic Ocean during summer, *J Geophys Res-Atmos*, 106, 32111-32123, <https://doi.org/10.1029/2000jd900426>, 2001.
- 1045 Zieger, P., Fierz-Schmidhauser, R., Gysel, M., Ström, J., Henne, S., Yttri, K. E., Baltensperger, U., and Weingartner, E.: Effects of relative humidity on aerosol light scattering in the Arctic, *Atmos. Chem. Phys.*, 10, 3875-3890, <https://doi.org/10.5194/acp-10-3875-2010>, 2010.