

Impact of formulations of the homogeneous nucleation rate on ice nucleation events in cirrus

Peter Spichtinger¹, Patrik Marschalik^{1,2}, and Manuel Baumgartner^{1,3,4}

¹ Institute for Atmospheric Physics, Johannes Gutenberg University Mainz, Germany

² now at DB Systel GmbH, Berlin, Germany

³ Zentrum für Datenverarbeitung, Johannes Gutenberg University Mainz, Germany

⁴ now at German Weather Service (DWD), Offenbach, Germany

Correspondence: Peter Spichtinger (peter.spichtinger@uni-mainz.de)

Abstract. Homogeneous freezing of solution droplets is an important pathway of ice formation in the tropopause region. The nucleation rate can be parameterised as a function of water activity, based on empirical fits and some assumptions on the underlying properties of super-cooled water, although a general theory is missing. It is not clear how nucleation events are influenced by the exact formulation of the nucleation rate or even their inherent uncertainty. In this study we investigate the formulation of the nucleation rate of homogeneous freezing of solution droplets (1) to link the formulation to the nucleation rate of pure water droplets, (2) to derive a robust and simple formulation of the nucleation rate, and (3) to determine the impact of variations in the formulation on nucleation events. The nucleation rate can be adjusted and the formulation can be simplified to a threshold description. We use a state-of-the-art bulk ice microphysics model to investigate nucleation events as driven by constant cooling rates; the key variables are the final ice crystal number concentration and the maximum supersaturation during the event. The nucleation events are sensitive to the slope of the nucleation rate but only weakly affected by changes in its absolute value. This leads to the conclusion that details of the nucleation rate are less important for simulating ice nucleation in bulk models as long as the main feature of the nucleation rate (i.e. its slope) is represented sufficiently well. The weak sensitivity on the absolute values of the nucleation rate suggests that the amount of available solution droplets also does not crucially affect nucleation events. The use of only one distinct nucleation threshold function for analysis and model parameterisation should be reinvestigated, since it corresponds to a very high nucleation rate value, which is not reached in many nucleation events with low vertical updrafts. In contrast, the maximum supersaturation and thus the nucleation thresholds reached during an ice nucleation event depend on the vertical updraft velocity or cooling rate. This feature might explain some high supersaturation values during nucleation events in cloud chambers and suggests a reformulation of ice nucleation schemes used in coarse models based on a purely temperature-dependent nucleation threshold.

20 1 Introduction

Clouds are one of the most important components in the Earth-Atmosphere system. They influence the hydrological cycle and Earth's energy balance via interaction with radiation. Clouds can cool the system by partly scattering and reflecting incoming solar radiation (albedo effect) but also warm the atmosphere by absorbing and re-emitting thermal radiation as emitted by the

Earth's surface (greenhouse effect). While for liquid clouds a net cooling effect can be derived, the radiative effect for clouds
25 containing ice crystals is still under debate. In particular, for pure ice clouds (so-called cirrus clouds) at high altitudes in the
low temperature range ($T < 235$ K) albedo effect and greenhouse effect are of the same order of magnitude but admit different
signs, leading to different net-effects (see, e.g., Fusina et al., 2007; Joos et al., 2014; Gasparini et al., 2017). Thus, details
in microphysical properties of ice crystals might decide about a net warming or cooling of cirrus clouds, as can be seen in
former model studies (e.g. Zhang et al., 1999). A key aspect of ice crystals is their size which directly affects the scattering
30 and absorption of radiation. Smaller crystals scatter incoming solar light more effectively, thus the optical depth τ is directly
dependent on the size, as can be seen in the usual approximation (cf., e.g., Fu and Liou, 1993)

$$\tau = \text{IWC} \cdot \Delta z \cdot \left(a + \frac{b}{D_e} \right), \quad (1)$$

where D_e denotes the effective diameter of the crystal, IWC is the ice water content, Δz represents the vertical extent of the
cloud, and a, b are empirically derived constants. Since the available water vapour is mainly determined by thermodynamic
35 conditions, the pathway of ice nucleation often decides about the ice crystal number concentration in cirrus clouds and thus
their effective size (assuming a certain amount of available water vapour).

Ice crystals can be formed by very different nucleation processes, which can be grouped into two major pathways, namely *in situ*
and *liquid origin* ice formation (e.g. Krämer et al., 2016; Luebke et al., 2016; Wernli et al., 2016). The overall term *in situ*
formation refers to ice formation at humidities below water saturation, whereas *liquid origin formation* subsumes all formation
40 processes where cloud droplets are present and humidity is close to water saturation (e.g. freezing of cloud droplets), see the
definition in Wernli et al. (2016). It is well known, that the ice crystal number concentration varies crucially in dependence
on the underlying nucleation process, leading to potentially strong changes in the resulting radiative effect (see, e.g., Krämer
et al., 2020).

Despite of the availability of many observational data and laboratory experiments (e.g. Hoose and Möhler, 2012), and also the
45 development of new theoretical models (e.g. the soccer ball model, see Niedermeier et al., 2011), the details of ice nucleation
at the molecular scale are still unknown.

A special situation occurs for the so-called homogeneous freezing of super-cooled solution droplets (also short: homoge-
neous nucleation) at cold temperatures below 235 K. This process describes the spontaneous freezing of supercooled aqueous
solution particles containing a small amount of (usually inorganic) substances. Albeit also the details of this freezing process
50 are not completely understood on a molecular scale, reproducible laboratory experiments allowed the formulation of an em-
pirical fit for the nucleation rate (Koop et al., 2000). Such a fit bears inherent but maybe also unknown uncertainty, since we
have no generally accepted theory for comparison. Other fits or a change in the fit parameters might also lead to different
formulations of nucleation rates.

A priori, it is not clear how large the impact of the formulation of nucleation rates might be on simulating nucleation
55 events in models resolving nucleation events in time. This issue is the starting point of our investigation. We want to address
three different aspects. First, we want to link the former formulation by Koop et al. (2000) to recent findings on pure water
in order to formulate a consistent framework for our models. Second, we want to derive a robust and simple formulation

of the homogeneous nucleation rates, which can be used for analytical as well as numerical investigations. Third, we want to investigate the impact of variations of nucleation rates (based on the new formulation) on nucleation events, i.e. on the resulting ice crystal number concentrations.

From theory (e.g. Baumgartner and Spichtinger, 2019) and former idealized box model simulations (e.g. Kärcher and Lohmann, 2002; Ren and Mackenzie, 2005; Spichtinger and Gierens, 2009), we know that ice crystal numbers as produced in homogeneous nucleation events driven by a constant cooling rate (equivalent to a constant vertical velocity) crucially depend on several parameters and, thus, affect also the radiative properties of the formed ice cloud (see, e.g., calculations in Krämer et al., 2020; Joos et al., 2009). Therefore, it is of high importance to understand the impact of the formulation of nucleation rates on the resulting ice crystal number concentrations.

We emphasize that all our investigations are meant in a bulk-sense, i.e. only integrated quantities such as the ice crystal number and (total) ice crystal mass are considered. Using this approach, we consider the case of a newly forming cirrus cloud and do not focus on the freezing or forming details of single ice crystals.

The study is structured as follows. In the next section, we present the fit by Koop et al. (2000) and its empirical basis, as related to water theories. In section 3 we describe the simple model used for idealized simulations for testing the impact of different formulations of nucleation rates. In section 4 the more compact formulation of the nucleation rate along with several approximations is discussed. The consequences of using the proposed approximations are explored by idealized numerical simulations. In section 5 we investigate the impact of a recently proposed formulation of the saturation vapour pressure over super-cooled liquid water on the nucleation events (Nachbar et al., 2019). In section 6 a new formulation of the nucleation rate based on results for freezing of pure super-cooled water (Koop and Murray, 2016) is presented and its impact on the number concentration of nucleated ice crystals is discussed. In section 7 we investigate thresholds of ice nucleation as well as the peak values of supersaturation during nucleation events. Finally, we summarize the results and draw some conclusions in section 8.

2 Empirical fit of the nucleation rate

Nucleation events are investigated in the phase space spanned by temperature and water activity of the aqueous solution. The latter is defined as the ratio of saturation pressures of water vapour over the solution p_{sol} and pure water p_{liq} , as $a_w := \frac{p_{\text{sol}}}{p_{\text{liq}}}$. In this representation, the melting curve for different inorganic solutions turns out to be solely temperature dependent, i.e. $a_w^i(T) := a_w(T_m) = \frac{p_{\text{ice}}(T)}{p_{\text{liq}}(T)}$ (cf. Koop, 2015, his eq.(5)), where p_{ice} denotes the saturation vapour pressure over ice. The important insight here is that also the freezing/nucleation events collapse to a single line in the diagram (see Koop et al., 2000; Koop, 2004, 2015), which can be fitted by shifting the melting curve (deviation $\Delta a_w \sim 0.305$). This also means that the nucleation events do not depend on the solute, which is at least true for most inorganic substances (see, e.g., Koop, 2004). Thus, the nucleation rate can be solely parameterized as a function of $\Delta a_w = a_w - a_w^i$. For the fitting procedure in Koop et al. (2000), a polynomial of degree 3 is used and results in the formulation

$$J_{\text{sol}}(\Delta a_w) = 10^{p_3(\Delta a_w)} \quad \text{with} \quad p_3(x) = \sum_{k=0}^3 a_k x^k \quad (2)$$

90 of the homogeneous nucleation rate coefficient J_{sol} . The nucleation rate coefficient is used to formulate the probability of freezing of aqueous solution droplets. The fit was used in the spirit of the representation of the nucleation rate for pure water as derived by Pruppacher (1995). During this time, three water theories were available, and the nucleation rate (as a cubic polynomial) was chosen according to the stability limit hypothesis (e.g. Mishima and Stanley, 1998), leading to an unlimited increase in the rate (see, e.g., Pruppacher, 1995, his figure 3). However, meanwhile this water theory can be ruled out by
 95 experimental evidence, thus only the two other water theories remain (singularity-free hypothesis vs. liquid-liquid critical point, cf. Gallo et al., 2019, 2016), which do not admit an unlimited increase in nucleation rates of pure water (see, e.g., Koop and Murray, 2016). Thus, the heuristic basis for choosing a cubic polynomial as a fit is not valid anymore.

Note that for atmospheric relevant conditions, both remaining water theories produce essentially the same results. Only at very low temperatures $T < 150$ K, where highly viscous or even glassy states of water occur, a different behaviour is predicted.
 100 Such temperatures are not relevant for investigations of ice clouds in the tropopause region, where homogeneous freezing of solution droplets takes place. However, these theories provide the basis for the formulation of the saturation vapour pressure over supercooled water in the no man's land (Murphy and Koop, 2005), combining heat capacities of liquid water and amorphous ice.

Finally, using the assumption of solution droplets being in equilibrium with their environment and neglecting size effects,
 105 water activity equals the liquid water saturation ratio S_{liq} due to

$$a_w = \frac{p_{\text{sol}} \stackrel{\text{in eq.}}{=} p_v}{p_{\text{liq}}} = S_{\text{liq}} \quad (3)$$

where p_v denotes the partial water vapour pressure. Using this representation of a_w together with the ice saturation ratio $S_i = \frac{p_v}{p_{\text{ice}}}$, the computation

$$\begin{aligned} \Delta a_w &= a_w - a_w^i = \frac{p_v}{p_{\text{liq}}(T)} - \frac{p_{\text{ice}}(T)}{p_{\text{liq}}(T)} = (S_i - 1) \frac{p_{\text{ice}}(T)}{p_{\text{liq}}(T)} \\ &= (S_i - 1) a_w^i(T) \end{aligned} \quad (4)$$

110 shows that Δa_w only depends on the ice saturation ratio and temperature.

Note, that although recent measurements (Pathak et al., 2021) corroborate the procedure in the study by Murphy and Koop (2005), in a recent study by Nachbar et al. (2019) the combination of liquid water and amorphous ice is called into question, leading to a different formulation of the saturation vapour pressure over supercooled water and thus a different water activity. In the following investigations, we will also use this formulation in order to determine the sensitivity of the nucleation events on
 115 the choice of a saturation vapour pressure formulation. Note, that for each choice the water activity $a_w^i(T)$ must be recalculated.

3 Model description

We begin with the description of the governing equations for the relevant ice processes in a nucleation event, i.e. homogeneous nucleation and diffusional growth. Both processes are key for determining the properties of the nucleation event, such as the number of nucleated ice crystals and the evolution of the ice saturation ratio (e.g. its peak value). Of course, other processes

120 such as sedimentation and aggregation of ice crystals are important for the evolution of ice clouds, but usually act on longer time scales, e.g., when the particles are grown to larger sizes. Thus, we omit these processes and concentrate on nucleation and growth, as in former studies (e.g. Kärcher and Lohmann, 2002; Baumgartner and Spichtinger, 2019).

We formulate the model in terms of averaged quantities for ice crystal mass and number concentration (q_i, n_i), i.e. as a 2-moment scheme. Additionally, the saturation ratio with respect to hexagonal ice, $S_i = \frac{p_v}{p_{\text{ice}}(T)}$, is used, with the partial water vapour pressure p_v and the saturation water vapour pressure over hexagonal ice, $p_{\text{ice}}(T)$. Thus, the complete set of equations
 125 for an adiabatically ascending air parcel can be represented as

$$\dot{n}_i = \text{Nuc}_n \quad (5)$$

$$\dot{q}_i = \text{Nuc}_q + \text{Dep}_q \quad (6)$$

$$\dot{S}_i = \text{Cool} + \text{Dep}_s \quad (7)$$

$$130 \quad \dot{T} = \left. \frac{dT}{dt} \right|_{\text{adiabatic}} + \left. \frac{dT}{dt} \right|_{\text{diabatic}} = -\frac{g}{c_p} w + \frac{L}{c_p} \left. \frac{dq_i}{dt} \right|_{\text{phase}} = -\frac{g}{c_p} w + \frac{L}{c_p} (\text{Nuc}_q + \text{Dep}_q) \quad (8)$$

$$\dot{p} = \left. \frac{dp}{dt} \right|_{\text{adiabatic}} = -g\rho w, \quad (9)$$

including changes of temperature T and pressure p . In these equations, w denotes the vertical velocity of the air parcel, c_p is the specific heat capacity of dry air (assumed as a constant, see, Baumgartner et al., 2020), L denotes the (constant) latent heat of sublimation, and ρ is the air density. The assumption of an ideal gas is adopted for air and water vapour. The terms $\text{Nuc}_n, \text{Nuc}_q$
 135 denote changes due to nucleation, the terms $\text{Dep}_q, \text{Dep}_s$ describe changes due to diffusional growth of ice crystals. The term Cool denotes the impact of adiabatic expansion due to upward motion with velocity w , this is also reflected in the change of temperature and pressure, using adiabatic lapse rate and hydrostatic pressure, respectively. For temperature, we would have to consider diabatic changes due to latent heat release in phase changes.

Computing the total derivative of the saturation ratio using the representation $S_i = \frac{p q_v}{\varepsilon_0 p_{s,i}(T)}$, where ε_0 denotes the ratio of
 140 molar masses of water and dry air, together with the Clausius-Clapeyron equation yields

$$\text{Cool} = \frac{\partial S_i}{\partial T} \left. \frac{dT}{dt} \right|_{\text{adiabatic}} + \frac{\partial S_i}{\partial p} \left. \frac{dp}{dt} \right|_{\text{adiabatic}} = \left[\frac{Lg}{c_p R_v T^2} - \frac{g}{R_a T} \right] S_i w \quad (10)$$

and

$$\text{Dep}_s = \frac{\partial S_i}{\partial T} \left. \frac{dT}{dt} \right|_{\text{diabatic}} + \frac{\partial S_i}{\partial q_v} (\text{Nuc}_q + \text{Dep}_q) = - \left[\frac{L^2}{c_p R_v T^2} + \frac{1}{q_v} \right] S_i (\text{Nuc}_q + \text{Dep}_q). \quad (11)$$

To a good approximation, for cold temperatures the first term in the bracket in (11), which describes latent heat release due to
 145 phase changes, can be omitted. In the following, we will omit the evolution equations (8), (9) for temperature and pressure, i.e. we assume these as being constant during the nucleation event. Thus, we arrive at

$$\text{Dep}_s \approx - \frac{p}{\varepsilon_0 p_{s,i}} (\text{Nuc}_q + \text{Dep}_q) \quad (12)$$

As a result of assuming temperature and pressure as being constant, only the vertical velocity w is an external parameter for the supersaturation. For the terms $\text{Nuc}_x, \text{Dep}_x$ ($x = n, q, s$) we have to keep temperature and pressure as fixed parameters

150 $T = T_{\text{env}}, p = p_{\text{env}}$. This approach was also used in former investigations (see, e.g., Spreitzer et al., 2017; Baumgartner and Spichtinger, 2019).

The nucleation term can be described as

$$\text{Nuc}_n = J_{\text{nuc}} V_d n_a, \quad \text{Nuc}_q = m_0 \text{Nuc}_n \quad (13)$$

where, V_d is the mean volume of a supercooled solution droplet, n_a is the number concentration of solution droplets, and m_0 is
 155 the mean mass of a newly frozen solution droplet, which can be set to $m_0 = 10^{-16}$ kg. The nucleation rate for the homogeneous freezing of solution droplets is denoted by J_{nuc} . For comparison with former investigations (Kärcher and Lohmann, 2002; Spichtinger and Gierens, 2009), we set the number concentration of the background aerosol to a quite large value of $n_a \rho = 10^4 \text{ cm}^{-3} = 10^{10} \text{ m}^{-3}$; since the resulting ice crystal number concentration as produced in nucleation events is usually some orders of magnitude smaller, we do not have to care about a possible consumption of a major part fraction of solution droplets.
 160 We will later discuss the impact of this value in terms of nucleation events.

The diffusional growth of ice crystals is determined by the growth rate

$$\text{Dep}_q = n_i \cdot 4\pi D_v^* C G_v (S_i - 1) f_v \quad (14)$$

with the diffusion constant for water vapour in air $D_v^* = D_v(p, T) f_D$ as corrected by the factor f_D for the kinetic regime, the capacity of ice crystals, C , assuming columnar shape, the Howell factor $G_v(p, T)$ describing the impact of latent heat, and the
 165 ventilation correction f_v , respectively. Note, that the capacity also depends on the mean mass of the ice crystal ensemble, i.e. $C = C(\bar{m}) = C(n_i, q_i)$. The details of the formulation are given in appendix A.

Combining the expressions from above, the reduced system of equations reads

$$\dot{n}_i = \text{Nuc}_n \quad (15)$$

$$\dot{q}_i = \text{Nuc}_q + \text{Dep}_q \quad (16)$$

$$170 \quad \dot{S}_i = \left[\frac{Lg}{c_p R_v T^2} - \frac{g}{R_a T} \right] S_i w + \frac{p}{\varepsilon_0 p_{\text{si}}} (\text{Nuc}_q + \text{Dep}_q) \quad (17)$$

Remarks:

1. As shown in Spreitzer et al. (2017), it is possible to determine and characterize the steady states of the reduced system, which additionally includes sedimentation. This leads to a nonlinear oscillator with a bifurcation diagram, depending on the updraft velocity w , and on the temperature T .
- 175 2. The usefulness of this simple double moment scheme depends on the scales of the scenarios. We generally found good agreement with such parcel models and also on an LES scale (and even coarser resolution) with observations, more sophisticated models, and also theory (see, e.g., Spichtinger and Gierens, 2009; Spichtinger, 2014; Baumgartner et al., 2022).

4 Investigations of the nucleation rates

180 Investigations of ice clouds in the cold temperature regime ($T < 235$ K) need to include the nucleation process of homogeneous
freezing of aqueous solution droplets. As pointed out in section 1 the formulation by Koop et al. (2000) based on water activity
is a meaningful fit to experimental data. However, for theoretical investigations and the use in reduced order models, a simpler
but still accurate approximation would be helpful. In the following we present a way how to derive such an approximation
185 based on the original fit through measurements by Koop et al. (2000) in addition to recent observations for pure super-cooled
water.

4.1 Correction of the nucleation rate

In the study by Koop and Murray (2016) a parametrisation of the nucleation rate of pure supercooled water $J_{\text{pure liq}}(T)$ was
derived, based on recent measurements. Thus, in the context of homogeneous freezing of solution droplets, the nucleation rate
for pure water particles should coincide with the nucleation rate of solution droplets J_{sol} at water saturation, i.e. the condition

$$190 \quad J_{\text{sol}}(\Delta a_w^*) \stackrel{RH=1}{\equiv} J_{\text{pure liq}}(T) \quad (18)$$

should hold for a value Δa_w^* at water saturation, as claimed and already used in the study by Koop et al. (2000). However,
evaluating these two formulations of the nucleation rates at water saturation shows a similar qualitative behaviour down to
temperatures $T \sim 235$ K but a quantitative disagreement, see the blue and black curve in Figure 1. A reasonable requirement
is that the values of both formulations should match in the temperature range $235 \text{ K} \leq T \leq 240 \text{ K}$, since this range is relevant
195 for the freezing of pure water cloud droplets with reasonable sizes. This temperature range at water saturation is equivalent to
the range of water activity difference $0.27 \leq \Delta a_w \leq 0.31$. The offset between the curves and can be corrected by shifting the
logarithm of the nucleation rate for solution droplets by a constant value. The value of the shift was calculated by minimizing
the square distance between the curves in the respective temperature range. Thus, the corrected nucleation rate for aqueous
solution droplets reads as

$$200 \quad \log_{10}(J_{\text{sol,new}}(\Delta a_w)) = \log_{10}(J_{\text{sol}}(\Delta a_w)) + \delta \quad (19)$$

with $\delta = -1.522$. The nucleation rates are given in SI units (as used for all quantities throughout this study), i.e. $[J] = \text{m}^{-3} \text{s}^{-1}$.

Remarks:

- The nucleation rate of pure water droplets can be used for a direct parametrisation of the nucleation rate of aqueous
solution droplets. This will be carried out in section 6.1.
- 205 – The (new) disagreement (or small shift) of the rates solely stems from the comparison with the new formulation of Koop
and Murray (2016), since originally the nucleation rate for solution droplets was chosen in agreement with measurements
of nucleation rates for pure water droplets (Koop et al., 2000).
- In the following we will refer to the corrected nucleation rate as “reference” nucleation rate, since, to the best of our
knowledge, it provides the best and most recent fit for the homogenous nucleation rate of solution particles.




Figure 1. Nucleation rates for pure super-cooled water droplets (Koop and Murray, 2016, red) and aqueous solution droplets (Koop et al., 2000) at water saturation (i.e. infinitely dissolved); original values by Koop et al. (2000) in blue, shifted values ($\delta = -1.522$) in black (new reference nucleation rate $J_{\text{sol,new}}$). For the calculation, the saturation vapour pressure formulae by Murphy and Koop (2005) are used.

210 4.2 Nucleation rate as a function of T and S_i

The general strategy of the study is to represent the exponent of the nucleation rate by low order polynomials in a thermodynamic variable x , i.e.

$$J = 10^{p_n(x)}, \quad p_n(x) := \sum_{k=0}^n a_k x^k, \quad \deg p_n = n. \quad (20)$$

For instance, the formulation of the nucleation rate for aqueous solution droplets by Koop et al. (2000) is based on a polynomial
215 of degree three, i.e.

$$J_{\text{sol}}(\Delta a_w) = 10^{p_3(\Delta a_w)}, \quad p_3(x) = \sum_{k=0}^3 a_k x^k \quad (21)$$

using the thermodynamic quantity $x = \Delta a_w = a_w - a_w^i$.

Note, also the nucleation rate $J_{\text{pure liq}}$ for pure water droplets is based on the same structure, i.e. $\log_{10}(J_{\text{pure liq}})$ is a polynomial of order 6 in the thermodynamic variable T (cf. Koop and Murray, 2016). For analytical investigations of the homogeneous
220 nucleation, it is desirable to represent $\log_{10}(J)$ by a polynomial with low degree. As will be shown in the following, the

formulation

$$\log_{10}(J) \approx p_n(x), \quad n = \deg p_n \leq 2 \quad (22)$$

with a polynomial yields sufficient agreement with the reference. For analytical investigations (e.g. using asymptotic analysis) it is helpful to represent the nucleation rate using a threshold for the humidity to account for the explosive character of nucleation events as used in the analysis by Baumgartner and Spichtinger (2019). Thus, for the nucleation rate for super-cooled solution droplets we make the following ansatz

$$J = 10^{p_n(x)} = \exp(A(T)(S_i - S_c) + B(T)(S_i - S_c)^2) \quad (23)$$

where $S_c = S_c(T)$ is the temperature-dependent threshold value for the saturation ratio. Note, that the ansatz is consistent (or even equivalent) with condition (22). In order to describe J as a function of S_i and T we reformulate Δa_w as

$$\Delta a_w = (S_i - 1)a_w^i(T) = (S_i - S_c)a_w^i(T) + (S_c - 1)a_w^i(T) \quad (24)$$

using a threshold $S_c(T)$ that corresponds to a fixed value J_0 of the nucleation rate, i.e. $J(S_c(T), T) = J_0$. Taking the logarithm, this equality implies $p_n(x_0) = j_0 = \log_{10}(J_0)$ with $x = \Delta a_w$. As in former studies (see, e.g., Koop et al., 2000; Kärcher and Lohmann, 2002), we choose $J_0 = 10^{16} \text{ m}^{-3} \text{ s}^{-1} = 10^{10} \text{ cm}^{-3} \text{ s}^{-1}$. Note, that this choice for the parameterisation is quite arbitrary and has no strict physical interpretation, although one can argue with the cooling rates of the underlying experiments and thus with the probability of the freezing of droplets with a given volume within a certain predefined time interval (Koop et al., 2000).

Evaluating eq. (24) at $S_i = S_c$, we arrive at

$$p_n^{-1}(j_0) = x_0 = (S_i - S_c)a_w^i(T) + (S_c - 1)a_w^i(T) \quad (25)$$

$$\stackrel{S_i \equiv S_c}{=} (S_c - 1)a_w^i(T)$$

leading to a description of the threshold

$$S_c = \frac{1}{a_w^i(T)} p_n^{-1}(j_0) + 1 \quad \text{and} \quad (26)$$

$$\Delta a_w = (S_i - S_c)a_w^i(T) + p_n^{-1}(j_0)$$

if the polynomial $p_n(x)$ can be inverted in the relevant range $0.26 \leq \Delta a_w \leq 0.34$. Combining the equations from above, the nucleation rate can be represented as

$$\log_{10} J = p(\Delta a_w) = p_n((S_i - S_c)a_w^i(T) + p_n^{-1}(j_0)) \quad (27)$$

which is a threshold description using the thermodynamic variables S_i, T . This representation amounts to a reformulation of the original approximation, if the inverse function $p_n^{-1}(x)$ exists in the relevant range (i.e. $p_n(x)$ is strictly monotonic). In the following we consider the case of linear and quadratic polynomials, as determined by the ansatz (23).

1. Case of a linear polynomial $p_1(x) = a_0 + a_1x$

The inverse function of $p_1(x) = y$ is given by $p_1^{-1}(y) = \frac{y - a_0}{a_1}$ implying the threshold

$$S_c(T) = \frac{1}{a_w^i(T)} \frac{j_0 - a_0}{a_1} + 1. \quad (28)$$

250 Substituting eq. (28) into the expression (27) yields

$$\begin{aligned} \log_{10} J(S_i, T) &= j(S_i, T) = j_0 + a_1 a_w^i(T) (S_i - S_c(T)) \\ &= j_0 + A(T) (S_i - S_c(T)) \end{aligned} \quad (29)$$

where $A(T) = a_1 a_w^i(T)$. The coefficients a_0, a_1 can be determined in different ways, see section 4.3. Furthermore, approximations to the functions $A(T)$ and $S_c(T)$ can be investigated.

2. Case of a quadratic polynomial $p_2(x) = b_0 + b_1x + b_2x^2 = a(x - b)^2 + c$

255 Since a quadratic function is not strictly monotonic in general, inverting the quadratic polynomial leads to two functions, i.e.

$$p_2^{-1}(y) = b \pm \sqrt{\frac{y - c}{a}}. \quad (30)$$

If one solution can be ruled out (e.g. due to physical constraints) we can formulate

$$\log_{10} J = p_2((S_i - S_c(T)) a_w^i(T) + p_2^{-1}(j_0)) \quad (31)$$

260 using the threshold description

$$S_c(T) = \left(b \pm \sqrt{\frac{j_0 - c}{a}} \right) \frac{1}{a_w^i(T)} + 1 \quad (32)$$

Equivalently, we can derive a formulation

$$\begin{aligned} \log_{10} J &= c_0 + q_1(T) (S_i - S_c(T)) \\ &\quad + q_2(T) (S_i - S_c(T))^2 \end{aligned} \quad (33)$$

with appropriate functions q_1, q_2 , which might be useful for analytic investigations.

265 **Remark:** We will again use this quadratic ansatz for a direct approximation of the nucleation rate of pure water droplets (see section 6.1).

4.3 Linear polynomial fit for the nucleation rate

In this section we investigate approximations of the exponent of the nucleation rate of aqueous solution droplets J_{sol} and their impact on nucleation events in an idealised scenario. We concentrate on the reference formulation (Koop et al., 2000). Since
270 the polynomial $p_3(x)$ in the original formulation

$$J_{\text{sol}}(\Delta a_w) = 10^{p_3(\Delta a_w)}, \quad p_3(x) = \sum_{k=0}^3 a_k x^k \quad (34)$$

nearly behaves as a linear polynomial in the relevant range $0.26 \leq \Delta a_w \leq 0.34$, it can be easily approximated by a linear relation, i.e. $p_3(x) \approx b_0 + b_1 x$. For this we can use two different approaches: (i) using a least square fit to p_3 and (ii) a Taylor expansion at a prescribed value y_0 . While the first approach is just a fitting procedure in the relevant range $0.26 \leq \Delta a_w \leq 0.34$,
 275 the second approach relies on an a priori choice for the evaluation point $y_0 \in [0.26, 0.34]$ and it is not evident from the outset which value should be used to provide an accurate approximation. For this, we investigate the sensitivity of p_3 to a small perturbation $\varepsilon = y - y_0$, i.e. we consider

$$p_3(y) = p_3(y_0 + \varepsilon) = p_3(y_0) + \left. \frac{dp_3}{dx} \right|_{y_0} \varepsilon + \mathcal{O}(\varepsilon^2) \quad (35)$$

$$\approx b_{t0} + b_{t1} \cdot y = p_{t,y_0}(y) \quad (36)$$

280 with the coefficients

$$b_{t0} = p_3(y_0) - \left. \frac{dp_3}{dx} \right|_{y_0} \cdot y_0 \quad \text{and} \quad b_{t1} = \left. \frac{dp_3}{dx} \right|_{y_0}. \quad (37)$$

The Taylor approximation provides a range for the slope of the linear approximation; these values motivate the sensitivity analysis in section 4.5.2. In the relevant range $0.26 \leq y \leq 0.34$ for $y = \Delta a_w$ we obtain slopes in the range $221 \leq b_{t1} \leq 453$. This investigation gives us a hint about possible variations in the slope which will be used later for the sensitivity analysis in
 285 section 4.5.2.

In contrast, using a least square fitting routine for $0.26 \leq \Delta a_w \leq 0.34$ we obtain a linear function

$$p_{ls}(x) = b_{ls,0} + b_{ls,1} \cdot x \quad (38)$$

with $b_{ls,0} = -62.19267$ and $b_{ls,1} = 254.7749$. For the further investigations, we only use the linear fit from eq. (38). We observe that the linear fit $p_{ls}(x)$ best approximates p_3 close to the inflection point $x_{\text{infl}} \approx 0.30756$ (see figure 2, left panel).

290 For each linear approximation $p(x) = b_0 + b_1 \cdot x$ of $p_3(x)$, the exponent of the nucleation rate and the saturation ratio threshold become, as demonstrated in section 4.2,

$$j(S_i, T) = j_0 + \underbrace{b_1 a_w^i(T)}_{:=A(T)} (S_i - S_c(T)), \quad (39)$$

$$S_c(T) = \frac{1}{a_w^i(T)} \frac{j_0 - b_0}{b_1} + 1.$$

Since a_w^i is a rather complicated function of temperature, it is particularly useful in the context of analytical investigations to have simpler approximations of this quantity. This motivates to approximate a_w^i and its inverse $\frac{1}{a_w^i}$ in the relevant temperature
 295 range $190 \leq T \leq 230$ K by polynomials $q(T)$ of degree $\deg q \leq 2$. Similarly, we can approximate the nucleation threshold $S_c(T)$ by polynomials $s(T)$ of degree $\deg s \leq 2$. For the approximations we use a least square procedure within the temperature range $190 \leq T \leq 230$ K. The results are presented in figure 2 (middle and right panels). Note that the thresholds, either exact or approximate, are quite similar to the former approximations by Ren and Mackenzie (2005), while there is a larger difference to the approximation by Kärcher and Lohmann (2002).




Figure 2. Polynomial approximations of the nucleation rate (left), the ice water activity $a_w^i(T) = \frac{p_{ice}(T)}{p_{liq}(T)}$ (middle), and the saturation ratio threshold $S_c(T)$ (right panel), respectively. The right panel also includes the approximations by Kärcher and Lohmann (2002) and Ren and Mackenzie (2005).

300 Combining the approximations $q(T)$ and $s(T)$ yields the formulation

$$\begin{aligned}
 j(S_i, T) &= j_0 + b_1 q(T) (S_i - s(T)) \\
 &\approx j_0 + A(T) (S_i - S_c(T))
 \end{aligned}
 \tag{40}$$

of $\log_{10}(J)$. As can be seen in Figure 2, the nucleation threshold is accurately approximated by a linear relation (deviation is smaller than 0.3%). In former studies (e.g. Kärcher and Lohmann, 2002; Ren and Mackenzie, 2005) linear fits were derived for the nucleation thresholds; however, these fits deviate significantly more from the reference in comparison to ours (see figure 2).

305 The deviation depends on the respective formulation (or approximation) of a_w^i .

Also former studies use (empirical) linear approximation for the saturation threshold. Comparing these linear approximations with ours in Fig 2 it is evident that they deviate significantly.

4.4 Thresholds for prescribed nucleation rate values

The threshold description in section 4.3 was based on the choice $j_0 = 16$, corresponding to a nucleation rate $J = 10^{16} \text{ m}^{-3} \text{ s}^{-1}$.

310 As already mentioned, the choice of j_0 is quite arbitrary, and these high values of J are very often not reached in the numerical simulations (see section 4.5). For a better diagnostics of the nucleation events and the relative strength of nucleation events, we introduce a similar concept for nucleation thresholds, based on a prescribed nucleation rate value $J \sim 10^{x_0}$. For this purpose we use eq. (40) of the nucleation threshold based on the linear approximation of the nucleation rate with a fixed but arbitrary value $x_0 > 0$ for the nucleation rate value; hence, we can write

$$\begin{aligned}
 x_0 &= j(S_0, T) = j_0 + A(T) (S_0 - S_c(T)) \\
 \Leftrightarrow S_{cx0}(T) &= S_0 = \frac{x_0 - j_0}{A(T)} + S_c(T)
 \end{aligned}
 \tag{41}$$

where the function $A(T) = b_1 a_w^i(T)$ depends only on the linear approximation of J as stated in section 4.2. Note that obviously $S_{cx0}(T) = S_c(T)$ for $x_0 = j_0$. This leads to the formulation of the nucleation rate

$$j(S_i, T) = x_0 + A(T)(S_0 - S_{cx0}(T)). \quad (42)$$

with a general nucleation value x_0 and its associated threshold function $S_{cx0}(T)$. The threshold function is just shifted by
 320 the value $\frac{x_0 - j_0}{A(T)}$, i.e. the type of the threshold function remains the same. This formulation will be used for the theoretical investigations using small perturbations (see section 4.6)

4.5 Numerical simulations of nucleation events for different approximations

In the following we investigate the impact of our approximations of $\log_{10}(J)$ on nucleation events. The setup is as follows: We use the simple bulk ice physics model as described by the set of ODEs (15), (16), (17) in section 3. A nucleation event is ensured
 325 by assuming a constant vertical velocity, which directly translates into a constant adiabatic cooling of the air parcel and, thus, an initially increasing saturation ratio. Instead of changing the temperature adiabatically, we directly control the supersaturation as described in sec. 3; this allows us to control the nucleation event without the need to disentangle the different contributions of temperature and supersaturation.

The nucleation events always admit the same structure: Due to the supersaturation source $\sim wS_i$ with constant updraft w
 330 the variable S_i increases and the nucleation term produces ice crystals, which can grow by water vapour diffusion, constituting a sink for supersaturation. The peak value of S_i is reached once the source and sink of supersaturation balance. Afterwards the variable S_i decreases due to diffusional growth and thus shut off the nucleation term. The peak value depends crucially on the number of nucleated ice crystals that are needed, to balance the source for S_i by the diffusional growth (depending on the product of number concentration and mean radius of ice crystals). The number concentration of ice crystals produced in the
 335 nucleation event clearly depends on the vertical velocity w (source term) and the environmental conditions (diffusion depends on temperature and pressure). For details of the time evolution of nucleation events see appendix B.

4.5.1 Standard approximation

We compare the following four different representations of the nucleation rate using numerical simulations:

1. nucleation rate in the water activity formulation by Koop et al. (2000) with the correction as described in section 4.1
 340 (reference nucleation rate)
2. water activity approximated by the linear fit as described in section 4.3 (see eq. (38), linear regression)
3. nucleation rate as a function of S_i, T as described in section 4.2 based on the formulation

$$\log_{10} J = j_0 + A(T)(S_i - S_c(T)) \quad (43)$$

of the exponent of the nucleation rate. We compare the following two sets of approximations for $A(T)$ and $S_c(T)$:




Figure 3. Different approximations of nucleation rate for different temperatures (left: $T = 196$ K, middle: $T = 216$ K, right: $T = 236$ K). Black: Reference nucleation rate; red: linear fit to reference nucleation rate; blue: threshold description due to eq 43, using a linear approximation for a_w^i and a quadratic threshold function S_c ; green: threshold description due to eq. (43), using a constant for a_w^i and a linear threshold function S_c .

- 345 (a) a linear approximation for $A(T)$ and a quadratic approximation for $S_c(T)$,
 (b) a constant approximation for $A(T)$ and a linear approximation for $S_c(T)$.

These are specific cases, however arbitrary combinations of approximations for $A(T)$ and $S_c(T)$ might be used.

Figure 3 shows the approximated exponents of the nucleation rate together with the (corrected) reference formulation by Koop et al. (2000) for the three temperatures $T = 196, 216, 236$ K as functions of Δa_w . These temperatures are chosen for
 350 direct comparison with former studies (Kärcher and Lohmann, 2002; Spichtinger and Gierens, 2009). Evidently, the linear fit with respect to water activity is very close to the reference, and the same is true for the case of a linear function $A(T)$ and a quadratic approximation $S_c(T)$. For the simplest approximation (constant function $A(T)$ and linear approximation $S_c(T)$), larger deviations from the reference nucleation rate can be seen. At $T = 196$ K, there is a strong underestimation in the lower range of Δa_w , whereas for $T = 236$ K the underestimation is most pronounced for higher values of Δa_w (green vs. black
 355 curves). In both cases, we expect deviations in the number concentrations of nucleated ice crystals during the nucleation event and the maximum saturation ratio attained.

We investigate standard nucleation events in terms of (i) the resulting ice crystal number concentration at the end of the simulation as in former studies (e.g. Kärcher and Lohmann, 2002; Spichtinger and Gierens, 2009) and (ii) the maximum (peak) supersaturation, which was reached during the nucleation event. Although the latter is usually not considered, it is of interest
 360 for comparisons with real measurements, e.g. in cloud chambers.

Figure 4 shows the results of the numerical simulations, i.e. the number of nucleated ice crystals (left panel) and the maximum saturation ratio (right panel) at environmental pressure $p = 200$ hPa (the results are similar for other environmental conditions).




Figure 4. Comparison of different approximations of the nucleation rate by Koop et al. (2000) for standard nucleation events driven by a constant vertical velocity w . Left: ice crystal number concentration; right: maximum supersaturation.

Comparing the number of nucleated ice crystals as well as the maximum saturation ratio it is evident, that the difference between the reference calculation, based on the corrected nucleation rate by Koop et al. (2000), and the simulations using the approximated nucleation rates are rather small.

For almost all nucleation events, the deviation from the reference simulations is not larger than $\pm 15\%$. To assess these deviations one should keep in mind that measurements of ice crystal number concentrations are quite difficult and the uncertainties are usually larger than 15%. For instance, for the FSSP instrument, which was used in many flight campaigns (e.g. Voigt et al., 2017), the uncertainty is estimated by about $\sim 10\%$ (de Reus et al., 2009). Thus, the deviations in our simulations and the uncertainties of realistic measurements are roughly of the same order. This fact renders it presumably impossible to decide on the correctness of any of the different formulations and approximations of the nucleation rate based on the available observations.

Finally, we conclude that a linear approximation of the reference nucleation rate by Koop et al. (2000) is accurate enough to represent nucleation events in a physically meaningful way. Thus, we can use this description as well as the derived formulations of the nucleation rate as a function of temperature T and saturation ratio S_i in order to investigate which parameters of the nucleation rates significantly affect the outcome of nucleation events. This will be carried out in the next section.

4.5.2 Impact of the parameters of the linear approximation

Generally, we are interested in the impact of the formulation of the nucleation rate on nucleation events. The original parameterisation by Koop et al. (2000) is based on a cubic polynomial, which admits slopes in the range $221 \leq b \leq 453$, see Sect. 4.3. The linear approximation is sufficiently good for representing the “reference” rate; thus, we now use this simple linear repre-

sentation $\log_{10} J = b_0 + b_1 \cdot \Delta a_w$ in order to test the sensitivity of nucleation events on the two parameters b_0, b_1 . Parameter b_0 controls the absolute value of the nucleation rate while parameter b_1 accounts for its steepness, i.e. the slope.

In a first step, we investigate the impact of the slope of the nucleation rate given by coefficient b_1 . One should keep in mind that during the nucleation event the value of $\Delta a_w = (S_i - 1)a_w^i(T)$ is increasing as S_i increases, thus the exponent of the nucleation rate basically grows linearly. Consequently, an increase in the saturation ratio immediately translates into an increase in Δa_w , hence the abscissa in figure 5 may be thought of as representing saturation ratio. If high values of the nucleation rate are already reached at lower supersaturation values, the nucleation is triggered earlier in comparison to the reference scenario.

However, an earlier onset of ice nucleation implies that the newly nucleated ice crystals already start to grow by diffusion. Consequently, the growing ice crystals tend to decrease the saturation ratio and, if they are sufficiently numerous, prematurely stop the nucleation event. In this case, less ice crystals will nucleate and a smaller maximum saturation ratio will be reached compared to the reference. The opposite mechanism is expected for smaller values of the nucleation rate in comparison to the reference, i.e. higher ice crystals concentrations will occur together with larger maximal saturation ratio.




Figure 5. Artificial change in the slope of the linear function in the exponent of the nucleation rate. The fit to the reference curve is indicated by the green line (slope $b_1 \sim 250$); a reduced slope ($b_1 \sim 100$) is displayed in red, an enhanced slope ($b_1 \sim 500$) is displayed in blue.

In order to illustrate this mechanism more quantitatively, we artificially changed the slope of the linear function. The “reference” slope $b_1 \approx 255$ is either reduced to a value of $b_1 = 100$ or enhanced to a value $b_1 = 500$, which is motivated by the values of the Taylor approximation, derived in section 4.3. In both cases, the parameter b_0 of the linear function is adapted such that the inflection point of the polynomial $p_3(\Delta a_w)$ at $\Delta a_w \sim 0.311$ is met for better comparison with the reference simulations. The resulting nucleation rates are displayed in figure 5, while the number of nucleated ice crystals and the maximum ice saturation ratio during the nucleation event are summarized in figure 6: The left panel shows the concentrations of nucleated ice crystals and the right panel the maximum saturation ratio during the nucleation events.




Figure 6. Impact of the slope on the idealized nucleation events. Left: ice crystal number concentrations, right: maximum supersaturation values. The colours are chosen as in figure 5, i.e. red squares indicate reduced slope, and blue triangles indicate enhanced slope, respectively.

In case of the enhanced or reduced slope as indicated in figure 5 we exactly see the theoretically proposed behaviour in the ice crystal number concentration: the values are reduced for reduced slopes, and enhanced for enhanced slopes. The reductions are by up to a factor of 0.4, the enhancements are by up to a factor of 2.4, and the largest changes can be seen at the highest temperature $T = 236$ K.

In the right panel of Figure 6, a dependency on temperature and vertical velocity is seen. For very low vertical velocities, the maximum supersaturation behaves as expected, i.e. reduced values for the reduced slope and enhanced values for the enhanced slope. For very high vertical velocities, this behaviour is reversed, i.e. we see reduced values of $S_{i,\max}$ for enhanced slopes and enhanced values of $S_{i,\max}$ for reduced slopes. The transition slightly depends on the temperature. This can be explained as follows: For low vertical velocities, Δa_w (and thus the supersaturation) is always below the inflection point $\Delta a_w \sim 0.311$. Thus the nucleation rate is always smaller for the enhanced slope in comparison to the reference while it is always larger in comparison to the reference for the reduced slope. Therefore, in case of an enhanced slope the nucleation starts later compared to the reference. This leads to the behaviour as described above. However, beyond the inflection point the behaviour is reversed and thus the resulting maximum supersaturation is now enhanced for reduced slope and it is reduced for enhanced slope. The

inflection point is reached at different vertical velocities for different temperatures, i.e. for lower temperatures at lower values of w and for higher temperatures at higher values of w . Note, only the maximum supersaturation is affected upon Δa_w crossing the inflection point while no influence on the number concentration of ice crystals is seen.

After having varied the slope of the nucleation rate, we now turn to its absolute values and modify coefficient b_0 , which translates into a change of values of J by 10^{b_0} . In order to investigate the sensitivity, we add a constant value $\Delta b \in \{-6, -3, 3, 6\}$ to the coefficient b_0 , resulting in an increase or decrease in the absolute value of the nucleation rate by a factor of $10^{\Delta b}$. In figure 7 the results in terms of ice crystal number concentration and maximum supersaturation are displayed.




Figure 7. Comparison of ice crystal number concentrations (left panel) and maximum supersaturation (right panel) for absolute changes in the nucleation rate by a factor $10^{\Delta b}$ with $\Delta b \in \{-6, -3, 3, 6\}$.

Maybe surprisingly, the absolute values of the number concentrations of ice crystals in comparison to the reference formulation are not crucially affected (see figure 7, left panel), although some deviations occur (up to a factor of two). The strongest deviations can be seen for warm temperatures ($T = 236$ K) at very low vertical velocities. Overall, the relative deviations from the reference events in variables n_i and peak values of S_i are within the interval $[0.4, 2]$, but for vertical velocities in the range $w \geq 0.05$ m s⁻¹ the relative deviation is within the interval $[0.8, 1.4]$.

Comparing the influence of a scaling of the absolute values of the nucleation rate and the steepness of the rate, we conclude that the correct steepness of the nucleation rate is much more important than the absolute value of J . Even changes by orders of magnitude in the values of the nucleation rate has a minor impact on the number of nucleated ice crystals. A similar conclusion was also drawn in the theoretical study by Baumgartner and Spichtinger (2019). In that study, the authors investigated a slightly simplified system of equations by means of asymptotic analysis. The simplified system describes the temporal evolution of the number concentration of ice crystals and the saturation ratio and an approximate asymptotic solution was constructed. To

leading order, the approximate solution for the number concentration of ice crystals was completely independent of the precise values of the nucleation rate, but the steepness contributed directly. The only necessary condition on the values of the nucleation rate was that it attains large values, i.e. significantly larger than the other coefficients within the equations.

For the maximum supersaturation values, the impact of the absolute value of J is much more pronounced. As expected, upon reduction of the nucleation rate by a factor of $10^{\Delta b}$ with $\Delta b \in \{-6, -3\}$ the supersaturation reaches much higher values of S_i , until the values of the rescaled nucleation rate become large enough to initiate the nucleation of ice crystals. For the enhancement of the absolute values of the nucleation rate, the results are reversed: The maximum supersaturation is reduced, since the enhanced nucleation rate attains values that allow the production of ice crystals for smaller saturation ratios. This behaviour is represented in the right panel of figure 7.

Remark: This idealized enhancement of the nucleation rate can also be seen in the connection with the aerosol number concentration n_a . A change of n_a by some orders of magnitudes while no changes in J are applied has the same effect as changing the absolute value of the nucleation rate (or the parameter b_0 in the argument of the exponential function). Thus, a strong reduction or enhancement of the available solution droplets will only slightly change the amount of ice crystals in a nucleation event. Therefore, we can conclude that for a meaningful approximation of the nucleation rate the exact number concentration of available aerosols is also not crucial for the strength of the homogeneous nucleation event, but perhaps for the starting time of the event. Including size effects of the solution droplets might additionally change the picture quantitatively (see, e.g., Baumgartner et al., 2020).

4.6 Impact of perturbations in S_i and T on the nucleation rate

In this section we investigate the impact of changes in S_i and/or T on the nucleation rate by employing a perturbation analysis. A short explanation of this technique is given in Appendix D. In the real atmosphere, variations of the temperature due to dynamical processes will introduce such changes, e.g. such as from a passing or even breaking gravity wave. In numerical simulations, these variations (also often called fluctuations) are often artificially introduced (e.g. Jensen and Pfister, 2004). In any case, the impact of such changes is investigated using perturbation analysis (also called asymptotic analysis).

We start with the linear approximation of the nucleation rate as formulated in eq. (40) with $A(T) = b_1 a_w^i(T)$. We can estimate the usual values of the function $A(T)$ in the temperature range $190\text{ K} \leq T \leq 230\text{ K}$ using $0.51 \leq a_w^i(T) \leq 0.66$ such that $129 \leq A(T) \leq 169$. For a very simple but still sufficiently accurate constant approximation of $a_w^i(T)$ we can set $a_w^i = 0.574312$ (see fig. 2, pink line) such that $A(T) \approx A_0 = b_1 a_w^i = 146.32$. Finally we can state $A(T) = \mathcal{O}(\varepsilon^{-2})$ with the usual perturbation approach $\varepsilon \sim 0.1$, such that we set $A(T) = A^* \varepsilon^{-2}$ with $A^* = \mathcal{O}(1)$ as $\varepsilon \rightarrow 0$. For the non-dimensionalization of the threshold function in the linear approximation $S_c(T) \approx s_0 + s_1 T$ we have to estimate the order of the coefficients for the relevant temperature range. Using $190\text{ K} \leq T_{\text{ref}} \leq 230\text{ K}$ and the definition $T = T_{\text{ref}} \vartheta$ with the nondimensional temperature ϑ , we find

$$S_c(T) = s_c(\vartheta) = s_0 + s_1 T = s_0 + s_1 T_{\text{ref}} \vartheta = \sigma_0 - \sigma_1 \vartheta \quad (44)$$

with $\sigma_1 = -s_1 T_{\text{ref}}$. Obviously, $s_0 = \sigma_0 = 2.27697 = \mathcal{O}(1)$ and $0.66 \leq \sigma_1 \leq 0.8$ such that $\sigma_1 = \mathcal{O}(1)$. Using the simplest approximation $A(T) = A_0$ and $S_c(T) = s_0 + s_1 T$ for the general formulation of the threshold function S_{cx0} (cf. eq. (41)) we can
 465 simplify the expression as

$$\begin{aligned} S_{cx0}(T) &= \frac{x_0 - j_0}{A_0} + s_0 + s_1 T = \underbrace{\left(\frac{x_0 - j_0}{A_0} + s_0 \right)}_{=: s_{x0}} + s_1 T \\ &= s_{x0} + s_1 T. \end{aligned} \quad (45)$$

Using non-dimensionalization we end up with the following representation

$$S_{cx0}(T) = s_{cx0}(\vartheta) = s_{x0} + s_1 T = \sigma_{x0} - \sigma_1 \vartheta, \quad (46)$$

where $\sigma_{x0} = s_{x0}$, $\sigma_1 = -s_1 T_{\text{ref}}$. Finally, we use the estimation $A_0 = A^* \varepsilon^{-2}$ to obtain

$$470 \quad \sigma_{x0} = (A^*)^{-1} \varepsilon^2 (x_0 - j_0) + \sigma_0 = \delta + \sigma_0. \quad (47)$$

Since $j_0 = \mathcal{O}(\varepsilon^{-1})$ and $x_0 = \mathcal{O}(\varepsilon^\beta)$ with $\beta \geq -1$ we find $\sigma_{x0} = \delta + \sigma_0 = \mathcal{O}(\varepsilon) + \mathcal{O}(1) = \mathcal{O}(1)$. After non-dimensionalizing the argument in the nucleation rate, we can now investigate the response of the nucleation rate upon a perturbation (i) in saturation ratio (i.e. in the same way as the numerical simulations are set up), (ii) in temperature, and (iii) in adiabatic changes of temperature driving changes in the saturation ratio simultaneously. In reality, almost exclusively case (iii) is relevant.

475 First, we estimate the increase of J due to variations of S_i at a constant temperature $T = T_{\text{ref}}$. For this purpose we start at a given value of the saturation ratio S_i which corresponds to a certain threshold x_0 via the relation (41). We choose this value as a reference value $S_{\text{ref}} = S_{cx0}(T_{\text{ref}}) = s_{cx0}(1) = \sigma_0 - \sigma_1$; this corresponds to a reference value of the nucleation rate $J = J_{\text{ref}} = J_{\text{unit}} \cdot 10^{x_0}$ (with $J_{\text{unit}} = 1 \text{ m}^{-3} \text{ s}^{-1}$). Assuming the expansion

$$S_i = S_0 + \varepsilon S_1 + \varepsilon^2 S_2 + \varepsilon^3 S_3 + \mathcal{O}(\varepsilon^4) \quad (48)$$

480 for the saturation ratio where $S_0 = S_{\text{ref}} \sigma_{x0} - \sigma_1$ we investigate the impact of such a perturbation on the exponent j . Keeping the temperature fixed as in the numerical simulations we arrive at

$$\begin{aligned} j(S_i, T) &= j(s, t) = x_0 + A_0 (S_i - S_{\text{ref}}) \\ &= x_0 + A^* \varepsilon^{-2} (S_{\text{ref}} + \varepsilon S_1 + \varepsilon^2 S_2 + \varepsilon^3 S_3 + \mathcal{O}(\varepsilon^4) - S_{\text{ref}}) \\ &= x_0 + \varepsilon^{-1} A^* S_1 + A^* S_2 + \varepsilon A^* S_3 + \mathcal{O}(\varepsilon^2). \end{aligned} \quad (49)$$

We are interested in the relative change of the nucleation rates $\frac{J(S_i, T)}{J_{\text{ref}}}$, which translates into $j(S_i, 1) - j(S_{\text{ref}}, 1)$. By definition, we have $x_0 = j(S_{\text{ref}}, 1)$, thus we obtain

$$485 \quad j(S_i, 1) - j(S_{\text{ref}}, 1) = \varepsilon^{-1} A^* S_1 + A^* S_2 + \varepsilon A^* S_3 + \mathcal{O}(\varepsilon^2). \quad (50)$$

Inspecting eq. (50) it is evident, that a nonzero perturbation term S_α in eq. (48) is connected with the factor $\varepsilon^{\alpha-2}$, hence a change of order $\mathcal{O}(\varepsilon^\alpha)$ in supersaturation translates into a change of order $\mathcal{O}(\varepsilon^{\alpha-2})$ in the exponent of J . For instance, a change by $S \sim 0.01$ translates into a change of $\mathcal{O}(1)$ in j , thus in a change by a factor of 10 in the nucleation rate J

Second, we consider perturbations of temperature without changing the saturation ratio, although this might not happen
 490 in the atmosphere. Using the approach above with a constant reference value of saturation, i.e. $S_{\text{ref}} = s_{cx0}(1) = \sigma_0 - \sigma_1$ and
 temperature perturbations $\vartheta = 1 + \varepsilon\vartheta_1 + \varepsilon^2\vartheta_2 + \varepsilon^3\vartheta_3 + \mathcal{O}(\varepsilon^4)$ we find the following expression:

$$\begin{aligned}
 j(S_{\text{ref}}, \vartheta) &= x_0 + A_0 (S_{\text{ref}} \\
 &\quad - (\sigma_0 - \sigma_1 (1 + \varepsilon\vartheta_1 + \varepsilon^2\vartheta_2 + \varepsilon^3\vartheta_3 + \mathcal{O}(\varepsilon^4)))) \\
 &= x_0 + A^* \varepsilon^{-2} (\varepsilon\vartheta_1 + \varepsilon^2\sigma_1\vartheta_2 + \varepsilon^3\sigma_1\vartheta_3 + \mathcal{O}(\varepsilon^4)) \\
 &= x_0 + \varepsilon^{-1} A^* \sigma_1 \vartheta_1 + A^* \sigma_1 \vartheta_2 + \varepsilon A^* \sigma_1 \vartheta_3 + \mathcal{O}(\varepsilon^2).
 \end{aligned} \tag{51}$$

The relative change of the nucleation rate is then given by

$$\begin{aligned}
 j(S_{\text{ref}}, \vartheta) - j(S_{\text{ref}}, 1) &= x_0 + \varepsilon^{-1} A^* \sigma_1 \vartheta_1 \\
 &\quad + A^* \sigma_1 \vartheta_2 + A^* \sigma_1 \vartheta_3 \varepsilon + \mathcal{O}(\varepsilon^2).
 \end{aligned} \tag{52}$$

495 Thus, a temperature perturbation ϑ_α of order $\mathcal{O}(\varepsilon^\alpha)$ leads to a relative change in j of order $\mathcal{O}(\varepsilon^{\alpha-2})$. Note the sign of the
 perturbations, which turns into the opposite sign in the change of j . Because of the strictly monotonic decrease of the threshold
 function $S_{cx0}(T)$, a negative temperature change leads to a higher threshold and in turn to a lower nucleation rate at a given
 saturation ratio.

Instead of perturbing the saturation ratio and the temperature individually, these quantities are connected in the real world. To
 500 a good approximation, their joint variation is through an purely adiabatic change. Therefore, we finally investigate the impact
 of adiabatic temperature changes on the saturation ratio and in turn on the nucleation rate. For this purpose we have to consider
 the dependence of S_i on adiabatic temperature changes. We start with the cooling source term of the saturation ratio

$$dS_i = \left(\frac{1}{\kappa} - \frac{L}{R_v T} \right) S_i \frac{dT}{T}. \tag{53}$$

The term $\gamma(T) = \frac{1}{\kappa} - \frac{L}{R_v T}$ within the bracket admits the values $-28.8 \leq \gamma(T) \leq -23.2$ for $190 \text{ K} \leq T \leq 230 \text{ K}$ such that
 505 we find $\gamma(T) = \mathcal{O}(\varepsilon^{-1}) = \gamma^* \varepsilon^{-1}$ and $\gamma^* \sim -2.5 < 0$. Approximating the total differential in eq. (53) with finite differences
 $\Delta S_i, \Delta T$, we arrive at

$$\frac{\Delta S_i}{S_i} = \gamma^* \varepsilon^{-1} \frac{\Delta T}{T}. \tag{54}$$

We set as an approximation $S_i = S_{\text{ref}}$ and $T = T_{\text{ref}}$ such that we can set

$$\frac{\Delta S_i}{S_{\text{ref}}} = \varepsilon^l S_l + \mathcal{O}(\varepsilon^{l+1}) = \mathcal{O}(\varepsilon^l) \tag{55}$$

510 with $S_l = \mathcal{O}(1)$. We assume $l \geq 1$ since we do not consider changes of the saturation ratio of order $\mathcal{O}(1)$. The analogous
 expansion for the temperature reads

$$\frac{\Delta T}{T_{\text{ref}}} = \varepsilon^k \vartheta_k + \mathcal{O}(\varepsilon^{k+1}) = \mathcal{O}(\varepsilon^k) \quad \text{with } \vartheta_k = \mathcal{O}(1) \quad \forall k \geq 1. \tag{56}$$

Combining these expansions, Eq. (53) becomes

$$\begin{aligned}\frac{\Delta S_i}{S_{\text{ref}}} &= \gamma^* \varepsilon^{-1} \frac{\Delta T}{T_{\text{ref}}} = \gamma^* \varepsilon^{-1} (\varepsilon^k \vartheta_k + \mathcal{O}(\varepsilon^{k+1})) \\ &= \gamma^* t_k \varepsilon^{k-1} + \mathcal{O}(\varepsilon^k)\end{aligned}\quad (57)$$

515 or equivalently

$$\varepsilon^l S_l = \Delta S_i = S_{\text{ref}} \gamma^* \vartheta_k \varepsilon^{k-1} + \mathcal{O}(\varepsilon^k). \quad (58)$$

The only non-trivial balance is achieved for $l = k - 1$, i.e.

$$S_{k-1} = S_{\text{ref}} \gamma^* \vartheta_k \Leftrightarrow S_k = S_{\text{ref}} \gamma^* \vartheta_{k+1}. \quad (59)$$

Note that $l = k - 1 \geq 1$, i.e. we have to consider $k \geq 2$ for the perturbation of temperature. This is a meaningful restriction
520 since we are interested in small changes of temperature in the cold temperature regime, i.e. a change in the temperature in the order of $\sim 1\text{K}$ in physical units. Hence, we would not expect adiabatic temperature changes of order $\mathcal{O}(\varepsilon)$, corresponding to changes of order $\sim 10\text{K}$. Thus, we assume an asymptotic expansion

$$\vartheta = 1 + \varepsilon^2 \vartheta_2 + \varepsilon^3 \vartheta_3 + \mathcal{O}(\varepsilon^4) \quad (60)$$

for the temperature perturbation. We are generally interested in adiabatic expansions due to vertical upward motion, which
525 in turn leads to decreasing temperatures, hence we conclude $\vartheta_k < 0$ for $k \geq 2$. Since $\gamma^* < 0$, equation (59) leads to positive changes in the saturation ratio $s_k > 0$ for $\vartheta_k < 0$. Generally, warming due to adiabatic compression can be studied in the same way by setting $\vartheta_k > 0$.

Now we consider the nucleation rate in the formulation for arbitrary thresholds x_0 in the nucleation rate using $S_{\text{ref}} = \sigma_{x_0} - \sigma_1 = s_{cx_0}(1)$:

$$\begin{aligned}j(S_i, T) &= j(S_i, t) = x_0 + A^* \varepsilon^{-2} (S_i - s_{cx_0}(\vartheta)) \\ &= x_0 + A^* \varepsilon^{-2} (S_{\text{ref}} + \varepsilon S_1 + \varepsilon^2 S_2 + \varepsilon^3 S_3 \\ &\quad - (\sigma_{x_0} - \sigma_1 (1 + \varepsilon^2 \vartheta_2 + \varepsilon^3 \vartheta_3)) + \mathcal{O}(\varepsilon^4)) \\ &= x_0 + A^* \varepsilon^{-2} (\varepsilon S_1 + \varepsilon^2 S_2 + \varepsilon^3 S_3 + \varepsilon^2 \sigma_1 \vartheta_2 \\ &\quad + \varepsilon^3 \sigma_1 \vartheta_3 + \mathcal{O}(\varepsilon^4)) \\ 530 &= x_0 + A^* \varepsilon^{-2} (\varepsilon S_{\text{ref}} \gamma^* \vartheta_2 + \varepsilon^2 S_{\text{ref}} \gamma^* \vartheta_3 + \varepsilon^3 S_{\text{ref}} \gamma^* \vartheta_4 \\ &\quad + \varepsilon^2 \sigma_1 \vartheta_2 + \varepsilon^3 \sigma_1 \vartheta_3 + \mathcal{O}(\varepsilon^4)) \\ &= x_0 + A^* \varepsilon^{-2} (\varepsilon S_{\text{ref}} \gamma^* \vartheta_2 + \varepsilon^2 (S_{\text{ref}} \gamma^* \vartheta_3 + \sigma_1 \vartheta_2) \\ &\quad + \varepsilon^3 (S_{\text{ref}} \gamma^* \vartheta_4 + \sigma_1 \vartheta_3) + \mathcal{O}(\varepsilon^4)) \\ &= x_0 + A^* S_{\text{ref}} \gamma^* \vartheta_2 \varepsilon^{-1} + A^* (S_{\text{ref}} \gamma^* \vartheta_3 + \sigma_1 \vartheta_2) \\ &\quad + A^* (s_{\text{ref}} \gamma^* \vartheta_4 + \sigma_1 \vartheta_3) \varepsilon + \mathcal{O}(\varepsilon^2).\end{aligned}\quad (61)$$

Thus, for $k \geq 2$ we find terms of the form $A^* (S_{\text{ref}}\gamma^*\vartheta_{k+1} + \sigma_1\vartheta_k)\varepsilon^{k-2}$ of order $\mathcal{O}(\varepsilon^{k-2})$. Comparing the nucleation rates we find for the relative change

$$\begin{aligned} j(S_i, T) - j(S_{\text{ref}}, T_{\text{ref}}) &= j(S_i, t) - j(S_{\text{ref}}, 1) \\ &= A^* S_{\text{ref}}\gamma^*\vartheta_2\varepsilon^{-1} + A^* (S_{\text{ref}}\gamma^*\vartheta_3 + \sigma_1\vartheta_2) \\ &\quad + A^* (S_{\text{ref}}\gamma^*\vartheta_4 + \sigma_1\vartheta_3)\varepsilon + \mathcal{O}(\varepsilon^2). \end{aligned} \tag{62}$$

For the relative impact of these terms we use the estimations $\gamma^* < -2.3$ and $\sigma_1 \leq 0.8$. We have to distinguish two scenarios
535 for perturbations $\vartheta_k < 0$:

1. $\vartheta_k < 0$ for all $k \geq 2$. In this case we can assume

$$S_{\text{ref}}\gamma^*\vartheta_{k+1} + \sigma_1\vartheta_k > 0. \tag{63}$$

Therefore, an adiabatic temperature perturbation ϑ_k of order $\mathcal{O}(\varepsilon^k)$ ($k \geq 2$) leads to relative changes in j of order
540 $\mathcal{O}(\varepsilon^{k-3})$. Note, that the changes in saturation ratio are always dominant and larger than the changes in the threshold, which changes j by order $\mathcal{O}(\varepsilon^{k-2})$ in the opposite direction.

2. $\vartheta_k < 0$ and $\vartheta_{k+1} = 0$ for a distinct $k \geq 2$. In this case, the previously discussed temperature effect can be seen, i.e. the nucleation threshold is changed, leading to a reduction of the nucleation rate exponent. This effect is merely academic, since we have to switch off higher perturbations in temperature, which is quite unlikely.

One should keep in mind that we investigated the relative increase in the exponent of the nucleation rate. A relative change of
545 order $\mathcal{O}(\varepsilon^k)$ in the exponent translates into a relative change of order $\mathcal{O}(\exp(\varepsilon^k))$ in the nucleation rate J , thus ranging over several orders of magnitudes. For instance, in the first scenario changes of temperature of order $\sim 1\text{K}$ lead to changes in j of about ~ 10 , which in turn translate into a change of the nucleation rate J by a factor of $\exp 10 \sim 10^5$.

Overall, we can state that changes in S_i are most important for changing j , either stemming from adiabatic temperature changes or driven directly as in our numerical studies.

550 **5 Impact of saturation vapour pressure formulation**

Since the formulation of the nucleation rate by Koop et al. (2000) relies on the water activity, and thus on the function $a_w^i(T) = \frac{p_{\text{ice}}(T)}{p_{\text{liq}}(T)}$, the saturation vapour pressure over liquid water (i.e. in the no man's land) plays an important role. In this section we investigate the impact of choosing another formulation for $p_{\text{liq}}(T)$ on the nucleation rate and thus the nucleation events.

5.1 New representation of saturation water vapour

555 In the formulation by Murphy and Koop (2005) the extrapolation of the saturation vapour pressure into the no man's land of water's phase diagram is based on the assumption that the state of amorphous ice is thermodynamically equivalent to super-cooled liquid water. Therefore, the specific heat of liquid water can be extended in the super-cooled regime using measurements of amorphous ice. This leads to the established formulation in Murphy and Koop (2005).

560 Recently, a new representation of the saturation vapour pressure over super-cooled liquid water was proposed by Nachbar et al. (2019). In this study, the authors consider different states of water in the low temperature range. They conclude that amorphous ice is thermodynamically different from super-cooled water, thus they provide a different extrapolation for the saturation vapour pressure (Nachbar et al., 2019).

Although the deviation between the two curves is very small – even in the low temperature range less than 10% – its impact on saturation ratios as well as on the nucleation thresholds is quite large, as can be seen in figure 8.




Figure 8. Water saturation ($S_i = \frac{p_{liq}}{p_{ice}}$) and nucleation threshold (for $J = 10^{16} \text{ m}^{-3} \text{ s}^{-1}$) for different formulations of saturation vapour pressure over super-cooled water, Murphy and Koop (2005) vs. Nachbar et al. (2019)

565 The curves of water saturation as well as the nucleation thresholds are systematically shifted to higher values. In addition, the new curves have a more linear shape than the curves resulting from Murphy and Koop (2005). The ratio of the saturation pressures over ice and liquid (i.e. the functions $a_w^i(T)$ and $\frac{1}{a_w^l(T)}$) behave differently: $a_w^i(T)$ is much closer to a quadratic curve as can be seen in the left panel of figure 9. These new fits were used for the formulation of the approximated nucleation rate. Thus, we do not change the general approach for approximating the nucleation rate etc., we only use a different representation
570 of the function $a_w^i(T)$.




Figure 9. Left: Function $a_w^i(T) = \frac{p_{ice}(T)}{p_{liq}(T)}$ (black line) and polynomial approximations (red: quadratic, blue: linear, green: constant). Right: Nucleation threshold $S_c(T)$ (black line) and polynomial approximations (red: quadratic, blue: linear). Note that the former approximation by Kärcher and Lohmann (2002) (dark green) are now very close to the new formulation, whereas the fit by Ren and Mackenzie (2005) (turquoise) deviates significantly.

5.2 Numerical simulations of nucleation events

In figure 10 the results of the nucleation events using the new representation of the saturation vapour pressure due to Nachbar et al. (2019) are shown. As for former experiments, the ice crystal number concentration (left panel) and the maximum supersaturation values (right panel) are shown.

575 For the ice crystal number concentration, the impact of the new formulation of p_{liq} is small; the relative deviation from the reference simulations using the original vapour pressure formulation by Murphy and Koop (2005) is always smaller than 15%. The deviation increases with decreasing temperature and is most prominent for lower vertical updrafts ($w < 1 \text{ m s}^{-1}$).

For the maximum saturation ratio the change as compared to the reference simulations is much more prominent. As can be seen in figure 8 the nucleation thresholds for a value of $J = 10^{16} \text{ m}^{-3} \text{ s}^{-1}$ are increasing with decreasing temperature with a larger slope compared to the reference case. This behaviour can clearly be seen in the maximum supersaturation; for decreasing temperature the maximum supersaturation is increasing to higher values in comparison to the reference simulations. The increase does not depend on the vertical velocities.

Remark: At the moment it is not clear, which thermodynamic hypothesis and thus which resulting approximation for the saturation vapour pressure over liquid water is physically correct, although the formulation by Murphy and Koop (2005) seems to agree with recent measurements (Pathak et al., 2021). In particular, it is not clear if the formulation of Nachbar et al. (2019) can be extrapolated to values $T < 200 \text{ K}$. Thus, we cannot recommend to use a certain formulation.




Figure 10. Impact of the formulation of the saturation vapour pressure by Nachbar et al. (2019) on the idealized nucleation events. Left: ice crystal number concentrations, right: maximum supersaturation values. The relative differences in number concentrations are always smaller than 15%

6 Another approach to formulate the nucleation rate

Up to now we always employed the “reference” nucleation rate in our computations, i.e. the formulation as in Koop et al. (2000) but corrected by a constant offset, see Section 4.1, in order to match the nucleation rate for pure water droplets by Koop and Murray (2016) in a certain temperature range. In this section we take a different point of view assuming that we can just directly adopt the formulation by Koop and Murray (2016) for the nucleation rate of aqueous solution droplets, providing an exact match of both curves by definition. In the following we discuss the consequences of using such a direct approach in terms of nucleation events.

6.1 Direct fit to nucleation rate of pure water

In order to arrive at a direct fit, we assume that at water saturation, the freezing of pure water droplets should behave as the freezing of solution droplets at super-cooled states. To avoid a complicated reformulation of the formula from Koop and Murray (2016) in terms of the water activity Δa_w , we use a quadratic polynomial fit to the original formulation $J_{\text{hom}}(T)$ at water saturation, see Appendix C for details. Figure 11 presents the original data (black curve) together with the quadratic fit (red curve) and the corrected formulation of Koop et al. (2000) (blue curve) from section 4.1. In contrast to the (corrected) formulation by Koop et al. (2000) the nucleation rate reaches a maximum at $\Delta a_w \sim 0.345$ and decreases afterwards. As a result, there is a significant deviation between the two nucleation rates ($J_{\text{K2000}} > J_{\text{KM2016}}$) for the range $\Delta a_w > 0.32$. Thus, we can expect that for cold temperatures and/or high upward motions there will be large deviations in the produced ice crystal number concentrations within nucleation events.




Figure 11. Freezing rate of water droplets (Koop and Murray, 2016, black), a polynomial fit (red), and the nucleation rate of solution droplets (Koop et al., 2000, corrected, blue), all depending on Δa_w .

It should be kept in mind that the range of the parameterisation of the nucleation rate as given in Koop et al. (2000) is restricted to the interval $0.26 \leq \Delta a_w \leq 0.34$. As a result, it is not clear if the parameterisation works well for values $\Delta a_w > 0.34$. However, there are measurements (see Laksmono et al., 2015) for the freezing of pure water droplets that also show a kind of plateau at cold temperatures (corresponding to high values of Δa_w). Thus, for higher values $\Delta a_w > 0.34$ we use the value $J_{\text{KM2016}}(\Delta a_w) = J_{\text{KM2016}}(0.34)$ to (a) mimic the plateau in the measurements, and (b) avoid numerical issues in the simulations. Assuming that the nucleation rate does not depend on other quantities than water activity, it may now be used in numerical simulations of homogeneous nucleation events.

6.2 Numerical simulations of nucleation events

After having obtained the direct formulation of the nucleation rate of (Koop and Murray, 2016), we now investigate its impact on nucleation events using numerical simulations as before. For completeness, two different types of simulations are done: (1) Simulations using the standard formulation of p_{liq} by Murphy and Koop (2005) and (2) simulations using the new formulation of p_{liq} by Nachbar et al. (2019). The results of the simulations are shown in figure 12.

First we consider the ice crystal number concentrations (left panel). For low vertical updrafts, the values of n_i are only slightly affected in case of using the adapted nucleation rate. For higher vertical velocities, there is a reduction in the produced ice crystal number concentrations; this reduction increases with increasing vertical updrafts. This effect can be explained as




Figure 12. Impact of the direct formulation of the nucleation rate based on Koop and Murray (2016) on the idealized nucleation events. Black triangles and lines indicate the reference simulation, red squares and lines denote the use of the nucleation rate based on Koop and Murray (2016), and blue squares and lines represent the use of the nucleation rate based on Koop and Murray (2016) together with the saturation vapour pressure due to Nachbar et al. (2019). Left: ice crystal number concentrations, right: maximum supersaturation values.

follows. The nucleation rates differ significantly for higher values $\Delta a_w \geq 0.31$, i.e. the slope of the adapted rate is (much) smaller than the original nucleation rate by Koop et al. (2000). For higher updrafts, the supersaturation reaches higher values, which is equivalent to higher values of Δa_w . Thus, the nucleation rates differ for these high updraft events, and less ice crystals are produced for using the adapted nucleation rate. Apart from the influence at high vertical velocities, there is almost no difference in the ice crystal number concentrations between the nucleation events using the different formulations.

Considering the values of maximum supersaturation (right panel), there is a similar behaviour as for n_i . At low vertical velocities there is almost no difference between the reference nucleation rate and the newly adapted rate. In case of using the saturation vapour pressure according to Nachbar et al. (2019), the observed shift in the maximum supersaturation values stems from the increased difference between the values of the saturation vapor pressures at low temperatures, see Section 5. At higher updrafts ($w > 0.5 \text{ m s}^{-1}$), the maximum supersaturation values increase nonlinearly. For the coldest temperature ($T = 196 \text{ K}$) we note a dramatic increase up to very high values ($S_{i,\text{max}} \sim 1.8$). However, note that in all cases the values of the maximum supersaturation stays below water saturation, hence no liquid origin ice formation would occur.

7 Thresholds of ice nucleation

For the evaluation of measurements of ice clouds, the possible range of supersaturation is often estimated using the so-called Koop-line, i.e. the supersaturation threshold $S_c(T)$ which corresponds to a nucleation rate value $J = 10^{16} \text{ m}^{-3} \text{ s}^{-1} = 10^{10} \text{ cm}^{-3} \text{ s}^{-1}$. In many investigations (see, e.g., Krämer et al., 2009) this function is used as an upper bound for possible

635 values of S_i inside and also outside of ice clouds. However, from our investigations in this study so far, we have to carefully consider two different aspects from a purely theoretical point of view:

1. The nucleation threshold assigned to the frequently used value $j_0 = 16$ is arbitrary chosen; there is no convincing physical justification for using this particular value. In Koop et al. (2000) different values $J = 10^{j_0} \text{ m}^{-3} \text{ s}^{-1}$ with $j_0 \in [1, 17]$ are used, but for testing the impact of droplet sizes, they used the value $j_0 = 16$. Nucleation of ice crystals is not a switching
640 process, it occurs gradually and smooth, although the nucleation rates are very steep functions of the supersaturation. The size or strength of the nucleation event cannot be determined just by the maximum of the supersaturation; the amount of ice crystals as formed in the nucleation event is determined by the integral over the supersaturation curve (see, e.g., the discussion in Dinh et al., 2016). Thus, it is possible to form many crystals in lower updrafts even if the high nucleation threshold is not reached. From our simulations we observe that the peak supersaturation for nucleation
645 events depends crucially on the vertical velocity, i.e. on the temperature rate, which is prescribed during the event. This is quite obvious from the differential equation determining the change of S_i : The peak value is given by $\frac{dS_i}{dt} = 0$, i.e. when source and sink terms balance each other. Since the source includes the vertical velocity linearly, the dependence of the peak supersaturation on w is obvious, although not linear.
2. As described above in section 5, it is still not clear which formulation of the saturation vapor pressure is physically
650 correct. However, the use of the formulation by Nachbar et al. (2019) leads to a higher saturation vapour pressure and thus to a higher nucleation threshold, even for arbitrary values j_0 and its associated nucleation threshold $S_{cx0}(T)$.

Taking these two aspects into account, we can observe the following behaviour. In figure 13 (left panel) we compare the nucleation thresholds for the saturation vapour pressure according to Murphy and Koop (2005) for $j_0 = 10$ (red curve) and $j_0 = 16$ (dark blue curve) with the range of peak supersaturations for vertical velocities $0.01 \text{ m s}^{-1} \leq w \leq 2 \text{ m s}^{-1}$ (black
655 vertical bar) and the maximum value for a very unrealistic value $w = 10 \text{ m s}^{-1}$ (black crosses). For comparison, the well known Koop-line as fit and proposed by Kärcher and Lohmann (2002) is plotted (light blue curve). It is quite obvious, that for typical vertical velocity values the “classical” Koop-line is not reached, i.e. the peak supersaturation is below the threshold. Nevertheless, for strong cooling rates (very high vertical velocities), as are used in experiments in cloud chambers, high supersaturations are reached, which still partly remain below the Koop-line. If we change the saturation vapour pressure to the
660 formulation by Nachbar et al. (2019), the qualitative picture remains the same (right panel in fig 13): Even for high vertical updrafts the high nucleation rates are reached; for moderate and small updrafts, the peak supersaturation stays well below the classical nucleation threshold. However, the nucleation thresholds are generally shifted to higher values of supersaturation due to the different saturation vapour pressure formulation. It seems that these values fit better to the experiments in the AIDA cloud chamber as reported in Baumgartner et al. (2022) and Schneider et al. (2021). This might be interpreted as a hint that the
665 formulation by Nachbar et al. (2019) might be the more appropriate formulation for the saturation vapour pressure, although the formulation by Murphy and Koop (2005) agrees well with recent measurements (Pathak et al., 2021). In any case, one has to consider the impact of the cooling rate on the peak supersaturation in a nucleation event. Therefore, the use of the “Koop-line” in the currently applied way is misleading and does not correspond to the actual physics of nucleation events. Note, that the

temperature dependent threshold is used in some parameterisations of ice clouds in climate and numerical weather prediction models (see, e.g., Kärcher et al., 2006; Köhler and Seifert, 2015). A simple but albeit more realistic extension of such schemes would be a threshold depending on both, vertical velocity w and temperature T ; a 2D fit to the maximum supersaturation data from our simulations might be a first attempt into this direction.




Figure 13. Comparison of nucleation thresholds (red curve: $x_0 = 10$, blue curve $x_0 = 16$) and the classical “Koop-line” (light blue curve). The black vertical bars indicate the range of peak supersaturation ratios within the nucleation events computed using vertical velocities ranging from 0.01 m s^{-1} to 2 m s^{-1} . The black cross corresponds to the peak supersaturation ratio for the vertical velocity of 10 m s^{-1} . Left panel: Curves based on the water activity using the saturation vapour pressure formulation by Murphy and Koop (2005); right panel: the same for the saturation vapour pressure formulation by Nachbar et al. (2019).

Finally, we can also investigate the peak supersaturation values for the new empirical nucleation rate formulation, as derived in section 6.1. Generally, we see the same behaviour as for the reference simulations with a monotonic increase of peak supersaturation values with increasing vertical velocity (cf. figure 14). The use of the saturation vapour pressure formulation by Nachbar et al. (2019) additionally enhances the peak values as seen before. However, the peak values for cold temperatures and very high vertical velocities are strongly enhanced in comparison with the reference simulations. Also these high values are still in line with the measurements in the AIDA chamber as reported by Baumgartner et al. (2022) and Schneider et al. (2021).

680 8 Summary and outlook

We have investigated the impact of the representation of nucleation rates and diffusional growth on idealized nucleation events, as driven by a constant vertical updraft (i.e. a constant cooling rate). In a first step, we have investigated the original formulation




Figure 14. Same as in figure 13, but using the nucleation rate as empirically derived in section 6.1.

of the nucleation rate for homogeneous freezing of aqueous solution droplets in the formulation by Koop et al. (2000); for a better agreement with the nucleation rate of pure water droplets a simple shift could be applied. For analytical purposes and simple model calculations, a less complicated formulation is desired. We showed that a linear fit to the original formulation depending on the difference in water activity $\Delta a_w = a_w - a_w^i$ is accurate enough to reproduce the ice crystal number concentrations quantitatively. Based on this linearization approach, we derived a threshold formulation of the nucleation rate, which can be used for analytical investigations as already presented in Baumgartner and Spichtinger (2019). Again, the new formulations are good enough to represent nucleation events quantitatively as compared to the reference nucleation formulation.

Using the linear approximation as a starting point, we investigated the impact of different formulations on idealized nucleation events, changing the two relevant parameters (slope and constant offset). These investigations led to the first major results:

- The absolute values of the nucleation rate has only marginal impact on the resulting ice crystal number concentrations in a nucleation event. Even a scaling by up to six orders of magnitudes did not severely affect the resulting number concentrations. However, the maximum supersaturations changed, and the resulting deviations range up to few percent relative humidity. In addition, the time of nucleation onset is slightly shifted.
- The slope of the nucleation rate (or more precisely in the argument of the exponential function) has a much larger impact on the resulting nucleation event, and the ice crystal number concentration. Variations in the slope change the number concentrations in the nucleation events by up to a factor 2.5 (in both directions). Also, the maximum supersaturation is affected by a deviation of a few percent of relative humidity.

As a final conclusion of this part of our work, we can state that the shape of the nucleation rate is of high importance for the representation of the nucleation process, whereas the absolute strength of the rate is almost negligible, if the values are high enough. This shows that the nucleation process (homogeneous freezing of solution droplets) itself is a quite robust process, thus the accurate formulation is maybe less critical as we thought. Also the amount of available solution droplets as controlled
705 by the background aerosol does not affect the nucleation events itself; it can be seen as a scaling factor of the nucleation rate, in the same sense as in the sensitivity analysis of changing the absolute values of nucleation rates. As long as the amount of aerosol particles is some orders of magnitude larger than the ice crystal number concentration as predicted for a nucleation event, this does not play a role for the nucleation events, and we do not have to care about exhausting the reservoir of solution droplets.

710 We also investigated the impact of a recently published formulation of the water saturation pressure based on a thermodynamic assumption of different phases of water in the very low temperature range (Nachbar et al., 2019). This new formulation leads to changes in the function a_w^i , which directly affected the nucleation rate based on Δa_w . Following the derivations of the threshold description, approximations were constructed. The new resulting functions $a_w^i(T)$ and $S_c(T)$ can be accurately approximated with polynomials of smaller degrees, as compared to the standard formulation. The new formulation of p_{liq} only
715 marginally changed the resulting ice crystal number concentrations. However, the impact on the maximum supersaturations increased with decreasing temperature up to few percent of relative humidity. Overall, the two different representations of the saturation vapour pressure over liquid water produced very similar, even almost identical, results. Thus, a decision about the validity of a certain formulation must be left to extensive experimental measurements.

In a more speculative part of the study we adapted the nucleation rate of homogeneous freezing of pure water droplets (Koop
720 and Murray, 2016) as a new parameterisation for homogeneous freezing of aqueous solution droplets. This representation is quite similar for low values of Δa_w to the original formulation by Koop et al. (2000) and its approximations. However, for very high water activities (i.e. high supersaturations as driven by large vertical updrafts), there is a significant deviation from the reference nucleation rate. Thus, for some cases in the parameter space (high updrafts and low temperatures) there is a significant deviation in the number concentrations, and, more obvious, in the maximum supersaturations, which reach
725 almost water saturation in some cases. This approach showed that the shape of the nucleation rate is important for the resulting nucleation events; strong deviations of the shape from its reference affect the results of the nucleation event significantly. If this representation of the nucleation rate is more close to the physics of ice nucleation remains open, and might be an objective for experimental investigations.

Finally, we investigated the commonly used threshold for homogeneous nucleation (“Koop-line”) in the light of peak super-
730 saturation values during nucleation events. This threshold corresponds to a nucleation rate of $10^{16} \text{m}^{-3} \text{s}^{-1}$, but is only rarely reached during nucleation events. Nucleation itself starts usually at much lower values of S_i corresponding to lower values of the nucleation rate. The peak supersaturation during a nucleation event, characterised as an equilibrium between sources and sinks of supersaturation depends on temperature and vertical velocity. The peak supersaturation is a much more physical quantity to investigate the strength of a nucleation event. The peak supersaturation as diagnosed from the numerical simulations

735 might be a more physical representation of ice nucleation in coarse resolution models in comparison to the frequently used
nucleation threshold.

It should be emphasized that all the results and conclusions are meant in a bulk-sense, i.e. for a large collection of ice crystals
such as a newly forming cirrus cloud. If one is interested in the details of ice formation for a single or only a small number of
particles then all details of the nucleation rate might be equally important. In that respect, our study shows that homogeneous
740 cirrus formation is a robust physical process.

Appendix A: Model description - details

In this appendix, we present the details of the model as used for the numerical simulations of the nucleation events. Note, that
we use the mathematical (and also programming) notation of logarithms, i.e. \log denotes the natural logarithm (to base e).

Background aerosol

745 We assume for the aqueous solution droplets in the tropopause region a size distribution of lognormal type:

$$f_{\text{sol}}(r) = \frac{n_a}{\sqrt{2\pi} \log \sigma_r} \exp\left(-\frac{1}{2} \left(\frac{\log(r/r_{\text{sol}})}{\log \sigma_r}\right)^2\right) \frac{1}{r} \quad (\text{A1})$$

with a modal radius $r_{\text{sol}} = 75 \cdot 10^{-9}$ m, and a geometric standard deviation $\sigma_r = 1.5$. These values are adapted from the more
complex model by Spichtinger and Gierens (2009), using the fact that the dry aerosol population, as used in Spichtinger and
Gierens (2009), has grown to larger sizes by water vapour uptake (i.e. assuming Köhler theory, see, e.g. Köhler, 1936). The
750 mean volume of the solution droplets

$$V_d = V_{\text{sol}} = \frac{4}{3} \pi r_{\text{sol}}^3 \cdot \exp\left(\frac{9}{2} (\log \sigma_r)^2\right) \quad (\text{A2})$$

is calculated from the third moment of the lognormal distribution.

Mass distribution for ice crystals

For the ice crystals, we assume a mass distribution of lognormal type

$$755 \quad f(m) = \frac{n_i}{\sqrt{2\pi} \log \sigma_m} \exp\left(-\frac{1}{2} \left(\frac{\log(m/m_m)}{\log \sigma_m}\right)^2\right) \frac{1}{m} \quad (\text{A3})$$

with a parameter

$$r_0 = \exp\left((\log \sigma_m)^2\right), \quad \bar{m} = m_m \sqrt{r_0} = 3 \quad (\text{A4})$$

representing the width of the distribution as described in Spichtinger and Gierens (2009). This distribution is used for the
derivation of the rates in the ODE system for the mean quantities of ice mass and number concentration. The integration of

760 weighting functions of the type $m^k, k \in \mathbb{R}_+$ leads to general moments, which can be computed analytically:

$$\begin{aligned} \mu[m]_k &:= \int_0^\infty m^k f(m) dm = n_i \cdot m_m^k \exp\left(\frac{1}{2}(k \log \sigma_m)^2\right) \\ &= n_i \cdot \bar{m}^k r_0^{\frac{k(k-1)}{2}} \end{aligned} \quad (\text{A5})$$

Note, that for the averaged quantities we obtain $n_i = \mu[m]_0, q_i = \mu[m]_1$, respectively. Thus, we use a double moment scheme in our model.

Diffusion constant

765 For the diffusion of water vapour in dry air, we use the following expression

$$D_v = D_{v0} \left(\frac{T}{T_0}\right)^{1.94} \left(\frac{p_0}{p}\right) \quad (\text{A6})$$

which is an empirical fit to measurement data (Hall and Pruppacher, 1976). Note, that the valid temperature range is different in the book Pruppacher and Klett (2010) and in the original article Hall and Pruppacher (1976). For analytical investigations, a representation using a quadratic temperature dependence constitute a good approximation for a restricted temperature range.

770 For the kinetic correction we use the function

$$f_D(r, a, b) = \frac{1}{\frac{r}{r+a} + \frac{b}{r}} = \frac{r^2 + ar}{r^2 + br + ab} \quad (\text{A7})$$

whereas r denotes the radius of the ice crystal (using a bulk density of ice $\rho_b = 0.81 \text{ kg m}^{-3}$), and the parameters are given by

$$a = \lambda \cdot C_{\text{cunn}}, \quad b = \frac{4D_v}{\alpha_m \bar{c}_v} \quad (\text{A8})$$

using the mean free path of water molecules in air λ (acc. to Pruppacher and Klett, 2010), the Cunningham correction factor

775 $C_{\text{cunn}} = 0.7$, and the mean velocity of water molecules \bar{c}_v . We set the accommodation coefficient $\alpha_m = 0.5$ for comparison with former investigations (Kärcher and Lohmann, 2002); this value is also within the range as recommended in recent work by Skrotzki et al. (2013).

For representing the growth rates for the ensemble of ice crystals, by comparison with numerical integration we find that using a shifted mean mass $m_1 = c_1 \cdot \bar{m}$, $c_1 \approx 0.819$ in the kinetic correction function $f(r_1, a, b)$ is a good approximation.

780 Howell factor

Latent heat release due to phase changes during diffusional growth changes the surface temperature of the ice crystal. For taking this into account, we use the Howell factor

$$\begin{aligned} G_v &= \left[\left(\frac{L}{R_v T} - 1 \right) \frac{L}{T} \frac{D_v^*}{K_T^*} + \frac{R_v T}{p_{si}} \right]^{-1} \\ &\approx \left[\left(\frac{L}{R_v T} - 1 \right) \frac{L}{T} \frac{D_v}{K_T} + \frac{R_v T}{p_{si}} \right]^{-1}. \end{aligned} \quad (\text{A9})$$

In the approximation, we neglect the kinetic corrections for diffusion coefficient D_v and heat conductivity of air K_T .

785 Capacity of ice crystals

For ice crystals we assume spherical shape for small crystals and columnar shape for large crystals as in Spichtinger and Gierens (2009); thus the shape factor, or capacity, can be determined exactly using the electrostatic analogy (McDonald, 1963), using a prolate spheroid with semi axes a, b ; the capacity can be analytically expressed by

$$C = \frac{L \epsilon'}{\log\left(\frac{1+\epsilon'}{1-\epsilon'}\right)} \quad (\text{A10})$$

790 using the eccentricity $\epsilon' = \sqrt{1 - \left(\frac{b}{a}\right)^2}$ and the length L of the crystal, which in turn is a function of the crystal mass. Note, that the eccentricity changes with crystal growth since the aspect ratio is changing (see Spichtinger and Gierens, 2009, their eq. (17)) We find a very good approximation to the piece-wise definition of the capacity by Spichtinger and Gierens (2009) depending on the ice crystal mass

$$C(m) \approx a_1 \cdot m^{b_1} + a_2 \cdot m^{b_2} \quad (\text{A11})$$

795 with constants

$$\begin{aligned} a_1 &= 0.015755 \text{ m kg}^{\frac{1}{b_1}}, \quad b_1 = 0.3, \\ a_2 &= 0.33565 \text{ m kg}^{\frac{1}{b_2}}, \quad b_2 = 0.43. \end{aligned} \quad (\text{A12})$$

The representation of the capacity in the ice crystal ensemble is given by the integration, leading to general moments $\mu[m]_{b_i}$.

Ventilation correction

The empirical ventilation corrections usually depend on the use of two dimensionless numbers, i.e. the Schmidt number N_{Sc}
800 and the Reynolds number N_{Re}

$$N_{\text{Sc}} = \frac{\mu}{D_v \rho}, \quad N_{\text{Re}} = \frac{\rho}{\mu} v_t L \quad (\text{A13})$$

using the dynamic viscosity of air μ (e.g. Dixon, 2007). Thus, the size of the ice crystal L is influencing the Reynolds number via the product $v_t(m)L$, using the terminal velocity v_t for an ice crystal of mass m . The effect of ventilation, i.e. the additional uptake of water vapour by the airflow around the particle crucially depends on the shape of the particles. For columnar shaped
805 ice crystals, we adapt the empirical quadratic fit by Liu et al. (2003) to the simulation data (Ji and Wang, 1999) as follows

$$f_v = 1 + c_\chi \cdot \chi^2, \quad c_\chi = 0.14856, \quad \chi = N_{\text{Sc}}^{\frac{1}{3}} N_{\text{Re}}^{\frac{1}{2}} \quad (\text{A14})$$

For the formulation of the terminal velocity of columnar shaped ice crystals, $v_t(m)$, we use the formulation by Spichtinger and Gierens (2009), including also the correction for temperature and pressure, respectively. For representing the ensemble of ice crystals, by comparison with the numerical integration we find that using a shifted mean mass $m_2 = c_2 \cdot \bar{m}$, $c_2 = 1.5$ in the
810 formulation of the Reynolds number leads to a very good agreement.

Appendix B: Reference simulation results

In this section we report on the results of the reference simulations, using the corrected formulation of the nucleation rate for super-cooled aqueous solution droplets by Koop et al. (2000). For evaluating the quality of the simplified model, we compare the number concentration of ice crystals as obtained from standard nucleation events with results from literature, i.e. with a
815 model using sophisticated particle physics (Kärcher and Lohmann, 2002) and a complex bulk physics scheme (Spichtinger and Gierens, 2009). In figure B1 the results are represented for the temperatures $T = 196, 216, 236$ K at pressure $p = 200$ hPa, as prescribed in Kärcher and Lohmann (2002).




Figure B1. Comparison of ice crystal number concentrations as obtained for typical nucleation events from different models. Red squares: Particle model by Kärcher and Lohmann (2002), blue circles: complex two moment bulk scheme by Spichtinger and Gierens (2009), black line & triangles: simpler bulk model from this study, indicated as new reference

In comparison we see an overall good agreement of our simple model with the more sophisticated models (Kärcher and Lohmann, 2002; Spichtinger and Gierens, 2009). However, we have to remark here that the deviation in the results for temper-
820 ature $T = 236$ K at low vertical velocities is the result of the neglectance of the ventilation correction in the model by Kärcher and Lohmann (2002). In summary, our simplified approach compares very well with the results of the other studies.

In figure B2 a typical nucleation event is shown. Here, two different nucleation parameterizations are used, the reference by Koop et al. (2000) (black line) and the linear fit (red line). There are small differences in the time evolution of the variables

saturation ratio S_i (left panel), number concentration n_i (middle panel) and mean mass m (right panel), but in general there is
 825 the same behaviour in both cases.




Figure B2. Representative example for a typical nucleation event for temperature $T = 216$ K and pressure $p = 200$ hPa with a forcing of $w = 1$ m s⁻¹. Red line: reference nucleation rate after Koop et al. (2000), black line: nucleation rate approximated by linear function as given in eq. (38).

The source of supersaturation (i.e. cooling by vertical updraft and adiabatic expansion) leads to an increase in S_i until nucleation starts at about $t_{\text{start}} \sim 40$ s, i.e. at very low values of the nucleation rate. S_i is still increasing since the sink of depositional growth is not strong enough to reduce water vapour efficiently; thus, the ice crystal number concentration is further increasing due to permanent ice nucleation. At the peak supersaturation, source and sink of supersaturation are balanced
 830 ($t_{\text{peak}} \sim 110$ s); after this time, S_i is decreasing due to the dominant growth term. The number concentration does not change much from this time on but as long as the values of S_i are large enough, still ice nucleation takes place. At about $t \sim 125$ s the nucleation event is complete, no further nucleation takes place, since the nucleation rate is too small. Note that during the time interval $[t_{\text{start}}, t_{\text{peak}}]$ the mean mass m is almost constant (this feature is more prominent in the linear fit case), whereas for $t > t_{\text{peak}}$ the mass increases. For $t < t_{\text{peak}}$ the nucleation is dominant, thus diffusional growth just compensates the number increase
 835 by mass, whereas afterwards crystal growth is dominant over nucleation. This feature was already seen in former investigations, which leads to a model reduction for analytical investigations (Baumgartner and Spichtinger, 2019). The different nucleation parameterisations agree qualitatively for a nucleation event; however, the nonlinear reference rate leads to some variations. While for the linear fit case, the increase in n_i is approximately an exponential growth $n_i(t) \sim \exp(\alpha t)$, and in turn the mean mass is almost constant in the relevant time interval, for the reference case the change deviates slightly from exponential
 840 growth.

Note, that the thresholds of constant nucleation rates in figure B2 (left panel) can be calculated from eq. (39) using the respective values for j_0 (i.e. $j_0 \in \{10, 12, 14, 16\}$) in the formulation of the supersaturation threshold.

Appendix C: Simple fit for nucleation rate of pure water droplets

In Koop and Murray (2016) a polynomial of degree 6 is used for fitting the experimental values of the nucleation rate for pure
 845 super-cooled water. Since polynomials of high degree are difficult to evaluate numerically, we present fits with polynomials of
 lower degrees, which are still accurate in the relevant temperature range. The original formulation of the nucleation rate is

$$J_{\text{hom}}(T) = 10^{p_n(x)}, \quad p_n(x) = \sum_{i=0}^n c_i \cdot x^i, \quad x = T - T_m. \quad (\text{C1})$$

with a polynomial $p_n(x)$ of degree $n = 6$ using the melting temperature of pure water $T_m = 273.15\text{K}$. The coefficients c_i
 are reported in Koop and Murray (2016, table VII), where the nucleation rate is given in units $\text{cm}^{-3} \text{s}^{-1}$. We reformulate
 850 the nucleation rate in SI units (i.e. $[J] = \text{m}^{-3} \text{s}^{-1}$) by a factor of 10^6 and approximate the logarithmic values $\log_{10}(J)$ by
 polynomials of degree 2 and 4, respectively, i.e.

$$\begin{aligned} p_2(T) &= a_0 + a_1 \cdot T + a_2 \cdot T^2, \\ p_4(T) &= a_0 + a_1 \cdot T + a_2 \cdot T^2 + a_3 \cdot T^3 + a_4 \cdot T^4 \end{aligned} \quad (\text{C2})$$

the coefficients are given in table C1. For this purpose we use a least square fit for the temperature range $225 \leq T \leq 245\text{K}$,
 for which supercooled water droplets can still exist (see, e.g., figure 4 in Koop and Murray, 2016). In figure C1 (left panel) the
 855 approximations are shown in comparison with the original fit, while the ratio $r = \frac{p_i(T)}{p(x)}$ is shown in the right panel.




Figure C1. Polynomial fits of low degrees for the nucleation rate as given by Koop and Murray (2016). Left: Reference and fits $p_2(T)$, $p_4(T)$, right: ratio of reference and fits $p_2(T)$, $p_4(T)$

As can be seen the relative error for the polynomial fit $p_4(T)$ is less than 0.25%, while even for the quadratic fit $p_2(T)$ the
 error is smaller than 2%. For practical applications in the relevant temperature range $225 \leq T \leq 240\text{K}$ the quadratic fit might

be sufficient. If the original polynomial is used, a sophisticated evaluation of the polynomial is recommended (e.g. Horner scheme).

fit	a_0	a_1	a_2	a_3	a_4
$p_2(T)$	-5369.61	46.96750	-0.10236	-	-
$p_4(T)$	-848143.02	14534.5767	-93.481032	0.26745460	-0.0002872

Table C1. Coefficients for the polynomial fits of the nucleation rate by Koop and Murray (2016) as given in equation (C2).

860 Appendix D: Perturbation Analysis

The perturbation analysis or asymptotic analysis as applied in Section 4.6 is a well-known technique to investigate the impact of perturbations on a mathematical object such as a mathematical expression or the solution of an equation. A good general introduction is given in Holmes (2013) and an application in meteorology is explained in Klein (2010).

The basic idea is to introduce a small parameter ε , to expand the quantity of interest in powers of ε , i.e. $\varepsilon, \varepsilon^2, \dots$ and to substitute this expansion into the mathematical object of interest (see Eq. (48) for such an expansion). Since the resulting expression should hold for any value of ε and even in the limit $\varepsilon \rightarrow 0$, all the contributions from the various powers of ε may be considered individually. Given that the parameter ε is assumed as being small, effects that stem from terms with higher powers of ε will only have a small impact whereas effects with a lower power of ε will be dominant.

For practical applications it is common to also scale coefficients and parameters of the mathematical expression in powers of ε . This step ensures that the mutual magnitudes of the parameters stay consistent, even in the limit $\varepsilon \rightarrow 0$. This task involves usually some free choices, and is known as distinguished limit.

In the spirit of the works of Klein and Majda, the scaling of the parameters is often done by assuming $\varepsilon \sim 0.1$, i.e. substituting this value of ε into the scaling of the parameters yields a realistic value of the parameters (see, e.g., Hittmeir and Klein, 2017; Baumgartner and Spichtinger, 2019; Klein and Majda, 2006). As an example, the parameter $A(T) \approx A_0 = b_1 a_{w_0}^i \approx 149.32$ is written as $A_0 = A^* \varepsilon^{-2}$ with $A^* = \mathcal{O}(1)$ where the latter may be understood as A^* is independent of ε . With $A^* \approx 1.4932$ the value $\varepsilon \sim 0.1$ restores the original value of A_0 .

In essence, the goal is to determine the nonzero-parts of the expansion of the mathematical expression and the respective power of ε indicates how strong this contribution is. As an example, if the final expansion is found to be $\varepsilon^{-1}\omega_{-1} + \omega_0 + \varepsilon\omega_1 + \mathcal{O}(\varepsilon^2)$ then the term ω_{-1} will be dominant since this term is associated with the lowest power of ε . If the result would be the expansion $\omega_0 + \varepsilon\omega_1 + \mathcal{O}(\varepsilon^2)$ then we may conclude that ω_0 is the dominant part and all ω_k for $k \geq 1$ only contribute small corrections (since ε is small).

Another fruitful use of perturbation analysis is to allow an equation to determine the possible matching powers of ε , i.e. to answer the question: Which powers of ε are needed to achieve a balance in the equation at hand? As an example, from physical

885 considerations we know that equation (59) holds. As a consequence, after having expanded both sides of the equation in powers of ε , the expansions on both sides must agree (otherwise the equation would not hold). This is only possible if the powers of ε match, hence the appropriate powers for ε may be inferred.

Author contributions. PM carried out calculations and approximations of the nucleation rates. PS run the numerical simulations. MB and PS carried out the investigations using asymptotics. All authors were involved in the preparation of the manuscript and have read and approved the final manuscript version.

890 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. Patrik Marschalik and Peter Spichtinger acknowledge support by the German Bundesministerium für Bildung und Forschung (BMBF) within the HD(CP)² initiative, project S4 (01LK1216A). Manuel Baumgartner acknowledges support by the Deutsche Forschungsgemeinschaft (DFG) within the Transregional Collaborative Research Centre TRR165 Waves to Weather, (www.wavestoweather.de), project Z2. Peter Spichtinger acknowledges support by the DFG within the Transregional Collaborative Research Centre TRR301 TPChange, (tpchange.de), project B7. We thank Martina Krämer for fruitful discussions during her stay as a Gutenberg Research College fellow at JGU Mainz, and Katrin Amann-Winkel for providing us a study about recent measurements of supercooled water. Finally, we thank two anonymous reviewers for their helpful comments and suggestions which lead to a significantly improved manuscript.

References

- 900 Baumgartner, M. and Spichtinger, P.: Homogeneous nucleation from an asymptotic point of view, *Theoretical and Computational Fluid Dynamics*, 33, 83–106, <https://doi.org/10.1007/s00162-019-00484-0>, 2019.
- Baumgartner, M., Weigel, R., Harvey, A., Plöger, F., and Spichtinger, P.: Reappraising the appropriate calculation of a common meteorological quantity: potential temperature, *Atmospheric Chemistry and Physics*, 20, 15 585–15 616, <https://doi.org/10.5194/acp-20-15585-2020>, 2020.
- 905 Baumgartner, M., Rolf, C., Groöß, J.-U., Schneider, J., Schorr, T., Möhler, O., Spichtinger, P., and Krämer, M.: New investigations on homogeneous ice nucleation: the effects of water activity and water saturation formulations, *Atmospheric Chemistry and Physics*, 22, 65–91, <https://doi.org/10.5194/acp-22-65-2022>, 2022.
- de Reus, M., Borrmann, S., Bansemmer, A., Heymsfield, A. J., Weigel, R., Schiller, C., Mitev, V., Frey, W., Kunkel, D., Kuerten, A., Curtius, J., Sitnikov, N. M., Ulanovsky, A., and Ravegnani, F.: Evidence for ice particles in the tropical stratosphere from in-situ measurements, *Atmospheric Chemistry and Physics*, 9, 6775–6792, <https://doi.org/10.5194/acp-9-6775-2009>, 2009.
- 910 Dinh, T., Podglajen, A., Hertzog, A., Legras, B., and Plougonven, R.: Effect of gravity wave temperature fluctuations on homogeneous ice nucleation in the tropical tropopause layer, *Atmospheric Chemistry and Physics*, 16, 35–46, <https://doi.org/10.5194/acp-16-35-2016>, 2016.
- Dixon, J.: *The Shock Absorber Handbook*, Wiley, 2007.
- Fu, Q. and Liou, K.: Parameterization of the Radiative Properties of Cirrus Clouds, *J. Atmos. Sci.*, 50, 2008–2025, [https://doi.org/10.1175/1520-0469\(1993\)050<2008:POTRPO>2.0.CO;2](https://doi.org/10.1175/1520-0469(1993)050<2008:POTRPO>2.0.CO;2), 1993.
- 915 Fusina, F., Spichtinger, P., and Lohmann, U.: Impact of ice supersaturated regions and thin cirrus on radiation in the midlatitudes, *Journal of Geophysical Research*, 112, <https://doi.org/10.1029/2007jd008449>, 2007.
- Gallo, P., Amann-Winkel, K., Angell, C. A., Anisimov, M. A., Caupin, F., Chakravarty, C., Lascaris, E., Loerting, T., Panagiotopoulos, A. Z., Russo, J., Sellberg, J. A., Stanley, H. E., Tanaka, H., Vega, C., Xu, L., and Pettersson, L. G. M.: Water: A Tale of Two Liquids, *Chemical Reviews*, 116, 7463–7500, <https://doi.org/10.1021/acs.chemrev.5b00750>, 2016.
- 920 Gallo, P., Loerting, T., and Sciortino, F.: Supercooled water: A polymorphic liquid with a cornucopia of behaviors, *Journal of Chemical Physics*, 151, <https://doi.org/10.1063/1.5135706>, 2019.
- Gasparini, B., Munch, S., Poncet, L., Feldmann, M., and Lohmann, U.: Is increasing ice crystal sedimentation velocity in geoengineering simulations a good proxy for cirrus cloud seeding?, *Atmospheric Chemistry and Physics*, 17, 4871–4885, <https://doi.org/10.5194/acp-17-4871-2017>, 2017.
- 925 Hall, W. and Pruppacher, H.: Survival of Ice Particles Falling From Cirrus Clouds in Subsaturated Air, *Journal of the Atmospheric Sciences*, 33, 1995–2006, [https://doi.org/10.1175/1520-0469\(1976\)033<1995:TSOIPF>2.0.CO;2](https://doi.org/10.1175/1520-0469(1976)033<1995:TSOIPF>2.0.CO;2), 1976.
- Hittmeir, S. and Klein, R.: Asymptotics for moist deep convection I: refined scalings and self-sustaining updrafts, *Theoretical and Computational Fluid Dynamics*, <https://doi.org/10.1007/s00162-017-0443-z>, 2017.
- Holmes, M. H.: Introduction to Perturbation Methods, vol. 20 of *Texts in Applied Mathematics*, Springer-Verlag, New York, second edition
930 edn., <https://doi.org/10.1007/978-1-4614-5477-9>, 2013.
- Hoose, C. and Möhler, O.: Heterogeneous ice nucleation on atmospheric aerosols: a review of results from laboratory experiments, *Atmospheric Chemistry and Physics*, 12, 9817–9854, 2012.
- Jensen, E. and Pfister, L.: Transport and freeze-drying in the tropical tropopause layer, *Journal of Geophysical Research*, 109, <https://doi.org/10.1029/2003JD004022>, 2004.

- 935 Ji, W. and Wang, P.: Ventilation coefficients for falling ice crystals in the atmosphere at low-intermediate Reynolds numbers, *Journal of the Atmospheric Sciences*, 56, 829–836, [https://doi.org/10.1175/1520-0469\(1999\)056<0829:VCFFIC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1999)056<0829:VCFFIC>2.0.CO;2), 1999.
- Joos, H., Spichtinger, P., and Lohmann, U.: Orographic cirrus in a future climate, *Atmospheric Chemistry and Physics*, 9, 7825–7845, <https://doi.org/10.5194/acp-9-7825-2009>, 2009.
- Joos, H., Spichtinger, P., Reutter, P., and Fusina, F.: Influence of heterogeneous freezing on the microphysical and radiative properties of
940 orographic cirrus clouds, *Atmospheric Chemistry and Physics*, 14, 6835–6852, <https://doi.org/10.5194/acp-14-6835-2014>, 2014.
- Kärcher, B. and Lohmann, U.: A parameterization of cirrus cloud formation: Homogeneous freezing of supercooled aerosols, *Journal of Geophysical Research: Atmospheres*, 107, <https://doi.org/10.1029/2001JD000470>, 2002.
- Kärcher, B., Hendricks, J., and Lohmann, U.: Physically based parameterization of cirrus cloud formation for use in global atmospheric models, *Journal of Geophysical Research*, 111, 2006.
- 945 Klein, R.: Scale-Dependent Models for Atmospheric Flows, *Annual Review of Fluid Mechanics*, 42, 249–274, <https://doi.org/10.1146/annurev-fluid-121108-145537>, 2010.
- Klein, R. and Majda, A. J.: Systematic multiscale models for deep convection on mesoscales, *Theoretical and Computational Fluid Dynamics*, 20, 525–551, <https://doi.org/10.1007/s00162-006-0027-9>, 2006.
- Köhler, H.: The nucleus in and the growth of hygroscopic droplets, *Transactions of the Faraday Society*, 32, 1152–1161, 1936.
- 950 Koop, T.: Homogeneous ice nucleation in water and aqueous solutions, *Zeitschrift für Physikalische Chemie/International journal of research in physical chemistry and chemical physics*, 218, 1231–1258, 2004.
- Koop, T.: Atmospheric water, in: *Water: Fundamentals as the Basis for Understanding the Environment and Promoting Technology*, edited by Debenedetti, P. G., Ricci, M. A., and Bruni, F., vol. 187 of *Proceedings of the International School of Physics “Enrico Fermi”*, pp. 45–75, <https://doi.org/10.3254/978-1-61499-507-4-45>, 2015.
- 955 Koop, T. and Murray, B. J.: A physically constrained classical description of the homogeneous nucleation of ice in water, *Journal of Chemical Physics*, 145, <https://doi.org/10.1063/1.4962355>, 2016.
- Koop, T., Luo, B., Tsias, A., and Peter, T.: Water activity as the determinant for homogeneous ice nucleation in aqueous solutions, *Nature*, 406, 611–614, 2000.
- Krämer, M., Schiller, C., Afchine, A., Bauer, R., Gensch, I., Mangold, A., Schlicht, S., Spelten, N., Sitnikov, N., Borrmann, S., de Reus,
960 M., and Spichtinger, P.: Ice supersaturations and cirrus cloud crystal numbers, *Atmospheric Chemistry and Physics*, 9, 3505–3522, <https://doi.org/10.5194/acp-9-3505-2009>, 2009.
- Krämer, M., Rolf, C., Luebke, A., Afchine, A., Spelten, N., Costa, A., Meyer, J., Zoeger, M., Smith, J., Herman, R. L., Buchholz, B., Ebert, V., Baumgardner, D., Borrmann, S., Klingebiel, M., and Avallone, L.: A microphysics guide to cirrus clouds - Part 1: Cirrus types, *Atmospheric Chemistry and Physics*, 16, 3463–3483, <https://doi.org/10.5194/acp-16-3463-2016>, 2016.
- 965 Krämer, M., Rolf, C., Spelten, N., Afchine, A., Fahey, D., Jensen, E., Khaykin, S., Kuhn, T., Lawson, P., Lykov, A., Pan, L. L., Riese, M., Stroh, F., Thornberry, T., Wolf, V., Woods, S., Spichtinger, P., Quaas, J., and Sourdeval, O.: A microphysics guide to cirrus – Part 2: Climatologies of clouds and humidity from observations, *Atmospheric Chemistry and Physics*, 20, 12 569–12 608, <https://doi.org/10.5194/acp-20-12569-2020>, 2020.
- Köhler, C. G. and Seifert, A.: Identifying sensitivities for cirrus modelling using a two-moment two-mode bulk microphysics scheme, *Tellus
970 B: Chemical and Physical Meteorology*, 67, 24 494, <https://doi.org/10.3402/tellusb.v67.24494>, 2015.
- Laksmo, H., McQueen, T. A., Sellberg, J. A., Loh, N. D., Huang, C., Schlesinger, D., Sierra, R. G., Hampton, C. Y., Nordlund, D., Beye, M., Martin, A. V., Barty, A., Seibert, M. M., Messerschmidt, M., Williams, G. J., Boutet, S., Amann-Winkel, K., Loerting, T., Pettersson,

- L. G. M., Bogan, M. J., and Nilsson, A.: Anomalous Behavior of the Homogeneous Ice Nucleation Rate in "No-Man's Land", *The Journal of Physical Chemistry Letters*, 6, 2826–2832, <https://doi.org/10.1021/acs.jpcllett.5b01164>, 2015.
- 975 Liu, H., Wang, P., and Schlesinger, R.: A numerical study of cirrus clouds. Part I: Model description, *Journal of the Atmospheric Sciences*, 60, 1075–1084, [https://doi.org/10.1175/1520-0469\(2003\)60<1075:ANSOCC>2.0.CO;2](https://doi.org/10.1175/1520-0469(2003)60<1075:ANSOCC>2.0.CO;2), 2003.
- Luebke, A. E., Afchine, A., Costa, A., Grooss, J.-U., Meyer, J., Rolf, C., Spelten, N., Avallone, L. M., Baumgardner, D., and Kraemer, M.: The origin of midlatitude ice clouds and the resulting influence on their microphysical properties, *Atmospheric Chemistry and Physics*, 16, 5793–5809, <https://doi.org/10.5194/acp-16-5793-2016>, 2016.
- 980 McDonald, J.: Use of Electrostatic Analogy in Studies of Ice Crystal Growth, *Zeitschrift für Angewandte Mathematik und Physik*, 14, 610–&, <https://doi.org/10.1007/BF01601268>, 1963.
- Mishima, O. and Stanley, H.: The relationship between liquid, supercooled and glassy water, *Nature*, 396, 329–335, 1998.
- Murphy, D. and Koop, T.: Review of the vapour pressure of ice and supercooled water for atmospheric applications, *Quarterly Journal of the Royal Meteorological Society*, 131, 1539–1565, 2005.
- 985 Nachbar, M., Duft, D., and Leisner, T.: The vapor pressure of liquid and solid water phases at conditions relevant to the atmosphere, *Journal of Chemical Physics*, 151, <https://doi.org/10.1063/1.5100364>, 2019.
- Niedermeier, D., Shaw, R. A., Hartmann, S., Wex, H., Clauss, T., Voigtlaender, J., and Stratmann, F.: Heterogeneous ice nucleation: exploring the transition from stochastic to singular freezing behavior, *Atmospheric Chemistry and Physics*, 11, 8767–8775, <https://doi.org/10.5194/acp-11-8767-2011>, 2011.
- 990 Pathak, H., Späh, A., Esmaeildoost, N., Sellberg, J. A., Kim, K. H., Perakis, F., Amann-Winkel, K., Ladd-Parada, M., Koliyadu, J., Lane, T. J., Yang, C., Lemke, H. T., Oggenfuss, A. R., Johnson, P. J. M., Deng, Y., Zerdane, S., Mankowsky, R., Beaud, P., and Nilsson, A.: Enhancement and maximum in the isobaric specific-heat capacity measurements of deeply supercooled water using ultrafast calorimetry, *Proceedings of the National Academy of Sciences*, 118, e2018379 118, <https://doi.org/10.1073/pnas.2018379118>, 2021.
- Pruppacher, H. R.: A new look at homogeneous ice nucleation in supercooled water drops, *Journal of the Atmospheric Sciences*, 52, 1924–995 1933, 1995.
- Pruppacher, H. R. and Klett, J. D.: *Microphysics of Clouds and Precipitation*, vol. 18 of *Atmospheric and Oceanographic Sciences Library*, Kluwer Academic Publishers, Dordrecht, 2010.
- Ren, C. and Mackenzie, A. R.: Cirrus parametrization and the role of ice nuclei, *Quarterly Journal of the Royal Meteorological Society*, 131, 1585–1605, <https://doi.org/10.1256/qj.04.126>, 2005.
- 1000 Schneider, J., Höhler, K., Wagner, R., Saathoff, H., Schnaiter, M., Schorr, T., Steinke, I., Benz, S., Baumgartner, M., Rolf, C., Krämer, M., Leisner, T., and Möhler, O.: High homogeneous freezing onsets of sulfuric acid aerosol at cirrus temperatures, *Atmospheric Chemistry and Physics*, 21, 14403–14425, <https://doi.org/10.5194/acp-21-14403-2021>, 2021.
- Skrotzki, J., Connolly, P., Schnaiter, M., Saathoff, H., Moehler, O., Wagner, R., Niemand, M., Ebert, V., and Leisner, T.: The accommodation coefficient of water molecules on ice - cirrus cloud studies at the AIDA simulation chamber, *Atmospheric Chemistry and Physics*, 13, 4451–4466, <https://doi.org/10.5194/acp-13-4451-2013>, 2013.
- 1005 Spichtinger, P.: Shallow cirrus convection – a source for ice supersaturation, *Tellus A*, 66, 19937, <https://doi.org/10.3402/tellusa.v66.19937>, 2014.
- Spichtinger, P. and Gierens, K. M.: Modelling of cirrus clouds – Part 1a: Model description and validation, *Atmospheric Chemistry and Physics*, 9, 685–706, <https://doi.org/10.5194/acp-9-685-2009>, 2009.

- 1010 Spreitzer, E. J., Marschalik, M. P., and Spichtinger, P.: Subvisible cirrus clouds – a dynamical system approach, *Nonlinear Processes in Geophysics*, 24, 307–328, <https://doi.org/10.5194/npg-24-307-2017>, 2017.
- Voigt, C., Schumann, U., Minikin, A., Abdelmonem, A., Afchine, A., Borrmann, S., Boettcher, M., Bucuchholz, B., Bugliaro, L., Costa, A., Curtius, J., Dollner, M., Doernbrack, A., Dreiling, V., Ebert, V., Ehrlich, A., Fix, A., Forster, L., Frank, F., Fuetterer, D., Giez, A., Graf, K., Grooss, J.-U., Gross, S., Heimerl, K., Heinold, B., Hueneke, T., Jaervinen, E., Jurkat, T., Kaufmann, S., Kenntner, M., Klingebiel, M.,
- 1015 Klimach, T., Kohl, R., Kraemer, M., Krisna, T. C., Luebke, A., Mayer, B., Mertes, S., Molleker, S., Petzold, A., Pfeilsticker, K., Port, M., Rapp, M., Reutter, P., Rolf, C., Rose, D., Sauer, D., Schaefer, A., Schlage, R., Schnaiter, M., Schneider, J., Spelten, N., Spichtinger, P., Stock, P., Walser, A., Weigel, R., Weinzierl, B., Wendisch, M., Werner, F., Wernli, H., Wirth, M., Zahn, A., Ziereis, H., and Zoger, M.: ML-CIRRUS – The Airborne Experiment on Natural Cirrus and Contrail Cirrus with the High-Altitude Long-Range Research Aircraft HALO, *Bulletin of the American Meteorological Society*, 98, 271–288, <https://doi.org/10.1175/BAMS-D-15-00213.1>, 2017.
- 1020 Wernli, H., Boettcher, M., Joos, H., Miltenberger, A. K., and Spichtinger, P.: A trajectory-based classification of ERA-Interim ice clouds in the region of the North Atlantic storm track, *Geophysical Research Letters*, 43, 6657–6664, <https://doi.org/10.1002/2016GL068922>, 2016.
- Zhang, Y., Macke, A., and Albers, F.: Effect of crystal size spectrum and crystal shape on stratiform cirrus radiative forcing, *Atmospheric Research*, 52, 59–75, [https://doi.org/10.1016/S0169-8095\(99\)00026-5](https://doi.org/10.1016/S0169-8095(99)00026-5), 1999.