Inferring and evaluating satellite-based constraints on NOx emissions estimates in air quality simulations

James D. East^{1,2}, Barron H. Henderson³, Sergey L. Napelenok³, Shannon N. Koplitz³, Golam Sarwar³, Robert Gilliam³, Allen Lenzen⁴, Daniel Q. Tong⁵, R. Bradley Pierce⁴, Fernando Garcia-Menendez¹

¹Department of Civil, Construction, and Environmental Engineering, North Carolina State University, Raleigh, NC 27695, USA
 ²Oak Ridge Institute for Science and Education, Office of Research and Development, U.S. Environmental Protection Agency, Research Triangle Park, NC 27711, USA

³U.S. Environmental Protection Agency, Research Triangle Park, NC 27711, USA

⁴Space Science and Engineering Center, University of Wisconsin-Madison, Madison, WI 53706, USA ⁵Department of Atmospheric, Oceanic and Earth Sciences, George Mason University, Fairfax, VA 22030 USA

Correspondence to: Fernando Garcia-Menendez (f garcia@ncsu.edu)

5

Abstract. Satellite observations of tropospheric NO₂ columns can provide top-down observational constraints on emissions estimates of nitrogen oxides (NO_x). Mass-balance_based methods are often applied for this purpose, but do not isolate near-surface emissions from those aloft, such as lightning emissions. Here, we introduce an inverse modeling framework that couples satellite chemical data assimilation to a chemical transport model. and In the framework, infers-satellite-constrained emissions totals are inferred using model simulations with and without data assimilation using in the iterative finite-difference mass-balance method. The approach improves the finite-difference mass-balance inversion by isolating the near-surface emissions increment. We apply the framework to separately estimate lightning and anthropogenic NO_x emissions over the Northern Hemisphere for 2019. Using overlapping observations from the Ozone Monitoring Instrument (OMI) and the

- Tropospheric Monitoring Instrument (TROPOMI), we compare <u>separate</u> NO_x emissions inferences from these satellite instruments, as well as the impacts of emissions changes on modeled NO_2 and O_3 . OMI inferences of anthropogenic emissions consistently lead to larger emissions than TROPOMI inferences, attributed to a low bias in TROPOMI NO_2 retrievals. Updated lightning NO_x emissions from either satellite improve the chemical transport model's low tropospheric O_3 bias. The combined
- 25 lighting and anthropogenic emissions updates improve the model's ability to reproduce measured ozone by adjusting natural, long-range, and local pollution contributions. Thus, the framework informs and supports the design of domestic and international control strategies. Combined lightning and anthropogenic updates inferred from satellite observations can improve the model's ability to represent background and ground level O₃ concentrations, an ongoing policy consideration in the U.S. as domestic and international emissions control strategies evolve.

1 Introduction 30

Tropospheric nitrogen oxides (NO_x), nitric oxide (NO) and nitrogen dioxide (NO_2), harm human health (Anenberg et al., 2018; Murray et al., 2020) and play a key role in the formation of important secondary atmospheric pollutants, such as O_3 (Jacob, 2000). NO_x is emitted to the troposphere primarily by anthropogenic combustion processes, but natural sources, including soil, lightning, and wildfires, also contribute to the atmospheric NO_x budget (Jacob, 1999). Accurate NO_x emissions

- 35 are a critical component of local- to global-scale atmospheric chemistry simulations. On hemispheric scales, realistically representing the formation and intercontinental transport of O₃ with models requires adequate international global emissions inventories (Itahashi et al., 2020; Zhang et al., 2016; Zhang et al., 2008; Verstraeten et al., 2015; Mathur et al., 2017). In regional air quality simulations, which commonly rely on hemispheric or global models for chemical boundary conditions, the relative contribution of long-range pollutant transport to ground-level O₃ concentrations has grown in many areas as O₃ 40 precursor emissions have decreased in the U.S. and other developed high income countries (HICs) (McDuffie et al., 2020;
- Jaffe et al., 2018; Simon et al., 2015). As a result, air quality management policies, often informed by regional modeling, are strengthened by accurate and up-to-date global NO_x emissions inventories. However, compilation of bottom-up regional and global emissions inventories, developed from source- and location-specific emissions factors and activity data, is time- and labor-intensive, and can be hindered by limited data. As a result, bottom-up inventories lag emissions changes and are often
- incomplete. Although uUncertainties in bottom-up emissions estimates are particularly large for developing-lower-middle 45 income countries (McDuffie et al., 2020; Elguindi et al., 2020), they and remain significant for developed countriesHICs as well (Day et al., 2019).

Satellite observations of NO₂ can bridge temporal gaps in emissions estimates (Tong et al., 2016; Tong et al., 2015) and constrain uncertainty in emissions inventories through inverse modeling (e.g. Lamsal et al., 2011; Goldberg et al., 2021; de

- 50 Foy and Schauer, 2022). Several methods have been applied to develop top-down emissions estimates using satellite observations and atmospheric models, each carrying advantages and limitations (Elguindi et al., 2020). Adjoint-based methods can provide detailed precise emissions updates, but require significant computational resources (e.g. Qu et al., 2017; Qu et al., 2019; Muller and Stavrakou, 2005; Kurokawa et al., 2009; Cooper et al., 2017; Zhang et al., 2019; Wang et al., 2020b). Similarly, Kalman filtering and related approaches have been used but are computationally-intensive (e.g. Napelenok et al., 55 2008; Ding et al., 2020; Ding et al., 2015; Mijling and Van Der A, 2012; Miyazaki and Eskes, 2013; Miyazaki et al., 2017; Miyazaki et al., 2012a; Miyazaki et al., 2012b; Sekiya et al., 2021). Mass balance inversion approaches, which scale model emissions by directly comparing model estimates and satellite observations, were introduced by Martin et al. (2003), updated by Lamsal et al. (2011), and have been widely used in research and forecasting (e.g. Boersma et al., 2015; Itahashi et al., 2019; Li et al., 2018; Visser et al., 2019; Zhu et al., 2021; Cooper et al., 2017). Although lower computational costs allow the finitedifference mass-balance (FDMB) approach (Lamsal et al., 2011) to readily update emissions, the method is subject to an
- 60

emissions smearing effect (e.g. Cooper et al., 2017), which can cause emissions updates to be spatially misallocated. Since FDMB uses satellite observations directly, near-surface NO₂ bias cannot be isolated from biases in the middle and upper

troposphere, which obscures the surface emissions inference. Further, applications often rely on a single inversion from a single satellite, although available satellite products have been shown to have significant biases. For example, early versions

65 <u>of</u> the Tropospheric Monitoring Instrument (TROPOMI) NO₂ product showed a low bias in urban areas when compared against surface-basedground and airborne spectrometer measurements (Judd et al., 2020; Verhoelst et al., 2021) and the Ozone Monitoring Instrument (OMI) NO₂ product has been reported to differ with measurements by ±20% (Lamsal et al., 2014). The impact of biases in satellite-based NO₂ data on mass-balance inversions has not been fully explored despite the wide used of the method to scale NO_x emissions. Minimizing bias in anthropogenic emissions inferences and understanding the potential for them to propagate to emissions updates are needed to improve mass-balance-based inversions.

Here, we introduce a modeling framework that couples satellite chemical data assimilation to the Community Multiscale Air Quality model (CMAQ) and applies an iterative FDMB inversion to estimate NO_x emissions in the Northern Hemisphere. The framework provides observational constraints to improve emissions estimates in areas where emissions are highly uncertain, at a lower computational cost relative to adjoint- and Kalman-filter-based approaches. We apply the framework in an iterative assimilation to infer 2019 NO_x emissions, the first complete year in which OMI and TROPOMI records overlap. In contrast to traditional FDMB, which directly compares modeled and observed columns, our framework improves the FDMB

- method by first assimilating satellite-retrieved NO₂, and then performing the inversion by comparing model simulations with and without assimilation. In the assimilation step, updates to model concentrations are vertically allocated to model layers. As a result, Aassimilating the observed column allows the inversion framework to use only the near-surface portion of the model
- 80 <u>column in the FDMB inversion to target near-surface NO₂-, and minimizinge</u> influences from the upper troposphere <u>and</u>, extending the framework proposed by Lamsal et al. (2011). In addition, our analysis compares <u>independent</u> inversions using which separately use OMI and or TROPOMI NO₂ data. We show that the inverse emissions produced by this framework influence representation of intercontinental O₃ transport to the U.S., offering an opportunity to improve chemical boundary conditions in policy-relevant regional-scale air quality simulations.

85 2 Methods

We develop a framework to update NO_x emissions estimates using the CMAQ chemical transport air quality model (Byun and Schere, 2006), 3D-variational (3DVAR) chemical data assimilation (Sandu and Chai, 2011), and space-based NO₂ observations. We apply the framework to estimate 2019 lightning and anthropogenic NO_x emissions, and compare surfaceground- and space-based NO₂ observations to model simulations using the prior emission (inventory before the framework is applied) and posterior emissions (inventory after the framework is applied) to assess the impact of the updates. Figure 1 provides an overview of the framework, in which lightning NO_x (LNO_x) emissions and anthropogenic NO_x (ANO_x) emissions are updated separately.

75



Figure 1: NO_x emissions inversion framework. Lightning NOx (LNO_x) emissions are updated in the first step. Then, anthropogenic NO_x
 (ANO_x) emissions are updated iteratively. CMAQ boxes represent air quality simulations without chemical data assimilation, and CMAQ+3DVAR boxes represents air quality model simulations with chemical data assimilation. Satellite NO₂ retrievals and a background error covariance (BECOV) are inputs to the chemical data assimilation, described in Sect. 2.3. Red dotted lines around the inversion boxes represent the boundary of the inversion algorithmcorrespond to the red dotted lines in Fig. 3, which is details the inversion algorithm.ed in Fig. 3. Dashed black emissions input lines around the ANO_x inversion represent the iterative process. Iteration and convergence criteria are described in Sect. 2.6.

2.1 Satellite data

We use NO₂ tropospheric column observations from the National Aeronautics and Space Administration's (NASA's) OMI and the Royal Netherlands Meteorological Institute's (KNMI's) TROPOMI instruments in the inversion framework. TROPOMI was launched in October 2017 and provides 7.2×3.6 km² resolution NO₂ retrievals, upgraded to 5.6×3.6 km² resolution in August 2019 (Van Geffen et al., 2020; Veefkind et al., 2012). TROPOMI's sun-synchronous polar orbit crosses the equator at approximately 1:30 pm local time, allowing the instrument to achieve global coverage in one day. We assimilate the Level-2 tropospheric slant column retrieved from NASA's Earth Science Data Systems Program (<u>https://earthdata.nasa.gov/</u>). The data product is described in the Algorithm and Theoretical Basis Document (ATBD) for TROPOMI NO₂ (Van Geffen et al., 2019). We only consider TROPOMI observations with a quality flag greater than 0.5 and

- 110 cloud fraction lower than 30% in the assimilation, following data product recommendations (Eskes et al., 2019). We use the latest publicly available versions of the TROPOMI retrieval for 2019 (versions 1.2.2 to 1.3.2) at the time of the analysis. Version 1.3 introduced updates to cloud processing that decrease noisy hotspots and broadened the range of acceptable air mass factors (Eskes et al., 2021). Information about the updates applied in each version and the dates on which updates were applied is given in Eskes et al. (2021). A research version with an updated retrieval applied to 2019 observations has been
- 115 developed (Van Geffen et al., 2021), but was not yet standard and was not available at the time of this analysis. We discuss the impact of these latest updates in Sect. 3.3.

OMI, onboard the Aura satellite launched in 2004, provides tropospheric NO_2 vertical and slant column retrievals with a resolution of 13×24 km² near nadir in a sun-synchronous polar orbit with a local equator crossing time of 1:45 pm. Global coverage is achieved in two days. We use the NASA Goddard Space Flight Center (GSFC) Level-2 NO_2 product (Krotkov et al. 2019b). OMI was impacted by a row anomaly beginning in 2008, reducing the number of usable pixels in the OMI retrieval

120

125

al., 2019b). OMI was impacted by a row anomaly beginning in 2008, reducing the number of usable pixels in the OMI retrieval (Boersma et al., 2018). We include only pixels with cloud fraction lower than 30% and a summary quality flag of 0. Detailed information about the NO₂ data product is included in the OMI ATBD (Chance, 2002) and in Krotkov et al. (2019a).

A low bias has been noted in the versions of TROPOMI NO₂ used for this study (Judd et al., 2020; Verhoelst et al., 2021). Although TROPOMI NO₂ in 2019 has been reprocessed with retrieval version 2.3.1, resulting in an improvement of the bias (Eskes et al., 2021), these reprocessed datasets were not yet available at the time this analysis was conducted. Figure 2 compares TROPOMI and OMI tropospheric vertical column density (VCD) for 2019, regridded to the CMAQ grid used. For the VCDs shown in the figure, we remove the effect of the assumed vertical profile of NO₂ from the original satellite product by recalculating the VCDs with the NO₂ vertical profile simulated by CMAQ. In the Rresults, we discuss the low bias in TROPOMI data and explore its impact on emissions inversions.

130



Figure 2: 2019 annual average TROPOMI and OMI vertical NO₂ vertical column densities, with CMAQ NO₂ profiles applied, and the ratio between them. Column densities ratios are only shown for the grid cells where NO_x emissions updates are applied in the emissions inversion.

135 2.2 Hemispheric air quality modeling

Model simulations in the inversion framework were completed for January–December 2019 using CMAQ v5.3.2 (Appel et al., 2021; U.S. EPA, 2020). CMAQ has been used to simulated air quality over the Northern Hemisphere and shown to

adequately capture chemical composition against observations (Mathur et al., 2017). Model inputs and satellite observations are summarized in Table 1. Simulations, designed to capture continental-scale pollutant transport, cover the Northern

- 140 Hemisphere with 108 km horizontal grid spacing and a 44-layer vertical structure reaching 50 hPa (Mathur et al., 2017). The simulations use version CB6r3 of the Carbon Bond 6 chemical mechanism (Luecken et al., 2019), the AERO7 aerosol module (Xu et al., 2018), and updated halogen chemistry (Kang et al., 2021). Anthropogenic emissions are modeled using representative day-of-week emissions that change month-to-month. Representative-day emissions are created by averaging data from the prior emissions inventory on a day-of-week basis by month. For each day of the week and each month, there is
- 145 <u>a unique hourly emissions file that is used for every matching day of the week in that month. As a result, diurnal and weekly patterns are captured in the emissions, while daily variations that are specific to the prior emissions inventory year are averaged. The prior emissions inventory, relies on the best available emissions data at the time of the study. Anthropogenic emissions for North America are from the U.S. Environmental Protection Agency's (EPA) 2017 National Emissions Inventory (NEI) modeling platform (Adams, 2020). Emissions in China are for the year 2015 (Zhao et al., 2018) and emissions for the rest of</u>
- 150 the hemisphere are based on the Hemispheric Transport of Air Pollution (HTAP) version 2, projected from their original 2010 date to 2014 with scaling factors from the Community Emissions Data System (CEDS). To initialize the 2019 prior and posterior simulations and reduce the impact of chemical initial conditions on the results, we use a 1-year spin-up period not considered for the analyses. CMAQ model runs are driven by meteorology from a retrospective hemispheric simulation using the Weather Research and Forecasting (WRF) model (Skamarock et al., 2008) version 4.1.1 configured following Mathur et al. 2008) version 4.1.1

155 al. (2017) and Xing et al. (2015).

Table 1: Prior emissions and model inputs

Table 1. The emissions and model inputs					
Data	Year	Source			
Prior emissions (North America)	2017 EPA platform (v7.1)	(Adams, 2020)			
Prior emissions (China)	2015 Tsinghua University	(Zhao et al., 2018)			
Prior emissions (Rest of hemisphere)	HTAPv2 (2010) projected to 2014 using CEDS scaling factors	(Janssens-Maenhout et al., 2015; Hoesly et al., 2018)			
Prior emissions (LNO _x)	2017 GEIA*	(Price et al., 1997)			
Biomass burning emissions	2019 FINN*	(Wiedinmyer et al., 2011)			
Soil NO _x emissions	2018 CAMS* v2.1 with canopy reduction factor	(Granier et al., 2019)			
Biogenic emissions	2018 MEGAN*	(Guenther et al., 2006)			
Meteorology	2019 WRF v4.1.1	(Powers et al., 2017)			
Satellite observation year	2019 NO ₂ retrievals from OMI and TROPOMI				

* GEIA = Global Emissions Initiative; FINN = Fire Inventory from NCAR; CAMS = Community Atmosphere Modeling System; MEGAN = Model of Emissions of Gases and Aerosols from Nature

2.3 Chemical data assimilation in CMAQ

160

We adjust modeled NO₂ concentrations using satellite observations by coupling the CMAQ model to a data assimilation model, the National Centers for Environmental Prediction (NCEP) Gridpoint Statistical Interpolation (GSI) program version 3.3 (Shao et al., 2016). GSI performs three-dimensional variational (3DVAR) data assimilation by minimizing the cost function, *J*:

$$J = \frac{1}{2} [x^T \mathbf{B}^{-1} x + (H(x) - y)^T \mathbf{R}^{-1} (H(x) - y)], \qquad (1)$$

where y is the observation innovation y = y_o - H(x_b), x is the analysis increment x = x_a - x_b, x_a is the analysis field (NO₂ concentration after application of chemical data assimilation), x_b is the model background (the simulated NO₂ concentration before application of chemical data assimilation), y_o is the satellite observations, **B** is the background error covariance matrix, **R** is the observation error matrix, and *H* is the observation operator. To compute the difference between the model column (x_b) and the satellite column (y_o), the observation operator *H* is applied, which transforms the model background to the form of the satellite observations. For TROPOMI data, the averaging kernel is first converted to scattering weights as

$$w(z) = \mathbf{A}(z) \times \mathbf{M}_{total} , \qquad (2)$$

where $\mathbf{A}(z)$ is the vertically-resolved TROPOMI averaging kernel for level z, \mathbf{M}_{total} is the air mass factor provided with the satellite data, and w(z) are the vertically-resolved scattering weights. Scattering weights accompany the OMI NO₂ data

175 product, so this step is not needed to assimilate OMI data. Scattering weights are then applied to compute the model slant column as

$$\Omega_s^m = \sum_z \Omega_v^m(z) w(z), \ z \le z_{tropopause}, \tag{3}$$

where $\Omega_v^m(z)$ is the model partial vertical column in the troposphere, interpolated to the satellite grid, and Ω_s^m is the model 180 tropospheric slant column density (SCD). The difference between the modeled and observed slant columns, or the observation innovation y in Equation 1, is estimated as

$$\Omega_s' = \Omega_v^o \mathbf{M}_{\rm trop} - \Omega_s^m,\tag{4}$$

where Ω'_s is the analysis increment, Ω^o_v is the satellite tropospheric VCD, and \mathbf{M}_{trop} is the tropospheric air mass factor, 185 distributed with the satellite data. We eliminate the influence of the *a priori* satellite vertical profile by computing the analysis

increment with the model and observed SCD, which, unlike the VCD, does not rely on the *a priori* vertical NO₂ profile assumed by the satellite.

190

We compute **B** using the Generalized background error covariance matrix model (GENBE v2.0) (Descombes et al., 2015), which models background errors by comparing a free-running simulation and a simulation with either lightning or anthropogenic NO_x emissions perturbed. We use GENBE with the prior simulation and a simulation with a uniform -15% perturbation to LNO_x to create 3-dimensional background errors in the upper troposphere for the LNO_x assimilation. After updating LNO_x emissions (as described in Sect. 2.5), we create 3-dimensional background errors in the boundary layer for the anthropogenic NO_x assimilation by using GENBE with the LNO_x posterior simulation and a simulation with a -15% perturbation to surface anthropogenic NO_x emissions. Observation error **R** is provided with the satellite data.

195 Online coupling between GSI and CMAQ was developed in this study to perform the assimilation. At each model timestep in which a satellite observation is available, the CMAQ model simulation is paused and 3DVAR assimilation is performed. The CMAQ model state at that time step is used as x_b . After assimilation using 3DVAR within GSI, CMAQ returns to a freerunning mode and the new model state, x, is updated to more closely match the satellite observation. The difference in the monthly average NO₂ VCDs from the assimilation and no-assimilation runs is used in the inversion as $\Delta\Omega$.

200

2.4 Finite difference mass balance inversion

In the inversion framework developed, we iterate the approach of Lamsal et al. (2011). The FDMB process as applied here is summarized in Fig. 3. In the past, this approach has been used by directly comparing model and satellite columns (e.g. Itahashi et al., 2019; Cooper et al., 2017; Lamsal et al., 2011). We update modify the approach by first updating model concentrations with assimilationing the of satellite observations, and then updating the emissions using <u>AQ</u> the difference between the modeled VCD with and without assimilated satellite information. All updates are performed on a monthly average basis.

205



Figure 3: FDMB inversion. The red dashed line corresponds to the red dashed lines in Fig. 1, and the processes inside show additional details of the FDMB inversion. In this framework, the prior emissions (black box on the far left) are input to the CMAQ model. CMAQ simulations are performed with unperturbed prior emissions (100% arrow and E_{100}) and prior emissions with a -15% perturbation (85% arrow and E_{85}). The resulting modeled VCDs are Ω_{100} and Ω_{85} , respectively. These VCDs are used to compute the sensitivity, β (blue box). New emissions totals are calculated with FDMB (yellow box), using β , NO₂ VCD from a CMAQ simulation without assimilation (Ω), and NO₂ VCD from a CMAQ simulation with assimilation (Ω'). When iteration is used, the posterior emissions from the previous iteration are used as input to the CMAQ model to simulate new VCDs, Ω and Ω' .

In FDMB, following Lamsal et al. (2011), emissions changes are inferred through the relationship

$$\frac{\Delta E}{E} = \beta \frac{\Delta \Omega}{\Omega} \tag{5}$$

220 where ΔE is the inferred NO_x emissions change, *E* is the NO_x emissions prior, Ω is the model simulated NO₂ VCD without chemical data assimiliation, and $\Delta \Omega = \Omega_{assim} - \Omega$, the monthly average difference between the model simulated tropospheric NO₂ VCD with (Ω_{assim}) and without (Ω) chemical data assimilation. β is a unitless scaling parameter, the Jacobian, that linearly relates NO₂ VCD changes to NO_x emissions changes. β is calculated through finite differencing as

$$\beta = \frac{E'-E}{E} \frac{\Omega_{\rm E}}{\Omega_{E'} - \Omega_{\rm E}} \tag{6}$$

225

where E' is perturbed NOx emissions, Ω_E is the tropospheric NO₂ VCD simulated with model emissions E, and $\Omega_{E'}$ is the tropospheric NO₂ VCD simulated with model emissions E'. To estimate β , we use the same -15% perturbation used to create

background errors **B** in the boundary layer. Cooper et al. (2017) found that using perturbations ranging from 5% to 20% to calculate β changed posterior emissions estimates by less than 2% globally.

230

2.5 Inverse modeling NO_x emissions

In our framework, LNO_x emissions are updated first, separately from anthropogenic emissions. Due to the satellite instruments' sensitivity to NO₂ in the upper atmosphere (e.g. Eskes and Boersma, 2003), small model biases there can influence the total column comparison and adversely impact the anthropogenic emissions adjustment. By updating LNO_x emissions, we aim to decrease this bias and its impact on the ANO_x inversion. We compute the scaling parameter for lightning emissions, β_{LNOx} , using the -15% LNO_x perturbation simulations applied to create background errors for the upper troposphere. We then assimilate satellite NO₂ observations using the background errors for the upper troposphere and apply β_{LNOx} in a single inversion iteration using the full tropospheric VCD to compute spatially-varying LNO_x adjustment factors. Updates to LNO_x are calculated using monthly averages.

240

235

After LNO_x emissions are updated, ANO_x emissions are updated by iteratively applying a FDMB inversion independently for each month in 2019. Iterating the FDMB has been shown to improve emissions estimates compared to a single FDMB application (Cooper et al., 2017). In the FDMB iteration, each update to the emissions serves as the prior emissions for the subsequent iteration (represented as black dashed lines in Fig. 1). The number of iterations is determined based on the synthetic observation experiment described in Sect. 2.6. β is held constant during all ANO_x inversion iterations, and not recalculated

- each time, to prevent instability in β as changes in the column become smaller with subsequent iterations. In the ANO_x emissions inversion, we only consider grid cells in which local anthropogenic NO_x emissions likely significantly contribute to the satellite-observed NO₂ column by only including grid cells in which anthropogenic NO_x emissions comprise at least 50% of total NO_x emissions following Lamsal et al. (2011), population density is greater than 15,000 people km⁻² (CIESIN, 2018), modeled cloud cover is less than 30%, and the local time is 1:00 or 2:00 pm (OMI and TROPOMI overpass times). Only emissions in grid cells meeting these criteria are adjusted. Table S1 describes each simulation performed for the LNO_x and
 - ANO_x inversions.

The FDMB method assumes emissions impacts are local (i.e., emissions in one grid cell do not affect VCD amounts in neighboring grid cells). This assumption is most valid when NO_x lifetime to chemical losses is shorter than NO_x transport time to neighboring grid cells, which is typical near the surface in coarse resolution models (Martin et al., 2003), such as the one used in this study. However, the assumption is less realistic at finer resolutions and in the upper troposphere, where the lifetime of NO₂ is longer than at the surface and NO₂ concentrations are not directly impacted by coincident near-surface emissions. Even at coarse resolution (e.g., 100-km grid spacing), emissions smearing effects, which occur when the FDMB assumption of local emissions effects is incorrect and emissions are inappropriately adjusted, can appear due to NO_x transport, reservoir species, and chemical feedbacks (Turner et al., 2012; Cooper et al., 2017). Traditional FDMB, which directly compares

260 modeled and remotely sensed columns, cannot address this effect. Assimilating the satellite VCD introduces an additional

complication. The horizontal length scales (on the order of several hundreds of kilometers) used in the background error extend beyond the grid cell horizontal dimensions (nominally 108km) in the middle and upper troposphere and, as a result, NO₂ changes introduced by assimilation ($\Delta\Omega$) do not have a local relationship with surface emissions directly below. In our work, assimilating the observed column information, instead of directly comparing modeled and satellite retrieved VCDs, allows the

265

275

analysis to be restricted to the lower troposphere, mitigating both the misallocation errors of FDMB and the effect of horizontal length scales extending beyond the grid cell dimension. To that end, we limit the anthropogenic emissions analysis to the lowest 20 model layers, which is nominally from the ground to ~720 hPa over non-mountainous terrain in the summer, and use that partial column to calculate $\Delta\Omega$ in the FDMB inversion. $\Delta\Omega$ for a single month above and below the threshold is illustrated in Fig. S1. By applying this cutoff, we focus the inversion on surface anthropogenic NO_x.

270 2.6 Inversion system testing

We conduct a synthetic observation experiment to evaluate the ability of the inversion system to constrain emissions to a known perturbation. Artificial NO₂ observations were generated from CMAQ simulations with unperturbed emissions and NO_x emissions reduced by 15%. As expected, assimilating the synthetic observations derived from a simulation with unperturbed emissions results in an analysis increment of zero. The results of an iterative emissions inversion based on the synthetic observations derived from the simulation with perturbed emissions are shown in Fig. S2. Across Northern Hemisphere regions, the normalized mean error (NME) relative to the known perturbed emissions, and the rate at which it changes, decrease with subsequent iterations. The NME is minimized after 7–9 iterations, depending on the region. In all subsequent results, emissions inferences made with 8 iterations of the inversion system are shown and analyzed. Convergence of the inversion in different global regions adds confidence to the system's ability to constrain real-world emissions.

280 **3. Results**

3.1 Lightning NO_x emissions updates

Assimilation of retrievals from either satellite increases LNO_x emissions across all seasons, relative to the prior emissions (monthly climatology from GEIA), with largest changes occurring during the summer (Figs. S3 and S4). Applying 2019 OMI data increases total LNO_x emissions in 2019 by 20% over the GEIA climatology, while assimilation of TROPOMI data increases LNO_x emissions by 24%. The emissions increases inferred by both satellite products are driven by NO₂ increases in the mid and upper troposphere due to assimilation, with changes near the surface negligible in comparison. Increases in background areas with small NO₂ column totals and subsequent LNO_x increases in these areas suggest a low bias in modeled background NO₂ relative to observations from both satellites. A low bias agrees with the findings reported by other model-satellite NO₂ comparisons (Silvern et al., 2019; Qu et al., 2021; Goldberg et al., 2017). The LNO_x emissions adjustments inferred here decrease the differences between modeled and satellite-derived NO₂ in the upper troposphere and decrease the bias that differences in the upper troposphere can introduce to the subsequent ANO_x inversion.

3.2 Impact of assimilation on modeled NO₂ vertical column density

Figure 4 shows the change to CMAQ-modeled tropospheric VCD ($\Delta\Omega$) caused by assimilating NO₂ observations from OMI or TROPOMI with background errors for the boundary layer, before applying any emissions adjustments. In Fig. 4, and

- 295 throughout the results, $\Delta\Omega$ reflects differences near the surface (as described in Sect. 2.5). Assimilating OMI NO₂ data generally increases modeled NO₂ columns near populated areas in China, India, and the U.S. In contrast, assimilating TROPOMI NO₂ data decreases modeled NO₂ columns more widely across the Northern Hemisphere. The changes brought about by assimilating satellite data are larger during the winter and fall, and smaller in the spring and summer, when NO_x lifetime is shortest and NO₂ columns are smaller. During the winter in northeast China, where the assimilation impacts are
- 300 most apparent, the seasonal average change due to assimilation reaches 1.8×10^{15} molecules cm⁻² for OMI and -2.8×10^{15} molecules cm⁻² for TROPOMI. The direction of $\Delta\Omega$ after assimilation of OMI data is more heterogeneous and shows a stronger seasonality, while $\Delta\Omega$ based on assimilating TROPOMI data is consistently negative. Over Europe, $\Delta\Omega$ after assimilating OMI observations is close to zero in warm months and negative in colder seasons. Assimilating satellite-observed NO₂ increases the NO₂ levels modeled over the ocean and less-populous areas, such as the Sahara, with low NO_x emissions and small NO₂
- 305 column amounts.



Figure 4: Seasonal NO₂ VCD change ($\Delta\Omega$) from CMAQ simulation using prior emissions after assimilating OMI or TROPOMI tropospheric NO₂ observations and modeling atmospheric composition with prior NO_x emissions. $\Delta\Omega$ shown for winter (DJF), spring (MAM), summer (JJA), and fall (SON).

Over polluted areas, the direction of $\Delta\Omega$ for the TROPOMI or OMI data assimilations tends to differ. This discrepancy is likely due to the low bias in TROPOMI-derived tropospheric NO₂ columns, which has been reported to be approximately 10% over the U.S., Europe, and India, and greater than 20% over China when compared with the OMI Quality Assurance for

- 315 Essential Climate Variables (QA4ECV) retrieval (Van Geffen et al., 2021; Verhoelst et al., 2021; Wang et al., 2020a; Li et al., 2021). Over background areas, the analysis increments that results from assimilation of observations from both satellites generally agree. The consistency suggests a low bias in modeled background NO₂ concentrations and also agrees with the low bias in CMAQ-modeled free tropospheric NO₂ reported by Goldberg et al. (2017). Such a bias can contribute to the positive analysis increment over background areas. However, NO₂ columns observed in these regions may be smaller than the retrieval accuracy of 0.7 × 10¹⁵ molecules cm⁻² (Van Geffen et al., 2019), reducing confidence in the analysis increment at these locations. In the anthropogenic emissions inversion, our filtering criteria exclude from the analysis background areas which
- locations. In the anthropogenic emissions i are more likely to have low VCD amounts.

3.3 Emissions inversion

- 325 Season-average β values, relating NO₂ vertical column differences to anthropogenic near-surface NO_x emissions updates, are shown in Fig. S5. Based on our criteria for gridcell inclusion in the inversion, described in Sect. 2.5, we consider 13% of the grid cells in the domain, which represent 88% of prior anthropogenic NO_x emissions. Seasonal domain-average values range from 1.33 to 1.66, and are lower in the winter and higher in the summer. A β value less than 1.0 results in an emissions update that is smaller than the VCD change, while a β greater than 1.0 has the opposite effect. β tends to be less than 1.0 in polluted regions during colder months and larger during warmer months and in less polluted regions, although many grid cells which are less polluted are not considered in the analysis. The scaling factors are smallest over China, and larger over the U.S., India, Mexico, and Europe. The differences among regions stem from local differences in NO_x lifetime and transport. In Indonesia and sub-Saharan Africa, lower emissions and a small response from tropospheric VCD to anthropogenic emissions
- 335 10, following Cooper et al. (2017). Scaling factors estimated here are larger than the 1.16 global-average previously reported by Lamsal et al. (2011). However, in Lamsal et al. (2011) modeled NO₂ vertical columns were sampled at the morning SCanning Imaging Absorption spectroMeter for Atmospheric CHartographY (SCIAMACHY) overpass time, rather than the afternoon OMI or TROPOMI overpass times, and β tends to be closer to 1.0 during the morning in regions with high NO_x emissions (Li and Wang, 2019). Li and Wang (2019) show that over rural regions with lower NO_x concentrations, β is larger

perturbations can lead to large β values. To prevent overly large or small β values, we constrain the factor to between 0.1 and

340 at the OMI or TROPOMI overpass window than at the SCIAMACHY overpass window, suggesting a larger overall β for

analyses based on OMI or TROPOMI products should be expected. Additionally, NO_x emissions have decreased considerably in several regions of the Northern Hemisphere after the Lamsal study was conducted (2011), including the U.S. (Tong et al., 2015) and China (Miyazaki et al., 2017), which has changed the sensitivity of NO_2 VCDs to NO_x emissions (Qu et al., 2021; Silvern et al., 2019).

345

Annual bottom-up prior ANO_x emissions estimates are shown in Fig. 5. Season-average ANO_x emissions inferences from the inversions based on OMI and TROPOMI observations are shown in Fig. 6. The use of OMI observations generally tends to increase emissions in most industrialized nations outside of Europe. NO_x emissions increases driven by OMI observations are largest in winter and spring and smaller, or slight decreases, in summer and fall. In contrast, the use of TROPOMI retrievals tends to drive a decrease in NO_x emissions across all seasons and continents, with the largest impacts in the summer and

350 smallest in the spring. The largest emissions changes based on both OMI and TROPOMI retrievals are in Northeast China during the winter. Over India, OMI inferred changes are concentrated in Central India where prior emissions are lower, while the largest changes inferred from TROPOMI are in the Northern, Eastern, and Southern zones, where prior emissions are highest. Relative NO_x emissions changes driven by TROPOMI observations tend to be small over dense urban areas, with more uniform decreases over cells with lower emissions.



355

Figure 5: 2019 prior anthropogenic NO_x emissions totals. Data sources are described in Table 1.



Figure 6: Season-average NO_x emissions changes from inverse modeling updates based on OMI and TROPOMI observations. Emissions changes are shown for winter (DJF), spring (MAM), summer (JJA), and fall (SON).

ANOx emissions totals and inferred changes are explored for China, India, Europe, Mexico, and the U.S. (Fig. 7). We also show 2019 NO_x emissions totals from the Copernicus Atmosphere Monitoring Service (CAMS) bottom-up emissions inventory (Granier et al., 2019), and the NASA Tropospheric Chemical Reanalysis products 2 (TCR-2) satellite-inferred
inventory (Miyazaki et al., 2019, 2020). TCR-2 top-down NOx emissions are constrained using satellite observations of NO₂, CO, O₃, and SO₂ at a resolution of 1.125°× 1.125° and are further described in (Miyazaki et al., 2017). CAMS anthropogenic NO_x emissions are based on the Emissions Database for Global Atmospheric Research (EDGAR version 5.3) estimates for 2015 (Crippa et al., 2020) projected to 2019 using CEDS scaling factors, and are provided at 0.1°× 0.1°. Both datasets provide monthly anthropogenic NO_x totals. Except for Europe, assimilation toward OMI retrievals increases annual emissions totals
in the regions analyzed, while using TROPOMI retrievals decreases them. TCR-2 NO_x emission estimates are larger than the prior emissions used by our inverse modeling framework, except for India, while CAMS totals are lower than the prior emission estimates and similar to TROPOMI inferred emissions. Across the regions considered, TROPOMI infers an average annual decrease to NO_x emissions from the regions of -33%, while OMI infers a +9% increase. In Europe, the only region where the sign of the inferred changes match, use of OMI retrievals results in a -1% change, while applying TROPOMI observations

- 375 leads to a -36% decrease in NO_x emissions. The largest total changes are inferred in the highest emitting region, China, while the greatest relative changes, -41% inferred with TROPOMI, are for India, where emissions are highly uncertain. Changes inferred with OMI observations over the U.S. are greater than $1,200 \times 10^3$ short tons NO_x as NO₂ per year, but smaller than the difference between our prior U.S. emissions estimates and TCR-2 or CAMS estimates. A change of -3,000 ×10³ short tons NO_x as NO₂ emitted annually in the U.S. as inferred by TROPOMI, over 30% of the prior emissions, differs significantly from
- 380 National Emissions Inventory estimates but leads to a total close to that of the 2019 CAMS inventory.



Figure 7: Prior and satellite-inferred 2019 anthropogenic NOx emissions in select global regions. Top plot shows total emissions (as NO₂) from prior emissions estimates, inference with OMI or TROPOMI observations (OMI and TROPOMI posterior), and CAMS or TCR-2 inventories in the U.S., China, India, Mexico, and Europe. The bottom plot shows the percent change ($\Delta E NO_x$) inferred with OMI or TROPOMI data, relative to prior emission estimates, for each region.

Across the months simulated, inferences using OMI retrievals consistently lead to higher NO_x emissions than using TROPOMI retrievals. Figure 8 shows monthly NO_x emission totals and inferred changes for several global regions. The magnitude of changes is generally smallest in summer months and largest in winter months for both OMI and TROPOMI inferred emissions. Monthly prior emissions totals lay between the OMI and TROPOMI inferences, except for summertime emissions in India and Mexico where both satellite inferences decrease NO_x emissions. Over Europe, both satellite products infer a decrease during the winter, although fewer valid satellite pixels due to snow cover at high latitudes and longer winter NO_2 atmospheric lifetimes may influence the inference.





395 Figure 8: Monthly prior and satellite-inferred anthropogenic NO_x emissions in 2019 in select global regions. Total monthly emission (as NO_2) from prior emission estimates, inference with OMI or TROPOMI observations (OMI and TROPOMI posterior), and CAMS or TCR-2 inventories in the U.S., China, India, Mexico, and Europe are shown. Percent change ($\Delta E NOx$) inferred with OMI or TROPOMI data, relative to prior emission estimates, for each region are shown by the purple and orange bars.

- Based on reported NO_x emissions trends (McDuffie et al., 2020; U.S. EPA, 2022a), changes from the prior emissions inventory (Table 1) to 2019 are expected. Relative to the prior emissions, significant decreases in NO_x emissions in China, estimated for 2015 in the prior inventory, and smaller reductions in Europe and North America, reported for 2014 and 2017 in the prior inventory, respectively, should be anticipated. TROPOMI inferred emissions reflect the direction anticipated for these changes, but with larger than expected magnitudes. For example, the 28% decrease in anthropogenic NO_x emissions over
- 405 China between 2015 and 2019 inferred from the TROPOMI observations is substantially larger than the 8% decrease estimated between 2015 and 2017 by the Community Emissions Data System (McDuffie et al., 2020). Bottom-up estimates indicate that anthropogenic NO_x emissions in the U.S. have decreased through 2019 (U.S. EPA, 2022a). Although the direction of the



emissions change inferred from TROPOMI agrees with the trend in bottom-up estimates, its magnitude is larger than expected. An underestimate of U.S. emissions in winter in the prior inventory when compared with OMI inferences contrasts with field

- study results reporting no bias in Northeastern U.S. winter emissions estimates (Jaegle et al., 2018; Salmon et al., 2018). In 410 India, bottom-up emissions inventories report sustained growth of NO_x emissions (Kurokawa and Ohara, 2020; McDuffie et al., 2020) and NO₂ levels observed by OMI have been increasing since 2005 (Goldberg et al., 2021; Cooper et al., 2022). The decrease in anthropogenic NO_x emissions inferred by TROPOMI observations contrasts with these trends in bottom-up estimates and OMI observations.
- 415 The low bias known to affect TROPOMI NO₂ observations influences the results of the emissions inversion, which targets grid cells with high emissions, likely leading to larger than expected inferred emissions decreases. We conduct an inversion using the reprocessed TROPOMI NO₂ version 2.3.1 (Van Geffen et al., 2021) to infer NO_x emissions for January 2019, and find that the updated data increases the TROPOMI posterior inference by 17% over the U.S. and 4% in China relative to version 1.2.2, but still differs from significantly from that obtained using OMI observations (Figs. S10 and S11). The 420 differences between emissions inferred by OMI and TROPOMI observations highlight the importance of ongoing efforts to harmonize OMI and TROPOMI NO2 retrieval algorithms, such as the NASA Multi-Decadal Nitrogen Dioxide and Derived Products from Satellites (MINDS) (Lamsal et al., 2020) and the QA4ECV (Boersma et al., 2017) datasets.

In addition to smearing effects, coarse resolution models can artificially alter nonlinear NO₂ chemistry, leading to biases in inferences of NO_x emissions from satellite NO₂ columns (Valin et al., 2011; Sekiya et al., 2021; Lamsal et al., 2011). Higher

- resolution simulations can better resolve β and reduce biases caused by nonlinear chemistry. Additional errors in the emissions 425 estimates may be associated with emissions from non-anthropogenic NO_x sources. Although the emissions inversion targets anthropogenic sources only, changes in NO₂ columns observed by the satellite instruments driven by natural NO_x emissions processes may not be captured by in the air quality model simulations and subsequently lead to biased anthropogenic emissions inferences (Li et al., 2021).
- 430

The emissions resulting from the inverse modeling framework are comparable to CAMS and TCR-2 2019 emissions estimates in several ways. In the U.S., China, and Europe, the magnitudes of ANO_x emissions from OMI retrievals are comparable to TCR-2 NO_x emissions estimates and exhibit similar monthly patterns. Annual NO_x emissions inferred from OMI observations are also relatively similar to TCR-2 estimates for India and Mexico, although monthly emissions patterns differ. Unlike TCR-2 emissions estimates, which are is also constrained by OMI NO₂ observations, 2019 CAMS emissions 435 estimates are projected from 2015 bottom-up data. However, CAMS estimates provide a representation of anticipated

- emissions trends. In all regions considered, CAMS NO_x emissions estimates are close to the TROPOMI inference annual totals and lower than the prior emissions, OMI inferences, and TCR-2 estimates, potentially suggesting that global NO_x emissions have not decreased as much as anticipated by the CAMS inventory projections.
 - 18

440 3.4 Impacts of emissions updates on modeled NO₂ and O₃

We evaluate and compare the CMAQ simulations' ability to reproduce observed pollutant concentrations when driven with NO_x emissions estimates from the prior inventory and those derived by the inverse modeling framework. Figure 9 compares 2019 OMI and TROPOMI NO₂ VCD retrievals with modeled NO₂ VCDs using the prior emissions with no updates, LNO_x emissions updates, and LNO_x and ANO_x emissions updates. Satellite-based LNO_x emissions updates improve CMAQ

- 445 model performance—correlation coefficient (R), normalized mean error (NME), and normalized mean bias (NMB)—when evaluated against tropospheric VCD retrievals, relative to model performance with the prior emissions. OMI-inferred ANO_x emissions updates further improve CMAQ model performance evaluated against VCD retrievals, decreasing NMB from -20% to -5% and NME from 38% to 28%. Model performance is improved by using OMI data in the inverse modeling framework across all seasons (Figs. S6-S9). Although LNO_x emissions updates derived from TROPOMI observations improve model bias
- 450 and error relative to the CMAQ simulation using prior emission estimates, TROPOMI-inferred anthropogenic emissions do not, except during summer months (Figs. 9 and S6-S9). The lack of significant improvements in CMAQ-simulated NO₂ VCDs after applying the emissions inversion with TROPOMI NO₂ retrievals prior to the version 2.3.1 update (Van Geffen et al., 2021) may be associated with changing chemical regimes that are not captured in the emissions inversion process.
- Changes in modeled VCD due to assimilation and the emission inferences calculated in the TROPOMI ANO_x inversion 455 exceed the emissions perturbation and VCD changes used to calculate β . For example, over the eastern U.S., the -15% emissions perturbation used to calculate β leads to VCD changes of -15% on average in winter, but assimilating TROPOMI retrievals leads to VCD changes ($\Delta\Omega$) of -19% on average in the winter, with individual changes exceeding -30%. Modeled NO_x chemistry and NO₂ vertical profiles after assimilating TROPOMI retrievals may be different than those used in the calculation of β . As a result, assimilating TROPOMI retrievals in the ANOx inversion may lead to modeled NO₂ vertical
- 460 profiles which are inconsistent with the precalculated β used in the FDMB relationship and less reliable subsequent emissions inferences. In contrast, the magnitude of VCD changes due to assimilating OMI retrievals over the eastern U.S. in winter is 8%, well within the magnitude of the VCD changes used to precalculate β . This highlights the importance of applying a β sensitivity valid for the magnitude of anticipated emissions changes in FDMB inversions and the potential consequences of relying on satellite-derived retrievals with pre-existing biases in emissions inversions.

465



Figure 9: Impact of NOx emissions updates on modeled NO₂ VCDs. Plots compare 2019 season-average CMAQ-modeled NO₂ VCD at each model grid cell in which NOx emissions were updated by the inverse modeling framework against OMI and TROPOMI tropospheric NO₂ VCD retrievals averaged in each model grid cell. Modeled NO₂ VCD using prior emissions (Prior), inferred LNOx emissions (LNOx posterior), and inferred lightning and anthropogenic NOx emissions (ANOx posterior) are each compared with NO₂ VCD retrievals. Top row plots compare retrievals and modeled VCD based on OMI observations, while bottom row plots compare retrievals and modeled VCD based on TROPOMI observations. Linear regression line, correlation coefficient (R), normalized mean error (NME), and normalized mean bias (NMB), relative to tropospheric NO₂ VCD retrievals, are shown for each CMAQ simulation.

475 Comparing CMAQ-modeled O₃ to ozonesonde measurements from the World Ozone and Ultraviolet Radiation Data Centre (WOUDC) network shows the impacts updating LNO_x emissions on simulated tropospheric O₃ (Fig. 10). Above 300 hPa, the model is biased low, but neither update has a major impact on this bias. However, within the free troposphere, the effects of LNO_x emissions updates are larger. LNO_x satellite-inferred emissions from both satellites increase O₃ and subsequently improve the model's low O₃ bias across all seasons, with the strongest effect in the summer. This suggests a low background NO₂ in our prior simulation, consistent with several studies demonstrating that models underestimate background NO₂ (Goldberg et al., 2017; Qu et al., 2021; Silvern et al., 2019).



485

Figure 10: Ozonesonde observations from the WOUDC network and impact of lightning emissions inferences on modeled ozone. Left plot shows sonde observations averaged in each season and total number of launches per season. The NMB is shown for the prior emissions simulation. Plots on the right show the decrease in the NMB, relative to the prior simulation, for simulations with LNO_x emissions updated with OMI and TROPOMI data.

Comparisons of CMAQ-modeled NO2 and O3 concentrations with ground-level measurements highlight the challenges of 490 reproducing local air quality with a coarse scale model, but suggest potential to improve model performance with satellitederived NO_x emissions updates. Table 2 shows statistics evaluating modeled ground-level daily average NO₂ and maximum 8-hour O_3 concentrations over the U.S. against observations from 1,218 monitoring sites in the Air Quality System (AQS) (U.S. EPA, 2022b), excluding near-road monitors for which the gridded NO₂ fields are not representative of. Statistics for each season are included in Tables S2 and S3. There is a significant low bias in CMAQ-predicted ground-level NO₂ concentrations 495 compared with monitoring site measurements, likely due to the model's coarse grid resolution and the aggregation of NO_2 monitors within urban areas with high NOx emissions and large concentration gradients. CMAQ simulations at higher horizontal resolution do not show the same bias against NO₂ surface observations (Toro et al., 2021). Agreement between modeled and observed ground-level NO₂ concentrations is improved by using OMI-inferred NO_x emissions, compared with the prior emissions simulation, in particular during winter and spring months. Model performance evaluated against ground-

- 500 level O₃ measurements improves to a smaller extent with OMI-inferred NO_x emissions during winter and spring months. The use of TROPOMI-inferred emissions has mixed impacts on CMAQ performance against observed ground-level NO₂ and O₃ concentrations, leading to limited gains in seasonal R and some seasonal biases and errors, but also less agreement with observations for other seasonal statistics. In the U.S., the network of ground-based air quality observations is relatively large. However, in some regions where emissions uncertainties are expected to be especially high, ground-based observations are
- 505 significantly limited and less accessible. Assessing the impact of emissions updated against ground-based observations in these regions, although a challenge, would provide further evaluation of the inversion framework in locations where satellite retrievals have the largest potential to provide important constraints to emissions estimates.

Table 2: CMAQ model performance evaluated against daily average NO₂ (DA NO₂) and maximum 8-hour O₃ concentrations (MDA8 O₃) observed in 2019 by AQS monitoring sites in the U.S. Near-road monitors are not considered. Statistics are shown for simulations using prior emissions (Prior), lightning and anthropogenic NOx emissions inferred with OMI data (OMI-inferred), and lightning and anthropogenic NOx emissions inferred with TROPOMI data (TROPOMI-inferred). Coefficient of determination (R), normalized mean error (NME), and normalized mean bias (NMB), relative to AQS observations, are estimated for each CMAQ simulation.

Pollutant	NO _x emissions	R	NME	NMB
MDA8 O ₃	Prior	0.65	15.9%	-1.4%
	OMI-inferred	0.68	15.4%	3.4%
	TROPOMI-inferred	0.68	15.3%	-3.3%
DA NO ₂	Prior	0.45	62.2%	-56.9%
	OMI-inferred	0.52	57.4%	-49.6%
	TROPOMI-inferred	0.45	71.7%	-69.9%

3.5 Impacts of emissions updates on long-range O₃ transport

- 510 Global NO_x emissions estimates affect model simulations of long-range air pollution transport. To explore these impacts, we examine the response of CMAQ-modeled transpacific O₃ to the inverse modeling framework's NO_x emissions updates. Figure 11 shows season-average changes in simulated free-tropospheric O₃ over the North Pacific Ocean resulting from the use of OMI- and TROPOMI-inferred ANO_x emissions, relative to the emissions simulation with LNO_x emissions updated. As expected, the emissions inversions lead to O₃ variations that follows NO_x emissions changes inferred for each satellite's
- 515 observations, with OMI inferences resulting in higher O₃ concentrations and TROPOMI inferences resulting in lower O₃ concentrations over the North Pacific Ocean. Season-average differences with respect to the prior emissions simulation are as large as +1.8 ppb in winter, using OMI-based updates, and -1.9 ppb in spring, using TROPOMI-based updates. Combined with transpacific wind patterns, the effects of the NO_X emissions inversions on modeled O₃ suggest potential implications of uncertain Asian emissions estimates for U.S. air quality management and emphasize the impacts of biases in satellite retrievals
- 520 in inverse modeling systems.



Figure 11: Changes 2019 season-average free-tropospheric O_3 concentrations (averaged between 750-250 hPa) simulated over the North Pacific Ocean using lightning and anthropogenic NO_x emissions inferred with OMI or TROPOMI observations, relative to simulation using prior ANO_x emissions and updated LNO_x emissions. Differences are shown for winter (DJF), spring (MAM), summer (JJA), and fall (SON). Arrows depict season-average free-tropospheric winds (750-250 hPa). Star marker indicates location of Trinidad Head, California.

At the Trinidad Head, California, a location where atmospheric composition is relatively unaffected by local emissions sources and responsive to transpacific pollution transport (Fig. 11), differences in modeled daily average free-tropospheric O₃ concentrations can reach +5 ppb or -3 ppb. Figure 12 compares CMAQ-modeled vertical O₃ profiles to observations from 39 ozonesondes launched in at Trinidad Head in 2019 (WOUDC, 2019). Relative to the CMAQ simulation using prior emissions,

- 530 NO_x emissions updates inferred from OMI and TROPOMI data can improve the model's ability to reproduce ozonesonde O_3 distributions measured from the site, in particular during winter and spring when the discrepancies between modeled and observed concentrations are largest. These impacts on modeled vertical O_3 profiles are largely driven by changes the modeling framework's updates to lightning NO_x emissions. The inferred LNO_x increases from each satellite improve O_3 biases, while subsequent anthropogenic updates have smaller impacts, suggesting that biases in O_3 could be driven by background NO_2
- 535 composition in the model and not solely by long-range transport resulting from anthropogenic emissions.



Figure 12: Season-average vertical O₃ concentration profiles modeled by CMAQ and measured by ozonesondes launched at Trinidad Head, California, in 2019. Vertical distributions are shown for simulations using prior emissions (Prior), lightning and anthropogenic NOx emissions inferred with OMI data (OMI-inferred), and lightning and anthropogenic NOx emissions inferred with TROPOMI data (TROPOMI-inferred). Modeled season-average profiles are shown during winter (DJF), spring (MAM), summer (JJA), and fall (SON) for days and times matching ozonesonde launches. Shading around sonde and prior emissions profiles represent the maximum and minimum O₃ at each pressure level. Map shows location of the Trinidad Head launch site.

4 Conclusions

In this study, we describe a satellite chemical data assimilation and inverse emissions modeling framework based on the 545 CMAQ hemispheric air quality modeling platform. In the framework, data assimilation adjusts modeled NO₂ concentrations online using satellite retrievals of tropospheric NO₂ VCDs. The NO₂ column changes drive the FDMB inversion, resulting in satellite-constrained top-down emissions estimates. Here, we implement the framework in a NO_x emissions inversion to separately update 2019 Northern Hemisphere lightning and anthropogenic NO_x emissions estimates using NO₂ products from the OMI and TROPOMI satellite instruments. Relative to the modeling platform's prior emissions derived from regional and 550 global emissions inventories, updates inferred using OMI and TROPOMI observations change average anthropogenic NO_x emissions by -41% to +12% in China, the U.S., India, Europe, and Mexico. Evaluated against ground-based NO₂ observations recorded over the U.S. in 2019, the model performs best when using OMI-updated emissions, although a low bias in CMAO predictions using prior emissions persists into simulations with satellite data assimilation. Compared with U.S ground-based O₃ observations, satellite-inferred emissions have mixed impacts on model performance, improving agreement with the measurements during certain months. LNO_x emissions inferences improve modeled O₃ when compared against ozonesonde 555 observations across the Northern Hemisphere. The framework's NO_x emissions updates also affect model estimates of trans-Pacific O₃ transport, a source of growing concern in the U.S., with changes ranging from -3 ppb to +5 ppb in simulated O₃ at a remote West Coast site resulting from use of satellite-inferred emissions.

The modeling framework presented has several limitations. <u>The computational cost is greater than that of traditional</u> 560 <u>FDMB inversions due to the assimilation step. However, the computational burden is comparable or less than other satellite</u> assimilation methods such as Kalman-Filter and adjoint 4D-variational approaches. In addition, the framework requires <u>minimal code changes to the underlying CTM, so inverse estimates will improve as the underlying air quality model is updated</u> with little additional effort needed to implement this framework. The global coverage of instruments on polar-orbiting satellites, such as Aura and Sentinel-5P, makes the emissions inversions possible but does not allow satellite observations to

- inform diurnal emissions variations. Upcoming geostationary satellite missions, including GEMS, TEMPO, and Sentinel-4, 565 will provide this capability. Our approach, which balances computational costs and precision in the inversion, is subject to several assumptions. As in all mass-balance based approaches, our method fully attributes the change in the VCD to emissions changes. To the extent that column differences are due to chemistry or transport and not emissions, this assumption introduces error into mass-balance inversions, including the inversion implemented in our framework. Large changes to the model
- concentrations resulting from the chemical data assimilation may invalidate assumptions in the subsequent FDMB inversion, 570 leading to biases in the inferred emissions. The FDMB inversion treats each grid cell independently and cannot relate NO₂ column changes in one grid cell to emissions in another. Although emissions smearing in the approach is mitigated by only analyzing the lower portion of the model column, our emissions changes may be less precise than targeted assimilation methods, such as 4DVAR adjoint-based methods. Further, coarse grid resolution exacerbates biases in modeled NO₂ columns
- 575 (Valin et al., 2011) and inferred NO_x emissions (Sekiya et al., 2021). The air quality model used here does not include stratospheric chemistry, which could affect comparisons against NO2 retrievals. Nevertheless, the framework shows the potential to improved air quality model predictions using satellite-derived emissions updates, in particular for regions with highly uncertain emissions inventories.
- Emissions inversions based on satellite observations can provide valuable information for air quality modeling by 580 addressing the gaps in bottom-up emissions inventories. However, our analysis shows that such inversions and subsequent air quality simulations can be strongly influenced by uncertainties and biases in the satellite data products used. In the analysis conducted, NO_x emissions inferred from TROPOMI observations appear biased low when assessed against those inferred from OMI data and surface and concentration measurements. The bias is consistent with recent research showing a low bias in TROPOMI v1.2 and v1.3 tropospheric columns (Judd et al., 2020; Verhoelst et al., 2021; Li et al., 2021; Van Geffen et al.,
- 585 2021). The results highlight the importance of efforts to develop robust and consistent satellite data products for use in air quality modeling evaluation, assimilation, and emissions inversions. Ongoing efforts to this end include the MINDS (Lamsal et al., 2020) and the QA4ECV (Boersma et al., 2017) projects. This study also emphasizes the need for longer-term satellite data assimilation and comparisons of established and new satellite data products. The framework introduced here can serve a generalized tool with applications beyond those explored in this study, and allows new satellite data products to be incorporated as they become available. As satellite data products evolve and advance, the emissions inferred by the framework will improve.

590

595

Data and code availability. NO_x emissions data derived from this research are available from the authors upon request. Level-2 satellite retrievals are available from NASA's Goddard Earth Sciences Data and Information Services Center for OMI (https://disc.gsfc.nasa.gov/datasets/OMNO2 003/summary) and TROPOMI 1 version (https://disc.gsfc.nasa.gov/datasets/S5P L2 NO2 1/summary). TROPOMI retrievals reprocessed to version 2.3.1 are available through the Sentinel-5P data portal (https://data-portal.s5p-pal.com/). WOUDC ozonesonde data, including data at

the Trinidad California available WOUDC Head, launch site, are through at

(<u>https://woudc.org/data/dataset_info.php?id=ozonesonde</u>). Hourly AQS O₃ and NO₂ observations are available from EPA's Air Data website (<u>https://aqs.epa.gov/aqsweb/airdata/download_files.html</u>) (U.S. EPA, 2022b). GSI code is available via

600 <u>https://dtcenter.org/community-code/gridpoint-statistical-interpolation-gsi/download</u> (DTC, 2018) CMAQ source code is freely available via <u>https://github.com/usepa/cmaq.git</u> and via the EPA (U.S. EPA, 2020).

Author contributions. JDE, BHH, SLN, SNK, and FGM designed the study. RBP, BHH, and DQT provided initial conceptualization for the research. JDE conducted the air quality simulations, data assimilation, emissions inversions, and data

- 605 analysis. AL developed the OMI and TROPOMI observation operators in GSI. BHH and JDE developed the CMAQ and GSI coupling. RG performed the meteorological modeling. GS implemented the halogen chemistry in the CMAQ model. JDE, BHH, SLN, SNK, RBP, and FGM participated in data analysis and discussions. JDE wrote the paper with input from all co-authors.
- 610 Competing interests. The authors declare that they have no conflict of interest.

Disclaimer. The views expressed in this article are those of the authors and do not necessarily represent the views or policies of the U.S. Environmental Protection Agency.

- 615 Acknowledgments. J. D. East was supported, in part, by an appointment to the Research Participation Program at the Office of Research and Development, U.S. Environmental Protection Agency, administered by the Oak Ridge Institute for Science and Education through an interagency agreement between the U.S. Department of Energy and the EPA. D. Q. Tong and R. B. Pierce acknowledge financial support from the NASA Health and Air Quality Applied Sciences Team (HAQAST) Tiger Team under award number 80NSSC21K0427. We gratefully acknowledge the free availability and use of observational data sets
- 620 from AQS and WOUDC; remote sensing retrievals from OMI and TROPOMI; and global emission inventories from CAMS, TCR-2, and GEIA. Comments by Heather Simon and Tanya Spero at the U.S. EPA served to strengthen this manuscript.

References

Adams, E.: 2017 v1 NEI Emissions Modeling Platform (Premerged CMAQ-ready Emissions) (V1), UNC Dataverse [dataset], doi:10.15139/S3/TCR6BB, 2020.

625 Anenberg, S. C., Henze, D. K., Tinney, V., Kinney, P. L., Raich, W., Fann, N., Malley, C. S., Roman, H., Lamsal, L., Duncan, B., Martin, R. V., van Donkelaar, A., Brauer, M., Doherty, R., Jonson, J. E., Davila, Y., Sudo, K., and Kuylenstierna, J. C. I.: Estimates of the Global Burden of Ambient PM2.5, Ozone, and NO2 on Asthma Incidence and Emergency Room Visits, Environ Health Persp, 126, 10.1289/Ehp3766, 2018.

Appel, K. W., Bash, J. O., Fahey, K. M., Foley, K. M., Gilliam, R. C., Hogrefe, C., Hutzell, W. T., Kang, D. W., Mathur, R., Murphy, B.
 N., Napelenok, S. L., Nolte, C. G., Pleim, J. E., Pouliot, G. A., Pye, H. O. T., Ran, L. M., Roselle, S. J., Sarwar, G., Schwede, D. B., Sidi, F. I., Spero, T. L., and Wong, D. C.: The Community Multiscale Air Quality (CMAQ) model versions 5.3 and 5.3.1: system updates and evaluation, Geosci Model Dev, 14, 2867-2897, 10.5194/gmd-14-2867-2021, 2021.

Boersma, K. F., Vinken, G. C. M., and Tournadre, J.: Ships going slow in reducing their NOx emissions: changes in 2005-2012 ship exhaust inferred from satellite measurements over Europe, Environ Res Lett, 10, 10.1088/1748-9326/10/7/074007, 2015.

- 635 Boersma, K. F., Eskes, H., Richter, A., De Smedt, I., Lorente, A., Beirle, S., Van Geffen, J., Peters, E., Van Roozendael, M., and Wagner, T.: QA4ECV NO2 tropospheric and stratospheric vertical column data from OMI (Version 1.1) [dataset], 10.21944/qa4ecv-no2-omiv1.1, 2017.
 - Boersma, K. F., Eskes, H. J., Richter, A., De Smedt, I., Lorente, A., Beirle, S., van Geffen, J. H. G. M., Zara, M., Peters, E., Van Roozendael, M., Wagner, T., Maasakkers, J. D., van der A, R. J., Nightingale, J., De Rudder, A., Irie, H., Pinardi, G., Lambert, J. C., and Compernolle, S. C.: Improving algorithms and uncertainty estimates for satellite NO2 retrievals: results from the quality assurance for

640

- the essential climate variables (QA4ECV) project, Atmos Meas Tech, 11, 6651-6678, 10.5194/amt-11-6651-2018, 2018.
 Byun, D. and Schere, K. L.: Review of the governing equations, computational algorithms, and other components of the models-3 Community Multiscale Air Quality (CMAQ) modeling system, Appl Mech Rev, 59, 51-77, 10.1115/1.2128636, 2006.
- Chance, K. (Ed.) OMI Algorithm Theoretical Basis Document. OMI Trace Gas Algorithms., Version 2.0, Smithsonian Astrophysical
 Observatory, Cambridge, MA, USA, 2002.

CIESIN: Center for International Earth Science Information Network - CIESIN - Columbia University. Gridded Population of the World, Version 4 (GPWv4): Population Density, Revision 10 [dataset], 2018.

Cooper, M., Martin, R. V., Padmanabhan, A., and Henze, D. K.: Comparing mass balance and adjoint methods for inverse modeling of nitrogen dioxide columns for global nitrogen oxide emissions, J Geophys Res-Atmos, 122, 4718-4734, 10.1002/2016jd025985, 2017.

- 650 Cooper, M. J., Martin, R. V., Hammer, M. S., Levelt, P. F., Veefkind, P., Lamsal, L. N., Krotkov, N. A., Brook, J. R., and McLinden, C. A.: Global fine-scale changes in ambient NO2 during COVID-19 lockdowns, Nature, 601, 380-+, 10.1038/s41586-021-04229-0, 2022.
 - Crippa, M., Solazzo, E., Huang, G. L., Guizzardi, D., Koffi, E., Muntean, M., Schieberle, C., Friedrich, R., and Janssens-Maenhout, G.: High resolution temporal profiles in the Emissions Database for Global Atmospheric Research, Sci Data, 7, 10.1038/s41597-020-0462-2, 2020.
- 655 Day, M., Pouliot, G., Hunt, S., Baker, K. R., Beardsley, M., Frost, G., Mobley, D., Simon, H., Henderson, B. B., Yelverton, T., and Rao, V.: Reflecting on progress since the 2005 NARSTO emissions inventory report, J Air Waste Manage, 69, 1023-1048, 10.1080/10962247.2019.1629363, 2019.
 - de Foy, B. and Schauer, J. J.: An improved understanding of NOx emissions in South Asian megacities using TROPOMI NO2 retrievals, Environ Res Lett, 17, <u>https://doi.org/10.1088/1748-9326/ac48b4</u>, 2022.
- 660 Descombes, G., Auligné, T., Vandenberghe, F., Barker, D. M., and Barré, J.: Generalized background error covariance matrix model (GEN_BE v2.0), Geosci. Model Dev., 8, 669-696, 10.5194/gmd-8-669-2015, 2015.
 - Developmental Testbed Center (DTC): Gridpoint Statistical Interpolation (GSI), https://dtcenter.org/community-code/gridpoint-statisticalinterpolation-gsi/download, 2018.
- Ding, J., van der A, R. J., Mijling, B., Levelt, P. F., and Hao, N.: NOx emission estimates during the 2014 Youth Olympic Games in Nanjing,
 Atmos Chem Phys, 15, 9399-9412, 10.5194/acp-15-9399-2015, 2015.
- Ding, J., van der A, R. J., Eskes, H. J., Mijling, B., Stavrakou, T., van Geffen, J. H. G. M., and Veefkind, J. P.: NOx Emissions Reduction and Rebound in China Due to the COVID-19 Crisis, Geophys Res Lett, 47, 10.1029/2020GL089912, 2020.

 Elguindi, N., Granier, C., Stavrakou, T., Darras, S., Bauwens, M., Cao, H., Chen, C., van der Gon, H. A. C. D., Dubovik, O., Fu, T. M., Henze, D. K., Jiang, Z., Keita, S., Kuenen, J. J. P., Kurokawa, J., Liousse, C., Miyazaki, K., Muller, J. F., Qu, Z., Solmon, F., and Zheng, B.: Intercomparison of Magnitudes and Trends in Anthropogenic Surface Emissions From Bottom-Up Inventories, Top-Down

Estimates, and Emission Scenarios, Earths Future, 8, 10.1029/2020EF001520, 2020.

- Eskes, H. J. and Boersma, K. F.: Averaging kernels for DOAS total-column satellite retrievals, Atmos Chem Phys, 3, 1285-1291, DOI 10.5194/acp-3-1285-2003, 2003.
- Eskes, H. J., Eichmann, K. U., Lambert, J. C., Loyola, D., Veefkind, J. P., Dehn, A., and Zehner, C.: S5P/TROPOMI NO2 Level 2 Product Readme File Sentinel-5P Mission Performance Centre 2021.
- Eskes, H. J., van Geffen, J., Boersma, K. F., Eichmann, K. U., Apituley, A., Pedergnana, M., Sneep, M., veefkind, J. P., and Diego, L.: S5P/TROPOMI Level-2 Product User Manual Nitrogen Dioxide, Royal Netherlands Meteorological InsituteS5P-KNMI-L2-0021-MA, 2019.
- Goldberg, D. L., Lamsal, L. N., Loughner, C. P., Swartz, W. H., Lu, Z. F., and Streets, D. G.: A high-resolution and observationally
 constrained OMI NO2 satellite retrieval, Atmos Chem Phys, 17, 11403-11421, 10.5194/acp-17-11403-2017, 2017.
 - Goldberg, D. L., Anenberg, S. C., Lu, Z. F., Streets, D. G., Lamsal, L. N., McDuffie, E. E., and Smith, S. J.: Urban NOx emissions around the world declined faster than anticipated between 2005 and 2019, Environ Res Lett, 16, 10.1088/1748-9326/ac2c34, 2021.
- Granier, C., S. Darras, H. Denier van der Gon, J. Doubalova, N. Elguindi, B. Galle, M. Gauss, M. Guevara, J.-P. Jalkanen, J. Kuenen, C. Liousse, B. Quack, D. Simpson, and Sindelarova, K.: The CAMS global and regional emissions (April 2019 version), Copernicus
 Atmosphere Monitoring Service, 10.24380/d0bn-kx16, 2019.
 - Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), Atmos. Chem. Phys., 6, 3181-3210, 10.5194/acp-6-3181-2006, 2006.

- Hoesly, R. M., Smith, S. J., Feng, L. Y., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M.,
 Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J., Li, M., Liu, L., Lu, Z. F., Moura, M. C. P., O'Rourke, P. R., and Zhang, Q.:
 Historical (1750-2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS),
 Geosci Model Dev, 11, 369-408, 10.5194/gmd-11-369-2018, 2018.
- Itahashi, S., Mathur, R., Hogrefe, C., Napelenok, S. L., and Zhang, Y.: Modeling stratospheric intrusion and trans-Pacific transport on tropospheric ozone using hemispheric CMAQ during April 2010-Part 2: Examination of emission impacts based on the higher-order decoupled direct method, Atmos Chem Phys, 20, 3397-3413, 10.5194/acp-20-3397-2020, 2020.
- Itahashi, S., Yumimoto, K., Kurokawa, J. I., Morino, Y., Nagashima, T., Miyazaki, K., Maki, T., and Ohara, T.: Inverse estimation of NOx emissions over China and India 2005?2016: contrasting recent trends and future perspectives, Environ Res Lett, 14, 10.1088/1748-9326/ab4d7f, 2019.
- Jacob, D. J.: Global Budget of Nitrogen Oxides, in: Introduction to Atmospheric Chemistry, 1st ed., Princeton University Press, Princeton, NJ, 213, 1999.
- Jacob, D. J.: Heterogeneous chemistry and tropospheric ozone, Atmos Environ, 34, 2131-2159, Doi 10.1016/S1352-2310(99)00462-8, 2000.
 Jaegle, L., Shah, V., Thornton, J. A., Lopez-Hilfiker, F. D., Lee, B. H., McDuffie, E. E., Fibiger, D., Brown, S. S., Veres, P., Sparks, T. L., Ebben, C. J., Wooldridge, P. J., Kenagy, H. S., Cohen, R. C., Weinheimer, A. J., Campos, T. L., Montzka, D. D., Digangi, J. P., Wolfe, G. M., Hanisco, T., Schroder, J. C., Campuzano-Jost, P., Day, D. A., Jimenez, J. L., Sullivan, A. P., Guo, H., and Weber, R. J.: Nitrogen Oxides Emissions, Chemistry, Deposition, and Export Over the Northeast United States During the WINTER Aircraft Campaign, J Geophys Res-Atmos, 123, 12368-12393, 10.1029/2018jd029133, 2018.
 - Jaffe, D. A., Cooper, O. R., Fiore, A. M., Henderson, B. H., Tonnesen, G. S., Russell, A. G., Henze, D. K., Langford, A. O., Lin, M. Y., and Moore, T.: Scientific assessment of background ozone over the US: Implications for air quality management, Elementa-Sci Anthrop, 6, 10.1525/elementa.309, 2018.
- 710 Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Dentener, F., Muntean, M., Pouliot, G., Keating, T., Zhang, Q., Kurokawa, J., Wankmuller, R., van der Gon, H. D., Kuenen, J. J. P., Klimont, Z., Frost, G., Darras, S., Koffi, B., and Li, M.: HTAP_v2.2: a mosaic of regional and global emission grid maps for 2008 and 2010 to study hemispheric transport of air pollution, Atmos Chem Phys, 15, 11411-11432, 10.5194/acp-15-11411-2015, 2015.
- Judd, L. M., Al-Saadi, J. A., Szykman, J. J., Valin, L. C., Janz, S. J., Kowalewski, M. G., Eskes, H. J., Veefkind, J. P., Cede, A., Mueller,
 M., Gebetsberger, M., Swap, R., Pierce, R. B., Nowlan, C. R., Abad, G. G., Nehrir, A., and Williams, D.: Evaluating Sentinel-5P TROPOMI tropospheric NO2 column densities with airborne and Pandora spectrometers near New York City and Long Island Sound, Atmos Meas Tech, 13, 6113-6140, 10.5194/amt-13-6113-2020, 2020.
 - Kang, D., Willison, J., Sarwar, G., Madden, M., Hogrefe, C., Mathur, R., Gantt, B., and Saiz-Lopez, A.: Improving the Characterization of Natural Emissions in CMAQ, EM Magazine, 30-36, 2021.
- 720 Krotkov, N. A., Lamsal, L. N., Marchenko, S. V., and Swartz, W. H.: OMNO2 README Document. Data Product Version 4.0., NASA/Goddard Space Flight Center, 2019a.
 - Krotkov, N. A., Lamsal, L. N., Marchenko, S. V., J.Bucsela, E., Swartz, W. H., Joiner, J., and team, O. c.: OMI/Aura Nitrogen Dioxide (NO2) Total and Tropospheric Column 1-orbit L2 Swath 13x24 km V003, Goddard Earth Sciences Data and Information Services Center (GES DISC) [dataset], 10.5067/Aura/OMI/DATA2017, 2019b.
- 725 Kurokawa, J. and Ohara, T.: Long-term historical trends in air pollutant emissions in Asia: Regional Emission inventory in ASia (REAS) version 3, Atmos Chem Phys, 20, 12761-12793, 10.5194/acp-20-12761-2020, 2020.
 - Kurokawa, J., Yumimoto, K., Uno, I., and Ohara, T.: Adjoint inverse modeling of NOx emissions over eastern China using satellite observations of NO2 vertical column densities, Atmos Environ, 43, 1878-1887, 10.1016/j.atmosenv.2008.12.030, 2009.
- Lamsal, L. N., Martin, R. V., Padmanabhan, A., van Donkelaar, A., Zhang, Q., Sioris, C. E., Chance, K., Kurosu, T. P., and Newchurch, M.
 J.: Application of satellite observations for timely updates to global anthropogenic NOx emission inventories, Geophys Res Lett, 38, 10.1029/2010gl046476, 2011.
- Lamsal, L. N., Nickolay A. Krotkov, Sergey V. Marchenko, Joanna Joiner, Luke Oman, Alexander Vasilkov, Bradford Fisher, Wenhan Qin, Eun-Su Yang, Zachary Fasnacht, Sungyeon Choi, Peter Leonard, and Haffner, D.: OMI/Aura NO2 Tropospheric, Stratospheric & Total Columns MINDS 1-Orbit L2 Swath 13 km x 24 km, Goddard Earth Sciences Data and Information Services Center (GES DISC)
 [dataset], 10.5067/MEASURES/MINDS/DATA201, 2020.
- Lamsal, L. N., Krotkov, N. A., Celarier, E. A., Swartz, W. H., Pickering, K. E., Bucsela, E. J., Gleason, J. F., Martin, R. V., Philip, S., Irie, H., Cede, A., Herman, J., Weinheimer, A., Szykman, J. J., and Knepp, T. N.: Evaluation of OMI operational standard NO2 column retrievals using in situ and surface-based NO2 observations, Atmos Chem Phys, 14, 11587-11609, 10.5194/acp-14-11587-2014, 2014.
- Li, J. F. and Wang, Y. H.: Inferring the anthropogenic NOx emission trend over the United States during 2003-2017 from satellite observations: was there a flattening of the emission trend after the Great Recession?, Atmos Chem Phys, 19, 15339-15352, 10.5194/acp-19-15339-2019, 2019.
 - Li, M., Klimont, Z., Zhang, Q., Martin, R. V., Zheng, B., Heyes, C., Cofala, J., Zhang, Y. X., and He, K. B.: Comparison and evaluation of anthropogenic emissions of SO2 and NOx over China, Atmos Chem Phys, 18, 3433-3456, 10.5194/acp-18-3433-2018, 2018.
 - 28

- Li, M., McDonald, B. C., McKeen, S. A., Eskes, H., Levelt, P., Francoeur, C., Harkins, C., He, J., Barth, M., Henze, D. K., Bela, M. M.,
 Trainer, M., Gouw, J. A., and Frost, G. J.: Assessment of Updated Fuel-Based Emissions Inventories Over the Contiguous United States
 Using TROPOMI NO2 Retrievals, J Geophys Res-Atmos, 126, 10.1029/2021JD035484, 2021.
 - Luecken, D. J., Yarwood, G., and Hutzell, W. T.: Multipollutant modeling of ozone, reactive nitrogen and HAPs across the continental US with CMAQ-CB6, Atmos Environ, 201, 62-72, 10.1016/j.atmosenv.2018.11.060, 2019.
- Martin, R. V., Jacob, D. J., Chance, K., Kurosu, T. P., Palmer, P. I., and Evans, M. J.: Global inventory of nitrogen oxide emissions
 constrained by space-based observations of NO2 columns, J Geophys Res-Atmos, 108, 10.1029/2003jd003453, 2003.
- Mathur, R., Xing, J., Gilliam, R., Sarwar, G., Hogrefe, C., Pleim, J., Pouliot, G., Roselle, S., Spero, T. L., Wong, D. C., and Young, J.: Extending the Community Multiscale Air Quality (CMAQ) modeling system to hemispheric scales: overview of process considerations and initial applications, Atmos Chem Phys, 17, 12449-12474, 10.5194/acp-17-12449-2017, 2017.
- McDuffie, E. E., Smith, S. J., O'Rourke, P., Tibrewal, K., Venkataraman, C., Marais, E. A., Zheng, B., Crippa, M., Brauer, M., and Martin,
 R. V.: A global anthropogenic emission inventory of atmospheric pollutants from sector- and fuel-specific sources (1970-2017): an application of the Community Emissions Data System (CEDS), Earth Syst Sci Data, 12, 3413-3442, 10.5194/essd-12-3413-2020, 2020.
 - Mijling, B. and van der A, R. J.: Using daily satellite observations to estimate emissions of short-lived air pollutants on a mesoscopic scale, J Geophys Res-Atmos, 117, 10.1029/2012jd017817, 2012.
- Miyazaki, K. and Eskes, H.: Constraints on surface NOx emissions by assimilating satellite observations of multiple species, Geophys Res Lett, 40, 4745-4750, 10.1002/grl.50894, 2013.
 - Miyazaki, K., Eskes, H. J., and Sudo, K.: Global NOx emission estimates derived from an assimilation of OMI tropospheric NO2 columns, Atmos Chem Phys, 12, 2263-2288, 10.5194/acp-12-2263-2012, 2012a.
 - Miyazaki, K., Eskes, H., Sudo, K., Boersma, K. F., Bowman, K., and Kanaya, Y.: Decadal changes in global surface NOx emissions from multi-constituent satellite data assimilation, Atmos Chem Phys, 17, 807-837, 10.5194/acp-17-807-2017, 2017.
- 765 Miyazaki, K., Eskes, H. J., Sudo, K., Takigawa, M., van Weele, M., and Boersma, K. F.: Simultaneous assimilation of satellite NO2, O-3, CO, and HNO3 data for the analysis of tropospheric chemical composition and emissions, Atmos Chem Phys, 12, 9545-9579, 10.5194/acp-12-9545-2012, 2012b.
 - Miyazaki, K., Bowman, K., Sekiya, T., Eskes, H., Boersma, F., Worden, H., Livesey, N., Payne, V. H., Sudo, K., Kanaya, Y., Takigawa, M., and Ogochi, K.: Chemical Reanalysis Products [dataset], 2019.
- 770 Miyazaki, K., Bowman, K., Sekiya, T., Eskes, H., Boersma, F., Worden, H., Livesey, N., Payne, V. H., Sudo, K., Kanaya, Y., Takigawa, M., and Ogochi, K.: Updated tropospheric chemistry reanalysis and emission estimates, TCR-2, for 2005-2018, Earth Syst Sci Data, 12, 2223-2259, 10.5194/essd-12-2223-2020, 2020.
 - Muller, J. F. and Stavrakou, T.: Inversion of CO and NOx emissions using the adjoint of the IMAGES model, Atmos Chem Phys, 5, 1157-1186, 2005.
- 775 Murray, C. J. L., Aravkin, A. Y., Zheng, P., Abbafati, C., Abbas, K. M., Abbasi-Kangevari, M., Abd-Allah, F., Abdelalim, A., Abdollahi, M., Abdollahpour, I., Abegaz, K. H., Abolhassani, H., Aboyans, V., Abreu, L. G., Abrigo, M. R. M., and Factors, G. R.: Global burden of 87 risk factors in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019, Lancet, 396, 1223-1249, 10.1016/S0140-6736(20)30752-2, 2020.
- Napelenok, S. L., Pinder, R. W., Gilliland, A. B., and Martin, R. V.: A method for evaluating spatially-resolved NOx emissions using Kalman filter inversion, direct sensitivities, and space-based NO2 observations, Atmos Chem Phys, 8, 5603-5614, DOI 10.5194/acp-8-5603-2008, 2008.
 - Powers, J. G., Klemp, J. B., Skamarock, W. C., Davis, C. A., Dudhia, J., Gill, D. O., Coen, J. L., Gochis, D. J., Ahmadov, R., Peckham, S. E., Grell, G. A., Michalakes, J., Trahan, S., Benjamin, S. G., Alexander, C. R., Dimego, G. J., Wang, W., Schwartz, C. S., Romine, G. S., Liu, Z. Q., Snyder, C., Chen, F., Barlage, M. J., Yu, W., and Duda, M. G.: THE WEATHER RESEARCH AND FORECASTING
- 785 MODEL Overview, System Efforts, and Future Directions, B Am Meteorol Soc, 98, 1717-1737, 10.1175/Bams-D-15-00308.1, 2017. Price, C., Penner, J., and Prather, M.: NOx from lightning .1. Global distribution based on lightning physics, J Geophys Res-Atmos, 102, 5929-5941, Doi 10.1029/96jd03504, 1997.
 - Qu, Z., Henze, D. K., Theys, N., Wang, J., and Wang, W.: Hybrid Mass Balance/4D-Var Joint Inversion of NOx and SO2 Emissions in East Asia, J Geophys Res-Atmos, 124, 8203-8224, 10.1029/2018jd030240, 2019.
- 790 Qu, Z., Henze, D. K., Capps, S. L., Wang, Y., Xu, X. G., Wang, J., and Keller, M.: Monthly top-down NOx emissions for China (2005-2012): A hybrid inversion method and trend analysis, J Geophys Res-Atmos, 122, 4600-4625, 10.1002/2016jd025852, 2017.
 - Qu, Z., Jacob, D. J., Silvern, R. F., Shah, V., Campbell, P. C., Valin, L. C., and Murray, L. T.: US COVID-19 Shutdown Demonstrates Importance of Background NO2 in Inferring NOx Emissions From Satellite NO2 Observations, Geophys Res Lett, 48, 10.1029/2021GL092783, 2021.
- 795 Salmon, O. E., Shepson, P. B., Ren, X., He, H., Hall, D. L., Dickerson, R. R., Stirm, B. H., Brown, S. S., Fibiger, D. L., McDuffie, E. E., Campos, T. L., Gurney, K. R., and Thornton, J. A.: Top-Down Estimates of NOx and CO Emissions From Washington, DC-Baltimore During the WINTER Campaign, J Geophys Res-Atmos, 123, 7705-7724, 10.1029/2018jd028539, 2018.
 - Sandu, A. and Chai, T. F.: Chemical Data Assimilation-An Overview, Atmosphere-Basel, 2, 426-463, 10.3390/atmos2030426, 2011.

- Sekiya, T., Miyazaki, K., Ogochi, K., Sudo, K., Takigawa, M., Eskes, H., and Boersma, K. F.: Impacts of Horizontal Resolution on Global
 Data Assimilation of Satellite Measurements for Tropospheric Chemistry Analysis, J Adv Model Earth Sy, 13, 10.1029/2020MS002180, 2021.
 - Shao, H., Derber, J., Huang, X. Y., Hu, M., Newman, K., Stark, D., Lueken, M., Zhou, C. H., Nance, L., Kuo, Y. H., and Brown, B.: Bridging Research to Operations Transitions: Status and Plans of Community GSI, B Am Meteorol Soc, 97, 1427-1440, 10.1175/Bams-D-13-00245.1, 2016.
- 805 Silvern, R. F., Jacob, D. J., Mickley, L. J., Sulprizio, M. P., Travis, K. R., Marais, E. A., Cohen, R. C., Laughner, J. L., Choi, S., Joiner, J., and Lamsal, L. N.: Using satellite observations of tropospheric NO2 columns to infer long-term trends in US NOx emissions: the importance of accounting for the free tropospheric NO2 background, Atmos Chem Phys, 19, 8863-8878, 10.5194/acp-19-8863-2019, 2019.
- Simon, H., Reff, A., Wells, B., Xing, J., and Frank, N.: Ozone Trends Across the United States over a Period of Decreasing NOx and VOC
 Emissions, Environ Sci Technol, 49, 186-195, 10.1021/es504514z, 2015.
- Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Duda, M. G., Huang, X. Y., Wang, W., and Powers, J. G.: A Description of the Advanced Research WRF Version 3, NCAR Tech Note NCAR/TN 475 STR, University Corporation for Atmospheric Research, 125, 10.5065/D68S4MVH, 2008.
- Tong, D. Q., Lamsal, L., Pan, L., Ding, C., Kim, H., Lee, P., Chai, T. F., Pickering, K. E., and Stajner, I.: Long-term NOx trends over large cities in the United States during the great recession: Comparison of satellite retrievals, ground observations, and emission inventories, Atmos Environ, 107, 70-84, 10.1016/j.atmosenv.2015.01.035, 2015.
 - Tong, D. Q., Pan, L., Chen, W. W., Lamsal, L., Lee, P., Tang, Y. H., Kim, H., Kondragunta, S., and Stajner, I.: Impact of the 2008 Global Recession on air quality over the United States: Implications for surface ozone levels from changes in NOx emissions, Geophys Res Lett, 43, 9280-9288, 10.1002/2016gl069885, 2016.
- 820 Toro, C., Foley, K., Simon, H., Henderson, B., Baker, K. R., Eyth, A., Timin, B., Appel, W., Luecken, D., Beardsley, M., Sonntag, D., Possiel, N., and Roberts, S.: Evaluation of 15 years of modeled atmospheric oxidized nitrogen compounds across the contiguous United States, Elementa-Sci Anthrop, 9, 10.1525/elementa.2020.00158, 2021.
 - Turner, A. J., Henze, D. K., Martin, R. V., and Hakami, A.: The spatial extent of source influences on modeled column concentrations of short-lived species, Geophys Res Lett, 39, 10.1029/2012gl051832, 2012.
- 825 U.S. Environmental Protection Agency (U.S. EPA): Community Multiscale Air Quality (CMAQ) model version 5.3.2, 10.5281/zenodo.4081737, 2020.
 - U.S. Environmental Protection Agency (U.S. EPA): Air Pollutant Emissions Trends Data, https://www.epa.gov/air-emissions-inventories/air-pollutant-emissions-trends-data, 2022a.
 - U.S. Environmental Protection Agency (U.S. EPA): Air Quality System (AQS), https://www.epa.gov/aqs, 2022b.
- 830 Valin, L. C., Russell, A. R., Hudman, R. C., and Cohen, R. C.: Effects of model resolution on the interpretation of satellite NO2 observations, Atmos Chem Phys, 11, 11647-11655, 10.5194/acp-11-11647-2011, 2011.
 - van Geffen, J., Eskes, H. J., Boersma, K. F., Maasakkers, J. D., and veefkind, J. P.: TROPOMI ATBD of the total and tropospheric NO2 data products, Royal Netherlands Meteorological Institute, 2019.
- van Geffen, J., Boersma, K. F., Eskesl, H., Sneep, M., ter Linden, M., Zara, M., and Veefkind, J. P.: S5P TROPOMI NO2 slant column retrieval: method, stability, uncertainties and comparisons with OMI, Atmos Meas Tech, 13, 1315-1335, 10.5194/amt-13-1315-2020, 2020.
 - van Geffen, J., Eskes, H., Compernolle, S., Pinardi, G., Verhoelst, T., Lambert, J. C., Sneep, M., Ter Linden, M., Ludewig, A., Boersma, F., and Veefkind, J. P.: Sentinel-5P TROPOMI NO2 retrieval: impact of version v2.2 improvements and comparisons with OMI and ground-based data, Atmospheric Measurement Techniques Discussions, [preprint]. In review., 10.5194/amt-2021-329, 2021.
- 840 Veefkind, J. P., Aben, I., McMullan, K., Forster, H., de Vries, J., Otter, G., Claas, J., Eskes, H. J., de Haan, J. F., Kleipool, Q., van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser, H., and Levelt, P. F.: TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications, Remote Sens Environ, 120, 70-83, 10.1016/j.rse.2011.09.027, 2012.
- Verhoelst, T., Compernolle, S., Pinardi, G., Lambert, J. C., Eskes, H. J., Eichmann, K. U., Fjaeraa, A. M., Granville, J., Niemeijer, S., Cede,
 A., Tiefengraber, M., Hendrick, F., Pazmino, A., Bais, A., Bazureau, A., Boersma, K. F., Bognar, K., Dehn, A., Donner, S., Elokhov,
 A., Gebetsberger, M., Goutail, F., de la Mora, M. G., Gruzdev, A., Gratsea, M., Hansen, G. H., Irie, H., Jepsen, N., Kanaya, Y.,
 Karagkiozidis, D., Kivi, R., Kreher, K., Levelt, P. F., Liu, C., Muller, M., Comas, M. N., Piters, A. J. M., Pommereau, J. P., Portafaix,
 T., Prados-Roman, C., Puentedura, O., Querel, R., Remmers, J., Richter, A., Rimmer, J., Cardenas, C. R., de Miguel, L. S., Sinyakov,
 V. P., Stremme, W., Strong, K., Van Roozendael, M., Veefkind, J. P., Wagner, T., Wittrock, F., Gonzalez, M. Y., and Zehner, C.:
 Ground-based validation of the Copernicus Sentinel-5P TROPOMI NO2 measurements with the NDACC ZSL-DOAS, MAX-DOAS
- 850 Ground-based validation of the Copernicus Sentinel-5P TROPOMI NO2 measurements with the NDACC ZSL-DOAS, MAX-DOAS and Pandonia global networks, Atmos Meas Tech, 14, 481-510, 10.5194/amt-14-481-2021, 2021.
 - Verstraeten, W. W., Neu, J. L., Williams, J. E., Bowman, K. W., Worden, J. R., and Boersma, K. F.: Rapid increases in tropospheric ozone production and export from China, Nat Geosci, 8, 690-+, 10.1038/Ngeo2493, 2015.

- Visser, A. J., Boersma, K. F., Ganzeveld, L. N., and Krol, M. C.: European NOx emissions in WRF-Chem derived from OMI: impacts on summertime surface ozone, Atmos Chem Phys, 19, 11821-11841, 10.5194/acp-19-11821-2019, 2019.
 - Wang, C. J., Wang, T., Wang, P. C., and Rakitin, V.: Comparison and Validation of TROPOMI and OMI NO2 Observations over China, Atmosphere-Basel, 11, 10.3390/atmos11060636, 2020a.
- Wang, Y., Wang, J., Xu, X. G., Henze, D. K., Qu, Z., and Yang, K.: Inverse modeling of SO2 and NOx emissions over China using multisensor satellite data Part 1: Formulation and sensitivity analysis, Atmos Chem Phys, 20, 6631-6650, 10.5194/acp-20-6631-2020, 2020b.
- Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning, Geosci. Model Dev., 4, 625-641, 10.5194/gmd-4-625-2011, 2011.
- WOUDC: World Ozone and Ultraviolet Data Center OzoneSonde data [dataset], 10.14287/10000008, 2019.
- 865 Xing, J., Mathur, R., Pleim, J., Hogrefe, C., Gan, C. M., Wong, D. C., and Wei, C.: Can a coupled meteorology-chemistry model reproduce the historical trend in aerosol direct radiative effects over the Northern Hemisphere?, Atmos Chem Phys, 15, 9997-10018, 10.5194/acp-15-9997-2015, 2015.
- Xu, L., Pye, H. O. T., He, J., Chen, Y. L., Murphy, B. N., and Ng, N. L.: Experimental and model estimates of the contributions from biogenic monoterpenes and sesquiterpenes to secondary organic aerosol in the southeastern United States, Atmos Chem Phys, 18, 12613-12637, 10.5194/acp-18-12613-2018, 2018.
- Zhang, L., Jacob, D. J., Boersma, K. F., Jaffe, D. A., Olson, J. R., Bowman, K. W., Worden, J. R., Thompson, A. M., Avery, M. A., Cohen, R. C., Dibb, J. E., Flock, F. M., Fuelberg, H. E., Huey, L. G., McMillan, W. W., Singh, H. B., and Weinheimer, A. J.: Transpacific transport of ozone pollution and the effect of recent Asian emission increases on air quality in North America: an integrated analysis using satellite, aircraft, ozonesonde, and surface observations, Atmos Chem Phys, 8, 6117-6136, DOI 10.5194/acp-8-6117-2008, 2008.
- 875 Zhang, X., Jones, D. B. A., Keller, M., Walker, T. W., Jiang, Z., Henze, D. K., Worden, H. M., Bourassa, A. E., Degenstein, D. A., and Rochon, Y. J.: Quantifying Emissions of CO and NOX Using Observations From MOPITT, OMI, TES, and OSIRIS, J Geophys Res-Atmos, 124, 1170-1193, 10.1029/2018jd028670, 2019.
 - Zhang, Y. Q., Cooper, O. R., Gaudel, A., Thompson, A. M., Nedelec, P., Ogino, S. Y., and West, J. J.: Tropospheric ozone change from 1980 to 2010 dominated by equatorward redistribution of emissions, Nat Geosci, 9, 875-+, 10.1038/Ngeo2827, 2016.
- 2015, P. Natl Acad Sci USA, 115, 12401-12406, 10.1073/pnas.1812955115, 2018.
 2015, P. Natl Acad Sci USA, 115, 12401-12406, 10.1073/pnas.1812955115, 2018.
- Zhu, Y. Z., Hu, Q. H., Gao, M., Zhao, C., Zhang, C. X., Liu, T., Tian, Y., Yan, L., Su, W. J., Hong, X. H., and Liu, C.: Quantifying Contributions of Local Emissions and Regional Transport to NOX in Beijing Using TROPOMI Constrained WRF-Chem Simulation, Remote Sens-Basel, 13, 10.3390/rs13091798, 2021.